

1 Short Report

2 **Midbrain dopamine neurons compute inferred and**
3 **cached value prediction errors in a common framework**

4
5 Brian Sadacca ^{1*}, Joshua L. Jones ¹, and Geoffrey Schoenbaum ^{1,2,3*}

- 6 1. Intramural Research program of the National Institute on Drug Abuse, NIH;
7 2. Department of Anatomy and Neurobiology, University of Maryland School of
8 Medicine;
9 3. Department of Neuroscience, Johns Hopkins School of Medicine.

10

11 Words in abstract: 149

12 Words in body: 3223

13 Tables: 0

14 Figures: 3

15 Supplemental Files: 1 (3 figs)

16 Keywords: VTA; dopamine; prediction-error; TDRL; inference; single unit

17

18 * Corresponding authors:

19 Brian Sadacca (brian.sadacca@nih.gov)

20 Geoffrey Schoenbaum (geoffrey.schoenbaum@nih.gov).

21

22 Abstract

23 Midbrain dopamine neurons have been proposed to signal reward prediction errors as
24 defined in temporal difference (TD) learning algorithms. While these models have been
25 extremely powerful in interpreting dopamine activity, they typically do not use value derived
26 through inference in computing errors. This is important because much real world behavior
27 – and thus many opportunities for error-driven learning – is based on such predictions. Here,
28 we show that error-signaling rat dopamine neurons respond to the inferred, model-based
29 value of cues that have not been paired with reward and do so in the same framework as
30 they track the putative cached value of cues previously paired with reward. This suggests
31 that dopamine neurons access a wider variety of information than contemplated by standard
32 TD models and that, while their firing conforms to predictions of TD models in some cases,
33 they may not be restricted to signaling errors from TD predictions.

34 Impact Statement: Midbrain dopamine neurons have been repeatedly shown to signal
35 reward prediction errors based on direct experience. However much real world behavior -
36 and thus many of our opportunities for error-driven learning - is based on inference. Here,
37 we show that the firing of dopamine neurons reflects inferred and directly experience value
38 similarly.

39 Introduction

40 Midbrain dopamine neurons have been proposed to signal the reward prediction errors
41 defined in temporal difference (TD) learning algorithms (1, 2). This proposal was initially
42 based on observations that these neurons fired more strongly to unpredicted than to
43 predicted reward, suppressed firing on omission of a predicted reward, and developed firing
44 to reward-paired cues with learning (3). Further work has shown that phasic activity in
45 dopamine neurons obeys formal predictions for such TD error signals under more complex
46 conditions (4-10), including in tasks such as blocking and conditioned inhibition, in which
47 experimental conditions are arranged to precisely distinguish between prediction error
48 signals and other possible explanations of such activity. These studies have confirmed that
49 the neural correlates correspond closely to the theoretical accounts. Indeed careful work in
50 monkeys has shown that the activity provides a quantitative match with the error signal
51 described in the TD model (9). With the advent of optogenetic techniques and Cre-driver
52 lines in rats, it has also been shown in rats that artificially stimulating or inhibiting likely
53 dopamine neurons for very brief periods is sufficient to restore the associative learning
54 driven by endogenous positive or negative prediction errors (11, 12), suggesting that phasic
55 activity in dopamine neurons can act like a prediction error, at least in some downstream
56 targets and behavioral paradigms (13, 14).

57 But is it a TD prediction error? The errors in TD models, at least as they have been
58 applied to interpret the firing of dopamine neurons, are based on the value that has been
59 assigned to events based on direct experience (13-16). This so-called *cached* value is pre-
60 computed and resides in the predictive event, be it a primary reward or reward-predicting
61 cue. The value is calculated free of any other predictive information about the environment,

62 defining this class of algorithms as *model-free*. While these TD models have been
63 extremely powerful in interpreting the activity of dopamine neurons in tasks in which value is
64 based on experienced reward, they are unable to compute prediction errors elicited in
65 situations in which value is derived through inference rather than through direct experience.
66 Deriving a value (or a prediction) based on inference – meaning to deduce from an
67 understanding of the relationships amongst environmental stimuli and events - is a hallmark
68 of a second class of algorithms, termed *model-based* (17).

69 This distinction between algorithms is important because much of our real world
70 behavior, and thus many of our opportunities for error-driven learning, is based on this
71 model-based inference (17, 18). Rarely does one engage in value-based behavior that is
72 simply a pure repetition of prior learning; typically moderating or mitigating information
73 acquired in other situations, separate from the original learning, influences our decision-
74 making. This basic concept is operationalized (and the effect of such *inferred* or *model-*
75 *based* value, isolated) in sensory pre-conditioning (19). In this task, animals first learn that
76 one innocuous cue (cue A) predicts another (cue B), in the absence of reward, and then
77 later learn that the second cue (B) is a reliable predictor of reward. Cue A has not been
78 directly paired with reward and thus it has not had any opportunity to acquire any cached
79 value (13, 15). As a result, cue A cannot elicit a TD prediction error, despite the fact that it
80 has value as defined by the animal's responding and its ability to modulate error-driven
81 learning (20). If TD models are an accurate and complete description of the information
82 contained in dopamine neuron activity, then cue A should not elicit dopamine neuron firing,
83 at least not above the level of a control cue.

84 **Results**

85 To test this, we trained 14 rats with recording electrodes implanted in the ventral
86 tegmental area (VTA) in a sensory-preconditioning task. In the first phase, rats learned to
87 associate two pairs of environmental cues (A->B; C->D) in the absence of reward. As there
88 was no reward, rats showed no significant responding at the food cup and no differences in
89 responding during the different cues (ANOVA, $F_{3, 55} = .7$, $p = 0.52$; Figure 1A). In the
90 second phase, rats learned that the second cue of one pair (B) predicted reward and the
91 other (D) did not; learning was reflected in an increase in responding at the food cup during
92 presentation of B (ANOVA, main effect of cue: $F_{1, 163} = 280.1$, $p < 0.001$, main effect of
93 session: $F_{5, 163} = 9.7$, interaction: $F_{5, 163} = 10.81$, $p < 0.001$; Figure 1B). Finally, in the third
94 phase, the rats were presented again with the four cues, first a reminder of cue B and D's
95 reward contingency followed by an unrewarded probe test of responding to cues A and C.
96 As expected, the rats responded at the food cup significantly more during presentation of A,
97 the cue that predicted B, than during presentation of C, the cue that predicted D (ANOVA,
98 main effect of cue: $F_{1, 167} = 8.7$, $p < 0.001$, main effect of trial: $F_{5, 167} = 6.08$, $p < 0.001$,
99 interaction: $F_{5, 167} = 2.07$, $p = 0.07$; Figure 1C).

100 Single unit activity was recorded in the VTA throughout training. To identify putative
101 dopamine neurons, we used a recently developed, optogenetically-validated strategy that
102 classifies VTA neurons on the basis of their response dynamics during Pavlovian
103 conditioning. In published work (21, 22), this strategy identified VTA dopamine neurons (i.e.
104 neurons expressing Cre under the control of the promoter for the dopamine transporter) with
105 near perfect fidelity. Here we applied this same analysis to the mean normalized responses
106 of all VTA neurons recorded during conditioning and reminder sessions ($n = 632$; Figure 2A).
107 We extracted the major modes of variation among the neurons with principal components
108 analysis (PCA; Figure 2B) and then performed hierarchical clustering on those PCs (Figure

109 2C). This analysis successfully extracted the 3 previously described VTA response types
110 from our data (Figure 2D): neurons with sustained excitation to cues and reward (putative
111 GABAergic), neurons with phasic excitation to cue onset and reward onset (putative
112 dopaminergic), and neurons with sustained inhibition to cue and reward (unknown). We
113 then assessed the responses of the putative dopamine neurons ($n = 304$) to the cue and
114 reward over the course of conditioning. Consistent with their classification, we found
115 changes in firing during conditioning that were in accord with signaling of reward prediction
116 errors. Specifically, early in conditioning, these neurons' maximal response occurred just
117 after reward delivery (Figure 2E, top trace), whereas late in conditioning, the maximal
118 response occurred just after onset of the cue predicting that reward (Figure 2E, lower
119 traces). As a result, the difference in activity at the time of cue onset versus reward
120 increased significantly from the start to the end of conditioning ($r_{302} = 0.24$, $p < 0.01$; Figure
121 2F), consistent with signaling of TD prediction errors (13, 14).

122 Having established that putative dopamine neurons identified in this manner exhibit
123 firing during conditioning consistent with signaling of TD errors, we next examined activity in
124 neurons recorded in just the probe test. We again identified these neurons by their pattern
125 of firing to the reward predictive cue ($n = 102$; Figure 3A-D). As before, this analysis
126 identified a group of cells with strong phasic responses to B, the cue that had been directly
127 paired with reward ($n = 52$). **While this response generalized somewhat to D, the**
128 **control cue that had been presented without reward during conditioning sessions,**
129 **these neurons fired significantly more during the first second of B than to D ($t_{51} =$**
130 **4.40, $p < 0.001$, black versus gray lines, respectively, with shading for SEM, Figure 3E).**

131 However, in addition to this expected pattern of firing, these cells also had strong
132 phasic responses to both preconditioned cues (blue and red traces, A and C, respectively,

133 with shading for SEM, Figure 3E). While the common element of these responses could
134 reflect novelty or salience, since these cues had not been presented for a number of days,
135 or perhaps generalization from conditioning to B, the actual phasic neural response was
136 significantly stronger for A, the cue that predicted the reward-paired cue, than for C, the
137 preconditioned control cue ($t_{51} = 5.02$, $p < 0.001$). This difference cannot be explained on
138 the basis of novelty, salience, or generalization, since A and C were treated similarly. Nor
139 can it be explained by direct experience with reward, because A was never paired with
140 reward, and it was only paired with B before conditioning. Thus, the phasic response in
141 these putative dopamine neurons appeared to be influenced by inferred value of cue A.
142 **Interestingly, the neural response was perhaps somewhat better at discriminating A**
143 **from C (Figure 3E, bottom panel) than B from D (Figure 3E, top panel), perhaps**
144 **reflecting the differences in training between A and C, which were only presented a**
145 **few times in unrewarded sessions, versus B and D, which were presented many times**
146 **across several days of conditioning. Despite this**, the influence of the inferred value of
147 cue A on the firing of these neurons during the first second of their cue response was
148 strongly and significantly correlated with the influence of value on these neurons firing at the
149 onset of B, the cue directly paired with reward ($r_{50} = 0.63$, $p < 0.001$; Figure 3F). Notably
150 this was also true for a handful of neurons ($n = 4$) that exhibited the classic wide, polyphasic
151 waveforms traditionally used to identify dopamine neurons (Figure 3F, filled circles, see
152 Figure 3-figure supplement 1 for PSTH's). This relationship in the initial phasic response to
153 the cues did not reach significance in the other two neural subtypes identified by the
154 clustering analysis (see Figure 3-figure supplements 2 and 3 for analyses of tonically-
155 modulated neurons).

156 Beyond their phasic responses at the start of the cues and to reward, the putative
157 dopamine neurons also exhibited another notable feature: the average response of these
158 neurons throughout cues A and C was above baseline, and this sustained firing was
159 significantly higher to cue A than C (Figure 3E, final 9s of cues, $t_{51} = 2.56$, $p < 0.05$). This
160 elevated firing may be a sign of dopamine's reported ability to anticipate proximity to reward
161 or to signal state value (23, 24), if our rats' expectation of reward delivery is based on
162 knowing that progression to the offset of cue A should lead to the subsequent presentation
163 of cue B and then reward. Importantly, in our design, reward is presented during B rather
164 than at its termination. This would explain why this pattern of sustained firing is not present
165 throughout cue B (Figure 3E, $t_{51} = 1.09$, $p = 0.278$). Interestingly this pattern of sustained and
166 differential firing to A (vs C) and not to B (vs D) in the putative dopamine neurons is the
167 mirror image of firing in neurons classified as tonically excited, which showed relatively
168 modest changes in sustained firing to A and C and much larger increases in firing to B (see
169 Figure 3-figure supplement 2). This relationship would be consistent with recent proposals
170 that these neurons, thought to be GABAergic (21), exert tonic inhibition to suppress the
171 firing of dopamine neurons (22).

172 **Discussion**

173 Here we report that VTA dopamine neurons, identified based either on traditional
174 waveform criteria or through an optogenetically-validated clustering analysis of their
175 response properties to a conditioned cue, exhibited phasic cue-evoked responses that were
176 influenced by inferred value. These responses were observed even though the critical cue
177 had no prior history of direct pairing with any rewarding event. In addition, they were greater
178 than responses to a control cue that was treated similarly and thus had similar levels of
179 salience or novelty or generalized value, all variables that have been proposed to explain
180 phasic activity in other settings that appeared to be at odds with the standard explanation of
181 phasic dopamine activity (25-27). These data show that the phasic activity of dopamine
182 neurons can reflect information about value that is not contemplated by TD models, at least
183 as they have been applied to understand the phasic firing of these neurons (13-16).

184 Our finding is consistent with a number of recent reports, suggesting that dopamine
185 neurons are likely to access more complex information than is available to standard TD
186 models. For example, dopamine neurons in the rat VTA utilize input from the orbitofrontal
187 cortex to disambiguate states that are not easily distinguished via external information in
188 order to more accurately calculate prediction errors (28). While this result does not require
189 the use of inference in calculating errors, merely access to state information, it suggests that
190 dopamine neurons have access to a major source of this information, given the central role
191 of the orbitofrontal cortex in inference-based behavior (29).

192 The phasic activity of dopamine neurons has also been shown to track the value of
193 one cue after changes in the value of an associated cue (30). Again these data suggest that
194 dopamine neurons have access to higher order information, which could be described as

195 inference. Indeed these authors describe their results in terms of inference; however, as
196 they note in their discussion (30, *final paragraph of discussion*), the inference seen in their
197 task may differ from that shown here in that it does not require access to model-based
198 information, but could instead be based on direct, 'cached' value from earlier training
199 sessions.

200 Finally elevated dopamine has also been found using microdialysis during an
201 aversive version of the sensory preconditioning task used here (31). However the use of an
202 aversive paradigm, a measurement technique with low temporal resolution, and the lack of
203 control conditions to confirm signaling of cached value errors make it difficult to apply these
204 results to address the very specific proposal that phasic changes in the firing of dopamine
205 neurons signal TD prediction errors in appetitive paradigms.

206 Our study addresses the limitations of these best available prior reports. We are
207 recording phasic activity of dopamine neurons at their source. We have identified dopamine
208 neurons by two different classification schemes, an old one that has been used repeatedly
209 across labs and species to identify error-signaling dopamine neurons (3-5, 7, 32-35), as well
210 as a new, optogenetically-validated approach that has identified error signals in mice (21)
211 and is favored by those that dislike the use of waveform criteria (36). We used a carefully
212 designed behavioral preparation in which our critical cue of interest has no prior history of
213 association with reward, thus unique firing to this cue cannot be explained as any sort of
214 cached value (see '*A comment on the basis of responding to the preconditioned cue*' in the
215 *behavioral Methods for further information*). Further this appetitive task includes two
216 important control cues: one designed to rule out explanations based on generalization and
217 salience (cue C) and another designed to reveal cached value prediction errors (cue B).
218 The inclusion of this cue in particular is important because it allows us to assess the

219 relationship between traditional error signals and any influence of inferred value on the firing
220 of the dopamine neurons.

221 The close relationship in the firing of the dopamine neurons to B, the cue directly
222 paired with reward, and A, the cue that predicts reward only through B, suggests that
223 whatever is ultimately signaled when a cue with inferred, model based value is presented
224 may be similar to what the same neurons signal in response to the unexpected appearance
225 of a cue that has been directly paired with reward. While this might be explained as error
226 signaling to B, calculated from TD models, and error signaling to A, calculated from
227 something beyond TD models, this solution is cumbersome, particularly given that inferred
228 and experienced value are actually confounded for a cue directly paired with reward (a fact
229 illustrated by the normal efficacy of reinforcer devaluation at changing conditioned
230 responding (37)). A more parsimonious explanation is that dopamine neurons, unlike
231 standard TD models, have access to a wide variety of information when computing expected
232 value. And that while their firing may conform to what is expected for errors calculated from
233 TD models in some special cases, they may not be signaling TD derived errors. Such a
234 suggestion aligns nicely with recent proposals that dopamine neurons signal errors based
235 on changes in economic utility (38), and it would be consistent with data presented in
236 abstract form suggesting that cue-evoked dopamine release in nucleus accumbens is
237 sensitive to devaluation of a paired reward (39), though it contradicts data only just
238 published from a similar study in which cue-evoked release was not immediately altered
239 when reward value was manipulated via salt depletion (40). This variability in
240 correspondence between our unit data and evidence from studies of dopamine efflux in
241 accumbens may reflect to the different dynamics of the two processes or it may indicate

242 some specificity with regard to the information content of the dopaminergic afferents in
243 accumbens versus other areas.

244 Finally it is important to explicitly note that the general proposal that phasic changes
245 in dopamine are a TD error signal incorporates two very separate sets of predictions. One
246 set, most relevant to the single unit correlates that form the basis of this hypothesis,
247 concerns the information used to construct the error signals. This is obviously the part of
248 the question we have addressed in the current study. That is, do dopaminergic errors reflect
249 only model-free information derived from TD systems or do they also incorporate the
250 predictions of non-TD, model-based systems? We believe our data favor the latter position.

251 The second set of predictions, not addressed by our study, concerns what the
252 dopaminergic errors do downstream. Do they act only to stamp in the so-called cached
253 values that are acquired through learning in TD models or do they act more broadly to
254 facilitate increases in the strength of associative representations in a way that is orthogonal
255 to distinctions between the systems, model-free or model based, in which those
256 representations reside? The latter role would be more in accord with earlier learning theory
257 accounts that viewed prediction errors as acting on the strength of associative
258 representations (13, 41, 42). Importantly the answer to the second question is formally
259 separate from the answer to the first. In other words, phasic changes in dopamine may
260 reflect model-based information and yet only act to support model-free, cached-value
261 learning. Or phasic changes in dopamine could act more broadly, supporting both model-
262 free and model-based learning, even if they only reflected value predictions from the former
263 system.

264 Notably the jury remains out on what sort of learning the brief phasic changes in
265 dopamine thought to signal prediction errors serve to support. In support of dopamine's role
266 in supporting model-based learning, prediction errors observed in ventral striatal target
267 regions seem to reflect both model based and model free information (43). Further studies
268 have shown that elevated dopamine levels, either observed (44) or directly manipulated (45,
269 46), bias subjects towards making model-based decisions, as do changes to dopaminergic
270 gene expression (47). While suggestive, these studies do not directly distinguish the effects
271 of phasic changes in dopamine neuron firing or release from the effects of slower tonic
272 changes. Such tonic changes may play a very different role from the phasic error signals
273 observed in single unit activity (24, 48). Further these studies do not isolate the effects of
274 errors themselves, independent from other confounding variables. Such isolation and
275 specificity can be achieved using Cre-driver lines in rodent species, and as noted earlier,
276 there is now strong evidence from such studies that brief, phasic changes in dopamine
277 neuron firing can act like a prediction error, at least in some downstream targets and
278 behavioral paradigms (11, 12). However the behavior supported by the artificially induced
279 prediction errors in these experiments may be either model-based or model-free (or a
280 mixture). A more definitive answer to this question will require these approaches to be
281 married to paradigms that distinguish these two types of learning.

282

283 **Materials and Methods**

284 **Subjects:** 14 adult Long-Evans rats (10 male, 4 female weighing 275–325 g on arrival) were
285 individually housed and given ad libitum access to food and water, except during behavioral
286 training and testing, which which they received 15 minutes of ad-lib water access following
287 each training session. Rats were maintained on a 12-h light/dark cycle and trained and
288 tested during the light cycle. Experiments were performed at the National Institute on Drug
289 Abuse Intramural Research Program, in accordance with NIH guidelines. The number of
290 subjects was chosen to have sufficient power to assess learning on the final test-day (18),
291 and to gather a sufficient number of isolated neurons (>100) for subsequent analysis on the
292 final test day.

293 **Apparatus:** Behavioral training and testing were conducted in standard behavior boxes
294 with commercially-available equipment (Coulbourn Instruments, Allentown, PA). A recessed
295 dipper was placed in the center of the right wall approximately 2 cm above the floor. The
296 dipper was mounted outside the behavior chamber and delivered 40 ul of flavored milk
297 (Nestle) per dipper elevation. Auditory cues (tone, siren, 2 Hz clicker, white noise) calibrated
298 to ~65 dB were used during the behavioral testing.

299 **Surgical procedures:** Rats underwent surgery for implantation of chronic recording
300 electrode arrays. Rats were anesthetized with isoflurane and placed in a standard
301 stereotaxic device. The scalp was excised, and holes were bored in the skull for the
302 insertion of ground screws and electrodes. Multi-electrode bundles [16 nichrome microwires
303 attached to a microdrive] were inserted 0.5 mm above dorsal VTA [anteroposterior (AP) 5.4
304 mm and mediolateral (ML) 0.8 mm relative to bregma (Paxinos and Watson, 1998); and
305 dorsoventral (DV) 7.0 mm from dura]. In 3 rats, microwire electrodes were also implanted

306 0.5 mm above ipsilateral orbitofrontal cortex [AP 3.2 mm and ML 3.0 mm relative to bregma
307 (Paxinos and Watson, 1998); and DV 4.0 mm from the dura], and in 2 other rats, microwire
308 electrodes were also implanted 0.5 mm above ipsilateral ventral striatum [AP 1.0 mm and
309 ML 3.0 mm relative to bregma (Paxinos and Watson, 1998); and DV 6.0 mm from the dura].
310 Once in place, the assemblies were cemented to the skull using dental acrylic. Six rats also
311 received infusions of 1.0ul of AAV5-DIO-HMD4 into central VTA [anteroposterior (AP) 5.4
312 mm and mediolateral (ML) 0.8 mm relative to bregma, and 8.1 mm below dura]; there were
313 no effects of this treatment on any of the results we have reported.

314 **Behavioral Training:** Rats began sensory preconditioning 2 weeks after electrode
315 implantation. The sensory preconditioning procedure consisted of three phases, of similar
316 design to a prior study (20).

317 *Preconditioning:* Rats were shaped to retrieve a liquid reward from a fluid dipper over three
318 sessions; each session consisted of twenty deliveries of 40 ul of flavored milk. After this
319 shaping, rats underwent 2 days of preconditioning. Each day of preconditioning, rats
320 received twelve trials in which two pairs of auditory cues (A->B and C->D) were presented
321 sequentially, with no delay between cues, six times each, in a blocked design. Cues were
322 each 10s long, the inter-trial intervals varied from 3 to 6 min, and the order the blocks
323 alternated across days. Cues A and C were white noise or clicker (counterbalanced), and
324 cues B and D were siren or tone, (counterbalanced).

325 *Conditioning:* After preconditioning, rats underwent conditioning. Each day, rats received a
326 single training session, consisting of six trials of cue B paired with the flavored milk reward
327 and six trials of D paired with no reward. The flavored milk reward was presented three
328 times via the dipper in the food cup at 1, 4, and 7s into the 10s presentation of cue B. Cue

329 D was presented for 10s without reward. The two cues were presented in 3-trial blocks,
330 counterbalanced. The inter-trial intervals varied between 3 and 6 min. Ten rats were given
331 6 days of conditioning, while two were advanced to the probe test after 5 days due the
332 presence of putative wide waveform neurons at the beginning of the sixth conditioning day.
333 There was no difference between these groups in the final test.

334 *Probe test:* After conditioning, the rats underwent a single probe test, which consisted of
335 three reminder trials of B paired with reward and three trials of D unpaired interleaved,
336 followed by presentation of cues A and C, alone, six times each, without reward, with the
337 presentation order counterbalanced across animals. In six animals A and C trials were
338 blocked, while in 8 animals they were interleaved; both groups showed all reported effects
339 and were merged. Cue durations, timing of reward, and inter-trial intervals were as above.

340 *A comment on the basis of responding to the preconditioned cue:* We have interpreted our
341 sensory preconditioning effect in terms of an associative chaining or value inference
342 mechanism. An alternative account, which has been employed in other recent studies using
343 similar procedures (49, 50), is that the conditioned responding to cue A results from
344 mediated learning that occurs during the second phase of the experimental procedure (51).
345 Briefly, this account suggests that following the initial pairings of A and B, subsequent
346 presentations of B for conditioning activate a representation of A in memory within a
347 relatively close temporal contiguity with the delivery of sucrose, resulting in the
348 representation of A becoming directly associated with this reward. If this were to occur, then
349 at test, the subsequent conditioned responding to A might reflect the cue's direct association
350 with sucrose, rather than requiring B to bridge the experiences of A and sucrose.

351 While there is significant evidence within the literature for the phenomenon of mediated
352 learning (reviewed in 52, 53), several features of our behavioral design were chosen to bias
353 strongly against the operation of this mechanism.

354 First, we used forward ($A \rightarrow B$) rather than simultaneous (AB) or backward pairings ($B \rightarrow A$) of
355 the pre-conditioned and conditioned cues. This is important because mediated learning in
356 rodents has been suggested to operate primarily when A and B are presented
357 simultaneously (51) or as the serial compound $B \rightarrow A$ (i.e. backward sensory preconditioning;
358 52). The reason for this is intuitive because either of these temporal arrangements
359 maximizes the chances that B will evoke a representation of A during the conditioning phase
360 and concurrent with reward delivery, an arrangement that obvious benefits in maximizing the
361 ability of an evoked representation of A to become directly associated with reward.

362 Our design avoids this by initially presenting A and B as the serial compound $A \rightarrow B$, an
363 arrangement which, as far as B is concerned, parallels a backward conditioning procedure.
364 Backward cue-outcome pairings have generally been shown to yield weak, if any, excitatory
365 conditioning (54). For this reason, it is widely accepted that $A \rightarrow B$ pairings will render B
366 relatively ineffective at subsequently conjuring up a memory of A, thus making the
367 contribution of mediated learning insubstantial (55).

368 Second, the amount of training given in Phase 2 of conditioning, with B-reward pairings, was
369 also designed to discourage mediated learning. As noted above, the presentation of B in
370 conditioning should activate a representation of A in memory. However, with repeated
371 presentations of B without A, the representation of A evoked by B will also extinguish. For
372 this purpose, we present 3 times as many B (not A) trials in the conditioning phase as we

373 present A and B pairings, further undermining the likelihood that an evoked A representation
374 will be maintained.

375 In conclusion, we believe our implementation of these specific behavioral parameters should
376 largely eliminate any potential contribution of mediated learning to the sensory
377 preconditioning effect in our particular design, and favor the parsimonious interpretation of
378 the sensory preconditioning effect in terms of an associative chaining or value inference
379 mechanism. We would note that this interpretation is supported by our own prior report that
380 OFC inactivation at probe test in this exact paradigm abolishes responding to A and has no
381 effect on responding to B (20), since mediated learning is basically simple conditioning and
382 OFC manipulations typically have no effect on expression of previously acquired conditioned
383 responding (56, 57).

384 **Electrophysiology:** Neural signals were collected from the VTA during each behavioral
385 session. Differential recordings were fed into a parallel processor capable of digitizing 16-to-
386 32 signals at 40 kHz simultaneously (Plexon). Discriminable action potentials of <3:1
387 signal/noise ratio were isolated on-line from each signal using an amplitude criterion in
388 cooperation with a template algorithm. Discriminations were checked continuously
389 throughout each session. Time-stamped records of stimulus onset and neuronal spikes
390 were saved digitally, as were all sampled spike waveforms and the discrimination file. Off-
391 line re-analysis incorporating 3D cluster-cutting techniques confirmed and corrected on-line
392 discriminations. Except where explicitly noted, all neurons identified via off-line sorting were
393 included in each analysis.

394 **Statistical analyses:** Raw data were processed with Matlab to extract food cup entries and
395 spike-timing relative to cue-onset. Entries were converted to a response measure: the

396 percentage of time rats spent with their head in the food cup during cue presentation as
397 measured by an infrared photo beam positioned at the front of the food cup. Spike times
398 were binned and analyzed as specified below. In comparing cue-evoked to reward-evoked
399 activity, bins spanning the first 500ms of each period were analyzed. In comparing
400 response differences evoked by different cues, bins spanning the first 1s of cue-evoked
401 activity were analyzed. For all statistical tests, an alpha level of 0.05 was used.

402 **AUC calculation:** As per prior reports (21), we normalized the firing rate of individual
403 neurons by comparing the histogram of spike counts during each bin of spiking activity
404 (100ms, test bins from each trial for a cue, at a particular time post-stimulus) against a
405 histogram of baseline (100ms) bins, from all trials for that cue. The ROC in question is
406 calculated by (1) normalizing all test and baseline bin counts, such that the minimum bin
407 count was 0 and the maximal bin count was 1, (2) sliding a discrimination threshold across
408 each histogram of bins, from 0 to 1 in .01 steps, such that fraction of test bins identified
409 above the threshold was a 'true positive' rate and the fraction of baseline bins above the
410 threshold was a 'false negative' rate for an ROC curve. The area under this curve was then
411 estimated by trapezoidal numerical estimation, with an auROC below .5 being indicative of
412 inhibition, and an auROC above .5 being indicative of excitation above baseline.

413 **Classification of dopamine neurons by response dynamics:** In order to isolate VTA
414 neuron response types shown to be indicative of putative dopaminergic and GABAergic
415 genetic identities (21), we took the auROC normalized responses of neurons during their
416 response to the cue predictive of reward (cue B), and performed a simple classification to
417 separate neural responses. We first performed principal components analysis on a matrix of
418 neural responses during cue B and reward presentation (neuron-by-time) to simplify the
419 neural dynamics to the 3 most descriptive ways in which neurons differed. We then

420 classified this description of the neural population (first 3 principal components) with a
421 simple unsupervised hierarchical clustering algorithm, finding the similarity (Euclidean
422 distance) between all pairs of neurons in principal components space, and iteratively
423 grouping the neurons them into larger and larger clusters on the basis of their similarity (i.e.
424 agglomerative complete-linkage clustering). A distance-criterion was then set to extract
425 exactly 3 clusters from this hierarchical tree.

426 **Classification of dopamine neurons by waveform:** Neurons were screened for wide
427 waveform and amplitude characteristics, calculated on their mean action-potential across a
428 recording session. Neurons were identified as dopaminergic if their negative half-width
429 exceeded a standard criterion (450 μ s) and the ratio of (max) positive to (min) negative
430 voltage deflections was greater than zero (3, 28, 32, 33). Four such neurons were identified
431 as wide waveform across the probe sessions.

432 **Histology:** After the final recording session, rats were euthanized and perfused first with
433 PBS and then 4% formalin in PBS. Electrolytic lesions (1 mA for 10 s) made just before
434 perfusion were examined in fixed, 0.05 mm coronal slices stained with cresyl violet.
435 Anatomical localization for each recording was verified on the basis of histology, stereotaxic
436 coordinates of initial positioning, and recording notes.

437 **Acknowledgments**

438 This work was supported by the Intramural Research Program at the National Institute on
439 Drug Abuse. The opinions expressed in this article are the authors' own and do not reflect
440 the view of the NIH/DHHS.

441

442 **Financial Disclosures**

443 The authors declare no competing financial interests.

444 **References**

- 445 1. W. Schultz, P. Dayan, P. R. Montague, A neural substrate for prediction and reward.
446 *Science* **275**, 1593-1599 (1997).
- 447 2. R. S. Sutton, Learning to predict by the method of temporal difference. *Machine*
448 *Learning* **3**, 9-44 (1988).
- 449 3. J. Mirenowicz, W. Schultz, Importance of unpredictability for reward responses in
450 primate dopamine neurons. *Journal of Neurophysiology* **72**, 1024-1027 (1994).
- 451 4. P. Waelti, A. Dickinson, W. Schultz, Dopamine responses comply with basic
452 assumptions of formal learning theory. *Nature* **412**, 43-48 (2001).
- 453 5. P. N. Tobler, A. Dickinson, W. Schultz, Coding of predicted reward omission by
454 dopamine neurons in a conditioned inhibition paradigm. *Journal of Neuroscience* **23**,
455 10402-10410 (2003).
- 456 6. A. Lak, W. R. Stauffer, W. Schultz, Dopamine prediction error responses integrate
457 subjective value from different reward dimensions. *Proceedings of the National*
458 *Academy of Science* **111**, 2342-2348 (2014).
- 459 7. W.-X. Pan, R. Schmidt, J. R. Wickens, B. I. Hyland, Dopamine cells respond to
460 predicted events during classical conditioning: evidence for eligibility traces in the
461 reward-learning network. *Journal of Neuroscience* **25**, 6235-6242 (2005).
- 462 8. A. S. Hart, R. B. Rutledge, P. W. Glimcher, P. E. Phillips, Phasic dopamine release in
463 the rat nucleus accumbens symmetrically encodes a reward prediction error term.
464 *Journal of Neuroscience* **34**, 698-704 (2014).
- 465 9. H. M. Bayer, P. Glimcher, Midbrain dopamine neurons encode a quantitative reward
466 prediction error signal. *Neuron* **47**, 129-141 (2005).
- 467 10. J. R. Hollerman, W. Schultz, Dopamine neurons report an error in the temporal
468 prediction of reward during learning. *Nature Neuroscience* **1**, 304-309 (1998).
- 469 11. E. E. Steinberg *et al.*, A causal link between prediction errors, dopamine neurons and
470 learning. *Nature Neuroscience* **16**, 966-973 (2013).
- 471 12. C. Y. Chang *et al.*, Brief optogenetic inhibition of VTA dopamine neurons mimics the
472 effects of endogenous negative prediction errors during Pavlovian over-expectation.
473 *Nature Neuroscience* **19**, 111-116 (2016).
- 474 13. P. W. Glimcher, Understanding dopamine and reinforcement learning: The dopamine
475 reward prediction error hypothesis. *Proceedings of the National Academy of Science*
476 **108**, 15647-15654 (2011).
- 477 14. W. Schultz, Getting Formal with Dopamine and Reward. *Neuron* **36**, 241-263 (2002).
- 478 15. Y. Niv, G. Schoenbaum, Dialogues on prediction errors. *Trends in Cognitive Sciences*
479 **12**, 265-272 (2008).
- 480 16. J. J. Clark, N. G. Hollon, P. E. M. Phillips, Pavlovian valuation systems in learning
481 and decision making. *Current Opinion in Neurobiology* **22**, 1054-1061 (2012).
- 482 17. N. D. Daw, Y. Niv, P. Dayan, Uncertainty-based competition between prefrontal and
483 dorsolateral striatal systems for behavioral control. *Nature Neuroscience* **8**, 1704-
484 1711 (2005).
- 485 18. B. B. Doll, D. A. Simon, N. D. Daw, The ubiquity of model-based reinforcer learning.
486 *Current Opinion in Neurobiology* **22**, 1075-1081 (2012).
- 487 19. W. J. Brogden, Sensory pre-conditioning. *Journal of Experimental Psychology* **25**,
488 323-332 (1939).

- 489 20. J. L. Jones *et al.*, Orbitofrontal cortex supports behavior and learning using inferred
490 but not cached values. *Science* **338**, 953-956 (2012).
- 491 21. J. Y. Cohen, S. Haesler, L. Vong, B. B. Lowell, N. Uchida, Neuron-type-specific
492 signals for reward and punishment in the ventral tegmental area. *Nature* **482**, 85-88
493 (2012).
- 494 22. N. Eshel *et al.*, Arithmetic and local circuitry underlying dopamine prediction errors.
495 *Nature* **525**, 243-246 (2015).
- 496 23. M. W. Howe, P. L. Tierney, S. G. Sandberg, P. E. Phillips, A. M. Graybiel, Prolonged
497 dopamine signalling in striatum signals proximity and value of distant rewards. *Nature*
498 **500**, 575-579 (2013).
- 499 24. A. A. Hamid *et al.*, Mesolimbic dopamine signals the value of work. *Nature*
500 *Neuroscience* **19**, 117-126 (2016).
- 501 25. S. Kakade, P. Dayan, Dopamine: generalization and bonuses. *Neural Networks* **15**,
502 549-559 (2002).
- 503 26. E. S. Bromberg-Martin, M. Matsumoto, O. Hikosaka, Dopamine in motivational
504 control: rewarding, aversive and alerting. *Neuron* **68**, 815-834 (2010).
- 505 27. M. Matsumoto, O. Hikosaka, Two types of dopamine neuron distinctly convey positive
506 and negative motivational signals. *Nature* **459**, 837-841 (2009).
- 507 28. Y. K. Takahashi *et al.*, Expectancy-related changes in firing of dopamine neurons
508 depend on orbitofrontal cortex. *Nature Neuroscience* **14**, 1590-1597 (2011).
- 509 29. T. A. Stalnaker, N. K. Cooch, G. Schoenbaum, What the orbitofrontal cortex does not
510 do. *Nature Neuroscience* **18**, 620-627 (2015).
- 511 30. E. S. Bromberg-Martin, M. Matsumoto, S. Hong, O. Hikosaka, A pallidus-habenula-
512 dopamine pathway signals inferred stimulus values. *Journal of Neurophysiology* **104**,
513 1068-1076 (2010).
- 514 31. A. M. J. Young, R. G. Ahier, R. L. Upton, M. H. Joseph, J. A. Gray, Increased
515 extracellular dopamine in the nucleus accumbens of the rat during associative
516 learning of neutral stimuli. *Neuroscience* **83**, 1175-1183 (1998).
- 517 32. M. R. Roesch, D. J. Calu, G. Schoenbaum, Dopamine neurons encode the better
518 option in rats deciding between differently delayed or sized rewards. *Nature*
519 *Neuroscience* **10**, 1615-1624 (2007).
- 520 33. Y. S. Jo, J. Lee, S. J. Mizumori, Effects of prefrontal cortical inactivation on neural
521 activity in the ventral tegmental area. *Journal of Neuroscience* **33**, 8159-8171 (2013).
- 522 34. G. Morris, A. Nevet, D. Arkadir, E. Vaadia, H. Bergman, Midbrain dopamine neurons
523 encode decisions for future action. *Nature Neuroscience* **9**, 1057-1063 (2006).
- 524 35. J. Xin, R. M. Costa, Start/stop signals emerge in nigrostriatal circuits during sequence
525 learning. *Nature* **466**, 457-462 (2010).
- 526 36. E. B. Margolis, H. Lock, G. O. Hjelmstad, H. L. Fields, The ventral tegmental area
527 revisited: Is there an electrophysiological marker for dopaminergic neurons? *Journal*
528 *of Physiology* **577**, 907-924 (2006).
- 529 37. P. C. Holland, J. J. Straub, Differential effects of two ways of devaluing the
530 unconditioned stimulus after Pavlovian appetitive conditioning. *Journal of*
531 *Experimental Psychology: Animal Behavior Processes* **5**, 65-78 (1979).
- 532 38. W. Schultz, R. M. Carelli, R. M. Wightman, Phasic dopamine signals: from subjective
533 reward value to formal economic utility. *Current opinion in behavioral sciences* **5**,
534 147-154 (2015).

- 535 39. V. Martinez, M. E. Walton, J. O. Gan, P. E. M. Phillips, Phasic dopamine transmission
 536 following state-based reinforcer devaluation in a dual-reward detection task. *Society*
 537 *for Neuroscience Abstracts*, (2008).
- 538 40. J. J. Cone *et al.*, Physiological state gates acquisition and expression of mesolimbic
 539 reward prediction signals. *Proceedings of the National Academy of Science* **113**,
 540 1943-1948 (2016).
- 541 41. R. R. Bush, F. Mosteller, A mathematical model for simple learning. *Psychological*
 542 *Review* **58**, 313-323 (1951).
- 543 42. R. A. Rescorla, A. R. Wagner, in *Classical Conditioning II: Current Research and*
 544 *Theory*, A. H. Black, W. F. Prokasy, Eds. (Appleton-Century-Crofts, New York, 1972),
 545 pp. 64-99.
- 546 43. N. D. Daw, S. J. Gershman, B. Seymour, P. Dayan, R. J. Dolan, Model-based
 547 influences on humans' choices and striatal prediction errors. *Neuron* **69**, 1204-1215
 548 (2011).
- 549 44. L. Deserno *et al.*, Ventral striatal dopamine reflects behavioral and neural signatures
 550 of model-based control during sequential decision making. *Proceedings of the*
 551 *National Academy of Sciences of the United States of America* **112**, 1595-1600
 552 (2015).
- 553 45. K. Wunderlich, P. Smittenaar, R. J. Dolan, Dopamine enhances model-based over
 554 model-free choice behavior. *Neuron* **75**, 418-424 (2012).
- 555 46. M. E. Sharp, K. Foerde, N. D. Daw, D. Shohamy, Dopamine selectively remediates
 556 'model-based' reward learning: a computational approach. *Brain : a journal of*
 557 *neurology*, (2015).
- 558 47. B. B. Doll, K. G. Bath, N. D. Daw, M. J. Frank, Variability in Dopamine Genes
 559 Dissociates Model-Based and Model-Free Reinforcement Learning. *The Journal of*
 560 *neuroscience : the official journal of the Society for Neuroscience* **36**, 1211-1222
 561 (2016).
- 562 48. Y. Niv, N. D. Daw, D. Joel, P. Dayan, Tonic dopamine: opportunity costs and the
 563 control of response vigor. *Psychopharmacology* **191**, 507-520 (2007).
- 564 49. Z. Kurth-Nelson, G. Barnes, D. Sejdinovic, R. Dolan, P. Dayan, Temporal structure in
 565 associative retrieval. *eLife* **4**, (2015).
- 566 50. G. E. Wimmer, N. D. Daw, D. Shohamy, Generalization of value in reinforcement
 567 learning by humans. *The European journal of neuroscience* **35**, 1092-1104 (2012).
- 568 51. R. A. Rescorla, L. Freberg, The extinction of within-compound flavor associations.
 569 *Learning and Motivation* **9**, 411-427 (1978).
- 570 52. J. Ward-Robinson, G. Hall, Backward sensory preconditioning. *Journal of*
 571 *Experimental Psychology: Animal Behavior Processes* **22**, 395-404 (1996).
- 572 53. P. C. Holland, Event representation in Pavlovian conditioning: image and action.
 573 *Cognition* **37**, 105-131 (1990).
- 574 54. N. J. Mackintosh, *The Psychology of Animal Learning*. (Academic Press, London,
 575 1974).
- 576 55. G. Hall, Learning about associatively activated representations: Implications for
 577 acquired equivalence and perceptual learning. *Animal Learning & Behavior* **24**, 233-
 578 255 (1996).
- 579 56. Y. Takahashi *et al.*, The orbitofrontal cortex and ventral tegmental area are necessary
 580 for learning from unexpected outcomes. *Neuron* **62**, 269-280 (2009).
- 581 57. Y. K. Takahashi *et al.*, Neural estimates of imagined outcomes in the orbitofrontal
 582 cortex drive behavior and learning. *Neuron* **80**, 507-518 (2013).

584 **Figure Legends**

585

586 **Figure 1: Rats infer the value of cues during sensory preconditioning.** Panels
587 illustrate the task design and show the percentage of time spent in the food cup during
588 presentation of the cues during each of the three phases of training. In the 'preconditioning'
589 phase (**A**) rats learn to associate auditory cues in the absence of reinforcement; during this
590 phase there is minimal food cup responding (ANOVA, $F(3, 55) = .7, p = 0.52$). In
591 subsequent 'conditioning' (**B**), rats learn to associate one of the cues (B) with reward;
592 conditioned responding at the food cup during B increases across sessions (ANOVA, main
593 effect of cue: $F(1, 163) = 280.1, p < 0.001$, main effect of session: $F(5, 163) = 9.7$,
594 interaction: $F(5, 163) = 10.81, p < 0.001$). In a final 'probe' test (**C**), rats are presented with
595 each of the 4 auditory cues; conditioned responding at the food cup is maintained to B and
596 is also now evident during presentation of A, the cue that had been paired with B in the
597 preconditioning phase (ANOVA, main effect of cue: $F(1, 167) = 8.7, p < 0.001$, main effect
598 of trial: $F(5, 167) = 6.08, p < 0.001$, interaction: $F(5, 167) = 2.07, p = 0.07$).

599

600 **Figure 2: VTA dopamine neurons exhibit firing to a reward-paired cue that is**
601 **consistent with TD error signaling.** We recorded 632 neurons across all days of
602 conditioning and the final reminder session. (**A**) Normalized responses (AUC) are displayed
603 for each neuron, sorted by the classification algorithm applied by Cohen, Uchida and
604 colleagues (21). The first three principal components (PCs) were extracted, to find the
605 major modes of this population's response (**B**), then hierarchical agglomerative clustering
606 was used on those PCs to identify similar neural responses; groups identified are
607 highlighted in color (**C**); The mean group response of each of the populations identified are
608 displayed (**D**); in accordance with previous results (21) we found populations undergoing
609 sustained excitation, phasic excitation, and sustained inhibition. Consistent with
610 identification as putative dopamine neurons, the average (AUC) response to cue B from the
611 phasic group on each day of conditioning exhibited a peak response that was highest to
612 reward early in conditioning and migrated to earlier cue onset across conditioning (**E-F**,
613 $r(302) = 0.24, p < 0.01$). This change in firing is in accordance with signaling of a TD error.

614

615 **Figure 3: VTA dopamine neurons exhibit firing to a pre-conditioned cue that is not**
 616 **consistent with TD error signaling.** We recorded 102 neurons during the probe test. AUC
 617 normalized neural responses were classified with a hierarchical clustering as in Figure 2 (**A-**
 618 **D**) in order to identify putative dopamine neurons ($n = 52$). In addition, we also identified 4
 619 neurons based on traditional waveform criteria. While the classified putative dopamine
 620 neurons showed firing to all cues, they exhibited the largest responses at the onset of B, the
 621 reward-paired cue (significantly above responding to D, $t(51) = 4.40$, $p < 0.001$), and to A,
 622 the cue that had been paired with B in the preconditioning phase (significantly above
 623 responding to control cue C, $t(51) = 5.02$, $p < 0.001$) (**E-F**). Further, the activity elicited by
 624 these two cues was strongly correlated (**F**), suggesting that dopamine neurons code errors
 625 elicited by these two types of cues in a common framework (correlation between B-D and A-
 626 C, $r(50) = 0.63$, $p < 0.001$).

627

628 **Figure 3-figure supplement 1: neural responses from phasic and tonic wide-**
 629 **waveform neurons.** (**A**) raster plot of 18 trials of cue responses, resorted according to cue,
 630 for phasic responding wide waveform neuron. (**B**) baseline subtracted mean responses of
 631 panel **A** for cues B and D. (**C**) baseline subtracted mean responses to of panel **A** for cues B
 632 and D (**D**) raster plot of 18 trials of cue responses, resorted according to cue, for tonic
 633 excited wide waveform neuron. (**E**) baseline subtracted mean responses of panel **D** for
 634 cues B and D. (**F**) baseline subtracted mean responses to of panel **D** for cues B and D

635

636 **Figure 3-figure supplement 2: neural responses from 39 neurons classified as**
 637 **tonically excited by cue B.** (**A**) baseline subtracted, mean responses of all neurons to
 638 cues B and D, \pm SEM (**B**) baseline subtracted, mean responses of all neurons to cues A
 639 and C, \pm SEM (**C**) histogram of differences in neural responding to cached value (B-D) for
 640 all tonically excited neurons for the first second of cue response; there was no significant
 641 difference ($t(38) = 0.37$, $p = 0.71$) between responses to cue B and D (**D**) histogram of
 642 differences in neural responding to inferred value (A-C) for all tonically excited neurons for
 643 the first second of cue response; there was a significant difference between early responses

644 to cue A and C ($t(38) = 2.9, p < 0.01$), **(E)** histogram of differences in neural responding to
 645 cached value (B-D) for all tonically excited neurons for the final nine seconds of cue
 646 response; neurons fired significantly more to cue B than D ($t(38) = 6.3, p > 0.001$) **(F)**
 647 histogram of differences in neural responding to inferred value (A-C) for all tonically excited
 648 neurons for the last nine seconds of cue response; there was a smaller but significant
 649 difference between responses to cue A and C ($t(38) = 2.4, p < 0.05$) **(G)** scatter of individual
 650 responses to cached vs inferred value (i.e. data from panel C vs panel D); while there was a
 651 positive relationship, the correlation was not significant ($r(37) = 0.26, p = 0.11$).

652

653 **Figure 3-figure supplement 3: neural responses from 11 neurons classified as**
 654 **tonically inhibited by cue B.** **(A)** baseline subtracted, mean responses of all neurons to
 655 cues B and D, +/- SEM **(B)** baseline subtracted, mean responses of all neurons to cues A
 656 and C, +/- SEM **(C)** histogram of differences in neural responding to cached value (B-D) for
 657 all tonically inhibited neurons for the first second of cue response; there was no significant
 658 difference ($t(10) = -1.56, p = 0.15$) between responses to cue B and D **(D)** histogram of
 659 differences in neural responding to inferred value (A-C) for all tonically inhibited neurons for
 660 the first second of cue response; there was no significant difference ($t(10) = 0.99, p = 0.34$)
 661 between responses to cue A and C. **(E)** histogram of differences in neural responding to
 662 cached value (B-D) for all tonically inhibited neurons for the last nine seconds of cue
 663 response; there was no significant difference ($t(10) = -1.6, p = 0.14$) between responses to
 664 cue B and D **(F)** histogram of differences in neural responding to inferred value (A-C) for all
 665 tonically inhibited neurons for the last nine second of cue response; there was no significant
 666 difference ($t(10) = 0.03; p = 0.98$) between responses to cue A and C **(G)** scatter of
 667 individual responses to cached vs inferred value (i.e. data from panel C vs panel D); while
 668 there was a positive relationship, the correlation was not significant ($r(9) = 0.46, p = 0.15$).

Figure 1




Figure 2




Figure 3

