1
2
3 **Synaptic memory requires CaMKII**
4

5 Wucheng Tao^{1,2}, Joel Lee^{2#}, Xiumin Chen^{2#}, Javier Diaz- Alonso^{2,5}, Jing Zhou³,
6 Samuel Pleasure³ and Roger A. Nicoll^{2,4*}

7 ¹Key Laboratory of Brain Aging and Neurodegenerative Diseases, School of
8 Basic Medical Sciences, Fujian Medical University, Fuzhou 350108, Fujian
9 Province, China.

10 ²Departments of Cellular and Molecular Pharmacology, ³Neurology, and

11 ⁴Physiology, University of California, San Francisco, CA

12 ⁵Present address: Department of Neuroscience, University of California, Irvine,
13 CA

14
15 [#]contributed equally

16
17 *Correspondence should be addressed to:

18 Roger A. Nicoll

19 Department of Cellular and Molecular Pharmacology

20 University of California

21 San Francisco, CA 94158

22 Email: roger.nicoll@ucsf.edu
23
24

25 Submission: July 12, 2020
26
27
28
29
30

SUMMARY

Long-term potentiation (LTP) is arguably the most compelling cellular model for learning and memory. While the mechanisms underlying the induction of LTP (“learning”) are well understood, the maintenance of LTP (“memory”) has remained contentious over the last 20 years. Here, we find that CaMKII contributes to synaptic transmission and is required LTP maintenance. Acute inhibition of CaMKII erases LTP and transient inhibition of CaMKII enhances subsequent LTP. These findings strongly support the role of CaMKII as a molecular storage devise.

INTRODUCTION

The mechanism by which the brain stores information has fascinated neuroscientists for well over a century. Two general ideas have emerged. The first proposes that information is held by ongoing activity in neuron ensembles. This, indeed, underlies short-term working memory (D'Esposito and Postle, 2015; Inagaki et al., 2019; Wang, 2001). However, it is clear that such a mechanism cannot account for long lasting, enduring memories. For instance, it is well established that patients who recover from prolonged periods of isoelectric EEG and brain stem silence following barbiturate overdose have no detectable cognitive deficits, e.g., (Bird and Plum, 1968). The second and generally accepted view, first enunciated by Cajal over a century ago (Cajal, 1911), is the modification of neuronal connections. How these changes might be stored has remained a mystery.

Over the past few decades, Ca^{2+} -calmodulin-dependent kinase II (CaMKII) has emerged as a very attractive molecular candidate for information storage (Bayer and Schulman, 2019; Bhattacharyya et al., 2020; Hell, 2014; Kennedy, 2013; Lisman et al., 2002; Lisman et al., 2012; Lisman and Goldring, 1988). This

dodecameric kinase is activated by $\text{Ca}^{2+}/\text{CaM}$ resulting in autophosphorylation and the activity of the kinase remains after the removal of Ca^{2+} (Miller and Kennedy, 1986), a state referred to as autonomy. Furthermore, evidence suggests that the enzyme can undergo activation-triggered subunit exchange such that subunits that did not experience the original Ca^{2+} transient are phosphorylated, a mechanism postulated to allow for continued phosphorylation in the face of protein turnover (Bhattacharyya et al., 2020; Bhattacharyya et al., 2016; Stratton et al., 2014) Thus, CaMKII has many attractive biochemical features expected for a memory molecule. What is the physiological evidence for such a role?

Long-term potentiation (LTP), in which brief high frequency synaptic stimulation results in a lasting increase in synaptic strength, is an attractive cellular model for learning and memory (Choquet, 2018; Collingridge et al., 2004; Huganir and Nicoll, 2013; Malinow and Malenka, 2002; Nicoll, 2017). A role for CaMKII in LTP is compelling. Pharmacologically blocking CaMKII (Malenka et al., 1989; Malinow et al., 1989; Otmakhov et al., 1997) or genetically deleting CaMKII (Giese et al., 1998; Incontro et al., 2018; Silva et al., 1992) blocks LTP and constitutively active CaMKII both mimics and occludes LTP (Lledo et al., 1995; Pettit et al., 1994; Pi et al., 2010; Poncer et al., 2002). However, despite the attractive biochemical properties of CaMKII, whether CaMKII is involved in the *maintenance* of LTP and, by extension, memory storage remains problematic.

There are two predictions if CaMKII is responsible for LTP maintenance and synaptic memory (Lisman, 2017; Sanhueza and Lisman, 2013). First, transiently blocking CaMKII *after* the induction of LTP should cause a long-lasting erasure. Attempts at reversing LTP have failed on numerous occasions (Buard et al., 2010; Chen et al., 2001; Malinow et al., 1989; Murakoshi et al., 2017; Otmakhov et al., 1997), but see (Feng, 1995). Second, if CaMKII is involved in synaptic memory, one would expect it to leave a lasting trace at synapses (Lisman, 2017; Sanhueza et al., 2011; Sanhueza and Lisman, 2013). Thus, silencing CaMKII

93 should reduce synaptic transmission. Interestingly, the CaMKII/NMDAR complex
94 does exist under basal conditions (Leonard et al., 1999). However, the genetic
95 deletion of CaMKII (Achterberg et al., 2014; Giese et al., 1998; Silva et al., 1992),
96 but see (Hinds et al., 1998), or pharmacological inhibition of CaMKII does not
97 affect baseline synaptic responses (Buard et al., 2010; Chen et al., 2001; Feng,
98 1995; Malinow et al., 1989; Murakoshi et al., 2017; Otmakhov et al., 1997; Wang
99 and Kelly, 1996). Finally, using a FRET-based CaMKII sensor, CaMKII activation
100 in spines after LTP induction persists for only ~1 min. (Lee et al., 2009). Thus,
101 physiological support for a role of CaMKII in maintaining LTP or synaptic
102 transmission is lacking.

103
104 Recent findings have prompted us to reevaluate CaMKII's role in the
105 maintenance of LTP. Deleting CaMKII with CRISPR (Incontro et al., 2018) or
106 expressing a peptide inhibitor of CaMKII (Goold and Nicoll, 2010; Sanhueza et al.,
107 2011) cause a substantial reduction in synaptic transmission. Furthermore,
108 studies using a membrane permeable peptide inhibitor of CaMKII (tatCN21 or
109 antCN27) have reported a lasting depression in synaptic transmission (Barcomb
110 et al., 2016; Gouet et al., 2012; Sanhueza et al., 2011; Sanhueza et al., 2007)
111 and evidence for a reduction in LTP maintenance (Sanhueza et al., 2011;
112 Sanhueza et al., 2007).

113
114 To reevaluate the role of CaMKII in synaptic memory we have used two different
115 rapidly acting and reversible CaMKII inhibitors. As would be expected if LTP
116 contributes to synaptic memory, inhibition of CaMKII depresses synaptic
117 transmission and the properties of this persistent action of CaMKII are
118 remarkably similar to those of LTP. Furthermore, we demonstrate that applying
119 these inhibitors *after* inducing LTP fully reverses LTP, indicating that CaMKII is
120 required for the persistence of LTP. Thus, our findings strongly support CaMKII
121 as a molecular storage devise. Reasons for why the present results differ from
122 most previous studies are discussed.

RESULTS

Blocking CaMKII inhibits synaptic transmission

Previous studies established, both with the expression of inhibitory peptides (Goold and Nicoll, 2010; Sanhueza et al., 2011) or CRISPR deletion of CaMKII (Incontro et al., 2018) that CaMKII contributes ~50% to AMPAR responses. In order to study the basis for this activity as well as the role of CaMKII in LTP maintenance, rapid and reversible block of CaMKII is essential. Since peptide inhibitors have proved most effective, this requires making these peptides membrane permeable either by cell-penetrating peptides (CPPs) (e.g., tat) or by protein lipidation (e.g., myristoylation). Delivery of peptides into cells has profound therapeutic potential and therefore has received a great deal of study (Allen et al., 2018; LeCher et al., 2017; Nelson et al., 2007; Patel et al., 2019). The effectiveness of CPPs is hotly debated, because, although these peptides clearly enter the cell by endocytosis, it is not certain to what degree these peptides actually have access to the cytoplasm. This appears not to be an issue with lipid modification.

As an initial assay we bathed acute slices in a peptide inhibitor and measured the AMPAR/NMDAR ratio. We would expect the ratio to be reduced by ~50%, since expressing these peptides had no effect on NMDAR currents. We first tested the effectiveness of tatCN21 (5 μ M) (Barcomb et al., 2016; Buard et al., 2010; Sanhueza et al., 2011) in reducing the AMPAR/NMDAR ratio (Fig. S1). In agreement with previous reports, we failed to see an effect on AMPAR responses. Since higher concentrations of tatCN21 have significant nonspecific effects (Barcomb et al., 2016), we turned to other peptides. Previous work (Goold and Nicoll, 2010) showed that prolonged (1-2 days) incubation with myr-CN27 (10 μ M) caused a ~50% reduction in the AMPAR/NMDAR ratio. We, therefore, repeated this experiment with shorter incubations (1-2 hours) and a lower concentration (1 μ M) and found that myr-CN27 reduced the ratio by ~50% (Fig. S1/Figure1-figure suple1). What might account for the lack of effect of tatCN21?

To address this, we repeated the experiment using Myr-CN21 (5 μ M) and found that it reduced the ratio by ~50% (Fig. S1/ Figure1-figure suple1), suggesting that lipidation is a more effective mode of delivery than CPPs. We, therefore, selected myr-CN27 for our studies.

Application of myr-CN27 to acute slices (1 μ M) caused a highly reproducible slowly developing depression of synaptic AMPAR mediated EPSCs, which stabilized after approximately 40 min. (Fig. 1A and B). A number of experiments were carried out to ensure that the effect we observe with myr-CN27 is due to the specific inhibition of CaMKII. First, in many experiments we measured the size of the NMDAR EPSC, by depolarizing the cell to +40 mV before and after the application of the peptide. We found that myr-CN27 had no effect on the NMDAR EPSC (Fig. 1A, red circles, Fig. 1B). Second, it is important to note that the magnitude of the decrease in transmission seen with myr-CN27 is similar to that reported with the genetic deletion of CaMKII α (Incontro et al., 2018) suggesting that myr-CN27 fully silences CaMKII. Finally, we examined the effects of myr-CN27 in cells lacking CaMKII α (Fig. 1C and D). CaMKII α was deleted using CRISPR (Incontro et al., 2018). We simultaneously recorded from control cells and cells lacking CaMKII α in slice culture. In the absence of CaMKII α (green circle) synaptic responses were ~50% of control (black circles), as expected. Application of myr-CN27 depressed controls cells to the level recorded in cells lacking CaMKII α (Fig. 1C and D, black circles), but it had no additional effect on cells lacking CaMKII α (green circles) (n = 6). These results demonstrate that the effect of CN27 is fully explained by its selective and complete silencing of CaMKII α .

Previous studies have relied on field potential recording and thus we repeated our experiments with this approach (Fig. 1E). We still recorded a depression, but in this case the depression took tens of minutes to develop (black symbols). We interleaved these experiments with whole cell recording and confirmed that the depression is considerably faster (blue symbols). We interpret the difference in

time course to the fact that whole cell recordings are made from superficial cells, while field potential responses are generated by populations of neurons throughout the depth of the slice. Penetration of peptides into the slice is expected to be slow.

We next examined the effects of AIP, a peptide inhibitor designed from the autoinhibitory domain. We first expressed AIP in neurons in slice culture and compared AMPAR and NMDAR responses to those in simultaneously recorded neighboring control cells (Fig. 2A). Similar to the effect of expressing CN27 (Goold and Nicoll, 2010) or applying myr-CN27 (see above), there was a selective ~50% reduction in the AMPAR EPSC, but no change in the NMDAR response. Given these positive results, we bathed acute slices in myr-AIP (20 μ M) and measured the AMPAR/NMDAR ratio (Fig. 2B) ($n = 12$). The ratio was significantly reduced. Acute application of myr-AIP in slice culture caused a slowly developing depression in synaptic activity (Fig. 2C, black circles) ($n = 6$) and this effect is due to the selective inhibition of CaMKII, because interleaved recording from cells in which CaMKII α had been deleted (Fig. 2C, green circles), myr-AIP no longer had any effect ($n = 5$). These findings show that myr-CN27 and myr-AIP selectively and completely inhibit CaMKII, both in acute slices and in slice culture. It is interesting that, while CaMKII is expressed presynaptically and reported to effect transmitter release (Benfenati et al., 1992; Hinds et al., 2003), there was no presynaptic effect of the peptide inhibitors.

The time course of inhibition for both myr-CN27 and myr-AIP is slow, requiring 10's of minutes. Is this due to the slow access of these peptides to the cell interior, or is it due to the slow reversal of the action of CaMKII? To address this, we turned to a novel light inducible inhibitor of CaMKII, in which the inhibitory peptide AIP2 (Ishida et al., 1998) is linked to the LOV-J α helix domain of Phototropin (referred to as paAIP2) (Murakoshi et al., 2017). Blue light rapidly exposes the AIP2 peptide, which then returns to its closed inactive conformation in approximately 40 s following light exposure. We expressed paAIP2 in

individual neurons with biolistics, which, unlike the myristoylated peptides, has the advantage of limiting the manipulation to individual postsynaptic cells. The synaptic properties of neurons expressing paAIP2 were not different from the properties of neighboring control cells (Fig. S2A and B/ Figure3-figure suple1.2A and B) (n = 16). Exposure of these cells to blue light resulted in a slowly developing inhibition of AMPAR responses (Fig. 3A) (n = 12), similar to the kinetics of myr-CN27 and myr-AIP. The blue light had no effect on simultaneously recorded control cells (Fig. 3A, black circles) or in cells co-expressing paAIP2 with CRISPR to delete CaMKII α (Fig. S3/ Figure3-figure suple2) (n = 5), establishing the specificity of paAIP2. These results with paAIP2 suggest that the slow time course is not due to slow access, but rather due to the slow reversal of the action of CaMKII.

The origin of the constitutive action of CaMKII

The persistent CaMKII activity has two possible origins. First, it could represent ongoing constitutive Ca²⁺ activation of CaMKII from various sources. It seems unlikely that resting Ca²⁺ levels (Maravall et al., 2000) stimulate CaMKII, since the sensitivity of CaMKII to Ca²⁺ requires considerably higher levels of Ca²⁺ (Schulman, 1984). To directly test whether constitutive CaMKII activity is Ca²⁺ dependent, we loaded cells with the Ca²⁺ chelator BAPTA (15 mM). However, myr-CN27 retained its depressive effect (Fig. S4A/Figure4-figure suple1.A). This finding also rules out spontaneous background NMDAR activity as a source for the constitutive CaMKII activity. In addition, acute application of the NMDAR antagonist APV, has little effect on AMPAR EPSCs (Incontro et al., 2018).

The constitutive activity might reflect an LTP process induced by slice preparation, which is known to depolarize cells and release glutamate. To address this possibility, we pretreated animals with a high dose of MK-801, which is known to block *in vivo* NMDAR responses (Davies et al., 1988). To verify that NMDARs were blocked by this procedure, we recorded NMDAR EPSCs from

slices from these animals and found that the NMDAR EPSC was blocked (Fig. S4B/Figure4-figure suple1.B). Having confirmed the absence of NMDAR responses the slices were perfused with MK-801. Under such conditions myr-CN27 still exerted its inhibitory effect (Fig. S4C/ Figure4-figure suple1.C).

Second, if CaMKII is involved in synaptic memory, one would expect that LTP acquired when the animal was alive, would contribute to synaptic transmission, as previously proposed (Lisman, 2017; Sanhueza and Lisman, 2013). If this is the case, there are a number of predictions. One would expect that the transient inhibition of CaMKII should result in long lasting inhibition, akin to the resetting of a molecular switch. In a number of cells we monitored the depression over time following the removal of myr-CN27 and found no recovery (Fig. 4A). One might argue that myr-CN27 fails to washout of the cells. To circumvent this possibility, we repeated these experiments with paAIP2 whose action rapidly reverses upon terminating the blue light (~40 s). Again, we failed to observe any recovery (Fig. 4B). We confirmed that this was not due to the continued action of the inhibitor, by demonstrating that LTP could be induced following light termination (see below, Fig. 5).

If the constitutive activity resulted from prior NMDAR-dependent LTP, one would expect that this inhibitory action should be absent in cells in which NMDARs have been deleted embryonically. We carried out *in utero* electroporation to express cre recombinase in neurons in GluN1 floxed mice. Transfected cells were recorded at P16-20 in acute slices. We first carried out paired recordings in the AMPAR inhibitor CNQX to ensure that cre expressing neurons lacked NMDAR EPSCs (Fig. 4C, left panel). Application of myr-AIP had no effect in these cells (Fig. 4C, middle and right panels, green circles and traces), whereas it had its normal inhibitory effect in interleaved control cells (Fig. 4C, black circles and traces), further supporting the conclusion that the persistent CaMKII activity represents prior LTP. A caveat to these experiments is that the synaptic action of CaMKII requires its binding to the GluN2B C-terminus of the NMDAR. Thus, the

loss of GluN2B in the GluN1 lacking neurons could equally explain the loss of effect of myr-CN27. To address this concern we repeated these experiments, replacing GluN1 with a pore dead GluN1 mutant (N812W) (Amin et al., 2017). While this mutant assembles normally with GluN2 subunits and traffics normally to the membrane, it generates little current (Amin et al., 2017). We carried out *in utero* electroporation in GluN1 floxed mice to transfect Cre and GluN1(N812W). Acute slices were made at P14-18. The NMDAR EPSC in transfected neurons was severely reduced, when compared to neighboring control cells recorded in CNQX (Fig. 4D, left panel). The depressant effect of myr-CN27 recorded in these transfected cells was absent (Fig. 4D, middle and right panels). These results suggest that in behaving animals, cells that lack NMDAR function, but with GluN2B intact, Ca^{2+} signaling independent of the NMDAR appears incapable of activating CaMKII, at least in regard to its action on synaptic function.

Since LTP is saturable, one might expect that when constitutive activity is *transiently* silenced, the magnitude of LTP should be larger than that in control conditions. To test this prediction, we transfected paAIP2 *in utero* and prepared acute slices at days P15-25. The slices were exposed to blue light for 40 minutes and then simultaneous recordings were made from a transfected cell and a control cell (Fig. 5A). In a typical experiment (Fig. 5B), as well as in the summary of all experiments (Fig. 5D), the size of baseline EPSCs in paAIP2 expressing cells following blue light exposure was, as expected, approximately half of that of the control neighboring cells. LTP was then induced in both cells. To compare the magnitude of LTP in the two cells we normalized the baseline EPSCs to 100%. As is clear, both in the single experiment (Fig. 5C) and in the summary (Fig. 5E), the magnitude of LTP in cells expressing paAIP2 is approximately twice that of control cells. This interaction between the constitutive potentiation and LTP is consistent with their sharing the same underlying mechanism. Taken together these results strongly support the conclusion that while the animal is alive, LTP is acquired and leaves a synaptic memory trace.

Blocking CaMKII erases LTP

The experiments presented thus far are all consistent with the notion that synapses acquire LTP that contributes to synaptic transmission. If this is correct, then inhibiting CaMKII *after* LTP induction must reverse the potentiation. Yet, as discussed in the introduction, this experiment has failed on numerous occasions (Buard et al., 2010; Chen et al., 2001; Malinow et al., 1989; Murakoshi et al., 2017; Otmakhov et al., 1997), but see (Feng, 1995). Thus, it was mandatory that we revisit this crucial experiment. We therefore carried out a “two pathway experiment”, in which the responses from two independent pathways were recorded from a single cell (Fig. 6A). Following the recording of baseline responses, LTP was induced in one of the pathways (Fig. 6B, red circles) by pairing depolarization of the cell to 0 mV with continued synaptic stimulation, while in the other control pathway, stimulation was stopped during the depolarization (Fig. 6B, black circles). Once LTP had been established, myr-CN27 was applied and the effect on both pathways recorded. Myr-CN27 caused a complete reversal of established LTP and as expected, a ~50% reduction in the control pathway (Fig. 6B). A summary of all the experiments (Fig. 6B, n = 11) demonstrates that in the presence of myr-CN27 the LTP pathway and control pathway converge to approximately 50% of the control. This experiment establishes the essential role of CaMKII in maintaining LTP. A previous field potential study using tatCN21 (5 μ M) failed to observe a reversal of LTP (Buard et al., 2010). We, therefore, repeated our two pathway experiment using field potential recording. Again, we observed that myr-CN27 (1 μ M) caused a slow reduction the EPSPs in the control pathway and a reversal of LTP (Fig.S5/ Figure6-figure suple1). Give that tatCN21 (5 μ M) had no effect on synaptic transmission (Buard et al., 2010) (Fig. S1/ Figure1-figure suple1), it is not surprising that it failed to reverse LTP in the previous study (Buard et al., 2010).

DISCUSSION

The role of CaMKII in the maintenance of LTP and information storage has been one of the most vexing issues in the field of synaptic plasticity. On the one hand, the biochemical properties of CaMKII have for decades made this molecule an extremely attractive candidate for molecular storage (Bhattacharyya et al., 2020; Coultrap and Bayer, 2012; Hell, 2014; Lisman et al., 2002; Lisman et al., 2012). On the other hand, numerous physiological experiments over the years have failed to support the role of CaMKII in the *maintenance* of LTP and by extension memory. To reevaluate the physiological role of CaMKII in synaptic memory, we have used two classes of CaMKII inhibitory peptides. With these inhibitors we find that CaMKII is required for the maintenance of LTP and provide evidence that LTP acquired while the animal was alive leaves a lasting synaptic memory trace.

The first class of inhibitory peptide we used is derived from an endogenous CaMKII inhibitory protein referred to as CaMKIINtide (CN27) (Chang et al., 1998; Goold and Nicoll, 2010; Pellicena and Schulman, 2014; Vest et al., 2007). The second class of inhibitory peptide is derived from the autoinhibitory domain of CaMKII (Bayer et al., 2001; Leonard et al., 1999). Although these inhibitory peptides were thought to bind to separate sites (S and T sites), recent structural studies of the binding of GluN2B and other interacting peptides to CaMKII indicate that these peptides use similar interactions to bind across the substrate binding pocket of the CaMKII active site (Özden et al., 2020). Thus, at present it is not possible to use these two classes of inhibitors to distinguish between the disruption of GluN2B binding and the blockade of kinase activity. These peptides included a myristoylated version of CN27 (myr-CN27), AIP (myr-AIP) and a recently developed photoactivatable peptide inhibitor (paAIP2) (Murakoshi et al., 2017). Both classes of peptides block kinase activity with high affinity (Chang et al., 2001; Ishida et al., 1998), but also interfere with the binding of CaMKII to GluN2B (Sanhueza et al., 2011; Vest et al., 2007).

Our experiments clearly establish that inhibitory peptides fully reverse LTP. How long is CaMKII responsible for maintaining LTP? Experiments addressing the role of constitutive CaMKII in the maintenance of LTP are constrained by the length of recording; approximately 1 hour for whole cell and a few hours for field potentials. However, if LTP is the substrate for memory, it should leave a lasting memory trace at synapses, acquired while the animal is alive (Lisman, 2017; Sanhueza and Lisman, 2013). A number of studies have reported a lasting depression following transient inhibition of CaMKII (Barcomb et al., 2016; Gouet et al., 2012; Sanhueza et al., 2011; Sanhueza et al., 2007). Furthermore, even under basal conditions, CaMKII is found in isolated PSDs (Petersen et al., 2003; Strack et al., 1997a) and synaptic puncta (Bayer et al., 2006) and this is in the autophosphorylated state (Strack et al., 1997b; Trinidad et al., 2005). As discussed above all three inhibitory peptides (myr-CN27, myr-AIP and paAIP2) depress synaptic transmission and this depression requires the presence of CaMKII. A number of trivial explanations, such as ongoing constitutive Ca^{2+} activation of the enzyme, were excluded.

We then focused on the possibility that this constitutive action of CaMKII reflects prior LTP. First, the finding that the inhibition does not affect NMDAR responses is consistent with an LTP mechanism, since LTP preferentially enhances AMPAR responses (reviewed in (Nicoll, 2017)). Second, there should be no recovery from the transient inhibition of CaMKII, analogous to the resetting of a molecular switch. This is, indeed, the case. Third, cells in which the NMDAR has been deleted embryonically should be devoid of the constitutive activity. Again, this is the case. This is unlikely to be due to the loss of GluN2B subunit, because cells expressing a pore dead NMDAR mutant, in which assembles with GluN2B, had little CaMKII constitutive activity. This finding suggests that non-NMDAR sources of Ca^{2+} in the behaving animal are incapable of triggering CaMKII-dependent enhancement in synaptic transmission. This tight linking of NMDAR sources of Ca^{2+} to CaMKII is critical, because it prevents the degrading of the Hebbian nature of plasticity. Finally, it is well established that LTP is saturable. Thus, if

constitutively active CaMKII represents LTP, removing this component should allow for larger LTP. We find that in cells in which CaMKII is transiently blocked LTP is approximately twice as large as in neighboring control cells. Taken together these results suggest that constitutive CaMKII represents LTP acquired at synapses while the animal was alive, thus supporting a role for CaMKII in synaptic memory.

Based on the present results the term “basal synaptic transmission” needs to be reevaluated. It has generally been assumed that the synapses studied in a hippocampal slice, in which much of the afferent drive from multiple inputs have been removed in the slicing, are at a ground “basal” state. The present results indicate that the synaptic currents we measure are actually maintained by a persistent enhancement acquired prior to slicing.

It should be pointed out that the contribution of CaMKII to synaptic transmission is unlikely to be a simple addition to the overall preexisting excitatory drive onto the cell. It has long been argued that such a scenario would be unstable and quickly saturate (Bienenstock et al., 1982; Cooper and Bear, 2012; Fusi et al., 2005; Morrison et al., 2008; Toyozumi et al., 2014; Turrigiano, 2008). A number of well-established mechanisms exist to maintain stability, including heterosynaptic depression (Scanziani et al., 1996), NMDAR-dependent long-term depression (Malenka and Bear, 2004) and on a longer time scale cell wide homeostasis (Turrigiano, 2008). Thus, it is envisaged that the CaMKII memory trace is embedded in a network of synapses with no overall net change in the excitatory drive onto the cell or network.

It is striking to compare the action of paAIP2 on LTP induction (Murakoshi et al., 2017) and on synaptic transmission (present study). A brief light exposure (~1 min.), delivered immediately inducing LTP, is sufficient to prevent LTP (Murakoshi et al., 2017). This finding indicates that the blocking of CaMKII activation by paAIP2 is very rapid. In contrast, the effect on synaptic transmission

and the reversal of LTP required 10's of minutes. What might explain the dramatic difference in the kinetics of inhibition of LTP induction compared to LTP maintenance and constitutive CaMKII? There is a well-accepted sequence of events for the role of CaMKII in the induction of LTP (Coultrap and Bayer, 2012; Hell, 2014; Lisman et al., 2012; Nicoll, 2017). Following NMDAR activation Ca^{2+} binds to CaM, which then binds to CaMKII activating the enzyme. This causes the autophosphorylation of T286 and the translocation of cytosolic CaMKII to the PSD where it binds to the C-tail of GluN2B. It seems reasonable to expect that under baseline conditions much of CaMKII and paAIP2 are freely diffusible in the spine cytoplasm, as are phosphatases. The activation of paAIP2 would quickly inhibit CaMKII autophosphorylation and its recruitment to the PSD and phosphatases would quickly reverse its effect. During baseline transmission and during the maintenance of LTP a fraction of CaMKII is bound to GluN2B. Evidence suggests that the CaMKII/GluN2B complex is protected from phosphatases (Cheriyana et al., 2011; Lisman and Raghavachari, 2015; Mullasseril et al., 2007). Thus, reversing action of this sequestered CaMKII would be more difficult than preventing phosphorylation of cytosolic CaMKII.

The present results could shed light on the underlying mechanism of long-term depression (LTD). Although this topic remains contentious (Collingridge et al., 2010; Dore et al., 2016; Goodell et al., 2017; Lisman and Zhabotinsky, 2001; Malenka and Bear, 2004; Stein et al., 2021; Wong and Gray, 2018), the long-held view is that LTD is mediated by the activation of phosphatases, in particular protein phosphatase 1 (PP1) (Mulkey et al., 1994; Mulkey et al., 1993). One of the limitations to this model is the lack of evidence that synaptic transmission is maintained by kinase activity. Our demonstration that CaMKII maintains synaptic transmission provides a very simple model for LTD., i.e., a depotentiation or reversal of this prior LTP. One of the features of LTD is that it takes many minutes for its induction. Our finding that it takes many minutes for CaMKII inhibitors to depress synaptic transmission provides an explanation for this property. A formal model for the interplay between CaMKII and PP1 was

proposed some time ago (Lisman and Zhabotinsky, 2001). The absence of LTD in CaMKII knockout mice (Coultrap et al., 2014; Stevens et al., 1994) support such a model, but does not rule out other models.

What might account for the failure of previous studies to detect a component of synaptic transmission driven by constitutive CaMKII (Achterberg et al., 2014; Buard et al., 2010; Chen et al., 2001; Feng, 1995; Giese et al., 1998; Malinow et al., 1989; Murakoshi et al., 2017; Otmakhov et al., 1997; Silva et al., 1992; Wang and Kelly, 1996), but see (Hinds et al., 1998), or to detect a role of constitutive CaMKII in maintaining LTP (Buard et al., 2010; Chen et al., 2001; Malinow et al., 1989; Murakoshi et al., 2017; Otmakhov et al., 1997), but see (Feng, 1995)? We suggest three factors. The first concerns the duration of peptide application, which in our hands takes 10's of minutes to act, especially with field potential recording. The second concerns the concentration and time. As discussed above it is more difficult to reverse the constitutive action of CaMKII, than it is to block its activation. Finally, the membrane permeabilizing agent is important. In our hands myristoylation was more effective and specific than the cell-penetrating peptide, tat.

In summary our results overcome many of the obstacles that have prevented embracing CaMKII as a molecular storage device. Specifically, our results show that transient inhibition of CaMKII results in a lasting depression of synaptic transmission with properties consistent with the erasure of prior LTP acquired while the animal was alive. Furthermore, the finding that CaMKII inhibition *after* the induction of LTP reverses LTP, establishes its role in LTP maintenance. Thus, these physiological results compliment the rich biochemical literature on CaMKII, making CaMKII a particularly attractive molecular storage device. It is important to note, that our results have focused on LTP and it remains open as to whether CaMKII actually stores memories. Recent experiments (Rossetti et al., 2017) using the viral expression of the catalytically dead CaMKII K42M mutant, presumed to act in a dominant negative manner, support its role in memory.

METHODS

Animals

All the experimental procedures on animals were approved by the UCSF Animal Care and Use Committee, BUA # BU002466-04C. For acute slice recordings, typically we use 5-10 animals to obtain complete dataset; for culture slice recordings, we usually used 5-7 animals to obtain complete dataset.

Experimental constructs and chemical agents

The paAIP2 plasmid was obtained from Dr. Ryohei Yasuda and had been characterized (Murakoshi et al., 2017). The CRISPR construct targeting at CaMKII α was previously characterized (Incontro et al., 2018). For biolistic experiments, all the plasmids were expressed in pCAGGS vector, which contains an internal ribosome entry site (IRES) followed by the fluorophore GFP. myr-CaMKIIntide (myr-CN27) and myr-AIP were purchased from Calbiochem. Inc (catalog#, 208921; catalog#, 189482). MK-801 maleate was purchased from HelloBio. Inc (catalog#, HB30004). Myr-CN21 was custom ordered from Elim Biopharm, Inc by myristoylating the N-terminal of CN21:

KRPPKLGQIGRSKRVIEDDR amino acid sequence.

Slice culture and biolistic transfection

Hippocampal cultured slices are obtained from 6-8 days old rats (Stoppini et al., 1991). Biolistic transfection was done 1 day after sectioning, by using a Helio Gene Gun with 1 μ m DNA-coated gold particles (Bio-Rad). Slices were maintained at 34°C and the medium was changed every two days. Typically, slices are used for electrophysiological recording 6-8 days after transfection, except for CRISPR experiments, in which slices are maintained for another week before recording.

In utero electroporation

E15.5 pregnant mice were anesthetized with 2.5% isoflurane in O₂ and injected with buprenorphine for analgesia. The lateral ventricles of embryos were injected with 1 µl mix plasmid DNA (1 µg/µl) with beveled micropipette. Each embryo was electoporated with 5×50 ms, 35-40 V pulse.

Acute slice preparation

Mice of 2-3 weeks of age were anesthetized with 4% isoflurane, decapitated, and the brain dissected free. The whole brain was sliced into 300 µm slices in cutting solution as described (Granger et al., 2013); recovery at 34 °C for half an hour and then stored at room temperature. Solutions were continuously gassed with 95% O₂/5% CO₂.

Photostimulation

Blue light pulses (0.1 Hz, 1s, 20 mW/cm²) were generated by 473 nm blue DPSS laser (Shanghai Laser&Optics Century, BL473T8-300FC). The blue light from a laser was delivered through a optical patch cable connected to the optical fibers. Light pulses were controlled by a Master-8 (A. M. P. I). The blue light is applied for 1s duration with interval of 10s.

Injection of MK-801

MK-801 was dissolved in saline solution and administered as a single injection i.p. (10 mg/kg body weight) one hour before brain slicing.

Electrophysiological recording

Whole-cell voltage clamp recordings were obtained from either wild type cells or fluorescent transfected pyramidal cell in CA1 region of hippocampus (Schnell et al., 2002). Where indicated dual recordings were made from control and transfected cells. Pyramidal neurons were identified by location and morphology. All recordings were made at 20-25 °C. Internal solution (in mM): 135 CsMeSO₄, 8 NaCl, 10 HEPES, 5 QX314-Cl, 4 Mg-ATP, 0.3 Na-GTP, 0.3 EGTA, 0.1 spermine.

Osmolarity was adjusted to 290-295 mOsm and pH was buffered at 7.3-7.4. External solution (mM): 119 NaCl, 2.5 KCl, 4 CaCl₂, 4 MgCl₂, 1 NaH₂PO₄, 26.2 NaHCO₃, 11 Glucose, bubbled continuously with 95% O₂/5% CO₂. For field recordings, the internal solution was 3 M NaCl with a large opening pipette tip. Synaptic currents were evoked every 10 s with bipolar stimulating electrodes placed in *s. radiatum*. To record EPSCs, picrotoxin (100 μ M) was added to the external solution; for recording of AMPAR EPSCs, the cell membrane was held at -70 mV, while for NMDAR EPSCs, it was held at +40 mV; LTP are induced by stimulating at 2 Hz for 90 seconds while clamping the cell at 0 mV. For two independent synaptic pathway experiments, two bipolar stimulating electrodes were positioned to each side of the recording electrode with a distance of around 100 μ m and alternately stimulated every 20 s. Current responses were collected with a Multiclamp 700B amplifier (Axon Instruments), filtered at 2 kHz, and digitized at 10 kHz. Cells with series resistance larger than 20 M Ω were excluded from analysis.

Statistical analysis

Data analysis was carried out in Igor Pro (Wavemetrics), Excel (Microsoft), and GraphPad Prism (GraphPad Software). All paired recording data were analyzed statistically with a Wilcoxon Signed Rank Test for paired data. For unpaired data, a Mann-Whitney U test was used. Statistical parameters including the types of the statistical tests used, exact value of n, precision measures (mean $\pm$ SEM) and statistical significance are reported in the figure legends. All statistical tests performed were two-sided, and with all tests a p-value of < 0.05 was considered statistically significant. All error bars represent standard error of the mean.

ACKNOWLEDGMENTS

We thank Dr. R. Yasuda for the paAIP2 plasmids, Dr. Wei Lu for contributing the data for Fig. 3A and Dr. Lonnie Wullmuth for GluN1 pore dead plasmid (GluN1(N812W)). We thank members of the Nicoll lab for comments on the manuscript. We appreciate the technical assistance from Eric Dang. This research was supported by grant MH070957.

Figure legends

Figure 1. CaMKII inhibitor myr-CN27 inhibits AMPAR synaptic transmission through CaMKII α . **(A)** Time course of myr-CN27 (1 μ M) inhibition on AMPAR EPSCs and NMDAR EPSCs in acute slices. **(B)** Summary graphs showing myr-CN27 inhibited AMPAR EPSCs (before drug: 63.9 ± 12 pA; after drug: 27.6 ± 3.9 pA; $n = 10$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test), but not NMDAR EPSCs (before drug: 43.2 ± 13.7 pA; after drug: 42 ± 13.4 pA; $n = 5$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). **(C)** Time course of the effect of myr-CN27 on AMPAR EPSCs in wt cells and simultaneously recorded CRISPR-CaMKII α transfected cells, normalized to wt baseline (from culture slices). While myr-CN27 inhibited AMPAR EPSCs in wt cells, it had no effect on CRISPR-CaMKII α transfected cells ($n = 6$ $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). **(D)** Sample traces showing that myr-CN27 inhibited AMPAR EPSCs in control cells, but had no effect in CRISPR- CaMKII α transfected cells. Black traces are control cell, green traces are transfected cell. Mean $\pm$ s.e.m. $**P < 0.01$. **(E)** A comparison of the reduction of synaptic transmission in field and whole cell recording. For the field recording (black, $n=8$) after a 60 min stable baseline, 1 μ M myr-CN27 was applied to the slice for 100 min. For the whole cell recording (blue, $n = 7$), the drug was applied after a 10 min stable baseline. In both, the response was reduced to 50% of baseline. Example traces for both are shown taken at the timepoints indicated by 1 and 2. The field EPSPs shown on the bottom right have scale bars vertical 0.1 mV and 5 ms horizontal. The whole

cell recording shown bottom left have scale bars 50 pA vertical and 20 ms horizontal. Both recordings are done in WT slices with the whole cell recordings done in slice culture DIV3-14 while the fields were done in acute slices from P15-28 mice.

Figure 2. CaMKII inhibitor AIP inhibits AMPAR synaptic transmission through CaMKII α . (A) In culture slices, transfection of AIP reduced AMPAR EPSCs (left, $n = 10$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test) but not NMDAR EPSCs (right, $n = 10$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). Sample traces show the effects of AIP on AMPAR EPSCs and NMDAR EPSCs. (B) AMPA/NMDA ratios compared to WT ($n = 34$) are reduced after myr-AIP (20 μ M) treatment ($n = 12$). Scale bar = 50 pA vertical and 30 ms horizontal. (C) Left, Summary of the effect of myr-AIP (20 μ M) on AMPAR EPSCs in wt cells ($n = 6$) and interleaved CRISPR-CaMKII α transfected cells ($n = 5$) from culture slices, normalized to each cell's baseline. The difference between control and CRISPR-CaMKII α transfected cells at 30 min: $p < 0.01$, Mann-Whitney U test. Right, sample traces showing myr-AIP inhibition of AMPAR EPSC in control cells, but not in CRISPR- CaMKII α transfected cells. Black traces are control cell, green are transfected cell. Mean $\pm$ s.e.m.

Figure 3. Light-activated CaMKII inhibitor paAIP2 inhibits AMPAR synaptic transmission through CaMKII α . (A) Time course of paAIP2 effect on AMPAR and NMDAR synaptic transmission in culture slices. Blue light exposure inhibited AMPAR synaptic transmission in paAIP2 expressing cells (green circles), but not in simultaneously recorded control cells (black circles) or NMDAR synaptic transmission (red circles). (B) Summary data showing that paAIP2 inhibited AMPAR synaptic transmission (before light: 51 ± 13.8 pA; after light: 26.5 ± 7.2 pA; $n = 12$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test), but had no effect on NMDAR synaptic transmission (before light: 31.2 ± 7.6 pA; after light: 29.8 ± 7.7

pA; $n = 5$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). Mean $\pm$ s.e.m. $**P < 0.01$.

Figure 4. The depression caused by transient inhibition of CaMKII is long lasting and is absent in cells lacking NMDARs. (A) In acute slices, time course showing that following myr-CN27 application AMPAR synaptic transmission remains depressed (The difference between before and after myr-CN27: $n = 4$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test). **(B)** In culture slices, time course showing that following blue light exposure on paAIP2 expressing cells AMPAR synaptic transmission remains depressed (The difference between before and after blue light exposure: $n = 5$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test). **(C)** Left panel, paired recordings from acute slices of control cells and those expressing Cre in GluN1 floxed mice ($n = 5$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test); Middle panel, time course of the myr-AIP effect on AMPAR synaptic transmission in control cells ($n = 5$) and interleaved cells expressing Cre in GluN1 floxed mice ($n = 5$). $p < 0.01$, Mann-Whitney U test. Right panel, sample traces showing myr-AIP inhibited AMPAR synaptic transmission in control cell (black traces), but not in a GluN1KO cell (green traces). Mean $\pm$ s.e.m. **(D)** Left panel, paired recordings from acute slices of control cells and those expressing Cre and N812W (pore dead, PD) in GluN1 floxed mice ($n = 6$, $p < 0.05$, two-tailed Wilcoxon Signed Rank Test); Middle panel, the action of Myr-CN27 is not rescued by expressing a GluN1 pore dead mutant. $n = 8$; Right panel, sample traces showing myr-CN27 inhibited AMPAR synaptic transmission in control cell (black traces), but not in a GluN1KO+ PD cell (green traces).

Figure 5. Transient inhibition of CaMKII enhances subsequent LTP. (A) Timeline of experimental procedure. Following in utero electroporation acute slices were prepared at P15-P25. Slices were then exposed to blue light for 40 minutes. Paired recordings were then made from a control cell (black circles) and a paAIP2 expressing cell (green circles). **(B)** One representative experiment

showing that prior paAIP2 activation reduced baseline AMPAR EPSCs, but LTP was larger than that recorded in the neighboring control cell. **(C)** Normalized representative experiment showing that prior paAIP2 activation increased LTP. **(D)** Summary data showing that prior paAIP2 activation reduced baseline AMPAR EPSCs, but enhanced LTP (n = 5). (E) Normalized summary data showing that prior paAIP2 activation doubles the size of LTP. n = 5, p < 0.01, two-tailed Wilcoxon Signed Rank Test. Mean ± s.e.m.

Figure 6. Myr-CN27 erases LTP in acute slices. **(A)** cartoon diagram of two pathway experiment. To record the response from two independent pathways, two bipolar stimulating electrodes were positioned to either side of the recorded cell with a distance of around 100 µm. Stimuli were applied alternately every 20 s. **(B)** A sample experiment showing that myr-CN27 inhibited the LTP pathway more strongly than the control pathway. **(C)** Left, summary data showing that myr-CN27 reduced the control pathway (black circles) 50%, while completely reversing LTP (red circles) (The difference between control and LTP pathway at 60 min: n = 11, p > 0.05, two-tailed Wilcoxon Signed Rank Test). Responses are mean ± s.e.m. Right, sample traces showing the effect of myr-CN27 on AMPA EPSCs in control and LTP pathway. LTP is induced by 2 Hz stimulation for 90 s, while holding the cell at 0 mV.

Supplemental fig. legends

Figure S1/ Figure 1-figure suple1. ^[1-3]SEP The effect of various CaMKII peptide inhibitors on the AMPAR/NMDAR ratio. ^[1-3]SEP AMPAR/NMDAR ratios recorded in acute slices made from WT mice incubated without inhibitor treatment (WT) (n = 34), with 5 µM tatCN21 (n = 10), with 1 µM myr-CN27 (n = 47), and with 5 µM myr-CN21 (n = 9). Sample traces are shown at top. ^[1-3]SEP Scale bar = 50 pA vertical and 30 ms horizontal p < 0.01, Mann-Whitney U test.

Figure S2/ Figure 3-figure suple1. Characterization of synaptic transmission in control cells and paAIP2 transfected cells from culture slices, before and after blue light exposure. (A, B) transfection of paAIP2 had no effect on AMPAR synaptic transmission (control, 96.4 ± 19.7 pA; paAIP2, 96.1 ± 21.3 pA; $n = 16$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test), or NMDAR synaptic transmission (control, 81.5 ± 10.1 pA; paAIP2, 89.7 ± 16.6 pA; $n = 11$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). **(C, D)** blue light exposure reduced AMPAR synaptic transmission in paAIP2 transfected cells (control, 69.1 ± 11.1 pA; paAIP2, 31.7 ± 7.1 pA; $n = 8$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test), but have no effect on NMDAR synaptic transmission (control, 86.5 ± 23.7 pA; paAIP2, 78 ± 20.6 pA; $n = 8$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). Mean $\pm$ s.e.m.

Figure S3/ Figure 3-figure suple2. CRISPR deletion of CaMKII α prevents paAIP2 effect on AMPAR synaptic transmission in culture slices. Blue light exposure on transfected cells (with co-expression of paAIP2 and CRISPR-CaMKII α) did not inhibit AMPAR EPSCs ($n = 5$, $p > 0.05$, two-tailed Wilcoxon Signed Rank Test). Mean $\pm$ s.e.m.

Figure S4/ Figure 4-figure suple1. Origin of constitutive CaMKII effect on synaptic transmission in acute slices. (A) Loading cells with 15 mM BAPTA did not alter the CN-27 induced depression of AMPAR synaptic transmission ($n = 5$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test). **(B)** Preinjection of MK801 (i.p. 10 mg/kg body weight) one hour before slicing greatly depressed the NMDAR EPSC resulting in a large increase in the AMPAR/NMDAR ratio (control: 0.72 ± 0.11 , $n = 5$; MK-801: 4.55 ± 0.58 , $n = 5$; $p < 0.01$, Mann-Whitney U test). **(C)** With preinjection of MK-801 one hour before slicing, and in the presence of MK-801 (100 μ M), myr-CN27 inhibited AMPAR synaptic transmission ($n = 5$, $p < 0.01$, two-tailed Wilcoxon Signed Rank Test).

Figure S5/ Figure 6-figure suple1. Reversal of LTP and reduction of synaptic transmission with application of 1 μ M myr-CN27. After a 55 min stable baseline in both pathways (Control = Black, LTP = Red, both n = 1), a train of 100 Hz for 1 s stimuli was applied 4 times with an interval of 20 s. Following the stimulus, the LTP pathway exhibited a rapid increase in fEPSP slope that plateaued at ~140% of baseline. The control pathway did not exhibit this increase. After a stable baseline was achieved for the LTP pathway, 1 μ M myr-CN27 was applied. Following application, both pathways exhibited a decrease in fEPSP slope with the LTP pathway exhibiting a greater decrease. Example traces for both pathways are displayed below the graph and taken at the indicated timepoints (1, 2, 3). Traces are colored according to the pathway they correspond to. The scale bars are vertical 0.1 mV and 5 ms horizontal. Recording done in WT acute slices from P15-28 mice.

Figure1-Source Data File1. CaMKII inhibitor myr-CN27 inhibits AMPAR synaptic transmission through CaMKII α . In this dataset, the results of the effects of myr-CN27 on AMPAR EPSCs and NMDAR EPSCs are included.

Figure2-Source Data File1. CaMKII inhibitor AIP inhibits AMPAR synaptic transmission through CaMKII α . In this dataset, the results of the effects of AIP on AMPAR EPSCs and NMDAR EPSCs are included.

Figure3-Source Data File1. Light-activated CaMKII inhibitor paAIP2 inhibits AMPAR synaptic transmission through CaMKII α . In this dataset, the results of the effects of paAIP2 on AMPAR EPSCs and NMDAR EPSCs are included.

Figure 4-Source Data File1. The depression caused by transient inhibition of CaMKII is long lasting and is absent in cells lacking NMDARs. In this dataset, the results of the effects of myr-CN27 and paAIP2 on AMPAR EPSCs over time are included.

Figure 5-Source Data File1. Transient inhibition of CaMKII enhances subsequent LTP. In this dataset, the results of the effects of LTP induction on control cells and pre-inhibition cells are included.

Figure 6-Source Data File1. Myr-CN27 erases LTP in acute slices. In this dataset, the results of the effects of myr-CN27 on LTP and control pathways are included.

References

- Achterberg, K.G., Buitendijk, G.H., Kool, M.J., Goorden, S.M., Post, L., Slump, D.E., Silva, A.J., van Woerden, G.M., Kushner, S.A., and Elgersma, Y. (2014). Temporal and region-specific requirements of alphaCaMKII in spatial and contextual learning. *J Neurosci* 34, 11180-11187.
- Allen, J.K., Brock, D.J., Kondow-McConaghy, H.M., and Pellois, J.P. (2018). Efficient Delivery of Macromolecules into Human Cells by Improving the Endosomal Escape Activity of Cell-Penetrating Peptides: Lessons Learned from dfTAT and its Analogs. *Biomolecules* 8.
- Amin, J.B., Salussolia, C.L., Chan, K., Regan, M.C., Dai, J., Zhou, H.X., Furukawa, H., Bowen, M.E., and Wollmuth, L.P. (2017). Divergent roles of a peripheral transmembrane segment in AMPA and NMDA receptors. *J Gen Physiol* 149, 661-680.
- Barcomb, K., Hell, J.W., Benke, T.A., and Bayer, K.U. (2016). The CaMKII/GluN2B Protein Interaction Maintains Synaptic Strength. *J Biol Chem* 291, 16082-16089.
- Bayer, K.U., De Koninck, P., Leonard, A.S., Hell, J.W., and Schulman, H. (2001). Interaction with the NMDA receptor locks CaMKII in an active conformation. *Nature* 411, 801-805.
- Bayer, K.U., LeBel, E., McDonald, G.L., O'Leary, H., Schulman, H., and De Koninck, P. (2006). Transition from reversible to persistent binding of CaMKII to postsynaptic sites and NR2B. *J Neurosci* 26, 1164-1174.
- Bayer, K.U., and Schulman, H. (2019). CaM Kinase: Still Inspiring at 40. *Neuron* 103, 380-394.
- Benfenati, F., Valtorta, F., Rubenstein, J.L., Gorelick, F.S., Greengard, P., and Czernik, A.J. (1992). Synaptic vesicle-associated Ca²⁺/calmodulin-dependent protein kinase II is a binding protein for synapsin I. *Nature* 359, 417-420.
- Bhattacharyya, M., Karandur, D., and Kuriyan, J. (2020). Structural Insights into the Regulation of Ca(2+)/Calmodulin-Dependent Protein Kinase II (CaMKII). *Cold Spring Harb Perspect Biol* 12.
- Bhattacharyya, M., Stratton, M.M., Going, C.C., McSpadden, E.D., Huang, Y., Susa, A.C., Elleman, A., Cao, Y.M., Pappireddi, N., Burkhardt, P., *et al.* (2016).

805 Molecular mechanism of activation-triggered subunit exchange in
806 Ca(2+)/calmodulin-dependent protein kinase II. *Elife* 5.

807 Bienenstock, E.L., Cooper, L.N., and Munro, P.W. (1982). Theory for the
808 development of neuron selectivity: orientation specificity and binocular interaction
809 in visual cortex. *J Neurosci* 2, 32-48.

810 Bird, T.D., and Plum, F. (1968). Recovery from barbiturate overdose coma with a
811 prolonged isoelectric electroencephalogram. *Neurology* 18, 456-460.

812 Buard, I., Coultrap, S.J., Freund, R.K., Lee, Y.S., Dell'Acqua, M.L., Silva, A.J.,
813 and Bayer, K.U. (2010). CaMKII "autonomy" is required for initiating but not for
814 maintaining neuronal long-term information storage. *J Neurosci* 30, 8214-8220.

815 Cajal, S.R. (1911). *Histologie du Systeme Nerveux de l'Homme et des Vertebres*.
816 Translated by N. Swanson and L. W. Swanson, Oxford University Press, New
817 York.

818 Chang, B.H., Mukherji, S., and Soderling, T.R. (1998). Characterization of a
819 calmodulin kinase II inhibitor protein in brain. *Proc Natl Acad Sci U S A* 95,
820 10890-10895.

821 Chang, B.H., Mukherji, S., and Soderling, T.R. (2001). Calcium/calmodulin-
822 dependent protein kinase II inhibitor protein: localization of isoforms in rat brain.
823 *Neuroscience* 102, 767-777.

824 Chen, H.X., Otmakhov, N., Strack, S., Colbran, R.J., and Lisman, J.E. (2001). Is
825 persistent activity of calcium/calmodulin-dependent kinase required for the
826 maintenance of LTP? *J Neurophysiol* 85, 1368-1376.

827 Cheriyan, J., Kumar, P., Mayadevi, M., Surolia, A., and Omkumar, R.V. (2011).
828 Calcium/calmodulin dependent protein kinase II bound to NMDA receptor 2B
829 subunit exhibits increased ATP affinity and attenuated dephosphorylation. *PLoS*
830 *One* 6, e16495.

831 Choquet, D. (2018). Linking Nanoscale Dynamics of AMPA Receptor
832 Organization to Plasticity of Excitatory Synapses and Learning. *J Neurosci* 38,
833 9318-9329.

834 Collingridge, G.L., Isaac, J.T., and Wang, Y.T. (2004). Receptor trafficking and
835 synaptic plasticity. *Nat Rev Neurosci* 5, 952-962.

836 Collingridge, G.L., Peineau, S., Howland, J.G., and Wang, Y.T. (2010). Long-
837 term depression in the CNS. *Nat Rev Neurosci* 11, 459-473.

838 Cooper, L.N., and Bear, M.F. (2012). The BCM theory of synapse modification at
839 30: interaction of theory with experiment. *Nat Rev Neurosci* 13, 798-810.

840 Coultrap, S.J., and Bayer, K.U. (2012). CaMKII regulation in information
841 processing and storage. *Trends Neurosci* 35, 607-618.

842 Coultrap, S.J., Freund, R.K., O'Leary, H., Sanderson, J.L., Roche, K.W.,
843 Dell'Acqua, M.L., and Bayer, K.U. (2014). Autonomous CaMKII mediates both
844 LTP and LTD using a mechanism for differential substrate site selection. *Cell*
845 *Rep* 6, 431-437.

846 D'Esposito, M., and Postle, B.R. (2015). The cognitive neuroscience of working
847 memory. *Annu Rev Psychol* 66, 115-142.

848 Davies, S.N., Martin, D., Millar, J.D., Aram, J.A., Church, J., and Lodge, D.
849 (1988). Differences in results from in vivo and in vitro studies on the use-

dependency of N-methylaspartate antagonism by MK-801 and other phencyclidine receptor ligands. *Eur J Pharmacol* 145, 141-151.

Dore, K., Aow, J., and Malinow, R. (2016). The Emergence of NMDA Receptor Metabotropic Function: Insights from Imaging. *Front Synaptic Neurosci* 8, 20.

Feng, T.P. (1995). The involvement of PKC and multifunctional CaM kinase II of the postsynaptic neuron in induction and maintenance of long-term potentiation. *Prog Brain Res* 105, 55-63.

Fusi, S., Drew, P.J., and Abbott, L.F. (2005). Cascade models of synaptically stored memories. *Neuron* 45, 599-611.

Giese, K.P., Fedorov, N.B., Filipkowski, R.K., and Silva, A.J. (1998). Autophosphorylation at Thr286 of the alpha calcium-calmodulin kinase II in LTP and learning. *Science* 279, 870-873.

Goodell, D.J., Zaegel, V., Coultrap, S.J., Hell, J.W., and Bayer, K.U. (2017). DAPK1 Mediates LTD by Making CaMKII/GluN2B Binding LTP Specific. *Cell Rep* 19, 2231-2243.

Goold, C.P., and Nicoll, R.A. (2010). Single-cell optogenetic excitation drives homeostatic synaptic depression. *Neuron* 68, 512-528.

Gouet, C., Aburto, B., Vergara, C., and Sanhueza, M. (2012). On the mechanism of synaptic depression induced by CaMKIIN, an endogenous inhibitor of CaMKII. *PLoS One* 7, e49293.

Granger, A.J., Shi, Y., Lu, W., Cerpas, M., and Nicoll, R.A. (2013). LTP requires a reserve pool of glutamate receptors independent of subunit type. *Nature* 493, 495-500.

Hell, J.W. (2014). CaMKII: claiming center stage in postsynaptic function and organization. *Neuron* 81, 249-265.

Hinds, H.L., Goussakov, I., Nakazawa, K., Tonegawa, S., and Bolshakov, V.Y. (2003). Essential function of alpha-calcium/calmodulin-dependent protein kinase II in neurotransmitter release at a glutamatergic central synapse. *Proc Natl Acad Sci U S A* 100, 4275-4280.

Hinds, H.L., Tonegawa, S., and Malinow, R. (1998). CA1 long-term potentiation is diminished but present in hippocampal slices from alpha-CaMKII mutant mice. *Learn Mem* 5, 344-354.

Huganir, R.L., and Nicoll, R.A. (2013). AMPARs and synaptic plasticity: the last 25 years. *Neuron* 80, 704-717.

Inagaki, H.K., Fontolan, L., Romani, S., and Svoboda, K. (2019). Discrete attractor dynamics underlies persistent activity in the frontal cortex. *Nature* 566, 212-217.

Incontro, S., Diaz-Alonso, J., Iafrati, J., Vieira, M., Asensio, C.S., Sohal, V.S., Roche, K.W., Bender, K.J., and Nicoll, R.A. (2018). The CaMKII/NMDA receptor complex controls hippocampal synaptic transmission by kinase-dependent and independent mechanisms. *Nat Commun* 9, 2069.

Ishida, A., Shigeri, Y., Tatsu, Y., Uegaki, K., Kameshita, I., Okuno, S., Kitani, T., Yumoto, N., and Fujisawa, H. (1998). Critical amino acid residues of AIP, a highly specific inhibitory peptide of calmodulin-dependent protein kinase II. *FEBS Lett* 427, 115-118.

895 Kennedy, M.B. (2013). Synaptic Signaling in Learning and Memory. Cold Spring
 896 Harb Perspect Biol 8, a016824.
 897 LeCher, J.C., Nowak, S.J., and McMurry, J.L. (2017). Breaking in and busting
 898 out: cell-penetrating peptides and the endosomal escape problem. Biomol
 899 Concepts 8, 131-141.
 900 Lee, S.J., Escobedo-Lozoya, Y., Szatmari, E.M., and Yasuda, R. (2009).
 901 Activation of CaMKII in single dendritic spines during long-term potentiation.
 902 Nature 458, 299-304.
 903 Leonard, A.S., Lim, I.A., Hemsworth, D.E., Horne, M.C., and Hell, J.W. (1999).
 904 Calcium/calmodulin-dependent protein kinase II is associated with the N-methyl-
 905 D-aspartate receptor. Proc Natl Acad Sci U S A 96, 3239-3244.
 906 Lisman, J. (2017). Criteria for identifying the molecular basis of the engram
 907 (CaMKII, PKMzeta). Mol Brain 10, 55.
 908 Lisman, J., and Raghavachari, S. (2015). Biochemical principles underlying the
 909 stable maintenance of LTP by the CaMKII/NMDAR complex. Brain Res 1621, 51-
 910 61.
 911 Lisman, J., Schulman, H., and Cline, H. (2002). The molecular basis of CaMKII
 912 function in synaptic and behavioural memory. Nat Rev Neurosci 3, 175-190.
 913 Lisman, J., Yasuda, R., and Raghavachari, S. (2012). Mechanisms of CaMKII
 914 action in long-term potentiation. Nat Rev Neurosci 13, 169-182.
 915 Lisman, J.E., and Goldring, M.A. (1988). Feasibility of long-term storage of
 916 graded information by the Ca²⁺/calmodulin-dependent protein kinase molecules
 917 of the postsynaptic density. Proc Natl Acad Sci U S A 85, 5320-5324.
 918 Lisman, J.E., and Zhabotinsky, A.M. (2001). A model of synaptic memory: a
 919 CaMKII/PP1 switch that potentiates transmission by organizing an AMPA
 920 receptor anchoring assembly. Neuron 31, 191-201.
 921 Lledo, P.M., Hjelmstad, G.O., Mukherji, S., Soderling, T.R., Malenka, R.C., and
 922 Nicoll, R.A. (1995). Calcium/calmodulin-dependent kinase II and long-term
 923 potentiation enhance synaptic transmission by the same mechanism. Proc Natl
 924 Acad Sci U S A 92, 11175-11179.
 925 Malenka, R.C., and Bear, M.F. (2004). LTP and LTD: an embarrassment of
 926 riches. Neuron 44, 5-21.
 927 Malenka, R.C., Kauer, J.A., Perkel, D.J., Mauk, M.D., Kelly, P.T., Nicoll, R.A.,
 928 and Waxham, M.N. (1989). An essential role for postsynaptic calmodulin and
 929 protein kinase activity in long-term potentiation. Nature 340, 554-557.
 930 Malinow, R., and Malenka, R.C. (2002). AMPA receptor trafficking and synaptic
 931 plasticity. Annu Rev Neurosci 25, 103-126.
 932 Malinow, R., Schulman, H., and Tsien, R.W. (1989). Inhibition of postsynaptic
 933 PKC or CaMKII blocks induction but not expression of LTP. Science 245, 862-
 934 866.
 935 Maravall, M., Mainen, Z.F., Sabatini, B.L., and Svoboda, K. (2000). Estimating
 936 intracellular calcium concentrations and buffering without wavelength ratioing.
 937 Biophys J 78, 2655-2667.
 938 Miller, S.G., and Kennedy, M.B. (1986). Regulation of brain type II
 939 Ca²⁺/calmodulin-dependent protein kinase by autophosphorylation: a Ca²⁺-
 940 triggered molecular switch. Cell 44, 861-870.

941 Morrison, A., Diesmann, M., and Gerstner, W. (2008). Phenomenological models
 942 of synaptic plasticity based on spike timing. *Biol Cybern* 98, 459-478.
 943 Mulkey, R.M., Endo, S., Shenolikar, S., and Malenka, R.C. (1994). Involvement
 944 of a calcineurin/inhibitor-1 phosphatase cascade in hippocampal long-term
 945 depression. *Nature* 369, 486-488.
 946 Mulkey, R.M., Herron, C.E., and Malenka, R.C. (1993). An essential role for
 947 protein phosphatases in hippocampal long-term depression. *Science* 261, 1051-
 948 1055.
 949 Mullasseril, P., Dosemeci, A., Lisman, J.E., and Griffith, L.C. (2007). A structural
 950 mechanism for maintaining the 'on-state' of the CaMKII memory switch in the
 951 post-synaptic density. *J Neurochem* 103, 357-364.
 952 Murakoshi, H., Shin, M.E., Parra-Bueno, P., Szatmari, E.M., Shibata, A.C.E., and
 953 Yasuda, R. (2017). Kinetics of Endogenous CaMKII Required for Synaptic
 954 Plasticity Revealed by Optogenetic Kinase Inhibitor. *Neuron* 94, 690.
 955 Nelson, A.R., Borland, L., Allbritton, N.L., and Sims, C.E. (2007). Myristoyl-based
 956 transport of peptides into living cells. *Biochemistry* 46, 14771-14781.
 957 Nicoll, R.A. (2017). A Brief History of Long-Term Potentiation. *Neuron* 93, 281-
 958 290.
 959 Otmakhov, N., Griffith, L.C., and Lisman, J.E. (1997). Postsynaptic inhibitors of
 960 calcium/calmodulin-dependent protein kinase type II block induction but not
 961 maintenance of pairing-induced long-term potentiation. *J Neurosci* 17, 5357-5365.
 962 Özden, C., Santos, N., Agnello, E., Gaubitz, C., Lapinskas, E., and al., e. (2020).
 963 CaMKII binds both substrates and effectors at the active site bioRxiv:
 964 <https://doi.org/10.1101/20201025354241>.
 965 Patel, S.G., Sayers, E.J., He, L., Narayan, R., Williams, T.L., Mills, E.M.,
 966 Allemann, R.K., Luk, L.Y.P., Jones, A.T., and Tsai, Y.H. (2019). Cell-penetrating
 967 peptide sequence and modification dependent uptake and subcellular distribution
 968 of green florescent protein in different cell lines. *Sci Rep* 9, 6298.
 969 Pellicena, P., and Schulman, H. (2014). CaMKII inhibitors: from research tools to
 970 therapeutic agents. *Front Pharmacol* 5, 21.
 971 Petersen, J.D., Chen, X., Vinade, L., Dosemeci, A., Lisman, J.E., and Reese, T.S.
 972 (2003). Distribution of postsynaptic density (PSD)-95 and Ca²⁺/calmodulin-
 973 dependent protein kinase II at the PSD. *J Neurosci* 23, 11270-11278.
 974 Pettit, D.L., Perlman, S., and Malinow, R. (1994). Potentiated transmission and
 975 prevention of further LTP by increased CaMKII activity in postsynaptic
 976 hippocampal slice neurons. *Science* 266, 1881-1885.
 977 Pi, H.J., Otmakhov, N., Lemelin, D., De Koninck, P., and Lisman, J. (2010).
 978 Autonomous CaMKII can promote either long-term potentiation or long-term
 979 depression, depending on the state of T305/T306 phosphorylation. *J Neurosci* 30,
 980 8704-8709.
 981 Poncer, J.C., Esteban, J.A., and Malinow, R. (2002). Multiple mechanisms for the
 982 potentiation of AMPA receptor-mediated transmission by alpha-Ca²⁺/calmodulin-
 983 dependent protein kinase II. *J Neurosci* 22, 4406-4411.
 984 Rossetti, T., Banerjee, S., Kim, C., Leubner, M., Lamar, C., Gupta, P., Lee, B.,
 985 Neve, R., and Lisman, J. (2017). Memory Erasure Experiments Indicate a Critical
 986 Role of CaMKII in Memory Storage. *Neuron* 96, 207-216 e202.

987 Sanhueza, M., Fernandez-Villalobos, G., Stein, I.S., Kasumova, G., Zhang, P.,
 988 Bayer, K.U., Otmakhov, N., Hell, J.W., and Lisman, J. (2011). Role of the
 989 CaMKII/NMDA receptor complex in the maintenance of synaptic strength. *J*
 990 *Neurosci* 31, 9170-9178.
 991 Sanhueza, M., and Lisman, J. (2013). The CaMKII/NMDAR complex as a
 992 molecular memory. *Mol Brain* 6, 10.
 993 Sanhueza, M., McIntyre, C.C., and Lisman, J.E. (2007). Reversal of synaptic
 994 memory by Ca²⁺/calmodulin-dependent protein kinase II inhibitor. *J Neurosci* 27,
 995 5190-5199.
 996 Scanziani, M., Malenka, R.C., and Nicoll, R.A. (1996). Role of intercellular
 997 interactions in heterosynaptic long-term depression. *Nature* 380, 446-450.
 998 Schnell, E., Sizemore, M., Karimzadegan, S., Chen, L., Brecht, D.S., and Nicoll,
 999 R.A. (2002). Direct interactions between PSD-95 and stargazin control synaptic
 1000 AMPA receptor number. *Proc Natl Acad Sci U S A* 99, 13902-13907.
 1001 Schulman, H. (1984). Phosphorylation of microtubule-associated proteins by a
 1002 Ca²⁺/calmodulin-dependent protein kinase. *J Cell Biol* 99, 11-19.
 1003 Silva, A.J., Stevens, C.F., Tonegawa, S., and Wang, Y. (1992). Deficient
 1004 hippocampal long-term potentiation in alpha-calcium-calmodulin kinase II mutant
 1005 mice. *Science* 257, 201-206.
 1006 Stein, I.S., Park, D.K., Claiborne, N., and Zito, K. (2021). Non-ionotropic NMDA
 1007 receptor signaling gates bidirectional structural plasticity of dendritic spines. *Cell*
 1008 *Rep* 34, 108664.
 1009 Stevens, C.F., Tonegawa, S., and Wang, Y. (1994). The role of calcium-
 1010 calmodulin kinase II in three forms of synaptic plasticity. *Curr Biol* 4, 687-693.
 1011 Stoppini, L., Buchs, P.A., and Muller, D. (1991). A simple method for organotypic
 1012 cultures of nervous tissue. *J Neurosci Methods* 37, 173-182.
 1013 Strack, S., Barban, M.A., Wadzinski, B.E., and Colbran, R.J. (1997a). Differential
 1014 inactivation of postsynaptic density-associated and soluble Ca²⁺/calmodulin-
 1015 dependent protein kinase II by protein phosphatases 1 and 2A. *J Neurochem* 68,
 1016 2119-2128.
 1017 Strack, S., Choi, S., Lovinger, D.M., and Colbran, R.J. (1997b). Translocation of
 1018 autophosphorylated calcium/calmodulin-dependent protein kinase II to the
 1019 postsynaptic density. *J Biol Chem* 272, 13467-13470.
 1020 Stratton, M., Lee, I.H., Bhattacharyya, M., Christensen, S.M., Chao, L.H.,
 1021 Schulman, H., Groves, J.T., and Kuriyan, J. (2014). Activation-triggered subunit
 1022 exchange between CaMKII holoenzymes facilitates the spread of kinase activity.
 1023 *Elife* 3, e01610.
 1024 Toyozumi, T., Kaneko, M., Stryker, M.P., and Miller, K.D. (2014). Modeling the
 1025 dynamic interaction of Hebbian and homeostatic plasticity. *Neuron* 84, 497-510.
 1026 Trinidad, J.C., Thalhammer, A., Specht, C.G., Schoepfer, R., and Burlingame,
 1027 A.L. (2005). Phosphorylation state of postsynaptic density proteins. *J Neurochem*
 1028 92, 1306-1316.
 1029 Turrigiano, G.G. (2008). The self-tuning neuron: synaptic scaling of excitatory
 1030 synapses. *Cell* 135, 422-435.
 1031 Vest, R.S., Davies, K.D., O'Leary, H., Port, J.D., and Bayer, K.U. (2007). Dual
 1032 mechanism of a natural CaMKII inhibitor. *Mol Biol Cell* 18, 5024-5033.

1033 Wang, J.H., and Kelly, P.T. (1996). The balance between postsynaptic Ca(2+)-
1034 dependent protein kinase and phosphatase activities controlling synaptic strength.
1035 Learn Mem 3, 170-181.
1036 Wang, X.J. (2001). Synaptic reverberation underlying mnemonic persistent
1037 activity. Trends Neurosci 24, 455-463.
1038 Wong, J.M., and Gray, J.A. (2018). Long-Term Depression Is Independent of
1039 GluN2 Subunit Composition. J Neurosci 38, 4462-4470.
1040

Fig.1

Fig. S1

WT vs Various Peptides AMPA/NMDA Ratio

Fig.2

Fig.3

Fig. S2

Before light

A

B

After light

C

D

Fig. S3

Fig.4

Fig. S4

Fig.5

Fig.6

Fig. S5