1 Molecular determinants of phase separation for *Drosophila* DNA replication licensing factors

2
3 Matthew W. Parker^{1#}, Jonchee Kao^{2#}, Alvin Huang^{2#}, James M. Berger³, Michael R. Botchan²

4
5 [#]These authors contributed equally to this work.

6 ¹Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, Texas 75235,
7 USA.

8 ²Department of Molecular and Cell Biology, University of California Berkeley, Berkeley, California
9 94720, USA.

10 ³Department of Biophysics and Biophysical Chemistry, Johns Hopkins School of Medicine, Baltimore,
11 Maryland 21205, USA.

12 *Correspondence should be addressed to: matthew.parker@utsouthwestern.edu

13
14 **Keywords:** DNA replication, liquid phase condensates, intrinsically disordered proteins, DNA binding
15 proteins

ABSTRACT

Liquid-liquid phase separation (LLPS) of intrinsically disordered regions (IDRs) in proteins can drive the formation of membraneless compartments in cells. Phase-separated structures enrich for specific partner proteins and exclude others. Previously, we showed that the IDRs of metazoan DNA replication initiators drive DNA-dependent phase separation *in vitro* and chromosome binding *in vivo*, and that initiator condensates selectively recruit replication-specific partner proteins (Parker et al., 2019). How initiator IDRs facilitate LLPS and maintain compositional specificity is unknown. Here, using *D. melanogaster* (*Dm*) Cdt1 as a model initiation factor, we show that phase separation results from a synergy between electrostatic DNA-bridging interactions and hydrophobic inter-IDR contacts. Both sets of interactions depend on sequence composition (but not sequence order), are resistant to 1,6-hexanediol, and do not depend on aromaticity. These findings demonstrate that distinct sets of interactions drive condensate formation and specificity across different phase-separating systems and advance efforts to predict IDR LLPS propensity and partner selection *a priori*.

INTRODUCTION

The internal environment of a cell is compartmentalized both by membrane-bound organelles and by protein-rich and protein/nucleic acid-rich compartments that lack an enclosing membrane. These membraneless compartments are commonly called biomolecular condensates and form through an ability of their constituent factors to undergo liquid-liquid phase separation (**LLPS**) (reviewed in (Banani et al., 2017)). Cellular bodies that form by LLPS are often spherical due to surface tension minimization across the phase boundary (Elbaum-Garfinkle, 2019). Sphericity, however, is not a defining feature of biomolecular condensates and many variables can contribute to droplet (de)formation, including the physical parameters of the liquid phase (e.g., surface tension, viscosity and size) and external variables (e.g., application of a force and surface interactions). Non-spherical condensates often nucleate from or assemble along relatively rigid intracellular scaffolds. This effect is seen with cellular signaling complexes (Banjade and Rosen, 2014; Li et al., 2012; Su et al., 2016) and tight junctions (Beutel et al., 2019), which spread with the dimensions of the plasma membrane. Likewise, the chromatin-associated synaptonemal complex (Rog et al., 2017) and the perichromosomal layer (Booth and Earnshaw, 2017) are predicted to form a protein-rich phase that

contours with the more rigid nature of their substrate, mitotic chromosomes (Batty and Gerlich, 2019; Goloborodko et al., 2016; Houlard et al., 2015; Sun et al., 2018).

We previously discovered that the factors responsible for initiating DNA replication in metazoans, in particular the fly homologs of the Origin Recognition Complex (ORC) and Cdc6, as well as fly and human Cdt1, undergo DNA-dependent LLPS at physiological salt and protein concentrations *in vitro* (Parker et al., 2019). Recent studies have also demonstrated that human Orc1 and Cdc6 can phase-separate in the presence of DNA (Hossain et al., 2021). In live cells, initiators first associate with chromatin in mitosis where they appear to uniformly coat, or “wet”, anaphase chromosomes (Baldinger and Gossen, 2009; Kara et al., 2015; Parker et al., 2019; Sonnevile et al., 2012). These cytological observations are consistent with expectations for a protein that undergoes a condensation reaction with a relatively rigid intracellular scaffolding partner. The ability to condense on DNA is conferred by a metazoan-specific, N-terminal intrinsically disordered region (IDR) present in Orc1, Cdc6, and Cdt1. For Orc1, this region is required for binding chromatin in tissue culture cells and for viability in flies (Parker et al., 2019). *In vitro* assays show that the Orc1 IDR is dispensable for ATP-dependent DNA-binding by *D. melanogaster* ORC and loading of the Mcm2-7 replicative helicase (Schmidt and Bleichert, 2020). However, at physiological ionic strength and protein concentrations, initiator IDRs drive initiator coalescence into a condensed phase that both supports ATP-dependent recruitment of Mcm2-7 and can be controlled by CDK-dependent phosphorylation (Parker et al., 2019), an event that directly regulates initiator mechanism (Findeisen et al., 1999; Lee et al., 2012). Together, the available data suggest that LLPS facilitates the formation of an enriched layer of initiation factors that coats mitotic chromosomes to enhance the kinetics of helicase loading, a process that is completed within a relatively narrow window of the eukaryotic cell cycle (Dimitrova et al., 2002; Méndez and Stillman, 2000; Okuno et al., 2001).

How metazoan initiator IDRs facilitate DNA-dependent LLPS at a molecular level is unknown. Protein multivalency underlies the formation of intermolecular interaction networks that can drive LLPS. In many phase-separating systems, functional multivalency is contributed by the presence of one or more IDRs (Li et al., 2012). Although phase-separating IDR sequences vary between systems, some sequence features capable of supporting LLPS have begun to emerge. For example, many phase-

separating IDRs have low amino acid sequence complexity (so-called low complexity domains (LCDs)) and are highly enriched for a few select amino acids (e.g., the nucleolar protein FIB1 (Feric et al., 2016), the P-granule protein Laf-1 (Elbaum-Garfinkle et al., 2015), the stress granule protein hnRNPA1 (Molliex et al., 2015), the transcription factor EWS (Chong et al., 2018), and the transcriptional coactivator MED1 (Sabari et al., 2018)). However, there exist important exceptions to this correlation, including phase separation by the intrinsically disordered Nephrin Intracellular Domain (NICD) (Pak et al., 2016) and by metazoan replication initiators (Parker et al., 2019), both of which possess high sequence-complexity IDRs. Thus, LCDs are but a subtype of phase-separating IDR.

Literature reports of phase-separating sequences are increasing rapidly, yet relatively few of these studies define the molecular basis for LLPS. An important goal of the field is to understand and ultimately predict the extent to which different disordered protein regions utilize a generally shared or individually distinctive set of cohesive interactions to drive phase separation. Interestingly, many condensates (Table 1) dissolve upon treatment with the aliphatic alcohol 1,6-hexanediol (1,6-HD), a compound that permeabilizes the nuclear pore by disrupting weak hydrophobic Phe-Gly ('FG')-repeat interactions that are prevalent in the assembly (Patel et al., 2007; Shulga and Goldfarb, 2003). The broad sensitivity of protein LLPS to 1,6-HD could be taken to indicate that phase-separation mechanisms are generalizable, at least for certain IDR classes (Kroschwald et al., 2017). Consistently, IDR sequence aromaticity has been shown to be an essential feature in a variety of phase-separating systems (Chiu et al., 2020; Chong et al., 2018; Lin et al., 2017; Nott et al., 2015; Pak et al., 2016; Qamar et al., 2018; Wang et al., 2018). Nevertheless, other interaction types appear to predominate in other LLPS systems. For example, charged residues can facilitate phase separation either through homomeric protein-protein interactions (Elbaum-Garfinkle et al., 2015; Mitrea et al., 2018; Nott et al., 2015) or through heteromeric protein-protein (Ferrolino et al., 2018; Pak et al., 2016) and protein-nucleic acid interactions (Lin et al., 2015; Zhang et al., 2017). We note also that membraneless organelles *in vivo* have a particular compositional bias that often underpins their utility, implying that there exists a molecular 'grammar' or 'sorting code' that accepts a particular subset of partner factors while excluding inappropriate factors. Consistent with this idea, we have observed that initiator IDRs can co-recruit pathway-specific partner proteins but not other types of

phase-separating sequences (such as human Fused in Sarcoma (FUS)) (Parker et al., 2019). An understanding of the molecular rules that govern IDR sorting mechanisms is still in its infancy.

Here, we use *Drosophila melanogaster* Cdt1 as a model to dissect the molecular basis for DNA-dependent LLPS by metazoan replication initiator proteins. We find that phase separation by Cdt1 is unaffected by treatment with 1,6-hexanediol and that sequence aromatics, which represent < 3% of the total initiator IDR amino acids, are dispensable for condensate formation. We show that protein-DNA interactions are electrostatic in nature and contribute an adhesive force to LLPS, with DNA functioning as a counterion bridge. These interactions synergize with cohesive intermolecular IDR-IDR interactions, which are driven by the hydrophobic effect. Using mutagenesis, we demonstrate that intermolecular IDR-IDR interactions serve as the primary driving force behind initiator phase separation, with DNA-mediated bridging interactions playing a secondary but still facilitative role. Collectively, these studies provide a detailed picture of the biochemical mechanisms underlying IDR LLPS in replication initiation factors and reinforce the concept that biomolecular condensates can form by a variety of different molecular interaction types.

RESULTS

Initiator liquid-liquid phase separation is resistant to treatment with 1,6-hexanediol and does not require aromatic residues.

The IDRs of the metazoan replication initiation factors are necessary and sufficient for DNA-dependent LLPS (Parker et al., 2019). To understand the mechanisms that promote initiator condensation, we first set out to identify similarities in amino acid sequence composition between initiator IDRs and the IDRs of other condensate forming proteins. We generated a sequence heatmap to compare the fractional representation of each amino acid in the IDRs of Orc1 (residues 187-549), Cdc6 (residues 1-246), and Cdt1 (residues 1-297) with the IDRs of a small suite of known condensate-forming proteins (human FUS (residues 1-293), Ddx4 (residues 1-260), hnRNPA1 (residues 178-372), DNA-directed RNA polymerase II subunit RPB1 (residues 1531-1970), and Mediator of RNA polymerase II transcription subunit 1 (MED1) (residues 948-1574), *C. elegans* Laf-1 (residues 1-203), and mouse BuGZ (residues 63-495)) (Figure 1A). This analysis revealed major differences in the

amino acid composition of initiator IDRs as compared to the other proteins analyzed, demonstrating that initiator IDRs represent an independent class of sequences. The most striking difference was the conspicuous absence of any one or more highly enriched residue within the initiator IDRs (dark red = fraction composition > 15%). All other sequences in the comparison suite possess at least one highly enriched residue, most often serine, proline, and/or glycine. By contrast, initiator IDR composition is distributed more uniformly across all amino acids, apart from histidine, cysteine, and tryptophan side chains (which are weakly represented amongst all IDRs assessed), and methionine, glycine and aromatic residues (which are weakly represented in initiators but not necessarily other IDRs). Extending our heatmap analysis to a more diverse set of phase-separating disordered sequences (Figure 1 – figure supplement 1A) revealed a similar trend, with most sequences being highly enriched in a small subset of amino acids. However, there are a minority of IDRs that show a more unbiased sequence composition similar to that of replication initiators, including the human Nephhrin Intracellular Domain (NICD) (Pak et al., 2016), TACC3 (So et al., 2019), and Tau (Kanaan et al., 2020).

The broad utilization of amino acid sequence space by initiator IDRs suggested that these sequences have higher complexity than many other phase separating proteins. To quantitatively assess complexity, we calculated the informational entropy of each IDR sequence. In these calculations, entropy represents the amount of information stored in a given linear sequence of amino acids as derived from the number of observed occurrences of each of the twenty amino acids, and has a theoretical range of 0-3 (Wootton and Federhen, 1993) (Figure 1-figure supplement 1B). This analysis revealed that initiator IDRs have a higher sequence complexity (score = 2.7) than most other condensate-forming IDRs analyzed (14 of 17 total sequences). The human phase separating proteins Ddx4 (score = 2.7) (Figure 1A), NICD (score = 2.8), TACC3 (score = 2.7) and Tau (score = 2.7) have similar complexity scores while still maintaining a unique sequence composition (Figure 1-figure supplement 1A). To contextualize these complexity calculations, we calculated the sequence complexity of all predicted disordered (n = 3178) and ordered (n = 10,884) sequences longer than 150 amino acids within the *Drosophila* proteome. Complexity scores for both classes of sequences adopted a negatively skewed distribution, with ordered sequences narrowly centered around 2.9 and disordered sequences more broadly distributed around a score of 2.7. (Figure 1-figure supplement 1C).

These data reveal that phase separating sequences show high variability in complexity and that initiator-type IDRs are amongst the most complex phase-separating sequences. The unique sequence features (i.e., composition and complexity) of initiator IDRs was a first indication that initiator LLPS might proceed through a mechanism distinct from many other model phase-separating proteins.

The aliphatic alcohol 1,6-hexanediol (1,6-HD) can inhibit the self-assembly and phase separation of compositionally distinct classes of disordered sequences (see [Table 1](#) references). This behavior suggests that even for different IDR sequences, many use similar types of molecular interactions to drive LLPS. We therefore tested whether condensates formed from initiator IDRs are likewise sensitive to 1,6-HD. Initiator LLPS was directly visualized by fluorescence microscopy by individually mixing ORC, Cdc6, or Cdt1 with a Cy5-labeled 60 bp dsDNA (Cy5-dsDNA). LLPS by all three proteins proved to be resistant to treatment with 10% 1,6-HD ([Figure 1B](#) and [Figure 1-figure supplement 2](#)), although 1,6-HD did lead to morphological changes in initiator/DNA condensates, including droplet clustering and, for ORC, a reduction in overall size. To confirm the microscopy results, we assessed the effect of 1,6-HD on Cdt1 partitioning into a condensed phase by a solution depletion assay. In this approach, LLPS is induced by the addition of dsDNA (60 bp), the denser phase separated material is pelleted by centrifugation, and LLPS is assessed by protein depletion from the supernatant. Titrations of 1,6-HD from 0-10% (w/v) showed no ability to inhibit Cdt1 phase separation ([Figure 1C](#)), confirming that initiator phase separation is insensitive to this reagent.

Table 1: List of proteins that form liquid phase condensates sensitive to 1,6-hexanediol.

Protein	Reference(s)
TDP43	(Babinchak et al., 2019; Schmidt et al., 2019)
hnRNPA1	(Molliex et al., 2015)
Tau	(Wegmann et al., 2018)
Huntingtin protein exon 1	(Peskett et al., 2018)
Rnq1	(Kroschwald et al., 2015)
BRD4	(Sabari et al., 2018)
MED1	(Cho et al., 2018; Sabari et al., 2018)
Hp1	(Strom et al., 2017)
hnRNPA2	(Lin et al., 2016)
TACC3	(So et al., 2019)
FUS	(Kroschwald et al., 2017)
RPB1	(Boehning et al., 2018)

Chromosome passenger complex	(Trivedi et al., 2019)
Synaptonemal complex	(Rog et al., 2017)

Aromatic residues can play a critical role in driving protein LLPS, where they contribute multivalent π - π and π -cation interactions (Figure 1D) (Chiu et al., 2020; Chong et al., 2018; Lin et al., 2017; Nott et al., 2015; Pak et al., 2016; Qamar et al., 2018; Vernon et al., 2018; Wang et al., 2018). However, the resistance of initiator LLPS to treatment with 1,6-HD (Figure 1B-C), as well as the low (< 3%) aromatic residue content of initiator IDRs, suggested that these interactions may not be a primary driving force for phase separation. The IDR of Cdt1 contains only eight aromatic residues, all phenylalanine, that are relatively equally distributed throughout the region (Figure 1E). To test whether initiator LLPS is influenced by these aromatic residues, we mutated all phenylalanines in the Cdt1 IDR to leucine (Cdt1^{Phe→Leu}) and assessed phase separation by both the depletion assay (Figure 1F) and fluorescence microscopy (Figure 1G, top panel). The replacement of Phe with Leu abolishes aromaticity in the IDR while maintaining a comparable level of hydrophobicity (phenylalanine hydrophobicity index = 2.8, leucine hydrophobicity index = 3.8, (Kyte and Doolittle, 1982)). Interestingly, the loss of aromaticity had no detectable effect on DNA-induced phase separation propensity. Mutating Cdt1's aromatic residues to alanine (Cdt1^{Phe→Ala}) similarly did not block phase separation (Figure 1G, bottom panel). Together, these data show that initiator phase separation does not rely on aromatic residue-mediated interactions, consistent with its ability to resist treatment with 1,6-HD.

Initiator IDR electrostatics promote coacervation with polyanionic scaffolds.

The DNA dependency of initiator LLPS suggested that initiator droplets form by coacervation, a process in which two oppositely charged polymers drive the formation of a condensed phase (Sing and Perry, 2020). We therefore set out to test the sensitivity of Cdt1 LLPS to salt. Using the depletion assay, we measured phase separation at different concentrations of potassium glutamate (KGlu; 75, 150 and 300 mM) (Figure 2A). We observed Cdt1 LLPS at or below physiological levels of salt (150 mM) but a loss of condensation above this value. These data demonstrate that electrostatic interactions are important for driving DNA-dependent phase separation by initiator proteins; however, it remained unclear whether specific Cdt1/DNA interactions, such as major- and minor-

groove binding or ring-stacking interactions, might also be necessary for LLPS. To address this question, we assessed whether other anionic polyelectrolytes of distinct chemical composition could drive condensation. Double-stranded (ds) and single-stranded DNA (ssDNA), dsRNA, heparin, and poly-glutamate (each at 30 $\mu\text{g/mL}$) were combined with 2 μM Cdt1 and phase separation was assessed with the depletion assay (Figure 2B). All polyanions tested showed an equal propensity to induce Cdt1 phase separation, strongly supporting the notion that non-specific electrostatic interactions are a significant driving force underlying DNA-induced initiator phase separation.

In principle, two mechanisms could account for the dependence of Cdt1 LLPS on the presence of a counterion (Figure 2C). First, a counterion may function as an intermolecular bridge to contribute adhesive forces that help promote initiator LLPS. Alternatively, electrostatic repulsion between the positive charges on the Cdt1 IDR might prevent the spontaneous formation of inter-IDR interactions and LLPS, and a counterion could help neutralize this repulsive force. To distinguish between these mechanisms, we assayed phase separation by Cdt1 in the presence of mono- and multivalent counterions, with net charge ranging from -1 (KH_2PO_4) to -3 (K_2HPO_4) (Figure 2D). We reasoned that if counterions simply reduce electrostatic repulsion, then the titration of a monovalent salt would be sufficient to shield the IDR's positive charge and drive phase separation. However, if counterions provide an intermolecular "bridge" between basic residues within the Cdt1 IDR, then multivalent salts would be required to drive phase separation. Using the depletion assay we found that monobasic potassium phosphate (pH 7.5) was unable to induce phase separation of Cdt1 up to the highest concentration tested (625 mM). Conversely, we observed significant depletion of Cdt1 at the highest concentration of dibasic potassium phosphate (625 mM, pH 7.5), as well as for the three highest concentrations of tribasic potassium phosphate (400, 500 and 625 mM, pH 7.5). eGFP-tagged Cdt1 (eGFP-Cdt1) and fluorescence microscopy were used to confirm the depletion assay results. In the presence of either 625 mM mono- or dibasic potassium phosphate, eGFP-Cdt1 appeared monodisperse; however, tribasic potassium phosphate induced formation of a concentrated eGFP-Cdt1 species. Interestingly, this species was morphologically distinct from the round droplets of Cdt1 that are observed in the presence of DNA and appeared more akin to aggregates than to droplets. These data indicate that the physiochemistry of a complexing polyanion can shape the properties of initiator condensates, consistent with previous observations for the complex coacervation of

poly(proline-arginine) peptides (Boeynaems et al., 2019). Altogether, our findings strongly indicate that counterions contribute an adhesive force to initiator self-assembly, bridging between the IDRs of different Cdt1 protomers.

In addition to overall net charge, recent theoretical and experimental work has demonstrated that the patterning of charged residues within disordered domains can fine-tune phase separation propensity (Lin et al., 2018; Nott et al., 2015; Pak et al., 2016; Paloni et al., 2020). We assessed the distribution of charged residues within the Cdt1 IDR by calculating the net charge per residue (NCPR) over a 5-residue window (Figure 2E, top). Ionic residues within the Cdt1 IDR cluster into local regions with net positive and net negative charge. We predicted that if charge distribution were important for initiator phase separation, then charge patterning would be conserved across metazoan initiator IDRs. To test this idea, we performed a kappa value analysis of initiator IDRs. Kappa is a parameter between 0-1 that quantitatively describes the degree of mixing between oppositely charged residues within a sequence (well-mixed samples have a low kappa) and serves as a predictor of IDR conformational state (extended or collapsed) (Das and Pappu, 2013). Given the absence of sequence identity across metazoan initiator IDRs (Cdt1, Orc1, and Cdc6), we were surprised to find that the kappa values for metazoan initiator IDRs are similar (range 0.20-0.29), as are other physiochemical parameters (e.g., the overall fraction of charged residues, or FCR (range 0.22-0.32) and the isoelectric point, or pI (range 9.4-11.1)). None of these properties are preserved in budding yeast initiation factors (Table 2), which lack IDRs and do not undergo LLPS (Parker et al., 2019).

Table 2: Charged residue properties of metazoan and yeast initiator IDRs.

Species	Protein	FCR	Kappa	pI
<i>Fruit fly</i>	Orc1	0.30	0.23	10.2
	Cdc6	0.31	0.25	9.4
	Cdt1	0.30	0.20	10.1
<i>Human</i>	Orc1	0.31	0.21	10.6
	Cdc6	0.25	0.21	10.6
	Cdt1	0.28	0.26	10.6
<i>Frog</i>	Orc1	0.32	0.29	9.7
	Cdc6	0.25	0.21	10.8
	Cdt1	0.30	0.20	10.1
<i>Zebrafish</i>	Orc1	0.29	0.30	9.8

	Cdc6	0.22	0.20	11.1
	Cdt1	0.29	0.20	9.9
	Orc1	0.48	0.43	4.7
<i>Budding yeast</i>	Cdc6	0.32	0.59	6.3
	Cdt1 (no IDR)	na	na	na
	Orc1	0.32	0.37	10.6
<i>Fission yeast</i>	Cdc6	0.21	0.21	10.2
	Cdt1 (no IDR)	na	na	na

To directly test whether the distribution of charged residues within the Cdt1 IDR are important for its ability to form condensates, we produced a Cdt1 variant, Cdt1^{Uniform}, in which charged residues (K, R, D, and E) were uniformly distributed across the IDR sequence ($\kappa = 0.06$), but where sequence composition and pI were maintained as per the wild-type protein. A plot of NCPR for Cdt1^{Uniform} demonstrates the loss of regions with high or low local net charge (Figure 2E, bottom). To quantitatively compare the phase separation propensity of Cdt1^{Uniform} with that of wild-type Cdt1, we generated phase diagrams for these proteins, assaying the ability of Cy5-dsDNA to induce phase separation over a range of protein (40-5000 nM) and KGlutamate (50-300 mM) concentrations that span physiological conditions (*in vivo*, [Cdt1] $\approx$ 100 nM (Parker et al., 2019) and [salt] $\approx$ 150 mM (Andersen, 2013; Park et al., 2016)) (Figure 2F-G). Wild-type Cdt1 phase separation was observed down to 200 nM protein at [KGlutamate] = 150 mM. Higher concentrations of Cdt1 resulted in phase separation at all concentrations of KGlutamate tested (Figure 2F). Surprisingly, Cdt1^{Uniform} was still highly effective at inducing phase separation, although its critical concentration at [KGlutamate] = 150 mM was reduced to 1 μ M and it was more sensitive to increasing concentrations of KGlutamate (Figure 2G). Quantitation of mean droplet intensity at 5 μ M protein concentrations revealed a maximal Cy5-dsDNA signal at 150 mM KGlutamate for both proteins and an overall reduction in droplet intensity for Cdt1^{Uniform} (Figure 2G and Figure 2-figure supplement 1). These results demonstrate that charge patterning in Cdt1 is not strictly required for phase separation, but that the charged residue distribution observed in the wild-type sequence favors lower LLPS critical concentrations and confers increased resistance to higher salt concentrations.

The dispensability of charged residue patterning for phase separation was initially surprising and prompted us to consider alternative roles for the observed native distribution of ionizable amino

acids. A possible answer was gleaned from observations made during the purification of Cdt1^{Uniform}. All proteins purified for this study were subject to a final polishing step over a size exclusion column, and it is at this step that we noticed a substantial difference in the elution profile of Cdt1 versus Cdt1^{Uniform}. Wild-type Cdt1 eluted from an analytical sizing column at a volume consistent with its physical parameters (size and globularity) and had no contaminating nucleic acid (A260/A280 = 0.56, 100% protein by mass) (Figure 2H, black trace). Conversely, Cdt1^{Uniform} eluted considerably earlier in a much broader peak, indicating that it possesses a more extended conformation compared to the native protein, and it co-eluted with a small fraction of nucleic acid (A260/A280 = 0.79, or 98% protein by mass) (Figure 2H, red trace). Additionally, a significant fraction of the Cdt1^{Uniform} protein eluted in the column's void volume, suggesting that this mutant shows higher aggregation propensity than wild-type Cdt1. Together, these results indicate that the distribution of charged residues within the Cdt1 IDR may encode some type of higher-order structural or conformational information that also regulates nucleic-acid binding capacity.

Intermolecular IDR-IDR interactions can drive LLPS in the presence of crowding reagent.

A parsimonious mechanism for LLPS by Cdt1 would rely exclusively on adhesive interactions (i.e., interactions between two dissimilar molecules) between the initiators and DNA (Figure 3A, top). However, it remained to be determined whether direct, inter-IDR contacts might provide an additional set of cohesive interactions (i.e., interactions between two similar molecules) that also promote LLPS (Figure 3A, bottom). To probe this issue, we asked whether Cdt1 could form condensates in the absence of DNA (or any other appropriate counterion) by the addition of a crowding reagent (Figure 3B). Using the depletion assay, we titrated PEG-3350 (from 0-12.5% w/v in 2.5% increments) against a fixed concentration of Cdt1 and assayed protein depletion from the supernatant. We observed a statistically significant reduction of Cdt1 from the supernatant starting at the lowest concentration of PEG-3350 (2.5%) that progressed as PEG concentrations were increased to a near-complete depletion of Cdt1 from the supernatant at the highest value tested (12.5% PEG-3350). To confirm that the PEG-induced loss of protein in the depletion assay represented phase separation (as opposed to precipitation), fluorescence microscopy was used to visualize mixtures of eGFP-Cdt1 and PEG-3350. Mixtures of Cdt1 and 4% PEG-3350 were sufficient to induce droplet

formation (Figure 3C, bottom panel). Thus, in addition to poly-anions such as DNA, intermolecular Cdt1-Cdt1 interactions also provide a driving force for LLPS.

The ability of Cdt1 to self-associate prompted us to look for specific motifs within its IDR that could promote such interactions. As a first step, three different 100-amino acid segments were removed from the Cdt1 IDR (Figure 3D) and assessed for how the deletions impacted PEG-induced Cdt1 phase separation (Figure 3E). Deletion of IDR residues 1-100 (Cdt1 Δ 1-100) or 101-200 (Cdt1 Δ 101-200) had no discernible effect on Cdt1's propensity to phase separate in the presence of PEG. However, we observed a modest reduction in Cdt1 partitioning (~50%) into the condensed phase when residues 201-297 (Cdt1 Δ 201-297) were deleted (Figure 3F). Although there is a predicted coiled-coil domain within the final third of the Cdt1 IDR (residues 196-223) that was suggestive of a possible self-association motif (Figure 3D, bottom), the deletion of this region alone had no impact on Cdt1's condensation properties (Figure 3-figure supplement 1).

Protein multivalency is a key property of proteins that undergo LLPS and frequently can arise from disordered sequences that contain short repetitive motifs (Banani et al., 2017). In the absence of obvious structural features to explain Cdt1 self-association, we attempted to identify repetitive motifs within the Cdt1 IDR – particularly within the 201-297 region – that might explain Cdt1 self-assembly. A prototypic example of repetitive IDR elements are those contained within human FUS, whose IDR contains multiple [S/G]YG motifs that are important for self-association (Kato et al., 2012; Luo et al., 2018). These repeats, in addition to the protein's poly-glycine sequences, can be visualized diagrammatically by a Dotplot (Madeira et al., 2019) (Figure 3G, top). In this plot, a sequence is plotted against itself and a dot is placed at each point where a repetitive match of a specific length and threshold is found. We used this approach to search for repetitive sequence elements within the Cdt1 IDR. However, in stark contrast to FUS and many other condensate-forming proteins (Figure 3-figure supplement 2A-F), we observed no repetitive motif(s) within the Cdt1 IDR (Figure 3G, bottom). This result led us to hypothesize that Cdt1 lacks repetitive sequences that would help promote self-association, and that it is the IDR amino acid composition alone that contains the necessary components to promote inter-protomer interactions and phase separation. To test this idea, we synthesized and produced a Cdt1 variant, Cdt1^{ScrIDR}, where the order of the N-terminal IDR amino

acids (residues 1-297) were randomly scrambled and then fused back to the wild-type C-terminal portion of the protein (residues 298-743). Dot plot analysis of the Cdt1 IDR versus that of the scrambled variant shows that no sequence motifs were inadvertently retained in Cdt1^{ScrIDR} (Figure 3H). The depletion assay was then used to ask whether this protein was still capable of phase separation. Surprisingly, Cdt1^{ScrIDR} showed robust phase separation propensity in the presence of a crowding agent, demonstrating that Cdt1^{ScrIDR} retains the self-associative capacity observed for wild-type Cdt1 (Figure 3I). We also assessed whether Cdt1^{ScrIDR} could undergo DNA-induced LLPS by assessing phase separation in the presence of Cy5-dsDNA over a range of protein and KGlutamate concentrations (Figure 3J). At [KGlutamate] = 150 mM, the Cdt1^{ScrIDR} critical concentration (200 nM) was equivalent to that of wild-type Cdt1 (Figure 2F) and, in fact, Cdt1^{ScrIDR} was able to phase separate over a broader set of experimental conditions. Notably, Cdt1^{ScrIDR} condensates were also able to recruit native eGFP-Cdt1, although the efficiency of recruitment was lower than that observed with wild-type Cdt1 condensates (Figure 3-figure supplement 3A-B). Collectively, these data demonstrate that it is the overall amino acid composition of initiator IDR sequences, not sequence order, that is essential for Cdt1's self-associative capacity and that composition alone can promote heteromeric inter-IDR interactions. However, the differences observed in the ability of eGFP-Cdt1 to be recruited into condensates formed by wild-type Cdt1 and Cdt1^{ScrIDR} also suggest that there exist specific sequence patterns that can fine tune IDR association that have yet to be identified.

Branched hydrophobics are disproportionately enriched in the Cdt1 IDR and drive LLPS.

The Cdt1 IDR contains alternating blocks of net-positive and net-negative charge (Figure 2E). We reasoned that these interspersed blocks of opposite charge might facilitate inter-IDR interactions in the presence of a crowding reagent to drive phase separation; other condensate-forming proteins, such as *C. elegans* Laf1 (Elbaum-Garfinkle et al., 2015) and Ddx4 (Nott et al., 2015), have been shown to phase-separate in such a manner. To test this idea, we compared the salt-sensitivity of PEG-induced Cdt1 LLPS to DNA-induced condensation (Figure 4A). At 75 mM and 150 mM KGlutamate, both DNA and PEG were comparable in their ability to induce Cdt1 phase separation. By contrast, salt concentrations above 150 mM KGlutamate abolished DNA-induced phase-separation by Cdt1 but had no effect on PEG-induced LLPS. This result demonstrates that charge-charge interactions are not a major contributor to the inter-IDR interactions that drive PEG-induced phase separation. To test

whether salt-insensitive inter-IDR interactions are a generally conserved feature of initiator phase separation, we assayed DNA and PEG-induced phase separation of both purified *Drosophila* ORC and Cdc6 at 150 mM and 300 mM KGlutamate concentrations (Figure 4-figure supplement 1). As seen for Cdt1, we observed that at 150 mM KGlutamate, either DNA or PEG are sufficient to induce Cdc6 and ORC phase separation, suggesting a role for both DNA-bridging interactions and direct inter-IDR interactions in metazoan initiator LLPS. Additionally, we found that DNA-induced phase separation is sensitive to 300 mM KGlutamate while PEG-induced phase separation largely is not. These data support a generally conserved mechanism of phase separation for metazoan Cdt1, Cdc6 and ORC.

We next set out to test whether initiator inter-IDR interactions might be driven by the hydrophobic effect. We first asked whether initiator IDRs are uniquely enriched in hydrophobic residues. However, when we compared the hydropathy of initiator IDRs to all ordered and disordered segments longer than 100 amino acids within the *D. melanogaster* proteome (14,002 ordered domains and 4957 disordered domains), we found that initiator IDRs were closer to the hydropathy of the median IDR (average of -1.04 for Orc1, Cdc6, and Cdt1) than to either the lower (-1.24) or upper (-0.80) quartile of the data (Figure 4B). This result led us to take a more unbiased approach and compare initiator IDR sequence composition (fraction average of each amino acid across Orc1, Cdc6, and Cdt1 IDRs) to the average composition of all ordered and disordered domains longer than 100 amino acids (Figure 4C). We found that initiator IDRs are more highly enriched in both the three most hydrophobic residues (isoleucine, valine and leucine) and the three least hydrophobic residues (glutamate, lysine and arginine) compared to other disordered domains (Figure 4C, bottom). We also compared the total fraction of Leu, Ile, and Val (fraction-L,I,V) residues in the initiators to all ordered and disordered domains proteome wide. We found that the initiator fraction-L,I,V was within the top 25% of all disordered domains (Figure 4D). By comparison, initiator IDRs are relatively deficient in glycine and, to a lesser extent, uncharged polar residues, compared to other disordered sequences (Figure 4C, bottom).

The higher fraction of branched hydrophobic residues within initiator IDRs prompted us to test whether these amino acid types promote cohesive inter-IDR interactions that drive Cdt1 self-association. The fraction-L,I,V in the Cdt1 IDR was first calculated over a five-residue sliding

window to determine whether hydrophobic residues were uniformly distributed throughout the sequence or clustered into discrete motifs targetable by mutagenesis (Figure 4E, top panel). Overall, the Cdt1 IDR bears an average fraction-L,I,V of 0.19 – or approximately one Leu, Ile, or Val for every 5 residues – and the hydrophobic residues are generally well-distributed throughout the sequence. There are five 5-residue windows where the fraction-L,I,V reaches 0.6, and each third of the IDR (residues 1-100, 101-200, 201-297) contains at least one of these regions. A fraction-L,I,V = 0.8 occurs only once, between residues 244-248. Interestingly, the C-terminal third of the Cdt1 IDR (residues 201-297) contributes more strongly to PEG-induced LLPS than either the first (1-100) or middle third (101-200) of the sequence (Figure 3E), suggesting that the 244-248 sequence may contribute to the greater cohesiveness of this region. A higher-precision analysis of Cdt1's hydrophobic residue distribution showed that 67% of Leu, Ile, and Val residues reside in isolation, 22% are immediately adjacent to another Leu, Ile, or Val residue, and 11% are within tri-peptides composed of the three amino acids.

Because sequence analysis shows that hydrophobic residues are broadly distributed across the length of the Cdt1 IDR, we were interested to test whether these amino acids contribute to PEG-induced LLPS in an accumulative manner. Two Cdt1 IDR mutants were constructed to probe this issue, one with half of all Ile and Leu residues mutated to alanine (Cdt1^{Ile/Leu(0.5)}) and another with all Ile and Leu residues mutated (Cdt1^{Ile/Leu(1.0)}). Ile and Leu were chosen as targets, as these are the two most highly enriched hydrophobic residues in initiator IDRs compared to disordered sequences across the proteome. To assess phase separation potential, PEG was titrated (from 0-12.5% in 2.5% increments) against each mutant and their disappearance monitored from the supernatant by depletion assay (Figure 4F). The phase separation propensity of both variants was significantly disrupted. At the highest concentration of PEG (12.5%), we observed only a modest, insignificant loss of Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} (27% and 19%, respectively) from the supernatant, compared to nearly a 100% loss for wild-type Cdt1 (Figure 3B). These data establish that, relative to the average disordered domain, initiator IDRs are enriched in branched hydrophobic residues and these amino acid types mediate cohesive inter-IDR interactions that promote phase separation by Cdt1 under conditions of molecular crowding.

Inter-IDR hydrophobic interactions are necessary for DNA-induced LLPS.

Our results have shown that both DNA-bridging interactions (mediated by electrostatics) and inter-IDR interactions (mediated by hydrophobic residues) are important for phase separation by Cdt1. However, the extent to which these two types of interactions synergize remained unclear. We therefore set out to determine whether inter-IDR interactions promote phase separation by Cdt1 in the presence of DNA. During the purification of Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} we noticed that both proteins showed a normal affinity for heparin (Figure 5-figure supplement 1), indicating that the mutants might bind DNA with wild-type-like affinity. This assumption was confirmed by electrophoretic mobility shift assay, which showed that the DNA-binding affinity of Cdt1^{Ile/Leu(0.5)} ($K_{d, app} = 67$ nM) and Cdt1^{Ile/Leu(1.0)} ($K_{d, app} = 106$ nM) is similar to that of the wild-type protein ($K_{d, app} = 69$ nM) (Figure 5A). These data establish Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} as separation-of-function mutants that are capable of binding DNA (Figure 5A) but that are deficient in their ability to participate in inter-IDR interactions (Figure 4).

We next utilized these mutants to ask whether DNA bridging by itself is sufficient to drive Cdt1 LLPS, or whether inter-IDR cohesive interactions are also required. We first used the depletion assay to compare DNA-induced phase separation of Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} vs. the wild-type protein (Figure 5B). While wild-type Cdt1 showed near complete partitioning into the condensed phase (8% protein retention in the supernatant), 63% of Cdt1^{Ile/Leu(0.5)}, and 73% of Cdt1^{Ile/Leu(1.0)} remained in the supernatant, demonstrating a marked reduction in DNA-induced phase separation (Figure 5C). To gain a quantitative understanding of the DNA-dependent phase separation propensity of these mutants, we generated phase diagrams for Cdt1^{Ile/Leu(0.5)} (Figure 5D) and Cdt1^{Ile/Leu(1.0)} (Figure 5E) to compare with wild-type protein (Figure 2F). We observed a 25-fold reduction in the critical concentration for both mutants. Furthermore, the droplets which formed at 5 μ M Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} and [KGlutamate] = 150 mM showed a greater than 60% reduction in mean droplet Cy5-dsDNA signal intensity compared to that of the wild-type protein (Figure 5F). Overall, these data reveal a striking reduction in the ability of Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} to undergo DNA-induced phase separation, despite possessing apparently normal DNA binding activity, demonstrating that cohesive inter-IDR interactions play a uniquely critical role in driving the phase separation propensity of Cdt1.

DISCUSSION

In previous work, we discovered that several factors used to initiate DNA replication in metazoans – in particular the Orc1 subunit of ORC, as well as Cdc6 and Cdt1 – possess an N-terminal IDR that promotes protein phase separation upon binding DNA (Parker et al., 2019). When these initiator proteins are combined and DNA is added, they form co-mingled condensates that exclude non-cognate phase-separating proteins and are active for the ATP-dependent recruitment of the Mcm2-7 helicase. The biophysical mechanisms underlying LLPS by replication initiators have remained unknown.

Here, we have used *Drosophila* Cdt1 as a model system to dissect the molecular basis for DNA-dependent phase separation by initiation factors. Our studies show that this condensation mechanism has several distinctive features compared to other proteins characterized to date that also undergo LLPS. We find that two types of intermolecular interactions synergize with each other to drive phase separation by Cdt1. One set of interactions occur between Cdt1 and DNA. These associations appear to be non-specific and electrostatic in nature, with DNA serving as a polyanion that bridges the cationic IDRs of multiple Cdt1 molecules (Figure 2 and Figure 5G). A second set of interactions that occur between the IDRs of different Cdt1 protomers were also identified; these appear to be salt-insensitive and primarily hydrophobic in nature (Figure 4 and Figure 5G). Importantly, we find that inter-IDR interactions can drive Cdt1, ORC and Cdc6 LLPS in the absence of DNA (Figure 3 and Figure 4 – figure supplement 1). When we abolish these interactions in Cdt1, DNA is insufficient to induce phase separation (Figure 5). Thus, inter-IDR interactions provide the primary force behind initiator LLPS, with electrostatic DNA bridging interactions contributing an additional adhesive force.

In the absence of a crowding reagent, DNA is required to induce initiator phase separation at physiological salt and protein concentrations (Parker et al., 2019). This dependency suggests that DNA serves as an essential nucleating factor that determines where Cdt1 can form condensates, ensuring that initiator LLPS is restricted to a chromatin context. Interestingly, other anionic

biopolymers, such as RNA and heparin, are also capable of inducing phase separation by Cdt1. This type of scaffold ‘promiscuity’ has been observed in non-initiator systems that undergo LLPS, such as for the nucleolar protein fibrillarin (Feric et al., 2016). Given that the cytosol contains multiple anionic biopolymers, future work will need to answer how replication initiators are specifically targeted to chromatin, a question that is similarly relevant for RNA-dependent cellular bodies (Berry et al., 2015; Elbaum-Garfinkle et al., 2015; Feric et al., 2016; Guillén-Boixet et al., 2020; Mitrea et al., 2016; Molliex et al., 2015; Van Treeck et al., 2018). The selection of an appropriate nucleic acid scaffold across different phase-separating systems is likely to rely on both general mechanisms, such as the sequestration of alternative, inappropriate scaffolds by proteins with different or non-existent LLPS properties, and by system-specific mechanisms, such as a requirement for particular sequences or secondary structure (Langdon et al., 2018; Maharana et al., 2018).

A growing body of work has demonstrated the importance of aromatic residues in driving protein phase separation (Chiu et al., 2020; Chong et al., 2018; Lin et al., 2017; Nott et al., 2015; Pak et al., 2016; Qamar et al., 2018; Wang et al., 2018), along with a related role for π - π and π -cation interactions (Vernon et al., 2018). In parallel, many studies have shown that condensates formed through aromatic residue interactions are readily dissolved by 1,6-HD (Table 1). Cdt1 LLPS is both resistant to treatment with 1,6-HD and does not rely on aromatic residues within its IDR for phase separation (Figure 1). ORC and Cdc6 phase-separation is also resistant to 1,6-HD (Figure 1-figure supplement 2). Although we have not explicitly investigated the role of aromatic residues within these proteins, we note that the fraction aromatic residues of Orc1 (0.019) and Cdc6 (0.012) is lower than that of Cdt1 (0.027), suggesting that aromaticity is dispensable for LLPS by replication initiation factors in general. Similarly, charged residues have an established role in facilitating homomeric inter-IDR interactions that lead to phase separation (Elbaum-Garfinkle et al., 2015; Mitrea et al., 2018; Nott et al., 2015), yet the inter-IDR interactions that drive initiator phase separation (Figure 4 and Figure 4 – figure supplement 1) are salt insensitive.

Instead of relying on aromatic-mediated π - π / π -cation interactions or charge-charge interactions, our data show that interactions between Cdt1 IDRs are instead facilitated by branched hydrophobic residues. Although hydrophobic sequences have been shown to be important for the phase separation

of other factors, such as the extracellular matrix protein elastin (Quiroz and Chilkoti, 2015; Reichheld et al., 2017; Yeo et al., 2011) and the Nephrin intracellular domain (NICD) (Pak et al., 2016), important mechanistic distinctions exist with respect to metazoan replication initiators. For example, both NICD and Cdt1 bear an identical fraction of charged residues (FCR = 31%) and rely on a counterion for phase separation. However, inter-IDR interactions within NICD appear to rely primarily on sequence aromatics (11% aromatic content), with hydrophobic residues playing only a supporting role in the self-association of this protein (Pak et al., 2016). By contrast, Elastin is more like metazoan replication initiators in that it is relatively devoid of aromatic residues (3.9%) and contains a similar content of branched hydrophobic amino acids that underly phase separation (elastin and Cdt1 fraction-L,I,V = 0.21 and 0.19, respectively); however, Elastin also differs from initiators in that it readily forms condensates in the absence of a counterion. Elastin's hydrophobic sequences are additionally contained within highly repetitive "VPGVG" and "GLG" sequences, and scrambling or shuffling these motifs negatively impacts elastin phase-separation (Pepe et al., 2008, 2005; Toonkool et al., 2001). For their part, initiator IDRs lack such repeats and (for Cdt1 at least) LLPS is insensitive to sequence order. We speculate that the inter-IDR interactions which occur in initiators are weaker than those of Elastin due to the non-repetitive, distributed nature of their hydrophobic residues, a property that necessitates additional counterion-mediated interactions to promote Cdt1 self-assembly.

Understanding the distinctive molecular interactions of different phase-separating systems is essential for developing *in silico* approaches for predicting the cellular partitioning of a given IDR from primary sequence alone. Such efforts will ultimately require an understanding of both the specific and general features of condensates' scaffolding components (e.g., repetitive motifs vs. amino acid composition). The present work highlights the importance of the general sequence features of initiator IDRs (as opposed to strict amino acid order) in promoting LLPS. Strikingly, we find that the Cdt1 IDR can be fully scrambled without losing the ability to form condensates (Figure 3I-J) and that condensates formed from the scrambled variant can recruit wild-type Cdt1 into droplets (Figure 3 – figure supplement 3). Regions of NICD can similarly be scrambled without losing the ability to form cellular condensates (Pak et al., 2016), as can the LCDs of certain transcription factors without affecting promoter targeting (Brodsky et al., 2020). These studies demonstrate the importance of

sequence composition is not only driving phase separation but also in facilitating interactions with partner-proteins, and is consistent with recent work showing that proteins with similar functional annotation have disordered sequences with similar, evolutionarily conserved physical-chemical features (pI, composition, FCR, kappa, etc.) (Zarin et al., 2019). Our observations involving the co-condensation properties of Cdt1, ORC and Cdc6 – three proteins with IDRs of similar amino acid composition but no direct sequence homology (Parker et al., 2019) – provide experimental support for this “like-recruits-like” concept. Future work will need to address whether heteromeric inter-IDR interactions (e.g. Orc1^{IDR}-Cdc^{IDR}) are indeed governed strictly by sequence composition or whether linear sequence motifs facilitate specific interactions, as was recently suggested for human Orc1 and Cdc6 (Hossain et al., 2021). Further, while metazoan Cdt1 orthologs have no identifiable sequence identity across their disordered domains (Figure 5 – figure supplement 2A-B), they do have a similar amino acid composition (Parker et al., 2019). This led us to predict that the capacity to phase-separate was likely conserved across metazoan initiators and we confirmed this for human Cdt1 (Parker et al., 2019). These predictions were further validated for human Orc1 and Cdc6 with the demonstration that these proteins form condensates in the presence of DNA (Hossain et al., 2021). These results suggest that certain classes of disordered domains can have conserved functionality in the absence of linear sequence identity.

Although amino acid composition is the primary determinant underpinning LLPS by replication initiators, short sequence motifs can nonetheless play a critical role in modulating condensation behavior and controlling biological function. Sequence information within the Orc1 and Cdt1 IDR's is high, and while saltatory leaps seem to have occurred between phyla (Figure 5 – figure supplement 2B), the conservation of sequence homology within the *Drosophila* genus (Figure 5 – figure supplement 2C and (Parker et al., 2019)), over millions of years, is remarkable and indicates an evolutionary pressure on sequence order for unknown function. A conserved sequence feature within the IDRs of Orc1, Cdt1, and Cdc6 are CDK phosphorylation sites which are known to regulate replication initiation *in vivo* (Findeisen et al., 1999; Lee et al., 2012). We have previously shown that the phosphorylation of these sites – which are distributed throughout the initiator IDRs – by CDKs abrogates the ability of the *Drosophila* initiators to form condensates (Parker et al., 2019), and a recent finding shows that phosphorylation impedes inter-IDR interactions between human ORC1

and CDC6 as well (Hossain et al., 2021). Notably, consensus motifs for CDK-dependent phosphorylation (full site = [S/T]PX[R/K] and minimal site = [S/T]P) are conserved in a majority of metazoan Cdt1s (Figure 5 – figure supplement 2D) and in all sequenced *Drosophilidae* Cdt1 orthologs (Figure 5 – figure supplement 2E), leading us to predict that phospho-tunable phase separation is a broadly conserved mechanism for regulating metazoan replication licensing. Given that a scrambled IDR can still support LLPS *in vitro* through both self-interactions and cross-interactions with a wild-type IDR sequence (Figure 3), it will be useful to determine whether such a construct (with or without the native CDK sequences) can support normal kinetics of chromatin association and MCM recruitment and/or cell viability. Such studies, along with deletion and more targeted mutagenesis efforts, will be important to define how the plasticity of IDR sequences can be fine-tuned to elicit specific timing and partnering responses.

Defining the molecular ‘grammar’ that encodes LLPS in metazoan initiators likewise has implications for understanding the impact of initiator phase separation *in vivo*. Admittedly, neither this nor our previous study (Parker et al., 2019) provides unequivocal evidence for initiator phase separation in cells. It is also unclear what function such concentrated initiator assemblies might have, such as improving the kinetics or efficiency of helicase loading (as we have suggested here and previously (Parker et al., 2019)), or something else entirely, such as maintaining the appropriate nuclear levels of the licensing factors (as proposed in (Hossain et al., 2021)). We note that our system lacks the convenience of identifying initiator condensation morphologically (i.e., by the presence of round cellular foci), as mitotic chromosomes – the stage of initiator assembly – are relatively rigid structures built from a central proteinaceous scaffold through an energy-consuming reaction (Earnshaw and Laemmli, 1983; Gibcus et al., 2018; Hirano and Mitchison, 1994; Kim et al., 2020). Thus, we predict that initiators condense on the surface of mitotic chromosomes without altering their shape. Consistently, work in multiple model organisms has revealed a switch-like transition in initiator localization during anaphase, at which point initiators begin to coat chromosomes (Baldinger and Gossen, 2009; Kara et al., 2015; Parker et al., 2019; Sonnevile et al., 2012). We predict that phosphorylation-dependent control of initiator self-assembly underlies this behavior and produces a dense, replication-competent zone of chromatin-bound initiators poised to drive the precipitous, genome-wide loading of the Mcm2-7 complex in late mitosis (Dimitrova et al., 2002; Méndez and

Stillman, 2000; Okuno et al., 2001). This moment in the cell cycle is an opportune time to prepare for replication, as conflicts with the transcriptional machinery are minimized (Palozola et al., 2017) and the chromatin substrate is, relative to interphase, uniformly compacted (Ou et al., 2017). The identification of Cdt1 mutants that selectively block phase separation, but not DNA binding (Figure 5), affords an opportunity to directly investigate the physical nature of these global initiator interactions *in vivo*.

Beyond helicase loading and licensing, initiator LLPS might also serve to prime replisome assembly. Following helicase loading, Mcm2-7 is phosphorylated by the Dbf4-dependent kinase (DDK), a heterodimer of Cdc7 and Dbf4 (Deegan et al., 2016; Sheu and Stillman, 2010; Yeeles et al., 2015). Intriguingly, the *Drosophila* homolog of Dbf4, Chiffon, contains multiple IDRs, including a large internal IDR (residues 903-1209) that possesses striking compositional similarity to initiator-type IDRs (Figure 5-figure supplement 3). The identification of an IDR in Chiffon with compositional similarity to the initiator-type IDRs suggests that such sequences may be present in chromatin-associated factors beyond replication licensing components. Indeed, the capacity to undergo DNA-dependent phase separation may have broad utility for chromatin-localized factors and their respective cellular pathways. Thus, our future work will aim to develop algorithms that accurately identify proteins proteome-wide that possess disordered domains with compositional homology to initiator IDRs and to understand how these sequences impact protein functional dynamics.

In summary, the present work provides a detailed view of the DNA-dependent liquid-liquid phase separation mechanism for *Drosophila* Cdt1. Due to a high-level of IDR compositional homology and an ability to form co-mingled phases (Parker et al., 2019), the mechanism we describe for Cdt1 phase-separation likely extends to the other replication initiation factors, ORC and Cdc6. These studies set the stage for investigating the physiological significance of initiator self-assembly in replication licensing and for identifying other sequences in the proteome that possess IDRs capable of associating with chromatin and initiation factors alike.

MATERIALS AND METHODS

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Recombinant DNA reagent	2Cc-T	QB3 Macrolab (UC Berkeley)	RRID:Addgene_37237	Ligation independent cloning (LIC); <i>E. coli</i> expression vector
Recombinant DNA reagent	1GFP	QB3 Macrolab (UC Berkeley)	RRID:Addgene_29663	Ligation independent cloning (LIC); <i>E. coli</i> expression vector
Peptide, recombinant protein	TEV	QB3 Macrolab (UC Berkeley)		Used at 1/10 (weight/weight) TEV/substrate ratio
Strain, strain background (<i>Escherichia coli</i>)	Rosetta 2(DE3) pLysS	QB3 Macrolab (UC Berkeley)		Chemically competent cells

Protein production and purification. Wild-type and mutant Cdt1 coding sequences (Table 3) were cloned into the QB3 Macrolab vector 2Cc-T to append a tobacco etch virus (TEV) protease-cleavable C-terminal hexa-histidine (His6)-maltose binding protein (MBP) tag. To produce eGFP-tagged Cdt1, the wild-type Cdt1 coding sequence was cloned into QB3 Macrolab vector 1GFP to append an N-terminal His6-eGFP tag. All Cdt1 variants were produced from Rosetta 2(DE3) pLysS cells. Cells were grown in 2xYT broth at 37°C to an OD₆₀₀ = 0.8 and after a 15 min incubation in an ice bath induced with 1 mM isopropyl β-d-thiogalactoside (IPTG). After growth overnight at 16°C, cells were harvested by centrifugation and cell pellets frozen at -80°C until further processing.

Cell pellets were resuspended in Lysis Buffer (20 mM Tris pH 7.5, 500 mM NaCl, 30 mM Imidazole, 10% glycerol, 200 μM PMSF, 1x cOmplete EDTA-free Protease Inhibitor Cocktail (Sigma-Aldrich), 1 mM BME and 0.1 mg/mL lysozyme) and lysed by sonication. Lysates were clarified by centrifugation

at 30,000 xg for 1 hr and filtered through a 0.45 μ m bottle-top filter unit (Nalgene Rapid-Flow, ThermoFisher). Lysates were then passed over a 5 mL HisTrap HP column (GE Healthcare) and washed with 10 column volumes (CV) of Nickel Wash Buffer (20 mM Tris pH 7.5, 1 M NaCl, 30 mM Imidazole, 10% glycerol, 200 μ M PMSF, 1 mM BME) followed by 5 CV of Low Salt Nickel Wash Buffer (20 mM Tris pH 7.5, 150 mM NaCl, 30 mM Imidazole, 10% glycerol, 200 μ M PMSF and 1 mM BME). Protein was eluted from the column with 5 CV of Nickel Elution Buffer (20 mM Tris pH 7.5, 150 mM NaCl, 500 mM Imidazole, 10% glycerol, 200 μ M PMSF and 1 mM BME). Eluted protein was then loaded onto a 5 mL HiTrap Heparin HP column (GE Healthcare), washed with 5 CV of Heparin Wash Buffer (20 mM Tris pH 7.5, 150 mM NaCl, 10% glycerol, 200 μ M PMSF and 1 mM BME) and eluted with a linear gradient of increasing salt from 150 mM – 1 M NaCl (20 mM Tris pH 7.5, 10% glycerol, 200 μ M PMSF and 1 mM BME). Fractions containing Cdt1 were pooled and TEV protease (QB3 Macrolab) added at a 1:20 w/w ratio of protein:TEV and incubated overnight at 4°C. A second nickel affinity step was used to remove TEV, uncleaved protein, and the free His6-MBP tag. Finally, the sample was concentrated to 2 mL and loaded onto a HiPrep 16/60 Sephacryl S-300 HR column (GE Healthcare) equilibrated and run in Sizing Buffer (50 mM HEPES pH 7.5, 300 mM KGlutamate, 10% glycerol and 1 mM BME). Peak fractions were collected, concentrated in a 30K Amicon Ultra-15 concentrator (Millipore), flash frozen in liquid nitrogen, and stored at -80°C. The same protocol was used to purify GFP-tagged Cdt1 with the exception that TEV protease was not added, and the second nickel affinity step was omitted.

D. melanogaster ORC and Cdc6 were purified as previously described (Parker et al., 2019).

Fluorescence microscopy. All microscopy assays with DNA utilized “Cy5-dsDNA”, a sixty base pair duplex of sequence:

5'-GAAGCTAGACTTAGGTGTCATATTGAACCTACTATGCCGAAGTACGAGCTATAAAC-3' that had a 5' Cy5 label (IDT or Eurofins). Duplex DNA was annealed at 100 μ M in 50 mM Tris pH 7.5, 50 mM KCl. Samples were prepared by mixing protein (10 μ M Cdt1 and Cdc6 and 5 μ M ORC unless otherwise noted in the text) prepared in Protein Buffer (50 mM HEPES pH 7.5, 300 mM KGlutamate, 10% glycerol, 1 mM BME) with an equal volume of equimolar Cy5-dsDNA prepared in Dilution Buffer (50 mM HEPES pH 7.5, 10% glycerol, 1 mM BME), were mixed thoroughly by pipetting and

incubated for 2 mins at room temperature. 7 μ L of sample was spotted onto a glass slide, a coverslip placed on top, and imaged with a 60x oil objective using confocal fluorescence microscopy. Images were processed in FIJI and quantitation performed by thresholding for Cy5 signal intensity and then calculating the mean Cy5 and/or eGFP intensity within droplets. The effect of 1,6-hexanediol was assessed in a similar fashion except that protein stocks (10 μ M Cdt1, 10 μ M Cdc6 and 5 μ M ORC) were prepared in Protein Buffer with 20% 1,6-hexanediol and then diluted 1:1 with Cy5-dsDNA in Dilution Buffer (final [Cy5-dsDNA] = 5 μ M and 1,6-hexanediol = 10%). eGFP-Cdt1 phase separation was assessed in the presence of PEG-3350 (Sigma 202444-250G) by mixing equal volumes of 10 μ M eGFP-Cdt1 in Protein Buffer and 8% PEG dissolved in Dilution Buffer. Experiments were completed in triplicate and at least three fields of view imaged for each experiment.

Protein depletion LLPS assay. All depletion assays with DNA utilized “dsDNA”, a sixty base pair duplex with the same sequence as Cy5-dsDNA. Duplex DNA was annealed at 100 μ M in 50 mM Tris pH 7.5, 50 mM KCl. Samples were prepared by mixing protein (4 μ M Cdt1 unless otherwise noted in the text) prepared in Protein Buffer (50 mM HEPES pH 7.5, 300 mM KGlutamate, 10% glycerol, 1 mM BME) with an equal volume of equimolar dsDNA prepared in Dilution Buffer (50 mM HEPES pH 7.5, 10% glycerol, 1 mM BME), were mixed by gently flicking the tube and incubated for 30 mins. Subsequently, the dense phase-separated material was pelleted by centrifugation (10 min at 16,000 xg) and the supernatant removed to assess for protein retention by SDS-PAGE analysis and Coomassie staining. Every assay included a load control that had not been centrifuged. Samples were run on 4-20% gradient gels (BioRad 4561096).

The same protocol was used for comparing the efficiency of different polyanionic scaffolds at inducing Cdt1 phase separation except that 0.06 mg/mL of each polyanion was mixed with an equal volume of 4 μ M Cdt1. The polyanions included dsDNA (same sequence as above), ssDNA (same sequence as dsDNA), heparin (Sigma H3149), polyglutamate (Sigma P4886), as well as dsRNA prepared using the MEGAscript RNAi Kit (Life Technologies) from a 422 base pair PCR fragment derived from the *D. melanogaster* DPOA2 gene. For depletion assays completed in the presence of crowding reagent, PEG-3350 (Sigma 202444-250G) was used in all cases and phase separation assessed by mixing equal volumes of 16 μ M Cdt1 in Protein Buffer and PEG-3350 dissolved in Dilution Buffer

to reach a final concentration of 0-12.5% PEG-3350. In these experiments, the samples were diluted 4-fold with water prior to analysis by SDS-PAGE to reduce migration artifacts induced by high concentrations of PEG-3350. For depletion assays with monovalent and multivalent phosphate counterions, concentrated stocks of monobasic, dibasic and tribasic potassium phosphate were prepared in Dilution Buffer and the pH adjusted to 7.5 prior to use. Experiments were completed in triplicate and band quantitation was completed using FIJI (Schindelin et al., 2012).

Generation of phase diagrams for Cdt1 and Cdt1 variants. A dilution series of Cdt1 (30 μ M, 6 μ M, 1.2 μ M and 0.24 μ M) and Cy5-dsDNA (50 μ M, 15 μ M, 3 μ M and 0.6 μ M) was prepared in Protein Buffer and Dilution Buffer, respectively. These stocks were then combined in a matrix with buffer containing 50 mM HEPES pH 7.5, 10% glycerol and variable concentration of KGlutamate to reach final [KGlutamate] = 300 mM, 150 mM, 100 mM or 0 mM. Samples were prepared on ice in microcentrifuge tubes by first combining DNA with buffer and then adding protein. The samples were gently mixed by flicking the tube and then briefly spun prior to adding 15 μ L of each sample to a CellVis 384-well glass bottom multiwell plate (P384-1.5H-N). Imaging of each sample matrix was automated using a 40x air objective (NA = 0.75) on a Nikon Ti2-E spinning disk confocal microscope (Yokogawa X1) with 561 nm laser light illumination. For each sample, a 6-micron z-stack (0.5 μ m spacing) was collected for a 3x3 field of view (1024x1024 pixels with 5% overlap between adjacent images) which were stitched together using Nikon NIS Elements software. Two replicate experiments were completed for each protein on different days which revealed qualitatively equivalent results and a quantitative change in mean signal intensity of < 10% between replicate samples. A single replicate was used for further processing, analysis and illustration.

Processing and analysis of phase diagram data was completed exclusively using Nikon NIS Elements software. First, a max intensity projection was generated and then a median filter (kernel = 5) applied. The detection of phase separated particles was automated in NIS Elements. Specifically, an algorithm was applied that identified foci with intensity greater than 230 (a.u.) and an area > 0.3 μ m². Phase separating conditions were defined as those producing > 2 foci (as defined above) per field of view. Foci intensities were directly compared between conditions and across samples by calculating the

mean intensity of each focus in each large image (3x3 field of view). For visualization, a 75 μm x 75 μm region was cropped from each 3x3 field of view.

DNA-binding electrophoretic mobility shift assays (EMSAs). DNA-binding assays were completed with dsDNA containing a 5' IRDye800 label. Protein was titrated from 16 nM to 1 μM in the presence of 20 nM IRDye800-dsDNA in Assay Buffer (50 mM HEPES pH 7.5, 150 mM KGlutamate, 10% glycerol, 1 mM BME). The samples were mixed by pipetting and incubated for 45 mins at room temperature. 5 μL of each sample was run on a 7.5% PAGE gel (BioRad 4561026) pre-run at 100 volts for 1 hour and gels imaged on an Odyssey imaging system (LI-COR) through detection of IRDye800.

Sequence analysis and bioinformatics. Amino acid heatmaps and sequence complexity were calculated in Python using the Seaborn and SciPy (Virtanen et al., 2020) modules, respectively. EMBOSS dotmatcher was used to generate sequence dotplots with parameters "Window size" = 15 and "Threshold" = 32 (Rice et al., 2000). Coiled-coil predictions were made using the Parcoil2 server (McDonnell et al., 2006).

A custom Python script was used to calculate proteome-wide statistics on ordered and disordered domains (https://github.com/mwparkerlab/Parkeretal2021_Cdt1LLPS.git). Briefly, IUPRED2 (Mészáros et al., 2018) disorder prediction was run on every protein in the *D. melanogaster* reference proteome (release UP000000803_7227.fasta) and the per-residue disorder scores were smoothed by applying a moving average over a twenty-residue window. Sequences of contiguous disorder and order longer than 100 amino acids were extracted from the data and the sequence hydropathy (Kyte and Doolittle, 1982) and amino acid composition were calculated for each region. The average fraction of each of the twenty amino acids in all ordered and disordered domains was calculated from the composition of all ordered and disordered domains longer than 100 residues. Sequence hydropathy and composition were separately calculated for the initiator IDRs, and included Orc1 residues 187-549, Cdc6 residues 1-246 and Cdt1 residues 1-297, resulting in three values that were averaged to generate initiator hydropathy and initiator composition. The difference in amino acid composition between initiators and all other ordered and disordered protein regions proteome-wide was calculated by subtracting the proteome-wide values from the initiator values.

782

783 *Analytical size exclusion and heparin affinity chromatography.* To assess how charged residue
 784 distribution impacts the conformation of the Cdt1 IDR, we compared the size exclusion
 785 chromatography elution profiles of Cdt1 and Cdt1^{Uniform}. 50 μ L of 6 μ M Cdt1 or Cdt1^{Uniform} was
 786 prepared in Protein Buffer and then loaded and run on a Superose 5/150 GL sizing column (GE
 787 Healthcare) pre-equilibrated and run in Protein Buffer. We compared the heparin binding profiles of
 788 Cdt1, Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)} by injecting 50 μ L of 8 μ M protein in Protein Buffer onto a 1 mL
 789 HiTrap Heparin HP column (GE Healthcare) and eluting with a 10 CV linear gradient from 150 mM
 790 – 1 M NaCl in 20 mM Tris pH 7.5, 10% glycerol, 1 mM BME.

791

792 **Table 3: IDR sequences of the proteins used in this study.**

793 All IDR sequences were fused back to wild-type Cdt1 residues 298-743.

<i>DmCdt1</i> (WT)	ATGCCCCAGCCATCGGTAGCTGCATTTTTCACAAACCGCAAACGCGCCGCTTGGATGATGCTATCAGTATCAAGAACAGGCGTTTGG TGGAAACCGCTGAAACCGTCTCTCTCGCTCGGCCCTTCCCAGTTGCCAGCCGCGACAGGATGCGGATCTAGACACCTGAAGGC GGGGGCCAGGGCATGCGTACCCGATCCGGACGCACTGCCGACTAATTGTACCGCCGCTCAAGAGAGCAAAAAGACACCGGC TGCCGCCAAGATGGAGCCACATCAAGCAGCCCAAGCTGGTGCAATTCATTAAGGGCACTCTGTGCCCAGGAAACAGGCTCA GTCCAGTAAGCTGGACGAGGAGAGCTGCAGCAGTCGTGCGCCATAAGCGAGCACACGCCCAAGGTTAACTTCACCATCACAAGCCA GCAGAATGCGGACAATGTGCAGCGTGGCCTGCGCACACCCACCAAGCAGATCCTCAAGGATGCCTCGCCGATCAAGGCGGATCTCCG CCGTCAGCTCACTTTCGACGAGGTAAAAACGAAGGTATCGCGGAGTGCCAAGCTGCAGGAACCTCAAGGCAGTGTGGCCCTTAAGGC GGGCTCGAGCAGAAGCGCAAGGAGCAGGAGAGCGCAACAGGAACTCCGCGACGCTGGCCCTCCCCATCGAAGTCCAAGATGA GCGTCGAGCTCAAGGAATTCGACACAATCGAACTGGAGGTGCTTATAAGCCCTTTGAAGACCTTCAAGACTCCCAAAAAATACCGC CACCCACCCCGACAAACATGAGCTTATGTGCGCGCTCACACTGACGTCTCAAGCGCCTTCTCTTCAGTCCGGCCAAAAATGGATC TCCTGTCAAATTGGTGGAG
<i>DmCdt1</i> ^{Phe→Leu}	ATGGCACAGCCTAGCGTCGCGCACTTTTAACCAATCGTAAGCGTGCCGCTTAGATGACGCCATTTGATTAAAGATCGCCGCTCTGG TTGAGCCCGCGGAAACAGTCAGCCCGGCAAGTGCCCGCTCGCAGTTGCCCGCCGGGATCAGGACGCAGATTTAGATACATTAAG CTGCGGCCACAGGTATGCGTACCCGTTCTGGGCGTACGGCCGCTTTGATTGTGACAGCGGCCAAGAGTCGAAAAAAGAGCCCCAG CTGTGCTAAAAATGGAACCATATCAAGCAACGCAAACTGGTCCAGTTAATCAAAAAAGGAACCTGAGCCCGCGTAAACAGCTC AGTCAAGTAAGTTGGATGAGGAAGAGTTGCAGCAGTCGTGACGCTATCCGAACACACCCCAAGGTGAATCTGACCATCACTAGTC AGCAAAACGCGGATAACGTTACGCGGCACTGCGTACCCCAACCAACAAATTTTGAAGGATGCCTCTCCAATTAAGGCAGATCTGC GTCGTGAGTACGTTAGACGAGGTAAGACCAAGGTGAGTCGCTCAGCAAGCTGCAGGAAGTGAAGGCTGTATTAGCTCTGAAGG CAGCCTTGGAACAGAAGCGCAAGGAGCAAGAAGCAACGCAACGCAAGCTGCGGACGCGGCGCGTCCGCAAGCAAGTCTAAGATG TCGGTACAACCTAAAGAATTAGACACTATTGAGTTAGAGGTCTTATCTCTCGCTGAAAAGCTTGAAGACCCCACTAAGATTCGCG CGCAACCCCTGATAAACACGAGTTAATGTCTCCCGTCATACCGAGCTATCGAAGCGCCTTTATTGAGTCTCGGAAGAACGGATC CCCAGTAAAAATTGGTCGAG
<i>DmCdt1</i> ^{Phe→Ala}	ATGGCACAGCCTCGGTAGCGGACGCGCCACGAATCGCAAGCGTGCCGCTTTAGACGATGCAATCAGTATTAAGAATCGCCGCTCTT GTAGAGCTCGGGAGACGCTCTCTCTGCAAGTGCTCTTCTCAGCTTCCAGCGGAGATCAGGACGCGGACTTAGATACACTGAAG

	<p>GCTGCCGAACCGAATGCGTACACGTTACAGGTCGCACTGCCGCTCTTATCGTGACAGCGCCAGGAGTCTAAAAAACTCCA GCAGCAGCAAAGATGGAACTCATATCAAGCAGCCAAACTGGTACAGGCGATCAAAAAGGGCACTCTGTCCCCACGCAAGCAAGC TCAATCGAGCAAAGTTAGACGAGGAGGAGTTGCAGCAGTCTAGCGCCATTTCGGAACATACTCCAAAAGTAAACGCGACGATCACTTC TCAACAAAATGCCGATAATGTCCAGCGTGGGTTACGCACGCCCTACAAAACAGATTTTGAAGGACGCTAGCCCTATTAAAGCCGATCT TCGTGCTCAACTGACCGCTGATGAAGTGAAGACCAAGGTAAGCCGTTCCGCAAACTGCAAGAACTTAAAGCCGTGCTGGCTTTAAA AGCGGCTTTGGAACAAAAGCGTAAAGAGCAGGAAGAGCGTAACCGTAAGTTGCGTGACGCAGGACCAAGCCCTTCAAAGTCCAAGA TGAGCGTACAACCTAAGGAAGCTGATACTATTGAATTGGAGGTAAGTCTCGCCTCTTAAGACTGCCAAGACGCCCAAAAAATCC CGCCCCCACACCTGACAAGCATGAACCTATGAGCCACGCCACACGGATGTGTCCAAGCGTTTGTCTGCCAGCCCCGCTAAGAAGC GATCCCCGGTCAAACCTGGTAGAG</p> <p>MAQPSVAAAATNRKRAALDDAISIKNRRLVEPAETVSPASAPSQLPAGDQDADLDTLKAATGMRTSRGRTARLIVTAAQESKKKTPAAA KMPEHIKQPKLVQAIKKGLTSPRKQAQSSKLDEEELQSSAISEHTPKVNFITTSQQNADNVQRGLRTPTKQJLKDASPIKADLRRLQTADDEV KTKVRSRAKLQELKAVLALKAALQKKEQEERNRKLDRAGSPSKSKMSVQLKEADTIELEVLIPLKTKTPTKIPPTPDKHELMSPRHDT VSKRLASPKNKSPVKLVE</p>
<i>DmCdt1</i> ^{Uniform}	<p>ATGGCTCAACCTTCTGTCAAAGCTGCATTCTTTACCCGCGAAAAACAAGCGCGGCACTTGGCGATCTCAAAGATTAACCTTGGCGAAG TGCCCGCAAAACTGTTTCTCCAGACGCTTCAAAAGCTCGTGAACCTAGTCAACTTCCGAAAGCAGGACAGGCTTTGACTGACCGTGA GAAACTGGCGGACGCCACAGGCAAGATGACTTCGGGTCGCGAAACCGCCGATAAGTTAATCGTAACGCGTCCAAACAATCACGTG AAACACCCGCCGCAAGGCGAGATATGCCGCATATTGAGAAGCGCGAACCCCTGGTGCAAGTTTCAATAGGGGACCTTAGACTCCCCTC GTGAGCAAAAAGCCGAGTCTTCTCTTTGAAACAACAGTCGCGTGAAGATTACGCCATTAAGTCACATACGCCCGTGAACAAATTCGG CGAAACAATCACAAGTGACCAAAAGCAAAACGCCAACGTACAGCGTGAGAAGGGCTTGACGCCGACACAAAAGGACATTTTAGCCT CACGCGAACCTATCAAGGCCTTGCAACTGACTTTTAAGGTGACTGATCGTGAAGTCTCATCTGCGAACTTCAATTAGCGGTTCTGAA GGCGAAGCCTTAGATGCAGCTCTGCAAAACAAAACCTTGCAGGTGCGGAGCCCAAAAGTCCAGTAGTGACATGCGAAGGTTCA GTTACGTGAGTTCACAATCAAACCTGGTTTGTATTTACCAGACAAGTTGCGTGAAACTTTCCTCAACCAAAAATCCGCCACCCACG CCCCGCGAGGATAAGCATTTAATGAGTCCCCACAAAACCGTTTCACTGCGCGAACTGTTCAAAATCCGACCTTGCTAATGGAAGTAAGC CGGTACGCGAGCTTGTCT</p> <p>MAQPSVKAFFTRENKAADLAISKINLREVPKTVSPDASKAREPSQLPKAGQALTDREKLAAATGKMTSGRETADKLIVTAAKQRETPAA KADMPHIKQREPLVQFIKGLTDSPREQKAQSSLLKQSSREDSAIKSHTPVNFRETITSDQKQNAVQREKGLTPQTQKILASREPIKALQTF KVTDREVSSAKLQAVLKREALDALQKQNLGAREPKSPSSDMKSVQLKEFTIKVLISPKLRETFTPTKIPPTPDKHELMSPRHDTVSKRLFS LFKSDPANGSKPVREL</p>
<i>DmCdt1</i> ^{Δ1-100}	<p>ATGCTGGTGCAATTCAATAAAAAGGGCACTCTGTGCGCCAGGAAACAGGCTCAGTCCAGTAAGCTGGACGAGGAGGAGCTGCAGCAG TCGTGCGCCATAAGCGAGCACACGCCCAAGGTTAACTTACCATTACACAAGCCAGCAGAAATGCGGACAATGTGCAGCGTGGCTGCGC ACACCCACCAAGCAGATCCTCAAGGATGCCTCGCCGATCAAGCGGATCTCCGCCGTCAGCTCACTTTCGACGAGGTAAAAACGAAG GTATCGCGGAGTGCCAAGCTGCAGGAATCAAGGCAGTGCTGCGCCCTTAAGCGCGCGCTCGAGCAGAAGCGCAAGGAGCAGGAGGA GGCACAACAGGAACTCCGCGACGCTGGCCCCCTCCCATCGAAGTCCAAGATGAGCGTGCACTCAAGGAATTCGACACAATCGAACT GGAGGTGCTTATAAGCCCTTTGAAGACCTTCAAGACTCCCAAAAAATACCGCCACCCACCCCGGACAAAACATGAGCTTATGTGCGCG CGTCACTAGTACGCTCTCAAGCGCCTTCTCTTCACTCGGCCAAAAATGGATCTCCTGTCAAATTTGGTGGAG</p> <p>MLVQFIKKGLTSPRKQAQSSKLDEEELQSSAISEHTPKVNFITTSQQNADNVQRGLRTPTKQJLKDASPIKADLRRLQTFDEVKTKVRSRAKL QELKAVLALKAALQKKEQEERNRKLDRAGSPSKSKMSVQLKEFDTIELEVLIPLKTFKTPTKIPPTPDKHELMSPRHDTVSKRLFS KNGSPVKLVE</p>
<i>DmCdt1</i> ^{Δ101-200}	<p>ATGGCCAGCCATCGGTAGCTGCATTTTCAAAAACCGCAACGCGCGCCTTGGATGATGCTATCAGTATCAAGAACAGGCGTTTGG TGGAACCCGCTGAAACCGTCTCTCTGCCTCGGCCCTTCCAGTTGCCAGCGCGGACACAGGATGCGGATCTAGACACCTGAAGGC GGCGGCCACGGGATGCGTACCCGATCCGGACGCACTGCCGACTAATTGTACCGCGCGCTCAAGAGAGCAAAAAGAAGACACCGGC TGCCGCCAAGATGGAGCCACACATCAAGCAGCCCAAGGCCCTTAAGCGCGCGCTCGAGCAGAAGCGCAAGGAGCAGGAGGAGCGCA ACAGGAACTCCGCGACGCTGGCCCCCTCCCATCGAAGTCCAAGATGAGCGTGCACTCAAGGAATTCGACACAATCGAACTGGAGG TGCTTATAAGCCCTTTGAAGACCTTCAAGACTCCCAAAAAATACCGCCACCCACCCCGGACAAAACATGAGCTTATGTGCGCGCTCA CACTGACGTCTCAAGCGCCTTCTCTTCACTCGGCCAAAAATGGATCTCCTGTCAAATTTGGTGGAG</p> <p>MAQPSVAAFFTNRKRAALDDAISIKNRRLVEPAETVSPASAPSQLPAGDQDADLDTLKAATGMRTSRGRTARLIVTAAQESKKKTPAAAK MEPHIKQPKALKAALQKKEQEERNRKLDRAGSPSKSKMSVQLKEFDTIELEVLIPLKTFKTPTKIPPTPDKHELMSPRHDTVSKRLFS PAKNGSPVKLVE</p>
<i>DmCdt1</i> ^{Δ201-297}	<p>ATGGCCAGCCATCGGTAGCTGCATTTTCAAAAACCGCAACGCGCGCCTTGGATGATGCTATCAGTATCAAGAACAGGCGTTTGG TGGAACCCGCTGAAACCGTCTCTCTGCCTCGGCCCTTCCAGTTGCCAGCGCGGACACAGGATGCGGATCTAGACACCTGAAGGC GGCGGCCACGGGATGCGTACCCGATCCGGACGCACTGCCGACTAATTGTACCGCGCGCTCAAGAGAGCAAAAAGAAGACACCGGC TGCCGCCAAGATGGAGCCACACATCAAGCAGCCCAAGCTGGTGCAATTCAATAAAAAGGGCACTCTGTGCGCCAGGAAACAGGCTCA GTCCAGTAAGCTGGACGAGGAGGAGCTGCAGCAGTCTGCGGCATAAGCGAGCACACGCCCAAGGTTAACTTCAACATCACAAGCCA GCAGAATGCGGACAATGTGCAGCGTGGCCTGCGCACACCCACCAAGCAGATCCTCAAGGATGCCTCGCCGATCAAGGCGGATCTCCG</p>

	<p>CCGTCAGCTCACTTTCGACGAGGTAAAAACGAAGGTATCGCGGAGTGCCAAGCTGCAGGAAGCTCAAGGCAGTGCTG</p> <p>MAQPSVAFFFTNRKRAALDDAISIKNRRLVEPAETVSPASAPSQLPAGDQDADLDTLKAATGMRTSRGRTARLIVTAAQESKKKTPAAAK</p> <p>MEPHIKQPKLVQFIKKGTLSPRKQAQSSKLDEEELQSSAISEHTPKVNFITTSQQNADNVQRGLRTPTKQILKDasPIKADLRRQLTFDEVKT</p> <p>KVSRSAKLQELKAVL</p>
<i>DmCdt1</i> ^{ΔCoiled-coil}	<p>ATGGCCCAGCCATCGGTAGTCATTTTTACAAACCGCAAACGCGCGCCTTGGATGATGCTATCAGTATCAAGAACAGGCGTTTGG</p> <p>TGGAACCCGCTGAAACCGTCTCTCCTGCTCGGCCCTTCCCAGTTGCCAGCGCGGACCAGGATGCGGATCTAGACACCTGAAGGC</p> <p>GGCGGCCACGGCATGCGTACCCGATCCGGACGCACTGCCGACTAATTGTACCGCGCTCAAGAGAGCAAAAAGAAGACACCGGC</p> <p>TGCCGCAAGATGGAGCCACACATCAAGCAGCCCAAGCTGGTGAATTCATTAAGGCACTCTGTCGCCAGGAAACAGGCTCA</p> <p>GTCCAGTAAGCTGGACGAGGAGGAGCTGCAGAGTCGTCGCCATAAGCGAGCACACGCCCAAGGTTAACTTACCATCACAAGCCA</p> <p>GCAGAATGCGGACAATGTGCAGCGTGGCCTGCGCACACCCACCAAGCAGATCCTCAAGGATGCTCGCCGATCAAGGCGGATCTCCG</p> <p>CCGTAGCTCACTTTCGACGAGGTAAAAACGAAGGTATCGCGGAGTGCCAAGCTGCAGGAAGGCCCTCCCATCGAAGTCCAAGAT</p> <p>GAGCGTGCAGCTCAAGGAATTCGACACAATCGAACTGGAGGTGCTTATAAGCCCTTTGAAGACCTTCAAGACTCCCAAAAAATACC</p> <p>GCCACCCACCCGGACAAACATGAGCTTATGTCGCGCGCTCACACTGACGCTCTCAAGCGCCTTCTTTCAGTCGGGCCAAAAATGGA</p> <p>TCTCTGTCAAATTGGTGGAG</p> <p>MAQPSVAFFFTNRKRAALDDAISIKNRRLVEPAETVSPASAPSQLPAGDQDADLDTLKAATGMRTSRGRTARLIVTAAQESKKKTPAAAK</p> <p>MEPHIKQPKLVQFIKKGTLSPRKQAQSSKLDEEELQSSAISEHTPKVNFITTSQQNADNVQRGLRTPTKQILKDasPIKADLRRQLTFDEVKT</p> <p>KVSRSAKLQEGSPSKSMVQLKEFDITIEVLISPLTKFTPTKIPPTPDKHELMSPRHTDVSKRLLFSPAKNGSPVKLVE</p>
<i>DmCdt1</i> ^{ScRIDR}	<p>ATGTTAACAGAGGATTTGCAAAAGTTTACCCCGCCAAAGCGCGGGATTACGACAAGTCTTGAAAAACGAGTCCCGCCGCTCACAG</p> <p>AAACCTATCTCGCAATTGCCCAAGGATATAAAAAAGAGCACCGCCGACGGAGCAAGAGGCCGAGTCAAAAGCACAGGCCGCGGA</p> <p>ACTTGAGAAAGATCCTCAGCTTGACTCAATCCGTGCGTTGGTAAACGCCCCGAGGTCAAAATTTACAAACAGCGTCGCCAAGCTGTT</p> <p>GCAGCGGAGAGTTAGAGGCTGGGAAGGAAGCAGGTTTGACCGCTGTTAAGGACAAGGCGCAGAGGAAGCCCGCGTTCACCCAA</p> <p>GGATTGCTGAATTTGACATCCGCTCGCGCTCAGCCGTGCGGAGCGTGAGTAAGTTATCCCGCTGATTGATGCAAAAGCTGTGGAA</p> <p>ACTTTATTAGATACAAAGACCGCCCTACTGCCCGGCCATTAAAGCCGCTATGAACAAGCAGCCACGCGACCATAAGACCCAACAA</p> <p>ACCCCGCCCAACCAAGGAGCGCGCATCCGCACACGTATGAAGCTATCCAAGCAGAGGCCGCTACTCACAGCAGCCGCAACACAG</p> <p>TTACACCTTCAAAGGGGAAGTGGAGTTACCAAGAGCAAAACAAAGCTTGCGACTGCAAAACCGCGCAGGTTAAACCCAGATGTTG</p> <p>AGTGACTTGTGAGGCCTGAACACACTATGCTTACAAAGAATAGCCCCGCACGCACGATCCAAGTGAGTCAATTTTCGCGTCCGAAC</p> <p>TCTTGGTAGATGATAAGTTTGCCTTGGCCCTGAGCAAGGTCAAGATCCATCTGAGTTTCGATAATTCCTGCCCGTAGTCAAGTCTCG</p> <p>TTAGTCTTAGACATCGCG</p> <p>MLTEDLQKFTPPKRGITTSLEKTPAASQKPIQLPKDTKKKAPTEQEAEVKAPGRELEKDPQLDSIRRLVNAPKVKISQRRQAVAAGELE</p> <p>AGKEAGLTAVKDKGAEARRSPKDSLNFDIRSRAVASVSKLSPLIDAKLLETLLDTKTARTAPAIPKPRMNKQPRDHKTQQRQPKPEAIR</p> <p>RMKAIQAEAAATHSSRKQFTFKGEVELPKSQKLATANPAQVKTQMLSDLSEPEHTMLTKNSPARTIQVSQFSRPNFLVDDKFALALSKVKIH</p> <p>LSFDNSLPVVKSSLVLDIA</p>
<i>DmCdt1</i> ^{Ile/Leu(0.5)}	<p>ATGGCCCAGCCCTCCGTTGCTGCGTTTTTTACAAACCGCAAGCGTGCTGCAGCGGATGATGCCGCGAGTATCAAGAACCGCGCCTTG</p> <p>TGGAGCCAGCGGAAACCGTGTGCGCAGCCTCTGCACCTTCCCAAGCTCCAGCAGGTGACCAGGATGCTGATTGGACACGGCAAGG</p> <p>CCGCCGCCACTGGCATGCGTACGCGCTCCGGACGCACGGCTCGTTAGCTGTAACCGCTGCTCAAGAGTCCAAAAGAAAAACACCTG</p> <p>CAGCAGCAAAGATGGAACCGCACATCAAAACAACCGAAGGCCCTCAATTTCGCAAAAAAAGGGACACTTTCGCCCCGTAAACAAGCC</p> <p>CAGTCATCCAAGGCTGACGAAGAGGAGCTTCAACAATCTTCGCAATTAGTGAACACACGCCTAAGGTGAATTTCACTGCTACTTCAC</p> <p>AACAAAATGCAGATAATGTACAGCGGAGCTCGCACTCCAAACAAAACAGCGCTTAAAGATGCCAGTCCCATTAAGCAGATGCTC</p> <p>GCCGCCAACTTACTTTGACGAGGTAAAGACGAAGGTGAGCCGTTCGCGAAGGCGCAGGAGCTTAAAGCTGTGGCCGCCCTGAAGG</p> <p>CCGCGGTGAGCAGAAACGCAAGAACAGGAGGAGCGCAATCGTAAGCTTCGTGATGCGGGGCCGTGCGCAAGCAAAATCGAAGATG</p> <p>AGTGTGCAAGCAAAGGAGTTTGATACAGCTGAGCTGGAGGTGCGCAATTTACCGCTGAAGACGTTTAAAGACCCGACCAAGCTCCG</p> <p>CCACCCACCCAGATAAGCATGAGGCGATGTCGCTCGCCACACAGAGCTGAGCAAGCGTTTAGCATTTTCGCTGCCAAAAATGGA</p> <p>AGTCCCGTTAAACTGGTTGAA</p> <p>MAQPSVAFFFTNRKRAADDAASIKNRRLVEPAETVSPASAPSQLPAGDQDADLDTAKAAATGMRTSRGRTARLAVTAAQESKKKTPAA</p> <p>AKMEPHIKQPKAVQFAKKGTLSPRKQAQSSKADEEELQSSAISEHTPKVNFITTSQQNADNVQRGARTPTKQALKDasPIKADARRQLTF</p> <p>DEVKTKVSRSAKQELKAVAALKAEEQKRKEQEERNRKLRDAGSPSPSKSMVQAKEFDTAEELEVAISPLTKFTPTKAPPTPDKHEAMS</p> <p>PRHTDVSKRLAFSPAKNGSPVKLVE</p>
<i>DmCdt1</i> ^{Ile/Leu(1.0)}	<p>ATGGCGCAACCATCGGTGGCGCCTTCTTTACAAACCGTAAACGCGCGGCAGCCGACGATGCTGCGAGTGCGAAGAATCGCCGTGCC</p> <p>GTGGAACAGCAGAGACCGTATCACCTGCGCTCTGCGCCTTCTCAGGCACCGCGGGAGACCAAGATGCTGATGCAGACACCGCAAAAG</p> <p>GCAGCGCTACGGGATGCGCACGCGTTCGGTGCCTAGTCCGCTGCTGCGCTTACTGCCGCCAAGAGAGTAAAAAAGACGCCA</p> <p>GCTGTGCTAAAAATGGAGCCGACGCAAAACAGCCTAAAGCAGTGCAGTTTTCGCAAAAAAGGCACCGCAAGTCTCGCAAGCAAGC</p> <p>CCAGTCTAGTAAGGCTGACGAGGAAGAGGCTCAACAATCGTCTGCGCCTCCGAACATACTCCGAAGGTTAATTTTACGGCAACGTC</p> <p>ACAGCAGAATGCCGATAATGTCCAGCGTGGCGCTCGCACCCCAACGAAGCAGGCTGCAAAAGATGCCAGCCAGCGAAAGCCGATG</p> <p>CACGCGCTCAGGCAACATTGATGAGGTCAAGACCAAGGTATCACGCTCGGCCAAAGCGCAAGAAGCTAAGGCGGTAGCGGCAGCC</p>

	AAAGCGGCTGCCGAACAAAAACGTAAGGAGCAGGAGGAGCGTAATCGCAAAGCTCGCGATGCGGGTCCATCACCAGTAAATCAAA GATGTCTGTACAGGCGAAAGAATTGACACGGCGGAAGCCGAAGTAGCTGCCTCCCCAGCGAAGACTTTTAAGACTCTACAAAAGC ACCGCCACCTACTCCAGACAAGCATGAGGCTATGTACCGCGTCACACCGACGTCTCCAAGCGTGCTGCATTCTCACCTGCCAAGAAC GGATCTCCAGTAAAAGCTGTCGAG
	MAQPSVAFFTNRKRAAADDAASAKNRRAVEPAETVSPASAPSQAPAGDQDADADTAKAAATGMRTRSGRTARAAVTAAQESKKKTPA AAKMEPHAKQPKAVQFAKKGTASPRKQAQSSKADEEEAQSSAASEHTPKVNFTATSQQNADNVQRGARTPTKQAAKDASPAKADARR QATFDEVKTKVSRSAKAQEAQAAKAAAEQKRKEQEERNRKARDAGPSPSKMSVQAKEFDTAEEVAASPAKTFKTPKAPPTPD KHEAMSPRHDTVSKRAAFSPAKNGSPVKAVE

ACKNOWLEDGEMENTS

We thank past and present members of the Berger and Botchan labs for helpful discussion and advice. This work was supported by an NIH NRSA postdoctoral fellowship (F32GM116393, to M.W.P.), by the UC Berkeley Jesse Rabinowitz Award (to J.A.K.), by the NCI (R01CA030490, to J.M.B. and M.R.B.) and by the NIGMS (R01GM141045-01, to J.M.B. and M.R.B.). M.W.P. is the Cecil H. and Ida Green Endowed Scholar in Biomedical Computational Science. Research in his lab is supported in part by CPRIT (RR200070) and the Welch foundation (I-2074-20210327).

FIGURES, FIGURE TITLES AND LEGENDS

FIGURE 1

Figure 1: Initiator phase separation is insensitive to 1,6-hexanediol and does not require aromatic residues. A) Heatmap representation of the amino acid composition of the three *Drosophila* initiation factors (Cdt1, Orc1 and Cdc6) compared to other phase separating disordered sequences. B) Confocal fluorescence microscopy images of condensates formed with 5 μ M Cy5-dsDNA and 5 μ M Cdt1 in the absence (top) and presence (bottom) of 1,6-hexanediol. C) 2 μ M Cdt1 was combined with 2 μ M dsDNA and phase separation was assessed with a depletion assay in the presence of increasing concentrations (w/v) of 1,6-HD. Phase separation is indicated by protein loss. D) Aromatic residues contribute to the phase separation of many condensate systems through their ability to participate in π - π and π -cation interactions. E) Schematic of the *D. melanogaster* Cdt1 protein (top). Cdt1 contains two winged-helix domains (WHd, blue) and an N-terminal disordered domain (black line). The N-terminal IDR contains multiple aromatic residues distributed throughout its length (bottom). F) Depletion assay results assessing phase separation of the Cdt1 aromatic mutant, Cdt1^{Phe→Leu}. For this construct all phenylalanine residues have been mutated to leucine. “L” is a load control, “-” is in the absence of DNA, and “+” is in the presence of DNA. G) Confocal fluorescence microscopy images of condensates formed with 5 μ M Cy5-dsDNA and either Cdt1^{Phe→Leu} (top) or Cdt1^{Phe→Ala} (bottom). Gel and microscopy images are representative from three independent experiments.

FIGURE 2

Figure 2: DNA contributes adhesive interactions to Cdt1 LLPS by acting as a counterion bridge. A) DNA-induced phase separation was assessed by the depletion assay in buffer containing 75, 150 or 300 mM potassium glutamate (KGlut). Cdt1 LLPS was observed at 75 and 150 mM KGlut, but not at 300 mM KGlut. B) Multiple anionic polyelectrolytes were assessed for their ability to induce Cdt1 phase separation by the depletion assay. “L” indicates a load control and “-” is in the absence of any anion polymer. dsDNA, ssDNA, dsRNA, heparin and poly-Glutamate (poly-Glu) all effectively induced Cdt1 LLPS. A t-test was used to calculate whether a significant change in pelleting resulted from the addition of polyanion compared to the negative (“-”) control. C) Two models explaining the ability of DNA to induce Cdt1 phase separation. (Top) DNA (black/red) functions as a counterion bridge to facilitate inter-IDR (blue/red) interactions. (Bottom) DNA neutralizes inter-IDR electrostatic repulsion to drive self-assembly. D) Monovalent (KH_2PO_4) and multivalent (K_2HPO_4 and K_3PO_4) phosphate counterions were assessed for the ability to induce Cdt1 phase separation by the depletion assay and fluorescence microscopy. For the line plot (left), the fraction of Cdt1 retained in the depletion assay was quantitated and plotted against counterion concentration (mM). For the microscopy experiment (right), 5 μM eGFP-Cdt1 was combined with 2 M KCl, 625 mM KH_2PO_4 , 625 mM K_2HPO_4 or 625 mM K_3PO_4 (top to bottom) and assessed for droplet formation by confocal fluorescence microscopy. Concentrated and highly networked species were observed only in the presence of K_3PO_4 . E) Plot of the net charge per residue (NCPR) over a five-residue sliding window for Cdt1 and the charged residue variant, Cdt1^{Uniform}. F-G) Phase diagrams for stoichiometric

combinations of Cy5-dsDNA and Cdt1 (F) or Cdt1^{Uniform} (G) at different protein and KGlutamate concentrations. Tiles bordered in green show conditions where phase separation was observed. Each tile is 75x75 μm . The mean signal intensity for Cy5-dsDNA within each droplet was quantitated for 5 μM Cdt1 (grey) and Cdt1^{Uniform} (red) across the four salt concentrations. All pairwise comparisons are significantly different ($p < 0.05$ as determined by t-test) . H) Analytical size exclusion chromatography analysis of Cdt1 and Cdt1^{Uniform}. Cdt1^{Uniform} adopts a more extended conformation in solution as evidenced by a lower retention volume and co-purifies with nucleic acid ($260/280 = 0.56$ and 0.79 for Cdt1 and Cdt1^{Uniform}, respectively). “*” = $p < 0.05$, t-test. Gel and microscopy images are representative from three independent experiments.

FIGURE 3

Figure 3: Inter-IDR cohesive interactions drive Cdt1 phase separation in the absence of DNA. A) (Top) DNA (black/red) can induce Cdt1 (black/blue) phase separation by acting as a counterion bridge. (Bottom) To what extent homotypic inter-IDR interactions affect phase separation was unknown. B) Depletion assay to assess phase separation of 8 μ M Cdt1 in the presence of increasing concentrations of PEG-3350 (0-12% w/v). C) Phase separation of 5 μ M eGFP-labeled Cdt1 was assessed by confocal fluorescence microscopy in the absence (top) and presence (bottom) of 4% PEG-3350. D) Schematic representation of the domain architecture of *D. melanogaster* Cdt1 (top) with coiled-coil domain prediction for the Cdt1 N-terminal IDR (bottom). The position of the three approximately 100 amino acid deletions used in Figure 3E are indicated. E) Cdt1 Δ 1-100 (left), Cdt1 Δ 101-200 (middle), and Cdt1 Δ 201-297 (right) were assayed for the ability to undergo PEG-induced phase separation by the depletion assay. “L” is a load control, “-” is in the absence of PEG, and “+” is in the presence of PEG. F) Quantitation of depletion assay results presented in Figure 3E. A t-test was used to calculate whether a significant change in pelleting occurred in the presence compared to the absence of PEG. G) Dotplot analysis (EMBOSS Dotmatcher, (Madeira et al., 2019)) of repetitive elements in the human FUS intrinsically disordered domain (top) and Cdt1 N-terminal IDR (bottom). The region representing Cdt1 amino acids 201-297 are highlighted with a grey background. The main diagonal represents sequence homology with itself and diagonals off the main diagonal represent repetitive motifs. Many repetitive motifs are apparent in FUS but are lacking in Cdt1. H) Dotplot analysis of the Cdt1^{ScrIDR}, a Cdt1 variant with N-terminal IDR residues randomly scrambled, versus

the wild-type Cdt1 IDR. I) Depletion assay to assess phase separation of Cdt1^{ScrIDR}. “L” is a load control, “-” is in the absence of PEG, and “+” is in the presence of PEG. J) Phase diagram for stoichiometric amounts of Cdt1^{ScrIDR} and Cy5-dsDNA at variable protein and KGlutamate concentrations. Tiles bordered in green show conditions where phase separation was observed. Each tile is 75x75 μm . “*” = $p < 0.05$, t-test. Gel and microscopy images are representative from three independent experiments.

FIGURE 4

Figure 4: Branched hydrophobic residues underlie Cdt1 homotypic inter-IDR interactions. A) The depletion assay was used to assess the salt-sensitivity of PEG-induced phase separation side-by-side with DNA-induced phase separation. “LOAD” is a load control, “-” is in the absence of DNA and PEG, “D” indicates the addition of DNA and “P” indicates the addition of PEG-3350. Cdt1 phase separation was assessed at 75, 150 and 300 mM KGlutamate (“KGlut”). B) Comparison of initiator IDR hydropathy (dashed line indicates the average hydropathy of the Orc1, Cdc6 and Cdt1 IDRs) to the hydropathy of predicted ordered domains (“Order”) and disordered domains (“Disorder”) proteome wide. C) Percent difference in initiator IDR amino acid composition relative to all predicted ordered (top) and disordered (bottom) domains proteome wide. The amino acids are

ordered from high to low hydropathy (left to right) and those with a positive hydropathy value are indicated by a green background. D) Comparison of initiator IDR fraction Leu, Ile and Val (dashed line indicates the average fraction-L,I,V of the Orc1, Cdc6 and Cdt1 IDRs) to the fraction-L,I,V of all predicted ordered domains (“Order”) and disordered domains (“Disorder”) proteome wide. E) To assess the distribution of branched hydrophobic residues, the fraction-L,I,V was calculated over a 5-residue sliding window for the Cdt1 IDR (top). Two mutants were made with either half (Cdt1^{Ile/Leu(0.5)}) or all (Cdt1^{Ile/Leu(1.0)}) of Leu and Ile residues mutated to Ala, and the distribution of L,I,V was calculated for each (middle and bottom, respectively). F) The depletion assay was used to assess the phase separation capacity of Cdt1^{Ile/Leu(0.5)} (top gel) and Cdt1^{Ile/Leu(1.0)} (bottom gel) in the presence of increasing concentrations of PEG-3350 (0-12% w/v). The fraction of protein retained at each concentration of PEG-3350 was quantitated. A t-test was used to calculate whether the different PEG concentrations resulted a significant change in pelleting relative to the no PEG control. No significant (“NS”) dose-dependent trend was identified for the effect of PEG on Cdt1^{Ile/Leu(0.5)} or Cdt1^{Ile/Leu(1.0)} phase separation. Gel images are representative from three independent experiments.

Figure 5: Homotypic inter-IDR interactions underlie DNA-induced initiator phase separation. A) An electrophoretic mobility shift assay (EMSA) was used to assess the binding affinity of Cdt1 (left), Cdt1^{Ile/Leu(0.5)} (middle) and Cdt1^{Ile/Leu(1.0)} (right) for dsDNA. Binding was assessed with 10 nM IRDye800-labeled dsDNA and a titration of Cdt1 from 16 to 1000 nM. Each variant showed comparable affinity to wild-type Cdt1. B) Depletion assay results assessing DNA-induced phase separation of Cdt1 (left), Cdt1^{Ile/Leu(0.5)} (middle) and Cdt1^{Ile/Leu(1.0)} (right). “L” is a load control, “-” is in the absence of DNA, and “+” is in the presence of DNA. C) Quantitation of the depletion assay results presented in Figure 5B. D-E) Phase diagram for stoichiometric amounts of Cdt1^{Ile/Leu(0.5)} (D) or Cdt1^{Ile/Leu(1.0)} (E) and Cy5-dsDNA at variable protein and KGlutamate concentrations. Tiles bordered in green show conditions where phase separation was observed. Each tile is 75x75 μ m. F) Violin plot representation of the mean droplet intensity of Cy5-dsDNA at 5 μ M protein concentration and variable KGlutamate concentrations for wild-type (white), Cdt1^{Ile/Leu(0.5)} (orange) or Cdt1^{Ile/Leu(1.0)} (red). G) Our data support a

model where, in the absence of DNA, inter-IDR electrostatic repulsion prevents spontaneous phase separation. The addition of DNA and resulting IDR-DNA interactions neutralize the initiator IDR charge which drives self-assembly through synergistic heteromeric DNA-bridging interactions and homomeric IDR-IDR hydrophobic interactions. “*” = $p < 0.05$, t-test. Gel and microscopy images are representative from three independent experiments.

1006

1007

SUPPLEMENTAL FIGURES AND LEGENDS

FIGURE 1—figure supplement 1

Figure 1—figure supplement 1: Heatmap analysis and complexity calculations for phase-separating disordered domains. A) Heatmap analysis of the amino acid composition of various phase-separating disordered sequences. Initiator IDRs are listed first and the remaining sequences are ordered according to their complexity score. B) Theoretical sequences with the maximum and minimum complexity score. C) Complexity scores for all predicted ordered (grey) and disordered (red) domains across the *D. melanogaster* proteome were calculated and plotted as a histogram. These visualizations are expanded in the right panels with the initiator complexity scores indicated by a dashed line.

FIGURE 1—figure supplement 2

Figure 1—figure supplement 2: *D. melanogaster* ORC and Cdc6 phase separation is resistant to treatment with 1,6-hexanediol. A) 2.5 μM ORC was combined with 2.5 μM Cy5-dsDNA and phase separation assessed by confocal fluorescence microscopy in the absence (left) and presence (right) of 10% 1,6-hexanediol. B) 5 μM Cdc6 was combined with 5 μM Cy5-dsDNA and phase separation assessed by confocal fluorescence microscopy in the absence (left) and presence (right) of 10% 1,6-hexanediol.

FIGURE 2–figure supplement 1

Figure 2–figure supplement 1: Quantitative comparison of DNA enrichment in condensates formed by 5 μM wild-type and variant Cdt1s at 50, 100, 150 and 300 mM KGlutamate concentrations. Images from phase diagram experiments were used to calculate the mean Cy5-dsDNA signal in condensates and individual condensate values were plotted as a scatter plot for each experimental condition. Each dot represents a different condensate. Red horizontal bars indicate mean intensity overall.

FIGURE 3—figure supplement 1

Figure 3—figure supplement 1: Depletion assay analysis of PEG-induced phase separation propensity for a Cdt1 variant lacking a predicted coiled-coil region. “L” is a load control, “-” is in the absence of PEG, and “+” is in the presence of PEG.

FIGURE 3—figure supplement 2

Figure 3 – figure supplement 2: Dotplot analysis of various phase-separating IDRs to identify the presence of repetitive motifs. EMBOSS Dotmatcher (Madeira et al., 2019) was used to assess the presence of repetitive elements in the IDRs of A) Laf1, B) Ddx4, C) hnRNPA1, D) RPB1, E) MED1, and F) BuGZ. All proteins analyzed contain short repetitive motifs.

FIGURE 3—figure supplement 3

Figure 3—figure supplement 3: Cdt1^{ScrIDR} condensates can recruit wild-type Cdt1. A) Droplets were prepared with 5 μ M Cy5-dsDNA and either 5 μ M wild-type Cdt1 (left two columns) or 5 μ M Cdt1^{ScrIDR} (right two columns) and 1 μ M eGFP-tagged Cdt1 was subsequently added to each reaction. Samples were imaged by fluorescence microscopy. B) Quantitation of panel A). The level of eGFP-Cdt1 recruitment was assessed by calculating green signal intensity within the droplets containing Cy5-dsDNA and either Cdt1 or Cdt1^{ScrIDR}.

FIGURE 4-figure supplement 1

Figure 4-figure supplement 1: DNA-induced phase separation of ORC and Cdc6 is sensitive to salt but PEG-induced LLPS is not. The depletion assay was used to assess the salt-sensitivity of PEG-induced phase separation side-by-side with DNA-induced phase separation. Phase separation of ORC (2 μ M) and Cdc6 (4 μ M) was assessed at 150 and 300 mM KGlutamate (“KGlut”). “LOAD” is a load control, “-” is in the absence of DNA and PEG.

FIGURE 5—figure supplement 1

Figure 5—figure supplement 1: Heparin binding propensity of Cdt1 and the Cdt1 branched hydrophobic residue variants, Cdt1^{Ile/Leu(0.5)} and Cdt1^{Ile/Leu(1.0)}. Cdt1 (top) and Cdt1 variants (middle and bottom) were applied to a heparin column and eluted off with a linear gradient of 150 mM – 1 M NaCl. All three proteins elute off at similar salt concentrations.

FIGURE 5—figure supplement 2

Figure 5—figure supplement 2: Analysis of Cdt1 IDR conservation amongst *Drosophilidae* and metazoans. A) Domain architecture of *D. melanogaster* Cdt1. B) Sequence identity analysis for eleven metazoan Cdt1 orthologs. The blue regions correspond to the globular, winged helix domains (WHd) of the protein. C) Sequence identity analysis for forty *Drosophilidae* Cdt1 orthologs. D) Scatter plot analysis of the number of minimal ([S/T]P) or full ([S/T]PX[R/K]) CDK/Cyc consensus motifs in metazoan IDRs versus IDR length. E) Scatter plot analysis of the number of minimal ([S/T]P) or full ([S/T]PX[R/K]) CDK/Cyc consensus motifs in *Drosophilidae* IDRs versus IDR length.

FIGURE 5—figure supplement 3

Figure 5—figure supplement 3: Heatmap analysis of the amino acid composition of initiator IDRs (Cdt1, Orc1, Cdc6) and of an internal IDR from Chiffon, the fly homolog of Dbf4. These four proteins have similar amino acid sequence signatures, suggesting that Chiffon may undergo a similar DNA-dependent phase separation reaction or may co-partition into initiator condensates.

REFERENCES

- Andersen OS. 2013. Cellular Electrolyte Metabolism Encyclopedia of Metalloproteins. Springer New York. pp. 580–587.
- Babinchak WM, Haider R, Dumm BK, Sarkar P, Surewicz K, Choi JK, Surewicz WK. 2019. The role of liquid-liquid phase separation in aggregation of the TDP-43 low-complexity domain. *J Biol Chem*. doi:10.1074/jbc.RA118.007222
- Baldinger T, Gossen M. 2009. Binding of Drosophila ORC proteins to anaphase chromosomes requires cessation of mitotic cyclin-dependent kinase activity. *Mol Cell Biol* **29**:140–149. doi:10.1128/MCB.00981-08
- Banani SF, Lee HO, Hyman AA, Rosen MK. 2017. Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**:285–298. doi:10.1038/nrm.2017.7
- Banjade S, Rosen MK. 2014. Phase transitions of multivalent proteins can promote clustering of membrane receptors. *Elife*. doi:10.7554/eLife.04123
- Batty P, Gerlich DW. 2019. Mitotic Chromosome Mechanics: How Cells Segregate Their Genome. *Trends Cell Biol*. doi:10.1016/j.tcb.2019.05.007
- Berry J, Weber SC, Vaidya N, Haataja M, Brangwynne CP, Weitz DA. 2015. RNA transcription modulates phase transition-driven nuclear body assembly. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1509317112
- Beutel O, Maraschini R, Pombo-García K, Martin-Lemaitre C, Honigsmann A. 2019. Phase Separation of Zonula Occludens Proteins Drives Formation of Tight Junctions. *Cell*. doi:10.1016/j.cell.2019.10.011
- Boehning M, Dugast-Darzacq C, Rankovic M, Hansen AS, Yu T, Marie-Nelly H, McSwiggen DT, Kokic G, Dailey GM, Cramer P, Darzacq X, Zweckstetter M. 2018. RNA polymerase II clustering through carboxy-terminal domain phase separation. *Nat Struct Mol Biol*. doi:10.1038/s41594-018-0112-y
- Boeynaems S, Holehouse AS, Weinhardt V, Kovacs D, Van Lindt J, Larabell C, Bosch L Van Den, Das R, Tompa PS, Pappu R V., Gitler AD. 2019. Spontaneous driving forces give rise to protein–RNA condensates with coexisting phases and complex material properties. *Proc*

1222 *Natl Acad Sci U S A*. doi:10.1073/pnas.1821038116

1223 Booth DG, Earnshaw WC. 2017. Ki-67 and the Chromosome Periphery Compartment in Mitosis.

1224 *Trends Cell Biol*. doi:10.1016/j.tcb.2017.08.001

1225 Brodsky S, Jana T, Mittelman K, Chapal M, Kumar DK, Carmi M, Barkai N. 2020. Intrinsically

1226 Disordered Regions Direct Transcription Factor In Vivo Binding Specificity. *Mol Cell*.

1227 doi:10.1016/j.molcel.2020.05.032

1228 Chiu YP, Sun YC, Qiu DC, Lin YH, Chen YQ, Kuo JC, Huang J rong. 2020. Liquid-liquid phase

1229 separation and extracellular multivalent interactions in the tale of galectin-3. *Nat Commun*.

1230 doi:10.1038/s41467-020-15007-3

1231 Cho WK, Spille JH, Hecht M, Lee C, Li C, Grube V, Cisse II. 2018. Mediator and RNA

1232 polymerase II clusters associate in transcription-dependent condensates. *Science (80-)*.

1233 doi:10.1126/science.aar4199

1234 Chong SS, Dugast-Darzacq C, Liu Z, Dong P, Dailey GM, Cattoglio C, Heckert A, Banala S,

1235 Lavis L, Darzacq X, Tjian R. 2018. Imaging dynamic and selective low-complexity domain

1236 interactions that control gene transcription. *Science (80-)* **361**:378-+.

1237 doi:10.1126/science.aar2555;ear2555

1238 Deegan TD, Yeeles JT, Diffley JF. 2016. Phosphopeptide binding by Sld3 links Dbf4-dependent

1239 kinase to MCM replicative helicase activation . *EMBO J*. doi:10.15252/emj.201593552

1240 Dimitrova DS, Prokhorova TA, Blow JJ, Todorov IT, Gilbert DM. 2002. Mammalian nuclei

1241 become licensed for DNA replication during late telophase. *J Cell Sci*.

1242 Earnshaw WC, Laemmli UK. 1983. Architecture of metaphase chromosomes and chromosome

1243 scaffolds. *J Cell Biol* **96**:84. doi:10.1083/JCB.96.1.84

1244 Elbaum-Garfinkle S. 2019. Matter over mind: Liquid phase separation and neurodegeneration. *J*

1245 *Biol Chem*. doi:10.1074/jbc.REV118.001188

1246 Elbaum-Garfinkle S, Kim Y, Szczepaniak K, Chen CC, Eckmann CR, Myong S, Brangwynne

1247 CP. 2015. The disordered P granule protein LAF-1 drives phase separation into droplets

1248 with tunable viscosity and dynamics. *Proc Natl Acad Sci U S A* **112**:7189–7194.

1249 doi:10.1073/pnas.1504822112

1250 Feric M, Vaidya N, Harmon TS, Mitrea DM, Zhu L, Richardson TM, Kriwacki RW, Pappu R

1251 V., Brangwynne CP. 2016. Coexisting Liquid Phases Underlie Nucleolar Subcompartments.

1252 *Cell*. doi:10.1016/j.cell.2016.04.047

1253 Ferrolino MC, Mitrea DM, Michael JR, Kriwacki RW. 2018. Compositional adaptability in
 1254 NPM1-SURF6 scaffolding networks enabled by dynamic switching of phase separation
 1255 mechanisms. *Nat Commun* 2018 91 **9**:1–11. doi:10.1038/s41467-018-07530-1

1256 Findeisen M, El-Denary M, Kapitza T, Graf R, Strausfeld U. 1999. Cyclin A-dependent kinase
 1257 activity affects chromatin binding of ORC, Cdc6, and MCM in egg extracts of *Xenopus*
 1258 *laevis*. *Eur J Biochem* **264**:415–426.

1259 Gibcus JH, Samejima K, Goloborodko A, Samejima I, Naumova N, Nuebler J, Kanemaki MT,
 1260 Xie L, Paulson JR, Earnshaw WC, Mirny LA, Dekker J. 2018. A pathway for mitotic
 1261 chromosome formation. *Science* (80-) **359**.
 1262 doi:10.1126/SCIENCE.AAO6135/SUPPL_FILE/PAP.PDF

1263 Goloborodko A, Imakaev M V., Marko JF, Mirny L. 2016. Compaction and segregation of sister
 1264 chromatids via active loop extrusion. *Elife*. doi:10.7554/eLife.14864

1265 Guillén-Boixet J, Kopach A, Holehouse AS, Wittmann S, Jahnel M, Schlübler R, Kim K,
 1266 Trussina IREA, Wang J, Mateju D, Poser I, Maharana S, Ruer-Gruß M, Richter D, Zhang
 1267 X, Chang YT, Guck J, Honigmann A, Mahamid J, Hyman AA, Pappu R V., Alberti S,
 1268 Franzmann TM. 2020. RNA-Induced Conformational Switching and Clustering of G3BP
 1269 Drive Stress Granule Assembly by Condensation. *Cell*. doi:10.1016/j.cell.2020.03.049

1270 Hirano T, Mitchison TJ. 1994. A heterodimeric coiled-coil protein required for mitotic
 1271 chromosome condensation in vitro. *Cell* **79**:449–458. doi:10.1016/0092-8674(94)90254-2

1272 Hossain M, Bhalla K, Stillman B. 2021. Multiple, short protein binding motifs in ORC1 and
 1273 CDC6 control the initiation of DNA replication. *Mol Cell*.
 1274 doi:10.1016/j.molcel.2021.03.003

1275 Houlard M, Godwin J, Metson J, Lee J, Hirano T, Nasmyth K. 2015. Condensin confers the
 1276 longitudinal rigidity of chromosomes. *Nat Cell Biol*. doi:10.1038/ncb3167

1277 Kanaan NM, Hamel C, Grabinski T, Combs B. 2020. Liquid-liquid phase separation induces
 1278 pathogenic tau conformations in vitro. *Nat Commun* 2020 111 **11**:1–16.
 1279 doi:10.1038/s41467-020-16580-3

1280 Kara N, Hossain M, Prasanth SG, Stillman B. 2015. Orc1 binding to mitotic chromosomes
 1281 precedes spatial patterning during G₁ phase and assembly of the origin
 1282 recognition complex in human cells. *J Biol Chem*. doi:10.1074/jbc.M114.625012

1283 Kim E, Kerssemakers J, Shaltiel IA, Haering CH, Dekker C. 2020. DNA-loop extruding

condensin complexes can traverse one another. *Nat* 2020 5797799 **579**:438–442.
doi:10.1038/s41586-020-2067-5

Kroschwald S, Maharana S, Mateju D, Malinowska L, Nüske E, Poser I, Richter D, Alberti S.
2015. Promiscuous interactions and protein disaggregases determine the material state of
stress-inducible RNP granules. *Elife*. doi:10.7554/eLife.06807

Kroschwald S, Maharana S, Simon A. 2017. Hexanediol: a chemical probe to investigate the
material properties of membrane-less compartments. *Matters*.
doi:10.19185/matters.201702000010

Kyte J, Doolittle RF. 1982. A simple method for displaying the hydropathic character of a
protein. *J Mol Biol*. doi:10.1016/0022-2836(82)90515-0

Langdon EM, Qiu Y, Niaki AG, McLaughlin GA, Weidmann CA, Gerbich TM, Smith JA,
Crutchley JM, Termini CM, Weeks KM, Myong S, Gladfelter AS. 2018. mRNA structure
determines specificity of a polyQ-driven phase separation. *Science* (80-).
doi:10.1126/science.aar7432

Lee KY, Bang SW, Yoon SW, Lee SH, Yoon JB, Hwang DS. 2012. Phosphorylation of ORC2
protein dissociates origin recognition complex from chromatin and replication origins. *J
Biol Chem* **287**:11891–11898. doi:10.1074/jbc.M111.338467

Li P, Banjade S, Cheng HC, Kim S, Chen B, Guo L, Llaguno M, Hollingsworth J V, King DS,
Banani SF, Russo PS, Jiang QX, Nixon BT, Rosen MK. 2012. Phase transitions in the
assembly of multivalent signalling proteins. *Nature* **483**:336–340. doi:10.1038/nature10879

Lin Y, Currie SL, Rosen MK. 2017. Intrinsically disordered sequences enable modulation of
protein phase separation through distributed tyrosine motifs. *J Biol Chem* **292**:19110–
19120. doi:10.1074/jbc.M117.800466

Lin Y, Mori E, Kato M, Xiang S, Wu L, Kwon I, McKnight SL. 2016. Toxic PR Poly-Dipeptides
Encoded by the C9orf72 Repeat Expansion Target LC Domain Polymers. *Cell*.
doi:10.1016/j.cell.2016.10.003

Lin Y, Protter DSW, Rosen MK, Parker R. 2015. Formation and Maturation of Phase-Separated
Liquid Droplets by RNA-Binding Proteins. *Mol Cell*. doi:10.1016/j.molcel.2015.08.018

Lin YH, Forman-Kay JD, Chan HS. 2018. Theories for Sequence-Dependent Phase Behaviors of
Biomolecular Condensates. *Biochemistry*. doi:10.1021/acs.biochem.8b00058

Maharana S, Wang J, Papadopoulos DK, Richter D, Pozniakovsky A, Poser I, Bickle M, Rizk S,

1315 Guillén-Boixet J, Franzmann TM, Jahnel M, Marrone L, Chang YT, Sterneckert J,
 1316 Tomancak P, Hyman AA, Alberti S. 2018. RNA buffers the phase separation behavior of
 1317 prion-like RNA binding proteins. *Science* (80-). doi:10.1126/science.aar7366
 1318 Méndez J, Stillman B. 2000. Chromatin Association of Human Origin Recognition Complex,
 1319 Cdc6, and Minichromosome Maintenance Proteins during the Cell Cycle: Assembly of
 1320 Prereplication Complexes in Late Mitosis. *Mol Cell Biol*. doi:10.1128/mcb.20.22.8602-
 1321 8612.2000
 1322 Mészáros B, Erdős G, Dosztányi Z. 2018. IUPred2A: Context-dependent prediction of protein
 1323 disorder as a function of redox state and protein binding. *Nucleic Acids Res*.
 1324 doi:10.1093/nar/gky384
 1325 Mitrea DM, Cika JA, Guy CS, Ban D, Banerjee PR, Stanley CB, Nourse A, Deniz AA, Kriwacki
 1326 RW. 2016. Nucleophosmin integrates within the nucleolus via multi-modal interactions
 1327 with proteins displaying R-rich linear motifs and rRNA. *Elife*. doi:10.7554/elife.13571
 1328 Mitrea DM, Cika JA, Stanley CB, Nourse A, Onuchic PL, Banerjee PR, Phillips AH, Park CG,
 1329 Deniz AA, Kriwacki RW. 2018. Self-interaction of NPM1 modulates multiple mechanisms
 1330 of liquid–liquid phase separation. *Nat Commun* 2018 91 9:1–13. doi:10.1038/s41467-018-
 1331 03255-3
 1332 Molliex A, Temirov J, Lee J, Coughlin M, Kanagaraj AP, Kim HJ, Mittag T, Taylor JP. 2015.
 1333 Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and
 1334 Drives Pathological Fibrillization. *Cell*. doi:10.1016/j.cell.2015.09.015
 1335 Nott TJ, Petsalaki E, Farber P, Jervis D, Fussner E, Plochowietz A, Craggs TD, Bazett-Jones DP,
 1336 Pawson T, Forman-Kay JD, Baldwin AJ. 2015. Phase transition of a disordered nuage
 1337 protein generates environmentally responsive membraneless organelles. *Mol Cell* 57:936–
 1338 947. doi:10.1016/j.molcel.2015.01.013
 1339 Okuno Y, McNairn AJ, den Elzen N, Pines J, Gilbert DM. 2001. Stability, chromatin association
 1340 and functional activity of mammalian pre-replication complex proteins during the cell cycle.
 1341 *EMBO J* 20:4263–4277. doi:10.1093/emboj/20.15.4263
 1342 Ou HD, Phan S, Deerinck TJ, Thor A, Ellisman MH, O’Shea CC. 2017. ChromEMT:
 1343 Visualizing 3D chromatin structure and compaction in interphase and mitotic cells. *Science*
 1344 (80-). doi:10.1126/science.aag0025
 1345 Pak CW, Kosno M, Holehouse AS, Padrick SB, Mittal A, Ali R, Yunus AA, Liu DR, Pappu R

1346 V, Rosen MK. 2016. Sequence Determinants of Intracellular Phase Separation by Complex
 1347 Coacervation of a Disordered Protein. *Mol Cell* **63**:72–85.
 1348 doi:10.1016/j.molcel.2016.05.042
 1349 Paloni M, Bailly R, Ciandrini L, Barducci A. 2020. Unraveling Molecular Interactions in Liquid-
 1350 Liquid Phase-Separation of Disordered Proteins by Atomistic Simulations. *J Phys Chem B*.
 1351 doi:10.1021/acs.jpcc.0c06288
 1352 Palozola KC, Donahue G, Liu H, Grant GR, Becker JS, Cote A, Yu H, Raj A, Zaret KS. 2017.
 1353 Mitotic transcription and waves of gene reactivation during mitotic exit. *Science* (80-).
 1354 doi:10.1126/science.aal4671
 1355 Park JO, Rubin SA, Xu YF, Amador-Noguez D, Fan J, Shlomi T, Rabinowitz JD. 2016.
 1356 Metabolite concentrations, fluxes and free energies imply efficient enzyme usage. *Nat*
 1357 *Chem Biol* 2016 127 **12**:482–489. doi:10.1038/nchembio.2077
 1358 Parker MW, Bell M, Mir M, Kao JA, Darzacq X, Botchan MR, Berger JM. 2019. A new class of
 1359 disordered elements controls DNA replication through initiator self-assembly. *Elife*.
 1360 doi:10.7554/eLife.48562
 1361 Patel SS, Belmont BJ, Sante JM, Rexach MF. 2007. Natively Unfolded Nucleoporins Gate
 1362 Protein Diffusion across the Nuclear Pore Complex. *Cell*. doi:10.1016/j.cell.2007.01.044
 1363 Pepe A, Flamia R, Guerra D, Quaglino D, Bochicchio B, Pasquali Ronchetti I, Tamburro AM.
 1364 2008. Exon 26-coded polypeptide: An isolated hydrophobic domain of human tropoelastin
 1365 able to self-assemble in vitro. *Matrix Biol*. doi:10.1016/j.matbio.2008.02.006
 1366 Pepe A, Guerra D, Bochicchio B, Quaglino D, Gheduzzi D, Pasquali Ronchetti I, Tamburro AM.
 1367 2005. Dissection of human tropoelastin: Supramolecular organization of polypeptide
 1368 sequences coded by particular exons. *Matrix Biol*. doi:10.1016/j.matbio.2005.01.004
 1369 Peskett TR, Rau F, O’Driscoll J, Patani R, Lowe AR, Saibil HR. 2018. A Liquid to Solid Phase
 1370 Transition Underlying Pathological Huntingtin Exon1 Aggregation. *Mol Cell*.
 1371 doi:10.1016/j.molcel.2018.04.007
 1372 Qamar S, Wang G, Randle SJ, Ruggeri FS, Varela JA, Lin JQ, Phillips EC, Miyashita A,
 1373 Williams D, Strohl F, Meadows W, Ferry R, Dardov VJ, Tartaglia GG, Farrer LA,
 1374 Kaminski Schierle GS, Kaminski CF, Holt CE, Fraser PE, Schmitt-Ulms G, Klenerman D,
 1375 Knowles T, Vendruscolo M, St George-Hyslop P. 2018. FUS Phase Separation Is
 1376 Modulated by a Molecular Chaperone and Methylation of Arginine Cation- π Interactions.

1377 *Cell* **173**:720-734 e15. doi:10.1016/j.cell.2018.03.056

1378 Quiroz FG, Chilkoti A. 2015. Sequence heuristics to encode phase behaviour in intrinsically
 1379 disordered protein polymers. *Nat Mater.* doi:10.1038/nmat4418

1380 Reichheld SE, Muiznieks LD, Keeley FW, Sharpe S. 2017. Direct observation of structure and
 1381 dynamics during phase separation of an elastomeric protein. *Proc Natl Acad Sci U S A.*
 1382 doi:10.1073/pnas.1701877114

1383 Rice P, Longden L, Bleasby A. 2000. EMBOSS: The European Molecular Biology Open
 1384 Software Suite. *Trends Genet.* doi:10.1016/S0168-9525(00)02024-2

1385 Rog O, Köhler S, Dernburg AF. 2017. The synaptonemal complex has liquid crystalline
 1386 properties and spatially regulates meiotic recombination factors. *Elife.*
 1387 doi:10.7554/eLife.21455

1388 Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett
 1389 NM, Zamudio A V., Manteiga JC, Li CH, Guo YE, Day DS, Schuijers J, Vasile E, Malik S,
 1390 Hnisz D, Lee TI, Cisse II, Roeder RG, Sharp PA, Chakraborty AK, Young RA. 2018.
 1391 Coactivator condensation at super-enhancers links phase separation and gene control.
 1392 *Science* (80-). doi:10.1126/science.aar3958

1393 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
 1394 Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,
 1395 Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.
 1396 *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

1397 Schmidt HB, Barreau A, Rohatgi R. 2019. Phase separation-deficient TDP43 remains functional
 1398 in splicing. *Nat Commun.* doi:10.1038/s41467-019-12740-2

1399 Schmidt JM, Bleichert F. 2020. Structural mechanism for replication origin binding and
 1400 remodeling by a metazoan origin recognition complex and its co-loader Cdc6. *Nat*
 1401 *Commun.* doi:10.1038/s41467-020-18067-7

1402 Sheu YJ, Stillman B. 2010. The Dbf4-Cdc7 kinase promotes S phase by alleviating an inhibitory
 1403 activity in Mcm4. *Nature* **463**:113–117. doi:10.1038/nature08647

1404 Shulga N, Goldfarb DS. 2003. Binding Dynamics of Structural Nucleoporins Govern Nuclear
 1405 Pore Complex Permeability and May Mediate Channel Gating. *Mol Cell Biol.*
 1406 doi:10.1128/mcb.23.2.534-542.2003

1407 Sing CE, Perry SL. 2020. Recent progress in the science of complex coacervation. *Soft Matter.*

doi:10.1039/d0sm00001a

So C, Seres KB, Steyer AM, Mönnich E, Clift D, Pejkovska A, Möbius W, Schuh M. 2019. A liquid-like spindle domain promotes acentrosomal spindle assembly in mammalian oocytes. *Science* (80-). doi:10.1126/science.aat9557

Sonneville R, Querenet M, Craig A, Gartner A, Blow JJ. 2012. The dynamics of replication licensing in live *Caenorhabditis elegans* embryos. *J Cell Biol* **196**:233–246. doi:10.1083/jcb.201110080

Strom AR, Emelyanov A V, Mir M, Fyodorov D V, Darzacq X, Karpen GH. 2017. Phase separation drives heterochromatin domain formation. *Nature* **547**:241–245. doi:10.1038/nature22989

Su X, Ditlev JA, Hui E, Xing W, Banjade S, Okrut J, King DS, Taunton J, Rosen MK, Vale RD. 2016. Phase separation of signaling molecules promotes T cell receptor signal transduction. *Science* (80-) **352**:595–599. doi:10.1126/science.aad9964

Sun M, Biggs R, Hornick J, Marko JF. 2018. Condensin controls mitotic chromosome stiffness and stability without forming a structurally contiguous scaffold. *Chromosom Res.* doi:10.1007/s10577-018-9584-1

Toonkool P, Jensen SA, Maxwell AL, Weiss AS. 2001. Hydrophobic Domains of Human Tropoelastin Interact in a Context-dependent Manner. *J Biol Chem.* doi:10.1074/jbc.M107920200

Trivedi P, Palomba F, Niedzialkowska E, Digman MA, Gratton E, Stukenberg PT. 2019. The inner centromere is a biomolecular condensate scaffolded by the chromosomal passenger complex. *Nat Cell Biol.* doi:10.1038/s41556-019-0376-4

Van Treeck B, Protter DSW, Matheny T, Khong A, Link CD, Parker R. 2018. RNA self-assembly contributes to stress granule formation and defining the stress granule transcriptome. *Proc Natl Acad Sci U S A.* doi:10.1073/pnas.1800038115

Vernon RM, Chong PA, Tsang B, Kim TH, Bah A, Farber P, Lin H, Forman-Kay JD. 2018. Pi-Pi contacts are an overlooked protein feature relevant to phase separation. *Elife* **7**. doi:10.7554/eLife.31486

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ, Feng Y, Moore

1439 EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris
 1440 CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P, Vijaykumar A, Bardelli A
 1441 Pietro, Rothberg A, Hilboll A, Kloeckner A, Scopatz A, Lee A, Rokem A, Woods CN,
 1442 Fulton C, Masson C, Häggström C, Fitzgerald C, Nicholson DA, Hagen DR, Pasechnik D
 1443 V., Olivetti E, Martin E, Wieser E, Silva F, Lenders F, Wilhelm F, Young G, Price GA,
 1444 Ingold GL, Allen GE, Lee GR, Audren H, Probst I, Dietrich JP, Silterra J, Webber JT,
 1445 Slavič J, Nothman J, Buchner J, Kulick J, Schönberger JL, de Miranda Cardoso JV, Reimer
 1446 J, Harrington J, Rodríguez JLC, Nunez-Iglesias J, Kuczynski J, Tritz K, Thoma M,
 1447 Newville M, Kümmerer M, Bolingbroke M, Tartre M, Pak M, Smith NJ, Nowaczyk N,
 1448 Shebanov N, Pavlyk O, Brodtkorb PA, Lee P, McGibbon RT, Feldbauer R, Lewis S, Tygier
 1449 S, Sievert S, Vigna S, Peterson S, More S, Pudlik T, Oshima T, Pingel TJ, Robitaille TP,
 1450 Spura T, Jones TR, Cera T, Leslie T, Zito T, Krauss T, Upadhyay U, Halchenko YO,
 1451 Vázquez-Baeza Y. 2020. SciPy 1.0: fundamental algorithms for scientific computing in
 1452 Python. *Nat Methods* **17**:261–272. doi:10.1038/s41592-019-0686-2
 1453 Wang J, Choi JM, Holehouse AS, Lee HO, Zhang X, Jahnel M, Maharana S, Lemaitre R,
 1454 Pozniakovsky A, Drechsel D, Poser I, Pappu R V., Alberti S, Hyman AA. 2018. A
 1455 Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA
 1456 Binding Proteins. *Cell*. doi:10.1016/j.cell.2018.06.006
 1457 Wegmann S, Eftekharzadeh B, Tepper K, Zoltowska KM, Bennett RE, Dujardin S, Laskowski
 1458 PR, MacKenzie D, Kamath T, Commins C, Vanderburg C, Roe AD, Fan Z, Molliex AM,
 1459 Hernandez-Vega A, Muller D, Hyman AA, Mandelkow E, Taylor JP, Hyman BT. 2018.
 1460 Tau protein liquid–liquid phase separation can initiate tau aggregation. *EMBO J*.
 1461 doi:10.15252/embj.201798049
 1462 Wootton JC, Federhen S. 1993. Statistics of local complexity in amino acid sequences and
 1463 sequence databases. *Comput Chem* **17**:149–163.
 1464 Yeeles JT, Deegan TD, Janska A, Early A, Diffley JF. 2015. Regulated eukaryotic DNA
 1465 replication origin firing with purified proteins. *Nature* **519**:431–435.
 1466 doi:10.1038/nature14285
 1467 Yeo GC, Keeley FW, Weiss AS. 2011. Coacervation of tropoelastin. *Adv Colloid Interface Sci*.
 1468 doi:10.1016/j.cis.2010.10.003
 1469 Zarin T, Strome B, Nguyen Ba AN, Alberti S, Forman-Kay JD, Moses AM. 2019. Proteome-

1470 wide signatures of function in highly diverged intrinsically disordered regions. *Elife*.
1471 doi:10.7554/eLife.46883
1472 Zhang X, Lin Y, Eschmann NA, Zhou H, Rauch JN, Hernandez I, Guzman E, Kosik KS, Han S.
1473 2017. RNA stores tau reversibly in complex coacervates. *PLoS Biol* **15**:e2002183.
1474 doi:10.1371/journal.pbio.2002183
1475 ~~20211203, 9:23AM~~