**Host ecology regulates interspecies recombination in bacteria of the genus
*Campylobacter***

Evangelos Mourkas¹, Koji Yahara², Sion C. Bayliss¹, Jessica K. Calland¹, Håkan Johansson³,
Leonardos Mageiros¹, Zilia Y. Muñoz-Ramirez⁴, Grant Fitcher¹, Guillaume Méric^{1#},
Matthew D. Hitchings⁵, Santiago Sandoval-Motta⁴, Javier Torres⁴, Keith A. Jolley⁶, Martin
C. J. Maiden⁶, Patrik Ellström⁷, Jonas Waldenström³, Ben Pascoe¹, Samuel K. Sheppard^{1,6*}

¹The Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath BA2
7AY, UK; ²Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo, 162-
8640, Japan; ³Centre for Ecology and Evolution in Microbial Model Systems, Linnaeus University, Kalmar, 391
82, Sweden; ⁴Unidad de Investigacion en Enfermedades Infecciosas, UMAE Pediatria, Instituto Mexicano del
Seguro Social; ⁵Swansea University Medical School, Swansea University, Singleton Park, Swansea, SA2 8PP,
UK; ⁶Department of Zoology, University of Oxford, South Parks Road, Oxford, OX1 3PS, UK; ⁷Department of
Medical Sciences, Zoonosis Science Centre, Uppsala University, Uppsala, Sweden;

*Corresponding authors: Samuel K. Sheppard; s.k.sheppard@bath.ac.uk

[#]Present address: Cambridge Baker Systems Genomics Initiative, Baker Heart and Diabetes Institute, 75
Commercial Rd, Melbourne 3004, Victoria, Australia; Department of Infectious Diseases, Central Clinical
School, Monash University, Melbourne, Victoria 3004, Australia.

Keywords: *Campylobacter*, genus, species, niche, adaptation, host, evolution.

Running title: Comparative genomics of the *Campylobacter* genus

Abstract

Horizontal gene transfer (HGT) can allow traits that have evolved in one bacterial species to transfer to another. This has potential to rapidly promote new adaptive trajectories such as zoonotic transfer or antimicrobial resistance. However, for this to occur requires gaps to align in barriers to recombination within a given time frame. Chief among these barriers is the physical separation of species with distinct ecologies in separate niches. Within the genus *Campylobacter* there are species with divergent ecologies, from rarely isolated single host specialists to multi-host generalist species that are among the most common global causes of human bacterial gastroenteritis. Here, by characterising these contrasting ecologies, we can quantify HGT among sympatric and allopatric species in natural populations. Analysing recipient and donor population ancestry among genomes from 30 *Campylobacter* species we show that cohabitation in the same host can lead to a 6-fold increase in HGT between species. This accounts for up to 30% of all SNPs within a given species and identifies highly recombinogenic genes with functions including host adaptation and antimicrobial resistance. As described in some animal and plant species, ecological factors are a major evolutionary force for speciation in bacteria and changes to the host landscape can promote partial convergence of distinct species through HGT.

Introduction

It is well established that bacteria do not conform to a strict clonal model of reproduction but engage in regular horizontal gene transfer (HGT) [1]. This lateral exchange of DNA can confer new functionality on recipient genomes, potentially promoting novel adaptive trajectories such as colonization of a new host or the emergence of pathogenicity [2]. In some cases, gene flow can occur at such magnitude, even between different species [3, 4], that one may question why disparate lineages do not merge and why distinct bacterial species exist at all [5]. An answer to this lies in considering the successive processes that enable genes from one strain to establish in an entirely new genetic background.

The probability of HGT is governed by the interaction of multiple factors, including exposure to DNA, the susceptibility of the recipient genome to DNA uptake, and the impact of recombined DNA on the recipient strain. These factors can be broadly defined in three functional phases and HGT can only occur when gaps align in each successive ecological, mechanistic and adaptive barriers within a given time frame (Figure 1). In the first phase, the quantity of DNA available to recipient strains is determined by ecological factors such as the distribution, prevalence and interactions of donor and recipient bacteria, as well as the capacity for free DNA to be disseminated among species/strains. In the second phase, there are mechanistic barriers to HGT imposed by the homology dependence of recombination [6] or other factors promoting DNA specificity - such as restriction-modification, CRISPR interference or antiphage systems [7–11] - that can act as a defence against the uptake of foreign DNA (mechanistic barriers) [12, 13]. Finally, the effect that HGT has on the fitness of the recipient cell in a given selective environment (adaptive barrier) will determine if the recombinant genotype survives for subsequent generations [2, 14].

Understanding how ecology maintains, and potentially confines, distinct strains and species has become increasingly important in the light of global challenges such as the emergence and spread of zoonotic pathogens [15]. A typical approach to investigating this is to consider spillover of particular strains or clones from one host to another (clonal transmission). This is an important phenomenon and may be influenced by anthropogenic change, such as habitat encroachment or agricultural intensification [16]. However, in many cases, important phenotypes, including antimicrobial resistance (AMR) [17–19], can be conferred by relatively few genes. In such cases, it may be important to consider how cohabiting strains and species can potentially draw genes from a common pangenome pool [20–23] and how

genes, rather than clones, can transition between segregated populations (gene pool transmission). To investigate the impact of ecological segregation (ecological barriers) on this gene pool transmission, in natural populations, requires quantification of HGT among sympatric and allopatric bacteria.

Species within the genus *Campylobacter* are an ideal subject for considering how ecology influences the maintenance of genetically distinct species for several reasons. First, *Campylobacter* are a common component of the commensal gut microbiota of reptiles [24, 25], birds [26, 27] and mammals [28] but, being microaerophilic, do not survive well outside of the host. This creates island populations that have some degree of ecological isolation. Second, because at least 12 species have been identified as human pathogens [29] and *C. jejuni* and *C. coli* among the most common global causes of bacterial gastroenteritis [30], large numbers of isolate genomes have been sequenced from potential reservoir hosts as part of public health source tracking programs [31, 32]. Third, within the genus there are species and strains that inhabit one or multiple hosts (ecological specialists and generalists [16, 26, 33–37]). As a single host can simultaneously carry multiple lineages [38], possibly occupying different sub-niches within that host [39], there is potential to compare allopatric and sympatric populations. Finally, high magnitude interspecies admixture (introgression) between *C. jejuni* and *C. coli* isolated from agricultural animals suggests that host ecology plays a role in the maintenance of species [40–43].

Here, we quantify HGT among 600 genomes from 30 *Campylobacter* species using a ‘chromosome painting’ approach [44–46] to characterize shared ancestry among donor and recipient populations. Specifically, we investigate the role of ecological barriers to interspecies gene flow. By identifying recombining species pairs within the same and different hosts we can describe interactions where co-localization enhances gene flow, quantify the impact of ecological barriers in these populations and distinguish highly recombinogenic genes that are found in multiple genetic backgrounds. This provides information about the evolutionary forces that gave rise to species and the extent to which ecological barriers maintain them as discrete entities.

Results

Host restricted and host generalist *Campylobacter* species

Isolate genomes were taken from publicly available databases to represent diversity within the genus *Campylobacter*, including environmental isolates from the closely related *Arcobacter* and *Sulfurospirillum* species to provide phylogenetic context within the *Campylobacteraceae* family (Figure 2a–figure supplement 1). In total, there were 631 isolates from 30 different *Campylobacter* species (Figure 2a) and 64 different sources, isolated from 31 different countries between 1964 and 2016 (Supplementary File 1). Among the isolates, 361 were *C. jejuni* and *C. coli* and could be classified according to 31 Clonal Complexes (CCs) based upon sharing four or more alleles at seven housekeeping genes defined by multi-locus sequence typing (MLST) (Supplementary File 1) [47] and were representative of known diversity in both species [16, 33]. The obligate human commensal and pathogen *C. concisus* (n=106 isolates), comprised 2 genomospecies (GSI, n=32 and GSII, n=74), as previously described [48] (Supplementary File 1). The collection also included more than 52 *C. fetus* isolate genomes, including 3 subspecies: *C. fetus subsp. fetus* (n=8), *C. fetus subsp. venerealis* (n=23) and *C. fetus subsps. testudinum* (n=21) (Supplementary File 1) [49]. Two clades were observed in *C. lari* (Figure 2a–figure supplement 2) which could correspond to previously described subspecies based on 16S rRNA sequencing [50].

A maximum-likelihood phylogeny of the *Campylobacter* genus was reconstructed on a gene-by-gene concatenated sequence alignment of 820 gene families shared by >95% of all isolates, with a core genome of 903,753 base pairs (Figure 2a). The phylogeny included species which appear to be restricted to one host or environment, including *C. iguanorium* [51] and *C. geochelonis* [52] (reptiles), *C. lanienae* [53] (pigs), *C. hepaticus* [54] (chicken liver), *C. lari* group [55] (marine birds and environment) and *C. pinnipediorum* [56] (seals) species, most of which were discovered recently (Figure 2a–figure supplement 3). There was no evidence that phylogeography was reflected in the observed population structure for *Campylobacter* isolates from multiple hosts and countries. (Figure 2–figure supplement 4). This is unsurprising as it is well known that host associated genetic variation transcends phylogeographic structuring in *Campylobacter* [35]. While some low-level local gene flow can be identified within a given country [57], this is vastly outweighed by recombination within particular host niches [36], particularly in small isolate collections such as those for some of the species in this study.

Host restricted species had lower diversity possibly linked to low sample numbers, with *C. hepaticus* having the lowest diversity (Figure 2–figure supplement 2) with 8/10 genomes associated with isolates from the same outbreak [54]. For other species there was evidence of a broad host range (ecological generalists) (Figure 2b). For example, highly structured *C. jejuni* and *C. coli* isolates were sampled from seven and six host sources respectively (Figure 2–figure supplements 2-3, Supplementary File 1). For *C. fetus* there was distinct separation between mammal-associated *C. fetus* subsp *fetus* and *C. fetus* subsp *venerealis* and reptile-associated *C. fetus* subsp *testudinum* (Figure 2–figure supplement 2) as previously described [49]. Unsurprisingly, a large proportion of the isolates in this study were from humans, likely reflecting intensive sampling. *C. jejuni* (27.52%; n=60/218), *C. coli* (14.68%; n=32/218) and *C. concisus* (44.5%; n=97/218) were all common among human clinical samples. However, less common species were also present, with nearly half of all *Campylobacter* species (44.83%, n=13/29) isolated from humans at least once (Figure 2b, Supplementary File 1). Agricultural animals were also a common source accounting for more than 1/3 of the isolates (38.35%; 242/631), with 10/30 *Campylobacter* species isolated from more than one source (Figure 2b, Supplementary File 1).

Evidence of interspecies recombination in the core and accessory genome

Genome size varied between 1.40 and 2.51 Mb (Figure 3–figure supplement 1) (mean 1.73) and the number of genes (per isolate) ranged from 1,293 to 2,170 (mean 1,675) (Figure 3a–figure supplement 2). The pangenome for the genus comprised 15,649 unique genes, found in at least one of the 631 isolates (Figure 3b–source data 1), with 820 genes (5.24 % of the pangenome) shared by >95% of all isolates (core genome), across 30 species (Figure 3b–source data 1). We excluded species with fewer than 3 isolates in subsequent analysis. For the remaining 15 species the core genome ranged in size from 1,116 genes in *C. lari* to 1,700 in *C. geochelonis* (Figure 3a right panel–source data 1). Differences were also noted in the size of accessory genomes, with *C. concisus* (mean: 981 genes), *C. hyointestinalis* (mean: 946 genes), *C. showae* (mean: 1,160 genes), *C. geochelonis* (mean: 1,021 genes) and *C. fetus* (mean: 912 genes) containing the highest average number of accessory genes (Figure 3a left panel–source data 1). Functional annotation of all 14,829 accessory genes showed that 71% (10,561) encoded hypothetical proteins of unknown function due to the lack of homology with well-characterized genes (Figure 3–figure supplement 3) [58]. Remaining genes were related to metabolism, DNA modification, transporters, virulence, inner

membrane/periplasmic, adhesion, regulators, metal transport and antimicrobial resistance (Figure 3–figure supplement 3).

To further understand genetic differentiation within and between species, we generated genus-wide similarity matrices for the core and accessory genomes (Figure 3c–d–source data 1). For the core genome, pairwise average nucleotide identity (ANI) was calculated for shared genes in all possible genome pairs (Figure 3c–source data 1) using FastANI [59]. On average, isolates of the same species shared >95% similarity (Figure 3c–source data 1), with decreasing genetic similarity (between 85% and 90%) over greater phylogenetic distances. The number of core genome SNPs ranged from 983 to 230,264 for the 15 *Campylobacter* species with ≥ 3 isolates in our dataset, with *C. coli* and *C. concisus* having the greatest mean SNP numbers (Figure 3–figure supplement 4a) indicating considerable diversity within these species. In contrast *C. hepaticus* and *C. geochelonis* had low mean SNP numbers with 986 and 4,310, respectively. This is likely related to low sample numbers with isolates either sampled in close proximity [52] or from a single outbreak [54].

The core genome similarity matrix provided initial evidence of interspecies gene flow (introgression). This can be observed as elevated nucleotide identity between *C. jejuni* and clade 1 *C. coli* (Figure 3c–source data 1), consistent with previous studies [40, 42, 43]. Further evidence of introgression came from pairwise ANI comparison of genus-wide core genes, in all isolates of the 15 major *Campylobacter* species, to the *C. jejuni* genome (Figure 3–figure supplement 4b). In the absence of gene flow, isolates from the two species should have an approximately unimodal ANI distribution reflecting accumulation of mutations throughout the genome. This was largely the case but for some species, low nucleotide divergence suggested recent recombination with *C. jejuni*. There was also evidence of interspecies accessory genome recombination. Presence/absence patterns in the accessory genome matrix show considerable accessory gene sharing among several species that was inconsistent with the phylogeny (Figure 3d–source data 1). This is well illustrated in *C. lanienae* where much of the accessory genome was shared with other *Campylobacter* species (Figure 3d–source data 1).

Enhanced interspecies recombination among cohabiting species.

For *Campylobacter* inhabiting different host species there is a physical barrier to HGT. However, when there is niche overlap, interspecies recombination can occur, for example

between *C. jejuni* and *C. coli* inhabiting livestock [33, 40, 42]. To understand the extent to which inhabiting different hosts impedes interspecies gene flow we quantified recombination among *Campylobacter* species where isolates originated from same host (x_1, y) and different hosts (x_2, y) (Figure 4a).

ChromoPainterV2 software was used to infer tracts of DNA donated from multiple donor groups, belonging to the same CC but isolated from different hosts to recipient groups (Materials and Methods). Among 27 combinations of multiple donor groups and recipient groups, overall, there were more recombining SNPs within hosts than between hosts (Figure 4b) and for 10/27 species pairs there was evidence of enhanced within species recombination ($x_1 \rightarrow y > x_2 \rightarrow y$; Figure 4c). To assess the robustness of the analysis we included the effect of randomization and repeated the analysis by assigning random hosts for every strain (Figure 4-figure supplement 1). In the 10 pair species comparisons where $x_1 \rightarrow y > x_2 \rightarrow y$, we detected 174,594 within-host recombining SNPs (mapped to 473 genes; 28.8% of NCTC11168 genes) and 109,564 between-host recombining SNPs (mapped to 395 genes; 24.05% of NCTC11168 genes). From the 473 within-host recombining genes, 45 genes contained the highest number (>95th percentile) of recombining SNPs (Figure 4-figure supplements 2–3, Supplementary File 2). These genes have diverse inferred functions including metabolism, cell wall biogenesis, DNA modification, transcription, and translation (Supplementary File 2).

Interspecies recombination was observed for isolates sampled from chickens between generalist lineages CC21 and CC45 (donors; *C. jejuni*) and generalist CC828 (recipient; *C. coli*). These lineages appear to have high recombination to mutation (r/m) ratio as inferred by ClonalFrameML (Supplementary File 3). DNA from generalist *C. jejuni* CC45 was introduced into three *Campylobacter* species, including *C. hepaticus* (chicken), *C. concisus* GSI and GSII (clinical) and *C. ureolyticus* (clinical) (Figure 4c-figure supplement 2–3, Supplementary File 4). Clonal complex 45 had the highest r/m ratio from all other lineages or species involved in the comparisons (Supplementary File 3). There was increased recombination in genomes sampled from cattle between *C. jejuni* CC61 (donor; *C. jejuni*) and *C. fetus* and *C. hyointestinalis* (recipients) with 71.75% of all within-host recombining SNPs from all 10 comparisons detected in these two pairs (Figure 4c-figure supplement 2–3, Supplementary File 4). Agricultural associated *C. jejuni* CC61 and *C. fetus* subsp. *venerealis* involved in these comparisons were among the lineages and subspecies with the highest r/m ratios (Supplementary File 3). The cattle-associated CC61 has previously been described as

highly recombinant, and has been associated with rapid clonal expansion and adaptation in cattle [16].

The within-host mobilome

Bacteria inhabiting the same niche may benefit from functionality conferred by similar gene combinations. Recombination can promote the dissemination of adaptive genetic elements among different bacterial species. Therefore, we postulated that the genes that recombine most among species (>95th percentile) will include those that are potentially beneficial in multiple genetic backgrounds. To investigate this, we quantified mobility within the genome identifying recombining SNPs found in more than one species comparison (Figure 5a). These SNPs mapped to 337 genes (20.52% of the NCTC11168 genes; 2.15% of the pangenome) (Figure 5a, Supplementary File 5). We found that 32 of those genes (9.49%) have also been found on plasmids (Supplementary File 5). A total of 16 genes showed elevated within-host interspecies recombination in more than five species pairs (Figure 5c, Supplementary File 5). Genes included *cmeA* and *cmeB* which are part of the predominant efflux pump CmeABC system in *Campylobacter*. Sequence variation in the drug-binding pocket of the *cmeB* gene has been linked to increased efflux function leading to resistance to multiple drugs [60]. Many of the same antimicrobial classes are used in human and veterinary medicine and this may be linked to selection for AMR *Campylobacter*, that are commonly isolated from livestock [61]. To investigate this further, we compared the genomes of all 631 isolates in our dataset to 8,762 known antibiotic resistance genes from the Comprehensive Antibiotic Resistance Database (CARD) [62], ResFinder [63] and the National Center for Biotechnology Information (NCBI) databases. Homology (>75%) was found for 42 AMR determinants associated with multi-drug efflux pumps, aminoglycosides, tetracyclines and β -lactams (Figure 5b–figure supplement 1–source data 1). Species that contained >40% isolates from livestock, including *C. jejuni*, *C. coli*, *C. lanienae*, *C. hepaticus*, *C. hyointestinalis* and *C. fetus* contained far more AMR determinants (Figure 5d–figure supplement 1–source data 1). AMR genes are often collocated in the genome [64] and our analysis revealed several gene clusters (Figure 5–figure supplement 2) that have been described in previous studies [64, 65]. These findings are consistent with HGT-mediated circulation of AMR genes among different *Campylobacter* species and support hypotheses that ecology drives gene pool transmission [2, 64].

Campylobacter host transmission and virulence have been linked with biofilm formation and changes into surface polysaccharides [66, 67]. The *carB* gene showed elevated within-host interspecies recombination in eight species pair comparisons (Figure 5c, Supplementary File 5). This gene encodes a carbamoylphosphate synthase that has been associated with biosynthesis of substrates for many polysaccharides and is known to contain transposon insertion sites upstream of its genomic position [67]. Other genes with elevated within-host interspecies transfer (>7 species pairs) included *typA* (Figure 5c, Supplementary File 5), a translator regulator for GTPase and *gltX* (Figure 5c), a *glutamate-tRNA ligase*, promoting survival under stress conditions [68, 69]. Other genes included *gidA* and *hydB* associated with virulence [70] and hydrogenase enzyme activity (respiratory pathway in *C. concisus*, 69), respectively. By considering genes that overcome barriers to interspecies recombination and establish in multiple new genetic backgrounds, it may be possible to infer important phenotypes that allow bacteria to adapt to different hosts and environments.

Discussion

Phylogenetic reconstruction of the genus *Campylobacter* revealed a highly structured population. Distinct core genome clustering largely supported known classification for species, subspecies (*C. fetus*, [49]), genomospecies (*C. concisus*, [48]) and clades (*C. coli* [42]). Also consistent with previous studies, certain species are principally associated with a specific host niche. For example, *C. fetus subsp testudinum*, *C. iguanorium*, *C. geochelonis* were only sampled from reptile species, and *C. pinnipediorum* was only sampled from seals. However, for several species there was clear evidence for host generalism, including *C. jejuni*, *C. coli* and *C. lari*, all of which were sampled from multiple hosts [26, 72]. It is clear that the hosts with the greatest diversity of *Campylobacter* species were agricultural animals (and humans) (Figure 2–figure supplement 3). While this undoubtedly reflects oversampling of these sources to some extent, the cohabitation of species in the same host niche potentially provides opportunities for interspecies HGT.

Initial evidence of interspecies gene flow came from comparison of average nucleotide identity (ANI) and the accessory genome gene presence/absence for all isolates. In each case, patterns of genetic similarity largely mirrored the phylogeny. However, consistent with previous studies [40], there was clear evidence of elevated homologous and non-homologous recombination between some species. For example, core genome ANI was higher between *C. jejuni* and *C. coli* clade 1, compared to other *C. coli* clades (Figure 3c–source data 1). The

evidence for non-homologous gene sharing was even more striking with accessory genome sharing across considerable genetic distances (Figure 3d—source data 1), exemplified by *C. lanienae* which shares accessory genes with most other *Campylobacter* species.

To quantify the extent to which ecological barriers influenced interspecies gene flow, it was necessary to focus on donor-recipient species pairs where there was evidence of elevated HGT in the same (sympatry) compared to different (allopatry) hosts. Perhaps unsurprisingly, this was not the case for all species comparisons. Interacting factors could lead to genetic isolation even when species inhabit the same host. First, rather than being a single niche, the host represents a collection of subniches with varying degrees of differentiation. For example, gut-associated bacteria in the same intestinal tract have been shown to occupy different microniches [73] and more striking segregation may be expected between *C. hepaticus* inhabits the liver in poultry [54] and gut-dwelling *C. jejuni* and *C. coli* in the same host. Second, there is potential for the resident microbiota to influence the colonization potential of different *Campylobacter* species and therefore the opportunity for genetic exchange, for example through succession [74] and inhibition of transient species by residents, as seen in some other bacteria [75–77] in humans.

Continued exposition of the microecology of subniches is important but for 10 species comparisons there was clear evidence of enhanced within-host gene flow allowing quantitative analysis of ecological barriers to gene flow. Specifically, there was on average a 3-fold increase in recombination among species pairs inhabiting the same host. In some cases, this was greater, with 5-6 times more recombination among cohabiting species *C. jejuni* and *C. hointestinalis/C. fetus* in cattle. In absolute terms, this equates to approximately 30% of all recorded SNPs in the recipient species being the result of introgression. To place this in context, if greater than half (51%) of the recorded SNPs resulted from interspecies recombination then the forces of species convergence would be greater than those that maintain distinct species. If maintained over time, these relative rates could lead to progressive genetic convergence unless countered by strong genome-wide natural selection against introgressed DNA.

Quantitative SNP-based comparisons clearly ignore one very important factor. Specifically, that recombined genes that do not reduce the fitness of the recipient genome (provide an adaptive advantage) will remain in the population while others will be purged through natural

selection. Therefore, by identifying genomic hotspots of recombination and the putative function of genes that recombine between species it is possible to understand more about micro-niche segregation and the host adapted gene pool. Of the 35 genes with evidence of enhanced within host HGT in ≥ 5 species pairs, several were linked to functions associated with proliferation in, and exploitation of, the host. For example, the *carB* gene, encoding the large subunit of carbamoylphosphatase associated with polysaccharide biosynthesis, recombined in eight cohabiting species pairs and is potentially linked to enhanced virulence and growth [67]. In addition, other highly mobile genes, including *typA* and *gltX* are associated with survival and proliferation in stress conditions [68, 69], and *hydB* is linked to NiFe hydrogenase and nickel uptake that is essential for the survival of *C. jejuni* in the gut of birds and mammals [78].

Some genes showed evidence of elevated recombination in a specific host species. For example, the *glmS* and *napA* genes in cohabiting *Campylobacter* species in cattle. In many bacteria, analogs of *glmS* have multiple downstream integration specific sites (Tn7) [79] which may explain the mobility of this gene. Explaining the mobility of *napA* is less straightforward, but this gene is known to encode a nitrate reductase in *Campylobacter* [80] in microaerobic conditions which may be ecologically significant as the accumulation of nitrate in slurry, straw and drainage water can be potentially toxic to livestock mammals [81].

Factors such as host physiology, diet, and metabolism undoubtedly impose selection pressures upon resident bacteria and the horizontal acquisition of genes provides a possible vehicle for adaptation. However, the widespread use of antimicrobials by humans, pets and livestock production [82, 83], provides another major ecological barrier to niche colonization. We found that *gyrA* was among the most recombinogenic genes in *Campylobacter* in chickens. This is important as a single mutation in this gene is known to confer resistance to ciprofloxacin [84]. While the rising trend in fluoroquinolones resistance in *Campylobacter* from humans and livestock [85] may result from spontaneous independent mutations, it is likely that it is accelerated by HGT. However, there is currently no clear evidence for the transfer of resistant versions of *gyrA*. Interspecies recombination of AMR genes has been observed between *C. jejuni* and *C. coli* isolates from multiple sources including livestock, human and sewage [64]. Consistent with this, we found AMR genes present in strains from 12 *Campylobacter* species in multiple hosts (Figure 5–figure supplement 2). In some cases, strains from phylogenetically closely related species (*C. fetus* and *C. hyointestinalis*) isolated

from cattle, shared the same AMR gene cluster (*tet44* and *ant(6)-Ib*) described before in *C. fetus* subsp. *fetus* [65], indicating the circulation of colocalized AMR genes among related species and host niche gene pools. Strikingly, the efflux pump genes *cmeA* and *cmeB*, associated with multidrug resistance (MDR) were highly mobile among *Campylobacter* species with evidence of elevated within host interspecies recombination in >7 species pairs. Furthermore, the *gltX* gene, which when phosphorylated by protein kinases promotes MDR [69], was also among the most introgressed genes. While a deeper understanding of gene interactions, epistasis and epigenetics would be needed to prove that the lateral acquisition of AMR genes promotes niche adaptation, these data do suggest that HGT may facilitate colonization of antimicrobial-rich host environments, potentially favouring the spread of genes into multiple genetic backgrounds.

In conclusion, we show that species within the genus *Campylobacter* include those that are host restricted as well as host generalists. When species cohabit in the same host, ecological barriers to recombination can be perforated leading to considerable introgression between species. While the magnitude of introgression varies, potentially reflecting microniche structure with the host, there is clear evidence that ecology is important in maintaining genetically distinct species. This parallels evolution in some interbreeding eukaryotes, such as Darwin's Finches, where fluctuating environmental conditions can change the selection pressures acting on species inhabiting distinct niches, potentially favouring hybrids [86, 87]. Consistent with this, the host landscape is changing for *Campylobacter*, with intensively reared livestock now constituting 60-70% of bird and mammal biomass on earth respectively [88]. This creates opportunities for species to be brought together in new adaptive landscapes and for genes to be tested from multiple genetic backgrounds. By understanding the ecology of niche segregation and the genetics of bacterial adaptation we can hope to improve strategies and interventions to reduce the risk of zoonotic transmission and the spread of problematic genes among species.

Materials & Methods

Isolate genomes

A total of 631 *Campylobacter*, 17 *Arcobacter*, seven *Sulfurospirillum* and five *Helicobacter* genomes were assembled from previously published datasets (Supplementary File 1). Isolates were sampled from clinical cases of campylobacteriosis and faeces of chickens, ruminants, wild birds, wild mammals, pets and environmental sources. Genomes and related metadata

were uploaded and archived in the BIGS database [89]. Quality control was performed based on the genome size, number of contigs, N50 and N95 contig length using the integrated tools in BIGS database. All assembled contigs were further screened for contamination and completeness using CheckM [90] (Supplementary File 1). All assembled genomes can be downloaded from FigShare (doi: 10.6084/m9.figshare.15061017). Comparative genomics analyses focused on the *Campylobacter* genomes representing 30 species including: *C. avium* (n=1); *C. coli* (n=143); *C. concisus* (n=106); *C. corcagiensis* (n=1); *C. cuniculorum* (n=2); *C. curvus* (n=2); *C. fetus* (n=52); *C. geochelonis* (n=3); *C. gracilis* (n=2); *C. helveticus* (n=1); *C. hepaticus* (n=10); *C. hominis* (n=1); *C. hyointestinalis* (n=16); *C. iguanorium* (n=3); *C. insulaenigrae* (n=1); *C. jejuni* (n=218); *C. lanienae* (n=26); *C. lari* (n=13); *C. mucosalis* (n=1); *C. ornithocola* (n=1); *C. peloridis* (n=1); *C. pinnipediorum* (n=9); *C. rectus* (n=1); *C. showae* (n=3); *C. sputorum* (n=1); *C. subantarcticus* (n=3); *C. upsaliensis* (n=3); *C. ureolyticus* (n=4); *C. volucris* (n=2); *Campylobacter sp* (n=1) (Supplementary File 1). Genomes belonging to *C. jejuni* and *C. coli* species were selected to represent a wide range of hosts, sequence types, and clonal complexes and reflect the known population structure for these two species. For other *Campylobacter* species, all genomes that were publicly available at the time of this study were included in the analysis. (Supplementary File 1).

Pangenome characterization and phylogenetic analysis

Sequence data were analysed using PIRATE, a fast and scalable pangenomics tool which allows for orthologue gene clustering in divergent bacterial species [91]. Genomes were annotated in Prokka [92], using a genus database comprising well annotated *C. jejuni* strains NCTC11168, 81116, 81-176 and M1, and plasmids pTet and pVir in addition to the already existing databases used by Prokka [92]. Briefly, annotated genomes were used as input for PIRATE. Non-redundant representative sequences were produced using CD-HIT and the longest sequence was used as a reference for sequence similarity interrogation using BLAST/DIAMOND. Gene orthologues were defined as “gene families” and were clustered in different MCL thresholds, from 10 to 98 % sequence identity (10, 20, 30, 40, 50, 60, 70, 80, 90, 95, 98). Higher MCL thresholds were used to identify allelic variation within different loci. An inflation value of 4 was used to increase the granularity of MCL clustering between gene families. BLAST high-scoring pairs with a reciprocal minimum length of 90% of the query/subject sequence were excluded from MCL clustering to reduce the number of spurious associations between distantly related or conserved genes [93]. This information was used to generate gene presence/absence and allelic variation matrices. A core gene-by-gene multiple

sequence alignment [89] was generated using MAFFT [94] comprising genes shared >95% of isolates. Phylogenetic trees, based on core gene-by-gene alignments, were reconstructed using the maximum-likelihood algorithm implemented in RAxML v8.2.11 [95] with GTRGAMMA as substitution model.

Quantifying core and accessory genome variation

The degree of genetic differentiation between species was investigated gene-by-gene as in previous studies [40, 96] by calculating the average nucleotide identity (ANI) of all 631 *Campylobacter* genomes using FastANI v.1.0 [59]. The analysis generated a lower triangular matrix with the lowest ANI value at 75% (as computed by FastANI). A comparable gene presence/absence matrix was produced using PIRATE and was further used to generate a heatmap of accessory genome similarity based upon gene presence or absence. Subsequently, all *Campylobacter* genomes were screened for the presence of antimicrobial resistance genes against the CARD [62], ResFinder [63] and NCBI databases. All *Campylobacter* genomes were further screened for the presence of phage, conjugative elements and plasmid DNA using publicly available online databases to investigate the effect of other transfer mechanisms. First, we used the PHAge Search Tool Enhanced Release (PHASTER) [97] to identify and annotate prophage sequences within our genomes. A total of 86% (254/297) of the genomes used in chromosome painting were found to have DNA sequence of phage origin. Second, we used Iceberg 2.0 [98] for the detection of integrative and conjugative elements, identifying 32 ICEs in 19% (56/297) of the genomes used in the chromosome painting analysis. Finally, we used MOB-suite software for clustering, reconstruction and typing of plasmids from draft assemblies [99, 100]. A positive hit was defined when a gene had >75% nucleotide identity over >50% of the sequence length showing that 32 genes identified in the recombination analysis have also been located on plasmids. A gene presence/absence matrix for every antimicrobial resistance gene was generated for every genome. Genomes carrying AMR genes were screened to characterize the location of adjacent genes using SnapGene software (GSL Biotech; available at snapgene.com), as previously described [64]. The number of core SNPs was identified using SNP-sites (v2.3.2) [101].

Inference of recombination

Each combination of a recipient group and multiple donor groups (belonging to the same CC but isolated from different hosts) was selected to compare the extent of interspecies

recombination into the recipient genomes. Each donor group consisted of 8 isolates to avoid the influence of difference in sample size on estimation of the extent of interspecies recombination. Each recipient group included at least 4 isolates. We excluded *C. jejuni* and *C. coli* clade 1 genomes isolated from seals and water, as these most likely represent spillover events and not true host segregated populations. Briefly, we conducted a pairwise genome alignment between reference genome NCTC11168 and one of the strains included in the donor-recipient analysis using progressiveMauve [102]. This enabled the construction of positional homology alignments for all genomes regardless gene content and genome rearrangements, which were then combined into a multiple whole-genome alignment, as previously described [103]. ChromoPainterV2 software was used to calculate the amount of DNA sequence that is donated from a donor to a recipient group [45]. Briefly, for each donor-recipient pair, SNPs in which >90% recipient individuals had recombined with the donor group were considered in the analysis. These SNPs were mapped to genomic regions and specific genes were identified. A total of 258,444 (96.83%) recombining SNPs mapped to 558 genes of the NCTC11168 reference strain with >90% probability of copying from a donor to a recipient strain. Genes containing the highest number of recombining SNPs were considered for subsequent analyses (>95th percentile) (Supplementary File 2). ClonalFrameML[104] was used to infer the relative number of substitutions introduced by recombination (r) and mutation (m) as the ratio r/m as previously described [16].

Data availability

Genomes sequenced as part of other studies are archived on the Short Read Archive associated with BioProject accessions: PRJNA176480, PRJNA177352, PRJNA342755, PRJNA345429, PRJNA312235, PRJNA415188, PRJNA524300, PRJNA528879, PRJNA529798, PRJNA575343, PRJNA524315 and PRJNA689604. Additional genomes were also downloaded from NCBI [105] and pubMLST (<http://pubmlst.org/campylobacter>). Contiguous assemblies of all genome sequences compared are available at the public data repository Figshare (doi: 10.6084/m9.figshare.15061017) and individual project and accession numbers can be found in Supplementary File 1.

References

1. Smith JM, Dowson CG, Spratt BG. Localized sex in bacteria. *Nature* 1991; **349**: 29–31.
2. Sheppard SK, Guttman DS, Fitzgerald JR. Population genomics of bacterial host

adaptation. *Nat Rev Genet* 2018; **19**: 549–565.

3. Shapiro BJ, Leducq J-B, Mallet J. What Is Speciation? *PLOS Genet* 2016; **12**: e1005860.
4. Doolittle WF, Zhaxybayeva O. On the origin of prokaryotic species. *Genome Res* 2009; **19**: 744–756.
5. Doolittle WF, Papke RT. Genomics and the bacterial species problem. *Genome Biol* 2006; **7**: 116.
6. Fraser C, Hanage WP, Spratt BG. Recombination and the nature of bacterial speciation. *Science* 2007; **315**: 476–80.
7. Budroni S, Siena E, Hotopp JCD, Seib KL, Serruto D, Nofroni C, et al. *Neisseria meningitidis* is structured in clades associated with restriction modification systems that modulate homologous recombination. *Proc Natl Acad Sci* 2011; **108**: 4494–4499.
8. Oliveira PH, Touchon M, Rocha EPC. Regulation of genetic flux between bacteria by restriction–modification systems. *Proc Natl Acad Sci* 2016; **113**: 5658–5663.
9. Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, et al. Systematic discovery of antiphage defense systems in the microbial pangenome. *Science* (80-) 2018; **359**: eaar4120.
10. Nandi T, Holden MTG, Didelot X, Mehershahi K, Boddey JA, Beacham I, et al. *Burkholderia pseudomallei* sequencing identifies genomic clades with distinct recombination, accessory, and epigenetic profiles. *Genome Res* 2015; **25**: 129–141.
11. Marraffini LA, Sontheimer EJ. CRISPR Interference Limits Horizontal Gene Transfer in Staphylococci by Targeting DNA. *Science* (80-) 2008; **322**: 1843–1845.
12. Thomas CM, Nielsen KM. Mechanisms of, and Barriers to, Horizontal Gene Transfer between Bacteria. *Nat Rev Microbiol* 2005; **3**: 711–721.
13. Eggleston AK, Mitchell AH, West SC. In Vitro Reconstitution of the Late Steps of Genetic Recombination in *E. coli*. *Cell* 1997; **89**: 607–617.
14. Zhu P, van der Ende A, Falush D, Brieske N, Morelli G, Linz B, et al. Fit genotypes and escape variants of subgroup III *Neisseria meningitidis* during three pandemics of epidemic meningitis. *Proc Natl Acad Sci* 2001; **98**: 5234–5239.
15. Boni MF, Lemey P, Jiang X, Lam TT-Y, Perry BW, Castoe TA, et al. Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. *Nat Microbiol* 2020; **5**: 1408–1417.
16. Mourkas E, Taylor AJ, Méric G, Bayliss SC, Pascoe B, Mageiros L, et al. Agricultural intensification and the evolution of host specialism in the enteric pathogen

- 553 *Campylobacter jejuni*. *Proc Natl Acad Sci* 2020; **117**: 11018–11028.
- 554 17. Johnson AP, Woodford N. Global spread of antibiotic resistance: the example of New
555 Delhi metallo- β -lactamase (NDM)-mediated carbapenem resistance. *J Med Microbiol*
556 2013; **62**: 499–513.
- 557 18. Schwarz S, Johnson AP. Transferable resistance to colistin: a new but old threat. *J*
558 *Antimicrob Chemother* 2016; **71**: 2066–70.
- 559 19. Baker KS, Dallman TJ, Field N, Childs T, Mitchell H, Day M, et al. Horizontal
560 antimicrobial resistance transfer drives epidemics of multiple *Shigella* species. *Nat*
561 *Commun* 2018; **9**: 1462.
- 562 20. Young JPW. Bacteria Are Smartphones and Mobile Genes Are Apps. *Trends*
563 *Microbiol* 2016; **24**: 931–932.
- 564 21. McInerney JO, Whelan FJ, Domingo-Sananes MR, McNally A, O’Connell MJ.
565 Pangenomes and Selection: The Public Goods Hypothesis. *The Pangenome*. 2020.
566 Springer International Publishing, Cham, pp 151–167.
- 567 22. Vos M, Eyre-Walker A. Are pangenomes adaptive or not? *Nat Microbiol* 2017; **2**:
568 1576.
- 569 23. Werren JH. Selfish genetic elements, genetic conflict, and evolutionary innovation.
570 *Proc Natl Acad Sci* 2011; **108**: 10863–10870.
- 571 24. Giacomelli M, Piccirillo A. Pet reptiles as potential reservoir of *Campylobacter*
572 species with zoonotic potential: TABLE 1: *Vet Rec* 2014; **174**: 479.2-479.
- 573 25. Fitzgerald C, Tu Z chao, Patrick M, Stiles T, Lawson AJ, Santovenia M, et al.
574 *Campylobacter fetus* subsp. *testudinum* subsp. nov., isolated from humans and reptiles.
575 *Int J Syst Evol Microbiol* 2014; **64**: 2944–2948.
- 576 26. Griekspoor P, Colles FM, McCarthy ND, Hansbro PM, Ashhurst-Smith C, Olsen B, et
577 al. Marked host specificity and lack of phylogeographic population structure of
578 *Campylobacter jejuni* in wild birds. *Mol Ecol* 2013; **22**: 1463–1472.
- 579 27. Atterby C, Mourkas E, Méric G, Pascoe B, Wang H, Waldenström J, et al. The
580 Potential of Isolation Source to Predict Colonization in Avian Hosts: A Case Study in
581 *Campylobacter jejuni* Strains From Three Bird Species. *Front Microbiol* 2018; **9**.
- 582 28. Leatherbarrow AJH, Griffiths R, Hart CA, Kemp R, Williams NJ, Diggle PJ, et al.
583 *Campylobacter lari*: genotype and antibiotic resistance of isolates from cattle, wildlife
584 and water in an area of mixed dairy farmland in the United Kingdom. *Environ*
585 *Microbiol* 2007; **9**: 1772–1779.
- 586 29. Man SM. The clinical importance of emerging *Campylobacter* species. *Nat Rev*

- 587 *Gastroenterol Hepatol* 2011; **8**: 669–685.
- 588 30. Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global Epidemiology of
589 *Campylobacter* Infection. *Clin Microbiol Rev* 2015; **28**: 687–720.
- 590 31. Sheppard SK, Dallas JF, MacRae M, McCarthy ND, Sproston EL, Gormley FJ, et al.
591 *Campylobacter* genotypes from food animals, environmental sources and clinical
592 disease in Scotland 2005/6. *Int J Food Microbiol* 2009; **134**: 96–103.
- 593 32. Sheppard SK, Dallas JF, Strachan NJC, MacRae M, McCarthy ND, Wilson DJ, et al.
594 *Campylobacter* genotyping to determine the source of human infection. *Clin Infect Dis*
595 2009; **48**: 1072–8.
- 596 33. Sheppard SK, Colles FM, McCarthy ND, Strachan NJC, Ogden ID, Forbes KJ, et al.
597 Niche segregation and genetic structure of *Campylobacter jejuni* populations from
598 wild and agricultural host species. *Mol Ecol* 2011; **20**: 3484–90.
- 599 34. Dearlove BL, Cody AJ, Pascoe B, Méric G, Wilson DJ, Sheppard SK. Rapid host
600 switching in generalist *Campylobacter* strains erodes the signal for tracing human
601 infections. *ISME J* 2016; **10**: 721–729.
- 602 35. Sheppard SK, Colles F, Richardson J, Cody AJ, Elson R, Lawson A, et al. Host
603 Association of *Campylobacter* Genotypes Transcends Geographic Variation. *Appl*
604 *Environ Microbiol* 2010; **76**: 5269–5277.
- 605 36. Sheppard SK, Cheng L, Méric G, de Haan CPA, Llarena A-K, Marttinen P, et al.
606 Cryptic ecology among host generalist *Campylobacter jejuni* in domestic animals. *Mol*
607 *Ecol* 2014; **23**: 2442–2451.
- 608 37. Woodcock DJ, Krusche P, Strachan NJC, Forbes KJ, Cohan FM, Méric G, et al.
609 Genomic plasticity and rapid host switching can promote the evolution of generalism:
610 a case study in the zoonotic pathogen *Campylobacter*. *Sci Rep* 2017; **7**: 9650.
- 611 38. Colles FM, Dingle KE, Cody AJ, Maiden MCJ. Comparison of *Campylobacter*
612 Populations in Wild Geese with Those in Starlings and Free-Range Poultry on the
613 Same Farm. *Appl Environ Microbiol* 2008; **74**: 3583–3590.
- 614 39. Colles FM, McCarthy ND, Bliss CM, Layton R, Maiden MCJ. The long-term
615 dynamics of *Campylobacter* colonizing a free-range broiler breeder flock: an
616 observational study. *Environ Microbiol* 2015; **17**: 938–946.
- 617 40. Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Méric G, et al. Progressive
618 genome-wide introgression in agricultural *Campylobacter coli*. *Mol Ecol* 2013; **22**:
619 1051–64.
- 620 41. Taylor AJ, Méric G, Yahara K, Pascoe B, Mageiros L, Mourkas E, et al. Cross-species

- evolutionary rewiring in the enteric bacterium *Campylobacter*; *bioRxiv* 2021; 2021.03.15.435406.
42. Sheppard SK, McCarthy ND, Falush D, Maiden MCJ. Convergence of *Campylobacter* Species: Implications for Bacterial Evolution. *Science* (80-) 2008; **320**: 237–239.
 43. Sheppard SK, McCarthy ND, Jolley KA, Maiden MCJ. Introgression in the genus *Campylobacter*: generation and spread of mosaic alleles. *Microbiology* 2011; **157**: 1066–1074.
 44. Thorell K, Yahara K, Berthenet E, Lawson DJ, Mikhail J, Kato I, et al. Rapid evolution of distinct *Helicobacter pylori* subpopulations in the Americas. *PLOS Genet* 2017; **13**: e1006730.
 45. Lawson DJ, Hellenthal G, Myers S, Falush D. Inference of Population Structure using Dense Haplotype Data. *PLoS Genet* 2012; **8**: e1002453.
 46. Yahara K, Furuta Y, Oshima K, Yoshida M, Azuma T, Hattori M, et al. Chromosome Painting In Silico in a Bacterial Species Reveals Fine Population Structure. *Mol Biol Evol* 2013; **30**: 1454–1464.
 47. Dingle KE, Colles FM, Wareing DR, Ure R, Fox AJ, Bolton FE, et al. Multilocus sequence typing system for *Campylobacter jejuni*. *J Clin Microbiol* 2001; **39**: 14–23.
 48. Kirk KF, Méric G, Nielsen HL, Pascoe B, Sheppard SK, Thorlacius-Ussing O, et al. Molecular epidemiology and comparative genomics of *Campylobacter concisus* strains from saliva, faeces and gut mucosal biopsies in inflammatory bowel disease. *Sci Rep* 2018; **8**: 1902.
 49. Iraola G, Forster SC, Kumar N, Lehours P, Bekal S, García-Peña FJ, et al. Distinct *Campylobacter fetus* lineages adapted as livestock pathogens and human pathobionts in the intestinal microbiota. *Nat Commun* 2017; **8**: 1367.
 50. Debruyne L, On SLW, De Brandt E, Vandamme P. Novel *Campylobacter lari*-like bacteria from humans and molluscs: description of *Campylobacter peloridis* sp. nov., *Campylobacter lari* subsp. *concheus* subsp. nov. and *Campylobacter lari* subsp. *lari* subsp. nov. *Int J Syst Evol Microbiol* 2009; **59**: 1126–32.
 51. Gilbert MJ, Kik M, Miller WG, Duim B, Wagenaar JA. *Campylobacter iguaniorum* sp. nov., isolated from reptiles. *Int J Syst Evol Microbiol* 2015; **65**: 975–982.
 52. Piccirillo A, Niero G, Calleros L, Pérez R, Naya H, Iraola G. *Campylobacter geochelonis* sp. nov. isolated from the western Hermann's tortoise (*Testudo hermanni hermanni*). *Int J Syst Evol Microbiol* 2016; **66**: 3468–3476.
 53. Logan MJ, Burnens A, Linton D, Lawson AJ, Stanley J. *Campylobacter lanienae* sp.

- nov., a new species isolated from workers in an abattoir. *Int J Syst Evol Microbiol* 2000; **50**: 865–872.
54. Van TTH, Elshagmani E, Gor MC, Scott PC, Moore RJ. *Campylobacter hepaticus* sp. nov., isolated from chickens with spotty liver disease. *Int J Syst Evol Microbiol* 2016; **66**: 4518–4524.
55. Miller WG, Yee E, Chapman MH, Smith TPL, Bono JL, Huynh S, et al. Comparative genomics of the *Campylobacter lari* group. *Genome Biol Evol* 2014; **6**: 3252–66.
56. Gilbert MJ, Miller WG, Leger JS, Chapman MH, Timmerman AJ, Duim B, et al. *Campylobacter pinnipediorum* sp. nov., isolated from pinnipeds, comprising *Campylobacter pinnipediorum* subsp. *pinnipediorum* subsp. nov. and *Campylobacter pinnipediorum* subsp. *caledonicus* subsp. nov. *Int J Syst Evol Microbiol* 2017; **67**: 1961–1968.
57. Pascoe B, Méric G, Yahara K, Wimalaratna H, Murray S, Hitchings MD, et al. Local genes for local bacteria: Evidence of allopatry in the genomes of transatlantic *Campylobacter* populations. *Mol Ecol* 2017; **26**: 4497–4508.
58. Pascoe B, Williams LK, Calland JK, Meric G, Hitchings MD, Dyer M, et al. Domestication of *Campylobacter jejuni* NCTC 11168. *Microb genomics* 2019; **5**.
59. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018; **9**: 5114.
60. Yao H, Shen Z, Wang Y, Deng F, Liu D, Naren G, et al. Emergence of a Potent Multidrug Efflux Pump Variant That Enhances *Campylobacter* Resistance to Multiple Antibiotics. *MBio* 2016; **7**: 1–11.
61. Livermore DM. Introduction: the challenge of multiresistance. *Int J Antimicrob Agents* 2007; **29**: S1–S7.
62. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res* 2017; **45**: D566–D573.
63. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 2012; **67**: 2640–2644.
64. Mourkas E, Florez-Cuadrado D, Pascoe B, Calland JK, Bayliss SC, Mageiros L, et al. Gene pool transmission of multidrug resistance among *Campylobacter* from livestock, sewage and human disease. *Environ Microbiol* 2019; **21**: 4597–4613.

- 689 65. Abril C, Brodard I, Perreten V. Two Novel Antibiotic Resistance Genes, tet(44) and
690 ant(6)-Ib, Are Located within a Transferable Pathogenicity Island in *Campylobacter*
691 *fetus* subsp. *fetus*. *Antimicrob Agents Chemother* 2010; **54**: 3052–3055.
- 692 66. Szymanski CM, Logan SM, Linton D, Wren BW. *Campylobacter* – a tale of two
693 protein glycosylation systems. *Trends Microbiol* 2003; **11**: 233–238.
- 694 67. McLennan MK, Ringoir DD, Frirdich E, Svensson SL, Wells DH, Jarrell H, et al.
695 *Campylobacter jejuni* Biofilms Up-Regulated in the Absence of the Stringent
696 Response Utilize a Calcofluor White-Reactive Polysaccharide. *J Bacteriol* 2008; **190**:
697 1097–1107.
- 698 68. Margus T, Remm M, Tenson T. Phylogenetic distribution of translational GTPases in
699 bacteria. *BMC Genomics* 2007; **8**: 1–18.
- 700 69. Semanjski M, Germain E, Bratl K, Kiessling A, Gerdes K, Macek B. The kinases
701 HipA and HipA7 phosphorylate different substrate pools in *Escherichia coli* to
702 promote multidrug tolerance. *Sci Signal* 2018; **11**.
- 703 70. Mikheil DM, Shippy DC, Eakley NM, Okwumabua OE, Fadl AA. Deletion of gene
704 encoding methyltransferase (*gidB*) confers high-level antimicrobial resistance in
705 *Salmonella*. *J Antibiot (Tokyo)* 2012; **65**: 185–192.
- 706 71. Benoit SL, Maier RJ. Site-directed mutagenesis of *Campylobacter concisus* respiratory
707 genes provides insight into the pathogen’s growth requirements. *Sci Rep* 2018; **8**:
708 14203.
- 709 72. Cody AJ, McCarthy ND, Bray JE, Wimalarathna HML, Colles FM, Jansen van
710 Rensburg MJ, et al. Wild bird-associated *Campylobacter jejuni* isolates are a
711 consistent source of human disease, in Oxfordshire, United Kingdom. *Environ*
712 *Microbiol Rep* 2015; **7**: 782–788.
- 713 73. Hayashi H, Takahashi R, Nishi T, Sakamoto M, Benno Y. Molecular analysis of
714 jejunal, ileal, caecal and recto-sigmoidal human colonic microbiota using 16S rRNA
715 gene libraries and terminal restriction fragment length polymorphism. *J Med Microbiol*
716 2005; **54**: 1093–1101.
- 717 74. Lu J, Idris U, Harmon B, Hofacre C, Maurer JJ, Lee MD. Diversity and succession of
718 the intestinal bacterial community of the maturing broiler chicken. *Appl Environ*
719 *Microbiol* 2003; **69**: 6816–24.
- 720 75. Stecher B, Chaffron S, Käppeli R, Hapfelmeier S, Friedrich S, Weber TC, et al. Like
721 will to like: abundances of closely related species can predict susceptibility to
722 intestinal colonization by pathogenic and commensal bacteria. *PLoS Pathog* 2010; **6**:

- 723 e1000711.
- 724 76. van Elsas JD, Chiurazzi M, Mallon CA, Elhottova D, Kristufek V, Salles JF. Microbial
725 diversity determines the invasion of soil by a bacterial pathogen. *Proc Natl Acad Sci U*
726 *S A* 2012; **109**: 1159–64.
- 727 77. Nowrouzian FL, Wold AE, Adlerberth I. *Escherichia coli* strains belonging to
728 phylogenetic group B2 have superior capacity to persist in the intestinal microflora of
729 infants. *J Infect Dis* 2005; **191**: 1078–83.
- 730 78. Howlett RM, Hughes BM, Hitchcock A, Kelly DJ. Hydrogenase activity in the
731 foodborne pathogen *Campylobacter jejuni* depends upon a novel ABC-type nickel
732 transporter (NikZYXWV) and is SlyD-independent. *Microbiology* 2012; **158**: 1645–
733 1655.
- 734 79. Choi K-H. Applications of Transposon-Based Gene Delivery System in Bacteria. *J*
735 *Microbiol Biotechnol* 2009; **19**: 217–28.
- 736 80. Pittman MS, Elvers KT, Lee L, Jones MA, Poole RK, Park SF, et al. Growth of
737 *Campylobacter jejuni* on nitrate and nitrite: electron transport to NapA and NrfA via
738 NrfH and distinct roles for NrfA and the globin Cgb in protection against nitrosative
739 stress. *Mol Microbiol* 2007; **63**: 575–590.
- 740 81. Alexander J, Benford D, Cockburn A, Cravedi J, Dogliotti E, Domenico A Di, et al.
741 Nitrite as undesirable substances in animal feed - Scientific Opinion of the Panel on
742 Contaminants in the Food Chain. *EFSA J* 2009; **7**: 1–47.
- 743 82. Teuber M. Veterinary use and antibiotic resistance. *Curr Opin Microbiol* 2001; **4**:
744 493–499.
- 745 83. Price LB, Koch BJ, Hungate BA. Ominous projections for global antibiotic use in
746 food-animal production. *Proc Natl Acad Sci U S A* 2015; **112**: 5554–5.
- 747 84. Luo N, Sahin O, Lin J, Michel LO, Zhang Q. In Vivo Selection of *Campylobacter*
748 Isolates with High Levels of Fluoroquinolone Resistance Associated with *gyrA*
749 Mutations and the Function of the CmeABC Efflux Pump. *Antimicrob Agents*
750 *Chemother* 2003; **47**: 390–394.
- 751 85. Sproston EL, Wimalarathna HML, Sheppard SK. Trends in fluoroquinolone resistance
752 in *Campylobacter*. *Microb Genomics* 2018; **4**: 1–8.
- 753 86. Mallet J. Hybrid speciation. *Nature* 2007; **446**: 279–283.
- 754 87. Grant PR, Grant BR. Hybridization of Bird Species. *Science (80-)* 1992; **256**: 193–
755 197.
- 756 88. Bar-On YM, Phillips R, Milo R. The biomass distribution on Earth. *Proc Natl Acad*

757 *Sci* 2018; **115**: 6506–6511.

758 89. Sheppard SK, Jolley KA, Maiden MCJ. A Gene-By-Gene Approach to Bacterial
759 Population Genomics: Whole Genome MLST of *Campylobacter*. *Genes (Basel)* 2012;
760 **3**: 261–277.

761 90. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing
762 the quality of microbial genomes recovered from isolates, single cells, and
763 metagenomes. *Genome Res* 2015; **25**: 1043–1055.

764 91. Bayliss SC, Thorpe HA, Coyle NM, Sheppard SK, Feil EJ. PIRATE: A fast and
765 scalable pangenomics toolbox for clustering diverged orthologues in bacteria.
766 *Gigascience* 2019; **8**: 1–9.

767 92. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014; **30**:
768 2068–2069.

769 93. Sahl JW, Caporaso JG, Rasko DA, Keim P. The large-scale blast score ratio (LS-BSR)
770 pipeline: a method to rapidly compare genetic content between bacterial genomes.
771 *PeerJ* 2014; **2**: e332.

772 94. Katoh K. MAFFT: a novel method for rapid multiple sequence alignment based on fast
773 Fourier transform. *Nucleic Acids Res* 2002; **30**: 3059–3066.

774 95. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
775 large phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.

776 96. Didelot X, Achtman M, Parkhill J, Thomson NR, Falush D. A bimodal pattern of
777 relatedness between the *Salmonella* Paratyphi A and Typhi genomes: convergence or
778 divergence by homologous recombination? *Genome Res* 2007; **17**: 61–8.

779 97. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better,
780 faster version of the PHAST phage search tool. *Nucleic Acids Res* 2016; **44**: W16–
781 W21.

782 98. Liu M, Li X, Xie Y, Bi D, Sun J, Li J, et al. ICEberg 2.0: an updated database of
783 bacterial integrative and conjugative elements. *Nucleic Acids Res* 2019; **47**: D660–
784 D665.

785 99. Robertson J, Nash JHE. MOB-suite: software tools for clustering, reconstruction and
786 typing of plasmids from draft assemblies. *Microb genomics* 2018; **4**.

787 100. Robertson J, Bessonov K, Schonfeld J, Nash JHE. Universal whole-sequence-based
788 plasmid typing and its utility to prediction of host range and epidemiological
789 surveillance. *Microb Genomics* 2020; **6**: 1–12.

790 101. Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, et al. SNP-sites:

rapid efficient extraction of SNPs from multi-FASTA alignments. *Microb Genomics* 2016; **2**: e000056.

102. Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with Gene Gain, Loss and Rearrangement. *PLoS One* 2010; **5**: e11147.

103. Yahara K, Nakayama S, Shimuta K, Lee K, Morita M, Kawahata T, et al. Genomic surveillance of *Neisseria gonorrhoeae* to investigate the distribution and evolution of antimicrobial-resistance determinants and lineages. *Microb Genomics* 2018; **4**: 1–3.

104. Didelot X, Wilson DJ. ClonalFrameML: Efficient Inference of Recombination in Whole Bacterial Genomes. *PLOS Comput Biol* 2015; **11**: e1004041.

105. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 2016; **44**: D7–D19.

Acknowledgements

This work was supported by Wellcome Trust grants 088786/C/09/Z and Medical Research Council (MRC) grants MR/M501608/1 and MR/L015080/1 awarded to S.K.S.

Competing interests

The authors declare no competing interests.

Figure legends

Figure 1. Barriers to horizontal gene transfer in bacteria. A series of barriers must be surmounted for DNA to transmit from one species to another. These are broadly defined in three categories. At a given time, alignment of holes in successive barriers is necessary for HGT to occur. Here we focus on ecological barriers that are influenced by multiple factors that reflect the physical isolation of bacteria in separate niches.

Figure 2. Population structure and host ecology in the genus *Campylobacter*. **a**, Phylogenetic tree of 631 *Campylobacter* isolates from 30 species reconstructed using a gene-by-gene concatenated alignment of 820 core genes (shared by >95% of isolates) and an approximation of the maximum-likelihood algorithm (ML) implemented in RAxML. The species name is indicated adjacent to the associated sequence cluster. The scale bar indicates the estimated number of substitutions per site. **b**, Isolation source of *Campylobacter* species with $n \geq 3$ isolates.

Figure 3. Core and accessory genome variation in the genus *Campylobacter*. **a**, Overall distribution of the total number of accessory genes (left) and core genes (right) per isolate for each *Campylobacter* species (where $n \geq 3$ isolates). The number of accessory genes is shown as boxplots (min to max). **b**, Venn diagram of pangenomes among different *Campylobacter* species ($n \geq 9$). The number of core genes shared by all species is illustrated in the center. **c**, Pairwise average nucleotide identity comparison calculated for all 631 *Campylobacter* isolates based upon 820 core genes shared by $>95\%$ of isolates. ANI values $<75\%$ are not calculated by FastANI [59]. **d**, Pairwise accessory genome similarity based upon gene presence or absence at 2,168 non-core loci. The heatmaps coloring ranges from yellow (minimum) to red (maximum). The matrices are ordered according to the phylogenetic tree presented in figure 2a. Different colours correspond to *Campylobacter* species with ≥ 3 isolates.

Figure 4. Elevated within-host interspecies recombination and donor-recipient comparisons. **a**, A hypothesis depicting the relationships between *Campylobacter* species, *C. jejuni* (x_1, x_2) and *C. coli* (y), when found in the same or in different hosts. **b**, Number of recombining SNPs within and between host as inferred by chromosome painting analysis for all donor recipient species comparisons. The error bar represents the standard error of the mean (SEM). **c**, The figure shows the number of donated SNPs in 10 donor-recipient pair species comparisons. The proportion (%) of recombining SNPs with $>90\%$ probability of copying from a donor to a recipient genome is illustrated in the y axis. All donor groups are shown in the x axis. All coloured boxes correspond to comparison where donors and recipients are found in the same host.

Figure 5. The mobilome of the *Campylobacter* genus. **a**, The graph illustrates the proportion of recombining genes in 10 different species comparisons. The number of species pairs in which the gene was found to recombine is shown on the x axis and the number of genes in each category is given on the y axis and. The exact number of genes found in each group comparison is shown on the top of each box. **b**, Number of *Campylobacter* species harbouring AMR genes that belong to efflux pumps and four different antibiotic classes which are shown on the x axis. **c**, The circos plot indicates the 16 genes involved in recombination in >5 donor-recipient pair species comparisons. Gene matches are indicated by joining lines, coloured differently for each gene. Gene names are shown around the perimeter for each *Campylobacter* species. **d**, The circos plot indicates the sharing of AMR genes

associated with efflux pumps and four antibiotic classes among *Campylobacter* species. Presence of at least one gene (not necessarily the same gene) conferring resistance to a specific antibiotic class is indicated by joining lines, coloured differently for each drug class. Efflux pumps (i), β -lactams (ii), tetracyclines (iii), aminoglycosides (iv) and lincosamides (v) are shown around the perimeter for each *Campylobacter* species.

Figure 2–figure supplement 1. Population structure of the *Campylobacteraceae* family.

Phylogenetic tree of 506 isolates that belong to the *Campylobacteraceae* family with *Helicobacter pylori* used as an outgroup. Different colors correspond to main species with number of isolates greater than three. The Tree was reconstructed using a gene-by-gene concatenated alignment of 799 core genes shared by >95% by all isolates and an approximation of the maximum-likelihood algorithm (ML) implemented in RAxML. The scale bar indicates the estimated number of substitutions per site.

Figure 2–figure supplement 2. Core genome species trees. Single-species trees for nine *Campylobacter* species with >4 isolates demonstrating the diversity for among species. The scale bars indicate the estimated number of substitutions per site. (*) The scale for the tree corresponding to *C. hepaticus* is 10 times smaller than the rest.

Figure 2–figure supplement 3. Overview of host-associations of *Campylobacter* species.

Abundance and diversity of 631 *Campylobacter* isolates in each host and environment. Different colours correspond to main species with number of isolates $\geq$ three. The number of isolates is shown on the y axis while the various isolation sources on the x axis.

Figure 2–figure supplement 4. Core genome species trees. Single-species trees for *C. jejuni*, *C. coli* and *C. fetus* species which contain isolates from multiple hosts and countries. The scale bars indicate the estimated number of substitutions per site.

Figure 3–figure supplement 1. Genome size variation of the *Campylobacter* genus. The frequency distribution of the genome size of all *Campylobacter* genomes used in this study is shown as a histogram. The number of genomes is shown on the y axis while the genome size (in bp) on the x axis.

Figure 3–figure supplement 2. Gene variation in the genus *Campylobacter*. Overall distribution of the total number of genes per isolate for each *Campylobacter* species (where $n \geq 3$ isolates). The number of genes is shown as boxplots (min to max).

Figure 3–figure supplement 3. Accessory gene function in all main *Campylobacter* species. The different gene functions are depicted on the y axis, while the number of shared accessory genes on the x axis. Different colours corresponding to different *Campylobacter* species.

Figure 3–figure supplement 4. Core genome allelic variation and the effect of recombination. **a**, Number of SNPs per genome of the main *Campylobacter* species (where $n \geq 3$ isolates) in the core genome alignment. The horizontal line in each plot represents the mean value while the upper and lower lines the standard deviation. **b**, Average nucleotide identity for pairwise comparisons of 820 core genes for 605 genomes of 15 main *Campylobacter* species. Different colours corresponding to different *Campylobacter* species.

Figure 4-figure supplement 1. Probability of the recipient genomes sharing DNA with each donor groups is illustrated as box whiskers (white) for every donor-recipient comparison for all 10 pairs that supported our hypothesis. The analysis where the host data were randomized across all isolates is illustrated as box whiskers (red). The probability of copying DNA from a donor to a recipient genome is shown on the y axis. The midline in the box whiskers indicates the mean and the error bars the standard deviation.

Figure 4-figure supplement 2. Genome position of genes containing recombining SNPs. Genes and their corresponding number of recombining SNPs, inferred by Chromosome Painting analysis for all 10 species comparisons, and mapped to the NCTC11168 reference genome. Genes from within-host (red) and between-host (white) pair comparisons are shown for each comparison. Donors are isolates from chicken (triangle), cattle (square), wild bird (cross), pig (star), clinical (circle) and water (snowflake) samples. The dashed line indicates the 95th percentile for every individual group comparison.

Figure 4-figure supplement 3. Genes ranked in ascending order of the number of recombining SNPs they contain as inferred by Chromosome Painting analysis for all ten species comparisons. Genes from within-host (red) and between-host (white) are shown for

each comparison. Donors are isolates from chicken (triangle), cattle (square), wild bird (cross), pig (star), clinical (circle) and water (snowflake) samples.

Figure 5-figure supplement 1. Presence of antimicrobial resistance genes in the *Campylobacter* genus. The phylogenetic tree was reconstructed using a gene-by-gene concatenated alignment of 820 core and soft-core genes and an approximation of the maximum-likelihood algorithm (ML) implemented in RAxML. The designated colour scheme was used for each species in the first column. The second column indicates whether the strain is isolated from an agricultural animal (grey). Remaining columns indicate presence of AMR genes (black). The scale represents the number of substitutions per site.

Figure 5-figure supplement 2. Genetic organization of AMR genes in *Campylobacter*. The presence of each AMR gene, highlighted in different colours, is shown for representative genomes from *C. jejuni*, *C. coli*, *C. lanienae*, *C. hyointestinalis* and *C. fetus subspecies fetus* sampled from different agricultural animals. The number of isolate genomes containing each genomic arrangement is indicated in parenthesis.

Supplementary File 1. Isolate information about the genomes used in this study.

Supplementary File 2. Within-host highly (>95th percentile) recombining genes.

Supplementary File 3. Recombination parameters as calculated by ClonalFrameML.

Supplementary File 4. Quantifying recombination between co-habiting species using ChromoPainter.

Supplementary File 5. Genes involved in interspecies recombination in 10 species comparisons.

Figure 3–source data 1. This file contains the numerical values on which the graphs in Figure 3 are based.

Figure 5–source data 1. This file contains the numerical values on which the graphs in Figure 5b-d are based.

Host ecology regulates interspecies recombination in bacteria of the genus *Campylobacter*

Evangelos Mourkas¹, Koji Yahara², Sion C. Bayliss¹, Jessica K. Calland¹, Håkan Johansson³, Leonardos Mageiros¹, Zilia Y. Muñoz-Ramirez⁴, Grant Fitcher¹, Guillaume Méric^{1#}, Matthew D. Hitchings⁵, Santiago Sandoval-Motta⁴, Javier Torres⁴, Keith A. Jolley⁶, Martin C. J. Maiden⁶, Patrik Ellström⁷, Jonas Waldenström³, Ben Pascoe^{1,8}, Samuel K. Sheppard^{1,6*}

¹The Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK; ²Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo, 162-8640, Japan; ³Centre for Ecology and Evolution in Microbial Model Systems, Linnaeus University, Kalmar, 391 82, Sweden; ⁴Unidad de Investigacion en Enfermedades Infecciosas, UMAE Pediatria, Instituto Mexicano del Seguro Social; ⁵Swansea University Medical School, Swansea University, Singleton Park, Swansea, SA2 8PP, UK; ⁶Department of Zoology, University of Oxford, South Parks Road, Oxford, OX1 3PS, UK; ⁷Department of Medical Sciences, Zoonosis Science Centre, Uppsala University, Uppsala, Sweden; ⁸Faculty of Veterinary Medicine, Chiang Mai University, Chiang Mai 50100, Thailand

*Corresponding authors: Samuel K. Sheppard; s.k.sheppard@bath.ac.uk

[#]Present address: Cambridge Baker Systems Genomics Initiative, Baker Heart and Diabetes Institute, 75 Commercial Rd, Melbourne 3004, Victoria, Australia; Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria 3004, Australia.

Keywords: *Campylobacter*, genus, species, niche, adaptation, host, evolution.

Running title: Comparative genomics of the *Campylobacter* genus

Abstract

Horizontal gene transfer (HGT) can allow traits that have evolved in one bacterial species to transfer to another. This has potential to rapidly promote new adaptive trajectories such as zoonotic transfer or antimicrobial resistance. However, for this to occur requires gaps to align in barriers to recombination within a given time frame. Chief among these barriers is the physical separation of species with distinct ecologies in separate niches. Within the genus *Campylobacter* there are species with divergent ecologies, from rarely isolated single host specialists to multi-host generalist species that are among the most common global causes of human bacterial gastroenteritis. Here, by characterising these contrasting ecologies, we can quantify HGT among sympatric and allopatric species in natural populations. Analysing recipient and donor population ancestry among genomes from 30 *Campylobacter* species we show that cohabitation in the same host can lead to a 6-fold increase in HGT between species. This accounts for up to 30% of all SNPs within a given species and identifies highly recombinogenic genes with functions including host adaptation and antimicrobial resistance. As described in some animal and plant species, ecological factors are a major evolutionary force for speciation in bacteria and changes to the host landscape can promote partial convergence of distinct species through HGT.

Introduction

It is well established that bacteria do not conform to a strict clonal model of reproduction but engage in regular horizontal gene transfer (HGT) [1]. This lateral exchange of DNA can confer new functionality on recipient genomes, potentially promoting novel adaptive trajectories such as colonization of a new host or the emergence of pathogenicity [2]. In some cases, gene flow can occur at such magnitude, even between different species [3, 4], that one may question why disparate lineages do not merge and why distinct bacterial species exist at all [5]. An answer to this lies in considering the successive processes that enable genes from one strain to establish in an entirely new genetic background.

The probability of HGT is governed by the interaction of multiple factors, including exposure to DNA, the susceptibility of the recipient genome to DNA uptake, and the impact of recombined DNA on the recipient strain. These factors can be broadly defined in three functional phases and HGT can only occur when gaps align in each successive ecological, mechanistic and adaptive barriers within a given time frame (Figure 1). In the first phase, the quantity of DNA available to recipient strains is determined by ecological factors such as the distribution, prevalence and interactions of donor and recipient bacteria, as well as the capacity for free DNA to be disseminated among species/strains. In the second phase, there are mechanistic barriers to HGT imposed by the homology dependence of recombination [6] or other factors promoting DNA specificity - such as restriction-modification, CRISPR interference or antiphage systems [7–11] - that can act as a defence against the uptake of foreign DNA (mechanistic barriers) [12, 13]. Finally, the effect that HGT has on the fitness of the recipient cell in a given selective environment (adaptive barrier) will determine if the recombinant genotype survives for subsequent generations [2, 14].

Understanding how ecology maintains, and potentially confines, distinct strains and species has become increasingly important in the light of global challenges such as the emergence and spread of zoonotic pathogens [15]. A typical approach to investigating this is to consider spillover of particular strains or clones from one host to another (clonal transmission). This is an important phenomenon and may be influenced by anthropogenic change, such as habitat encroachment or agricultural intensification [16]. However, in many cases, important phenotypes, including antimicrobial resistance (AMR) [17–19], can be conferred by relatively few genes. In such cases, it may be important to consider how cohabiting strains and species can potentially draw genes from a common pangenome pool [20–23] and how

genes, rather than clones, can transition between segregated populations (gene pool transmission). To investigate the impact of ecological segregation (ecological barriers) on this gene pool transmission, in natural populations, requires quantification of HGT among sympatric and allopatric bacteria.

Species within the genus *Campylobacter* are an ideal subject for considering how ecology influences the maintenance of genetically distinct species for several reasons. First, *Campylobacter* are a common component of the commensal gut microbiota of reptiles [24, 25], birds [26, 27] and mammals [28] but, being microaerophilic, do not survive well outside of the host. This creates island populations that have some degree of ecological isolation. Second, because at least 12 species have been identified as human pathogens [29] and *C. jejuni* and *C. coli* among the most common global causes of bacterial gastroenteritis [30], large numbers of isolate genomes have been sequenced from potential reservoir hosts as part of public health source tracking programs [31, 32]. Third, within the genus there are species and strains that inhabit one or multiple hosts (ecological specialists and generalists [16, 26, 33–37]). As a single host can simultaneously carry multiple lineages [38], possibly occupying different sub-niches within that host [39], there is potential to compare allopatric and sympatric populations. Finally, high magnitude interspecies admixture (introgression) between *C. jejuni* and *C. coli* isolated from agricultural animals suggests that host ecology plays a role in the maintenance of species [40–43].

Here, we quantify HGT among 600 genomes from 30 *Campylobacter* species using a ‘chromosome painting’ approach [44–46] to characterize shared ancestry among donor and recipient populations. Specifically, we investigate the role of ecological barriers to interspecies gene flow. By identifying recombining species pairs within the same and different hosts we can describe interactions where co-localization enhances gene flow, quantify the impact of ecological barriers in these populations and distinguish highly recombinogenic genes that are found in multiple genetic backgrounds. This provides information about the evolutionary forces that gave rise to species and the extent to which ecological barriers maintain them as discrete entities.

Results

Host restricted and host generalist *Campylobacter* species

Isolate genomes were taken from publicly available databases to represent diversity within the genus *Campylobacter*, including environmental isolates from the closely related *Arcobacter* and *Sulfurospirillum* species to provide phylogenetic context within the *Campylobacteraceae* family (Figure 2a–figure supplement 1). In total, there were 631 isolates from 30 different *Campylobacter* species (Figure 2a) and 64 different sources, isolated from 31 different countries between 1964 and 2016 (Supplementary File 1). Among the isolates, 361 were *C. jejuni* and *C. coli* and could be classified according to 31 Clonal Complexes (CCs) based upon sharing four or more alleles at seven housekeeping genes defined by multi-locus sequence typing (MLST) (Supplementary File 1) [47] and were representative of known diversity in both species [16, 33]. The obligate human commensal and pathogen *C. concisus* (n=106 isolates), comprised 2 genomospecies (GSI, n=32 and GSII, n=74), as previously described [48] (Supplementary File 1). The collection also included 52 *C. fetus* isolate genomes, including 3 subspecies: *C. fetus subsp. fetus* (n=8), *C. fetus subsp. venerealis* (n=23) and *C. fetus subsp. testudinum* (n=21) (Supplementary File 1) [49]. Two clades were observed in *C. lari* (Figure 2a–figure supplement 2) which could correspond to previously described subspecies based on 16S rRNA sequencing [50].

A maximum-likelihood phylogeny of the *Campylobacter* genus was reconstructed on a gene-by-gene concatenated sequence alignment of 820 gene families shared by >95% of all isolates, with a core genome of 903,753 base pairs (Figure 2a). The phylogeny included species which appear to be restricted to one host or environment, including *C. iguanorium* [51] and *C. geochelonis* [52] (reptiles), *C. lanienae* [53] (pigs), *C. hepaticus* [54] (chicken liver), *C. lari* group [55] (marine birds and environment) and *C. pinnipediorum* [56] (seals) species, most of which were discovered recently (Figure 2a–figure supplement 3). There was no evidence that phylogeography was reflected in the observed population structure for *Campylobacter* isolates from multiple hosts and countries. (Figure 2–figure supplement 4). This is unsurprising as it is well known that host associated genetic variation transcends phylogeographic structuring in *Campylobacter* [35]. While some low-level local gene flow can be identified within a given country [57], this is vastly outweighed by recombination within particular host niches [36], particularly in small isolate collections such as those for some of the species in this study.

Host restricted species had lower diversity possibly linked to low sample numbers, with *C. hepaticus* having the lowest diversity (Figure 2–figure supplement 2) with 8/10 genomes

associated with isolates from the same outbreak [54]. For other species there was evidence of a broad host range (ecological generalists) (Figure 2b). For example, highly structured *C. jejuni* and *C. coli* isolates were sampled from seven and six host sources respectively (Figure 2-figure supplements 2-3, Supplementary File 1). For *C. fetus* there was distinct separation between mammal-associated *C. fetus* subsp *fetus* and *C. fetus* subsp *venerealis* and reptile-associated *C. fetus* subsp *testudinum* (Figure 2-figure supplement 2) as previously described [49]. Unsurprisingly, a large proportion of the isolates in this study were from humans, likely reflecting intensive sampling. *C. jejuni* (27.52%; n=60/218), *C. coli* (14.68%; n=32/218) and *C. concisus* (44.5%; n=97/218) were all common among human clinical samples. However, less common species were also present, with nearly half of all *Campylobacter* species (44.83%, n=13/29) isolated from humans at least once (Figure 2b, Supplementary File 1). Agricultural animals were also a common source accounting for more than 1/3 of the isolates (38.35%; 242/631), with 10/30 *Campylobacter* species isolated from more than one source (Figure 2b, Supplementary File 1).

Evidence of interspecies recombination in the core and accessory genome

Genome size varied between 1.40 and 2.51 Mb (Figure 3-figure supplement 1) (mean 1.73) and the number of genes (per isolate) ranged from 1,293 to 2,170 (mean 1,675) (Figure 3a-figure supplement 2). The pangenome for the genus comprised 15,649 unique genes, found in at least one of the 631 isolates (Figure 3b-source data 1), with 820 genes (5.24 % of the pangenome) shared by >95% of all isolates (core genome), across 30 species (Figure 3b-source data 1). We excluded species with fewer than 3 isolates in subsequent analysis. For the remaining 15 species the core genome ranged in size from 1,116 genes in *C. lari* to 1,700 in *C. geochelonis* (Figure 3a right panel-source data 1). Differences were also noted in the size of accessory genomes, with *C. concisus* (mean: 981 genes), *C. hyointestinalis* (mean: 946 genes), *C. showae* (mean: 1,160 genes), *C. geochelonis* (mean: 1,021 genes) and *C. fetus* (mean: 912 genes) containing the highest average number of accessory genes (Figure 3a left panel-source data 1). Functional annotation of all 14,829 accessory genes showed that 71% (10,561) encoded hypothetical proteins of unknown function due to the lack of homology with well-characterized genes (Figure 3-figure supplement 3) [58]. Remaining genes were related to metabolism, DNA modification, transporters, virulence, inner membrane/periplasmic, adhesion, regulators, metal transport and antimicrobial resistance (Figure 3-figure supplement 3).

To further understand genetic differentiation within and between species, we generated genus-wide similarity matrices for the core and accessory genomes (Figure 3c-d–source data 1). For the core genome, pairwise average nucleotide identity (ANI) was calculated for shared genes in all possible genome pairs (Figure 3c–source data 1) using FastANI [59]. On average, isolates of the same species shared >95% similarity (Figure 3c–source data 1), with decreasing genetic similarity (between 85% and 90%) over greater phylogenetic distances. The number of core genome SNPs ranged from 983 to 230,264 for the 15 *Campylobacter* species with ≥ 3 isolates in our dataset, with *C. coli* and *C. concisus* having the greatest mean SNP numbers (Figure 3–figure supplement 4a) indicating considerable diversity within these species. In contrast *C. hepaticus* and *C. geochelonis* had low mean SNP numbers with 986 and 4,310, respectively. This is likely related to low sample numbers with isolates either sampled in close proximity [52] or from a single outbreak [54].

The core genome similarity matrix provided initial evidence of interspecies gene flow (introgression). This can be observed as elevated nucleotide identity between *C. jejuni* and clade 1 *C. coli* (Figure 3c–source data 1), consistent with previous studies [40, 42, 43]. Further evidence of introgression came from pairwise ANI comparison of genus-wide core genes, in all isolates of the 15 major *Campylobacter* species, to the *C. jejuni* genome (Figure 3–figure supplement 4b). In the absence of gene flow, isolates from the two species should have an approximately unimodal ANI distribution reflecting accumulation of mutations throughout the genome. This was largely the case but for some species, low nucleotide divergence suggested recent recombination with *C. jejuni*. There was also evidence of interspecies accessory genome recombination. Presence/absence patterns in the accessory genome matrix show considerable accessory gene sharing among several species that was inconsistent with the phylogeny (Figure 3d–source data 1). This is well illustrated in *C. lanienae* where much of the accessory genome was shared with other *Campylobacter* species (Figure 3d–source data 1).

Enhanced interspecies recombination among cohabiting species.

For *Campylobacter* inhabiting different host species there is a physical barrier to HGT. However, when there is niche overlap, interspecies recombination can occur, for example between *C. jejuni* and *C. coli* inhabiting livestock [33, 40, 42]. To understand the extent to which inhabiting different hosts impedes interspecies gene flow we quantified recombination

among *Campylobacter* species where isolates originated from same host (x_1, y) and different hosts (x_2, y) (Figure 4a).

ChromoPainterV2 software was used to infer tracts of DNA donated from multiple donor groups, belonging to the same CC but isolated from different hosts to recipient groups (Materials and Methods). Among 27 combinations of multiple donor groups and recipient groups, overall, there were more recombining SNPs within hosts than between hosts (Figure 4b) and for 10/27 species pairs there was evidence of enhanced within species recombination ($x_1 \rightarrow y > x_2 \rightarrow y$; Figure 4c). To assess the robustness of the analysis we included the effect of randomization and repeated the analysis by assigning random hosts for every strain (Figure 4-figure supplement 1). In the 10 pair species comparisons where $x_1 \rightarrow y > x_2 \rightarrow y$, we detected 174,594 within-host recombining SNPs (mapped to 473 genes; 28.8% of NCTC11168 genes) and 109,564 between-host recombining SNPs (mapped to 395 genes; 24.05% of NCTC11168 genes). From the 473 within-host recombining genes, 45 genes contained the highest number (>95th percentile) of recombining SNPs (Figure 4-figure supplements 2–3, Supplementary File 2). These genes have diverse inferred functions including metabolism, cell wall biogenesis, DNA modification, transcription, and translation (Supplementary File 2).

Interspecies recombination was observed for isolates sampled from chickens between generalist lineages CC21 and CC45 (donors; *C. jejuni*) and generalist CC828 (recipient; *C. coli*). These lineages appear to have high recombination to mutation (r/m) ratio as inferred by ClonalFrameML (Supplementary File 3). DNA from generalist *C. jejuni* CC45 was introduced into three *Campylobacter* species, including *C. hepaticus* (chicken), *C. concisus* GSI and GSII (clinical) and *C. ureolyticus* (clinical) (Figure 4c-figure supplement 2–3, Supplementary File 4). Clonal complex 45 had the highest r/m ratio from all other lineages or species involved in the comparisons (Supplementary File 3). There was increased recombination in genomes sampled from cattle between *C. jejuni* CC61 (donor; *C. jejuni*) and *C. fetus* and *C. hyointestinalis* (recipients) with 71.75% of all within-host recombining SNPs from all 10 comparisons detected in these two pairs (Figure 4c-figure supplement 2–3, Supplementary File 4). Agricultural associated *C. jejuni* CC61 and *C. fetus* subsp. *venerealis* involved in these comparisons were among the lineages and subspecies with the highest r/m ratios (Supplementary File 3). The cattle-associated CC61 has previously been described as highly recombinant, and has been associated with rapid clonal expansion and adaptation in cattle [16].

The within-host mobilome

Bacteria inhabiting the same niche may benefit from functionality conferred by similar gene combinations. Recombination can promote the dissemination of adaptive genetic elements among different bacterial species. Therefore, we postulated that the genes that recombine most among species (>95th percentile) will include those that are potentially beneficial in multiple genetic backgrounds. To investigate this, we quantified mobility within the genome identifying recombining SNPs found in more than one species comparison (Figure 5a). These SNPs mapped to 337 genes (20.52% of the NCTC11168 genes; 2.15% of the pangenome) (Figure 5a, Supplementary File 5). We found that 32 of those genes (9.49%) have also been found on plasmids (Supplementary File 5). A total of 16 genes showed elevated within-host interspecies recombination in more than five species pairs (Figure 5c, Supplementary File 5). Genes included *cmeA* and *cmeB* which are part of the predominant efflux pump CmeABC system in *Campylobacter*. Sequence variation in the drug-binding pocket of the *cmeB* gene has been linked to increased efflux function leading to resistance to multiple drugs [60]. Many of the same antimicrobial classes are used in human and veterinary medicine and this may be linked to selection for AMR *Campylobacter*, that are commonly isolated from livestock [61]. To investigate this further, we compared the genomes of all 631 isolates in our dataset to 8,762 known antibiotic resistance genes from the Comprehensive Antibiotic Resistance Database (CARD) [62], ResFinder [63] and the National Center for Biotechnology Information (NCBI) databases. Homology (>75%) was found for 42 AMR determinants associated with multi-drug efflux pumps, aminoglycosides, tetracyclines and β -lactams (Figure 5b–figure supplement 1–source data 1). Species that contained >40% isolates from livestock, including *C. jejuni*, *C. coli*, *C. lanienae*, *C. hepaticus*, *C. hyointestinalis* and *C. fetus* contained far more AMR determinants (Figure 5d–figure supplement 1–source data 1). AMR genes are often collocated in the genome [64] and our analysis revealed several gene clusters (Figure 5–figure supplement 2) that have been described in previous studies [64, 65]. These findings are consistent with HGT-mediated circulation of AMR genes among different *Campylobacter* species and support hypotheses that ecology drives gene pool transmission [2, 64].

Campylobacter host transmission and virulence have been linked with biofilm formation and changes into surface polysaccharides [66, 67]. The *carB* gene showed elevated within-host interspecies recombination in eight species pair comparisons (Figure 5c, Supplementary File

5). This gene encodes a carbamoylphosphate synthase that has been associated with biosynthesis of substrates for many polysaccharides and is known to contain transposon insertion sites upstream of its genomic position [67]. Other genes with elevated within-host interspecies transfer (>7 species pairs) included *typA* (Figure 5c, Supplementary File 5), a translator regulator for GTPase and *gltX* (Figure 5c), a *glutamate-tRNA ligase*, promoting survival under stress conditions [68, 69]. Other genes included *gidA* and *hydB* associated with virulence [70] and hydrogenase enzyme activity (respiratory pathway in *C. concisus*, 69), respectively. By considering genes that overcome barriers to interspecies recombination and establish in multiple new genetic backgrounds, it may be possible to infer important phenotypes that allow bacteria to adapt to different hosts and environments.

Discussion

Phylogenetic reconstruction of the genus *Campylobacter* revealed a highly structured population. Distinct core genome clustering largely supported known classification for species, subspecies (*C. fetus*, [49]), genomospecies (*C. concisus*, [48]) and clades (*C. coli* [42]). Also consistent with previous studies, certain species are principally associated with a specific host niche. For example, *C. fetus subsp testudinum*, *C. iguanorium*, *C. geochelonis* were only sampled from reptile species, and *C. pinnipediorum* was only sampled from seals. However, for several species there was clear evidence for host generalism, including *C. jejuni*, *C. coli* and *C. lari*, all of which were sampled from multiple hosts [26, 72]. It is clear that the hosts with the greatest diversity of *Campylobacter* species were agricultural animals (and humans) (Figure 2–figure supplement 3). While this undoubtedly reflects oversampling of these sources to some extent, the cohabitation of species in the same host niche potentially provides opportunities for interspecies HGT.

Initial evidence of interspecies gene flow came from comparison of average nucleotide identity (ANI) and the accessory genome gene presence/absence for all isolates. In each case, patterns of genetic similarity largely mirrored the phylogeny. However, consistent with previous studies [40], there was clear evidence of elevated homologous and non-homologous recombination between some species. For example, core genome ANI was higher between *C. jejuni* and *C. coli* clade 1, compared to other *C. coli* clades (Figure 3c–source data 1). The evidence for non-homologous gene sharing was even more striking with accessory genome sharing across considerable genetic distances (Figure 3d–source data 1), exemplified by *C. lanienae* which shares accessory genes with most other *Campylobacter* species.

To quantify the extent to which ecological barriers influenced interspecies gene flow, it was necessary to focus on donor-recipient species pairs where there was evidence of elevated HGT in the same (sympatry) compared to different (allopatry) hosts. Perhaps unsurprisingly, this was not the case for all species comparisons. Interacting factors could lead to genetic isolation even when species inhabit the same host. First, rather than being a single niche, the host represents a collection of subniches with varying degrees of differentiation. For example, gut-associated bacteria in the same intestinal tract have been shown to occupy different microniches [73] and more striking segregation may be expected between *C. hepaticus* inhabits the liver in poultry [54] and gut-dwelling *C. jejuni* and *C. coli* in the same host. Second, there is potential for the resident microbiota to influence the colonization potential of different *Campylobacter* species and therefore the opportunity for genetic exchange, for example through succession [74] and inhibition of transient species by residents, as seen in some other bacteria [75–77] in humans.

Continued exposition of the microecology of subniches is important but for 10 species comparisons there was clear evidence of enhanced within-host gene flow allowing quantitative analysis of ecological barriers to gene flow. Specifically, there was on average a 3-fold increase in recombination among species pairs inhabiting the same host. In some cases, this was greater, with 5-6 times more recombination among cohabiting species *C. jejuni* and *C. hyointestinalis/C. fetus* in cattle. In absolute terms, this equates to approximately 30% of all recorded SNPs in the recipient species being the result of introgression. To place this in context, if greater than half (51%) of the recorded SNPs resulted from interspecies recombination then the forces of species convergence would be greater than those that maintain distinct species. If maintained over time, these relative rates could lead to progressive genetic convergence unless countered by strong genome-wide natural selection against introgressed DNA.

Quantitative SNP-based comparisons clearly ignore one very important factor. Specifically, that recombined genes that do not reduce the fitness of the recipient genome (provide an adaptive advantage) will remain in the population while others will be purged through natural selection. Therefore, by identifying genomic hotspots of recombination and the putative function of genes that recombine between species it is possible to understand more about micro-niche segregation and the host adapted gene pool. Of the 35 genes with evidence of

enhanced within host HGT in ≥ 5 species pairs, several were linked to functions associated with proliferation in, and exploitation of, the host. For example, the *carB* gene, encoding the large subunit of carbamoylphosphatase associated with polysaccharide biosynthesis, recombined in eight cohabiting species pairs and is potentially linked to enhanced virulence and growth [67]. In addition, other highly mobile genes, including *typA* and *gltX* are associated with survival and proliferation in stress conditions [68, 69], and *hydB* is linked to NiFe hydrogenase and nickel uptake that is essential for the survival of *C. jejuni* in the gut of birds and mammals [78].

Some genes showed evidence of elevated recombination in a specific host species. For example, the *glmS* and *napA* genes in cohabiting *Campylobacter* species in cattle. In many bacteria, analogs of *glmS* have multiple downstream integration specific sites (Tn7) [79] which may explain the mobility of this gene. Explaining the mobility of *napA* is less straightforward, but this gene is known to encode a nitrate reductase in *Campylobacter* [80] in microaerobic conditions which may be ecologically significant as the accumulation of nitrate in slurry, straw and drainage water can be potentially toxic to livestock mammals [81].

Factors such as host physiology, diet, and metabolism undoubtedly impose selection pressures upon resident bacteria and the horizontal acquisition of genes provides a possible vehicle for adaptation. However, the widespread use of antimicrobials by humans, pets and livestock production [82, 83], provides another major ecological barrier to niche colonization. We found that *gyrA* was among the most recombinogenic genes in *Campylobacter* in chickens. This is important as a single mutation in this gene is known to confer resistance to ciprofloxacin [84]. While the rising trend in fluoroquinolones resistance in *Campylobacter* from humans and livestock [85] may result from spontaneous independent mutations, it is likely that it is accelerated by HGT. However, there is currently no clear evidence for the transfer of resistant versions of *gyrA*. Interspecies recombination of AMR genes has been observed between *C. jejuni* and *C. coli* isolates from multiple sources including livestock, human and sewage [64]. Consistent with this, we found AMR genes present in strains from 12 *Campylobacter* species in multiple hosts (Figure 5–figure supplement 2). In some cases, strains from phylogenetically closely related species (*C. fetus* and *C. hyointestinalis*) isolated from cattle, shared the same AMR gene cluster (*tet44* and *ant(6)-Ib*) described before in *C. fetus* subsp. *fetus* [65], indicating the circulation of colocalized AMR genes among related species and host niche gene pools. Strikingly, the efflux pump genes *cmeA* and *cmeB*,

associated with multidrug resistance (MDR) were highly mobile among *Campylobacter* species with evidence of elevated within host interspecies recombination in >7 species pairs. Furthermore, the *gltX* gene, which when phosphorylated by protein kinases promotes MDR [69], was also among the most introgressed genes. While a deeper understanding of gene interactions, epistasis and epigenetics would be needed to prove that the lateral acquisition of AMR genes promotes niche adaptation, these data do suggest that HGT may facilitate colonization of antimicrobial-rich host environments, potentially favouring the spread of genes into multiple genetic backgrounds.

In conclusion, we show that species within the genus *Campylobacter* include those that are host restricted as well as host generalists. When species cohabit in the same host, ecological barriers to recombination can be perforated leading to considerable introgression between species. While the magnitude of introgression varies, potentially reflecting microniche structure with the host, there is clear evidence that ecology is important in maintaining genetically distinct species. This parallels evolution in some interbreeding eukaryotes, such as Darwin's Finches, where fluctuating environmental conditions can change the selection pressures acting on species inhabiting distinct niches, potentially favouring hybrids [86, 87]. Consistent with this, the host landscape is changing for *Campylobacter*, with intensively reared livestock now constituting 60-70% of bird and mammal biomass on earth respectively [88]. This creates opportunities for species to be brought together in new adaptive landscapes and for genes to be tested from multiple genetic backgrounds. By understanding the ecology of niche segregation and the genetics of bacterial adaptation we can hope to improve strategies and interventions to reduce the risk of zoonotic transmission and the spread of problematic genes among species.

Materials & Methods

Isolate genomes

A total of 631 *Campylobacter*, 17 *Arcobacter*, seven *Sulfurospirillum* and five *Helicobacter* genomes were assembled from previously published datasets (Supplementary File 1). Isolates were sampled from clinical cases of campylobacteriosis and faeces of chickens, ruminants, wild birds, wild mammals, pets and environmental sources. Genomes and related metadata were uploaded and archived in the BIGS database [89]. Quality control was performed based on the genome size, number of contigs, N50 and N95 contig length using the integrated tools in BIGS database. All assembled contigs were further screened for contamination and

completeness using CheckM [90] (Supplementary File 1). All assembled genomes can be downloaded from FigShare (doi: 10.6084/m9.figshare.15061017). Comparative genomics analyses focused on the *Campylobacter* genomes representing 30 species including: *C. avium* (n=1); *C. coli* (n=143); *C. concisus* (n=106); *C. corcagiensis* (n=1); *C. cuniculorum* (n=2); *C. curvus* (n=2); *C. fetus* (n=52); *C. geochelonis* (n=3); *C. gracilis* (n=2); *C. helveticus* (n=1); *C. hepaticus* (n=10); *C. hominis* (n=1); *C. hyointestinalis* (n=16); *C. iguanorium* (n=3); *C. insulaenigrae* (n=1); *C. jejuni* (n=218); *C. lanienae* (n=26); *C. lari* (n=13); *C. mucosalis* (n=1); *C. ornithocola* (n=1); *C. peloridis* (n=1); *C. pinnipediorum* (n=9); *C. rectus* (n=1); *C. showae* (n=3); *C. sputorum* (n=1); *C. subantarcticus* (n=3); *C. upsaliensis* (n=3); *C. ureolyticus* (n=4); *C. volucris* (n=2); *Campylobacter sp* (n=1) (Supplementary File 1). Genomes belonging to *C. jejuni* and *C. coli* species were selected to represent a wide range of hosts, sequence types, and clonal complexes and reflect the known population structure for these two species. For other *Campylobacter* species, all genomes that were publicly available at the time of this study were included in the analysis. (Supplementary File 1).

Pangenome characterization and phylogenetic analysis

Sequence data were analysed using PIRATE, a fast and scalable pangenomics tool which allows for orthologue gene clustering in divergent bacterial species [91]. Genomes were annotated in Prokka [92], using a genus database comprising well annotated *C. jejuni* strains NCTC11168, 81116, 81-176 and M1, and plasmids pTet and pVir in addition to the already existing databases used by Prokka [92]. Briefly, annotated genomes were used as input for PIRATE. Non-redundant representative sequences were produced using CD-HIT and the longest sequence was used as a reference for sequence similarity interrogation using BLAST/DIAMOND. Gene orthologues were defined as “gene families” and were clustered in different MCL thresholds, from 10 to 98 % sequence identity (10, 20, 30, 40, 50, 60, 70, 80, 90, 95, 98). Higher MCL thresholds were used to identify allelic variation within different loci. An inflation value of 4 was used to increase the granularity of MCL clustering between gene families. BLAST high-scoring pairs with a reciprocal minimum length of 90% of the query/subject sequence were excluded from MCL clustering to reduce the number of spurious associations between distantly related or conserved genes [93]. This information was used to generate gene presence/absence and allelic variation matrices. A core gene-by-gene multiple sequence alignment [89] was generated using MAFFT [94] comprising genes shared >95% of isolates. Phylogenetic trees, based on core gene-by-gene alignments, were reconstructed

using the maximum-likelihood algorithm implemented in RAxML v8.2.11 [95] with GTRGAMMA as substitution model.

Quantifying core and accessory genome variation

The degree of genetic differentiation between species was investigated gene-by-gene as in previous studies [40, 96] by calculating the average nucleotide identity (ANI) of all 631 *Campylobacter* genomes using FastANI v.1.0 [59]. The analysis generated a lower triangular matrix with the lowest ANI value at 75% (as computed by FastANI). A comparable gene presence/absence matrix was produced using PIRATE and was further used to generate a heatmap of accessory genome similarity based upon gene presence or absence. Subsequently, all *Campylobacter* genomes were screened for the presence of antimicrobial resistance genes against the CARD [62], ResFinder [63] and NCBI databases. All *Campylobacter* genomes were further screened for the presence of phage, conjugative elements and plasmid DNA using publicly available online databases to investigate the effect of other transfer mechanisms. First, we used the PHAge Search Tool Enhanced Release (PHASTER) [97] to identify and annotate prophage sequences within our genomes. A total of 86% (254/297) of the genomes used in chromosome painting were found to have DNA sequence of phage origin. Second, we used Iceberg 2.0 [98] for the detection of integrative and conjugative elements, identifying 32 ICEs in 19% (56/297) of the genomes used in the chromosome painting analysis. Finally, we used MOB-suite software for clustering, reconstruction and typing of plasmids from draft assemblies [99, 100]. A positive hit was defined when a gene had >75% nucleotide identity over >50% of the sequence length showing that 32 genes identified in the recombination analysis have also been located on plasmids. A gene presence/absence matrix for every antimicrobial resistance gene was generated for every genome. Genomes carrying AMR genes were screened to characterize the location of adjacent genes using SnapGene software (GSL Biotech; available at snapgene.com), as previously described [64]. The number of core SNPs was identified using SNP-sites (v2.3.2) [101].

Inference of recombination

Each combination of a recipient group and multiple donor groups (belonging to the same CC but isolated from different hosts) was selected to compare the extent of interspecies recombination into the recipient genomes. Each donor group consisted of 8 isolates to avoid the influence of difference in sample size on estimation of the extent of interspecies

recombination. Each recipient group included at least 4 isolates. We excluded *C. jejuni* and *C. coli* clade 1 genomes isolated from seals and water, as these most likely represent spillover events and not true host segregated populations. Briefly, we conducted a pairwise genome alignment between reference genome NCTC11168 and one of the strains included in the donor-recipient analysis using progressiveMauve [102]. This enabled the construction of positional homology alignments for all genomes regardless gene content and genome rearrangements, which were then combined into a multiple whole-genome alignment, as previously described [103]. ChromoPainterV2 software was used to calculate the amount of DNA sequence that is donated from a donor to a recipient group [45]. Briefly, for each donor-recipient pair, SNPs in which >90% recipient individuals had recombined with the donor group were considered in the analysis. These SNPs were mapped to genomic regions and specific genes were identified. A total of 258,444 (96.83%) recombining SNPs mapped to 558 genes of the NCTC11168 reference strain with >90% probability of copying from a donor to a recipient strain. Genes containing the highest number of recombining SNPs were considered for subsequent analyses (>95th percentile) (Supplementary File 2). ClonalFrameML[104] was used to infer the relative number of substitutions introduced by recombination (r) and mutation (m) as the ratio r/m as previously described [16].

Data availability

Genomes sequenced as part of other studies are archived on the Short Read Archive associated with BioProject accessions: PRJNA176480, PRJNA177352, PRJNA342755, PRJNA345429, PRJNA312235, PRJNA415188, PRJNA524300, PRJNA528879, PRJNA529798, PRJNA575343, PRJNA524315 and PRJNA689604. Additional genomes were also downloaded from NCBI [105] and pubMLST (<http://pubmlst.org/campylobacter>). Contiguous assemblies of all genome sequences compared are available at the public data repository Figshare (doi: 10.6084/m9.figshare.15061017) and individual project and accession numbers can be found in Supplementary File 1.

References

1. Smith JM, Dowson CG, Spratt BG. Localized sex in bacteria. *Nature* 1991; **349**: 29–31.
2. Sheppard SK, Guttman DS, Fitzgerald JR. Population genomics of bacterial host adaptation. *Nat Rev Genet* 2018; **19**: 549–565.
3. Shapiro BJ, Leducq J-B, Mallet J. What Is Speciation? *PLOS Genet* 2016; **12**:

521 e1005860.

522 4. Doolittle WF, Zhaxybayeva O. On the origin of prokaryotic species. *Genome Res*
523 2009; **19**: 744–756.

524 5. Doolittle WF, Papke RT. Genomics and the bacterial species problem. *Genome Biol*
525 2006; **7**: 116.

526 6. Fraser C, Hanage WP, Spratt BG. Recombination and the nature of bacterial
527 speciation. *Science* 2007; **315**: 476–80.

528 7. Budroni S, Siena E, Hotopp JCD, Seib KL, Serruto D, Nofroni C, et al. *Neisseria*
529 *meningitidis* is structured in clades associated with restriction modification systems
530 that modulate homologous recombination. *Proc Natl Acad Sci* 2011; **108**: 4494–4499.

531 8. Oliveira PH, Touchon M, Rocha EPC. Regulation of genetic flux between bacteria by
532 restriction–modification systems. *Proc Natl Acad Sci* 2016; **113**: 5658–5663.

533 9. Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, et al. Systematic
534 discovery of antiphage defense systems in the microbial pangenome. *Science* (80-)
535 2018; **359**: eaar4120.

536 10. Nandi T, Holden MTG, Didelot X, Mehershahi K, Boddey JA, Beacham I, et al.
537 *Burkholderia pseudomallei* sequencing identifies genomic clades with distinct
538 recombination, accessory, and epigenetic profiles. *Genome Res* 2015; **25**: 129–141.

539 11. Marraffini LA, Sontheimer EJ. CRISPR Interference Limits Horizontal Gene Transfer
540 in Staphylococci by Targeting DNA. *Science* (80-) 2008; **322**: 1843–1845.

541 12. Thomas CM, Nielsen KM. Mechanisms of, and Barriers to, Horizontal Gene Transfer
542 between Bacteria. *Nat Rev Microbiol* 2005; **3**: 711–721.

543 13. Eggleston AK, Mitchell AH, West SC. In Vitro Reconstitution of the Late Steps of
544 Genetic Recombination in *E. coli*. *Cell* 1997; **89**: 607–617.

545 14. Zhu P, van der Ende A, Falush D, Brieske N, Morelli G, Linz B, et al. Fit genotypes
546 and escape variants of subgroup III *Neisseria meningitidis* during three pandemics of
547 epidemic meningitis. *Proc Natl Acad Sci* 2001; **98**: 5234–5239.

548 15. Boni MF, Lemey P, Jiang X, Lam TT-Y, Perry BW, Castoe TA, et al. Evolutionary
549 origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19
550 pandemic. *Nat Microbiol* 2020; **5**: 1408–1417.

551 16. Mourkas E, Taylor AJ, Méric G, Bayliss SC, Pascoe B, Mageiros L, et al. Agricultural
552 intensification and the evolution of host specialism in the enteric pathogen
553 *Campylobacter jejuni*. *Proc Natl Acad Sci* 2020; **117**: 11018–11028.

554 17. Johnson AP, Woodford N. Global spread of antibiotic resistance: the example of New

- 555 Delhi metallo- β -lactamase (NDM)-mediated carbapenem resistance. *J Med Microbiol*
556 2013; **62**: 499–513.
- 557 18. Schwarz S, Johnson AP. Transferable resistance to colistin: a new but old threat. *J*
558 *Antimicrob Chemother* 2016; **71**: 2066–70.
- 559 19. Baker KS, Dallman TJ, Field N, Childs T, Mitchell H, Day M, et al. Horizontal
560 antimicrobial resistance transfer drives epidemics of multiple *Shigella* species. *Nat*
561 *Commun* 2018; **9**: 1462.
- 562 20. Young JPW. Bacteria Are Smartphones and Mobile Genes Are Apps. *Trends*
563 *Microbiol* 2016; **24**: 931–932.
- 564 21. McInerney JO, Whelan FJ, Domingo-Sananes MR, McNally A, O’Connell MJ.
565 Pangenomes and Selection: The Public Goods Hypothesis. *The Pangenome*. 2020.
566 Springer International Publishing, Cham, pp 151–167.
- 567 22. Vos M, Eyre-Walker A. Are pangenomes adaptive or not? *Nat Microbiol* 2017; **2**:
568 1576.
- 569 23. Werren JH. Selfish genetic elements, genetic conflict, and evolutionary innovation.
570 *Proc Natl Acad Sci* 2011; **108**: 10863–10870.
- 571 24. Giacomelli M, Piccirillo A. Pet reptiles as potential reservoir of *Campylobacter*
572 species with zoonotic potential: TABLE 1: *Vet Rec* 2014; **174**: 479.2-479.
- 573 25. Fitzgerald C, Tu Z chao, Patrick M, Stiles T, Lawson AJ, Santovenia M, et al.
574 *Campylobacter fetus* subsp. *testudinum* subsp. nov., isolated from humans and reptiles.
575 *Int J Syst Evol Microbiol* 2014; **64**: 2944–2948.
- 576 26. Griekspoor P, Colles FM, McCarthy ND, Hansbro PM, Ashhurst-Smith C, Olsen B, et
577 al. Marked host specificity and lack of phylogeographic population structure of
578 *Campylobacter jejuni* in wild birds. *Mol Ecol* 2013; **22**: 1463–1472.
- 579 27. Atterby C, Mourkas E, Méric G, Pascoe B, Wang H, Waldenström J, et al. The
580 Potential of Isolation Source to Predict Colonization in Avian Hosts: A Case Study in
581 *Campylobacter jejuni* Strains From Three Bird Species. *Front Microbiol* 2018; **9**.
- 582 28. Leatherbarrow AJH, Griffiths R, Hart CA, Kemp R, Williams NJ, Diggle PJ, et al.
583 *Campylobacter lari*: genotype and antibiotic resistance of isolates from cattle, wildlife
584 and water in an area of mixed dairy farmland in the United Kingdom. *Environ*
585 *Microbiol* 2007; **9**: 1772–1779.
- 586 29. Man SM. The clinical importance of emerging *Campylobacter* species. *Nat Rev*
587 *Gastroenterol Hepatol* 2011; **8**: 669–685.
- 588 30. Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global Epidemiology of

- Campylobacter* Infection. *Clin Microbiol Rev* 2015; **28**: 687–720.
31. Sheppard SK, Dallas JF, MacRae M, McCarthy ND, Sproston EL, Gormley FJ, et al. *Campylobacter* genotypes from food animals, environmental sources and clinical disease in Scotland 2005/6. *Int J Food Microbiol* 2009; **134**: 96–103.
 32. Sheppard SK, Dallas JF, Strachan NJC, MacRae M, McCarthy ND, Wilson DJ, et al. *Campylobacter* genotyping to determine the source of human infection. *Clin Infect Dis* 2009; **48**: 1072–8.
 33. Sheppard SK, Colles FM, McCarthy ND, Strachan NJC, Ogden ID, Forbes KJ, et al. Niche segregation and genetic structure of *Campylobacter jejuni* populations from wild and agricultural host species. *Mol Ecol* 2011; **20**: 3484–90.
 34. Dearlove BL, Cody AJ, Pascoe B, Méric G, Wilson DJ, Sheppard SK. Rapid host switching in generalist *Campylobacter* strains erodes the signal for tracing human infections. *ISME J* 2016; **10**: 721–729.
 35. Sheppard SK, Colles F, Richardson J, Cody AJ, Elson R, Lawson A, et al. Host Association of *Campylobacter* Genotypes Transcends Geographic Variation. *Appl Environ Microbiol* 2010; **76**: 5269–5277.
 36. Sheppard SK, Cheng L, Méric G, de Haan CPA, Llarena A-K, Marttinen P, et al. Cryptic ecology among host generalist *Campylobacter jejuni* in domestic animals. *Mol Ecol* 2014; **23**: 2442–2451.
 37. Woodcock DJ, Krusche P, Strachan NJC, Forbes KJ, Cohan FM, Méric G, et al. Genomic plasticity and rapid host switching can promote the evolution of generalism: a case study in the zoonotic pathogen *Campylobacter*. *Sci Rep* 2017; **7**: 9650.
 38. Colles FM, Dingle KE, Cody AJ, Maiden MCJ. Comparison of *Campylobacter* Populations in Wild Geese with Those in Starlings and Free-Range Poultry on the Same Farm. *Appl Environ Microbiol* 2008; **74**: 3583–3590.
 39. Colles FM, McCarthy ND, Bliss CM, Layton R, Maiden MCJ. The long-term dynamics of *Campylobacter* colonizing a free-range broiler breeder flock: an observational study. *Environ Microbiol* 2015; **17**: 938–946.
 40. Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Méric G, et al. Progressive genome-wide introgression in agricultural *Campylobacter coli*. *Mol Ecol* 2013; **22**: 1051–64.
 41. Taylor AJ, Méric G, Yahara K, Pascoe B, Mageiros L, Mourkas E, et al. Cross-species evolutionary rewiring in the enteric bacterium *Campylobacter*; *bioRxiv* 2021; 2021.03.15.435406.

42. Sheppard SK, McCarthy ND, Falush D, Maiden MCJ. Convergence of *Campylobacter* Species: Implications for Bacterial Evolution. *Science* (80-) 2008; **320**: 237–239.
43. Sheppard SK, McCarthy ND, Jolley KA, Maiden MCJ. Introgression in the genus *Campylobacter*: generation and spread of mosaic alleles. *Microbiology* 2011; **157**: 1066–1074.
44. Thorell K, Yahara K, Berthenet E, Lawson DJ, Mikhail J, Kato I, et al. Rapid evolution of distinct *Helicobacter pylori* subpopulations in the Americas. *PLOS Genet* 2017; **13**: e1006730.
45. Lawson DJ, Hellenthal G, Myers S, Falush D. Inference of Population Structure using Dense Haplotype Data. *PLoS Genet* 2012; **8**: e1002453.
46. Yahara K, Furuta Y, Oshima K, Yoshida M, Azuma T, Hattori M, et al. Chromosome Painting In Silico in a Bacterial Species Reveals Fine Population Structure. *Mol Biol Evol* 2013; **30**: 1454–1464.
47. Dingle KE, Colles FM, Wareing DR, Ure R, Fox AJ, Bolton FE, et al. Multilocus sequence typing system for *Campylobacter jejuni*. *J Clin Microbiol* 2001; **39**: 14–23.
48. Kirk KF, Méric G, Nielsen HL, Pascoe B, Sheppard SK, Thorlacius-Ussing O, et al. Molecular epidemiology and comparative genomics of *Campylobacter concisus* strains from saliva, faeces and gut mucosal biopsies in inflammatory bowel disease. *Sci Rep* 2018; **8**: 1902.
49. Iraola G, Forster SC, Kumar N, Lehours P, Bekal S, García-Peña FJ, et al. Distinct *Campylobacter fetus* lineages adapted as livestock pathogens and human pathobionts in the intestinal microbiota. *Nat Commun* 2017; **8**: 1367.
50. Debruyne L, On SLW, De Brandt E, Vandamme P. Novel *Campylobacter lari*-like bacteria from humans and molluscs: description of *Campylobacter peloridis* sp. nov., *Campylobacter lari* subsp. *concheus* subsp. nov. and *Campylobacter lari* subsp. *lari* subsp. nov. *Int J Syst Evol Microbiol* 2009; **59**: 1126–32.
51. Gilbert MJ, Kik M, Miller WG, Duim B, Wagenaar JA. *Campylobacter iguaniorum* sp. nov., isolated from reptiles. *Int J Syst Evol Microbiol* 2015; **65**: 975–982.
52. Piccirillo A, Niero G, Calleros L, Pérez R, Naya H, Iraola G. *Campylobacter geocheilonis* sp. nov. isolated from the western Hermann's tortoise (*Testudo hermanni hermanni*). *Int J Syst Evol Microbiol* 2016; **66**: 3468–3476.
53. Logan JMJ, Burnens A, Linton D, Lawson AJ, Stanley J. *Campylobacter lanienae* sp. nov., a new species isolated from workers in an abattoir. *Int J Syst Evol Microbiol* 2000; **50**: 865–872.

54. Van TTH, Elshagmani E, Gor MC, Scott PC, Moore RJ. *Campylobacter hepaticus* sp. nov., isolated from chickens with spotty liver disease. *Int J Syst Evol Microbiol* 2016; **66**: 4518–4524.
55. Miller WG, Yee E, Chapman MH, Smith TPL, Bono JL, Huynh S, et al. Comparative genomics of the *Campylobacter lari* group. *Genome Biol Evol* 2014; **6**: 3252–66.
56. Gilbert MJ, Miller WG, Leger JS, Chapman MH, Timmerman AJ, Duim B, et al. *Campylobacter pinnipediorum* sp. nov., isolated from pinnipeds, comprising *Campylobacter pinnipediorum* subsp. *pinnipediorum* subsp. nov. and *Campylobacter pinnipediorum* subsp. *caledonicus* subsp. nov. *Int J Syst Evol Microbiol* 2017; **67**: 1961–1968.
57. Pascoe B, Méric G, Yahara K, Wimalaratna H, Murray S, Hitchings MD, et al. Local genes for local bacteria: Evidence of allopatry in the genomes of transatlantic *Campylobacter* populations. *Mol Ecol* 2017; **26**: 4497–4508.
58. Pascoe B, Williams LK, Calland JK, Méric G, Hitchings MD, Dyer M, et al. Domestication of *Campylobacter jejuni* NCTC 11168. *Microb genomics* 2019; **5**.
59. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018; **9**: 5114.
60. Yao H, Shen Z, Wang Y, Deng F, Liu D, Naren G, et al. Emergence of a Potent Multidrug Efflux Pump Variant That Enhances *Campylobacter* Resistance to Multiple Antibiotics. *MBio* 2016; **7**: 1–11.
61. Livermore DM. Introduction: the challenge of multiresistance. *Int J Antimicrob Agents* 2007; **29**: S1–S7.
62. Jia B, Raphenya AR, Alcock B, Wagglechner N, Guo P, Tsang KK, et al. CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res* 2017; **45**: D566–D573.
63. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 2012; **67**: 2640–2644.
64. Mourkas E, Florez-Cuadrado D, Pascoe B, Calland JK, Bayliss SC, Mageiros L, et al. Gene pool transmission of multidrug resistance among *Campylobacter* from livestock, sewage and human disease. *Environ Microbiol* 2019; **21**: 4597–4613.
65. Abril C, Brodard I, Perreten V. Two Novel Antibiotic Resistance Genes, tet(44) and ant(6)-Ib, Are Located within a Transferable Pathogenicity Island in *Campylobacter*

- fetus* subsp. *fetus*. *Antimicrob Agents Chemother* 2010; **54**: 3052–3055.
66. Szymanski CM, Logan SM, Linton D, Wren BW. *Campylobacter* – a tale of two protein glycosylation systems. *Trends Microbiol* 2003; **11**: 233–238.
 67. McLennan MK, Ringoir DD, Frirdich E, Svensson SL, Wells DH, Jarrell H, et al. *Campylobacter jejuni* Biofilms Up-Regulated in the Absence of the Stringent Response Utilize a Calcofluor White-Reactive Polysaccharide. *J Bacteriol* 2008; **190**: 1097–1107.
 68. Margus T, Remm M, Tenson T. Phylogenetic distribution of translational GTPases in bacteria. *BMC Genomics* 2007; **8**: 1–18.
 69. Semanjski M, Germain E, Bratl K, Kiessling A, Gerdes K, Macek B. The kinases HipA and HipA7 phosphorylate different substrate pools in *Escherichia coli* to promote multidrug tolerance. *Sci Signal* 2018; **11**.
 70. Mikheil DM, Shippy DC, Eakley NM, Okwumabua OE, Fadl AA. Deletion of gene encoding methyltransferase (*gidB*) confers high-level antimicrobial resistance in *Salmonella*. *J Antibiot (Tokyo)* 2012; **65**: 185–192.
 71. Benoit SL, Maier RJ. Site-directed mutagenesis of *Campylobacter concisus* respiratory genes provides insight into the pathogen's growth requirements. *Sci Rep* 2018; **8**: 14203.
 72. Cody AJ, McCarthy ND, Bray JE, Wimalarathna HML, Colles FM, Jansen van Rensburg MJ, et al. Wild bird-associated *Campylobacter jejuni* isolates are a consistent source of human disease, in Oxfordshire, United Kingdom. *Environ Microbiol Rep* 2015; **7**: 782–788.
 73. Hayashi H, Takahashi R, Nishi T, Sakamoto M, Benno Y. Molecular analysis of jejunal, ileal, caecal and recto-sigmoidal human colonic microbiota using 16S rRNA gene libraries and terminal restriction fragment length polymorphism. *J Med Microbiol* 2005; **54**: 1093–1101.
 74. Lu J, Idris U, Harmon B, Hofacre C, Maurer JJ, Lee MD. Diversity and succession of the intestinal bacterial community of the maturing broiler chicken. *Appl Environ Microbiol* 2003; **69**: 6816–24.
 75. Stecher B, Chaffron S, Käppeli R, Hapfelmeier S, Friedrich S, Weber TC, et al. Like will to like: abundances of closely related species can predict susceptibility to intestinal colonization by pathogenic and commensal bacteria. *PLoS Pathog* 2010; **6**: e1000711.
 76. van Elsas JD, Chiurazzi M, Mallon CA, Elhottova D, Kristufek V, Salles JF. Microbial

- diversity determines the invasion of soil by a bacterial pathogen. *Proc Natl Acad Sci U S A* 2012; **109**: 1159–64.
77. Nowrouzian FL, Wold AE, Adlerberth I. *Escherichia coli* strains belonging to phylogenetic group B2 have superior capacity to persist in the intestinal microflora of infants. *J Infect Dis* 2005; **191**: 1078–83.
78. Howlett RM, Hughes BM, Hitchcock A, Kelly DJ. Hydrogenase activity in the foodborne pathogen *Campylobacter jejuni* depends upon a novel ABC-type nickel transporter (NikZYXWV) and is SlyD-independent. *Microbiology* 2012; **158**: 1645–1655.
79. Choi K-H. Applications of Transposon-Based Gene Delivery System in Bacteria. *J Microbiol Biotechnol* 2009; **19**: 217–28.
80. Pittman MS, Elvers KT, Lee L, Jones MA, Poole RK, Park SF, et al. Growth of *Campylobacter jejuni* on nitrate and nitrite: electron transport to NapA and NrfA via NrfH and distinct roles for NrfA and the globin Cgb in protection against nitrosative stress. *Mol Microbiol* 2007; **63**: 575–590.
81. Alexander J, Benford D, Cockburn A, Cravedi J, Dogliotti E, Domenico A Di, et al. Nitrite as undesirable substances in animal feed - Scientific Opinion of the Panel on Contaminants in the Food Chain. *EFSA J* 2009; **7**: 1–47.
82. Teuber M. Veterinary use and antibiotic resistance. *Curr Opin Microbiol* 2001; **4**: 493–499.
83. Price LB, Koch BJ, Hungate BA. Ominous projections for global antibiotic use in food-animal production. *Proc Natl Acad Sci U S A* 2015; **112**: 5554–5.
84. Luo N, Sahin O, Lin J, Michel LO, Zhang Q. In Vivo Selection of *Campylobacter* Isolates with High Levels of Fluoroquinolone Resistance Associated with *gyrA* Mutations and the Function of the CmeABC Efflux Pump. *Antimicrob Agents Chemother* 2003; **47**: 390–394.
85. Sproston EL, Wimalarathna HML, Sheppard SK. Trends in fluoroquinolone resistance in *Campylobacter*. *Microb Genomics* 2018; **4**: 1–8.
86. Mallet J. Hybrid speciation. *Nature* 2007; **446**: 279–283.
87. Grant PR, Grant BR. Hybridization of Bird Species. *Science* (80-) 1992; **256**: 193–197.
88. Bar-On YM, Phillips R, Milo R. The biomass distribution on Earth. *Proc Natl Acad Sci* 2018; **115**: 6506–6511.
89. Sheppard SK, Jolley KA, Maiden MCJ. A Gene-By-Gene Approach to Bacterial

- Population Genomics: Whole Genome MLST of *Campylobacter*. *Genes (Basel)* 2012; **3**: 261–277.
90. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 2015; **25**: 1043–1055.
91. Bayliss SC, Thorpe HA, Coyle NM, Sheppard SK, Feil EJ. PIRATE: A fast and scalable pangenomics toolbox for clustering diverged orthologues in bacteria. *Gigascience* 2019; **8**: 1–9.
92. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014; **30**: 2068–2069.
93. Sahl JW, Caporaso JG, Rasko DA, Keim P. The large-scale blast score ratio (LS-BSR) pipeline: a method to rapidly compare genetic content between bacterial genomes. *PeerJ* 2014; **2**: e332.
94. Katoh K. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 2002; **30**: 3059–3066.
95. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.
96. Didelot X, Achtman M, Parkhill J, Thomson NR, Falush D. A bimodal pattern of relatedness between the *Salmonella* Paratyphi A and Typhi genomes: convergence or divergence by homologous recombination? *Genome Res* 2007; **17**: 61–8.
97. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 2016; **44**: W16–W21.
98. Liu M, Li X, Xie Y, Bi D, Sun J, Li J, et al. ICEberg 2.0: an updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res* 2019; **47**: D660–D665.
99. Robertson J, Nash JHE. MOB-suite: software tools for clustering, reconstruction and typing of plasmids from draft assemblies. *Microb genomics* 2018; **4**.
100. Robertson J, Bessonov K, Schonfeld J, Nash JHE. Universal whole-sequence-based plasmid typing and its utility to prediction of host range and epidemiological surveillance. *Microb Genomics* 2020; **6**: 1–12.
101. Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, et al. SNP-sites: rapid efficient extraction of SNPs from multi-FASTA alignments. *Microb Genomics* 2016; **2**: e000056.

102. Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with Gene Gain, Loss and Rearrangement. *PLoS One* 2010; **5**: e11147.
103. Yahara K, Nakayama S, Shimuta K, Lee K, Morita M, Kawahata T, et al. Genomic surveillance of *Neisseria gonorrhoeae* to investigate the distribution and evolution of antimicrobial-resistance determinants and lineages. *Microb Genomics* 2018; **4**: 1–3.
104. Didelot X, Wilson DJ. ClonalFrameML: Efficient Inference of Recombination in Whole Bacterial Genomes. *PLOS Comput Biol* 2015; **11**: e1004041.
105. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 2016; **44**: D7–D19.

Acknowledgements

This work was supported by Wellcome Trust grants 088786/C/09/Z and Medical Research Council (MRC) grants MR/M501608/1 and MR/L015080/1 awarded to S.K.S. The computational calculations were performed at the Human Genome Center at the Institute of Medical Science (University of Tokyo) and at the National Institute of Genetics.

Competing interests

The authors declare no competing interests.

Figure legends

Figure 1. Barriers to horizontal gene transfer in bacteria. A series of barriers must be surmounted for DNA to transmit from one species to another. These are broadly defined in three categories. At a given time, alignment of holes in successive barriers is necessary for HGT to occur. Here we focus on ecological barriers that are influenced by multiple factors that reflect the physical isolation of bacteria in separate niches.

Figure 2. Population structure and host ecology in the genus *Campylobacter*. **a**, Phylogenetic tree of 631 *Campylobacter* isolates from 30 species reconstructed using a gene-by-gene concatenated alignment of 820 core genes (shared by >95% of isolates) and an approximation of the maximum-likelihood algorithm (ML) implemented in RAxML. The species name is indicated adjacent to the associated sequence cluster. The scale bar indicates the estimated number of substitutions per site. **b**, Isolation source of *Campylobacter* species with $n \geq 3$ isolates.

Figure 3. Core and accessory genome variation in the genus *Campylobacter*. **a**, Overall distribution of the total number of accessory genes (left) and core genes (right) per isolate for each *Campylobacter* species (where $n \geq 3$ isolates). The number of accessory genes is shown as boxplots (min to max). **b**, Venn diagram of pangenomes among different *Campylobacter* species ($n \geq 9$). The number of core genes shared by all species is illustrated in the center. **c**, Pairwise average nucleotide identity comparison calculated for all 631 *Campylobacter* isolates based upon 820 core genes shared by $>95\%$ of isolates. ANI values $<75\%$ are not calculated by FastANI [59]. **d**, Pairwise accessory genome similarity based upon gene presence or absence at 2,168 non-core loci. The heatmaps coloring ranges from yellow (minimum) to red (maximum). The matrices are ordered according to the phylogenetic tree presented in figure 2a. Different colours correspond to *Campylobacter* species with ≥ 3 isolates.

Figure 4. Elevated within-host interspecies recombination and donor-recipient comparisons. **a**, A hypothesis depicting the relationships between *Campylobacter* species, *C. jejuni* (x_1, x_2) and *C. coli* (y), when found in the same or in different hosts. **b**, Number of recombining SNPs within and between host as inferred by chromosome painting analysis for all donor recipient species comparisons. The error bar represents the standard error of the mean (SEM). **c**, The figure shows the number of donated SNPs in 10 donor-recipient pair species comparisons. The proportion (%) of recombining SNPs with $>90\%$ probability of copying from a donor to a recipient genome is illustrated in the y axis. All donor groups are shown in the x axis. All coloured boxes correspond to comparison where donors and recipients are found in the same host.

Figure 5. The mobilome of the *Campylobacter* genus. **a**, The graph illustrates the proportion of recombining genes in 10 different species comparisons. The number of species pairs in which the gene was found to recombine is shown on the x axis and the number of genes in each category is given on the y axis and. The exact number of genes found in each group comparison is shown on the top of each box. **b**, Number of *Campylobacter* species harbouring AMR genes that belong to efflux pumps and four different antibiotic classes which are shown on the x axis. **c**, The circos plot indicates the 16 genes involved in recombination in >5 donor-recipient pair species comparisons. Gene matches are indicated by joining lines, coloured differently for each gene. Gene names are shown around the perimeter for each *Campylobacter* species. **d**, The circos plot indicates the sharing of AMR genes

associated with efflux pumps and four antibiotic classes among *Campylobacter* species. Presence of at least one gene (not necessarily the same gene) conferring resistance to a specific antibiotic class is indicated by joining lines, coloured differently for each drug class. Efflux pumps (i), β -lactams (ii), tetracyclines (iii), aminoglycosides (iv) and lincosamides (v) are shown around the perimeter for each *Campylobacter* species.

Figure 2–figure supplement 1. Population structure of the *Campylobacteraceae* family.

Phylogenetic tree of 506 isolates that belong to the *Campylobacteraceae* family with *Helicobacter pylori* used as an outgroup. Different colors correspond to main species with number of isolates greater than three. The Tree was reconstructed using a gene-by-gene concatenated alignment of 799 core genes shared by >95% by all isolates and an approximation of the maximum-likelihood algorithm (ML) implemented in RAxML. The scale bar indicates the estimated number of substitutions per site.

Figure 2–figure supplement 2. Core genome species trees. Single-species trees for nine *Campylobacter* species with >4 isolates demonstrating the diversity for among species. The scale bars indicate the estimated number of substitutions per site. (*) The scale for the tree corresponding to *C. hepaticus* is 10 times smaller than the rest.

Figure 2–figure supplement 3. Overview of host-associations of *Campylobacter* species.

Abundance and diversity of 631 *Campylobacter* isolates in each host and environment. Different colours correspond to main species with number of isolates $\geq$ three. The number of isolates is shown on the y axis while the various isolation sources on the x axis.

Figure 2–figure supplement 4. Core genome species trees. Single-species trees for *C. jejuni*, *C. coli* and *C. fetus* species which contain isolates from multiple hosts and countries. The scale bars indicate the estimated number of substitutions per site.

Figure 3–figure supplement 1. Genome size variation of the *Campylobacter* genus. The frequency distribution of the genome size of all *Campylobacter* genomes used in this study is shown as a histogram. The number of genomes is shown on the y axis while the genome size (in bp) on the x axis.

Figure 3–figure supplement 2. Gene variation in the genus *Campylobacter*. Overall distribution of the total number of genes per isolate for each *Campylobacter* species (where $n \geq 3$ isolates). The number of genes is shown as boxplots (min to max).

Figure 3–figure supplement 3. Accessory gene function in all main *Campylobacter* species. The different gene functions are depicted on the y axis, while the number of shared accessory genes on the x axis. Different colours corresponding to different *Campylobacter* species.

Figure 3–figure supplement 4. Core genome allelic variation and the effect of recombination. **a**, Number of SNPs per genome of the main *Campylobacter* species (where $n \geq 3$ isolates) in the core genome alignment. The horizontal line in each plot represents the mean value while the upper and lower lines the standard deviation. **b**, Average nucleotide identity for pairwise comparisons of 820 core genes for 605 genomes of 15 main *Campylobacter* species. Different colours corresponding to different *Campylobacter* species.

Figure 4-figure supplement 1. Probability of the recipient genomes sharing DNA with each donor groups is illustrated as box whiskers (white) for every donor-recipient comparison for all 10 pairs that supported our hypothesis. The analysis where the host data were randomized across all isolates is illustrated as box whiskers (red). The probability of copying DNA from a donor to a recipient genome is shown on the y axis. The midline in the box whiskers indicates the mean and the error bars the standard deviation.

Figure 4-figure supplement 2. Genome position of genes containing recombining SNPs. Genes and their corresponding number of recombining SNPs, inferred by Chromosome Painting analysis for all 10 species comparisons, and mapped to the NCTC11168 reference genome. Genes from within-host (red) and between-host (white) pair comparisons are shown for each comparison. Donors are isolates from chicken (triangle), cattle (square), wild bird (cross), pig (star), clinical (circle) and water (snowflake) samples. The dashed line indicates the 95th percentile for every individual group comparison.

Figure 4-figure supplement 3. Genes ranked in ascending order of the number of recombining SNPs they contain as inferred by Chromosome Painting analysis for all ten species comparisons. Genes from within-host (red) and between-host (white) are shown for

each comparison. Donors are isolates from chicken (triangle), cattle (square), wild bird (cross), pig (star), clinical (circle) and water (snowflake) samples.

Figure 5-figure supplement 1. Presence of antimicrobial resistance genes in the *Campylobacter* genus. The phylogenetic tree was reconstructed using a gene-by-gene concatenated alignment of 820 core and soft-core genes and an approximation of the maximum-likelihood algorithm (ML) implemented in RAxML. The designated colour scheme was used for each species in the first column. The second column indicates whether the strain is isolated from an agricultural animal (grey). Remaining columns indicate presence of AMR genes (black). The scale represents the number of substitutions per site.

Figure 5-figure supplement 2. Genetic organization of AMR genes in *Campylobacter*. The presence of each AMR gene, highlighted in different colours, is shown for representative genomes from *C. jejuni*, *C. coli*, *C. lanienae*, *C. hyointestinalis* and *C. fetus subspecies fetus* sampled from different agricultural animals. The number of isolate genomes containing each genomic arrangement is indicated in parenthesis.

Supplementary File 1. Isolate information about the genomes used in this study.

Supplementary File 2. Within-host highly (>95th percentile) recombining genes.

Supplementary File 3. Recombination parameters as calculated by ClonalFrameML.

Supplementary File 4. Quantifying recombination between co-habiting species using ChromoPainter.

Supplementary File 5. Genes involved in interspecies recombination in 10 species comparisons.

Figure 3–source data 1. This file contains the numerical values on which the graphs in Figure 3 are based.

Figure 5–source data 1. This file contains the numerical values on which the graphs in Figure 5b-d are based.

ECOLOGICAL BARRIER

Host biogeography,
distribution, prevalence
behaviour and interactions.
Sub-niche segregation.

MECHANISTIC BARRIER

Factors promoting DNA
specificity including
homology dependence of
recombination, restriction-
modification systems.

ADAPTIVE BARRIER

Fitness of the recombinant
genotype in a given niche.

a **b**

a **b** **c**

***C. jejuni* (n=218)**

***C. coli* (n=143)**

***C. concisus* (n=106)**

***C. fetus* (n=52)**

***C. lanienae* (n=26)**

***C. hyointestinalis* (n=16)**

***C. lari* (n=13)**

C. hepaticus* (n=10)

***C. pinnipediorum* (n=9)**

C. jejuni

C. coli

C. fetus

a **b**

C. jejuni

C. coli

C. lanienae

C. hyointestinalis

C. fetus subs fetus