1 **Structural dynamics determine voltage and pH gating in**
2 **human voltage-gated proton channel**

3
4 Shuo Han, Sophia Peng, Joshua Vance, Kimberly Tran, Nhu Do, Nauy Bui, Zhenhua Gui and

5 Shizhen Wang

6 Department of Cell Biology and Biophysics, School of Biological and Chemical Sciences

7 University of Missouri-Kansas City, Kansas City MO 64110 USA

8
9 Correspondence should be addressed to SW (wangshizhen@umkc.edu)

10

Abstract

Voltage-gated ion channels are key players of electrical signaling in cells. As a unique subfamily, voltage-gated proton (Hv) channels are standalone voltage sensors without separate ion conductive pores. Hv channels are gated by both voltage and transmembrane proton gradient (i.e. ΔpH), serving as acid extruders in most cells. Amongst their many functions, Hv channels are known for regulating the intracellular pH of human spermatozoa and compensating for the charge and pH imbalances caused by NADPH oxidases in phagocytes. Like the canonical voltage sensors, Hv channels are a bundle of 4 helices (named S1 through S4), with the S4 segment carrying 3 positively charged Arg residues. Extensive structural and electrophysiological studies on voltage-gated ion channels, in general, agree on an outwards movement of the S4 segment upon activating voltage, but the real-time conformational transitions are still unattainable. With purified human voltage-gated proton (hHv1) channels reconstituted in liposomes, we have examined its conformational dynamics, including the S4 segment at different voltage and pHs using single-molecule fluorescence resonance energy transfer (smFRET). Here, we provide the first glimpse of real-time conformational trajectories of the hHv1 voltage sensor and show that both voltage and pH gradient shift the conformational dynamics of the S4 segment to control channel gating. Our results indicate that the S4 segment transits among 3 major conformational states and kinetic analysis suggest that only the transitions between the inward and outward conformations are highly dependent on voltage and pH changes. Our smFRET studies uncover the stochastic conformational dynamics of S4 and demonstrate how voltage and pH shift its conformational distributions to regulate channel gating. Altogether, we propose a kinetic model that explains the mechanisms underlying voltage and pH gating in Hv channels, which may also serve as a general framework for understanding the voltage sensing and gating in other voltage-gated ion channels.

Introduction

Voltage-gated cation channels mediate the electrical excitability of many cells, such as cardiomyocytes, muscle and nerve cells. As a unique subfamily, voltage-gated proton (Hv) channels are solely composed of voltage sensors without the separate pore-forming domains typically found in canonical voltage-gated cation channels (1, 2). Hv channel voltage sensors are hourglass-shaped with a bundle of 4 helices (named S1 through S4), like those of the voltage-gated cation channels (3, 4). The S4 segment of the voltage-gated cation channels is highly conserved and contains multiple positively charged residues to sense membrane voltage(5). High resolution structures of the voltage sensors from different organisms, including Hv channels, largely agree on a model where the charge carrying S4 segment is driven outward by depolarization voltage – a movement that is then transduced to gate the channel pore(6). Despite many years of research, the conformational trajectories of the voltage sensor, as well as the underlying mechanisms by which voltage and pH drive these conformational transitions in Hv channels, remain unclear and, sometimes even controversial. For example, accessibility assays showed that only the middle Arg in the S4 segment of the Hv channel switches its accessibility upon voltage activation(7), supported by the crystal structure of a voltage-sensitive phosphatase and the electron paramagnetic resonance data of the human Hv channel (hHv1)(8, 9). However, the gating currents of Hv channels and many other voltage-gated cation channels suggested approximately 3 Arg residues move across the membrane barrier upon gating transition(10, 11). Furthermore, a unique function-defining feature of Hv channels is that their voltage dependence is shifted by proton concentrations at the intra- and extracellular sides (i.e transmembrane pH gradient) (12). As a result, Hv channels are only activated when the proton electrochemical gradient is directed outwards(13). However, it is still unclear how pH modulates the voltage sensitivity of the Hv channels. Titratable residues were proposed to be involved in the pH regulation of Hv channels and H168 has been identified as a critical intracellular pH sensor in hHv1 channels (14, 15). In addition, the most recent studies indicated that both pH

gradient and voltage alter the dynamics of the Hv channel gating currents that reflect the movements of the S4 segment (16, 17).

To understand the structural basis underlying voltage and pH gating in the hHv1 channel, we utilized the single-molecule fluorescence resonance energy transfer (smFRET) approach that directly visualized the real-time conformational transitions and dynamics of hHv1 channels in lipid environments. In the present work, smFRET measurements were performed at the voltages and pH gradients that the gating status of hHv1 channels has been unambiguously defined. Our results indicate that both voltage and pH gradient modify the conformational dynamics of the S4 segment in hHv1 channels. Our studies provide not only a mechanistic rationale for the voltage-pH interplay in determining hHv1 channel gating but also structural insights into voltage sensing in other voltage-gated cation channels.

Results

1. Conformational dynamics of the hHv1 channels driven by voltages

The codon-optimized hHv1 WT channels with an N-terminal 6x histidine tag were expressed and purified from *E. coli* host cells, exhibiting a single, symmetric peak in size exclusion chromatography profile (Figure 1-figure supplement 1A). The hHv1 WT protein was reconstituted into liposomes (POPE/POPG=3/1, w/w, protein/lipid = 1/200, w/w) and its channel function was examined by liposome flux assay (Figure 1-figure supplement 1A). Our results indicated that the purified hHv1 WT protein retained its native proton channel function (Figure 1-figure supplement 1C).

To implement smFRET studies on the hHv1 channel, we mutated the two intrinsic Cys107 and Cys249 residues into Ser, then introduced cysteine mutations, one pair at a time, at different sites of the hHv1 channel. The goal was to uncover the relative movements among the 4 transmembrane segments at different pHs/voltages by measuring the real-time FRET changes between the donor and acceptor fluorophores conjugated with the cysteine residues

87 which were introduced at different sites. These movements reported by FRET were mapped
88 onto the available Hv channel structures (3, 4, 18) to reconstruct the conformational transitions
89 and their dynamic behaviors (Figure 1A). The hHv1 channel mutants were expressed, purified,
90 and then labeled by the chemically modified Cy3/Cy5 FRET fluorophore pair with improved
91 photostabilities (19). The fluorophore-labeled hHv1 channel proteins were reconstituted into
92 liposomes (POPE/POPG=3/1, w/w, 5 mg/ml) with an extremely low protein lipid ratio of 1:4,000
93 (w/w), so that most liposomes were either empty or only contained one hHv1 channel.
94 Moreover, hHv1 channels in liposomes exist mainly as monomers because the concentration of
95 hHv1 proteins during reconstitution is only 38 nM, while the dissociation constant of
96 homodimeric hHv1 proteins is $\sim 3.2 \mu\text{M}$ (8). The orientation of the hHv1 channel in
97 proteoliposomes was controlled by incubating the samples on the PEG passivated coverslip
98 surface coating with anti-His-tag antibodies (MA1-21315-BTIN, Invitrogen), so that only those
99 with the N-terminal His-tag of the hHv1 facing outside were retained for smFRET imaging. Thus,
100 the defined electrical potential was generated with K^+ gradients and K^+ ionophore valinomycin
101 and applied to the hHv1 channel in liposomes (Figure 1A). The smFRET imaging was
102 performed on a customized TIRF microscope with a dual-view beam splitter and EMCCD
103 camera as described previously(20, 21), and the raw movies collected without any corrections
104 were processed by the SPARTAN software to identify individual molecules, extract
105 donor/acceptor intensities and calculate FRET (22). The liposome voltages were calculated with
106 the Nernst equation and switched by changing the extraliposomal K^+ concentrations (Figure
107 1A). Our initial smFRET measurements did not detect any significant voltage-dependent FRET
108 changes at the voltage-sensing S4 segment (Figure 1-figure supplement 2A). We then realized
109 that the proton uptake through the hHv1 channel will decrease the extracellular pH of hHv1
110 channels (i.e the intraliposomal pH) and start to deactivate them in less than 1 min (Figure 1-
111 figure supplement 2B). As a result, the duration of the activated conformations was too short to
112 be captured. To circumvent this problem, we introduced the N214R mutation, which was shown

113 to significantly reduce the outward H^+ currents without altering the voltage dependence of hHv1
114 channels (10, 11). We confirmed that hHv1 N214R proteins reconstituted into liposomes exhibit
115 no detectable proton uptake within 30 min (Figure 1-figure supplement 2B, upper left panel).
116 The hHv1 channel proteins for smFRET studies contain multiple mutations with attached
117 fluorophores. Our liposome flux assay data indicated that, despite these modifications, most
118 hHv1 proteins without the N214R mutation remained functional to mediate proton uptake
119 (Figure 1-figure supplement 2B and C). On the N214R background, substantial voltage-
120 dependent FRET changes were detected at both the K169C-Q194C and K125C-S224C labeling
121 sites, which exhibited reverse FRET changes upon activating voltages (Figure 1A, Figure 1-
122 figure supplement 3A). SmFRET traces from the K169-Q194 labeling sites, exhibited large and
123 spontaneous transitions among three major FRET states with centers at 0.24, 0.52 and 0.78,
124 respectively, at both the resting (-85 mV) and activating voltages (Figure 1B, Figure 1-figure
125 supplement 4A). For all these traces, the donor and acceptor fluorescence intensities were
126 strongly anticorrelated, indicating that the FRET changes truly resulted from the structural
127 changes rather than fluorophore photophysics (Fig 1B, Figure 1-figure supplement 4A). We
128 fitted the histogram from the smFRET data collected at all voltages with 1-5 FRET states and
129 the SSE (sum of squares due to errors) of Gaussian fits decreases >95% with 3 FRET states
130 (Figure 1-figure supplement 3B). Therefore, we performed kinetic analysis using a model
131 containing low (F_L), medium (F_M), and high (F_H) FRET states. We also introduced the F_B FRET
132 state for some blinking or bleaching events that the SPARTAN software occasionally failed to
133 exclude (Figure 1F). However, the occupancies of the F_B state were extremely low and
134 transitions to the F_B state were rare, therefore they were ignored in the kinetic analyses follow-
135 up. The smFRET traces were then idealized using the Maximum Point Likelihood (MPL)
136 algorithm (23), which allowed us to fix the centers of FRET states and analyze multiple datasets
137 collected at different conditions. The FRET state occupancies calculated from several hundreds
138 of hHv1 channels at different voltages showed that the weak (0 mV) and strong activating (120

mV) voltages significantly enriched the low FRET-0.24 state, while diminishing the medium and high FRET state (Figure 1C). Shifts of the FRET state occupancies indicated that activating voltage altered the conformational distributions of the S4 segment by enriching those at the extracellular side. The results were further validated by the smFRET data collected from the K125C-S224C sites (Figure 1A), which also exhibited large, spontaneous FRET transitions among 3 FRET states with centers at 0.27, 0.6 and 0.9 (Figure 1D, Figure 1-figure supplement 4B). The activating voltages (0 and 120 mV), in sharp contrast, promoted the high FRET 0.9 state at the K125C-S224C labeling sites (Figure 1E). The voltage-dependent changes in FRET state occupancies were also reflected in the transition density plots, which showed that activating voltages promoted the transitions between the low FRET state at the K169C-Q194C sites and the high FRET state at the K125C-S224C sites (Figure 1G and H). Our smFRET data, from both the K169C-Q194C and K125C-S224C labeling sites, consistently indicated that the S4 segment of hHv1 undergoes spontaneous transitions at equilibrium state under all voltages, and the activating voltage stabilizes the S4 segment at the extracellular side.

2. Global conformational changes of the hHv1 channels at resting and activated voltages

To uncover the global conformational changes of hHv1 channels, we systematically examined the FRET distributions at multiple labeling sites, one pair at a time, at different voltages. The smFRET traces of all these labeling sites were analyzed with the 3+1 FRET state model using the MPL algorithm (Figure 1F), except for the K125C-K169C sites, which fit better with a 2+1 FRET state model. Among the labeling sites we examined, FRET state occupancies of the S98C-Q194C sites, which reflect the movement of S1 vs S4, demonstrated similar voltage-dependent changes as that of K169C-Q194C sites, although not statistically significant (Figure 2). However, at all other sites including the Q194C-S219C (S4 itself), S98C-N132C (S1 vs S2) and K125C-K169C (S1 vs S3), FRET state occupancies did not exhibit any significant shifts that could be potentially associated with voltage gating (Figure 2). These data suggested that voltage primarily acts on the voltage sensing S4 segment to promote its outward

conformations, while the S1, S2 and S3 segments probably do not undergo significant relative movements between each other during gating transition.

Our studies were performed on the hHv1 channels at equilibrium state, as shown by the FRET contour maps, the FRET distributions did not change over time (Figure 1-figure supplement 3A). However, the S4 segment appeared to be highly dynamic and can sample all conformational states, at all voltages from -85 to 120 mV. These structural transitions potentially underlie the stochastic gating transitions found in voltage-gated ion channels. At a strong resting voltage of -85 mV, hHv1 channels should mostly be stabilized at the closed state. At a strong activating voltage of 120 mV, however, hHv1 channels may not necessarily reach the full opening state. A previous electrophysiological study indicated that the maximum open probability recorded from hHv1 channels in human eosinophils, even at the optimal pH gradient, is ~ only 0.6 (24). Therefore, incomplete gating transitions may lead to the relatively small changes in FRET state occupancies observed in our studies.

3. pH gates hHv1 channels by modifying the dynamics of the S4 segment

To understand how pH shifts the voltage sensitivity of the hHv1 channel, smFRET measurements were performed on the K125C-S224C and K169C-Q194C sites at different pHs (Figure 3A). At 0 mV, lowering the intracellular pH (i.e the extraliposomal pH) to 6.5 remarkably enhanced the high FRET 0.9 state and decreased the low FRET 0.27 state occupancies (Figure 3A, left). Reverse changes in FRET occupancies were observed at the K169C-Q194C labeling sites, where the low FRET state (F0.24) increased and the high FRET state (F0.78) decreased upon intracellular pH 6.5 (Figure 3B, left panels). FRET occupancy data from both labeling sites support that lowering the intracellular pH enriches the outward conformations of the S4 segment, consistent with its effect to shift the voltage dependence of hHv1 channels negatively, therefore promoting openings (12). In sharp contrast, changing both the intra- and extracellular pHs symmetrically only shifted the FRET occupancies slightly (Figure 3A and B, middle panels), agreeing well with the electrophysiological results that show the transmembrane pH gradient

shifts the voltage dependence of hHv1 channels (12). To further confirm our results, we introduced the H168Q background mutation, which was previously reported to attenuate the sensitivity of hHv1 channels to intracellular pH (15). The results indicated that FRET distributions at both labeling sites no longer responded to intracellular pH changes (Figure 3A and B, right panels). Our data showed that pH gating of the hHv1 channels originates from its effect on the conformational dynamics of the S4 segment.

To elucidate the pH voltage interplay in determining hHv1 channel gating, we examined the changes in FRET distributions induced by voltages at intracellular pH 8.5, which shifts the voltage dependence of hHv1 channels positively. Consistently, voltage-dependent enrichments of the high FRET 0.9 state at the K125C-SS24C sites were almost abolished (Figure 4A, right and middle panels). The potential of activating voltage in enriching the low FRET 0.24 state was also significantly attenuated at the K169C-Q194C sites (Figure 4B, right and middle panels). As shown in Figure 3, the H168Q mutation not only abolished the dependence of FRET distributions on intracellular pH but also enriched the high FRET 0.9 state at the K125C-S224C sites and the low FRET 0.24 state at the K169C labeling sites, at all intracellular pHs tested. These data propose that the H168Q mutation may stabilize the outward conformations of the S4 segment, therefore more negative voltage may be required to drive the S4 segment towards the resting state. Indeed, we found that the enrichment of the low FRET 0.27 populations correlating with channel closure was only observed on the K125C-S224C labeling sites at a more negative voltage (-200 mV) on the H168Q background, unlike those on the WT background that can be observed on the less negative voltage of -85 mV (Figure 4A, left and right panels). These data suggest that the H168 residue may be involved in sensing intracellular pH through direct or indirect interactions with the S4 segment by stabilizing its inward resting conformations. However, only very minor effects were observed on the K169C-Q194C labeling site, perhaps due to the local impacts by the K169C mutations and conjugated fluorophores (Figure 4B, right panel).

4. The structural transitions are dependent on the voltage/pH

To provide mechanistic insights into the voltage and pH gating in hHv1 channels, we performed kinetic analysis using the 3+1 state model as described previously with the MPL algorithm (Figure 1F). The smFRET traces at different voltage/pH conditions were idealized and the rate constants of transitions among the low, medium, and FRET states were obtained. We then calculated the equilibrium constants (K_{eq}) of different transitions as ratios of the forward (K_F) to reverse (K_R) transition rates. At both labeling sites, under the same pH, only the transitions between the low and high FRET states (i.e LH) exhibited voltage-dependent changes (Figure 5A). Activating voltage drove the equilibrium towards the high FRET state at the K125C-S224C sites and towards the low FRET state at the K169C-Q194C labeling sites (Figure 5A). At 0 mV and extracellular pH 7.5, again only the equilibrium of the LH transitions was remarkably shifted by intracellular pH, lowering it to 6.5 enriched the high FRET state at the K125C-S224C sites and the low FRET state at the K169C-Q194C labeling sites (Figure 5A). These data match well with functional results, i.e lowering intracellular pH promotes hHv1 channel openings. We further showed that the H168Q mutation impaired the sensitivity of hHv1 channels to intracellular pH by diminishing the pH-dependent changes in K_{eq} of LH transitions. Based on these kinetic data, we propose a kinetic model that explained voltage pH gating in hHv1 channels (Figure 5B). Based on our smFRET data, the S4 segment may transit among 3 positions across the membrane, i.e the intracellular side (I), middle (M), and extracellular side (E). The I conformation has 2 populations with their equilibrium depending on the protonation states of the intracellular pH sensor H168 and extracellular pH sensor that is yet to be identified (Figure 5B). Activating voltage mainly shifts the I-E transition by driving the S4 segment from the intracellular side (I conformation) directly to the extracellular side (E conformation), thereby promoting channel openings.

In summary, our results suggest that both pH and voltage act on the S4 segment of the hHv1 voltage sensor to modulate channel gating, which is consistent with the most recent gating

current recordings(17). In hHv1 channels, the mobilizing voltage of the S4 segment is probably much more negative, like those from canonical voltage-gated ion channels. However, the pH sensors at the intra (H168) and the extracellular sides may physically interact with the S4 segment, directly or indirectly, to stabilize it at the inward resting conformations. The molecular couplings between the S4 segment and pH sensors are dependent upon the protonation state of the H168 and other pH sensing residues. Thus, the stability of the inward resting conformation (i.e I) is tuned by pH gradient, which in turn shifts the equilibrium of LH transition that is driven by voltages to open or close hHv1 channels.

Discussion

Structural alignments of the voltage sensors from different organisms or even the same channels largely agree on a model where the positively charged S4 segment is driven outwards by depolarization – a movement utilized by pore modules to control ion permeation(6, 9, 25). However, the details of this movement are far from clear and, different movements have been described for different voltage sensors. For example, the S4 segment moves up for 3 helical turns in a voltage-gated sodium channel NavAb (25) while it moves up for only 1 helical turn in a voltage sensitive phosphatase(9). Unfortunately, most atomic structures of voltage-gated ion channels were obtained at 0 mV voltage with their gating states unclear, some even at the conformations that may not be physiologically relevant (26). Thus, the potential conformational changes of the voltage sensors inferred from the structural alignments remain to be functionally defined. We performed quantitative analysis on the real-time conformational transitions and dynamics of the hHv1 voltage sensor in lipid environments, at the voltages and pHs where the hHv1 channel functional status has been clearly defined. Our results showed that both voltage and pH act on the S4 segment in hHv1 channels to regulate channel gating. With smFRET imaging, we revealed the spontaneous, large conformational transitions of the S4 segment in hHv1 channels, which perhaps are the structural basis of the gating current noise (27). The

gating currents of Hv channels suggested multiple closed states (10, 28). Consistently, our data showed that the S4 segment in hHv1 channels transits among at least 3 conformations and it seems that only the transitions between the inward and outward conformations correlate with voltage and pH gating. Although hHv1 channels at deactivated and activated states are NOT available, long-time simulations have provided models of 'deactivated' and 'activated' conformations (18). FRET values were predicted from these hHv1 structural models using the FRET Positioning and Screening (FPS) software and then compared with our data (29). From the 'deactivated' to 'activated' conformations, FPS software predicted the FRET changes from 0.28 to 0.57 at the K125C-S224C labeling sites, and from 0.72 to 0.45 at the K169C-Q194C labeling sites. The predicted FRET matched well with 2 of the 3 FRET centers identified by our smFRET studies, i.e 0.27 and 0.6 at the K125C-S224C sites, and 0.78 and 0.52 at the K169C-Q194C sites. Interestingly, the unmatched FRET 0.9 state of K125C-S224C and FRET 0.24 state of K169C-Q194C sites, represented the same activated conformation (i.e the E conformation in Fig. 5b). Our results imply that the simulation study perhaps did not reach the fully activated conformations.

A previous study by Li et al showed that the oligomeric state of purified hHv1 channels is concentration-dependent with an apparent dissociation constant of 3.2 μ M (8). For smFRET samples, the hHv1 channels were reconstituted at an extremely low protein lipid ratio of 1:4,000 (w/w) therefore the hHv1 concentration in liposome reconstitution was only ~38 nM, almost 100x lower than its dissociation constant. As a result, most hHv1 channels imaged were monomers and very few homodimers were excluded because they contained multiple donor and acceptor fluorophores, therefore exhibiting multiple bleaching steps. Electrophysiological results indicated that both monomeric and homodimeric hHv1 channels are functional, but the dimeric hHv1 channel exhibits strong intersubunit cooperativity by showing sigmoidal gating dynamics (30). Therefore, it would be very interesting to further examine how subunit interactions alter the structural dynamics of hHv1 channels and, thus their gating kinetics. In the present work, we

introduced the N214R background mutation to abolish the proton uptake into liposomes driven by positive voltages (Fig 1 supplement figure 2B) that could change the extracellular pH environment of hHv1 channels. However, due to the technical challenge, we were not able to directly measure or even better, monitor the inside pHs and voltages of the liposomes that contain each individual hHv1 molecule being imaged. This limitation may impact the smFRET data collected at 0 mV with intra- and extraliposomal pH of 7.5 and 8.5 (Fig 3A and B, left panels, Fig 4A and B, middle panels). Under the above voltage/pH conditions, protons may flow out of the liposomes following the pH gradient through hHv1 channels that may still be activated, since the N214R mutation does not abolish the inward proton currents (10, 11). The increase in extracellular pH of hHv1 channels due to proton efflux could attenuate, but probably does not cancel the effects of intracellular pH 8.5, since enrichments of the FRET states representing resting conformations were still observed under the pH voltage condition (Fig 3A and B, left panels, Fig 4A and B, middle panels).

In conclusion, the unique voltage and pH gating in the hHv1 channel is based on the spontaneous, voltage-dependent structural transitions of the S4 segment among multiple conformations. The intra- and extracellular pHs modify the stability of inward resting conformations by altering the protonation states of the pH sensing residues that may directly or indirectly interact with the S4 segment in hHv1 channels. The gating model we proposed also explains the most recent patch-clamp fluorometry data and gating current from Hv channels (Figure 5B), which showed that the conformation transitions of the S4 segment are dependent on both intra- and extracellular pHs (17, 31).

Abbreviations:

POPE, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine; POPG, 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-(1'-rac-glycerol); TCEP, tris(2-carboxyethyl)phosphine; ACMA, 9-Amino-6-Chloro-2-Methoxyacridine; CCCP, carbonyl cyanide m-chlorophenyl hydrazine; 2-GBI, 2-

guanidinobenzimidazole; NADPH, reduced nicotinamide adenine dinucleotide phosphate; NMDG, N-Methyl-D-glucamine; ROS, reactive oxygen species; PCD, protococatechuate 3,4-Dioxygenase; PCA, protocatechuic acid; COT, cyclooctatetraene; Hepes, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; Tris, 2-Amino-2-(hydroxymethyl)propane-1,3-diol; EMCCD, electron multiplying charge coupled device.

Acknowledgments

This work was funded by NIH grant 1R15GM137215-01 (SW) and the startup fund of UMKC. We would like to thank Decker Gates for his help on protein preparation and data analysis; Dr. Lejla Zubcevic at the Kansas University Medical Center and Dr. Xiaolan Yao at the University of Missouri-Kansas City for their help in reviewing and revising the manuscript.

METHODS

1. Protein expression, purification and fluorophore labeling

The hHv1 cDNA was codon-optimized and synthesized by Genscript Inc, then inserted into the pET28a(+) vector between the NdeI and BamHI restriction sites. The resulting hHv1 protein contained a 6x Histidine tag at N- terminus, followed by a thrombin protease cutting site. All mutations were introduced into hHv1 by site-direct mutagenesis (Agilent Inc.) and confirmed by DNA sequencing. The hHv1 proteins were expressed in *E. coli* BL21 (DE3) host cells and purified as that described by Li et al(8). In brief, the hHv1 proteins were purified by metal affinity chromatography and then loaded onto a Superdex-200 size-exclusion column with a running buffer containing 20 mM Tris, 150 mM NaCl, 1 mM Fos-Choline 12 and 1 mM TCEP, pH8.0. For smFRET imaging samples, the proteins were subjected for buffer exchange using a 5 mL Hi-Trap desalting column to labeling buffer containing 20 mM Hepes, 150 NaCl, 1 mM Fos-Choline 12 (n-Dodecylphosphocholine), pH 7.0, and then mixed with Cy3 and Cy5 c5 maleimide(19) in a 1:1 mixture at a protein fluorophore molar ratio of 1:6. The labeling reactions were performed under 4 °C for 3 hours, then the free fluorophores were completely removed by performing a metal affinity chromatography and a size exclusion chromatography in a row. All the proteins were either stored at -80 °C for later use or reconstituted immediately.

2. hHv1 liposome reconstitution

The lipids dissolved in chloroform containing POPE/POPG (3:1, w/w, Avanti Polar Lipids Inc.) were dried in clean glass tubes with argon gas first, and then in a vacuum desiccator for 4 hours to evaporate the organic solvent completely. The dried lipids were resuspended in reconstitution buffer containing 20 mM Hepes, 5 mM KCl, 150 mM NaCl, pH7.5 (for smFRET imaging) or 20 mM Hepes, 150 mM KCl, 0.05 mM NaCl, pH7.5 (for liposome flux assay). The liposomes were formed by sonication and then destabilized with 10 mM Fos-choline 12 (Anatrace Inc). For liposome flux assay, the purified protein was mixed with the lipid solution at a protein lipid ratio of 1:200 (w/w), and the hHv1 proteoliposomes were formed by detergent

removal using Bio-beads SM2 (Bioad Inc). For smFRET imaging, the proteins were mixed with lipids at a ratio of 1:4000 (w/w) and the detergents were removed by dialyzing against reconstitution buffer containing 1 mM TCEP at the volume ratio of 1:500 for at least 3 times, each time over 12 hours.

3. Liposome fluorescence flux assay

The K⁺ gradient between inside and outside of the liposomes was established by diluting the liposomes in the extraliposomal buffer containing 20 mM Hepes, 150 mM NMDG, pH7.5. The liposomes were incubated with 2 μM of ACMA fluorescence probes for ~5 min, then ACMA fluorescence was measured using a 96 well plate reader (FluoStar, Ex/Em = 390nm/460nm) for ~5 min. After valinomycin was added at a final concentration of 0.45 μM, the fluorescence measurements were resumed with the same optical setting for ~40 min. Proton-specific ionophore CCCP was used as the positive control, and empty liposomes without hHv1 channels were used as negative controls. The fluorescence liposome flux data were processed following the method of Su et al (32). In brief, the hHv1 channel activities were calculated from the fluorescence readings normalized between 0 and 1 using the following equation:

$$A = (F_0 - F_{\text{val}})/(F_0 - F_{\text{cccp}})$$

where F_0 , F_{val} and F_{cccp} were the steady-state ACMA fluorescence at initial, after adding valinomycin and CCCP, respectively. The relative activities of the hHv1 channel were normalized against the hHv1 WT proteoliposomes included in every batch of assays.

4. Single-molecule imaging and data analysis

Flow chambers for smFRET imaging were prepared following the protocol of Joo et al (33). An objective-based TIRF built on a Nikon TE-2000U inverted microscope (TE-2000s) with 100x APO TIRF NA1.49 objective lens, 532 nm and 640 nm lasers, was used for single-molecule imaging. Donor and acceptor emissions were separated by W-view Gemini beam splitter with chromatic aberration correction (Hamamatsu Inc.) carrying the 638 nm long-pass beam splitter, then cleaned by 585/65 nm and 700/75 nm bandwidth filters (Chroma Inc.). The images were

collected by an ImagEM X2 EMCCD camera (Hamamatsu Inc.). The liposomes containing fluorophore-labeled hHv1 channels were retained on the PEGylated surface coated with biotinylated Histag antibodies (1:200 dilution, ThermoFisher). Fluorophores were excited by a 532 nm laser ($\sim 1.0 \text{ W/cm}^2$) and time-lapse movies were collected at 10 frames per second (i.e., time resolution of 100 ms). The 640 nm laser ($\sim 1.0 \text{ W/cm}^2$) was only used to confirm the existence of acceptor fluorophores when overall FRET was very low. Typical recording time was ~ 3 min, with half bleaching time being ~ 1 min. All imaging buffers contained ~ 3 mM Trolox, 5 mM PCA, and $15 \mu\text{g}/\mu\text{L}$ of PCD to enhance the photostability of the fluorophores (34, 35). For symmetrical pH conditions, β -escin at a final concentration of $50 \mu\text{M}$ was used to permeabilize liposomes(36). At least 3 batches of independent smFRET imaging were performed for each sample/condition. The movies were imported into the SPARTAN software directly without any corrections (22). The donor and acceptor channels were aligned using 2D maps generated from TetraSpeck fluorescent beads (T7279, Invitrogen). The molecules were identified as point-spread functions using the threshold method with a window size of 7 pixels. The smFRET traces were extracted and then preselected using the Autotrace function of the SPARTAN software with criteria setting as FRET Lifetime > 50 frames, donor/acceptor correlation coefficient between -1.1 and 0.5 , Signal-to-Noise ratio > 8 , background noise level < 70 , Cy3 blinks < 4 and overlap molecules removed (22). The resulting traces were further picked manually following the criteria described in the previous studies (37, 38). The FRET efficiency E was calculated with the following equation:

$$E = (I_A - I_D \cdot 0.07) / (I_A + I_D)$$

Where I_A and I_D are intensities of the donor and acceptor fluorophores. The cross-talk value is 0.07 , determined from a protein sample containing the donor fluorophore only.

Overall, there were ~ 500 molecules per field of view with ~ 100 molecules containing both acceptor and donor fluorophores. The bin size of all histograms and contour maps was 0.03 and FRET contour plots were generated from the smFRET data of the first 5s . For kinetic analysis, a

410 kinetic model containing 4 FRET states was used, with one state assigned to bleaching or
411 blinking events as FRET close to 0 (Figure 1F). The FRET peak centers were fixed for all
412 smFRET traces from the same labeling sites, with the K125C-S224C sites as 0.27, 0.6 and 0.9
413 and the K169C-Q194C site as 0.24, 0.52, and 0.78. These peak centers were determined by
414 Gaussian fit to the FRET histograms from all smFRET traces of the same labeling sites (Figure
415 1-figure supplement 3B). The smFRET traces were idealized using the Maximum Point
416 Likelihood (MPL) algorithm included in the SPARTAN software, which allows model constraints
417 and directly optimizes rate constants obtained at different experimental conditions (23).

Figures

Figure 1. The voltage-dependent conformation dynamics of the hHv1 channel S4 segment revealed by smFRET

A. Left panel, the cartoon of the hHv1 NMR structure (5oqk). The K125-S224 (blue) and K169-Q194C (red) labeling sites were highlighted by spheres; Right panel, the sample immobilization configuration for smFRET imaging. The PEG passivated surface with 2% PEG-biotin (Btn-Peg) binds the biotinylated anti-Histag antibodies (Btn-HisAb) via neutravidin (NA), then the hHv1 liposomes with N-terminal 6*Histag (6xHis) facing the outside were selectively retained for smFRET imaging. The liposome voltages were controlled by the K^+ gradient across liposomes in the presence of K^+ ionophore valinomycin (Val). B, D. Representative smFRET traces between the K169C-Q194C (B) and K125C-224C (D) labeling sites under -85 mV (resting), 0 mV (weak) and 120 mV (strong activating) voltages. The green and pink lines are donor and acceptor fluorescence intensities; the blue and red lines are the real and idealized FRET. C, E. FRET histograms and state occupancies of the smFRET data from the K169C-Q194C (C) and K125C-224C (E) labeling sites under different voltages. State occupancy data presented as mean $\pm$ s.e, N is the number of smFRET traces. Unpaired t-tests were performed to examine significance levels of voltage-induced FRET state occupancy changes, in comparison to the occupancy of the same FRET state at -85 mV, with * and ** indicating $p<0.05$ and $p<0.01$. F. The structure model used for kinetic analysis of smFRET traces, contains 4 FRET states, including low (F_L), medium (F_M), high (F_H), and bleaching/blinking states (F_B). However, the occupancies of the F_B state are

extremely low and the transitions to the F_B state are very rare at all labeling sites and voltage/pH conditions, therefore the F_B state was ignored in the following kinetic analyses. G, H. Transition density plots from the smFRET data at the K169C-Q194C (G) and K125C-224C (H) labeling sites under different voltages.

Figure 2. Global conformational changes of hHv1 channels driven by voltages

FRET histograms and state occupancies were calculated from the smFRET data at the S98C-Q194C, Q194C-S219C, S98C-N132C, and K125C-K169C labeling sites on the N214R background. All data were analyzed with the 4 state kinetic model as shown in Fig 1F, except for the K125C-K169C sites with a 3 state kinetic model containing $F_{0.66}$, $F_{0.4}$ and the F_B state for bleaching/blinking events. The state occupancy data presented as mean $\pm$ s.e, N is the number of smFRET traces. Unpaired t-tests were performed to examine the significance levels of voltage-induced FRET state occupancy changes, in comparison to the occupancy of the same FRET state at -85 mV, with * and ** indicating $p<0.05$ and $p<0.01$.

Figure 3. The structural dynamics of the S4 segment are pH-dependent

FRET histograms and state occupancies of the smFRET data from the K125C-S224C (A), K169C-Q194C (B) of the hHv1 channels on the N214R mutation background, in liposomes under 0 mV with asymmetrical (left and right panels), symmetrical (middle) pHs on the WT (left, middle panels), or H168Q mutation backgrounds. State occupancy data presented as mean $\pm$ s.e, N is the number of smFRET traces. Unpaired t-tests were performed to examine the significance levels of pH-induced FRET state occupancy

changes, in comparison to the occupancy of the same FRET state at pH 6.5, with * and ** indicating $p < 0.05$ and $p < 0.01$.

Figure 4. The structural dynamics of the S4 segment are determined by voltage pH interplay

FRET histograms and state occupancies of the smFRET data from the K125C-S224C (A) and K169C-Q194C (B) labeling sites of the hHv1 channels with N214R mutation at different voltage/pH conditions on the WT (left, middle panels) and H168Q mutation backgrounds (right panels). State occupancy data presented as $\text{mean} \pm \text{s.e.}$, N is the number of smFRET traces. Unpaired t-tests were performed to examine the significance levels of voltage-induced FRET state occupancy changes, in comparison to the occupancy of the same FRET state at -85 mV or -200 mV on the H168Q mutation background, with * and ** indicating $p < 0.05$ and $p < 0.01$.

Figure 5. The structural basis underlying voltage pH gating of hHv1 channels

A. Voltage and pH-dependent changes in equilibrium constants (K_{eq}), as the ratio of forward (K_F) and reverse (K_R) transition rates. The transition rates were calculated from the smFRET traces at different voltage/pH conditions idealized using the MPL algorithm with the 4 FRET state kinetic model. B. The structural model to explain the voltage and pH gating in hHv1 channels. The S4 segment of hHv1 channels has 3 distinct conformations according to their positions in membranes, i.e the intracellular (I), middle (M) and extracellular (E) conformations. The I conformation has 2 populations with one interacting with the intracellular (H168) and/or extracellular pH sensors (?), directly or indirectly, depending on their protonation states. Decoupling of the S4 segment from pH

488 sensors allows the spontaneous transitions towards the M and E conformations, with
489 activating voltage mainly shifting the equilibrium of I-E transitions.

Supplement Figures

Figure 1- figure supplement 1. The purified hHv1 protein functions as a proton channel

A. SDS-PAGE and size exclusion chromatography profiles of the hHv1 channel WT protein expressed and purified from *E. coli* host cells. B. The liposome fluorescence flux assay to determine channel activities of the hHv1 channel. The liposome voltage is generated by the K^+ gradient across the liposomes in the presence of K^+ ionophore valinomycin (Val). Proton uptake through the hHv1 channel into liposomes quenches the ACMA fluorescences. C. Liposome flux assay of the proton uptake mediated by hHv1 channel reconstituted into liposomes (POPE/POPG = 3/1, protein/lipid = 1/200, w/w). Data presented as mean $\pm$ s.e, n=3. The arrows mark the time points when 0.45 μ M valinomycin (blue) and 1 μ M CCCP (green) were added.

Figure 1- figure supplement 2. The N214R mutation abolishes hHv1 channel proton inhibition by preventing proton uptake into the liposomes

A. Contour maps of the smFRET data between the hHv1 K125C-S224C sites on the WT background at different voltages. B. The liposome flux assays of hHv1 protein samples for smFRET studies. The fluorophore-labeled protein samples for smFRET studies, except for WT and N214R mutant, were reconstituted into liposomes (POPE/POPG = 3/1) at a protein lipid ratio of 1:200 (w/w). The arrows mark the time points when the 0.45 μ M valinomycin (blue) and 1 μ M CCCP (green) were added. Data presented as mean $\pm$ s.e, n=3. C. Relative channel activities of the fluorophore-labeled hHv1 mutant proteins subjected to smFRET studies, without the N214R mutation,

determined by liposome flux assay. All liposome flux assay data were normalized by the WT hHv1 liposome samples of the same batch. Data presented as mean $\pm$ s.e, n=3.

Figure 1- figure supplement 3. The voltage induced conformational changes of the hHv1 channels revealed by smFRET

A. FRET contour maps of the K169C-Q194C and K125C-S224C labeling sites on the N214R mutation background at different voltages. N is the number of traces. B. SSE (sum of squared errors) for 1~5 Gaussian fits to FRET histograms of the K169C-Q194C and K125C-S224C labeling sites.

Figure 1- figure supplement 4. Representative smFRET traces of the K169C-Q194C (A) and K125C-S224C (B) labeling sites on the N214R mutation background at different voltages

The green and pink lines are donor and acceptor fluorescence intensities, the blue and red lines are the real and idealized FRET.

- 530 1. I. S. Ramsey, M. M. Moran, J. A. Chong, D. E. Clapham, A voltage-gated proton-
531 selective channel lacking the pore domain. *Nature* **440**, 1213-1216 (2006).
- 532 2. M. Sasaki, M. Takagi, Y. Okamura, A voltage sensor-domain protein is a voltage-gated
533 proton channel. *Science* **312**, 589-592 (2006).
- 534 3. K. Takeshita *et al.*, X-ray crystal structure of voltage-gated proton channel. *Nat Struct*
535 *Mol Biol* **21**, 352-357 (2014).
- 536 4. M. Bayrhuber *et al.*, Nuclear Magnetic Resonance Solution Structure and Functional
537 Behavior of the Human Proton Channel. *Biochemistry* **58**, 4017-4027 (2019).
- 538 5. C. Gonzalez, H. P. Koch, B. M. Drum, H. P. Larsson, Strong cooperativity between
539 subunits in voltage-gated proton channels. *Nat Struct Mol Biol* **17**, 51-56 (2010).
- 540 6. J. Guo *et al.*, Structure of the voltage-gated two-pore channel TPC1 from Arabidopsis
541 thaliana. *Nature* **531**, 196-201 (2016).
- 542 7. C. Gonzalez, S. Rebolledo, M. E. Perez, H. P. Larsson, Molecular mechanism of voltage
543 sensing in voltage-gated proton channels. *J Gen Physiol* **141**, 275-285 (2013).
- 544 8. Q. Li *et al.*, Resting state of the human proton channel dimer in a lipid bilayer. *Proc Natl*
545 *Acad Sci U S A* **112**, E5926-5935 (2015).
- 546 9. Q. Li *et al.*, Structural mechanism of voltage-dependent gating in an isolated voltage-
547 sensing domain. *Nat Struct Mol Biol* **21**, 244-252 (2014).
- 548 10. E. M. Carmona *et al.*, Gating charge displacement in a monomeric voltage-gated proton
549 (Hv1) channel. *Proc Natl Acad Sci U S A* **115**, 9240-9245 (2018).
- 550 11. V. De La Rosa, I. S. Ramsey, Gating Currents in the Hv1 Proton Channel. *Biophys J*
551 **114**, 2844-2854 (2018).
- 552 12. V. V. Cherny, V. S. Markin, T. E. DeCoursey, The voltage-activated hydrogen ion
553 conductance in rat alveolar epithelial cells is determined by the pH gradient. *J Gen*
554 *Physiol* **105**, 861-896 (1995).
- 555 13. T. E. DeCoursey, Voltage-gated proton channels: molecular biology, physiology, and
556 pathophysiology of the H(V) family. *Physiol Rev* **93**, 599-652 (2013).
- 557 14. V. V. Cherny *et al.*, Tryptophan 207 is crucial to the unique properties of the human
558 voltage-gated proton channel, hHV1. *J Gen Physiol* **146**, 343-356 (2015).
- 559 15. V. V. Cherny, D. Morgan, S. Thomas, S. M. E. Smith, T. E. DeCoursey, Histidine(168) is
560 crucial for DeltapH-dependent gating of the human voltage-gated proton channel, hHV1.
561 *J Gen Physiol* **150**, 851-862 (2018).
- 562 16. T. E. DeCoursey, Voltage and pH sensing by the voltage-gated proton channel, HV1. *J*
563 *R Soc Interface* **15**, (2018).
- 564 17. E. M. Carmona *et al.*, The voltage sensor is responsible for DeltapH dependence in Hv1
565 channels. *Proc Natl Acad Sci U S A* **118**, (2021).
- 566 18. A. D. Geragotelis *et al.*, Voltage-dependent structural models of the human Hv1 proton
567 channel from long-timescale molecular dynamics simulations. *Proc Natl Acad Sci U S A*
568 **117**, 13490-13498 (2020).
- 569 19. Q. Zheng *et al.*, Electronic tuning of self-healing fluorophores for live-cell and single-
570 molecule imaging. *Chem Sci* **8**, 755-762 (2017).
- 571 20. S. Wang *et al.*, Potassium channel selectivity filter dynamics revealed by single-molecule
572 FRET. *Nat Chem Biol* **15**, 377-383 (2019).
- 573 21. S. Wang, R. Vafabakhsh, W. F. Borschel, T. Ha, C. G. Nichols, Structural dynamics of
574 potassium-channel gating revealed by single-molecule FRET. *Nat Struct Mol Biol* **23**, 31-
575 36 (2016).
- 576 22. M. F. Juetten *et al.*, Single-molecule imaging of non-equilibrium molecular ensembles on
577 the millisecond timescale. *Nat Methods* **13**, 341-344 (2016).

23. F. Qin, A. Auerbach, F. Sachs, A direct optimization approach to hidden Markov modeling for single channel kinetics. *Biophys J* **79**, 1915-1927 (2000).
24. V. V. Cherny, R. Murphy, V. Sokolov, R. A. Levis, T. E. DeCoursey, Properties of single voltage-gated proton channels in human eosinophils estimated by noise analysis and by direct measurement. *J Gen Physiol* **121**, 615-628 (2003).
25. G. Wisedchaisri *et al.*, Resting-State Structure and Gating Mechanism of a Voltage-Gated Sodium Channel. *Cell* **178**, 993-1003 e1012 (2019).
26. C. A. Villalba-Galea, W. Sandtner, D. M. Starace, F. Bezanilla, S4-based voltage sensors have three major conformations. *Proc Natl Acad Sci U S A* **105**, 17600-17607 (2008).
27. D. Sigg, E. Stefani, F. Bezanilla, Gating current noise produced by elementary transitions in Shaker potassium channels. *Science* **264**, 578-582 (1994).
28. T. E. DeCoursey, Gating currents indicate complex gating of voltage-gated proton channels. *Proc Natl Acad Sci U S A* **115**, 9057-9059 (2018).
29. S. Kalinin *et al.*, A toolkit and benchmark study for FRET-restrained high-precision structural modeling. *Nat Methods* **9**, 1218-1225 (2012).
30. S. M. Smith, T. E. DeCoursey, Consequences of dimerization of the voltage-gated proton channel. *Prog Mol Biol Transl Sci* **117**, 335-360 (2013).
31. T. M. Schladt, T. K. Berger, Voltage and pH difference across the membrane control the S4 voltage-sensor motion of the Hv1 proton channel. *Sci Rep* **10**, 21293 (2020).
32. Z. Su, E. C. Brown, W. Wang, R. MacKinnon, Novel cell-free high-throughput screening method for pharmacological tools targeting K⁺ channels. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 5748-5753 (2016).
33. C. Joo, T. Ha, Preparing sample chambers for single-molecule FRET. *Cold Spring Harbor protocols* **2012**, 1104-1108 (2012).
34. C. E. Aitken, R. A. Marshall, J. D. Puglisi, An oxygen scavenging system for improvement of dye stability in single-molecule fluorescence experiments. *Biophysical journal* **94**, 1826-1835 (2008).
35. R. Dave, D. S. Terry, J. B. Munro, S. C. Blanchard, Mitigating unwanted photophysical processes for improved single-molecule fluorescence imaging. *Biophysical journal* **96**, 2371-2381 (2009).
36. J. S. Fan, P. Palade, Perforated patch recording with beta-escin. *Pflügers Archiv : European journal of physiology* **436**, 1021-1023 (1998).
37. S. Wang, R. Vafabakhsh, W. F. Borschel, T. Ha, C. G. Nichols, Structural dynamics of potassium-channel gating revealed by single-molecule FRET. *Nature structural & molecular biology* **23**, 31-36 (2016).
38. S. Wang, J. B. Brettmann, C. G. Nichols, in *Potassium Channels*. (Springer, 2018), pp. 163-180.