1 ***Neural correlates and determinants of approach-avoidance conflict***
2 ***in the prelimbic prefrontal cortex***

3
4 Fernandez-Leon, J.A.^{#1,2}; Engelke, D.S.^{#1}; Aquino-Miranda, G.^{#1}; Goodson, A.^{1,3};
5 Rasheed, M.N.¹; Do-Monte, F.H.^{1,3*}.
6
7
8
9

10 ¹ Dept. of Neurobiology & Anatomy, The University of Texas Health Science Center,
11 Houston, TX77030, USA;

12
13 ² Current Affiliation: Exact Sciences Faculty, CIFICEN (UNCPBA-CONICET-CICPBA)
14 & INTIA (UNCPBA-CICPBA), Tandil 7000, Buenos Aires, Argentina;

15
16 ³ Rice University, Houston, TX77030, USA
17

18 [#] These authors contributed equally to this work
19

20 *Corresponding author:

21 Fabricio H. Do Monte, DVM, PhD.
22 Department of Neurobiology and Anatomy, McGovern Medical School,
23 The University of Texas Health Science Center at Houston
24 6431 Fannin Street, Room 7.166, Houston, Texas 77030
25 Phone: 713-500-5613
26 Email: fabricio.h.domonte@uth.tmc.edu
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **Keywords:** Prelimbic prefrontal cortex, electrophysiology, optogenetics, fear, reward,
42 decision-making.
43
44

ABSTRACT

The recollection of environmental cues associated with threat or reward allows animals to select the most appropriate behavioral responses. Neurons in the prelimbic cortex (PL) respond to both threat- and reward-associated cues. However, it remains unknown whether PL regulates threat-avoidance vs. reward-approaching responses when an animals' decision depends on previously associated memories. Using a conflict model in which male Long-Evans rats retrieve memories of shock- and food-paired cues, we observed two distinct phenotypes during conflict: i) rats that continued to press a lever for food (*Pressers*); and ii) rats that exhibited a complete suppression in food seeking (*Non-pressers*). Single-unit recordings revealed that increased risk-taking behavior in *Pressers* is associated with persistent food-cue responses in PL, and reduced spontaneous activity in PL glutamatergic (PL^{GLUT}) neurons during conflict. Activating PL^{GLUT} neurons in *Pressers* attenuated food-seeking responses in a neutral context, whereas inhibiting PL^{GLUT} neurons in *Non-pressers* reduced defensive responses and increased food approaching during conflict. Our results establish a causal role for PL^{GLUT} neurons in mediating individual variability in memory-based risky decision making by regulating threat-avoidance vs. reward-approach behaviors.

72 INTRODUCTION

73 The brain's ability to identify and discriminate cues associated with threat or reward
74 allows organisms to respond appropriately to changes in the environment ([Schultz,](#)
75 [2015; Hu, 2016](#)). Animals respond to threatening cues with a series of defensive
76 behaviors including avoidance responses that decrease their chances of being
77 exposed to aversive outcomes ([McNaughton and Corr, 2014; Krypotos et al., 2015;](#)
78 [Cain, 2019](#)). In contrast, reward cues have attractive and motivational properties that
79 elicit approach behavior ([Robinson and Flagel, 2009; Morales and Berridge, 2020](#)).
80 When animals are exposed to threat and reward cues simultaneously, an approach-
81 avoidance conflict emerges, and decision-making processes are recruited to resolve
82 the situation ([Kirlic et al., 2017; Barker et al., 2019](#)). While many studies have
83 investigated the neural mechanisms that control threat-avoidance and reward-
84 approach independently of each other, it is unclear how the brain uses previously
85 learned information to regulate the opposing behavioral drives of avoiding threats and
86 seeking rewards during a conflict situation.

87 Neurons in the prelimbic (PL) subregion of the medial prefrontal cortex (mPFC)
88 change their firing rates in response to cues that predict either threat or reward ([Baeg](#)
89 [et al., 2001; Burgos-Robles et al., 2009; Burgos-Robles et al., 2013; Moorman and](#)
90 [Aston-Jones, 2015; Dejean et al., 2016; Otis et al., 2017](#)). Accordingly, activity in PL
91 neurons is necessary for the retrieval of both food- and threat-associated memories
92 ([Sierra-Mercado et al., 2011; Courtin et al., 2014; Sangha et al., 2014; Do-Monte et](#)
93 [al., 2015; Otis et al., 2017](#)). PL neurons are reciprocally connected with the
94 basolateral nucleus of the amygdala (BLA) ([McDonald, 1991; Vertes, 2004](#)), a region
95 implicated in the detection of threats or rewards ([Amir et al., 2015; Namburi et al.,](#)
96 [2015; Beyeler et al., 2016; Zhang et al., 2020](#)). During a risky foraging task in rats,
97 dynamic modifications in the activity of PL and BLA neurons correlate with the
98 detection of imminent threats and the defensive readiness for action ([Kim et al., 2018;](#)

Kyriazi et al., 2020). In addition, during a modified Pavlovian cue discrimination task involving footshocks as punishment, increased activity in the BLA-PL pathway is sufficient and necessary for the expression of freezing responses (Burgos-Robles et al., 2017), a passive form of defensive behavior. Conversely, inhibitory signaling in PL neurons correlates with threat avoidance (Diehl et al., 2018), an active form of defensive behavior. While these studies suggest a potential role of PL during motivational conflict involving states of certainty (i.e., imminent threats), it is unknown whether changes in PL activity underlie the behavioral variability in approach-avoidance responses under states of uncertainty, when animal's decision depends entirely on the retrieval of previously associated memories. It is also unclear whether PL activity is necessary to coordinate appropriate behavioral responses during conflict, and if so, which sub-types of PL neurons govern the competing demands of approaching rewards vs. avoiding potential threats.

To address these questions, we designed an approach-avoidance conflict test that assesses the ability of rats to remember cues previously associated with either food or footshocks to make a behavioral decision. Using a combination of optogenetics and single-unit recordings, we investigated rats' individual variability in reward seeking and defensive responses during the conflict test and correlated their behaviors (e.g. freezing, avoidance, risk-assessment) with the firing rate of photoidentified glutamatergic and GABAergic neurons in PL. We then examined the role of PL neurons in risky decision-making by optogenetically manipulating PL activity with high temporal resolution and cell-type specificity during the conflict test.

RESULTS

Rats show individual variability in reward-seeking and defensive responses during the approach-avoidance conflict test.

125 To investigate the motivational conflict between approaching rewards and avoiding
126 potential threats, we established a behavioral model in which rats need to balance
127 food seeking with conditioned defensive responses based on their memories of
128 previously acquired cues. Food-restricted rats (18 g of chow per day) were initially
129 placed in an operant box and trained to press a lever for sucrose in the presence of
130 audiovisual cues that signaled the availability of food. Each lever press during the
131 audiovisual cue presentation resulted in the delivery of a sucrose pellet into a nearby
132 dish (see Methods for details). When rats reached 50% of discrimination during cued
133 food-seeking, they began lever-pressing for sucrose preferentially during the
134 audiovisual cues (**Figure 1 - Figure supplement 1A-B**). During the habituation day,
135 rats were placed in an odor arena and familiarized with the food cues and the neutral
136 odor amyl acetate (see Methods for details). Next, to pair the odor cue with an
137 aversive stimulus, rats were exposed to an olfactory threat conditioning training (Day
138 1). Animals were placed in an operant box (conditioning box; **Fig. 1A left**) previously
139 connected to an olfactometer and habituated to one odor presentation (amyl acetate,
140 30 s) without footshock, followed by five odor presentations of the same odor that co-
141 terminated with an electrical footshock (0.7 mA, 1 s duration, 270-390 s inter-trial
142 intervals, **Fig. 1A far-left**). Food cues (30 s duration) were presented during the odor
143 intervals to assess how threat conditioning alters lever press responses. Rats showed
144 robust defensive responses during the threat conditioning training, as evidenced by
145 an increase in freezing (Shapiro Wilk normality test, $p < 0.001$, Friedman Test,
146 Friedman statistic = 84.08, $p < 0.001$; Dunn's *post-hoc*, $p < 0.001$) during the
147 conditioned odor presentation (**Fig. 1B**), and a decrease in lever presses (**Fig. 1C**,
148 Shapiro Wilk normality test, $p < 0.001$, Friedman Test, Friedman statistic = 35.11, $p <$
149 0.001 ; Dunn's *post-hoc*, $p < 0.001$) and an increase in the latency to press the lever
150 (**Fig. 1D**, Shapiro Wilk normality test, $p < 0.001$, Friedman Test, Friedman statistic =
151 29.45, $p < 0.001$; Dunn's *post-hoc*, $p < 0.001$) during the presentation of the food

cues across trials. After rats have acquired the reward and threat associations, they were returned to the same odor arena in which they were previously habituated and exposed to a test session (Day 2) (**Fig 1A, right**). The test session consisted of three different phases: (i) a *Reward Phase*, in which only the audiovisual cues signaling the availability of food were presented; (ii) an *Odor Phase*, in which only the conditioned odor was presented, and a (iii) *Conflict phase*, in which both the food cues and the conditioned odor were presented simultaneously (**Fig 1A, far-right**).

During the reward phase, rats spent ~40% of the time in the food area and pressed the lever for food in ~95% of the food cue trials, without exhibiting significant defensive behaviors (**Fig. 1E-J**). Introduction of the shock-paired odor during the odor phase reduced the percentage of time rats spent in the food area to ~15% (Shapiro Wilk normality test, $p < 0.05$, Friedman Test, Friedman statistic = 32.19, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$) and increased defensive behaviors characterized by freezing (Shapiro Wilk normality test, $p < 0.05$, Friedman Test, Friedman statistic = 40.46, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$), avoidance (Shapiro Wilk normality test, $p < 0.05$, Friedman Test, Friedman statistic = 31.67, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$), and risk-assessment responses (Shapiro Wilk normality test, $p < 0.05$, Friedman Test, Friedman statistic = 29.86, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$; **Fig. 1E-H**). These defensive behaviors were attenuated by the introduction of food cues during the conflict phase, as evidenced by a reduction in the percentage of time avoiding the conditioned odor (**Fig. 1F**, Dunn's *post-hoc*, $p = 0.0031$) and an increase in the percentage of time approaching the food area (**Fig. 1H**, Dunn's *post-hoc*, $p < 0.001$). This indicates that the concomitant presentation of food cues and shock-paired odor induced a behavioral conflict in the animals. Interestingly, when we analyzed the percentage of rewarded presses during the conflict phase (**Fig. 1J**), two behavioral phenotypes emerged: *i*) rats that continued to press the lever for food in the presence of the threatening odor (*Pressers*, **Fig. 1K**); and *ii*) rats that showed a

complete suppression in lever presses in the presence of the threatening odor (*Non-pressers*, **Fig. 1L**). We then separated the animals into two different groups based on whether the animals pressed the lever or not during the conflict phase and compared their behaviors during the entire test session (**Fig. 1K-R, Video 1**). *Pressers* and *Non-pressers* exhibited similar behavioral responses during the reward phase (all p 's > 0.05 , see table S1). However, during the odor and the conflict phases, *Pressers* showed a lower percentage of time exhibiting freezing (**Fig. 1M**, two-way repeated measures ANOVA, interaction - $F_{(2, 60)} = 29.54$ $p < 0.001$, Bonferroni *post-hoc* test - odor phase, $p < 0.001$; conflict phase, $p < 0.001$) and avoidance responses (**Fig. 1N**, Two-way repeated measures ANOVA, interaction- $F_{(2, 60)} = 23.27$, $p < 0.001$, Bonferroni *post-hoc* test - odor phase, $p < 0.001$; conflict phase, $p < 0.001$), and a greater percentage of time approaching the food area (**Fig. 1P**, two-way repeated measures ANOVA, interaction- $F_{(2, 60)} = 22.49$, $p < 0.001$, Bonferroni *post-hoc* test - odor phase, $p = 0.0453$; conflict phase, $p < 0.001$) when compared to *Non-pressers* during both the odor and the conflict phases. A principal component analysis (PCA) showed that PC1 explained most of the variance of the data ($> 60\%$), with latency to press the lever and percentage of time in the food area being the two variables that most contributed to PC1 (0.54 and 0.52, respectively, see Methods for details). Because these two variables are directly associated with lever presses, the PCA results support our binary classification of rats into *Pressers* and *Non-pressers* based on whether they pressed the lever or not during the conflict phase.

Subsequent behavioral analyses demonstrated that these two individual phenotypes were not due to prior differences in reward-seeking motivation or odor-shock association because *Pressers* and *Non-pressers* showed similar lever pressing rates during the cued food-seeking training (**Figure 1 - Figure supplement 1A-B**; Two-way repeated measures ANOVA, interaction- $F_{(1, 55)} = 0.1065$, $p = 0.7454$) and threat conditioning phase (**Figure 1 - Figure supplement 1C**; Two-way repeated

measures ANOVA, $F_{(1,51)} = 0.265$, $p = 0.608$), as well as the same freezing levels (Two-way repeated measures ANOVA, $F_{(1,51)} = 3.737$, $p = 0.058$) and maximum speed (Two-way repeated measures ANOVA, $F_{(1,51)} = 6.538e007$, $p = 0.999$) in response to the shock-paired odor during the threat conditioning phase (**Figure 1 - Figure supplement 1D-E**). The two phenotypes might not be attributed to prior differences in the relative salience of the odor and the audiovisual cues because *Pressers* and *Non-pressers* spent the same time investigating the odor and exhibited the same response latency to the audiovisual cues during the pre-conditioning period (**Figure 1 - Figure supplement 1F-G**; Welch's t-test, $t = 0.41$, $p = 0.683$ and Welch's t-test, $t = 0.61$, $p = 0.538$, respectively). Although *Non-pressers* exhibited higher freezing levels during pre-odor trials 3 and 4 of the threat conditioning phase (**Figure 1 - Figure supplement 1H**, $F_{(5, 250)} = 3.038$, $p = 0.011$, Bonferroni's post-hoc $p < 0.05$), freezing responses before the first food cue and odor presentation were the same during the test day, indicating similar contextual discrimination between the two groups (**Figure 1 - Figure supplement 1I**, Shapiro Wilk normality test, $p < 0.05$, Mann-Whitney test, $U = 248$, $p = 0.113$). A minute-by-minute analysis during the odor phase demonstrated that behavioral differences in freezing (Two-way repeated measures ANOVA, Group $F_{(1,50)} = 13.07$, $p < 0.001$; Interaction, $F_{(9,450)} = 1.327$, $p = 0.220$), avoidance (Group $F_{(1,50)} = 20.31$, $p < 0.001$; Interaction, $F_{(9,450)} = 2.109$, $p = 0.027$, Bonferroni post-hoc min 1 vs. min 10 $p > 0.999$), and time spent in the food area (Group $F_{(1, 50)} = 117.5$, $p = 0.001$; Interaction, $F_{(9, 450)} = 0.573$, $p = 0.819$) between *Pressers* and *Non-pressers* were already observed in the beginning of the odor phase, and these behaviors remained constant in both groups across the entire duration of the session (**Figure 1 - Figure supplement 1J-L**), ruling out the possibility that the group differences were caused by extinction of the odor-shock association in *Pressers*.

Together, our results demonstrate that our conflict model is a suitable paradigm to investigate the interactions between reward- and threat-associated memories. Given that rats exhibit individual differences in food seeking and defensive responses during the test session, we next took advantage of the two observed phenotypes to examine the neuronal correlates of risk-taking (*Pressers*) and risk-avoiding (*Non-pressers*) behaviors in PL neurons.

PL neurons respond differently to reward cues in *Pressers* vs. *Non-pressers* during the conflict test.

To investigate the role of PL neurons in regulating food-approach and threat-avoidance responses, we performed single-unit recordings across the different phases of the conflict test (**Fig. 2A**). We aligned the activity of PL neurons to the onset of the food cues during the reward phase and tracked the firing rates of the same cells during the conflict phase. Using the behavioral classification shown in **Fig. 1J**, we separated the animals into *Pressers* or *Non-pressers* and compared changes in PL activity in response to food cues during the reward and conflict phases (**Fig. 2B-V**). When PL activity was time-locked to the onset of the food cues during the reward phase, *Pressers* showed a higher number of food-cue responsive neurons than *Non-pressers* (**Fig. 2C-D vs. Fig. 2M-N**; Fisher Exact Test, 33% in *Pressers* vs. 21% in *Non-pressers*, $p = 0.0418$), with a similar proportion of excitatory and inhibitory responses between the two groups (Fisher Exact Test, $p = 0.073$ for excitatory, $p = 0.571$ for inhibitory). During the conflict phase, both *Pressers* and *Non-pressers* showed a significant reduction in the number of food-cue responsive neurons (**Fig. 2C-D vs. 2M-N**; Fisher Exact Test, from 33% to 14% in *Pressers*, $p < 0.001$; from 21% to 6% in *Non-pressers*, $p = 0.0086$), as well as in the magnitude of excitatory food-cue responses compared to the reward phase (**Fig. 2E inset and Fig.**

20 inset, Shapiro-Wilk normality test, $p < 0.001$, Wilcoxon test, *Pressers* – $W = -824$, $p < 0.001$; *Non-pressers* – $W = -37$, $p = 0.032$). In addition, the percentage of reduction in the number of responsive cells across the phases was similar between *Pressers* and *Non-pressers* (**Fig. 2C vs. Fig. 2M**, Fisher Exact Test, 47 out of 232 neurons for *Pressers*, 13 out of 89 neurons for *Non-pressers*, $p = 0.427$), suggesting that PL neurons can distinguish between reward and conflict situations (**Fig. 2G vs. 2H and 2Q vs. 2R**).

Using a temporal frequency separation of the food cue responses into transient (< 600 ms duration) and sustained (≥ 900 ms duration) activity ([Bezdudnaya et al., 2006](#)), we revealed that *Pressers* display a higher proportion of sustained excitatory responses during the reward phase, when compared to *Non-pressers* (**Fig. 2F vs. 2P**, 50 % in *Pressers* vs. 10% in *Non-pressers*, Fisher Exact Test, $p = 0.032$). In addition, *Pressers* showed a higher magnitude of inhibitory food-cue responses during the reward phase when compared to *Non-pressers* (**Fig. 2I blue bar inset vs. 2S blue bar inset**, Shapiro-Wilk normality test, $p < 0.001$, Mann-Whitney test, $U = 50$, $p = 0.0045$) and, in contrast to *Non-pressers*, such responses were attenuated during the conflict phase (**Fig. 2I inset vs. Fig. 2S inset**, *Pressers*, Shapiro-Wilk normality test, $p < 0.001$, Wilcoxon test – $W = 367$, $p < 0.001$; *Non-pressers*, Paired Student's t -test, $t = 0.59$, $p = 0.569$). A correlation analysis during the conflict phase revealed that food cue-evoked excitatory PL activity in *Pressers* was inversely correlated with lever press latency, indicating that the higher the firing rate of food-cue responsive cells after the onset of the audiovisual cues, the quicker the animals searched for rewards (**Figure 2 - Figure supplement 1A-B**).

Next, we time-locked the activity of PL neurons to the onset of the food cues during the conflict phase. Both *Pressers* and *Non-pressers* show the same percentage of food-cue responsive neurons (**Figure 2 - Figure supplement 2A-B vs. Figure 2 - Figure supplement 2K-L**, 28% in *Pressers* vs. 20% in *Non-pressers*,

Fisher Exact Test, $p = 0.391$) and the same magnitude of excitatory food-cue responses during the conflict phase (**Figure 2 - Figure supplement 2C, E, F vs. Figure 2 - Figure supplement 2M, O, P**; area under the curve, *Pressers* vs. *Non-pressers*, Shapiro-Wilk normality test, $p < 0.001$, Mann-Whitney, $U = 107$, $p = 0.123$). However, in *Pressers*, 42% of excitatory food-cue responses showed sustained activity during the conflict phase whereas such responses were completely absent in *Non-pressers* (**Figure 2 - Figure supplement 2D vs. Figure 2 - Figure supplement 2N**, Fisher Exact Test, $p = 0.018$).

In addition to food cue responses, we observed a significant proportion of PL neurons that changed their firing rates in response to lever presses (23%) or rewarded food dish entries (16%, **Figure 2 - Figure supplement 3A-N**). A longitudinal tracking of PL activity throughout the reward phase demonstrated that most PL responsive neurons changed their activities selectively to food cues, lever presses, or food dish entries, with a smaller number of cells responding during two or more of these events (**Figure 2 - Figure supplement 3O**). An analysis of PL activity at a random timepoint (e.g., 1 second before the food cue onset) resulted in less than 3% of responsive cells, indicating that the proportion of PL neurons that responded to food cues, lever presses, and food dish entries was different from the proportion obtained by chance (Fisher Exact Test, all p 's < 0.05 , see table S1). These observations suggest that PL neurons exhibit a heterogeneous pattern of activity during reward-seeking behavior, consistent with a recent study using calcium imaging recordings from PL neurons in head-fixed mice ([Grant et al., 2021](#)).

To further explore whether changes in activity dynamics of PL neurons differ between *Pressers* and *Non-pressers*, we compared the spontaneous firing rate of the neurons before vs. after each phase of the test session (**Figure 2 - Figure supplement 4A**). While *Pressers* showed the same proportion of neurons excited and inhibited across the different phases (Fisher Exact Test, all p 's > 0.05 , see table

S1), *Non-pressers* exhibited a significant increase in the proportion of neurons excited during the conflict phase (**Figure 2 - Figure supplement 4C**, Fisher Exact Test, $p = 0.015$ compared to odor phase, $p = 0.059$ compared to *Pressers*). This suggests that increased spontaneous activity in PL neurons during the conflict phase may be associated with the complete suppression in lever presses observed in *Non-pressers* (**Fig. 1J**). Collectively, these results suggest that differences in the number and magnitude of excitatory food cue responses, as well as in the spontaneous activity of PL neurons during the conflict test, may contribute to the individual differences in risky decision making observed between the two behavioral phenotypes.

Different subsets of PL neurons signal freezing, avoidance, and risk-assessment behaviors in both *Pressers* and *Non-pressers*.

To investigate whether PL activity correlates with the expression of distinct defensive behaviors during the test session, we used a pose estimation algorithm (DeepLabCut, see Methods for details) to identify the onset of freezing, avoidance or risk-assessment responses and align these time points with the activity of PL neurons. We found that a small percentage of PL neurons changed their firing rates during the onset of freezing (**Fig. 3A**), avoidance (**Fig. 3B**), or risk-assessment (**Fig. 3C**) behaviors in both *Pressers* and *Non-pressers*, with a similar proportion of excitatory and inhibitory responses being observed in the two groups (**Fig. 3A-I**). Interestingly, most PL responsive neurons (80%) changed their activities exclusively during the onset of one of these three behaviors, with a reduced number of avoidance-responsive cells also responding during the onset of risk-assessment behavior (**Fig. 3J-M**). Moreover, a smaller fraction of PL neurons changed their firing rates 600 ms before the onset of either freezing, avoidance or risk-assessment responses in both *Pressers* and *Non-pressers* (**Figure 3 - Figure supplement 1A-M**), indicating that

some PL neurons can anticipate an animal's defensive behavior during the test. Overall, these results suggest that different subsets of PL neurons signal distinct behavioral outcomes during a conflict situation, with only a reduced number of PL neurons encoding the aversive salience of environmental cues independently of the defensive response expressed by the animal.

***Pressers* and *Non-pressers* show significant differences in delta and theta oscillations in PL**

Previous studies have shown that oscillations in mPFC neuronal activity at different frequency bands correlate with distinct behavioral states in both rodents and humans (Narayanan et al., 2013; Harris and Gordon, 2015). Neural oscillations in the mPFC emerge from the network of excitatory and inhibitory synaptic connections and are thought to contribute to neural communication when subjects engage in reward and threat memory tasks (Hyman et al., 2011; Likhtik and Paz, 2015; Park and Moghaddam, 2017; Widge et al., 2019). To investigate whether *Pressers* and *Non-pressers* show significant differences in PL oscillations during conflict, we recorded local field potentials (LFPs) from PL neurons and calculated the average of power spectral density (PSD) at different frequencies across the test session. After comparing the PSD contribution for each frequency range in *Pressers* and *Non-pressers*, we observed that most of the signal originated from the delta (0-4 Hz) and theta (4-10 Hz) bands, with a much smaller contribution coming from the alpha (10-14 Hz), beta (14-35 Hz), and gamma (>35 Hz) frequencies (**Fig. 4A**). We therefore focused our analyses on these two bands and found that *Pressers* displayed increased power in the delta band, whereas *Non-pressers* exhibited increased power in the theta band during the three phases of the test session (**Fig. 4B-C**, Welch's t-test of Area Under the Curve, all p's < 0.001, see table S1). Differences between *Pressers* and *Non-pressers* were also observed in the time-frequency domain

through changes in the log of PSD for delta and theta bands across the different phases (**Fig. 4D-E**, Paired Student's t-test of Area Under the Curve, all p 's < 0.001, see table S1). These results indicate that phenotypic differences in approach-avoidance conflict are associated with distinct oscillatory frequencies in PL.

In pressers, PL^{GLUT} neurons show reduced spontaneous activity during the conflict phase.

The rodent mPFC, including PL, is primarily composed of excitatory glutamatergic cells that correspond to 75-85% of the neurons in this area. In contrast, inhibitory GABAergic interneurons comprise 15-25% of the local neurons (Santana et al., 2004; Gabbott et al., 2005). Previous studies have shown that PL glutamatergic (PL^{GLUT}) neurons are necessary for the retrieval of conditioned threat responses (Domonte et al., 2015), whereas PL GABAergic (PL^{GABA}) neurons are implicated in both the encoding and the retrieval of threat associations by regulating the firing rate of PL^{GLUT} neurons (Courtin et al., 2014; Cummings and Clem, 2020). In addition, during foraging in a safe context, food-associated cues activate both PL^{GLUT} and PL^{GABA} neurons (Burgos-Robles et al., 2013; Gaykema et al., 2014), and inactivation of PL^{GLUT} neurons may increase or reduce conditioned food-seeking responses depending on the specific downstream projections that are being modulated (Otis et al., 2017). While these studies suggest a role for both PL^{GLUT} and PL^{GABA} neurons in the regulation of threat and food-seeking responses in isolation, it remains unexplored how these two subsets of PL neurons regulate the trade-off between seeking rewards and avoiding potential threats during a conflict situation. To address this question, we combined single-unit recordings with optogenetics to track the neuronal activity of photoidentified PL^{GLUT} and PL^{GABA} neurons during the test session.

392 For photoidentification of PL^{GLUT} neurons, we injected into PL a viral vector
393 (AAV-CaMKII α -hChR2-(H134R)-eYFP) with a gene promoter (CaMKII α) that favors
394 the expression of the light-activated cation channel channelrhodopsin (ChR2) in
395 PL^{GLUT} neurons. This CaMKII α labeling approach has been successfully used in
396 previous studies ([Gradinaru et al., 2009](#); [Tye et al., 2011](#)) and was validated here for
397 PL neurons by showing a lack of immunocolabeling between the viral vector and the
398 GABAergic marker GAD67 (**Fig. 5A**). Rats expressing ChR2 selectively in PL^{GLUT}
399 neurons were implanted with an optrode into the same region for optogenetic-
400 mediated identification of PL^{GLUT} neurons at the end of the behavioral session (**Fig.**
401 **5B**). Among the recorded PL cells, 36 out of 104 neurons (n = 5 rats) showed short-
402 latency responses (< 6 ms) and high spike reliability (Fano Factor ratio > 1) to laser
403 illumination and were classified as PL^{GLUT} neurons (**Fig. 5C-D** and Methods). The < 6
404 ms criterion was defined by using the triangle method detection ([Zack et al., 1977](#)) to
405 identify the cluster division in the histogram distribution of response latencies (**Fig. 5C**
406 and Methods). The < 6 ms criterion was similar or stricter than the response latency
407 criterion used in previous photoidentification studies *in vivo* ([Lima et al., 2009](#); [Cohen](#)
408 [et al., 2012](#); [Burgos-Robles et al., 2017](#); [Allsop et al., 2018](#)). Photoactivation of
409 PL^{GLUT} neurons can lead to indirect activation of synaptically connected neurons in
410 the same cortical region, but these indirect responses to laser illumination take longer
411 than 9 ms to occur ([Lima et al., 2009](#)). For photoidentification of PL^{GABA} neurons, we
412 injected into PL a viral vector (AAV-mDlx-ChR2-mCherry) with a gene promoter
413 (mDlx) that favors the expression of ChR2 in PL^{GABA} neurons. This mDlx labeling
414 approach has been successfully used in previous studies ([Dimidschstein et al., 2016](#);
415 [Sun et al., 2020](#)), and was validated here for PL neurons by using two different
416 methods: an immunohistochemical approach that resulted in significant
417 immunocolabeling between the viral vector and the GABAergic marker GAD67 (**Fig.**
418 **5E**), and an *in situ* hybridization approach which confirmed that ~88% of the cells

labeled with the viral vector also expressed the GABAergic marker vGAT (**Figure 5 - Figure supplement 1A-B**). Rats expressing ChR2 selectively in PL^{GABA} neurons were implanted with an optrode into the same region for optogenetic-mediated identification of PL^{GABA} neurons at the end of the behavioral session (**Fig. 5F**). Among the recorded PL cells, 69 out of 338 neurons (n = 19 rats) showed short-latency responses (< 6 ms) and high spike reliability (Fano Factor ratio > 1) to laser illumination and were classified as PL^{GABA} neurons (**Fig. 5G-H and Methods**).

After separating the photoidentified cells into PL^{GLUT} and PL^{GABA} neurons, we aligned their activities to the onset of the food cues and compared changes in firing rates from the reward to the conflict phase in *Pressers* (**Fig. 5I**). We observed that the proportions of excitatory and inhibitory food cue responses for PL^{GLUT} and PL^{GABA} neurons were similar when comparing between the reward and the conflict phases as well as within each one of the phases (**Fig. 5J-K**, Fisher Exact Test, all p's > 0.05, see table S1). Next, we analyzed the spontaneous activity of PL^{GLUT} and PL^{GABA} neurons and compared changes in their firing rates across the different phases of the test session (**Fig. 5L**). We found that the average firing rate of PL^{GLUT} neurons remained the same across the different phases of the test (~5 Hz; **Fig. 5M**, One-way repeated measures ANOVA, $F_{(2.03, 69.02)} = 1.204$, $p = 0.306$), with most of the cells (57%) changing their activities in more than one session (**Fig. 5N**). An average firing rate analysis across phases demonstrated that the activity of PL^{GLUT} neurons didn't change significantly from the reward to the odor phase (Fisher Exact Test, all p's > 0.05, see table S1), but was inhibited from the odor to the conflict phase when *Pressers* resumed searching for food (**Fig. 5O**, Fisher Exact Test, odor vs conflict, $p = 0.0046$). Similar to PL^{GLUT} neurons, the average firing rate of PL^{GABA} neurons also remained the same across the different phases of the test (~8 Hz, **Fig. 5P**, One-way repeated measures ANOVA, $F_{(1.164, 79.17)} = 0.013$, $p = 0.935$), with most of the cells (62%) changing their activities in more than one session (**Fig. 5Q**). However, in

contrast to PL^{GLUT} neurons, a group analysis of the firing rates of PL^{GABA} neurons did not reveal significant differences across the phases (**Fig. 5R**, Fisher Exact Test, all p 's > 0.05, see table S1). Because PL is comprised of different subpopulations of interneurons that inhibit each other during food seeking or defensive responses (Gaykema et al., 2014; Cummings and Clem, 2020), we cannot rule out the possibility that distinct subsets of PL^{GABA} neurons were preferentially recruited during each one of the phases.

To evaluate how the spontaneous activity of the same PL neurons changed during the test session, we tracked the firing rate of PL^{GLUT} and PL^{GABA} neurons across the different phases. We found that all PL^{GLUT} neurons that were either excited or inhibited during the reward phase responded in opposite direction or did not change their activities during the odor phase (**Figure 5 - Figure supplement 2A-B**), suggesting the existence of distinct subpopulations of PL^{GLUT} neurons that encode reward and threat-related information differently in our task. In contrast, no significant differences in the proportions of excitation and inhibition were observed in PL^{GABA} neurons during the transition from reward to odor phase nor during the transition from odor to conflict phase for both subsets of PL neurons (**Figure 5 - Figure supplement 2C-F**). Furthermore, both PL^{GLUT} and PL^{GABA} neurons showed the same proportion of excitatory and inhibitory responses before or after the onset of freezing, avoidance, or risk-assessment behaviors (**Figure 5 - Figure supplement 3A-F**, Fisher Exact Test, all p 's > 0.05, see table S1). These results indicate that both glutamatergic and GABAergic neurons in PL may contribute to the expression of distinct defensive responses during conflict. Together, our data suggest that a significant proportion of PL^{GLUT} neurons are inhibited when rat's behavior transitions from increased defensive responses during the conditioned odor phase to increased food-seeking responses during the conflict phase.

Photoactivation of PL^{GLUT}, but not PL^{GABA} neurons, suppresses reward-seeking responses.

To further establish whether changes in the activity of PL neurons can alter cue-triggered food-seeking responses, we used an optogenetic approach to selectively activate either PL^{GLUT} or PL^{GABA} neurons during a cued food-seeking test in a neutral context. We initially infused either the viral vector AAV-CaMKII α -ChR2-eYFP (**Fig. 6A**) or AAV-mDlx-ChR2-mCherry (**Fig. 6E**) into PL and implanted an optrode into the same region to examine how photoactivation of PL^{GLUT} or PL^{GABA} neurons change local activity. Laser illumination of PL^{GLUT} somata increased the firing rate of most responsive PL neurons (9 out of 20 neurons, 45%), with some neurons showing reduced activity (6 out of 20 neurons, 30%, **Fig. 6B-D**). Neurons that increased their activities showed shorter response latencies (3.31 ± 1.03 ms) compared to neurons that reduced their activities (21.0 ± 3.74 ms) when analyzed in short bins of 1 ms, suggesting direct responses (i.e., opsin-mediated) versus indirect responses (i.e., multi-synaptic), respectively. Conversely, although some PL^{GABA} neurons showed increased activity right after the laser onset (revealed by short bins of 1 ms, as shown in **Fig. 5E-H**), illumination of PL^{GABA} somata reduced the firing rate of all responsive PL neurons when analyzing the entire duration of the train (16 out of 22 neurons, 73%; **Fig. 6F-H**), indicating a suppression in local activity.

After investigating the local effects of photoactivating either PL^{GLUT} and PL^{GABA} neurons, we infused another set of animals with the same viral vectors in PL and implanted bilateral optical fibers into the same region to manipulate PL activity during the cued food-seeking test (**Fig. 6I-J**). Rats expressing only eYFP in PL were used to control for any nonspecific effects of viral transduction or laser heating. To assess the effects of PL photoactivation on lever presses, we alternated two trials of food cues with the laser on vs. laser off conditions in a total of 12 trials (**Fig. 6K-L**).

Photoactivation of PL^{GLUT} (CaMKII-ChR2), but not PL^{GABA} (mDlx-ChR2) neurons,

reduced the frequency of lever presses (**Fig. 6M**, Two-way Repeated measures ANOVA, $F_{(10,180)} = 7.009$, $p < 0.001$; Bonferroni post-hoc, CaMKII-ChR2 vs. Control, all laser on periods - $p < 0.001$, mDlx-ChR2 vs. Control, all laser on periods - $p > 0.05$) and increased the latency for the first press after the cue onset (**Fig. 6N**, $F_{(10,180)} = 9.931$, $p < 0.001$; Bonferroni post-hoc, CaMKII-ChR2 vs. Control, all laser on periods - $p < 0.001$, mDlx-ChR2 vs. Control, all laser on periods - $p > 0.05$), when compared to control group. The diminished behavioral effect observed during the third block of laser on could be the result of conformational changes in the opsin (e.g. photobleaching) or temporary depletion of synaptic vesicles following repeated laser illumination, as previously reported (Kittellmann et al., 2013; Stahlberg et al., 2019).

Photoactivation of either PL^{GLUT} or PL^{GABA} neurons did not induce freezing behavior (**Fig. 6O**, $F_{(10,180)} = 1.124$, $p = 0.346$). These results are consistent with our electrophysiological recordings in **Fig. 5O** showing that increased inhibition in the firing rate of PL^{GLUT} neurons correlates with augmented reward-seeking responses during conflict. Overall, these findings suggest that increasing the activity of PL^{GLUT} neurons is sufficient to suppress cued reward-seeking responses in a neutral context.

Photoinhibition of PL^{GLUT} neurons in *Non-pressers* reduces freezing responses and increases food approaching during conflict.

Our electrophysiological experiments in **Fig. 5O** demonstrate that PL^{GLUT} neurons are inhibited when rats' behavior transitions from defensive responses in the odor phase to food-seeking responses in the conflict phase. In addition, our photoactivation experiments in **Fig. 6K-O** indicate that increasing the activity of PL^{GLUT} neurons suppresses cued reward-seeking behavior in rats that are pressing a lever for food. We therefore hypothesized that photoinhibition PL^{GLUT} neurons during conflict would attenuate defensive behaviors and rescue food-seeking responses in *Non-pressers*. To test this hypothesis, we injected a group of rats with the viral vector

527 AAV-CaMKII α -eNpHR-eYFP (or AAV-CaMKII α -eYFP) into PL to express the
528 inhibitory opsin halorhodopsin (or eYFP control) selectively in PL^{GLUT} neurons (**Fig.**
529 **7A**). Rats were initially exposed to a cued food-seeking test to assess the effects of
530 photoinhibition of PL^{GLUT} neurons on food-seeking responses in a neutral context. We
531 observed that photoinhibition of PL^{GLUT} neurons had no effect on lever pressing rate
532 (Two-way repeated measures ANOVA, $F_{(5,110)} = 1.336$, $p = 0.254$), latency to press
533 the lever ($F_{(5,110)} = 0.637$, $p = 0.671$) or freezing ($F_{(5,95)} = 1.395$, $p = 0.231$) responses
534 before threat conditioning (**Figure 7 - Figure supplement 1A-E**).

535 Animals were then threat conditioned as in **Fig. 1** and on the following day
536 exposed to the odor arena for a test session. During the conflict phase, the first pair
537 of food cues was used to classify the animals into *Pressers* and *Non-pressers*,
538 whereas the subsequent pairs of food cues were alternated between laser on and
539 laser off conditions to assess the effects of illumination of PL^{GLUT} neurons on
540 approach-avoidance responses (**Fig. 7B-C**). Remarkably, photoinhibition of PL^{GLUT}
541 neurons (CaMKII-eNpHR, Shapiro-Wilk normality test, all p 's < 0.05, see table S1) in
542 *Non-pressers* reduced the percentage of time rats spent freezing (**Fig. 7D**, Wilcoxon
543 test, $W = -64$, laser off vs. laser on, $p = 0.0020$, Mann-Whitney Test, $U = 18$ Control
544 vs. CaMKII-eNpHR, $p = 0.319$) and avoiding the odor area (**Fig. 7E**, Wilcoxon test, W
545 $= -21$, laser off vs. laser on, $p = 0.031$; Mann-Whitney Test, $U = 19.5$ Control vs.
546 CaMKII-eNpHR, $p = 0.365$), and increased the percentage of time rats spent
547 approaching the food area (**Fig 7F**, Wilcoxon test, $W = 21$, laser off vs. laser on, $p =$
548 0.031 ; Mann-Whitney Test, $U = 17$ Control vs. CaMKII-eNpHR, $p = 0.221$) during the
549 food cue presentation, when compared to the eYFP-control group (Wilcoxon test,
550 Freezing: $W = 3$, laser off vs. laser on, $p = 0.812$; avoidance: $W = 3$, laser off vs. laser
551 on, $p = 0.500$, time in food area: $W = -3$, laser off vs. laser on, $p = 0.500$). Despite the
552 increase in food approaching behavior, photoinhibition of the same cells had no effect
553 on the percentage of rewarded lever presses (**Fig. 7G**, Wilcoxon test, $W = 6$, laser off

554 vs. laser on, $p = 0.250$; Mann-Whitney Test, $U = 22.5$ Control vs. CaMKII-eNpHR, $p =$
555 0.697) or latency to press the lever (**Fig. 7H**, Wilcoxon test, $W = -10$, laser off vs.
556 laser on, $p = 0.125$; Mann-Whitney Test, $U = 21$ Control vs. CaMKII-eNpHR, $p =$
557 0.357), when compared to the control group (Wilcoxon test, rewarded lever presses:
558 $W = -1$, laser off vs. laser on, $p > 0.999$; latency to press the lever: $W = 1$, laser off vs.
559 laser on, $p > 0.999$).

560 In another subset of *Non-pressers* (**Fig. 7I-K**), photoactivation of PL^{GABA}
561 neurons (mDlx-ChR2, Shapiro-Wilk normality test, all p 's < 0.05 , see table S1) did not
562 alter freezing (**Fig. 7L**, Wilcoxon test, $W = 18$, laser off vs. laser on, $p = 0.156$; Mann-
563 Whitney Test, $U = 14$ Control vs. mDlx-ChR2, $p > 0.999$), avoidance (**Fig. 7M**,
564 Wilcoxon test, $W = 4$, laser off vs. laser on, $p = 0.500$; Mann-Whitney Test, $U = 12$
565 Control vs. mDlx-ChR2, $p = 0.772$) time in food area (**Fig. 7N**, Wilcoxon test, $W = -2$,
566 laser off vs. laser on, $p = 0.750$; Mann-Whitney Test, $U = 11$ Control vs. mDlx-ChR2,
567 $p = 0.660$) rewarded lever presses (**Fig. 7O**, Wilcoxon test, $W = 3$, laser off vs. laser
568 on, $p = 0.500$; Mann-Whitney Test, $U = 13.5$ Control vs. mDlx-ChR2, $p > 0.999$) and
569 latency to press the lever (**Fig. 7P**, Wilcoxon test, $W = -3$, laser off vs. laser on, $p =$
570 0.500 ; Mann-Whitney Test, $U = 10$ Control vs. mDlx-ChR2, $p = 0.490$), when
571 compared to the control group (eYFP-control virus, white bars, $n = 4$, Wilcoxon test,
572 freezing: $W = -2$, laser off vs. laser on, $p = 0.875$; avoidance, $W = 0$, $p > 0.999$, time in
573 food area: $W = 3$, $p = 0.500$, rewarded lever presses: $W = 1$, $p = 0.500$, latency to
574 press the lever: all animals reached maximum latency). Photoactivation of PL^{GABA}
575 neurons in *Pressers* also did not affect defensive responses and food seeking
576 behavior during the conflict test (**Figure 7 - Figure supplement 2A-H**, Repeated
577 measures ANOVA, all p 's > 0.05 , see table S1). Taken together, these results
578 demonstrate that reduced activity in PL^{GLUT} neurons during conflict situations
579 decreases defensive responses and biases rats' behavior toward food seeking.

582 **DISCUSSION**

583 Using a novel approach-avoidance conflict test, we identified two distinct behavioral
584 phenotypes during the combined presentation of reward- and threat-paired cues: *i)*
585 rats that continued to press a lever for food (*Pressers*), and *ii)* rats that exhibited a
586 complete suppression in food-seeking responses (*Non-pressers*). Single-unit
587 recordings revealed that PL neurons regulate reward-approach vs. threat-avoidance
588 responses during situations of uncertainty, when rats use previously associated
589 memories to guide their decisions. We observed that increased risk-taking behavior in
590 *Pressers* was associated with a larger number of food-cue responses in PL neurons,
591 which showed sustained excitatory activity that persisted during the conflict phase,
592 when compared to *Non-pressers*. In addition, PL^{GLUT} neurons showed reduced
593 spontaneous activity during risky reward seeking and photoactivation of these cells in
594 a neutral context was sufficient to suppress lever press responses. Accordingly,
595 photoinhibition of PL^{GLUT} neurons at the onset of the food cues in *Non-pressers*
596 reduced defensive responses and increased food-approaching during the conflict
597 phase, consistent with our observation that a small fraction of PL neurons changed
598 their activity at the onset of freezing, avoidance or risk-assessment responses.
599 Altogether, these results suggest that under memory-based conflict situations,
600 reduced or increased activity in PL^{GLUT} neurons can favor the behavioral expression
601 of food-approaching or threat-avoidance responses, respectively.

602 During our approach-avoidance conflict test, *Pressers* and *Non-pressers*
603 showed similar levels of lever pressing before the conflict phase (e.g., cued food-
604 seeking training, threat conditioning, and reward phases). This observation suggests
605 that these two individual phenotypes most likely emerged during the test session and
606 were independent of prior differences in sucrose preference or food-seeking
607 motivation. Similarly, because both groups exhibited the same percentage of freezing

to the shock-paired odor during the olfactory threat conditioning session, the increased defensive behaviors and the reduced food-seeking responses observed in *Non-pressers* during the test session were unlikely due to higher acquisition of conditioned threat responses. Furthermore, other internal factors such as shock sensitivity or pain tolerance cannot be accounted for the individual differences observed in our experiments because both groups reacted equally to the unconditioned stimulus (i.e., velocity measured as maximum speed after the footshocks) and, different from other conflict tasks using footshocks as a punishment during the conflict test (Geller, 1960; Vogel et al., 1971; Oberrauch et al., 2019), in our model rats were not exposed to footshocks during the conflict phase. Therefore, the most plausible interpretation for the behavioral differences observed in our task is that *Pressers* and *Non-pressers* have allocated distinct motivational significance to the food- or shock-paired cues during the test session.

Individual differences in risky decision making have also been reported in other studies using rodent models of behavioral conflict involving footshock punishment (Simon et al., 2009; Jean-Richard-Dit-Bressel et al., 2019; Bravo-Rivera et al., 2021), reversal learning (Bari et al., 2010), or variations in reward probability (Ainslie, 1975; St Onge and Floresco, 2009; Dellu-Hagedorn et al., 2018), although the neural mechanisms underlying such differences are less clear. Evidence indicates that some of the neurobiological bases of individual variation in stimulus-reward response depend on differences in dopamine levels in subcortical circuits (Tomie et al., 2000; Flagel et al., 2007; Flagel et al., 2011), which are regulated by top-down mechanisms involving the mPFC (Ferenczi et al., 2016; Haight et al., 2017; Serrano-Barroso et al., 2019). Accordingly, our neural correlate analyses of risk-taking vs. risk-avoiding behaviors in the PL subregion of the mPFC revealed some clear differences between the two phenotypes, suggesting that PL neurons participate in behavioral selection when rats' decision depends on the conflicting memories of reward and threat. Both

635 *Pressers* and *Non-Pressers* showed a reduction in the number and magnitude of food-
636 cue responses from reward to conflict phases, indicating that PL neurons can
637 differentiate between situations involving motivational conflict and those that do not.

638 One intriguing finding in our study was the observation that *Pressers* showed a
639 larger number of sustained excitatory food-cue responses during the conflict phase,
640 when compared to *Non-pressers*. Because PL neurons are known for encoding the
641 value of reward-predictive cues (Sharpe and Killcross, 2015; Otis et al., 2017), the
642 increase in the number and magnitude of food cue responses observed in *Pressers*
643 might result in a greater allocation of attention to reward cues, which would explain
644 the persistent reward-seeking responses observed in this group during motivational
645 conflict. In support of this interpretation, reward-paired cues can acquire motivational
646 salience in some subjects and become sufficient to elicit reward-seeking responses in
647 both rodents (Robinson and Flagel, 2009; Robinson et al., 2014) and humans (Smith
648 et al., 2011; Jensen and Walter, 2014). Consistently, *Pressers* also showed a larger
649 number of food cue responses in PL before the conflict phase (i.e., reward phase),
650 although the percentage of rewarded presses and the latency to press the lever
651 during the reward phase were similar between the two groups.

652 Another possible interpretation for the differences in food-cue responses in
653 *Pressers* and *Non-pressers* is the reduced excitatory food cue responses in *Non-*
654 *pressers*, which may be mediated by cue-evoked activity in inhibitory inputs to PL
655 during the conflict phase. While the source of this inhibition is unclear, a potential
656 candidate are GABAergic neurons in the ventral tegmental area (VTA^{GABA}), which
657 correspond to 35% of the cells in this region and send significant projections to PL
658 (Nair-Roberts et al., 2008; Breton et al., 2019). Previous studies have shown that
659 VTA^{GABA} neurons change their firing rates in response to reward-predicting cues
660 (Cohen et al., 2012), and chemogenetic activation of these cells suppress the activity
661 of local dopaminergic neurons (van Zessen et al., 2012), reduces cue-evoked

662 sucrose-seeking responses (Wakabayashi et al., 2019), and induces conditioned
663 place aversion in rodents (Tan et al., 2012). Future studies need to determine
664 whether this regulation of rewarding and aversive responses by VTA^{GABA} neurons can
665 also be attributed to their long-range inhibitory projections to PL neurons, particularly
666 during conflict situations.

667 Differences in risk-taking and risk-avoiding behaviors were also reflected on
668 LFP frequencies in PL neurons in the beginning of the test session, with *Pressers* and
669 *Non-pressers* displaying increased power in the delta or theta bands, respectively.
670 These findings are in corroboration with previous studies showing that increased
671 delta power activity in the mPFC is associated with both reward seeking and
672 preparatory attention (Horst and Laubach, 2013; Totah et al., 2013; Emmons et al.,
673 2016), whereas augmented theta power in the mPFC or synchronized theta activity
674 between mPFC and BLA is correlated with the expression of avoidance responses or
675 the consolidation of threat memories, respectively (Popa et al., 2010; Padilla-Coreano
676 et al., 2019). More specifically, increased synchrony between mPFC and BLA activity
677 in the theta frequency range has been reported for animals that successfully
678 differentiate between aversive and safe cues (or environments) during a differential
679 threat conditioning task (or an open field arena) (Likhtik et al., 2014; Stujenske et al.,
680 2014). In addition, prior studies have shown that 4 Hz LFP oscillations in the mPFC
681 and BLA were strongly synchronized during conditioned freezing episodes (Courtin et
682 al., 2014; Dejean et al., 2016; Karalis et al., 2016), and these sustained 4 Hz
683 oscillations in the mPFC were independent of hippocampal low-theta oscillations,
684 suggesting that they were internally generated in the mPFC during the expression of
685 freezing behavior (Karalis et al., 2016). Consistent with these findings, in our study
686 *Non-pressers* showed increased theta activity and marked 4 Hz oscillations in PL
687 neurons, which were associated with better discrimination between reward and threat

cues and increased freezing responses during the test session, when compared to *Pressers*.

Increased risk-taking behavior in *Pressers* was associated with a higher number of PL^{GLUT} neurons showing reduced spontaneous activity during the conflict phase. In contrast, risk-avoiding responses in *Non-pressers* were associated with increased spontaneous activity during conflict. While this set of results suggest that distinct patterns of PL activity are associated with risk-taking or risk-avoiding behaviors in conflict situations, our optogenetic manipulation provided a causal role for PL^{GLUT} neurons in the regulation of approach-avoidance conflict. For instance, the reduction in food-seeking responses during photoactivation of PL^{GLUT} neurons indicates that increased activity in PL pyramidal cells is sufficient to recapitulate the reward-seeking suppression observed during conflict. Our findings agree with previous studies showing that increased activity in mPFC neurons, including PL, attenuates reward-seeking responses in a neutral context ([Berglind et al., 2007](#); [Chen et al., 2013](#); [Ferenczi et al., 2016](#); but see: [Warthen et al., 2016](#)), an effect that has been attributed, at least in part, to downstream projections to the paraventricular nucleus of the thalamus (PVT) ([Otis et al., 2017](#)). Notably, PVT neurons are necessary for the retrieval of both reward- and threat-associated memories ([for a review see: Do Monte et al., 2016; Millan et al., 2017; McGinty and Otis, 2020; Penzo and Gao, 2021](#)), and activity in PVT neurons has recently been shown to be associated with the regulation of approach-avoidance responses during situations of conflict ([Choi and McNally, 2017](#); [Choi et al., 2019](#); [Engelke et al., 2021](#)), suggesting a potential target by which PL glutamatergic neurons may exert their effects.

Considering that *Pressers* showed a higher number of sustained excitatory food-cue responses than *Non-pressers*, it is counterintuitive that photoactivation of PL^{GLUT} neurons during the food cue onset resulted in reduced food-seeking responses. However, it is important to note that our optogenetic manipulation not only

altered the activity of food-cue responsive neurons, but mostly the global activity of other PL^{GLUT} neurons. Thus, it is possible that increased activity in the firing rate of PL^{GLUT} neurons may result in reduced signal-to-noise ratio during the food cue onset (Kroener et al., 2009; McGinley et al., 2015), and consequently decreased food-seeking responses. In contrast, we speculate that by reducing their spontaneous firing rates during conflict situations, PL^{GLUT} neurons become more likely to fire in response to food cues due to an increase in the signal-to-noise ratio, thereby resulting in persistent reward-seeking responses during the conflict phase as we propose in our schematic in Fig. 8.

Additionally, our findings showing that inactivation of PL^{GLUT} neurons increases food-approaching responses in *Non-pressers* suggest that PL activity is indispensable to inhibit reward pursuit in the presence of threat-associated cues. The lack of effects on lever pressing indicate that other parallel brain regions may be modulating the suppression of operant lever-press responses during conflict. Alternatively, photoinhibition of PL^{GLUT} neurons was not large enough to produce a more global effect on risky behavior (i.e., completely restore lever presses). Collectively, these results add to a growing literature indicating that PL neurons are necessary to guide appropriate food-seeking behavior in tasks that rely on discrimination among environmental cues (Marquis et al., 2007; Sangha et al., 2014; Moorman and Aston-Jones, 2015) or decision-making tasks involving risk of punishment in which animals need to: i) adapt choice behavior during shifts in risk contingencies (Orsini et al., 2018), ii) regulate behavioral flexibility (Radke et al., 2015; Capuzzo and Floresco, 2020), or iii) suppress reward seeking in response to conditioned aversive stimuli (Kim et al., 2017; Piantadosi et al., 2020). Moreover, our results are in accordance with previous findings demonstrating that inactivation of PL neurons, or their inputs from BLA, increases risk-taking behavior in a conflict task in

741 which rats needed to refrain from consuming sucrose to avoid a footshock (Burgos-
742 Robles et al., 2017; Verharen et al., 2019).

743 Previous studies have shown that PL neurons fire in response to shock-paired
744 cues and such activity is highly correlated with the expression of freezing responses
745 (Burgos-Robles et al., 2009; Sotres-Bayon et al., 2012; Kim et al., 2013; Courtin et
746 al., 2014). Adding to these findings, our recordings demonstrated that the activity of a
747 small number of PL neurons changed immediately before or after the onset of
748 freezing responses, with the same proportion of freezing-responsive cells being
749 classified as PL^{GLUT} or PL^{GABA} neurons (~6-14%). At first sight, the lack of effects on
750 freezing behavior following optogenetic activation of PL^{GLUT} neurons seems at odds
751 with our recordings. It also seems to disagree with previous studies showing that
752 electrical stimulation or optogenetic induction of 4Hz oscillations in PL increases
753 conditioned freezing responses (Vidal-Gonzalez et al., 2006; Courtin et al., 2014) by
754 synchronizing the neural activity between PL and BLA regions (Karalis et al., 2016).
755 However, one important difference between our study and others is that
756 photoactivation of PL^{GLUT} neurons in our experiments was performed in naïve rats, in
757 the absence of shock-paired cues. Thus, the increased freezing responses following
758 PL activation reported in previous studies appear to be dependent on the
759 preexistence of a conditioned threat memory.

760 Overall, our results outline the neural correlates of risk-taking and risk-avoiding
761 behaviors in PL and reveal an important role for PL^{GLUT} neurons in coordinating
762 memory-based risky decision-making during conflict situations. Further studies will
763 focus on identifying the PL downstream/upstream circuits that regulate reward-
764 approaching and threat-avoidance responses, as well as the potential genetic and
765 epigenetic factors that could contribute to the observed behavioral phenotypes.
766 Elucidating the underlying mechanisms that mediate risk-taking vs. risk-avoiding
767 responses during situations of uncertainty may help to provide understanding of

response selection and adaptive behaviors, and may have clinical relevance to many psychiatric disorders (Aupperle and Paulus, 2010; Kirlic et al., 2017). Whereas persistent avoidance of presumed threats is the cardinal symptom of anxiety disorders (Treanor and Barry, 2017), seeking reward despite negative consequences is a hallmark of both eating and substance use disorders in humans (Volkow et al., 2012).

ACKNOWLEDGEMENTS

We thank Dr. Roger Janz for helping us with the packaging of the AAV-mDLX-ChR2-mCherry viral construct, Ryia Albert and Sharon Gordon for their technical assistance, and Dr. Robson Scheffer-Teixeira for his support with the statistical analyses. We also thank current and former members of the Do Monte and Quirk Labs for their valuable comments on the manuscript, and the Mind the Graph team for creating the schematic drawings presented in the manuscript. This work was supported by NIH grants R00-MH105549 and R01-MH120136, a Brain & Behavior Research Foundation grant (NARSAD Young Investigator), and a Rising STARS Award from UT System to F.H.D-M.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to F.H.D-M.

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (Rat, male)	<i>Rattus Norvegicus</i>	Charles River	Strain code: 006	Male Long-Evans hooded
Recombinant DNA reagent	AAV-CaMKII α -eNpHR-eYFP	University of North Carolina Viral Vector Core	AAV Stock Vector Karl Deisseroth	3rd Gen Opto Inhibition: eNpHR 3.0
Recombinant DNA reagent	AAV-CaMKII α -hChR2(H134R)-eYFP	University of North Carolina Viral Vector Core	AAV Stock Vector Karl Deisseroth	Opto Excitation: ChR2
Recombinant DNA reagent	AAV-CaMKII α -eYFP	University of North Carolina Viral Vector Core	AAV Stock Vector Karl Deisseroth	Control Fluorophores
Recombinant DNA reagent	pAAV-mDlx-ChR2-mCherry-Fishell-3	Addgene and University of North Carolina Viral Vector Core	Addgene plasmid #83898	Packaged by UNC vector core (Serotype 5)
Antibody	Anti-GAD67 raised in mouse polyclonal	Millipore-Sigma	Cat No. MAB5406B	1:400
Antibody	Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) polyclonal	Abcam	Cat No. ab150113	1:200
Antibody	Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) polyclonal	Abcam	Cat No. ab150116	1:200
Other	Opal 520	Akoya Biosciences	Cat No. FP1487001KT	1:1000 Fluorescent dye
Other	Opal 620	Akoya Biosciences	Cat No. FP1495001KT	1:1000 Fluorescent dye
Sequence based reagent	RNA scope probe mCherry	Advanced Cell Diagnostics	Cat No. 431201-C3	
Sequence based reagent	RNA scope probe vGAT	Advanced Cell Diagnostics	Cat No. 424541	
Commercial Assay or Kit	RNAscope Multiplex Fluorescent Detection Kit v2	Advanced Cell Diagnostics	Cat No. 323110	
Chemical compound, drug	VECTASHIELD® Antifade Mounting Medium with DAPI	Vectorlabs	Cat No. H-1200-10	
Software, algorithm	Matlab	Mathworks	RRID:SCR_001622	https://it.mathworks.com/products/matlab.html
Software, algorithm	NeuroExplorer	Plexon	RRID:SCR_001818	https://plexon.com/products/neuroexplorer/
Software, algorithm	Offline Sorter	Plexon	RRID:SCR_000012	https://plexon.com/products/offline-sorter/
Software, algorithm	CinePlex Behavioral Research System	Plexon		https://plexon.com/plexon-systems/cineplex-behavioral-research-system/
Software, algorithm	GraphPad Prism	GraphPad	RRID:SCR_002798)	https://www.graphpad.com/scientific-software/prism/

799 **Animals.** All experimental procedures were approved by the Center for Laboratory
800 Animal Medicine and Care of The University of Texas Health Science Center at
801 Houston. The National Institutes of Health guidelines for the care and use of
802 laboratory animals were strictly followed to minimize any potential discomfort and
803 suffering of the animals. Male Long-Evans hooded adult rats (Charles Rivers
804 Laboratories) with 3-5 months of age and weighing 300-450 g at the time of the
805 experiment were used. Rats were single-housed and after a 3-day acclimation period
806 handled and trained to press a lever for sucrose as described below. Animals were
807 kept in a 12-hour light/12-hour dark cycle (light from 7:00 to 19:00) and maintained on
808 a restricted diet of 18 g of standard laboratory rat chow provided daily at the end of
809 experimentation. Animals were given *ad libitum* access to water. Animals' weights
810 were monitored weekly to ensure all animals maintained their weight under food
811 restriction. During pre- and post- surgery phases, animals were given *ad libitum*
812 access to food for a total of 7 days.

813 **Surgeries.** Rats were anaesthetized with 5% isoflurane in an induction chamber.
814 Animals were positioned in a stereotaxic frame (Kopf Instruments) and anesthesia
815 was maintained with 2.5% isoflurane delivered through a facemask. A heating pad
816 was positioned below the body of the animal and both temperature and respiration
817 were monitored during the entire surgery. Veterinary lubricant ointment was applied
818 on the eyes to avoid dryness during the surgery. Animals received a subcutaneous
819 injection of the local anesthetic bupivacaine (0.25%, 0.3 ml) at the incision site. Iodine
820 and ethanol (70%) were alternately applied for asepsis of the incision site. The
821 surgery procedures varied according to the type of implantation/injection (see below).
822 For injection-only surgeries, the incision was stitched after the injection by using
823 surgical suture (Nylon, 3-0). For implantation surgeries, the implants were fixed to the
824 skull using C&B metabond (Parkell), ortho acrylic cement, and four to six anchoring

screws. After surgery, animals received a subcutaneous injection of meloxicam (1 mg/Kg) and a topical triple antibiotic was applied to the incision area.

Viral vector injection. Viral injections were performed using a microsyringe (SGE, 0.5 μ l) with an injection rate of 0.04 μ l/min plus an additional waiting time of 12 min to avoid back-flow. The adeno-associated virus (AAV) was bilaterally injected at a volume of 0.4 μ l per side. The AAV-CaMKII α -eNpHR-eYFP vector was used to inhibit glutamatergic neurons, whereas AAV-mDlx-ChR2-mCherry or AAV-CaMKII α -ChR2-eYFP vectors were used to activate either GABAergic or glutamatergic neurons, respectively. The use of mDlx or CaMKII α promoters enabled transgene expression favoring either GABAergic or glutamatergic neurons, as previously shown ([Gradinaru et al., 2009](#); [Tye et al., 2011](#); [Dimidschstein et al., 2016](#); [Sun et al., 2020](#)) and was confirmed by our immunohistochemical and RNAscope assessment (**Figure 5 - Figure supplement 1**). The viral construct AAV-CaMKII α -eYFP was used to control for any nonspecific effects of viral infection or laser heating. All plasmid or viral vectors were obtained from Addgene or University of North Carolina Viral Vector Core. For implantation of optrodes, the following coordinates from bregma were used for virus injection: PL, +2.7 mm AP, $\pm$ 0.7 mm ML, -3.8 mm DV at a 0° angle. For PL soma illumination, an optical fiber (0.39 NA, 200 nm core, Inper) was implanted in each hemisphere targeting PL neurons using the following coordinates from bregma: +2.7 mm AP; $\pm$ 1.5 mm ML; -4.0 mm DV at a 15° angle.

Single-unit electrodes. An array of 16 or 32 microwires was unilaterally implanted targeting the PL using the following coordinates from bregma: +2.7 mm AP, $\pm$ 0.8 mm ML, -3.9 mm DV. Three different electrode configurations were used: i) 32-channel silicon probes (Buzsaki32-CM32 or A1x32-5mm-25-177-CM32, Neuro Nexus Technologies, USA), ii) Micro-Wire Arrays (MWA) of 16 or 32 channels (Bio-Signal Technologies Ltd, USA); or iii) custom designed electrodes with 2x8 grid with 150 μ m

of space between wires, 200 μ of space between rows, with 35 μ m diameter wires (Innovative Neurophysiology Inc., USA). For photoidentification of GABAergic or glutamatergic neurons, a Hermes 32 channels optrode array was used (200 nm core, Bio-Signal Technologies Ltd). Optrodes were unilaterally implanted at the same coordinates described above after the infusion of 0.6 μ l of AAV-mDlx-ChR2-mCherry or AAV-CaMKII α -ChR2-eYFP vectors. In all cases, the ground wire was wrapped around a grounding screw previously anchored into the skull. Two insulated metal hooks were implanted bilaterally into the cement to allow firmly attachment of the array connector to the cable during recording.

Odor preparation. A 99% amyl acetate solution (Sigma Aldrich) was diluted in propylene glycol (Bluewater Chemgroup, Inc) to a 10% solution and presented to the rats during the different stages and phases of the olfactory threat conditioning test. A customized olfactometer (Med Associates) was used to control the flow of air into the animal's chamber. Before being mixed with the 10% amyl acetate solution, the air was passed through a desiccant and a charcoal filter to remove any moisture and odors, and was finally rehydrated with distilled water before being delivered into the chamber through a thermoplastic PVC-based tube (Tygon) attached to an odor port located in the odor area.

Behavioral Tasks

Lever-press training. Rats were placed in a plexiglass, standard operant box (34 cm high x 25 cm wide x 23 cm deep, Med Associates), and trained to press a lever for sucrose on a fixed ratio of one pellet for each press. Next, animals were trained in a variable interval schedule of reinforcement that was gradually reduced across the days (one pellet every 15 s, 30 s, or 60 s) until the animals reached a minimum criterion of 10 presses/min. All sessions lasted 30 min and were performed on consecutive days. Sucrose pellet delivery, variable intervals, and session duration

were controlled by an automated system (ANY-maze, Stoelting). Lever-press training lasted approximately one week, after which animals were assigned to surgery or cued food-seeking training. A small number of rats failed to reach the lever press criteria and were excluded from the experiments (< 3%).

Cued food-seeking training. Rats previously trained to press a lever for sucrose were trained to learn that each lever press in the presence of an audiovisual cue (tone: 3 KHz, 75 dB; light: yellow, 2.8 W; 30 s duration) resulted in the delivery of a sucrose pellet into a nearby dish. Reward cue conditioning also took place in the standard operant boxes. While the light cue helps to direct the animals toward the lever during the beginning of the training phase, the tone assures that the animals will not miss the presentation during the trial and provides the temporal precision required for single-unit recordings. After ~4 consecutive days of training (24 trials per day, pseudorandom intertrial interval of ~120 s, 60 min session), rats learned to discriminate the food-associated cue as indicated by a significant increase in press rate during the presence of the audiovisual cues, when compared to the 30 s immediately before the cue onset (cue-off, see **Figure 1 - Figure supplement 1A**). The cued food-seeking training was completed when animals reached 50% of discriminability index (presses during cue-on period minus presses during cue-off period divided by the total number of presses).

After the cued food-seeking training was completed, rats with single-unit electrodes were exposed to an additional training session in which the audiovisual cue ceased immediately after the animals pressed the lever and a single sucrose pellet was delivered into the dish. This extra training reduced the rat's response to a single press and dish entry per cue, thereby enabling us to correlate each food-seeking event with the neuronal firing rate by avoiding overlapping between consecutive events (e.g., lever presses). The single-pellet training took place in the same plexiglass rectangular arena subsequently used for the odor test (40 cm high x

905 60 cm wide x 26 cm deep, Med Associates, see schematic in **Fig. 1A right**). The
906 arena consisted of a hidden area (40 cm high x 20 cm wide x 26 cm deep) separated
907 from an open area by a plexiglass division. An 8-cm slot located in the center of the
908 division enabled the animal to transition between both sides of the arena. For
909 behavioral quantification, the open area was subdivided into a center area and a food
910 area (40 cm high x 12 cm wide x 26 cm deep), the latter containing a lever, a dish,
911 and an external feeder similar to the food-seeking operant box.

912 Habituation day. Animals were placed in the odor arena and exposed to 12
913 audiovisual cues (30 s duration, pseudorandom inter-trial intervals of between 25-40
914 s) followed by 10 min of presentation to the neutral odor alone (10% amyl acetate)
915 and an additional 12 audiovisual cues similar to the first cues but in the presence of
916 the neutral odor delivered in the food area of the arena. Each lever press in the
917 presence of the audiovisual cue resulted in the delivery of a sucrose pellet into the
918 dish, and the audiovisual cue ceased immediately after the animal pressed the lever.

919 Threat conditioning day. One day after the habituation day, rats were placed in a
920 plexiglass, standard operant box similar to the cued food-seeking training box, but
921 with the grid floor previously attached to a shock generator system. Rats were
922 habituated to one non-reinforced odor presentation (10% amyl acetate, 30 s duration)
923 followed by five odor presentations that coterminated with a foot shock (0.7 mA, 1 s
924 duration, 258-318 s inter-trial intervals). An olfactometer system was used to deliver
925 the odor into the box (see odor preparation session above), whereas an exhaustor
926 system was used to remove it during the intertrial intervals. Between each odor
927 presentation, audiovisual cues (30 s duration) signaling the availability of sucrose
928 were presented to the animals. Each lever press during the audiovisual cues resulted
929 in the delivery of a sucrose pellet into the dish. Shock grids and floor trays were
930 cleaned with 70% ethanol between each rat. No rats were excluded from the
931 analyses due to distinct levels of freezing following the threat conditioning session.

Test day. One day after the threat conditioning session, rats were returned to the same arena used during the habituation and exposed to the exact same protocol. The first phase of the test session was called *reward phase* (12 min) and the animals were exposed to 12 food cues. The second phase was called *odor phase* and the animals were exposed to 10 min of conditioned odor (10% amyl acetate) alone. The last phase was called *conflict phase* (12 min) and the animals were exposed to 12 food cues in the presence of the conditioned odor. An odor dispersion sensor (200B miniPID, Aurora Scientific) revealed that the odor took approximately 2.21 ± 0.28 s to reach detectable concentrations (56 particles per billion; Punter, 1983) in the arena after the olfactometer onset, and approximately 19.59 ± 0.97 s to be completely removed from the arena after the olfactometer offset and concomitant activation of the exhaustor fan. Due to the low temporal resolution to control the delivery of the odor in the arena, the odor was maintained constant during the entire duration of the odor and conflict phases. In order to press the lever for sucrose during the conflict phase, rats had to approach the conditioned odor presented in the food area. After the end of the conflict phase, the odor was extracted from the arena with the exhaustor fan and the floor and walls of the arena were cleaned thoroughly with 70% ethanol solution.

Behavioral tracking. Both the standard operant boxes and the testing arenas were equipped with video cameras and a behavior tracking software (ANY-maze, Stoelting) which were used to record the animal's behavior and control the delivery of sucrose, foot shock, tone, light and odor in the apparatuses. Avoidance responses were characterized by the time spent in the hidden area of the arena. Freezing responses were characterized by the complete absence of movements except those needed for respiration. Risk-assessment responses were characterized by a body stretching movement to peep out toward the food area while in the hidden area and were used as a measure of risk-assessment behavior (Blanchard et al., 2011).

959 For single-unit recording analyses, the detection of freezing, avoidance and
960 risk assessment behaviors were performed using the open source tool DeepLabCut,
961 a machine learning software that tracks complex patterns of behavior from videos
962 ([Mathis et al., 2018](#)). After a video has been analyzed, the data was saved to a .csv
963 file that contained the x and y location of each rat's body part in pixels, as well as the
964 analysis of the expected accuracy (i.e., likelihood) of the tracked positions across
965 time. After DeepLabCut has calculated the positions and the likelihood, we used three
966 different Python codes to identify each one of the three behaviors. For freezing
967 behavior, the code used DeepLabCut's position data and determined if the rat was
968 still for more than 500 ms. The animal was considered to be still if the position in
969 question was within 1.05 pixels of each other. For avoidance behavior, the code used
970 DeepLabCut's position data to determine the location of the rat in the arena and
971 based on the center of its head to identify when the animals entered the hidden area
972 of the arena. Finally, for risk assessment behavior, the code used DeepLabCut's
973 position data to identify the nose, ears, center of the head, and spine to determine
974 whether the rat was located in the hidden area of the arena with its body stretched
975 and the head looking through the open division of the apparatus. Each of these codes
976 generated a .xlsx file that contained the onset and the total duration of each
977 behavioral episode (see single-unit analyses section for more details).

978 **Optogenetic stimulation during behavior.** Bilateral optical cables (200 μ m core,
979 0.37 NA, 2.5 mm ceramic ferrule, Inper) were connected to a blue laser (diode-
980 pumped solid-state, 473 nm, 150 mW output, OptoEngine) or a yellow laser (diode-
981 pumped solid-state, 593.5 nm, 300 mW output, OptoEngine) by using a patch cord
982 (200 μ m, 0.39 NA, FC/PC connector, Inper) through a dual rotary joint (200 μ m core,
983 Doric lenses). During the stimulation, the optical cables were coupled to the
984 previously implanted optical fibers by using a ceramic sleeve (2.5 mm, Precision
985 Fiber Products). An optogenetic interface (Ami-2, Stoelting) and an electrical

stimulator (Master 9, A.M.P. Instruments) were used to control the onset of the laser, pulse width, train duration, and frequency. The power density estimated at the tip of the optical fiber was 7-10 mW for illumination of PL somata (PM-100D, Power Energy Meter, Thor Labs).

Single-unit recording. A 64-channels neuronal data acquisition system (Omniplex, Plexon) integrated with a high-resolution video-tracking system (Cineplex, Plexon) was used for electrophysiological recordings from freely behaving animals. Both videos and neuronal recordings were combined within the same file, thereby facilitating the correlation of behavior with neuronal activity. An electrical isolation, Faraday cage was made and connected to the grounding port of the data acquisition system. The system was connected to the head-mounted electrode/optrode by using a digital headstage cable (32 channels, Plexon), a motorized carrousel commutator (Plexon), and a digital headstage processor (Plexon). Rats were habituated to the headstage cable daily for approximately one week before the beginning of the experiments. Extracellular waveforms exceeding a voltage threshold were band-pass filtered (500 - 5,000 Hz), digitized at 40 KHz, and stored onto disk. Automated processing was performed using a valley-seeking scan algorithm and then visually evaluated using sort quality metrics (Offline Sorter, Plexon, see single-unit analyses section below).

Photoidentification of PL neurons during recordings. During neuronal photoidentification, we recorded from rats expressing channelrhodopsin (ChR2) in PL neurons previously implanted with an optrode in the same region. An optical cable connected to a blue laser was attached to the headstage cable and coupled to the previously implanted optical fiber by using a ceramic sleeve. At the end of the behavioral session, 10 trains of 10-s blue laser pulses (5 ms pulse width, 5 Hz) were delivered by a Master-9 programmable pulse stimulator, which also sent flags to the data acquisition system to mark the time of the laser events.

Neurons were considered to be responsive to photoactivation if they showed a significant increase in firing rate above baseline (20 ms, Z-score > 3.29, $p < 0.001$) and higher reliability within the 6 ms after laser, similar to previous studies (Lima et al., 2009; Pi et al., 2013; Burgos-Robles et al., 2017; Engelke et al., 2021). To identify the threshold separation for the frequency distribution of response latencies to laser illumination, we implemented the Triangle Method detection (Zack et al., 1977). This calculation is particularly effective for left-skewed distributions as in our sample. We considered bins with response latency values from 0 to 12 ms and excluded those with larger values as they would most likely reflect indirect stimulation via collateral activity. We computed the distance normal to the line along with the minimum and maximum values in the histogram. The threshold was defined as the maximum distance between the histogram and the line (i.e. a normalized level within 0 and 1), which in our analysis resulted in 5.8 ms (rounded to 6 ms bin). In addition, to measure the reliability of neural responses to photoactivation, we calculated the Fano Factor (FF), defined as the variance-to-mean ratio of spike counts (Churchland et al., 2011), to characterize the variability of neuronal responses 6 ms before (FF before) and 6 ms after (FF after) the laser pulses for each train of illumination (10 trains of 50 laser pulses, 200 ms pulse interval). When the variance in the counts equals the mean count, FF was equal to 1. Afterwards, we computed the 'overall FF ratio' between the 'FF after' divided by the 'FF before' to compute the reliability of each cell to the laser onsets. Only neurons showing an overall FF higher than 1, which indicates reliable responses to laser illumination compared to baseline, were included as photoidentified cells.

A small number of laser-generated photoelectric artifacts (~10% of the channels in less than 10% of the rats) were observed during the photoactivation. However, they were easily distinguished from the action potentials by their descending voltage signals of high amplitude, pulse shapes distinct from the regular

waveforms, isolated spatial distribution in the principal component analyses, and occurrence restricted to the period of laser activation, which resulted in lack of activity during the behavioral session.

Optogenetic manipulation of PL neurons during behavior. During the cued food-seeking test, rats expressing ChR2 or eNpHR in PL were bilaterally illuminated in the same region by using a blue (5 ms pulse width, 5 Hz for CaMKII α or 20 Hz for mDlx) or a yellow laser (constant illumination), respectively. The laser was activated at cue onset and persisted throughout the entire 30 s of the audiovisual cue presentation.

Rats were exposed to two consecutive cues with laser off followed by two consecutive cues with laser on in a total of 12 cues (pseudorandom inter-trial intervals of between 25-40 s). To assess the effects of PL illumination on rat's defensive behavior, PL neurons of rats expressing ChR2 or eNpHR were bilaterally illuminated during 6 distinct epochs of 30 s during the *odor phase* by using a blue (5 ms pulse width, 20 Hz) or a yellow laser (constant illumination), respectively. To assess the effects of PL illumination on food-seeking responses during the conflict phase, rats were exposed to two consecutive cues with laser off followed by two consecutive cues with laser on in a total of 12 cues (pseudorandom inter-trial intervals of between 25-40 s). The laser was activated at cue onset and persisted on until the animal pressed the bar or the 30 s of the audiovisual cue was completed.

Histology. Animals were transcardially perfused with KPBS followed by 10% buffered formalin. Brains were processed for histology as previously described ([Domonte et al., 2013](#)). At the end of the recording sessions, a micro-lesion was made by passing anodal current (0.3 mA for 15 s) through the active wires to deposit iron in the tissue. After perfusion, brains were extracted from the skull and stored in a 30% sucrose/ 6% ferrocyanide solution to stain the iron deposits. Only rats with the presence of eYFP or mCherry labeling and the track of the electrode wires or optical fiber tips located exclusively in PL were included in the statistical analyses.

Immunohistochemistry. Rats previously infused with AAV-mDlx-ChR2-mCherry or AAV-CaMKII α -ChR2-eYFP were transcardially perfused with 300 ml of KPBS followed by 500 ml of 4% paraformaldehyde. Brains were removed from the skull, transferred to a 20% sucrose solution in KPBS for 24 h, and stored in a 30% sucrose solution in KPBS for another 24 h. Next, coronal PL sections (40 μ m thick) were cut in a cryostat (CM 1860, Leica), blocked in 20% normal goat serum and 0.3% Triton X-100 in KPBS at room temperature for 1 h. For identification of GABAergic neurons, PL sections were incubated with anti-GAD67 serum raised in mouse (1:400; Millipore-Sigma) at 4°C for 48 h. After sections were washed in KPBS for 5 times, sections were incubated with a secondary anti-mouse antibody (1:200, Alexa Fluor 488 or Alexa Fluor 594, Abcam) for 2 h. Sections were washed with KPBS, mounted in Superfrost Plus slides, and coverslipped with anti-fading mounting medium (Vectashield, Vectorlabs). Images were generated by using a microscope (Nikon, Eclipse NiE Fully Motorized Upright Microscope) equipped with a fluorescent lamp (X-Cite, 120 LED) and a digital camera (Andor Zyla 4.2 PLUS sCMOS).

***In situ* hybridization.** Single molecule fluorescent *in situ* hybridization (RNAscope Multiplex Fluorescent Detection Kit v2, Advanced Cell Diagnostics) was used following the manufacturer protocol for fixed-frozen brains sample. Brain samples were sectioned at a thickness of 20 μ m in a cryostat (CM1860, Leica). Sections were collected onto superfrost plus slides (Fisher Scientific) and transferred to a -80°C freezer. To prepare for the assay, brain sections were serially dehydrated with EtOH (50%, 75%, and 100%, each for 5 min) and then incubated in hydrogen peroxide for 10 min. Target retrieval was performed with RNAscope target retrieval reagents at 99°C for 5 min. The sections were then pretreated with Protease III (RNAscope) for 40 min at 40°C. RNAscope probes (Advanced Cell Diagnostics) for mCherry (Cat No. 431201-C3) and vGAT (Cat No. 424541) were hybridized at 40°C for 2h, serially amplified, and revealed with horseradish peroxidase, Opal Dye/TSA Plus fluorophore

(Akoya Biosciences), and horseradish peroxidase blocker. Sections were covered with anti-fading mounting medium with DAPI (Vectashield, Vectorlabs) and kept in the refrigerator. Images were generated by using an epifluorescent microscope (Nikon, Eclipse NiE Fully Motorized Upright Microscope) equipped with a fluorescent lamp (X-Cite, 120 LED) and a digital camera (Andor Zyla 4.2 PLUS sCMOS). Expression of mCherry mRNA (red, Opal 620) and GAD67 mRNA (green, Opal 520) was determined by using an automated fluorescent threshold detector (NIS-Elements). Colabeled cells were manually counted by an experimenter by measuring either the percentage of mCherry positive neurons in PL that were also labeled with GAD67, or the percentage of GAD67 positive neurons in PL that were also labeled with mCherry.

Data analyses.

Behavioral quantification and statistical analysis. Rats were recorded with digital video cameras (Logitech C920) and behavioral responses were measured by using an automated video-tracking system (ANY-maze) or machine learning (DeepLabCut). Presses per minute were calculated by measuring the number of presses during the 30 s cue multiplied by two. All graphics and numerical values reported in the figures are presented as mean $\pm$ s.e.m. Given that the different phases of the test have different duration, we have normalized the data in percentage to be able to compare the behavior of the animals across the different phases of the test. Shapiro Wilk normality test was performed before all the statistical analyses to determine parametrical or non-parametrical statistical tests, and reported in the Results section when the values showed non-normal distribution. For normal data, statistical significance was determined with paired Student's t test, Welch's t test, repeated-measures ANOVA followed by Bonferroni post-hoc comparisons (Prism 7), and Z-test or Fisher Exact Test, as indicated in the Results section, figure legends, and Table 1S. For non-parametric data, Wilcoxon test (paired groups) or Mann-Whitney test

(unpaired groups) were performed. Effects sizes for pairwise comparisons or ANOVA were calculated by using Cohen's d or partial eta squared, respectively (Cohen, 1988; Richardson, 2011). Effect sizes < 0.2 were considered small, effects sizes between 0.2 and 0.8 were considered medium, and effect sizes > 0.8 were considered large (Cohen, 1988). Sample size (n) was based on estimations by power analysis with a level of significance of 0.05 and an expected effect size of 0.8.

Principal component analyses on behaviors. The principal component analysis (PCA) method (Jolliffe, 2002) was implemented to further understand the relationships between approaching, avoidance, freezing, and latency to press during the conflict phase for both *Pressers* and *Non-pressers*. This method was used to reduce the dimensionality of our multivariate data whilst preserving as much of the relevant information as possible. Briefly, we performed a linear transformation of the data to a new coordinate system such that the new set of variables, the principal components, were linear functions of the original variables. These variables were also uncorrelated, and the greatest variance by any projection of the data came to lie on the first coordinate, the second greatest variance on the second coordinate, and so on. Afterwards, we analyzed the explained variance (%) of the first two PCs (i.e. PC1 and PC2) together with the relevance of each analyzed behavior to define which variables better explained the differences between *Pressers* and *Non-pressers* (see Results section for additional details).

Single-unit analyses. Single-units were selected based on three principal components and waveform features such as valley-to-peak and amplitude measurements. The principal-component scores for unsorted waveforms were computed and plotted in a three (or two) dimensional principal-component space. Clusters containing similar valid waveforms were manually defined. After manually clustering similar valid waveforms, a group of spikes were considered from a single neuron if the waveforms formed a discrete, isolated, cluster in the principal-component space. A Commercial

software (NeuroExplorer, NEXT Technologies) combined with Matlab (MathWorks) scripts, and Python scripting were implemented to calculate the spontaneous firing rate, changes in neural activity in response to food cues, lever presses, food dish entries, as well as the neural correlates of freezing, avoidance, and risk-assessment behaviors. The spontaneous firing rate was calculated by comparing the frequency of spike trains during the last 30 s of the food-seeking phase, odor phase, or conflict phase against the 30 s prior to the beginning of each session. Food cue, lever press, and food dish responses were calculated by implementing Matlab scripts as Z-scores normalized to 20 pre-cue bins of 300 ms. Neurons showing a Z-score > 2.58 ($p < 0.01$) during the first two-bins following the onset of the aligned event were classified as excitatory responses, whereas neurons showing a Z-score < -1.96 ($p < 0.05$) during the same first two-bins were classified as inhibitory responses. A temporal frequency separation was used to classify the food cue responses according to the pattern of activity, similar to a previous study ([Bezudnaya et al., 2006](#)). Neurons showing a transient increase in firing rate (< 600 ms duration) were classified as transient activity, whereas neurons showing a sustained increase in firing rate (≥ 900 ms duration) were classified as sustained activity (Z-score > 2.58 during the first 3 seconds after food cue onset).

To analyze freezing, avoidance, and risk-assessment responses, the time onsets for each behavior were filtered by selecting only the events that lasted more than 1 second and were not preceded by the same behavior during the previous 6 s (baseline). The final list of time onsets was entered into the single-unit recording files to create the events and temporally align them with the neuronal recordings. To increase the number of events during our analyses, we combined the behavioral responses emitted during the odor and conflict phases. Only animals exhibiting at least 6 events for each behavior were included. We used an interval criterium of 600 ms to select the neurons that responded close in time to the onset of the analyzed

1175 behavior (before or after), thereby avoiding potential neural activity contamination
1176 caused by other types of behavioral responses.

1177 Data availability

1178 All the data that support the findings presented in this study are available in the
1179 source data file in the supplementary material.

1180

1182 **Figure 1. Rats show individual variability in reward-seeking responses during an**
 1183 **approach-avoidance conflict test. (A)** Schematic and timeline of the approach-avoidance
 1184 conflict test. **(B-D)** Rats exhibited an increase in the percentage of time freezing (Shapiro-
 1185 Wilk normality test, $p < 0.001$, Friedman Test, Friedman statistic = 84.08, $p < 0.001$, Dunn's
 1186 *post-hoc*, $p < 0.001$) and a reduction in lever presses (Shapiro-Wilk normality test, $p <$
 1187 0.001 , Friedman Test, Friedman statistic = 35.11, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$)
 1188 with a higher latency to press the lever (Shapiro-Wilk normality test, $p < 0.001$, Friedman
 1189 Test, Friedman statistic = 29.45, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$) during the olfactory
 1190 threat conditioning session on day 1 ($n = 32$), when compared to before the shock. **(E-G)**
 1191 Patterns of defensive responses and food seeking during the different phases (reward,
 1192 odor, conflict) of the test session on day 2. Rats showed an increase in defensive
 1193 responses characterized by an augment in the percentage of time exhibiting (E) freezing
 1194 (Shapiro-Wilk normality test, $p < 0.05$, Friedman Test, Friedman statistic = 40.46, $p < 0.001$,
 1195 Dunn's *post-hoc*, $p < 0.001$), (F) avoidance (Shapiro-Wilk normality test, $p < 0.05$, Friedman
 1196 Test, Friedman statistic = 31.67, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$) and (G) risk-
 1197 assessment (Shapiro-Wilk normality test, $p < 0.05$, Friedman Test, Friedman statistic =
 1198 29.86, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$) ; and a decrease in the (H) percentage of
 1199 time spent in the food area (Shapiro-Wilk normality test, $p < 0.05$, Friedman Test, Friedman
 1200 statistic = 32.19, $p < 0.001$, Dunn's *post-hoc*, $p < 0.001$) during the odor presentation,
 1201 when compared to the reward phase. Rats' defensive responses were significantly
 1202 attenuated during the conflict phase as evidenced by a reduction in the percentage of time
 1203 (F) avoiding the odor ($p = 0.0031$) and an increase in the percentage of time (H)
 1204 approaching the food area ($p < 0.001$) , when compared to the odor phase. **(I-J)** Two
 1205 different behavioral phenotypes emerged during the conflict phase: rats that continued to
 1206 press the lever (*Pressers*, green circle, $n = 25$) and rats that showed a complete
 1207 suppression in lever pressing (*Non-pressers*, purple circle, $n = 7$). Rewarded presses were
 1208 calculated as the percentage of the 12 cue trials in which rats pressed the lever. **(K-L)**
 1209 Representative tracks and heatmaps of time spent in each compartment of the arena for a
 1210 (K) *Presser* or a (L) *Non-pressers* rat during the test session. **(M-R)** Patterns of defensive
 1211 responses and food seeking during the different phases (reward, odor, conflict) of the test
 1212 session on day 2 after separating the animals into *Pressers* and *Non-pressers*. When
 1213 compared to *Non-pressers*, *Pressers* showed reduced defensive responses characterized
 1214 by an attenuation in the percentage of time exhibiting (M) freezing ($F_{(2, 60)} = 29.54$, $p <$
 1215 0.001) and (N) avoidance responses ($F_{(2, 60)} = 23.27$, $p < 0.001$), and an augment in the
 1216 percentage of time (P) approaching the food area ($F_{(2, 60)} = 22.49$, $p < 0.001$) during both
 1217 the odor and the conflict phases (Bonferroni post-hoc test - odor phase, $p = 0.0453$; conflict
 1218 phase, $p < 0.001$). (Q) *Non-pressers* showed increased latency to press the lever during
 1219 the conflict phase when compared to the reward phase or to *Pressers* in the same phase
 1220 ($F_{(1, 30)} = 55.14$, $p < 0.001$, Bonferroni post-hoc test – all p 's < 0.001). (R) The percentage of
 1221 rewarded trials was used as a binary criterium for group classification. Data shown as
 1222 mean $\pm$ SEM. One-way or two-way ANOVA repeated measures followed by Bonferroni
 1223 *post-hoc* test, all * p 's < 0.05 compared to the same group during the reward phase, all &
 1224 p 's < 0.05 compared to the same group during the odor phase, all \$ p 's < 0.05 compared to
 1225 *Pressers* during the same phase. All statistical analysis details are presented in table S1.
 1226 See also Figure 1 - Figure supplement 1 and Video 1.

Figure 2. PL neurons respond differently to reward cues in *Pressers* vs. *Non-Pressers* during conflict (A) Diagram of the electrode placements in PL. (B) Schematic and timeline of PL recordings for food cue responses during of the approach-avoidance conflict test (12 food cues per phase). (C) Pie charts showing changes in PL firing rate in response to food cues during reward (left) vs. conflict (right) phases for *Pressers* (n = 237 neurons from 25 rats, Fisher Exact Test, responsive during reward phase: n = 79, responsive during conflict phase: n = 32, $p < 0.001$; excited during reward phase: n = 48, excited during conflict phase: n = 25, $p = 0.0049$; inhibited during reward phase: n = 31, inhibited during conflict phase: n = 7, $p < 0.001$). (D) Heatmap of Z-scored neural activities for PL neurons selected during reward phase and tracked to conflict phase. (E) Average peri-stimulus time histograms (PSTHs) for all PL neurons showing excitatory food cue responses (Z-score > 2.58 , dotted line) during reward (red line) compared to the same cells during conflict (gray line). E, Inset: Differences in the positive area under the curve (AUC) between the two phases (Shapiro-Wilk normality test, $p < 0.001$; Wilcoxon test, $W = -824$, excitatory responses reward phase vs. conflict phase, $p < 0.001$). (F) Pie charts showing the percentage of sustained vs. transient excitatory food-cue responses in PL neurons during the reward phase with the same neurons tracked during the conflict phase. (G-H) Representative PSTHs for a PL neuron showing excitatory responses to food cues during the (G) reward phase vs. the same neuron during the (H) conflict phase. (I) Average PSTHs for all PL neurons showing inhibitory food cue responses (Z-score < -1.96 , dotted line) during reward (blue line) compared to the same cells during conflict (gray line). I, Inset: Differences in the negative AUC between the two phases (Shapiro-Wilk normality test, $p < 0.001$; Wilcoxon test, $W = 367$, inhibitory responses reward phase vs. conflict phase, $p < 0.001$). (J) Pie charts showing the percentage of sustained vs. transient inhibitory food-cue responses in PL neurons during the reward phase with the same neurons tracked during the conflict phase. (K-L) Representative PSTHs for a PL neuron showing inhibitory responses to food cues during the reward phase (K) vs. the same neuron during the conflict phase (L). (M) Pie charts showing changes in PL firing rate in response to food cues during reward (left) vs. conflict (right) phases for *Non-pressers* (n= 89 neurons from 7 rats; Fisher Exact Test, responsive during reward phase: n = 19, responsive during conflict phase: n = 6, $p < 0.0086$; excited during reward phase: n = 10, excited during conflict phase: n = 4, $p = 0.162$; inhibited during reward phase: n = 9, inhibited during conflict phase: n = 2, $p = 0.057$). (N-O) Same as D-E, but for *Non-pressers*. O, Inset: Differences in the positive AUC between the two phases (Paired Student's t-test, $t = 2.34$, $p = 0.043$). (P-S) Same as F-I, but for *Non-pressers*. S, Inset: Differences in the negative AUC between the two phases (Paired Student's t-test, $t = 0.59$, $p = 0.569$) (T-V) Same as J-L, but for *Non-pressers*. The threshold used to identify significant differences per neurons was Z-score > 2.58 for excitation and Z-score < -1.96 for inhibition. Legend: cc, corpus callosum, CG1, anterior cingulate cortex; IL, infralimbic cortex. All statistical analysis details are presented in table S1. See also Supplementary Figs. 2-5.

Figure 3. PL activity correlates with the onset of freezing, avoidance or risk-assessment behaviors in both *Pressers* and *Non-pressers*. (A-C) Both *Pressers* and *Non-pressers* showed the same number and proportion of excitatory and inhibitory PL responses during the onset of (A) freezing (Fisher Exact Test, responsive neurons in *Pressers*: 22 neurons, in *Non-pressers*: 15 neurons, $p = 0.462$), (B) avoidance (Fisher Exact Test, responsive neurons in *Pressers*: 43 neurons, in *Non-pressers*: 9 neurons, $p = 0.999$) or (C) risk-assessment (RA, Fisher Exact Test, responsive neurons in *Pressers*: 12 neurons, in *Non-pressers*: 6 neurons, $p = 0.318$) behaviors. (D-F) Representative PSTHs for distinct PL neurons showing excitatory responses at the onset of (D) freezing, (E) avoidance or (F) risk-assessment behaviors. (G-I) Representative PSTHs for distinct PL neurons showing inhibitory responses at the onset of freezing (G), avoidance (H) or risk-assessment (I) behaviors. (J) Venn Diagram showing the percentage of all PL responsive neurons (29 out of 88 neurons) by behavior. Most of the responsive neurons responded selectively at the onset of one of the behaviors. (K-M) Heatmap of Z-scored neural activities for PL neurons selected at the onset of freezing (K), avoidance (L) or risk-assessment behavior (M) with the same cells tracked during the other behaviors. The threshold used to identify significant differences per neurons was Z-score > 2.58 for excitation and Z-score < -1.96 for inhibition. n.s. = non-significant. All statistical analysis details are presented in table S1. See also Figure 3 - Figure supplement 1.

Figure 4. Pressers and Non-pressers show significant differences in PL oscillations during the test session. (A) Power spectral density (PSD) contribution at different frequency bands. **(B-C)** Average of PSD (%) in the (B) delta (0-4 Hz) or (C) theta (4-10 Hz) bands in *Pressers* (green line, n = 25 rats) and *Non-pressers* (purple line, n = 7 rats) during the (left) reward, (center) odor, and (right) conflict phases of the test session. *Pressers* showed increased power in the delta band, whereas *Non-pressers* showed increased power in the theta band during the three phases of the test session (Unpaired Student's t-test comparing *Pressers* vs. *Non-pressers*, all p's < 0.001) . **(D-E)** Representative time-frequency spectrogram showing changes in the log of PSD (dB) for delta and theta bands in (D) *Pressers* and (E) *Non-pressers* across the different phases of the session. All statistical analysis details are presented in table S1.

Figure 5. In pressers, PL^{GLUT} neurons show reduced spontaneous activity during the conflict phase. (A) Top, Schematic of viral infusion. Bottom, Representative immunohistochemical micrograph showing lack of colabeling (white arrows) between the ChR2 viral construct (green, AAV-CaMKII-ChR2-eYFP) and the GABA marker GAD67 (red), confirming that the use of a CaMKII promoter enables transgene expression favoring PL glutamatergic neurons. Scale bars: 25 μ m. **(B-D)** Photoidentification of PL^{GLUT} neurons. (B) Frequency histogram showing the latency of response to laser illumination for PL neurons (n = 36 photoidentified PL^{GLUT} neurons out of 104 recorded cells). Triangle method detection of cluster distribution revealed a separation of latency frequencies at 6 ms (see details in Methods section). (C) Cells with photoresponse latencies < 6 ms (identified as the first bin with Z-score > 3.29, p < 0.001, red dotted line) and high spike reliability during the 6 ms (FF, Fano Factor ratio > 1 compared to pre-laser baseline) were classified as PL^{GLUT} neurons (see details in Methods section). (D) Raster plot and peristimulus time histogram showing a representative PL^{GLUT} neuron responding to a 5 Hz train of laser stimulation. **(E)** Top, Schematic of viral infusion. Bottom, Representative immunohistochemical micrograph showing colabeling (white arrows) between the ChR2 viral construct (red, AAV-mDlx- ChR2-mCherry) and the GABA marker GAD67 (green), confirming that the use of a mDlx promoter enables transgene expression favoring PL GABA neurons. Scale bars: 25 μ m. **(F- H)** Photoidentification of PL^{GABA} neurons. (F) Frequency histogram showing the latency of response to laser illumination for PL neurons (n = 69 photoidentified PL^{GABA} neurons out of 338 recorded neurons). Triangle method detection of cluster distribution revealed a separation of latency frequencies at 6 ms (see details in Methods section). (G) Cells with photoresponse latencies < 6 ms (identified as the first bin with Z-score > 3.29, p < 0.001, red dotted line) and high spike reliability during the 6 ms (FF, Fano Factor ratio > 1 compared to pre-laser baseline) were classified as PL^{GABA} neurons (see details in Methods section). (H) Raster plot and peristimulus time histogram showing a representative PL^{GABA} neuron responding to a 5Hz train of laser stimulation. Vertical blue bars: laser onset. Bins of 1 ms. **(I)** Timeline of PL recordings for food cue responses in *Pressers* during test (12 food cues per phase). **(J-K)** Stacked bar showing the percentage of (J) PL^{GLUT} neurons or (K) PL^{GABA} neurons that changed their firing rates in response to food cues from the reward phase to the conflict phase. No significant differences were observed across the phases (Fisher Exact Test, all p's > 0.05; n.s. = non-significant). **(L)** Timeline of PL recordings for spontaneous activity in *Pressers* during test. **(M)** Average firing rate of PL^{GLUT} neurons across the different phases of test. **(N)** Venn diagram showing the percentage of responsive PL^{GLUT} neurons (28 out of 36 neurons) by events. **(O)** Stacked bar showing the percentage of PL^{GLUT} neurons that changed their spontaneous firing rates across the different phases of the test. PL^{GLUT} neurons did not change their firing rates from the reward to the odor phase (Fisher Exact Test, inhibited in Reward phase: 10 neurons, inhibited in Odor phase: 3 neurons, p = 0.063), but were subsequently inhibited from the odor to the conflict phase (Fisher Exact Test, inhibited in Odor phase: 3 neurons, inhibited in Conflict phase: 14 neurons, p = 0.0046). **(P)** Average firing rate of PL^{GABA} neurons across the different phases of test. **(Q)** Venn diagram showing the percentage of responsive PL^{GABA} neurons (57 out of 69 neurons) by events. **(R)** Stacked bar showing the percentage of PL^{GABA} neurons that changed their spontaneous firing rates across the different phases of the test. No significant differences were observed across the phases (Fisher Exact Test, all p's > 0.05; n.s. = non-significant). All statistical analysis details are presented in table S1. See also Supplementary Figs. 7, 8, and 9.

Figure 6. Photoactivation of PL^{GLUT}, but not PL^{GABA}, neurons suppresses reward seeking in a neutral context. (A) Schematic of viral infusion and recordings in PL. **(B)** Changes in PL firing rate with illumination of PL^{GLUT} neurons in rats expressing AAV-CaMKII-ChR2-eYFP in PL (n= 20 neurons). **(C)** Average PSTH of PL neurons that were excited during laser illumination of PL^{GLUT} neurons. **(D)** Raster plot and peri-stimulus time histogram (PSTH) of representative PL neuron showing excitatory responses to illumination in rats expressing AAV-CaMKII-ChR2-eYFP in PL. **(E)** Schematic of viral infusion and recordings in PL. **(F)** Changes in PL firing rate with illumination of PL^{GABA} neurons in rats expressing AAV-mDlx-ChR2-mCherry in PL (n= 22 neurons). **(G)** Average PSTH of PL neurons that were inhibited during laser illumination of PL^{GABA} neurons. **(H)** Raster plot and PSTH of representative PL neuron showing inhibitory responses to illumination in rats expressing AAV-mDlx-ChR2-mCherry in PL. **(I-J)** Representative micrograph showing the expression of (I) CaMKII-ChR2-eYFP or (J) mDlx-ChR2-mCherry in PL and schematic of optical fiber location (gray dots) in the same region (compressed across different antero-posterior levels of PL). Green or red areas represent the minimum (dark) and the maximum (light) viral expression into the PL. **(K-L)** Schematic and timeline of PL photoactivation during the cued food seeking test in a neutral context. **(M-N)** Optogenetic activation of PL^{GLUT} neurons (CaMKII-ChR2, dark red circles, n = 7), but not PL^{GABA} neurons (mDlx-ChR2, gold circles, n = 17), reduced the (M) frequency of lever presses ($F_{(10, 180)} = 7.009$, $p < 0.001$, Bonferroni *post-hoc*, CaMKII-ChR2 vs. Control, all laser on periods - $p < 0.01$; mDlx-ChR2 vs. Control, all laser on periods - $p > 0.05$) and increased (N) the latency for the first press ($F_{(10, 180)} = 9.931$, $p < 0.001$, CaMKII-ChR2 vs. Control, all laser on periods - Bonferroni *post-hoc*, $p < 0.001$; mDlx-ChR2 vs. Control, all laser on periods, $p > 0.05$). **(O)** Optogenetic activation of PL neurons did not alter freezing behavior ($F_{(10, 180)} = 1.124$, $p = 0.346$). Blue shaded area represents laser-on trials (PL^{GLUT}: 5 Hz, PL^{GABA}: 20 Hz; 5ms pulse width, 7-10 mW, 30 s duration). Data shown as mean $\pm$ SEM. Each circle represents the average of two consecutive trials. Two-way repeated-measures ANOVA followed by Bonferroni *post-hoc* test. All * p 's < 0.05 . All statistical analysis details are presented in table S1.

Figure 7. Photoinhibition of PL^{GLUT} neurons during conflict reduces freezing and increases food approaching in *Non-pressers*. (A) Schematic of AAV-CaMKII-eNpHR–eYFP virus infusion in PL and location of optical fibers (gray dots) in the same region (compressed across different antero-posterior levels of PL). Green areas represent the minimum (dark) and the maximum (light) viral expression into the PL. (B–C) Schematic and timeline of the approach-avoidance conflict test during optogenetic inhibition of PL^{GLUT} neurons. (D–H) Photoinhibition of PL^{GLUT} neurons (CaMKII-eNpHR, red bars, n = 11) during the conflict test reduced the percentage of time rats spent (D) freezing (Wilcoxon test, W = -64, laser off vs. laser on, p = 0.0020, Mann-Whitney Test, U = 18 Control vs. CaMKII-eNpHR, p = 0.319) and (E) avoiding the odor area (Wilcoxon test, W = -21, laser off vs. laser on, p = 0.031; Mann-Whitney Test, U = 19.5 Control vs. CaMKII-eNpHR, p = 0.365), and increased the percentage of time rats spent in the (F) food area (Wilcoxon test, W = 21, laser off vs. laser on, p = 0.031; Mann-Whitney Test, U = 17 Control vs. CaMKII-eNpHR, p = 0.221) during the conflict test without altering (G) the number of lever presses (Wilcoxon test, W = 6, laser off vs. laser on, p = 0.250; Mann-Whitney Test, U=22.5 Control vs. CaMKII-eNpHR, p=0.697) and (H) the latency to press (Wilcoxon test, W = -10, laser off vs. laser on, p = 0.125; Mann-Whitney Test, U = 21 Control vs. CaMKII-eNpHR, p = 0.357). Laser stimulation did not alter behaviors in controls (eYFP-control virus, white bars, n= 5, Wilcoxon test, Freezing: W = 3, p = 0.812, avoidance: W = 3, p = 0.500, food area: W = -3, p = 0.500, lever presses: W = -1, p = 0.999, latency to press: W = 1, p = 0.999). For all Shapiro-Wilk normality test, p < 0.05. (I) Schematic of AAV-mDlx-ChR2–mCherry virus infusion in PL and location of optical fibers (gray dots) in the same region (compressed across different antero-posterior levels of PL). Red areas represent the minimum (dark) and the maximum (light) viral expression into the PL (J–K) Schematic and timeline of the approach-avoidance conflict test during optogenetic activation of PL^{GABA} neurons. (L–P) Photoactivation of PL^{GABA} neurons during the conflict test did not alter rats' behavior in the mDlx-ChR2 group (gold bars, n= 7) or in the control group (eYFP-control virus, white bars, n = 4, Wilcoxon and Mann-Whitney tests, all p's > 0.05). For all Shapiro-Wilk normality test, p < 0.05. PL neurons were illuminated from cue onset until the animals pressed the lever or from cue onset until the end of the 30 s cues if the animals didn't press the lever (PL^{GLUT}: 5 Hz, PL^{GABA}: 20 Hz; 5 ms pulse width, 7-10 mW). Data shown as mean ± SEM. Each bar represents the average of six trials alternated in blocks of 2. All * p's < 0.05. All statistical analysis details are presented in table S1. See also Supplementary Fig 10 and 11.

Figure 8. Schematic showing differences in food-cue responses and spontaneous firing rate of PL neurons in *Pressers* and *Non-pressers*. Left, *Pressers* showed reduced spontaneous firing rate and increased food-cue responses in PL neurons during the conflict test, which may have resulted in higher signal-to-noise ratio and increased food-seeking responses. Right, *Non-pressers* showed increased spontaneous firing rate and reduced food-cue responses in PL neurons during the conflict test, which may have resulted in lower signal-to-noise ratio and reduced food-seeking responses.

Figure 1 - Figure supplement 1. Pressers and Non-pressers showed similar behavioral responses during cued food-seeking training and olfactory threat conditioning. (A) *Pressers* and *Non-pressers* learned to discriminate equally the food cue off from the food cue on periods during the training phase (Group, $F_{(1, 55)} = 2.166$; $p = 0.146$, Time, $F_{(1, 55)} = 238.7$, $p < 0.001$, Interaction $F_{(1, 55)} = 0.106$, $p = 0.745$, Bonferroni *post-hoc* for early vs. late $p < 0.001$). Discriminability was calculated as the number of lever presses during the cue on period (30 s) minus the number of lever presses during the cue off period (30 s before cue) divided by the total number of presses during both periods. (B) *Pressers* and *Non-pressers* showed the same lever press rates during the cue off and cue on periods across the cued food-seeking training ($F_{(3, 153)} = 1.28$, $p = 0.283$). (C-E) *Pressers* and *Non-pressers* exhibited the same (C) lever press rates ($F_{(1, 51)} = 0.265$, $p = 0.608$), (D) freezing levels ($F_{(1, 51)} = 3.737$, $p = 0.058$), and (E) maximum speed ($F_{(1, 51)} = 6.538e007$, $p = 0.999$) during the olfactory threat conditioning training. (F-G) *Pressers* and *Non-pressers* spent the same (F) time investigating the odor (Unpaired Student's t-test, $t = 0.43$, $p = 0.665$) and showed the same (G) latency to press the lever (Unpaired Student's t-test, $t = 0.55$, $p = 0.578$) before the first odor-shock pairing. (H) *Non-pressers* showed higher contextual freezing during the third and fourth pre-odor trials (30 s before odor onset), compared to *Pressers* ($F_{(5, 250)} = 3.038$, $p = 0.011$; Bonferroni *post-hoc* test, third pre-odor trial: $p = 0.0214$; fourth pre-odor trial: $p = 0.015$). (I-L) During the test session on Day 2, *Pressers* and *Non-pressers* showed the same levels of (I) contextual freezing before the first food cue and odor onset (Shapiro-Wilk normality test, $p < 0.001$, Mann Whitney, $U = 248$, $p = 0.113$). The behavioral responses were different between groups but constant across the session for (J) freezing (Group $F_{(1, 50)} = 13.07$, $p < 0.001$; Interaction, $F_{(9, 450)} = 1.327$, $p = 0.220$), (K) avoidance (Group $F_{(1, 50)} = 20.31$, $p < 0.001$; Interaction, $F_{(9, 450)} = 2.109$, $p = 0.027$, Bonferroni *post-hoc* min 1 vs. min 10 $p > 0.999$) and (L) time in food area (Group $F_{(1, 50)} = 117.5$, $p = 0.001$; Interaction, $F_{(9, 450)} = 0.573$, $p = 0.819$). Combined data for all *Pressers* and *Non-pressers* used across the experiments. Data shown as mean $\pm$ SEM. Two-way ANOVA repeated measures followed by Bonferroni *post-hoc* test, * $p < 0.05$ for group comparison. All statistical analysis details are presented in table S1.

Figure 2 - Figure supplement 1. Correlation between food cue-evoked PL activity and lever press latency during the conflict phase in *Pressers*. **(A)** Scatter plot showing a significant inverse correlation between the normalized firing rate (Z-score) of excitatory food-cue responsive neurons and the latency to press the lever after the food cue onset during the conflict phase ($r = -0.103$, $p = 0.020$). **(B)** Scatter plot showing lack of correlation between the normalized firing rate (Z-score) of inhibitory food-cue responsive neurons and the latency to press the lever after the food cue onset during the conflict phase ($r = 0.038$, $p = 0.451$). Each data point denotes the averaged Z-scored response of one neuron until the animal pressed the lever (Y axis) vs. the respective latency to press during each food-cue presentation (X axis). The black lines represent the linear regression of the data points, fitting the first degree polynomial to the data (see Methods for details). The equation for the linear fitting in the excitatory case is $y = 0.0098 \cdot x + 1.1$ and for inhibitory cases is $y = 0.0018 \cdot x - 0.79$.

Figure 2 - Figure supplement 2. Changes in PL responses to reward cues selected during the conflict phase for *Pressers* and *Non-pressers*. (A) Pie charts showing changes in PL firing rate in response to food cues during conflict (left) vs. reward (right) phases for *Pressers* (n = 237 neurons from 25 rats, Fisher Exact Test, responsive during conflict phase: n = 68, responsive during reward phase: n = 34, p < 0.001, excitatory during conflict phase: n = 36, excitatory during reward phase: n = 26, p = 0.22; inhibitory during conflict phase: n = 32, inhibitory during reward phase: n = 8, p < 0.001). (B) Heatmap of Z-scored neural activities for PL neurons selected during conflict phase and tracked to reward phase. (C) Average peri-stimulus time histograms (PSTHs) for all PL neurons showing excitatory food cue responses (Z-score > 2.58, dotted line) during conflict (orange line) compared to the same cells during reward (gray line). C, Inset: Differences in the positive area under the curve (AUC) between the two phases (Wilcoxon test, W = 122, excitatory responses reward phase vs. conflict phase, p = 0.346). (D) Pie charts showing the percentage of sustained vs. transient excitatory food-cue responses in PL neurons during the conflict phase with the same neurons tracked back during the reward phase. (E-F) Representative PSTHs for a PL neuron showing excitatory responses to food cues during the (E) conflict phase vs. the same neuron during the (F) reward phase. (G) Average PSTHs for all PL neurons showing inhibitory food cue responses (Z-score < -1.96, dotted line) during conflict (blue line) compared to the same cells during reward (gray line). G, Inset: Differences in the negative AUC between the two phases (Wilcoxon test, W = 266, inhibitory responses reward phase vs. conflict phase, p = 0.011). (H) Pie charts showing the percentage of sustained vs. transient inhibitory food-cue responses in PL neurons during the conflict phase with the same neurons tracked back during the reward phase. (I-J) Representative PSTHs for a PL neuron showing inhibitory responses to food cues during the (I) conflict phase vs. the same neuron during the (J) reward phase. (K) Pie charts showing changes in PL firing rate in response to food cues during conflict (left) vs. reward (right) phases for *Non-Pressers* (n = 89 neurons from 7 rats, Fisher Exact Test, responsive during conflict phase: n = 18, responsive during reward phase: n = 6, p = 0.0129, excitatory during conflict phase: n = 9, excitatory during reward phase: n = 4, p = 0.2486; inhibitory during conflict phase: n = 9, inhibitory during reward phase: n = 2, p = 0.0573). (L-M) Same as B-C, but for *Non-Pressers*. M, Inset: Differences in the positive AUC between the two phases (Wilcoxon test, W = 19, excitatory responses reward phase vs. conflict phase, p = 0.300). (N) Pie charts showing the percentage of sustained vs. transient excitatory food-cue responses in PL neurons during the conflict phase with the same neurons tracked back during the reward phase. (O-Q) Same as E-G, but for *Non-pressers*. Q, Inset: Differences in the negative AUC between the two phases (Wilcoxon test, W = 11, excitatory responses reward phase vs. conflict phase, p = 0.570). (R) Pie charts showing the percentage of sustained vs. transient inhibitory food-cue responses in PL neurons during the conflict phase with the same neurons tracked back during the reward phase. (S-T) Same as I-J, but for *Non-pressers*. For all Shapiro-Wilk normality test, p < 0.05. All statistical analysis details are presented in table S1.

Figure 2 - Figure supplement 3. Distinct subpopulations of PL neurons change their firing rates in response to food cues, lever presses, and dish entries. (A) Schematic of the recordings during cued lever presses. **(B)** Pie charts showing the percentage of lever press responsive neurons during the reward phase. **(C)** Heatmap showing the normalized firing rate of individual PL neurons time-locked for lever presses. **(D)** Average peri-stimulus time histogram (PSTH) of all PL neurons showing excitatory lever-press responses. **(E)** Raster plot and PSTH of a representative PL neuron showing excitatory lever-press responses. **(F)** Average PSTH of all PL neurons showing inhibitory lever-press responses. **(G)** Raster plot and PSTH of a representative PL neuron showing inhibitory lever-press responses. **(H)** Schematic of the recordings during rewarded food dish entries. **(I)** Pie charts showing the percentage of dish entry responsive neurons during the reward phase (n = 237 neurons). **(J)** Heatmap showing the normalized firing rate of individual PL neurons time-locked for rewarded food dish entries. **(K)** Average PSTH of all PL neurons showing excitatory dish entry responses. **(L)** Raster plot and PSTH of a representative PL neuron showing excitatory dish entry responses. **(M)** Average PSTH of all PL neurons showing inhibitory dish entry responses. **(N)** Raster plot and PSTH of a representative PL neuron showing inhibitory dish entry responses. **(O)** Venn Diagram showing the percentage of all responsive neurons (120 out of 237) by events. Most of the responsive neurons (59%) responded exclusively to one of the events. n = 237 neurons from 25 rats.

Figure 2 - Figure supplement 4. Changes in PL spontaneous activity in *Pressers* and *Non- Pressers* across the different phases of the test session. (A) Timeline of PL recordings for spontaneous activity during test. **(B-C)** Stacked bars showing the percentage of PL neurons that changed their spontaneous firing rates across the different phases of the test in both *Pressers* (B) and *Non-pressers* (C). *Non-pressers* showed a significant increase in the proportion of excited neurons from the odor to the conflict phase (Fisher Exact Test, excited from Reward phase to Odor phase: 14 neurons, excited from Odor phase to Conflict phase: 28 neurons, $p = 0.015$. n.s. = non-significant, * $p < 0.05$. All statistical analysis details are presented in table S1.

Figure 3 - Figure supplement 1. PL activity anticipates the onset of freezing, avoidance or risk-assessment behaviors in both *Pressers* and *Non-pressers*. (A-C) Both *Pressers* and *Non-pressers* showed the same number and proportion of excitatory and inhibitory PL responses 600 ms before the onset of (A) freezing (Fisher Exact Test, $p = 0.34$), (B) avoidance (Fisher Exact Test, $p = 0.1249$) or (C) risk-assessment (RA, Fisher Exact Test, $p = 0.8168$) behaviors. **(D-F)** Representative PSTHs for distinct PL neurons showing excitatory responses 600 ms before the onset of (D) freezing, (E) avoidance or (F) risk- assessment behaviors. **(G-I)** Representative PSTHs for distinct PL neurons showing inhibitory responses 600 ms before the onset of freezing (G), avoidance (H) or risk- assessment (I) behaviors. **(J)** Venn Diagram showing the percentage of all PL responsive neurons (12 out of 88 neurons) by behavior. Most of the responsive neurons responded selectively at the onset of one of the behaviors. **(K-M)** Heatmap of Z-scored neural activities for PL neurons selected at 600 ms before the onset of freezing (K), avoidance (L) or risk-assessment behavior (M) with the same cells tracked during the other behaviors. The threshold used to identify significant differences per neurons was Z-score > 2.58 for excitation and Z-score < -1.96 for inhibition. Stack bar values were compared using Fisher Exact Test, n.s. = non-significant. All statistical analysis details are presented in table S1.

Figure 5 - Figure supplement 1. Validation of the mDlx promoter used for viral vector targeting of GABAergic neurons in PL. (A) Schematic of AAV-mDlx-ChR2-mCherry infusion and fluorescent *in situ* hybridization in PL. **(B, left)** Representative micrograph of PL cells expressing mRNA for mCherry (red label), the GABAergic marker vGAT (green label), or both (white arrowheads). Blue label: DAPI. **(B, right)** Quantification of the total of cells expressing mDlx-mCherry that were also labeled with vGAT (white bar), or cells labeled with vGAT that also expressed mDlx-mCherry (black bar). n = 3 rats.

Figure 5 - Figure supplement 2. Changes in the spontaneous firing rate of PL^{GLUT} and PL^{GABA} neurons across the different phases of the test session. (A) Timeline of PL recordings for changes in spontaneous activity from reward to odor phase. **(B-C)** Pie charts showing the proportion of (B) PL^{GLUT} neurons (n = 36) and (C) PL^{GABA} neurons (n = 69) that changed their spontaneous firing rates from baseline to reward phase (central pie charts, green borders) and from reward phase to odor phase (peripheral pie charts, yellow borders) in *Pressers*. All PL^{GLUT} neurons that were excited or inhibited during the reward phase responded in opposite directions or did not respond during the odor phase. **(D)** Timeline of PL recordings for changes in spontaneous activity from odor to conflict phase. **(E-F)** Pie charts showing the proportion of (E) PL^{GLUT} neurons and (F) PL^{GABA} neurons that changed their spontaneous firing rates from reward to odor phase (central pie charts, yellow borders) and from odor phase to conflict phase (peripheral pie charts, blue borders) in *Pressers*. For all the pie charts the activity of the same cells was tracked across the session.

Figure 5 - Figure supplement 3. Changes in the firing rate of PL^{GLUT} and PL^{GABA} neurons before or after the onset of freezing, avoidance or risk-assessment behaviors. (A-F) Both PL^{GLUT} (n = 36) and PL^{GABA} (n = 69) neurons showed the same number and proportion of excitatory and inhibitory responses when aligned at 600 ms before the onset of (A) freezing (Fisher Exact Test, excitatory responses p = 0.549; inhibitory p = 0.270), (B) avoidance (excitatory responses p = 0.6582; inhibitory p >0.999), or (C) risk- assessment (excitatory responses p = 0.919; inhibitory p >0.999) behaviors, as well as when aligned at 600 ms after the onset of (D) freezing (excitatory responses p > 0.999; inhibitory p = 0.688), (E) avoidance (excitatory responses p = 0.0544; inhibitory p = 0.768) or (F) risk-assessment (excitatory responses p = 0.741; inhibitory p = 0.490) behaviors. The threshold used to identify significant differences per neurons was Z-score > 2.58 for excitation and Z-score < -1.96 for inhibition. Stack bar values were compared using Fisher Exact Test, n.s. = non-significant. All statistical analysis details are presented in table S1.

Figure 7 - Figure supplement 1. Photoinhibition of PL^{GLUT} neurons did not affect cued food-seeking responses in a neutral context. (A-B) Schematic and timeline of PL photoinhibition during the cued food-seeking test in a neutral context. **(C-D)** Optogenetic inhibition of PL^{GLUT} neurons (CaMKII-ChR2, dark red circles, n = 12) had no effect on (C) frequency of lever presses ($F_{(5,110)} = 1.336$, $p = 0.254$), (D) latency for the first press ($F_{(5,110)} = 0.637$, $p = 0.671$), or (E) freezing responses ($F_{(5,95)} = 1.395$, $p = 0.231$), when compared to the control group (eYFP-control virus, white circles, n = 12). Yellow laser illumination (7- 10 mW) was delivered for 30 s at cue onset. Data shown as mean $\pm$ SEM. Each circle represents the average of two consecutive trials. Two-way repeated-measures ANOVA followed by Bonferroni *post-hoc* test. All statistical analysis details are presented in table S1.

Figure 7 - Figure supplement 2. Photoinhibition of PL^{GABA} neurons in *Pressers* does not alter defensive responses and food seeking during conflict (A) Schematic of AAV-mDlx- ChR2-mCherry virus infusion in PL. **(B-C)** Schematic and timeline of the approach-avoidance conflict test during optogenetic activation of PL^{GABA} neurons. **(D-H)** Photoactivation of PL^{GABA} neurons during the conflict test did not alter rats' behavior (mDlx- ChR2, gold bars, n = 8, Control, white bars, n = 5; Repeated measures ANOVA, Freezing: $F_{(1, 11)} = 2.186$, $p = 0.167$; Avoidance: $F_{(1, 11)} = 0.9568$, $p = 0.349$; Time in food area: $F_{(1, 11)} = 1.798$, $p = 0.207$; Rewarded Presses: Shapiro-Wilk normality test, $p < 0.05$, Wilcoxon test, off vs. on - $W = 19$, $p = 0.312$, Mann-Whitney, mDlx-ChR2 vs. control - $U=18$, $p = 0.673$; Latency to press: $F_{(1, 11)} = 1.038$, $p = 0.330$). PL neurons were illuminated from cue onset until the animals pressed the lever or from cue onset until the end of the 30 s cues if the animals didn't press the lever (PL^{GABA}: 20 Hz; 5 ms pulse width, 7-10 mW). Data shown as mean $\pm$ SEM. Each bar represents the average of six trials alternated in blocks of 2. All statistical analysis details are presented in table S1.

1610 **Video 1. Presser vs Non-presser animals during approach-avoidance conflict task.**
1611 Representative video comparing the different behavioral strategies during the conflict phase.
1612 Animals' body were label using DeepLabCut software and dots were used to track the animal
1613 position, freezing and risk assessment.

REFERENCES

- Ainslie G (1975) Specious reward: a behavioral theory of impulsiveness and impulse control. *Psychol Bull* 82:463-496.
- Allsop SA et al. (2018) Corticoamygdala Transfer of Socially Derived Information Gates Observational Learning. *Cell* 173:1329-1342 e1318.
- Amir A, Lee SC, Headley DB, Herzallah MM, Pare D (2015) Amygdala Signaling during Foraging in a Hazardous Environment. *J Neurosci* 35:12994-13005.
- Aupperle RL, Paulus MP (2010) Neural systems underlying approach and avoidance in anxiety disorders. *Dialogues Clin Neurosci* 12:517-531.
- Baeg EH, Kim YB, Jang J, Kim HT, Mook-Jung I, Jung MW (2001) Fast spiking and regular spiking neural correlates of fear conditioning in the medial prefrontal cortex of the rat. *Cereb Cortex* 11:441-451.
- Bari A, Theobald DE, Caprioli D, Mar AC, Aidoo-Micah A, Dalley JW, Robbins TW (2010) Serotonin modulates sensitivity to reward and negative feedback in a probabilistic reversal learning task in rats. *Neuropsychopharmacology* 35:1290-1301.
- Barker TV, Buzzell GA, Fox NA (2019) Approach, avoidance, and the detection of conflict in the development of behavioral inhibition. *New Ideas Psychol* 53:2-12.
- Berglind WJ, See RE, Fuchs RA, Ghee SM, Whitfield TW, Jr., Miller SW, McGinty JF (2007) A BDNF infusion into the medial prefrontal cortex suppresses cocaine seeking in rats. *Eur J Neurosci* 26:757-766.
- Beyeler A, Namburi P, Globor GF, Simonnet C, Calhoun GG, Conyers GF, Luck R, Wildes CP, Tye KM (2016) Divergent Routing of Positive and Negative Information from the Amygdala during Memory Retrieval. *Neuron* 90:348-361.
- Bezudnaya T, Cano M, Bereshpolova Y, Stoelzel CR, Alonso JM, Swadlow HA (2006) Thalamic burst mode and inattention in the awake LGNd. *Neuron* 49:421-432.
- Blanchard DC, Griebel G, Pobbe R, Blanchard RJ (2011) Risk assessment as an evolved threat detection and analysis process. *Neurosci Biobehav Rev* 35:991-998.
- Bravo-Rivera H, Rubio Arzola P, Caban-Murillo A, Velez-Aviles AN, Ayala-Rosario SN, Quirk GJ (2021) Characterizing Different Strategies for Resolving Approach-Avoidance Conflict. *Front Neurosci* 15:608922.
- Breton JM, Charbit AR, Snyder BJ, Fong PTK, Dias EV, Himmels P, Lock H, Margolis EB (2019) Relative contributions and mapping of ventral tegmental area dopamine and GABA neurons by projection target in the rat. *J Comp Neurol* 527:916-941.
- Burgos-Robles A, Vidal-Gonzalez I, Quirk GJ (2009) Sustained conditioned responses in prelimbic prefrontal neurons are correlated with fear expression and extinction failure. *J Neurosci* 29:8474-8482.
- Burgos-Robles A, Bravo-Rivera H, Quirk GJ (2013) Prelimbic and infralimbic neurons signal distinct aspects of appetitive instrumental behavior. *PLoS One* 8:e57575.
- Burgos-Robles A, Kimchi EY, Izadmehr EM, Porzenheim MJ, Ramos-Guasp WA, Nieh EH, Felix-Ortiz AC, Namburi P, Leppla CA, Presbrey KN, Anandalingam KK, Pagan-Rivera PA, Anahtar M, Beyeler A, Tye KM (2017) Amygdala inputs to prefrontal cortex guide behavior amid conflicting cues of reward and punishment. *Nat Neurosci*.
- Cain CK (2019) Avoidance Problems Reconsidered. *Curr Opin Behav Sci* 26:9-17.
- Capuzzo G, Floresco SB (2020) Prelimbic and Infralimbic Prefrontal Regulation of Active and Inhibitory Avoidance and Reward-Seeking. *J Neurosci* 40:4773-4787.
- Chen BT, Yau HJ, Hatch C, Kusumoto-Yoshida I, Cho SL, Hopf FW, Bonci A (2013) Rescuing cocaine-induced prefrontal cortex hypoactivity prevents compulsive cocaine seeking. *Nature* 496:359-362.

Choi EA, McNally GP (2017) Paraventricular Thalamus Balances Danger and Reward. *J Neurosci* 37:3018-3029.

Choi EA, Jean-Richard-Dit-Bressel P, Clifford CWG, McNally GP (2019) Paraventricular Thalamus Controls Behavior during Motivational Conflict. *J Neurosci* 39:4945-4958.

Churchland AK, Kiani R, Chaudhuri R, Wang XJ, Pouget A, Shadlen MN (2011) Variance as a signature of neural computations during decision making. *Neuron* 69:818-831.

Cohen J (1988) *Statistical Power Analysis for the Behavioral Sciences* (2nd edition). Hillsdale: Erlbaum Associates.

Cohen JY, Haesler S, Vong L, Lowell BB, Uchida N (2012) Neuron-type-specific signals for reward and punishment in the ventral tegmental area. *Nature* 482:85-88.

Courtin J, Chaudun F, Rozeske RR, Karalis N, Gonzalez-Campo C, Wurtz H, Abdi A, Baufreton J, Bienvenu TC, Herry C (2014) Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. *Nature* 505:92-96.

Cummings KA, Clem RL (2020) Prefrontal somatostatin interneurons encode fear memory. *Nat Neurosci* 23:61-74.

Dejean C, Courtin J, Karalis N, Chaudun F, Wurtz H, Bienvenu TC, Herry C (2016) Prefrontal neuronal assemblies temporally control fear behaviour. *Nature* 535:420-424.

Dellu-Hagedorn F, Rivalan M, Fitoussi A, De Deurwaerdere P (2018) Inter-individual differences in the impulsive/compulsive dimension: deciphering related dopaminergic and serotonergic metabolisms at rest. *Philos Trans R Soc Lond B Biol Sci* 373.

Diehl MM, Bravo-Rivera C, Rodriguez-Romaguera J, Pagan-Rivera PA, Burgos-Robles A, Roman-Ortiz C, Quirk GJ (2018) Active avoidance requires inhibitory signaling in the rodent prelimbic prefrontal cortex. *Elife* 7.

Dimidschstein J et al. (2016) A viral strategy for targeting and manipulating interneurons across vertebrate species. *Nat Neurosci* 19:1743-1749.

Do-Monte FH, Quinones-Laracuente K, Quirk GJ (2015) A temporal shift in the circuits mediating retrieval of fear memory. *Nature* 519:460-463.

Do-Monte FH, Rodriguez-Romaguera J, Rosas-Vidal LE, Quirk GJ (2013) Deep brain stimulation of the ventral striatum increases BDNF in the fear extinction circuit. *Front Behav Neurosci* 7:102.

Do Monte FH, Quirk GJ, Li B, Penzo MA (2016) Retrieving fear memories, as time goes by. *Mol Psychiatry* 21:1027-1036.

Emmons EB, Ruggiero RN, Kelley RM, Parker KL, Narayanan NS (2016) Corticostriatal Field Potentials Are Modulated at Delta and Theta Frequencies during Interval-Timing Task in Rodents. *Front Psychol* 7:459.

Engelke DS, Zhang XO, O'Malley JJ, Fernandez-Leon JA, Li S, Kirouac GJ, Beierlein M, Do-Monte FH (2021) A hypothalamic-thalamostriatal circuit that controls approach-avoidance conflict in rats. *Nat Commun* 12:2517.

Ferenczi EA, Zalocusky KA, Liston C, Grosenick L, Warden MR, Amatya D, Katovich K, Mehta H, Patenaude B, Ramakrishnan C, Kalanithi P, Etkin A, Knutson B, Glover GH, Deisseroth K (2016) Prefrontal cortical regulation of brainwide circuit dynamics and reward-related behavior. *Science* 351:aac9698.

Flagel SB, Watson SJ, Robinson TE, Akil H (2007) Individual differences in the propensity to approach signals vs goals promote different adaptations in the dopamine system of rats. *Psychopharmacology (Berl)* 191:599-607.

Flagel SB, Clark JJ, Robinson TE, Mayo L, Czuj A, Willuhn I, Akers CA, Clinton SM, Phillips PE, Akil H (2011) A selective role for dopamine in stimulus-reward learning. *Nature* 469:53-57.

Gabbott PL, Warner TA, Jays PR, Salway P, Busby SJ (2005) Prefrontal cortex in the rat: projections to subcortical autonomic, motor, and limbic centers. *J Comp Neurol* 492:145-177.

- Gaykema RP, Nguyen XM, Boehret JM, Lambeth PS, Joy-Gaba J, Warthen DM, Scott MM (2014) Characterization of excitatory and inhibitory neuron activation in the mouse medial prefrontal cortex following palatable food ingestion and food driven exploratory behavior. *Front Neuroanat* 8:60.
- Geller I (1960) The acquisition and extinction of conditioned suppression as a function of the base-line reinforcer. *J Exp Anal Behav* 3:235-240.
- Gradinaru V, Mogri M, Thompson KR, Henderson JM, Deisseroth K (2009) Optical deconstruction of parkinsonian neural circuitry. *Science* 324:354-359.
- Grant RI, Doncheck EM, Vollmer KM, Winston KT, Romanova EV, Siegler PN, Holman H, Bowen CW, Otis JM (2021) Specialized coding patterns among dorsomedial prefrontal neuronal ensembles predict conditioned reward seeking. *Elife* 10.
- Haight JL, Fuller ZL, Fraser KM, Flagel SB (2017) A food-predictive cue attributed with incentive salience engages subcortical afferents and efferents of the paraventricular nucleus of the thalamus. *Neuroscience* 340:135-152.
- Harris AZ, Gordon JA (2015) Long-range neural synchrony in behavior. *Annu Rev Neurosci* 38:171-194.
- Horst NK, Laubach M (2013) Reward-related activity in the medial prefrontal cortex is driven by consumption. *Front Neurosci* 7:56.
- Hu H (2016) Reward and Aversion. *Annu Rev Neurosci* 39:297-324.
- Hyman JM, Hasselmo ME, Seamans JK (2011) What is the Functional Relevance of Prefrontal Cortex Entrainment to Hippocampal Theta Rhythms? *Front Neurosci* 5:24.
- Jean-Richard-Dit-Bressel P, Ma C, Bradfield LA, Killcross S, McNally GP (2019) Punishment insensitivity emerges from impaired contingency detection, not aversion insensitivity or reward dominance. *Elife* 8.
- Jensen J, Walter H (2014) Incentive motivational salience and the human brain. *Restor Neurol Neurosci* 32:141-147.
- Jolliffe IT (2002) Principal Component Analysis. Springer Series in Statistics 2nd edition.
- Karalis N, Dejean C, Chaudun F, Khoder S, Rozeske RR, Wurtz H, Bagur S, Benchenane K, Sirota A, Courtin J, Herry C (2016) 4-Hz oscillations synchronize prefrontal-amygdala circuits during fear behavior. *Nat Neurosci*.
- Kim CK, Ye L, Jennings JH, Pichamoorthy N, Tang DD, Yoo AW, Ramakrishnan C, Deisseroth K (2017) Molecular and Circuit-Dynamical Identification of Top-Down Neural Mechanisms for Restraint of Reward Seeking. *Cell* 170:1013-1027 e1014.
- Kim EJ, Kim N, Kim HT, Choi JS (2013) The prelimbic cortex is critical for context-dependent fear expression. *Front Behav Neurosci* 7:73.
- Kim EJ, Kong MS, Park SG, Mizumori SJY, Cho J, Kim JJ (2018) Dynamic coding of predatory information between the prelimbic cortex and lateral amygdala in foraging rats. *Sci Adv* 4:eaar7328.
- Kirlic N, Young J, Aupperle RL (2017) Animal to human translational paradigms relevant for approach avoidance conflict decision making. *Behav Res Ther* 96:14-29.
- Kittelmann M, Liewald JF, Hegemann J, Schultheis C, Brauner M, Steuer Costa W, Wabnig S, Eimer S, Gottschalk A (2013) In vivo synaptic recovery following optogenetic hyperstimulation. *Proc Natl Acad Sci U S A* 110:E3007-3016.
- Kroener S, Chandler LJ, Phillips PE, Seamans JK (2009) Dopamine modulates persistent synaptic activity and enhances the signal-to-noise ratio in the prefrontal cortex. *PLoS One* 4:e6507.
- Krypotos AM, Eftting M, Kindt M, Beckers T (2015) Avoidance learning: a review of theoretical models and recent developments. *Front Behav Neurosci* 9:189.
- Kyriazi P, Headley DB, Pare D (2020) Different Multidimensional Representations across the Amygdalo-Prefrontal Network during an Approach-Avoidance Task. *Neuron* 107:717-730 e715.

1765 Likhtik E, Paz R (2015) Amygdala-prefrontal interactions in (mal)adaptive learning. *Trends*
 1766 *Neurosci* 38:158-166.
 1767 Likhtik E, Stujenske JM, Topiwala MA, Harris AZ, Gordon JA (2014) Prefrontal entrainment of
 1768 amygdala activity signals safety in learned fear and innate anxiety. *Nat Neurosci* 17:106-
 1769 113.
 1770 Lima SQ, Hromadka T, Znamenskiy P, Zador AM (2009) PINP: a new method of tagging
 1771 neuronal populations for identification during in vivo electrophysiological recording. *PLoS*
 1772 *One* 4:e6099.
 1773 Marquis JP, Killcross S, Haddon JE (2007) Inactivation of the prelimbic, but not infralimbic,
 1774 prefrontal cortex impairs the contextual control of response conflict in rats. *Eur J*
 1775 *Neurosci* 25:559-566.
 1776 Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018)
 1777 DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.
 1778 *Nat Neurosci* 21:1281-1289.
 1779 McDonald AJ (1991) Organization of amygdaloid projections to the prefrontal cortex and
 1780 associated striatum in the rat. *Neuroscience* 44:1-14.
 1781 McGinley MJ, David SV, McCormick DA (2015) Cortical Membrane Potential Signature of
 1782 Optimal States for Sensory Signal Detection. *Neuron* 87:179-192.
 1783 McGinty JF, Otis JM (2020) Heterogeneity in the Paraventricular Thalamus: The Traffic Light of
 1784 Motivated Behaviors. *Front Behav Neurosci* 14:590528.
 1785 McNaughton N, Corr PJ (2014) Approach, avoidance, and their conflict: the problem of
 1786 anchoring. *Front Syst Neurosci* 8:124.
 1787 Millan EZ, Ong Z, McNally GP (2017) Paraventricular thalamus: Gateway to feeding, appetitive
 1788 motivation, and drug addiction. *Prog Brain Res* 235:113-137.
 1789 Moorman DE, Aston-Jones G (2015) Prefrontal neurons encode context-based response
 1790 execution and inhibition in reward seeking and extinction. *Proc Natl Acad Sci U S A*
 1791 112:9472-9477.
 1792 Morales I, Berridge KC (2020) 'Liking' and 'wanting' in eating and food reward: Brain
 1793 mechanisms and clinical implications. *Physiol Behav* 227:113152.
 1794 Nair-Roberts RG, Chatelain-Badie SD, Benson E, White-Cooper H, Bolam JP, Ungless MA
 1795 (2008) Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons
 1796 in the ventral tegmental area, substantia nigra and retrorubral field in the rat.
 1797 *Neuroscience* 152:1024-1031.
 1798 Namburi P, Beyeler A, Yorozu S, Calhoon GG, Halbert SA, Wichmann R, Holden SS, Mertens
 1799 KL, Anahtar M, Felix-Ortiz AC, Wickersham IR, Gray JM, Tye KM (2015) A circuit
 1800 mechanism for differentiating positive and negative associations. *Nature* 520:675-678.
 1801 Narayanan NS, Cavanagh JF, Frank MJ, Laubach M (2013) Common medial frontal
 1802 mechanisms of adaptive control in humans and rodents. *Nat Neurosci* 16:1888-1895.
 1803 Oberrauch S, Sigrist H, Sautter E, Gerster S, Bach DR, Pryce CR (2019) Establishing operant
 1804 conflict tests for the translational study of anxiety in mice. *Psychopharmacology (Berl)*
 1805 236:2527-2541.
 1806 Orsini CA, Heshmati SC, Garman TS, Wall SC, Bizon JL, Setlow B (2018) Contributions of
 1807 medial prefrontal cortex to decision making involving risk of punishment.
 1808 *Neuropharmacology* 139:205-216.
 1809 Otis JM, Nambodiri VM, Matan AM, Voets ES, Mohorn EP, Kosyk O, McHenry JA, Robinson
 1810 JE, Resendez SL, Rossi MA, Stuber GD (2017) Prefrontal cortex output circuits guide
 1811 reward seeking through divergent cue encoding. *Nature* 543:103-107.
 1812 Padilla-Coreano N, Canetta S, Mikofsky RM, Alway E, Passecker J, Myroshnychenko MV,
 1813 Garcia-Garcia AL, Warren R, Teboul E, Blackman DR, Morton MP, Hupalo S, Tye KM,
 1814 Kellendonk C, Kupferschmidt DA, Gordon JA (2019) Hippocampal-Prefrontal Theta
 1815 Transmission Regulates Avoidance Behavior. *Neuron* 104:601-610 e604.

1816 Park J, Moghaddam B (2017) Risk of punishment influences discrete and coordinated encoding
 1817 of reward-guided actions by prefrontal cortex and VTA neurons. *Elife* 6.
 1818 Penzo MA, Gao C (2021) The paraventricular nucleus of the thalamus: an integrative node
 1819 underlying homeostatic behavior. *Trends Neurosci*.
 1820 Pi HJ, Hangya B, Kvitsiani D, Sanders JI, Huang ZJ, Kepecs A (2013) Cortical interneurons that
 1821 specialize in disinhibitory control. *Nature* 503:521-524.
 1822 Piantadosi PT, Yeates DCM, Floresco SB (2020) Prefrontal cortical and nucleus accumbens
 1823 contributions to discriminative conditioned suppression of reward-seeking. *Learn Mem*
 1824 27:429-440.
 1825 Popa D, Duvarci S, Popescu AT, Lena C, Pare D (2010) Coherent amygdalocortical theta
 1826 promotes fear memory consolidation during paradoxical sleep. *Proc Natl Acad Sci U S A*
 1827 107:6516-6519.
 1828 Punter PH (1983) Measurement of human olfactory thresholds for several groups of structurally
 1829 related compounds. *Chem Senses* 7:215-235.
 1830 Radke AK, Nakazawa K, Holmes A (2015) Cortical GluN2B deletion attenuates punished
 1831 suppression of food reward-seeking. *Psychopharmacology (Berl)* 232:3753-3761.
 1832 Richardson JTE (2011) Eta squared and partial eta squared as measures of effect size in
 1833 educational research. *Educational Research Review* 6:135-147.
 1834 Robinson TE, Flagel SB (2009) Dissociating the predictive and incentive motivational properties
 1835 of reward-related cues through the study of individual differences. *Biol Psychiatry*
 1836 65:869-873.
 1837 Robinson TE, Yager LM, Cogan ES, Saunders BT (2014) On the motivational properties of
 1838 reward cues: Individual differences. *Neuropharmacology* 76 Pt B:450-459.
 1839 Sangha S, Robinson PD, Greba Q, Davies DA, Howland JG (2014) Alterations in reward, fear
 1840 and safety cue discrimination after inactivation of the rat prelimbic and infralimbic
 1841 cortices. *Neuropsychopharmacology* 39:2405-2413.
 1842 Santana N, Bortolozzi A, Serrats J, Mengod G, Artigas F (2004) Expression of serotonin1A and
 1843 serotonin2A receptors in pyramidal and GABAergic neurons of the rat prefrontal cortex.
 1844 *Cereb Cortex* 14:1100-1109.
 1845 Schultz W (2015) Neuronal Reward and Decision Signals: From Theories to Data. *Physiol Rev*
 1846 95:853-951.
 1847 Serrano-Barroso A, Vargas JP, Diaz E, O'Donnell P, Lopez JC (2019) Sign and goal tracker rats
 1848 process differently the incentive salience of a conditioned stimulus. *PLoS One*
 1849 14:e0223109.
 1850 Sharpe MJ, Killcross S (2015) The prelimbic cortex directs attention toward predictive cues
 1851 during fear learning. *Learn Mem* 22:289-293.
 1852 Sierra-Mercado D, Padilla-Coreano N, Quirk GJ (2011) Dissociable roles of prelimbic and
 1853 infralimbic cortices, ventral hippocampus, and basolateral amygdala in the expression
 1854 and extinction of conditioned fear. *Neuropsychopharmacology* 36:529-538.
 1855 Simon NW, Gilbert RJ, Mayse JD, Bizon JL, Setlow B (2009) Balancing risk and reward: a rat
 1856 model of risky decision making. *Neuropsychopharmacology* 34:2208-2217.
 1857 Smith KS, Berridge KC, Aldridge JW (2011) Disentangling pleasure from incentive salience and
 1858 learning signals in brain reward circuitry. *Proc Natl Acad Sci U S A* 108:E255-264.
 1859 Sotres-Bayon F, Sierra-Mercado D, Pardilla-Delgado E, Quirk GJ (2012) Gating of fear in
 1860 prelimbic cortex by hippocampal and amygdala inputs. *Neuron* 76:804-812.
 1861 St Onge JR, Floresco SB (2009) Dopaminergic modulation of risk-based decision making.
 1862 *Neuropsychopharmacology* 34:681-697.
 1863 Stahlberg MA, Ramakrishnan C, Willig KI, Boyden ES, Deisseroth K, Dean C (2019)
 1864 Investigating the feasibility of channelrhodopsin variants for nanoscale optogenetics.
 1865 *Neurophotonics* 6:015007.

- Stujenske JM, Likhtik E, Topiwala MA, Gordon JA (2014) Fear and safety engage competing patterns of theta-gamma coupling in the basolateral amygdala. *Neuron* 83:919-933.
- Sun X, Bernstein MJ, Meng M, Rao S, Sorensen AT, Yao L, Zhang X, Anikeeva PO, Lin Y (2020) Functionally Distinct Neuronal Ensembles within the Memory Engram. *Cell* 181:410-423 e417.
- Tan KR, Yvon C, Turiault M, Mirzabekov JJ, Doeber J, Labouebe G, Deisseroth K, Tye KM, Luscher C (2012) GABA neurons of the VTA drive conditioned place aversion. *Neuron* 73:1173-1183.
- Tomie A, Aguado AS, Pohorecky LA, Benjamin D (2000) Individual differences in pavlovian autoshaping of lever pressing in rats predict stress-induced corticosterone release and mesolimbic levels of monoamines. *Pharmacol Biochem Behav* 65:509-517.
- Total NK, Jackson ME, Moghaddam B (2013) Preparatory attention relies on dynamic interactions between prelimbic cortex and anterior cingulate cortex. *Cereb Cortex* 23:729-738.
- Treanor M, Barry TJ (2017) Treatment of avoidance behavior as an adjunct to exposure therapy: Insights from modern learning theory. *Behav Res Ther* 96:30-36.
- Tye KM, Prakash R, Kim SY, Fenno LE, Grosenick L, Zarabi H, Thompson KR, Gradinaru V, Ramakrishnan C, Deisseroth K (2011) Amygdala circuitry mediating reversible and bidirectional control of anxiety. *Nature* 471:358-362.
- van Zessen R, Phillips JL, Budygin EA, Stuber GD (2012) Activation of VTA GABA neurons disrupts reward consumption. *Neuron* 73:1184-1194.
- Verharen JPH, van den Heuvel MW, Luijendijk M, Vanderschuren L, Adan RAH (2019) Corticolimbic Mechanisms of Behavioral Inhibition under Threat of Punishment. *J Neurosci* 39:4353-4364.
- Vertes RP (2004) Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse* 51:32-58.
- Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ (2006) Microstimulation reveals opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear. *Learn Mem* 13:728-733.
- Vogel JR, Beer B, Clody DE (1971) A simple and reliable conflict procedure for testing anti-anxiety agents. *Psychopharmacologia* 21:1-7.
- Volkow ND, Wang GJ, Fowler JS, Tomasi D, Baler R (2012) Food and drug reward: overlapping circuits in human obesity and addiction. *Curr Top Behav Neurosci* 11:1-24.
- Wakabayashi KT, Feja M, Baidur AN, Bruno MJ, Bhimani RV, Park J, Hausknecht K, Shen RY, Haj-Dahmane S, Bass CE (2019) Chemogenetic activation of ventral tegmental area GABA neurons, but not mesoaccumbal GABA terminals, disrupts responding to reward-predictive cues. *Neuropsychopharmacology* 44:372-380.
- Warthen DM, Lambeth PS, Ottolini M, Shi Y, Barker BS, Gaykema RP, Newmyer BA, Joy-Gaba J, Ohmura Y, Perez-Reyes E, Guler AD, Patel MK, Scott MM (2016) Activation of Pyramidal Neurons in Mouse Medial Prefrontal Cortex Enhances Food-Seeking Behavior While Reducing Impulsivity in the Absence of an Effect on Food Intake. *Front Behav Neurosci* 10:63.
- Widge AS, Heilbronner SR, Hayden BY (2019) Prefrontal cortex and cognitive control: new insights from human electrophysiology. *F1000Res* 8.
- Zack GW, Rogers WE, Latt SA (1977) Automatic measurement of sister chromatid exchange frequency. *J Histochem Cytochem* 25:741-753.
- Zhang X, Kim J, Tonegawa S (2020) Amygdala Reward Neurons Form and Store Fear Extinction Memory. *Neuron* 105:1077-1093 e1077.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

Figure 6

FIGURE 7

Figure 7

FIGURE 8

SUPPLEMENTARY FIGURE 1

SUPPLEMENTARY FIGURE 2

SUPPLEMENTARY FIGURE 3

Pressers

Non-pressers

SUPPLEMENTARY FIGURE 4

SUPPLEMENTARY FIGURE 5

SUPPLEMENTARY FIGURE 6

SUPPLEMENTARY FIGURE 7

SUPPLEMENTARY FIGURE 8

SUPPLEMENTARY FIGURE 9

Selected at 600 ms before behavioral onset

Selected at 600 ms after behavioral onset

SUPPLEMENTARY FIGURE 10

SUPPLEMENTARY FIGURE 11