Title: Cnidarian hair cell development illuminates an ancient role for the class IV POU transcription factor in defining mechanoreceptor identity

Authors:

Ethan Ozment¹ (etozment@uark.edu)

Arianna N. Tamvacakis¹ (ariannat@uark.edu)

Jianhong Zhou¹ (jxz011@uark.edu)

Pablo Yamild Rosiles-Loeza² (pablo.rosiles@cinvestav.mx)

Esteban Elías Escobar-Hernandez² (esteban.escobar@cinvestav.mx)

Selene L. Fernandez-Valverde² (selene.fernandez@cinvestav.mx)

Nagayasu Nakanishi^{1*} (nnakanis@uark.edu)

¹Department of Biological Sciences, University of Arkansas, 850 West Dickson St. Fayetteville, Arkansas, 72701

²Unidad de Genómica Avanzada (Langebio), Centro de Investigación y de Estudios Avanzados del IPN, Irapuato, México

* Correspondence:

Nagayasu Nakanishi

nnakanis@uark.edu

Running title: Cnidarian mechanoreceptor evo-devo

Key words: Cnidaria, mechanoreceptor, evo-devo, cell differentiation, *Nematostella*

Abstract:

Although specialized mechanosensory cells are found across animal phylogeny, early evolutionary histories of mechanoreceptor development remain enigmatic. Cnidaria (e.g. sea anemones and jellyfishes) is the sister group to well-studied Bilateria (e.g. flies and vertebrates), and has two mechanosensory cell types – a lineage-specific sensory-effector known as the cnidocyte, and a classical mechanosensory neuron referred to as the hair cell. While developmental genetics of cnidocytes is increasingly understood, genes essential for cnidarian hair cell development are unknown. Here we show that the class IV POU homeodomain transcription factor (POU-IV) – an indispensable regulator of mechanosensory cell differentiation in Bilateria and cnidocyte differentiation in Cnidaria – controls hair cell development in the sea anemone cnidarian *Nematostella vectensis*. *N. vectensis* POU-IV is postmitotically expressed in tentacular hair cells, and is necessary for development of the apical mechanosensory apparatus, but not of neurites, in hair cells. Moreover, it binds to deeply conserved DNA recognition elements, and turns on a unique set of effector genes – including the transmembrane-receptor-encoding gene *polycystin 1* – specifically in hair cells. Our results suggest that POU-IV directs differentiation of cnidarian hair cells and cnidocytes via distinct gene regulatory mechanisms, and support an evolutionarily ancient role for POU-IV in defining the mature state of mechanosensory neurons.

Introduction:

One of the most fundamental sensory cell types that emerged in animal evolution is the mechanosensory cell – the specialized sensory epithelial cell that transduces mechanical stimuli (e.g. water vibration) into internal signals. These signals are then communicated, usually via the nervous system, to effector cells (e.g. muscle cells) to elicit behavioral and/or physiological responses of the organism. Indeed, specialized mechanosensory cells are found across diverse animal lineages, from vertebrate hair cells, cephalopod angular acceleration receptors, to statocyst cells of cnidarian jellyfish and ctenophores. Typically, a mechanosensory cell bears an apical mechanosensory apparatus consisting of a single non-motile cilium surrounded by a circle of rigid microvilli with actin rootlets (i.e. stereovilli, or stereocilia), and extends basal neuronal processes that connect to the nervous system (reviewed in (Beisel et al., 2008, Budelmann, 1989, Manley and Ladher, 2008)).

The structure of animal mechanosensory cells is not uniform, however (reviewed in (Bezares-Calderon et al., 2020)). For instance, insect and cephalopod mechanosensory cells lack stereovilli (Jarman, 2002, Budelmann, 1989), while the apical mechanosensory apparatus of vertebrate hair cells is differently shaped, having a cilium on one side of a group of stereovilli of graded lengths, with the stereovilli next to the cilium being the longest (Fain, 2003). The observed morphological diversity in mechanosensory cells of distantly related animals has led to a fundamental question in animal mechanoreceptor evolution: whether the diversity evolved by divergence from a common ancestral form (Beisel et al., 2008, Jørgensen, 1989, Schlosser, 2020), or by independent evolution (Coffin et al., 2004, Holland, 2005). Addressing this question requires an understanding of the mechanisms of mechanoreceptor development across disparate groups of animals.

Developmental genetics of mechanosensory cells has been extensively studied in bilaterian models such as vertebrates and flies (reviewed in (Schlosser, 2020, Boekhoff-Falk, 2005, Beisel et al., 2008)). Yet, relatively little is known about the genetics of mechanoreceptor development in non-bilaterian, early-evolving animal groups such as Cnidaria (e.g. jellyfish, corals and sea anemones), Ctenophora (combjellies), Placozoa and Porifera (sponges), the knowledge of which is key to defining the ancestral conditions for mechanoreceptor development basal to Bilateria. This baseline knowledge, in turn, is necessary for reconstructing how mechanoreceptors diversified in each lineage. In this paper, we focus on investigating the

development of a fundamental, yet understudied, mechanosensory cell type of Cnidaria – the concentric hair cell.

Cnidaria is the sister group to Bilateria (Medina et al., 2001, Putnam et al., 2007, Hejnol et al., 2009, Erwin et al., 2011), and has two broad classes of mechanosensory cells – cnidocytes (Brinkmann et al., 1996)) and concentric hair cells ((Arkett et al., 1988, Oliver and Thurm, 1996, Holtman and Thurm, 2001); not to be confused with vertebrate hair cells) - that are characterized by an apical mechanosensory apparatus consisting of a single cilium surrounded by a ring of stereovilli. The cnidocyte is the phylum-defining stinging cell type, and additionally contains a cnidarian-specific exocytotic organelle called the cnida (plural: cnidae) which is made up of a capsule enclosing a coiled tubule (reviewed in (Thomas and Edwards, 1991, Lesh-Laurie and Suchy, 1991, Fautin and Mariscal, 1991)). Cnidocytes are abundant in the ectodermal epithelium of cnidarian tentacles, and, upon perceiving mechanical stimuli, discharge cnidae by rapidly everting the coiled tubule to pierce nearby animals for defense and/or prey capture. There is no structural or functional evidence that the cnidocyte transmits sensory information to other cells, but firing of cnidae is thought to be modulated by neurons that innervate cnidocytes through chemical synapses (Westfall, 2004). Thus, the cnidocyte is a cnidarian-specific mechanosensory cell type that – uniquely among animal mechanosensory cells – functions as an effector cell.

The cnidarian hair cell, on the other hand, represents the classical mechanosensory cell type with dedicated sensory-neuronal function. Hair cells are integrated within the ectodermal epithelium of mechanosensory structures, such as gravity-sensors of jellyfishes and tentacles of hydroids and corals (Horridge, 1969, Lyons, 1973, Tardent and Schmid, 1972, Singla, 1975, Hundgen and Biela, 1982). Structurally, the cnidarian hair cell exhibits the stereotypical mechanosensory neuron-like morphology described above, including the apical mechanosensory apparatus and basal neurites that become part of the basiepithelial nerve plexus (Horridge, 1969, Singla, 1975, Singla, 1983, Hundgen and Biela, 1982). Upon stimulation, the hair cells communicate mechanosensory information to other cells by converting mechanical stimuli into internal electrical signals (Arkett et al., 1988, Oliver and Thurm, 1996), and are thought to generate highly coordinated response behaviors such as righting and feeding. Similar to vertebrate hair cells, hair cells of jellyfish gravity-sensors are sensitive to sound and can be lost due to noise trauma (Sole et al., 2016). Cnidarian hair cells show morphological and functional

characteristics that parallel those of mechanosensory cells in other animal lineages, consistent with a deep evolutionary origin or convergent origins.

Although genetics of cnidocyte development is increasingly understood (e.g. (Babonis and Martindale, 2017, Richards and Rentzsch, 2015, Richards and Rentzsch, 2014, Wolenski et al., 2013)), that of cnidarian hair cell development remains poorly known. This knowledge gap severely limits our ability to reconstruct the evolutionary histories of mechanoreceptor development within Cnidaria and across the basal branches of the animal tree. A previous study has shown that the class IV POU homeodomain transcription factor (POU-IV or Brn-3)-encoding gene is expressed in the hair-cell-bearing mechanosensory organ called the touch plate in moon jellyfish *Aurelia sp.1* (Nakanishi et al., 2010), consistent with a role in cnidarian hair cell development. Yet, the function of POU-IV in cnidarian hair cell development, if any, remains undefined. As the first step towards elucidating the genetic mechanism of cnidarian hair cell development, here we dissect the role of POU-IV in the development of mechanosensory hair cells using the genetically tractable sea anemone cnidarian model *Nematostella vectensis*.

POU-IV is shared by all extant animal groups except for Ctenophora (comb jellies), indicative of early emergence in animal evolution (Gold et al., 2014). POU-IV is absent in choanoflagellates, although the class II POU-like gene has been reported to be present in *Mylnosiga fluctuans* indicative of a premetazoan origin of POU transcription factors (Lopez-Escardo et al., 2019). As in other POU proteins, POU-IV is characterized by having a bipartite DNA binding domain consisting of the N-terminal POU-specific domain and the C-terminal POU homeodomain (reviewed in (Herr and Cleary, 1995)). In Bilateria, POU-IV-binding DNA elements are POU-IV-class-specific and conserved; mammalian POU-IV proteins Brn3.0 (Brn-3a or POU4F1) and Brn3.2 (Brn-3b or POU4F2) and *C. elegans* POU-IV protein Unc-86 bind to a highly symmetrical core sequence AT(A/T)A(T/A)T(A/T)AT (Gruber et al., 1997). In bilaterian animal models such as *C. elegans*, POU-IV is known to function as a terminal selector - a transcription factor that determines mature cell identity via direct regulation of effector genes (reviewed in (Leyva-Diaz et al., 2020)). The cell type whose identity is defined by POU-IV across bilaterian lineages is the mechanosensory cell. In humans, mutations at one of the *pou-iv* loci – Brn-3c (Brn3.1 or POU4F3) – have been linked to autosomal dominant hearing loss (Vahava et al., 1998), and in Brn-3c knockout mice, auditory and vestibular hair cells fail to complete differentiation (Erkman et al., 1996, Xiang et al., 1997b) and are lost by cell death (Xiang et al.,

1998). Likewise, in *C. elegans*, the *pou-iv* ortholog (*unc-86*) regulates differentiation of mechanosensory touch cells (Chalfie and Sulston, 1981, Chalfie and Au, 1989, Finney and Ruvkun, 1990, Duggan et al., 1998). In addition to its role in mechanoreceptor differentiation, POU-IV defines the identity of olfactory chemosensory neurons in *Drosophila* (Clyne et al., 1999), as well as retinal ganglion cells (Brn-3b; (Erkman et al., 1996, Gan et al., 1996) and subsets of sensory and CNS neurons in mice (Brn-3a; (Serrano-Saiz et al., 2018, McEvilly et al., 1996, Xiang et al., 1996). In Cnidaria, POU-IV is expressed not only in the developing mechanoreceptor of *Aurelia sp.1* (Nakanishi et al., 2010) as described above, but also in the statocysts of the freshwater hydrozoan jellyfish *Craspedacusta sowerbii* (Hroudova et al., 2012). Also, POU-IV is required for postmitotic differentiation of cnidocytes, as well as *elav::mOrange* neurons, in *Nematostella vectensis* (Tourniere et al., 2020). Consistent with cnidarian POU-IV being a terminal selector, a genome-wide analysis of differential gene expression between POU-IV knockout mutant *N. vectensis* and their siblings indicates that POU-IV controls the expression of effector genes that define mature neural identity, such as those involved in ion channel activity (Tourniere et al., 2020). However, it remains unclear if cnidarian POU-IV *directly* regulates effector gene expression, as expected for a terminal selector. Furthermore, although POU-IV recognition element-like sequences have been previously identified in the *N. vectensis* genome based on sequence similarity to bilaterian POU-IV-binding motifs (Sebe-Pedros et al., 2018), cnidarian POU-IV recognition elements have not been experimentally defined, and consequently, whether the conservation of POU-IV-binding sequence extends beyond Bilateria remains vague.

Sea anemones together with corals form the clade Anthozoa, which is sister to the Medusozoa – a group characterized by typically having a pelagic medusa (jellyfish) stage - consisting of Staurozoa, Hydrozoa, Scyphozoa and Cubozoa (Collins et al., 2006, Zapata et al., 2015). Sea anemones have multicellular mechanosensory structures, known as the hair bundle mechanoreceptors, in the ectoderm of the oral feeding tentacles (Mire-Thibodeaux and Watson, 1994, Mire and Watson, 1997, Watson et al., 1997). A hair bundle mechanoreceptor consists of a central sensory cell surrounded by peripheral support cells (Figure 1 – Figure supplement 1). The central sensory cell exhibits morphological hallmarks of concentric hair cells, with an apical cilium surrounded by stereovilli, and basal neurites. Support cells contribute stereovilli or microvilli that encircle the apical ciliary-stereovillar structure of the central hair cell. The cilium and stereovilli of the central cell and stereovilli/microvilli of support cells are interconnected by

lateral linkages; in addition, extracellular linkages have been observed between the tips of stereovilli/microvilli of support cells, resembling the tip links of vertebrate mechanosensory hair cells (Watson et al., 1997). The apical sensory apparatus, or the hair bundle, of the mechanoreceptor thus consists of the cilium and stereovilli of the central hair cell and the peripheral stereovilli/microvilli of support cells (Mire and Watson, 1997). We note that in the literature, the support cells of hair bundle mechanoreceptors are sometimes referred to as hair cells (e.g. (Mire and Watson, 1997)). In this paper, in accordance with the morphological definition of cnidarian hair cells, a *hair cell* refers to the central sensory cell of the hair bundle mechanoreceptor, and a *support cell* refers to the cell that abuts the central sensory cell and contributes peripheral stereovilli/microvilli to the hair bundle.

Figure 1 - Figure supplement 1: Diagram of the hair bundle mechanoreceptor of sea anemones.

A cilium (ci) extends from a central hair cell (hc). It is encircled by stereovilli (sv), also produced by the hair cell, as well as smaller-diameter microvilli (mv) contributed by surrounding support cells (su), forming a cone-shaped structure. Abbreviations: ne neurites; nu nucleus

In this report, we use the starlet sea anemone *N. vectensis* to investigate the role of POU-IV in the development of cnidarian hair cells. *N. vectensis* is a convenient model for studies of

mechanisms of cnidarian development because of the availability of the genome sequence (Putnam et al., 2007) and a wide range of powerful molecular genetic tools including CRISPR-Cas9 genome editing (Ikmi et al., 2014, Nakanishi and Martindale, 2018). During embryogenesis, *N. vectensis* gastrulates by invagination to form an embryo consisting of ectoderm and endoderm separated by the extracellular matrix known as the mesoglea (Kraus and Technau, 2006, Magie et al., 2007). The blastopore becomes the mouth/anus (“oral”) opening of the animal (Fritzenwanker et al., 2007, Lee et al., 2007). The embryo develops into a free-swimming, ciliated planula larva, which transforms into a polyp with circum-oral tentacles that house mechanosensory hair cells in the ectoderm (Figure 1 - Figure supplement 2; (Nakanishi et al., 2012, Watson et al., 2009)). The polyp then grows and reaches sexual maturity. Previous studies have indicated that hair bundles of *N. vectensis* polyps are indeed sensitive to movement of surrounding water (Watson et al., 2009), and that stereovilli/microvilli of hair bundles express TRP (Transient Receptor Potential)-like cation channels (Mahoney et al., 2011) and a putative extracellular linkage component cadherin 23 (Watson et al., 2008). In the present work, we provide evidence that POU-IV regulates postmitotic differentiation of hair cells by directly activating effector genes that define mature cell identity.

Figure 1 - Figure supplement 2: Life cycle transition in the sea anemone cnidarian *Nematostella vectensis*

Confocal sections of *Nematostella vectensis* at metamorphosis from a free-swimming planula (A; 3-5 days-post-fertilization (dpf)), through the tentacle-bud stage (B; 5-7 dpf), into a primary polyp (C; 7-10 dpf). Filamentous actin is labeled with phalloidin (Pha). All panels show side

views of the animal with the oral pole facing up. A: The planula consists of ectoderm (ec) and endoderm (en) separated by an extracellular matrix, and develops muscle fibers (ms) in the endoderm. B: At the tentacle-bud stage, four tentacle primordia known as the tentacle buds (tb) emerge in the circumoral ectoderm. C: Four primary tentacles (te) and the body column then elongate along the oral-aboral axis, forming a primary polyp. Scale bar: 50 μ m

Results:

Sea anemone hair cell has an apical cilium surrounded by a circle of stereovilli and extends basal neuronal processes.

We first examined the structure of hair cells in the oral tentacles of the sea anemone *Nematostella vectensis* at the primary polyp stage by light and electron microscopy. We used phalloidin to label F-actin enriched in stereovilli of hair cells, and the lipophilic dye DiI to label the plasma membrane of hair cells. Hair cells are an epithelial cell type whose cell body is pear-shaped and occurs exclusively in the superficial stratum of pseudostratified ectoderm in oral tentacles (Figure 1). The hair cell has an apical cilium surrounded by 8 large-diameter stereovilli that extend actin filament-containing rootlets into the cytoplasm (Figure 1B-I). In primary polyps, the apical cilium is 10-15 μ m-long; stereovilli are 3-5 μ m-long and 200-400 nm in diameter; stereovillar rootlets are 2-3 μ m-long. Electron-lucent vesicles ranging from 50-100 nm in diameter are abundant in the cytoplasm of a hair cell (Figure 1I). Stereovilli of a hair cell are encircled by smaller-diameter microvilli (80-150 nm) contributed by adjacent support cells that are enriched in electron-dense vacuoles in the apical cytoplasm (Figure 1E-I). This multicellular apical sensory apparatus, consisting of the cilium and stereovilli of the hair cell surrounded by stereovilli/microvilli of support cells, constitutes the hair bundle (Mire and Watson, 1997). A subset of cnidocytes – nematocytes but not anthozoan-specific spirocytes – forms a morphologically similar apical mechanosensory apparatus known as the ciliary cone (Fautin and Mariscal, 1991); however, the ciliary cone of tentacular nematocytes in *N. vectensis* is less pronounced than that of hair cells, and consists of a single cilium surrounded by short microvilli (2-2.5 μ m long) that lack actin rootlets (Figure 1 - Figure supplement 3). Basally, a hair cell extends thin neuronal processes that likely form synapses with the tentacular nerve net and/or longitudinal muscle fibers located at the base of the ectodermal epithelium alongside mesoglea (Figure 1C, E, G).

Figure 1: Morphology of sea anemone hair cells.

A-F: Confocal sections of oral tentacles of *N. vectensis* at the primary polyp stage. Filamentous actin is labeled with phalloidin (Pha), and nuclei are labeled with DAPI (dapi). DiI is used to label cell membrane of a subset of hair cells. In A, the distal end of the tentacle is to the top, and in B-I, the apical surface of the ectodermal epithelium is to the top. A: sections through the tentacle. Numerous hair bundles (arrowheads) are evident on the tentacle surface. B: sections through the hair bundles at the tentacle tip, showing stereovilli (st) and their prominent rootlets (rl) of central hair cells. C-D: sections through a DiI-labeled hair cell (hc) at the tentacle tip. Note that the hair cell has an apical cilium (ci) surrounded at its base by stereovilli (st), and basally extended thin neurites (ne). An empty space within the cell body shows the location of a nucleus (nu), as evidenced by DAPI staining (D). E-H: sections through a DiI-labeled hair cell (hc) located near the tip of a tentacle. Arrowheads in F point to microvilli of the mechanoreceptor hair bundle contributed by peripheral support cells, which are DiI-negative. I: Electron microscopic section of an apical region of the tentacular ectodermal epithelium of *N. vectensis* polyp, showing a hair cell (hc) and a support cell (su). The hair cell has stereovilli that extend dense filaments into the cytoplasm, forming 2-3 μm -long rootlets (rl_{st}), as well as numerous clear vesicles (ve), while the support cell has apical microvilli (mv) and electron-dense vacuoles (va). Abbreviations: ms muscle fibers; rl_{ci} ciliary rootlet; ec ectoderm. Scale bar: 10 μm (A-F); 2 μm (inset in B); 500 nm (G)

Figure 1 - Figure supplement 3: Morphology of nematocytes in the tentacles of *Nematostella vectensis* polyps.

A, B: Confocal sections of *Nematostella vectensis* primary polyp, labeled with an antibody against tyrosinated β -tubulin (“tyrTub”). Filamentous actin is labeled with phalloidin (Pha), and mature cnidocysts (cnidae) are labeled with a high concentration of DAPI in the presence of EDTA (Szczepanek et al., 2002). The section shows the ectoderm at the tentacle tip, with the epithelial surface facing up. Note that nematocytes (nm) have an apical cilium (ci) surrounded by microvilli (mv) without rootlets. Stereovilli (st) and their prominent rootlets (rl) of hair cells are also shown. C: An electron micrograph of a section of *N. vectensis* primary polyp, showing an apical structure of a cnida(cn)-containing nematocyte in the tentacular ectoderm. Microvilli (mv) without rootlets occur on the apical cell surface. Scale bar: 10 μ m (A, B); 500 nm (C)

Hair cells commence development at metamorphosis in the sea anemone.

We next sought to determine the timing of hair cell development by using phalloidin to label stereovilli – the morphological hallmark of hair cells in *N. vectensis*. We never found stereovilli in the circumoral ectoderm during planula development (Figure 1 - Figure supplement 4A, B). However, pronounced stereovilli became evident in the circumoral ectoderm at the tentacle-bud stage (Figure 1 - Figure supplement 4C, D). These observations suggest that the hair cell is a postembryonic cell type that does not initiate development until metamorphosis in *N. vectensis*.

Figure 1 - Figure supplement 4: Hair cell development begins in the ectoderm of tentacle primordia at metamorphosis in sea anemones.

Confocal sections of *N. vectensis* at the late planula (A, B) and tentacle-bud (C, D) stages. Filamentous actin is labeled with phalloidin (Pha), and nuclei are labeled with DAPI (dapi). All panels show side views of animals with the blastopore/mouth facing up. A and C show longitudinal sections through the center, and B and D show surface ectoderm of tentacular primordia boxed in A and C, respectively. Note that 3-5 μm -long stereovilli characteristic of hair cells become evident at the tentacle bud stage (arrowheads in D), indicative of hair cell differentiation. Scale bar: 50 μm (A, C); 10 μm (B, D)

Class IV POU transcription factor is postmitotically expressed in hair cells in the sea anemone.

The *N. vectensis* genome contains a single gene that encodes the class IV POU homeodomain transcription factor (Nv160868; (Tourniere et al., 2020, Nakanishi et al., 2010, Gold et al., 2014)), termed as *NvPOU4* by (Tourniere et al., 2020); in this paper, we will simplify the

nomenclature by referring to *POU-IV/POU4/Brn3/unc-86* gene as *pou-iv* and its protein product as POU-IV. It has been previously shown that *pou-iv* mRNA is strongly expressed in circum-oral ectoderm during metamorphosis in *N. vectensis* (Tourniere et al., 2020), consistent with a role in tentacular morphogenesis. Although gene expression analysis using a transgenic reporter line has indicated that *pou-iv* is expressed in cnidocytes throughout the body including those in the tentacles (Tourniere et al., 2020), whether *pou-iv* is expressed in mechanosensory hair cells is not known. To address this, we first developed a rabbit polyclonal antibody against an N-terminal, non-DNA-binding region of the *Nematostella vectensis* POU-IV based on the amino acid sequence predicted from *pou-iv* cDNA (see Materials and Methods). As detailed in the next section, specificity of the antibody was confirmed by western blot analysis using *pou-iv* mutants and their wildtype siblings. In addition, immunostaining and *in situ* hybridization experiments showed that the pattern of anti-POU-IV immunoreactivity paralleled that of *pou-iv* mRNA expression (Figure 2 – Figure supplement 1), further supporting the specificity of the antibody. We therefore used immunostaining with the anti-POU-IV to analyze the expression pattern of POU-IV in developing oral tentacles in *N. vectensis* at metamorphosis. We found that POU-IV protein localized to nuclei of differentiating and differentiated hair cells, but not to those of support cells, in the ectoderm of developing tentacles (Figure 2A-L). In addition, we confirmed POU-IV expression in cnidocytes (Figure 2 - Figure supplement 2), consistent with the previous report (Tourniere et al., 2020). Nuclear labeling by the anti-POU-IV was abolished when the antibody was preadsorbed with the POU-IV antigen prior to immunostaining (Figure 2 - Figure supplement 3), evidencing that the antibody reacts with nuclear POU-IV.

We then carried out EdU pulse labeling experiments to test whether any of the POU-IV-expressing cells in the tentacular ectoderm were at S-phase and thus proliferative. As observed for *pou-iv* transcript-expressing cells during embryogenesis (Tourniere et al., 2020), we found that none of the POU-IV-expressing epithelial cells in the developing tentacles examined (n>220 cells across 3 tentacle-bud-stage animals and 8 primary polyps) incorporated EdU (e.g. Figure 2M-P), indicative of their postmitotic cell-cycle status. Taken together, the gene expression pattern suggests a role for POU-IV in postmitotic development of mechanosensory hair cells, as well as cnidocytes, in the tentacles of *N. vectensis* polyps.

Figure 2: POU-IV is postmitotically expressed in hair cells of tentacular ectoderm at metamorphosis in the sea anemone.

Confocal sections of *N. vectensis* at metamorphosis, labeled with antibodies against POU-IV, and/or tyrosinated β -tubulin ("tyrTub"). Filamentous actin is labeled with phalloidin (Pha), and nuclei are labeled with DAPI (dapi). Proliferative cells are labeled by the thymidine analog EdU. A shows a section through the presumptive tentacle primordia with the blastopore/mouth facing up. E, I, M-P show sections through developing oral tentacles with the distal end of the tentacle facing up; M-P are tangential sections of tentacles at the level of the surface ectoderm and parts of the endoderm (en). B-D, F-H, and J-L are magnified views of the boxed regions in A, E and I,

respectively, with the apical epithelial surface facing up. A-D: late planula. E-H: tentacle-bud. I-P: primary polyp. At the late planula stage prior to hair cell differentiation, POU-IV-positive nuclei are primarily localized at the basal and middle layers of the ectoderm of presumptive tentacle primordia (arrowheads in B-D); few POU-IV-positive nuclei are detectable at the superficial stratum. At the tentacle-bud stage, hair cells with pronounced stereovilli (st) and POU-IV-positive nuclei begin to develop in the superficial stratum of the ectodermal epithelium in tentacle primordia (arrowheads in F-H). POU-IV-positive nuclei in the superficial layer specifically occur in hair cells (white arrowheads in J-L) and not in adjacent support cells (purple arrowheads in J-L). The inset in J shows a magnified view of stereovilli (st) of a POU-IV-positive hair cell; note the presence of stereovillar rootlets (rl). In addition to hair cells, cnidocytes express POU-IV in the tentacular ectoderm (Figure 2 - Figure supplement 2; (Tournier et al., 2020)). POU-IV-positive cells are EdU-negative (arrowheads in I-L), evidencing their postmitotic cell-cycle status. Abbreviations: ec ectoderm; en endoderm. Scale bar: 10 μ m (A-P); 2 μ m (inset in J)

Figure 2 – Figure supplement 1: The pattern of immunoreactivity with an anti-POU-IV antibody recapitulates that of *pou-iv* mRNA expression.

Confocal sections of oral tentacles of *Nematostella vectensis* at the primary polyp stage, labeled with an anti-POU-IV antibody (“POU-IV protein”; A-D), or an antisense riboprobe against *pou-*

377 *iv* mRNA (“*pou-iv* mRNA”; E-H). Microtubules are labeled by antibodies against tyrosinated δ -
 378 tubulin (“*tyrTub*”; A, B) or acetylated δ -tubulin (“*acTub*”; E, F). Nuclei are labeled with DAPI
 379 (dapi). All panels show sections through oral tentacles with the distal end of the tentacle facing
 380 up. Anti-POU-IV immunoreactivity occurs in a subset of apical nuclei (white arrowhead in A-D)
 381 and the majority of basal nuclei (grey arrowhead in A-D) within the pseudostratified ectodermal
 382 epithelium (ec), suggesting nuclear localization of POU-IV proteins in these cells. Consistent
 383 with the pattern of Anti-POU-IV immunoreactivity, *in situ* hybridization signals of *pou-iv*
 384 mRNA occur in a subset of ectodermal epithelial cells whose cell bodies are located apically
 385 (white arrowhead in E-H), and in those ectodermal cells whose cell bodies are localized basally
 386 (bracketed in E-H). Anti-POU-IV immunoreactivity is absent in nuclei of tentacular endodermal
 387 cells (en in A-D), and *in situ* hybridization signals for *pou-iv* mRNA are not detectable in these
 388 cells (en in E-H), indicating the lack of *pou-iv* expression in tentacular endoderm. Scale bar: 10
 389 μ m
 390

392 **Figure 2 - Figure supplement 2: POU-IV localizes to the nuclei of cnidocytes in tentacular**
 393 **ectoderm of the sea anemone.**

394 Confocal sections of an oral tentacle of *N. vectensis* at the primary polyp stage, labeled with
 395 antibodies against POU-IV and/or minicollagen 3 (“*Ncol3*”; (Zenkert et al., 2011)). Nuclei are
 396 labeled with DAPI (dapi). Arrowheads show POU-IV positive nuclei of *Ncol3*-positive
 397 cnidocytes that reside in the tentacular ectoderm. Scale bar: 10 μ m
 398

Figure 2 – Figure supplement 3: Immunostaining with a preadsorbed anti-POU-IV antibody.

Confocal sections of oral tentacles of *Nematostella vectensis* at the primary polyp stage, labeled with an anti-POU-IV antibody (“anti-POU-IV”; A-C) and an anti-POU-IV antibody preadsorbed with the POU-IV antigen (CQPTVSESQFDKPFETPSPINamide) used to generate the antibody (“Preadsorbed anti-POU-IV”; D-F). Nuclei are labeled with DAPI (dapi). In all panels, the distal end of the tentacle is to the top, and sections are at the level of surface ectoderm. Arrowheads indicate nuclear immunoreactivity that is abolished when the preadsorbed antibody is used, indicating that the anti-POU-IV reacts with nuclear POU-IV. Scale bar: 10 μ m

Generation of POU-IV mutant sea anemones

To investigate the function of POU-IV in hair cell development in *N. vectensis*, we generated a *pou-iv* mutant line by CRISPR-Cas9-mediated mutagenesis. First, a cocktail containing *pou-iv*-specific single guide RNAs (sgRNAs) and the endonuclease Cas9 protein was injected into fertilized eggs to produce founder (F0) animals. Multiple sgRNAs were designed to cleave

flanking sites of the coding region of the *pou-iv* locus (Figure 3A; Figure 3 - Figure supplement 1). Large deletions were readily confirmed by genotyping PCR using genomic DNA extracted from single CRISPR-injected embryos (Figure 3 - Figure supplement 1). DNA sequencing of mutant bands confirmed that excision of both POU- and homeo-domains could be induced by this approach. F0 animals were raised and crossed with wildtype animals, in order to generate F1 heterozygous animals carrying a *pou-iv* knockout allele. Mutant allele carriers were identified by genotyping individual F1 polyps. One of the mutant alleles, which will be here referred to as *pou-iv⁻*, had a 705bp deletion that removed most of the POU domain (i.e. all but the first four N-terminal residues) and all of the homeodomain at the *pou-iv* locus (Figure 3B; Figure 3 - Figure supplement 2). This mutant allele differs from the previously generated *NvPOU4⁻* allele which harbors a frameshift mutation (31 bp deletion) at the start of the POU-domain-encoding sequence (Tourniere et al., 2020). F1 *pou-iv* +/- heterozygotes were subsequently crossed with each other to produce F2 offspring, a quarter of which, on average, were *pou-iv* -/- mutants. *pou-iv* -/- mutants were identified by PCR-based genotyping methods (Figure 3B, C) using genomic DNA extracted from polyp tentacles (Ikmi et al., 2014) or from pieces of tissue isolated from early embryos (Nakanishi and Martindale, 2018, Silva and Nakanishi, 2019). Western blotting with the anti-POU-IV has confirmed that *pou-iv* -/- polyps express mutant POU-IV lacking DNA-binding domains (18.7kDa), but not wildtype POU-IV (35.2kDa) (Figure 3D), validating the specificity of the antibody.

expected size for mutant POU-IV lacking DNA-binding domains (18.7kDa) is detectable in mutant (-/-) extracts, but not in wildtype (+/+) extracts (purple arrowhead). The band just below the expected size of the wildtype POU-IV occur in both wildtype and mutant protein extracts, and therefore represents non-POU-IV protein(s) that are immunoreactive with the anti-POU-IV antibody.

Figure 3 – Source data 1: An original gel image used to generate Figure 3C and the original image with relevant lanes labelled.

Figure 3 – Source data 2: An original western blot image used to generate Figure 3D (top; anti-*N. vectensis* POU-IV) and the original image with relevant lanes labelled.

Figure 3 – Source data 3: An original western blot image used to generate Figure 3D (bottom; anti-acetylated α -tubulin) and the original image with relevant lanes labelled.

Figure 3 – Figure supplement 1: Generation of *pou-iv* F0 mosaic mutants by CRISPR-Cas9-mediated mutagenesis in *N. vectensis*.

A schematic view of the *pou-iv* locus (left), and genomic DNA PCR results of an uninjected wildtype embryo (“WT”) and an F0 embryo injected with locus-specific sgRNAs and Cas9 (“*pou-iv* F0”) (right). A blue bar shows the predicted translation start site, and a red bar shows the predicted translation termination site. The orange and yellow highlighted regions are POU- and Homeo- DNA-binding domains, respectively. Purple arrowheads show sgRNA target sites. Blue arrows mark regions targeted in the PCR analysis shown to the right. Note that genomic PCR of the WT embryo shows expected sizes of PCR fragments (1495 bp for primary PCR (“a”), and 1263 bp for secondary nested PCR (“b”)), while F0 embryos show additional bands of smaller sizes (arrowheads), indicating that targeted deletions of different sizes have occurred

mosaically in each embryo. DNA sequencing of the <500 bp band (lower arrowhead) indicated that this mutant allele harbored 981 bp deletion encompassing the POU- and homeo-domain-encoding regions.

Figure 3 – Figure supplement 1 - Source data 1: An original gel image used to generate Figure 3 – Figure supplement 1 (right) and the original image with relevant lanes labelled.

Figure 3 – Figure supplement 2: Sequence alignment of wildtype and mutant *pou-iv* alleles.

A: Diagrams of the *pou-iv* locus and the disrupted mutant allele (*pou-iv⁻*). Blue bars show predicted translation start sites; red bars show predicted translation termination sites. In the schematic depicting the *pou-iv* locus, filled boxes indicate exons, and the regions that encode the POU- and homeo-domains are highlighted in orange (“POU”) and yellow (“HD”), respectively. In the schematic of the *pou-iv⁻* allele, deletion mutation is boxed in orange dotted lines. B: Alignment of nucleotide and translated amino acid sequences of wildtype (“+”) and mutant (“*pou-iv⁻*”) alleles boxed in A. POU- and homeo-domains are boxed in orange and yellow, respectively. Predicted translation termination sites are boxed in red, and 705 bp deletion mutation is boxed in dotted blue lines. Note that all but the first four residues of the POU domain and the entire homeodomain are deleted in the *pou-iv⁻* allele.

POU-IV is necessary for touch-response behavior of tentacles in the sea anemone.

If POU-IV indeed plays a key role in postmitotic differentiation of mechanosensory hair cells, mechanosensitive behaviors of oral tentacles are expected to be perturbed in *pou-iv* null mutants. We tested this prediction by using F2 *pou-iv* ^{-/-} mutants and their heterozygous and wildtype siblings. In wildtype polyps, oral tentacles typically respond to touch stimuli by local contraction of longitudinal muscles. Strong and repeated touch stimuli of tentacles lead to excitation of longitudinal muscles in the body column, causing the tentacles to retract into the body column. In this study, a hair held in a tungsten needle holder was quickly depressed on the distal portion

of each tentacle, and the presence/absence of the touch-response behavior of tentacles was scored for each animal. 100% of the F2 *pou-iv* *+/+* wildtype animals that were examined (n=44) contracted at least one tentacle in response to touch (Figure 4A, B; Figure 4 – video 1). In contrast, we observed that only 35% of the F2 *pou-iv* *-/-* knockout animals (n=40) showed any sign of tentacular retraction in response to touch; 65% of the knockout mutants exhibited no discernable tentacular response to tactile stimuli (Figure 4C, D; Figure 4 – video 2). The majority of F2 *pou-iv* *+/-* heterozygotes (87%, n=62) showed touch-induced, tentacular responses; 13% did not show touch responses, suggestive of dose-dependent effects of POU-IV expression on the mechanosensory behavior. The reduced tentacular response to touch in *pou-iv* *-/-* mutants is not due to the inability of tentacular muscles to contract, as *pou-iv* *-/-* mutants responded to crushed brine shrimp extract by contracting tentacles (100%, n=8 animals; Figure 4 - Figure supplement 1; Figure 4 - video 3, 4). Hence, *pou-iv* is specifically required for the touch-sensitive behavior of oral tentacles in *N. vectensis*, consistent with POU-IV having a role in regulating the development of the mechanosensory hair cells.

Figure 4: POU-IV is essential for touch-response behavior in the sea anemone.

A-D: Behavior of wildtype (F2 *pou-iv* +/+, A, B) and mutant (F2 *pou-iv* -/-, C, D) *N. vectensis* polyps in response to tactile stimuli to their oral tentacles. A hair held in a tungsten needle holder was used to touch the distal portion of each tentacle. Animals before (A, C) and after (B, D) tentacle touch are shown. Tactile stimuli to tentacles elicit tentacular retraction in the wildtype individual (100%, n=44; A, B). In contrast, the majority of *pou-iv* homozygous mutants were touch-insensitive (65%, n=40; B, D); only 35% of the animals showed any contractile response to touch stimuli. Arrowheads in B point to retracted tentacles. Scale bar: 1 mm

Figure 4 – Figure supplement 1: *pou-iv* mutants respond to brine shrimp extract by tentacular contraction and pharyngeal protrusion.

A-D: Behavior of wildtype (F2 *pou-iv* +/+, A, B) and mutant (F2 *pou-iv* -/-, C, D) *N. vectensis* polyps in response to exposure to *Artemia* extract. Both wildtype and mutant animals contracted tentacles (arrowheads in B and D), and protruded the pharynx (ph) within one minute of exposure to the extract (F2 *pou-iv* +/+, 100%, n=3; F2 *pou-iv* -/-, 100%, n=9). Scale bar: 1 mm

Figure 4 - Video 1: Touch-sensitive behavior of a wildtype (F2 *pou-iv* +/+) polyp.

Figure 4 - Video 2: Touch-sensitive behavior of a *pou-iv* mutant (F2 *pou-iv* -/-) polyp.

Figure 4 - Video 3: Behavior of a wildtype (F2 *pou-iv* +/+) polyp upon exposure to brine shrimp extract.

Figure 4 - Video 4: Behavior of a *pou-iv* mutant (F2 *pou-iv* -/-) polyp upon exposure to brine shrimp extract.

POU-IV is necessary for normal development of hair cells in the sea anemone.

To understand the structural basis of touch-insensitivity in *pou-iv* null mutants, we examined the morphology of tentacular cells in *pou-iv* null mutants and their heterozygous and wildtype siblings by light and confocal microscopy. At the primary polyp stage, F-actin labeling by phalloidin showed that the longitudinal muscles in the tentacle of F2 *pou-iv* $-/-$ mutants developed normally (Figure 5A, D, G), consistent with the behavioral evidence demonstrating the ability of the mutant tentacles to contract in response to the brine shrimp extract. We have confirmed the previous finding that mature cnidocytes with cnidae fail to develop in *pou-iv* knockout mutants (Figure 5C, F, I; (Tourniere et al., 2020)). In addition, we found that mature hair cells with stereovillar rootlets were lacking in the tentacles of F2 *pou-iv* $-/-$ polyps (n=6 animals), while mature hair cells formed normally in tentacles of *pou-iv* $+/-$ (n=6 animals) and *pou-iv* $+/+$ (n=3 animals) siblings (Figure 5A-I). Ciliary cone-like structures lacking stereovillar rootlets occurred in *pou-iv* $-/-$ mutants (Figure 5G-I), raising the possibility that hair cells might undergo partial differentiation in *pou-iv* $-/-$ mutants.

Electron microscopic observations confirmed these findings. Stereovillar rootlets and cnidae were absent in the tentacles of F2 *pou-iv* $-/-$ polyps (n=2 animals) but were present in the tentacles of their wildtype siblings (n=2 animals) (Figure 5J-L; Figure 5 - Figure supplement 1, 2). We also confirmed by electron microscopy the presence of a hair-cell-like cell that has an apical ciliary cone without stereovillar rootlets, surrounded by support cells with characteristic electron-dense vacuoles that contribute microvilli to the ciliary cone in *pou-iv* $-/-$ mutants (Figure 5K, L); ciliary rootlets were observed in these hair-cell-like cells in *pou-iv* $-/-$ mutants (Figure 5 – Figure supplement 3).

The lack of cnidae is consistent with the inability of *pou-iv* null mutants to capture prey as previously reported (Tourniere et al., 2020), but cannot explain the lack of tentacular contraction in response to touch. Stereovillar rootlets provide stereovilli with structural resilience against physical damage and are necessary for normal mechanosensitivity in vertebrate hair cells (Kitajiri et al., 2010). We therefore suggest that touch-insensitivity of oral tentacles in *pou-iv* null mutants results, at least in part, from the failure of hair cells to generate structurally robust apical mechanosensory apparatus (see Discussion).

Figure 5: POU-IV is necessary for hair cell differentiation in the sea anemone.

A-I: Confocal sections of oral tentacles of wildtype (F2 *pou-iv* +/+, A-C), heterozygous (F2 *pou-iv* +/-, D-F) and homozygous *pou-iv* mutant (F2 *pou-iv* -/-, G-I) *N. vectensis* polyps. Filamentous actin is labeled with phalloidin (Pha), and nuclei are labeled with DAPI (dapi). In all panels, the distal end of the tentacle is to the top. A, D, G: sections through the tentacle. B, C, E, F, H, I: sections through hair bundles/ciliary cones at the tip of tentacles. Ciliary cones occur on the epithelial surface of the tentacle regardless of the genotype (arrowheads in A, D, G). However, stereovilli (st) with rootlets (rl) characteristic of mechanosensory hair cells are observed in wildtype (B) and heterozygous (E) siblings, but not in homozygous *pou-iv* mutants whose ciliary cones contain microvilli without prominent actin rootlets (mv in H). Arrowheads in C and F indicate spaces occupied by cnidocysts in wildtype and heterozygous siblings, respectively, which are absent in *pou-iv* homozygous mutants (I; Figure 5 - Figure supplement 1). J-L: Electron microscopical sections of a hair cell of a F2 *pou-iv* +/+ polyp (J) and an epithelial cell with hair-cell-like morphologies in a F2 *pou-iv* -/- polyp (K, L). In all panels, apical cell surfaces face up. K and L are sections of the same cell at different levels. The hair cell-like epithelial cell of the mutant has a central apical cilium surrounded by a collar of rootlet-less microvilli (mv in K, L), which are encircled by microvilli of the adjacent support cells (mv_s in L), forming a ciliary cone. It also has numerous clear vesicles (ve in L) in the cytoplasm, characteristic of hair cells (ve in J; Figure 1G). Support cells of mutants are morphologically indistinguishable from those of wildtype animals, having characteristic large electron-dense vacuoles (va in L) in addition to

apical microvilli (mv_s in L) that contribute to the ciliary cone/hair bundle. Consistent with light microscopy data (A-C, G-I), stereovillar rootlets (rl_{st}) are absent in the F2 *pou-iv* $-/-$ polyp, but are present in hair cells of their wildtype siblings (J). In K and L, regions of apical cytoplasm where stereovillar rootlets would normally be observed are boxed with dotted purple lines. Abbreviations: ms muscle fibers; ec ectoderm; st stereovilli; ci cilium; rl_{st} stereovillar rootlets. Scale bar: 10 μ m (A-I); 2 μ m (insets in B, E, H); 500 nm (J-L)

Figure 5 – Figure supplement 1: *pou-iv* mutants lack mature cnidocytes.

Electron microscopic sections of tentacular ectoderm of a F2 *pou-iv* $+/+$ polyp (A) and F2 *pou-iv* $-/-$ polyp (B). Note the occurrence of numerous cnidae – nematocysts (nc) and spirocysts (sp) – in A that are absent in B. Abbreviations: nu nucleus; ec ectoderm; en endoderm; ms muscle fibers; mg mesoglea. Scale bar: 2 μ m

Figure 5 – Figure supplement 2: F2 *pou-iv* wildtype siblings develop hair cells with stereovillar rootlets.

Electron microscopic sections of hair cells of a F2 *pou-iv* +/+ polyp. In all panels, apical cell surfaces face up. Note the presence of stereovillar rootlets (arrowheads). Abbreviations: st stereovilli; ci cilium. Scale bar: 500 nm

Figure 5 – Figure supplement 3: Hair-cell-like cells of *pou-iv* mutants have ciliary rootlets.

Electron microscopic sections of a hair-cell-like cell of a F2 *pou-iv* *-/-* polyp, characterized by ciliary-cone-associated microvilli (mv). The apical cell surface faces up. Note the presence of a ciliary rootlet (rl_{ci}). Abbreviations: ve vesicles. Scale bar: 500 nm

POU-IV is necessary for maturation, but not initial differentiation or survival, of hair cells in the sea anemone.

The lack of functional hair cells in *pou-iv* *-/-* mutants is consistent with POU-IV having a necessary role in initial differentiation and/or maturation of hair cells. In order to more precisely define the functional role of POU-IV in hair cell development, we investigated the morphological and molecular characteristics of epithelial cells expressing the mutant form of POU-IV, which we refer to as POU-IV(-), in tentacular ectoderm of *pou-iv* *-/-* mutants. Because the epitope that the anti-POU-IV antibody reacts with is intact in the protein encoded by the *pou-iv* allele (Figure 3A, B, D), it was possible to use immunostaining with the anti-POU-IV to localize POU-IV(-) in *pou-iv* *-/-* mutants. A number of epithelial cells in the tentacular ectoderm were found to express POU-IV(-) (Figure 6A-C). In contrast to the primarily nuclear localization

of POU-IV in wildtype animals (cf. Figure 2), however, POU-IV(-) is distributed throughout the cytoplasm of POU-IV(-)-expressing cells in *pou-iv* ^{-/-} mutants (Figure 6A-F), presumably due to the lack of nuclear localization signal (located at the N-terminal end of the homeodomain (Sack et al., 1996)) in POU-IV(-) (Figure 3B). We found that the epithelial cells bearing apical ciliary cones in *pou-iv* ^{-/-} mutants expressed POU-IV(-) (Figure 6D-F) and therefore could represent partially differentiated hair cells that failed to undergo maturation. Alternatively, as ciliary cones characterize nematocytes in wildtype *N. vectensis*, it was possible that these ciliary cone-bearing epithelial cells in *pou-iv* ^{-/-} mutants were immature nematocytes without cnidae.

To clarify the identity of ciliary cone-bearing epithelial cells in *pou-iv* ^{-/-} mutants, we used an antibody against a pan-cnidocyte differentiation marker minicollagen 3 (Ncol3; (Babonis and Martindale, 2017, Zenkert et al., 2011)) to label immature cnidocytes. It was previously shown that Ncol3 was expressed in a subset of ectodermal epithelial cells of *pou-iv* knockout mutants despite the lack of mature cnidae, indicating that immature cnidocytes are present in *pou-iv* mutants and that *pou-iv* is not necessary for initial differentiation of cnidocytes (Tourniere et al., 2020). By using immunostaining with an anti-Ncol3, we confirmed that Ncol3-positive immature cnidocytes in *pou-iv* ^{-/-} mutants indeed expressed POU-IV(-) (Figure 6G-I). However, none of the Ncol3-positive immature cnidocytes in *pou-iv* ^{-/-} mutants had distinct apical ciliary cones (e.g. Figure 6J-L), suggesting that ciliary cone-bearing epithelial cells in *pou-iv* ^{-/-} mutants represent immature hair cells, and not immature nematocytes. Thus, hair cells appear to be present in their immature, yet morphologically differentiated, form in *pou-iv* ^{-/-} mutants. The presence of partially differentiated hair cells in *pou-iv* ^{-/-} mutants supports the hypothesis that POU-IV regulates maturation, but not initial differentiation, of hair cells in *N. vectensis*.

As discussed above, the absence of stereovillar rootlets in hair cells of *pou-iv* ^{-/-} mutants may underlie the observed touch-insensitivity of the mutants. It was also possible that these immature hair cells failed to extend basal neurites to form normal mechanosensory neural circuits. To examine this possibility, we visualized the morphology of immature hair cells in *pou-iv* ^{-/-} mutants by using a *pou-iv::kaede* transgenic reporter construct, in which the 3.2 kb genomic sequence upstream of the start codon of the *pou-iv* gene was placed in front of the Kaede fluorescent protein-encoding gene (Ando et al., 2002). We first confirmed that the *pou-iv::kaede* reporter construct indeed drove the expression of Kaede in POU-IV-positive cell types - hair cells and cnidocytes - in tentacular ectoderm of wildtype animals, recapitulating the

endogenous POU-IV expression pattern (Figure 6 - Figure supplement 1). Interestingly, we unexpectedly found that cnidocytes, in addition to hair cells, had basal neurite-like processes (Figure 6 - Figure supplement 1I-L), which has never been reported in cnidarian literature to our knowledge. We then injected *pou-iv::kaede* plasmids into *pou-iv* F2 zygotes, which were allowed to develop into primary polyps, and subsequently carried out immunostaining with antibodies against Kaede and Ncol3. Animals lacking mature cnidae based on Ncol3 staining were assumed to be *pou-iv* ^{-/-} mutants. In these presumptive mutants, Kaede-positive immature hair cells were readily identifiable based on morphology and position; their cell bodies were pear-shaped and located in the superficial stratum of the tentacular ectoderm, some of which contained apical microvilli that are organized into a ciliary cone-like microvillar structure (Figure 6M, N). These immature hair cells, however, developed morphologically normal basal neurites (Figure 6O), indicating that *pou-iv* is not necessary for neurite extension in hair cells. Neither is *pou-iv* required for the development of basal neurite-like processes in cnidocytes; basal processes were observed in Ncol3-positive immature cnidocytes (Figure 6 - Figure supplement 2).

In mice, one of the *pou-iv* paralogs - *brn3c* – is thought to be required for survival of hair cells because the number of apoptotic cells increases in the inner ear sensory epithelia in *Brn-3c* null mutant mice (Xiang et al., 1998). We have therefore tested whether *pou-iv* regulates hair cell survival in *N. vectensis*, by carrying out the terminal deoxynucleotidyl transferase dUTP nick end-labeling (TUNEL) assay in *pou-iv* ^{-/-} mutants. We found that none of the POU-IV(-)-expressing epithelial cells examined in the tentacular ectoderm (n= 100 cells across 5 primary polyps) had TUNEL-positive, pyknotic nuclei indicative of apoptotic DNA fragmentation, although TUNEL-positive nuclear fragments were commonly observed in the endoderm (Figure 6P-R). Thus, in sea anemones, POU-IV does not appear to be directly involved in the survival of hair cells.

Figure 6: POU-IV is necessary for maturation of hair cells in the sea anemone.

Confocal sections of oral tentacles in F2 *pou-iv*^{-/-} *N. vectensis* polyps, labeled with antibodies against tyrosinated β -tubulin (“tyrTub”), minicollagen 3 (“Ncol3”; (Zenkert et al., 2011)) mutant POU-IV (“POU-IV(-)”), and/or Kaede fluorescent protein (“pou-iv::kaede”). DNA fragmentation is labeled by TUNEL. Filamentous actin is labeled with phalloidin (Pha), and nuclei are labeled with DAPI (dapi). In all panels, the apical surface of the tentacular ectodermal epithelium is to the top. A-C: sections through developing oral tentacles with the distal end of the tentacle facing up. Arrowheads point to a subset of POU-IV(-)-expressing epithelial cells, which are abundant in the tentacular ectoderm (ec). Note the cytoplasmic distribution of the POU-IV(-) likely resulting from the lack of nuclear localization signal. D-F: sections showing ciliary cone microvilli (mv)-bearing cells. Ciliary cone-bearing epithelial cells express POU-IV(-) (arrowheads). G-I: sections at the level of surface ectoderm of developing oral tentacles with the distal end of the tentacle facing up. A subset of POU-IV(-)-expressing cells are Ncol3-positive (arrowheads), representing immature cnidocytes. J-L: sections showing an immature cnidocyte which expresses POU-IV(-) and Ncol-3. Note that the cell bears apical microvilli (mv) that do not form a ciliary cone. M-O: sections showing immature hair cells in F2 *pou-iv*^{-/-} *N. vectensis* polyps injected with *pou-iv::kaede* construct. Note the presence of ciliary cone microvilli (mv) and basal neurites (ne). P-R: sections through tentacles with the distal end facing up. White arrowheads point to nuclei of POU-IV(-)-expressing ectodermal epithelial cells, which are TUNEL-negative. TUNEL-positive, pyknotic nuclei are observed in the endoderm (purple arrowheads). Abbreviations: ec ectoderm; en endoderm; ms muscle fiber. Scale bar: 10 μ m

Figure 6 – Figure supplement 1: *pou-iv::Kaede* reporter construct drives transgene expression in hair cells and cnidocytes.

Confocal sections of oral tentacles of *N. vectensis* polyps injected with *pou-iv::kaede* construct, labeled with antibodies against Kaede (“*pou-iv::kaede*”) and minicollagen 3 (“Ncol3”).

Filamentous actin is labeled with phalloidin (Pha), and nuclei are labeled with DAPI (dapi).

Panels A-H show tangential sections at the level of the surface ectoderm. Panels I-L show a side view of a cnidocyte (cn), with the apical cell surface facing up. *pou-iv::Kaede* expression occurs

in hair cells (hc in A-D) and cnidocytes (cn in A-L). Arrowheads in A-C indicate stereovilli of

hair cells. Cnidocytes have basally localized nuclei (nu) and extend basal neurite-like processes

(ne). Scale bar: 10 μm

Figure 6 – Figure supplement 2: Cnidocytes in *pou-iv* null mutants develop basal processes.

Confocal sections of a cnidocyte (cn) in the body column of a F2 *pou-iv* +/+ polyp injected with *pou-iv::kaede* construct and labeled with antibodies against Kaede (“*pou-iv::kaede*”) and minicollagen 3 (“Ncol3”). Nuclei are labeled with DAPI (dapi). Panels show tangential sections at the level of the basal ectoderm. Note that the cnidocyte extends neurite-like processes (ne). Scale bar: 10 μm

POU-IV-binding motifs are conserved across Cnidaria and Bilateria.

The evidence presented above thus indicates that in *N. vectensis*, POU-IV is involved in the maturation of mechanosensory hair cells – in addition to that of cnidocytes (Tourniere et al., 2020). How, then, does POU-IV control the development of these two distinct mechanosensory cell types? One possibility is that the POU-IV transcription factor regulates the expression of a shared set of genes critical for differentiation of both cell types. Given that both hair cells and cnidocytes are mechanosensory, POU-IV might control the expression of the same set of mechanotransduction genes in these cell types. Another possibility is that POU-IV regulates the expression of distinct sets of genes in different neural cell types, actively driving the differentiation of the two mechanosensory cell types.

To begin to address this question, we identified genome-wide binding sites for POU-IV by chromatin immunoprecipitation sequencing (ChIP-seq) using the antibody against *N. vectensis* POU-IV. We used adult *N. vectensis* for this experiment, because 1) neurogenesis continues through adulthood (e.g. (Havrilak et al., 2021)), and 2) we needed over 1 gram of tissue samples, which was more difficult to obtain from other developmental stages including primary polyps. We sequenced anti-POU-IV immunoprecipitated DNA and input DNA, and mapped the reads to the *N. vectensis* genome (Putnam et al., 2007). We identified 12,972 genomic sites that were enriched in ChIP DNA (i.e. ChIP peaks) (Figure 7 – Source data 1). We

then performed a *de novo* motif search and motif enrichment analysis, and found three motifs *rwrwaatmatgcattattaatatt* (motif 1; $E=5.2e-075$), *rmataaataatgcattattatky* (motif 2; $E=1.2e-052$) and *tkcataaataattatgmm* (motif 3; $E=4.8e-36$) that were enriched towards the center of ChIP peaks ($p=3.5e-368$ and $p=1.0e-138$, respectively) (Figure 7A). When we compared these three motifs against the Jaspar database (Fornes et al., 2020), we discovered that they showed significant sequence similarity to *Homo sapiens* POU4F1, POU4F2 and POU4F3 binding motifs (Figure 7B; $p < 10^{-5}$, $p < 10^{-3}$ and $p < 10^{-3}$, respectively), indicative of deep conservation of POU-IV-binding motifs across Cnidaria and Bilateria. Indeed, the motifs we have identified contain the sequence AT(A/T)ATT(A/T)AT (shown in bold in motif sequences above), which is nearly identical to the core recognition sequence of bilaterian POU-IV, AT(A/T)A(T/A)T(A/T)AT (Gruber et al., 1997). In addition, the preference of GC residues 5' to the core recognition sequence is evident in motifs 1 and 2 (underlined in motif sequences above), and in bilaterian POU-IV binding sequences (Gruber et al., 1997), and therefore appears to be conserved. We tested the ability of POU-IV to bind to the core recognition motif-like sequences by electrophoretic mobility shift assays (EMSAs), and confirmed that they were indeed essential for DNA recognition by POU-IV (Figure 7C). We infer that in the last common ancestor of Cnidaria and Bilateria, POU-IV bound to the consensus DNA element GCAT(A/T)ATT(A/T)AT to regulate gene expression.

Figure 7: POU-IV-binding motifs are conserved across Cnidaria and Bilateria.

A: Motifs enriched in *Nematostella vectensis* POU-IV ChIP-seq peaks. B: *Homo sapiens* POU motifs resulting from sequence alignment and comparison against the Jaspar database. The p-value reported corresponds to the highest p value for any of the three *Nematostella vectensis*

POU4 motifs found. C: Electrophoretic mobility shift assay (EMSA) using purified *N. vectensis* POU-IV protein and a 50 bp DNA probe containing the conserved core motif *CATTATTAAT*. Note that retardation of probe migration occurs in the presence of POU-IV protein (blue arrowhead; lane 2), indicative of formation of the protein-DNA complex. Retardation is inhibited in the presence of an unlabeled competitor probe (“Specific competitor”; lane 3). Removal of the motif sequence in the probe (“Probe without motif”) abolishes retardation of probe migration by POU-IV (lane 4), demonstrating that the motif is necessary for formation of the protein-DNA complex. The mobility of the probe is further decreased in the presence of the anti-POU-IV antibody (purple arrowhead; lane 5), confirming that the protein bound to the probe is POU-IV.

Figure 7 – Source data 1: List of 12,972 genome-wide binding sites for POU-IV.

Figure 7 – Source data 2: An original gel image used to generate Figure 7C and the original image with relevant lanes labelled.

Downstream target genes of POU-IV are enriched with effector genes likely involved in neural function in the sea anemone.

We next sought to identify downstream target genes of POU-IV, based on the criteria that a target gene has at least one POU-IV ChIP peak within the gene locus which includes the promoter region – 350 bp upstream and 100 bp downstream of the transcription start site – and the gene body. Using this criterion, we found a total of 4188 candidate POU-IV downstream target genes (Supplementary File 1). We then examined which of these candidate POU-IV target genes were activated/repressed by POU-IV, using publicly available transcriptome data from *NvPOU4* mutant polyps and their siblings (Tourniere et al., 2020). Re-analysis of the transcriptome data identified 577 genes that were downregulated in *NvPOU4* mutants relative to their siblings (Supplementary File 2), and 657 genes that were upregulated in the mutants (Supplementary File 3) (adjusted p-value <0.01). Consistent with the previous report (Tourniere et al., 2020), Gene Ontology terms overrepresented in genes downregulated in *NvPOU4* mutants included those related to nervous system function such as “synaptic transmission” and “detection of stimulus” (Supplementary File 4). GO terms overrepresented in genes upregulated in mutants included “endoplasmic reticulum”, as identified by Tourniere et al (Tourniere et al., 2020), as well as a number of additional ones, such as “proteolysis” and “activation of signaling protein

activity involved in unfolded protein response” (Supplementary File 5). Out of the 577 genes downregulated in NvPOU4 mutants relative to their siblings, 293 were POU-IV target genes (Supplementary File 6), while out of the 657 genes upregulated in NvPOU4 mutants (Supplementary File 7), 178 were POU-IV target genes; we assume that the former represent genes that are directly activated by POU-IV, while the latter represent those directly repressed by POU-IV. Among the POU-IV-repressed genes is the *pou-iv* gene itself, indicating that POU-IV negatively regulates its own expression. Gene Ontology analysis found that 84 GO terms were overrepresented in the 293 genes directly activated by POU-IV, which include terms related to nervous system function such as “synaptic transmission” (p-adjusted<0.05) (Supplementary File 8). No GO terms were significantly overrepresented in the 178 genes directly repressed by POU-IV (p-adjusted<0.05).

POU-IV regulates the expression of the hair-cell-specific effector gene *polycystin 1* in the sea anemone.

To shed light on the mechanism by which POU-IV regulates hair cell maturation, we assessed which genes were directly activated by POU-IV in hair cells. Among the 577 genes significantly downregulated in NvPOU4 mutants relative to their siblings is a transmembrane receptor-encoding *polycystin-like* gene (JGI ID: 135278). By using *in situ* hybridization, we found that this gene was specifically expressed in tentacular epithelial cells whose cell bodies were located in the superficial stratum of the pseudostratified epithelium, resembling the hair cell (Figure 8A-F). We discovered by RTPCR that this gene and another *polycystin-like* gene (JGI ID: 218539) upstream – which was also one of the 577 genes significantly downregulated in NvPOU4 mutants relative to their siblings – together constitute a single *polycystin-like* gene. The transcript of the *polycystin-like* is 11,279 bases long, and encodes a protein that is 3457 amino acids long (Figure 8 - Figure supplement 1; GenBank accession number, OK338071). ChIP-seq data show that there are two POU-IV-binding motifs around the transcription start site of this locus (Figure 8G), suggesting that the *polycystin-like* gene is directly regulated by POU-IV.

We predicted the structure of the *polycystin-like* based on sequence similarity to known *polycystin 1* proteins. Transmembrane-spanning regions were predicted by using Phyre2 (Kelley et al., 2015) and based on the alignment with human and *Fugu* PKD1 sequences (GenBank accession numbers AAC37576 and AAB86683, respectively). Non-transmembrane-spanning

regions were predicted by using NCBI conserved domain search with default Blast search parameters. The *N. vectensis polycystin 1-like* protein was predicted to have a *Polycystin 1* (PKD1)-like domain structure, containing the extracellular PKD domain and REJ (receptor for egg jelly) module that are uniquely shared by Polycystin 1 proteins (Moy et al., 1996, Bycroft et al., 1999), the extracellular WSC (cell wall integrity and stress component) carbohydrate-binding domain, the intracellular PLAT (polycystin-1, lipoxigenase and alpha toxin) domain (Bateman and Sandford, 1999), the extracellular TOP (tetragonal opening for polycystins) domain (Grieben et al., 2017), and 11 transmembrane domains (Sandford et al., 1997) (Figure 8 - Figure supplement 1). However, unlike vertebrate Polycystin 1, Leucine rich repeat, C-type lectin, and LDL-like domains that reside in the N-terminal extracellular tail and a coiled-coil domain at the C-terminal intracellular tail were not identifiable. The last six transmembrane domains of Polycystin 1 are thought to be homologous to transient receptor potential (TRP) cation channels including Polycystin 2 (PKD2) (Mochizuki et al., 1996); in addition, the TOP domain is shared across Polycystin 1 and 2 (Grieben et al., 2017). We therefore generated an amino acid sequence alignment of the TOP domain and transmembrane domains of Polycystin 1 and Polycystin 2 proteins, and used it to carry out maximum likelihood phylogeny estimation. The results robustly placed the newly discovered *Nematostella vectensis polycystin-1-like* within the Polycystin 1 group (Figure 8 - Figure supplement 2). We therefore designate this gene as *Nematostella vectensis polycystin 1*.

To better resolve the cell type in which *polycystin 1* is expressed, we generated a reporter construct using 3704bp sequence encompassing the two POU-IV-binding motifs and upstream promoter region of the gene (scaffold 353:49,338-53,412). We injected this construct into wildtype zygotes and confirmed reporter gene expression specifically in hair cells at the primary polyp stage (Figure 8H-K). In addition, we have validated by *in situ* hybridization that *polycystin1* expression is lost in *pou-iv* ^{-/-} mutants (n=5) but not in their siblings (n=3) (Figure 8L-S). Taken together, these results suggest that *polycystin1* is directly activated by POU-IV in hair cells. To our knowledge, *polycystin 1* represents the first molecular marker specific to cnidarian hair cells.

Figure 8: POU-IV activates the expression of *polycystin 1* specifically in hair cells.

A-F: Confocal sections of primary polyps labeled with an antisense riboprobe against *polycystin1* transcript (“pkd1”) and antibodies against acetylated β -tubulin (“acTub”) and minicollagen 3 (“Ncol3”; (Zenkert et al., 2011)). Nuclei are labeled with DAPI (“dapi”). A-C are side views of the animal with the oral opening facing up. Expression of *polycystin1* occurs exclusively in the ectoderm of the oral tentacles (te). D-F are side views of a *polycystin1*-expressing epithelial cell (arrowhead) in the tentacular ectoderm (ec) with its apical surface facing up. Note that the cell body is localized apically and lacks minicollagen 3 expression. G: A schematic of the *polycystin1* locus, showing the distribution of POU-IV ChIP DNA (“POU-IV”) and input DNA from adult polyps. JGI gene models (“Genes”) and the revised gene model based on RTPCR (“Revised”) and the locations of the consensus POU-IV-binding motif - AT(A/T)ATT(A/T)AT - are numbered as I and II. X-axis shows the position along the genomic scaffold, and Y-axis

shows the number of reads. * shows an exon whose sequence is missing in the publicly available *N. vectensis* genome (v.1.0; (Putnam et al., 2007)). H-K: Confocal sections of an oral tentacle of a primary polyp injected with *polycystin1::Kaede* construct, labeled with an antibody against Kaede (“pkd1::kaede”). Filamentous actin is labeled with phalloidin (Pha). The apical surface of the tentacular ectodermal epithelium is to the top. Note that the Kaede-positive cell (arrowhead) has an apical cilium (ci) and stereovilli (st), a central nucleus (nu), and basal neurites (ne), exhibiting morphological hallmarks of a hair cell. No other cell types were found to be Kaede-positive. L-S: Confocal sections of a homozygous *pou-iv* mutant (“F2 *pou-iv* -/-“, P-S) and its sibling control (F2 *pou-iv* +/+ or *pou-iv* +/-, “F2 sibling ctrl”, L-O) at the primary polyp stage, labeled with an antisense riboprobe against *polycystin1* transcript (“pkd1”) and antibodies against acetylated β -tubulin (“acTub”) and minicollagen 3 (“Ncol3”; (Zenkert et al., 2011)). Panels show side views of the animal with the oral opening facing up. Animals lacking mature cnidocysts based on Ncol3 staining were assumed to be *pou-iv* -/- mutants; animals with mature cnidocysts were assumed to be *pou-iv* +/+ or *pou-iv* +/- . O and S are magnified views of tentacles boxed in M and Q, respectively. Note that *polycystin1* expression in tentacular ectoderm (arrowhead in M; O) is absent in the POU-IV null mutant (arrowhead in Q; S), demonstrating that POU-IV is necessary for *polycystin1* expression. Abbreviations: en endoderm; cn cnidocyst; nu nucleus; ms muscle fiber. Scale bar: 50 μ m (A-C, L-S); 10 μ m (D-F, H-K)

Figure 8 – Figure supplement 1: The predicted protein structure of the *Nematostella vectensis* polycystin 1-like.

The predicted protein structure of *Nematostella vectensis* polycystin 1-like. Extracellular domains are shown in blue, the intracellular domains in orange, and the transmembrane domains in purple. Transmembrane-spanning regions were predicted by using Phyre2 (Kelley et al., 2015) and based on alignment with human and *Fugu* PKD1 sequences. Non-transmembrane-spanning regions were predicted by using NCBI conserved domain search with default Blast search parameters. The PKD domain and the REJ module are specific to Polycystin 1 (PKD1) proteins. Abbreviations: WSC, cell wall integrity and stress response component; REJ, receptor for egg jelly; GPCR, G-protein coupled receptor; TM, transmembrane domain; PLAT, polycystin-1, lipooxygenase and alpha toxin; TOP, tetragonal opening for polycystins

Figure 8 – Figure supplement 1 – Source Data 1: Partial cDNA sequence of *Nematostella vectensis polycystin 1-like*, in which the start codon and stop codon are highlighted in green and red, respectively.

Figure 8 – Figure supplement 1 – Source Data 2: Tanslated amino acid sequence of *Nematostella vectensis polycystin 1-like*, in which conserved domains are highlighted in light blue for transmembrane-spanning regions and in purple for other domains.

Figure 8 – Figure supplement 2: *Nematostella vectensis polycystin 1-like* belongs to the Polycystin 1/PKD1 group.

Unrooted maximum likelihood phylogeny of Polycystin protein families based on the alignment of Polycystin 1 (PKD1) and Polycystin 2 (PKD2) protein sequences that span the conserved TOP

and transmembrane domains over 323 amino acid sites. NCBI accession numbers are shown with the name of each sequence. Bootstrap support values are shown at each node except when lower than 50%. The unit of the branch length is the number of substitutions per site. Note the strong support for the placement of *Nematostella vectensis polycystin 1-like* (*Nematostella vectensis* PKD1, highlighted in purple) within the PKD1 group, and for that of *N. vectensis* PKD2-like within the PKD2 group. The JGI ID for *N. vectensis* PKD2 (polycystin 2)-like is 160849.

Figure 8 – Figure supplement 2 – Source Data 1: Alignment of *polycystin 1* and *polycystin 2* amino acid sequences used to construct phylogeny shown in Figure 8 – Figure supplement 2.

POU-IV controls the maturation of hair cells and cnidocytes via distinct gene regulatory mechanisms.

Next, we have utilized publicly available single-cell transcriptome data from *N. vectensis* wildtype adults (Sebe-Pedros et al., 2018) to uncover additional candidate genes that are directly activated by POU-IV in hair cells. Both *polycystin-like* gene models (JGI IDs: 135278 and 218539) that are part of the newly discovered *polycystin 1* are uniquely represented in one of Sebe-Pedros et al.’s transcriptomically defined adult cell types (“c79”) referred to as the metacells (Sebe-Pedros et al., 2018). We have therefore deduced that the adult metacell c79 represents the hair cell, which enabled identification of additional POU-IV target genes activated in hair cells. Out of the 293 genes directly activated by POU-IV, we found a total of 32 genes that were represented in the adult metacell c79 (the presumptive hair cell) (Supplementary File 9). They include *potassium channel-like* (NVE ID: 12832; no JGI ID), *GABA receptor-like* (JGI gene ID: 98897), and *polycystin 2* (JGI gene ID: 160849; Figure 8 - Figure supplement 2), in addition to *polycystin 1* identified above. 3 of the 32 genes – *coagulation factor/neuropilin-like* (JGI gene ID: 202575), *CD59 glycoprotein-like* (NVE ID: 735; no JGI ID), and *polycystin 1* – are not found in any other metacell. No transcription factor-encoding genes were found to be activated by POU-IV in hair cells. Gene Ontology analysis found that 5 GO terms were overrepresented in POU-IV activated genes in hair cells (p-adjusted<0.05), including “potassium ion transmembrane transport” and “sensory perception of sound” (Supplementary File 10). In contrast, out of the 178 genes that are repressed by POU-IV, only two genes – *pou-iv* itself and

peptidylglycine alpha-amidating monooxygenase-like (JGI gene ID: 172604) – are represented in the adult metacell c79 (Supplementary File 9). These results are consistent with the hypothesis that POU-IV controls the maturation of hair cells by activating a cell-type-specific combination of effector genes that confer hair cell identity.

We have taken a similar approach to examine how POU-IV regulates the maturation of cnidocytes. Sebe-Pedro et al. categorized cnidocytes into eight transcriptomically distinct metacells (c1-c8), out of which c8 expresses *pou-iv* (Sebe-Pedros et al., 2018). Of the 293 genes activated by POU-IV, we found 3 genes in c8, which consisted of two *transmembrane propyl 4-hydroxylase-like* genes (JGI gene IDs: 239358 and 118978) and a *serine transporter-like* gene (JGI gene ID: 238501) (Supplementary File 11). *serine transporter-like* and one of the *hydroxylase-like* genes (JGI gene IDs: 239358) are represented specifically in cnidocyte metacells and not in others (Sebe-Pedros et al., 2018). As in hair cells, transcription factor-encoding genes were not found to be activated by POU-IV in cnidocytes. Of the 178 genes that are repressed by POU-IV, 13 genes were found in the cnidocyte metacell c8, which included genes encoding zinc finger and Sox transcription factors (Supplementary File 11) and showed significant enrichment of the GO term “transcription from RNA polymerase II promoter” (p-adjusted<0.05; Supplementary File 12). Importantly, we found no overlap in genes activated by POU-IV between hair cells and cnidocytes, which indicates that POU-IV controls the maturation of hair cells and cnidocytes by regulating the expression of distinct sets of genes. In addition, POU-IV may have a more significant role as a leaky repressor – to fine-tune gene expression levels – in cnidocytes than in hair cells, as the proportion of POU-IV-repressed genes relative to the total number of POU-IV targets represented in a given metacell was substantially higher in the cnidocyte (13/16; 81.25%) than in the hair cell (3/35; 8.6%). This pattern of gene regulation by POU-IV appears to be specific to cnidocytes. The proportion of POU-IV-repressed genes relative to the total number of POU-IV targets represented in non-cnidocyte, POU-IV-positive adult metacells – namely, c63, c64, c65, c66, c75, c76, and c102, in addition to c79 – was low, ranging from 2.2% to 13%, while that in POU-IV-positive, adult cnidocyte metacells - c100 that expresses spirocyte-specific *minicollagen* (JGI gene ID: 81941) and c101 that expresses nematocyte-specific *minicollagen1* (JGI gene ID: 211803) (Sebe-Pedros et al., 2018) – was 83.9% in c100 and 82.1% in c101 (Supplementary File 13), similar to the cnidocyte metacell c8. Taken together, these data suggest that POU-IV directs the maturation of cnidocytes not only by

activating a unique combination of effector genes, but also by negatively controlling the expression levels of a larger number of genes, including those encoding transcriptional regulators, in a leaky manner. Thus, the gene regulatory mechanisms by which POU-IV orchestrates the differentiation of hair cells and cnidocytes appear to be remarkably distinct.

Discussion:

In this paper, we identified the class IV POU homeodomain transcription factor (POU-IV) as an essential developmental regulator of cnidarian mechanosensory hair cells. This is the first discovery of a gene regulatory factor necessary for the development of classical mechanosensory neurons – that transmit mechanosensory information to other cells to elicit coordinated behavior – from a non-bilaterian, early-evolving animal group. Using the starlet sea anemone *Nematostella vectensis*, we have shown that POU-IV is postmitotically expressed in hair cells in the ectodermal epithelium of feeding tentacles during development. In addition, we have found that null mutation of *pou-iv* renders the animal unable to respond to tactile stimulation to its tentacles, and results in the loss of stereovillar rootlets, but not of neurites, in hair cells. Furthermore, we have presented evidence that POU-IV binds to deeply conserved DNA sequence motifs, and directly activates the expression of a unique combination of effector genes, but not transcription factor-encoding genes, specifically in hair cells. Among the POU-IV target effector genes we discovered the first cnidarian hair cell-specific molecular marker, *polycystin1*, which encodes a transmembrane receptor of the TRP channel superfamily. The results suggest that POU-IV plays a necessary role in regulating the maturation of mechanosensory hair cells in the sea anemone by directly activating the expression of cell-type-specific effector genes. Our findings strongly support POU-IV being the terminal selector of hair cell identity in the sea anemone.

Several lines of evidence indicate that POU-IV is specifically involved in the maturation, and not progenitor proliferation, initial differentiation, or survival, of hair cells in the tentacular ectoderm in *N. vectensis*. First, POU-IV is postmitotically expressed in hair cells in tentacular ectoderm, and thus is unlikely to have roles in proliferation or generation of their progenitor cells. Second, in POU-IV null mutants, we have found ciliary-cone-bearing epithelial cells that resemble hair cells in morphology and position within the epithelium; these cells are characterized by having an apical cilium surrounded by a circle of microvilli, a pear-shaped cell

body located in the superficial stratum of the pseudostratified epithelium, and basal neurites. None of the ciliary-cone-bearing epithelial cells express the pan-cnidocyte marker minicollagen 3, suggesting that the ciliary-cone-bearing cells in POU-IV null mutants do not represent partially differentiated nematocytes. The existence of differentiated hair cells in POU-IV null mutants implies that POU-IV is not involved in the initial differentiation of hair cells. However, the hair-cell-like cells of POU-IV null mutants failed to form a mature apical mechanosensory apparatus with stereovillar rootlets, indicating that POU-IV is essential for maturation of hair cells. Lastly, we have found no evidence that the epithelial cells expressing the non-functional form of POU-IV protein in POU-IV null mutants undergo programmed cell death in the tentacular ectoderm. Thus, POU-IV does not seem to be required for the survival of hair cells in the tentacles. Taken together, these data support the hypothesis that POU-IV regulates the maturation, but not progenitor proliferation, initial differentiation, or survival, of mechanosensory hair cells in the sea anemone.

The loss of stereovillar rootlets in hair cells in *pou-iv* mutants suggests that the POU-IV transcription factor regulates the expression of genes that are involved in stereovillar development. Given that stereovillar rootlets consist of actin filaments, actin-binding proteins may be regarded as potential regulators of stereovillar rootlet formation in hair cells. Among the identified POU-IV target genes expressed in hair cells is *polycystin 1*, which encodes a large transmembrane receptor with multiple extracellular and intracellular domains and TRP-channel-like transmembrane domains. Interestingly, its mouse homolog (PC-1) colocalizes with F-actin in inner ear hair cell stereovilli and is necessary for maintenance of stereovillar structure and normal hearing (Steigelman et al., 2011). In addition, PC-1 has been shown to regulate actin cytoskeleton reorganization in canine kidney epithelial cells (Boca et al., 2007). If the function of Polycystin 1 in modulating the organization of actin cytoskeleton is broadly conserved, *N. vectensis* Polycystin 1 might control the structural integrity of stereovilli in hair cells through its interaction with F-actin. POU-IV may therefore direct stereovillar development in cnidarian hair cells by activating *polycystin 1*. Functional analysis of *N. vectensis polycystin 1* to evaluate its role in stereovillar development is warranted.

We have proposed that the lack of tentacular response to tactile stimuli in *pou-iv* mutants is due to the loss of structural rigidity in the apical mechanosensory apparatus – stereovilli, in particular – of hair cells, resulting from the failure of hair cells to form stereovillar rootlets. We

note, however, that it could additionally be because of the functional defects in mechanotransduction channels. POU-IV is known to directly activate the expression of the mechanotransduction channel-encoding gene, *mec-4*, that is necessary for touch-cell function in *C. elegans* (Duggan et al., 1998). The Polycystin 1 protein discussed above contains transmembrane domains that are homologous to the transient receptor potential (TRP) calcium channel. If this channel is involved in mechanotransduction, the loss of *polycystin 1* expression in *pou-iv* mutants would directly lead to loss of mechanotransduction channel function. This hypothesis may be evaluated by specifically examining the role of the channel-encoding region of *N. vectensis polycystin 1* in mechanotransduction.

Alternatively, the loss of *polycystin 1* expression may indirectly perturb channel function. In epithelial cells of vertebrate kidneys, Polycystin 1 interacts with the calcium ion channel Polycystin 2 to form a complex that functions as a fluid flow sensor with Polycystin 1 acting as a mechanosensitive cell surface receptor and Polycystin 2 as an ion-permeant pore (reviewed in (Delmas, 2004)). The *Nematostella vectensis* genome encodes *polycystin 2* (Figure 8 - Figure supplement 2), which is co-expressed with *polycystin 1* in the adult metacell c79 (i.e. the hair cell) (Supplementary File 9; (Sebe-Pedros et al., 2018)). If these two proteins form a mechanically gated ion channel complex in hair cells as in vertebrate kidney epithelial cells, the loss of expression of *polycystin 1* would perturb the ability of the complex to sense mechanical stimuli, resulting in defects in channel function. To explore this hypothesis, the important next step will be to assess whether Polycystin 1 and 2 form a complex in *N. vectensis*.

We note that, although our findings are consistent with the hypothesis that cnidarian hair cells function as mechanosensors, we do not rule out the possibility that cnidarian hair cells might be multimodal sensory cells; they might have additional functions as chemoreceptors and/or photoreceptors. Indeed, hair cells of the sea anemone *Haliplanella luciae* have been reported to respond to N-acetylated sugars by elongating their stereovilli (Mirethibodeaux and Watson, 1994, Watson and Roberts, 1995), indicative of chemosensory function. The metacell c79 (the presumptive hair cell) of adult *N. vectensis* expresses several G-protein-coupled receptor (GPCR)-encoding genes (Supplementary File 14; (Sebe-Pedros et al., 2018), some of which might encode chemosensory receptors; none were found to encode opsins. Functional analyses of these GPCRs may shed light on the molecular basis of sensory multimodality in cnidarian hair cells.

Our results indicate that the role for POU-IV in mechanoreceptor development is broadly conserved across Cnidaria and Bilateria. As mentioned above, one of the vertebrate *pou-iv* paralogs (*Brn3c*) is necessary for maturation and survival of inner ear hair cells in mice (Xiang et al., 1998, Xiang et al., 1997a, Erkman et al., 1996). Likewise, in *C. elegans*, differentiation of mechanosensory touch cells requires a *pou-iv* ortholog *unc-86* (Chalfie and Sulston, 1981, Chalfie and Au, 1989, Finney and Ruvkun, 1990, Duggan et al., 1998, Chalfie et al., 1981). In Cnidaria, *pou-iv* is expressed in mechanosensory organs in scyphozoan and hydrozoan jellyfish (Nakanishi et al., 2010, Hroudova et al., 2012), and is necessary for differentiation of the lineage-specific mechanosensory-effector cell, the cnidocyte, in *N. vectensis* (Tournier et al., 2020). In this report, we have demonstrated that *pou-iv* has an essential role in the maturation of the classical mechanosensory neuron of Cnidaria – the hair cell – using *N. vectensis*. These comparative data show that POU-IV-dependent regulation of mechanosensory cell differentiation is pervasive across Cnidaria and Bilateria, and likely predates their divergence. How early the role of POU-IV in mechanoreceptor differentiation emerged in animal evolution remains unresolved, and requires comparative data from placozoans and sponges, which are wanting.

We note, however, that POU-IV has a broad role in the differentiation of multiple neural cell types across Cnidaria and Bilateria. In *N. vectensis*, POU-IV expression is not restricted to mechanosensory hair cells and cnidocytes, but also found in RFamidergic neurons and *NvElav1*-positive endodermal neurons (Tournier et al., 2020). Likewise in Bilateria, POU-IV regulates the differentiation of a variety of neural cell types beyond mechanosensory cells, including chemosensory neurons in insects (Clyne et al., 1999) and photosensory neurons in vertebrates (retinal ganglion cells; e.g. Erkman et al., 1996, Gan et al., 1996)). Therefore, it seems plausible that POU-IV was ancestrally involved in the differentiation of multiple neural cell types in addition to mechanosensory cells.

Interestingly, POU-IV is required for normal development of stereovilli in hair cells in both sea anemones (this study) and mice (Xiang et al., 1998), raising the possibility that POU-IV controlled the formation of the apical sensory apparatus of mechanosensory cells in the last common ancestor of Cnidaria and Bilateria, potentially via regulation of *polycystin 1*. Alternatively, the essential role for POU-IV in stereovillar formation in mechanosensory cells could have evolved independently in Cnidaria and vertebrates. Comparative studies of the

mechanism of stereovillar formation across sea anemones and vertebrates, along with mechanistic studies of POU-IV gene function in phylogenetically informative taxa, such as medusozoan cnidarians and acoel bilaterians, are needed to evaluate these alternative hypotheses.

Regulatory factors acting upstream of POU-IV in cnidarian hair cell development remain unknown. In Bilateria, members of the Atonal basic-loop-helix-loop-helix (bHLH) transcription factor family appear to have a conserved role in positive regulation of POU-IV expression (reviewed in (Leyva-Diaz et al., 2020)). For instance, vertebrate *atonal* genes act upstream of *pou-iv* genes to drive the differentiation of inner ear hair cells (Ikeda et al., 2015, Masuda et al., 2011, Yu et al., 2021) and retinal ganglion cells (Liu et al., 2001). Similarly, *C. elegans atonal* ortholog *lin-32* controls the expression of the *pou-iv* ortholog *unc-86* in touch sensory neurons (Baumeister et al., 1996). Although cnidarians lack unambiguous *atonal* orthologs, they have divergent bHLH genes that belong to the Atonal superfamily, which consists of Atonal and related bHLH gene families including Neurogenin, and NeuroD (Gyoja et al., 2012, Simionato et al., 2007). Whether these *atonal-like* bHLH factors function upstream of POU-IV in the context of cnidarian hair cell development needs to be assessed, as it may provide insights into the evolution of gene regulatory mechanisms underpinning mechanoreceptor development across Cnidaria and Bilateria.

In light of new comparative data reported herein emerges a model of mechanosensory cell differentiation in the last common ancestor of Cnidaria and Bilateria. We assume that the embryo of the Cnidaria-Bilateria ancestor had neurogenic ectoderm (Nakanishi et al., 2012). During late embryogenesis or postembryonic development of this ancestor, mechanosensory cell progenitors differentiated into postmitotic sensory cells in the ectoderm, extending apical cilia and basal neurites. These postmitotic sensory cells expressed the terminal selector POU-IV, which translocated to the cell nuclei, and bound to the DNA recognition motif GCAT(A/T)ATT(A/T)AT (i.e. the consensus motif across Bilateria and Cnidaria) associated with target genes in these cells. This activated the expression of effector genes, possibly including *polycystin 1*, whose protein products generated mechanoreceptor-specific structures necessary for mechanosensory function, such as the apical mechanosensory apparatus consisting of a cilium surrounded by a ring of stereovilli. The mature identity of the mechanosensory cell was thereby established.

Following the divergence of Cnidaria and Bilateria, POU-IV may have been recruited for the evolution of the lineage-specific mechanosensory effector – the cnidocyte – in Cnidaria, as *pou-iv* is essential for cnidocyte development in *N. vectensis* (Tourniere et al., 2020). How POU-IV would have come to direct cnidocyte development remains unclear. Given that both hair cells and cnidocytes are mechanosensory cell types, it seems reasonable to expect that an ancestral POU-IV gene regulatory network that defined mechanosensory cell identity was repurposed for the emergence of cnidocytes, and should be shared across these two cell types. However, we found no evidence in support of this hypothesis; instead, our results suggest that POU-IV controls distinct sets of genes in each cell type. One possible evolutionary scenario to account for this observation is that POU-IV initially activated the same battery of effector genes in the ancestral cnidocytes and hair cells, but POU-IV target genes diverged substantially during cnidarian evolution so that they no longer overlap between the two cell types. Another possibility is that POU-IV regulated a unique set of genes in the ancestral cnidocytes when POU-IV became part of the cnidocyte gene regulatory network. This possibility seems conceivable if POU-IV expression was activated in epigenetically distinct cell lineages, so that between the cnidocyte lineage and the hair cell lineage 1) POU-IV cooperated with different co-factors, and/or 2) the accessibility of POU-IV target genes differed, which would result in differential expression of POU-IV target genes. Evidence from bilaterian models such as *C. elegans* and mice indicates that POU-IV cooperates with a range of different co-factors to define distinct neural identities (reviewed in (Leyva-Diaz et al., 2020), suggesting an important role for POU-IV cofactors in the diversification of neural cell types. Whether evolution of POU-IV cofactors played a role in the evolution of cnidocytes remains to be tested. Investigation into the mechanism by which POU-IV activates distinct sets of genes across cnidocytes and hair cells will be the critical next step for shedding light on how POU-IV may have contributed to the evolution of the novel mechanosensory cell type of Cnidaria.

Materials and Methods:

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (<i>Nematostella vectensis</i>)	F1 <i>pou-iv</i> +/-	This paper	Nv F1 <i>pou-iv</i> +/-, this paper	Nakanishi lab, University of Arkansas
Antibody	POU-IV/Brn-3; rabbit; polyclonal	This paper	RRID:AB_2895562	(1:200) dilution used; Nakanishi lab, University of Arkansas

1164

1165 **Animal culture**

1166 *Nematostella vectensis* were cultured as previously described (Fritzenwanker and Technau, 2002,
1167 Hand and Uhlinger, 1992).

1168 **RNA extraction, cDNA synthesis and gene cloning**

1169 Total RNA was extracted from a mixture of planulae and primary polyps using TRIzol (Thermo
1170 Fisher Scientific). cDNAs were synthesized using the SMARTer RACE cDNA Amplification
1171 Kit (Cat. No. 634858; Takara, Mountain View, CA, USA). *In silico* predicted *pou-iv* gene
1172 sequence was retrieved from the Joint Genome Institute genome database (*Nematostella*
1173 *vectensis* v1.0, protein ID 160868; <http://genome.jgi-psf.org/Nemve1/Nemve1.home.html>). 5'
1174 and 3' RACE PCR experiments were conducted, following manufacturer's recommendations, in
1175 order to confirm gene prediction. Gene specific primer sequences used for RACE PCR are: 3'
1176 RACE Forward 5'-CGATGTCGGGTCCGCGCTTGCACATTG-3'; 5' RACE Forward 5'-
1177 GCCGCGCCGATAGACGTGCGTTTACG-3'. RACE PCR fragments were cloned into a
1178 pCR4-TOPO TA vector using the TOPO TA Cloning kit (Cat. No. K457501; ThermoFisher
1179 Scientific), and sequenced at Genewiz, New Jersey.

1180 The *polycystin 1* cDNA sequence (GenBank accession number: OK338071) was obtained
1181 by subcloning small overlapping gene fragments (1.5-4kb). Gene fragments were generated by
1182 RTPCR using RACE-ready cDNAs as templates. Gene specific primer sequences used to
1183 amplify *polycystin 1* cDNA are listed in Supplementary File 15. The PCR products were then

1184 cloned into a pCR4-TOPO TA vector using the TOPO TA Cloning kit (Cat. No. K457501;
1185 ThermoFisher Scientific), and sequenced at Eurofins Genomics, Kentucky.

1186 **Generation of an antibody against *N. vectensis* POU-IV**

1187 An antibody against a synthetic peptide CQPTVSESQFDKPFETPSPINamide corresponding in
1188 amino acid sequence to N-terminal QPTVSESQFDKPFETPSPIN of *N. vectensis* POU-IV
1189 (Figure 4A) was generated in rabbit (YenZym Antibodies, LLC). TBLASTN search of the
1190 antigen sequence against the *N. vectensis* genome
1191 (http://metazoa.ensembl.org/Nematostella_vectensis/Info/Index) yielded a single hit at the *pou-iv*
1192 locus ([NEMVEscaffold_16:1069268-1069327](#)); there were no significant matches to other loci.
1193 Following immunization, the resultant antiserum was affinity purified with the
1194 CQPTVSESQFDKPFETPSPINamide peptide.

1195 **CRISPR-Cas9-mediated mutagenesis**

1196 20 nt-long sgRNA target sites were manually identified in the *N. vectensis pou-iv* genomic locus.
1197 To minimize off-target effects, target sites that had 17 bp-or-higher sequence identity elsewhere
1198 in the genome (*Nematostella vectensis* v1.0;
1199 <http://genome.jgi.doe.gov/Nemvel/Nemvel.home.html>) were excluded. Selected target
1200 sequences were as follows.

1201 5'- CTACGATGCGCACGATATTT-3' (Cr1)

1202 5'- ACGAGAGCTGGAATGGTTCG-3' (Cr2)

1203 5'- TAAACGCACGTCTATCGGCG-3' (Cr3)

1204 5'- AATAATGGACATCTACGCCG-3' (Cr4)

1205 The sgRNA species were synthesized *in vitro* (Synthego), and mixed at equal
1206 concentrations. The sgRNA mix and Cas9 endonuclease (PNA Bio, PC15111, Thousand Oaks,
1207 CA, USA) were co-injected into fertilized eggs at concentrations of 500 ng/μl and 1000 ng/μl,
1208 respectively.

1209 **Genotyping of embryos and polyps**

1210 Genomic DNA from single embryos or from tentacles of single polyps was extracted by using a
1211 published protocol (Ikmi et al., 2014), and the targeted genomic locus was amplified by nested
1212 PCR. Primer sequences used for nested genomic PCR are: "1" Forward 5'-
1213 CGAATTCCTCTGCAATAATCACTGATCG-3', "1" Reverse 5'-
1214 CTCGTTGGCAGGTGCGGAAAGAG-3', "2" Forward 5'-

1215 CGTTCGACTTCATTTCCGCTCGTC-3', "2" Reverse 5'-
1216 CGGAAGTTAACGTCGTTAATGCGAAGG-3'. To determine the sequence of mutant alleles,
1217 PCR products from genomic DNA extracted from F1 mutant polyps were gel-purified, cloned
1218 and sequenced by using a standard procedure. Using the sequence information of the *pou-iv*-
1219 mutant allele, genotyping primers for F2 animals were designed as follows (Figure 4B).
1220 Forward 5'- CGTTCGACTTCATTTCCGCTCGTC-3'
1221 Reverse (1), 5'- GCCGCGCCGATAGACGTGCGTTTACG-3' (*pou-iv*+ -specific; expected size
1222 of PCR product, 689 bp)
1223 Reverse (2), 5'- CGGAAGTTAACGTCGTTAATGCGAAGG-3' (expected size of PCR
1224 product: *pou-iv*+, 1312 bp; *pou-iv*-, 558 bp)

1225 **Behavioral analysis**

1226 Animals were selected for behavior analyses if they were 10-16dpf, unfed, had reached the
1227 primary polyp stage, and had two or more tentacles present. Animals were only tested if their
1228 tentacles protruded from their bodies at time of testing initiation. All behavior experiments were
1229 performed with the experimenter blind to the animal's genotype until after testing was completed.
1230 Animals were allowed to rest for at least two hours between tests. Behavioral analyses were
1231 performed under a Zeiss Stemi 508 microscope with Nikon DSL-4 camera attachment.

1232 To examine response to touch, a hair attached to a microdissection needle holder (Roboz
1233 Surgical Instrument Co., Gaithersburg, MD, USA) was pressed briefly on the distal end of each
1234 tentacle. The stimulus was applied once more to remaining unretracted tentacles, to ensure that a
1235 tentacle was not missed during the first stimulus. The number of primary polyps that retracted
1236 one or more tentacles upon touch was counted. If any other part of the body was touched
1237 accidentally during tentacle stimulation, data for that animal was discarded and the trial was
1238 repeated two hours or more after the previous test.

1239 Chemosensory response of primary polyps to *Artemia* chemical cues was analyzed as
1240 follows. *Artemia* shrimp extract was made from 1 day old *Artemia* brine shrimp, ground with a
1241 micropestle (USA Scientific) in 1/3 artificial sea water (Instant Ocean), at a concentration of
1242 approximately 1 shrimp per 1 microliter. 2 microliters of shrimp extract was applied with an
1243 Eppendorf pipette to the head and tentacle area of each sea anemone. The animal was observed
1244 for 1 minute to examine the occurrence of tentacular retraction.

1245 **CM-DiI labeling**

The lipophilic tracer CM-DiI (Molecular Probes, C7000) was used to label the cell membrane of a subset of mature hair cells of the polyp tentacles. Primary polyps were incubated in 1/3 seawater with 10 μ M CM-DiI for 1 hour at room temperature. The labelled polyps were rinsed in fresh 1/3 seawater, and were anesthetized in 2.43% MgCl_2 for 20 minutes. They were then fixed in 4% formaldehyde for 1 hour at room temperature. Specimens were washed in PBSTr (1xPBS + 0.5% Triton-X100) for 1 hour to permeabilize the tissue, before labeling filamentous actin and nuclei with AlexaFluor 488-conjugated phalloidin (1:25, Molecular Probes A12379) and the fluorescent dye 4',6-diamidino-2-phenylindole (DAPI; 1:1,000, Molecular Probes), respectively.

EdU pulse labeling

Primary polyps were incubated in 1/3 seawater containing 200 μ M of the thymidine analogue, EdU (Click-iT EdU AlexaFluor 488 cell proliferation kit, C10337, Molecular Probes), for 20 minutes to label S-phase nuclei. Following washes in fresh 1/3 seawater, the polyps were immediately fixed as described previously (Martindale et al., 2004, Nakanishi et al., 2012), and immunohistochemistry was then carried out as described below. Following the immunohistochemistry procedure, fluorescent labelling of incorporated EdU was conducted according to the manufacture's recommendations prior to DAPI labelling.

Western blotting

3–4-week-old polyps were lysed in AT buffer (20 mM Hepes pH7.6, 16.8 mM $\text{Na}_4\text{P}_2\text{O}_7$, 10 mM NaF, 1 mM Na_3VO_4 , 0.5 mM DTT, 0.5 mM EDTA, 0.5mM EGTA, 20% glycerol, 1% Triton X-100, and protease inhibitor cocktail (Sigma)) on ice with a plastic pestle in a microcentrifuge tube until there were no large fragments. The mixture was then sonicated with a Branson Digital Sonifier 3 times with the setting of 0.5 s on 1 s off for 10 s at an amplitude of 10%. NaCl was added to the lysate to a final concentration of 150 mM. The samples were centrifuged at 21,000 g for 20 min at 4 °C. The supernatant was transferred to a new centrifuge tube and the pellet was discarded. Protein concentration of the supernatant was determined by Bradford Reagent (Sigma). The proteins were then separated on a 12% SDS-PAGE (40 μ g protein/lane), transferred to a PVDF membrane (Amersham Hybond; 0.2 μ m). After blocking with the Odyssey Blocking Buffer (TBS) for 30 min at room temperature, the membrane was incubated with an anti-POU-IV polyclonal antibody (rabbit, 1:1000) at 4 °C overnight. The membranes were then washed extensively with TBST and incubated with 1:10,000 IRDye 800CW donkey anti-rabbit IgG at

room temperature for 1 h. After washing, protein bands were visualized on a LI-COR (9120) Imaging System. Anti-tubulin (T6793 Sigma) was used as a loading control.

Chromatin immunoprecipitation and Sequencing (ChIP-seq)

Adult animals (~1.2 g wet weight) were harvested and washed with PBS twice. The animals were treated with 2 % formaldehyde in PBS for 12 min at room temperature, and the cross-linking reagent was quenched with 0.125 M glycine for 5 min at room temperature. After washing with PBS twice, the crosslinked samples were resuspended in 10 ml buffer1 (50 mM Hepes, pH 7.5, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% NP-40, 0.25% Triton X-100, 1 mM DTT, and protease inhibitors (Sigma)) and lysed with 10 slow strokes of a tight-fitting pestle (type B) in a 15 ml *Dounce* homogenizer. The lysate was centrifuged at 500 g for 5 min at 4 °C, and the resulting pellet was resuspended in 10 ml Buffer1 and homogenized as described above. The homogenization processes were repeated 1-2 more times. In the last homogenization, the lysate was centrifuged at 2000 g for 10 min at 4 °C, and the pellet (nuclei) was resuspended in 4 ml SDS lysis buffer (50 mM Tris-HCl, pH 8.0, 10 mM EDTA, 1% SDS, and protease inhibitors). The chromatin was sheared to 200-500 bp fragments by sonicating the samples 12 times (1" on and 1" off for 1 min) at an amplitude of 50% with a Branson Digital Sonifier. The sonicated samples were centrifuged at 21,000 g for 10 min at 4 °C and then diluted 10X with CHIP dilution buffer (17.7 mM Tris-HCl, pH 8.0, 167 mM NaCl, 1.2 mM EDTA, 1.1% Triton X-100, 0.01% SDS, and protease inhibitors). After the lysate was cleared with Protein A and G magnetic beads (Cell Signaling), 50 µl of the cleared sample was set aside as input DNA, and 5 ml of lysate was incubated with 10 µg anti-Brn3 Rabbit polyclonal antibody, which was conjugated to 30 µl of protein A+G magnetic beads. After incubation at 4 °C overnight, the beads were washed 3 times with 1 ml of Low salt buffer (20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), 3 times with 1 ml of High salt buffer (20mM Tris-HCl, pH 8.0, 500 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.01% SDS), 3 times with 1 ml of LiCl wash buffer (10 mM Tris-HCl, pH 8.0, 0.25 M LiCl, 0.5% NP-40, 0.5% sodium dexycolate, and 1 mM EDTA), and 3 times with 1 ml of TE buffer (10 mM Tris-HCl, pH 8.0, and 1 mM EDTA). The chromatin was eluted in SDS Elution buffer (50 mM Tris-HCl, pH8.0, 1% SDS, and 1mM EDTA), followed by reverse cross-linking at 65 °C overnight. After being treated with RNase A (1 mg/ml) at 37 °C for 1 h and then with protease K (0.2 mg/ml) at 45 °C for 1 h, the DNA fragments were purified with QIAquick Spin columns (QIAGEN) and the

purified DNA samples were quantified by Qubit4 (ThermoFisher). 20 ng of the immunoprecipitated DNA or input DNA was used to generate a library with the NEBNext Ultra II DNA Library kit following the manufacturer's protocol. Libraries were initially quantified by Qubit4 and the size profiles were determined by TapeStation (Agilent) and then quantified by qPCR (NEBNext Library Quant Kit) for high-throughput sequencing. Four biological replicates of libraries of immunoprecipitated DNA and the input DNA were pooled in equimolar ratio and the pooled libraries were sequenced on a DNBseq Sequencing platform (BGI, China) for PE 100 bp.

Expression and purification of POU-IV protein

cDNA encoding POU-IV was inserted into a modified PET28a plasmid in which POU-IV was expressed under a 2X Flag tag and a tobacco etch virus (TEV) protease cleavage site by PCR using forward primer 5'-GATGACAAGGGAGGTGGATCCATGAACCGGGACGGATTGCTTAAC-3' and reverse primer 5'-GGTGGTGGTGGTGGCTCGAGTCAATGTACGGAGAACTTCATTCTC-3'. The construct was transformed into BL21 (DE3) cells (C2530, NEB). After the transformed cells were grown in LB medium to 0.6 at OD600, the expression of the protein was induced by 1mM of Isopropyl β -D-1-thiogalactopyranoside (IPTG) at 30 °C for 5 h. The cells were lysed by sonication in Lysis buffer (20 mM Tris pH7.5, 150 mM NaCl, 1% TritonX-100, 10% glycerol, 1 mM EDTA and protease inhibitor (P8340, Sigma)). The lysate was cleared by centrifugation at 30,000 g for 30 min at 4 °C, and the supernatant was incubated with anti-Flag M2 Affinity Gel (A2220, Sigma) overnight at 4 °C. After washing with Wash Buffer (20 mM Tris pH7.5, 150 mM NaCl, 0.5% TritonX-100, and 1 mM EDTA), the bound proteins were eluted with Elution buffer [50 mM Tris pH 7.5, 30 mM NaCl, and 0.25 mg/ml 3X Flag peptide (F4799, Sigma)]. The buffer for the eluted protein was changed to 20 mM Tris pH7.5, and 100 mM NaCl using an Amicon Ultra Centricon with 10 kDa cut-off. The purified protein was stored at -80 °C for further use.

Electrophoretic mobility shift assay (EMSA)

The biotin-labeled DNA oligonucleotides with or without motif were purchased from Integrated DNA Technologies. For the experiment shown in Figure 8C, the probe sequence with motif was 5'-AAACAAAGATTCTAAGCATCC***ATTATTA***ATATACATCCCTAGAAAAAATC-3' (motif in bold and italics; scaffold 353:52091-52140,

<https://mycocosm.jgi.doe.gov/Nemve1/Nemve1.home.html>), and that without motif was 5'-

ATCGAAAACAAAGATTCTAAGCATCATACATCCCTAGAAAAAATCTCCGC-3'.

The two complementary strands were annealed together by mixing equivalent molar amounts of each oligonucleotide, heating at 95 °C for 5 min, and slow cooling on bench to room temperature.

Gel mobility shift assay was carried out using Gelshift Chemiluminescent EMSA kit (#37341, Active Motif) with modifications. Briefly, 0.25 µg POU-IV protein, 20 fmol biotin-labeled probes with or without motif were incubated in Binding Buffer (10 mM Tris pH7.4, 50 mM KCl, 2 mM MgCl₂, 1 mM EDTA, 1 mM DTT, 5% glycerol, 4 µg/ml BSA, and 0.125ug/µl salmon sperm DNA) in a total volume of 20 µl at RT for 30 min. For the competition, unlabeled probe was added to the reaction mixture at 300-fold molar excess of the biotin labeled probe. For supershift assay, 2 µg POU-IV antibody was incubated with POU-IV protein for 1 h at room temperature before the biotin-labeled probe was added. The DNA-protein complexes were separated with a 5% nondenaturing polyacrylamide gel in 0.5X TBE buffer at 100 V for 1 h. The probes were then transferred to a positively charged Nylon Membrane (Nytran SPC, Cytiva) in 0.5X TBE buffer at 380 mA for 30 min at 4 °C. After cross-linking the transferred probes to the membrane by CL-1000 Ultraviolet Crosslinker (UVP) at 120 mJ/cm² for 1 min, the membrane was incubated with HRP-conjugated streptavidin, and the chemiluminescence of the biotin-labeled probes were detected with ECL HRP substrate in X-ray film.

Immunofluorescence, *in situ* hybridization, and TUNEL

Animal fixation and immunohistochemistry were performed as previously described (Martindale et al., 2004, Nakanishi et al., 2012). For immunohistochemistry, we used primary antibodies against POU-IV (rabbit, 1:200), minicollagen 3 (guinea pig, 1:200; (Zenkert et al., 2011)), Kaede (rabbit; 1:500; Medical & Biological Laboratories, PM012M), and tyrosinated β -tubulin (mouse, 1:500, Sigma T9028), and secondary antibodies conjugated to AlexaFluor 568 (1:200, Molecular Probes A-11031 (anti-mouse) or A-11036 (anti-rabbit)) or AlexaFluor 647 (1:200, Molecular Probes A-21236 (anti-mouse) or A-21245 (anti-rabbit)). Nuclei were labeled using the fluorescent dye DAPI (1:1,000, Molecular Probes D1306), and filamentous actin was labeled using AlexaFluor 488-conjugated phalloidin (1:25, Molecular Probes A12379). For *in situ* hybridization, antisense digoxigenin-labeled riboprobes against *Nematostella vectensis pou-iv* and *polycystin 1* were synthesized from 5' and 3' RACE products, respectively (MEGAscript transcription kit; Ambion), and were used at a final concentration of 1 ng/µl. TUNEL assay was

carried out after immunostaining, by using In Situ Cell Death Detection Kit (TMR red cat no. 1684795, Roche, Indianapolis, IN, USA) according to manufacturer's recommendation. Specimens were mounted in ProlongGold antifade reagent (Molecular Probes, P36930). Fluorescent images were recorded using a Leica SP5 Confocal Microscope or a Zeiss LSM900. Images were viewed using ImageJ.

Transmission Electron Microscopy

1-4 week old primary polyps were anesthetized in 2.43% MgCl₂ for 20 minutes, and then fixed in 2.5% glutaraldehyde and 0.1 M cacodylate buffer at 4°C overnight. Fixed polyps were washed four times in 0.1 M cacodylate buffer for 10 min, and post-fixed for one hour in 0.1 M cacodylate buffer and 1% OsO₄. Specimens were rinsed with five 5 min washes of 0.1 M cacodylate buffer, followed by dehydration in a graded ethanol series consisting of 15 min washes in 30%, 50%, 70%, 80%, and 95% ethanol, followed by two 15 min washes in 100% ethanol. Dehydrated polyps were placed in a 1:1 solution of ethanol/Spurr's resin and left in a desiccator for 1 hr. The ethanol/resin mixture was replaced with a 100% resin solution, and polyps were left in a desiccator overnight. Samples were then transferred to flat-embedding molds filled with 100% Spurr's resin and placed in an oven at 70°C for 14 hours.

Blocks containing embedded polyps were trimmed with a razor blade and cut into ultra-thin sections using a diamond knife on a Sorval Porter-Blum ultramicrotome. Sections were transferred to carbon/formvar coated copper grids, which were then stained with 2% uranyl acetate and lead citrate and viewed on a JEOL JEM-1011 transmission electron microscope at 100 kV.

Generation of *kaede* transgenic animals

The *pou-iv::kaede* and *pkd1::kaede* transgenic animals were produced by I-SceI-mediated transgenesis as described previously (Renfer et al., 2010) with modifications. To generate *pou-iv::kaede* plasmid, 3199 bp genomic sequence upstream of the start codon of the *Nematostella vectensis pou-iv* (scaffold 16: 1065408-1068606; <https://mycocosm.jgi.doe.gov/Nemve1/Nemve1.home.html>) was cloned in front of the open reading frame of the Kaede gene (Ando et al., 2002) by FastCloning (Li et al., 2011). To generate *pkd1::kaede* plasmid, 3704 bp genomic sequence upstream of the 5th base in exon 3 of the *Nematostella vectensis polycystin 1* (scaffold 353: 49524-53227; <https://mycocosm.jgi.doe.gov/Nemve1/Nemve1.home.html>) was cloned in front of the open

reading frame of the Kaede gene. The plasmid was digested with I-SceI for 15-30 minutes at 37 °C and injected into zygotes at 50 ng/ µl. The injected animals were raised to primary polyps, and Kaede was visualized by using an anti-Kaede antibody.

Phylogenetic analysis

Sequence alignment and phylogenetic analyses were performed on the Geneious Prime platform (v.2019.2.). *polycystin 1* and *polycystin 2* sequences were retrieved from GenBank at the NCBI website (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>), either directly or via the protein BLAST search using the *N. vectensis* sequences as queries. Peptide sequences were aligned with MUSCLE (v3.7) [57] configured for highest accuracy (MUSCLE with default settings). After alignment, ambiguous regions (i.e. containing gaps and/or poorly aligned) were manually removed. The final alignment spanned the conserved TOP and transmembrane domains over 323 amino acid sites (Figure 8 – Figure supplement 2 – Source Data 1). Phylogenetic trees were reconstructed using the maximum likelihood method implemented in the PhyML program [58]. The WAG substitution model [59] was selected assuming an estimated proportion of invariant sites and 4 gamma-distributed rate categories to account for rate heterogeneity across sites. The gamma shape parameter was estimated directly from the data. Reliability for internal branches of maximum likelihood trees was assessed using the bootstrapping method (100 bootstrap replicates).

RNA-seq data analysis

The accession number from the RNA-seq data used in this study is E-MTAB-8658. The raw fastq files were filtered for low quality reads using Trimmomatic v0.39 (SLIDINGWINDOW:4:15, MINLEN:60, HEADCROP:10) (Bolger et al., 2014). Filtered reads were aligned to the *Nemastostella vectensis* genome (ENA accession: GCA_000209225) using STAR v 2.7.5a (sjdbOverhang: 99) (Dobin et al., 2013). The alignment files were processed using Samtools v1.9 (Danecek et al., 2021) and reads on genes were counted using HTSeq v0.12.4 (-t gene) (Anders et al., 2015). Genome annotation reported by Fredman, et al. (Fredman et al., 2013) was used. The differential expression analysis and normalization were performed in R, using the DESeq2 (Love et al., 2014) and Apeglm (Zhu et al., 2019) packages.

ChIP-seq data analysis

ChIP-seq data are available at the BioProject database (accession number: PRJNA767103). The raw data was trimmed using Trimmomatic v0.39 (Bolger et al., 2014). Reads were aligned to the

Nemastostella vectensis genome (ENA accession: GCA_000209225) using STAR v 2.7.5a (Dobin et al., 2013) and alignment files were processed using Samtools v1.9 (Danecek et al., 2021). Peak calling was performed in the aligned readings with MACS2 (Zhang et al., 2008). The quality of the peaks in the replicates (n=3) was checked using phantompeakqualtools (Landt et al., 2012). To improve sensitivity and specificity of peak calling, and identify consensus regions of the multiple replicates, we used Multiple Sample Peak Calling (MSPC: -r tec -w 1e-4 -s 1e-8) (Jalili et al., 2015). A *de novo* motif search and motif enrichment was performed using RSAT local-word-analysis (Thomas-Chollier et al., 2012). The motif comparison tool TomTom (Gupta et al., 2007) was used to search enriched motifs against the Jaspar database (Fornes et al., 2020).

The scripts for RNA-Seq and ChIP-seq analysis are available here:

[\[https://github.com/pyrosiles197/POU-IV_analysis\]](https://github.com/pyrosiles197/POU-IV_analysis).

Acknowledgements:

We thank Suat Özbek for providing us with the antibodies against minicollagens, and Sakura Rieck for helping with the behavioral analysis of *pou-iv* mutants. We are also grateful to Betty Martin and Mourad Benamara at the Arkansas Nano & Bio Materials Characterization Facility for their assistance with confocal microscopy and transmission electron microscopy. Finally, we would like to thank reviewers for comments on the earlier version of the manuscript, which greatly improved the manuscript.

Competing interests:

No competing interests declared.

Funding:

This work was supported by Arkansas Bioscience Institute, University of Arkansas, and National Science Foundation Grant No.1931154.

References:

ANDERS, S., PYL, P. T. & HUBER, W. 2015. HTSeq-a Python framework to work with high-throughput sequencing data. *Bioinformatics*, 31, 166-169.

- 1461 ANDO, R., HAMA, H., YAMAMOTO-HINO, M., MIZUNO, H. & MIYAWAKI, A. 2002. An
1462 optical marker based on the UV-induced green-to-red photoconversion of a fluorescent
1463 protein. *Proceedings of the National Academy of Sciences of the United States of America*,
1464 99, 12651-12656.
- 1465 ARKETT, S. A., MACKIE, G. O. & MEECH, R. W. 1988. Hair cell mechanoreception in the
1466 jellyfish *Aglantha digitale*. *Journal of Experimental Biology*, 135, 329-342.
- 1467 BABONIS, L. S. & MARTINDALE, M. Q. 2017. PaxA, but not PaxC, is required for cnidocyte
1468 development in the sea anemone *Nematostella vectensis*. *Evodevo*, 8.
- 1469 BATEMAN, A. & SANDFORD, R. 1999. The PLAT domain: a new piece in the PKD1 puzzle.
1470 *Current Biology*, 9, R588-R590.
- 1471 BAUMEISTER, R., LIU, Y. X. & RUVKUN, G. 1996. Lineage-specific regulators couple cell
1472 lineage asymmetry to the transcription of the *Caenorhabditis elegans* POU gene *unc-86*
1473 during neurogenesis. *Genes & Development*, 10, 1395-1410.
- 1474 BEISEL, K. W., HE, D., HALLWORTH, R. & FRITZSCH, B. 2008. 3.05 - Genetics of
1475 Mechanoreceptor Evolution and Development. In: MASLAND, R. H., ALBRIGHT, T.
1476 D., ALBRIGHT, T. D., MASLAND, R. H., DALLOS, P., OERTEL, D., FIRESTEIN, S.,
1477 BEAUCHAMP, G. K., CATHERINE BUSHNELL, M., BASBAUM, A. I., KAAS, J. H.
1478 & GARDNER, E. P. (eds.) *The Senses: A Comprehensive Reference*. New York:
1479 Academic Press.
- 1480 BEZARES-CALDERON, L. A., BERGER, J. & JEKELY, G. 2020. Diversity of cilia-based
1481 mechanosensory systems and their functions in marine animal behaviour. *Philosophical*
1482 *Transactions of the Royal Society B-Biological Sciences*, 375.
- 1483 BOCA, M., D'AMATO, L., DISTEFANO, G., POLISHCHUK, R. S., GERMINO, G. G. &
1484 BOLETTA, A. 2007. Polycystin-1 induces cell migration by regulating
1485 phosphatidylinositol 3-kinase-dependent cytoskeletal Rearrangements and GSK3 beta-
1486 dependent cell-cell mechanical adhesions. *Molecular Biology of the Cell*, 18, 4050-4061.
- 1487 BOEKHOFF-FALK, G. 2005. Hearing in *Drosophila*: Development of Johnston's organ and
1488 emerging parallels to vertebrate ear development. *Developmental Dynamics*, 232, 550-
1489 558.
- 1490 BOLGER, A. M., LOHSE, M. & USADEL, B. 2014. Trimmomatic: a flexible trimmer for
1491 Illumina sequence data. *Bioinformatics*, 30, 2114-2120.
- 1492 BRINKMANN, M., OLIVER, D. & THURM, U. 1996. Mechanoelectric transduction in
1493 nematocytes of a hydropolyp (Corynidae). *Journal of Comparative Physiology A Sensory*
1494 *Neural and Behavioral Physiology*, 178, 125-138.
- 1495 BUDELMANN, B. U. 1989. Hydrodynamic receptor systems in invertebrates. *The*
1496 *mechanosensory lateral line*. New York: Springer-Verlag.
- 1497 BYCROFT, M., BATEMAN, A., CLARKE, J., HAMILL, S. J., SANDFORD, R., THOMAS, R.
1498 L. & CHOTHIA, C. 1999. The structure of a PKD domain from polycystin-1:
1499 Implications for polycystic kidney disease. *Embo Journal*, 18, 297-305.
- 1500 CHALFIE, M. & AU, M. 1989. Genetic control of differentiation of the *Caenorhabditis elegans*
1501 touch receptor neurons. *Science*, 243, 1027-33.
- 1502 CHALFIE, M., HORVITZ, H. R. & SULSTON, J. E. 1981. Mutations that lead to reiterations in
1503 the cell lineages of *C. elegans*. *Cell*, 24, 59-69.
- 1504 CHALFIE, M. & SULSTON, J. 1981. Developmental genetics of the mechanosensory neurons
1505 of *Caenorhabditis elegans*. *Dev Biol*, 82, 358-70.

1506 CLYNE, P. J., CERTTEL, S. J., DE BRUYNE, M., ZASLAVSKY, L., JOHNSON, W. A. &
 1507 CARLSON, J. R. 1999. The odor specificities of a subset of olfactory receptor neurons
 1508 are governed by Acj6, a POU-domain transcription factor. *Neuron*, 22, 339-47.
 1509 COFFIN, A., KELLEY, M., MANLEY, G. A. & POPPER, A. N. 2004. Evolution of sensory
 1510 hair cells. *Springer Handbook of Auditory Research*, 22, 55-94.
 1511 COLLINS, A. G., SCHUCHERT, P., MARQUES, A. C., JANKOWSKI, T., MEDINA, M. &
 1512 SCHIERWATER, B. 2006. Medusozoan phylogeny and character evolution clarified by
 1513 new large and small subunit rDNA data and an assessment of the utility of phylogenetic
 1514 mixture models. *Syst Biol*, 55, 97-115.
 1515 DANECEK, P., BONFIELD, J. K., LIDDLE, J., MARSHALL, J., OHAN, V., POLLARD, M.
 1516 O., WHITWHAM, A., KEANE, T., MCCARTHY, S. A., DAVIES, R. M. & LI, H. 2021.
 1517 Twelve years of SAMtools and BCFtools. *Gigascience*, 10.
 1518 DELMAS, P. 2004. Polycystins: From mechanosensation to gene regulation. *Cell*, 118, 145-148.
 1519 DOBIN, A., DAVIS, C. A., SCHLESINGER, F., DRENKOW, J., ZALESKI, C., JHA, S.,
 1520 BATUT, P., CHAISSON, M. & GINGERAS, T. R. 2013. STAR: ultrafast universal
 1521 RNA-seq aligner. *Bioinformatics*, 29, 15-21.
 1522 DUGGAN, A., MA, C. & CHALFIE, M. 1998. Regulation of touch receptor differentiation by
 1523 the *Caenorhabditis elegans* mec-3 and unc-86 genes. *Development*, 125, 4107-19.
 1524 ERKMAN, L., MCEVILLY, R. J., LUO, L., RYAN, A. K., HOOSHMAND, F., O'CONNELL,
 1525 S. M., KEITHLEY, E. M., RAPAPORT, D. H., RYAN, A. F. & ROSENFELD, M. G.
 1526 1996. Role of transcription factors Brn-3.1 and Brn-3.2 in auditory and visual system
 1527 development. *Nature*, 381, 603-6.
 1528 ERWIN, D. H., LAFLAMME, M., TWEEDT, S. M., SPERLING, E. A., PISANI, D. &
 1529 PETERSON, K. J. 2011. The Cambrian Conundrum: Early Divergence and Later
 1530 Ecological Success in the Early History of Animals. *Science*, 334, 1091-1097.
 1531 FAIN, G. L. 2003. *Sensory Transduction*, Sunderland, Massachusetts U.S.A., Sinauer Associates,
 1532 Inc.
 1533 FAUTIN, D. G. & MARISCAL, R. N. 1991. Cnidaria: Anthozoa. In: HARRISON, F. W. (ed.)
 1534 *Microscopic anatomy of invertebrates*. New York: Wiley-Liss.
 1535 FINNEY, M. & RUVKUN, G. 1990. The unc-86 gene product couples cell lineage and cell
 1536 identity in *C. elegans*. *Cell*, 63, 895-905.
 1537 FORNES, O., CASTRO-MONDRAGON, J. A., KHAN, A., VAN DER LEE, R., ZHANG, X.,
 1538 RICHMOND, P. A., MODI, B. P., CORREARD, S., GHEORGHE, M., BARANASIC,
 1539 D., SANTANA-GARCIA, W., TAN, G., CHENEY, J., BALLESTER, B., PARCY, F.,
 1540 SANDELIN, A., LENHARD, B., WASSERMAN, W. W. & MATHELIER, A. 2020.
 1541 JASPAR 2020: update of the open-access database of transcription factor binding profiles.
 1542 *Nucleic Acids Research*, 48, D87-D92.
 1543 FREDMAN, D., SCHWAIGER, M., RENTZSCH, F. & TECHNAU, U. 2013. *Nematostella*
 1544 *vectensis* transcriptome and gene models v2.0. *figshare*.
 1545 FRITZENWANKER, J. H., GENIKHOVICH, G., KRAUS, Y. & TECHNAU, U. 2007. Early
 1546 development and axis specification in the sea anemone *Nematostella vectensis*.
 1547 *Developmental Biology*, 310, 264-279.
 1548 FRITZENWANKER, J. H. & TECHNAU, U. 2002. Induction of gametogenesis in the basal
 1549 cnidarian *Nematostella vectensis*(Anthozoa). *Dev Genes Evol*, 212, 99-103.

1550 GAN, L., XIANG, M., ZHOU, L., WAGNER, D. S., KLEIN, W. H. & NATHANS, J. 1996.
 1551 POU domain factor Brn-3b is required for the development of a large set of retinal
 1552 ganglion cells. *Proc Natl Acad Sci U S A*, 93, 3920-5.
 1553 GOLD, D. A., GATES, R. D. & JACOBS, D. K. 2014. The Early Expansion and Evolutionary
 1554 Dynamics of POU Class Genes. *Molecular Biology and Evolution*, 31, 3136-3147.
 1555 GRIEBEN, M., PIKE, A. C. W., SHINTRE, C. A., VENTURI, E., EL-AJOUS, S., TESSITORE,
 1556 A., SHRESTHA, L., MUKHOPADHYAY, S., MAHAJAN, P., CHALK, R., BURGESS-
 1557 BROWN, N. A., SITSAPESAN, R., HUISKONEN, J. T. & CARPENTER, E. P. 2017.
 1558 Structure of the polycystic kidney disease TRP channel Polycystin-2 (PC2). *Nature*
 1559 *Structural & Molecular Biology*, 24, 114-+.
 1560 GRUBER, C. A., RHEE, J. M., GLEIBERMAN, A. & TURNER, E. E. 1997. POU domain
 1561 factors of the Brn-3 class recognize functional DNA elements which are distinctive,
 1562 symmetrical, and highly conserved in evolution. *Molecular and Cellular Biology*, 17,
 1563 2391-2400.
 1564 GUPTA, S., STAMATOYANNOPOULOS, J. A., BAILEY, T. L. & NOBLE, W. S. 2007.
 1565 Quantifying similarity between motifs. *Genome Biology*, 8.
 1566 GYOJA, F., KAWASHIMA, T. & SATOH, N. 2012. A genomewide survey of bHLH
 1567 transcription factors in the coral *Acropora digitifera* identifies three novel orthologous
 1568 families, pearl, amber, and peridot. *Development Genes and Evolution*, 222, 63-76.
 1569 HAND, C. & UHLINGER, K. R. 1992. The Culture, Sexual and Asexual Reproduction, and
 1570 Growth of the Sea-Anemone *Nematostella Vectensis*. *Biological Bulletin*, 182, 169-176.
 1571 HAVRILAK, J. A., AL-SHAER, L., BABAN, N., AKINCI, N. & LAYDEN, M. J. 2021.
 1572 Characterization of the dynamics and variability of neuronal subtype responses during
 1573 growth, degrowth, and regeneration of *Nematostella vectensis*. *Bmc Biology*, 19.
 1574 HEJNOL, A., OBST, M., STAMATAKIS, A., OTT, M., ROUSE, G. W., EDGECOMBE, G. D.,
 1575 MARTINEZ, P., BAGUNA, J., BAILLY, X., JONDELIUS, U., WIENS, M., MULLER,
 1576 W. E. G., SEAVER, E., WHEELER, W. C., MARTINDALE, M. Q., GIRIBET, G. &
 1577 DUNN, C. W. 2009. Assessing the root of bilaterian animals with scalable phylogenomic
 1578 methods. *Proceedings of the Royal Society B-Biological Sciences*, 276, 4261-4270.
 1579 HERR, W. & CLEARY, M. A. 1995. The Pou Domain - Versatility in Transcriptional
 1580 Regulation by a Flexible 2-in-One DNA-Binding Domain. *Genes & Development*, 9,
 1581 1679-1693.
 1582 HOLLAND, L. Z. 2005. Non-neural ectoderm is really neural: Evolution of developmental
 1583 patterning mechanisms in the non-neural ectoderm of chordates and the problem of
 1584 sensory cell homologies. *Journal of Experimental Zoology Part B Molecular and*
 1585 *Developmental Evolution*, 304B, 304-323.
 1586 HOLTMAN, M. & THURM, U. 2001. Variations of concentric hair cells in a cnidarian sensory
 1587 epithelium (*Coryne tubulosa*). *Journal of Comparative Neurology*, 432, 550-563.
 1588 HORRIDGE, G. A. 1969. Statocysts of medusae and evolution of stereocilia. *Tissue & Cell*, 1,
 1589 341-353.
 1590 HROUDOVA, M., VOJTA, P., STRNAD, H., KREJCIK, Z., RIDL, J., PACES, J., VLCEK, C.
 1591 & PACES, V. 2012. Diversity, Phylogeny and Expression Patterns of Pou and Six
 1592 Homeodomain Transcription Factors in Hydrozoan Jellyfish *Craspedacusta sowerbyi*.
 1593 *PLoS ONE*, 7, e36420, 1-11.
 1594 HUNDGEN, M. & BIELA, C. 1982. Fine structure of touch-plates in the scyphomedusan
 1595 *Aurelia aurita*. *Journal of ultrastructure research*, 80, 178-184.

IKEDA, R., PAK, K., CHAVEZ, E. & RYAN, A. F. 2015. Transcription Factors with Conserved Binding Sites Near ATOH1 on the POU4F3 Gene Enhance the Induction of Cochlear Hair Cells. *Molecular Neurobiology*, 51, 672-684.

IKMI, A., MCKINNEY, S. A., DELVENTHAL, K. M. & GIBSON, M. C. 2014. TALEN and CRISPR/Cas9-mediated genome editing in the early-branching metazoan *Nematostella vectensis*. *Nature Communications*, 5.

JALILI, V., MATTEUCCI, M., MASSEROLI, M. & MORELLI, M. J. 2015. Using combined evidence from replicates to evaluate ChIP-seq peaks. *Bioinformatics*, 31, 2761-2769.

JARMAN, A. P. 2002. Studies of mechanosensation using the fly. *Human Molecular Genetics*, 11, 1215-1218.

JØRGENSEN, J. M. Evolution of Octavolateralis Sensory Cells. 1989 New York, NY. Springer New York, 115-145.

KELLEY, L. A., MEZULIS, S., YATES, C. M., WASS, M. N. & STERNBERG, M. J. E. 2015. The Phyre2 web portal for protein modeling, prediction and analysis. *Nature Protocols*, 10, 845-858.

KITAJIRI, S., SAKAMOTO, T., BELYANTSEVA, I. A., GOODYEAR, R. J., STEPANYAN, R., FUJIWARA, I., BIRD, J. E., RIAZUDDIN, S., RIAZUDDIN, S., AHMED, Z. M., HINSHAW, J. E., SELLERS, J., BARTLES, J. R., HAMMER, J. A., RICHARDSON, G. P., GRIFFITH, A. J., FROLENKOV, G. I. & FRIEDMAN, T. B. 2010. Actin-Bundling Protein TRIOBP Forms Resilient Rootlets of Hair Cell Stereocilia Essential for Hearing. *Cell*, 141, 786-798.

KRAUS, Y. & TECHNAU, U. 2006. Gastrulation in the sea anemone *Nematostella vectensis* occurs by invagination and immigration: an ultrastructural study. *Dev Genes Evol*, 216, 119-32.

LANDT, S. G., MARINOV, G. K., KUNDAJE, A., KHERADPOUR, P., PAULI, F., BATZOGLOU, S., BERNSTEIN, B. E., BICKEL, P., BROWN, J. B., CAYTING, P., CHEN, Y. W., DESALVO, G., EPSTEIN, C., FISHER-AYLOR, K. I., EUSKIRCHEN, G., GERSTEIN, M., GERTZ, J., HARTEMINK, A. J., HOFFMAN, M. M., IYER, V. R., JUNG, Y. L., KARMAKAR, S., KELLIS, M., KHARCHENKO, P. V., LI, Q. H., LIU, T., LIU, X. S., MA, L. J., MILOSAVLJEVIC, A., MYERS, R. M., PARK, P. J., PAZIN, M. J., PERRY, M. D., RAHA, D., REDDY, T. E., ROZOWSKY, J., SHORESH, N., SIDOW, A., SLATTERY, M., STAMATOYANNOPOULOS, J. A., TOLSTORUKOV, M. Y., WHITE, K. P., XI, S., FARNHAM, P. J., LIEB, J. D., WOLD, B. J. & SNYDER, M. 2012. ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome Research*, 22, 1813-1831.

LEE, P. N., KUMBUREGAMA, S., MARLOW, H. Q., MARTINDALE, M. Q. & WIKRAMANAYAKE, A. H. 2007. Asymmetric developmental potential along the animal-vegetal axis in the anthozoan cnidarian, *Nematostella vectensis*, is mediated by Dishevelled. *Developmental Biology*, 310, 169-186.

LESH-LAURIE, G. E. & SUCHY, P. E. 1991. Cnidaria: Scyphozoa and Cubozoa. In: HARRISON, F. W. (ed.) *Microscopic anatomy of invertebrates*. New York: Wiley-Liss.

LEYVA-DIAZ, E., MASOUDI, N., SERRANO-SAIZ, E., GLENWINKEL, L. & HOBERT, O. 2020. Brn3/POU-IV-type POU homeobox genes-Paradigmatic regulators of neuronal identity across phylogeny. *Wiley Interdisciplinary Reviews-Developmental Biology*, 9.

1640 LI, C. K., WEN, A. Y., SHEN, B. C., LU, J., HUANG, Y. & CHANG, Y. C. 2011. FastCloning:
 1641 a highly simplified, purification-free, sequence- and ligation-independent PCR cloning
 1642 method. *Bmc Biotechnology*, 11.
 1643 LIU, W., MO, Z. Q. & XIANG, M. Q. 2001. The Ath5 proneural genes function upstream of
 1644 Brn3 POU domain transcription factor genes to promote retinal ganglion cell
 1645 development. *Proceedings of the National Academy of Sciences of the United States of*
 1646 *America*, 98, 1649-1654.
 1647 LOPEZ-ESCARDO, D., GRAU-BOVE, X., GUILLAUMET-ADKINS, A., GUT, M.,
 1648 SIERACKI, M. E. & RUIZ-TRILLO, I. 2019. Reconstruction of protein domain
 1649 evolution using single-cell amplified genomes of uncultured choanoflagellates sheds light
 1650 on the origin of animals. *Philosophical Transactions of the Royal Society B-Biological*
 1651 *Sciences*, 374.
 1652 LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and
 1653 dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15.
 1654 LYONS, K. M. 1973. Collar Cells in Planula and Adult Tentacle Ectoderm of Solitary Coral
 1655 Balanophyllia-Regia (Anthozoa Eupsammiidae). *Zeitschrift Fur Zellforschung Und*
 1656 *Mikroskopische Anatomie*, 145, 57-74.
 1657 MAGIE, C. R., DALY, M. & MARTINDALE, M. Q. 2007. Gastrulation in the cnidarian
 1658 *Nematostella vectensis* occurs via invagination not ingression. *Developmental Biology*,
 1659 305, 483-497.
 1660 MAHONEY, J. L., GRAUGNARD, E. M., MIRE, P. & WATSON, G. M. 2011. Evidence for
 1661 involvement of TRPA1 in the detection of vibrations by hair bundle mechanoreceptors in
 1662 sea anemones. *Journal of Comparative Physiology A Sensory Neural and Behavioral*
 1663 *Physiology*, 197, 729-742.
 1664 MANLEY, G. A. & LADHER, R. 2008. 3.01 - Phylogeny and Evolution of Ciliated
 1665 Mechanoreceptor Cells. In: MASLAND, R. H., ALBRIGHT, T. D., ALBRIGHT, T. D.,
 1666 MASLAND, R. H., DALLOS, P., OERTEL, D., FIRESTEIN, S., BEAUCHAMP, G. K.,
 1667 CATHERINE BUSHNELL, M., BASBAUM, A. I., KAAS, J. H. & GARDNER, E. P.
 1668 (eds.) *The Senses: A Comprehensive Reference*. New York: Academic Press.
 1669 MARTINDALE, M. Q., PANG, K. & FINNERTY, J. R. 2004. Investigating the origins of
 1670 triploblasty: 'mesodermal' gene expression in a diploblastic animal, the sea anemone
 1671 *Nematostella vectensis* (phylum, Cnidaria; class, Anthozoa). *Development*, 131, 2463-74.
 1672 MASUDA, M., DULON, D., PAK, K., MULLEN, L. M., LI, Y., ERKMAN, L. & RYAN, A. F.
 1673 2011. Regulation of Pou4f3 Gene Expression in Hair Cells by 5' DNA in Mice.
 1674 *Neuroscience*, 197, 48-64.
 1675 MCEVILLY, R. J., ERKMAN, L., LUO, L., SAWCHENKO, P. E., RYAN, A. F. &
 1676 ROSENFELD, M. G. 1996. Requirement for Brn-3.0 In differentiation and survival of
 1677 sensory and motor neurons. *Nature*, 384, 574-577.
 1678 MEDINA, M., COLLINS, A. G., SILBERMAN, J. D. & SOGIN, M. L. 2001. Evaluating
 1679 hypotheses of basal animal phylogeny using complete sequences of large and small
 1680 subunit rRNA. *Proceedings of the National Academy of Sciences of the United States of*
 1681 *America*, 98, 9707-9712.
 1682 MIRE, P. & WATSON, G. M. 1997. Mechanotransduction of hair bundles arising from
 1683 multicellular complexes in anemones. *Hearing Research*, 113, 224-234.

- MIRE-THIBODEAUX, P. & WATSON, G. M. 1994. Morphodynamic hair bundles arising from sensory cell/supporting cell complexes frequency-tune nematocyst discharges in sea anemones. *Journal of Experimental Zoology*, 268, 282-292.
- MIRETHIBODEAUX, P. & WATSON, G. M. 1994. Morphodynamic Hair Bundles Arising from Sensory Cell/Supporting Cell Complexes Frequency-Tune Nematocyst Discharge in Sea-Anemones. *Journal of Experimental Zoology*, 268, 282-292.
- MOCHIZUKI, T., WU, G. Q., HAYASHI, T., XENOPHONTOS, S. L., VELDHUISEN, B., SARIS, J. J., REYNOLDS, D. M., CAI, Y. Q., GABOW, P. A., PIERIDES, A., KIMBERLING, W. J., BREUNING, M. H., DELTAS, C. C., PETERS, D. J. M. & SOMLO, S. 1996. PKD2, a gene for polycystic kidney disease that encodes an integral membrane protein. *Science*, 272, 1339-1342.
- MOY, G. W., MENDOZA, L. M., SCHULZ, J. R., SWANSON, W. J., GLABE, C. G. & VACQUIER, V. D. 1996. The sea urchin sperm receptor for egg jelly is a modular protein with extensive homology to the human polycystic kidney disease protein, PKD1. *Journal of Cell Biology*, 133, 809-817.
- NAKANISHI, N. & MARTINDALE, M. Q. 2018. CRISPR knockouts reveal an endogenous role for ancient neuropeptides in regulating developmental timing in a sea anemone. *eLife*, 7, e39742.
- NAKANISHI, N., RENFER, E., TECHNAU, U. & RENTZSCH, F. 2012. Nervous systems of the sea anemone *Nematostella vectensis* are generated by ectoderm and endoderm and shaped by distinct mechanisms. *Development*, 139, 347-357.
- NAKANISHI, N., YUAN, D., HARTENSTEIN, V. & JACOBS, D. K. 2010. Evolutionary origin of rhopalia: insights from cellular-level analyses of Otx and POU expression patterns in the developing rhopalial nervous system. *Evolution & Development*, 12, 404-415.
- OLIVER, D. & THURM, U. 1996. Mechanoreception by concentric hair cells of hydropolyps. In: ELSNER, N. & SCHNITZLER, H. U. (eds.) *Gottingen Neurobiology Report*. Stuttgart: Georg Thieme.
- PUTNAM, N. H., SRIVASTAVA, M., HELLSTEN, U., DIRKS, B., CHAPMAN, J., SALAMOV, A., TERRY, A., SHAPIRO, H., LINDQUIST, E., KAPITONOV, V. V., JURKA, J., GENIKHOVICH, G., GRIGORIEV, I. V., LUCAS, S. M., STEELE, R. E., FINNERTY, J. R., TECHNAU, U., MARTINDALE, M. Q. & ROKHSAR, D. S. 2007. Sea anemone genome reveals ancestral eumetazoan gene repertoire and genomic organization. *Science*, 317, 86-94.
- RENER, E., AMON-HASSENZAHN, A., STEINMETZ, P. R. H. & TECHNAU, U. 2010. A muscle-specific transgenic reporter line of the sea anemone, *Nematostella vectensis*. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 104-108.
- RICHARDS, G. S. & RENTZSCH, F. 2014. Transgenic analysis of a SoxB gene reveals neural progenitor cells in the cnidarian *Nematostella vectensis*. *Development*, 141, 4681-4689.
- RICHARDS, G. S. & RENTZSCH, F. 2015. Regulation of *Nematostella* neural progenitors by SoxB, Notch and bHLH genes. *Development*, 142, 3332-3342.
- SANDFORD, R., SGOTTO, B., APARICIO, S., BRENNER, S., VAUDIN, M., WILSON, R. K., CHISSOE, S., PEPIN, K., BATEMAN, A., CHOTHIA, C., HUGHES, J. & HARRIS, P. 1997. Comparative analysis of the polycystic kidney disease 1 (PKD1) gene reveals an

1729 integral membrane glycoprotein with multiple evolutionary conserved domains. *Human*
1730 *Molecular Genetics*, 6, 1483-1489.

1731 SCHLOSSER, G. 2020. 2.17 - Evolution of Hair Cells1. In: FRITZSCH, B. (ed.) *The Senses: A*
1732 *Comprehensive Reference (Second Edition)*. Oxford: Elsevier.

1733 SEBE-PEDROS, A., SAUDEMONT, B., CHOMSKY, E., PLESSIER, F., MAILHE, M. P.,
1734 RENNO, J., LOE-MIE, Y., LIFSHITZ, A., MUKAMEL, Z., SCHMUTZ, S., NOVAULT,
1735 S., STEINMETZ, P. R. H., SPITZ, F., TANAY, A. & MARLOW, H. 2018. Cnidarian
1736 Cell Type Diversity and Regulation Revealed by Whole-Organism Single-Cell RNA-Seq.
1737 *Cell*, 173, 1520-+.

1738 SERRANO-SAIZ, E., LEYVA-DIAZ, E., DE LA CRUZ, E. & HOBERT, O. 2018. BRN3-type
1739 POU Homeobox Genes Maintain the Identity of Mature Postmitotic Neurons in
1740 Nematodes and Mice. *Curr Biol*, 28, 2813-2823 e2.

1741 SILVA, M. A. P. & NAKANISHI, N. 2019. Genotyping of Sea Anemone during Early
1742 Development. *Jove-Journal of Visualized Experiments*.

1743 SIMIONATO, E., LEDENT, V., RICHARDS, G., THOMAS-CHOLLIER, M., KERNER, P.,
1744 COORNAERT, D., DEGNAN, B. M. & VERVOORT, M. 2007. Origin and
1745 diversification of the basic helix-loop-helix gene family in metazoans: insights from
1746 comparative genomics. *Bmc Evolutionary Biology*, 7.

1747 SINGLA, C. L. 1975. Statocysts of Hydromedusae. *Cell and Tissue Research*, 158, 391-407.

1748 SINGLA, C. L. 1983. Fine-Structure of the Sensory Receptors of Aglantha-Digitale
1749 (Hydromedusae, Trachylina). *Cell and Tissue Research*, 231, 415-425.

1750 SOCK, E., ENDERICH, J., ROSENFELD, M. G. & WEGNER, M. 1996. Identification of the
1751 nuclear localization signal of the POU domain protein Tst-1/Oct6. *Journal of Biological*
1752 *Chemistry*, 271, 17512-17518.

1753 SOLE, M., LENOIR, M., FONTUNO, J. M., DURFORT, M., VAN DER SCHAAR, M. &
1754 ANDRE, M. 2016. Evidence of Cnidarians sensitivity to sound after exposure to low
1755 frequency noise underwater sources. *Scientific Reports*, 6.

1756 STEIGELMAN, K. A., LELLI, A., WU, X. D., GAO, J. G., LIN, S., PIONTEK, K.,
1757 WODARCZYK, C., BOLETTA, A., KIM, H., QIAN, F., GERMINO, G., GELEOC, G.
1758 S. G., HOLT, J. R. & ZUO, J. 2011. Polycystin-1 Is Required for Stereocilia Structure
1759 But Not for Mechanotransduction in Inner Ear Hair Cells. *Journal of Neuroscience*, 31,
1760 12241-12250.

1761 TARDENT, P. & SCHMID, V. 1972. Ultrastructure of Mechanoreceptors of Polyp Coryne-
1762 Pintneri (Hydrozoa, Athecata). *Experimental Cell Research*, 72, 265-&.

1763 THOMAS, M. & EDWARDS, N. 1991. Cnidaria: Hydrozoa. In: FW, H. (ed.) *Microscopic*
1764 *anatomy of invertebrates*. New York: Wiley-Liss.

1765 TOURNIERE, O., DOLAN, D., RICHARDS, G. S., SUNAGAR, K., COLUMBUS-SHENKAR,
1766 Y. Y., MORAN, Y. & RENTZSCH, F. 2020. NvPOU4/Brain3 Functions as a Terminal
1767 Selector Gene in the Nervous System of the Cnidarian Nematostella vectensis. *Cell Rep*,
1768 30, 4473-4489 e5.

1769 VAHAVA, O., MORELL, R., LYNCH, E. D., WEISS, S., KAGAN, M. E., AHITUV, N.,
1770 MORROW, J. E., LEE, M. K., SKVORAK, A. B., MORTON, C. C., BLUMENFELD,
1771 A., FRYDMAN, M., FRIEDMAN, T. B., KING, M. C. & AVRAHAM, K. B. 1998.
1772 Mutation in transcription factor POU4F3 associated with inherited progressive hearing
1773 loss in humans. *Science*, 279, 1950-1954.

1774 WATSON, G. M., MIRE, P. & HUDSON, R. R. 1997. Hair bundles of sea anemones as a model
 1775 system for vertebrate hair bundles. *Hearing Research*, 107, 53-66.
 1776 WATSON, G. M., MIRE, P. & KINLER, K. M. 2009. Mechanosensitivity in the model sea
 1777 anemone *Nematostella vectensis*. *Marine Biology (Berlin)*, 156, 2129-2137.
 1778 WATSON, G. M., PHAM, L., GRAUGNARD, E. M. & MIRE, P. 2008. Cadherin 23-like
 1779 polypeptide in hair bundle mechanoreceptors of sea anemones. *Journal of Comparative*
 1780 *Physiology A Sensory Neural and Behavioral Physiology*, 194, 811-820.
 1781 WATSON, G. M. & ROBERTS, J. 1995. Chemoreceptor-Mediated Polymerization and
 1782 Depolymerization of Actin in Hair Bundles of Sea-Anemones. *Cell Motility and the*
 1783 *Cytoskeleton*, 30, 208-220.
 1784 WESTFALL, J. A. 2004. Neural pathways and innervation of cnidocytes in tentacles of sea
 1785 anemones. *Hydrobiologia*, 530-, 117-121.
 1786 WOLENSKI, F. S., BRADHAM, C. A., FINNERTY, J. R. & GILMORE, T. D. 2013. NF-kappa
 1787 B is required for cnidocyte development in the sea anemone *Nematostella vectensis*.
 1788 *Developmental Biology*, 373, 205-215.
 1789 XIANG, M., GAN, L., LI, D., CHEN, Z. Y., ZHOU, L., O'MALLEY, B. W., JR., KLEIN, W. &
 1790 NATHANS, J. 1997a. Essential role of POU-domain factor Brn-3c in auditory and
 1791 vestibular hair cell development. *Proc Natl Acad Sci U S A*, 94, 9445-50.
 1792 XIANG, M., GAN, L., LI, D., ZHOU, L., CHEN, Z. Y., WAGNER, D., O'MALLEY, B. W., JR.,
 1793 KLEIN, W. & NATHANS, J. 1997b. Role of the Brn-3 family of POU-domain genes in
 1794 the development of the auditory/vestibular, somatosensory, and visual systems. *Cold*
 1795 *Spring Harb Symp Quant Biol*, 62, 325-36.
 1796 XIANG, M. Q., GAN, L., ZHOU, L. J., KLEIN, W. H. & NATHANS, J. 1996. Targeted
 1797 deletion of the mouse POU domain gene Brn-3a causes a selective loss of neurons in the
 1798 brainstem and trigeminal ganglion, uncoordinated limb movement, and impaired suckling.
 1799 *Proceedings of the National Academy of Sciences of the United States of America*, 93,
 1800 11950-11955.
 1801 XIANG, M. Q., GAO, W. Q., HASSON, T. & SHIN, J. J. 1998. Requirement for Brn-3c in
 1802 maturation and survival, but not in fate determination of inner ear hair cells. *Development*,
 1803 125, 3935-3946.
 1804 YU, H. Z. V., TAO, L. T., LLAMAS, J., WANG, X. Z., NGUYEN, J. D., TRECEK, T. &
 1805 SEGIL, N. 2021. POU4F3 pioneer activity enables ATOH1 to drive diverse
 1806 mechanoreceptor differentiation through a feed-forward epigenetic mechanism.
 1807 *Proceedings of the National Academy of Sciences of the United States of America*, 118.
 1808 ZAPATA, F., GOETZ, F. E., SMITH, S. A., HOWISON, M., SIEBERT, S., CHURCH, S. H.,
 1809 SANDERS, S. M., AMES, C. L., MCFADDEN, C. S., FRANCE, S. C., DALY, M.,
 1810 COLLINS, A. G., HADDOCK, S. H. D., DUNN, C. W. & CARTWRIGHT, P. 2015.
 1811 Phylogenomic Analyses Support Traditional Relationships within Cnidaria. *Plos One*, 10.
 1812 ZENKERT, C., TAKAHASHI, T., DIESNER, M. O. & OZBEK, S. 2011. Morphological and
 1813 Molecular Analysis of the *Nematostella vectensis* Cnidom. *Plos One*, 6.
 1814 ZHANG, Y., LIU, T., MEYER, C. A., EECKHOUTE, J., JOHNSON, D. S., BERNSTEIN, B. E.,
 1815 NUSSBAUM, C., MYERS, R. M., BROWN, M., LI, W. & LIU, X. S. 2008. Model-
 1816 based Analysis of ChIP-Seq (MACS). *Genome Biology*, 9.
 1817 ZHU, A. Q., IBRAHIM, J. G. & LOVE, M. I. 2019. Heavy-tailed prior distributions for
 1818 sequence count data: removing the noise and preserving large differences. *Bioinformatics*,
 1819 35, 2084-2092.

1820
1821 **Supplementary Files:**
1822 **Supplementary File 1: List of 4,188 candidate POU-IV downstream target genes.**
1823 **Supplementary File 2: List of 577 genes significantly downregulated in NvPOU4 mutants**
1824 **relative to their siblings.**
1825 **Supplementary File 3: List of 657 genes significantly upregulated in NvPOU4 mutants**
1826 **relative to their siblings.**
1827 **Supplementary File 4: List of Gene Ontology terms overrepresented in genes**
1828 **downregulated in NvPOU4 mutants relative to their siblings.**
1829 **Supplementary File 5: List of Gene Ontology terms overrepresented in genes upregulated**
1830 **in NvPOU4 mutants relative to their siblings.**
1831 **Supplementary File 6: List of 293 POU-IV-activated genes.**
1832 **Supplementary File 7: List of 178 POU-IV-repressed genes.**
1833 **Supplementary File 8: List of Gene Ontology terms overrepresented in POU-IV-activated**
1834 **genes.**
1835 **Supplementary File 9: List of POU-IV downstream target genes represented in the adult**
1836 **metacell c79 (hair cell).**
1837 **Supplementary File 10: List of Gene Ontology terms overrepresented in POU-IV-activated**
1838 **genes represented in the adult metacell c79 (hair cell).**
1839 **Supplementary File 11: List of POU-IV downstream target genes represented in the**
1840 **cnidocyte metacell c8.**
1841 **Supplementary File 12: List of Gene Ontology terms overrepresented in POU-IV-repressed**
1842 **genes represented in the cnidocyte metacell c8.**
1843 **Supplementary File 13: Lists of POU-IV downstream target genes represented in POU-IV**
1844 **positive adult metacells c63, c64, c65, c66, c75, c76, c100, c101 and c102.**
1845 **Supplementary File 14: List of GPCR-encoding genes in the adult metacell c79 (hair cell).**
1846 **Supplementary File 15: List of gene specific primer sequences used to amplify *polycystin 1***
1847 **cDNA.**

Planula

A

Pha

Tentacle-bud

B

Primary polyp

C

Figure 2

A

Nv genome
scaffold 16

B **C** **D**

Nv genome
scaffold 16

a

1495 bp

b

1263 bp

pou-iv F0

WT

1

2

1

2

A

Nv genome
scaffold 16

B

	Before touch	After touch
F2 pou-iv +/+	A	B 100% (n=44)
F2 pou-iv -/-	C	D 65% (n=40)

F2 pou-iv +/+

F2 pou-iv -/-

F2 pou-iv -/-

mv

ve

rl_{ci}

A

B

C

Figure 8

PKD 1

PKD 2

0.6