Quantitative Mapping of Keratin Networks in 3D

Reinhard Windoffer, Nicole Schwarz, Sungjun Yoon, Teodora Piskova, Michael Scholkemper, Johannes Stegmaier, Andrea Bönsch, Jacopo Di Russo, Rudolf E. Leube

* Correspondence:

Corresponding Author

rleube@ukaachen.de, rwindoffer@ukaachen.de

Keywords: cytoskeleton, intermediate filament, keratin, retinal pigment epithelium, microscopy, 3D network, image analysis, 3D visualization, virtual reality

Abstract

Mechanobiology requires precise quantitative information on processes taking place in specific 3D microenvironments. Connecting the abundance of microscopical, molecular, biochemical and cell mechanical data with defined topologies has turned out to be extremely difficult. Establishing such structural and functional 3D maps needed for biophysical modeling is a particular challenge for the cytoskeleton, which consists of long and interwoven filamentous polymers coordinating subcellular processes and interactions of cells with their environment. To date, useful tools are available for the segmentation and modeling of actin filaments and microtubules but comprehensive tools for the mapping of intermediate filament organization are still lacking. In this work, we describe a workflow to model and examine the complete 3D arrangement of the keratin intermediate filament cytoskeleton in canine, murine and human epithelial cells both, in vitro and in vivo. Numerical models are derived from confocal Airyscan high resolution 3D imaging of fluorescence-tagged keratin filaments. They are interrogated and annotated at different length scales using different modes of visualization including immersive virtual reality. In this way, information is provided on network organization at the subcellular level including mesh arrangement, density and isotropic configuration as well as details on filament morphology such as bundling, curvature and orientation. We show that the comparison of these parameters helps to identify, in quantitative terms, similarities and differences of keratin network organization in epithelial cell types defining subcellular domains, notably basal, apical, lateral and perinuclear systems. The described approach and the presented data are pivotal for generating mechanobiological models that can be experimentally tested.

1 Introduction

Quantification of cellular properties is a central task of cell biology (Howard, 2014; Liberali and Pelkmans, 2012; Lidke, 2017; Lippincott-Schwartz, 2014; Piston, 2014; Roca-Cusachs et al., 2017). It is necessary for scientific analysis, testing and model building. The structural diversity of eukaryotic cells supports their complex functions and is essential to provide local microenvironments, which facilitate and enhance the probability of highly specialized molecular interactions and reactions. The cytoskeleton is a key determinant of the necessary spatial organization. It consists of filamentous polymers, which assemble into microtubules, actin filaments and intermediate filaments. The resulting networks are interlinked and connected to intra- and

extracellular targets, including cytoplasmic organelles and specialized plasma membrane sites. In this way, they act as structural and mechanical connectors between extra- and intracellular forces. The participating structures and polypeptides have been referred to as the mechanobiome (Kothari et al., 2019; Parajón et al., 2021; Wang et al., 2014), which defines the mechanical setting of a cell. The properties of several mechanobiome components have already been quantified including their turnover and mechanical features at different levels. Comprehensive quantitative information on entire cytoskeletal networks, however, is missing for most components and, consequently, only little is known about the distribution, local composition, mesh size, orientation and mechanical properties of cytoskeletal networks at the subcellular level. Therefore, there is an urgent need to describe cytoskeletal network architectures in quantitative terms at subcellular resolution. To this end, we present a novel approach to create the complete digital representation of keratin intermediate filament networks in three dimensions (3D), providing maps and quantitative measurements of the entire keratin network in cultured cells and in cells within their native tissue context.

The intermediate filament cytoskeleton consists of polypeptides, which are expressed in a tissue- and context-specific manner. The largest group of intermediate filaments, the keratin intermediate filaments are the main cytoplasmic networks in epithelial tissues. They confer, together with their associated desmosomal cell-cell and hemidesmosomal cell-extracellular matrix adhesions, mechanical resilience (Hatzfeld et al., 2017; te Molder et al., 2021). The human genome contains 54 keratin genes, which are expressed in epithelia and epidermal appendages in a cell type-specific pattern (Jacob et al., 2018). Keratin filaments are highly flexible and extensible (Block et al., 2015). They form hollow tubes with remarkable variability in local diameter and ultrastructure (Weber et al., 2021). Keratin filaments generate branched bundles of different thickness forming anisotropic networks formed (Beil et al., 2005; Martin et al., 2016). The 3D organization of keratin networks differs profoundly between different epithelia: E.g., the simple epithelial cells lining the intestine contain a well-defined lateral and apical network, keratinocytes contain a dense 3D network encompassing the entire cytoplasm, and hepatocytes have a prominent submembranous network (Schwarz et al., 2015; Strnad et al., 2008). Network organization is furthermore affected by mechanical stress (Beriault et al., 2012; Felder et al., 2008; Fois et al., 2013; Karsch et al., 2020; Lutz et al., 2020; Quinlan et al., 2017) and differentiation-status (Coch and Leube, 2016; Iwatsuki and Suda, 2010; Leube et al., 2017; Quinlan et al., 2017; Tateishi et al., 2017). In addition, individual proteins such as Ndel1 and plectin have been shown to regulate keratin network organization (Kim et al., 2021; Moch et al., 2016; Osmanagic-Myers et al., 2006). Knowledge of the 3D keratin network organization in single epithelial cells is therefore important to understand and characterize the particular functional status of a given cell within its complex tissue context under different conditions.

Segmentation of cellular elements is a key task in biological image analysis. Its application to specific structures, however, can be quite difficult and requires appropriate adjustment. Networks of curvilinear objects such as keratin networks are ubiquitous ranging from cytoskeletal filaments at the micro- to nanoscale to blood vessels or neuronal networks at the tissue and organ level (Özdemir and Reski, 2021). Filament detection systems have been developed for vascular (Imran et al., 2019; Moccia et al., 2018; Samuel and Veeramalai, 2020) and neuronal networks (Acciai et al., 2016; Magliaro et al., 2019; Quan et al., 2016). But these tools have not been optimized for subcellular structures and are specialized to detect branched, tree-like structures. The fishnet-like organization of keratin networks poses novel challenges, which are further confounded by the variably spaced branching points connecting filaments of different thickness and curvature.

A wide range of specialized tools are available for segmentation of cytoskeletal actin (Alioscha-Perez et al., 2016; Nowak et al., 2020), which also work in 3D (Haspinger et al., 2021). Further tools have been developed for the segmentation of microtubules (Faulkner et al., 2017) and their prokaryotic homologue FtsZ (Liu et al., 2019; Özdemir and Reski, 2021). Recently, a tool was described that is suitable for segmentation of cytoskeletal networks including keratin networks using data obtained by electron microscopy and super-resolution fluorescence microscopy (Flormann et al., 2021). This tool, however, lacks 3D capability and does not capture information on filament thickness. Another segmentation tool for filamentous networks in electronmicroscopic images was also recently reported (Dimchev et al., 2021). Until now, however, no tools are available that segment 3D fluorescence image data of keratin filament networks and transform them into a numerical graph model.

To segment keratin filament networks and generate data structures with defined nodes and internodal regions (= segments) we used the open-source program TSOAX. TSOAX has been applied for 3D segmentation of actin networks using fluorescence images of yeast cells (Xu et al., 2019). It is an implementation of SOAX (Xu et al., 2015, 2011), which was developed from the Stretching Open Active Contours (SOAC) algorithm (Li et al., 2010, 2009b, 2009a). SOACs are open curves, which start at segment tips and delineate the filament centers. We now show that TSOAX can be used to reliably segment keratin networks in entire cells. TSOAX also provides density information of the identified keratin filaments. However, the association of nodes and segments is not detected satisfactorily as the generated TSOAX snakes are not separated consistently at filament nodes and therefore hinder statistical analysis. We therefore developed additional tools (KerNet) for optimal network representation of a node-segment structure. The combination of the established segmentation algorithms (TSOAX) and the additional KerNet analysis tools helped to create accurate numerical representations of keratin network topology.

The developed image analysis tools were applied to confocal airyscan high-resolution fluorescence image stacks of keratin networks in single cells either in cultured epithelial monolayers or within their native tissue context. We show that the image analysis tools provide cell type-specific information on filament mesh characteristics as well as information on filament thickness, directionality and connectivity in 3D. Cinematic rendering and immersive visualization make analysis of network heterogeneity accessible at a hitherto unknown quality and precision of 3D reconstruction.

2 RESULTS

2.1 Recording of the keratin intermediate filament network in 3D

In a first step, we wanted to generate high-resolution fluorescence data that would be suitable for image analysis of single filaments and filament bundles and would represent keratin filament networks in different epithelial cell types. We selected three non-carcinogenic epithelial cell types: (i) Canine kidney epithelial MDCK cells producing YFP-tagged keratin 8 (subclone H9) were chosen as representatives of polarized simple epithelial cells. MDCK cells form a one-layered coherent epithelium with the cuboidal to cylindrical cells attached to each other by highly-organized junctional complexes. Intracellularly, the keratin filaments have a complex arrangement consisting of a dense subapical network, perinuclear and radial keratin filaments and interdesmosomal filaments below the plasma membrane (Quinlan et al., 2017). Immunoblot analysis of total cell lysates revealed comparable amounts of YFP-tagged and endogenous keratin 8 (**Figure 1-figure supplement 1**). (ii) Spontaneously immortalized human epidermal keratinocytes expressing YFP-tagged keratin 5

(HaCaT B9) were chosen as representatives of squamous epithelial cells -(Boukamp et al., 1988; Moch et al., 2013). They contain pancytoplasmic keratin filament networks whose 3D organization is only partially elucidated (Moch et al., 2020). (iii) To examine the 3D organization of the keratin intermediate filament network in its native tissue context, we made use of the recently described knock-in mouse line expressing only YFP-tagged keratin 8 but no wild-type keratin 8. Retinal epithelium cells (RPE) were selected for the analysis because of their well-delineated cytoplasmic network (see below). High resolution airyscan confocal image stacks were prepared from each cell type.

2.2 Transformation of 3D fluorescence recordings of keratin filaments into digital network models

The recorded image stacks were transformed into Euclidean 3D maps using TSOAX and the KerNet software (**Figures 1, 2; Figure 1-figure supplement 2**). In these maps, curved segments that are connected by nodes describe the entire keratin networks of single cells. The segments represent keratin filament bundles of different thickness and length. These segments are defined as 3D polygonal chains of vertices, each of which has a specific xyz position and brightness. The first and last vertices of a segment are defined as nodes. Nodes are either shared by adjacent connecting segments within a filament or are positioned at the end of a filament in case of loose ends. Note that keratin filaments consist of multiple consecutive segments. To validate the quality of the numerical transformation, all vertices were plotted onto the maximum intensity projection of the fluorescence image stacks. **Figure 1A** shows the maximum intensity projection of the fluorescence in an MDCK cell as an example highlighting selected regions of interest at higher magnification. Separating the regions above and below the nucleus (apical and basal, respectively) furthermore allowed to distinguish the different keratin network structures in these subcellular domains. Visual inspection showed a high degree of overlap between the recorded fluorescence and the derived numerical positions of vertices (dotted lines) at all subcellular locations.

To obtain more detailed comparisons of the calculated vertices with the fluorescence data, vertices were first plotted in 3D and their brightness was encoded as filament diameter. The 3D plots were then convolved by Gaussian blurring to simulate the microscope's point spread function. **Figure 1B** shows the very good match of the convolved vertices in terms of position and brightness in comparison to the maximum intensity projection of the fluorescence image. To assess the overall reliability of the digital transformation, the validation steps were applied to all cells selected for the study (**Figure 1-figure supplement 3**). Additional slice-by-slice comparisons are presented exemplarily in **Video 1** further demonstrating that the recorded and derived numerical 3D representations of the keratin filament network are in agreement.

For close inspection of the keratin filament network in its subcellular complexity, we used cinematic rendering. The rendering displays the network as a true 3D model including the brightness of filaments which are shown as tubes with different degrees of thickness thus providing information on the segments in a 3D-viewer. To distinguish individual segments, each segment was randomly color-coded (**Figure 1C**).

2.3 Comparison of 3D cinematic keratin filament renderings in MDCK, HaCaT and RPE cells

Renderings of keratin filament networks in MDCK, HaCaT and RPE cells are exemplarily shown from different views in **Figure 2** and **Videos 2-4**. Interactive 3D renderings of all cells are available at kernet.rwth-aachen.de. Furthermore, immersive visualization was used to provide unrivaled impressions of the spatial arrangements of the keratin network in stereoscopic 3D (**Figure 2-figure supplement 1**). The multidimensional visualizations revealed hallmark features of the different keratin filament networks.

Thus, an apical and a basal keratin filament network domain can be distinguished in MDCK cells (**Video 2**). The major difference between both domains are the thicker keratin bundles in the basal domain. Each subcellular network is characterized by a regular mesh pattern. The lateral keratin filament network is more heterogenous and dominated by filaments running between apical and basal. Circumferential subcortical keratin filaments are mainly restricted to the apical and basal network domains. In the cell interior, keratin filaments form a network around the nucleus. It is best developed on the basal side with highly bundled filaments, many of which are rather straight.

HaCaT cells are extremely flat and do not show a clearly separated apical and basal network and lack a distinct lateral network (**Video 3**). A prominent feature of their keratin filament network are long filament bundles that often run in parallel. They are absent above and below the nucleus but enclose the nucleus laterally.

The keratin network is less dense in RPE cells than in the other two cell types (**Video 4**). It is prominent in the cytoplasmatic apical domain, which faces the photoreceptor cells in the retina. It is also very well developed around the nucleus. In contrast to MDCK cells, the perinuclear network contains thicker filaments at the apical than the basal side. The basal cytoplasmatic domain is mostly devoid of keratin filaments. Some filaments run through the cytoplasm from basal to apical.

In conclusion, rendering of the keratin filament networks in three cell types revealed very different arrangements. The numerical data representation now allows detailed further analyses of distinct features. Especially with the help of VR, classification of network domains becomes feasible.

2.4 Comparison of keratin filament brightness distribution patterns

The numerical network representations were used to quantify keratin filament properties. A prominent property, which was already graphically encoded in the 3D-models, is filament brightness as a measure of filament thickness. The assumption is that filament thickness is a reliable indicator of filament bundling which can now be easily quantified throughout entire cells. We will therefore use the terms brightness and thickness interchangeably. The histograms in **Figure 3A-C** depict the spread of filament segment brightness in MDCK, HaCaT and RPE cells. The quantifications show that filament segments with a medium brightness dominated in MDCK cells with a near Gaussian distribution, whereas thin filaments were prevalent in HaCaT and RPE cells. Thin filaments were most abundant in HaCaT cells.

Maps of filament brightness are shown in **Figure 3D-F** for selected examples of each cell type at low and high magnification. In each case, thin filaments were numerous in the cell periphery. This may be due to the increased *de novo* single filament formation in these regions (Windoffer et al., 2004, 2006, Moch et al., 2013). Some areas, such as those shown at the right margin of the MDCK cell in **Figure 3D**, however, presented rather thick keratin filament bundles, which may represent matured interdesmosomal keratin filaments (Quinlan et al., 2017). The most heterogeneous network in terms

of filament thickness was encountered in MDCK cells. The animation in **Video 2** clearly shows that filaments in the basal network of the cell are thicker than that in the apical network. In HaCaT cells, enrichment of thin filaments was visible around the nucleus (**Figure 3E**) whereas thicker filaments were detectable in the cytoplasm, many of which appeared to be arranged in parallel (see also **Video 4**). In contrast, thick filament bundles surrounded the nucleus of RPE cells (**Figure 3F**; **Video 5**). Furthermore, thick bundles were more frequent above the nucleus, i.e., at the part of the cell facing the photoreceptors. Together, the analyses of the digitalized networks revealed major differences in the extent and distribution of keratin filament bundling suggesting that the keratin network is organized to withstand and cope with different types of mechanical stress.

2.5 Number and length of keratin filament segments and overall keratin filament length in single cells

The number and length of individual keratin filament segments are key measurements of network organization. Significantly more segments were detected in HaCaT cells than in the other cell types (**Figure 4A**). The average segment length was shortest in MDCK cells ($0.80 \pm 0.04 \mu\text{m}$), longest in RPE cells ($0.88 \pm 0.05 \mu\text{m}$) and intermediate in HaCaT cells ($0.83 \pm 0.13 \mu\text{m}$) (**Figure 4B**). The combined length of all segments per cell (**Figure 4C**) was similar in MDCK ($1.63 \pm 0.41 \text{ mm}$) and RPE cells ($1.57 \pm 0.48 \text{ mm}$) but was more than twice longer in HaCaT cells ($4.40 \pm 1.53 \text{ mm}$). To see whether this difference was due to different cell dimensions, the cell volumes were determined for all three cell types. Significant differences could be detected, i.e., $4.65 \pm 1.13 \mu\text{m}^3$ for MDCK, $9.16 \pm 6.10 \mu\text{m}^3$ for HaCaT and $2.73 \pm 0.95 \mu\text{m}^3$ for RPE (**Figure 4-figure supplement 1A Suppl. Figure 5A**). Taking the different cell volumes into account, filament density, i.e. filament length per volume, was comparable in HaCaT ($604.82 \pm 304.20 \mu\text{m}/\mu\text{m}^3$) and RPE cells ($585.60 \pm 63.97 \mu\text{m}/\mu\text{m}^3$), while it was considerably lower in MDCK cells ($355.24 \pm 56.49 \mu\text{m}/\mu\text{m}^3$) (**Figure 4D**). Given that the human body has $\sim 0.14 \times 10^{12}$ epidermal cells (Bianconi et al., 2013), the entire length of epidermal keratin filament bundles in an individual should be $\sim 600\,000 \text{ km}$. Assuming a bundling rate of 19 filaments per filament bundle (Hémonnot et al., 2016) there are $\sim 1.14 \times 10^7 \text{ km}$ of single keratin filaments in the epidermis of a human body.

2.6 Estimation of cellular keratin content

It is now possible to estimate the amount of keratin from our observations and to compare it to the previously estimated $8.07 \pm 1.88 \text{ pg}$ keratin in single basal keratinocytes of the murine epidermis (Feng et al., 2013). To this end, we take the experimentally determined mass per length (MPL) of keratin filaments of 25 kDa/nm (range: $19\text{--}30 \text{ kDa/nm}$; (Herrmann et al., 1999)) and a bundling factor (Bf) of 19 (Hémonnot et al., 2016) into account. As shown in the previous section (2.5), the total length of keratin filament bundles (L_{tot}) in HaCaT cells is $\sim 4.4 \text{ mm}$. By using the following formula ($\text{MPL} * L_{\text{tot}} * \text{Bf} / N_{\text{a}}$ [N_{a} = Avogadro constant]), the amount of keratin per cell is 3.47 pg .

An alternative estimate can be obtained by considering the dimensions and mass of the tetrameric subunit. A keratin tetramer has a length of 44 nm (Parry et al., 2007) and consists primarily of two keratin 5 and two keratin 14 polypeptides with a combined molecular weight of $228\,165 \text{ Da}$ in keratinocytes. Thus, 22.72 tetramers make up a $1 \mu\text{m}$ long protofilament. It has been previously shown that ~ 6 protofilaments are contained within a 10 nm keratin filament (Weber et al., 2021). Thus, a $1 \mu\text{m}$ -long keratin filament has a MW of $1.24\text{E}+08 \text{ Da}$. Assuming a bundling factor of 19 and a total keratin filament bundle length of $\sim 4.4 \text{ mm}$ in HaCaT keratinocytes, the total MW of keratin/cell is $2.60\text{E}+12 \text{ Da}$ corresponding to a mass of $\sim 4.3 \text{ pg/cell}$.

Performing the same type of calculation we deduced a keratin content of 1.3 pg or 1.4 pg per MDCK cell and 1.2 or 1.4 pg for single RPE cells.

2.7 Branching and intersection of keratin filaments

Next, we analyzed branching and intersection of keratin filament bundles. -Different types of branching could be envisioned depending on the number of segments per node. Three segments per node would define a Y-shaped branch, whereas four segments would define an X-shaped intersection of two filaments (**Figure 1-figure supplement 2C**). More than four branches would result in a star-like configuration. Our analyses show that the Y-shaped organization (three segments per node) was by far the most frequent configuration (~60%) followed by the X-shaped (four segment per node) arrangement (~30%) and rarely higher degrees of branching (~10%) (**Figure 5**).

2.8 Bending of keratin filaments

Bending of keratin intermediate filaments is of interest as it may provide information on material properties of filaments and may indicate whether filaments are under longitudinal compression or tension. First, we calculated the distance/length ratio of segments (**Figure 6A**). The highest ratios were observed in MDCK (0.92 ± 0.09) and HaCaT (0.92 ± 0.08), which were significantly higher than in RPE cells (0.90 ± 0.10) indicating that the filaments in RPE cells are less straight. This interpretation was confirmed by curvature determination of segments (**Figure 6B**). The mean curvature of segments was significantly higher in RPE cells ($3.30 \pm 0.15 \mu\text{m}^{-1}$) than in MDCK cells ($2.54 \pm 0.11 \mu\text{m}^{-1}$) and HaCaT cells ($2.97 \pm 0.72 \mu\text{m}^{-1}$). The curvature distribution in HaCaT cells was more widely spread than in the other two cell types. Maps of curvature distribution are shown from two different perspectives in **Figure 6C-E** revealing complex patterns. We did not see a correlation between filament bundle thickness and curvature (pairwise Pearson correlation coefficient in MDCK = -0.10, HaCaT = 0.08, RPE = 0.14). Finally, we determined the apparent persistence length of keratin filament segments being aware that the resulting values are difficult to interpret, since segments are not free to move but are connected to other segments within the network. **Figure 4-figure supplement 1B** shows that the calculated values (~2.6 μm) do not differ significantly between the three cell types. As expected, they are higher than the experimentally determined persistence length of single filaments (cf. (Block et al., 2015)).

2.9 Keratin filament segment orientation

To calculate how evenly the segments are orientated in space, the azimuth (= horizontal orientation) and the elevation (= vertical orientation) of segments were calculated for each cell (**Figure 7-figure supplements 1,2**). The histograms in **Figure 7A-F** show examples of the azimuth and elevation distribution in single MDCK, HaCaT and RPE cells. Comparing the determined values with the expected values for segments orientated evenly in all directions, shows that the azimuth distribution is quite even in MDCK and RPE cells but not in HaCaT cells. All cells have an uneven distribution of vertical orientation, indicating that segments are preferentially oriented horizontally. Compilations of the analyses for all cells (**Figure 7G, H**) confirmed the single cell data and affirmed that the azimuth distribution in HaCaT is more oriented than in MDCK or RPE cells. Furthermore, the elevation

orientation is more directed in MDCK and HaCaT cells than in RPE cells. Mapping the horizontal segment orientation in selected cells (**Figure 7I-L**) reveals a random pattern in MDCK and RPE cells and highlights parallel-arranged filaments in HaCaT cells.

To obtain insight about the relationship between segment directionality and cell shape, the orientation of the segments was analyzed with respect to the cell center. To this end, all segments were translocated to the cell center and their spatial distribution was plotted in radial 2D histograms. As expected, MDCK and RPE cells showed a homogenous distribution of horizontal segment orientation in the x/y plane while HaCaT cells were more polarized (**Figure 8A-C**). In the vertical direction (y/z plane) horizontally oriented segments predominate in MDCK and HaCaT cells while segments are more uniformly oriented in RPE cells (**Figure 8D-F**).

To compare the 3D-isotropy of entire keratin filament networks a sum vector of all segments was constructed for each cell. To this end, the orientation vectors of segments were translocated to the cell center as described above and mapped with the length of each segment. **Figure 8G-I** shows the resulting sum vectors in each plane together with corresponding views of the segmented keratin filament networks. The sum vectors in MDCK and RPE cells are relatively small, indicating an isotropic orientation of the respective keratin networks. In contrast, the sum vectors in HaCaT cells are much longer, indicating a higher degree of anisotropy of the keratin network (**Figure 8K**).

3 DISCUSSION

With this work we provide, for the first time, experimentally-derived numerical 3D representations of entire keratin networks in eukaryotic cells. 3D fluorescence data were used to define segments that are connected by nodes. This enabled us to compare local properties of keratin intermediate filament networks such as filament thickness, curvature and orientation in different cell types *in vitro* and *in vivo*. It further facilitated comprehensive comparisons of entire networks by quantifying parameters such as entire filament length, degree of filament connectivity and spatial network orientation.

A critical aspect of the work was evaluation of the quantification process. Since synthetic 3D data of cellular keratin networks were lacking, the quality of segmentation was evaluated by visual inspection of superimposed imaging and segmentation data demonstrating that the location and brightness of filaments were mapped with high precision and reliability. The subsequent allocation of nodes and segments was visually assessed in rendered 3D models (cinematic rendering) again revealing agreement of the assignments. Nevertheless, the numerical representation has its limitations in accuracy. Proper segment and node detection was particularly challenging in HaCaT cells because of their complex network geometry in combination with low filament bundling. Extreme bending at the single filament level with bending radii of 100 nm (Weber et al., 2021) are not detected in the present set-up due to the resolution limit of the airyscan. Mapping of keratin filaments at the cell edges was also difficult in some regions, mostly because of the manual assignment of cell borders in the confluent cell layers with tightly attached and partially overlapping neighboring cells. In cultured cells, mixing of fluorescent and non-fluorescent cells could overcome this problem. Segmentation quality may be further improved in the future by using other and more powerful super-resolution techniques than the airyscan technology used in the present work.

A major advance of the current report is the fabrication of numerical models of the keratin filament network in 3D. The rendered 3D models can be thoroughly inspected with the help of 3D display devices such as 3D screens, virtual reality-headsets (HMDs) or Cave Automatic Virtual Environments (CAVEs). The benefits of such immersive environments have been demonstrated

(Cruz-Neira et al., 1993; Laha et al., 2013). Three factors thereby improve the domain scientists' spatial judgment and thus the task performance during the visual analysis: (i) an enlarged field of regard compared to standard desktop systems reduces clutter during data exploration, (ii) stereoscopy provides supporting depth cues, and (iii) head tracking provides enhanced motion parallax. Motion parallax is a monocular depth cue which allows observers to perceive the depth of the visualized cell from their own motion. The filaments close to the observers are perceived to be moving faster than filaments in the distance.

The derived digital network models furthermore allow comparisons between cells with similar or dissimilar differentiation status (**Figure 9**). As a proof of concept, we paradigmatically investigated three different epithelial cell types finding significant differences in filament bundling, segment length, filament density, filament curvature and filament orientation. These parameters are a crucial and first step to build mechanical models of keratin filament networks in different cell types and functional contexts. The following list shows potential implications of the extracted features for our understanding of keratin filament network properties and functions:

- *Brightness of keratin filament segments*: Brightness levels correspond to filaments thickness and is therefore a measure of keratin filament bundling. The observed variation in filament bundling in the different cell types may therefore be a consequence of differences in the local force balance. This force balance is the result of intrinsic cell type-specific properties such as cell shape and arrangement of the actomyosin system and is impacted by forces imposed by forces imposed by neighboring cells and the extracellular matrix. This is supported by the observed differences in bundling of the apical, basal and perinuclear keratin filament networks. However, the variation in filament thickness may also reflect other properties such as the propensity of the network to reorganize, e.g., at different times of the cell cycle. Correlating the derived image data with information on mechanical keratin filament properties obtained by laser cutting experiments, force-sensor measurements or traction force analyses may help to better understand the relationship between filament bundling and mechanics.

- *Length of keratin filament segments*: Keratin filament segment length is an indirect measure of mesh size and may therefore describe another crucial mechanical property of the network. We were surprised to find that the segment length varied only by 10% between the different cell types and thus showed less variation than filament brightness or curvature. This means that the mesh size is fixed within a relatively small range, although the number of meshes per volume differed significantly between the different cell types. A conclusion of this observation is that mesh size may be determined by intrinsic keratin filament properties such as elastic modulus or branching capacity. However time-lapse fluorescence analyses (e.g., (Wöll et al., 2005)) have shown that small meshes can merge and enlarge by the combination of nodes from two segments, forming thicker filament bundles. Conversely, large meshes are subdivided into small meshes by recruitment of segments into a node. A self-regulatory mechanism may thus control the balance between enlargement and reduction of individual meshes.

- *Branching of keratin filament segments*: Experimental and mechanistic evidence has been provided for keratin filament branching (Beil et al., 2005; Feng and Coulombe, 2015; Lee et al., 2012; Martin et al., 2016; Nafeey et al., 2016). The fact that only two types of branching/intersections were detected in this study, points to a limited repertoire of connectivity. Furthermore, the Y- and X-shaped configurations may be fundamentally different. The Y-shaped joining of three segments may be linked to the recently proposed bundling mechanism involving electrostatic and hydrophobic interactions (Haimov et al., 2020). On the other hand, the X-shaped intersections may be related to

the disulfide bond mediated interactions reported for keratins 5/14 (Lee et al., 2012). In this instance, preventing these links resulted in altered keratin filament network dynamics and epidermal tissue homeostasis (Feng and Coulombe, 2015; Guo et al., 2020).

- Curvature of keratin filament segments: Bending of keratin filaments provides important information on whether keratins are stretched, relaxed or compressed. Keratin intermediate filaments are highly elastic (Kreplak et al., 2005; Wagner et al., 2007; Yamada et al., 2003). They can be extended under tension forming straight filaments and buckle under compression (= negative tension) (Fudge et al., 2008). The measurements of segment curvature can therefore be taken as indicators of local keratin filament tension. Straight segments were almost completely absent in all cells suggesting overall negative tension of the keratin networks. The observation that filament bundle thickness and curvature are not correlated suggests that the 10 nm filaments are only loosely connected in keratin bundles. The degree of filament bending, however, differed between the different cell types indicating cell type-specific differences in intrinsic force balance and may suggest that the different networks retain different degrees of flexibility before they become stretched to function as mechanical counterbalances to strain. Curvature of keratin filaments also allowed to estimate the persistence length of network-embedded keratin filaments, i.e. the minimum length between two points on the polymer where they become independent of one another. The value of ~2 µm was only slightly higher than what was reported for other intermediate filaments including keratin filaments (Block et al., 2015; Wagner et al., 2007). This is low in comparison to actin filaments (~10 µm) (De La Cruz and Gardel, 2015; Isambert et al., 1995; Takatsuki et al., 2014) and microtubules (~5000 µm) (Hawkins et al., 2010). This result emphasizes the maintained high flexibility of keratin filaments despite their connectivity through nodes.

- Orientation of keratin filaments: To understand the response of the keratin network to internal and external forces, it is important to have measures of filament orientation. We suggest that uniformly radial orientation of keratin filament segments indicates that cells experience mechanical forces that are homogenous in all directions. In contrast, a preferential orientation in one direction indicates that keratin filaments align in response to mechanical cues that may be induced by unevenly distributed cell-cell contacts, polarized cell-extracellular matrix adhesions or contractile activity. Our data therefore indicate that HaCaT cells are subjected to mechanical strain (uneven azimuth distribution and large sum vector), whereas MDCK and RPE are subjected to little mechanical strain (uniform azimuth distribution and small sum vector). Fine mapping of the distribution patterns of hemidesmosomes and desmosomes may help to understand the differences and to initiate mechanical modelling, improving previous models of keratin network, morphogenesis, dynamics and mechanics (Beil et al., 2009; Dallon et al., 2019; Latorre et al., 2018; Portet, 2013; Portet et al., 2019, 2015).

The observed high degree of segment property variability must be taken as an indication of the amazing plasticity of the keratin filament system. –The node/segment arrangement revealed a unique 3D structure that can be modified in multiple ways affecting bundling, flexibility and local cross-talk. Keratin isotypes, posttranslational modifications, external and internal forces, and interactions with other cytoskeletal and cellular components may determine these properties and thereby act as modulators of network structure at different length and time scales.

4 MATERIALS AND METHODS

4.1 Cell culture

Madin-Darby Canine Kidney (MDCK) cells were stably transfected with a plasmid encoding human full-length keratin 8 with an EYFP tag at its carboxyterminus (Windoffer et al., 2004; recloned into

pEYFP-N1 (Clontech) with BamHI and EcoRI). MDCK cells were obtained from DKFZ (Deutsches Krebsforschungszentrum). Resulting cell clone H9 was used for the analyses described in this paper. Cells were grown in Dulbecco's Modified Eagle Medium (DMEM, Sigma-Aldrich) with 10% (v/v) fetal calf serum (Pan Biotech) including 700 µg/ml G418 (Sigma-Aldrich) for selection. Vital MDCK cells were imaged after 3 days when the monolayer had reached complete confluence.

Immortalized human HaCaT keratinocyte cell clone B10 expressing EYFP-tagged human keratin 5 (HK5-EYFP) has been described (Moch et al., 2013). HaCaT cells were obtained from the Fusenig lab, which first isolated and described this cell line (Boukamp et al., 1988). The cells were grown in DMEM containing l-alanyl-glutamine (Sigma-Aldrich) supplemented with 10% (v/v) fetal calf serum (SeraPlus; PAN Biotech) and passaged as described (Moch et al., 2013). Vital HaCaT cells were imaged 1-2 days after reaching confluence corresponding to 5-7 after seeding. Both cell lines were tested by PCR test to be mycoplasma free

4.2 Preparation of retinal pigment epithelium

Retinas were prepared from homozygous keratin 8-YFP knock-in mice, which synthesize fluorescence-tagged keratin 8 from the modified endogenous gene locus (Schwarz et al., 2015). Mice were sacrificed by cervical dislocation and eyes were enucleated using curved forceps. Eyeballs were then fixed with 2% paraformaldehyde in rotation at 4°C for 4 h. For retina flat-mount preparation, the eye balls were dissected as previously described (Clayton and Bishop, 2011). Briefly, cornea and lens were removed and the eyecup was cut into a multileaved flower shape. The neural retina was removed and the remaining tissue was mounted on a microscope slide with Mowiol (Sigma-Aldrich).

4.3 Image recording

To obtain statistically relevant data, ten image data sets were recorded of each cell type with a Zeiss LSM710 using airyscan mode (Carl Zeiss, Jena, Germany). An oil immersion Plan-Apochromat 63x/1.4-NA, DIC M27 objective (Carl Zeiss) was used. Fluorescence was detected with an 488 nm argon laser (module LGK 7872 ML8) and filter set BP 495-550 using 2.5% power. Optimal factory settings for z-stacks were used resulting in a voxel size of $xy = 0.044$ to 0.065 µm and $z = 0.17$ to 0.2 µm. No further image optimization was used. To assure that only single cells were analyzed, the fluorescence in individual cells was cropped from confluent MDCK monolayers and retinal pigment epithelium (RPE) using Fiji standard tools. Cell borders were manually defined for each z-slice and the cell exterior was set to a grey value of 0. Wild-type HaCaT and HK5-EYFP producing HaCaT clone B10 cells were mixed prior to imaging to prevent overlap of fluorescence signals in neighbouring cells.

4.4 Segmentation of image stacks

TSOAX software (Xu et al., 2019, 2015, 2011) was used for the segmentation of filaments from the recorded 3D image stacks (**Figure 1-figure supplement 2A**). TSOAX facilitates quantitative analysis of network dynamics of multi-dimensional biopolymer networks imaged by various microscopic imaging modalities. The underlying methods of TSOAX include multiple Stretching Open Active Contour (SOAC) models for extraction and a combined local and global graph-matching framework (Li et al., 2010, 2009b, 2009a). For optimal results, the default TSOAX parameters were used, with the exception of 'external-factor' which was set to 2 based on internal testing. Since computation time can be extremely long, TSOAX was executed in parallel instances on the supercomputer CLAIX at RWTH Compute Cluster (<https://www.its.rwth-aachen.de/go/id/eucm>).

474

475 4.5 Calculation of a segment/node-based network model

476 To get a perfect network representation it is necessary to divide the network into defined pieces (= segments) that do not cross each other, and can be connected at their ends by nodes. Segmentations using TSOAX defines polygonal chains with n vertices (= snakes). Each vertex is defined in x, y, z and has an associated density (= fluorescence intensity). However, the snakes are not associated with each other via nodes and may cross each other. (**Figure 1-figure supplement 2B**). To obtain a comprehensive and coherent numerical network definition from snakes, KerNet software was developed. It generates a graph representation consisting of segments (~ edges) and nodes. To compute such a connected network representation, vertices of a snake that lie within $\epsilon = 1.1 \times$ voxelsize distance of one another are combined to form a node. If more than two vertices are within ϵ distance, a candidate set is created. Any vertex, which is at most ϵ away from a vertex in the candidate set, is added to the set until this is no longer possible. Then all vertices in the candidate set are combined to form a single node. Finally, any start or end vertex of a segment is a node. If two nodes are connected by a part of a snake then a segment connects the two nodes in the network. In contrast to classical edges, which are defined as lines connecting the nodes, each segment is a list of vertices (x,y,z) and their associated densities. The segments therefore inform about the trajectory of the snake between its two endpoints and the density progression along the segment. To summarize, nodes represent the first and last vertex of a segment and mark the connection between segments whose endpoints are close (**Figure 1-figure supplement 2C**). All segments and nodes together represent a Euclidean 3D network map of a given cell.

495 4.6 Validation of segmentation

496 For validation of the segmentation, the original fluorescence image stacks were compared with the segmented 3D data. All vertices retrieved from KerNet were plotted into the original image stack. 497 The z-values of the vertices were rounded to the nearest z-slice and all segment vertices were drawn 498 as discs with a diameter proportional to their brightness. To simulate the blurriness of the raw 499 images, the vertex stacks were 3D-Gauss filtered using Fiji (values 5,5,3). 500

501 4.7 Cinematic rendering of segments

502 For inspection of the 3D network representation, the xyz-position data of segments were exported 503 into OBJ files (<http://www.martinreddy.net/gfx/3d/OBJ.spec>), which, in turn, can be imported into 504 3D rendering programs. For the representation of segment brightness as renderable tubes, density 505 properties of segments were compiled into 3ds Max (Autodesk) script files that translate segment 506 brightness into tube thickness, resulting in renderable splines for each segment. In some instances, 507 random colors were assigned to each segment (**Figure 1-figure supplement 2D**). Finally, 3ds Max 508 rendering was used for the generation of stills and videos.

509 4.8 Virtual reality

510 For further visual inspection of the keratin networks, a basic Virtual Reality (VR) application was 511 designed. The VR application was implemented using Unreal Engine 4 512 (<https://www.unrealengine.com/en-US/>). It displays the datasets of cells as 3D models. The 513 individual filaments are visualized as tubes. Their brightness is encoded in the tubes' thickness. The 514 resulting structure of interconnected tubes can then be visually inspected in VR, allowing a detailed 515 view on the digitalized subcellular network in any magnification. Individual filaments can be

interactively assigned to one of several distinct classes, pre-defined by the domain expert. Color-coding is then used to visually differentiate the classes in the immersive visualization.

During our research, we used the HTC Vive Pro headset to manually classify the filaments of a cell. Afterwards, the aixCAVE at RWTH Aachen University was used to collaboratively discuss and adjust the classification.

4.9 Numerical analysis and visualization

Matlab routines were used for further analysis of network properties. Up to 150 000 vertices may be present per cell. Segment length, distance between nodes, bending, curvature and apparent persistence length of segments were calculated. Bending was defined as the ratio between the shortest distance between the segment endpoints and the actual segment length. The curvature was calculated as the 2nd derivative of the smoothed tangent vector with respect to the arc length (Sternberg, 2012). Since persistence length can only be calculated in free moving segments and the segments are restricted in their movement by the nodes in our situation, we calculated an apparent persistence length as described (Doi and Edwards, 1990).

Statistical analyses were performed with Matlab and GraphPad Prism software. For non-Gaussian distribution, Mann-Whitney test was used. Values are expressed as the mean $\pm$ SD. * represents a p-value of $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$. The differences were considered significant when $p < 0.05$. Non-significant differences were indicated by n.s.

To visualize segment properties such as length or curvature, the respective values were plotted and color-coded at the vertex location resulting in heat maps showing the distribution of the selected property. Here, Fiji scripts were used to plot color-coded properties into xyz coordinates of vertices.

For calculation of cell size, a threshold was used, that separated the background fluorescence inside and outside of cells.

4.10 Directional vectors

The directionality of segments was calculated for the keratin network of single cells in two ways :

(i) Since the start and end points of segments are interchangeable, segments with an orientation difference of 180 degrees are equally orientated. Normalized histograms were plotted for azimuth and elevation from 0-180 degrees. Next, we calculated the deviation of the histogram from a histogram with an even distribution of orientations where each degree has the same frequency. Finally, the difference between the segment histogram and the evenly distributed histogram was calculated as the sum of the differences in each bin.

(ii) Keeping the segments in their original orientation we analyzed the position of segments in relation to the cell center. The cell center was defined as the mean of the mean positions of all segments in xyz. It was then determined whether the start point or end point of each segment was closer to the cell center. The segment was then moved to the cell center such that the closer point was placed to the cell center. In this way, all segments originate at the cell center and direct outwards. The resulting radial histograms were plotted to depict the distribution of segments in xy and yz directions.

554 Treating each segment as a vector extending from the cell center to the segment's end, the sum vector
555 of all segments was calculated and plotted.

556 4.11 Software, original and processed data

557 Software, original and processed data are available at <http://kernet.rwth-aachen.de/> and
558 <https://github.com/VRGroupRWTH/Zytoskelett>.

559

560 5 Citations

561

562 Acciai L, Soda P, Iannello G. 2016. Automated Neuron Tracing Methods: An Updated Account.
563 *Neuroinform* **14**:353–367. doi:10.1007/s12021-016-9310-0

564 Alioscha-Perez M, Benadiba C, Goossens K, Kasas S, Dietler G, Willaert R, Sahli H. 2016. A Robust
565 Actin Filaments Image Analysis Framework. *PLOS Computational Biology* **12**:e1005063.
566 doi:10.1371/journal.pcbi.1005063

567 Beil M, Braxmeier H, Fleischer F, Schmidt V, Walther P. 2005. Quantitative analysis of keratin
568 filament networks in scanning electron microscopy images of cancer cells. *Journal of*
569 *Microscopy* **220**:84–95. doi:10.1111/j.1365-2818.2005.01505.x

570 Beil M, Lück S, Fleischer F, Portet S, Arendt W, Schmidt V. 2009. Simulating the formation of
571 keratin filament networks by a piecewise-deterministic Markov process. *Journal of*
572 *Theoretical Biology* **256**:518–532. doi:10.1016/j.jtbi.2008.09.044

573 Beriault DR, Haddad O, McCuaig JV, Robinson ZJ, Russell D, Lane EB, Fudge DS. 2012. The
574 Mechanical Behavior of Mutant K14-R125P Keratin Bundles and Networks in NEB-1
575 Keratinocytes. *PLoS One* **7**:e31320. doi:10.1371/journal.pone.0031320

576 Bianconi E, Piovesan A, Facchin F, Beraudi A, Casadei R, Frabetti F, Vitale L, Pelleri MC, Tassani
577 S, Piva F, Perez-Amodio S, Strippoli P, Canaider S. 2013. An estimation of the number of
578 cells in the human body. *Annals of Human Biology* **40**:463–471.
579 doi:10.3109/03014460.2013.807878

580 Block J, Schroeder V, Pawelzyk P, Willenbacher N, Köster S. 2015. Physical properties of
581 cytoplasmic intermediate filaments. *Biochim Biophys Acta* **1853**:3053–3064.
582 doi:10.1016/j.bbamcr.2015.05.009

583 Boukamp P, Petrussevska RT, Breitkreutz D, Hornung J, Markham A, Fusenig NE. 1988. Normal
584 keratinization in a spontaneously immortalized aneuploid human keratinocyte cell line.
585 *Journal of Cell Biology* **106**:761–771. doi:10.1083/jcb.106.3.761

586 Claybon A, Bishop AJR. 2011. Dissection of a Mouse Eye for a Whole Mount of the Retinal Pigment
587 Epithelium. *JoVE (Journal of Visualized Experiments)* e2563. doi:10.3791/2563

588 Coch RA, Leube RE. 2016. Intermediate Filaments and Polarization in the Intestinal Epithelium.
589 *Cells* **5**:32. doi:10.3390/cells5030032

590 Cruz-Neira C, Sandin DJ, DeFanti TA. 1993. Surround-Screen Projection-Based Virtual Reality: The
591 Design and Implementation of the CAVEIn_ Proc. 20th Annu. Conf. Comput. Graph.
592 Interactive Techn. pp. 135–142.

- 593 Dallon JC, Leduc C, Etienne-Manneville S, Portet S. 2019. Stochastic modeling reveals how motor
594 protein and filament properties affect intermediate filament transport. *Journal of Theoretical*
595 *Biology* **464**:132–148. doi:10.1016/j.jtbi.2018.12.022
- 596 De La Cruz EM, Gardel ML. 2015. Actin Mechanics and Fragmentation. *J Biol Chem* **290**:17137–
597 17144. doi:10.1074/jbc.R115.636472
- 598 Dimchev G, Amiri B, Fäßler F, Falcke M, Schur FK. 2021. Computational toolbox for ultrastructural
599 quantitative analysis of filament networks in cryo-ET data. *bioRxiv* 2021.05.25.445599.
600 doi:10.1101/2021.05.25.445599
- 601 Doi M, Edwards SF. 1990. The Theory of Polymer Dynamics, International Series of Monographs on
602 Physics. Oxford Science Publications.
- 603 Faulkner C, Zhou J, Evrard A, Bourdais G, MacLean D, Häweker H, Eckes P, Robatzek S. 2017. An
604 automated quantitative image analysis tool for the identification of microtubule patterns in
605 plants. *Traffic* **18**:683–693. doi:10.1111/tra.12505
- 606 Felder E, Siebenbrunner M, Busch T, Fois G, Miklavc P, Walther P, Dietl P. 2008. Mechanical strain
607 of alveolar type II cells in culture: changes in the transcellular cytokeratin network and
608 adaptations. *Am J Physiol Lung Cell Mol Physiol* **295**:L849–L857.
609 doi:10.1152/ajplung.00503.2007
- 610 Feng X, Coulombe PA. 2015. A role for disulfide bonding in keratin intermediate filament
611 organization and dynamics in skin keratinocytes. *J Cell Biol* **209**:59–72.
612 doi:10.1083/jcb.201408079
- 613 Feng X, Zhang H, Margolick JB, Coulombe PA. 2013. Keratin intracellular concentration revisited:
614 Implications for keratin function in surface epithelia. *J Invest Dermatol* **133**:850–853.
615 doi:10.1038/jid.2012.397
- 616 Flormann DAD, Schu M, Terriac E, Thalla D, Kainka L, Koch M, Gad AKB, Lautenschläger F.
617 2021. A novel universal algorithm for filament network tracing and cytoskeleton analysis.
618 *FASEB J* **35**:e21582. doi:10.1096/fj.202100048R
- 619 Fois G, Weimer M, Busch T, Felder ET, Oswald F, Wichert G von, Seufferlein T, Dietl P, Felder E.
620 2013. Effects of keratin phosphorylation on the mechanical properties of keratin filaments in
621 living cells. *The FASEB Journal* **27**:1322–1329. doi:10.1096/fj.12-215632
- 622 Fudge D, Russell D, Beriault D, Moore W, Lane EB, Vogl AW. 2008. The Intermediate Filament
623 Network in Cultured Human Keratinocytes Is Remarkably Extensible and Resilient. *PLOS*
624 *ONE* **3**:e2327. doi:10.1371/journal.pone.0002327
- 625 Guo Y, Redmond CJ, Leacock KA, Brovkina MV, Ji S, Jaskula-Ranga V, Coulombe PA. 2020.
626 Keratin 14-dependent disulfides regulate epidermal homeostasis and barrier function via 14-
627 3-3σ and YAP1. *eLife* **9**:e53165. doi:10.7554/eLife.53165
- 628 Haspinger DC, Klinge S, Holzapfel GA. 2021. Numerical analysis of the impact of cytoskeletal actin
629 filament density alterations onto the diffusive vesicle-mediated cell transport. *PLOS*
630 *Computational Biology* **17**:e1008784. doi:10.1371/journal.pcbi.1008784
- 631 Hatzfeld M, Keil R, Magin TM. 2017. Desmosomes and Intermediate Filaments: Their
632 Consequences for Tissue Mechanics. *Cold Spring Harb Perspect Biol* **9**:a029157.
633 doi:10.1101/cshperspect.a029157

- 634 Hawkins T, Mirigian M, Selcuk Yasar M, Ross JL. 2010. Mechanics of microtubules. *Journal of*
635 *Biomechanics*, Special Issue on Cell Mechanobiology **43**:23–30.
636 doi:10.1016/j.jbiomech.2009.09.005
- 637 Hémonnot CYJ, Reinhardt J, Saldanha O, Patommel J, Graceffa R, Weinhausen B, Burghammer M,
638 Schroer CG, Köster S. 2016a. X-rays Reveal the Internal Structure of Keratin Bundles in
639 Whole Cells. *ACS Nano* **10**:3553–3561. doi:10.1021/acsnano.5b07871
- 640 Hémonnot CYJ, Reinhardt J, Saldanha O, Patommel J, Graceffa R, Weinhausen B, Burghammer M,
641 Schroer CG, Köster S. 2016b. X-rays Reveal the Internal Structure of Keratin Bundles in
642 Whole Cells. *ACS Nano* **10**:3553–3561. doi:10.1021/acsnano.5b07871
- 643 Herrmann H, Häner M, Brettel M, Ku N-O, Aebi U. 1999. Characterization of distinct early assembly
644 units of different intermediate filament proteins1 Edited by W. Baumeister. *Journal of*
645 *Molecular Biology* **286**:1403–1420. doi:10.1006/jmbi.1999.2528
- 646 Howard J. 2014. Quantitative cell biology: the essential role of theory. *MBoC* **25**:3438–3440.
647 doi:10.1091/mbc.e14-02-0715
- 648 Imran A, Li J, Pei Y, Yang J-J, Wang Q. 2019. Comparative Analysis of Vessel Segmentation
649 Techniques in Retinal Images. *IEEE Access* **7**:114862–114887.
650 doi:10.1109/ACCESS.2019.2935912
- 651 Isambert H, Venier P, Maggs AC, Fattoum A, Kassab R, Pantaloni D, Carlier M-F. 1995. Flexibility
652 of Actin Filaments Derived from Thermal Fluctuations: EFFECT OF BOUND
653 NUCLEOTIDE, PHALLOIDIN, AND MUSCLE REGULATORY PROTEINS*. *Journal of*
654 *Biological Chemistry* **270**:11437–11444. doi:10.1074/jbc.270.19.11437
- 655 Iwatsuki H, Suda M. 2010. Seven Kinds of Intermediate Filament Networks in the Cytoplasm of
656 Polarized Cells: Structure and Function. *Acta Histochemica Et Cytochemica* **43**:19–31.
657 doi:10.1267/ahc.10009
- 658 Jacob JT, Coulombe PA, Kwan R, Omary MB. 2018. Types I and II Keratin Intermediate Filaments.
659 *Cold Spring Harb Perspect Biol* **10**:a018275. doi:10.1101/cshperspect.a018275
- 660 Karsch S, Büchau F, Magin TM, Janshoff A. 2020. An intact keratin network is crucial for
661 mechanical integrity and barrier function in keratinocyte cell sheets. *Cell Mol Life Sci*
662 **77**:4397–4411. doi:10.1007/s00018-019-03424-7
- 663 Kim Y-B, Hlavaty D, Maycock J, Lechler T. 2021. Roles for Ndel1 in keratin organization and
664 desmosome function. *MBoC* mbc.E21-02-0087. doi:10.1091/mbc.E21-02-0087
- 665 Kothari P, Johnson C, Sandone C, Iglesias PA, Robinson DN. 2019. How the mechanobiome drives
666 cell behavior, viewed through the lens of control theory. *J Cell Sci* **132**:jcs234476.
667 doi:10.1242/jcs.234476
- 668 Kreplak L, Bär H, Leterrier JF, Herrmann H, Aebi U. 2005. Exploring the Mechanical Behavior of
669 Single Intermediate Filaments. *Journal of Molecular Biology* **354**:569–577.
670 doi:10.1016/j.jmb.2005.09.092
- 671 Laha B, Bowman DA, Schiffbauer JD. 2013. Validation of the MR Simulation Approach for
672 Evaluating the Effects of Immersion on Visual Analysis of Volume Data. *IEEE Transactions*
673 *on Visualization and Computer Graphics* **19**:529–538. doi:10.1109/TVCG.2013.43
- 674 Latorre E, Kale S, Casares L, Gómez-González M, Uroz M, Valon L, Nair RV, Garreta E, Montserrat
675 N, del Campo A, Ladoux B, Arroyo M, Trepas X. 2018. Active superelasticity in three-

- 676 dimensional epithelia of controlled shape. *Nature* **563**:203–208. doi:10.1038/s41586-018-
677 0671-4
- 678 Lee C-H, Kim M-S, Chung BM, Leahy DJ, Coulombe PA. 2012. Structural basis for heteromeric
679 assembly and perinuclear organization of keratin filaments. *Nat Struct Mol Biol*
680 **19**:10.1038/nsmb.2330. doi:10.1038/nsmb.2330
- 681 Leube RE, Moch M, Windoffer R. 2017. Intracellular Motility of Intermediate Filaments. *Cold*
682 *Spring Harb Perspect Biol* **9**:a021980. doi:10.1101/cshperspect.a021980
- 683 Li H, Shen T, Smith MB, Fujiwara I, Vavylonis D, Huang X. 2009a. Automated actin filament
684 segmentation, tracking and tip elongation measurements based on open active contour
685 models2009 IEEE International Symposium on Biomedical Imaging: From Nano to Macro.
686 Presented at the 2009 IEEE International Symposium on Biomedical Imaging: From Nano to
687 Macro. pp. 1302–1305. doi:10.1109/ISBI.2009.5193303
- 688 Li H, Shen T, Vavylonis D, Huang X. 2010. Actin Filament Segmentation Using Spatiotemporal
689 Active-Surface and Active-Contour Models In: Jiang T, Navab N, Pluim JPW, Viergever
690 MA, editors. Medical Image Computing and Computer-Assisted Intervention – MICCAI
691 2010, Lecture Notes in Computer Science. Berlin, Heidelberg: Springer. pp. 86–94.
692 doi:10.1007/978-3-642-15705-9_11
- 693 Li H, Shen T, Vavylonis D, Huang X. 2009b. Actin Filament Tracking Based on Particle Filters and
694 Stretching Open Active Contour Models In: Yang G-Z, Hawkes D, Rueckert D, Noble A,
695 Taylor C, editors. Medical Image Computing and Computer-Assisted Intervention – MICCAI
696 2009, Lecture Notes in Computer Science. Berlin, Heidelberg: Springer. pp. 673–681.
697 doi:10.1007/978-3-642-04271-3_82
- 698 Liberali P, Pelkmans L. 2012. Towards quantitative cell biology. *Nat Cell Biol* **14**:1233–1233.
699 doi:10.1038/ncb2648
- 700 Lidke DS. 2017. Quantitative cell biology: uniting disciplines to understand the cell. *Mol Biol Cell*
701 **28**:3133. doi:10.1091/mbc.E17-09-0551
- 702 Lippincott-Schwartz J. 2014. Quantitative cell biology: transforming the conceptual, theoretical,
703 instrumental, and methodological approaches to cell biology. *Mol Biol Cell* **25**:3437.
704 doi:10.1091/mbc.E14-08-1297
- 705 Liu Y, Kolagunda A, Treible W, Nedo A, Caplan J, Kambhamettu C. 2019. Intersection to Overpass:
706 Instance Segmentation on Filamentous Structures With an Orientation-Aware Neural
707 Network and Terminus Pairing Algorithm2019 IEEE/CVF Conference on Computer Vision
708 and Pattern Recognition Workshops (CVPRW). Presented at the 2019 IEEE/CVF Conference
709 on Computer Vision and Pattern Recognition Workshops (CVPRW). pp. 125–133.
710 doi:10.1109/CVPRW.2019.00021
- 711 Lutz A, Jung D, Diem K, Fauler M, Port F, Gottschalk K, Felder E. 2020. Acute effects of cell stretch
712 on keratin filaments in A549 lung cells. *The FASEB Journal* **34**:11227–11242.
713 doi:https://doi.org/10.1096/fj.201903160RR
- 714 Magliaro C, Callara AL, Vanello N, Ahluwalia A. 2019. Gotta Trace ‘em All: A Mini-Review on
715 Tools and Procedures for Segmenting Single Neurons Toward Deciphering the Structural
716 Connectome. *Front Bioeng Biotechnol* **0**. doi:10.3389/fbioe.2019.00202

- 717 Martin I, Moch M, Neckernuss T, Paschke S, Herrmann H, Marti O. 2016. Both monovalent cations
718 and plectin are potent modulators of mechanical properties of keratin K8/K18 networks. *Soft*
719 *Matter* **12**:6964–6974. doi:10.1039/C6SM00977H
- 720 Moccia S, De Momi E, El Hadji S, Mattos LS. 2018. Blood vessel segmentation algorithms —
721 Review of methods, datasets and evaluation metrics. *Computer Methods and Programs in*
722 *Biomedicine* **158**:71–91. doi:10.1016/j.cmpb.2018.02.001
- 723 Moch M, Herberich G, Aach T, Leube RE, Windoffer R. 2013. Measuring the regulation of keratin
724 filament network dynamics. *Proc Natl Acad Sci U S A* **110**:10664–10669.
725 doi:10.1073/pnas.1306020110
- 726 Moch M, Schwarz N, Windoffer R, Leube RE. 2020. The keratin–desmosome scaffold: pivotal role
727 of desmosomes for keratin network morphogenesis. *Cell Mol Life Sci* **77**:543–558.
728 doi:10.1007/s00018-019-03198-y
- 729 Moch M, Windoffer R, Schwarz N, Pohl R, Omenzetter A, Schnakenberg U, Herb F, Chaisaowong
730 K, Merhof D, Ramms L, Fabris G, Hoffmann B, Merkel R, Leube RE. 2016. Effects of
731 Plectin Depletion on Keratin Network Dynamics and Organization. *PLoS One* **11**:e0149106.
732 doi:10.1371/journal.pone.0149106
- 733 Nafeey S, Martin I, Felder T, Walther P, Felder E. 2016. Branching of keratin intermediate filaments.
734 *Journal of Structural Biology* **194**:415–422. doi:10.1016/j.jsb.2016.03.023
- 735 Nowak J, Gennermann K, Persson S, Nikoloski Z. 2020. CytoSeg 2.0: automated extraction of actin
736 filaments. *Bioinformatics* **36**:2950–2951. doi:10.1093/bioinformatics/btaa035
- 737 Osmanagic-Myers S, Gregor M, Walko G, Burgstaller G, Reipert S, Wiche G. 2006. Plectin-
738 controlled keratin cytoarchitecture affects MAP kinases involved in cellular stress response
739 and migration. *J Cell Biol* **174**:557–568. doi:10.1083/jcb.200605172
- 740 Özdemir B, Reski R. 2021. Automated and semi-automated enhancement, segmentation and tracing
741 of cytoskeletal networks in microscopic images: A review. *Computational and Structural*
742 *Biotechnology Journal* **19**:2106–2120. doi:10.1016/j.csbj.2021.04.019
- 743 Parajón E, Surcel A, Robinson DN. 2021. The mechanobiome: a goldmine for cancer therapeutics.
744 *American Journal of Physiology-Cell Physiology* **320**:C306–C323.
745 doi:10.1152/ajpcell.00409.2020
- 746 Parry DAD, Strelkov SV, Burkhard P, Aebl U, Herrmann H. 2007. Towards a molecular description
747 of intermediate filament structure and assembly. *Experimental Cell Research, Special Issue -*
748 *Intermediate Filaments* **313**:2204–2216. doi:10.1016/j.yexcr.2007.04.009
- 749 Piston DW. 2014. Biophysical Journal Special Issue: Focus on Quantitative Cell Biology. *Biophys J*
750 **107**:E01. doi:10.1016/j.bpj.2014.10.038
- 751 Portet S. 2013. Dynamics of in vitro intermediate filament length distributions. *Journal of*
752 *Theoretical Biology* **332**:20–29. doi:10.1016/j.jtbi.2013.04.004
- 753 Portet S, Leduc C, Etienne-Manneville S, Dallon J. 2019. Deciphering the transport of elastic
754 filaments by antagonistic motor proteins. *Phys Rev E* **99**:042414.
755 doi:10.1103/PhysRevE.99.042414
- 756 Portet S, Madzvamuse A, Chung A, Leube RE, Windoffer R. 2015. Keratin Dynamics: Modeling the
757 Interplay between Turnover and Transport. *PLoS One* **10**:e0121090.
758 doi:10.1371/journal.pone.0121090

- 759 Quan T, Zhou H, Li J, Li S, Li A, Li Y, Lv X, Luo Q, Gong H, Zeng S. 2016. NeuroGPS-Tree:
760 automatic reconstruction of large-scale neuronal populations with dense neurites. *Nature*
761 *Methods* **13**:51–54. doi:10.1038/nmeth.3662
- 762 Quinlan RA, Schwarz N, Windoffer R, Richardson C, Hawkins T, Broussard JA, Green KJ, Leube
763 RE. 2017. A rim-and-spoke hypothesis to explain the biomechanical roles for cytoplasmic
764 intermediate filament networks. *J Cell Sci* **130**:3437–3445. doi:10.1242/jcs.202168
- 765 Roca-Cusachs P, Conte V, Trepas X. 2017. Quantifying forces in cell biology. *Nat Cell Biol* **19**:742–
766 751. doi:10.1038/ncb3564
- 767 Samuel PM, Veeramalai T. 2020. Review on retinal blood vessel segmentation - an algorithmic
768 perspective. *International Journal of Biomedical Engineering and Technology* **34**:75–105.
769 doi:10.1504/IJBET.2020.110362
- 770 Schwarz N, Windoffer R, Magin TM, Leube RE. 2015. Dissection of keratin network formation,
771 turnover and reorganization in living murine embryos. *Scientific Reports* **5**:9007.
772 doi:10.1038/srep09007
- 773 Sternberg S. 2012. Curvature in Mathematics and Physics, 1st ed. Dover Publications, Inc.
- 774 Strnad P, Stumppner C, Zatloukal K, Denk H. 2008. Intermediate filament cytoskeleton of the liver in
775 health and disease. *Histochem Cell Biol* **129**:735–749. doi:10.1007/s00418-008-0431-x
- 776 Takatsuki H, Bengtsson E, Månsson A. 2014. Persistence length of fascin-cross-linked actin filament
777 bundles in solution and the in vitro motility assay. *Biochimica et Biophysica Acta (BBA) -*
778 *General Subjects* **1840**:1933–1942. doi:10.1016/j.bbagen.2014.01.012
- 779 Tateishi K, Nishida T, Inoue K, Tsukita S. 2017. Three-dimensional Organization of Layered Apical
780 Cytoskeletal Networks Associated with Mouse Airway Tissue Development. *Sci Rep*
781 **7**:43783. doi:10.1038/srep43783
- 782 te Molder L, Kreft M, Heemskerk N, Schuring J, de Pereda JM, Wilhelmsen K, Sonnenberg A. 2021.
783 EGFR-dependent tyrosine phosphorylation of integrin β 4 is not required for downstream
784 signaling events in cancer cell lines. *Sci Rep* **11**:8675. doi:10.1038/s41598-021-88134-6
- 785 Wagner OI, Rammensee S, Korde N, Wen Q, Leterrier J-F, Janmey PA. 2007. Softness, strength and
786 self-repair in intermediate filament networks. *Exp Cell Res* **313**:2228–2235.
787 doi:10.1016/j.yexcr.2007.04.025
- 788 Wang J, Lü D, Mao D, Long M. 2014. Mechanomics: an emerging field between biology and
789 biomechanics. *Protein Cell* **5**:518–531. doi:10.1007/s13238-014-0057-9
- 790 Weber MS, Eibauer M, Sivagurunathan S, Magin TM, Goldman RD, Medalia O. 2021. Structural
791 heterogeneity of cellular K5/K14 filaments as revealed by cryo-electron microscopy. *eLife*
792 **10**:e70307. doi:10.7554/eLife.70307
- 793 Wöll S, Windoffer R, Leube RE. 2005. Dissection of keratin dynamics: different contributions of the
794 actin and microtubule systems. *European Journal of Cell Biology* **84**:311–328.
795 doi:10.1016/j.ejcb.2004.12.004
- 796 Xu T, Langouras C, Koudehi MA, Vos BE, Wang N, Koenderink GH, Huang X, Vavylonis D. 2019.
797 Automated Tracking of Biopolymer Growth and Network Deformation with TSOAX.
798 *Scientific Reports* **9**:1717. doi:10.1038/s41598-018-37182-6
- 799 Xu T, Li H, Shen T, Ojic N, Vavylonis D, Huang X. 2011. Extraction and analysis of actin networks
800 based on Open Active Contour models2011 IEEE International Symposium on Biomedical

Imaging: From Nano to Macro. Presented at the 2011 IEEE International Symposium on Biomedical Imaging: From Nano to Macro. pp. 1334–1340. doi:10.1109/ISBI.2011.5872647

Xu T, Vavylonis D, Tsai F-C, Koenderink GH, Nie W, Yusuf E, Lee I-J, Wu J-Q, Huang X. 2015. SOAX: A software for quantification of 3D biopolymer networks. *Scientific Reports* **5**:9081. doi:10.1038/srep09081

Yamada S, Wirtz D, Coulombe PA. 2003. The mechanical properties of simple epithelial keratins 8 and 18: discriminating between interfacial and bulk elasticities. *J Struct Biol* **143**:45–55. doi:10.1016/s1047-8477(03)00101-1

6 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

7 ETHICS STATEMENT

n.a.

8 Author Contributions

- Conceptualization: RW, JDR, REL
- Methodology: RW
- Software: RW, MS, JS, AB
- Validation: RW
- Formal analysis: RW
- Investigation: RW, NS, SY, TP
- Resources: RW
- Writing—original draft: RW, REL
- Writing—review & editing: RW, REL
- Visualization: RW, AB
- Supervision: RW
- Project administration: RW, REL
- Funding acquisition: RW, REL

9 Funding

The work was supported by the German Research Council (LE566/18-2; WI173/8-2; GRK2415/363055819).

Calculations were performed with computing resources granted by RWTH Aachen University under project rwth0452.

J.D.R and T.P. were supported by a grant from the Interdisciplinary Centre for Clinical Research within the faculty of Medicine at the RWTH Aachen University.

10 Acknowledgments

We gratefully acknowledge the helpful discussions and advice of Drs. Michael Schaub (RWTH Aachen University), Charlotte Lorenz and Sarah Köster (Göttingen University), Rudolf Merkel (FZ Jülich). We also thank Michael Anhuth and Till Petersen-Krauß (RWTH Aachen University) for the implementation of the VR application and Ursula Wilhelm (MOCA) for providing the immunoblot.

10.1 Figure Legends

FIGURE 1 Evaluation of 3D keratin filament network segmentation. The pictures show three different numerical representations of the same fluorescence data set detecting the keratin filament network in an MDCK cell expressing fluorescently labelled keratin 8. The cell is part of a confluent monolayer and the fluorescence of the single cell shown was manually excised from an image stack consisting of 32 slices (voxel size: $xy = 66 \text{ nm}$, $z = 182 \text{ nm}$). This cropped fluorescence was used to calculate the vertex positions and intensity of all keratin filaments. **(A)** The top picture shows an overlay of the recorded fluorescence in red as a maximum intensity projection (mip) and the corresponding calculated xy positions of vertices of all segments as green dots. The enlargements below (corresponding regions demarcated by boxes) show maximum intensity projections of focal planes above and below the nucleus (apical and basal) at left and two regions of interest (roi 1, roi 2) depicting maximum intensity projections of all focal planes in the cell periphery. **(B)** presents an overlay of the keratin fluorescence and corresponding vertices that were plotted in 3D with a diameter relative to their brightness and were subsequently subjected to 3D Gaussian blurring to simulate the microscope's blur. The enlargements correspond to those shown at left in **(A)** presenting the same regions of interest and focal planes. The top panel of **Video 1** presents an animated comparison of the fluorescence recording, segmentation and overlay in each focal plane. **(C)** depicts a cinematic rendering of segments. Each calculated segment is rendered in 3D with a thickness corresponding to the brightness of the original filament. The color-coding helps to identify and distinguish individual segments. The regions of interest correspond to the same single planes as those shown in **(A)**. White arrows demarcate thin filaments, blue arrows thick filaments.

Figure 1-figure supplement 1. Immunoblot of a total lysate of a transgenic MDCK cell clone detecting endogenous wild-type keratin 8 (Krt8) and eYFP-tagged keratin 8 (Krt8-EYFP). Cells were lysed in 40 mM Tris-Cl (pH), suspended in Laemmli buffer at 4°C and separated by SDS-polyacrylamide gel electrophoresis. After transfer onto polyvinylidene fluoride Immobilon-P membrane (Millipore), keratin 8 epitopes were detected with a rat monoclonal anti-keratin 8 (TROMA-1) hybridoma cell supernatant (Developmental Studies Hybridoma Bank) and visualized with the help of secondary antibodies and the ECL system. The position and molecular weight of co-electrophoresed markers are indicated at left, the star denotes an immunoreactive band of unknown identity.

Figure 1-figure supplement 2. Exemplary representation of major steps in the workflow. All data are in 3D. **(A)** Maximum intensity projection of a small region of interest taken from an airyscan image of fluorescently labelled keratin filaments. **(B)** 'Snakes' prepared by TSOAX segmentation. White dots represent vertices and orange dots represent start and end vertices of 'snakes'. Colored lines distinguish individual 'snakes'. Note that 'snakes' are not well defined since not all 'snakes' have an end at intersections (red arrows). Therefore, the start and end points of snakes are not defined and snakes may span over several intersections. **(C)** In the final segmentation, segments (colored

lines) and nodes (red dots) are defined. No intersections of segments exist. Each segment has a start and end point (= node), and segments are connected by nodes. Y-shaped (blue arrow) or X-shaped (red arrow) branchings occur. **(D)** The cinematic rendering shows segments as tubes. Thickness represents mean brightness.

Figure 1-figure supplement 3. Validation of segmentation. **(A)** Maximum intensity projections of airyscan recordings of 10 cells each (left, MDCK; middle, HaCaT; right, RPE. **(B)** The pictures show the xy-position and brightness of segment vertices after 3D-Gaussian blurring. The resulting convolutions mimic those of microscopic imaging. **(C)** The merged images were prepared by overlay of the first and second row above to compare original imaging data with segmentation. **(D)** The picture shows all digital representations prepared from focal planes of airyscan image stacks of single cells. The xy-position of all segments is plotted in their corresponding slice position. Corresponding Video 1 shows an animated version of the data.

FIGURE 2 Comparison of 3D cinematic rendering of keratin filaments in three different epithelial cell types. The position and intensity of fluorescence recordings were transferred into a numerical 3D-representation of individual segments. The thickness of the segments corresponds to the fluorescence brightness in the original data. **(A)** shows cinematic rendering of keratin filaments of an MDCK cell expressing fluorescently labelled keratin 8. **Video 2** presents the 3D-animation of the cell. **(B)** Depicts the cinematic rendering of keratin filaments in a HaCaT keratinocyte expressing fluorescently labelled keratin 5 (animation in **Video 3**). **(C)** Illustrates cinematic rendering of keratin filaments in a single RPE cell of a homozygous keratin 8-YFP knock-in mouse (animation in **Video 4**).

Figure 2-figure supplement 1. Visual inspection of the keratin network of a single cell in a virtual reality environment. The viewer can move freely within the network which appears to her/him as a true 3D stereoscopic structure. Classification of segments is done interactively in this environment by color coding.

FIGURE 3 Mapping of keratin filament brightness in MDCK, HaCaT and RPE cells. **(A-C)** The histograms display the distribution of relative segment brightness for each cell type. The maximum brightness was normalized to 100 in all cases. For each segment, the mean brightness was calculated and the determined brightness levels of all segments in 10 cells were pooled for each diagram. **(D-F)** The maps depict the segment brightness in a representative MDCK (left), HaCaT (middle) and RPE (right) cell (top panel; enlarged boxed area in bottom panel). The colour code represents relative brightness.

FIGURE 4 Measurements of keratin filaments in MDCK, HaCaT and RPE cells. **(A-D)** The whisker box plots depict the number of segments per cell (A), the mean segment length (B), the entire length of the keratin filament network within a single cell (C) and the filament length per μm^3 (D). 10 cells were analyzed for each cell type. **(E-G)** show color-coded keratin filament segment lengths for each cell type. The top panel presents the results for entire cells, the bottom panel the results for the boxed areas at higher magnification.

Figure 4-figure supplement 1. Boxplots of cell volumes estimated from fluorescence image stacks **(A)** and calculated virtual persistence length of segments **(B)** in MDCK, HaCaT and RPE cells.

927

928 **FIGURE 5** Calculation of keratin filament node types based on segment/node ratios in MDCK,
 929 HaCaT and RPE cells. A segment/node ratio of 3 refers to Y-shaped branching, a ratio of 4 to X-
 930 shaped branchings. Higher values refer to star-like branchings. The histograms depict the relative
 931 distribution of node types.

932 **FIGURE 6** Measurements of keratin filament segment bending in MDCK, HaCaT and RPE cells.
 933 (A). The whisker box plot shows the distance/length ratio of keratin segments. (B) The curvature of
 934 keratin filaments in the three cell types is shown as a whisker box plot. (C-E) Color-coded
 935 representation of the keratin network depicting the curvature of keratin filament segments in entire
 936 cells (top) and selected regions of interest (boxed areas) at higher magnification (bottom).

937 **FIGURE 7** Orientation of keratin filament segments in MDCK, HaCaT, and RPE cells. (A-C) The
 938 histograms show the normalized distribution of azimuth (= horizontal orientation) of keratin filament
 939 segments in single cells. Since the start and end points are interchangeable, the histogram is restricted
 940 to 180 degrees. The horizontal line represents the expected histogram levels if all angles were equally
 941 distributed. (D-F) The histograms show the elevation (= vertical orientation) of keratin filament
 942 segments. The horizontal line demarcates the expected histogram values, if elevation of keratin
 943 filament segments would be uniform. (G, H) Whisker box plot depicting the deviation of segment
 944 orientations from uniform unbiased distributions. The differences in azimuth (G) and elevation (H)
 945 were calculated for 10 MDCK, 10 HaCaT and 10 RPE cells. (I-L) The color-coded maps depict the
 946 azimuth of segments in entire cells (top; MDCK, left; HaCaT, middle; RPE, right) and selected
 947 regions of interest (boxed areas; bottom).

948 **Figure 7-figure supplement 1. Schematic illustration of keratin filament segment (green)**
 949 **orientation in 3D space.** The orientation of a given segment between start point (green dot) and end
 950 point (yellow dot) is calculated as the angles of azimuth and elevation.

951

952 **Figure 7-figure supplement 2. Histograms of keratin filament orientation** in MDCK, HaCaT, and
 953 RPE cells. (A) The normalized histograms show the normalized distribution of azimuth (= horizontal
 954 orientation) of keratin filament segments in 10 MDCK, HaCaT, and RPE cells, respectively. Since
 955 the segments have no defined start and end point, the histograms are restricted to 180 degree. The
 956 horizontal line represents the value, if all angles were uniformly distributed. (B) The normalized
 957 histograms show the elevation (= vertical orientation) of keratin filament segments in 10 MDCK,
 958 HaCaT, and RPE cells. The horizontal line delineates the value, if all angles were uniformly
 959 distributed.

960

961 **FIGURE 8** Analysis of segment orientation in MDCK, HaCaT and RPE cells. (A-F) The blue
 962 polygons represent radial histograms of the orientation of keratin filament segments in relation to the
 963 cell center of single cells. Horizontal orientation (A-C) and the vertical orientation (D-F) are shown.
 964 Each keratin segment in each cell was moved to the cell center and vectors pointing in the same
 965 directions were added. Thus, the longer the blue lines are, the more vectors point in that direction.
 966 (G-I) The red arrows represent the sum vectors of the direction and length of all segments in each
 967 cell. (K) Whisker box plots depicting the length of the main vectors in all three cell types.

968

969 **FIGURE 9** Schematic drawing of the reconstructed keratin network in the three cell types
 970 investigated.

971

972

973 10.2 Video legends

974

975 **Video 1.** Validation of keratin filament segmentation in an MDCK, HaCat and RPE cell. The
976 animation shows the original fluorescence image stacks (left), the segmented data including the
977 associated brightness (middle) and the overlays of both (right).

978

979 **Video 2.** Cinematic rendering of the reconstructed keratin filament network of an MDCK cell
980 growing in a confluent monolayer. The tubes represent keratin bundles with different thickness. At
981 the beginning of the video the segments are randomly color-coded.

982

983 **Video 3.** Cinematic rendering of the numerical reconstructed keratin network of a HaCaT cell
984 growing in a confluent monolayer. The tubes represent keratin bundles with different thickness. At
985 the beginning of the video the segments are randomly color-coded.

986

987 **Video 4.** Cinematic rendering of the numerical reconstructed keratin network of an RPE cell. The
988 tubes represent keratin bundles with different thickness. At the beginning of the video the segments
989 are randomly color-coded.

990

991

992

relative brightness

segments/cell

segment length

length/cell

length/ μm^3

filament length

MDCK

HaCaT

RPE

MDCK **HaCaT** **RPE**

orientation

MDCK

HaCaT

RPE

segment orientation

MDCK

HaCaT

RPE

Σ length vectors/10

MDCK

HaCaT

photoreceptors

RPE

Bruch's membrane

lysate

kDa

100

72

55

25

Krt8-eYFP

Krt8

*

cell volume

persistence length

A

filament azimuth

MDCK

B

filament elevation

MDCK