A guanosine tetraphosphate (ppGpp) mediated brake on photosynthesis is required for acclimation to nitrogen limitation in Arabidopsis.

Shanna Romand¹, Hela Abdelkefi^{1,2}, Cecile Lecampion¹, Mohamed Belaroussi¹, Melanie Dussenne¹, Brigitte Ksas⁶, Sylvie Citerne³, José Caius^{4,5}, Stefano D'Alessandro¹, Hatem Fakhfakh^{2,6}, Stefano Caffarri¹, Michel Havaux⁷, Ben Field^{1*}

¹Aix-Marseille University, CEA, CNRS, BIAM, LGBP Team, Marseille, France F-13009

²University of Tunis El Manar, Faculty of Sciences of Tunis, Laboratory of Molecular Genetics, Immunology and Biotechnology, 2092 Tunis, Tunisia

³Institut Jean-Pierre Bourgin, UMR1318 INRA-AgroParisTech, INRAE Centre de Versailles-Grignon, Université Paris-Saclay, Versailles, France

⁴ Université Paris-Saclay, CNRS, INRAE, Univ Evry, Institute of Plant Sciences Paris-Saclay (IPS2), 91405, Orsay, France.

⁵Université de Paris, CNRS, INRAE, Institute of Plant Sciences Paris-Saclay (IPS2), 91405, Orsay, France

⁶University of Carthage, Faculty of Sciences of Bizerte, Tunisia

⁷Aix-Marseille University, CEA, CNRS, BIAM, SAVE Team, Saint-Paul-lez-Durance, France F-13108

*Corresponding author, ben.field@univ-amu.fr

Abstract

Guanosine pentaphosphate and tetraphosphate (together referred to as ppGpp) are hyperphosphorylated nucleotides found in bacteria and the chloroplasts of plants and algae. In plants and algae artificial ppGpp accumulation can inhibit chloroplast gene expression, and influence photosynthesis, nutrient remobilisation, growth, and immunity. However, it is so far unknown whether ppGpp is required for abiotic stress acclimation in plants. Here, we demonstrate that ppGpp biosynthesis is necessary for acclimation to nitrogen starvation in *Arabidopsis*. We show that ppGpp is required for remodeling the photosynthetic electron transport chain to downregulate photosynthetic activity and for protection against oxidative stress. Furthermore, we demonstrate that ppGpp is required for coupling chloroplastic and nuclear gene expression during nitrogen starvation. Altogether, our work indicates that ppGpp is a pivotal regulator of chloroplast activity for stress acclimation in plants.

Introduction

Plants cannot easily escape harsh environmental fluctuations, and so their survival hinges on facing each threat. To this end plants have developed intricate stress perception and response mechanisms (Devireddy et al., 2021), where the chloroplast is recognized as both a major signaling hub and a target for acclimation (Kleine et al., 2021). Likely candidates for regulating chloroplast stress signaling are the hyperphosphorylated nucleotides guanosine pentaphosphate and tetraphosphate (together referred to as ppGpp) that are synthesized from ATP and GDP/GTP by chloroplast localized enzymes of the RelA SpoT Homologue (RSH) family (Boniecka et al., 2017; Field, 2018). In bacteria, where ppGpp was originally discovered, a considerable body of work indicates that ppGpp and related nucleotides interact directly with specific effector enzymes to regulate growth rate and promote stress acclimation (Ronneau and Hallez, 2019; Bange et al., 2021; Anderson et al., 2021). In plants, ppGpp signaling is less well understood both at the physiological and mechanistic levels. ppGpp levels increase transiently in diverse plants in response to treatment with a range of abiotic stresses and stress related hormones (abscisic acid, jasmonate and ethylene) (Takahashi et al., 2004; Ihara et al., 2015). The accumulation of ppGpp is controlled by the ppGpp synthesis and hydrolysis activity of three families of chloroplast targeted RSH enzyme, RSH1, RSH2/3 and RSH4/CRSH (Atkinson et al., 2011; Ito et al., 2017; Avilan et al., 2019). In the angiosperm *Arabidopsis thaliana* (Arabidopsis), where ppGpp signaling is best characterized, RSH1 lacks ppGpp synthase activity and acts as the main ppGpp hydrolase (Sugliani et al., 2016), the closely related RSH2 and RSH3 act as the major ppGpp synthases during the day (Mizusawa et al., 2008; Maekawa et al., 2015; Sugliani et al., 2016), and the calcium activated RSH (CRSH) is responsible for ppGpp synthesis at night and in response to darkness (Ihara et al., 2015; Ono et al., 2020). RSH2 and RSH3 are bifunctional ppGpp synthase / hydrolase enzymes, and their hydrolase activity was recently shown to be necessary for constraining CRSH-mediated ppGpp production at night (Ono et al., 2020). The artificial accumulation of ppGpp itself has been shown to repress the expression of certain chloroplast genes *in vivo* via the inhibition of chloroplast transcription (Yamburenko et al., 2015; Maekawa et al., 2015; Sugliani et al., 2016; Ono et al., 2020). The chloroplastic effectors of ppGpp signaling in plants have not yet been identified. Nevertheless, ppGpp is able to specifically repress activity of the chloroplast guanylate kinase, suggesting the existence of a ppGpp effector mechanism that is found in many firmicute bacteria (Nomura et al., 2014; Field, 2018; Bange et al., 2021). According to the firmicute model, guanylate kinase inhibition causes a drop in GTP concentration, which in turn results in reduced transcription

from rRNA-coding genes where GTP is usually the initiating nucleotide. Over accumulation of ppGpp in the chloroplast by the over-expression of RSH3 or bacterial ppGpp synthase domains has also shown that ppGpp may act as a conserved repressor of photosynthesis in both land plants (Maekawa et al., 2015; Sugliani et al., 2016; Honoki et al., 2018; Harchouni et al., 2021) and algae (Avilan et al., 2021). Interestingly, the artificial accumulation of ppGpp also appears to protect plants against nitrogen deprivation (Maekawa et al., 2015; Honoki et al., 2018). In addition, a few studies have now demonstrated that ppGpp is required for different physiological processes; these include the regulation of plant growth and development (Sugliani et al., 2016; Ono et al., 2020), plant immunity (Abdelkefi et al., 2018), and photosynthesis under standard growth conditions (Sugliani et al., 2016). However, there are so far no demonstrations that ppGpp is required for abiotic stress acclimation in plants.

Here, we looked at the role of ppGpp in acclimation to nitrogen deprivation. ppGpp was discovered in bacteria during research into the acclimation of *Escherichia coli* to amino acid limitation (Cashel and Gallant, 1969). We reasoned that a similar role was likely to be maintained in plants due to a shared and fundamental requirement for amino acids and the nitrogen to synthesize them. In particular, chloroplasts, and the photosynthetic machinery within, represent a major nitrogen store that must be remobilized during periods of nitrogen limitation and during senescence. We found that ppGpp biosynthesis by RSH2 and RSH3 is necessary for acclimation to nitrogen starvation in Arabidopsis. During nitrogen starvation ppGpp acts by remodeling the photosynthetic electron transport chain to downregulate photosynthetic activity, and by protecting against oxidative stress and tissue damage. Furthermore, we show that ppGpp couples chloroplastic and nuclear gene expression during nitrogen starvation. Overall, our work indicates that ppGpp is a pivotal regulator of chloroplast activity with a photoprotective role and that ppGpp signaling is required for the acclimation of plants to harsh environmental conditions.

Results

ppGpp is required for acclimation to nitrogen deprivation

To determine whether ppGpp plays a significant physiological role during nitrogen starvation we grew a series of *RSH* lines on a nitrogen limiting medium that imposes a progressive nitrogen starvation (Fig. 1A). The *RSH* lines were previously shown to accumulate lower (OX:RSH1, *RSH* quadruple mutant [*rsh_{QM}*]) or higher (*rsh1-1*) amounts of ppGpp under standard growth conditions (Sugliani et al., 2016). Growth arrest occurred for all lines 9-10 days after sowing on nitrogen limiting media and was followed by the production of anthocyanins and loss of chlorophyll (Fig. 1A, Fig. 1- figure supplement 1). Strikingly, we observed that the low ppGpp lines OX:RSH1 and *rsh_{QM}* showed significantly higher rates of cotyledon death than in the corresponding wild-type plants (Fig. 1A, B). Interestingly, between 16 and 22 days we also saw a greater rate of new leaf initiation in the ppGpp deficient lines (Fig. 1A).

We reasoned that the increased cotyledon death in the low ppGpp lines could be due to overproduction of reactive-oxygen species (ROS). We therefore imaged the autoluminescence of lipid peroxides, a signature of ROS accumulation (Fig. 1C). Under nitrogen-deprivation conditions strong autoluminescence was observed only in the OX:RSH1 and *rsh_{QM}* low ppGpp lines, while high ppGpp *rsh1-1* plants showed almost no autoluminescence (Fig. 1C). Quantification of lipid peroxidation products (hydroxy-octadecatrienoic acids [HOTEs], the oxidation products of linolenic acid, the major fatty acid in Arabidopsis leaves) by HPLC further supported these findings: HOTEs increased in the wild-type in response to nitrogen deprivation, and this increase was significantly larger in the ppGpp deficient lines (Fig. 1D). These results indicate that nitrogen deprivation promotes ROS accumulation, and that ppGpp is required to prevent overaccumulation of ROS, oxidative stress and death of cotyledons under these conditions.

Figure 1. ppGpp is required for acclimation to nitrogen deprivation. (A) Images of seedlings grown on nitrogen replete (+N) or nitrogen limiting (-N) medium for 8 (D8), 12 (D12) and 22 (D22) days. Scale, 3 mm. WT1 (Col-0) and WT2 (*qrt1-2*) are the wild type for OX:RSH1 and *rsh_{QM}* / *rsh1-1* respectively. See Figure 1–figure supplement 1 for additional images. (B) Percentage of plants with dead cotyledons (completely white with collapsed tissue) for different genotypes grown on -N medium for 22 days. Three pooled experimental replicates, median +/- 95% CI, n = 285 - 298 seedlings per genotype. (C) Bioluminescence emission from lipid peroxides in seedlings grown on -N medium for 16 days. Inset graph shows median number of luminescent seedlings +/- 95% CI, n = 100 seedlings. Scale, 1.1 cm. (D) Quantification of HOTE from seedlings grown in -N for 12 days or +N for 8 days where the developmental stage is similar. Mean +/- 95% CI, n = 4 experimental replicates. Concentrations of ppGpp and GTP in wild-type plants were determined (E) at the indicated time points during growth on +N and -N medium and (F) in different genotypes after 12 days of growth in -N medium (equivalent to 10 days in the experiment in panel E). The ppGpp/GTP ratio is shown in Figure 1–figure supplement 2. Mean +/- 95% CI, n = 4 experimental replicates. Statistical tests shown against respective wild-type controls, * $P < 0.05$, ** $P < 0.01$. Source data and statistical test reports are shown in Figure 1–source data 1.

ppGpp levels increase at an early stage of nitrogen deprivation

In bacteria and in plants ppGpp is known to peak in response to stress before stabilizing at lower levels (Varik et al., 2017; Takahashi et al., 2004a; Ihara et al., 2015). To determine the kinetics of changes in ppGpp concentration during nitrogen deprivation we quantified ppGpp levels in wild-type plants at different timepoints during growth on nitrogen limiting media. We observed a peak in ppGpp levels after 8 days on nitrogen limiting medium that was not observed in plants of the same age grown on nitrogen replete medium (Fig. 1E). The ppGpp concentration decreased and stabilised after 12 days on nitrogen limiting medium. In contrast, the level of GTP decreased throughout the timecourse. We next determined how (p)ppGpp levels were affected in each of the *RSH* lines during growth on nitrogen limiting medium (Fig. 1F). As anticipated we found that ppGpp levels in OX:RSH1 and *rsh_{QM}* lines were lower than in wild-type plants, while they were higher in *rshI-1* plants. Strikingly, we found that the lines with low ppGpp levels (OX:RSH1 and *rsh_{QM}*) maintained higher levels of GTP than in the wild-type controls or *rshI-1* (Fig. 1F), resulting in dramatic differences in the ppGpp/GTP ratio (Fig. 1- figure supplement 2). Therefore, nitrogen limitation leads to an early increase in ppGpp levels and is followed by a progressive diminution of the GTP pool that is dependent on ppGpp accumulation. These results suggest that ppGpp inhibits GTP biosynthesis, perhaps through inhibition of guanylate kinase (Nomura et al., 2014), and supports the hypothesis that plants have a ppGpp effector mechanism that operates through purine metabolism as found in many bacteria.

Nitrogen deprivation promotes a ppGpp-dependent drop in photosynthetic activity

Under standard growth conditions artificial production of ppGpp has been shown to downregulate photosynthetic activity in Arabidopsis (Sugliani et al., 2016; Maekawa et al., 2015). Defective downregulation of photosynthesis during nitrogen starvation could cause the over-accumulation of ROS and lipid peroxides observed in the low ppGpp lines due to oversaturation of photosynthesis in the presence of diminished metabolic sinks. We therefore measured different photosynthetic parameters in the wild type and *RSH* lines during growth on nitrogen limiting medium. A decrease in the maximum quantum yield (Fv/Fm) of photosystem II (PSII) caused by an increase in basal fluorescence (Fo, see Materials and Methods for formula) was observed in the cotyledons and first set of true leaves of wild-type seedlings (Fig. 2A-B, Fig. 2- figure supplement 1, Fig. 2- source data 1). Remarkably, this decrease was almost completely suppressed in the low ppGpp *rsh_{QM}* and OX:RSH1 lines, and was enhanced in the high ppGpp *rshI-1* mutant (Fig. 2A, 2B, Fig. 2- figure supplement 2A).

181 Analysis of *rsh2-1* and *rsh3-1* single mutants, as well as the complementation of an *rsh2-1*
 182 *rsh3-1* double mutant with the genomic version of *RSH3*, indicates that the RSH3 enzyme is
 183 the major ppGpp synthase responsible for driving the Fv/Fm decrease under nitrogen
 184 starvation conditions (Fig. 2- figure supplement 2B). Likewise, the enhanced Fv/Fm decrease
 185 in the high ppGpp *rsh1-1* mutant and complementation of the *rsh1-1* phenotype by expression
 186 of the genomic *RSH1* indicate that the ppGpp hydrolase RSH1 acts antagonistically to RSH3,
 187 probably by constraining ppGpp accumulation during nitrogen deprivation (Fig. 1F. 2A, 2B,
 188 Fig. 2- figure supplement 2C).

189 **Figure 2. Nitrogen deprivation promotes a ppGpp-dependent drop in photosynthetic capacity.**
 190 Seedlings were grown 8 days on nitrogen replete media (+N) or 12 days on nitrogen limiting (-N)
 191 media and (A) the maximal yield of PSII (Fv/Fm) measured by fluorescence imaging individual
 192 seedlings. Median +/- 95% CI, n = 95-100 seedlings. (B) Fv/Fm images of whole seedlings grown on
 193 +N and -N media for 12 days. (C) Relative ETR measurements in different lines grown 8 days on +N
 194 media or 12 days on -N media. Median +/- 95% CI, n = 95-100 seedlings. (D) Fv/Fm time-courses
 195 from seedlings grown on -N media and transferred to three different light intensities (photosynthetic
 196 photon flux density, 10, 80 and 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$) after 6 days. Median +/- 95% CI, n = 95-100
 197 seedlings. Tests shown against respective wild-type controls, * $P < 0.05$, ** $P < 0.01$. Additional
 198 supporting data is presented in Figure 2- figure supplements 2-4. Source data and statistical test
 199 reports are shown in Figure 2- source data 1.

While unlikely, it is possible that RSH enzymes could influence Fv/Fm via mechanisms that do not require ppGpp synthesis or hydrolysis. To exclude this possibility we conditionally overexpressed a heterologous chloroplast targeted ppGpp hydrolase from *Drosophila melanogaster* (Metazoan SpoT homolog, MESH)(Sugliani et al., 2016) (Fig. 2- figure supplement 2D). We observed that ppGpp depletion by MESH suppresses the Fv/Fm decrease in response to nitrogen limitation, confirming a specific role for ppGpp in this process.

Our experiments were carried out on plate-grown seedlings and in the presence of exogenous sugar in the media. To test the robustness of our observations to different experimental set-ups we exposed mature plants grown in quartz sand to nitrogen starvation conditions (Fig. 2- figure supplement 2E). We observed responses from the wild type, *rsh_{QM}* and *rshI-1* mutant that were consistent with those observed for plate grown seedlings, confirming the robustness of the ppGpp-dependent decrease in Fv/Fm in response to nitrogen limitation.

Fv/Fm measurements provide information principally on PSII. To understand the state of the entire photosynthetic electron transport chain we next measured the relative rate of electron transport (ETR). We found that nitrogen deprivation led to a large decrease in ETR in wild-type plants (Fig. 2C, Fig. 2- figure supplement 4A). The *RSH* lines showed similar ETRs to the wild type under nitrogen replete conditions. However, nitrogen deprivation led to large differences in the *RSH* lines compared to the wild type: a low ETR was observed in the high ppGpp *rshI-1* mutant, and a substantially higher ETR in the low ppGpp *rsh_{QM}* and OX:*RSH1* lines. Induction of the MESH ppGpp hydrolase also prevented much of the decrease in ETR observed under nitrogen deprivation, again indicating that this phenomenon is dependent on the activity of RSH enzymes and ppGpp accumulation (Fig. 2- figure supplement 4B).

The ppGpp-dependent decline in photosynthesis occurs even at low light fluences

The ppGpp-dependent drop in Fv/Fm during nitrogen deprivation is reminiscent of photoinhibition, a process whose rate is proportional to light intensity (Tyystjarvi and Aro, 1996). We therefore asked whether the ppGpp-dependent decrease in Fv/Fm is similarly dependent on the excitation pressure on the photosystems. Strikingly, under nitrogen deprivation the Fv/Fm of the wild type decreased to a similar minimum regardless of the light intensity (photosynthetic photon flux density; low light, 10 $\mu\text{mol m}^{-2}\text{s}^{-1}$; growth light, 80 $\mu\text{mol m}^{-2}\text{s}^{-1}$; and higher light, 150 $\mu\text{mol m}^{-2}\text{s}^{-1}$)(Fig. 2D). However, the rate of the Fv/Fm

decrease was proportional to light intensity, showing a slower decrease to the minimum under low light than under normal light or high light. A similar phenomenon was observed in the high ppGpp *rsh1-1* line, except that the Fv/Fm dropped at a faster rate and to a considerably lower level. In contrast, in the low ppGpp *rsh_{QM}* mutant we observed almost no decrease in Fv/Fm, regardless of the light intensity tested. The insensitivity of the *rsh_{QM}* mutant indicates that the process is completely dependent on ppGpp biosynthesis. Furthermore, the response of the wild-type and even stronger response of the *rsh1-1* mutant at all light fluences indicate that ppGpp-initiated regulation of Fv/Fm is triggered in response to nitrogen deprivation rather than to changes in the excitation status of the photosynthetic electron transport chain (Fig. 2D).

Figure 3. ppGpp-dependent alterations in the photosynthetic machinery during nitrogen deficiency. (A) Immunoblots showing evolution in abundance of the indicated proteins in seedlings grown in nitrogen replete (+N) or nitrogen limiting (-N) media for the indicated number of days. RbcL was revealed by Coomassie Brilliant Blue. Equal quantities of total proteins were loaded and PBA1, a subunit of the proteasome, was used as a protein normalization control (B) Immunoblots showing the abundance of the indicated proteins in purified thylakoid membranes from seedlings grown in +N for 8 days or -N for 12 days. RbcL was revealed by Coomassie Brilliant Blue staining. Equal quantities of total chlorophyll were loaded. Immunoblots after ppGpp-depletion by induction of chloroplastic MESH are shown in Figure 3- supplement 1. (C) Chlorophyll a/b ratios in extracts from seedlings subjected to -N for 12 days. Means $\pm$ 95% CI, data from four experimental replicates. (D) Emission spectrum of chlorophyll fluorescence at 77°K between 660 and 720 nm, normalized to the PSII peak at 685 nm. The full spectra are shown in Figure 2- supplement 2. Measurements were made on seedlings grown in +N for 8 days or -N for 12 days. Means $\pm$ 95% CI; data from four experimental replicates. Statistical tests, ** $P < 0.01$. Uncropped immunoblots are available in Figure 3- source data 1 for 3A and source data 2 for 3B. Numeric data and statistical test reports are shown in Figure 3- source data 3.

ppGpp is required for the timely degradation of photosynthetic proteins

In order to better understand the changes in the photosynthetic machinery that underlie the reduction of photosynthetic capacity during nitrogen starvation we analyzed the abundance of representative photosynthetic proteins in the wild type and the low ppGpp line OX:RSH1 (Fig. 3A). In the wild type, we observed a decrease in proteins representative of nearly all the major photosynthetic complexes analysed (PsbA, Lhcb1, PsbO, PetA, PsaD, Lhca1, RBCL) and PTOX analyzed between 8 and 16 days of growth under nitrogen deprivation. The decrease in photosynthetic proteins was defective in OX:RSH1 plants, which showed a marked delay for the decrease in abundance of PsbA, PsbO, PetA, PTOX, PsaD and RBCL. This delay was especially visible after 12 days on nitrogen limiting medium. On the contrary, changes in Lhcb1 and Lhca1, the nuclear encoded light harvesting proteins of PSII and PSI, were indistinguishable between wild-type and OX:RSH1 plants.

In contrast to the downregulation of many components of the photosynthetic chain during nitrogen deprivation, mitochondrial activity appeared to increase. The mitochondrial marker protein COXII, a subunit of the mitochondrial complex IV, increased to reach a maximum on day 12 of nitrogen deprivation. Interestingly, the COXII maximum was higher in the wild type than in OX:RSH1 plants suggesting increased ppGpp levels could also positively influence mitochondrial activity. PsbS, a key component of non-photochemical quenching, and SAG12, a marker of senescence, showed similar patterns of accumulation in WT and OX:RSH1 plants.

We next extended our analysis from OX:RSH1 to the other RSH lines to confirm that ppGpp is responsible for the observed changes in photosynthetic proteins during nitrogen deprivation (Fig. 3B). In this experiment we normalized to total chlorophyll to exclude possible artefacts linked to normalization on total protein, yet we obtained broadly similar results. PetA and RBCL, the proteins with the greatest differences in OX:RSH1 during the timecourse, were more abundant in the low ppGpp lines OX:RSH1 and *rsh_{QM}* at day 12 under nitrogen deprivation. The same proteins were also less abundant than the wild type in the high ppGpp line *rsh1-1*. Interestingly, PsbA accumulation did not appear different between the lines, contrasting with results from the OX:RSH1 time course. However, this difference is likely linked to the chlorophyll normalization because Lhcb1 levels were higher in the wild type and *rsh1-1* mutant, indicating a similar direction of change for the ratio of PSII core to antenna.

Finally, to unequivocally link these changes to ppGpp we analyzed representative protein levels in induced MESH ppGpp hydrolase lines, which again showed the ppGpp is required for diminution of PetA and RBCL levels during nitrogen deprivation (Fig. 3- figure supplement 1). Altogether our immunoblotting experiments indicate the ppGpp is required for the timely reduction in abundance of key photosynthetic proteins during nitrogen starvation, and that proteins such as PsbO (PSII core), PetA (Cyt b₆f) and RBCL (Rubisco) appear to be preferentially affected.

Nitrogen starvation promotes ppGpp-dependent energetic uncoupling of PSII antenna

The ppGpp dependent drop in Fv/Fm and drop in the PsbA / Lhcb1 ratio suggest that ppGpp specifically remodels the structure of the PSII complex during nitrogen deprivation. Supporting this idea, we also found a ppGpp dependent drop in the Chl a/b ratio (Fig. 3C), indicating a substantial decrease in PSII RC, which lacks Chl b, to Chl b rich PSII antenna. An increase in relative PSII antenna abundance might be accompanied by the energetic uncoupling from the PSII RC. Low temperature chlorophyll fluorescence spectra confirmed this hypothesis. During nitrogen deprivation there was a shift of the PSII antenna emission peak towards lower wavelengths in the wild type, indicating an energetic decoupling from PSII RC (Crepin et al., 2016)(Fig. 3D, Fig. 3- figure supplement 2). This shift is mainly ppGpp dependent because it was smaller in the low ppGpp OX:RSH1 and *rsh_{QM}* lines, and much stronger in the high ppGpp *rsh1-1* line. Therefore, multiple approaches show that ppGpp is necessary for PSII remodeling and decreasing the excitation pressure on the photosystems during nitrogen starvation.

ppGpp plays a major role during acclimation to nitrogen deprivation

Our results show that defects in the capacity of plants to accumulate ppGpp lead to aberrant photosynthesis and stress phenotypes under nitrogen deprivation. To determine the extent and specificity of the impact of ppGpp on cellular processes we analyzed global nuclear and organellar transcript abundance in wild-type and OX:RSH1 plants.

Nitrogen deprivation caused a massive alteration in transcript levels in both the wild type and OX:RSH1, significantly affecting the accumulation of about 15,000 transcripts (Fig. 4A, Fig. 4- source data 1). For the wild type there was substantial overlap (80%) with differentially accumulating transcripts recently identified in plants grown under chronic nitrogen limitation for 60 days (Luo et al., 2020). Strikingly, we observed large increases in *RSH2* and *RSH3*

transcript levels in response to nitrogen deprivation (Fig 4B). These were accompanied by a decrease in *RSH1* transcript levels, suggesting a coordinated upregulation of ppGpp biosynthetic capacity. The expression profile of OX:RSH1 plants showed clear differences to the wild type, especially under nitrogen deprivation. Under control conditions OX:RSH1 plants had a very similar gene expression profile to the wild type, with the differential accumulation of only 150 transcripts (Fig. 4A, Fig. 4- source data 1). However, under nitrogen deprivation we observed the differential accumulation of 2700 transcripts in OX:RSH1. The greater deregulation of the OX:RSH1 transcript profile under nitrogen deprivation indicates that ppGpp plays a major and specific role in the acclimation response.

Analysis of enrichment for gene ontology terms corroborated our findings that low ppGpp plants are unable to properly acclimate to nitrogen deprivation. OX:RSH1 plants accumulated more transcripts associated with stress, including oxidative stress than the wild type (Fig. 4C, Fig. 4- source data 2). We also observed a significant decrease in the abundance of transcripts for genes involved in photosynthesis and chloroplast activity. Notably, these downregulated genes were all nucleus encoded (Fig. 4C).

Figure 4. ppGpp plays a major role during acclimation to nitrogen deprivation. RNA-seq experiments were performed on WT and OX:RSH1 seedlings grown 8 days on nitrogen replete media (+N) or 12 days on nitrogen limiting (-N) media, n=3 experimental replicates. **(A)** Venn diagram for transcripts showing differential accumulation for each of three comparisons. All differentially accumulating transcripts are listed in Figure 4- source data 1. **(B)** RNA-seq transcript levels for the four *RSH* genes in the WT, +N vs -N. Mean +/- 95% CI, ** $P<0.01$. **(C)** Enriched gene ontology terms among significantly up- and down-regulated transcripts in OX:RSH1 vs WT under -N. The ten most significant terms are shown, point size is proportional to gene number. FDR, false discovery rate. The full GO analysis is presented in Figure 4- source data 2. Source data and statistical test reports are shown in Figure 4- source data 3.

ppGpp is required for global downregulation of chloroplast gene expression

We next turned to the effects of nitrogen deprivation on organellar gene expression. Nitrogen deprivation leads to a strong decrease in the levels of all chloroplast transcripts except *PsbN* in the wild type (Fig. 5A, Fig. 5- source data 1). The OX:RSH1 mutant showed minor differences to the wild type under normal growth conditions. However, under nitrogen starvation the vast majority of chloroplast transcripts (120 of 133 transcripts analyzed) were

present at significantly higher levels in OX:RSH1 than in the wild type, indicating that ppGpp is required for globally reducing chloroplast gene expression. In contrast to the monodirectional response of the chloroplast to nitrogen deprivation, we found that mitochondrial transcripts showed diverse responses in the wild type (Fig. 5B, Fig. 5- source data 2). Significantly, mitochondrial transcript levels in the OX:RSH1 line were highly similar to the wild type under control and nitrogen deprivation conditions. Taken together, these results indicate that ppGpp is required for the downregulation of chloroplast gene expression during nitrogen starvation and demonstrate that ppGpp acts specifically within the chloroplast.

Figure 5. ppGpp is required for the downregulation of chloroplast gene expression during nitrogen deficiency. (A) The differential expression of chloroplast transcripts ordered along the chloroplast genome (Figure 5- source data 1). (B) The differential expression of mitochondrial transcripts ordered along the mitochondrial genome (Figure 5- source data 2). (C) Relative transcript levels in OX:RSH1 versus the wild-type control for nuclear (black) and chloroplast (green) genes encoding subunits of the photosystem II complex. Solid colors indicate significant changes in expression ($P<0.05$), transparent colors indicate non-significant changes. Source data for 5C is shown in Figure 5- source data 3.

ppGpp is required for coordinating chloroplast and nuclear gene expression

Our gene expression analysis showed the ppGpp depletion during nitrogen starvation appeared to have opposite effects on transcript abundance for nuclear encoded chloroplast genes and chloroplast encoded genes. Many chloroplast protein complexes contain subunits encoded by both the nuclear and chloroplast genomes. Notably, in the wild-type chloroplast protein complexes showed coordinate downregulation of both nuclear and chloroplast

transcripts in response to nitrogen deprivation (Fig. 5- figure supplement 1). However, ppGpp depletion caused widespread mis-regulation of transcript abundance for complex subunits (Fig 5C- source data 3). For the PSII complex there was little difference between OX:RSH1 and the wild type under nitrogen replete conditions. However, in OX:RSH1 nitrogen deprivation caused the large scale uncoupling of the coordination of gene expression between the nuclear and chloroplast genomes: all but one of the transcripts for nuclear-encoded subunits of PSII were downregulated, while all the transcripts for chloroplast-encoded subunits of PSII were upregulated (Fig. 5C). Similar genome uncoupling was observed in OX:RSH1 for transcripts encoding subunits of the chloroplastic Cyt b_6f , PSI, ATP synthase, NDH, TIC, transcription and translation complexes (Fig. 5- source data 3). These results indicate first that nitrogen deprivation promotes the coordinated downregulation of genes encoding chloroplastic proteins in the nuclear and chloroplastic genomes, and second that ppGpp plays a pivotal role in coupling gene expression between nuclear and chloroplast genomes under nitrogen limitation.

Discussion

Nitrogen deprivation has long been known to cause a drop in photosynthetic capacity in plants (Terashima and Evans, 1988; Nunes et al., 1993; Verhoeven et al., 1997; Lu and Zhang, 2000; Garai and Tripathy, 2018). Here we show that ppGpp signaling plays a major role in this process. We demonstrate that nitrogen starvation leads to an early and transient increase in ppGpp levels (Fig. 1E), and that the capacity to synthesize and accumulate ppGpp is then required for protecting plants against excess ROS accumulation, tissue damage and stress (Fig. 1A-D, 4C). We show that ppGpp is likely to mediate the acclimation response by affecting the ppGpp/GTP ratio (Fig. 1- figure supplement 2), promoting the downregulation of photosynthetic capacity (Fig. 2A-D, 4C, 5A) and remodeling the photosynthetic machinery (Fig. 3A-D, 5C). Finally, we show that ppGpp may function by specifically downregulating chloroplast transcript abundance during nitrogen deprivation to maintain an equilibrium between chloroplast and nuclear gene expression (Fig. 5A-C, Fig. 5- source data 1). ppGpp was initially discovered thanks to the identification of bacterial mutants that were unable to downregulate rRNA transcription during amino acid starvation (Stent and Brenner, 1961; Cashel and Gallant, 1969). Our results here therefore indicate that the fundamental involvement of ppGpp signaling in acclimation to nitrogen deprivation is likely to have been maintained despite the large evolutionary distance separating bacteria from plants that includes major shifts including the domestication of the cyanobacterial ancestor of the chloroplast, and the different regulatory logic and signaling networks of photosynthetic eukaryotes.

The physiological relevance of ppGpp signaling in plants

Depletion or removal of ppGpp is usually necessary for directly establishing the implication of ppGpp in a physiological process. Up to now a handful of studies in plants have directly implicated ppGpp physiological processes by showing a requirement for ppGpp synthesis (Sugliani et al., 2016; Abdelkefi et al., 2018; Honoki et al., 2018). We now add to these studies by using multiple approaches, including three different ppGpp depletion methods (OX:RSH1, *rsh_{QM}* and MESH), to demonstrate that ppGpp is required for acclimation to nitrogen deprivation. Previous work hinted that ppGpp may play such a role by showing that plants ectopically over-accumulating ppGpp appear able to better withstand transfer to media lacking nitrogen (Maekawa et al., 2015; Honoki et al., 2018). Notably, Honoki et al. (2018) also used a similar *rsh2 rsh3* double mutant in their experiments though, apart from a delay in Rubisco degradation, did not observe clear-cut differences from the wild type in terms of

photosynthetic activity after transfer to media lacking nitrogen. The reasons for this are not clear, but it could be related to differences in experimental set up such as the timepoints for analysis, how nitrogen starvation was imposed, or the different RSH3 mutant allele used.

Our findings further reinforce the evidence that RSH enzymes function in the chloroplast. GFP fusion experiments have shown that RSH enzymes are localised to the chloroplasts (Maekawa et al., 2015), and the pleiotropic phenotypes of RSH3 overexpressing plants can be suppressed by chloroplast expression of a ppGpp hydrolase (Sugliani et al., 2016). However, some proteins, such as ATP synthase subunit beta-3, show low levels of dual targeting that are not detectable using full length fluorescent protein fusions (Sharma et al., 2019). Coverage of the full mitochondrial transcriptome here shows that overexpression of RSH1 has almost no effect on the large-scale changes in mitochondrial transcript abundance that occur in response to nitrogen deprivation (Fig. 5B). These results therefore confirm that RSH enzymes operate within the chloroplast.

New insights into ppGpp signaling in planta

The kinetics of changes in ppGpp concentration during nitrogen deprivation provide new information on how ppGpp acts *in planta*. We observed a transient peak in ppGpp levels after eight days of growth on nitrogen limiting medium (Fig. 1E). Similar transient peaks of ppGpp accumulation were also observed in plants subjected to different stresses (Takahashi et al., 2004; Ihara et al., 2015; Ono et al., 2020), and are observed in bacterial cells (Varik et al., 2017). Notably, the peak in ppGpp that we observed in response to nitrogen deprivation occurs prior to major changes in Fv/Fm (Fig. 2D), protein levels (Fig. 3A), or anthocyanin accumulation (Fig. 1- figure supplement 1). These factors, along with the progressive nature of the nitrogen starvation suggest that a signaling mechanism activates ppGpp biosynthesis relatively soon after perception of nitrogen supply limitation. The upregulation of *RSH2* and *RSH3* expression (Fig. 4B) indicates that there is at least a transcriptional component to this response. However, additional layers of regulation are likely, including for example the allosteric activation of RSH enzymes by interaction with other proteins or small molecules as found for bacterial RSH (Ronneau and Hallez, 2019; Irving and Corrigan, 2018). After the initial peak the ppGpp concentration then drops to levels not very different to those observed in plants grown in nitrogen replete conditions (Fig. 1E). Furthermore, OX:RSH1 plants only show slightly lower levels of ppGpp than the wild type despite showing very similar phenotypes to the *rsh_{QM}* mutant (Fig. 1F). Together these observations indicate that ppGpp

continues to exert an effect after peaking, suggesting the activation of irreversible processes or increased sensitivity to the presence of ppGpp. The second of these possibilities is supported by the phenotype of the high ppGpp *rsh1-1* mutant which displays much stronger photosynthesis phenotypes than wild-type plants despite a relatively modest increase in ppGpp levels (Fig. 1F, 2A-D, 3A-D).

GTP is a key nucleotide in ppGpp signaling because it is both the substrate for ppGpp biosynthesis and enzymes in its biosynthetic pathway such as guanylate kinase are potential ppGpp effectors (Field, 2018). We observed a striking decrease in GTP levels during nitrogen deprivation (Fig. 1E). Remarkably GTP levels were closely and inversely correlated to ppGpp levels in the wild type and the *RSH* lines (Fig. 1E, F). GTP is present at levels almost three orders of magnitude higher than ppGpp, so the relationship between ppGpp and GTP cannot be simply explained by the consumption of GTP for ppGpp biosynthesis. Therefore, ppGpp accumulation drives the depletion of the cellular GTP pools by other means, which could for example include the inhibition of enzymes involved in chloroplastic GTP biosynthesis such as guanylate kinase (Nomura et al., 2014). Even more remarkable is that the ppGpp-driven GTP depletion occurs specifically under nitrogen deprivation because accumulation of ppGpp to high levels in OX:RSH3 plants does not affect the GTP pool under normal growth conditions (Bartoli et al., 2020). The reduction in GTP levels may contribute to ppGpp signaling in two ways. First, reduced GTP levels are likely to directly contribute to the observed ppGpp-dependent reduction in chloroplast transcription in a similar fashion to ppGpp-signaling in firmicute bacteria (Bange et al., 2021). Second, the increase in the ppGpp / GTP ratio will significantly augment the action of ppGpp on effectors where ppGpp competes with GTP for binding. GTP-binding effectors are common in bacteria, and many are conserved in chloroplasts, particularly those involved in translation and ribosome biogenesis (Pausch et al., 2018; Field, 2018). We propose that the increase in the ppGpp / GTP ratio during nitrogen limitation is may also explain how ppGpp maintains its influence despite decreasing in absolute amounts.

The impact of ppGpp on photosynthesis under nitrogen deprivation

Our work also reveals the wide-ranging physiological effects of ppGpp on photosynthetic activity during nitrogen deprivation. These effects are reminiscent of the photosynthetic changes that occur during senescence (Krieger-Liszkay et al., 2019), and are remarkably similar to those found using the ectopic over-accumulation of ppGpp in Arabidopsis, moss

and algae (Maekawa et al., 2015; Sugliani et al., 2016; Imamura et al., 2018; Honoki et al., 2018; Avilan et al., 2021; Harchouni et al., 2021). However, compared to these studies our more detailed analysis here highlights first the global effect of ppGpp on chloroplast transcript abundance (Fig. 5A), and second the strong effect of ppGpp on certain photosynthetic proteins such as PsbO from PSII, PetA from Cyt b₆f, and RbcL (Fig. 3A, B). ppGpp controls the abundance of these proteins via the downregulation of chloroplast transcript abundance (Fig. 5A), reducing the quantity of transcripts available for translation as well as translation capacity via a likely reduction in rRNA transcription. The downregulation of transcript abundance may involve a GTP concentration dependent inhibition of transcription as discussed above. However, the strong reduction in subunits from complexes that are usually stable like Cyt b₆f suggests that more active processes could also be involved. This idea is supported by previous work showing the high stability of Cyt b₆f in tobacco (Hojka et al., 2014), the selective loss of Cyt b₆f during natural senescence (Krieger-Liszkay et al., 2019; Roberts et al., 1987), and the protease-mediated degradation mechanism for Cyt b₆f and Rubisco that occurs under nitrogen limitation in *Chlamydomonas* (Majeran et al., 2000; Wei et al., 2014). Another argument supporting the idea that ppGpp may act on protein stability is the contrast between reduced transcript and stable protein levels for the nuclear encoded photosynthetic genes Lhcb1 and Lhca1 in low ppGpp plants (Fig. 3A-B, Fig. 5C, Fig. 5- source data 3). This contrast suggests that ppGpp could promote the degradation of these proteins during nitrogen limitation.

PSII architecture was also modified in a ppGpp-dependent manner during nitrogen limitation with an increase in the ratio of PSII antenna to RC (Fig. 3A-D), causing an increase in basal fluorescence and contributing to the decrease in PSII maximal efficiency (Fv/Fm) (Fig. 2A, D). While this phenomenon has similarities to photoinhibition, our data suggest that ppGpp does not increase PSII RC photoinactivation, but instead inhibits the resynthesis of PSII RC via the inhibition of chloroplast transcription. Due to the rapid degradation of photoinactivated PSII RC this leads to an increase in the ratio of PSII antenna to RC.

ppGpp is implicated in photoprotection during nitrogen limitation

During nitrogen deficiency plants must remobilize nitrogen to drive acclimation processes. At the same time, growth arrest leads to a strong decrease in demand for the products of photosynthesis which has the potential to lead to overexcitation and ROS production. Therefore, in addition to remobilizing nitrogen, the plant must also downsize the

photosynthetic machinery in a controlled fashion to avoid producing damaging levels of ROS. Here, we show that ppGpp promotes a reduction in the photosynthetic machinery, a major nitrogen store, and at the same time plays a role in protecting against excess ROS production. We show that ROS increases in the wild type in response to nitrogen deprivation, and that defects in ppGpp biosynthesis lead to greater increases in ROS and cell death (Fig. 1B, D). The limited ROS increase in the wild type is in agreement with a recent study also showing ROS accumulation in the leaves of nitrogen starved plants (Safi et al., 2021). Chloroplasts are the major source of ROS in photosynthetic plant leaves (Rogers and Munné-Bosch, 2016; Domínguez and Cejudo, 2021). The role for ppGpp signaling in the prevention of ROS over-accumulation and tissue death during nitrogen limitation (Fig. 1A-D) is therefore very likely related to its role in downregulating photosynthetic activity (Fig. 2, Fig. 2- figure supplement 2, Fig. 2- figure supplement 4). As part of this process, ppGpp signaling alters the stoichiometry of PSII components, reducing excitation pressure on PSII and thus electron flow in the photosynthetic chain. Note that the reduced Rubisco levels we observe under nitrogen limitation (Fig. 3A, 3B) suggest that PSI may be acceptor side limited due to reduced activity of the Calvin-Benson cycle, which could lead to increased superoxide formation by reduction of molecular oxygen at PSI. Indeed, others have proposed that the reduction in levels of active PSII RC during photoinhibition can protect PSI against overexcitation damage (Tikkanen et al., 2014). Interestingly, the ppGpp-mediated alteration of PSII architecture appears to be independent of the global reduction in photosynthetic proteins during nitrogen limitation: photosynthetic proteins are downregulated in low ppGpp plants without major alterations in PSII activity (Fig. 3A-B, Fig. 2C-D). On the basis of these results, we suggest that one of the functions of ppGpp signaling may therefore be to optimize nitrogen remobilization, notably via the degradation of the most abundant leaf protein Rubisco, modulating at the same time the photosynthetic machinery to prevent overexcitation and photodamage. The most straightforward explanation based on the evidence presented here is that ppGpp acts through a general and coordinated down-regulation of chloroplast gene expression, and further evidence would be needed to determine whether ppGpp is also able to play a more specific role in remodeling photosynthesis.

We demonstrate a new physiological role for ppGpp signaling and offer new insights into the molecular mechanisms underlying its action. The wide range of stresses known to induce ppGpp accumulation (Takahashi et al., 2004) and the association between stress and overexcitation of the photosynthetic electron transport chain (Bechtold and Field, 2018),

580 together suggest that ppGpp signaling may allow plants to anticipate and acclimate to harsh
581 environmental conditions that would otherwise lead to overexcitation of the photosynthetic
582 machinery and oxidative stress. At the same time, our work also highlights important
583 questions about ppGpp function that are not yet fully answered: how does nitrogen limitation
584 activate ppGpp signaling? does ppGpp target a single process (transcription) or several? and
585 does ppGpp signaling interact with known photoprotective pathways (Pinnola and Bassi,
586 2018; Malnoë, 2018)?
587

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (<i>Arabidopsis thaliana</i>)	Col-0	Nottingham Arabidopsis Stock Centre (NASC)	WT1, Columbia, N1093 (NASC)	
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2</i>	Nottingham Arabidopsis Stock Centre (NASC), Coppenhaver et al. (2000)	WT2, N8846 (NASC)	Col-3 ecotype
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/rsh1-1</i>	Nottingham Arabidopsis Stock Centre (NASC), Sugliani et al. (2016), Sessions et al. (2002)	TDNA insertion SAIL_391_E11, N818025 (NASC)	<i>RSH</i> mutant
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/rsh2-1</i>	Nottingham Arabidopsis Stock Centre (NASC), Sugliani et al. (2016), Sessions et al. (2002)	TDNA insertion SAIL_305_B12, N814119 (NASC)	<i>RSH</i> mutant
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/rsh3-1</i>	Nottingham Arabidopsis Stock Centre (NASC), Sugliani et al. (2016), Sessions et al. (2002)	TDNA insertion SAIL_99_G05, N862398 (NASC)	<i>RSH</i> mutant
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/rsh2-1 rsh3-1</i>	Sugliani et al. (2016)	DM-23	<i>RSH</i> mutant
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/rsh_{QM}</i>	Sugliani et al. (2016)	<i>rsh1-1, rsh2-1, rsh3-1, crsh-ami / Qmaii</i>	<i>RSH</i> mutant
Genetic reagent (<i>Arabidopsis thaliana</i>)	Col-0/ OX:RSH1	Sugliani et al. (2016)	OX:RSH1-GFP (10.4)	overexpression line
Genetic reagent	<i>qrt1-2/rsh2-1 rsh3-1</i>	Sugliani et al.	C11	complementation

(<i>Arabidopsis thaliana</i>)	pRSH3:RSH3	(2016)		line
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/</i> <i>rsh2-1 rsh3-1</i> pRSH3:RSH3	Sugliani et al. (2016)	C43	complementation line
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/</i> <i>rsh2-1 rsh3-1</i> pRSH3:RSH3	Sugliani et al. (2016)	C131	complementation line
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/</i> <i>rsh2-1 rsh3-1</i> pRSH3:RSH3	Sugliani et al. (2016)	CX3	complementation line
Genetic reagent (<i>Arabidopsis thaliana</i>)	Col-0/ pOP:MESH	Sugliani et al. (2016)	MESH	DEX inducible MESH
Genetic reagent (<i>Arabidopsis thaliana</i>)	<i>qrt1-2/</i> <i>rsh1-1</i> pRSH1:RSH1	This study.	C110	Materials and methods: Creation of <i>rsh1-1</i> complementation lines
antibody	anti-COXII (rabbit polyclonal)	Agrisera	Ref. AS04 053A	dilution (1:2000)
antibody	anti-LHCA1 (rabbit polyclonal)	Agrisera	Ref. AS01 005	dilution (1:2000)
antibody	anti-LHCB1 (rabbit polyclonal)	Agrisera	Ref. AS01 004	dilution (1:2000)
antibody	anti-PBA1 (rabbit polyclonal)	Abcam	Ref. ab98861	dilution (1:2000)
antibody	anti-PetA (rabbit polyclonal)	Agrisera	Ref. AS08 306	dilution (1:2000)
antibody	anti-PsaD (rabbit polyclonal)	Agrisera	Ref. AS04 046	dilution (1:2000)
antibody	anti-PsbA (rabbit polyclonal)	Agrisera	Ref. AS05 084	dilution (1:2000)
antibody	anti-PsbO (rabbit polyclonal)	Agrisera	Ref. AS05 092	dilution (1:2000)
antibody	anti-PsbS (rabbit polyclonal)	Agrisera	Ref. AS09 533	dilution (1:1000)
antibody	anti-PTOX (rabbit	Uniplastomic	Kindly provided by	dilution (1:2000)

	polyclonal)		X.Johnson	
antibody	anti-SAG12 (rabbit polyclonal)	Agrisera	Ref. AS14 2771	dilution (1:2000)
chemical compound, drug	13C-ppGpp	Kindly provided by J.Bartoli and E.Bouveret		internal standard for ppGpp quantification
chemical compound, drug	13C-GTP	Sigma-Aldrich	710687	internal standard for GTP quantification
chemical compound, drug	15-HEDE	Cayman Chemical	Item No. 37700	internal standard for HOTE quantification
commercial assay or kit	Oasis WAX 1 cc Vac Cartridge	Waters	Ref. 186002491	Nucleotide quantification
chemical compound, drug	Nucleozol	Macherey Nagel	Ref. 740404.200	RNA extraction
chemical compound, drug	4-bromoanisole	Sigma-Aldrich	B56501	RNA extraction
commercial assay or kit	Clean & Concentrator™-25 kit	Zymo Research	Cat. No. : R1017	RNA extraction
commercial assay or kit	Ribo-Zero rRNA Removal Kit (Plant)	Illumina	Ref. MRZPL116	RNA treatment
other	Open FluorCam	Photon System Instruments	FC 800-O/2020-GFP	Chlorophyll fluorescence
software, algorithm	R	R Core Team (2020)		Data analysis
software, algorithm	ggplot2 package	Wickham (2009)		Data analysis
software, algorithm	Rmisc package	Hope (2013)		Data analysis
software, algorithm	boot package	Canty and Ripley, (2021); Davison and Hinkley, (1997)		Data analysis
software, algorithm	rstatix package	Kassambara (2021)		Data analysis
software, algorithm	Rcompanion package	Salvatore (2021)		Data analysis
software, algorithm	prepare_gene_ontology.pl script	This study		Data analysis, available at: https://github.com/c

				ecile- lecampion/gene- ontology-analysis- and-graph
--	--	--	--	--

589
590

Plant materials and growth conditions

Arabidopsis thaliana mutant lines, overexpressors and complementation lines and respective wild-type controls are listed in the Key Resources table. In each experiment the seeds for each line were derived from the same batch of plants grown together. For growth on plates, seeds were surface sterilized with 75% ethanol, rinsed with 100% ethanol, dried and placed in a ten by ten grid pattern on square culture plates containing 45 ml of nitrogen replete (+N) (0.5X Murashige and Skoog salts [Caisson Labs], 1% sucrose, 0.5 g L⁻¹ MES and 0.4% Phytigel [Merck Sigma-Aldrich], adjusted to pH 5.7 with KOH) or nitrogen limiting (-N) (+N medium diluted 1/25 in 0.5X Murashige and Skoog medium without nitrogen [Caisson Labs], 0.5 g/L MES and 0.4% Phytigel [Merck Sigma-Aldrich], adjusted to pH 5.7 with KOH) growth medium (for detailed medium composition see Supplementary File 1). Plates were placed at 4°C for 2 days in the darkness, and then transferred to a 16 h light (at 22°C) / 8 h darkness (at 19.5°C) photoperiod with 80 μmol photons m⁻² s⁻¹ lighting. For growth in quartz sand, seeds were germinated in soil, and then transferred into pots containing quartz sand at 7 days after germination. The plants were then grown under a 8 h light (at 22°C)/16 h dark (at 19.5°C) photoperiod at 18°C / 22°C with 110 μmol photons m⁻² s⁻¹ lighting. Plants were treated weekly with a complete nutrient solution. After 50 days the pots were rinsed, and then treated weekly with 0.5X Murashige and Skoog medium with or without nitrogen for three weeks.

Creation of *rsh1-1* complementation lines

The 9.1 kb genomic *RSH1* sequence including the 3' untranslated region, 5' untranslated region, and 3 kb of upstream sequence containing the promoter was amplified from *Arabidopsis* genomic DNA using Phusion polymerase (New England Biolabs) using flanking primers and then the primers RSH1-F (5'-TCCGTCTTGTCTGAATCAGCT-3') and RSH1-R (5'-TTTCTAGATTTACTTTGGTTTTGTCCA-3') with attB1 / attB2 adapters. The resulting PCR product was then introduced by Invitrogen BP Gateway recombination (Thermofisher) into pDONR207. The entry clone was confirmed by sequencing and recombined by Invitrogen LR Gateway recombination into the binary vector pGWB410 (Nakagawa et al., 2007) which carries a kanamycin resistance cassette for selection in plants. The resulting construct was transferred into *Agrobacterium tumefaciens* (strain GV3101) and used to transform *rsh1-1* plants by floral dipping.

Autoluminescence imaging and quantification of HOTEs

Peroxidated lipids were visualized by autoluminescence imaging as described previously (Birtic et al., 2011) and similar results were observed in four biological replicates. The intensity of the leaf autoluminescence signal is proportional to the amount of lipid peroxides present in the sample, the slow spontaneous decomposition of which produces luminescent species. For quantification of HOTEs, lipids were extracted from about 300 mg of seedlings in methanol / chloroform and analyzed by HPLC-UV as detailed elsewhere (Montillet et al., 2004; Shumbe et al., 2017).

Nucleotide triphosphate quantification

Nucleotides were extracted from about 150 mg of seedlings, and quantified by HPLC-MS/MS using stable isotope labelled ppGpp and GTP standards as described previously (Bartoli et al., 2020).

Chlorophyll fluorescence measurements

Plants were dark adapted for 20 min and chlorophyll fluorescence was measured in a Fluorcam FC 800-O imaging fluorometer (Photon System Instruments). PSII maximum quantum yield (F_v/F_m) was calculated as $(F_m - F_o)/F_m$. For relative electron transfer rate (ETR) measurements plants were exposed to 2 min periods of increasing actinic light intensity, with PSII quantum yield measurements at the end of each period. Relative ETR was calculated as quantum yield X light intensity (photosynthetic photon flux density). Photosynthetic photon flux density was measured using a Universal Light Meter 500 (ULM-500, Walz). All experiments on photosynthetic parameters were repeated two to five times with similar results. Steady state 77K chlorophyll fluorescence measurements were obtained from frozen seedling powder suspended in 85% (w/v) glycerol, 10 mM HEPES, pH 7.5 as described previously (Galka et al., 2012).

Protein extraction and immunoblotting

Protein extraction and immunoblotting were performed on 100 bulked seedlings as described previously (Sugliani et al., 2016), with the addition of a protein precipitation step with 20% TCA after protein extraction. The precipitation step was necessary to remove compounds in nitrogen-limited seedlings that interfered with the bicinchoninic acid assay for determining protein concentration. The following primary antibodies were used against CoxII (Agrisera; polyclonal; ref. AS04 053A), Lhca1 (Agrisera; polyclonal, ref. AS01 005), Lhcb1 (Agrisera; polyclonal, ref. AS01 004), Pba1 (Abcam; polyclonal; ref. ab98861), PetA (Agrisera;

polyclonal, ref. AS08 306), PsaD (Agrisera; polyclonal; ref. AS04 046), PsbA (Agrisera; polyclonal, ref. AS05 084), PsbO (Agrisera; polyclonal; ref. AS05 092), PsbS (Agrisera; polyclonal, ref. AS09 533), PTOX (Uniplastomic, Biviers, France; kindly provided by Xenie Johnson), SAG12 (Agrisera; polyclonal; ref. AS14 2771).

For the analysis of thylakoidal proteins, 100 bulked seedlings were harvested and then ground in liquid nitrogen. Proteins were extracted in an extraction buffer adapted from Pesaresi et al. (2011) (0.4 M sucrose, 10 mM NaCl, 5 mM MgCl₂, 10 mM tricine KOH pH 7.5, 100 mM ascorbate, 0.2 mM PMSF, 5 mM aminocaproic acid). Samples were centrifuged at 1000g for 5 min at 4°C, then the pellets were resuspended in extraction buffer. The samples were centrifuged again at 1000g for 5 min at 4°C, and the resulting chloroplast-enriched pellets suspended in lysis buffer as previously described (Chen and Schröder, 2016) (10 mM Tricine -NaOH pH 7.8) and incubated for 30 min. The samples were centrifuged at 5000g for 5 min at 4°C, and the pellets containing the membrane fraction were re-suspended in lysis buffer. Samples were centrifuged at 1000g for 5 min at 4°C, then the pellets were resuspended in extraction buffer (100 mM Tris pH 6.8, 20% glycerol, 10% SDS). The samples were heated to 40°C for 5 min and then centrifuged at 1300 g for 10 min at room temperature. The supernatant was recovered, and the samples were normalized on the amount of chlorophyll.

Chlorophyll quantification

Pigments were extracted in 80% acetone then separated and quantified by HPLC-UV as described previously (Campoli et al., 2009).

RNA sequencing

RNA sequencing was performed with three biological replicates on 100 bulked seedlings per line grown on +N medium for 8 days or -N medium for 12 days. RNA was extracted from frozen seedling powder with Nucleozol (Macherey-Nagel) with 4-bromoanisole to reduce DNA and anthocyanin contamination. Total RNA was cleaned and concentrated using RNA Clean & ConcentratorTM-25 (Zymo Research) according to the manufacturer's instructions. Genomic DNA was removed by treatment with DNase. RNA-seq libraries were constructed by the POPS platform (IPS2) using the TruSeq Stranded mRNA library prep kit (Illumina) with RiboZero plant (Illumina). Libraries were sequenced in single-end (SE) with a read length of 75 bases for each read on a NextSeq500 (Illumina). Approximately 30 million reads by sample were generated. Adapter sequences and bases with a Q-Score below 20 were

trimmed out from reads using Trimmomatic (v0.36) (Bolger et al., 2014) and reads shorter than 30 bases after trimming were discarded. Reads corresponding to rRNA sequences were removed using sortMeRNA (v2.1) (Kopylova et al., 2012) against the silva-bac-16s-id90, silva-bac-23s-id98, silva-euk-18s-id95 and silva-euk-28s-id98 databases. Read quality checks were performed using FastQC (Version 0.11.5)(Andrews, 2010). The raw data (fastq) was then aligned against the Arabidopsis transcriptome (Araport11_cdna_20160703_representative_gene_model.fa) concatenated with non-coding RNA (TAIR10.ncrna.fa) using Bowtie2 (version 2.2.9)(Langmead and Salzberg, 2012). Default parameters were used. Reads were counted using a modified version of a command line previously described (Van Verk et al., 2013). Differential expression analysis was performed with SARTools (version 1.7.3) (Varet et al., 2016) using edgeR. Study details and Fastq files were deposited at the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>) under accession number PRJEB46181. Workflow and analysis reports for RNAseq data analysis are provided in Supplementary File 2.

Data analysis

The majority of analysis was conducted in R (R Core Team, 2020) and custom annotated R markdown scripts are provided in Supplementary File 3. Graphs were produced using the package ggplot2 (Wickham, 2009) with confidence intervals determined for normally distributed data using the Rmisc package (Hope, 2013), and non-parametric data with bootstrap confidence interval calculation using the boot package (Canty and Ripley, 2021; Davison and Hinkley, 1997). Individual plants were counted as biological replicates, and experiments were repeated at least two times with similar results. For the timecourse in Fig. 3A and the nitrogen deprivation performed on mature plants in Fig. 2- figure supplement 2E single experimental replicates were performed. Statistical analyses were performed on normally distributed data using ANOVA with posthoc Tukey test and non-parametric data using the Kruskal-Wallis test with posthoc Dunn test in the rstatix package (Kassambara, 2021). For categorical data (Fig. 1B,C) significance was calculated using a proportion test and posthoc Fisher test using the Rstatix and Rcompanion packages (Kassambara, 2021; Salvatore, 2021). All statistical tests included adjustments for multiple comparisons. GO enrichment analysis for the RNAseq data was performed using a custom prepare_gene_ontology.pl script (<https://github.com/cecile-lecampion/gene-ontology-analysis-and-graph>, commit c170f8e90323f3061d44a87fcb76d6b0e8b00f63) which automatically uses PANTHER and REVIGO for the identification and simplification of

726 enriched GO terms according to the procedure proposed by Bonnot et al. (2019). Results were
727 plotted using ggplot2.

Acknowledgements

We thank colleagues at the LGBP, Xenie Johnson and Wojciech Nawrocki for critical discussion of the manuscript. We thank Julia Bartoli and Emmanuelle Bouveret for supplying isotope labelled ppGpp. Nucleotide measurements were performed on the IJPB Plant Observatory technological platform, and transcriptomics on the POPS platform which are both supported by Saclay Plant Sciences-SPS (ANR-17-EUR-0007). The work was funded by Agence Nationale de la Recherche (ANR-17-CE13-0005).

Competing interests

The authors declare no competing interests.

References

- Abdelkefi, H., Sugliani, M., Ke, H., Harchouni, S., Soubigou-Taconnat, L., Citerne, S., Mouille, G., Fakhfakh, H., Robaglia, C., and Field, B. (2018). Guanosine tetraphosphate modulates salicylic acid signalling and the resistance of *Arabidopsis thaliana* to Turnip mosaic virus. *Mol. Plant Pathol.* 19: 634–646.
- Anderson, B.W., Fung, D.K., and Wang, J.D. (2021). Regulatory Themes and Variations by the Stress-Signaling Nucleotide Alarmones (p)ppGpp in Bacteria. *Annu. Rev. Genet.* 55: doi: 10.1146/annurev-genet-021821-025827.
- Andrews, S. (2010). FastQC. A quality control tool for high throughput sequence data. Version 0.11.5 <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Atkinson, G.C., Tenson, T., and Hauryliuk, V. (2011). The RelA/SpoT homolog (RSH) superfamily: distribution and functional evolution of ppGpp synthetases and hydrolases across the tree of life. *PLoS One* 6: e23479.
- Avilan, L., Lebrun, R., Puppo, C., Citerne, S., Cuiné, S., Li-Beisson, Y., Menand, B., Field, B., and Gontero, B. (2021). ppGpp influences protein protection, growth and photosynthesis in *Phaeodactylum tricornutum*. *New Phytol.* 230: 1517–1532.
- Avilan, L., Puppo, C., Villain, A., Bouveret, E., Menand, B., Field, B., and Gontero, B. (2019). RSH enzyme diversity for (p)ppGpp metabolism in *Phaeodactylum tricornutum* and other diatoms. *Sci. Rep.* 9: 1–11.
- Bange, G., Brodersen, D.E., Liuzzi, A., and Steinchen, W. (2021). Two P or Not Two P: Understanding Regulation by the Bacterial Second Messengers (p)ppGpp. *Annu. Rev. Microbiol.*
- Bartoli, J., Citerne, S., Mouille, G., Bouveret, E., and Field, B. (2020). Quantification of guanosine triphosphate and tetraphosphate in plants and algae using stable isotope-labelled internal standards. *Talanta* 219: 121261.
- Bechtold, U. and Field, B. (2018). Molecular mechanisms controlling plant growth during abiotic stress. *J. Exp. Bot.* 69: 2753–2758.
- Birtic, S., Ksas, B., Genty, B., Mueller, M.J., Triantaphylidès, C., and Havaux, M. (2011). Using spontaneous photon emission to image lipid oxidation patterns in plant tissues. *Plant J.* 67: 1103–1115.
- Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinforma. Oxf. Engl.* 30: 2114–2120.
- Boniecka, J., Prusinska, J., Dabrowska, G.B., and Goc, A. (2017). Within and beyond the stringent response-RSH and (p)ppGpp in plants. *Planta* 246: 817–842.
- Bonnot, T., Gillard, M.B., and Nagel, D.H. (2019). A Simple Protocol for Informative Visualization of Enriched Gene Ontology Terms. *Bio-Protoc.*: e3429–e3429.
- Campoli, C., Caffarri, S., Svensson, J.T., Bassi, R., Stanca, A.M., Cattivelli, L., and Crosatti, C. (2009). Parallel pigment and transcriptomic analysis of four barley albina and

777 xantha mutants reveals the complex network of the chloroplast-dependent metabolism.
778 Plant Mol. Biol. 71: 173–191.

779 Canty, A. and Ripley, B. (2021). boot: Bootstrap R (S-Plus) Functions. Version 1.3-28
780 <https://cran.r-project.org/web/packages/boot/index.html>.

781 Cashel, M. and Gallant, J. (1969). Two compounds implicated in the function of the RC gene
782 of *Escherichia coli*. Nature 221: 838.

783 Chen, Y. and Schröder, W. (2016). Comparison of methods for extracting thylakoid
784 membranes of *Arabidopsis* plants. Physiol. Plant. 156: 3–12.

785 Copenhaver, G.P., Keith, K.C., and Preuss, D. (2000). Tetrad analysis in higher plants. A
786 budding technology. Plant Physiol. 124: 7–16.

787 Crepin, A., Santabarbara, S., and Caffarri, S. (2016). Biochemical and Spectroscopic
788 Characterization of Highly Stable Photosystem II Supercomplexes from *Arabidopsis*.
789 J. Biol. Chem. 291: 19157–19171.

790 Davison, A.C. and Hinkley, D.V. (1997). Bootstrap Methods and their Application 1st
791 edition. (Cambridge University Press).

792 Devireddy, A.R., Zandalinas, S.I., Fichman, Y., and Mittler, R. (2021). Integration of reactive
793 oxygen species and hormone signaling during abiotic stress. Plant J. 105: 459–476.

794 Domínguez, F. and Cejudo, F.J. (2021). Chloroplast dismantling in leaf senescence. J. Exp.
795 Bot. 72: 5905–5918.

796 Field, B. (2018). Green magic: regulation of the chloroplast stress response by (p)ppGpp in
797 plants and algae. J. Exp. Bot. 69: 2797–2807.

798 Galka, P., Santabarbara, S., Khuong, T.T.H., Degand, H., Morsomme, P., Jennings, R.C.,
799 Boekema, E.J., and Caffarri, S. (2012). Functional analyses of the plant photosystem
800 I-light-harvesting complex II supercomplex reveal that light-harvesting complex II
801 loosely bound to photosystem II is a very efficient antenna for photosystem I in state
802 II. Plant Cell 24: 2963–2978.

803 Garai, S. and Tripathy, B.C. (2018). Alleviation of Nitrogen and Sulfur Deficiency and
804 Enhancement of Photosynthesis in *Arabidopsis thaliana* by Overexpression of
805 Uroporphyrinogen III Methyltransferase (UPM1). Front. Plant Sci. 0.

806 Harchouni, S., England, S., Vieu, J., Aouane, A., Citerne, S., Legeret, B., Li-Beisson, Y.,
807 Menand, B., and Field, B. (2021). Guanosine tetraphosphate (ppGpp) accumulation
808 inhibits chloroplast gene expression and promotes super grana formation in the moss
809 *Physcomitrium* (*Physcomitrella*) patens. bioRxiv: 2021.01.06.425534.

810 Hojka, M., Thiele, W., Tóth, S.Z., Lein, W., Bock, R., and Schöttler, M.A. (2014). Inducible
811 Repression of Nuclear-Encoded Subunits of the Cytochrome b6f Complex in Tobacco
812 Reveals an Extraordinarily Long Lifetime of the Complex. Plant Physiol. 165: 1632–
813 1646.

814 Honoki, R., Ono, S., Oikawa, A., Saito, K., and Masuda, S. (2018). Significance of
815 accumulation of the alarmone (p)ppGpp in chloroplasts for controlling photosynthesis
816 and metabolite balance during nitrogen starvation in Arabidopsis. *Photosynth Res* 135:
817 299–308.

818 Hope, R. (2013). Rmisc: Ryan Miscellaneous. Version 1.5 [https://cran.r-](https://cran.r-project.org/package=Rmisc)
819 [project.org/package=Rmisc](https://cran.r-project.org/package=Rmisc).

820 Ihara, Y., Ohta, H., and Masuda, S. (2015). A highly sensitive quantification method for the
821 accumulation of alarmone ppGpp in Arabidopsis thaliana using UPLC-ESI-qMS/MS.
822 *J Plant Res* 128: 511–8.

823 Imamura, S., Nomura, Y., Takemura, T., Pancha, I., Taki, K., Toguchi, K., Tozawa, Y., and
824 Tanaka, K. (2018). The checkpoint kinase TOR (target of rapamycin) regulates
825 expression of a nuclear-encoded chloroplast RelA-SpoT homolog (RSH) and
826 modulates chloroplast ribosomal RNA synthesis in a unicellular red alga. *Plant J.* 94:
827 327–339.

828 Irving, S.E. and Corrigan, R.M. (2018). Triggering the stringent response: signals responsible
829 for activating (p)ppGpp synthesis in bacteria. *Microbiol. Read. Engl.* 164: 268–276.

830 Ito, D., Ihara, Y., Nishihara, H., and Masuda, S. (2017). Phylogenetic analysis of proteins
831 involved in the stringent response in plant cells. *J Plant Res* 130: 625–634.

832 Kassambara, A. (2021). Rstatix: Pipe-Friendly Framework for Basic Statistical Tests. Version
833 0.7.0 <https://CRAN.R-project.org/package=rstatix>.

834 Kleine, T. et al. (2021). Acclimation in plants – the Green Hub consortium. *Plant J.* 106: 23–
835 40.

836 Kopylova, E., Noé, L., and Touzet, H. (2012). SortMeRNA: fast and accurate filtering of
837 ribosomal RNAs in metatranscriptomic data. *Bioinforma. Oxf. Engl.* 28: 3211–3217.

838 Krieger-Liszkay, A., Krupinska, K., and Shimakawa, G. (2019). The impact of photosynthesis
839 on initiation of leaf senescence. *Physiol. Plant.* 166: 148–164.

840 Langmead, B. and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat.*
841 *Methods* 9: 357–359.

842 Lu, C. and Zhang, J. (2000). Photosynthetic CO₂ assimilation, chlorophyll fluorescence and
843 photoinhibition as affected by nitrogen deficiency in maize plants. *Plant Sci.* 151:
844 135–143.

845 Luo, J., Havé, M., Clément, G., Tellier, F., Balliau, T., Launay-Avon, A., Guérard, F., Zivy,
846 M., and Masclaux-Daubresse, C. (2020). Integrating multiple omics to identify
847 common and specific molecular changes occurring in Arabidopsis under chronic
848 nitrate and sulfate limitations. *J. Exp. Bot.* 71: 6471–6490.

849 Maekawa, M., Honoki, R., Ihara, Y., Sato, R., Oikawa, A., Kanno, Y., Ohta, H., Seo, M.,
850 Saito, K., and Masuda, S. (2015). Impact of the plastidial stringent response in plant
851 growth and stress responses. *Nat. Plants* 1: 15167.

852 Majeran, W., Wollman, F.-A., and Vallon, O. (2000). Evidence for a Role of ClpP in the
853 Degradation of the Chloroplast Cytochrome b6f Complex. *Plant Cell* 12: 137–150.

854 Malnoë, A. (2018). Photoinhibition or photoprotection of photosynthesis? Update on the
855 (newly termed) sustained quenching component qH. *Environ. Exp. Bot.* 154: 123–133.

856 Mizusawa, K., Masuda, S., and Ohta, H. (2008). Expression profiling of four RelA/SpoT-like
857 proteins, homologues of bacterial stringent factors, in *Arabidopsis thaliana*. *Planta*
858 228: 553–62.

859 Montillet, J.-L., Cacas, J.-L., Garnier, L., Montané, M.-H., Douki, T., Bessoule, J.-J.,
860 Polkowska-Kowalczyk, L., Maciejewska, U., Agnel, J.-P., Vial, A., and
861 Triantaphylidès, C. (2004). The upstream oxylipin profile of *Arabidopsis thaliana*: a
862 tool to scan for oxidative stresses. *Plant J.* 40: 439–451.

863 Nakagawa, T. et al. (2007). Improved Gateway binary vectors: high-performance vectors for
864 creation of fusion constructs in transgenic analysis of plants. *Biosci. Biotechnol.*
865 *Biochem.* 71: 2095–2100.

866 Nomura, Y., Izumi, A., Fukunaga, Y., Kusumi, K., Iba, K., Watanabe, S., Nakahira, Y.,
867 Weber, A.P.M., Nozawa, A., and Tozawa, Y. (2014). Diversity in Guanosine 3',5'-
868 Bisdiphosphate (ppGpp) Sensitivity among Guanylate Kinases of Bacteria and Plants
869 *. *J. Biol. Chem.* 289: 15631–15641.

870 Nunes, M.A., Ramalho, J., and Dias, M.A. (1993). Effect of Nitrogen Supply on the
871 Photosynthetic Performance of Leaves from Coffee Plants Exposed to Bright Light. *J.*
872 *Exp. Bot.* 44: 893–899.

873 Ono, S., Suzuki, S., Ito, D., Tagawa, S., Shiina, T., and Masuda, S. (2020). Plastidial
874 (p)ppGpp Synthesis by the Ca²⁺-Dependent RelA–SpoT Homolog Regulates the
875 Adaptation of Chloroplast Gene Expression to Darkness in *Arabidopsis*. *Plant Cell*
876 *Physiol.* 61: 2077–2086.

877 Pausch, P., Steinchen, W., Wieland, M., Klaus, T., Freibert, S.-A., Altegoer, F., Wilson, D.N.,
878 and Bange, G. (2018). Structural basis for (p)ppGpp-mediated inhibition of the
879 GTPase RbgA. *J. Biol. Chem.* 293: 19699–19709.

880 Pesaresi, P. (2011). Studying translation in *Arabidopsis* chloroplasts. *Methods Mol Biol* 774:
881 209–24.

882 Pinnola, A. and Bassi, R. (2018). Molecular mechanisms involved in plant photoprotection.
883 *Biochem. Soc. Trans.* 46: 467–482.

884 R Core Team (2020). R: A language and environment for statistical computing. R Foundation
885 for Statistical Computing, Vienna, Austria. R 4.1.0 [http://www.r-](http://www.r-project.org/index.html)
886 [project.org/index.html](http://www.r-project.org/index.html).

887 Roberts, D.R., Thompson, J.E., Dumbroff, E.B., Gepstein, S., and Mattoo, A.K. (1987).
888 Differential changes in the synthesis and steady-state levels of thylakoid proteins
889 during bean leaf senescence. *Plant Mol. Biol.* 9: 343–353.

890 Rogers, H. and Munné-Bosch, S. (2016). Production and Scavenging of Reactive Oxygen
891 Species and Redox Signaling during Leaf and Flower Senescence: Similar But
892 Different1[OPEN]. *Plant Physiol.* 171: 1560–1568.

893 Ronneau, S. and Hallez, R. (2019). Make and break the alarmone: regulation of (p)ppGpp
894 synthetase/hydrolase enzymes in bacteria. *FEMS Microbiol. Rev.* 43: 389–400.

895 Safi, A. et al. (2021). GARP transcription factors repress Arabidopsis nitrogen starvation
896 response via ROS-dependent and -independent pathways. *J. Exp. Bot.* 72: 3881–3901.

897 Salvatore, M. (2021). rcompanion: functions to support extension education program
898 evaluation in R. Version 2.4.1 [https://cran.r-](https://cran.r-project.org/web/packages/rcompanion/index.html)
899 [project.org/web/packages/rcompanion/index.html](https://cran.r-project.org/web/packages/rcompanion/index.html).

900 Sessions, A. et al. (2002). A high-throughput Arabidopsis reverse genetics system. *Plant Cell*
901 14: 2985–2994.

902 Sharma, M., Kretschmer, C., Lampe, C., Stuttmann, J., and Klösgen, R.B. (2019). Targeting
903 specificity of nuclear-encoded organelle proteins with a self-assembling split-
904 fluorescent protein toolkit. *J. Cell Sci.* 132: jcs230839.

905 Shumbe, L., D'Alessandro, S., Shao, N., Chevalier, A., Ksas, B., Bock, R., and Havaux, M.
906 (2017). METHYLENE BLUE SENSITIVITY 1 (MBS1) is required for acclimation of
907 Arabidopsis to singlet oxygen and acts downstream of β -cyclocitral. *Plant Cell*
908 *Environ.* 40: 216–226.

909 Stent, G.S. and Brenner, S. (1961). A genetic locus for the regulation of ribonucleic acid
910 synthesis. *Proc. Natl. Acad. Sci. U. S. A.* 47: 2005–2014.

911 Sugliani, M., Abdelkefi, H., Ke, H., Bouveret, E., Robaglia, C., Caffarri, S., and Field, B.
912 (2016). An Ancient Bacterial Signaling Pathway Regulates Chloroplast Function to
913 Influence Growth and Development in Arabidopsis. *Plant Cell* 28: 661–679.

914 Takahashi, K., Kasai, K., and Ochi, K. (2004). Identification of the bacterial alarmone
915 guanosine 5'-diphosphate 3'-diphosphate (ppGpp) in plants. *Proc Natl Acad Sci USA*
916 101: 4320–4324.

917 Terashima, I. and Evans, J.R. (1988). Effects of Light and Nitrogen Nutrition on the
918 Organization of the Photosynthetic Apparatus in Spinach. *Plant Cell Physiol.* 29: 143–
919 155.

920 Tikkanen, M., Mekala, N.R., and Aro, E.-M. (2014). Photosystem II photoinhibition-repair
921 cycle protects Photosystem I from irreversible damage. *Biochim. Biophys. Acta BBA*
922 - Bioenerg. 1837: 210–215.

923 Tyystjarvi, E. and Aro, E.M. (1996). The rate constant of photoinhibition, measured in
924 lincomycin-treated leaves, is directly proportional to light intensity. *Proc. Natl. Acad.*
925 *Sci.* 93: 2213–2218.

926 Van Verk, M.C., Hickman, R., Pieterse, C.M.J., and Van Wees, S.C.M. (2013). RNA-Seq:
927 revelation of the messengers. *Trends Plant Sci.* 18: 175–179.

928 Varet, H., Brillet-Guéguen, L., Coppée, J.-Y., and Dillies, M.-A. (2016). SARTools: A
929 DESeq2- and EdgeR-Based R Pipeline for Comprehensive Differential Analysis of
930 RNA-Seq Data. *PLoS ONE* 11: e0157022.

931 Varik, V., Oliveira, S.R.A., Hauryliuk, V., and Tenson, T. (2017). HPLC-based quantification
932 of bacterial housekeeping nucleotides and alarmone messengers ppGpp and pppGpp.
933 *Sci. Rep.* 7: 11022.

934 Verhoeven, A.S., Demmig-Adams, B., and Adams III, W.W. (1997). Enhanced Employment
935 of the Xanthophyll Cycle and Thermal Energy Dissipation in Spinach Exposed to
936 High Light and N Stress. *Plant Physiol.* 113: 817–824.

937 Wei, L., Derrien, B., Gautier, A., Houille-Vernes, L., Boulouis, A., Saint-Marcoux, D.,
938 Malnoë, A., Rappaport, F., de Vitry, C., Vallon, O., Choquet, Y., and Wollman, F.-A.
939 (2014). Nitric Oxide–Triggered Remodeling of Chloroplast Bioenergetics and
940 Thylakoid Proteins upon Nitrogen Starvation in *Chlamydomonas reinhardtii*[W]. *Plant*
941 *Cell* 26: 353–372.

942 Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag: New
943 York).

944 Yamburenko, M.V., Zubo, Y.O., and Borner, T. (2015). Absciscic acid affects transcription of
945 chloroplast genes via protein phosphatase 2C-dependent activation of nuclear genes:
946 repression by guanosine-3'-5'-bisdiphosphate and activation by sigma factor 5. *Plant J*
947 82: 1030–1041.

948

949

List of figures, figure supplements, source data and supplementary information

Figure 1. ppGpp is required for acclimation to nitrogen deprivation.

Figure 1- figure supplement 1. Extended timecourse of nitrogen deprivation.

Figure 1- figure supplement 2. GTP/ppGpp ratios in RSH lines.

Figure 1- source data 1. Source data and statistical test reports for Figure 1 and supplements.

Figure 2. Nitrogen deprivation promotes a ppGpp-dependent drop in photosynthetic capacity.

Figure 2- figure supplement 1. Changes in additional photosynthetic parameters (F_o , F_m and F_o / F_m) for panel 2A.

Figure 2- figure supplement 2. Role of ppGpp in the nitrogen deprivation induced decrease in PSII maximal yield.

Figure 2- figure supplement 3. Changes in additional photosynthetic parameters (F_o , F_m and F_o / F_m) for Figure 2- Figure supplement 2.

Figure 2- figure supplement 4. ppGpp is required for reducing ETR during nitrogen deprivation.

Figure 2- source data 1. Source data and statistical test reports for Figure 2 and supplements.

Figure 3. ppGpp-dependent alterations in the photosynthetic machinery during nitrogen deficiency.

Figure 3- figure supplement 1. ppGpp depletion by MESH also affects abundance of chloroplast proteins.

Figure 3- figure supplement 2. Full 77K chlorophyll fluorescence spectra under nitrogen deprivation.

Figure 3- source data 1. Uncropped immunoblots for 3A.

Figure 3- source data 2. Uncropped immunoblots for 3B.

Figure 3- source data 3. Source data and statistical test reports for Figure 3 and supplements.

Figure 4. ppGpp plays a major role during acclimation to nitrogen deprivation.

Figure 4- source data 1. Differentially accumulating transcripts.

Figure 4- source data 2. GO enrichment analysis.

Figure 4- source data 3. Source data and statistical test reports for Figure 4B.

Figure 5. ppGpp is required for the downregulation of chloroplast gene expression during nitrogen deficiency.

984 Figure 5- source data 1. Chloroplast gene expression.
985 Figure 5- source data 2. Mitochondrial gene expression.
986 Figure 5- source data 3. Source data for Figure 5C including extended figures for multiple
987 photosynthetic complexes.
988
989 Supplementary file 1. Media composition
990 Supplementary file 2. RNA-seq analysis reports. Related to Figure 4.
991 Supplementary file 3. R markdown scripts. Related to all figures.
992

Figure 1- figure supplement 1. Extended timecourse of nitrogen deprivation.
 Images of seedlings grown on nitrogen limiting medium for the indicated number of days. Scale, 3 mm.

1000

1001

1002

1003

1004

1005

1006

1007

Figure 1- figure supplement 2. GTP/ppGpp ratios in RSH lines. The ratio of ppGpp to GTP concentration in seedlings grown for ten days on nitrogen limiting media. Mean +/- 95% CI, n = 4 experimental replicates. Statistical tests shown against respective wild-type controls, * $P<0.05$, ** $P<0.01$.

Figure 2- figure supplement 1. Changes in additional photosynthetic parameters (F_o , F_m and F_o / F_m) for lines shown in panel 2A. Seedlings were grown 8 days on nitrogen replete media (+N) or 12 days on nitrogen limiting (-N) media. Median +/- 95% CI, n = 95-100 seedlings. Statistical tests shown against respective wild-type controls, * $P < 0.05$, ** $P < 0.01$, Kruskal-Wallis test with post-hoc Dunn tests.

Figure 2- figure supplement 3. Changes in additional photosynthetic parameters (Fo, Fm and Fo / Fm) for Figure 2- Figure supplement 2. (A) In seedlings of indicated lines grown 8 days on nitrogen replete media (+N) or 12 days on nitrogen limiting (-N) media. **(B)(C)(D)** In seedlings of indicated lines grown 12 days on -N. C110 is an *rsh1-1* complementation line complemented with the full length *RSH1* (*pRSH1:RSH1*). pOP:MESH is a dexamethasone (DEX) inducible line that expresses a chloroplast targeted ppGpp hydrolase MESH. **(E)** In mature plants grown on quartz sand supplemented with nitrogen replete media (+N) or nitrogen free (-N) media. Median +/- 95% CI, n = 95-100 seedlings (A-D), 6 plants (E). Statistical tests shown against respective wild-type controls, * $P < 0.05$, ** $P < 0.01$. Kruskal-Wallis test with post-hoc Dunn tests (A-D), one-way Anova with post-hoc Tukey test (E).

1036
1037
1038
1039

1040
1041
1042
1043
1044
1045
1046
1047
1048

Figure 2- figure supplement 4. ppGpp is required for reducing ETR during nitrogen deprivation. Relative ETR measurements in different lines grown 8 days on +N media (left) or 12 days on -N media (right). (A) A comparison between OX:RSH1 and the wild-type control. (B) ETR in the dexamethasone (DEX) inducible pOP:MESH line grown on non-inducing medium (0 μM DEX), or inducing medium (10 μM DEX). MESH is a chloroplast targeted ppGpp hydrolase. Median $\pm$ 95% CI, n = 95-100 seedlings.

Figure 3- figure supplement 1. ppGpp depletion by MESH also affects abundance of chloroplast proteins. Immunoblots showing the abundance of the indicated proteins in extracts from 100 bulked seedlings grown in -N for 12 days from two experimental replicates. MESH seedlings were induced by inclusion of dexamethasone (DEX) or not in the growth medium. RbcL was revealed by Coomassie Brilliant Blue. Equal quantities of total protein were loaded. Uncropped immunoblots are shown in Figure 3- source data 3.

1058

1059

1060

1061

1062

1063

1064

1065

1066

Figure 3- figure supplement 2. Full 77K chlorophyll fluorescence spectra under nitrogen deprivation. Emission spectrum of chlorophyll fluorescence at 77°K normalized to the PSII peak at 685 nm. Measurements made on 8 or 12 day-old seedlings grown under +N or -N. Means $\pm$ 95% CI; data from four experimental replicates.

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

50