

**Desiccation resistance differences in *Drosophila* species can be largely explained by variations in cuticular hydrocarbons**

Zinan Wang<sup>1,2,\*</sup>, Joseph P. Receveur<sup>1,2,3</sup>, Jian Pu<sup>1,4</sup>, Haosu Cong<sup>1</sup>, Cole Richards<sup>1</sup>, Muxuan Liang<sup>5</sup>, and Henry Chung<sup>1,2,\*</sup>

<sup>1</sup> Department of Entomology, Michigan State University, East Lansing, MI 48824, USA

<sup>2</sup> Ecology, Evolution, and Behavior Program, Michigan State University, East Lansing, MI 48824, USA

<sup>3</sup> Institute for Genome Sciences, University of Maryland, Baltimore, MD, 21201, USA

<sup>4</sup> College of Agriculture, Sichuan Agricultural University, Chengdu, Sichuan 611130, China

<sup>5</sup> Department of Biostatistics, University of Florida, Gainesville, FL 32611, USA

\* Correspondence: [wangzina@msu.edu](mailto:wangzina@msu.edu) or [hwchung@msu.edu](mailto:hwchung@msu.edu)

**Keywords:** water balance, methyl-branched cuticular hydrocarbons, physiological mechanism, random forest

## Significance

As our planet is becoming more arid due to global warming, preventing dehydration is key to the survival of insects, an essential part of our ecosystem. However, factors that determine how insects may evolve resistance to desiccation are relatively unknown. Using *Drosophila* species from diverse habitats, we showed that variations in the composition of cuticular hydrocarbons (CHCs), a hydrophobic layer found on insects to prevent evaporative water loss, can largely explain desiccation resistance differences. In addition, the evolution of longer methyl-branched CHCs (mbCHCs), underlies the evolution of higher desiccation resistance in this genus. Considering the ubiquity and the diverse compositions of CHCs across insect species, we suggest that evolutionary changes in CHC components may be a common mechanism in insects during the evolution of desiccation resistance and adaptation to various environments.

## Abstract

Maintaining water balance is a universal challenge for organisms living in terrestrial environments, especially for insects, which have essential roles in our ecosystem. Although the high surface area to volume ratio in insects makes them vulnerable to water loss, insects have evolved different levels of desiccation resistance to adapt to diverse environments. To withstand desiccation, insects use a lipid layer called cuticular hydrocarbons (CHCs) to reduce water evaporation from the body surface. It has long been hypothesized that the waterproofing capability of this CHC layer, which can confer different levels of desiccation resistance, depends on its chemical composition. However, it is unknown which CHC components are important contributors to desiccation resistance and how these components can determine differences in desiccation resistance. In this study, we used machine learning algorithms, correlation analyses, and synthetic CHCs to investigate how different CHC components affect desiccation resistance in 50 *Drosophila* and related species. We showed that desiccation resistance differences across these species can be largely explained by variation in CHC composition. In particular, length variation in a subset of CHCs, the methyl-branched CHCs (mbCHCs), is a key determinant of desiccation resistance. There is also a significant correlation between the evolution of longer mbCHCs and higher desiccation resistance in these species. Given that CHCs are almost ubiquitous in insects, we suggest that evolutionary changes in insect CHC components can be a general mechanism for the evolution of desiccation resistance and adaptation to diverse and changing environments.

## Introduction

Maintaining water balance is a universal challenge for organisms living in terrestrial environments where water constantly evaporates from the body surface (Hadley, 1994). Organisms such as insects have a high surface area to volume ratio due to their small sizes, rendering them vulnerable to water loss and desiccation (Gibbs, 2002b; Gibbs and Rajpurohit, 2010; K h sel et al., 2017). Studies using closely related insect species collected worldwide showed that phylogenetically related insect species can have very different levels of desiccation resistance and occupy very distinct habitats, while insects that are phylogenetically distant but dwelling in similar habitats could have similar levels of desiccation resistance (Kellermann et al., 2012; Li et al., 2022; Menzel et al., 2017; Rane et al., 2019). This suggests that extant species have evolved different levels of desiccation resistance to survive in their different habitats. However, as climate change accelerates the expansion of dryland (Huang et al., 2016) and changes aridity in many areas across the globe (Sherwood and Fu, 2014; Shi et al., 2021), it is less understood how insect species, which are integral parts of our ecosystems, can evolve higher levels of desiccation resistance to adapt to the more arid environments.

To determine how insects can adapt to desiccation stress, understanding how insects conserve water and prevent desiccation is crucial. In insects, cuticular water loss is the leading cause of desiccation (Chown et al., 2011; Gibbs and Matzkin, 2001; Wang et al., 2021). In an extreme example, cuticular water transpiration has been found to account for 97% of increased water loss in queens of the harvester ant *Pogonomyrmex barbatus* (Johnson and Gibbs, 2004). To conserve water and prevent desiccation, a general mechanism in insects is the use of a lipid layer on the epicuticle, named cuticular hydrocarbons (CHCs) (Gibbs et al., 2003; Gibbs and Matzkin, 2001). This CHC layer, which can contain more than 100 different compounds on the same individual, provides a hydrophobic barrier against evaporative water loss through the cuticle (Blomquist and Ginzl, 2021). This was first demonstrated when the physical or chemical removal of this layer using abrasive dust and various solvents resulted in increased water loss in various insect species in experiments published almost eight decades ago (Wigglesworth, 1945). In recent years, after the identification of a cytochrome P450 decarbonylase responsible for the synthesis of insect CHCs (Qiu et al., 2012), genetic manipulations of this gene in the fruitfly *Drosophila melanogaster* as well as in aphids and cockroaches, led to almost complete loss of CHCs and significant decrease in desiccation resistance (Chen et al., 2016; Chen et al., 2019; Qiu et al., 2012).

Variations in the composition of this CHC layer has been suggested to contribute to intraspecific variation in desiccation resistance in *D. melanogaster* (Rouault et al., 2004), the Mediterranean dung beetle *Onthophagus taurus* (Leeson et al., 2020), and the Argentine ant *Linepithema humile* (Buellesbach et al., 2018). This is supported by studies manipulating CHC composition either chemically or genetically, in different *Drosophila* species, producing different levels of desiccation resistance (Chiang et al., 2016; Ferveur et al., 2018; Gendron et al., 2014; Koto et al., 2019; Savage et al., 2021). Together, these studies suggest that different components in the CHC layer can influence desiccation resistance.

The varying ability of these CHC components to prevent desiccation depends on its chemical structure, which in turn determines its melting temperatures (Gibbs and Pomonis, 1995; Gibbs, 1998; 2002a). The melting temperature of the hydrocarbon is positively correlated with its water-proofing properties and contribution to desiccation resistance (Blomquist and Ginzl, 2021; Gibbs and Rajpurohit, 2010). In *Drosophila* and most other insects, CHCs range in lengths between approximately 21 to 50 carbons, and consist of linear alkanes (*n*-alkanes), alkenes (monoenes and dienes), and methyl-branched alkanes (Blomquist and Ginzl, 2021). Among them, *n*-alkanes have the highest melting temperature, followed by methyl-branched alkanes (mbCHCs), monoenes, and dienes. Increase in CHC chain length can also increase melting temperature and potentially leads to higher desiccation resistance (Gibbs and Pomonis, 1995). This is consistent with laboratory selection experiments in *D. melanogaster* selecting for increased desiccation resistance resulted in longer carbon-chain CHCs in the desiccation selected flies than the control flies (Gibbs et al., 1997; Kwan and Rundle, 2010). Longer CHC chain length is also correlated with climatic factors such as higher temperature and lower precipitation (Jezovit et al., 2017; Rouault et al., 2004). These climatic factors are associated with desiccation resistance in *Drosophila* species (Hoffmann, 2010; Hoffmann and Weeks, 2007; Kellermann et al., 2018).

While the studies mentioned above showed strong evidence that CHCs and variations in CHC compositions are associated with desiccation resistance in insects, no study has investigated the extent that CHC variation can determine desiccation resistance and whether we can identify the important CHC components that underlie variation in desiccation resistance across different species. Several large scale studies produced large datasets in determining the evolution of desiccation resistance across closely related *Drosophila* species (Kellermann et al., 2018; Kellermann et al., 2012; Matzkin et al., 2009) and ant species (Bujan et al., 2016; Hood and Tschinkel, 1990). Other studies produced large datasets in CHC variation across species (Khallaf et al., 2021; Menzel et al., 2017; Nunes et al., 2017; Van Oystaeyen et al., 2014) and focused on the communication aspects of CHCs. However, variations in CHCs and differences in desiccation resistance have not been experimentally connected in a phylogenetic framework. Analyzing CHC profiles and desiccation resistance across closely related species measured at the same condition can determine important CHC components that could predict various levels of desiccation resistance. Understanding the mechanisms underlying desiccation resistance and examining the evolution of these traits is key to predicting the species' responses in facing drier environments that could result from future climate change (Chown et al., 2010).

In this study, we used a cohort-based experimental design of 50 *Drosophila* and related species, and experimentally determined their CHC compositions and desiccation resistance under similar conditions. Using a random forest machine learning algorithm, we built decision trees connecting CHC components and desiccation resistance in the 50 species and tested their correlation. First, we determined that CHC variations can largely explain differences in desiccation resistance across these species. Second, we identified a subset of CHCs, the methyl-branched CHCs (mbCHCs), as the most important CHC components that can determine desiccation resistance across these species. Third, we



determined that the evolution of longer mbCHCs is also significantly correlated with evolution of higher desiccation resistance in *Drosophila* species. Taken together, our study showed that mbCHCs are a key predictive component of desiccation resistance in *Drosophila* species and suggests that evolution in CHC components can be a general mechanism for higher desiccation resistance and adaptation to more arid environments.

## Results

### A cohort-based experimental design for determining correlations between CHCs and desiccation resistance

To investigate how CHC variation across an evolutionary gradient affects desiccation resistance, for our experiments, we selected 46 *Drosophila* species representing both the *Sophophora* subgenus and the *Drosophila* subgenus, as well as three *Scaptodrosophila* species and one *Chymomyza* species, (**Figure 1-figure supplement 1**). As the assays for CHCs and desiccation resistance cannot be performed on the same individuals, we used a cohort-based experimental design (**Figure 1**) where subsets of each cohort (5-6 per species) were used for GC-MS determination of CHC profiles, desiccation resistance assays, and measuring body weight of the F1 progeny for each sex. Therefore, all measurements were performed at the cohort level.

Desiccation resistance varied across these 50 species with desert-dwelling species from the repleta group showing the highest desiccation resistance (e.g., *D. mojavensis* males:  $58.2 \pm 4.7$  hours) and species from melanogaster group showing the lowest desiccation resistance (e.g., *D. mauritiana* males:  $2.7 \pm 0.2$  hours) (**Figure 2, Figure 2-figure supplement 1**), consistent with the findings of previous research (Kellermann et al., 2012; Matzkin et al., 2009). GC-MS analyses of the CHCs in these 50 species detected five types of CHCs with different carbon-chain lengths and quantities, including linear alkanes (*n*-alkanes), methyl-branched alkanes (mbCHCs; the methyl branch is on the second carbon), monoenes (with one double bond), dienes (with two double bonds), and trienes (with three double bonds) (**Figure 2-figure supplement 2**). *n*-alkanes, which have the highest melting temperatures, were only present in species from the melanogaster group, some of which have the lowest desiccation resistance among the species tested in this study. This suggests that *n*-alkanes may not make a general contribution to desiccation resistance across *Drosophila* species. Trienes, which have the lowest melting temperatures, were only present in eight species with low to moderate quantities. The other three types of CHCs, the mbCHCs, monoenes and dienes, were observed in most species tested in our study.

### Higher relative quantities of CHCs are not primarily responsible for higher desiccation resistance

We first sought to determine if higher desiccation resistance in flies could be due to having higher amounts of CHCs. One caveat in the analysis between absolute CHC quantity and desiccation resistance is a possible correlation between CHC quantities and body size: species with larger body size may

possess higher quantities of CHCs. This can lead to potential biases when directly determining correlations between CHC quantities and desiccation resistance. Species with higher body weight can also have higher absolute water content and a lower surface area to volume ratio due to their larger size, which may also lead to higher desiccation resistance (Wang et al., 2021).

Our initial analyses showed a significant positive correlation (Females:  $r = 0.4$ ,  $P < 0.001$ ; Males:  $r = 0.5$ ,  $P < 0.001$ ) between body weight and desiccation (**Figure 3A**). We also found a significant positive correlation between body weight and total amount of CHCs (Female:  $r = 0.7$ ,  $P < 0.001$ ; Male:  $r = 0.7$ ,  $P < 0.001$ ) (**Figure 3B**), suggesting that variation in the body weight (or size) of different species may be a confound in determining the relationship between CHC quantity and desiccation resistance. To take this confound into consideration, we normalized the total CHC quantity by dividing by the body weight for each species. This shows that CHCs account for 0.02 to 0.5% of the total body weight of different species (**Figure 3-figure supplement 1**). Analyses between the normalized CHC quantity and desiccation resistance showed no correlation between these two variables in males (Male:  $P = 0.1$ ) and positive correlation in females (Female:  $r = 0.1$ ,  $P = 0.03$ ) (**Figure 3C**). The low value of  $r$  (0.1) indicates a weak correlation between CHC quantities and desiccation resistance in females. This suggests that having higher amounts of CHCs only has a limited contribution to higher desiccation resistance.

#### **mbCHCs are important determinants of desiccation resistance**

As total CHC quantity is not a main contributor to desiccation resistance in our study, we hypothesized that the composition of CHC profiles may be important for desiccation resistance. The CHC layer of each *Drosophila* species is composed of different CHC components, each with different levels of abundance, so we used “beta diversity” to represent this variation in the CHC composition of our sampled species, similar to what is used in microbiome analysis. This allows us to test the correlation between the compositional variation and desiccation resistance for the sampled *Drosophila* species. We then applied a permutational multivariate analysis of variance (PERMANOVA) to test whether the composition of CHCs differed across desiccation resistance (Anderson et al., 2006). Our results showed that the beta diversity of CHC compositions differs significantly across the increasing desiccation resistance ( $r^2 = 0.1$ ,  $P < 0.001$ ; **Figure 4-figure supplement 1**), suggesting that CHC composition is important for desiccation resistance. We further sought to identify individual components of CHC profiles that are important determinants of desiccation resistance. To test if variation in CHC components can be used to determine desiccation resistance, we applied a random forest regression method that uses decision trees to connect the variables of interest (Liaw and Wiener, 2002; Svetnik et al., 2003), e.g., CHC composition and desiccation resistance, and identified the importance of individual CHC components to predict desiccation resistance. In this analysis, we treated the CHC profiles of each species and sex as an individual dataset, giving us 100 datasets (50 species x 2 sexes) with 5 to 6 individual CHC profiles each. We then correlated the decision trees that generated from the CHC composition with desiccation resistance. The prediction process can identify key CHC components that are important contributors to desiccation resistance.

Random forest modeling of CHC composition was able to explain 85.5% of the variation in desiccation resistance (Out of Bag Estimate: Root Mean Square Error 'RMSE' = 4.5) (**Figure 4A**). Models built with a 70:30 training/test split ( $n = 382$ ,  $n = 164$ ) performed similarly to models built with out of bag estimate (RMSE = 5.4, **Figure 4-figure supplement 2**). Four mbCHCs were identified as having the highest contribution to predicting desiccation resistance in the regression model (listed in decreasing importance: 2MeC30, 2MeC28, 2MeC32, and 2MeC26), while many other CHCs did not substantially contribute to predicting desiccation resistance (**Figure 4B**). These results suggest that CHCs, and mbCHCs in particular, are important predictors of desiccation resistance in *Drosophila* species.

### **Longer mbCHCs are correlated with higher desiccation**

We further sought to determine how mbCHCs contribute to desiccation resistance. As each species produces different combinations of these mbCHCs that are correlated with each other (**Figure 5-figure supplement 1**), we did not perform correlation analyses on these mbCHCs with a single regression model due to multicollinearity issues (Belsley et al., 2005). Instead, we performed correlation analyses between each of the most important CHCs identified from the random forest modeling, 2MeC26, 2MeC28, 2MeC30, and 2MeC32, and desiccation resistance. We found that the quantity of 2MeC26 negatively correlates with desiccation resistance in males ( $r = -0.4$ ,  $P < 0.001$ ) but no significant correlation was identified in females ( $P = 0.1$ ) (**Figure 5A**). Significant positive correlations were observed between quantities of the longer mbCHCs and desiccation resistance for females (2MeC28:  $r = 0.4$ ,  $P < 0.001$ ; 2MeC30:  $r = 0.2$ ,  $P = 0.009$ ; 2MeC32:  $r = 0.3$ ,  $P < 0.001$ ) and males (2MeC28:  $r = 0.4$ ,  $P < 0.001$ ; 2MeC32:  $r = 0.4$ ,  $P < 0.001$ ) (**Figure 5B-D**). However, no significant correlation was observed between quantity of 2MeC30 and desiccation resistance in males (**Figure 5C**). These results suggested the production of longer mbCHCs could play a role in higher desiccation resistance.

### **Longer saturated CHCs can confer higher desiccation resistance in wild-type *D. melanogaster***

To investigate the above results suggesting that longer mbCHCs could underlie higher desiccation resistance in *Drosophila* species, we performed desiccation assays on *D. melanogaster* (*attP40* strain) coated with synthetic mbCHCs of different lengths (2MeC26, 2MeC28, and 2MeC30). The *D. melanogaster attP40* strain is a common lab strain used for transgenesis with a wild-type CHC profile. While previous studies showed that a mixture of 2MeC26, 2MeC28, and 2MeC30 can slightly increase desiccation resistance in *D. melanogaster* flies without CHCs, experiments were not performed on individual mbCHCs for desiccation resistance (Krupp et al., 2020). Our desiccation assays showed that the coating of 2MeC26 did not increase desiccation resistance in both sexes (*post hoc* comparison with Tukey's method followed by one-way ANOVA. Female:  $P = 0.07$ ; Male:  $P = 1.0$ ), while 2MeC30 significantly increased desiccation resistance in female *D. melanogaster* ( $t = 11.2$ ,  $P < 0.001$ ), and both 2MeC28 ( $t = 4.6$ ,  $P < 0.001$ ) and 2MeC30 ( $t = 6.9$ ,  $P < 0.001$ ) significantly increased desiccation resistance in male *D. melanogaster* (**Figure 5E**). There is a positive correlation with longer mbCHC length coated

and higher desiccation resistance (**Figure 5-figure supplement 2**), suggesting that having higher quantities of longer mbCHCs can lead to higher desiccation resistance. As a control, we repeated the same experiment with synthetic *n*-alkanes (C23, C25, C27, C29, and C31). Coating with *n*-alkanes on the *attP40* strain led to increases in desiccation resistance, which is positively correlated with carbon-chain length, similar to the mbCHCs experiments, suggesting that longer chain CHCs, with higher melting temperatures, can confer higher desiccation resistance when coated on the *attP40* strain (**Figure 5-figure supplement 3, Figure 5-figure supplement 4**).

The coating of synthetic mbCHCs and alkanes on the *attP40* strain showed that adding more CHCs to a strain with a wild-type CHC profile can increase desiccation resistance but did not show if these individual CHCs are sufficient to confer higher desiccation resistance. When these coating experiments (three mbCHCs, five *n*-alkanes) were individually repeated on *D. melanogaster* flies without CHCs (CHC-), we observed that while all mbCHCs and *n*-alkanes are able to confer higher desiccation resistance when coated on female CHC- flies, only a subset of these CHCs are able to confer resistance on male CHC- flies (**Figure 5-figure supplement 5**). In addition, only a positive correlation between longer mbCHCs and higher desiccation was observed in female CHC- flies. There is no correlation between mbCHC length and higher desiccation in male CHC- flies and no correlation between *n*-alkane length and higher desiccation resistance in female and male CHC- flies (**Figure 5-figure supplement 6, Figure 5-figure supplement 7**). We noted that coating mbCHCs and alkanes only produced a small increase in desiccation resistance in CHC- flies, compared to coating these CHCs on the *attP40* strain. This is similar to a previous study by Krupp et al. (2020) where the authors showed similar small increases in desiccation resistance when only *n*-alkanes or mbCHCs are coated on CHC- flies, compared to the larger increase in desiccation resistance when wild-type extracts of CHCs were coated. Taken together, these results suggest that CHCs with higher melting temperatures such as longer mbCHCs or *n*-alkanes contribute to higher desiccation resistance when in a blend of CHCs but not sufficient to confer high desiccation resistance by itself. This is consistent with the hypotheses from multiple papers that the cuticular lipid layer works as a blend or a mixture rather than as individual components (Gibbs, 1998; Menzel et al., 2019; Wigglesworth, 1945). Although we showed that both longer mbCHCs and *n*-alkanes can confer higher desiccation resistance when coated on *D. melanogaster* with a wild-type profile, as *n*-alkanes are mostly absent in the *Drosophila* genus, our experiments suggest a causal relationship between longer mbCHCs and higher desiccation resistance in *Drosophila*.

### **Evolution of mbCHCs underlies variation in desiccation resistance in *Drosophila* species**

We have shown that mbCHCs are important determinants of desiccation resistance in *Drosophila* species and experimentally coating insects with longer mbCHCs confers higher desiccation resistance. To further determine whether the evolution of mbCHCs underlies the variation in desiccation resistance in *Drosophila* species, we first examined the evolutionary trajectory of mbCHCs and then tested the correlation between mbCHCs and desiccation resistance with phylogenetic correction. We mapped the

evolution of mbCHC composition in the *Drosophila* genus using ancestral trait reconstruction and tested the phylogenetic signal using Pagel's  $\lambda$  (Freckleton et al., 2015; Goolsby et al., 2017). Ranging from 0 to 1, Pagel's  $\lambda$  measures the extent to which the variance of the trait can be explained by the phylogeny and therefore determines the degree of association between the trait evolution and the phylogeny (Lynch, 1991). If  $\lambda$  is closer to 1, the evolution of the trait has a higher association with the phylogeny. Phylogenetic signals in mbCHCs were detected for both females and males with  $\lambda = 0.75$  and 0.83, respectively (**Supplementary File 1a**). This suggests a moderate to strong association between mbCHC evolution and phylogeny. The derived ancestral state for mbCHCs in the *Drosophila* genus suggest that 2MeC28 and 2MeC30 are the major mbCHCs (**Figure 6-figure supplement 1**). The evolutionary trajectory of mbCHC composition showed repeated evolution in the lengths of mbCHCs, as well as their quantities. Independent evolution of longer mbCHCs was observed in four clades of species, including the willistoni, nasuta, and repleta groups, and the ananassae subgroup (including *D. ananassae*, *D. pseudoananassae*, and *D. bipectinata*), while shorter mbCHCs were mainly observed in several species in the melanogaster subgroup (e.g., *D. melanogaster*, *D. simulans*, and *D. teissieri*) (**Figure 2, Figure 6-figure supplement 1**).

Desiccation resistance also has a strong association with the phylogeny of *Drosophila* species (**Supplementary File 1b**) (Kellermann et al., 2018; Kellermann et al., 2012). We sought to determine if the variation of desiccation resistance could be explained by the evolution of mbCHCs in these species. We used a Phylogenetic Generalized Linear Square (PGLS) model to test for the correlation between mbCHCs and desiccation resistance when controlling the effects from the phylogenetic relationship of the species (Grafen, 1989; Mundry, 2014). Since mbCHCs are produced in a linear pathway with each species producing several different mbCHCs, we selected the mbCHC with the longest carbon-chain length in the species as the proxy to represent the lengths of mbCHCs in each species. We incorporated the quantity, length, and their interaction in the PGLS model for the longest mbCHC. The interaction term between the length and quantity of the longest mbCHC in the model can determine how the two variables combinatorically affect desiccation resistance. PGLS modeling showed, after correcting for phylogenetic effects, the higher quantity and longer length of the longest mbCHC both affect desiccation resistance (interaction term, Female:  $t = 3.5$ ,  $P < 0.001$ ; Male:  $t = 2.2$ ,  $P = 0.03$ ) (**Supplementary File 1c; Figure 6**). This suggests that the synthesis of mbCHCs with longer carbon-chain lengths could be a common mechanism underlying the evolution of higher desiccation resistance.

## Discussion

Reducing evaporative water loss through the cuticle using a layer of CHCs is one of the most important evolutionary innovations in insects that allows many species to survive and thrive in diverse and arid habitats. We showed that CHC composition can account for 85.5% of the variation in desiccation resistance in the 50 *Drosophila* and related species in this study. This suggests that CHC composition may be highly predictive of desiccation resistance. Algorithmic ranking in importance of CHC components

using a random forest machine learning model showed that mbCHCs have the highest contribution to determining desiccation resistance. Importantly, higher amounts of longer mbCHCs are important in desiccation resistance. This is consistent with previous studies showing that mbCHCs have higher melting temperatures than the other commonly present types of CHCs (monoenes and dienes) and longer CHC leads to higher melting temperature (Gibbs and Pomonis, 1995). Our overall analyses support the conclusion that the evolution of longer mbCHCs is correlated with higher desiccation resistance in our dataset, but we observed minor differences between males and females (**Figures 3 and 5**). In particular, while higher quantities of 2MeC28 and 2MeC32 are both positively correlated with higher desiccation resistance in both males and females (**Figure 5B, D**), there are differences between the sexes regarding 2MeC26 and 2MeC30 (**Figure 5A, C**). There could be several underlying reasons for this. Firstly, across our dataset, CHCs are sexually dimorphic in many species, leading to overall differences in the correlation analyses between the sexes. Secondly, other physiological differences, such as size differences, may contribute to the differences between males and females. In many species such as *D. melanogaster*, females are larger than males and this may contribute to the higher basal desiccation resistance (when CHCs are taken out of the equation) of female CHC- flies ( $3.6 \pm 0.1$  h) compared to that of male CHC- flies ( $2.1 \pm 0.1$  h) (**Figure 5E**). While these are minor differences in the overall correlation analyses between males and females, the synthetic CHC coating experiments showed that coating of longer mbCHCs confers higher desiccation resistance in both males and females.

### **mbCHCs and desiccation resistance**

Previous studies showed support for mbCHCs in desiccation resistance in *Drosophila*. RNAi knockdown of a methyl-branched specific fatty acid synthase (*mFAS*) in *D. serrata* eliminates almost all mbCHCs and leads to significant decrease in desiccation resistance which could be partially rescued with synthetic mbCHCs (Gendron et al., 2014). Its sibling species, the rainforest desiccation sensitive *D. birchii*, have only trace amounts of mbCHCs (Howard et al., 2003) and is unable to evolve desiccation resistance despite strong laboratory selection over many generations (Hoffmann et al., 2003). The presence of mbCHCs in most *Drosophila* and related species leads to another question: Is this a general rule for insects using mbCHCs to develop desiccation resistance? A high proportion of very-long-chain mbCHCs with a single methyl branch have been reported in many insect species from different taxa that dwell in desert environments, such as the desert tenebrionid beetle *Eleodes armata* (Hadley, 1977), the desert ants *Cataglyphis niger* and *Pogonomyrmex barbatus* (Johnson and Gibbs, 2004; Soroker and Hefetz, 2000), and the desert locust *Schistocerca gregaria* (Heifetz et al., 1998). Although a small proportion of *n*-alkanes was also reported in some of these species, this suggests the use of mbCHCs to minimize water evaporation may be a common mechanism for insects in extremely dry environments.

Ancestral trait reconstruction showed that the derived ancestral status for mbCHCs in the *Drosophila* genus are 2MeC28 and 2MeC30 as the major mbCHCs (**Figure 6-figure supplement 1**). As there are many extant species that do not have 2MeC30, this suggests that the ability to synthesize

longer mbCHCs may be lost during evolution, especially for species that do not inhabit arid environments and have low levels of desiccation resistance such as *D. melanogaster* and *D. biarmipes*. On the contrary, for species that dwell in extremely arid environments such as *D. mojavensis* and *D. arizonae*, higher quantities of 2MeC30 and an even longer mbCHC, 2MeC32, are observed. The observation is consistent with the results from our PGLS modeling, showing significant correlation between having higher quantity of longer mbCHCs and higher desiccation resistance in the *Drosophila* genus.

### **Constraints in evolution of CHCs and desiccation resistance**

Why are *n*-alkanes, which have the highest melting temperature and potentially better water proofing properties, not prevalent in species in our studies? A hypothesis that could explain this is the competition on the common precursors between the synthesis of linear unsaturated CHCs (monoenes and dienes) and *n*-alkanes. In insects, the synthesis of all CHCs start from the fatty acyl CoA synthesis pathway. The pathway is split where a cytosolic Fatty Acid Synthase (cFAS) will synthesize all the linear CHCs (alkanes, monoenes, dienes), while mFAS will synthesize mbCHCs (**Figure 5-figure supplement 8**) (Chung and Carroll, 2015; Holze et al., 2021). The pathway suggests that the synthesis of alkanes, monoenes, and dienes are in competition. As many monoenes and dienes function in *Drosophila* as contact pheromones in regulating different behaviors such as mating and aggression (Blomquist and Ginzl, 2021; Chertemps et al., 2007; Chertemps et al., 2006; Chung and Carroll, 2015; Krupp et al., 2008; Wang et al., 2011), the synthesis for these CHCs may compete with the synthesis of *n*-alkanes. In contrast, the synthesis of mbCHCs is in another part of the pathway. This suggests that there are potential constraints in the roles of *n*-alkanes in desiccation resistance in *Drosophila* due to competition with CHCs which function as signaling molecules, while the synthesis of mbCHCs is relatively unaffected by other products of the pathway (**Figure 5-figure supplement 8**) (Chung and Carroll, 2015). We speculate that in *Drosophila* and related species, the use of mbCHCs for modulating desiccation resistance may avoid the conflict between surviving desiccation and chemical signaling.

If there are no biosynthetic constraints to the synthesis of mbCHCs, would the evolution of longer mbCHCs be a general mechanism in the evolution of higher desiccation for species adapting to more arid climates? We suggest that this may be the case for many species, but for some species that use the shorter mbCHCs such as 2MeC26 and 2MeC28 as signaling molecules, such as *D. serrata* (Chenoweth and Blows, 2005; Gendron et al., 2014) and the longhorned beetle, *Mallodon dasystemus* (Spikes et al., 2010), natural selection and sexual selection may have opposing effects on the lengths of these mbCHCs. In *D. serrata*, having higher amounts of 2MeC26 leads to higher mating success in male *D. serrata* (Gendron et al., 2014). *Northern* populations of this species produce lower amounts of 2MeC26 but higher amounts of 2MeC30, due to a polymorphism in a fatty acyl-CoA reductase gene that shifts to an overall production of longer CHCs. As CHCs are sexually monomorphic in *D. serrata*, this polymorphism led to lower mating success in male *northern* flies and higher desiccation resistance in female *northern* flies compared to their common counterparts (Rusuwa et al., 2022). The sexually antagonistic effect due

to the functions of the different mbCHCs may affect their evolutionary response to desiccation in many species.

### **CHC variation and desiccation resistance across insects**

How much of what we learnt in this study is applicable broadly to natural populations of *Drosophila* and other insect species? The use of fly strains from the stock center reared on a common cornmeal artificial medium and the same abiotic factors rather than wild-caught flies allowed us to standardize this large-scale experiment. Previous studies have shown that diet (Fedina et al., 2012), temperature (Rajpurohit et al., 2021), and mating status (Everaerts et al., 2010) can affect CHC profiles in *D. melanogaster*, therefore the CHC profiles that we obtained in our studies may not be representative of the CHC variations present in some of these species in their natural habitats. However, these differences in CHC profiles are mostly quantitative, i.e., different quantities of the same CHCs, rather than different CHCs. Similarly, CHC variations between different natural populations of the same *Drosophila* species are also largely quantitative than qualitative (Dembeck et al., 2015; Rusuwa et al., 2022). Therefore, stock center strains would be representative of the qualitative CHC variations between species and our conclusions we reach in this study may be extended to wild populations. The use of standardized conditions also preclude the selection of species that require specialized media and conditions (e.g., species from the Hawaiian radiation and mycophagus species). While these are some of the limitations of our study, the standardized conditions allow us to reduce the number of confounds in our analyses for the correlation between CHC variation and desiccation resistance. As our study focused on a broad sampling of *Drosophila* and related species from different habitats spanning millions of years of evolution, the overall interpretation of the data and the broad conclusion that we reached regarding the correlation between the evolution of mbCHCs and desiccation resistance is likely to extend to natural populations of the *Drosophila* genus.

Constraints in the evolution of CHCs due to their roles in many insect species as contact pheromones are likely to affect their evolution in response to desiccation resistance. In this study, we showed that the evolution of mbCHCs are major determinants of desiccation resistance across *Drosophila* species, as *n*-alkanes shared the same branch in the fatty acyl-CoA pathway as monoenes and dienes (**Figure 5-figure supplement 8**), many of which function as contact pheromones in *Drosophila* (Ferveur, 2005). In other insect orders such as Lepidoptera, where volatile non-CHC pheromones are predominantly used for chemical communication (Ando et al., 2004), constraints on each class of CHCs may be different from *Drosophila*. Likewise, across eusocial hymenopterans (Van Oystaeyen et al., 2014), saturated CHCs such as *n*-alkanes and mbCHCs are used as queen pheromones and are likely to be under constraints in the evolution of desiccation resistance in these species. Finally, in some rare examples such as dragonflies where very long-chain methyl ketones and aldehydes are the major cuticular wax components (Futahashi et al., 2019), the evolution of CHCs may not be a major factor in the evolution of desiccation resistance. While we limit our conclusions that the



synthesis of mbCHCs with longer carbon-chain lengths is a common mechanism underlying the evolution of higher desiccation resistance to *Drosophila*, the evolution of cuticular components that modulate the melting temperature of the cuticular lipid layer can reduce water loss and increase desiccation resistance. We suggest that the evolution of CHC components in insects can be largely predictive of desiccation resistance between closely related species.

## Materials and Methods

### *Drosophila* species and strains

In this study, 46 *Drosophila* species, as well as three *Scaptodrosophila* species and one *Chymomyza* species were either obtained from the National Drosophila Species Stock Center (NDSSC) or were gifts from various colleagues. Details are listed in **Supplementary File 1d**. The *D. melanogaster* *y w; attP40* strain and UAS-*Cyp4g1* RNAi strain were obtained from the Perrimon lab and the Bloomington *Drosophila* Stock Center, respectively. The 5'*mFAS*-GAL4 strain was constructed by cloning the oenocyte enhancer fragment of the *mFAS/FASN2* (CG3524) gene into the GAL4 vector *pBPGUw* (Addgene #17575) using primers 5'CG3524-TOPO-F (5'-CACCCCGCGGCGTGTATTGAACC-3' and 5'CG3524-TOPO-R (5'-CTTGTTGCGCAGACAGACTG-3'), before injecting into the *D. melanogaster* strain 51C and integrated into the genome using the *PhiC31* integrase system. All species were reared on standard cornmeal medium (Flystuff 66-121 Nutri-Fly® Bloomington Formulation). The phylogenetic relationship of all 50 species in this study was adapted from Finet et al. (2021). CHC- *D. melanogaster* were generated by crossing the 5'*mFAS*-GAL4 driver line (which expresses GAL4 in adult oenocytes) with the UAS-*Cyp4g1* RNAi strain (**Figure 5-figure supplement 9**).

### Experimental design

To investigate the contribution of cuticular hydrocarbons to desiccation resistance, a cohort-based design was used. For each species, five to six cohorts were established and three measurements were conducted for each sex of the F1 progeny, including desiccation resistance, cuticular hydrocarbons, and body weight (**Figure 1**). Each cohort was treated as a biological replicate. Each cohort in each species was established by pooling five females and five males on a standard cornmeal medium in the environmental chambers set at 25°C and a 12L:12D photoperiod. To maximize the food and spatial availability and minimize competition between the F1 progenies (Mueller, 1988), the parent flies in each cohort were transferred to fresh food after 5 days. F1 progeny flies were collected daily following emergence, separated by sex, and maintained on fresh cornmeal medium. All flies used for measurements were four- to five-days-old.

### Desiccation resistance assays

Desiccation resistance assays were performed in a randomized and blinded manner with the setup consistent with a previously published protocol (**Figure 1**) (Gendron et al., 2014). Briefly, in each cohort,

ten adults of the same sex were subjected to the assay setup containing 10 grams of silica gel (S7500-1KG, Sigma-Aldrich, St. Louis, MO). After the assays were assembled, all the setups were randomly arranged with a number assigned. Mortality of the flies was recorded hourly after two hours. For each cohort, one vial was scored and the average time in hours until all flies died was recorded as desiccation resistance. Desiccation resistance assays for the 50 *Drosophila* and related species were conducted at 25°C and with a 12L:12D photoperiod, while the desiccation assays for the CHC coating experiments were conducted at 27°C (GAL4/UAS) and a 12L:12D photoperiod.

### **Cuticular hydrocarbon analyses**

Cuticular hydrocarbons (CHCs) were extracted and analyzed using GC-MS following previously published protocols (Lamb et al., 2020; Savage et al., 2021). Five flies of the same sex (four- to five-days-old) from each cohort were soaked for 10 min in 200 µl hexane containing hexacosane (C26; 25 ng/ul) as an internal standard. Extracts were directly analyzed by GC-MS (7890A, Agilent Technologies Inc., Santa Clara, CA) using a DB-17ht column (Agilent Technologies Inc., Santa Clara, CA). To identify the CHC composition, we first compared retention times and mass spectra to an authentic standard mixture (C7-C40) (Supelco® 49452-U, Sigma-Aldrich, St. Louis, MO) with CHC samples. The types of CHCs, including methyl-branched alkanes, monoenes, dienes, and trienes, were then identified by a combination of their specific fragment ions on the side of functional groups (methyl branch or double bonds), retention times relative to linear hydrocarbon standards, and the m/z value of the molecular ion. The position of methyl branch in mbCHCs was determined using the protocol described in Carlson et al. (1998), while the position of double bonds were not determined in this study. Each CHC peak was quantified using its comparison with the peak area of the internal standard (C26) and represented as nanogram per fly (ng/fly). Because we observed a biased integration of peak areas on longer CHCs in running the standard mixture (**Figure 1-figure supplement 2**), we corrected the peak areas of CHCs based on the carbon-chain lengths using the integration from the standard mixture.

### **Body weight measurement**

Body weight was incorporated in the data collection and analysis. The body weight was determined as the difference between the Eppendorf tube containing five to ten flies of the same sex and the same empty Eppendorf tube.

### **Coating of synthetic compounds**

To coat each of synthetic mbCHCs and *n*-alkanes on *D. melanogaster*, including 2MeC26, 2MeC28, 2MeC30, C23, C25, C27, C29, and C31, we first added 200 µL hexane containing 300 ng/µL of each CHC in a 2-mL glass vial and then used a nitrogen evaporator (BT1603 G-Biosciences®) to evaporate the hexane with only mbCHCs precipitate at the bottom of the vial. For the control group, 200 µL hexane was added. After the hexane was evaporated, ten flies of the same sex were transferred into each vial,

following shaking on a Vortex for 20 seconds on, 20 seconds off, and 20 seconds on, following previous protocols (Billeter et al., 2009; Gendron et al., 2014). The flies were then directly subjected to desiccation assays. Synthetic mbCHCs were kindly provided by Dr. Jocelyn Millar (University of California, Riverside) and synthetic *n*-alkanes were ordered from Sigma-Aldrich (C23 #91447-1G, C25 #286931-1G, C27 #51559-250MG, C29 #74156-250MG, and C31 #51529-250MG).

## Statistics

All analyses were conducted in R (Version 4.1). Correlation analyses were conducted with Pearson's method using '*cor.test*' function. The dependent variables were log-transformed to better conform with assumptions of normality. Variance in CHC beta diversity across desiccation resistance was determined using PERMANOVA using the '*vegan*' package (Anderson et al., 2006). The random forest regression analysis was performed using the '*ranger*' and '*randomForest*' packages (Liaw and Wiener, 2002; Wright and Ziegler, 2015). The random forest regression models were built using both Out Of Bag estimate and test/training sets (70:30 split). To determine how useful each CHC variable is in the prediction of desiccation resistance in the random forest regression analysis, the importance of top predictor CHCs was quantified using permutation importance (Altmann et al., 2010). Desiccation resistance in flies coated with different mbCHCs were determined using one-way ANOVA at  $\alpha = 0.05$ . *Post hoc* comparison was further conducted using Tukey's method. The ancestral trait reconstruction and estimation of phylogenetic signal, Pagel's  $\lambda$ , for mbCHC composition and desiccation resistance was determined using '*Phytools*', '*Picante*', and '*Rphylopars*' packages (Goolsby et al., 2017; Kembel et al., 2010; Revell, 2012). The PGLS analysis was conducted using generalized least squares fit by maximum likelihood and the covariance structure between species was used under a Brownian motion process of evolution. The PGLS analyses were conducted using '*GLS*' function in '*ape*' package (Paradis and Schliep, 2019).

## ACKNOWLEDGEMENTS

We thank Dr. Jocelyn Millar for guidance and advice, as well as the synthetic compounds used in this study. We also thank Ye Ma, Eliana Giannetti, Taylor Hori, and Zhuo Chen for technical assistance, Yushan Zhang for assistance with figure visualization, and the National Drosophila Species Stock Center for fly stocks of these different *Drosophila* species. This work is supported by a National Science Foundation grant (2054773) to H. Chung.

## DATA ACCESSIBILITY

Data and code are available in the supplementary materials.

## DECLARATION OF INTERESTS

The authors declare no competing interests.

## 525 REFERENCES

- 526 Altmann, A., Toloşi, L., Sander, O., and Lengauer, T. (2010). Permutation importance: a corrected feature  
527 importance measure. *Bioinformatics* 26, 1340-1347.
- 528 Anderson, M.J., Ellingsen, K.E., and McArdle, B.H. (2006). Multivariate dispersion as a measure of beta  
529 diversity. *Ecol. Lett.* 9, 683-693.
- 530 Ando, T., Inomata, S.-i., and Yamamoto, M. (2004). Lepidopteran sex pheromones. In *The chemistry of*  
531 *pheromones and other semiochemicals I*, S. Schulz, ed. (Springer), pp. 51-96. doi.org/10.1007/b95449.
- 532 Belsley, D.A., Kuh, E., and Welsch, R.E. (2005). *Regression diagnostics: Identifying influential data and*  
533 *sources of collinearity* (John Wiley & Sons).
- 534 Billeter, J.-C., Atallah, J., Krupp, J.J., Millar, J.G., and Levine, J.D. (2009). Specialized cells tag sexual  
535 and species identity in *Drosophila melanogaster*. *Nature* 461, 987.
- 536 Blomquist, G.J., and Ginzl, M.D. (2021). Chemical ecology, biochemistry, and molecular biology of  
537 insect hydrocarbons. *Annu. Rev. Entomol.* 66, 45-60.
- 538 Buellesbach, J., Whyte, B.A., Cash, E., Gibson, J.D., Scheckel, K.J., Sandidge, R., and Tsutsui, N.D.  
539 (2018). Desiccation Resistance and Micro-Climate Adaptation: Cuticular Hydrocarbon Signatures of  
540 Different Argentine Ant Supercolonies Across California. *J Chem Ecol* 44, 1101-1114.
- 541 Bujan, J., Yanoviak, S.P., and Kaspari, M. (2016). Desiccation resistance in tropical insects: causes and  
542 mechanisms underlying variability in a Panama ant community. *Ecology and evolution* 6, 6282-6291.
- 543 Carlson, D.A., Bernier, U.R., and Sutton, B.D. (1998). Elution patterns from capillary GC for methyl-  
544 branched alkanes. *Journal of Chemical Ecology* 24, 1845-1865.
- 545 Chen, N., Fan, Y.-L., Bai, Y., Li, X.-d., Zhang, Z.-F., and Liu, T.-X. (2016). Cytochrome P450 gene,  
546 CYP4G51, modulates hydrocarbon production in the pea aphid, *Acyrtosiphon pisum*. *Insect Biochem.*  
547 *Mol. Biol.* 76, 84-94.
- 548 Chen, N., Pei, X.J., Li, S., Fan, Y.L., and Liu, T.X. (2019). Involvement of integument-rich *CYP4G19* in  
549 hydrocarbon biosynthesis and cuticular penetration resistance in *Blattella germanica* (L.). *Pest*  
550 *management science*.
- 551 Chenoweth, S.F., and Blows, M.W. (2005). Contrasting mutual sexual selection on homologous signal  
552 traits in *Drosophila serrata*. *The American Naturalist* 165, 281-289.
- 553 Chertemps, T., Duportets, L., Labeur, C., Ueda, R., Takahashi, K., Saigo, K., and Wicker-Thomas, C.  
554 (2007). A female-biased expressed elongase involved in long-chain hydrocarbon biosynthesis and  
555 courtship behavior in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences* 104,  
556 4273-4278.
- 557 Chertemps, T., Duportets, L., Labeur, C., Ueyama, M., and Wicker-Thomas, C. (2006). A female-specific  
558 desaturase gene responsible for diene hydrocarbon biosynthesis and courtship behaviour in *Drosophila*  
559 *melanogaster*. *Insect Molecular Biology* 15, 465-473.
- 560 Chiang, Y.N., Tan, K.J., Chung, H., Lavrynenko, O., Shevchenko, A., and Yew, J.Y. (2016). Steroid  
561 Hormone Signaling Is Essential for Pheromone Production and Oenocyte Survival. *PLoS Genet* 12,  
562 e1006126.

563 Chown, S.L., Hoffmann, A.A., Kristensen, T.N., Angilletta Jr, M.J., Stenseth, N.C., and Pertoldi, C. (2010).  
564 Adapting to climate change: a perspective from evolutionary physiology. *Climate research* 43, 3-15.

565 Chown, S.L., Sørensen, J.G., and Terblanche, J.S. (2011). Water loss in insects: an environmental  
566 change perspective. *Journal of Insect Physiology* 57, 1070-1084.

567 Chung, H., and Carroll, S.B. (2015). Wax, sex and the origin of species: Dual roles of insect cuticular  
568 hydrocarbons in adaptation and mating. *Bioessays* 37, 822-830.

569 Dembeck, L.M., Böröczky, K., Huang, W., Schal, C., Anholt, R.R., and Mackay, T.F. (2015). Genetic  
570 architecture of natural variation in cuticular hydrocarbon composition in *Drosophila melanogaster*. *eLife* 4,  
571 e09861.

572 Everaerts, C., Farine, J.-P., Cobb, M., and Ferveur, J.-F. (2010). *Drosophila* cuticular hydrocarbons  
573 revisited: mating status alters cuticular profiles. *PloS one* 5, e9607.

574 Fedina, T.Y., Kuo, T.-H., Dreisewerd, K., Dierick, H.A., Yew, J.Y., and Pletcher, S.D. (2012). Dietary  
575 effects on cuticular hydrocarbons and sexual attractiveness in *Drosophila*. *PloS one* 7, e49799.

576 Ferveur, J.-F. (2005). Cuticular hydrocarbons: their evolution and roles in *Drosophila* pheromonal  
577 communication. *Behavior genetics* 35, 279.

578 Ferveur, J.-F., Cortot, J., Rihani, K., Cobb, M., and Everaerts, C. (2018). Desiccation resistance: effect of  
579 cuticular hydrocarbons and water content in *Drosophila melanogaster* adults. *PeerJ* 6, e4318.

580 Finet, C., Kassner, V.A., Carvalho, A.B., Chung, H., Day, J.P., Day, S., Delaney, E.K., De Ré, F.C.,  
581 Dufour, H.D., and Dupim, E. (2021). Drosophila: resources for drosophilid phylogeny and systematics.  
582 *Genome biology and evolution* 13, evab179.

583 Freckleton, R.P., Harvey, P.H., and Pagel, M. (2015). Phylogenetic analysis and comparative data: a test  
584 and review of evidence. *The American Naturalist*.

585 Futahashi, R., Yamahama, Y., Kawaguchi, M., Mori, N., Ishii, D., Okude, G., Hirai, Y., Kawahara-Miki, R.,  
586 Yoshitake, K., and Yajima, S. (2019). Molecular basis of wax-based color change and UV reflection in  
587 dragonflies. *eLife* 8, e43045.

588 Gendron, C.M., Kuo, T.H., Harvanek, Z.M., Chung, B.Y., Yew, J.Y., Dierick, H.A., and Pletcher, S.D.  
589 (2014). *Drosophila* life span and physiology are modulated by sexual perception and reward. *Science* 343,  
590 544-548. 10.1126/science.1243339.

591 Gibbs, A., and Pomonis, J.G. (1995). Physical properties of insect cuticular hydrocarbons: the effects of  
592 chain length, methyl-branching and unsaturation. *Comparative Biochemistry and Physiology Part B:*  
593 *Biochemistry and Molecular Biology* 112, 243-249.

594 Gibbs, A.G. (1998). Water-proofing properties of cuticular lipids. *American Zoologist* 38, 471-482.

595 Gibbs, A.G. (2002a). Lipid melting and cuticular permeability: new insights into an old problem. *Journal of*  
596 *Insect Physiology* 48, 391-400.

597 Gibbs, A.G. (2002b). Water balance in desert *Drosophila*: lessons from non-charismatic microfauna.  
598 *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 133, 781-789.

599 Gibbs, A.G., Chippindale, A.K., and Rose, M.R. (1997). Physiological mechanisms of evolved desiccation  
600 resistance in *Drosophila melanogaster*. *Journal of Experimental Biology* 200, 1821-1832.

601 Gibbs, A.G., Fukuzato, F., and Matzkin, L.M. (2003). Evolution of water conservation mechanisms in  
602 *Drosophila*. *Journal of experimental biology* 206, 1183-1192.

603 Gibbs, A.G., and Matzkin, L.M. (2001). Evolution of water balance in the genus *Drosophila*. *Journal of*  
604 *Experimental Biology* 204, 2331-2338.

605 Gibbs, A.G., and Rajpurohit, S. (2010). Cuticular lipids and water balance. *Insect hydrocarbons: biology,*  
606 *biochemistry, and chemical ecology*, 100-120.

607 Goolsby, E.W., Bruggeman, J., and Ané, C. (2017). Rphylopars: fast multivariate phylogenetic  
608 comparative methods for missing data and within-species variation. *Methods in Ecology and Evolution* 8,  
609 22-27.

610 Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society of*  
611 *London. B, Biological Sciences* 326, 119-157.

612 Hadley, N.F. (1977). Epicuticular lipids of the desert tenebrionid beetle, *Eleodes armata*: seasonal and  
613 acclimatory effects on composition. *Insect Biochemistry* 7, 277-283.

614 Hadley, N.F. (1994). *Water relations of terrestrial arthropods* (CUP Archive).

615 Heifetz, Y., Miloslavski, I., Aizenshtat, Z., and Applebaum, S.W. (1998). Cuticular surface hydrocarbons of  
616 desert locust nymphs, *Schistocerca gregaria*, and their effect on phase behavior. *Journal of chemical*  
617 *ecology* 24, 1033-1047.

618 Hoffmann, A., Hallas, R., Dean, J., and Schiffer, M. (2003). Low potential for climatic stress adaptation in  
619 a rainforest *Drosophila* species. *Science* 301, 100-102.

620 Hoffmann, A.A. (2010). Physiological climatic limits in *Drosophila*: patterns and implications. *Journal of*  
621 *Experimental Biology* 213, 870-880.

622 Hoffmann, A.A., and Weeks, A.R. (2007). Climatic selection on genes and traits after a 100 year-old  
623 invasion: a critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern Australia.  
624 *Genetica* 129, 133-147.

625 Holze, H., Schrader, L., and Buellesbach, J. (2021). Advances in deciphering the genetic basis of insect  
626 cuticular hydrocarbon biosynthesis and variation. *Heredity* 126, 219-234.

627 Hood, W.G., and Tschinkel, W.R. (1990). Desiccation resistance in arboreal and terrestrial ants.  
628 *Physiological Entomology* 15, 23-35.

629 Howard, R.W., Jackson, L.L., Banse, H., and Blows, M.W. (2003). Cuticular hydrocarbons of *Drosophila*  
630 *birchii* and *D. serrata*: identification and role in mate choice in *D. serrata*. *Journal of chemical ecology* 29,  
631 961-976.

632 Huang, J., Yu, H., Guan, X., Wang, G., and Guo, R. (2016). Accelerated dryland expansion under climate  
633 change. *Nature Climate Change* 6, 166.

634 Jezovit, J.A., Levine, J.D., and Schneider, J. (2017). Phylogeny, environment and sexual communication  
635 across the *Drosophila* genus. *Journal of Experimental Biology* 220, 42-52.

636 Johnson, R.A., and Gibbs, A.G. (2004). Effect of mating stage on water balance, cuticular hydrocarbons  
637 and metabolism in the desert harvester ant, *Pogonomyrmex barbatus*. *Journal of insect Physiology* 50,  
638 943-953.

639 Kellermann, V., Hoffmann, A.A., Overgaard, J., Loeschcke, V., and Sgrò, C.M. (2018). Plasticity for  
640 desiccation tolerance across *Drosophila* species is affected by phylogeny and climate in complex ways.  
641 Proc. R. Soc. B 285, 20180048.

642 Kellermann, V., Loeschcke, V., Hoffmann, A.A., Kristensen, T.N., Fløjgaard, C., David, J.R., Svenning,  
643 J.C., and Overgaard, J. (2012). Phylogenetic constraints in key functional traits behind species' climate  
644 niches: patterns of desiccation and cold resistance across 95 *Drosophila* species. Evolution 66, 3377-  
645 3389.

646 Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., Blomberg, S.P.,  
647 and Webb, C.O. (2010). Picante: R tools for integrating phylogenies and ecology. Bioinformatics 26,  
648 1463-1464.

649 Khallaf, M.A., Cui, R., Weißflog, J., Erdogmus, M., Svatoš, A., Dweck, H.K., Valenzano, D.R., Hansson,  
650 B.S., and Knaden, M. (2021). Large-scale characterization of sex pheromone communication systems in  
651 *Drosophila*. Nature communications 12, 1-14.

652 Koto, A., Motoyama, N., Tahara, H., McGregor, S., Moriyama, M., Okabe, T., Miura, M., and Keller, L.  
653 (2019). Oxytocin/vasopressin-like peptide inotocin regulates cuticular hydrocarbon synthesis and water  
654 balancing in ants. Proceedings of the National Academy of Sciences 116, 5597-5606.

655 Krupp, J.J., Kent, C., Billeter, J.-C., Azanchi, R., So, A.K.-C., Schonfeld, J.A., Smith, B.P., Lucas, C., and  
656 Levine, J.D. (2008). Social experience modifies pheromone expression and mating behavior in male  
657 *Drosophila melanogaster*. Current Biology 18, 1373-1383.

658 Krupp, J.J., Nayal, K., Wong, A., Millar, J.G., and Levine, J.D. (2020). Desiccation resistance is an  
659 adaptive life-history trait dependent upon cuticular hydrocarbons, and influenced by mating status and  
660 temperature in *D. melanogaster*. Journal of Insect Physiology 121, 103990.

661 Kühsel, S., Brückner, A., Schmelzle, S., Heethoff, M., and Blüthgen, N. (2017). Surface area–volume  
662 ratios in insects. Insect science 24, 829-841.

663 Kwan, L., and Rundle, H.D. (2010). Adaptation to desiccation fails to generate pre-and postmating  
664 isolation in replicate *Drosophila melanogaster* laboratory populations. Evolution: International Journal of  
665 Organic Evolution 64, 710-723.

666 Lamb, A.M., Wang, Z., Simmer, P., Chung, H., and Wittkopp, P.J. (2020). ebony Affects Pigmentation  
667 Divergence and Cuticular Hydrocarbons in *Drosophila americana* and *D. novamexicana*. Frontiers in  
668 Ecology and Evolution 8. 10.3389/fevo.2020.00184.

669 Leeson, S.A., Kennington, W.J., Evans, T.A., and Simmons, L.W. (2020). Phenotypic plasticity but no  
670 adaptive divergence in cuticular hydrocarbons and desiccation resistance among translocated  
671 populations of dung beetles. Evolutionary Ecology 34, 929-944.

672 Li, F., Rane, R.V., Luria, V., Xiong, Z., Chen, J., Li, Z., Catullo, R.A., Griffin, P.C., Schiffer, M., and Pearce,  
673 S. (2022). Phylogenomic analyses of the genus *Drosophila* reveals genomic signals of climate adaptation.  
674 Molecular Ecology Resources 22, 1559-1581.

675 Liaw, A., and Wiener, M. (2002). Classification and regression by randomForest. R news 2, 18-22.

676 Lynch, M. (1991). Methods for the analysis of comparative data in evolutionary biology. Evolution 45,  
677 1065-1080.

678 Matzkin, L.M., Watts, T.D., and Markow, T.A. (2009). Evolution of stress resistance in *Drosophila*:  
679 interspecific variation in tolerance to desiccation and starvation. Functional Ecology 23, 521-527.

680 Menzel, F., Blaimer, B.B., and Schmitt, T. (2017). How do cuticular hydrocarbons evolve? Physiological  
681 constraints and climatic and biotic selection pressures act on a complex functional trait. *Proceedings of*  
682 *the Royal Society B: Biological Sciences* **284**, 20161727.

683 Menzel, F., Morsbach, S., Martens, J.H., Räder, P., Hadjaje, S., Poizat, M., and Abou, B. (2019).  
684 Communication versus waterproofing: the physics of insect cuticular hydrocarbons. *Journal of*  
685 *Experimental Biology* **222**.

686 Mueller, L.D. (1988). Density-dependent population growth and natural selection in food-limited  
687 environments: the *Drosophila* model. *The American Naturalist* **132**, 786-809.

688 Mundry, R. (2014). Statistical issues and assumptions of phylogenetic generalized least squares. In  
689 *Modern phylogenetic comparative methods and their application in evolutionary biology*, (Springer), pp.  
690 131-153.

691 Nunes, T.M., Oldroyd, B.P., Elias, L.G., Mateus, S., Turatti, I.C., and Lopes, N.P. (2017). Evolution of  
692 queen cuticular hydrocarbons and worker reproduction in stingless bees. *Nature Ecology & Evolution* **1**,  
693 1-3.

694 Paradis, E., and Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary  
695 analyses in R. *Bioinformatics* **35**, 526-528.

696 Qiu, Y., Tittiger, C., Wicker-Thomas, C., Le Goff, G., Young, S., Wajnberg, E., Fricaux, T., Taquet, N.,  
697 Blomquist, G.J., and Feyereisen, R. (2012). An insect-specific P450 oxidative decarboxylase for cuticular  
698 hydrocarbon biosynthesis. *Proceedings of the National Academy of Sciences* **109**, 14858-14863.

699 Rajpurohit, S., Vrkoslav, V., Hanus, R., Gibbs, A.G., Cvačka, J., and Schmidt, P.S. (2021). Post-eclosion  
700 temperature effects on insect cuticular hydrocarbon profiles. *Ecology and Evolution* **11**, 352-364.

701 Rane, R.V., Pearce, S.L., Li, F., Coppin, C., Schiffer, M., Shirriffs, J., Sgrò, C.M., Griffin, P.C., Zhang, G.,  
702 and Lee, S.F. (2019). Genomic changes associated with adaptation to arid environments in cactophilic  
703 *Drosophila* species. *BMC genomics* **20**, 52.

704 Revell, L.J. (2012). phytools: an R package for phylogenetic comparative biology (and other things).  
705 *Methods in ecology and evolution*, 217-223.

706 Rouault, J.-D., Marican, C., Wicker-Thomas, C., and Jallon, J.-M. (2004). Relations between cuticular  
707 hydrocarbon (HC) polymorphism, resistance against desiccation and breeding temperature; a model for  
708 HC evolution in *D. melanogaster* and *D. simulans*. In *Drosophila melanogaster, Drosophila simulans: So*  
709 *Similar, So Different*, (Springer), pp. 195-212.

710 Rusuwa, B.B., Chung, H., Allen, S.L., Frentiu, F.D., and Chenoweth, S.F. (2022). Natural variation at a  
711 single gene generates sexual antagonism across fitness components in *Drosophila*. *Current Biology* **32**,  
712 3161-3169.

713 Savage, B.A., Wang, Z., Chung, H., Masten, S., and Grieshop, M. (2021). An Ozonolysis Based Method  
714 and Applications for the Non-lethal Modification of Insect Cuticular Hydrocarbons. *Journal of Chemical*  
715 *Ecology*.

716 Sherwood, S., and Fu, Q. (2014). A drier future? *Science* **343**, 737-739.

717 Shi, H., Tian, H., Lange, S., Yang, J., Pan, S., Fu, B., and Reyer, C.P. (2021). Terrestrial biodiversity  
718 threatened by increasing global aridity velocity under high-level warming. *Proceedings of the National*  
719 *Academy of Sciences* **118**.



720 Soroker, V., and Hefetz, A. (2000). Hydrocarbon site of synthesis and circulation in the desert ant  
721 *Cataglyphis niger*. *Journal of Insect Physiology* **46**, 1097-1102.

722 Spikes, A.E., Paschen, M.A., Millar, J.G., Moreira, J.A., Hamel, P.B., Schiff, N.M., and Ginzel, M.D.  
723 (2010). First contact pheromone identified for a longhorned beetle (Coleoptera: Cerambycidae) in the  
724 subfamily Prioninae. *Journal of chemical ecology* **36**, 943-954.

725 Svetnik, V., Liaw, A., Tong, C., Culberson, J.C., Sheridan, R.P., and Feuston, B.P. (2003). Random forest:  
726 a classification and regression tool for compound classification and QSAR modeling. *Journal of Chemical*  
727 *Information and computer Sciences* **43**, 1947-1958.

728 Van Oystaeyen, A., Oliveira, R.C., Holman, L., van Zweden, J.S., Romero, C., Oi, C.A., d'Ettorre, P.,  
729 Khalesi, M., Billen, J., and Wäckers, F. (2014). Conserved class of queen pheromones stops social insect  
730 workers from reproducing. *Science* **343**, 287-290.

731 Wang, L., Han, X., Mehren, J., Hiroi, M., Billeter, J.-C., Miyamoto, T., Amrein, H., Levine, J.D., and  
732 Anderson, D.J. (2011). Hierarchical chemosensory regulation of male-male social interactions in  
733 *Drosophila*. *Nature neuroscience* **14**, 757-762.

734 Wang, Y., Ferveur, J.F., and Moussian, B. (2021). Eco-genetics of desiccation resistance in *Drosophila*.  
735 *Biological Reviews*.

736 Wigglesworth, V.B. (1945). Transpiration through the cuticle of insects. *Journal of Experimental Biology*  
737 **21**, 97-114.

738 Wright, M.N., and Ziegler, A. (2015). ranger: A fast implementation of random forests for high dimensional  
739 data in C++ and R. *arXiv preprint arXiv:1508.04409*.

740  
741

## FIGURE LEGENDS

**Figure 1. A cohort-based experimental design.** A cohort-based experimental design can facilitate determining the correlation between desiccation resistance, cuticular hydrocarbons (CHCs), and body weight. Five to six cohorts of each species was established.

**Figure 2. Desiccation resistance and CHC composition in 46 *Drosophila* species and 4 outgroup species.** Desiccation resistance and CHC composition of each species were plotted together. Males and females are shown separately. The shading intensity of each species represents the number of hours of desiccation resistance, while the size and color of each circle represent the type of CHC and its quantity. *n*-alkanes are only present in some species from the melanogaster group.

**Figure 3. Higher amounts of CHCs do not contribute to higher desiccation resistance. (A)** Body weight is positively correlated with desiccation resistance (Females:  $r = 0.4$ ,  $P < 0.001$ , Males:  $r = 0.5$ ,  $P < 0.001$ ). **(B)** Total amount of CHCs is correlated with having higher body weight (Females:  $r = 0.7$ ,  $P < 0.001$ , Males:  $r = 0.7$ ,  $P < 0.001$ ). **(C)** A weak positive correlation between desiccation resistance and CHCs as a percentage of body weight in females, while no correlation in males (Females:  $r = 0.1$ ,  $P = 0.03$ , Males:  $P = 0.1$ ). All correlation analyses were conducted using Pearson's method.

**Figure 4. CHC composition can be used to predict desiccation resistance (A)** Random Forest regression modeling of CHC abundance was able to explain 85.5% of the variation in time to desiccation with a Root Mean Square Error (RMSE) of 4.5. **(B)** The abundance of four mbCHCs, 2MeC30, 2MeC28, 2MeC32, and 2MeC26 has the highest importance to the desiccation resistance in the random forest regression model, while most of CHCs have less contribution to the accuracy of the model for desiccation resistance.

**Figure 5. Longer mbCHCs are associated with higher desiccation resistance. (A)** Quantity of 2MeC26 is negatively correlated with desiccation resistance in males, but no correlation in females (Females:  $P = 0.1$ , Males:  $r = -0.4$ ,  $P < 0.001$ ). **(B)** Quantity of 2MeC28 is positively correlated with desiccation resistance (Females:  $r = 0.4$ ,  $P < 0.001$ , Males:  $r = 0.4$ ,  $P < 0.001$ ). **(C)** Quantity of 2MeC30 is positively correlated with desiccation resistance in females, but no correlation in males (Females:  $r = 0.2$ ,  $P = 0.009$ , Males:  $P = 0.2$ ). **(D)** Quantity of 2MeC32 is positively correlated with desiccation resistance (Females:  $r = 0.3$ ,  $P < 0.001$ , Males:  $r = 0.4$ ,  $P < 0.001$ ). Correlations between the quantities of each mbCHC and desiccation resistance were determined using Pearson's method. **(E)** *D. melanogaster* coated with longer mbCHCs have higher desiccation resistance. Coating of synthetic mbCHCs on *D. melanogaster* showed that 2MeC26 does not influence desiccation resistance. 2MeC30 increases desiccation resistance in female *D. melanogaster* while both 2MeC28 and 2MeC30 in increases

desiccation resistance in *male D. melanogaster*. The bold horizontal line within each box plot corresponds to the median value, the box length to the interquartile range, and the lines emanating from the box (whiskers) extend to the smallest and largest observations. One-Way ANOVA showed significant differences between *D. melanogaster* flies coated with different mbCHCs (Female:  $F_{(3,189)} = 73.1$ ,  $P < 0.001$ , Male:  $F_{(3,191)} = 23.7$ ,  $P < 0.001$ ). *Post hoc* comparison was conducted using Tukey's method.

**Figure 6. Evolution of longer mbCHCs are significantly correlated with evolution of higher desiccation resistance.** Patterns in the normalized quantities of the longest mbCHCs and desiccation resistance for females (top) and males (bottom) were listed across the phylogeny of the 50 *Drosophila* and related species. PGLS analysis between the longest mbCHCs and desiccation resistance showed both the higher quantity and longer length of the longest mbCHC affect desiccation resistance (interaction term, Female:  $t = 3.5$ ,  $P < 0.001$ ; Male:  $t = 2.2$ ,  $P = 0.03$ ).

**Figure 1-figure supplement 1. The phylogeny of 50 *Drosophila* and related species used in this study.** 50 species representing 46 *Drosophila* species (18 from *Drosophila* subgenus, 28 from *Sophophora* subgenus) and four species (3 *Scaptodrosophila* species and 1 *Chymomyza* species) as outgroup were used in our study. The 46 *Drosophila* species represents some of the main groups in the *Drosophila* genus: (1) melanogaster group, (2) obscura group, (3) willistoni group, (4) nasuta group, (5) virilis group, and (6) repleta group.

**Figure 1-figure supplement 2. Chromatogram of the authentic standard *n*-alkane mixture which contains 34 *n*-alkanes from C7 to C40 at the same concentration.** The authentic standard *n*-alkane mixture (C7-C40; Supelco® 49452-U) was obtained from Sigma-Aldrich (St. Louis, MO). C7 – C14 were excluded in the chromatogram due to the solvent delay in the GC-MS program. The integration of peak areas on longer CHCs is biased to have lower values. To address this issue, we corrected the peak areas of CHCs based on the carbon-chain lengths using the integration from this standard *n*-alkane mixture.

**Figure 2-figure supplement 1. Desiccation resistance and body weight in 46 *Drosophila* species and 4 outgroup species.** Desiccation resistance and body weight are shown in separate graphs.

**Figure 2-figure supplement 2. The melting temperature of CHCs is determined by methyl group, double bonds, and carbon chain length.** Five types of CHCs are detected in this study. Due to their chemical structures, *n*-alkanes have the highest melting temperatures followed by mbCHCs, monoenes, dienes, and trienes. Increasing length of the carbon-chain can also increase melting temperature. The positions of double bonds in this figure are for illustrative purposes and were not determined in this study.

**Figure 3-figure supplement 1. CHC quantity as a percentage of body weight.** As *Drosophila* species largely vary in size, we normalized the total CHCs as a percentage of body weight. CHCs account from 0.02% to 0.5% of the total body weight of different species.

**Figure 4-figure supplement 1. The composition of CHCs significantly differed across the increasing desiccation resistance.** NMDS plot showing the beta diversity of CHCs differs significantly across the increasing desiccation resistance (PERMANOVA:  $r^2 = 0.1$ ,  $P < 0.001$ ), suggesting CHC composition could affect desiccation resistance in the 50 *Drosophila* and selected species.

**Figure 4-figure supplement 2. Cross-validation of the random forest regression model has a similar performance to the regression model using the full dataset.** A 70:30 training/test split in the dataset ( $n = 382$ ,  $n = 164$ ) was used to evaluate the performance of the random forest regression model. The out of bag estimate of the root mean square error (RMSE) in the split dataset is 5.4, which is similar to the RMSE using the full dataset (RNMSE = 4.5).

**Figure 5-figure supplement 1. Correlation between different mbCHCs in 50 *Drosophila* and related species.** Correlation between different mbCHCs was determined using Pearson's method. Significant correlations between each pairs of mbCHCs were labeled with stars following with the correlation coefficients. The lines on the diagonals represent the distribution of the data for each mbCHC. \*\*:  $P < 0.01$ ; \*\*\*:  $P < 0.001$ .

**Figure 5-figure supplement 2. The length of coated mbCHCs on *D. melanogaster attP40* flies is positively correlated with desiccation resistance in both females and males.** Pearson's correlation: Female,  $r = 0.7$ ,  $P < 0.001$ ; Male,  $r = 0.5$ ,  $P < 0.001$ .

**Figure 5-figure supplement 3. Desiccation resistance of *D. melanogaster attP40* flies coated with individual *n*-alkanes (C23, C25, C27, C29, and C31).** One-Way ANOVA was applied to test the differences in desiccation resistance between the coating of different *n*-alkanes (Female:  $F_{(5, 277)} = 17.1$ ,  $P < 0.001$ , Male:  $F_{(5, 287)} = 8.0$ ,  $P < 0.001$ ), following with *post hoc* comparison using Tukey's method at  $\alpha = 0.05$ . \*:  $P < 0.05$ ; \*\*:  $P < 0.01$ ; \*\*\*:  $P < 0.001$ . The bold horizontal line within each box plot corresponds to the median value, the box length to the interquartile range, and the lines emanating from the box (whiskers) extend to the smallest and largest observations. Numbers on top of the boxplots are the mean in each treatment.

**Figure 5-figure supplement 4. The length of coated *n*-alkanes on *D. melanogaster attP40* flies is positively correlated with desiccation resistance in both females and males.** Pearson's correlation: Female,  $r = 0.4$ ,  $P < 0.001$ ; Male,  $r = 0.1$ ,  $P = 0.02$ .

**Figure 5-figure supplement 5. Desiccation resistance of CHC- *D. melanogaster* flies coated with individual mbCHCs (2MeC26, 2MeC28, 2MeC30) or *n*-alkanes (C23, C25, C27, C29, and C31).** One-Way ANOVA was applied to test the difference of desiccation resistance between the coating of different mbCHCs and *n*-alkanes (Female:  $F_{(8, 417)} = 11.2$ ,  $P < 0.001$ , Male:  $F_{(8, 408)} = 7.1$ ,  $P < 0.001$ ), following with *post hoc* comparison using Tukey's method at  $\alpha = 0.05$ . \*:  $P < 0.05$ ; \*\*:  $P < 0.01$ ; \*\*\*:  $P < 0.001$ . The bold horizontal line within each box plot corresponds to the median value, the box length to the interquartile range, and the lines emanating from the box (whiskers) extend to the smallest and largest observations. Numbers on top of the boxplots are the mean in each treatment.

**Figure 5-figure supplement 6. The length of coated mbCHCs on CHC- flies has a weak positive correlation with desiccation resistance in females but did not correlate with the levels of desiccation resistance in males.** Pearson's correlation: Female,  $r = 0.2$   $P = 0.004$ ; Male,  $P = 0.9$ .

**Figure 5-figure supplement 7. The length of coated *n*-alkanes on CHC- flies did not correlate with desiccation resistance.** Pearson's correlation: Female,  $P = 0.4$ ; Male,  $P = 1.0$ .

**Figure 5-figure supplement 8. Pathway for the synthesis of branched CHCs (mbCHCs) and linear CHCs (*n*-alkanes, monoenes, dienes).** All CHCs are synthesized via the fatty acyl-CoA synthesis pathway from acetyl-CoA. The pathway splits early due to the action of two fatty acid synthetases (FAS) into the branched CHC pathway or the linear CHC pathway. The precursors of each pathway are modified by synthesis enzymes such as elongases (Elo), reductases (FAR), and the terminal P450 decarbonylase (CYP4G) into CHCs. For linear CHCs, additional enzymes such as desaturases (Desat) incorporate double bonds during synthesis. The diagram was modified from Chung and Carroll, 2015.

**Figure 5-figure supplement 9. GC-MS chromatograms of 5'*mFAS*-GAL4, UAS-*Cyp4g1*-RNAi, and the F1 offspring from 5'*mFAS*-GAL4 x UAS-*Cyp4g1*-RNAi.** CHC- *D. melanogaster* flies are generated by crossing 5'*mFAS*-GAL4 and UAS-*Cyp4g1*-RNAi, resulting in adult oenocyte specific knockdown of *Cyp4g1* and loss of CHC production. IS = internal standard spiked in for quantification (C26). cVA = *cis*-vaccenyl acetate, a lipid produced in the male ejaculatory bulb not affected by the RNAi knockdown of *Cyp4g1* in the oenocytes.

**Figure 6-figure supplement 1. Ancestral state reconstruction of mbCHC for the *Drosophila* genus.** Bar plots on the right were the derived ancestral state of female and male mbCHC profiles for the common ancestor of the *Drosophila* genus (A), the *Sophophora* subgenus (B), the *Drosophila* subgenus (C), the melanogaster group (D), the willistoni group (E), the nasuta group (F), the virilis group (G), the repleta group (H), the clade of *D. melanogaster*, *D. simulans*, and *D. mauritiana* (I), and the clade of *D.*

889 *mojavensis* and *D. arizonae* (J). 2MeC28 and 2MeC30 were the major CHCs in the derived ancestral  
890 state (A).  
891

892 **LEGENDS for Supplementary File 1**

893 **Supplementary File 1a.** Phylogenetic signals for mbCHCs in *Drosophila* species.

894

895 **Supplementary File 1b.** Phylogenetic signals for desiccation resistance in *Drosophila* species.

896

897 **Supplementary File 1c.** Summary of the Phylogenetic Generalized Linear Square (PGLS) models  
898 between the longest mbCHCs and desiccation resistance for females and males in 50 *Drosophila* and  
899 related species.

900

901 **Supplementary File 1d.** List of species used in this study.

902

903

**LEGENDS for Source Data and Code Files**

**Figure 2-5 - Source Data 1.** Data of normalized CHC quantities, body weight, averaged desiccation resistance, total CHC quantity, and percentage of CHC quantities per body weight.

**Figure 5 - Source data 2.** Desiccation resistance of *Drosophila melanogaster attP40* flies when coated with individual mbCHC of 2MeC26, 2MeC28, and 2MeC30.

**Figure 5 - Source data 3.** Desiccation resistance of *Drosophila melanogaster attP40* flies when coated with individual *n*-alkane of C23, C25, C27, C29, and C31.

**Figure 5 - Source data 4:** Desiccation resistance of *Drosophila melanogaster* CHC- flies when coated with individual CHC of 2MeC26, 2MeC28, 2MeC30, C23, C25, C27, C29, and C31.


**Figure 6 - source data 1:** Normalized quantities of mbCHCs, the length and normalized quantities of the longest mbCHC in each species.

**Source Code File 1.** Code for Pearson correlation analysis and ANOVA tests in Figure 3 and Figure 5

**Source Code File 2.** Code for NMDS plot and random forest analysis for Figure 4


**Source Code File 3.** Code for PGLS analyses in Figure 6






## Females







### Cuticular hydrocarbons (CHCs)






## Males

### Cuticular hydrocarbons (CHCs)







**A****B**




## Female




## Male

































**Female**




**Male**



## Female




## Male










**Female**






**Male**















Female

Male

*5'mFas-GAL4*



*UAS-Cyp4g1-RNAi*



*5'mFas-GAL4 >*  
*UAS-Cyp4g1-RNAi*

