

Glutamine catabolism supports amino acid biosynthesis and suppresses the integrated stress response to promote photoreceptor survival

Reviewed Preprint

v2 • April 15, 2025

Revised by authors

Reviewed Preprint

v1 • August 28, 2024

Moloy T Goswami, Eric Weh, Shubha Subramanya, Katherine M Weh, Hima Bindu Durumutla, Heather Hager, Nicholas Miller, Sraboni Chaudhury, Anthony Andren, Peter Sajjakulnukit, Li Zhang, Cagri G Besirli, Costas A Lyssiotis, Thomas J Wubben 

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States • Molecular and Developmental Biology Graduate Program, Cincinnati Children's Hospital Medical Center, Cincinnati, United States • Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, United States • Department of Internal Medicine, Division of Gastroenterology and Hepatology, University of Michigan, Ann Arbor, United States • Rogel Cancer Center, University of Michigan, Ann Arbor, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

Goswami and colleagues used rod-specific GLS1 (the gene encoding glutaminase 1) knockout mice to investigate the role of GLS1 in photoreceptor health when GLS1 was deleted from developing or adult photoreceptor cells. This study is **fundamental** as it shows the critical role of glutamine catabolism in photoreceptor cell health using in vivo model systems. The evidence supporting the authors' claims is **compelling**. The studies add new insight into how specific metabolites support vision.

<https://doi.org/10.7554/eLife.100747.2.sa4>

Abstract

Photoreceptor loss results in vision loss in many blinding diseases, and metabolic dysfunction underlies photoreceptor degeneration. So, exploiting photoreceptor metabolism is an attractive strategy to prevent vision loss. Yet, the metabolic pathways that maintain photoreceptor health remain largely unknown. Here, we investigated the dependence of photoreceptors on glutamine (Gln) catabolism. Gln is converted to glutamate via glutaminase (GLS), so mice lacking GLS in rod photoreceptors were generated to inhibit Gln catabolism. Loss of GLS produced rapid rod photoreceptor degeneration. In vivo metabolomic methodologies and metabolic supplementation identified Gln catabolism as critical for glutamate and aspartate biosynthesis. Concordant with this amino acid deprivation, the integrated stress response (ISR) was activated with protein synthesis attenuation, and inhibiting the ISR delayed photoreceptor loss. Furthermore, supplementing asparagine, which is synthesized from aspartate, delayed photoreceptor degeneration. Hence, Gln

catabolism is integral to photoreceptor health, and these data reveal a novel metabolic axis in these metabolically-demanding neurons.

Impact statement

For the first time, this work reveals the metabolic dependency of photoreceptors on glutamine catabolism *in vivo* and further demonstrates the flexibility of photoreceptors to utilize fuel sources beyond glucose.

Introduction

Photoreceptor (PR) death is the cause of vision loss in many retinal diseases. A paucity of effective therapies exist that prevent PR death, so there is an unmet need for therapeutics that improve or prolong PR survival. The retina has a significant energetic demand, driven in large part by PRs (Pan et al., 2021 [DOI](#)). The prodigious bioenergetic requirements of photoreceptors (PRs) are due to the need to conduct phototransduction and neurotransmission as well as manufacture the lipid- and protein-rich outer segment, which is shed daily and phagocytosed by the retinal pigment epithelium (RPE) (Ng et al., 2015 [DOI](#); Young, 1967 [DOI](#)). PRs have little reserve capacity to generate adenosine triphosphate (ATP) and as a result, are adversely affected by small changes in energy homeostasis. As such, metabolic dysfunction has been shown to underlie PR cell death (Bowne et al., 2006 [DOI](#); Du et al., 2015 [DOI](#); Hartong et al., 2008 [DOI](#); Kooragayala et al., 2015 [DOI](#); Pan et al., 2021 [DOI](#)), and exploiting PR metabolism to make these cells more robust to stress is an attractive neuroprotective strategy (Pan et al., 2021 [DOI](#)). Yet, beyond glucose, the metabolic pathways integral to PR health remain largely unknown. This is a critical knowledge gap as identification of these pathways is likely to reveal new strategies for therapeutic intervention (Duncan et al., 2018 [DOI](#)).

Glucose is central to PR metabolism as these cells utilize aerobic glycolysis, or the conversion of glucose to lactate despite the presence of oxygen, for the production of both energy and anabolic building blocks, similar to cancer cells (Ait-Ali et al., 2015 [DOI](#); Chinchore et al., 2017 [DOI](#); Kanow et al., 2017 [DOI](#); Petit et al., 2018 [DOI](#); Swarup et al., 2019 [DOI](#)). Previous studies have shown that genetic knockdown of enzymes key to aerobic glycolysis leads to PR dysfunction and death (Chinchore et al., 2017 [DOI](#); Petit et al., 2018 [DOI](#); Rajala, 2020 [DOI](#); Weh et al., 2020 [DOI](#); Wubben et al., 2017 [DOI](#)). However, like other metabolically-demanding cells, recent work has demonstrated that PRs have the flexibility to utilize fuel sources beyond glucose to meet their metabolic needs (Adler et al., 2014 [DOI](#); Daniele et al., 2022 [DOI](#); Du, Cleghorn, Contreras, Linton, et al., 2013; Grenell et al., 2019 [DOI](#); Joyal et al., 2016 [DOI](#); Xu et al., 2020 [DOI](#)).

Glutamine (Gln) is the most abundant amino acid found in the body and circulating within the blood (Yang et al., 2017 [DOI](#)), and many rapidly dividing cells, including cancer cells, which utilize aerobic glycolysis, depend on Gln for their survival and proliferation (Yang et al., 2017 [DOI](#)). Gln can serve as a substrate for multiple pathways, providing a carbon and nitrogen source for biosynthesis, energetics, and cellular reactive oxygen species (ROS) homeostasis (Adler et al., 2014 [DOI](#); Xu et al., 2020 [DOI](#); Yang et al., 2017 [DOI](#)). These features potentially make Gln an ideal alternative fuel source for PRs, whose bioenergetic demand rivals that of cancer cells despite being terminally differentiated (Ng et al., 2015 [DOI](#)). To this end, over half of the Gln in the retina is found in the outer retina, which is primarily composed of PRs (Du, Cleghorn, Contreras, Linton, et al., 2013; Macosko et al., 2015 [DOI](#); Voigt et al., 2020 [DOI](#)). *Ex vivo* experiments have shown that Gln can be a source of carbons for TCA cycle intermediates in the retina and contribute to amino acid biosynthesis (Du, Cleghorn, Contreras, Linton, et al., 2013; Grenell et al., 2019 [DOI](#); Tsantilas et al., 2021 [DOI](#); Xu et al., 2020 [DOI](#)). Additionally, Gln was demonstrated to support nicotinamide adenine dinucleotide phosphate (NADPH) generation in the absence of glucose *in vitro* in isolated PRs

(Adler et al., 2014 [↗](#)). Finally, in a genetically altered mouse model that disrupts glucose transport to the PRs, both Gln and its transporter were upregulated in the retina, implying that metabolism of this amino acid may be supporting PR survival when glucose is limiting (Swarup et al., 2019 [↗](#)).

Glutaminolysis is the process by which Gln is metabolized into TCA cycle intermediates for critical biosynthetic precursors. This process is initiated by the catabolism of Gln to glutamate (Glu) via one of two glutaminase enzymes: kidney-type glutaminase (GLS) or liver-type glutaminase (GLS2) (Yang et al., 2017 [↗](#)). Single-cell RNA sequencing data (GSE63473 and GSE142449) has demonstrated that GLS is the predominant isoform in the retina and PRs (Macosko et al., 2015 [↗](#); Voigt et al., 2020 [↗](#)). Additionally, it has been shown that glutaminase activity is at least two-fold higher in mitochondria-rich PR inner segments (Ross et al., 1987 [↗](#)). Since the inner segments of PRs are responsible for supporting the majority of energy production and metabolism, GLS activity is likely critical for metabolic functions in PRs.

Glutaminolysis via GLS is indispensable to the metabolism of many cancer cells (Altman et al., 2016 [↗](#); Yang et al., 2017 [↗](#)), and a variety of data show the similarities between cancer cell metabolism and retinal metabolism, and specifically, metabolism of the PRs in the outer retina (Chinchore et al., 2017 [↗](#); Du, Cleghorn, Contreras, Linton, et al., 2013; Ng et al., 2015 [↗](#); Rajala, 2020 [↗](#)). Therefore, we hypothesized that GLS-initiated Gln catabolism may also be essential to PR metabolism, function, and survival. Previous studies utilizing *in vitro* or *ex vivo* methods with whole retinas from wild-type mice or mice with genetic perturbations not confined to PRs (Adler et al., 2014 [↗](#); Du, Cleghorn, Contreras, Linton, et al., 2013; Grenell et al., 2019 [↗](#); Xu et al., 2020 [↗](#)) have provided a foundation for this governing hypothesis, but none have examined the role of Gln catabolism specifically in PRs *in vivo*. In this study, we generated a rod photoreceptor-specific knockout of GLS to comprehensively study the importance of GLS-driven Gln catabolism in PRs.

Results

Generation of a rod photoreceptor-specific, *Gls* knockout mouse

To confirm the single-cell RNA sequencing data (Macosko et al., 2015 [↗](#); Voigt et al., 2020 [↗](#)), we conducted real-time reverse transcription PCR (qRT-PCR) with primers specific for either *Gls* or *Gls2* (Supplementary File 1 [↗](#)). These data show that *Gls* expression is 14 times greater than that of *Gls2* in the mouse retina (Figure 1 – figure supplement 1A [↗](#)), indicating that GLS is indeed the predominant isoform in murine retina. Furthermore, retinal sections stained for GLS using immunofluorescence showed that GLS is expressed throughout the retina with enrichment in the PR inner segments (Figure 1 – figure supplement 1B [↗](#)). It has been shown that GLS activity is at least two-fold higher in the PR inner segments, which are rich in mitochondria (Ross et al., 1987 [↗](#)). Accordingly, GLS segregated to the mitochondrial-enriched fraction more than the cytosolic fraction in the retina (Figure 1 – figure supplement 1C [↗](#)).

We generated a rod PR-specific, *Gls* conditional knockout mouse to determine the role of GLS activity in PR survival and function. Animals homozygous for a floxed *Gls* allele and expressing a Cre-recombinase under the control of the rhodopsin promoter (*Gls^{fl/fl};Rho-Cre⁺*, cKO) as well as animals expressing only the Cre-recombinase (*Gls^{wt/wt};Rho-Cre⁺*, WT) were generated. Total retinal lysate was collected from cKO and WT animals at postnatal day 14 (P14) and GLS expression was measured (Figure 1A [↗](#)). These data show that cKO animals have significantly less GLS expression compared to WT animals. Whole eyes were collected at P14 and stained for GLS using immunofluorescence. Figure 1B [↗](#) shows significant loss of GLS expression in PR inner segments of cKO animals compared to WT animals. The remaining GLS in the inner segment layer is found within cone PRs, as shown in Figure 1B [↗](#) (arrows), confirming the knockout is specific to rod PRs. A lack of compensatory upregulation of *Gls2* in the retina was confirmed via qRT-PCR and in PRs using immunofluorescence (Figure 1 – figure supplement 1D-E [↗](#)).




Figure 1.

Rod photoreceptor-specific knockout of glutaminase (GLS) displays rapid retinal degeneration and increased markers of cell death

(A) Western blot analysis showing decreased GLS protein levels in the retina of WT and cKO mice at post-natal day 14 (P14). Quantitation of Western blot results for N=3-4 animals per group. (B) Representative images for immunofluorescence of P14 mouse retinas (N=3 animals per group) stained for GLS (red), cone opsin (green) and nuclei (DAPI, blue) in WT and cKO mice. White arrows indicate remaining GLS expression in cone photoreceptors. Left and middle panel scale bars are 40 μ m. Right image scale bar is 20 μ m. (C) OCT images detailing outer retinal changes in cKO mice over time compared to WT. Retinal structures are comparable to WT mice at P14 but rapidly thin with age. (D) Total retinal thickness, (E) ONL thickness and (F) IS/OS thickness as determined by OCT in WT and cKO mice over time. N=4-9 eyes per group. (G) Representative hematoxylin and eosin stained retinal sections from rod photoreceptor-specific *Glsl* conditional knockout (cKO) mice compared to wild-type (WT) mice at P14, P21 and P42. ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion cell layer. Scale bar is 40 μ m. (H) ONL cell counts as a percent of WT retinas at P14, P21 and P42. N=5-10 eyes per group. (I) Representative images of WT and cKO retinas stained to detect TUNEL-positive cells (green) at P21. Scale bar is 40 μ m. N=3-5 animals per group. (J) Quantitation of percent TUNEL-positive cells at P21 showing an increase in TUNEL-positive cells in cKO animals. N=3-5 animals per group. (K) qRT-PCR of genes related to cell death pathways including apoptosis (Apop.), necroptosis (Nec.), autophagy (Aut.) and ferroptosis (Ferrop.) in WT and cKO mice at P14. N=6 animals per group. Statistical differences in (A), (D), (E), (F), (H), (J) and (K) are based on an unpaired two-tailed Student's t-test where * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$. Data are presented as mean $\pm$ standard error of the mean. OCT: optical coherence tomography, OS: outer segment, IS: inner segment, ONL: outer nuclear layer, INL: inner nuclear layer, GCL: ganglion cell layer, TUNEL: terminal deoxynucleotidyl transferase dUTP nick and labeling.

Loss of GLS causes rapid PR degeneration

These data clearly show that GLS is significantly downregulated in cKO animals. To determine if loss of GLS affects PR survival, optical coherence tomography (OCT) was used to measure the *in vivo* thickness of each retinal layer at various timepoints in cKO and WT animals (**Figure 1C**). At P14, cKO and WT animals are indistinguishable in total retinal and outer nuclear layer (ONL) thickness (**Figures 1D** and **1E**). However, by P21 a significant loss in total retinal, ONL, and IS/OS thickness is observed (**Figures 1D–1F**) and cKO animals continue to experience loss of retinal, ONL and IS/OS thickness out to P84. Considering mice open their eyes around P14, we assessed if light exposure was contributing to PR degeneration. cKO mice were reared in the dark, but no change in the rate of ONL degeneration was observed when comparing dark-reared mice to those reared in 12-hour light/12-hour dark cyclic lighting conditions (**Figure 1 – figure supplement 2**). Histology at P14, P21, and P42 confirmed loss of PR cell bodies (**Figures 1G** and **1H**). As has been seen in other models of PR degeneration, glial fibrillary acidic protein (GFAP), which is a marker of stress-induced Müller glial cell activation, was increased in the retina of the cKO mouse at P21 and P42 (**Figure 1 – figure supplement 3**) (Grenell et al., 2019).

Next, to assess the cell death pathways contributing to PR degeneration in the cKO mouse, retinal sections from P21 animals were stained for TUNEL (**Figure 1I**). At P21, there is a significant increase in TUNEL positive outer retinal cells in cKO mice compared to WT mice (**Figure 1J**). qRT-PCR analysis of cell death-related gene expression at P14 (**Figure 1K**) demonstrated an increase in genes involved in apoptosis, necroptosis, and ferroptosis. These data coincide with the thought that apoptosis is the predominant mechanism of PR death in many retinal diseases, considering *Casp8* showed the greatest increase in expression in the cKO retina, but also that other cell death mechanisms can contribute to PR cell death (Wubben et al., 2016).

Retinas were stained for RHO expression using immunofluorescence, which indicated rod OSs were shorter after *Gl*s knockout (**Figure 1 – figure supplement 4A**). Because cKO mice demonstrated shorter OSs, transmission electron microscopy (TEM) was used to investigate the ultrastructure of cKO rod OSs (**Figures 1 – figure supplement 4B–C**). The rod OSs in the cKO mouse appeared largely normal with well-organized, stacked disc membranes that maintained interdigitation with the RPE similar to the WT mouse, just shorter than WT rod OSs.

A deficiency in GLS activity could lead to a significant decrease in Glu, the main neurotransmitter used by PRs. TEM was utilized to assess the ultrastructure of rod PR ribbon synapses, which appear structurally intact in the remaining rod PRs of the cKO retina at P21 (**Figure 1 – figure supplement 4D**). To further evaluate the synaptic connectivity between PRs and the inner retina, retinal sections from cKO and WT mice at P14 were labeled with wheat germ agglutinin (WGA), a plant lectin that binds N-acetylglucosamine and sialic acid residues, to highlight PR synaptic membranes and non-synaptic membranes (McLaughlin et al., 1980). Retinal sections from P14 WT and cKO mice stained with WGA did not demonstrate significant differences in the labeling of PR synaptic membranes in the outer plexiform layer (OPL) (**Figure 1 – figure supplement 5A**). Staining of P14 retinal sections from WT and cKO mice with an antibody against Bassoon, which labels the ribbon synapses of rods and cones in the OPL (Kutsyr et al., 2021), did not demonstrate differences in the synaptic connectivity between PRs and second-order neurons (**Figure 1 – figure supplement 5B**). Additionally, retinal sections at P14, P21, and P42 were stained with antibodies against major cell-types in the inner retina, such as amacrine, ganglion, and bipolar cells, and no significant alterations in immunofluorescent patterns to suggest an inner retinal developmental abnormality were observed between cKO and WT animals (**Figure 1 – figure supplement 5C**). Accordingly, inner retinal thinning was not observed until P42 (**Figure 1 – figure supplement 5D**), when approximately 25% of cells remained in the ONL (**Figure 1H**). This thinning of the inner retinal area is likely secondary to the primary PR degeneration similar to that seen in other mouse models of retinal degeneration (Ueta et al., 2012).

Loss of *GLs* in rod photoreceptors impairs retinal function

Loss of IS/OS thickness, shorter OSs, and loss of PR cell bodies can result in functional loss. Thus, electroretinography (ERG) analysis was performed on cKO and WT animals at P21 and P42 (**Figure 2C**). As expected, cKO animals show a significant loss in rod-driven scotopic a- and b-wave amplitudes at P21, which are further decreased by P42 (**Figure 2A-2C**). Interestingly, a significant loss in photopic b-wave amplitude was also found at P42, suggesting a cone PR defect (**Figure 2D**). Immunofluorescent staining found that cone outer segments shorten as rod degeneration progresses from P14 to P42 (**Figure 2E**) supporting secondary cone degeneration, which is a common phenotype associated with rod-mediated retinal degeneration (Caruso et al., 2020).

GLS is necessary for maintenance of mature photoreceptors

In mice expressing a Cre-recombinase under the control of the rhodopsin promoter, Cre-mediated excision of floxed genomic DNA has been observed as early as P7 (Le et al., 2006). To ensure the function of GLS is not restricted to this early phase of PR development and maturation and that it is critical to fully developed PRs as well, mice homozygous for a floxed *Glsl* allele (*Glsl^{fl/fl}*) and expressing an inducible Cre-recombinase under the control of the *Pde6g* promoter (*Glsl^{fl/fl};Pde6g-Cre^{ERT2}*) (Koch et al., 2015) as well as animals expressing only the inducible Cre-recombinase (*Glsl^{wt/wt};Pde6g-Cre^{ERT2}*) were generated. This allowed for induction of cre-recombinase activity in fully-mature PRs, specifically, by the administration of tamoxifen (TAM). Animals heterozygous for *Pde6g-Cre^{ERT2}* and homozygous for either the WT or floxed *Glsl* allele (IND-cKO) were generated, and TAM was administered intraperitoneally for 5 consecutive days starting at P22 (**Figure 3C**) as retinal development is typically considered complete by P21 (Zhou et al., 2021). Significant reduction of GLS in the retina of IND-cKO mice after TAM induction was confirmed via Western blot and in PRs with immunofluorescence analysis (**Figures 3A** and **3B**). Ten days after TAM induction, IND-cKO mice began to show a thinning of the IS/OS layer that preceded and then paralleled the rapid ONL degeneration observed on longitudinal OCT and histology (**Figures 3C-3G**). Hence, GLS is also critical for the survival of fully developed PRs.

Beyond survival, GLS is also critical for PR function. ERG analyses performed 10 days after TAM induction, a time prior to major structural changes in the outer retina, demonstrated statistically significant reductions in the IND-cKO scotopic a- and b-waves as compared to the WT (**Figure 3H**). Similarly, photopic ERG demonstrated statistically significant decreases in the b-wave of the IND-cKO retina (**Figure 3I**). These data suggest that GLS-driven Gln catabolism plays a significant role not only in rod PR survival but their function as well. Additionally, *Pde6g* is expressed by rods to a significant degree but also by cones (Voigt et al., 2020). Therefore, the IND-cKO likely knocks out GLS from both rods and cones, which is in accordance with the immunofluorescence image in **Figure 3B** where GLS is not observed in rod or cone inner segments unlike in **Figure 1B** where GLS remains in cones. Hence, the reduction in IND-cKO photopic b-wave may suggest that GLS-driven Gln catabolism in cones impairs their synaptic transmission.

GLS knockout does not alter nucleotide metabolism

Gln, and its breakdown product, Glu, have several fates in cellular metabolism (**Figure 4A**), which may underly the rapid and significant PR degeneration observed when GLS is knocked out of rod PRs. PRs require nucleotides to support phototransduction as well as transcriptional efforts for the continued synthesis of OSs. The catabolism of Gln to Glu by GLS produces an ammonium ion that can be used for the synthesis of nucleotides (Yoo et al., 2020). Additionally, inhibition of GLS has previously been demonstrated to inhibit pyrimidine and purine biosynthesis (Alkan et al., 2018; Okazaki et al., 2017). To explore the role of GLS-initiated Gln catabolism in nucleotide metabolism, *in vivo* targeted liquid chromatography-tandem mass spectrometry (LC-MS/MS)




Figure 2.

Loss of *Glis* in rod photoreceptors impairs retinal function

(A) Representative scotopic and photopic electroretinography (ERG) tracings for the rod photoreceptor-specific *Glis* conditional knockout mice (cKO) compared wild-type (WT) mice at P21 and P42. (B) ERG scotopic a-wave and (C) b-wave amplitudes in cKO mice compared to WT mice at P21 and P42. A flash intensity of 32 cd*s/m² was utilized. N=3-6 animals per group. (D) ERG photopic b-wave amplitudes in cKO mice compared to WT mice at P21 and P42. A flash intensity of 100 cd*s/m² was used. N=3-6 animals per group. (E) Representative images from staining of the cone-specific marker peanut agglutinin (PNA, green) and nuclei (DAPI, blue) in retinal sections from cKO mice compared to WT mice at P14, P21 and P42. Scale bars are 40 μm. Statistical differences in (B), (C) and (D) are based on an unpaired two-tailed Student's t-test where *P < 0.05, **P < 0.01 and ***P < 0.001. Data are presented as mean ± standard error of the mean. ONL, outer nuclear layer; IS, inner segments; OS, outer segments. N=3 per group.




Figure 3.

Rod photoreceptors require GLS for maintenance and maturation

Glsl^{fl/fl} mice carrying a tamoxifen (TAM)-inducible Cre-recombinase (IND-cKO) under the control of *Pde6g* (*Glsl^{fl/fl};Pde6g-Cre^{ERT2}*) compared to mice expressing only the inducible Cre-recombinase (*Glsl^{wt/wt};Pde6g-Cre^{ERT2}*; WT). Both IND-cKO and WT were administered tamoxifen for 5 days starting at P22. (A) Quantitation of Western blot results showing decreased GLS protein levels in the retina of IND-cKO animals at 10 days after tamoxifen induction compared to WT mice. HSP90 was used as a loading control. N=3-4 animals per group. (B) Representative GLS immunofluorescence (red) in IND-cKO mice compared to the wild-type (WT) mouse 10 days after tamoxifen induction. N=3 animals per group. Scale bar is 40 μ m. (C) OCT images detailing outer retinal changes in WT and IND-cKO animals over time. Total retinal thickness (D), outer nuclear layer (ONL) thickness (E) and inner segment/outer segment (IS/OS) thickness (F) as determined by OCT for 38 days post tamoxifen. N= 5-10 eyes per group. (G) Representative hematoxylin and eosin-stained retinal sections from IND-cKO mice compared to WT mice at 10 and 38 days after tamoxifen induction. N=3 animals per group. Scale bar is 40 μ m. ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion cell layer. Quantitation of scotopic (H) and photopic (I) ERG a- and b-waves in WT and IND-cKO retina 10 days post TAM. N>12 eyes per genotype. Statistical differences in (A), (D), (E), (F), (H), and (I) are based on an unpaired two-tailed Student's t-test where *P<0.05, **P<0.01 and ***P<0.001. Data are presented as mean $\pm$ standard error of the mean.

metabolomics were performed on P14 retina from WT and cKO animals (**Supplementary File 2**). No differences in key purine or pyrimidine metabolism intermediates, such as ribose 5-phosphate (R5P), inosine monophosphate (IMP), and uridine monophosphate (UMP), were observed between the WT and cKO retina (**Figure 4B**), suggesting that GLS-mediated catabolism of Gln is not a critical pathway for nucleotide synthesis in rod PRs.

Redox balance is altered upon knockout of GLS in rod photoreceptors

Gln can also be used for the generation of the antioxidant molecule glutathione as well as in a non-canonical NADPH generating pathway (Son et al., 2013; Yoo et al., 2020). Glutathione is a tripeptide of glutamate, glycine, and cysteine. Glutamate is a product of the GLS reaction, and previous studies have demonstrated that inhibiting GLS reduces glutathione levels and increases reactive oxygen species (Li et al., 2015). As such, the $\text{NADP}^+/\text{NADPH}$ ratio was statistically significantly increased by 16% in the P14 cKO retina as compared to WT (**Figure 4C**), and the relative abundance of oxidized glutathione (GSSG) was decreased in the P14 cKO retina as compared to WT (**Figure 4D**). While an increased $\text{NADP}^+/\text{NADPH}$ ratio in the cKO retina may imply that the abundance of GSSG should be increased in the cKO retina since reduced glutathione can be recovered from the oxidized form by the conversion of NADPH to NADP^+ , the levels of GSSG have been seen to be reduced in other models where GLS is pharmacologically inhibited or genetically knocked down (Daemen et al., 2018). The lower levels of GSSG may signal an overall reduction in the biosynthesis of glutathione considering the product of the GLS reaction, glutamate, is directly necessary for its synthesis as well as indirectly responsible for cysteine via the SLC7A11 (xCT) cystine/glutamate antiporter (Yoo et al., 2020). The expression of *Slc7a11* was reduced in the P14 cKO retina (**Figure 4E**). Hence, the biosynthesis of glutathione in rod PRs may be affected directly and indirectly when GLS is knocked out. In accordance with the antioxidants NADPH and glutathione potentially being reduced, the expression of the antioxidant enzymes *Sod1* and *Sod2*, which detoxify superoxide radicals, were increased in the P14 cKO retina (**Figure 4E**). These data suggest that GLS-mediated Gln catabolism regulates redox balance in rod PRs, and the altered redox balance that is a consequence of its knockout may be one factor contributing to PR degeneration in the GLS cKO mouse.

Relative abundance of TCA cycle intermediates and mitochondrial function mostly unchanged in *Gls* cKO retina

While the changes in certain redox elements were statistically significant, they were modest alterations and may not account in full for the significant and rapid PR degeneration noted in the GLS cKO mouse. Many cancer cells that utilize aerobic glycolysis rely on Gln to replenish TCA cycle intermediates, which maintain oxidative metabolism and provide biosynthetic precursors. Considering PRs utilize aerobic glycolysis, similar to cancer cells, and have significant bioenergetic demands, we next sought to determine if rod PRs also depend on GLS-initiated glutaminolysis for TCA cycle metabolites and oxidative metabolism. Interestingly, LC-MS/MS-based targeted metabolomics demonstrated very few changes in the relative pool sizes of TCA cycle metabolites in the cKO compared to WT retina at P14 with only malate showing a statistically significant decrease (**Figure 4F**). Since GLS is enriched in the PR mitochondria (**Figure 1 – figure supplement 1B**) and previous studies demonstrated that inhibiting GLS in certain cell lines reduces mitochondrial function (Alkan et al., 2018), mitochondrial stress tests were also performed on P14 cKO and WT retina, prior to PR loss, using the BaroFuse (Kamat et al., 2023). The basal oxygen consumption rate (OCR) as well as the changes in OCR in response to oligomycin or carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) were not statistically significantly different between WT and cKO retina (**Figures 4G** and **4H**). The expression of complexes involved in oxidative phosphorylation were also unchanged between the WT and cKO retina at P14 (**Figure 4I**).




Figure 4.

GLS cKO mice maintain levels of nucleotide, glycolytic and TCA cycle metabolites and mitochondrial function but demonstrate altered redox balance

(A) Schematic summarizing the biosynthetic and bioenergetic roles of glutamine. (B) Relative abundance of key intermediates in nucleotide metabolism in the retina of WT and cKO mice at P14 as determined by targeted metabolomics. N=6-7 animals per group. Relative abundance is the ion intensity normalized to the WT. (C) The NADP⁺/NADPH ratio, as determined by bioluminescence assay, is significantly increased in the cKO as compared to WT retina at P14. N=5-6 animals per group. (D) Relative abundance of GSSG in the retina of WT and cKO mice at P14, prior to PR degeneration, as determined by targeted metabolomics. N=6-7 animals per group. (E) qRT-PCR of genes related to redox homeostasis are significantly altered in cKO compared to WT mice. N=6 animals per group. (F) Relative abundance of metabolites in glycolysis and the TCA cycle in WT and cKO retina at P14. N=6-7 animals per group. (G) Mitochondrial stress test carried out on isolated WT and cKO retina at P14 using the BaroFuse. The baseline was established by perfusing the tissue for 90 min and then oligomycin, FCCP, and KCN were injected into the perfusate sequentially as indicated. (H) Comparison of the effects of oligomycin and FCCP on OCR in P14 WT and cKO retina. N=6-8 animals per group. (I) Western blot analysis and quantitation of the mitochondrial electron transport chain complexes show no differences between WT and cKO retina. N=3-4 animals per group. Fold change is in relation to WT. Statistical differences in (B-F), (H) and (I) are based on an unpaired two-tailed Student's t-test where *P<0.05. Data are presented as mean ± standard error of the mean. R5P: ribose 5-phosphate, IMP: inosine monophosphate, UMP: uridine monophosphate, GSSG: glutathione disulfide, F6P: fructose 6-phosphate, DHAP: dihydroxyacetone phosphate, 2PG: 2-phospho-D-glycerate, PEP: phosphoenolpyruvate, Pyr: pyruvate, Lac: lactate, Cit: citrate, α-KG: alpha-ketoglutarate, Suc: succinate, Mal: malate, Oligo: oligomycin, FCCP: carbonyl cyanide p-trifluoromethoxyphenylhydrazine, KCN: potassium cyanide, CI-NDUFB8: complex 1, NADH:ubiquinone oxidoreductase subunit B8, CII-SDHB: complex 2, succinate dehydrogenase complex iron sulfur subunit B, CIII-UQCRC2: complex 3, ubiquinol-cytochrome c reductase core protein 2, CV-ATP5A: complex 5, ATP synthase F1 subunit alpha, HSP90: heat shock protein 90.

As stated earlier, a Cre-recombinase under the control of the rhodopsin has been shown to be activated as early as P7 (Li et al., 2005). To circumvent this early knockout of GLS, rod PRs may rewire their metabolism, as has been seen in other conditional knockout mouse models with this Cre-recombinase system (Subramanya et al., 2023; Wubben et al., 2017), to utilize different fuel sources for the maintenance of the TCA cycle and mitochondrial function. To this end, the expression of genes involved in glycolysis, pyruvate metabolism, and the TCA cycle were examined in the P14 cKO and WT retina before PR degeneration. Numerous genes were significantly altered in the cKO retina across the metabolic pathways (**Figure 4 – figure supplement 1**). The expression of multiple genes in glycolysis and pyruvate metabolism were increased (**Figure 4 – figure supplement 1A**), possibly suggesting that rod PRs are stimulating glucose oxidation to maintain the TCA cycle and mitochondrial function similar to that seen in certain cancer cells when GLS is inhibited (Okazaki et al., 2017). In contrast, the only significant change in expression of these genes in eyecup tissue was *Pdk4*, with a significant decrease in cKO vs WT mice (**Figure 4 – figure supplement 2**). However, targeted metabolomics demonstrated only a minor change in glycolytic intermediates between the cKO and WT retina at P14 (**Figure 4F**). As targeted metabolomics at the P14 timepoint provides only a snapshot of the pool size, additional analyses with stable isotope tracing metabolomics are necessary to assess nutrient utilization, in this case glucose. Uniformly-labeled, $^{13}\text{C}_6$ -glucose was intraperitoneally injected in P14 WT and cKO mice, and the retina harvested 45 minutes later for metabolomic analysis via LC-MS/MS (Yuan et al., 2019). No change in the fractional labeling of glycolytic intermediates was observed between the WT and cKO retina, and more so, an increase in the fractional labeling of TCA cycle intermediates was not observed in the cKO retina (**Figures 5A, 5B** and **Figure 5 – figure supplement 1**). Hence, it is unlikely rod PRs are increasing glucose oxidation to maintain the levels of TCA cycle intermediates and mitochondrial function.

GLS cKO in rod photoreceptors decreases the utilization of Gln in the TCA cycle

We initially anticipated that GLS knockout in rod PRs would significantly alter the relative TCA cycle metabolite pools in the cKO retina, similar to what is seen in Gln-dependent cancer cells (Daemen et al., 2018; Yang et al., 2017), as PRs and cancer cells have common metabolic relationships (Du, Cleghorn, Contreras, Linton, et al., 2013; Ng et al., 2015; Rajala, 2020), and previous *ex vivo* studies demonstrated Gln can supplement the TCA cycle in the retina (Grenell et al., 2019; Tsantilas et al., 2021). However, as observed in **Figure 4F**, very few changes were observed in the relative pool sizes of TCA cycle metabolites in the cKO compared to WT retina. To delineate how Gln is metabolized, we used LC-MS/MS to trace the metabolic fate of uniformly-labeled $^{13}\text{C}_5$ -Gln in the WT and cKO retina *in vivo* at P14. Uniformly-labeled $^{13}\text{C}_5$ -Gln was intraperitoneally injected into the mice and the retina harvested 45 minutes later. In accordance with GLS being the enzyme that initiates glutaminolysis (Yang et al., 2017), the fractional labeling of glutamate, TCA cycle intermediates, and pyruvate was decreased in the P14 cKO retina as compared to WT (**Figures 5C, 5D** and **Figure 5 – figure supplement 2**). Previous *ex vivo* studies have also observed that isotopically labeled $^{13}\text{C}_5$ -Gln can contribute to m+3 pyruvate in the retina possibly via the decarboxylation of m+4 malate by malic enzyme (Grenell et al., 2019).

α -ketoglutarate (α -KG) is the main entry point for Gln into the TCA cycle (**Figure 5C**) and has been shown to be a key metabolite in Gln metabolism (Yang et al., 2017). *In vivo* stable isotope tracing demonstrated decreased incorporation of Gln carbons into α -KG in the cKO retina compared to the WT (**Figure 5D**). Furthermore, supplementation with α -KG has been shown to improve PR survival in mouse models of inherited retinal disease (Wert et al., 2020). So, we attempted to rescue the PR degeneration phenotype in the GLS cKO mouse by supplementing cKO animals with 10 mg/mL of α -KG in the drinking water starting at P4 (Rowe et al., 2021; Wert et




Figure 5.

Loss of GLS in rod photoreceptors has significant effects on TCA cycle metabolism with only partial rescue upon α -KG supplementation

(A) Schematic summarizing $^{13}\text{C}_6$ -glucose labeling in glycolytic and TCA cycle intermediates. (B) Fractional labeling of glycolytic and TCA cycle metabolites in the retina following intraperitoneal injection of $^{13}\text{C}_6$ -glucose in WT and cKO mice at P14. N=5-6 animals per group. (C) Schematic summarizing $^{13}\text{C}_5$ -Gln labeling in the TCA cycle. (D) Fractional labeling of TCA cycle metabolites in the retina following intraperitoneal injection of $^{13}\text{C}_5$ -Gln in WT and cKO mice at P14. N=6-11 animals per group. (E) ONL thickness in cKO mice at P22 as assessed by OCT following α -KG supplementation (10 mg/mL) or vehicle (water) in the drinking water from P4-P22. N=5 animals per group. Statistical differences in (B), (D) and (E) are based on an unpaired two-tailed Student's t-test where * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$. Data are presented as mean $\pm$ standard error of the mean. F6P: fructose 6-phosphate, FBP: fructose 1,6-bisphosphate, DHAP: dihydroxyacetone phosphate, 2PG: 2-phosphoglycolate, PEP: phosphoenolpyruvate, Pyr: pyruvate, Lac: lactate, Gln: glutamine, Glu: glutamate, α -KG: alpha-ketoglutarate, Suc: succinate, Fum: fumarate, Mal: malate, OAA: oxaloacetate, Cit: citrate, Asp: aspartate.

al., 2020 [\[1\]](#)). A small, but significant increase in ONL thickness was identified in α -KG-treated animals at P22 using OCT (**Figure 5E** [\[1\]](#)). These data further suggest that Gln's role in supporting the TCA cycle is not the major mechanism by which PRs utilize Gln to suppress PR apoptosis.

Gls cKO retina has reduced NEAA levels and an upregulated integrated stress response

Gln and GLS-derived Glu play a central role in the biosynthesis of several NEAAs (Yoo et al., 2020 [\[1\]](#)). So, the disruption of GLS-initiated Gln catabolism may be causing a disruption in available NEAAs for biomass production. To this end, targeted metabolomics on the cKO retina at P14 showed a significant increase in the substrate of GLS, Gln, and a significant decrease in its NEAA product Glu, further illustrating loss of GLS function (**Figures 6A** [\[1\]](#) and **6B** [\[1\]](#)). The NEAA Asp was also significantly reduced (**Figure 6A** [\[1\]](#)), which is consistent with previous studies that inhibited GLS in cancer cells (Alkan et al., 2018 [\[1\]](#); Okazaki et al., 2017 [\[1\]](#)). Notably, similar decreases in Glu and Asp were noted in IND-cKO compared to WT retina at a point prior to PR degeneration (**Figure 6 – figure supplement 1** [\[1\]](#)).

Deprivation of amino acids can activate the integrated stress response (ISR), which responds to an array of stressors to maintain cellular homeostasis and particularly, protein homeostasis. Acutely, ISR activation can protect cells by temporarily halting protein synthesis. However, chronic ISR activation and global protein synthesis inhibition can trigger apoptosis (Pakos-Zebrucka et al., 2016 [\[1\]](#)). ISR activation leads to phosphorylation of the alpha subunit of eukaryotic translation initiation factor 2 (eIF2 α), which reduces global protein synthesis while preferentially allowing for the translation of activating transcription factor 4 (ATF4) (Pakos-Zebrucka et al., 2016 [\[1\]](#)). Considering the reduced levels of NEAAs Glu and Asp, retinas from P14 GLS cKO animals were assayed for these ISR components using Western blotting (**Figures 6C** [\[1\]](#) and **6D** [\[1\]](#)), which demonstrated increased levels of both phosphorylated eIF2 α and total ATF4, suggesting ISR activation. To determine if global protein synthesis was affected in GLS cKO retinas, we applied the SUnSET (Surface SEnsing of Translation) method to quantify protein synthesis in WT and cKO retinas at P14 (Fort et al., 2022 [\[1\]](#)). These data show a significant decrease of puromycin incorporation into nascent polypeptide chains, indicating a decrease in global protein synthesis consistent with ISR activation (**Figures 6E** [\[1\]](#) and **6F** [\[1\]](#)). To determine if PR degeneration could be rescued by inhibiting the ISR, ISRIB (2.5 mg/kg) was administered systemically daily starting at P5. In WT mice, ISRIB treatment did not impact retinal anatomy as assessed by OCT at P21 (**Figure 6 – figure supplement 2A** [\[1\]](#)). Notably, at P21, ISRIB treatment in the cKO mouse significantly increased ONL thickness compared to vehicle using OCT (**Figure 6G** [\[1\]](#)) and histology (**Figure 6 – figure supplement 3A** [\[1\]](#)), and this effect was sustained at P28 (**Figure 6 – figure supplement 2B** [\[1\]](#)). These results suggest that ISR activation is downstream of Gln catabolism and contributes to PR degeneration in the cKO mouse retina.

Previous work has demonstrated that when certain cancer cell lines are faced with Gln deprivation, α -KG alone does not restore cell proliferation (Pavlova et al., 2018 [\[1\]](#); Son et al., 2013 [\[1\]](#)). Rather, NEAA supplementation is necessary to restore proliferation in these reports. Similarly, α -KG supplementation had an incomplete survival impact on PR survival, showing approximately 10% increase in ONL thickness in cKO animals (**Figure 5E** [\[1\]](#)). Since systemic or intravitreal glutamate supplementation has demonstrated retinal toxicity (Sisk & Kuwabara, 1985 [\[1\]](#)), and Asp concentration is normally low in plasma with poor cellular membrane permeability (Sullivan et al., 2018 [\[1\]](#)), we investigated other NEAA approaches to rescue the PR degeneration observed in the GLS cKO mouse. The potential of asparagine (Asn) to rescue PR degeneration was chosen due to the growing list of Asn-mediated cellular processes that improve cancer cell survival and growth under metabolic stress, such as Gln deprivation. These Asn-mediated cellular processes include inhibiting ISR-induced apoptosis (Zhang et al., 2014 [\[1\]](#)), (1) promoting protein synthesis (Pavlova et al., 2018 [\[1\]](#)), and maintaining Asp pools (Halbrook et al., 2022 [\[1\]](#)). All these processes may be relevant to the PR degeneration observed in the cKO mouse.




Figure 6.

GLS cKO retina demonstrates decreased nonessential amino acids, ISR activation, and decreased global protein synthesis

(A) Amino acids significantly altered in the *GLS* cKO mouse retina at P14. Relative abundance is the ion intensity relative to WT retina. N=6-7 animals per group. (B) Ratio of glutamine to glutamate in WT and cKO retina. N=6-7 animals per group. (C) Western blot of ISR proteins phospho-eIF1 α^{S51} (p-eIF2 α), total eIF2 α and ATF4 in WT and cKO mice. N=3-4 animals per group. (D) Quantitation of Western blot in Panel C. (E) Western blot of protein puromycinylation in the WT and cKO mouse retina at P14 harvested 30 min after systemic puromycin administration. (F) Quantitation of puromycin incorporation in WT and cKO retina. N=4-5 animals per group. (G) ONL thickness at P21 in cKO mice as assessed by OCT following intraperitoneal injection of ISRIB (2.5 mg/kg) or vehicle (50% PEG 400, 43.4% saline, 6.6% DMSO) from P5-P21. N=3-6 animals per group. (H) ONL thickness at P21 in cKO mice as assessed by OCT following intraperitoneal injection of Asn (200 mg/kg) or vehicle (PBS) from P5-P21. N=3-5 animals per group. Statistical differences in (A), (B), (D), (F), (G), and (H) are based on an unpaired two-tailed Student's t-test where *P < 0.05, **P < 0.01 and ***P < 0.001. Data are presented as mean $\pm$ standard error of the mean. Gln: glutamine, Glu: glutamate, Asp: aspartate, ONL: outer nuclear layer.

Following administration of systemic Asn (200 mg/kg) daily starting at P5, *in vivo* analysis of retinal structure via OCT and histology at P21 demonstrated a significant improvement in ONL thickness compared to animals treated with vehicle, with a 22-26% increase in ONL thickness in some retinal locations (**Figure 6H** [↗](#) and **Figure 6 – figure supplement 3B** [↗](#)).

Discussion

In this study, we demonstrate for the first time the importance of Gln catabolism in PR metabolism, function, and survival *in vivo*. Rod PRs lacking GLS demonstrated rapid and complete degeneration with concomitant functional loss. Mechanistically, lack of GLS-initiated Gln catabolism in rod PRs, specifically, resulted in changes in redox homeostasis and a decrease in the fractional labeling of TCA cycle intermediates from stable isotope-labeled Gln *in vivo*. However, Gln anaplerosis did not appear to be driving the TCA cycle as targeted metabolomics showed few changes in the relative levels of TCA cycle intermediates, and supplementation with α KG showed a modest rescue effect in PR degeneration. The NEAAs Glu and Asp were found to be significantly decreased in the cKO mouse retina, and in accordance with this AA deprivation, the ISR was upregulated with a reduction in global protein synthesis suggesting Gln catabolism in rod PRs plays a significant role in supporting biomass production via NEAA synthesis. Interestingly, supplementing cKO mice with Asn significantly rescued PR degeneration, revealing a novel link between Gln and Asn metabolism in PRs. This work further demonstrates that PRs have the flexibility to utilize fuel sources other than glucose to meet their metabolic needs and that Gln is a critical amino acid that supports PR cell biomass, redox balance, and survival.

The rapid degenerative phenotype observed with conditional deletion of *Gls* in rod PRs was unexpected considering glucose is viewed as the primary substrate for the retina and PRs (Swarup et al., 2019 [↗](#)). The *Gls* cKO mouse demonstrated comparable ONL thickness to WT controls at P14 with approximately 50% loss of PR cell bodies by P21 (**Figure 1 – figure supplement 3A-B** [↗](#)) and near complete PR degeneration by P84 (**Figure 1C** [↗](#)). In contrast, conditional deletion of numerous enzymes within central glucose metabolism, including hexokinase 2 (HK2) (Weh et al., 2020 [↗](#)), pyruvate kinase M2 (PKM2) (Wubben et al., 2017 [↗](#)), glucose transporter 1 (GLUT1) (Daniele et al., 2022 [↗](#)), mitochondrial pyruvate carrier 1 (MPC1) (Grenell et al., 2019 [↗](#)), and lactate dehydrogenase A (LDHA) (Rajala et al., 2023 [↗](#)), results in a slower, age-related degeneration. For example, loss of GLUT1 in the retina and rod PRs does not demonstrate 50% thinning of the ONL until approximately 4 months of age (Daniele et al., 2022 [↗](#)), and the others noted above demonstrate even slower outer retinal degeneration. It has also been shown that PRs can utilize fatty acid β -oxidation for energy (Joyal et al., 2016 [↗](#)). Interestingly, in retinas lacking the very-low-density lipoprotein receptor (VLDLR), which facilitates fatty acid uptake, the PRs remained largely intact. In any of these transgenic mouse models, PRs may use other transporters to take up fatty acids or glucose, rewire their metabolism, or utilize metabolic redundancies to maintain metabolic homeostasis and stave off degeneration (Subramanya et al., 2023 [↗](#); Wubben et al., 2017 [↗](#)). Our data show that any metabolic reprogramming that is occurring in the cKO mouse retina appears unable to significantly circumvent the significant and rapid PR degeneration suggesting the importance of Gln catabolism in rod PRs. Furthermore, inducing GLS knockdown in mature PRs also demonstrated rapid PR degeneration (**Figure 3** [↗](#)).

Traditionally, it has been thought that Gln maintains cell survival through Gln-derived α -KG and maintenance of the TCA cycle (Zhang et al., 2014 [↗](#)). (1) While stable isotope tracing with uniformly labeled Gln demonstrated a reduction in fractional labeling of the TCA cycle intermediates (**Figure 5D** [↗](#)), targeted metabolomics showed few changes in the relative levels of the intermediates (**Figure 4F** [↗](#)) and supplementation with α KG did not rescue PR degeneration to a large degree (**Figure 5E** [↗](#)). Gln is also required for the biosynthesis of NEAAs, and the NEAAs Glu and Asp were reduced nearly two-fold in the retina of cKO mouse (**Figure 6A** [↗](#)). Glu is the product of the GLS reaction so it is not surprising that Glu was substantially reduced. Glu is

involved in glutathione biosynthesis directly and indirectly, can be converted into α -KG to enter the TCA cycle, and is crucial for the biosynthesis of NEAAs including Asp (Yoo et al., 2020 [↗](#)). Alterations in redox homeostasis were observed in the retina of the cKO mouse in accordance with the reduced level of Glu (**Figures 4C [↗](#)–4E [↗](#)**). While these redox imbalances likely contribute to the PR degeneration observed, it is unlikely these small but statistically significant changes fully account for the rapid and complete PR degeneration. Future rescue studies with antioxidants, such as N-acetylcysteine, are needed to shed light on the role of redox imbalance in this novel transgenic mouse model.

Previous *ex vivo* studies demonstrated that Gln can contribute carbons to Asp synthesis via the TCA cycle and/or contribute a nitrogen via glutamate-oxaloacetate transaminases (Du, Cleghorn, Contreras, Lindsay, et al., 2013; Xu et al., 2020 [↗](#)), the latter of which have been shown to be critical in the Gln metabolic rewiring of cancer cells (Son et al., 2013 [↗](#)) and more recently, in rod PR metabolism, function, and survival (Subramanya et al., 2023 [↗](#)). Our data shows that Gln-derived carbons via Glu are entering the TCA cycle *in vivo* and contributing to the synthesis of Asp (**Figure 5 [↗](#)**), and this contribution is reduced in the cKO mouse retina. Based on the previous studies noted above, Gln may also be contributing to Asp synthesis via the glutamate-oxaloacetate transaminase catalyzed transfer of a nitrogen from Glu. Future studies using amine-labeled Gln are needed to dissect the contributions of Gln-derived carbon versus nitrogen to the synthesis of Asp in the retina *in vivo*.

Asp is a proteinogenic amino acid that has many biosynthetic roles (Alkan et al., 2018 [↗](#)), and *ex vivo* neural retina studies have suggested that the retina needs Asp to maintain its metabolic homeostasis (Li et al., 2020 [↗](#)). In accordance with this data, Asp was one of the few metabolites that was significantly reduced in the retina of cKO mouse prior to rapid PR degeneration (**Figure 6A [↗](#)**). Based on *ex vivo* studies, it has been postulated that Asp utilization in the retina is necessary to maintain aerobic glycolysis and mitochondrial metabolism through the recycling of NADH to NAD⁺, shuttling electrons into the mitochondria via the malate-aspartate shuttle, replenishing oxaloacetate for biosynthesis and/or producing pyruvate via malic enzyme (Li et al., 2020 [↗](#)). Our *in vivo* metabolomics data further support this last point as uniformly labeled Gln contributed to m+3 pyruvate in the retina possibly via the decarboxylation of m+4 malate by malic enzyme (**Figure 6D [↗](#)**). The m+4 isotopologue of malate could arise from m+4 Asp via glutamate-oxaloacetate transaminase (GOT) and malate dehydrogenase (MDH) as part of the malate-aspartate shuttle. Furthermore, the decarboxylation of malate by malic enzyme produces pyruvate and also NADPH. A previous study revealed high malic enzyme activity in the retina and suggested it is capable of producing NADPH in the retina (Winkler et al., 1986 [↗](#)). The observation that malate was reduced (**Figure 4F [↗](#)**) and the NADP⁺/NADPH ratio was increased (**Figure 4C [↗](#)**) in the cKO retina further supports a metabolic pathway in PRs where the Gln carbon skeleton is converted into Asp, then OAA, malate, and finally, pyruvate via the cytosolic enzymes of the malate-aspartate shuttle and malic enzyme. This non-canonical Gln metabolism pathway is also necessary to sustain cancer cell growth (Son et al., 2013 [↗](#)).

The m+4 isotopologue of oxaloacetate (OAA) can also be converted to m+3 pyruvate by phosphoenolpyruvate carboxykinase (PEPCK). A recent *ex vivo* study postulated that PRs can utilize different metabolic cycles, such as the Cahill cycle or mini-Krebs cycle, to uncouple glycolysis from oxidative phosphorylation (Chen et al., 2024 [↗](#)). These cycles are fueled by Gln and require PEPCK to replenish pyruvate from OAA. While the lack of m+4 citrate in both the P14 WT and cKO retina and decrease in m+2 citrate in the cKO retina (**Figure 5 – figure supplement 2 [↗](#)**) may further support the existence of these previously postulated metabolic cycles, which avoid the citrate synthase step of the TCA cycle as well as others, further studies are needed to confirm the activity of these pathways in the retina and specifically, the PRs.

Interestingly, we did not observe any significant changes in glycolysis or mitochondrial function in the cKO retina despite a reduction in the relative level of Asp. It is hypothesized that Asp catabolism sustains glycolysis and oxidative phosphorylation in the retina by recycling NADH to NAD^+ and shuttling electrons into the mitochondria via the malate-aspartate shuttle (Li et al., 2020). However, in the GLS cKO retina, the NAD^+/NADH ratio ($P = 0.125$) and its oft-used proxy, the pyruvate/lactate ratio ($P = 0.192$), were not statistically significantly altered. Compensatory metabolic rewiring, which has been seen in other conditional knockout mouse models using the same Cre-recombinase system (Subramanya et al., 2023; Wubben et al., 2017), may be maintaining the NAD^+/NADH ratio in the cKO retina at P14 when there are equal numbers of PRs in the WT and cKO retina. To this end, the expression of the *Mdh1* gene, which is a component of the malate-aspartate shuttle that recycles NADH to NAD^+ in the cytoplasm, was upregulated in the cKO retina (Figure 4-figure supplement 1). The conversion of dihydroxyacetone phosphate (DHAP) to glycerol 3-phosphate via glycerol 3-phosphate dehydrogenase is an alternative pathway to regenerate NAD^+ . DHAP levels were significantly reduced in the cKO retina possibly suggesting increased activity of this pathway (Figure 4F). Yet, the gene expression and activity of glycerol 3-phosphate dehydrogenase in the retina has been shown to be low (Adler & Klucznik, 1982; Voigt et al., 2020).

Asp is also the immediate precursor to Asn. Intracellular levels of Asn are the lowest among the NEAAs (Zhang et al., 2014), so it was not surprising that Asn was below the level of confident detection in our targeted metabolomics analyses on the WT or cKO retina. Asn had been shown to rescue cancer cells from ISR-induced apoptosis, increase protein synthesis during Gln deprivation, and protect the Asp pool (Halbrook et al., 2022; Pavlova et al., 2018; Zhang et al., 2014). As metabolic similarities exist between PRs and cancer cells (Du, Cleghorn, Contreras, Linton, et al., 2013; Ng et al., 2015), and the *Gls* cKO mouse retina demonstrated ISR activation and apoptosis (Figures 1G-1I and 6C-6D), decreased protein synthesis (Figure 6E), and decreased Asp (Figure 6A), Asn supplementation was explored and proved effective as observed in the rescue of PR degeneration in the cKO mouse (Figure 6H). These results offer the first evidence for a role of Asn downstream of Gln metabolism in PR survival, but further studies are necessary to define which of the Asn-mediated processes is crucial for PR neuroprotection following targeted *Gls* knockout. Of note, while Asn supplementation provided a greater PR neuroprotective effect than α -KG, the two supplements had different routes of administration with Asn being provided intraperitoneally as previously described (Xu et al., 2021) and α -KG being provided in the drinking water as had previously improved PR survival in a mouse model of inherited retinal disease (Wert et al., 2020). It is unclear if supplementing α -KG to GLS cKO animals intraperitoneally would further boost retinal protection. Yet, the relative abundance of α -KG and most other TCA cycle intermediates were unchanged between WT and cKO retinas suggesting Gln may not be driving the TCA cycle in PRs.

The ISR is activated with reduced global protein synthesis in the *Gls* cKO mouse retina (Figures 6C-6F) and inhibiting the ISR with ISRIB delayed PR degeneration (Figure 6G). ISR activation is a hallmark of neurodegenerative diseases including retinal degenerative diseases (Gorbatyuk et al., 2020). Chronic activation of the ISR and protein synthesis attenuation has been observed in a multitude of preclinical models of retinal degeneration and shown to contribute to PR degeneration (Gorbatyuk et al., 2020). Metabolic dysfunction and deprivation of key nutrients, such as glucose and amino acids, are not only known stressors that activate the ISR (Pakos-Zebrucka et al., 2016), but also underlying mechanisms of PR degeneration (Ait-Ali et al., 2015; Caruso et al., 2020; Pan et al., 2021). There is a paucity of studies that examine the link between metabolism and the ISR in retinal degenerative disease, which is a critical gap in our knowledge since identification of molecular and metabolic pathways triggering PR death is likely to reveal novel targets for therapeutic intervention. A recent study demonstrated that imbalanced pro-apoptotic ISR signaling contributes to deoxysphingolipid-mediated toxicity in retinal disease and identified potential therapeutic strategies, such as pharmacologic enhancement of ATF6 activity or treatment with ATF6-regulated neurotrophic factor MANF, that attenuate the associated

retinal degeneration (Rosarda et al., 2023 [DOI](#)). The novel transgenic mouse model of retinal degeneration described here provides a unique tool to obtain further insight on the nexus of metabolism, ISR activation, and protein synthesis attenuation in PR degeneration to identify pharmacologically and metabolically tractable nodes for therapeutic intervention.

Collectively, our results indicate that Gln is critical for maintaining the pools of key biosynthetic precursors, Glu and Asp, in rod PRs and disrupting Gln catabolism results in profound loss of PR function and survival in part secondary to an imbalance in ISR activation and protein synthesis attenuation. Glucose remains central in PR metabolism, but improving our understanding of other metabolic pathways that support PR function and survival and how these metabolic pathways connect with cell death mechanisms could be transformative for preventing PR degeneration and vision loss in a multitude of retinal degenerative diseases.

Ideas and Speculation

Beyond glucose, the metabolic pathways integral to photoreceptor health remain largely unknown. This is a critical knowledge gap as identification of these pathways is likely to reveal new strategies for therapeutic intervention. This work demonstrates that rod photoreceptors depend on glutamine catabolism and suggests a metabolic axis where glutamine catabolism in rod photoreceptors supports the production of aspartate and asparagine to promote anabolism and prevent signaling through the pro-apoptotic integrated stress response pathway. Considering activation of the integrated stress response is a hallmark of neurodegenerative diseases and metabolic dysfunction underlies photoreceptor degeneration, defining the pathways by which glutamine catabolism contributes to photoreceptor health is likely to identify nodes that may be targeted to make photoreceptors less vulnerable to stress.

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (<i>Mus musculus</i>)	<i>Gls^{fl/fl}</i>	PMID: 26778975		
Genetic reagent (<i>Mus musculus</i>)	Rho-cre	PMID: 16636658		
Genetic reagent (<i>Mus musculus</i>)	<i>Gls^{fl/fl};Rho-Cre⁺</i>	This study		
Genetic reagent (<i>Mus musculus</i>)	<i>Pde6g-Cre^{ERT2}</i>	PMID: 26301813		
Genetic reagent (<i>Mus musculus</i>)	<i>Gls^{wt/wt};Pde6g-Cre^{ERT2}</i>	This study		

Genetic reagent (<i>Mus musculus</i>)	<i>GLS^{fl/fl};Pde6g-Cre^{ERT2}</i>	This study		
Antibody	(Mouse monoclonal) anti-GLS	Proteintech	66265-1-Ig	1:200; immunofluorescence
Antibody	(Rabbit polyclonal) anti-GLS2	Abcam	Ab113509	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-Rhodopsin	Abcam	Ab5417	1:1,000; immunofluorescence
Antibody	(Goat polyclonal) anti-OPN1MW/LW	Santa Cruz Biotechnology	Sc-22117	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-BRN3A	Santa Cruz Biotechnology	Sc-8429	1:100; immunofluorescence
Antibody	(Rabbit polyclonal) anti-calretinin	Millipore Sigma	C7479	1:100; immunofluorescence
Antibody	(Rat monoclonal) anti-GFAP	Thermo Fisher	13-300	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-Chx10	Santa Cruz Biotechnology	Sc-365519	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-Bassoon	Enzo	SAP7F407	1:1,000; immunofluorescence
Antibody	Goat-anti-mouse Alexa 488	Invitrogen	A11001	1:1,000; immunofluorescence
Antibody	Donkey-anti-mouse Alexa 594	Jackson ImmunoResearch Laboratories	715-585-151	1:500; immunofluorescence
Antibody	Donkey-anti-rabbit Alexa 594	Jackson ImmunoResearch Laboratories	711-585-152	1:500; immunofluorescence
Antibody	Donkey-anti-goat Alexa 647	Invitrogen	A21447	1,2000; immunofluorescence

Antibody	Donkey-anti-rat Alexa 594	Invitrogen	A21209	1:500; immunofluorescence
Antibody	PNA Alexa 594 conjugate	Invitrogen	L32459	1:200; immunofluorescence
Antibody	WGA Alexa 594 conjugate	Invitrogen	W11262	1:1,000; immunofluorescence
Antibody	(Rabbit polyclonal) anti-KGA-specific GLS	Proteintech	20170-1-AP	1:1,000; western
Antibody	(Rabbit polyclonal) anti-GAC-specific GLS	Proteintech	19959-1-AP	1:1,000; western
Antibody	(Rabbit polyclonal) anti-GLS	Proteintech	12855-1-AP	1:1,000; western
Antibody	(Rabbit polyclonal) anti-VDAC	Cell Signaling Technology	4866	1:1,000; western
Antibody	(Mouse monoclonal) Anti-TIM23	BD Biosciences	611223	1:1,000; western
Antibody	(Mouse monoclonal) anti-HSP90	Cell Signaling Technology	4877	1:2,000; western
Antibody	Horse-anti-mouse-HRP	Cell Signaling Technology	7076	1:5,000; western
Antibody	Goat-anti-rabbit-HRP	Cell Signaling Technology	7074	1:5,000; western
Antibody	(Rabbit monoclonal) anti-eIF2 α	Cell Signaling Technology	5324	1:1,000; western
Antibody	(Rabbit monoclonal) anti-Phospho-eIF2 α	Cell Signaling Technology	3398	1:5,000; western

Antibody	(Rabbit polyclonal) anti-ATF4	Invitrogen	PA5-27576	1:1,000; western
Antibody	(Mouse monoclonal) anti-puromycin	Biolegend	381502	1:1,000; western
Antibody	(Mouse monoclonal) Total OXPHOS rodent antibody	Abcam	Ab110413	1:1,000; western

Methods and materials

Animals

All animals were treated in accordance with the Association for Research in Vision and Ophthalmology Statement for the Use of Animals in Ophthalmic and Vision Research. The protocol was approved by the University Committee on Use and Care of Animals of the University of Michigan (PRO00011133). All animals were housed under standard husbandry conditions at room temperature in 12-hour light/12-hour dark cycles unless explicitly stated in the text. Both male and female mice were used for all experiments. A transgenic mouse where *Gls* is selectively deleted from rod photoreceptors was created by crossing mice with Lox-P sites flanking exon 1 of the *Gls* gene (*Gls^{fl/fl}*, courtesy of Dr. Stephen Rayport, Columbia University) with Rho-Cre mice, in which Cre-recombinase expression is driven specifically in rod PRs. *Gls^{fl/fl}* and Rho-Cre mice have been previously described (Le et al., 2006 [DOI](#); Mingote et al., 2015 [DOI](#)). Animals were screened for the *rd8* mutation (Mattapallil et al., 2012 [DOI](#)). Alpha-ketoglutarate (α -KG, Millipore-Sigma, St. Louis, MO, USA, Cat # K1128) was provided to mice in their drinking water (10 mg/mL) starting at P4. Asparagine (200 mg/kg, Millipore-Sigma, Cat # A4159) or vehicle (PBS) was injected IP starting at P5. ISRIB (2.5 mg/kg, Cayman Chemical, Ann Arbor, MI, USA, Cat # 16258) or vehicle (50% PEG 400, 43.4% saline, 6.6% DMSO) (Halliday et al., 2015 [DOI](#)) was injected IP starting at P5. For inducible deletion of *Gls*, *Gls^{fl/fl}* mice were crossed to *Pde6g*-Cre^{ERT2} mice (courtesy of Dr. Stephen Tsang, Columbia University) (Koch et al., 2015 [DOI](#)). The Cre-recombinase was activated via IP injection of tamoxifen (Millipore-Sigma, Cat # T5648) at a concentration of 100 mg/kg bodyweight for 5 consecutive days. Whole retinas were extracted from animals using the “cut-and-pick” method as previously described (Winkler, 1981 [DOI](#)), being careful to remove any adherent ciliary body or RPE before processing. Total retina was then either snap frozen on dry ice (Western blotting, metabolomics), immersed in RNAlater (Qiagen, Hilden, Germany, Cat # 76104) for qRT-PCR, or used immediately (BaroFuse or NADP⁺/NADPH bioluminescent assay).

In vivo functional and structural assessment

Visual function was assessed as previously described (Weh et al., 2020 [DOI](#); Wubben et al., 2017 [DOI](#)). Mice were anesthetized using a mixture of ketamine/xylazine (90/10 mg/kg) and their pupils were dilated using 1% tropicamide and 2.5% phenylephrine ophthalmic drops. Retinal function was determined using a Diagnosys Celeris ERG system (Diagnosys LLC, Lowell, MA, USA) following overnight dark adaptation. *In vivo* retinal thickness was measured using the Envisu-R SD-OCT imager (Leica Microsystems Inc., Buffalo Grove, IL, USA). A 1.5 mm horizontal B-scan (1000 A-scans x 100 frames) and a 1.5 mm x 1.5 mm rectangular volume (1000 A-scans x 36 B-scans x 6 frames) were captured, registered and averaged using the built-in software, and analyzed using the Diver 1.0 software suite (Leica Microsystems). Images were segmented manually to determine

total retinal, outer nuclear layer, and combined inner segment/outer segment thickness. Measurements were taken at 15 points on a 9×9 grid and averaged as previously described (Weh et al., 2022 [DOI](#)).

Immunofluorescence

Mouse eyes were enucleated and immersed in 4% paraformaldehyde overnight before embedding in paraffin and sectioned at 6 mm thickness. Following standard protocols, sections were de-paraffinized and antigen retrieval performed as previously described (Weh et al., 2020 [DOI](#); Wubben et al., 2017 [DOI](#)). Sections were blocked with 1% bovine serum albumin (BSA, Millipore-Sigma, Cat # A9647) in 1X phosphate buffered saline (PBS, Thermo Fisher Scientific, Waltham, MA, USA, Cat # BP399) with 0.125% Tween 20 (Thermo Fisher Scientific, Cat # BP337) and 10% normal goat serum prior to incubating with primary antibody in 1% BSA and 1% normal goat serum overnight at 4°C. Slides were then washed, secondary antibody applied for 1 hour at room temperature before washing, and finally, counterstained with DAPI (Thermo Fisher Scientific, Cat # P36930). Images were obtained on a Leica DM6000 microscope with a 40X objective. The antibodies used for immunofluorescence are found in the key resources table.

TUNEL staining and cell counts

TUNEL staining was performed as previously described using the DeadEnd kit (Promega, Madison, WI, USA, Cat # G3250) (Wubben et al., 2017 [DOI](#); Wubben et al., 2020 [DOI](#)). TUNEL-positive cells were counted in a masked fashion and normalized to the total number of nuclei using a custom ImageJ macro (Busov & Besirli, 2014 [DOI](#); Wubben et al., 2017 [DOI](#); Wubben et al., 2020 [DOI](#)). Tissue sections through the plane of the optic nerve were also stained with hematoxylin and eosin and the total number of nuclei in the ONL were determined after normalization to inner retinal area (Wubben et al., 2020 [DOI](#)).

Sub-cellular fractionation

Whole retina was fractionated into cytosolic and post-cytosolic (mitochondrial enriched) fractions as previously described (Weh et al., 2020 [DOI](#)) using the Subcellular Protein Fractionation Kit for Tissues (Thermo Fisher Scientific, Cat # 87790). Both retinas from a single animal were pooled and homogenized with a Dounce homogenizer in cytoplasmic extraction buffer supplemented with protease (HaltTM Protease Inhibitor Cocktail, Thermo Fisher Scientific, Cat # 87786) and phosphatase (HaltTM Phosphatase Inhibitor Cocktail, Thermo Fisher Scientific, Cat # 78420) inhibitors. The retinal lysate was then centrifuged at 10,000 x relative centrifugal force (RCF) for 10 minutes at 4°C. The resulting supernatant was saved as the cytosolic fraction, and the resulting pellet was resuspended in RIPA lysis buffer (Thermo Fisher Scientific, Cat # 89900) that included protease and phosphatase inhibitors (Cell Signaling Technology, Danvers, MA, USA, Cat # 5872) and sonicated at 20% amplitude with 1s on/off pulse for 10s. The lysate was centrifuged for 10 minutes at 10,000 x RCF at 4°C. The resulting supernatant was saved as the mitochondrial enriched fraction. The percentage of GLS in each fraction was determined using Western blotting.

Western blotting

Immunoblots were performed as previously described (Weh et al., 2020 [DOI](#)). Protein estimation was performed using the PierceTM BCA kit (Thermo Fisher Scientific, Cat # 23225). Equivalent micrograms of protein from each sample were diluted using 4X Laemmli buffer (Bio-Rad, Hercules, CA, USA, Cat # 1610747) supplemented with β-mercaptoethanol (Millipore-Sigma, Cat # M6250) before heating at 95°C for 5 minutes and finally loaded onto 4-20% polyacrylamide gel (Bio-Rad, Cat # 4561094). Samples were then transferred to a PVDF membrane using the Trans-Blot[®] TurboTM Transfer System (Bio-Rad, Cat # 1704150). Membranes were blocked using 5% non-fat milk powder diluted in TBST (Tris-buffered Saline, Bio-Rad, Cat # 1706435, supplemented with 0.1% Tween-20, Thermo Fisher Scientific, Cat # 28320) for 4 hours at room temperature. Primary antibodies were diluted in 5% BSA and added to blots before incubating overnight at 4°C. Blots

were then washed and appropriate secondary antibody was added for 1 hour at room temperature. Chemiluminescence was performed using the SuperSignal™ West Dura/Femto Extended Duration Substrate (Thermo Fisher Scientific, Cat # 34075 and 34094) and the immunoblots were imaged with an Azure 600 imaging system (Azure Biosystems; Dublin, CA USA). All antibodies and dilutions used are found in the key resources table.

Quantitative real-time PCR

Total RNA was extracted from whole retina using the RNeasy Mini Kit (Qiagen, Cat # 74104) following the manufacturer's protocol. Isolated RNA was assayed for quantity and quality with a Nanodrop 1000 (Thermo Fisher Scientific) and 1 µg of RNA was used as input for cDNA synthesis using the RNA QuantiTect transcription kit (Qiagen, Cat # 205311). Approximately 100 ng of cDNA was used as a template for each qRT-PCR reaction using the PowerTrack SYBR Green supermix (Thermo Fisher Scientific, Cat # 46109) as previously described (Subramanya et al., 2023 [\[link\]](#)). The Ct values for *Actb* were used to determine relative transcript expression levels using the $2^{-\Delta\Delta Ct}$ method with a cycle threshold cutoff of 35 cycles for the presence of transcript. The geometric mean was used to normalize samples. Custom qRT-PCR primers (Supplementary File 1 [\[link\]](#)) designed to specifically detect spliced transcripts were used to determine transcript levels following the above protocol.

Metabolomics

For unlabeled targeted metabolomics, both retinas from a single animal were rinsed in PBS, combined, and snap-frozen on dry ice. Metabolites were extracted using 80% methanol at -80°C and an OMNI Bead Ruptor (OMNI International, Kennesaw, GA, USA, Cat # 19-050A). Lysates were centrifuged at 14,000 x RCF for 10 minutes at 4°C, and the supernatant stored at -80°C until being processed in the SpeedVac. The pellet from each sample was saved for protein estimation. To determine the protein concentration for each sample, 150 µL of 0.1 M NaOH was added to the pellet for 24 hours at 37°C. The sample was then vortexed and centrifuged at 5000 x RCF at room temperature. The protein estimation was performed as described above. The protein concentration of each sample was used to normalize amount of sample for lyophilization with a SpeedVac concentrator (Thermo Fisher Scientific, Cat # 13875355). Dried metabolite pellets were resuspended for liquid chromatography-coupled mass spectrometry (LC/MS) analysis using an Agilent Technologies Triple Quad 6470 instrument (Santa Clara, CA, USA) as previously described (Wubben et al., 2020 [\[link\]](#)). Previously published parameters were used for data collection (Yuan et al., 2012 [\[link\]](#)). Agilent MassHunter Workstation Quantitative Analysis Software (B0900) was used to process raw data. Additional statistical analyses were performed in Microsoft Excel. Each sample was normalized by the total intensity of all metabolites to reflect sample protein content. To obtain relative metabolites, each metabolite abundance level in each sample was divided by the mean of the abundance levels across all control samples.

To analyze the incorporation of non-radioactive stable isotope carbon-13 (^{13}C) into metabolites in central carbon metabolism and related pathways, mice were intraperitoneally injected with 2g/kg of uniformly labeled $^{13}\text{C}_6$ -glucose (Cambridge Isotope Laboratories, Cat # CLM-1396) or 300 mg/kg of uniformly labeled $^{13}\text{C}_5$ -glutamine (Cambridge Isotope Laboratories, Cat # CLM-1822-H) and retinas were harvested 45 minutes later and snap-frozen as described above. Metabolites were extracted as described above and data collected according to previously published protocols (Yuan et al., 2019 [\[link\]](#)).

Surface Sensing of Translation (SUnSET) Method

In vivo protein synthesis in whole retina from P14 WT and cKO mice was measured using the SUnSET protocol as previously described (Fort et al., 2022 [\[link\]](#)). Briefly, a stock solution of 40 mg/mL puromycin hydrochloride (Millipore-Sigma, Cat # P7255) was prepared in sterile 0.9% sodium chloride and injected intraperitoneally into mice at a final concentration of 200 mg/kg body

weight. Mice were sacrificed after 30 min and fresh retinas were harvested and lysed in RIPA buffer (Thermo Fisher Scientific, Cat # 89900) with protease and phosphatase inhibitors (Cell Signaling Technology, Cat # 5872). Protein quantitation was conducted with the BCA assay as described above and 10 μ g of protein was analyzed by western blot analysis using an anti-puromycin antibody.

BaroFuse

Oxygen consumption rate (OCR) was determined as previously described (Kamat et al., 2023 [DOI](#)). Briefly, a BaroFuse (Entox Sciences, Mercer Island, WA) was used to determine the OCR of freshly isolated retinas. Single, whole retinas were dissected into Hank's Balanced salt solution supplemented with 0.1 g/100 mL BSA (HBSS, Cytiva, Marlborough, MA, USA, Cat # SH30031FS). Perfusion media consisted of commercial Krebs-Ringer Bicarbonate buffer (KRB, Thermo Fisher Scientific, Cat # J67795.K2) supplemented with 0.1 g/100 mL fatty-acid free BSA (Millipore-Sigma, Cat # A9647) and 4.4 mM glucose, for a final concentration of 5.5 mM glucose (Millipore-Sigma, Cat # G8270). The oxygen and CO₂ concentration of the perfusion media is maintained by saturating the solution in a 21% oxygen, 5% CO₂ atmosphere, and the temperature was maintained at 37°C throughout the experiment. At various times throughout the experiment Oligomycin-A (Cayman Chemical, Ann Arbor, MI, USA, Cat # 11342), FCCP (Trifluoromethoxy carbonyl cyanide phenylhydrazide, Cayman Chemical, Cat # 15218), and KCN (Potassium Cyanide, Thermo Fisher, Cat # 012136) were added to the perfusion media through an injection port. A chamber without tissue was used as a negative control.

NAD⁺/NADH and NADP⁺/NADPH Measurements

The NAD⁺/NADH and NADP⁺/NADPH measurements were conducted using the NAD⁺/NADH-Glo Assay (Promega, Cat # G9071) and NADP⁺/NADPH-Glo Assay (Promega, Cat # G9081), respectively, following manufacturer's instructions in whole retina from P14 WT and cKO mice. Briefly, two fresh retina per mouse were harvested for each sample in 150 μ L of PBS/Bicarbonate/0.5% dodecyltrimethylammonium bromide (DTAB) buffer. DTAB is used in the buffer to preserve dinucleotide stability. The samples were sonicated (20% strength and 1 sec/10 iterations) to create a uniform suspension and diluted with 150 μ L of PBS/Bicarbonate/0.5% DTAB buffer (lysate). For measuring NAD⁺/NADP⁺, 150 μ L of lysate was mixed with 75 μ L of 0.4N HCl (Thermo Fisher Scientific, Cat # A144) and heated to 60°C for 15 minutes. The lysate mixture was cooled to room temperature for 10 min and then neutralized by adding 150 μ L 0.5M Trizma base (Millipore-Sigma, Cat # T1503). For measuring NADH/NADPH, the remaining 150 μ L of lysate was heated to 60°C for 15 minutes, cooled to room temperature for 10 min and 150 μ L 0.5M Trizma base was added to the sample. Following lysate preparation, 50 μ L of lysate was incubated at room temperature for 30 min with 50 μ L of either the NAD⁺/NADP⁺-Glo or NADH/NADPH-Glo detection reagent in a 96-well white walled tissue culture plate (Thermo Fisher Scientific, Cat # 3610). Luminescence was recorded using the Omega plate reader (BMG Labtech) and data are reported as a ratio of NAD⁺/NADH or NADP⁺/NADPH of N=3 animals in triplicate technical measurements per animal sample.

Statistical analysis

All data is presented as mean $\pm$ SEM. The significance of the difference between means was determined using a two-tailed student's *t*-test or one-way ANOVA in Excel or Prism 9.0. Results with a *p*-value $\leq$ 0.05 were considered significant.

Resource Availability

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Thomas Wubben (twubben@med.umich.edu).

Supplemental Information

Figure supplements




Figure 1 – figure supplement 1.

GLS is the predominant isoform in the mouse retina and the rod photoreceptor-specific *Gls* conditional knockout mouse does not demonstrate compensatory upregulation of the *Gls2* isoform

(A) qRT-PCR analysis of *Gls* and *Gls2* in P14 mouse retina. Relative expression was analyzed by comparative threshold cycle ($2^{-\Delta\Delta CT}$) method. Expression values were represented as fold change over *Gls* after normalization with β -actin. N=9 animals per group. (B) Representative GLS immunofluorescence in P21 WT animals. Scale bar is 40 μm. (C) Western blot analysis and quantitation of GLS in fractionated wild-type (WT) mouse retinas. TIM23 was used as a mitochondrial fraction marker and HSP90 as a cytosolic fraction marker. WR whole retina extract; C, cytosolic fraction; M, mitochondrial fraction. N=3 animals. (D) qRT-PCR analysis of *Gls2* mRNA expression in the retina at P14 demonstrates no change between the rod photoreceptor-specific *Gls* conditional knockout mouse (cKO) and the wild-type (WT) mouse where both *Gls* alleles are present. Data are normalized to β -actin. N=5-6 animals per group. (E) Representative GLS2 immunofluorescence (red) images in cKO mice at P21 shows no increase in the expression of GLS2 as compared WT mice. Scale bar is 40 μm. Statistical differences in (A), (B) and (D) are based on an unpaired two-tailed Student's t-test where **P<0.01 and ***P<0.001. Data are presented as mean $\pm$ standard error of the mean.

Figure 1 – figure supplement 2.

***Gls* conditional knockout mouse demonstrates similar degeneration under cyclic- and dark-reared conditions**

(A) Representative OCT images from the rod photoreceptor-specific *Gls* conditional knockout (cKO) mouse compared to wild-type (WT) mice at P21 and P28 housed with 12-hour light/12-hour dark cycles (cyclic-reared) or dark-reared. The white vertical bar represents the thickness of the outer nuclear layer (ONL). (B) Quantitation of ONL thickness as determined by OCT in cKO mice compared to WT mice at P21 and P28 under the different housing conditions. N=4-6 animals per group. Statistical differences in (B) are based on a One-way ANOVA with Tukey's post hoc test for multiple comparisons where *** $P < 0.001$. Data are presented as mean $\pm$ standard error of the mean.





Figure 1 – figure supplement 3.

Histology confirms increased Müller glial cell activation in rod photoreceptor-specific *Gls* conditional knockout mice

Representative images from GFAP staining indicate increased Müller glial cell activation in cKO compared to WT mice at P21 and P42. Scale bar is 40 μ m.






Figure 1 – figure supplement 4.

Rod photoreceptor-specific *Gls* conditional knockout mice have shorter outer segments

(A) Representative immunofluorescence images of rhodopsin (RHO) staining (green) in rod photoreceptor-specific *Gls* conditional knockout (cKO) mice compared to wild-type (WT) mice at P21. N=3 animals per group. Scale bar is 40 μ m. (B) Representative transmission electron microscopy (TEM) images of retinal sections from cKO mice compared to WT mice at P21. Images taken at 250X magnification and scale bar is 5 μ m. (C) TEM images of retinal sections from cKO mice compared to WT mice at P21 taken at 500X magnification and scale bar is 3 μ m. (D) TEM images of retinal sections from cKO and WT mice at P21 taken at 8000X and 6000X magnification, respectively, to show ribbon synapses. Intact ribbon synapses in WT rods are shown in yellow boxes. A structurally intact ribbon synapse in cKO rods is denoted by the yellow arrow. Scale bars are 500 nm.




Figure 1 – figure supplement 5.

Rod photoreceptor-specific *GIs* knockout does not alter synaptic connectivity between photoreceptors and second-order neurons

(A) Representative retinal sections stained with wheat germ agglutinin (WGA) to label photoreceptor synaptic and non-synaptic membranes in the rod photoreceptor-specific *GIs* conditional knockout (cKO) mice compared to wild-type (WT) mice at P14. OPL, outer plexiform layer; IPL, inner plexiform layer. N=3 animals per group. Scale bar is 40 μ m. (B) Representative retinal sections stained with an antibody against Bassoon to label ribbon synapses of rods and cones in the OPL in cKO and WT mice at P14. N=3 animals per group. Scale bar is 40 μ m. (C) Representative retinal sections from WT and cKO animals at P14, P21 and P42 are immunostained with antibodies against major cell-type specific markers (red): calretinin (amacrine cells), BRN3A (ganglion cells), CHX10 (bipolar cells). ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion cell layer. N=3 animals per group. Scale bars are 40 μ m. (D) Inner retinal area as a percent of WT retinas at P14, P21 and P42. N=5-10 eyes per group. Statistical differences in (D) are based on an unpaired two-tailed Student's t-test where ** $P < 0.01$. Data are presented as mean $\pm$ standard error of the mean.




Figure 4 – figure supplement 1.

Loss of *Gls* in rod photoreceptors alters the expression of retinal metabolism-related genes

qRT-PCR analysis of genes related to (A) glycolysis and pyruvate metabolism as well as (B) the TCA cycle in rod photoreceptor-specific *Gls* conditional knockout mice (cKO) compared to wild-type (WT) mice retina at P14. Expression values are represented as fold change over WT after normalization with β -actin. N=6 animals per group. Statistical differences in (A) and (B) are based on an unpaired two-tailed Student's t-test where * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$. Data are presented as mean $\pm$ standard error of the mean.




Figure 4 – figure supplement 2.

Loss of *Gls* in rod photoreceptors does not alter the expression of metabolism-related genes in eyecups

qRT-PCR analysis of genes related to (A) glycolysis and pyruvate metabolism as well as (B) the TCA cycle in rod photoreceptor-specific *Gls* conditional knockout mice (cKO) eyecups compared to wild-type (WT) mice eyecups at P14. Expression values are represented as fold change over WT after normalization with β -actin. N=3-4 eyecups per group. Statistical differences in (A) and (B) are based on an unpaired two-tailed Student's t-test where $**P < 0.01$. Data are presented as mean $\pm$ standard error of the mean.




Figure 5 – figure supplement 1.

Mass isotopologue distribution of $^{13}\text{C}_6$ -glucose in glycolysis and TCA cycle intermediates in rod photoreceptor-specific *GIs* conditional knockout (cKO) mice compared to wild-type (WT) mice

Fractional labeling of glycolytic and TCA cycle intermediates with uniformly-labeled, $^{13}\text{C}_6$ -glucose in retina from rod photoreceptor-specific *GIs* cKO mice compared to WT mice at P14. Statistical differences are based on an unpaired two-tailed Student's t-test where *P<0.05 and **P<0.01. Data are presented as mean $\pm$ standard error of the mean.

Figure 5 – figure supplement 2.

Mass isotopologue distribution of $^{13}\text{C}_5$ -Gln in TCA cycle intermediates in rod photoreceptor-specific *Gls* conditional knockout mice (cKO) compared to wild-type (WT) mice

Fractional labeling of TCA cycle intermediates with uniformly-labeled, $^{13}\text{C}_5$ -Gln in retina from rod photoreceptor-specific *Gls* cKO mice compared to WT mice at P14. Statistical differences are based on an unpaired two-tailed Student's t-test where * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$. Data are presented as mean $\pm$ standard error of the mean.




Figure 6 – figure supplement 1.

Amino acid levels are altered in the IND-cKO retina

Relative abundance of glutamine (Gln), glutamate (Glu), aspartate (Asp) and asparagine (Asn) in retinal tissue from WT and IND-cKO 10 days post tamoxifen (TAM), which is prior to PR degeneration. N=5 animals per group. Statistical differences are based on an unpaired Student's t-test where * $P < 0.05$. Data are presented as mean $\pm$ standard error of the mean.




Figure 6 – figure supplement 2.

ISRIB does not impact retinal anatomy in WT mice and prolongs PR survival in the cKO retina

(A) ISRIB (2.5 mg/kg) or vehicle (50% PEG 400, 43.4% saline and 6.6% DMSO) was delivered intraperitoneally in WT mice starting at P5. Total retinal (TRT), outer nuclear layer (ONL) and the inner segment/outer segment thickness were measured at P21 by OCT. N=3 animals per group. (B) ONL thickness at P28 in cKO mice as assessed by OCT following intraperitoneal injection of ISRIB (2.5 mg/kg) or vehicle (50% PEG 400, 43.4% saline and 6.6% DMSO) from P5-P28. N=4-5 animals per group. Statistical differences in (A) and (B) are based on an unpaired two-tail Student's t-test where *P < 0.05 and **P < 0.01. Data are presented as mean $\pm$ standard error of the mean.





Figure 6 – figure supplement 3.

Histology confirms ISRIB and Asn prolong photoreceptor survival in the GLS cKO retina

Representative hematoxylin and eosin stained retinal sections from rod photoreceptor-specific *Glrs* conditional knockout (cKO) mice at P21 following intraperitoneal injection with A) ISRIB (2.5 mg/kg) or vehicle (50% PEG 400, 43.4% saline, 6.6% DMSO) or B) Asn (200 mg/kg) or vehicle (PBS) from P5-P21. ONL, outer nuclear layer; INL, inner nuclear layer. Asn, asparagine. Scale bar is 50 μ m.



Supplementary files

Gene	Forward Primer	Reverse Primer
<i>Gls</i>	CGTTCCATGTTGGTCTTCCT	ATCACCGACTTCACCCCTTTG
<i>Gls2</i>	GACCGTGGTGAACCTGCTAT	ACCTCCAGGTGGTTGAACTG
<i>Actb</i>	AGCCATGTACGTAGCCATCC	CTCTCAGCTGTGGTGGTGAA
<i>Aco2</i>	CAACATGGGTGCAGAAATTG	GTGAGCCAAGTCAGGGGTAA
<i>Idh3a</i>	GAGGTTTTGCTGGTGGTGTT	TCCTCCTGGTCCTTGAATTG
<i>Idh3b</i>	ATCTGAGCGAGGTGCAGAAT	TACGTTGGCAAACAAATCCA
<i>Idh3g</i>	TGTAAGCTCCAACGCTGATG	CACTCCTGGCAGGCTCTTAC
<i>Ogdh</i>	ACATGGCACAGTGCATCATT	ACATCTGCAGAAACCGCTCT
<i>Suc1g1</i>	AGATTCCCTTGGTTGTGTGC	GGGTTGTTTGGTGAACCTGCT
<i>Suc1g2</i>	CTTTGGTGGGATCGTCAACT	AACAGCTTTCTTGGCTGCAT
<i>Sdha</i>	ACACAGACCTGGTGGAGACC	GGATGGGCTTGGAGTAATCA
<i>Sdhb</i>	ACTGGTGGAACGGAGACAAG	TTAAGCCAATGCTCGCTTCT
<i>Sdhc</i>	GGAGGGGTCTCTCTTTTTTG	AAGTGTCGGATCCCATTGAG
<i>Sdhd</i>	GATCCCTGCTGGGTACTTGA	AAGTAGCAAAGCCCAGCAAA
<i>Fh1</i>	AGCAATGCATATTGCTGCTG	CGCATACTGGACTTGCTGAA
<i>Mdh1</i>	GAAGCCCTGAAAGACGACAG	TCGACACGAACTCTCCCTCT
<i>Mdh1b</i>	CATAGCCACCACCCTGAAGT	TGAATCAGATCGCCTGTGAG
<i>Mdh2</i>	GCTTTGTCTTCTCCCTCGTG	CAAAGTCCTCGCCTTTCTTG
<i>Cs</i>	TGCCTAAGGATCCCATGTTC	TTCATCTCCGTCATGCCATA
<i>Acly</i>	GAAGCTGACCTTGCTGAACC	CTGCCTCCAATGATGAGGAT
<i>Pdha1</i>	GGGACGTCTGTTGAGAGAGC	TGTGTCCATGGTAGCGGTAA
<i>Pdha2</i>	TCGAAGCCATAGAAGCCAGT	AGGCATAGGGACATCAGCAC
<i>Hk2</i>	GGGACGACGGTACACTCAAT	GCCAGTGGTAAGGAGCTCTG
<i>Pdk1</i>	GGCGGCTTTGTGATTTGTAT	ACCTGAATCGGGGGATAAAC

<i>Pdk2</i>	AGGAAGTCAATGCCACCAAC	GAGGGCCACCATAATCTTGA
<i>Pdk3</i>	GTGGAGTCCCACCTTCGAAAA	AAAAGTGGCAGCCTCTCAAA
<i>Pdk4</i>	CCTTTGGCTGGTTTTGGTTA	CCTGCTTGGGATACACCAGT
<i>Pcx</i>	ATGTTGTGGACGTGGCAGTA	AATCGAAGGCTGCGTACAGT
<i>Bcat1</i>	TAGAATGTGCCGATCTGCTG	CTTTGGAAGGCTTCTTGACG
<i>Bcat2</i>	GGACCCATGAAGACGGAGTA	CCCGCTTCAATTCCTTCATA
<i>Me1</i>	GGGATTGCTCACTTGGTTGT	GTTTCATGGGCAAACACCTCT
<i>Me2</i>	TTGTGTTCCCTGCATGGTTA	ATACAGACGGGCAGACCAAC
<i>Sod1</i>	CCAGTGCAGGACCTCATTTT	CACCTTTGCCCAAGTCATCT
<i>Sod2</i>	CCGAGGAGAAGTACCACGAG	GCTTGATAGCCTCCAGCAAC
<i>Slc7a11</i>	CCCAGATATGCATCGTCCTT	CGTCTGAACCACTTGGGTTT
<i>Gch1</i>	CACCAAGGGATACCAGGAGA	AGCCAATATGGACCCTTCCT
<i>Chac1</i>	GTACGGCTCCCTAGTGTGGA	GTCTTCAAGGAGGGTCACCA
<i>Ptgs2</i>	AGAAGGAAATGGCTGCAGAA	GCTCGGCTTCCAGTATTGAG
<i>Atg5</i>	AGATGGACAGCTGCACACAC	GCTGGGGGACAATGCTAATA
<i>Sqstm1</i>	GCTCAGGAGGAGACGATGAC	AGAAACCCATGGACAGCATC
<i>Ripk1</i>	CCTGCTGGAGAAGACAGACC	CATCATCTTCCCCTCTTCCA
<i>Ripk3</i>	ACACGGCACTCCTTGGTATC	CCGAACTGTGCTTGGTCATA
<i>Casp3</i>	GGGCCTGTTGAACTGAAAAA	CCGTCCTTTGAATTTCTCCA
<i>Casp8</i>	CCTAGACTGCAACCGAGAGG	GCAGGCTCAAGTCATCTTCC
<i>Casp9</i>	GATGCTGTCCCCTATCAGGA	GGGACTGCAGGTCTTCAGAG

Supplementary File 1.

Primers utilized in qRT-PCR experiments to measure gene expression

Compound Name	Precursor Ion (Da)	Product Ion (Quantifier) (Da)	Product Ion (Qualifier) (Da)	Retention Time (min)	Collision Energy (Volts)
±-Mevalonolactone	189.1	59.1		2.91	6
2-2-Dimethyl Succinic acid	145.1	101.1		14	11
	145.1		127.1	14	8
2-3-Dihydroxybenzoic acid	153	109		14.5	15
	153		53.2	14.5	25
2-3-Dihydroxyisovalerate	133.1	75.1		10.8	10
	133.1		57.2	10.8	15
2-3-Pyridinedicarboxylic acid	166	122		14.9	6
	166		78.1	14.9	13
2-4-Quinolinediol	160	117.9		15.2	19
	160		42.2	15.2	29
2-Deoxyadenosine	250	134.05		6.5	12
	310.1	250		6.5	4
2-Deoxyadenosine 5-diphosphate	410	79.1		15.7	44

Supplementary File 2.

List of metabolites and their parameters detected by LC-MS/MS

	410		158.7	15.7	24
2-Deoxyadenosine 5-monophosphate	330	79		12.7	48
	330		134.05	12.7	28
2-Deoxycytidine	286.1	226		2	4
	286.1		93	2	20
2-Deoxycytidine 5-diphosphate	386	79		14.7	36
	386		97	14.7	20
2-Deoxycytidine 5-monophosphate	306	79		10.4	44
	306		194.9	10.4	14
2-Deoxy-D-glucose 6-phosphate	243	79.1		9.1	45
	243		97	9.1	14
2-Deoxy-D-ribose	193.1	59.2		1.4	16
	193.1		133	1.4	0
2-Deoxyguanosine	266.1	150		4	12
	266.1		108	4	36
2-Deoxyguanosine 5-diphosphate	426	79.1		15.4	48
	426		158.9	15.4	28
2-Deoxyguanosine 5-monophosphate	346	79.1		12.6	48
	346		97	12.6	36
2-Deoxyinosine	251.1	135		3.7	20
	251.1		108	3.7	42
2-Deoxyribose 5-phosphate	213	79.1		9.23	40
	213		96.9	9.23	12
2-Deoxyuridine	227.1	184		2.7	6
	227.1		42.2	2.7	20
2-Deoxyuridine 5-triphosphate	467	158.8		16.8	36
	467		368.9	16.8	20
2-Isopropylmalic acid	175.1	113		15.5	13
	175.1		115	15.5	13
2-Ketobutyrate	101	57.2		12.4	4
2-Methyl-1-butanol	87.1	43.3		9.6	5
2-Phosphoglyceric acid	185	97		14.8	14
	185		79	14.8	37
3-2-Hydroxyethylindole	160.1	130.06		15.4	14
	160.1		142.07	15.4	16
3-Dehydroshikimic acid	171	127		7.4	8
	171		109	7.4	17
3-Hydroxyanthranilic acid	152	108		12.1	12
	152		107	12.1	23

Supplementary File 2. (continued)

3-Hydroxy-DL-kynurenine	223.1	205.9		2.3	4
	223.1		162	2.3	10
3-Hydroxyphenylacetic acid	151	107.1		14.2	4
	151		65.2	14.2	26
3-Indoleacetic acid	174.1	130		16.5	7
	174.1		128	16.5	19
3-Methylglutaric acid	145.1	101.1		14.1	11
	145.1		41.3	14.1	52
4-Aminobenzoic acid	136	92		9.8	8
4-Guanidobutyric acid	144.1	102		1.4	8
	144.1		41.2	1.4	32
4-Hydroxybenzoic acid	137	93		12.1	14
	137		65.2	12.1	34
4-Hydroxy-L-glutamic acid	162	144		5.3	6
	162		72.1	5.3	16
4-Hydroxyphenyl-pyruvic acid	179	107		14.7	4
4-Methyl-2-oxovaleric acid	129.1	85.1		13.7	7
4-Pyridoxic acid	182	138		15.3	12
	182		108.1	15.3	20
4-Quinolinol	144	66.1		9.2	44
	144		116	9.2	28
5-Deoxy-5-(methylthio)adenosine	356.1	134		11.8	12
	356.1		296	11.8	4
5-Hydroxy-3-indoleacetic acid	190.1	144		13	21
	190.1		116	13	46
5-Methoxytryptamine	189.1	144		2.4	28
	189.1		174	2.4	12
6-Hydroxynicotinic acid	138	94.1		9	10
	138		42.3	9	27
Adenine	134	107		2.8	18
	134		92.1	2.8	20
Adenosine	266	134		6.4	10
	326.1	134		6.4	20
Adenosine 3-5-cyclic monophosphate	328	134		13.4	24
	328		79	13.4	48
Adenosine 5-diphosphate	426	159		15.4	28
	426		328	15.4	16
Adenosine 5-monophosphate	346	79		11.6	38
	346		97	11.6	24

Supplementary File 2. (continued)

Adenosine 5-triphosphate	506	408.1		17.2	22
	506		159	17.2	38
Adenylosuccinic acid	462	97		16.4	24
	462		134.05	16.4	48
Adipic acid	145.1	101.1		13.9	11
	145.1		83.1	13.9	12
AICAR	257.1	125		2.3	10
	257.1		42.2	2.3	44
Allantoin	157	42.2		1.3	16
	157		97	1.3	12
alpha-D(+)Mannose 1-phosphate	259	79		7.6	48
	259		97	7.6	14
alpha-D-Glucose-1-phosphate	259	79		7.6	28
	259		240.9	7.6	9
alpha-Ketoglutaric acid	145	101		14.3	5
	145		57.2	14.3	9
Arabinose-5-phosphate	229	79		8.4	36
	229		97	8.4	8
Argininosuccinic acid	289	88		4.6	36
	289		131.1	4.6	27
beta-Nicotinamide adenine dinucleotide	662.1	540		9.5	12
	662.1		327.9	9.5	36
beta-Nicotinamide mononucleotide	333	251.1		3.7	12
	333		135.1	3.7	36
Cellobiose	341.1	161		1.3	5
Chorismic acid	225	189		6.7	6
cis-Aconitic acid	173	129		15.5	4
	173		85.1	15.5	8
Citramalic acid	147	85.1		13.8	12
	147		87	13.8	14
Citric acid	191	111.01		14.9	10
	191		87.01	14.9	16
CoA	766.1	408.1		17.8	40
	766.1		686.2	17.8	40
Creatine	130.1	88.1		1.3	8
	130.1		41.2	1.3	36
Creatine phosphate	210	79		7	16
	210		96.9	7	4
Creatinine	112	41.2		1.35	24
	112		88.1	1.35	16
Cysteine	120	33.2		1.31	10
Cytidine	242.1	109		1.8	8

Supplementary File 2. (continued)

	242.1		42.2	1.8	16
Cytidine 5-diphosphate	402.01	79		14.4	48
	402.01		158.92	14.4	24
Cytidine 5-triphosphate	482	158.8		17	40
	482		79	17	40
Cytidine-5-monophosphate	322	79		9.7	44
	322		97	9.7	22
Cytosine	110	67.1		1.4	8
	110		42.2	1.4	12
D-+-Galactosamine	238.1	159.9		1.1	8
	238.1		100	1.1	6
Deoxyadenosine 5-triphosphate	490	391.9		17.2	24
	490		158.9	17.2	32
Deoxycytidine 5-triphosphate	466	158.9		16.8	28
	466		367.9	16.8	20
Deoxyguanosine 5-triphosphate	506	407.9		17.2	20
	506		158.8	17.2	32
Deoxythymidine 5-triphosphate	481	158.8		17.2	36
	481		383.1	17.2	20
D-erythro-Dihydrosphingosine	300.3	199		8.9	8
	300.3		282	8.9	6
D-Fructose 1,6-biphosphate	338.9	97		14.8	22
	338.9		241.01	14.8	12
D-Fructose 6-phosphate	259	79		7.6	48
	259		97	7.6	14
D-Gluconic acid	195.1	75.2		5.9	18
	195.1		129	5.9	10
D-Glucosamine 6-phosphate	258	79		1.6	48
	258		97	1.6	16
D-Glucose 6-phosphate	259	79		7.6	48
	259		97	7.6	14
Dihydroxyacetone phosphate	169	79		10.5	32
	169		97	10.5	6
DL-2-Aminoadipic acid	160.1	142		4.3	9
	160.1		116	4.3	12
DL-Glyceraldehyde 3-phosphate	169	79		11.7	28
	169		97	11.7	4
DL-Isocitric acid	191	111.02		15.1	11
	191		173	15.1	6

Supplementary File 2. (continued)

D-Maltose	341.1	161		1.3	5
	341.1		179	1.3	4
D-Mannose	179.1	89		1.3	4
	179.1		59.2	1.3	16
D-pantothenic acid	218	88		12.1	10
	218		146.08	12.1	14
D-Ribose 5-phosphate	229	79		8.5	48
	229		97	8.5	10
D-Ribulose 1,5-biphosphate	308.9	79		14.7	46
	308.9		97	14.7	22
D-Sedoheptulose-7-phosphate	289	96.9		8.2	18
	289		79	8.2	48
D-Xylose	149.05	89.02		1.2	4
	149.05		71.01	1.2	4
D-Xylulose-5-phosphate	229	79		8.5	36
	229		138.98	8.5	8
Epicatechin	289.1	245		12.2	11
	289.1		109	12.2	26
Flavin adenine dinucleotide	784.1	346		17	35
	784.1		437	17	30
Folinic acid	472.2	315		14.8	28
	472.2		343	14.8	20
Galactonic acid	195.1	75.1		5.9	18
	195.1		129	5.9	11
gamma-Aminobutyric acid	102	84.1		1.2	8
gamma-Glu-Cys	249.1	128		8.1	8
	249.1		171	8.1	9
GlcNAC	220	119		1.4	2
Glucoheptonic acid	225.1	129		5.8	11
	225.1		75.1	5.8	18
Glutathione Reduced	306.1	143		8.43	17
	306.1		128.1	8.43	17
Glyceric acid	105	75.1		5.9	8
	105		57.2	5.9	14
Glyoxylic acid	73	45.2		5.9	6
Guanine	150	133.02		2	12
	150		66.01	2	36
Guanosine	282.1	150		3.9	17
	282.1		132.9	3.9	32
Guanosine 3,5-cyclic monophosphate	344	150.042		12.2	24
	344		133.02	12.2	44

Supplementary File 2. (continued)

Guanosine 5-diphosphate	442	150		14.8	31
	442		158.9	14.8	20
Guanosine 5-triphosphate	522.3	158.9		17	36
	522.3		424	17	20
Homocitrate	205	125		15.4	10
	205		145	15.4	10
Hypoxanthine	135	92		2	17
	135		65.1	2	28
Indoline-2-carboxylate	162.1	116		16.1	16
	162.1		118	16.1	11
Inosine	267.1	135		3.3	22
	267.1		108	3.3	44
Inosine 5-diphosphate	427	158.9		14.5	28
	427		135	14.5	24
Inosine 5-monophosphate	347	79		11.1	44
	347		97	11.1	22
Inosine 5-triphosphate	507	409		17.2	19
	507		159	17.2	40
Isopentenyl pyrophosphate	245	209		12.8	4
Isopentyl acetate	129.1	85.2		14.6	6
	129.1		41.2	14.6	12
Itaconic acid	129	85.1		13.6	6
	129		41.3	13.6	12
Ketoisovaleric acid	115	71.2		16.56	4
Ketovaleric acid	115	71.2		14.5	4
Lactic acid	89	43.3		7.4	10
	89		45.3	7.4	9
L-Arabinose	149	89.1		1.4	4
	149		59.2	1.4	12
L-Arabitol	151	89		1.3	12
	151		71.02	1.3	16
L-Arginine	173.1	131		1.1	11
	173.1		156	1.1	8
L-asparagine	131	113		1.2	4
	131		70	1.2	12
L-Aspartic Acid	132	88		4.9	10
	132		71	4.9	14
L-Canavanine	175.1	74.04		1.1	10
	175.1		118	1.1	8
L-Carnitine	220.1	146		1.2	4
L-Citrulline	174.1	131		1.3	8
	174.1		42.2	1.3	48

Supplementary File 2. (continued)

L-Cystathionine	221.1	134		1.3	8
	221.1		120	1.3	9
L-Cystine	239	74.1		1.2	20
	239		120	1.2	4
L-Dihydroorotic acid	157	113		7.3	5
	157		42.2	7.3	30
L-Glutamic acid	146	102		4.5	11
	146		128	4.5	8
L-Glutamine	145.1	127		1.3	7
	145.1		109	1.3	10
L-Glutathione (oxidized)	611.1	306.07		13.6	24
	611.1		272.09	13.6	28
L-Histidine	154.1	137		1.1	12
	154.1		93.1	1.1	16
L-Homocysteine	134	116.9		1.4	8
L-Homocystine	267	132		1.5	8
	267		115	1.5	16
L-Homoserine	118	72.1		1.3	10
	118		55	1.3	16
L-Hydroxyglutaric acid	147	128.9		13.8	7
	147		85.1	13.8	13
Lipoamide	204.1	158		16.8	4
	204.1		64.1	16.8	20
L-Isoleucine	130	45		2.1	16
L-Kynurenine	207.1	190		3.9	4
	207.1		144	3.9	20
L-Leucine	130	45		2.1	16
L-Malic acid	133	115		13.8	8
	133		71.1	13.8	14
L-Methionine	148	47.2		1.8	12
	148		100	1.8	6
L-Phenylalanine	164.1	147		4.5	9
	164.1		103	4.5	15
L-Proline	114	68.1		1.3	12
	114		45.2	1.3	12
L-Serine	104	74.1		1.2	8
L-Sorbose	179.1	89		1.3	4
	179.1		71.02	1.3	14
L-Threonine	118	74.1		1.3	9
L-Tryptophan	203.1	116		7.9	14
	203.1		74.1	7.9	14
L-Tyrosine	180	119		2.3	15
	180		93	2.3	12
Maleic acid	115	71.2		12.8	7

Supplementary File 2. (continued)

	115		27.3	12.8	15
Malonic acid	103	59.2		12.8	7
	103		41.3	12.8	32
Melibiose	341.1	179		1.3	4
	341.1		89	1.3	13
Mevalonic acid	147.1	128.9		9.8	8
	147.1		103.1	9.8	11
Mevalonic acid 5-phosphate	227	79		14.8	40
	227		97	14.8	12
m-Hydroxybenzoic acid	137	93.1		13.4	10
	137		65.2	13.4	28
myo-Inositol	179	71.02		1.3	12
	239.1	179		1.3	0
N-Acetyl D-galactosamine	220.1	119		1.2	0
N-Acetyl-alpha-D-glucosamine 1-phosphate	300	97		8.7	16
	300		79	8.7	36
N-acetylaspartate	174	88.1		14	14
	174		58.1	14	22
N-acetylaspartylglutamate	303.1	285		16.1	6
	303.1		128	16.1	14
N-Acetyl-D-glucosamine 6-phosphate	300	97		8.8	18
	300		79	8.8	48
N-Acetylglutamic acid	188.1	128.04		14.1	10
	188.1		102	14.1	16
N-Acetylneuraminic acid	308.1	87		6.2	14
	308.1		169.9	6.2	10
NADH	664.1	397		15.4	32
NADPH	744.1	426.1		17.2	33
N-Carbamoyl-DL-aspartic acid	175	132		13.5	7
	175		88.1	13.5	19
N-Carbamyl-L-glutamic acid	189.1	146		13.6	7
	189.1		102.1	13.6	20
N-Formyl-L-Tyrosine	208.1	164		12.8	7
	208.1		107	12.8	12
Nicotinic acid	122	78.1		11.9	11
	122		51.2	11.9	32
Nicotinic acid mononucleotide	334	289.9		8.6	4
	334		79	8.6	40
o-Hydroxy hippuric acid	194	150		16.5	11
	194		121	16.5	22

Supplementary File 2. (continued)

o-Phospho-L-Serine	184	79		10.2	36
	184		97	10.2	13
O-Phosphorylethanolamine	140	79		1.7	14
	140		63	1.7	38
Orotic acid	155	42		9.1	26
	155		40	9.1	44
O-Succinyl-L-homoserine	218.1	117		6.5	7
	218.1		118	6.5	0
Oxamic acid	88	44.3		6.5	6
	88		42.3	6.5	12
Phenylpyruvic acid	163	91.1		17.2	6
Phosphoenolpyruvic acid	167	79		15.3	12
Prephenic acid	225	101		12.4	9
Pyridoxal 5 phosphate	246	79		14.8	30
Pyridoxal hydrochloride	166.1	138		2.4	12
	166.1		108	2.4	18
Pyridoxamine	167.1	121		1.1	22
	167.1		122	1.1	18
Pyridoxine	168.1	122.1		2.4	17
	168.1		166	2.4	9
Pyruvic acid	87	43.2		9.5	4
Quinic acid	191.1	85.1		6	25
	191.1		93.1	6	24
Riboflavin	375.1	255		12.6	14
	375.1		212	12.6	28
Ribonic acid gamma lactone	147	59.2		1.5	16
	147		99	1.5	8
S-2-Aminoethyl-L-cysteine	163.1	76.1		1.1	10
	163.1		33.2	1.1	28
S-5-Adenosyl-L-homocysteine	383.1	134		5.3	24
	383.1		248	5.3	13
Salicylic acid	137	93.1		16.8	17
	137		65.2	16.8	33
Shikimic acid	173	111		5.8	7
	173		93.1	5.8	16
Succinic acid	117	73.1		13	10
	117		99	13	8
Succinic semialdehyde	101	57.2		7.1	7
Taurine	124	80		1.3	24
Taurocholic acid	514	514		20.7	50
Thiamine	264.1	234		1.1	6

Supplementary File 2. (continued)

	264.1		148	1.1	18
Thymidine	241.1	42.2		5.5	16
	241.1		151	5.5	6
Thymidine 5-diphosphate	401.01	97		15.4	25
	401.01		79	15.4	44
Thymine	125	42.2		2.7	14
trans-4-Hydroxy-L-proline	130.1	128		1.3	12
	130.1		66.03	1.3	16
trans-Aconitic acid	173	129		15.55	4
	173		85.1	15.55	10
trans-trans Muconic acid	141	53.3		14.7	8
Trehalose	341	89.01		1.3	12
	401.1	341		1.3	10
Trehalose 6-phosphate	421.1	241		7.9	27
Uracil	111	42.2		1.7	14
Uric acid	167	124		5.5	13
	167		42.3	5.5	26
Uridine	243.1	200		2.3	6
	243.1		110	2.3	12
Uridine 5-diphosphate	403	158.9		14.8	28
	403		79	14.8	48
Uridine 5'-diphosphogalactose	565	322.9		14.3	24
	565		384.9	14.3	29
Uridine 5-diphosphoglucose	565	322.9		14.3	24
	565		384.9	14.3	28
Uridine 5-monophosphate	323	79.1		10.8	48
	323		97	10.8	24
Uridine 5-triphosphate	483	158.9		17	36
	483		385	17	20
Vanillic acid	167	123		12.6	9
	167		108	12.6	18
Xanthine	151	108		2.1	16
	151		42.2	2.1	24
Xanthosine	283.1	151		8.2	18
	283.1		108	8.2	40
Xylitol	151.1	89		1.3	8
	151.1		71	1.3	8

Supplementary File 2. (continued)

Acknowledgements

Funding for this research was supported by a NEI K08EY031757, a RPB Unrestricted Grant, and the Global Ophthalmology Awards Program (GOAP), a Bayer-sponsored initiative committed to supporting ophthalmic research across the world. This work utilized the Vision Research Core funded by P30 EY007003 from the National Eye Institute. C.A.L. was supported by the following grants: R37CA237421, R01CA248160, R01CA244931, and P01HL149633. C.G.B was supported by R01EY029675. We thank Dr. Stephen Rayport in the Department of Psychiatry, Columbia University, New York, NY, for providing the *Gls^{fllox/flox}* mice. We kindly thank Dr. Stephen Tsang in the Department of Ophthalmology, Columbia University, New York, NY, for the photoreceptor-specific, tamoxifen inducible mouse model (*Pde6g-Cre^{ERT2}*).

Additional information

Author Contributions

Moloy T. Goswami, Conceptualization, Formal Analysis, Investigation, Writing – review and editing; Eric Weh, Conceptualization, Formal Analysis, Investigation, Writing – original draft preparation, Writing – review and editing, Visualization, Supervision; Shubha Subramanya, Formal Analysis, Investigation, Writing – review and editing, Visualization; Katherine M. Weh, Writing – review and editing, Visualization; Hima Bindu Durumutla, Investigation; Heather Hager, Investigation; Nicholas Miller, Investigation; Sraboni Chaudhury, Investigation; Anthony Andren, Resources; Peter Sajjakulnukit, Resources; Li Zhang, Resources; Cagri G. Besirli, Writing – review and editing, Supervision; Costas A. Lyssiotis, Conceptualization, Methodology, Resources, Supervision; Thomas J. Wubben, Conceptualization, Methodology, Formal Analysis, Writing – original draft preparation, Writing – review and editing, Visualization, Supervision, Funding acquisition.

References

- Adler A. J., Klucznik K. M (1982) **Glycerol phosphate dehydrogenase in developing chick retina and brain** *J Neurochem* **38**:909–915 <https://doi.org/10.1111/j.1471-4159.1982.tb05329.x>
- Adler L. t., Chen C., Koutalos Y. (2014) **Mitochondria contribute to NADPH generation in mouse rod photoreceptors** *J Biol Chem* **289**:1519–1528 <https://doi.org/10.1074/jbc.M113.511295>
- Ait-Ali N., Fridlich R., Millet-Puel G., Clerin E., Delalande F., Jaillard C., Blond F., Perrocheau L., Reichman S., Byrne L. C., Olivier-Bandini A., Bellalou J., Moyse E., Bouillaud F., Nicol X., Dalkara D., van Dorsselaer A., Sahel J. A., Leveillard T. (2015) **Rod-derived cone viability factor promotes cone survival by stimulating aerobic glycolysis** *Cell* **161**:817–832 <https://doi.org/10.1016/j.cell.2015.03.023>
- Alkan H. F., Walter K. E., Luengo A., Madreiter-Sokolowski C. T., Stryeck S., Lau A. N., Al-Zoughbi W., Lewis C. A., Thomas C. J., Hoefler G., Graier W. F., Madl T., Vander Heiden M. G., Bogner-Strauss J. G (2018) **Cytosolic Aspartate Availability Determines Cell Survival When Glutamine Is Limiting** *Cell Metab* **28**:706–720 <https://doi.org/10.1016/j.cmet.2018.07.021>
- Altman B. J., Stine Z. E., Dang C. V (2016) **From Krebs to clinic: glutamine metabolism to cancer therapy** *Nat Rev Cancer* **16**:619–634 <https://doi.org/10.1038/nrc.2016.71>
- Bowne S. J., Liu Q., Sullivan L. S., Zhu J., Spellicy C. J., Rickman C. B., Pierce E. A., Daiger S. P (2006) **Why do mutations in the ubiquitously expressed housekeeping gene IMPDH1 cause retina-specific photoreceptor degeneration?** *Invest Ophthalmol Vis Sci* **47**:3754–3765 <https://doi.org/10.1167/iovs.06-0207>
- Busov B., Besirli C. G (2014) **Optimization of ImageJ for automated image analysis to assess for photoreceptor cell death in retinal tissue sections** *Investigative Ophthalmology & Visual Science* **55**:4365–4365
- Campos-Sandoval J. A., Martin-Rufian M., Cardona C., Lobo C., Penalver A., Marquez J (2015) **Glutaminases in brain: Multiple isoforms for many purposes** *Neurochem Int* **88**:1–5 <https://doi.org/10.1016/j.neuint.2015.03.006>
- Caruso S. M., Ryu J., Quinn P. M., Tsang S. H (2020) **Precision metabolome reprogramming for imprecision therapeutics in retinitis pigmentosa** *J Clin Invest* **130**:3971–3973 <https://doi.org/10.1172/JCI139239>
- Chen Y., Zizmare L., Calbiague V., Wang L., Yu S., Herberg F. W., Schmachtenberg O., Paquet-Durand F., Trautwein T (2024) **The retina uncouples glycolysis and oxidative phosphorylation via Cori-, Cahill-, and mini-Krebs-cycle** *eLife* **12**:RP91141 <https://doi.org/10.7554/eLife.91141.1>
- Chinchore Y., Begaj T., Wu D., Drokhlyansky E., Cepko C. L (2017) **Glycolytic reliance promotes anabolism in photoreceptors** *eLife* **6** <https://doi.org/10.7554/eLife.25946>
- Daemen A., Liu B., Song K., Kwong M., Gao M., Hong R., Nannini M., Peterson D., Liederer B. M., de la Cruz C., Sangaraju D., Jauchico A., Zhao X., Sandoval W., Hunsaker T., Firestein R., Latham

- S., Sampath D., Evangelista M., Hatzivassiliou G. (2018) **Pan-Cancer Metabolic Signature Predicts Co-Dependency on Glutaminase and De Novo Glutathione Synthesis Linked to a High-Mesenchymal Cell State** *Cell Metab* **28**:383–399 <https://doi.org/10.1016/j.cmet.2018.06.003>
- Daniele L. L., Han J. Y. S., Samuels I. S., Komirisetty R., Mehta N., McCord J. L., Yu M., Wang Y., Boesze-Battaglia K., Bell B. A., Du J., Peachey N. S., Philp N. J. (2022) **Glucose uptake by GLUT1 in photoreceptors is essential for outer segment renewal and rod photoreceptor survival** *FASEB J* **36**:e22428 <https://doi.org/10.1096/fj.202200369R>
- Du J., Cleghorn W., Contreras L., Linton J. D., Chan G. C., Chertov A. O., Saheki T., Govindaraju V., Sadilek M., Satrustegui J., Hurley J. B (2013) **Cytosolic reducing power preserves glutamate in retina** *Proc Natl Acad Sci U S A* **110**:18501–18506 <https://doi.org/10.1073/pnas.1311193110>
- Du J., Cleghorn W. M., Contreras L., Lindsay K., Rountree A. M., Chertov A. O., Turner S. J., Sahaboglu A., Linton J., Sadilek M., Satrustegui J., Sweet I. R., Paquet-Durand F., Hurley J. B (2013) **Inhibition of mitochondrial pyruvate transport by zaprinast causes massive accumulation of aspartate at the expense of glutamate in the retina** *J Biol Chem* **288**:36129–36140 <https://doi.org/10.1074/jbc.M113.507285>
- Du J., Linton J. D., Hurley J. B (2015) **Probing Metabolism in the Intact Retina Using Stable Isotope Tracers** *Methods Enzymol* **561**:149–170 <https://doi.org/10.1016/bs.mie.2015.04.002>
- Duncan J. L., Pierce E. A., Laster A. M., Daiger S. P., Birch D. G., Ash J. D., Iannaccone A., Flannery J. G., Sahel J. A., Zack D. J., Zarbin M. A., the Foundation Fighting Blindness Scientific Advisory, B. (2018) **Inherited Retinal Degenerations: Current Landscape and Knowledge Gaps** *Transl Vis Sci Technol* **7**:6 <https://doi.org/10.1167/tvst.7.4.6>
- Fort P. E., Losiewicz M. K., Elghazi L., Kong D., Cras-Meneur C., Fingar D. C., Kimball S. R., Rajala R. V. S., Smith A. J., Ali R. R., Abcouwer S. F., Gardner T. W (2022) **mTORC1 regulates high levels of protein synthesis in retinal ganglion cells of adult mice** *J Biol Chem* **298**:101944 <https://doi.org/10.1016/j.jbc.2022.101944>
- Gorbatyuk M. S., Starr C. R., Gorbatyuk O. S (2020) **Endoplasmic reticulum stress: New insights into the pathogenesis and treatment of retinal degenerative diseases** *Prog Retin Eye Res* **79**:100860 <https://doi.org/10.1016/j.preteyeres.2020.100860>
- Grenell A., Wang Y., Yam M., Swarup A., Dilan T. L., Hauer A., Linton J. D., Philp N. J., Gregor E., Zhu S., Shi Q., Murphy J., Guan T., Lohner D., Kolandaivelu S., Ramamurthy V., Goldberg A. F. X., Hurley J. B., Du J (2019) **Loss of MPC1 reprograms retinal metabolism to impair visual function** *Proc Natl Acad Sci U S A* **116**:3530–3535 <https://doi.org/10.1073/pnas.1812941116>
- Halbrook C. J., Thurston G., Boyer S., Anaraki C., Jimenez J. A., McCarthy A., Steele N. G., Kerk S. A., Hong H. S., Lin L., Law F. V., Felton C., Scipioni L., Sajjakulnukit P., Andren A., Beutel A. K., Singh R., Nelson B. S., Van Den Bergh F., Krall A. S., Mullen P. J., Zhang L., Batra S., Morton J. P., Stanger B. Z., Christofk H. R., Digman M. A., Beard D. A., Viale A., Zhang J., Crawford H. C., Pasca di Magliano M., Jorgensen C., Lyssiotis C. A. (2022) **Differential integrated stress response and asparagine production drive symbiosis and therapy resistance of pancreatic adenocarcinoma cells** *Nat Cancer* **3**:1386–1403 <https://doi.org/10.1038/s43018-022-00463-1>
- Halliday M., Radford H., Sekine Y., Moreno J., Verity N., le Quesne J., Ortori C. A., Barrett D. A., Fromont C., Fischer P. M., Harding H. P., Ron D., Mallucci G. R. (2015) **Partial restoration of protein synthesis rates by the small molecule ISRIB prevents neurodegeneration without pancreatic toxicity** *Cell Death Dis* **6**:e1672 <https://doi.org/10.1038/cddis.2015.49>

- Hartong D. T., Dange M., McGee T. L., Berson E. L., Dryja T. P., Colman R. F (2008) **Insights from retinitis pigmentosa into the roles of isocitrate dehydrogenases in the Krebs cycle** *Nat Genet* **40**:1230–1234 <https://doi.org/10.1038/ng.223>
- Joyal J. S., Sun Y., Gantner M. L., Shao Z., Evans L. P., Saba N., Fredrick T., Burnim S., Kim J. S., Patel G., Juan A. M., Hurst C. G., Hatton C. J., Cui Z., Pierce K. A., Bherer P., Aguilar E., Powner M. B., Vevis K., Boisvert M., Fu Z., Levy E., Fruttiger M., Packard A., Rezende F. A., Maranda B., Sapieha P., Chen J., Friedlander M., Clish C. B., Smith L. E (2016) **Retinal lipid and glucose metabolism dictates angiogenesis through the lipid sensor Ffar1** *Nat Med* **22**:439–445 <https://doi.org/10.1038/nm.4059>
- Kamat V., Grumbine M. K., Bao K., Mokate K., Khalil G., Cook D., Clearwater B., Hirst R., Harman J., Boeck M., Fu Z., Smith L. E. H., Goswami M., Wubben T. J., Walker E. M., Zhu J., Soleimanpour S. A., Scarlett J. M., Robbins B. M., Hass D., Hurley J. B., Sweet I. R (2023) **A versatile pumpless multi-channel fluidics system for maintenance and real-time functional assessment of tissue and cells** *Cell Rep Methods* **3**:100642 <https://doi.org/10.1016/j.crmeth.2023.100642>
- Kanow M. A., Giarmarco M. M., Jankowski C. S., Tsantilas K., Engel A. L., Du J., Linton J. D., Farnsworth C. C., Sloat S. R., Rountree A., Sweet I. R., Lindsay K. J., Parker E. D., Brockerhoff S. E., Sadilek M., Chao J. R., Hurley J. B (2017) **Biochemical adaptations of the retina and retinal pigment epithelium support a metabolic ecosystem in the vertebrate eye** *eLife* **6** <https://doi.org/10.7554/eLife.28899>
- Koch S. F., Tsai Y. T., Duong J. K., Wu W. H., Hsu C. W., Wu W. P., Bonet-Ponce L., Lin C. S., Tsang S. H (2015) **Halting progressive neurodegeneration in advanced retinitis pigmentosa** *J Clin Invest* **125**:3704–3713 <https://doi.org/10.1172/JCI82462>
- Kooragayala K., Gotoh N., Cogliati T., Nellissery J., Kaden T. R., French S., Balaban R., Li W., Covian R., Swaroop A (2015) **Quantification of Oxygen Consumption in Retina Ex Vivo Demonstrates Limited Reserve Capacity of Photoreceptor Mitochondria** *Invest Ophthalmol Vis Sci* **56**:8428–8436 <https://doi.org/10.1167/iovs.15-17901>
- Kutsyr O., Noailles A., Martinez-Gil N., Maestre-Carballa L., Martinez-Garcia M., Maneu V., Cuenca N., Lax P (2021) **Short-term high-fat feeding exacerbates degeneration in retinitis pigmentosa by promoting retinal oxidative stress and inflammation** *Proc Natl Acad Sci U S A* **118**:43 <https://doi.org/10.1073/pnas.2100566118>
- Le Y. Z., Zheng L., Zheng W., Ash J. D., Agbaga M. P., Zhu M., Anderson R. E (2006) **Mouse opsin promoter-directed Cre recombinase expression in transgenic mice** *Mol Vis* **12**:389–398 <https://www.ncbi.nlm.nih.gov/pubmed/16636658>
- Li B., Zhang T., Liu W., Wang Y., Xu R., Zeng S., Zhang R., Zhu S., Gillies M. C., Zhu L., Du J (2020) **Metabolic Features of Mouse and Human Retinas: Rods versus Cones, Macula versus Periphery** *Retina versus RPE. iScience* **23**:101672 <https://doi.org/10.1016/j.isci.2020.101672>
- Li J., Csibi A., Yang S., Hoffman G. R., Li C., Zhang E., Yu J. J., Blenis J (2015) **Synthetic lethality of combined glutaminase and Hsp90 inhibition in mTORC1-driven tumor cells** *Proc Natl Acad Sci U S A* **112**:E21–29 <https://doi.org/10.1073/pnas.1417015112>
- Li S., Chen D., Sauve Y., McCandless J., Chen Y. J., Chen C. K (2005) **Rhodopsin-iCre transgenic mouse line for Cre-mediated rod-specific gene targeting** *Genesis* **41**:73–80 <https://doi.org/10.1002/gene.20097>

- Macosko E. Z., Basu A., Satija R., Nemesh J., Shekhar K., Goldman M., Tirosh I., Bialas A. R., Kamitaki N., Martersteck E. M., Trombetta J. J., Weitz D. A., Sanes J. R., Shalek A. K., Regev A., McCarroll S. A (2015) **Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets** *Cell* **161**:1202–1214 <https://doi.org/10.1016/j.cell.2015.05.002>
- Mattapallil M. J., Wawrousek E. F., Chan C. C., Zhao H., Roychoudhury J., Ferguson T. A., Caspi R. R (2012) **The Rd8 mutation of the Crb1 gene is present in vendor lines of C57BL/6N mice and embryonic stem cells, and confounds ocular induced mutant phenotypes** *Invest Ophthalmol Vis Sci* **53**:2921–2927 <https://doi.org/10.1167/iovs.12-9662>
- McLaughlin B. J., Wood J. G., Gurd J. W (1980) **The localization of lectin binding sites during photoreceptor synaptogenesis in the chick retina** *Brain Res* **191**:345–357 [https://doi.org/10.1016/0006-8993\(80\)91286-x](https://doi.org/10.1016/0006-8993(80)91286-x)
- Mingote S., Masson J., Gellman C., Thomsen G. M., Lin C. S., Merker R. J., Gaisler-Salomon I., Wang Y., Ernst R., Hen R., Rayport S (2015) **Genetic Pharmacotherapy as an Early CNS Drug Development Strategy: Testing Glutaminase Inhibition for Schizophrenia Treatment in Adult Mice** *Front Syst Neurosci* **9**:165 <https://doi.org/10.3389/fnsys.2015.00165>
- Ng S. K., Wood J. P., Chidlow G., Han G., Kittipassorn T., Peet D. J., Casson R. J (2015) **Cancer-like metabolism of the mammalian retina** *Clin Exp Ophthalmol* **43**:367–376 <https://doi.org/10.1111/ceo.12462>
- Okazaki A., Gameiro P. A., Christodoulou D., Laviollette L., Schneider M., Chaves F., Stemmer-Rachamimov A., Yazinski S. A., Lee R., Stephanopoulos G., Zou L., Iliopoulos O (2017) **Glutaminase and poly(ADP-ribose) polymerase inhibitors suppress pyrimidine synthesis and VHL-deficient renal cancers** *J Clin Invest* **127**:1631–1645 <https://doi.org/10.1172/JCI87800>
- Pakos-Zebrucka K., Koryga I., Mnich K., Ljubic M., Samali A., Gorman A. M (2016) **The integrated stress response** *EMBO Rep* **17**:1374–1395 <https://doi.org/10.15252/embr.201642195>
- Pan W. W., Wubben T. J., Besirli C. G (2021) **Photoreceptor metabolic reprogramming: current understanding and therapeutic implications** *Commun Biol* **4**:245 <https://doi.org/10.1038/s42003-021-01765-3>
- Pavlova N. N., Hui S., Ghergurovich J. M., Fan J., Intlekofer A. M., White R. M., Rabinowitz J. D., Thompson C. B., Zhang J (2018) **As Extracellular Glutamine Levels Decline, Asparagine Becomes an Essential Amino Acid** *Cell Metab* **27**:428–438 <https://doi.org/10.1016/j.cmet.2017.12.006>
- Petit L., Ma S., Cipi J., Cheng S. Y., Zieger M., Hay N., Punzo C (2018) **Aerobic Glycolysis Is Essential for Normal Rod Function and Controls Secondary Cone Death in Retinitis Pigmentosa** *Cell Rep* **23**:2629–2642 <https://doi.org/10.1016/j.celrep.2018.04.111>
- Rajala A., Bhat M. A., Teel K., Gopinadhan Nair G. K., Purcell L., Rajala R. V. S (2023) **The function of lactate dehydrogenase A in retinal neurons: implications to retinal degenerative diseases** *PNAS Nexus* **2**:pgad038 <https://doi.org/10.1093/pnasnexus/pgad038>
- Rajala R. V. S (2020) **Aerobic Glycolysis in the Retina: Functional Roles of Pyruvate Kinase Isoforms** *Front Cell Dev Biol* **8**:266 <https://doi.org/10.3389/fcell.2020.00266>
- Rosarda J. D., Giles S., Harkins-Perry S., Mills E. A., Friedlander M., Wiseman R. L., Eade K. T (2023) **Imbalanced unfolded protein response signaling contributes to 1-**

deoxysphingolipid retinal toxicity *Nat Commun* **14**:4119 <https://doi.org/10.1038/s41467-023-39775-w>

Ross C. D., Bowers M., Godfrey D. A (1987) **Distribution of glutaminase activity in retinal layers of rat and guinea pig** *Brain Res* **401**:168–172 [https://doi.org/10.1016/0006-8993\(87\)91178-4](https://doi.org/10.1016/0006-8993(87)91178-4)

Rowe A. A., Patel P. D., Gordillo R., Wert K. J (2021) **Replenishment of TCA cycle intermediates provides photoreceptor resilience against neurodegeneration during progression of retinitis pigmentosa** *JCI Insight* **6**:17 <https://doi.org/10.1172/jci.insight.150898>

Sisk D. R., Kuwabara T (1985) **Histologic changes in the inner retina of albino rats following intravitreal injection of monosodium L-glutamate** *Graefes Arch Clin Exp Ophthalmol* **223**:250–258 <https://doi.org/10.1007/BF02153655>

Son J., Lyssiotis C. A., Ying H., Wang X., Hua S., Ligorio M., Perera R. M., Ferrone C. R., Mullarky E., Shyh-Chang N., Kang Y., Fleming J. B., Bardeesy N., Asara J. M., Haigis M. C., DePinho R. A., Cantley L. C., Kimmelman A. C (2013) **Glutamine supports pancreatic cancer growth through a KRAS-regulated metabolic pathway** *Nature* **496**:101–105 <https://doi.org/10.1038/nature12040>

Subramanya S., Goswami M. T., Miller N., Weh E., Chaudhury S., Zhang L., Andren A., Hager H., Weh K. M., Lyssiotis C. A., Besirli C. G., Wubben T. J (2023) **Rod photoreceptor-specific deletion of cytosolic aspartate aminotransferase, GOT1, causes retinal degeneration [Original Research]** *Frontiers in Ophthalmology* **3** <https://doi.org/10.3389/fopht.2023.1306019>

Sullivan L. B., Luengo A., Danai L. V., Bush L. N., Diehl F. F., Hosios A. M., Lau A. N., Elmiligy S., Malstrom S., Lewis C. A., Vander Heiden M. G (2018) **Aspartate is an endogenous metabolic limitation for tumour growth** *Nat Cell Biol* **20**:782–788 <https://doi.org/10.1038/s41556-018-0125-0>

Swarup A., Samuels I. S., Bell B. A., Han J. Y. S., Du J., Massenzio E., Abel E. D., Boesze-Battaglia K., Peachey N. S., Philp N. J (2019) **Modulating GLUT1 expression in retinal pigment epithelium decreases glucose levels in the retina: impact on photoreceptors and Muller glial cells** *Am J Physiol Cell Physiol* **316**:C121–C133 <https://doi.org/10.1152/ajpcell.00410.2018>

Tsantilas K. A., Cleghorn W. M., Bisbach C. M., Whitson J. A., Hass D. T., Robbins B. M., Sadilek M., Linton J. D., Rountree A. M., Valencia A. P., Sweetwyne M. T., Campbell M. D., Zhang H., Jankowski C. S. R., Sweet I. R., Marcinek D. J., Rabinovitch P. S., Hurley J. B (2021) **An Analysis of Metabolic Changes in the Retina and Retinal Pigment Epithelium of Aging Mice** *Invest Ophthalmol Vis Sci* **62**:20 <https://doi.org/10.1167/iovs.62.14.20>

Ueta T., Inoue T., Furukawa T., Tamaki Y., Nakagawa Y., Imai H., Yanagi Y (2012) **Glutathione peroxidase 4 is required for maturation of photoreceptor cells** *J Biol Chem* **287**:7675–7682 <https://doi.org/10.1074/jbc.M111.335174>

Voigt A. P., Whitmore S. S., Lessing N. D., DeLuca A. P., Tucker B. A., Stone E. M., Mullins R. F., Scheetz T. E (2020) **Spectacle: An interactive resource for ocular single-cell RNA sequencing data analysis** *Exp Eye Res* **200**:108204 <https://doi.org/10.1016/j.exer.2020.108204>

Weh E., Lutrzykowska Z., Smith A., Hager H., Pawar M., Wubben T. J., Besirli C. G (2020) **Hexokinase 2 is dispensable for photoreceptor development but is required for survival during aging and outer retinal stress** *Cell Death Dis* **11**:422 <https://doi.org/10.1038/s41419-020-2638-2>

- Weh E., Scott K., Wubben T. J., Besirli C. G (2022) **Dark-reared rd10 mice experience rapid photoreceptor degeneration with short exposure to room-light during in vivo retinal imaging** *Exp Eye Res* **215**:108913 <https://doi.org/10.1016/j.exer.2021.108913>
- Wert K. J., Velez G., Kanchustambham V. L., Shankar V., Evans L. P., Sengillo J. D., Zare R. N., Bassuk A. G., Tsang S. H., Mahajan V. B (2020) **Metabolite therapy guided by liquid biopsy proteomics delays retinal neurodegeneration** *EBioMedicine* **52**:102636 <https://doi.org/10.1016/j.ebiom.2020.102636>
- Winkler B. S (1981) **Glycolytic and oxidative metabolism in relation to retinal function** *J Gen Physiol* **77**:667–692 <https://doi.org/10.1085/jgp.77.6.667>
- Winkler B. S., DeSantis N., Solomon F (1986) **Multiple NADPH-producing pathways control glutathione (GSH) content in retina** *Exp Eye Res* **43**:829–847 [https://doi.org/10.1016/s0014-4835\(86\)80013-6](https://doi.org/10.1016/s0014-4835(86)80013-6)
- Wubben T. J., Besirli C. G., Zacks D. N (2016) **Pharmacotherapies for Retinal Detachment** *Ophthalmology* **123**:1553–1562 <https://doi.org/10.1016/j.ophtha.2016.02.040>
- Wubben T. J., Pawar M., Smith A., Toolan K., Hager H., Besirli C. G (2017) **Photoreceptor metabolic reprogramming provides survival advantage in acute stress while causing chronic degeneration** *Sci Rep* **7**:17863 <https://doi.org/10.1038/s41598-017-18098-z>
- Wubben T. J., Pawar M., Weh E., Smith A., Sajjakulnukit P., Zhang L., Dai L., Hager H., Pai M. P., Lyssiotis C. A., Besirli C. G (2020) **Small molecule activation of metabolic enzyme pyruvate kinase muscle isozyme 2, PKM2, circumvents photoreceptor apoptosis** *Sci Rep* **10**:2990 <https://doi.org/10.1038/s41598-020-59999-w>
- Xu R., Ritz B. K., Wang Y., Huang J., Zhao C., Gong K., Liu X., Du J (2020) **The retina and retinal pigment epithelium differ in nitrogen metabolism and are metabolically connected** *J Biol Chem* **295**:2324–2335 <https://doi.org/10.1074/jbc.RA119.011727>
- Xu Y., Shi T., Cui X., Yan L., Wang Q., Xu X., Zhao Q., Xu X., Tang Q. Q., Tang H., Pan D (2021) **Asparagine reinforces mTORC1 signaling to boost thermogenesis and glycolysis in adipose tissues** *EMBO J* **40**:e108069 <https://doi.org/10.15252/embj.2021108069>
- Yang L., Venneti S., Nagrath D (2017) **Glutaminolysis: A Hallmark of Cancer Metabolism** *Annu Rev Biomed Eng* **19**:163–194 <https://doi.org/10.1146/annurev-bioeng-071516-044546>
- Yoo H. C., Yu Y. C., Sung Y., Han J. M (2020) **Glutamine reliance in cell metabolism** *Exp Mol Med* **52**:1496–1516 <https://doi.org/10.1038/s12276-020-00504-8>
- Young R. W (1967) **The renewal of photoreceptor cell outer segments** *J Cell Biol* **33**:61–72 <https://doi.org/10.1083/jcb.33.1.61>
- Yuan M., Breitkopf S. B., Yang X., Asara J. M (2012) **A positive/negative ion-switching, targeted mass spectrometry-based metabolomics platform for bodily fluids, cells, and fresh and fixed tissue** *Nat Protoc* **7**:872–881 <https://doi.org/10.1038/nprot.2012.024>
- Yuan M., Kremer D. M., Huang H., Breitkopf S. B., Ben-Sahra I., Manning B. D., Lyssiotis C. A., Asara J. M (2019) **Ex vivo and in vivo stable isotope labelling of central carbon metabolism**

and related pathways with analysis by LC-MS/MS *Nat Protoc* **14**:313–330 <https://doi.org/10.1038/s41596-018-0102-x>

Zhang J., Fan J., Venneti S., Cross J. R., Takagi T., Bhinder B., Djaballah H., Kanai M., Cheng E. H., Judkins A. R., Pawel B., Baggs J., Cherry S., Rabinowitz J. D., Thompson C. B (2014) **Asparagine plays a critical role in regulating cellular adaptation to glutamine depletion** *Mol Cell* **56**:205–218 <https://doi.org/10.1016/j.molcel.2014.08.018>

Zhou M., Liu Y., Ma C (2021) **Distinct Nuclear Architecture of Photoreceptors and Light-Induced Behaviors in Different Strains of Mice** *Transl Vis Sci Technol* **10**:37 <https://doi.org/10.1167/tvst.10.2.37>

Author information

Moloy T Goswami*

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

* equal contribution

Eric Weh*

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

* equal contribution

Shubha Subramanya

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

Katherine M Weh

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

Hima Bindu Durumutla

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States, Molecular and Developmental Biology Graduate Program, Cincinnati Children's Hospital Medical Center, Cincinnati, United States

Heather Hager

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

Nicholas Miller

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

Sraboni Chaudhury

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

Anthony Andren

Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, United States

Peter Sajjakulnukit

Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, United States

Li Zhang

Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, United States

Cagri G Besirli

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

ORCID iD: [0000-0001-7154-2024](https://orcid.org/0000-0001-7154-2024)

Costas A Lyssiotis

Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, United States, Department of Internal Medicine, Division of Gastroenterology and Hepatology, University of Michigan, Ann Arbor, United States, Rogel Cancer Center, University of Michigan, Ann Arbor, United States

ORCID iD: [0000-0001-9309-6141](https://orcid.org/0000-0001-9309-6141)

Thomas J Wubben

Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, United States

For correspondence: twubben@med.umich.edu

Editors

Reviewing Editor

Zhongjie Fu

Boston Children's Hospital, Boston, United States of America

Senior Editor

Lois Smith

Boston Children's Hospital, Boston, United States of America

Reviewer #1 (Public review):

Summary:

The authors show for the first time that deleting GLS from rod photoreceptors results in the rapid death of these cells. The death of photoreceptor cells could result from loss of synaptic activity because of a decrease in glutamate, as has been shown in neurons, changes in redox balance, or nutrient deprivation.

Strengths:

The strength of this manuscript is that the author shows a similar phenotype in the mice when GLS was knocked out early in rod development or the adult rod. They showed that rapid cell death is through apoptosis, and there is an increase in the expression of genes responsive to oxidative stress.

Comments on revisions:

The authors addressed all of my concerns in their responses to reviewers.

<https://doi.org/10.7554/eLife.100747.2.sa3>

Reviewer #2 (Public review):**Summary:**

Photoreceptor neurons are crucial for vision, and discovering pathways necessary for photoreceptor health and survival can open new avenues for therapeutics. Studies have shown that metabolic dysfunction can cause photoreceptor degeneration and vision loss, but the metabolic pathways maintaining photoreceptor health are not well understood. This is a fundamental study that shows that glutamine catabolism is critical for photoreceptor cell health using in vivo model systems.

Strengths:

The data are compelling, and the consideration of potential confounding factors (such as glutaminase 2 expression) and additional experiments to examine the synaptic connectivity and inner retina added strength to this work. The authors were also careful not to overstate their claims, but to provide solid conclusions that fit the results and data provided in their study. The findings linking asparagine supplementation and the inhibition of the integrated stress response to glutamine catabolism within the rod photoreceptor cell are intriguing and innovative. Overall, the authors provide convincing data to highlight that photoreceptors utilize various fuel sources to meet their metabolic needs, and that glutamine is critical to these cells for their biomass, redox balance, function and survival.

<https://doi.org/10.7554/eLife.100747.2.sa2>

Reviewer #3 (Public review):**Summary:**

The authors explored the role of GLS, a glutaminase, which is an enzyme catalyzes the conversion of glutamine to glutamate, in rod photoreceptor function and survival. The loss of GLS was found to cause rapid autonomous death of rod photoreceptors.

Strengths:

Interesting and novel phenotype. Two types of cre-lines were rigorously used to knockout GLS gene in rods. Both of the conditional knockouts led to a similar phenotype, i.e. rod death. Histology and ERG were carefully done to characterize the loss of rods over specific ages. Necessary metabolomic study was performed and appreciated. Some rescue experiments were performed, and revealed possible mechanism.

Weaknesses:

No major weaknesses. Mechanism of GLS-loss induced rod death could be followed up in the future, and same for GLS's role in cones. Authors have addressed all minor points raised by this reviewer.

<https://doi.org/10.7554/eLife.100747.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public Review):

Summary:

The authors show for the first time that deleting GLS from rod photoreceptors results in the rapid death of these cells. The death of photoreceptor cells could result from loss of synaptic activity because of a decrease in glutamate, as has been shown in neurons, changes in redox balance, or nutrient deprivation.

Strengths:

The strength of this manuscript is that the author shows a similar phenotype in the mice when Glis was knocked out early in rod development or the adult rod. They showed that rapid cell death is through apoptosis, and there is an increase in the expression of genes responsive to oxidative stress.

We thank the reviewer for their time reviewing the manuscript and their comments regarding the potential mechanism(s) by which rod photoreceptors rapidly degenerate upon knockout of GLS.

Weaknesses:

In this manuscript, the authors show a "metabolic dependency of photoreceptors on glutamine catabolism in vivo". However, there is a potential bias in their thinking that glutamine metabolism in rods is similar to cancer cells where it feeds into the TCA cycle. They should consider that as in neurons, GLS1 activity provides glutamate for synaptic transmission. The modest rescue shown by providing α -ketoglutarate in the drinking water suggests that glutamine isn't a key metabolic substrate for rods when glucose is plentiful. The ERG studies performed on the iCre-Glsflox/flox mice showed a large decrease in the scotopic b wave at saturating flashes which could indicate a decrease in glutamate at the rod synapse as stated by the authors. While EM micrographs of wt and iCre-Glsflox/flox mice were shown for the outer retina at p14, the synapse of the rods needs to be examined by EM.

We agree with the reviewer that in the presence of sufficient glucose, it appears a lack of GLS-driven glutamine (Gln) catabolism does not drastically alter the levels of TCA cycle metabolites or mitochondrial function as we demonstrated in Figure 4, and supplementation with alpha-ketoglutarate improved outer nuclear layer thickness by only a small amount as observed in Figure 5e. Hence, as we stated in the Results and Discussion, at least in the mouse where Glis is selectively deleted from rod photoreceptors by crossing *Gls^{fl/fl}* mice with *Rho-Cre* mice (*Gls^{fl/fl}; Rho-Cre⁺*, cKO), Gln's role in supporting the TCA cycle is not the major mechanism by which rod photoreceptors utilize Gln to suppress apoptosis.

With regards to GLS-driven Gln catabolism providing glutamate (Glu) for synaptic transmission, we again agree with the reviewer that Glu is an important excitatory neurotransmitter, but it is also a key metabolite necessary for the synthesis of glutathione, amino acids, and proteins. As noted and discussed at length in the manuscript, a lack of GLS-driven Gln catabolism in rod photoreceptors leads to reduced levels of oxidized glutathione (Figure 4D) possibly signaling an overall reduction in the biosynthesis of glutathione as Glu is directly and indirectly responsible for its synthesis. Furthermore, Gln and GLS-derived Glu play a central role in the biosynthesis of several nonessential amino acids and proteins. To this end, we see a reduction in the level of Glu, which is the product of the GLS reaction and further confirms the loss of GLS function. We also noted a significant decrease in aspartate (Asp), which can be constructed from the carbons and nitrogens of Gln as discussed at length in the manuscript (Figure 6A). Finally, we noted a significant decrease in global protein synthesis in the cKO retina as compared to the wild-type animal as well (Figure 6E). Therefore, the data suggest that GLS-driven Gln catabolism is critical for amino acid metabolism and protein synthesis and to some degree redox balance; although, the small but statistically significant changes in oxidized glutathione, NADP/NADPH, and redox gene expression may not fully account for the rapid and complete photoreceptor degeneration observed. Future studies are necessary to shed light on the role of redox imbalance in this novel transgenic mouse model.

Glu also plays a role in synaptic transmission, and we considered this scenario as described in Figure 1 – figure supplement 5. Here, the synaptic connectivity between photoreceptors and the inner retina did not demonstrate significant differences in the labeling of photoreceptor synaptic membranes in the outer plexiform layer nor alterations in the labeling of a key protein (Bassoon) in ribbon synapses. These data suggest that the synaptic connectivity between photoreceptors and second-order neurons was unaltered at P14 in the cKO retina, which is the time just prior to rapid photoreceptor degeneration when Glu was shown to be decreased (Figure 6A).

With regards to the ERG changes noted in Figure 2, we agree with the reviewer that a large decrease was noted in the scotopic b-wave at P21 and P42 in the cKO. We also agree, that to obtain greater insight into these ERG changes, the ribbon synapse in EM images can be examined. The EM images shown in Figure 1 – figure supplement 4 are from P21, which coincide with the age at which the ERG changes were first noted and when significant photoreceptor degeneration has already occurred. These images were utilized to assess the ribbon synapse for the revised version of the manuscript. As now shown in Figure 1 – figure supplement 4D, ribbon synapses are intact in WT animals as denoted by the yellow boxes. Similarly, the ribbons (yellow arrows) appear structurally intact in the photoreceptors that remain in the P21 cKO retina. These results are in accordance with the lack of significant differences in the labeling of photoreceptor synaptic membranes in the outer plexiform layer as well as the lack of alterations in the labeling of a key protein (Bassoon) in ribbon synapses (Figure 1-figure supplement 5A and B). While we cannot fully rule out that the decrease in glutamate is altering synaptic transmission, our structural data suggests the synapses remain intact. These data have been added to the revised manuscript.

However, an even larger reduction in the scotopic a-wave was noted at these ages as well. In animal models that disrupt photoreceptor synaptic function (Dick et al. *Neuron*. 2003; Johnson et al. *J Neuroscience*. 2007; Haeseleer et al. *Nature Neuroscience*. 2004; Chang et al. *Vis Neurosci*. 2006), a more negative ERG pattern is typically observed with the b-wave altered to a much larger degree than the a-wave. Additionally, in these models that disrupt photoreceptor synaptic transmission, the overall structure of the retina with respect to thickness is maintained (Dick et al. *Neuron*. 2003) or noted to have modest changes in the outer plexiform layer within the first two months of age with the outer nuclear layer not significantly altered until 8-10 months of age (Haeseleer et al. *Nature Neuroscience*. 2004). In

contrast, a rapid decline in the outer nuclear layer thickness was observed in the cKO retina after P14 likely contributing to the ERG changes noted in Figure 2. Also, Gln is catabolized to Glu primarily by GLS as suggested by the approximately 50% reduction in Glu levels in the cKO retina (Figure 6A), but other enzymes are also capable of catabolizing Gln to Glu, so Glu levels in the rod photoreceptors are unlikely to be zero. Coupling this with the fact that rods are equipped with a self-sufficient Glu recollecting system at their synaptic terminals (Hasegawa et al. Neuron. 2006; Winkler et al. Vis Neurosci. 1999) and that GLS activity is at least two-fold higher in the photoreceptor inner segments, which support energy production and metabolism, than any other layer in the retina (Ross et al. Brain Res. 1987) suggests that altered synaptic transmission secondary to reduced levels of Glu likely does not account in full for the rapid and robust photoreceptor degeneration observed in the cKO retina.

The authors note that the outer segments are shorter but they do not address whether there is a decrease in the number of cones.

We have adjusted Figure 2E by removing the GLS staining to better highlight the secondary degeneration of cone outer segments, the main point of the Figure, as we had already shown that GLS was cleanly knocked out of rod photoreceptors in Figure 1. Furthermore, qualitatively the number of cones appears the same at P14, P21, and P42 between the WT and cKO, which is consistent with other retinal degeneration models, like *rd1* and *rd10*, where cones do not begin to die until all the rods have degenerated (Xue et al. eLife. 2021).

Rod-specific Glis ko mice with an inducible promoter were generated by crossing the Pde6g-CreERT2 and homozygous for either the WT or floxed Glis allele (IND-cKO). In Figure 3 the authors document that by western blots and antibody labeling the GLS1 expression is lost in the IND-cKO 10 days post tamoxifen. OCT images show a decrease in the thickness of the outer nuclear layer between 17 and 38 days post-TAM. Ergs should be performed on the animals at 10 and 30 days post TAM, before and after major structural changes in rod photoreceptor cells, to determine if changes in light-stimulated responses are observed. These studies could help to parse out the cause of photoreceptor cell death.

We agree with the reviewer that the IND-cKO is a useful tool to help parse out the cause of photoreceptor cell death in this model as well as shed light on the role of GLS-driven Gln catabolism in photoreceptor synaptic transmission as discussed at length above. Hence, ERG analyses were performed 10 days post TAM, before major structural changes in the ONL are observed. Interestingly, ERG demonstrated statistically significant reductions in the IND-cKO scotopic a- and b-waves as compared to the WT 10 days post TAM. Similarly, photopic ERG demonstrated statistically significant decreases in the b-wave of the IND-cKO retina. These data suggest that GLS-driven Gln catabolism plays a significant role not only in rod photoreceptor survival but their function as well. This data has been added to Figure 3H-I and discussed in the corresponding manuscript text.

To this end, as discussed below and added to Figure 6 – figure supplement 1, amino acid levels, including glutamate (Glu), are already reduced 10 days post TAM. Reductions in the level of Glu may impact synaptic transmission and as a result, the scotopic b-wave. However, as noted above, altered synaptic transmission secondary to reduced levels of Glu likely does not account in full for the rapid and robust photoreceptor degeneration observed in the cKO retina as the b-wave to a-wave ratio is not significantly altered in the IND-cKO retina as compared to the WT retina, suggesting GLS-driven Gln catabolism is impairing both to a similar degree.

Additionally, *Pde6g* is expressed by rods to a significant degree but also by cones (GSE63473, scRNAseq data). Therefore, the IND-cKO mouse likely knocks out GLS from both rods and cones, which is in accordance with the immunofluorescence image in Figure 3B where GLS is

not observed in rod or cone inner segments unlike in Figure 1B where GLS remains in cones. Hence, the reduction in photopic b-wave may be demonstrating that GLS-driven Gln catabolism in cones impairs synaptic transmission. As noted in our reply to reviewer #3's comments, we have generated mice lacking GLS in cone photoreceptors specifically and are currently elucidating the role of GLS in cone photoreceptor metabolism, function, and survival. These results will be published in a separate manuscript.

The studies in Figure 4 were all performed on iCre-Glsflox/flox and control mice at p14, why weren't the IND-cKO mice used for these studies since the findings would not be confounded by development?

To gain further insight into the role of GLS-driven Gln catabolism in the maintenance of rod photoreceptors as compared to their development/maturation, we conducted a targeted metabolomic analysis on IND-cKO and WT retinas 10 days post TAM. For the purpose of this manuscript, we have included data regarding changes in amino acid levels in Figure 6 – figure supplement 1. Specifically, levels of glutamate, aspartate and asparagine are all significantly decreased in the IND-cKO retina prior to PR degeneration, which demonstrates that similar to the GLS cKO mouse (i.e. iCre-Gls flox/flox), GLS-driven Gln catabolism is critical for amino acid biosynthesis in mature rod PRs as well.

In all rescue studies, the endpoint was an ONL thickness, which only addressed rod cell death. The authors should also determine whether there are small improvements in the ERG, which would distinguish the role of GLS in preventing oxidative stress.

Optical coherence tomography (OCT) provides a sensitive in vivo method to detect small changes in retinal thickness without potential artifacts incurred through histological processing. Considering the Gls cKO retina demonstrates significant and rapid photoreceptor degeneration, we wanted to assess pathways that may be critical to photoreceptor survival downstream of GLS-driven Gln catabolism using rescue experiments with pharmacologic treatment or metabolite supplementation. That said, disruption of GLS-driven Gln catabolism may also significantly alter rod photoreceptor function beyond that which is secondary to photoreceptor cell death as we have demonstrated in the IND-cKO animal for the revised version of this manuscript and discussed in a response above. Therefore, the IND-cKO model provides a unique tool to assess the impact of rescue studies on photoreceptor function as the functional changes occur prior to significant degeneration. Also, unlike the GLS cKO mouse (i.e. iCre-Gls flox/flox) where photoreceptor degeneration starts very early, impairing our ability to capture reliable and robust ERG measurements, the IND-cKO mice are older at the time of functional changes allowing for robust ERG measurements. While the rate of photoreceptor degeneration in both mouse models is similar and the levels of key amino acids are altered similarly in both models, the mechanisms of cell death in developing/maturing photoreceptors may be different than that in mature photoreceptors. Hence, before we can assess if similar rescue experiments impact photoreceptor function via ERG in the IND-cKO mouse, we need to thoroughly examine how these photoreceptors are dying. These experiments and results will be published in a separate manuscript in the future.

Reviewer #2 (Public Review):

Summary:

Photoreceptor neurons are crucial for vision, and discovering pathways necessary for photoreceptor health and survival can open new avenues for therapeutics. Studies have shown that metabolic dysfunction can cause photoreceptor degeneration and vision loss, but the metabolic pathways maintaining photoreceptor health are not well understood.

This is a fundamental study that shows that glutamine catabolism is critical for photoreceptor cell health using in vivo model systems.

Strengths:

The data are compelling, and the consideration of potential confounding factors (such as glutaminase 2 expression) and additional experiments to examine the synaptic connectivity and inner retina added strength to this work. The authors were also careful not to overstate their claims, but to provide solid conclusions that fit the results and data provided in their study. The findings linking asparagine supplementation and the inhibition of the integrated stress response to glutamine catabolism within the rod photoreceptor cell are intriguing and innovative. Overall, the authors provide convincing data to highlight that photoreceptors utilize various fuel sources to meet their metabolic needs, and that glutamine is critical to these cells for their biomass, redox balance, function, and survival.

We greatly appreciate the reviewer's thoughtful comments and time spent reviewing this manuscript.

Weaknesses:

Recent studies have explored the metabolic "crosstalk" that exists within the mammalian retina, where metabolites are transferred between the various retinal cells and the retinal pigment epithelium. It would be of interest to test whether the conditional knockout mice have changes in metabolism (via qPCR such as shown in Figure 4 - Supplemental Figure 1) within the retinal pigment epithelium that may be contributing to the authors' findings in the neural retina. Additionally, the authors have very compelling data to show that inhibition of eIF2 α or supplementation with asparagine can delay photoreceptor death via OCT measurements in their conditional knockout mouse model (Figure 6G, H). However, does inhibition of eIF2 α or asparagine adversely impact the WT retina? It would also be impactful to know whether this has a prolonged effect, or if it is short-term, as this would provide strength to potential therapeutic targeting of these pathways to maintain photoreceptor health.

We agree with the reviewer that metabolic communication in the outer retina is crucial to the function and survival of both photoreceptors and RPE. Therefore, we have performed qRT-PCR on eyecups from cKO and WT mice at P14, prior to photoreceptor degeneration. These data, now included in Figure 4 – figure supplement 2, show no significant changes in genes related to glycolysis, pyruvate metabolism and the TCA cycle in eyecups from cKO mice compared to WT mice at P14. The only exception is a significant decrease in *Pdk4* in cKO mouse eyecups compared to WT, which was not observed in retina samples.

Additionally, we have added data demonstrating that systemic treatment with ISRIB does not adversely impact the anatomy of the wild-type retina. Specifically, we performed OCT after 21 days of ISRIB treatment via intraperitoneal delivery in WT mice and show that total retinal, ONL and inner segment/outer segment thickness is unchanged compared to vehicle. These data are now included in Figure 6 – figure supplement 2A. We have also included data to suggest that the effect of ISRIB extends beyond P21 in the cKO mouse. This data, presented in Figure 6 – figure supplement 2B, shows that at P28, ISRIB continues to statistically significantly increase ONL thickness compared to vehicle in cKO animals.

Reviewer #3 (Public Review):

Summary:

The authors explored the role of GLS, a glutaminase, which is an enzyme that catalyzes the conversion of glutamine to glutamate, in rod photoreceptor function and survival. The loss of GLS was found to cause rapid autonomous death of rod photoreceptors.

Strengths:

Interesting and novel phenotype. Two types of cre-lines were rigorously used to knockout the Gls gene in rods. Both of the conditional knockouts led to a similar phenotype, i.e. rod death. Histology and ERG were carefully done to characterize the loss of rods over specific ages. A necessary metabolomic study was performed and appreciated. Some rescue experiments were performed and revealed possible mechanisms.

We thank the reviewer for their comments and appreciation of the methods utilized herein to address the role of GLS-driven Gln catabolism in rod photoreceptors.

Weaknesses:

No major weaknesses were identified. The mechanism of GLS-loss-induced rod death seems not fully elucidated by this study but could be followed up in the future, and the same for GLS's role in cones.

We agree with the reviewer that the downstream metabolic and molecular mechanisms by which Gln catabolism impacts rod photoreceptor health are not fully elucidated. Defining these mechanisms will advance our understanding of photoreceptor metabolism and identify therapeutic targets promoting photoreceptor resistance to stress. Future studies are underway to uncover these mechanisms. Additionally, while outside the scope of the current manuscript, we have generated mice lacking GLS in cone photoreceptors specifically and are currently elucidating the role of GLS in cone photoreceptor metabolism, function, and survival. These results will be published in a separate manuscript.

Reviewer #1 (Recommendations For The Authors):

(1) The results could start at line 135, but the first paragraph isn't necessary. The data is published and could be referred to in the introduction.

We appreciate the reviewer's suggestion to shorten the beginning of the Results section; however, we believe the supplementary data, which is described in these lines, confirms the scRNAseq gene expression data, while adding GLS expression and localization data within the retina. The scRNAseq data and its publication was noted in the introduction, so we removed the sentence in line 117-119 that restates these results to shorten this section. We also reduced redundancy by removing an introductory sentence to the second Results paragraph.

(2) "However, like other metabolically-demanding cells, recent work has demonstrated that PRs have the flexibility to utilize fuel sources beyond glucose to meet their metabolic needs (Adler et al., 2014; Du, Cleghorn, Contreras, Linton, et al., 2013; Grenell et al., 2019; Joyal et al., 2016; Xu et al., 2020)." The paper by Daniele et al. demonstrated that glucose is essential for maintaining the viability of rod photoreceptor cells.

We thank the reviewer for highlighting published literature, which we apologetically overlooked. The reference for Daniele et al. has now been included.

(3) "Single-cell RNA sequencing data has demonstrated that Gls is expressed throughout the human and mouse retina and much greater than Gls2 (Voigt et al., 2020). The authors should indicate the specific databases searched in Spectacle.

We appreciate the reviewer's attention to detail and have now included the references in the Introduction for GSE63473 from Macosko et al. and GSE142449 from Voigt et al., which were the databases we used in Spectacle to assess *Gls* levels in the mouse and human retina, respectively.

References:

(1) Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A, McCarroll SA. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell*. 2015 May 21;161(5):1202-1214. doi: 10.1016/j.cell.2015.05.002. PMID: 26000488; PMCID: PMC4481139.

(2) Voigt AP, Binkley E, Flamme-Wiese MJ, Zeng S, DeLuca AP, Scheetz TE, Tucker BA, Mullins RF, Stone EM. Single-Cell RNA Sequencing in Human Retinal Degeneration Reveals Distinct Glial Cell Populations. *Cells*. 2020 Feb 13;9(2):438. doi: 10.3390/cells9020438. PMID: 32069977; PMCID: PMC7072666.

(4) *The immunolabeling in Figure 2 looks like the images are overexposed, and the GlS antibody is labeling the outer segment, not just the inner segment of photoreceptors.*

We thank the reviewer for their comments regarding our immunofluorescence data. There was background staining of the outer segment in both the WT and cKO retina with decreased GLS staining in the inner segment of the cKO rod photoreceptors at P14 demonstrating loss of GLS in rod photoreceptors similar to Figure 1B. For Figure 2E, we have provided adjusted images with PNA staining only that better represent the secondary cone degeneration that occurs in the rod photoreceptor-specific *Gls* cKO, which is the take home point of Figure 2E.

(5) *The authors could use a glutamate antibody to compare it to GlS KO mice as done in Davanger, S., Ottersen, O.P. and Storm-Mathisen, J. (1991), Glutamate, GABA, and glycine in the human retina: An immunocytochemical investigation. J. Comp. Neurol., 311: 483-494. <https://doi.org/10.1002/cne.903110404>*

We appreciate the reviewer's suggestion to assess glutamate levels in the wild-type and *Gls* KO retina via antibody labeling. Our targeted metabolomics studies in Figure 6A provide quantitative evidence that glutamate, the product of the GLS-catalyzed reaction, is decreased as one would expect in that *Gls* KO retina. The antibody would add to these data by providing the localization of glutamate in the retina. With a rod photoreceptor-specific genetic KO, we would expect glutamate levels to be decreased in these cells. The antibody may also show that glutamate is not only decreased in the rod photoreceptor inner segment, where GLS predominates, but also in the synaptic terminal in accordance with the reviewer's concerns regarding the impact of GLS KO on synaptic transmission. We have addressed this concern at length above, adding TEM images of the ribbon synapses in the GLS KO retina, and ERG analyses from the IND-cKO animals prior to significant degeneration. In the end, we agree with the reviewer that reduced Glu levels in the GLS cKO retina may impact synaptic transmission to a degree, but the synapses remain intact based on immunofluorescence and TEM analyses and a negative ERG pattern is not observed in the GLS cKO (i.e. iCre-Gls flox/flox) or IND-cKO mouse. As noted above, the structure of the retina in models that disrupt photoreceptor synaptic transmission is maintained (Dick et al. *Neuron*. 2003) or noted to have modest changes within the first two months of age with the outer nuclear layer not significantly altered until 8-10 months of age (Haeseleer et al. *Nature Neuroscience*. 2004). So, the impact of the reduced Glu levels on synaptic transmission in the GLS KO retina are unlikely to account in full for the rapid and profound photoreceptor degeneration observed. That said, the IND-cKO mouse, which allows us to assess photoreceptor function prior to

significant degeneration unlike the GLS cKO mouse (i.e. iCre-Gls flox/flox), demonstrates GLS-driven Gln catabolism plays a significant role in photoreceptor function but still does not demonstrate a negative ERG pattern. Therefore, assessing Glu localization in this mouse model 10 days post TAM will be informative as to how GLS-driven Gln catabolism impacts photoreceptor function prior to degeneration. The IND-cKO mouse model is currently being extensively characterized for future publication.

Reviewer #2 (Recommendations For The Authors):

Main Concerns:

(1) The authors checked for Gls2 compensation at P14 in the mouse retina. However, this data would be more compelling with an additional timepoint, particularly at P21 which is used in many of their figures throughout the study.

We thank the reviewer for their suggestion. Figure 1-figure supplement 1D demonstrates no change in *Gls2* gene expression at P14 between the WT and cKO retina. With regards to the reviewer's concern, in Figure 1-figure supplement 1E of the original submission, we demonstrate that the expression of GLS2 is not increased in the cKO retina at P21 via immunofluorescence.

(2) Recent studies have explored the metabolic "crosstalk" that exists within the mammalian retina, where metabolites are transferred between the various retinal cells and the retinal pigment epithelium. It would be compelling to see whether the cKO mice have changes in metabolism (via qPCR such as shown in Supplementary Figure 1 for Figure 4) within the RPE that may be contributing to their findings in the neural retina. Additionally, mention of this crosstalk and how it may impact their results should be added to the discussion.

We appreciate the reviewer's concern for metabolism changes in the RPE of *Gls* cKO mice. In agreement with reviewer 2, we performed qRT-PCR on eyecups from cKO and WT mice at P14, prior to photoreceptor degeneration. These data, now included in Figure 4 – figure supplement 2, show no significant changes in genes related to glycolysis, pyruvate metabolism and the TCA cycle in eyecups from cKO mice compared to WT mice at P14. The only exception is a significant decrease in *Pdk4* in cKO mouse eyecups compared to WT, which was not observed in retina samples.

(3) The authors use a tamoxifen-inducible cKO model to support their findings in developed rods. However, in Figure 3A it appears that this model has a greater reduction in GLS compared to the Rho-cre mouse model. Can the authors discuss this? Is this cre more efficient at targeting rods or is it leaky and may have affected other retinal cells?

We thank the reviewer for pointing out this interesting result associated with using the *Pde6g*-Cre-ERT2 mouse line. *Pde6g* is expressed by rods to a significant degree but also by cones (GSE63473, scRNAseq data). Therefore, the IND-cKO mouse likely knocks out GLS from both rods and cones upon the TAM induction. To this end, the immunofluorescence image in Figure 3B shows GLS is knocked out in both rod or cone inner segments unlike in Figure 1B where GLS remains in cones when using the rod photoreceptor-specific, *Gls^{fl/fl} Rho-Cre⁺* mouse. As such, as the astute reviewer noted, the fact that Western blot demonstrates greater reduction in GLS protein content fits with the protein being knocked out of both rods and cones. We have added this note about the mouse model in the corresponding text.

(4) The authors have very compelling data to show that inhibition of eIF2α can delay photoreceptor death via OCT measurements in their cKO mouse model (Figure 6G). However, does ISRIB adversely impact the WT retina? WT vehicle and ISRIB should be

shown. It would also be compelling to know whether this has a prolonged effect, or if it is short-term (i.e. would the effect still be present at P42)?

We appreciate the reviewer's comments regarding antagonizing the effects of p-eIF2a to prolong photoreceptor survival in the *Gls* cKO retina. As described above, we have data demonstrating systemic treatment with ISRIB does not adversely impact the anatomy of the wild-type retina (Figure 6-figure supplement 2A). Specifically, we treated WT animals with daily intraperitoneal ISRIB starting at P5 and performed OCT at P21 to show that total retinal, ONL and the inner segment/outer segment thickness is unchanged compared to vehicle-treated WT animals. Additionally, we have included data demonstrating the photoreceptor neuroprotective effect of ISRIB treatment in the *Gls* cKO mouse extends beyond P21 in the cKO mouse (Figure 6-figure supplement 2B).

(5) For Figure 6H, same as point #4.

While we have not specifically assessed potential retinal toxicity secondary to systemic Asn supplementation, oral Asn supplementation (up to 100mg/kg/day) was provided to patients for 24 months and found to be well-tolerated (PMID:31123592). Allometric scaling of this dose to the mouse would yield a mouse dose of 1234 mg/kg/day, which is much greater than the 200mg/kg/day dose provided here (PMID: 27057123). Additionally, a 90-day toxicity study of Asn in rats demonstrated a no observed adverse effect level of 1.62g/kg bodyweight/day in males and 1.73g/kg bodyweight/day in females (PMID: 18508175). The lower dose in that study equates to a mouse dose of 3.2g/kg bodyweight/day, well above the mouse dose utilized in this report. As such, future studies should focus on a dose-response relationship with Asn supplementation, and as the reviewer suggested, determining the duration of effect with Asn supplementation.

(6) Some of the results section belongs in the introduction or discussion and can be moved.

We have addressed the reviewer's concern by moving some of the results to the discussion and removing statements in the results that were either noted in the Introduction or conferred in the Discussion.

Minor Concerns:

(1) Scale bar mentions in the figure legends use plural when only one is present, or in some cases are missing. A scale bar should be added to the OCT images if possible.

We appreciate the reviewer's attention to detail, and information regarding scale bars has been updated in the figure legends.

(2) For Figures 1I and J, the sample size changes when J is a quantification of I. Please correct.

We have corrected the sample size to be consistent between Figures 1I and J.

(3) In Figure 1 - Figure Supplement 3 the P42 timepoint is not mentioned in the legend. Please correct.

We have now included the P42 timepoint in the legend for in Figure 1 – Figure Supplement 3 as well as the manuscript text.

(4) In Figure 1 - Figure Supplement 5 the wrong P value is mentioned in the legend. Please correct.

We have corrected the P value in the legend for Figure 1 – Figure Supplement 5.

(5) Can the authors double-check their ERG light intensity settings? They seem high. Please confirm if they are correct.

We appreciate the reviewer's concern for ERG light intensity settings and have confirmed the settings used in the study were 32 cd*s/m² and 100 cd*s/m² for scotopic and photopic ERG recordings, respectively.

(6) The legend key in Figure 2A would be more helpful if the axis were present by the representative traces.

We thank the reviewer for the suggestion of adding axes to the ERG traces. Figure 2A has been updated to reflect this modification.

(7) Can the authors check that the error bars are present in Figure 5E?

We appreciate the reviewer's concern for error bars in Figure 5E, which are included in the figure. The standard error in this experiment is so small that the symbols overlap with the error bars.

Reviewer #3 (Recommendations For The Authors):

Suggestions for improved or additional experiments, data, or analyses.

(1) Figure 6: ISRIB seems to give the most dramatic rescue of cKO GLS in P21 rods. Does it completely prevent rod death? i.e. What's the ONL thickness of P21 WT control? What's the ISRIB rescue of an older cKO animal, say P35?

The ONL thickness of P21 WT control is on average 0.06 mm (Figure 1E), while the ONL thickness of the GLS cKO retina with ISRIB treatment at P21 is on average 0.044 mm. Therefore, rod death is not completely prevented with ISRIB but rather, rod photoreceptor survival is prolonged. As noted above, we have provided data to demonstrate that the photoreceptor neuroprotective effect of ISRIB lasts beyond P21 (Figure 6-figure supplement 2B).

(2) What's the mechanistic link between ISR and GLS beyond current speculation? Does GLS have other unknown functions beyond converting glutamine to glutamate? Any novel insights from GLS protein structure?

We thank the reviewer for this thoughtful question. It is certainly possible that GLS has other functions outside of its role in glutaminolysis. It is well known that other metabolic enzymes have moonlighting functions including hexokinase 2, which has been shown to be important in preventing intrinsic apoptosis through blocking the binding of pro-apoptotic proteins to the mitochondria. While not directly related to ISR, a single report suggests GLS functions non-canonically in Gln-deprived states, promoting mitochondrial fusion to suppress ROS production (PMID: 29934617). Investigating the moonlighting functions of metabolic enzymes is part of our ongoing research program and GLS is included in these studies.

(3) Just curious about GLS cKO in cones. Any similar phenotype?

We appreciate the reviewer's curiosity regarding *Gls* cKO in cones and this study is currently ongoing with a poster presented at ARVO 2024 (Subramanya et al; Glutaminase-driven glutamine catabolism supports cone photoreceptor metabolism, function, and structure. Invest. Ophthalmol. Vis. Sci. 2024;65(7):193) and a manuscript in preparation. As discussed above, GLS knock out in cones likely impacts their function, in accordance with the data presented at ARVO 2024.

Recommendations for improving the writing and presentation.

(1) In the Discussion, lines 458-466, it's incorrect to compare the importance of glucose metabolism to GLS-dependent pathway to photoreceptors in this way. An alternative explanation: glucose metabolism is so important that the system has many redundancies, e.g. HK1 exists in addition to HK2, thus single gene KO leads to no phenotype. The only fair comparison is nutrient deprivation, e.g. taking out glucose or glutamine from retina explants (Punzo et al., 2009).

The reviewer makes an excellent point. While we do not see an upregulation of GLS2 in the retina or rod PRs upon GLS knockout (Figure 1-figure supplement 1 D and E), loss of *Gls* in rod PRs does alter the expression of many metabolism-related genes (Figure 4-figure supplement 1). We alluded to these data and the reviewer's point in the second paragraph of the discussion: "In any of these transgenic mouse models, PRs may use other transporters to take up fatty acids or glucose or rewire their metabolism to maintain metabolic homeostasis and stave off degeneration (Subramanya et al., 2023; Wubben et al., 2017). Our data show that any metabolic reprogramming that is occurring in the cKO mouse retina appears unable to significantly circumvent the significant and rapid PR degeneration suggesting the importance of Gln catabolism in rod PRs. Furthermore, inducing GLS knockdown in mature PRs also demonstrated rapid PR degeneration (Figure 3)."

In the revised article, we have amended these sentences to include the importance of metabolic redundancies. "In any of these transgenic mouse models, PRs may use other transporters to take up fatty acids or glucose, rewire their metabolism, or utilize metabolic redundancies to maintain metabolic homeostasis and stave off degeneration (Subramanya et al., 2023; Wubben et al., 2017). Our data show that any metabolic reprogramming that is occurring in the cKO mouse retina appears unable to significantly circumvent the significant and rapid PR degeneration suggesting the importance of Gln catabolism in rod PRs. Furthermore, inducing GLS knockdown in mature PRs also demonstrated rapid PR degeneration (Figure 3)."

(2) Please discuss the mosaic activity of Rho-cre used in this study, as described in the original study (Le et al 2006). Line 221 (Li et al 2005) seems to be a different Rho-Cre created by a different group. Please make sure the citation is correct and consistent.

We apologize for the confusion and have corrected the reference on line 221 to Le et al, 2006. The reviewer is correct that the original report (Le et al. 2006) demonstrated a mosaic of Cre-mediated recombination in rod photoreceptors and rod bipolar cells in the mouse line that had the shorter (0.2 kb) mouse opsin promoter-controlled Cre. In contrast, this same report showed only Cre-mediated recombination in rod photoreceptors in another line that utilized a long (4.1 kb) mouse opsin promoter-controlled Cre. We have published using this latter promoter-controlled Cre recombinase in at least 5 different mouse models (Wubben et al. 2017; Weh et al. 2020; Weh et al. 2023; Subramanya et al. 2023; the current report), and in all these models, we observe clear and consistent knockout by immunofluorescence only in rod photoreceptors with residual protein in cones and no significant change in protein expression in the INL where bipolar cells reside. Western blots confirm the reduction in protein expression.

| (3) *The authors should provide representative images of retina cross-sections for key rescue data (Figure 6G&H).*

As requested by Reviewer 3, representative histology images of retina cross-sections for the ISRIB and Asn rescue experiments in *Gls* cKO mice at P21 are now included in the manuscript in Figure 6 – figure supplement 3.

| *Minor corrections to the text and figures.*

| (1) *Spell out Gln in the Abstract when used for the first time.*

We have included glutamine (Gln) in the abstract upon first use.

| (2) *Line 433, Figure 6G should be 6H.*

Thank you for the correction, the manuscript has been updated.

<https://doi.org/10.7554/eLife.100747.2.sa0>