

Synaptic cell adhesion molecule *Cdh6* identifies a class of sensory neurons with novel functions in colonic motility

Reviewed Preprint

v2 • February 21, 2025

Revised by authors

Reviewed Preprint

v1 • November 21, 2024

Julieta Gomez-Frittelli, Gabrielle Devienne, Lee Travis, Melinda A Kyloh, Xin Duan, Tim J Hibberd, Nick J Spencer, John R Huguenard, Julia A Kaltschmidt 

Department of Chemical Engineering, Stanford University, Stanford, United States • Wu Tsai Neurosciences Institute, Stanford University, Stanford, United States • Department of Neurology & Neurological Sciences, Stanford University, Stanford, United States • College of Medicine and Public Health, Flinders Health & Medical Research Institute, Flinders University, Adelaide, Australia • Department of Ophthalmology, School of Medicine, University of California San Francisco, San Francisco, United States • Department of Neurosurgery, Stanford University School of Medicine, Stanford, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study characterizes the molecular signatures and function of a type of enteric neuron (IPAN) in the mouse colon, identifying molecular markers (*Cdh6* and *Cdh8*) for these cells. A battery of **compelling** and comprehensive experimental findings suggests data from other species are likely translatable to mice, bridging the abundant literature from humans and other mammals into this experimentally tractable animal model. This work will be of interest to scientists studying the motor control of the colon and more generally the enteric neuromuscular system.

<https://doi.org/10.7554/eLife.101043.2.sa3>

Abstract

Intrinsic sensory neurons are an essential part of the enteric nervous system (ENS) and play a crucial role in gastrointestinal tract motility and digestion. Neuronal subtypes in the ENS have been distinguished by their electrophysiological properties, morphology, and expression of characteristic markers, notably neurotransmitters and neuropeptides. Here we investigated synaptic cell adhesion molecules as novel cell type markers in the ENS. Our work identifies two Type II classic cadherins, *Cdh6* and *Cdh8*, specific to sensory neurons in the mouse colon. We show that *Cdh6*⁺ neurons demonstrate all other distinguishing classifications of enteric sensory neurons including marker expression of *Calcb* and *Nmu*, Dogiel type II morphology and AH-type electrophysiology and *I_H* current. Optogenetic activation of *Cdh6*⁺ sensory neurons in distal colon evokes retrograde colonic motor complexes (CMCs), while pharmacologic blockade of rhythmicity-associated current *I_H* disrupts the spontaneous generation of CMCs. These findings provide the first demonstration

of selective activation of a single neurochemical and functional class of enteric neurons, and demonstrate a functional and critical role for sensory neurons in the generation of CMCs.

One-Sentence Summary

Intrinsic sensory neurons of the enteric nervous system in the mouse distal colon exclusively express synaptic cell adhesion molecules *Cdh6* and *Cdh8*, evoke retrograde colonic motor complexes (CMCs) when stimulated, and possess rhythmicity-associated I_H current, involved in producing spontaneous CMCs.

Introduction

Sensory signaling within the gastrointestinal (GI) tract plays a critical role in the autonomous regulation of digestion. The GI tract is the only internal organ system containing its own sensory neurons. Intrinsic primary afferent neurons (IPANs) detect relevant stimuli through chemo- and mechano-sensation and direct appropriate GI functions via downstream components of the enteric nervous system (ENS), including ascending and descending interneurons, and excitatory and inhibitory motor neurons (1 [↗](#)). These neuronal subtypes have begun to be distinguished morphologically, electrophysiologically, and by marker expression, classically especially of neurotransmitters (2 [↗](#), 3 [↗](#)), and together this information has provided an opening to characterize individual neuron subtype function within the GI tract.

Synaptic cell adhesion molecules define neuronal subtype connectivity within many regions of the CNS. Type II Cadherins are a family of synaptic cell adhesion molecules with combinatorial expression in multiple neural circuits of the CNS, including retina, limbic, olivonuclear, and auditory projection systems (4 [↗](#)–6 [↗](#)). Type II Cadherins bind homophilically by expression of the same cadherin at both the pre- and post-synapse, which stabilizes developing synapses between correct partners while incorrect synapses are pruned away (7 [↗](#), 8 [↗](#)). Recent RNA-Seq studies of human and mouse ENS have identified synaptic cell adhesion molecules, including Type II Cadherins, expressed in enteric neuronal subtypes (9 [↗](#)–11 [↗](#)). However, the specificity of their expression has yet to be harnessed to assess neuronal subtype-specific function in the ENS.

Here we identify Type II Cadherin, *Cdh6*, as a novel marker for IPANs of the colonic ENS. We demonstrate the sensory identity of *Cdh6* neurons by immunohistochemical, morphological and neurophysiological classification. *Cdh6* neurons express IPAN markers *Calcb* and *Nmu*. Sparse labeling of individual IPANs reveals they project mainly circumferentially and branch extensively in myenteric ganglia. Whole cell patch clamp recordings of sensory neurons *in situ* reveal action potential (AP) slow after-hyperpolarizations characteristic of IPANs, and hyperpolarization-activated cationic current (I_H), a rhythmicity indicator in thalamocortical and other systems (12 [↗](#)). Using a *Cdh6* genetic mouse model, we show that optogenetic activation of distal colon IPANs is sufficient to evoke retrograde CMCs, while pharmacologic block of I_H in IPANs disrupts colonic rhythmicity and reversibly abolishes spontaneous CMCs.

Results

Expression of the type II classic cadherin *Cdh6* in colonic IPANs

To identify Cadherins expressed in enteric neuronal subtypes in mouse, we screened recently published RNA-Seq data (10 [↗](#), 11 [↗](#)) for Classic Type II cadherin expression. *Cdh6* and *Cdh8* appeared to be restricted to IPAN subsets in both small intestine and colon (10 [↗](#), 11 [↗](#)). *Cdh9* was

previously identified in a separate population of IPANs in the small intestine (9–11). We validated *Cdh6* and *Cdh8* expression by RNAscope *in situ* hybridization in the myenteric plexus, which contains the enteric motility circuitry. *Cdh6* mRNA was expressed in $14.7 \pm 0.8\%$ of myenteric neurons in small intestine (jejunum) and in $6.8 \pm 0.3\%$ of myenteric neurons in distal colon (mean $\pm$ SEM) [Fig.1A–C]. *Cdh8* was almost exclusively co-expressed in *Cdh6*⁺ neurons, although at a much lower level of detection [Fig.1D–H]. We therefore focused our further analysis on *Cdh6*⁺ neurons.

To confirm IPAN identity of *Cdh6*⁺ neurons, we first established the differential expression of two putative and broadly used markers of IPANs, *Calcb* and *Nmu* (3, 9, 11). We found that all *Nmu*⁺ neurons co-express *Calcb* in both jejunum and distal colon [Fig.1I–K]. In contrast, only about half of *Calcb*⁺ neurons in the jejunum and two-thirds in the distal colon co-express *Nmu* [Fig.1L].

We next assessed co-expression of *Cdh6* with *Calcb* and *Nmu*. In the jejunum, we found that nearly all *Nmu*⁺ neurons and *Calcb*⁺ neurons express *Cdh6* [Fig.1P, U], though only about three-quarters of all *Cdh6*⁺ neurons express *Calcb* and only about half express *Nmu* [Fig.1O, T]. In contrast, in the distal colon, while *Cdh6* is only expressed in about two-thirds of all *Calcb*⁺ neurons [Fig.1U], nearly all *Cdh6*⁺ neurons express *Nmu* and *Calcb* [Fig.1O, T]. Taken together, our data show that in the myenteric plexus, *Cdh6* is expressed exclusively in *Calcb*⁺/*Nmu*⁺ IPANs in the mouse distal colon [Fig.1W, X].

Mouse colonic IPANs display AH-type electrophysiology and I_H current

We next assessed the electrophysiological properties of *Cdh6*⁺ IPANs. We focused our analysis on the colon, and for ease of neuron tracing, took advantage of a genetic strategy to sparsely label *Cdh6*⁺ neurons. Previous studies of Hb9:GFP transgenic mice have shown that due to the inserted transgene's proximity to *Cdh6*, *Cdh6*⁺ neurons can express eGFP (13). Hb9:GFP⁺ neurons were rare and projected extensively throughout the myenteric plexus [Fig.2A–C]. *In situ* hybridization confirmed eGFP expression was limited to a small fraction ($3.5 \pm 0.8\%$) of *Cdh6*⁺ colonic neurons [Fig.2D, F–H], and all eGFP⁺ neurons expressed *Cdh6* [Fig.2E].

We developed a protocol to perform whole-cell patch-clamp recordings (14) in Hb9:GFP⁺ colonic neurons in the distal colon. Membrane capacitance reflecting overall size of these cells was 32 ± 8.7 pF [Fig.3D]; their resting membrane potential (RMP) was -49.4 ± 2.9 mV [Fig.3C]. The input resistance (R_{in}) was 393 ± 54.7 M Ω [Fig.3E] as computed from the slope of the voltage-current (V-I) relationship. All patched neurons had large-amplitude action potentials (AP; 72 ± 2.5 mV, [Fig.3F, E]) with threshold of -26.4 ± 0.9 mV [Fig.3L] and a half width of 1.2 ± 0.1 ms [Fig.3H] elicited at rheobase (20 ± 4.5 pA, [Fig.3G]), each followed by an afterhyperpolarization (AHP= -67.7 ± 0.9 mV, [Fig.3K]). In addition, the first derivative of membrane voltage during the action potential (dV/dt, [Fig.3J]) exhibited an inflection during the repolarization phase, suggesting the presence of fast and slow AP repolarization mechanisms [Fig.3F]. Patched Hb9:GFP⁺ neurons generally could not sustain repetitive APs in response to long depolarizing current pulses. All patched neurons responded to hyperpolarizing current pulses with a time-dependent membrane potential sag (-6.3 ± 1.2 mV, [Fig.3B, M]), and a rebound depolarization following the hyperpolarization [Fig.3B, N].

Consistent with these findings of sag and rebound, voltage clamp step hyperpolarizations revealed I_H (hyperpolarization-activated current, [Fig 3.N]) and, upon repolarization, I_T (transient inward presumed Ca^{2+} current) (15) in terms of its kinetics [Fig 3.O] and steady-state inactivation [Fig.3O, P]. Taken together, these results show that Hb9:GFP⁺ / *Cdh6*⁺ distal colonic neurons have AP after-hyperpolarizing (AH)-type electrophysiology typical of IPANs, including rhythm generating currents I_H and I_T (16–18).




Fig. 1.

***Cdh6* expression overlaps with IPAN markers *Calcb* and *Nmu*.**

(A, B) Representative images of jejunum (A) and distal colon (B) myenteric plexus labeled with HuC/D (IHC) (blue) and *Cdh6* (RNA) (red). (C) Proportion of total HuC/D neurons positive for *Cdh6* (jejunum, $n = 9$; distal colon, $n = 9$). (D, E) As in (A, B) for HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and *Cdh8* (RNA) (green). (F) Proportion of *Cdh6*⁺ neurons positive for *Cdh8* (jejunum, $n = 8$; distal colon, $n = 8$). (G) Proportion of *Cdh8*⁺ neurons positive for *Cdh6* (jejunum, $n = 8$; distal colon, $n = 8$). (H) Proportion of total HuC/D neurons positive for *Cdh8* (jejunum, $n = 8$; distal colon, $n = 8$). (I, J) As in (A, B) for HuC/D (IHC) (blue), *Calcb* (RNA) (red), and *Nmu* (RNA) (green). (K) Proportion of *Nmu*⁺ neurons positive for *Calcb* (jejunum, $n = 5$; distal colon, $n = 5$). (L) Proportion of *Calcb*⁺ neurons positive for *Nmu* (jejunum, $n = 5$; distal colon, $n = 5$). (M, N) As in (A, B) for HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and *Nmu* (RNA) (green). (O) Proportion of *Cdh6*⁺ neurons positive for *Nmu* (jejunum, $n = 7$; distal colon, $n = 8$). (P) Proportion of *Nmu*⁺ neurons positive for *Cdh6* (jejunum, $n = 7$; distal colon, $n = 8$). (Q) Proportion of total HuC/D neurons positive for *Nmu* (jejunum, $n = 7$; distal colon, $n = 8$). (R, S) As in (A, B) for HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and *Calcb* (RNA) (green). (T) Proportion of *Cdh6*⁺ neurons positive for *Calcb* (jejunum, $n = 9$; distal colon, $n = 7$). (U) Proportion of *Calcb*⁺ neurons positive for *Cdh6* (jejunum, $n = 9$; distal colon, $n = 7$). (V) Proportion of total HuC/D neurons positive for *Calcb* (jejunum, $n = 9$; distal colon, $n = 9$). (W, X) Schematic of marker overlap in jejunum (W) and distal colon (X). Scale bar represents 100 μm for (A, B, I, J, M, N, R, S), 50 μm for (D, E). All charts (mean $\pm$ SEM). * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.




Fig. 2.

Hb9:GFP+ is expressed in a small proportion of *Cdh6*+ colon myenteric neurons.

(A, B) Representative images of Hb9:GFP+ distal colon myenteric plexus labeled with HuC/D (IHC) (magenta) and GFP (green). (C) Proportion of total distal colon HuC/D neurons positive for GFP (n = 3). (D) Proportion of distal colon eGFP+ neurons positive for *Cdh6* (n = 3). (E) Proportion of distal colon eGFP+ neurons positive for *Cdh6* (n = 3). (F-H) Representative images of Hb9:GFP+ distal colon myenteric plexus labeled with HuC/D (IHC) (blue), *Cdh6* (RNA) (red), and eGFP (RNA) (green). Scale bar represents 200 μm for (A, B), 50 μm for (E-G). All charts (mean ± SEM).




Fig. 3.

Hb9:GFP+ distal colon neurons have AH electrophysiological characteristics.

(A) IR videomicroscopy image of an Hb9:GFP distal colon neuron that presented a large soma located in a ganglion (scale bar, 10 μ m). (B) Current-clamp recordings of the same neuron in (A) obtained in response to application of current pulse (bottom traces) of -50 pA, and $+10$ pA. Note the presence of a sag and a post-hyperpolarization rebound depolarization. (C-E) Box-and-whisker plots of cellular properties of recorded neurons. (C) resting membrane potential (RMP), (D) capacitance (C_m), and (E) membrane resistance (R_m) ($N = 5$). (F) Averaged traces of the first spike (rheobase action potential) after a depolarization step of 1 s. (J) Averaged derivative traces of the first spike (rheobase action potential). An inflection on the repolarizing phase is observed in the first derivative (arrow). (G-I, K-N) Box-and-whisker plots of electrophysiological properties of recorded neurons; rheobase action potential (AP, G-I) (G) current threshold, (H) half-width, (I) amplitude, (K) afterhyperpolarization (AHP), and (L) threshold. (M, N) non-AP properties sag (mV) and rebound (mV). (O) H and T currents in recorded neurons. Top: example of currents obtained from voltage protocol. Bottom: 500 ms hyperpolarizations ranging from -90 to -45 for 500 ms followed by depolarizing to -40 mV. Hyperpolarizations evoked slowly activating inward current (H-Current, arrow), followed by a transient inward current upon post-conditioning step to -30 mV (T-current, arrow) Largest T and H currents were obtained with the most hyperpolarized potentials (red trace). (P) Normalized peak I_T plotted versus holding potential to obtain the I/I_{max} curve ($N = 3/3$). Scale bar represents 10 μ m for (A).

Colonic IPANs have Dogiel type II morphology and abundant projections throughout the myenteric plexus

To visualize the morphology and projections of individual patched Hb9:GFP+ neurons, we included biocytin in the internal solution for post-fixation single-cell tracing [Fig.4]. Patched Hb9:GFP+ neurons displayed Dogiel type II morphology (19), with large smooth cell somas and multiple branching neurites. Projections were mainly circumferential and extensively branched within myenteric ganglia. Thus, Hb9:GFP+ neurons display morphological features characteristic of IPANs (2, 19).

To further visualize the full extent of IPAN circuitry in the myenteric plexus, we intercrossed *Cdh6*^{CreER} (20) and LSL-tdTomato (Ai14) (21) mice and induced Cre expression at 5-8 weeks of age [Fig.5A, B]. We confirmed tdTomato+ labeling in myenteric *Cdh6*+ neurons [Fig.5E, F], representing about 5% of the total neuronal population [Fig.5G]. tdTomato+ neurons had large cell somas (major axis, $27.8 \pm 0.7 \mu\text{m}$; minor axis, $15.9 \pm 0.4 \mu\text{m}$) [Fig.5D]. All ganglia of the myenteric plexus were densely innervated by tdTomato+ fibers [Fig.5A, B], which also projected into the circular muscle. We noted additional tdTomato+ labeling of some putative longitudinal and circular muscle cells [Fig.5A, B]. Taken together, our data reveal an IPAN array that spans the entire motility circuitry of the colonic myenteric plexus.

Optogenetic activation of distal colon IPANs evokes CMCs

In thalamocortical relay neurons, I_H contributes to intrinsic slow rhythmic burst firing at 1-2Hz (22), but the function of I_H in colonic IPANs is not known. During colonic motor complexes (CMCs), large regions of the ENS oscillate in synchrony at 1-2Hz to generate traveling contractions along the colon (23). Recent calcium imaging studies have shown that IPANs participate, along with all other subtypes of enteric neurons, in this synchronized oscillatory firing (24). Furthermore, our electrophysiological studies confirm the presence of I_H in mouse colonic IPANs [Fig.3N]. However, the role of IPANs in spontaneous CMCs is not well understood.

To interrogate the functional role of *Cdh6*+ IPANs in CMCs, we performed *ex vivo* colonic contraction force recordings in conjunction with optogenetic activation. To express ChR2-eYFP in *Cdh6*+ cells, we intercrossed *Cdh6*^{CreER} and LSL-ChR2-eYFP (Ai32) (25) mice and induced Cre expression at 5-8 weeks of age [Fig.6A-C]. In *Cdh6*^{CreER}+;ChR2-eYFP+ colon preparations we observed spontaneous CMCs at regular intervals of about 3-5 minutes. Blue light stimulation of *Cdh6*+ IPANs in distal colon 60-90 seconds after a spontaneous CMC (“control” CMC) resulted in an evoked, premature CMC that began during stimulation (N = 17 stimulations; n = 5 mice) [Fig.6D, E]. Evoked CMCs traveled retrogradely from the distal to the proximal colon. They were similar to spontaneous CMCs in peak amplitude, area under the curve, and duration, though the contractile force (peak amplitude and AUC) was slightly weaker in the proximal colon [Fig.6F-H]. Blue light stimulation in proximal or mid colon failed to generate CMCs (n = 5/5, data not shown). In comparison, stimulation in control *Cdh6*^{CreER}-negative;ChR2-eYFP+ colons failed to evoke any CMCs (N = 28 stimulations; n = 7 mice) [Fig.S1].

CMCs have previously been shown to depend on nicotinic cholinergic transmission (24). We performed optogenetic stimulation in the presence of hexamethonium, a blocker of nicotinic cholinergic transmission. Spontaneous CMCs were abolished in hexamethonium, and CMCs could not be evoked by optogenetic stimulation [Fig.6I]. We conclude that activation of *Cdh6*+ distal colon IPANs evokes retrograde-traveling but otherwise characteristic and hexamethonium-sensitive CMCs.




Fig. 4.

Hb9:GFP+ distal colon neurons have circumferential branching projections.

(A-C) Representative images of Hb9:GFP+ distal colon myenteric plexus labeled with HuC/D (blue), streptavidin (red), and GFP (green). (D, E) Tracings of Hb9:GFP+ distal colon neurons filled with biocytin during whole cell patch clamp recording. (F) Image of patched and filled Hb9:GFP+ distal colon neuron traced in (E). (G, H) Inset of (F). Scale bar represents 40 μm for (A-C), 200 μm for (D-F), 100 μm for (G, H).




Fig. 5.

Cdh6^{CreER}+/ tdTomato+ neurons have Dogiel type II morphology.

(**A**, **B**) Representative images of *Cdh6^{CreER}+/tdTomato+* distal colon myenteric plexus labeled with HuC/D (IHC) (magenta) and tdTomato (IHC) (green). (**C**) Inset of (**A**). (**D**) Dimensions of tdTomato+ neurons (major and minor axes) ($N = 73$; $n = 3$). (**E**) Representative image of *Cdh6^{CreER}+/tdTomato+* distal colon myenteric plexus labeled with HuC/D (IHC) (blue), tdTomato (IHC) (red), and *Cdh6* (RNA) (green). Arrowheads indicate *Cdh6*+ / tdTomato+ cells; arrowhead, *Cdh6*+ / tdTomato-negative cell. (**F**) Proportion of *Cdh6*+ distal colon neurons positive for tdTomato ($n = 6$). (**G**) Proportion of total HuC/D neurons positive for tdTomato (proximal colon, $n = 5$; mid colon, $n = 5$; distal colon, $n = 10$). Scale bar represents 100 μm for all images. All charts (mean $\pm$ SEM).




Fig. 6.

Optogenetic stimulation of distal colonic *Cdh6*⁺ neurons evokes CMCs, while pharmacologic blockade of *I_H* abolishes spontaneous CMCs

(A, B) Representative images of *Cdh6*^{CreER+};*ChR2-eYFP*⁺ distal colon myenteric plexus labeled with HuC/D (magenta) and GFP (green). (C) Proportion of total distal colon HuC/D neurons positive for *ChR2-eYFP* ($n = 6$). (D) Representative force traces. Blue bars indicate timing of light stimulation. LEDs placed distal to distal hook. (E) CMC intervals recorded from force traces. Evoked (blue) intervals represent the time from the prior spontaneous CMC before stimulation to the evoked CMC following stimulation. Control (grey) intervals represent the time between the spontaneous CMC prior to stimulation and the previous spontaneous CMC ($n = 5$). Paired t test, one tailed. (F) CMC peak amplitude recorded from force traces. Evoked (blue) indicates the evoked CMC following stimulation. Grey (control) indicates the spontaneous CMC prior to stimulation ($n = 5$). Paired t test, two tailed. (G) CMC AUC (area under the curve). Evoked (blue) and control (grey) as in (F) ($n = 5$). Paired t test, two tailed. (H) CMC duration. Evoked (blue) and control (grey) as in (F) ($n = 5$). Paired t test, two tailed. (I) Representative force traces. Hex indicates addition of 300 μ M hexamethonium. Blue bars indicate timing of light stimulation. LEDs placed distal to distal hook ($n = 5/5$). (J) Representative force traces on tethered pellets. First arrowhead indicates addition of 10 μ M ZD7288. Second arrowhead indicates washout in Krebs. Third arrowhead indicates addition of 1 μ M TTX. ZD7288 abolished CMCs in both proximal and distal colon ($n = 6/6$, $p = 0.0022$, Fisher's exact test). Washout in Krebs restored CMCs in both proximal and distal colon ($n = 6/6$, $p = 0.0022$, Fisher's exact test). (K) As in (J). First arrowhead indicates addition of 2 mM CsCl. Second arrowhead indicates washout in Krebs. Third arrowhead indicates addition of 1 μ M TTX. Typical CMC production was impaired or altered by CsCl (proximal colon, $n = 5/6$, $p = 0.0152$; distal colon, $n = 6/6$, $p = 0.0022$, Fisher's exact test): increased frequency (proximal colon, $n = 5/6$, $p = 0.0152$; distal colon, $n = 6/6$, $p = 0.0022$, Fisher's exact test), decreased in amplitude (proximal colon, $n = 5/6$, $p = 0.0152$; distal colon, $n = 4/6$, $p = 0.0606$, Fisher's exact test); retrograde force (proximal colon, $n = 2/6$; distal colon, $n = 2/6$). Scale bar represents 100 μ m for (A, B). * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Blockade of I_H current in colonic IPANs disrupts CMC production

To determine whether I_H in IPANs may contribute to oscillatory firing driving CMCs, we measured colonic contraction force on a tethered pellet in the presence of I_H blockers ZD7288 or CsCl (26 [Fig. 6J](#), 27 [Fig. 6J](#)). ZD7288 blocks I_H in all IPANs, including *Cdh6*⁺ IPANs. Spontaneous CMCs were recorded in all preparations of both proximal and distal colon prior to drug application [Fig. 7]. Addition of 10 μ M ZD7288 to the recording chamber abolished spontaneous CMCs and washout of ZD7288 recovered spontaneous CMC activity [Fig. 6J [Fig. 6J](#), Fig. S2 [Fig. S2](#)]. Addition of 2mM CsCl also impaired or altered spontaneous rhythmic production of typical CMCs [Fig. 6K [Fig. 6K](#), Fig. S3 [Fig. S3](#)]; rhythmic contractions increased in frequency (proximal colon, $n = 5/6$; distal colon, $6/6$), decreased in amplitude (proximal colon, $n = 5/6$; distal colon, $n = 4/6$, $p = 0.0606$), or in some cases even included significant retrograde force components (proximal colon, $n = 2/6$; distal colon, $n = 2/6$). We conclude that pharmacologic blockade of I_H in IPANs impairs the production of CMCs in the mouse colon.

Discussion

Our study shows that in the myenteric plexus, *Cdh6* is expressed exclusively in *Calcb*⁺/*Nmu*⁺ IPANs located in the mouse distal colon, while in the small intestine, *Cdh6* is also expressed in some *Calcb*⁺/*Nmu*⁺ and *Calcb*⁺/*Nmu*⁺ neurons. We confirm the IPAN identity of *Cdh6*⁺ distal colonic neurons by electrophysiological recordings revealing AH-type signature, and single neuron tracings showing Dogiel type II morphology. Finally, we demonstrate that activation of IPANs in the distal colon evokes retrograde CMCs, while pharmacologic blockade of I_H , a rhythmicity associated current we show also present in mouse colonic IPANs, disrupts spontaneous CMC generation.

Together with *Cdh8*, which we show to be co-expressed with *Cdh6*, our study validates two new adhesion molecules specific to IPANs in the distal colon. Notably, these markers show a different expression pattern than another cadherin, *Cdh9*, which is exclusively expressed in the small intestine in mouse, not in the colon (9 [Fig. 9](#)). *Cdh9* is expressed in a subset of IPANs not expressing either *Calcb* or *Nmu* (3 [Fig. 3](#), 9 [Fig. 9](#), 11 [Fig. 11](#)). It is possible that in the small intestine *Cdh6* and *Cdh9* mark some of the same neurons. However, RNA-Seq data from two separate studies suggest this is unlikely (10 [Fig. 10](#), 11 [Fig. 11](#)). We conclude that *Cdh6*/*Cdh8* IPANs are a separate population from *Cdh9* IPANs. Finally, our study was limited to the myenteric plexus, containing motility circuitry. IPANs are also present in the submucosal plexus, and it would be interesting to investigate *Cdh6* expression and neuronal subtype in the SMP.

Though IPANs are positioned to initiate motility by activating other subtypes in enteric circuitry, their role in spontaneous CMCs has been debated, as CMCs can occur in the absence of luminal contents, without any apparent stimulus for IPANs to sense, or upon stimulation of nitrergic populations (2 [Fig. 2](#), 28 [Fig. 28](#)). Only recently has evidence emerged to suggest that IPANs participate in oscillatory rhythmic firing of the ENS during CMCs, and that activation of neuronal subtypes expressing calretinin, including IPANs, together can evoke CMCs (24 [Fig. 24](#), 29 [Fig. 29](#)). Our results demonstrate that excitement of IPANs alone in the distal colon is capable of producing retrograde CMCs. The exact mechanism causing CMC generation at an established and controlled frequency of once every few minutes, without evident stimulus, remains to be determined.

A major observation was that optogenetic stimulation of *Cdh6*⁺ neurons readily evoked CMCs from the distal colon, but never from the proximal colon. Failure of proximal colonic stimulation to evoke CMCs may reflect inhibition by IPAN recruitment of descending pathways; or conversely, high efficacy of stimulation from distal colon may reflect bias toward activation of ascending excitatory cholinergic pathways. Our classification of *Cdh6*⁺ neurons as IPANs in the distal colon

may not extend to the proximal colon; thus, we cannot extrapolate that *Cdh6*⁺ optogenetic activation in the colon is restricted to IPANs as in the distal colon. In addition, our stimulus paradigm in the proximal colon may not have activated enough neurons based on light density and illumination. In previous studies, optogenetic stimulation of calretinin-expressing neurons or choline acetyl transferase (ChAT)-expressing neurons (both of which include IPANs, interneurons and motor neurons) elicited anterograde CMCs regardless of stimulus location, including proximal colon (29, 30). Proximal colon stimulation of nitric oxide synthase-expressing neurons also elicited CMCs (28). These studies, as noted, each activated multiple neuronal classes employing the same neurotransmitter. How broad and non-specific activation of such disparate neuron classes readily evokes CMCs remains unclear.

Spontaneous and evoked CMCs were abolished in hexamethonium, confirming that CMC synchronous firing is dependent on nicotinic cholinergic transmission (24). This reinforces that nicotinic cholinergic transmission is required for greater activation and synchrony of the entire ENS motility network to generate CMCs.

Through our electrophysiological investigation of IPANs in the distal colon using voltage clamp, our work reveals the presence of two voltage gated ion conductances and their underlying currents, I_T and I_H . Slow AHP, I_H and I_T have been previously identified as distinguishing characteristics of IPANs in rat and guinea pig (16–18). Although prior studies noted I_H and proposed I_T in guinea pig Dogiel type II neurons (17), they have not previously been reported in studies of intact mouse distal colon myenteric plexus due to the conventional reliance on sharp electrode recordings in which voltage clamp is not possible (31). The presence of I_H and I_T in thalamocortical relay neurons and other cell types supports intrinsic rhythmicity (22, 32). It is possible that these two currents may similarly promote autonomous rhythmic activity in colonic IPANs.

I_H is conducted through hyperpolarization-activated cyclic nucleotide gated (HCN) channels (33). HCN channel family members HCN1 and HCN2 have been shown to be present in mouse distal colonic Dogiel type II neurons (16), and RNA-Seq ENS screens similarly indicate their expression in IPANs (10, 11). Knockout of HCN2 in mouse leads to a severe growth restriction phenotype due to malnutrition and GI dysmotility (34). Here we demonstrate that pharmacologic blockade of I_H , which we show to be present in IPANs, with two distinct HCN channel blockers ZD7288 and CsCl abolishes CMCs, an otherwise persistent and ongoing pattern of motor activity in the mouse colon. We speculate that I_H in colonic IPANs, as in thalamocortical neurons, plays a role in promoting either rhythmic oscillatory single neuron activity, or network burst firing, or both (22, 32). Blocking I_H may impair individual IPANs' ability to fire rhythmically, or the ability of IPANs to synchronize into a network burst firing mode. Failure of IPANs to fully activate and synchronize could prevent generation of both (1) synchronized rhythmic myenteric network activation of motility circuits, and (2) the resulting synchronized contractions that sum to much larger contractile forces during CMCs.

Type II cadherins are most commonly homophilic synaptic cell adhesion molecules (7, 8). *Cdh6* can also form heterodimers with *Cdh7*, *Cdh10*, and *Cdh14* (35). However, *Cdh10* and *Cdh14* are very lowly expressed in the colon by single cell RNA sequencing (10), and we performed RNAscope for *Cdh7* and did not observe any expression (data not shown). Restricted expression of two Type II cadherins, *Cdh6* and *Cdh8*, to mouse colonic IPANs raises the possibility of these cadherins supporting IPAN-IPAN synaptic connections. While broadly speaking, IPANs are not known to receive synaptic input and in fact have been characterized electrophysiologically by their lack thereof (36), some work has in fact suggested that AH-AH neuron interconnected pairs may exist (37). Immunohistochemical and electron microscopy investigation of synapses on enteric neurons further showed that calbindin-positive neurons, presumed IPANs, do receive synapses, though fewer than non-calbindin neurons, and some of those synapses were also calbindin-reactive (38). These observations informed a proposed "IPAN driver circuit" theory, in

which IPANs form an interconnected network of positive feedback to synchronize and amplify sensory signaling and thus activate large swaths of enteric circuitry (39 [↗](#)). In contrast, more recently, activation of large regions of the ENS has been suggested to be driven by interneuronal networks (40 [↗](#)). It is important to note that our study was unsuccessful in localizing Cdh6 protein via immunohistochemistry, to confirm protein expression in neurons or in Cdh6-cre-tdT+ muscle cells or visualize synaptic connections between neurons or connections to other cell types. Further investigations are necessary to determine whether synaptic adhesion molecules, such as *Cdh6* and *Cdh8*, may in fact support IPAN-IPAN synapses underlying an “IPAN driver circuit.”

Materials and methods

Mice

All procedures conformed to the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and were approved by the Stanford University Administrative Panel on Laboratory Animal Care. Mice were group housed up to a maximum of five adults per cage. Food and water were provided ad libitum and mice were maintained on a 12:12 LD cycle. Male and female mice were used in all experiments.

Wild-type C57BL/6J mice (#000664), Hb9:GFP mice (#005029), Ai14 (#007908), and Ai32 (#024109) (21 [↗](#)) mice were obtained from the Jackson Laboratory and from the Animal Resource Center (ARC) in Western Australia, with JAX heritage. Cdh6^{CreER} mice (#029428) (20 [↗](#)) were provided by Xin Duan (UCSF). Cdh6^{CreER} mice were crossed to Ai14 mice and Ai32 mice to generate mice heterozygous for each allele, termed Cdh6^{CreER};Ai14 and Cdh6^{CreER};Ai32. Tamoxifen (20 mg/mL in corn oil) was administered via oral gavage to a final dose of 2.5mg/10g mouse for five consecutive days beginning at 5-8 weeks of age. Induced mice were group housed for at least four weeks prior to experiments.

Adult male and female mice (Cdh6^{CreER};Ai32 and Cdh6^{CreER};Ai32) aged 16 to 19 weeks were euthanised by isoflurane inhalation overdose in accordance with Flinders Animal Welfare Committee guidelines (ethics approval #4004). The protocol for animal euthanasia is approved by the National Health and Medical Research Council (NHMRC) Australian code for the care and use of animal for scientific purposes (8th edition, 2013) and recommendations from the NHMRC Guidelines to promote the wellbeing of animals used for scientific purposes (2008).

Dissection

Mice were culled by CO₂ and cervical dislocation. Small intestine and colon were removed and flushed with ice cold PBS, then placed in a Sylgard-lined Petri dish with ice cold PBS for further dissection.

Wholemount preparations

Intestinal segments were prepared and fixed as in Gomez-Frittelli *et al* (41 [↗](#)). Briefly, each intestinal segment was opened along the mesentery border, pinned flat under light tension serosa-side up, and fixed in 4% PFA in PBS at 4°C with gentle rocking for 90 minutes. Segments were washed three times in PBS at 4°C for at least 10 minutes with gentle rocking. Muscularis was separated from the mucosa at one end of the segment with fine forceps for 2-3 mm, then pinned mucosa-side up in the dish. The mucosa was peeled away from the muscularis with fine forceps while the muscularis was gently held down in the dish with a cotton swab. For immunohistochemistry, segments were processed immediately or stored in PBS with 0.1% NaN₃ at 4°C until use. For RNAscope, muscularis segments were post-fixed in 4% PFA in PBS at 4°C with gentle rocking overnight, then washed three times in PBS at 4°C with gentle rocking for at least 10 minutes each wash before use.

Immunohistochemistry

Immunohistochemistry was performed as described previously (42). Briefly, muscularis wholemount tissue segments about 7mm x 7mm were incubated with primary antibodies in PBT (PBS with 1% BSA and 0.3% Triton-X 100) at 4°C with gentle rocking overnight, then washed three times in PBT for at least ten minutes each at room temperature with gentle shaking. Tissues were incubated in secondary antibodies in PBT for 2 hours with gentle shaking, washed twice in PBT, and twice in PBS, then mounted on Superfrost Plus slides with Fluoromount G medium (Southern Biotech). Primary antibodies: human anti-HuC/D (1:75k) (gift from Vanda Lennon); sheep anti-GFP (1:1k) (Biogenesis); rabbit anti-RFP (1:1k) (Rockland); rabbit anti-PGP9.5 (1:4k) (Abcam). Secondary antibodies: donkey anti-human AlexaFluor (AF)-647 (1:500); donkey anti-sheep AF-488 (1:1k); donkey anti-rabbit AF-488 (1:1k); streptavidin AF-546 (1:500).

RNAscope

In situ hybridization in combination with IHC was performed on muscularis wholemount tissues using the RNAscope Multiplex Fluorescent V2 Assay kit with RNA-Protein Co-detection Ancillary Kit [ACD], according to the manufacturer's instructions with modifications as previously described (43). Probes used were *Cdh6* (#519541), *Cdh8* (#485461), *Nmu* (#446831), *Calcb* (#425511), and *eGFP* (#400281).

Confocal imaging

Images were acquired on a Leica SP8 confocal microscope using a 20x (NA 0.75) oil objective at 1024 x 1024 pixel resolution. Tiled images (24-30 tiles) of Z-stacks (2.5µm between planes) were acquired and stitched together using the Navigator mode within LASX (Leica). Imaged regions were located away from the mesenteric border.

Image analysis and quantification

Image analysis was performed using ImageJ/FIJI (NIH, Bethesda, MD), as described previously (42). HuC/D images (z-stack individual planes) were blurred and thresholded; then maximally projected and total neurons counted using the Analyze Particles function. *Cdh6*^{CreER}, *Ai32* expression and RNAscope *in situ* hybridization and *Cdh6*^{CreER}, *Ai14* expression were counted manually. Cell tracing was performed in Imaris using Filament Tracer (Bitplane, Oxford Instruments).

Electrophysiological recordings

Whole cell patch clamp electrophysiological recordings of Hb9:GFP+ neurons were performed according to Osorio & Delmas (14), with modifications for recording from the distal colon. The protocol is described in brief below.

Tissue dissection and preparation

Mice aged 8-10 weeks were culled by CO₂ and cervical dislocation. The colon was removed and flushed with ice cold oxygenated Krebs solution [118 mM NaCl, 4.8 mM KCl, 1 mM NaH₂PO₄, 25 mM NaHCO₃, 1.2 mM MgCl₂, 2.5 mM CaCl₂ and 11 mM glucose, supplemented with scopolamine (2 M) and nicardipine (6 µM)], then placed in a Sylgard-lined Petri dish with ice cold oxygenated Krebs solution for further dissection. Krebs solution was changed out for fresh oxygenated solution every 5 minutes. Under a dissection microscope, the distal colon was pinned and the mucosa peeled away using fine forceps, leaving a few millimeters of mucosa along the edges of the tissue for pinning stability. The muscularis was then flipped over and re-pinned, serosa side up, and the longitudinal muscle carefully peeled away. The tissue was transferred to a custom 3D-printed recording chamber lined with a thin layer of clear Sylgard, and re-pinned under light

tension, with the myenteric plexus facing up. The tissue was kept at 32°C and was continuously perfused with oxygenated Krebs solution. Hb9:GFP+ neurons were visually identified within a ganglion under epifluorescence illumination with a 455nm LED (Thorlabs, M455L2) and a 470 (excitation)/525 (emission) nm wavelength filter set. A local perfusion of protease XIV (0.2% in Krebs) (Sigma, P5417) was applied on top of the targeted cell to digest any muscle fiber residue. A 1-2M Ω pipet with a trimmed arm hair glued to the tip was used to brush and clean the surface of the ganglion. Further cleaning with 1mg/mL collagenase (Worthington, CLS-4) 4mg/mL dispase (Sigma, D4693) in Krebs solution was also performed to expose the GFP neuron for patching.

Patching and recording

Patch pipettes (4–6 M Ω) pulled from borosilicate glass were filled with internal solution containing in mM: 144 K-gluconate, 3 MgCl₂, 0.5 EGTA, 10 HEPES, pH 7.2 (285/295 mOsm) and 2% biocytin (Millipore Sigma, B4261-100MG). Patch clamp recordings were collected with a Multiclamp 700A (Molecular Devices) amplifier, a Digidata 1440 digitizer and pClamp10.7 (Molecular Devices). Recordings were sampled and filtered at 10kHz. Passive properties analysis was performed using pClamp10.7. Analysis of action potentials was performed using a custom MATLAB (MathWorks) software. All recordings were performed at 32°C. Membrane potentials were not corrected for liquid junction potential. Immediately after whole cell configuration, the cell was maintained at -70mV and a short voltage clamp membrane test protocol consisting of 20 times 600ms, 10mV depolarization steps was performed to assess cell health and recording conditions. Recordings were performed in Hb9:GFP+ colonic neurons with an access resistance less than 30M Ω (16.69 $\pm$ 2.63 M Ω). Next, the current clamp mode was used to measure resting membrane potential (RMP), input resistance (R_{in}) and APs stimulated. Membrane potential was not adjusted from resting potential, and cells were depolarized by 1 second current pulses in 10pA increments until APs were triggered (rheobase). Finally, if the seal was still stable, a voltage clamp steady-state inactivation of T-current protocol was performed as previously described (44 [link](#)). In brief, a sequence of depolarization from -90 to -45mV for 500ms quickly followed by a depolarization to -40mV for 200ms. Tissues were then fixed and immunostained according to *Wholemout preparations* and *Immunohistochemistry* sections above.

Mechanical recordings & optical stimulation

Optogenetic stimulation experiments were performed as previously described (29 [link](#)). A 2.5 mm stainless steel rod was inserted through the lumen of the colon and mounted in an organ bath (120 * 40 * 12 mm; L*W*H) located on a heated base. Krebs solution (35.5-36°C) superfused the bath (~5 mL/min). Smooth muscle force was recorded via 4 evenly spaced hooks in the colonic muscularis externa, each linked to an isometric force transducer (Grass FT03C) by suture thread. Initial base resting tension was set between 0.5 - 1.0 g. Preamplified signals (Biomedical Engineering, Flinders University) were digitized by a PowerLab 16/35 (ADInstruments, Bella Vista, NSW, Australia) and recorded using LabChart 7 software (ADInstruments) on iMac computer. Post hoc analysis of the mechanical recordings was done using LabChart 8 software on PC.

For optical stimulation during mechanical recordings *in vitro*, two LEDs (emitting 470 nm λ photons; C470DA2432, Cree Inc., NC, USA) were used, driven by a variable power supply. The area of light emission from each LED was 240 μ m x 320 μ m (0.0768 mm²). To characterise LED function, light power density across a range of currents was measured 5 mm from the LED using a standard photodiode power sensor (S120C; Thorlabs, NJ, USA) and a power meter (Thorlabs, PM100USB). The stimulator panel within LabChart software was used to set parameters and manually trigger LED pulse trains via the 10V analogue output of the PowerLab and an ILD1 opto-isolator.

Intraluminal pellet CMC recordings

To record proximal and distal colon CMCs separately (45 [45](#), 46 [46](#)), full length colon was bisected halfway between the caeco-colonic junction and terminal rectum, creating equal length proximal and distal colon preparations. Each preparation was suspended vertically on a stainless-steel holder inside a glass, water jacketed organ bath containing Krebs solution (Fig. 6J, K [47](#)). A 2.7 mm diameter synthetic pellet (polymethyl methacrylate, “Perspex”) was placed inside the gut lumen and linked by stainless steel rod to a force transducer (MLT0420, ADInstruments), allowing measurement of both anterograde and retrograde propulsive forces on the pellet. Signals were amplified by bridge amplifier (FE224, ADInstruments), digitized at 1kHz (PLCF1, ADInstruments) and recorded using LabChart 8 software.

ZD7288 (73777, Sigma-Aldrich) was dissolved in water as stock solution at 10 mM. Caesium Chloride (C4036, Sigma-Aldrich) was dissolved in water as stock solution at 200 mM. Tetrodotoxin citrate (T-550, Alomone Labs) was dissolved in water as stock solution at 3 mM. Control, ZD7288, CsCl and washout periods were at least 30 mins; TTX was applied for at least 10 minutes.

Statistical analysis

Statistical tests and graphical representation of data were performed using Prism 9 software (GraphPad). Statistical comparisons were performed using paired t tests (one-tailed, CMC intervals; two-tailed, peak amplitude, AUC, duration) and Welch’s t test (marker colocalizations). Asterisks indicate significant differences.

Data and materials availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Supplementary figures




Fig. S1.

Optogenetic stimulation control.

(A) Representative force traces of control $Cdh6^{CreER};ChR2-eYFP+$ colon ($n = 5$). Blue bars indicate timing of light stimulation. LEDs placed distal to distal hook.




Fig. S2.

Pharmacologic blockade of I_H with ZD7288 abolishes spontaneous CMCs.

(A, B) Representative force traces from tethered pellet in proximal half (A) or distal half (B) of colon. Addition of 10 μ M ZD7288 (first arrowhead), followed by washout in Krebs (second arrowhead), and addition of 1 μ M TTX (third arrowhead). Scale bars represent 2 g force (vertical bars) and 10 minutes (horizontal bars) for all traces.




Fig. S3.

Pharmacologic blockade of I_H with CsCl impairs generation of CMCs.

(A, B) Representative force traces from tethered pellet in proximal half (A) or distal half (B) of colon. Addition of 2 mM CsCl (first arrowhead), followed by washout in Krebs (second arrowhead), and addition of 1 μ M TTX (third arrowhead). Scale bars represent 2 g force (vertical bars) and 10 minutes (horizontal bars) for all traces.

Acknowledgements

We thank the members of the Kaltschmidt laboratory for experimental advice and discussions, Vanda Lennon (Mayo Clinic) for the HuC/D primary antibody and Beatriz G. Robinson for 3D printing of organ chambers for electrophysiology.

Additional information

Funding

National Institute of General Medical Sciences of the National Institutes of Health Award T32GM120007 (JGF)

National Institutes of Health Grant R01 EY030138 (XD)

National Health and Medical Research Council (NHMRC) project grant 1156416 (NJS)

Australian Research Council (ARC) Discovery Project grant DP220100070 (NJS)

NINDS 5R01NS34774 (JRH)

Wu Tsai Neurosciences Institute, Stanford University (JAK)

Department of Neurosurgery, Stanford University (JAK)

National Institutes of Health Grant R21 HD110950 (JAK)

The Firmenich Foundation (JAK)

The Carol and Eugene Ludwig Family Foundation (JAK)

Stanford ADRC Developmental Project Grant (National Institutes of Health Grant P30AG066515) (JAK)

Author contributions

Conceptualization: JGF, JAK

Resources: NJS, JRH, JAK

Methodology: JGF, GD, XD, TJH

Investigation: JGF, GD, LT, MAK, TJH

Visualization: JGF, GD, TJH

Funding acquisition: NJS, JRH, JAK

Supervision: NJS, JRH, JAK

Writing – original draft: JGF, GD

Writing – review & editing: JGF, NJS, JRH, JAK

References

1. Fung C., Vanden Berghe P. 2020) **Functional circuits and signal processing in the enteric nervous system** *Cell. Mol. Life Sci* **77**:4505–4522
2. Nurgali K., Stebbing M. J., Furness J. B. 2004) **Correlation of electrophysiological and morphological characteristics of enteric neurons in the mouse colon** *J. Comp. Neurol* **468**:112–124
3. Qu Z.-D., Thacker M., Castelucci P., Bagyánszki M., Epstein M. L., Furness J. B. 2008) **Immunohistochemical analysis of neuron types in the mouse small intestine** *Cell Tissue Res* **334**:147–161
4. Suzuki S. C., Inoue T., Kimura Y., Tanaka T., Takeichi M. 1997) **Neuronal Circuits Are Subdivided by Differential Expression of Type-II Classic Cadherins in Postnatal Mouse Brains** *Mol. Cell. Neurosci* **9**:433–447
5. Duan X., Krishnaswamy A., Laboulaye M. A., Liu J., Peng Y.-R., Yamagata M., Toma K., Sanes J. R. 2018) **Cadherin Combinations Recruit Dendrites of Distinct Retinal Neurons to a Shared Interneuronal Scaffold** *Neuron* **99**:1145–1154
6. Honjo M., Tanihara H., Suzuki S., Tanaka T., Honda Y., Takeichi M. 2000) **Differential Expression of Cadherin Adhesion Receptors in Neural Retina of the Postnatal Mouse** *Invest. Ophthalmol. Vis. Sci* **41**:546–551
7. Yamagata M., Duan X., Sanes J. R. 2018) **Cadherins Interact With Synaptic Organizers to Promote Synaptic Differentiation** *Front. Mol. Neurosci* **11**
8. Basu R., Taylor M. R., Williams M. E. 2015) **The classic cadherins in synaptic specificity** *Cell Adhes. Migr* **9**:193–201
9. May-Zhang A. A., Tycksen E., Southard-Smith A. N., Deal K. K., Benthall J. T., Buehler D. P., Adam M., Simmons A. J., Monaghan J. R., Matlock B. K., Flaherty D. K., Potter S. S., Lau K. S., Southard-Smith E. M. 2021) **Combinatorial Transcriptional Profiling of Mouse and Human Enteric Neurons Identifies Shared and Disparate Subtypes In Situ** *Gastroenterology* **160**:755–770
10. Drokhlyansky E., Smillie C. S., Wittenberghe N. V., Ericsson M., Griffin G. K., Eraslan G., Dionne D., Cuoco M. S., Goder-Reiser M. N., Sharova T., Kuksenko O., Aguirre A. J., Boland G. M., Graham D., Rozenblatt-Rosen O., Xavier R. J., Regev A. 2020) **The Human and Mouse Enteric Nervous System at Single-Cell Resolution** *Cell* **182**:1606–1622
11. Morarach K., Mikhailova A., Knoflach V., Memic F., Kumar R., Li W., Ernfors P., Marklund U. 2021) **Diversification of molecularly defined myenteric neuron classes revealed by single-cell RNA sequencing** *Nat. Neurosci* **24**:34–46
12. Wahl-Schott C., Biel M. 2008) **HCN channels: Structure, cellular regulation and physiological function** *Cell. Mol. Life Sci* **66**:470
13. Laboulaye M. A., Duan X., Qiao M., Whitney I. E., Sanes J. R. 2018) **Mapping Transgene Insertion Sites Reveals Complex Interactions Between Mouse Transgenes and**

14. Osorio N., Delmas P. 2011) **Patch clamp recording from enteric neurons in situ** *Nat. Protoc* **6**:15–27
15. Huguenard J. R. 1996) **Low-threshold calcium currents in central nervous system neurons** *Annu. Rev. Physiol* **58**:329–348
16. Xiao J., Nguyen T. V., Ngui K., Strijbos P. J. L. M., Selmer I.-S., Neylon C. B., Furness J. B. 2004) **Molecular and functional analysis of hyperpolarisation-activated nucleotide-gated (HCN) channels in the enteric nervous system** *Neuroscience* **129**:603–614
17. Nurgali K., Nguyen T. V., Matsuyama H., Thacker M., Robbins H. L., Furness J. B. 2007) **Phenotypic changes of morphologically identified guinea-pig myenteric neurons following intestinal inflammation** *J. Physiol* **583**:593–609
18. Mao Y., Wang B., Kunze W. 2006) **Characterization of Myenteric Sensory Neurons in the Mouse Small Intestine** *J. Neurophysiol* **96**:998–1010
19. Furness J. B., Robbins H. L., Xiao J., Stebbing M. J., Nurgali K. 2004) **Projections and chemistry of Dogiel type II neurons in the mouse colon** *Cell Tissue Res* **317**:1–12
20. Kay J. N., la Huerta I. D., Kim I.-J., Zhang Y., Yamagata M., Chu M. W., Meister M., Sanes J. R. 2011) **Retinal Ganglion Cells with Distinct Directional Preferences Differ in Molecular Identity, Structure, and Central Projections** *J. Neurosci* **31**:7753–7762
21. Madisen L., Zwingman T. A., Sunkin S. M., Oh S. W., Zariwala H. A., Gu H., Ng L. L., Palmiter R. D., Hawrylycz M. J., Jones A. R., Lein E. S., Zeng H. 2010) **A robust and high-throughput Cre reporting and characterization system for the whole mouse brain** *Nat. Neurosci* **13**:133–140
22. McCormick D. A., Pape H. C. 1990) **Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones** *J. Physiol* **431**:291–318
23. Spencer N. J., Costa M., Hibberd T. J., Wood J. D. 2021) **Advances in colonic motor complexes in mice** *Am. J. Physiol.-Gastrointest. Liver Physiol* **320**:G12–G29
24. Hibberd T. J., Travis L., Wiklendt L., Costa M., Brookes S. J. H., Hu H., Keating D. J., Spencer N. J. 2018) **Synaptic activation of putative sensory neurons by hexamethonium-sensitive nerve pathways in mouse colon** *Am. J. Physiol. Gastrointest. Liver Physiol* **314**:G53–G64
25. Madisen L., Mao T., Koch H., Zhuo J., Berenyi A., Fujisawa S., Hsu Y.-W. A., Garcia A. J., Gu X., Zanella S., Kidney J., Gu H., Mao Y., Hooks B. M., Boyden E. S., Buzsáki G., Ramirez J. M., Jones A. R., Svoboda K., Han X., Turner E. E., Zeng H. 2012) **A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing** *Nat. Neurosci* **15**:793–802
26. Harris N. C., Constanti A. 1995) **Mechanism of block by ZD 7288 of the hyperpolarization-activated inward rectifying current in guinea pig substantia nigra neurons in vitro** *J. Neurophysiol* **74**:2366–2378
27. Galligan J. J., Tatsumi H., Shen K. Z., Surprenant A., North R. A. 1990) **Cation current activated by hyperpolarization (IH) in guinea pig enteric neurons** *Am. J. Physiol.-Gastrointest. Liver Physiol* **259**:G966–G972

28. Koh S. D., Drumm B. T., Lu H., Kim H. J., Ryoo S.-B., Kim H.-U., Lee J. Y., Rhee P.-L., Wang Q., Gould T. W., Heredia D., Perrino B. A., Hwang S. J., Ward S. M., Sanders K. M. 2022) **Propulsive colonic contractions are mediated by inhibition-driven poststimulus responses that originate in interstitial cells of Cajal** *Proc. Natl. Acad. Sci* **119**:e2123020119
29. Hibberd T. J., Feng J., Luo J., Yang P., Samineni V. K., Gereau R. W., Kelley N., Hu H., Spencer N. J. 2018) **Optogenetic Induction of Colonic Motility in Mice** *Gastroenterology* **155**:514–528
30. Efimov A., Hibberd T., Wang Y., Wu M., Zhang K., Ting K., Madhvapathy S., Lee M., Kim J., Kang J., Riahi M., Zhang H., Travis L., Govier E. J., Yang L., Kelly N., Huang Y., Vázquez-Guardado A., Spencer N. J., Rogers J. A. 2024) **Remote optogenetic control of the enteric nervous system and brain-gut axis in freely-behaving mice enabled by a wireless, battery-free optoelectronic device** *Biosens. Bioelectron* **116298**
31. Li Z., Boesmans W., Kazwiny Y., Hao M. M., Vanden Berghe P. 2022) **Simultaneous whole-cell patch-clamp and calcium imaging on myenteric neurons** *Am. J. Physiol.-Gastrointest. Liver Physiol* **323**:G341–G347
32. Pape H. C., McCormick D. A. 1989) **Noradrenaline and serotonin selectively modulate thalamic burst firing by enhancing a hyperpolarization-activated cation current** *Nature* **340**:715–718
33. Benarroch E. E. 2013) **HCN channels** *Neurology* **80**:304–310
34. Fisher D. W., Luu P., Agarwal N., Kurz J. E., Chetkovich D. M. 2018) **Loss of HCN2 leads to delayed gastrointestinal motility and reduced energy intake in mice** *PloS One* **13**:e0193012
35. Shimoyama Y., Tsujimoto G., Kitajima M., Natori M. 2000) **Identification of three human type-II classic cadherins and frequent heterophilic interactions between different subclasses of type-II classic cadherins** *Biochem. J* **349**:159
36. Hirst G. D. S., Holman M. E., Spence I. 1974) **Two types of neurones in the myenteric plexus of duodenum in the guinea-pig** *J. Physiol* **236**:303–326
37. Kunze W. A. A., Furness J. B., Bornstein J. C. 1993) **Simultaneous intracellular recordings from enteric neurons reveal that myenteric ah neurons transmit via slow excitatory postsynaptic potentials** *Neuroscience* **55**:685–694
38. Pompolo S., Furness J. B. 1988) **Ultrastructure and synaptic relationships of calbindin-reactive, Dogiel type II neurons, in myenteric ganglia of guinea-pig small intestine** *J. Neurocytol* **17**:771–782
39. Wood J. D., Johnson L. R., Ghishan F. K., Kaunitz J. D., Merchant J. L., Said H. M., Wood J. D. 2012) **Chapter 21 - Cellular Neurophysiology of Enteric Neurons** *Physiology of the Gastrointestinal Tract* Boston: Academic Press :629–669 <https://www.sciencedirect.com/science/article/pii/B978012382026600021X>
40. Barth B. B., Spencer N. J., Grill W. M., Spencer N. J., Costa M., Brierley S. M. 2022) **Activation of ENS Circuits in Mouse Colon: Coordination in the Mouse Colonic Motor Complex as a Robust, Distributed Control System** *The Enteric Nervous System II* Cham: Springer International Publishing :113–123
41. Gomez-Frittelli J., Hamnett R., Kaltschmidt J. A. 2023) **Comparison of wholemount dissection methods for neuronal subtype marker expression in the mouse myenteric plexus** *bioRxiv*

42. Hamnett R., Dershowitz L. B., Sampathkumar V., Wang Z., Gomez-Frittelli J., De Andrade V., Kasthuri N., Druckmann S., Kaltschmidt J. A. 2022) **Regional cytoarchitecture of the adult and developing mouse enteric nervous system** *Curr. Biol* **32**:4483–4492
43. Guyer R. A., Stavely R., Robertson K., Bhawe S., Mueller J. L., Picard N. M., Hotta R., Kaltschmidt J. A., Goldstein A. M. 2023) **Single-cell multiome sequencing clarifies enteric glial diversity and identifies an intraganglionic population poised for neurogenesis** *Cell Rep* **42**:112194
44. Huguenard J. R., Prince D. A. 1992) **A novel T-type current underlies prolonged Ca(2+)-dependent burst firing in GABAergic neurons of rat thalamic reticular nucleus** *J. Neurosci* **12**:3804–3817
45. Ishizawa M. 1984) **Propulsive Activity of the Isolated Segmental Colon of Guinea-Pig** *Jpn. J. Smooth Muscle Res* **20**:401–406
46. Hibberd T. J., Costa M., Smolilo D. J., Keightley L. J., Brookes S. J., Dinning P. G., Spencer N. J. 2022) **Mechanisms underlying initiation of propulsion in guinea pig distal colon** *Am. J. Physiol.-Gastrointest. Liver Physiol* **323**:G71–G87

Author information

Julieta Gomez-Frittelli

Department of Chemical Engineering, Stanford University, Stanford, United States, Wu Tsai Neurosciences Institute, Stanford University, Stanford, United States

Gabrielle Devienne

Wu Tsai Neurosciences Institute, Stanford University, Stanford, United States, Department of Neurology & Neurological Sciences, Stanford University, Stanford, United States

Lee Travis

College of Medicine and Public Health, Flinders Health & Medical Research Institute, Flinders University, Adelaide, Australia

Melinda A Kyloh

College of Medicine and Public Health, Flinders Health & Medical Research Institute, Flinders University, Adelaide, Australia

Xin Duan

Department of Ophthalmology, School of Medicine, University of California San Francisco, San Francisco, United States

Tim J Hibberd

College of Medicine and Public Health, Flinders Health & Medical Research Institute, Flinders University, Adelaide, Australia

Nick J Spencer

College of Medicine and Public Health, Flinders Health & Medical Research Institute, Flinders University, Adelaide, Australia

ORCID iD: [0000-0003-2190-5303](https://orcid.org/0000-0003-2190-5303)

John R Huguenard

Wu Tsai Neurosciences Institute, Stanford University, Stanford, United States, Department of Neurology & Neurological Sciences, Stanford University, Stanford, United States

ORCID iD: [0000-0002-6950-1191](https://orcid.org/0000-0002-6950-1191)

Julia A Kaltschmidt

Wu Tsai Neurosciences Institute, Stanford University, Stanford, United States, Department of Neurosurgery, Stanford University School of Medicine, Stanford, United States

ORCID iD: [0000-0002-2893-1793](https://orcid.org/0000-0002-2893-1793)

For correspondence: jukalts@stanford.edu

Editors

Reviewing Editor

Albert Cardona

University of Cambridge, Cambridge, United Kingdom

Senior Editor

Albert Cardona

University of Cambridge, Cambridge, United Kingdom

Reviewer #1 (Public review):

Summary:

In their manuscript, Gomez-Frittelli and colleagues characterize the expression of cadherin6 (and -8) in colonic IPANs of mice. Moreover, they found that these *cdh6*-expressing IPANs are capable of initiating colonic motor complexes in the distal colon, but not proximal and midcolon. They support their claim by morphological, electrophysiological and optogenetic, and pharmacological experiments.

Strengths:

The work is very impressive and involves several genetic models and state-of-the-art physiological setups including respective controls. It is a very well-written manuscript that truly contributes to our understanding of GI-motility and its anatomical and physiological basis. The authors were able to convincingly answer their research questions with a wide range of methods without overselling their results.

Weaknesses:

The authors put quite some emphasis on stating that *cdh6* is a synaptic protein (in the title and throughout the text), which interacts in a homophilic fashion. They deduct that *cdh6* might be involved in IPAN-IPAN synapses (line 247ff.). However, *Cdh6* does not only interact in synapses and is expressed by non-neuronal cells as well (see e.g., expression in the proximal tubuli of the kidney). Moreover, *cdh6* does not only build homodimers, but also heterodimers with *Chd9* as well as *Cdh7*, -10, and -14 (see e.g., Shimoyama et al. 2000, DOI: 10.1042/0264-6021:3490159). It would therefore be interesting to assess the expression pattern of *cdh6*-proteins using immunostainings in combination with synaptic markers to substantiate the authors' claim or at least add the possibility of cell-cell-interactions other than synapses to the discussion. Additionally, an immunostaining of *cdh6* would confirm if

the expression of tdTomato in smooth muscle cells of the *cdh6-creERT* model is valid or a leaky expression (false positive).

Comments on revisions:

The authors have updated their manuscript and have provided insights and discussions to my remarks.

<https://doi.org/10.7554/eLife.101043.2.sa2>

Reviewer #2 (Public review):

Summary:

Intrinsic primary afferent neurons are an interesting population of enteric neurons that transduce stimuli from the mucosa, initiate reflexive neurocircuitry involved in motor and secretory functions, and modulate gut immune responses. The morphology, neurochemical coding, and electrophysiological properties of these cells have been relatively well described in a long literature dating back to the late 1800's but questions remain regarding their roles in enteric neurocircuitry, potential subsets with unique functions, and contributions to disease. Here, the authors provide RNAscope, immunolabeling, electrophysiological, and organ function data characterizing IPANs in mice and suggest that *Cdh6* is an additional marker of these cells.

Strengths:

This paper would likely be of interest to the enteric neuroscience community and increases information regarding the properties of IPANs in mice. These data are useful and suggest that prior data from studies of IPANs in other species are likely translatable to mice.

Weaknesses:

Major weaknesses:

(1) The novelty of this study is relatively limited. The main point of novelty suggests an additional marker of IPANs (*Cdh6*) that would add to the known list of markers for these cells. How useful this would be is unclear. Other main findings basically confirm that IPANs in mice display the same classical characteristics that have been known for many years from studies in guinea pigs, rats, mice and humans.

(2) Critical controls are needed to support the optogenetic experiments. Control experiments are needed to show that ChR2 expression 1) does not change the baseline properties of the neurons, 2) that stimulation with the chosen intensity of light elicits physiologically relevant responses in those neurons, and 3) that stimulation via ChR2 elicits comparable responses in IPANs in the different gut regions focused on here. These essential controls remain absent in the study and limit confidence in the data derived from this model.

(3) The motor effects observed in optogenetic experiments are difficult to understand in the absence of good controls for optogenetic control of the proposed neuron population (discussed above). It remains unclear how stimulating IPANs in the distal colon would generate retrograde CMCs while stimulating IPANs in the proximal colon did nothing. Key controls confirming that the optogenetic stimulus was adequate, specific, and relevant are needed. In addition, better characterization of the *Cdh6*⁺ population of cells in both regions would be needed to understand the mechanisms underlying these effects.

(4) From the data shown, it is clear that expression driven by the *Cdh6CreERT2* driver is not confined to IPANs. There is obviously expression of GFP and ChR2 in smooth muscle cells. This is a major limitation for the physiological experiments that attempt to use this model to

specifically stimulate IPANs and assess changes in gut motor function. Better characterization of this model is needed and control experiments are necessary to assess whether functional ChR2 is expressed in cells beyond the proposed subtype of enteric IPANs.

(5) Some of the main conclusions of this study are overstated and claims of priority are made that are not true. For example, the authors state on lines 27-28 of the abstract that their findings provide the "first demonstration of selective activation of a single neurochemical and functional class of enteric neurons". This is certainly not true since Gould et al (AJP-GIL 2019) expressed ChR2 in nitrergic enteric neurons and showed that activating those cells disrupted CMC activity. In fact, prior work by the authors themselves (Hibberd et al Gastro 2018) showed that activating calretinin neurons with ChR2 evoked motor responses. Work by other groups has used chemogenetics and optogenetics to show effects of activating multiple other classes of neurons in the gut.

(6) The electrophysiological characterization of mouse IPANs is useful but is limited to a small subset of Cdh6+ neurons in the distal colon myenteric plexus. Therefore, it remains unclear how well the properties reported here might reflect those of other Cdh6+ IPANs in the same or different regions. Similarly, blocking IH with ZD7288 affects all IPANs and does not add specific information regarding the role of the proposed Cdh6+ subtype.

(7) The submucosal plexus (SMP) also contains enteric IPANs and these were not included in the analysis of Cdh6 expression. Whether or not the proposed IPAN marker Cdh6 would be useful for identifying or targeting those cells remains unclear.

[Editor's note: The Reviewing Editor considers that further controls requested from the reviewers have largely been provided already in prior publications by other groups, as they concern specifically tools published years ago but in a different tissue context. Hence the methodology used to deliver the results reported here fall within the standard practices in the field. The comprehensive, multi-technique approach to the results is compelling in and of itself, and ought to suffice, rendering this work reproducible and therefore a basis for further research.]

<https://doi.org/10.7554/eLife.101043.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

Summary:

In their manuscript, Gomez-Frittelli and colleagues characterize the expression of cadherin6 (and -8) in colonic IPANs of mice. Moreover, they found that these cdh6-expressing IPANs are capable of initiating colonic motor complexes in the distal colon, but not proximal and midcolon. They support their claim by morphological, electrophysiological, optogenetic, and pharmacological experiments.

Strengths:

The work is very impressive and involves several genetic models and state-of-the-art physiological setups including respective controls. It is a very well-written manuscript that truly contributes to our understanding of GI-motility and its anatomical and physiological basis. The authors were able to convincingly answer their research questions with a wide range of methods without overselling their results.

We greatly appreciate the reviewer's time, careful reading and support of our study.

Weaknesses:

*The authors put quite some emphasis on stating that *cdh6* is a synaptic protein (in the title and throughout the text), which interacts in a homophilic fashion. They deduct that *cdh6* might be involved in IPAN-IPAN synapses (line 247ff.). However, *Cdh6* does not only interact in synapses and is expressed by non-neuronal cells as well (see e.g., expression in the proximal tubuli of the kidney). Moreover, *cdh6* does not only build homodimers, but also heterodimers with *Chd9* as well as *Cdh7*, -10, and -14 (see e.g., Shimoyama et al. 2000, DOI: 10.1042/02646021:3490159). It would therefore be interesting to assess the expression pattern of *cdh6* proteins using immunostainings in combination with synaptic markers to substantiate the authors' claim or at least add the possibility of cell-cell interactions other than synapses to the discussion. Additionally, an immunostaining of *cdh6* would confirm if the expression of *tdTomato* in smooth muscle cells of the *cdh6-creERT* model is valid or a leaky expression (false positive).*

We agree with the reviewer that *Cdh6* could be mediating some other cell-cell interaction besides synapses between IPANs, and we noted it in the discussion. *Cdh6* primarily forms homodimers but, as the reviewer points out, has been known to also form heterodimers with some other cadherins. We performed RNAscope in the colonic myenteric plexus with *Cdh7* and found no expression (data not shown). *Cdh10* is suggested to have very low expression (Drokhlyansky et al., 2020), possibly in putative secretomotor vasodilator neurons, and *Cdh14* has not been assayed in any RNAseq screens. We attempted to visualize *Cdh6* protein via antibody staining (Duan et al., 2018) but our efforts did not result in sufficient signal or resolution to identify synapses in the ENS, which remain broadly challenging to assay. Similarly, immunostaining with *Cdh6* antibody was unable to confirm *Cdh6* protein in *tdT*-expressing muscle cells, or by RNAscope. We have addressed these caveats in the discussion section.

(1) E. Drokhlyansky, C. S. Smillie, N. V. Wittenberghe, M. Ericsson, G. K. Griffin, G. Eraslan, D. Dionne, M. S. Cuoco, M. N. Goder-Reiser, T. Sharova, O. Kuksenko, A. J. Aguirre, G. M. Boland, D. Graham, O. Rozenblatt-Rosen, R. J. Xavier, A. Regev, The Human and Mouse Enteric Nervous System at Single-Cell Resolution. *Cell* 182, 1606-1622.e23 (2020).

(2) X. Duan, A. Krishnaswamy, M. A. Laboulaye, J. Liu, Y.-R. Peng, M. Yamagata, K. Toma, J. R. Sanes, Cadherin Combinations Recruit Dendrites of Distinct Retinal Neurons to a Shared Interneuronal Scaffold. *Neuron* 99, 1145-1154.e6 (2018).

Reviewer #2 (Public review):

Summary:

*Intrinsic primary afferent neurons are an interesting population of enteric neurons that transduce stimuli from the mucosa, initiate reflexive neurocircuitry involved in motor and secretory functions, and modulate gut immune responses. The morphology, neurochemical coding, and electrophysiological properties of these cells have been relatively well described in a long literature dating back to the late 1800's but questions remain regarding their roles in enteric neurocircuitry, potential subsets with unique functions, and contributions to disease. Here, the authors provide RNAscope, immunolabeling, electrophysiological, and organ function data characterizing IPANs in mice and suggest that *Cdh6* is an additional marker of these cells.*

Strengths:

This paper would likely be of interest to a focused enteric neuroscience audience and increase information regarding the properties of IPANs in mice. These data are useful and suggest that prior data from studies of IPANs in other species are likely translatable to mice.

We appreciate the reviewer's support of our study and insightful critiques for its improvement.

Weaknesses:

The advance presented here beyond what is already known is minimal. Some of the core conclusions are overstated and there are multiple other major issues that limit enthusiasm. Key control experiments are lacking and data do not specifically address the properties of the proposed Cdh6+ population.

Major weaknesses:

(1) The novelty of this study is relatively low. The main point of novelty suggests an additional marker of IPANs (Cdh6) that would add to the known list of markers for these cells. How useful this would be is unclear. Other main findings basically confirm that IPANs in mice display the same classical characteristics that have been known for many years from studies in guinea pigs, rats, mice and humans.

We appreciate the already existing markers for IPANs in the ENS and the existing literature characterizing these neurons. The primary intent of this study was to use these well-established characteristics of IPANs in both mice and other species to characterize Cdh6-expressing neurons in the mouse myenteric plexus and confirm their classification as IPANs.

(2) Some of the main conclusions of this study are overstated and claims of priority are made that are not true. For example, the authors state in lines 27-28 of the abstract that their findings provide the "first demonstration of selective activation of a single neurochemical and functional class of enteric neurons". This is certainly not true since Gould et al (AJP-GIL 2019) expressed ChR2 in nitrergic enteric neurons and showed that activating those cells disrupted CMC activity. In fact, prior work by the authors themselves (Hibberd et al., Gastro 2018) showed that activating calretinin neurons with ChR2 evoked motor responses. Work by other groups has used chemogenetics and optogenetics to show the effects of activating multiple other classes of neurons in the gut.

We thank the reviewer for bringing up this important point and apologize if our wording was not clear. Whilst single neurochemical classes of enteric neurons have been manipulated to alter gut functions, all such instances to date do not represent manipulation of a single functional class of enteric neurons. In the given examples, multiple functional classes are activated utilizing the same neurotransmitter, as NOS and calretinin are each expressed to varying degrees across putative motor neurons, interneurons and IPANs. In contrast, Chd6 is restricted to IPANs and therefore this study is the first optogenetic investigation of enteric neurons from a single putative functional class. Our abstract and discussion emphasizes this point and differentiates this study from those previous.

(3) Critical controls are needed to support the optogenetic experiments. Control experiments are needed to show that ChR2 expression a) does not change the baseline properties of the neurons, b) that stimulation with the chosen intensity of light elicits physiologically relevant responses in those neurons, and c) that stimulation via ChR2 elicits comparable responses in IPANs in the different gut regions focused on here.

We completely agree controls are essential. However, our paper is not the first to express Chr2 in enteric neurons. Authors of our paper have shown in Hibberd *et al.* 2018 that expression of Chr2 in a heterogeneous population of myenteric neurons did not change network properties of the myenteric plexus. This was demonstrated in the lack of change in control CMC characteristics in mice expressing Chr2 under basal conditions (without blue light exposure). Regarding question (b), that it should be shown that stimulation with the chosen intensity of light elicits physiologically relevant responses in those neurons. We show the restricted expression of Chr2 in IPANs and that motor responses (to blue light) are blocked by selective nerve conduction blockade.

Regarding question (c), that our study should demonstrate that stimulation via Chr2 elicits comparable responses in IPANs in the different gut regions. We would not expect each region of the gut to behave comparably. This is because the different gut regions (i.e. proximal, mid, distal) are very different anatomically, as is anatomy of the myenteric plexus and myenteric ganglia between each region, including the density of IPANs within each ganglia, in addition to the presence of different patterns of electrical and mechanical activity [Spencer *et al.*, 2020]. Hence, it is difficult to expect that between regions stimulation of Chr2 should induce similar physiological responses. The motor output we record in our study (CMCs) is a unified motor program that involves the temporal coordination of hundreds of thousands of enteric neurons and a complex neural circuit that we have previously characterized [Spencer *et al.*, 2018]. But, never has any study until now been able to selectively stimulate a single functional class of enteric neurons (with light) to avoid indiscriminate activation of other classes of neurons.

(1) T. J. Hibberd, J. Feng, J. Luo, P. Yang, V. K. Samineni, R. W. Gereau, N. Kelley, H. Hu, N. J. Spencer, Optogenetic Induction of Colonic Motility in Mice. *Gastroenterology* 155, 514-528.e6 (2018).

(2) N. J. Spencer, L. Travis, L. Wiklendt, T. J. Hibberd, M. Costa, P. Dinning, H. Hu, Diversity of neurogenic smooth muscle electrical rhythmicity in mouse proximal colon. *American Journal of Physiology-Gastrointestinal and Liver Physiology* 318, G244–G253 (2020).

(3) N. J. Spencer, T. J. Hibberd, L. Travis, L. Wiklendt, M. Costa, H. Hu, S. J. Brookes, D. A. Wattchow, P. G. Dinning, D. J. Keating, J. Sorensen, Identification of a Rhythmic Firing Pattern in the Enteric Nervous System That Generates Rhythmic Electrical Activity in Smooth Muscle. *The Journal of Neuroscience* 38, 5507–5522 (2018).

(4) The electrophysiological characterization of mouse IPANs is useful but this is a basic characterization of any IPAN and really says nothing specifically about Cdh6+ neurons. The electrophysiological characterization was also only done in a small fraction of colonic IPANs, and it is not clear if these represent cell properties in the distal colon or proximal colon, and whether these properties might be extrapolated to IPANs in the different regions. Similarly, blocking IH with ZD7288 affects all IPANs and does not add specific information regarding the role of the proposed Cdh6+ subtype.

Our electrophysiological characterization was guided to be within a subset of Cdh6+ neurons by Hb9:GFP expression. As in the prior comment (1) above, we used these experiments to confirm classification of Cdh6+ (Hb9:GFP+) neurons in the distal colon as IPANs. We have clarified in the results and methods that these experiments were performed in the distal colon and agree that we cannot extrapolate that these properties are also representative of IPANs in the proximal colon. We apologize that this was confusing. Finally, we agree with the reviewer that ZD7288 affects all IPANs in the ENS and have clarified this in the text.

(5) Why SMP IPANs were not included in the analysis of Cdh6 expression is a little puzzling. IPANs are present in the SMP of the small intestine and colon, and it would be

useful to know if this proposed marker is also present in these cells.

We agree with the reviewer. In addition to characterizing Cdh6 in the myenteric plexus, it would be interesting to query if sensory neurons located within the SMP also express Cdh6. Our preliminary data (n=2) show ~6-12% tdT/Hu neurons in Cdh6-tdT ileum and colon (data not shown). We have added a sentence to the discussion.

(6) The emphasis on IH being a rhythmicity indicator seems a bit premature. There is no evidence to suggest that IH and IT are rhythm-generating currents in the ENS.

Regarding the statement there is no evidence to suggest that IH and IT are rhythm-generating currents in the ENS. We agree with the reviewer that evidence of rhythm generation by IH and IT in the ENS has not been explicitly confirmed. We are confident the reviewer agrees that an absence of evidence is not evidence of absence, although the presence of IH has been well described in enteric neurons. We have modified the text in the results to indicate more clearly that IH and IT are known to participate in rhythm generation in thalamocortical circuits, though their roles in the ENS remain unknown. Our discussion of the potential role of IH or IT in rhythm generation or oscillatory firing of the ENS is constrained to speculation in the discussion section of the text.

(7) As the authors point out in the introduction and discuss later on, Type II Cadherins such as Cdh6 bind homophillically to the same cadherin at both pre- and post-synapse. The apparent enrichment of Cdh6 in IPANs would suggest extensive expression in synaptic terminals that would also suggest extensive IPAN-IPAN connections unless other subtypes of neurons express this protein. Such synaptic connections are not typical of IPANs and raise the question of whether or not IPANs actually express the functional protein and if so, what might be its role. Not having this information limits the usefulness of this as a proposed marker.

We agree with the reviewer that the proposed IPAN-IPAN connection is novel although it has been proposed before (Kunze et al., 1993). As detailed in our response to Reviewer #1, we attempted to confirm Cdh6 protein expression, but were unsuccessful, due to insufficient signal and resolution. We therefore discuss potential IPAN interconnectivity in the discussion, in the context of contrasting literature.

(1) W. A. A. Kunze, J. B. Furness, J. C. Bornstein, Simultaneous intracellular recordings from enteric neurons reveal that myenteric ah neurons transmit via slow excitatory postsynaptic potentials. Neuroscience 55, 685–694 (1993).

(8) Experiments shown in Figures 6J and K use a tethered pellet to drive motor responses. By definition, these are not CMCs as stated by the authors.

The reviewer makes a valid criticism as to the terminology, since tethered pellet experiments do not record propagation. We believe the periodic bouts of propulsive force on the pellet is triggered by the same activity underlying the CMC. In our experience, these activities have similar periodicity, force and identical pharmacological properties. Consistent with this, we also tested full colons (n = 2) set up for typical CMC recordings by multiple force transducers, finding that CMCs were abolished by ZD7288, similar to fixed pellet recordings (data not shown).

(9) The data from the optogenetic experiments are difficult to understand. How would stimulating IPANs in the distal colon generate retrograde CMCs and stimulating IPANs in the proximal colon do nothing? Additional characterization of the Cdh6+ population of cells is needed to understand the mechanisms underlying these effects.

We agree that the different optogenetic responses in the proximal and distal colon are challenging to interpret, but perhaps not surprising in the wider context. It is not only possible that the different optogenetic responses in this study reflect regional differences in the Chd6⁺ neuronal populations, but also differences in neural circuits within these gut regions. A study some time ago by the authors showed that electrical stimulation of the proximal mouse colon was unable to evoke a retrograde (aborally) propagating CMC (Spencer, Bywater, 2002), but stimulation of the distal colon was readily able to. We concluded that at the oral lesion site there is a preferential bias of descending inhibitory nerve projections, since the ascending excitatory pathways have been cut off. In contrast, stimulation of the distal colon was readily able to activate an ascending excitatory neural pathway, and hence induce the complex CMC circuits required to generate an orally propagating CMC. Indeed, other recent studies have added to a growing body of evidence for significant differences in the behaviors and neural circuits of the two regions (Li et al., 2019, Costa et al., 2021a, Costa et al., 2021b, Nestor-Kalinoski et al., 2022). We have expanded this discussion.

- (1) N. J. Spencer, R. A. Bywater, Enteric nerve stimulation evokes a premature colonic migrating motor complex in mouse. *Neurogastroenterology & Motility* 14, 657–665 (2002).
- (2) Li Z, Hao MM, Van den Haute C, Baekelandt V, Boesmans W, Vanden Berghe P, Regional complexity in enteric neuron wiring reflects diversity of motility patterns in the mouse large intestine. *Elife* 8:e42914 (2019).
- (3) Costa M, Keightley LJ, Hibberd TJ, Wiklendt L, Dinning PG, Brookes SJ, Spencer NJ, Motor patterns in the proximal and distal mouse colon which underlie formation and propulsion of feces. *Neurogastroenterology & Motility* e14098 (2021a).
- (4) Costa M, Keightley LJ, Hibberd TJ, Wiklendt L, Smolilo DJ, Dinning PG, Brookes SJ, Spencer NJ, Characterization of alternating neurogenic motor patterns in mouse colon. *Neurogastroenterology & Motility* 33:e14047 (2021b).
- (5) Nestor-Kalinoski A, Smith-Edwards KM, Meerschaert K, Margiotta JF, Rajwa B, Davis BM, Howard MJ, Unique Neural Circuit Connectivity of Mouse Proximal, Middle, and Distal Colon Defines Regional Colonic Motor Patterns. *Cellular and Molecular Gastroenterology and Hepatology* 13:309-337.e303 (2022).

Recommendations for the Authors:

Reviewer #1 (Recommendations for the authors):

As mentioned above, immunolocalization of cdh6 would be helpful to substantiate the claims regarding IPAN-IPAN synapses.

As mentioned in our response to both reviewers' public reviews, we attempted to visualize Cdh6 protein via antibody staining (Duan et al., 2018), but our efforts did not result in sufficient signal or resolution to identify Cdh6⁺ synapses.

- (1) X. Duan, A. Krishnaswamy, M. A. Laboulaye, J. Liu, Y.-R. Peng, M. Yamagata, K. Toma, J. R. Sanes, Cadherin Combinations Recruit Dendrites of Distinct Retinal Neurons to a Shared Interneuronal Scaffold. *Neuron* 99, 1145-1154.e6 (2018).

Reviewer #2 (Recommendations for the authors):

(1) The authors repeatedly refer to IPANs as "sensory" neurons (e.g. in title, abstract, and introduction) but there is some debate regarding whether these cells are truly "sensory" because the information they convey never reaches sensory perception. This is why they

have classically been referred to as intrinsic primary afferent (IPAN) neurons. It would be more appropriate to stick with this terminology unless the authors have compelling data showing that information detected by IPANs reaches the sensory cortex.

We thank the reviewer for their comment, but respectfully disagree. The term “sensory neuron” is well established in the ENS. The first definitive proof that “sensory neurons” exist in the ENS was published in Kunze et al., 1995. We note that this paper did not use the word “IPAN” but used the term “sensory neuron”. Furthermore, mechanosensory neurons were published in Spencer and Smith (2004).

Regarding the reviewer’s comment that the authors would need compelling data showing that information detected by IPANs reaches the sensory cortex before the term “sensory neuron” should be valid, it is important to note that many sensory neurons do not provide direct information to the cortex.

(1) W. A. A. Kunze, J. C. Bornstein, J. B. Furness, Identification of sensory nerve cells in a peripheral organ (the intestine) of a mammal. *Neuroscience* 66, 1–4 (1995).

(2) N. J. Spencer, T. K. Smith, Mechanosensory S-neurons rather than AH-neurons appear to generate a rhythmic motor pattern in guinea-pig distal colon. *The Journal of Physiology* 558, 577–596 (2004).

(2) Important information regarding the gut region shown and other details are absent from many figure legends.

We apologize for this omission. We have updated the figure legends to include information on gut regions.

<https://doi.org/10.7554/eLife.101043.2.sa0>