

The penetration ring is a novel infection structure formed by the penetration peg for invading plant cell membrane in rice blast fungus

Reviewed Preprint

v2 • May 15, 2025

Revised by authors

Reviewed Preprint

v1 • September 20, 2024

Wenqin Fang, Xiaoyu Zai, Jia Chen, Yakubu Saddeeq Abubakar, Qiuqiu Wu, Zhenyu Fang, Xiuwei Huang, Xiang Gan, Daniel J Ebbale, Zonghua Wang , Wenhui Zheng 

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China • Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China • Fuzhou Institute of Oceanography, Minjiang University, Fuzhou, China • Department of Biochemistry, Faculty of Life Sciences, Ahmadu Bello University, Zaria, Nigeria • Department of Plant Pathology & Microbiology, Texas A&M University, College Station, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **valuable** study provides strong evidence for the development of a penetration ring during *Magnaporthe oryzae* infection and, supported by knockout and expression studies, shows that Ppe1 is involved in the virulence of the fungus. Although the authors demonstrated the close association of Ppe1 with the host plasma membrane, the work fell short in providing direct evidence for its role at the host-pathogen interface and the precise molecular function of the penetration ring. Therefore, the study presented strong structural and phenotypic characterization but remains **incomplete** regarding mechanistic insights of Ppe1.

<https://doi.org/10.7554/eLife.101209.2.sa2>

Abstract

Many fungal pathogens develop specialized infection structures such as appressoria to penetrate plant cells. However, it is not clear whether special structures are formed after cell wall penetration before invading host plasma membrane in hemibiotrophic pathogens. Here, we showed that a penetration ring consisting of Ppe1 secreted proteins is formed after appressorium-mediated cell wall penetration and remained at the base of penetration site after invading plant plasma membrane by the rice blast fungus *Magnaporthe oryzae*. The same persistent Ppe1 ring is formed after the penetration of neighboring cells by transpressoria. *PPE1* is specifically expressed during plant infection and the $\Delta ppe1$ mutant is defective in penetration and invasive growth. Blockage of penetration peg formation impedes the development of the Ppe1 ring. Close examinations showed that the penetration ring is formed at the collar of penetration pegs between the plant cell wall and plasma membrane and it is persistent as a fixed ring even after invasive hyphae invaded neighboring cells.

Furthermore, Ppe1 is a member of an expanded family of secreted proteins that are unique to fungal pathogens using extreme appressorium turgor for plant penetration. Other members of the Ppe1 family also localize to the penetration ring for anchoring on plasma membrane during plant infection. Taken together, a penetration ring consisting of a family of secreted proteins is formed between plant cell wall and plasma membrane, which may function as a novel physical structure at the interface between the tip of penetration pegs and plant plasma membrane before the differentiation of invasive hyphae.

Significance Statement

Like many other plant pathogens, the rice blast fungus forms melanized appressoria (specialized infection structures) to penetrate plant cells. In this study, we showed that a penetration ring is formed by penetration pegs after appressorium-mediated penetration of plant cell wall. This ring of secreted proteins is persistent at the point of penetration pegs invading plant plasma membrane to form invasive hyphae. Therefore, after the penetration of plant cell wall, the rice blast fungus forms a penetration ring that consists of a family of secreted proteins unique to pathogens using extreme appressorium turgor for penetration and may function as a physical structure for anchoring onto plant plasma membrane and developing invasive hyphae in penetrated cells.

Introduction

Magnaporthe oryzae, a filamentous ascomycete, is the causal agent of the devastating rice blast disease, which poses a significant threat to global rice production (1, 2). In addition to rice, *M. oryzae* is capable of infecting a wide range of grass species, including major cereal crops such as wheat, barley, oat, and millet, thus compromising global food security (3, 4). As one of the most important plant diseases, it is mainly controlled by planting resistant cultivars. Nevertheless, outbreaks of rice blast still occur throughout rice growing regions worldwide because of breakdown of resistance by new races of the pathogen. In addition, wheat blast caused by *M. oryzae* pathovar *Triticum* has recently spread from South America to Asia, threatening wheat production (5, 6).

Like many other foliar fungal pathogens, penetration of the cuticle and cell wall of host plants is the most critical infection step (7, 8). The infection cycle of *M. oryzae* starts with attachment of dispersing conidia to plant surface by releasing sport tip mucilage. Germ tubes emerging from attached conidia can sense surface hydrophobicity and hardness and develop highly specialized infection structures known as appressoria at the tip (1, 9). Mature dome-shaped appressoria are anchored on rice surface with appressorium mucilage (10) and heavily melanized by depositing a melanin layer between the cell wall and membrane as an impermeable barrier for turgor generation (11). Subsequently, the appressorium uses the extremely high turgor pressure to physically penetrate the cuticle and cell wall with a narrow penetration peg (7, 12, 13). As an hemibiotrophic pathogen, after the penetration of plant cell wall, the penetration peg does not penetrate plant plasma membrane but differentiates into primary invasive hyphae and invasive hyphae that are enveloped by plant-derived extra-invasive hyphal membrane (EIHM) in penetrated cells (7, 14). To date, it is not clear whether there is any special structure at the contact point between the penetration peg and plant plasma membrane before the differentiation of primary invasive hyphae. However, a biotrophic interfacial complex (BIC), a plant membrane-rich structure, is formed at the tip of the primary invasive hyphae for effector delivery (15, 16). Following proliferation within the first colonized cell, the tips of invasive hyphae differentiate into transpressoria, which subsequently form narrow infection pegs to traverse the

plasmodesmata (PD)-rich cell junctions, allowing entry into adjacent host cells (17, 18). The fungus continues this colonization strategy until disease symptoms manifest on infected rice plants.

To suppress plant defense responses and facilitate invasive growth, *M. oryzae* deploys effectors with various functions that can be delivered into host tissues or cells with two secretion systems during infection (19). Apoplastic effectors, such as Bas4 and Slp1, accumulate in the extracellular space between the fungal invasive hyphal cell wall and EIHM, outlining the entire invasive hyphae, are secreted through the conventional endoplasmic reticulum (ER)-Golgi-dependent pathway which is sensitive to brefeldin A (BFA) (19). Conversely, cytoplasmic effectors, including Pwl2, AvrPiz-t, and Bas1, accumulate in the BIC before being delivered into plant cells using a Golgi-independent pathway involving the exocyst complex and t-SNAREs, which is insensitive to BFA (19). To date, all the known proteinaceous effectors in *M. oryzae* belong to these two categories and localize to the extracellular space outlining invasive hyphae or the BICs and intracellular targets inside plant cells.

In a previous study, we identified a family of small secreted proteins that may function as effectors in *M. oryzae* (20). In this study, we found that one of them named as *PPE1* (for peg periphery-localized effector 1, MGG_16603) is specifically expressed during plant infection and is a crucial virulence factor. Surprisingly, Ppe1 preferentially localizes to the penetration site in a ring conformation that is described as the penetration ring below. The penetration ring consisting of Ppe1 proteins is formed after appressorium- and transpressorium-mediated plant penetration and remained at the point where penetration pegs transitioning into primary invasive hyphae. Ppe1 homologs are only present in fungal pathogens using extreme appressorium turgor for plant penetration. Other members of the Ppe1 family also localize to the penetration ring that is at the base of primary invasive hyphae outside plant plasma membrane. Taken together, a penetration ring consisting of secreted Ppe family proteins is formed between plant cell wall and plasma membrane, which may function as a novel physical structure at the interface between the tip of penetration pegs and plant plasma membrane before differentiation into invasive hyphae.

Results

Ppe1 is important for full virulence of *M. oryzae*

We previously identified 21 host-adapted effector genes in *M. oryzae* that may play critical roles during rice infection based on *in silico* analyses (20). To explore the functions of these effectors, we selected *PPE1* (MGG_16603) for further functional characterization because its high transcription level during early infection stages as reported (20, 21). We first assayed the expression profile of *PPE1* during infection by qRT-PCR. The result showed that its expression increased during the early stages of plant infection (24–36 hpi) and peaked at 36 hpi (Fig. 1A). Moreover, fluorescence signals of mCherry driven by the *PPE1* promoter indicated that its expression was only observed in matured appressoria and during infection (Fig. 1B). These results were consistent with the transcriptome data, suggesting that Ppe1 is functional during appressorium penetration and plant infection. Ppe1 encodes a 116-aa protein with typical characteristics of fungal effectors, including an N-terminal signal peptide and eight cysteine residues (Fig. S1A). The signal peptide of Ppe1 is functional in directing the secretion of yeast Suc2 invertase in a yeast signal trap system (Fig. S1B), suggesting that Ppe1 is a secreted protein.

To determine the biological functions of Ppe1 in *M. oryzae*, we generated a *PPE1* gene deletion mutant in the wild-type strain 70-15 (Fig. S1C). Deletion of *PPE1* had no obvious effect on the fungal growth, conidiation, and appressorium formation (Fig. S1D–G). However, in spray infection assays with rice leaves, the $\Delta ppe1$ mutant was significantly reduced in the number of disease lesions compared to the wild type (Fig. 1C). In wound infection assays, the $\Delta ppe1$ mutant also




Fig. 1

Ppe1 is important for full virulence in *M. oryzae*.

(A) The relative expression level of Ppe1 during infection of rice leaf sheaths assayed by qRT-PCR. (B) Fluorescence signals of the P_{PPE1} -mCherry transformant in penetration assays with rice leaf sheaths. Scale bars = 10 μ m. (C) Typical leaves of CO-39 seedlings sprayed with conidia of the wild type (70-15), *PPE1* gene deletion mutants ($\Delta ppe1-45$, $\Delta ppe1-101$) and complement strains ($\Delta ppe1-com-3$, $\Delta ppe1-com-8$) examined at 7 days post inoculation (dpi). Boxplot below shows the number of disease lesions per 5 cm of individual inoculated leaf. $n = 22, 23, 19, 19$, and 22 , respectively. (D) Wounded leaves of CO-39 drop-inoculated with the marked strains were examined at 10 dpi (upper panel) and measured for the lesion area (lower panel). $n = 10, 9, 7, 5$, and 4 respectively. (E) Rice leaf sheath penetration assays. Conidia suspensions from each strain were respectively injected into 4-5 weeks old rice leaf sheaths. Infection levels were observed at 32 hpi (hours post inoculation). The infection levels were grouped into four types: Type 1, appressorium formation without penetration; Type 2, having penetration peg or primary invasive hyphae; Type 3, having more than two branched invasive hyphae restricted in the first infected rice cell; Type 4, having invasive hyphae crossing to neighboring host cells. Scale bar = 10 μ m. Statistical analyses of the infection types and percentages are also shown. Significant differences were determined by one-way ANOVA. Asterisks indicated significant differences ($p < 0.0001$). ns indicated no significant difference ($p > 0.05$).

was significantly reduced in virulence and formed smaller lesions than the wild type (**Fig. 1D** [↗](#)). To confirm the role of *PPE1* in virulence, we generated a complemented strain *Δppe1*-com that had a copy of the *PPE1* gene ectopically integrated into its genome. In spray or wound infection assays, *Δppe1*-com was similar with the wild type in virulence (**Fig. 1C-D** [↗](#)), indicating a complete recovery in phenotypes. We then conducted rice leaf sheath penetration assays to observe the infection processes. At 32 hpi, over 91% of the appressoria formed by the wild type and the complemented strain had penetrated and developed invasive hyphae in rice cells.

Invasive hyphae in most of the penetrated cells (approximately 60%) had more than two branches, and approximately 10% of them had spread into the neighboring cells (**Fig. 1E** [↗](#)). In contrast, approximately 25% of the appressoria formed by the *Δppe1* mutant had no penetration. Although 53.5% of these mutant appressoria successfully penetrated into host cells, only 23% of *Δppe1* invasive hyphae had more than two branches and less than 4% had spread into adjacent cells (**Fig. 1E** [↗](#)). To further validate these observations, we generated a *PPE1* knockout in the wild-type reference strain Guy11. Similarly, we observed a significant decrease in the pathogenicity of the *Δppe1* mutants compared to Guy11 in the spray assay (**Fig. S2**). These results indicate that the *Δppe1* mutant has defects in appressorium penetration, invasive hyphal growth and subsequent spread of invasive hyphae into neighboring plant cells.

Ppe1 forms a ring at the base of appressoria during host penetration

To determine the localization of Ppe1 during infection, we expressed a full-length Ppe1-mCherry fusion protein under the control of *PPE1* promoter in both the wild type (70–15) and the *Δppe1* mutant. Rice leaf sheaths were then infected with these strains and the localization of Ppe1 was analyzed by confocal microscopy. In these transformants, Ppe1-mCherry signals were observed as a ring underneath the appressoria (likely at the site of the emergence of penetration pegs) at 22 hpi (**Fig. 2A** [↗](#), Video S1). The Ppe1-mCherry ring persisted after appressorial penetration and the ring-shaped fluorescence signal of the Ppe1-mCherry was still observed even after the formation of primary and secondary invasive hyphae (**Fig. 2B** [↗](#)). We further measured the size of this ring and found that the average diameter of the inner circle of the Ppe1-mCherry ring was 1.53 μm. The width of the Ppe1-mCherry ring varied from 1.79 to 4.78 μm, which is smaller than the width of appressoria (**Fig. 2C** [↗](#)) and known septin ring (5.9 μm) at the appressorium pore ([13](#) [↗](#)). Furthermore, the Ppe1-mCherry fusion protein could be detected by western blot analysis with proteins isolated from infected rice leave (**Fig. S3**). Because the Ppe1 ring always appears at the point of emergence of the penetration peg, we named this ring structure as penetration ring.

To confirm the localization of Ppe1, we also generated Ppe1-GFP transformants. Like Ppe1-mCherry, Ppe1-GFP also formed a ring at the penetration site underneath the appressorium (**Fig. 2D** [↗](#)). These results indicate that, irrespective of being fused with mCherry or GFP, Ppe1 localizes to the penetration ring. To avoid possible overexpression or unexpected effects due to ectopic integration of the fusion constructs, we generated a transformant from 70-15 strain in which mCherry was inserted right before the TAA stop codon of the endogenous *PPE1* gene. In the resulting transformant expressing the in situ *PPE1*-mCherry fusion construct, the penetration ring was also observed (**Fig. 2E** [↗](#)). These results further verify the localization of Ppe1 to the penetration ring during appressorial penetration.

Penetration ring is also formed during the penetration of neighboring cells by transpressoria

In *M. oryzae*, appressoria-like structures called transpressoria are formed by invasive hyphae on the wall boundary of the invaded cell in order to penetrate into neighboring cells ([17](#) [↗](#), [18](#) [↗](#)). Therefore, we observed localization of Ppe1 during transpressorium-mediated penetration of plant cells. In the *PPE1*-mCherry transformant, like during appressorial penetration, the




Fig. 2

Ppe1-mCherry forms a ring during appressorial penetration.

(A) A ring of Ppe1-mCherry signals at the penetration site was observed at 22 hpi in leaf sheath penetration assays with the *PPE1*-mCherry transformant. Upper panel indicates the Ppe1-mCherry expressed in wild-type strain (70-15). Lower panel indicates the Ppe1-mCherry expressed in $\Delta ppe1$ mutant. Scale bars = 2 μ m. (B) The Ppe1-mCherry penetration ring was persist even after the growth of primary and secondary invasive hyphae. Scale bars = 5 μ m. (C) The diameters of appressoria, Ppe1-mCherry penetration ring (ring), and its inner circle. $n=21, 65$ and 65 respectively. (D) The formation of Ppe1-GFP rings at the penetration site formed by the *PPE1*-GFP transformant. Scale bars = 2 μ m. (E) The in-situ *PPE1*-mCherry transformant also formed a fluorescent ring at the penetration site in leaf sheath penetration assays. Scale bar = 2 μ m. AP, Appressorium. Triangle arrows pointed to penetration rings.

penetration ring was observed during transpressoria-mediated penetration of neighboring cells (**Fig. 3A**). Similarly, we observed the penetration ring in the neighboring cells penetrated by invasive hyphae of the *PPE1*-GFP transformant (**Fig. 3B**). The Ppe1-mCherry or Ppe1-GFP ring (**Fig. 3A and B** – zoomed view) appeared to be at the transpressorial penetration point. These results indicate that the penetration ring is also formed during invasive hyphal cell-to-cell penetration. Furthermore, similar localization of Ppe1 to the penetration ring was observed in barley epidermal cells (**Fig. 3C**). Taken together, we conclude that Ppe1 localizes to the penetration ring during appressorium-mediated and transpressorium-mediated plant cell penetrations in *M. oryzae*.

Penetration ring is formed on the periphery of the penetration peg

In *M. oryzae*, appressorium-mediated penetration is a septin-dependent process in which septins aggregate and generate a hetero-oligomeric 5.9 μm ring structure at the appressorial pore to re-organize the F-actin cytoskeleton network to promote the emergence of penetration pegs (13). Following the penetration peg formation, septin ring undergoes further constriction into a smaller ring (22). To clarify the precise site of the penetration ring, we expressed Ppe1-mCherry (under the control of the *PPE1* promoter) in a strain expressing Sep3-GFP (Guy11 background) and then observed their spatiotemporal dynamics during rice leaf sheath infection. At 11 hpi, Sep3-GFP big ring was formed at the appressorium pore but the Ppe1-mCherry ring was not observed (**Fig. S4A**, Video S2). The Ppe1-mCherry ring signal was observed at 16 hpi, after the constriction of Sep3-GFP ring and the former was completely surrounding the latter (**Fig. S4B**). Subsequently (at 19 hpi), the Ppe1-mCherry ring (still surrounding the Sep3-GFP ring) became constricted (**Fig. 4A**, **Fig. S4C**, Video S3). However, on impenetrable hydrophobic coverslips, the Sep3-GFP ring was formed normally at the appressorium pore, but the Ppe1-mCherry ring and constriction of Sep3-GFP were not observed (**Fig. S5**). These results suggest that Ppe1-mCherry ring is formed after the formation of penetration pegs, outside the constricted Sep3-GFP ring.

Penetration pegs are rich in microfilaments, filasomes and microtubules (23), therefore, it can be labeled by Lifeact-GFP. Hence, we co-expressed Ppe1-mCherry with Lifeact-GFP in 70-15 and observed their positioning in penetration assays with rice leaf sheaths. The Ppe1-mCherry ring was also observed to surround the Lifeact-GFP signal (**Fig. 4B**, **Fig. S6**, Video S4), indicating that the penetration ring is localized at the periphery of penetration pegs during the initial infection stage and primary invasive hyphal development. To confirm this observation, we co-expressed the Ppe1-mCherry with cytosolic GFP in 70-15. Using a three-dimensional (3D) volume rendering image constructed by confocal optical z-series images, we found that the Ppe1-mCherry ring was situated around the narrow site between the appressorium and invasive hyphae labeled with GFP (**Fig. 4C**, **Fig. S7**).

A previous study showed that callose deposition is dynamic during *M. oryzae* infection and has three different forms: callose-rich papillae underneath appressorium penetration sites, callose collars formed at cell wall crossing sites around the invasive hyphae and callose deposition at plasmodesmata pit fields (24). To investigate whether the penetration ring locates outside the peg, we stained callose deposits with aniline blue to examine the possible positioning of the callose ring underneath the appressorial penetration site and Ppe1-mCherry. We discovered that the Ppe1-mCherry ring was surrounded by the callose ring, exhibiting partial colocalization and close attachment (**Fig. 4D**, Video S5). Furthermore, we examined the localization of Ppe1-mCherry in transgenic rice expressing GFP-OsPIP2 as a plasma membrane marker (25). Our data demonstrate that the penetration ring is tightly associated with the host plasma membrane (**Fig. S8**). Taken together, we conclude that the penetration ring locates at the periphery of the penetration peg and anchors the peg to the host plasma membrane during the initial appressorium-mediated host penetration.




Fig. 3

Penetration ring formation during invasive hyphal penetration into neighboring cells.

(A) Ppe1-mCherry forms rings at about 36 hpi. (B) Ppe1-GFP also forms rings similar to Ppe1-mCherry during penetration to neighboring cells. (C) Ppe1 ring formation is also observed during barley infection. Scale bars = 10 μ m. Arrows heads point to penetration rings at the penetration sites of invasive hyphae. White squares indicate zoomed regions.




Fig. 4

The penetration ring localizes on the periphery of penetration peg.

(A) Ppe1 ring surrounds the Sep3 constricted ring. Scale bar = 2 μm. (B) Ppe1 ring also surrounds the Lifeact ring. Scale bar = 1 μm. (C) Ppe1 ring is formed around the narrow site between an appressorium and invasive hypha. Three-dimensional (3D) volume rendering images were constructed by confocal optical z-series images captured at 0.35 μm intervals, with a total of 22 steps. (D) Ppe1 ring partly colocalized with callose ring at the apoplastic papilla. Scale bar = 1 μm. Hollow triangles indicate the site of penetration peg. Line graphs were generated at the directions pointed by the white arrows.

Formation of the penetration ring at the base of penetration pegs after transpressorium penetration

To determine the precise localization of the penetration ring during the transpressoria-mediated penetration, we expressed Ppe1-mCherry (under the control of the *PPE1* promoter) in a Sep5-GFP strain and then observed their spatiotemporal dynamics during rice leaf sheath infection. The results showed that Sep5-GFP first displayed a ring structure at the swollen part of the invasive hyphal tip (Fig. S9A), similar to the septin big ring at the appressorium pore. It then exhibited a collar structure at the constricted infection peg site (Fig. S9B-C), which is consistent with a previous report by Sakulkoo (26). Meanwhile, the Ppe1-mCherry ring or collar-like ring was formed outside the Sep5-GFP collar (Fig. S9B-C), indicating that the penetration ring formed during transpressoria-mediated cell-to-cell penetration is positioned outside the penetration peg. Taken together, we conclude that the penetration ring of *M. oryzae* is localized at the periphery of penetration pegs both during appressoria- and transpressoria-mediated host penetrations.

Penetration ring formation requires the initiation of penetration peg formation

To further characterize the relationship between the penetration ring and penetration peg, we observed the expression and localization of Ppe1-mCherry in $\Delta mst12$, $\Delta nox2$ and $\Delta nox1$ mutant strains that are defective in plant penetration (27, 28). In the $\Delta mst12$ mutant that is defective in both penetration initiation and peg formation, the Ppe1-mCherry fluorescence signal could not be detected. In the $\Delta nox2$ mutant that is unable to initiate penetration peg formation, the Ppe1-mCherry ring could not be formed but displayed a disorganized mass underneath the appressorium. Interestingly, in the $\Delta nox1$ mutant that forms normal penetration pegs but is defective in invasive hyphal elongation, the Ppe1-mCherry ring was formed during rice leaf sheath infection (Fig. 5A), indicating that penetration peg formation is a prerequisite for the emergence of the penetration ring. To confirm this observation, the localization of Ppe1-mCherry fusion proteins was observed after inducing appressorium formation on hydrophobic coverslips and cellophane membranes. Although melanized appressoria were formed, we failed to observe the Ppe1-mCherry ring on the impenetrable coverslips (Fig. 5B). However, the penetration ring is formed after the penetration of cellophane membranes (Fig. 5C). These results indicate that penetration is necessary for the formation of the penetration ring.

Other Ppe proteins also localize to the penetration ring

To determine whether other secreted proteins homologous to Ppe1 also localize to the penetration ring, we used the amino acid sequence of Ppe1 to carry out BLASTp search at the NCBI database to uncover the homolog proteins of Ppe1 and further analyzed their phylogenetic relationships. The Ppe1 homologs only exist in *Magnaporthe* and *Colletotrichum* species, such as the most common cereal phytopathogens *M. oryzae* and *C. graminicola* (Fig. 6A). Remarkably, these fungi are known to form melanized appressoria and deploy extremely high turgor pressure for plant penetration (8). To understand whether other members of the Ppe1 family (Ppe2, Ppe3, Ppe4, Ppe5, and Ppe6) from *M. oryzae* have similar localization as Ppe1, we tagged each of them with mCherry and expressed them individually in the 70-15 background under the control of their native promoters. Examination of their localizations during infection revealed that Ppe2-mCherry, Ppe3-mCherry and Ppe5-mCherry also exhibit a ring structure during appressorium-mediated host penetration, while the fluorescence signals of Ppe4-mCherry and Ppe6-mCherry were hardly observed, possibly due to their low expression (Fig. 6B). Moreover, when MoT-Ppe1 from *M. oryzae* pathovar *Triticum* was expressed in *M. oryzae* under the control of its native promoter, the characteristic penetration ring was also observed (Fig. 6B). All these results suggest that the secreted Ppe proteins accumulate at the penetration ring.

Fig. 5

Penetration ring formation requires the emergence of penetration peg.

(A) Ppe1 ring expression and localization in 70-15, $\Delta mst12$, $\Delta nox2$ and $\Delta nox1$ strains at 24 hpi. Conidia suspensions from each strain were inoculated on rice leaf sheaths and observed at 24 hpi. (B) Ppe1 ring could not be formed on impenetrable coverslips. Conidia from Ppe1-mCherry strain were inoculated on hydrophobic coverslips and observed at 24 hpi. (C) Ppe1 ring was observed at 38 hpi on penetrable cellophane membrane. Scale bars = 2 μ m.





Fig. 6

Other secreted Ppe proteins also localize to the penetration ring.

(A) Phylogenetic tree of Ppe1 constructed using neighbor-joining method. (B) Paralogs of Ppe1 in *M. oryzae* localize to the penetration ring. Ppe2-mCherry, Ppe3-mCherry and Ppe5-mCherry were driven by their respective native promoters. MoTPpe1-mCherry from *Magnaporthe oryzae* *Triticum* was expressed in *M. oryzae* (70-15), driven by its native promoter. Scale bars = 2 μ m. (C) Comparison of the localization of penetration ring with those of some previously reported fungal effectors. Bas4-GFP and Slp1-GFP uniformly outlined the extra-invasive hyphal membrane (EIHM), while Pwl2-mCherry and AvrPiz-t-GFP accumulated in the biotrophic interfacial complex (BIC). Scale bars = 10 μ m.



As controls, we also examined the localizations of Bas4, Slp1, Pwl2, and AvrPiz-t. Under the same experimental conditions, Bas4-GFP and Slp1-GFP were observed to uniformly accumulate within the extra-invasive hyphal membrane (EIHM), outlining the entire invasive hyphae, while Pwl2-mCherry and AvrPiz-t-GFP accumulated in the biotrophic interfacial complex (BIC) (**Fig. 6C**). which is consistent with previous reports (16, 29, 30). EIHM and BIC are specifically associated with invasive hyphae during biotrophic growth. In contrast, Ppe1 localized to the penetration ring at the periphery of penetration pegs (**Fig. 6C**). Therefore, unlike subcellular localizations of Bas4, Slp1, Pwl2, and AvrPiz-t to the extracellular space outlining invasive hyphae, BICs, or intracellular targets in plant cells, Ppe1 and its homologs localize to the penetration ring that may function as a novel physical structure at the interface between the tip of penetration pegs and plant plasma membrane before the differentiation of invasive hyphae.

Discussion

As the model plant pathogenic fungus, appressorium formation and appressorium-mediated penetration have been well-characterized in *M. oryzae* (18). The fungus uses extremely high turgor pressure generated in appressoria cemented to the surface with appressorium mucilage to penetrate underlying plant cells (12). As an hemibiotrophic pathogen, after piercing plant cell wall, the penetration peg does not penetrate the plasma membrane but differentiates into invasive hyphae enveloped in a layer of EIHM (16). In this study, we showed that a penetration ring of Ppe1 proteins is formed by the penetration peg on the surface of the plant plasma membrane before invading and differentiating into primary invasive hyphae (**Fig. 2** and **Fig. 7**). Other Ppe proteins including Ppe2, Ppe3, and Ppe5 proteins also localize to the penetration ring (**Fig. 6B**). Because the penetration ring is persistent at the point penetration pegs transitioning into invasive hyphae outside the plasma membrane even after the penetrated cells are filled with invasive hyphae, it may function as a novel physical structure for anchoring penetration pegs on the surface of plant plasma membrane after cell wall penetration (**Fig. 7**). It is also possible that this penetration ring functions as a collar that is associated with the differentiation of penetration pegs (on the surface of plasma membrane) into primary invasive hyphae enveloped in the EIHM (**Fig. 7**). Although this transition is not well characterized, it likely involves changes in cell polarity and cytoskeleton as well as cell wall modifications. Ppe1 and its homologs may be able to assemble into a protein complex in the penetration ring, which likely involves other proteins or components as facilitating factors. In *U. maydis*, the Stp secreted effectors are known to form a protein complex during plant infection (31).

The penetration ring is also formed after transpressoria-mediated penetration of neighboring plant cells (**Fig. 7**). Similarly, Ppe1 and its homologs form a ring after the penetration of cell wall but before differentiating into invasive hyphae enveloped with the EIHM. The penetration ring formed by transpressoria-mediated penetration is also persistent at the penetration site on the surface of plant plasma membrane in penetrated cells (**Fig. 3** and Fig. S5), suggesting that Ppe proteins are assembled into a complex that is not diffusible. In fact, similar penetration ring is observed after penetration of artificial membrane (**Fig. 5C**), which has no barrier to restrict the diffusion of secreted proteins, further indicating that the penetration ring of Ppe1 and other Ppe proteins is likely a structural feature or component for signaling the transition of penetration pegs to invasive hyphae. Although appressoria and transpressoria are different, the processes after the penetration of cell wall are the same in appressorium- and transpressorium-mediated plant penetration. Therefore, formation of the penetration ring in both appressorium- and transpressorium-mediated penetration is another evidence for it to be a structural element.

M. oryzae is known to secrete a repertoire of effectors that can be categorized into cytoplasmic and intracellular effectors (14). Ppe1 and its homologs are putative effectors that are described as HAG (host adaptation genes) effectors in an earlier study (20). Although it remains possible that Ppe1 and its homologs function as effectors during plant infection. However, in comparison




Fig. 7

A proposed working model for the discovery of penetration ring in *M. oryzae*.

Penetration ring forms at the periphery of peg and accumulates fungal secreted protein such as Ppe1 to promote *M. oryzae* invasion.

with the localizations of known effectors in *M. oryzae*, we found that the localization of Ppe1 to the penetration ring is unique and Ppe1 homologs have the same localization pattern (Fig. 6). Because of the persistent localization of Ppe1 and its homologs to the penetration ring, we think the penetration ring is likely a physical structure as described above. If Ppe1 and its homologs are effector proteins being secreted at this specific location, fluorescent signals of Ppe1 and its homologs will diffuse and disappear or diminish when effectors being translocated into plant cells. Furthermore, it is unlikely that secreted proteins localized to the penetration ring that has limited contact with the plant plasma membrane will function as effector proteins to interact with plant targets. Nevertheless, although unlikely, it remains possible that the penetration ring discovered in this study is a secretion site for secreted proteins involved in the transition of penetration pegs to invasive hyphae. After the penetration of plant cell wall, the penetration ring may not only function as a physical structure but also serve as an secretion site for the release of specific proteins to overcome plant immunity in early infection stages.

Taken together, in this study we observed the formation of the penetration ring that appears to be a structural element for signaling or triggering the differentiation of penetration pegs after the penetration of plant cell wall into invasive hyphae enveloped by the EIHM in *M. oryzae*. Based on the limited distribution of Ppe1 homologs, it is likely that only fungal pathogens using high appressorium turgor for plant penetration, such as *M. oryzae* and *Colletotrichum* species, developed the penetration ring. Therefore, it will be important to further characterize the functional relationship between the penetration ring and appressorium turgor as well as the transition of penetration pegs driven by turgor pressure into primary invasive hyphae (lower turgor pressure).

In summary, our findings have several significant implications. First, we believe that the discovery of the penetration ring as a novel physical structure associated with the differentiation of invasive hyphae represents a breakthrough in plant-pathogen interactions. Secondly, our study presents new role of the peg as a specialized platform for the deployment of secreted protein, in addition to its commonly known role as a physical penetration tool for the pathogen. Thirdly, we identified Ppe1 as a potential molecular target for controlling the devastating rice blast disease, as Ppe homologs are absent in plants and mammals. The primary challenge for future research lies in elucidating the pathogenesis associated with penetration rings. Achieving this will further our understanding of the mechanism of rice blast fungus-host interaction.

Materials and Methods

Strains and growth conditions

The *M. oryzae* strains 70-15 and Guy11 were used as the wild types in this study; all other fungal strains generated from the wild types are listed in Table S1 in the supplemental materials. All the fungal strains were grown on complete medium CM (CM: 10 g D-glucose, 2 g peptone, 1 g yeast extract, 1 g casamino acids, 50 mL 20 × nitrate salts (120 g/L NaNO₃, 10.4 g/L KCl, 30 g/L KH₂PO₄, and 10.4 g/L MgSO₄), 1 mL trace elements (0.1 g/L biotin, 0.1 g/L pyridoxine, 0.1 g/L thiamine, 0.1 g/L riboflavin, 0.1 g/L p-aminobenzoic acid and 0.1 g/L nicotinic acid), adjusted to pH 6.5 with NaOH, 15 g agar and distilled water added to 1 L) (32). Agar cultures were incubated at 26 °C with a 12-h light/12-h dark photoperiod for mycelial growth and conidiation. Liquid CM medium was used to harvest mycelia for extraction of genomic DNA and protoplast preparation. The bacterial strains *Escherichia coli* DH5α and *Agrobacterium tumefaciens* AGL1 were grown in Luria-Bertani (LB) medium at 37 °C and 28 °C, respectively.

Plant and growth conditions

Rice cultivar CO-39 (*Oryza sativa* cv. CO-39) was grown in a greenhouse at 26 and 70% relative humidity with a 14-h light/10-h dark photoperiod. Barley (*Hordeum vulgare* cv. Golden Promise) was grown at 23 with a 12-h light /12-h dark photoperiod.

Bioinformatic and phylogenetic analyses of Ppe1 homolog proteins

The Ppe1 was previously sequenced and the amino acid sequence established from our lab (33). Ppe1 signal peptide was predicted using SignalP 5.0 software. BLASTp was used to find Ppe1 homolog proteins at the NCBI database (<https://www.ncbi.nlm.nih.gov/>) and the representative homolog protein sequences were downloaded from the database. Phylogenetic tree was generated using MEGA7.0 software by neighbor-joining method.

Construction of recombinant DNA and *M. oryzae* transformation

The plasmids used in the study were generated by infusion cloning method based on homologous recombination using ClonExpress II one-step cloning kit or ClonExpress MultiS kit (Cat. No. C112-01 or C113-01, Vazyme, Nanjing, China). The primers used (Table S2 in the supplemental material) were specially designed to contain 18-22-bp overlap with adjoining sequences to allow the fragments assemble by homologous recombination. To observe the expression pattern and localization of *PPE1*, the *PPE1* promoter (1485 bp) fragment amplified from the genomic DNA of the wild type (70–15) was seamlessly cloned into the *KpnI/HindIII*-digested pKNT-mCherry/GFP fusion vectors (34) using the ClonExpress II one-step cloning kit (Cat. No. C112-01, Vazyme, Nanjing, China). For Ppe1-mCherry/GFP plasmid, the entire Ppe1 coding sequence with its native promoter but without stop codon was amplified and cloned into the *KpnI/HindIII*-digested vector pKNT-mCherry/GFP. The pKNT-AvrPizt-GFP and other plasmids used were constructed using the same method. The constructed plasmids were transformed into the fungal protoplasts via PEG-mediated transformation (35) and further verified by PCR screening and confocal microscopic examination of fluorescence signals. For situ fusion of Ppe1 with mCherry, the mCherry was inserted just before the Ppe1 stop codon and the construct was transformed by *Agrobacterium tumefaciens*-mediated (ATMT) transformation for homologous recombination (35).

Assays for the secretory function of the SP of Ppe1

To confirm the secretory function of the predicted signal peptide (SP), yeast signal trap system was performed as described previously (36). The predicted SP nucleotide sequences of Ppe1 (60 bp) and Pwl2 (63 bp) were respectively cloned into pSUC2 vector that carries a truncated invertase gene *SUC2*. The generated pSUC2-SP constructs and the empty pSUC2 vector were transformed into the yeast *suc2* mutant YTK12, respectively, and then their growth conditions analyzed on CMD-W (minus Trp) plates (0.67% yeast nitrogen base without amino acids, 0.075% tryptophan dropout supplement, 2% sucrose, 0.1% glucose, and 2% agar) and YPRAA medium plate (1% yeast extract, 2% peptone, 2% raffinose, and 2 µg/mL antimycin A) containing raffinose and lacking glucose. Transformants of YTK12 carrying the empty pSUC vector or pSUC2-Pwl2SP were used as negative and positive controls, respectively. Reduction of 2,3,5-triphenyltetrazolium chloride (TTC) to insoluble red-colored 1,3,5-triphenylformazan (TPF) was used as an indicator of the invertase enzymatic activity (37). Each of the above YTK12 transformants strain was inoculated in 5 mL of CMD-W liquid medium and incubated for 24 h at 30. The cells were then collected, washed and resuspended in 0.1% TTC in new centrifuge tubes and further incubated at 35 for 35 min. The color change was observed after 10 min of incubation at room temperature to detect the signal peptide secretion function.

Targeted gene deletion and complementation

Deletions of *NOX1*, *NOX2* and *MST12* genes were performed using Split-marker recombination method (38). The open reading frames (ORFs) of the various genes were replaced with a hygromycin phosphotransferase (*HPH*) resistance cassette, respectively. Briefly, about 1-kb upstream and downstream sequences of the targeted genes were amplified from the genomic DNA of the wild type (70–15) using specially designed primer pairs AF+AR and BF+BR, respectively. At the same time, the upstream and downstream sequences of the *HPH* with overlapping segments were amplified from pCB1003 plasmid using the primer pairs HYG/F + HY/R and YG/F + HYG/R, respectively. The corresponding upstream and downstream fragments were respectively ligated by splicing for overlap extension (SOE) PCR using the primer pairs AF+HY/R and YG/F+BR. The PCR products were purified and transformed into the *Ppe1*-mCherry strain protoplasts via a PEG-mediated transformation (35). Putative transformants were selected in the presence of hygromycin B and then confirmed by PCR using the primers OF+OR and UA+H853. PCR confirmations were repeated three times for consistency. All the primer sequences used are listed in Table S2 of the supplemental material.

For *PPE1* gene deletion, *HPH* sequence was inserted between the upstream and downstream sequences of the *PPE1* gene, replacing the ORF of the *PPE1* gene, in an ATMT1 plasmid using a ClonExpress MultiS One Step Cloning Kit (Cat. no. C113-01, Vazyme, Nanjing, China). The constructed plasmid was introduced into *Agrobacterium tumefaciens* AGL1 using freeze-thaw method (39). *Agrobacterium tumefaciens*-mediated transformation (ATMT) of the construct into the wild-type spores was then carried out. The transformants were selected on hygromycin B plates and confirmed by PCR. The mutants were further confirmed by Southern blot assay using DIG-High Prime DNA labeling and detection starter kit I (Cat. no. 11745832910, Roche, Mannheim, Germany). For complementation, the constructed plasmid pKNT-*Ppe1*-mCherry was transformed into the Δ *ppe1* mutant protoplast and further verified by PCR screening and confocal microscopic examination for fluorescence signal detection.

Live cell imaging

To monitor the expression and subcellular localization of *Ppe1* during plant infection, rice leaf sheath inoculation and live cell imaging of fluorescently labeled proteins were performed as described (15) with slight modifications. In brief, 5–7 cm long leaf sheath segments were cut from 4-week-old CO-39 rice seedlings, and the whole hollow spaces of the sheaths were filled with conidia suspensions (5×10^4 conidia/mL) using a 200 μ L pipette. The inoculated sheaths were placed horizontally (fixed with wet filter paper fragments) on an inverted 96-well PCR plate covered by one layer of wet filter paper and incubated in a sealed box whose inner bottom was lined with two layers of wet filter paper to keep the environment humid at 26. The epidermal parts of the inoculated sheaths were trimmed into thin and transparent samples and immediately imaged using an inverted Nikon A1 laser scanning confocal microscope (Nikon, Tokyo, Japan) at the indicated time points post inoculation. The 60 $\times$ /1.42 numerical aperture (NA) oil immersion objective lens was used for the imaging. The fluorescence and differential interference contrast (DIC) images were collected by confocal optical z-series acquired at 0.28 μ m intervals spanning the entire fluorescence signal depth between the top plane of the appressorium formed and the bottom plane of the rice leaf sheath cross sections. The fluorescence excitation/emission wavelengths were 488 nm/500–550 nm for GFP, 561 nm/570–620 nm for mCherry and 405 nm/425–475 nm for callose staining. The images were processed using NIS-Elements AR analysis software (Nikon, Tokyo, Japan). Unless otherwise stated in the figure legend, micrograph fluorescence represents maximum intensity projection of z-series (with a 0.28 μ m intervals) after removing extraneous z-planes that do not contain relevant information for a particular fluorescence. Three-dimensional (3D) volume rendering images and videos were generated from z-series in maximum intensity projection.

To observe the subcellular localizations of Ppe1 in conidia incubated on artificial hydrophobic surfaces and barley leaves, conidia suspensions (5×10^4 conidia/mL) were dropped on the surfaces of hydrophobic coverslips or on the back sides of barley leaves (5-day old plants) and incubated under humid and dark conditions at 26. For observation, the hydrophobic coverslips containing the appressoria were inverted, placed on 24mm×50 mm microscope glass slides and then observed under a scanning confocal microscope. Epidermal parts were removed from the infected barley leaves for observation. For penetration assay, cellophane was cut into round sheets of 4 cm in diameter and sterilized by autoclaving at 115 for 15 min. The sterile cellophane sheets were placed flat on CM plates and then the conidia suspensions (5×10^4 conidia/mL) were dropped on the surface of the cellophane and incubated at 26. Small pieces of the cellophane containing mycelia were cut for observation under a confocal microscope at 36–48 hpi.

Callose staining

For visualization of deposited callose, aniline blue (Cat. no. 28631-66-5, Sangon, Shanghai, China) staining was adopted (26 [↗](#)). Rice leaf sheaths were inoculated with conidia suspensions (3×10^4 conidia/mL). The inoculated sheaths were trimmed to thin and transparent samples at the indicated time points and stained with aniline blue (0.1mg/mL) solution prepared in 150 mM KH_2PO_4 (pH 9.5) for up to 30 min. The stained samples were imaged by using a laser scanning confocal microscope (Nikon, Japan).

Phenotypic analysis

For vegetative growth and conidiation assays, small mycelial blocks from each strain were cut from the edges of 4-day-old cultures and placed at the center of freshly prepared CM plates (90 mm) and incubated for ten days at 26. The colony diameters of the cultures were measured using vernier calipers and then photographs of the cultures were taken. The cultures were then washed with 5 mL sterilized double-distilled water, filtered through three layers of lens paper and the conidia counted using a hemocytometer under a light microscope. For conidia germination and appressoria formation, spore suspensions (1×10^5 conidia/mL) were incubated on hydrophobic coverslips (Cat. no. 12-545-101P, Fisherbrand™ Cover Glasses, USA) under humid and dark conditions at 26 for 24 h. The conidia were then observed under a light microscope for germination and formation of appressoria.

To investigate the pathogenicity of each strain, conidia spray, punch and leaf sheath inoculations were performed. Conidia suspensions (5×10^4 conidia/mL containing 0.02% Tween-20) were used to spray the leaf surfaces of 3-week-old CO-39 rice seedlings. The inoculated seedlings were incubated in a growth chamber at 25 with 90% humidity. The seedlings were first kept in the dark for 24 h, and then subjected to 12 h light / 12 h dark photoperiod for 7 days, after which photographs were taken and the number of disease lesions on the leaf surfaces counted. Punch inoculation assay was carried out according to a method previously described (30 [↗](#)). Briefly, 5-week-old CO-39 rice leaves were slightly wounded using mouse ear punch and 10 μL of conidia suspensions (5×10^5 conidia/mL containing 0.02% Tween-20) from the various strains were respectively dropped at the wounded regions. Then, both sides of each inoculated spot were sealed with scotch tape to sustain the spore suspension. The inoculated plants were kept in a growth chamber under the same conditions described above for 10 days. The disease lesions formed were scanned, and later analyzed using Adobe Photoshop software to measure the lesion areas. For leaf sheaths inoculation, conidia suspensions (1×10^5 conidia/mL) from the various strains were respectively injected into 4–5 weeks old CO-39 rice leaf sheaths. The epidermal parts of the infected leaves were trimmed to thin and transparent samples for observation under a light microscope at 32 hpi. At least 100 infected sites were observed using a Nikon A1 laser confocal microscope (Nikon, Tokyo, Japan).

qRT-PCR analysis

To analyze the expression level of *PPE1* during *M. oryzae* infection, rice leaf sheaths were cut from 4-to 5-week-old CO-39 rice seedlings and inoculated with conidia suspensions (2×10^5 conidia/mL) from the wild type (70–15). The epidermal parts of the inoculated sheaths were cut and collected at 12-, 36- and 48 hours post inoculation (hpi). At least 15 segments of the inoculated rice sheath samples were collected at each time point. Total RNAs were extracted from the leaf sheaths or rice seedling leaves using an Eastep Super Total RNA Extraction Kit (Cat. no. LS1040, Promega, Shanghai, China). Reverse transcription was used to generate cDNA using a reverse transcription kit (Cat. no. RR037B, Takara, Dalian, China). Quantitative PCR was finally conducted using a SYBR kit (Cat. no. RR067A, Takara, Dalian, China) using specific primers (Table S2). Fungal β -*TUBULIN* gene used as endogenous reference genes, and the expression levels were calculated using $2^{-\Delta\Delta Ct}$ method as previously reported (40 [40](#)). All qRT-PCR assays were repeated three times for each sample.

Statistical analyses

Statistical analyses were performed using GraphPad Prism7 (GraphPad Software, Inc., San Diego, CA). Statistically significant differences were determined by one-way ANOVA with two-sided least significant difference. **** indicated significant difference at $p < 0.0001$, ns indicated no significant difference.

Data Availability

The authors declare that all the data supporting the findings of this study are available within the paper and SI Appendix. Raw data and details of experimental procedures are available upon request from the corresponding authors.

Acknowledgements

We thank Prof. Nicholas J. Talbot and Prof. Barbara Valent for providing us with $\Delta sec5$ mutant and pBV591 plasmid respectively. We also thank Prof. Min Guo for providing us with Sep3/5-GFP strains and $\Delta sep3/4/5$ mutants. We also appreciate Profs. Guodong Lu, Jie Zhou, Huawei Zheng, Wei Tang, Ya Li, Xiaofeng Chen, and Huakun Zheng, as well as Drs Yijuan Hang, Meilian Chen and Yonghe Hong for their helpful suggestions. Thanks to Jin Zhang, Chunling Li and Yuping Fan's help. This research was supported by the National Key Research and Development Program of China (2023YFD1400200), the National Natural Science Foundation of China (32122071, 31601596), and the Natural Science Foundation of Fujian Province (2021J06015).

Additional files

Video S1. 3D visualization of *Ppe1-mCherry* ring formation at 22 hpi. [40](#)

Video S2. 3D visualization of *Ppe1-mCherry* and *Sep3-GFP* localizations at 11 hpi. [40](#)

Video S3. 3D visualization of *Ppe1-mCherry* ring surrounding the *Sep3-GFP* small ring at 19 hpi. [40](#)

Video S4. 3D visualization of *Ppe1-mCherry* ring surrounding the penetration peg at 22 hpi. [40](#)

Video S5. 3D visualization of *Ppe1-mCherry* ring and papilla callose ring at the penetration site at 22 hpi. [🔗](#)

Supplementary figures and tables




Fig. S1

Ppe1 is dispensable for mycelial growth, conidiation and appressorium formation in *M. oryzae*.

(A) Amino acid sequence of *M. oryzae* Ppe1. Signal peptides (green) were predicted by Signal-5. Cysteine residues are shown in red. (B) Detection of Ppe1 signal peptide secretion by yeast signal trap assay. The recombinant pSUC2-Ppe1SP and pSUC2-Pwl2SP constructs were transformed into the yeast strain YTK12 respectively. The pSUC2-Pwl2SP-expressing strain served as a positive control while pSUC2 empty vector was used as a negative control. (C) Targeted gene deletion strategy for *PPE1* in *M. oryzae* 70-15. Schematic map showing targeted disruption of *PPE1* gene via *Agrobacterium tumefaciens*-mediated transformation. The Southern blot result shows a successful replacement of *PPE1* by single insertion of hygromycin phosphotransferase (*HPH*) resistance cassette at *PPE1* ORF locus. (D) Colonies of the wild type (70-15), two independent *Δppe1* transformants, and complemented strains. (E) Statistical analyses of the colony diameters of each strain after being cultured on complete medium (CM) at 25 °C for 10 days. n = 9. (F) Statistical analyses of the number of conidia produced by each strain after being cultured on CM at 25 °C for 10 days. n = 17. (G) Appressorium formation by the indicated strain observed at 24 hours on artificial hydrophobic surfaces. Scale bar = 50 μm. Significant differences were determined by one-way ANOVA. ns stands for no significant difference (p>0.05).

Fig. S2

Ppe1 is essential for virulence in *M. oryzae* strain Guy11.

(A) Disease symptoms on leaves of CO-39 rice seedling inoculated with conidial suspension (5×10^4 spores/ml) from Guy11, $\Delta ppe1-21$ and $\Delta ppe1-24$. Photographs were taken 7 days after inoculation. (B) Quantification of the lesion number per 5 cm length of rice leaf. Asterisk represents significant difference. **, $p < 0.01$. ***, $p < 0.001$.





Fig. S3

Detection of Ppe1-mCherry fusion protein by western blot.

M, protein marker. CK, rice leaf sheaths injected with sterile double distilled water.






Fig. S4

Ppe1 ring is formed simultaneously with peg penetration.

(A-C) Spatiotemporal dynamics of Ppe1 ring and Sep3 ring during appressorium-mediated peg penetration into rice leaf sheaths. Scale bars = 2 μm . Hollow triangles show the sites of penetration peg. Line graphs were generated at the directions pointed by the white arrows.




Fig. S5

Spatiotemporal dynamics of Ppe1-mCherry and Sep3-GFP on hydrophobic coverslips.

Conidia were harvested from a strain co-expressing Ppe1-mCherry and Sep3-GFP and inoculated on hydrophobic coverslips and observed at each time point. Scale bars = 1 μ m.




Fig. S6

3D images of the co-expression of Ppe1-mCherry and Lifeact-GFP during primary invasive hyphal development.

AP, Appressorium. Arrow shows penetration ring.




Fig. S7

Co-expression of Ppe1-mCherry with cytoplasmic GFP at the early infection stage of *M. oryzae*.

(A) Ppe1-mCherry signal formed a tight ring-like structure closely associated with the base of the appressorium at 16 hpi. Scale bar = 5 μ m. (B) At 23 hpi, Ppe1-mCherry ring signal was still beneath the appressorium, and around the penetration peg which differentiated into the primary invasive hyphae. Scale bar = 10 μ m. AP, Appressorium. IH, invasive hyphae. hpi, hours post-inoculation. Arrows show penetration ring.




Fig. S8

The spatial positioning of Ppe1-mCherry ring and rice plasma membrane (marked by GFP-OsPIP2).

The zoomed images represent the sites pointed by the white arrows, respectively. Scale bar = 10 μm ; 2 μm for the zoomed images.




Fig. S9

Ppe1 ring was also observed during transpressorium-mediated peg penetration.

(A-D) The spatiotemporal dynamics of Ppe1-mCherry and Sep5-GFP during transpressorium-mediated cell-to-cell penetration. Red squares indicate the zoomed regions, respectively. Scale bars = 10 μm.

Strain	Description	Reference
70-15	Wild-type strain	(41)
mCherry _{PPE1 promoter}	pKNT-mCherry _{PPE1 promoter} transformant of 70-15	this study
Ppe1-mCherry	pKNT-MoPpe1-mCherry and pCB1532 co-transformant of 70-15	this study
Ppe1-GFP	pKNT-MoPpe1-GFP and pCB1532 co-transformant of 70-15	this study
Ppe1-mCherry _{situ insertion}	ATMT1-Ppe1-mCherry _{situ insertion} transformant of 70-15	this study
Sep3-GFP	Sep3-GFP transformant of Guy11	Provided by Prof. Guo Min
Sep5-GFP	Sep3-GFP transformant of Guy11	Provided by Prof. Guo Min
Ppe1-mCherry + Sep3-GFP	pKNT-Ppe1-mCherry transformant of Sep3-GFP strain	this study
Ppe1-mCherry + Sep5-GFP	pKNT-Ppe1-mCherry transformant of Sep5-GFP strain	this study
Ppe1-mCherry + Lifeact-GFP	pXY198-Lifeact-GFP transformant of Ppe1-mCherry strain	this study
Ppe1-mCherry + GFP	pKNT-RP27-GFP transformant of Ppe1-mCherry strain	this study
$\Delta ppe1-45/101$	<i>PPE1</i> deletion mutant of 70-15	this study
$\Delta ppe1$ -com-3/8	pKNT-Ppe1-mCherry transformant of $\Delta ppe1-45$ mutant	this study
$\Delta mst12$	<i>MST12</i> deletion mutant of Guy11	(27)
$\Delta nox1$	<i>NOX1</i> deletion mutant of Ppe1-mCherry strain	this study
$\Delta nox2$	<i>NOX2</i> deletion mutant of Ppe1-mCherry strain	this study
Ppe2-mCherry	pKNT-MGG_10732-mCherry transformant of 70-15	this study
Ppe3-mCherry	pKNT-MGG_16041-mCherry transformant of 70-15	this study
Ppe5-mCherry	pKNT-MGG_16379-mCherry transformant of	this study

Supplementary Table S1.

Fungal strains used in this study

	70-15	
MoTPpe1-mCherry	pKNT-MoTPpe1-mCherry transformant of 70-15	this study
PBV591	pBV591(Co-expression of Pwl2-mCherry and Bas4-GFP) transformant of 70-15	(16)
Slp1-GFP	pKNT-Slp1-mCherry transformant of 70-15	this study
AvrPiz-t-GFP	pKNT-AvrPiz-t-GFP transformant of 70-15	this study

Supplementary Table S1. (continued)

Primer Name	Primer Sequence (5'→3')
Construction of pSUC2 plasmid for confirming the secretory function of signal peptide (SP)	
pSUC2-Ppe1SP-F	TTTTAATTAAGAATTCATGCGTTCATCTACGATTATTCTG
pSUC2-Ppe1SP-R	CTATAGGGAGAACCTCGAGGGCGGCAACCAGGCCAGTG
pSUC2-Pwl2SP-F	TTTTAATTAAGAATTCATGAAATGCAACAACATCATCCTC
pSUC2-Pwl2SP-R	CTATAGGGAGAACCTCGAGGGCGGTTACAGTGGTCG
pSUC2-yF	ACAGATGTCGTTGTTCCAGA
Construction of plasmid for studying subcellular localization	
pKNT-Ppe1-promoter-F	AGGGAACAAAAGCTGGGTACCTCTTTCTCCCAGTCTCATC
pKNT-Ppe1-promoter-R	GCCCTTGCTCACCATAAGCTTGTTGACGCCAGGTCGTGAGA
pKNT-Ppe1-R	GCCCTTGCTCACCATAAGCTTTGCACATCCAATAAATCCAGCATGC
ATMT-mCherry _{situ} insertion -	GAATTCGAGCTCGGTACCGGCAACGACGACAAGAA

Supplementary Table S2.

List of Primers used in this study

AF	
ATMT-mCherry _{situ} insertion ⁻	GTCTGACTCTAGAGGATCCATTGCACATCCAATAAAATCCAGC
AR	
mCherry _{situ} insertion ^{-F}	ATGGTGAGCAAGGGCGAGGAG
mCherry _{situ} insertion ^{-R}	TTGACCTCCACTAGCTCCAGCCAAGCCCTACTTGTACAGCTCGTCC ATGCC
HYG/F	GGCTTGGCTGGAGCTAGTGGAGGTCAA
HYG/R	AACCCGCGGTCTGGCATCTACTCTATTC
ATMT-mCherry _{situ} insertion ⁻	GAATAGAGTAGATGCCGACCGCGGGTTGTTGAGCGGAATTGCCTG
BF	TTAT
ATMT-mCherry _{situ} insertion ⁻	GTCTGACTCTAGAGGATCCTTCGCTTGTTGTTCTTCG
BR	
pKNT-AvrPiz-t-M-F	AGGGAACAAAAGCTGGGTACCCCAACGTAGAGCGTATGCTGT
pKNT-AvrPiz-t-M-R	GCCCTTGCTCACCATAAGCTTTTGCGCTGAGCCTGAG
pCB1532-Slp1-G-F	ACTATAGGGCGAATTGGGTACCATTGAACAGTGCGATGCC
pCB1532-Slp1-G-R	CGCCCTTGCTCACCATAAGCTTGTTCTTGCAGATGGGGATG
GFP-sR	AGTTCACCTTGATGCCGTTC
mCherry-sR	TGGTCTTCTTCTGCATTACGG
pKNT-MoTPpe1-M-F	AGGGAACAAAAGCTGGGTACCTCTCTCAGTCTCATTCTACAGC
pKNT-MoTPpe1-M-R	GCCCTTGCTCACCATAAGCTTTGCACATCCAATAAAATCCAGC
pKNT-MGG_10732-M-F	AGGGAACAAAAGCTGGGTACCGCACGTACCTTTGGTGGTTGC
pKNT-MGG_10732-M-R	GCCCTTGCTCACCATAAGCTTAGCACATCCAACATGTCCAGG
pFGL1010M-MGG_16041-M-F	CATGATGATGCTCGAGAATTCACGCCGCTTTCTGGACTA
pFGL1010M-MGG_16041-M-R	CTCGCCCTTGCTCACGGTACCGACGCATCCAACAAACCCAGGAA
pFGL1010M-MGG_16379-M-F	CATGATGATGCTCGAGAATTCTATCGTTTCGCGGGTAGATTG
pFGL1010M-MGG_16379-M-R	CTCGCCCTTGCTCACGGTACCCTTTGCGCACCCAATGAATGC
Construction of gene deletion	
ATMT-Ppe1-AF	GAATTCGAGCTCGGTACCCACCTTTTCGTCCTCC
ATMT-Ppe1-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTGACGCCAGGTCTGTGAG

Supplementary Table S2. (continued)

ATMT-Ppe1-BF	GAATAGAGTAGATGCCGACCGCGGGTTAGCGGAATTGCCTGTTA
ATMT-Ppe1-BR	GTCGACTCTAGAGGATCCACTGCCGACCGTGACTA
Ppe1-OF	ATGCGTTCATCTACGATTATTCTG
Ppe1-OR	TGCACATCCAATAAATCCAG
Ppe1-UA	CATCATCGCCGTCCTCT
HYG/F	GGCTTGGCTGGAGCTAGTGGAGGTCAA
HY/R	GTATTGACCGATTCCCTGCGGTCCGAA
YG/F	GATGTAGGAGGGCGTGATATGTCCT
HYG/R	AACCCGCGGTCGGCATCTACTCTATTC
H853	GACAGACGTCGCGGTGAGTT
Nox1-AF	CACCAGATGCAGTAGTCAATT
Nox1-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTTCGTTAAAGAGCGTATG TC
Nox1-BF	GAATAGAGTAGATGCCGACCGCGGGTTGCCGAAATCTATTACGAC G
Nox1-BR	CCTGAATACTTCCAAGACACC
Nox1-OF	TCACCACCTGTTTATCCCG
Nox1-OR	GACACCAATACCAGTTCCAAT
Nox1-UA	CAGAGCCACGATGCGAAAA
Nox2-AF	CGATTGTGCCTGGATGA
Nox2-AR	TTGACCTCCACTAGCTCCAGCCAAGCCGTTGCTTGGTCGGTTGTC
Nox2-BF	GAATAGAGTAGATGCCGACCGCGGGTTGAAGCAAGCCTGGCAATA
Nox2-BR	CAAAAGCCTGTTTACGAGAT
Nox2-OF	AAGATGGACGCCGCTAA
Nox2-OR	AAACGATGAAGAAGTGGTGA
Nox2-UA	ACCTCCTTGACTCTTCCACC
qRT-PCR analysis	
Ppe1-qF	TTTGCTGCGACGCTGGTA
Ppe1-qR	GGAAAATCTGGGAATGGGT
Tublin-qF	TGACAACGAGGCTCTGTACG
Tublin-qR	CCATGAAGAAGTGCAGACGA

Supplementary Table S2. (continued)

References

1. Talbot NJ (2003) **On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*** *Annual review of microbiology* **57**:177–202
2. Dean R, et al. (2012) **The Top 10 fungal pathogens in molecular plant pathology** *Molecular plant pathology* **13**:414–430
3. Couch BC, et al. (2005) **Origins of host-specific populations of the blast pathogen *Magnaporthe oryzae* in crop domestication with subsequent expansion of pandemic clones on rice and weeds of rice** *Genetics* **170**:613–630
4. Langner T, Białas A, Kamoun S (2018) **The Blast Fungus Decoded: Genomes in Flux** *mBio* **9**:2
5. Ceresini PC, et al. (2018) **Wheat Blast: Past, Present, and Future** *Annual review of phytopathology* **56**:427–456
6. Malaker PK, et al. (2016) **First Report of Wheat Blast Caused by *Magnaporthe oryzae* Pathotype triticum in Bangladesh** *Plant Disease* **100**:2330–2330
7. Howard RJ, Valent B (1996) **Breaking and entering: host penetration by the fungal rice blast pathogen *Magnaporthe grisea*** *Annual review of microbiology* **50**:491–512
8. Talbot NJ (2019) **Appressoria** *Current biology: CB* **29**:R144–r146
9. Hamer JE, Howard RJ, Chumley FG, Valent B (1988) **A mechanism for surface attachment in spores of a plant pathogenic fungus** *Science (New York, N.y.)* **239**:288–290
10. Rocha RO, Elowsky C, Pham NTT, Wilson RA (2020) **Spermine-mediated tight sealing of the *Magnaporthe oryzae* appressorial pore-rice leaf surface interface** *Nature microbiology* **5**:1472–1480
11. Chumley FG, Valent BJMP-mI (1990) **Genetic analysis of melanin-deficient, nonpathogenic mutants of *Magnaporthe grisea*** *Molecular plant-microbe interactions* **3**:135–143
12. de Jong JC, McCormack BJ, Smirnov N, Talbot NJ (1997) **Glycerol generates turgor in rice blast** *Nature* **389**:244–244
13. Dagdas YF, et al. (2012) **Septin-mediated plant cell invasion by the rice blast fungus, *Magnaporthe oryzae*** *Science (New York, N.y.)* **336**:1590–1595
14. Fernandez J, Orth K (2018) **Rise of a Cereal Killer: The Biology of *Magnaporthe oryzae* Biotrophic Growth** *Trends in microbiology* **26**:582–597
15. Kankanala P, Czymmek K, Valent B (2007) **Roles for rice membrane dynamics and plasmodesmata during biotrophic invasion by the blast fungus** *The Plant cell* **19**:706–724
16. Khang CH, et al. (2010) **Translocation of *Magnaporthe oryzae* effectors into rice cells and their subsequent cell-to-cell movement** *The Plant cell* **22**:1388–1403

17. Cruz-Mireles N, Eseola AB, Osés-Ruiz M, Ryder LS, Talbot NJ (2021) **From appressorium to transpressorium-Defining the morphogenetic basis of host cell invasion by the rice blast fungus** *PLoS pathogens* **17**:e1009779
18. Ryder LS, et al. (2022) **The appressorium at a glance** *Journal of cell science* **135**:14
19. Giraldo MC, et al. (2013) **Two distinct secretion systems facilitate tissue invasion by the rice blast fungus *Magnaporthe oryzae*** *Nature communications* **4**:1996
20. Ebbole DJ, et al. (2021) **Evolution and Regulation of a Large Effector Family of *Pyricularia oryzae*** *Molecular plant-microbe interactions: MPMI* **34**:255–269
21. Yan X, et al. (2023) **The transcriptional landscape of plant infection by the rice blast fungus *Magnaporthe oryzae* reveals distinct families of temporally co-regulated and structurally conserved effectors** *The Plant cell* **35**:1360–1385
22. Eisermann I, Talbot NJ (2024) **Septin-dependent invasive growth by the rice blast fungus *Magnaporthe oryzae*** *Journal of Plant Diseases and Protection*
23. Bourett TM, Howard RJ (1992) **Actin in penetration pegs of the fungal rice blast pathogen, *Magnaporthe grisea*** *Protoplasma* **168**:20–26
24. Osés-Ruiz M, et al. (2021) **Appressorium-mediated plant infection by *Magnaporthe oryzae* is regulated by a Pmk1-dependent hierarchical transcriptional network** *Nature microbiology* **6**:1383–1397
25. Chen Z, et al. (2019) **Green Fluorescent Protein- and Discosoma sp. Red Fluorescent Protein-Tagged Organelle Marker Lines for Protein Subcellular Localization in Rice** *Frontiers in plant science* **10**:1421
26. Sakulkoo W, et al. (2018) **A single fungal MAP kinase controls plant cell-to-cell invasion by the rice blast fungus.** *Science (New York, N.y.)* **359**:1399–1403
27. Park G, Xue C, Zheng L, Lam S, Xu JR (2002) **MST12 regulates infectious growth but not appressorium formation in the rice blast fungus *Magnaporthe grisea*** *Molecular plant-microbe interactions: MPMI* **15**:183–192
28. Ryder LS, et al. (2013) **NADPH oxidases regulate septin-mediated cytoskeletal remodeling during plant infection by the rice blast fungus** *Proceedings of the National Academy of Sciences of the United States of America* **110**:3179–3184
29. Mentlak TA, et al. (2012) **Effector-mediated suppression of chitin-triggered immunity by *magnaporthe oryzae* is necessary for rice blast disease** *The Plant cell* **24**:322–335
30. Park CH, et al. (2012) **The *Magnaporthe oryzae* effector AvrPiz-t targets the RING E3 ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice** *The Plant cell* **24**:4748–4762
31. Ludwig N, et al. (2021) **A cell surface-exposed protein complex with an essential virulence function in *Ustilago maydis*** *Nature microbiology* **6**:722–730
32. Talbot NJ, Ebbole DJ, Hamer JE (1993) **Identification and characterization of MPG1, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*** *The Plant cell* **5**:1575–1590

33. Zhong Z, et al. (2018) **Population genomic analysis of the rice blast fungus reveals specific events associated with expansion of three main clades** *The ISME journal* **12**:1867–1878
 34. Chen X, et al. (2021) **Trehalose Phosphate Synthase Complex-Mediated Regulation of Trehalose 6-Phosphate Homeostasis Is Critical for Development and Pathogenesis in *Magnaporthe oryzae*** *mSystems* **6**:e0046221
 35. Khang CH, Park SY, Rho HS, Lee YH, Kang S (2006) **Filamentous Fungi (*Magnaporthe grisea* and *Fusarium oxysporum*). Methods in molecular biology (Clifton N.j** **344**:403–420
 36. Jiang C, et al. (2020) **An orphan protein of *Fusarium graminearum* modulates host immunity by mediating proteasomal degradation of TaSnRK1α** *Nature communications* **11**:4382
 37. Oh SK, et al. (2009) **In planta expression screens of *Phytophthora infestans* RXLR effectors reveal diverse phenotypes, including activation of the *Solanum bulbocastanum* disease resistance protein Rpi-blb2** *The Plant cell* **21**:2928–2947
 38. Kershaw MJ, Talbot NJ (2009) **Genome-wide functional analysis reveals that infection-associated fungal autophagy is necessary for rice blast disease** *Proceedings of the National Academy of Sciences of the United States of America* **106**:15967–15972
 39. Holsters M, et al. (1978) **Transfection and transformation of *Agrobacterium tumefaciens*** *Molecular & general genetics: MGG* **163**:181–187
 40. Livak KJ, Schmittgen TD (2001) **Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.** *Methods (San Diego Calif* **25**:402–408
 41. Dean RA, et al. (2005) **The genome sequence of the rice blast fungus *Magnaporthe grisea*** *Nature* **434**:980–986
- Dean R. A., Talbot N. J., Ebbole D. J., Farman M. L., Mitchell T. K., Orbach M. J., Thon M., Kulkarni R., Xu J. R., Pan H., Read N. D., Lee Y. H., Carbone I., Brown D., Oh Y. Y., Donofrio N., Jeong J. S., Soanes D. M., Djonovic S., Kolomiets E., Rehmeier C., Li W., Harding M., Kim S., Lebrun M. H., Bohnert H., Coughlan S., Butler J., Calvo S., Ma L. J., Nicol R., Purcell S., Nusbaum C., Galagan J. E., Birren B. W (2005) **The genome sequence of the rice blast fungus *Magnaporthe grisea*** *Nature* **434**:980
- Khang C. H., Berruyer R., Giraldo M. C., Kankanala P., Park S. Y., Czymmek K., Kang S., Valent B (2010) **Translocation of *Magnaporthe oryzae* Effectors into Rice Cells and Their Subsequent Cell-to-Cell Movement** *The Plant cell* **22**:1388
- Park G., Xue C., Zheng L., Lam S., Xu J. R (2002) **MST12 Regulates Infectious Growth But Not Appressorium Formation in the Rice Blast Fungus *Magnaporthe grisea*** *Molecular plant-microbe interactions: MPMI* **15**:183

Author information

Wenqin Fang

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Fuzhou Institute of Oceanography, Minjiang University, Fuzhou, China

Xiaoyu Zai

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

Jia Chen

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

Yakubu Saddeeq Abubakar

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Department of Biochemistry, Faculty of Life Sciences, Ahmadu Bello University, Zaria, Nigeria

Qiuqiu Wu

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

Zhenyu Fang

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

Xiuwei Huang

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

Xiang Gan

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

Daniel J Ebbole

Department of Plant Pathology & Microbiology, Texas A&M University, College Station, United States

Zonghua Wang

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Fuzhou Institute of Oceanography, Minjiang University, Fuzhou, China

For correspondence: wangzh@fafu.edu.cn

Wenhui Zheng

State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China, Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou, China

ORCID iD: [0009-0000-2767-3617](https://orcid.org/0009-0000-2767-3617)

For correspondence: wenhuiz@fafu.edu.cn

Editors

Reviewing Editor

Thabiso Motaung

University of Pretoria, Pretoria, South Africa

Senior Editor

Jürgen Kleine-Vehn

University of Freiburg, Freiburg, Germany

Joint Public Review:

This study presents novel insights into the formation and characterization of a penetration ring during host infection by *Magnaporthe oryzae*. Based on the solid genetic evidence and localization data, the authors demonstrate the structural presence of the penetration ring and the contribution of Ppe1 to fungal virulence. Nevertheless, the mechanisms through which the penetration ring influences host-pathogen interaction, including its potential function in effector translocation, remain only partially resolved. Further work using higher-resolution imaging and functional assays will help address this knowledge gap. Overall, the findings are valuable for advancing our understanding of plant-pathogen interactions, though important mechanistic questions remain open.

<https://doi.org/10.7554/eLife.101209.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

Summary:

This study focuses on characterizing a previously identified gene, encoding the secreted protein Ppe1, that may play a role in rice infection by the blast fungus Magnaporthe oryzae. Magnaporthe oryzae is a hemibiotrophic fungus that infects living host cells before causing disease. Infection begins with the development of a specialized infection cell, the appressorium, on the host leaf surface. The appressorium generates enormous internal turgor that acts on a thin penetration peg at the appressorial base, forcing it through the leaf cuticle. Once through this barrier, the peg elaborates into bulbous invasive hyphae that colonizes the first infected cell before moving to neighboring cells via plasmodesmata. During this initial biotrophic growth stage, invasive hyphae invaginate the host plasma membrane, which surrounds growing hyphae as the extra-invasive hyphae membrane (EIHM). To avoid detection, the fungus secretes apoplastic effectors into the EIHM matrix via the conventional ER-Golgi secretion pathway. The fungus also forms a plant-derived structure called the biotrophic interfacial complex (BIC) that receives cytoplasmic effectors through an unconventional secretion route before they are delivered into the host cell. Together, these secreted effector proteins act to evade or suppress host innate immune responses. Here the authors contribute to our understanding of M. oryzae infection biology by showing how Ppe1, which localizes to both the appressorial penetration peg and to the appressorial-like transpressoria associated with invasive hyphal movements into adjacent cells, maximizes host cell penetration and disease development and is thus a novel contributor to rice blast disease.

We sincerely appreciate the reviewer's thoughtful evaluation of our work. We are grateful for your recognition of Ppe1 as a novel contributor to *M. oryzae* infection biology and your insightful summary of its spatio-temporal localization and functional importance in host penetration. We also appreciate devoting your time to provide us with constructive feedback, which greatly strengthens our manuscript.

Strengths:

A major goal of M. oryzae research is to understand how the fungus causes disease, either by determining the physiological underpinnings of the fungal infection cycle or by identifying effectors and their host targets. Such new knowledge may point the way to novel mitigation strategies. Here, the authors make an interesting discovery that bridges both fungal physiology and effector biology research by showing how a secreted protein Ppe1, initially considered an effector with potential host targets, associates with its own penetration peg (and transpressoria) to facilitate host invasion. In a previous study, the authors had identified a small family of small secreted proteins that may function as effectors. Here they suggest Ppe1 (and, later in the manuscript, Ppe2/3/5) localizes outside the penetration peg when appressoria develops on surfaces that permit penetration, but not on artificial hard surfaces that prevent peg penetration. Deleting the PPE1 gene reduced (although did not abolish) penetration, and a fraction of those that penetrated developed invasive hyphae that were reduced in growth compared to WT. Using fluorescent markers, the authors show that Ppe1 forms a ring underneath appressoria, likely where the peg emerges, which remained after invasive hyphae had developed. The ring structure is smaller than the width of the appressorium and also lies within the septin ring known to form during peg development. This so-called penetration ring also formed at the transpressorial penetration point as invasive hyphae moved to adjacent cells. This structure is novel, and required for optimum penetration during

infection. Furthermore, Ppe1, which carries a functional signal peptide, may form on the periphery of the peg, together suggesting it is secreted and associated with the peg to facilitate penetration. Staining with aniline blue also suggests Ppe1 is outside the peg. Together, the strength of the work lies in identifying a novel appressorial penetration ring structure required for full virulence.

We are deeply grateful to the reviewer for the clear understanding and insightful evaluation of our work. Your recognition of the novel contribution and scientific merit of our study is both encouraging and motivating. We sincerely appreciate the time, expertise and constructive feedback dedicated to reviewing our manuscript, as the comments have been instrumental in enhancing the quality of this work.

Weaknesses:


The main weakness of the paper is that, although Ppe1 is associated with the peg and optimizes penetration, the function of Ppe1 is not known. The work starts off considering Ppe1 a secreted effector, then a facilitator of penetration by associating with the peg, but what role it plays here is only often speculated about. For example, the authors consider at various times that it may have a structural role, a signaling role orchestrating invasive hyphae development, or a tethering role between the peg and the invaginated host plasma membrane (called throughout the host cytoplasmic membrane, a novel term that is not explained). However, more effort should be expended to determine which of these alternative roles is the most likely. Otherwise, as it stands, the paper describes an interesting phenomenon (the appressorial ring) but provides no understanding of its function.

We sincerely appreciate the reviewer's comments. We have revised "host cytoplasmic membrane" to "host plasma membrane" throughout the manuscript for consistency. To further investigate the role of the Ppe1 in the interaction between *M. oryzae* and rice, we overexpressed *PPE1* in rice ZH11. A pCXUN-SP-GFP-Ppe1 vector containing a signal peptide and an N-terminal GFP tag was constructed (pCXUN-SP-GFP-Ppe1), and 35 GFP-PPE1-OX plants (T0) were subsequently obtained through *Agrobacterium*-mediated rice transformation. Subsequently, PCR and qRT-PCR validation were performed on the T0 transgenic plants. The PCR results showed that the inserted plasmid could be amplified from the genomic DNA extracted from the leaves of all the resulting T0 plants (Author response image 1A). qRT-PCR results indicated that most T0 transgenic plants could transcriptionally express *PPE1* (Author response image 1B). T0 plants with higher expression levels were selected for western blot analysis, which confirmed the presence of GFP-Ppe1 bands of the expected size (Author response image 1C). To further explore the targets of Ppe1 in rice, the leaf sheaths of T0 plants were inoculated with *M. oryzae* strain Guy11. Total proteins were extracted at 24 hours post-inoculation (hpi) and subjected to immunoprecipitation using GFP magnetic beads. Silver staining revealed more interacting protein bands in T0 plants compared to ZH11 and GFP-OX controls (Author response image 1D). These samples were then analyzed by mass spectrometry in which 331 rice proteins that potentially interact with Ppe1 were identified (Author response image 1E). Subsequently, yeast two-hybrid assays were performed on 13 putative interacting proteins with higher coverage, but no interaction was detected between Ppe1 and these proteins (Author response image 1F-G). Considering that the identification and functional validation of interacting proteins is a labor-intensive and time-consuming endeavor, we will focus our future efforts on in-depth studies of Ppe1's function in rice.

Author response image 1.

Screening of Ppe1 candidate targets in rice. (A) The determination of *GFP-PPE1* construct in transgenic rice. (B) The expression of *PPE1* transgenic rice (T0) was verified by qRT-PCR. (C)

Western blot analysis of Ppe1 expression in transgenic rice. (D) Rapid silver staining for detection of the purified proteins captured by the GFP-beads. (E) Venn diagram comparing the number of proteins captured in the different samples. (F) Identity of the potential targets of Ppe1 in rice. (G) Yeast two-hybrid assay showing negative interaction of Ppe1 with rice candidate proteins.




The inability to nail down the function of Ppe1 likely stems from two underlying assumptions with weak support. Firstly, the authors assume that Ppe1 is secreted and associated with the peg to form a penetration ring between the plant cell wall and cytoplasm membrane. However, the authors do not demonstrate it is secreted (for instance by blocking Ppe1 secretion and its association with the peg using brefeldin A).

To investigate the secretion pathway of Ppe1 in *M. oryzae*, we determined the inhibitory effects of Brefeldin A (BFA) on conventional ER-to-Golgi secretion in fungi as suggested by the reviewer. We inoculated rice leaf sheaths with conidia suspensions from the Ppe1-mCherry and PBV591 strains (containing a Pwl2-mCherry-NLS and Bas4-GFP co-expressing constructs) and treated them with BFA. We found that, even after exposure to BFA for 5 to 11 hours, the Ppe1-mCherry still formed its characteristic ring conformation (Author response image 2). Similarly, in the BFA-treated samples, the cytoplasmic effector Pwl2-mCherry accumulated at the BIC, while the apoplastic effector Bas4-GFP was retained in the invasive hyphae (Author response image 2). These results indicate that Ppe1 is not secreted through the conventional ER-Golgi secretion pathway.

Author response image 2.

The secretion of Ppe1 is not affected by BFA treatment. (A) and (B) The Ppe1-mCherry fluorescent signal was still observed both in the presence and absence of BFA. (C) Following BFA treatment, the secretion of the apoplastic effector Bas4-GFP was blocked while that of the cytoplasmic effector Pwl2-mCherry was not affected. The rice leaf sheath tissue was inoculated with 50 µg/mL BFA (0.1% DMSO) at 17 hpi. Images were captured at 22 hpi for A and 28 hpi for B and C. Scale bars = 10 µm.



Also, they do not sufficiently show that Ppe1 localizes on the periphery of the peg. This is because confocal microscopy is not powerful enough to see the peg. The association they are seeing (for example in Figure 4) shows localization to the bottom of the appressorium and around the primary hyphae, but the peg cannot be seen. Here, the authors will need to use SEM, perhaps in conjunction with gold labeling of Ppe1, to show it is associating with the peg and, indeed, is external to the peg (rather than internal, as a structural role in peg rigidity might predict). It would also be interesting to repeat the microscopy in Figure 4C but at much earlier time points, just as the peg is penetrating but before invasive hyphae have developed - Where is Ppe1 then? Finally, the authors speculate, but do not show, that Ppe1 anchors penetration pegs on the plant cytoplasm membrane. Doing so may require FM4-64 staining, as used in Figure 2 of Kankanala et al, 2007 (DOI: 10.1105/tpc.106.046300), to show connections between Ppe1 and host membranes. Note that the authors also do not show that the penetration ring is a platform for effector delivery, as speculated in the Discussion.


We sincerely appreciate the reviewer's valuable suggestion regarding SEM with immunogold labeling to precisely visualize Ppe1's association with penetration peg. While we fully acknowledge this would be an excellent approach, after consulting several experts in the field, we realized that the specialized equipment and technical expertise required for fungal immunogold-SEM are currently unavailable to us. We sincerely hope that the reviewer will understand this technical limitation.

To further strengthen our evidence for the role of Ppe1's in anchoring penetration peg to the plant plasma membrane, we provided new co-localization images of Ppe1 and penetration peg (Fig. S7). At 16 hours post-inoculation (hpi), when the penetration peg was just forming and prior to the development of invasive hyphae, the Ppe1-mCherry fluorescence forms a tight ring-like structure closely associated with the base of the appressorium. As at 23 hpi, the circular Ppe1-mCherry signal was still detectable beneath the appressorium, and around the penetration peg which differentiated into the primary invasive hyphae. Furthermore, we obtained 3D images of the strain expressing both Ppe1-mCherry and Lifeact-GFP during primary invasive hyphal development. The results revealed that Ppe1 forms a ring-like structure that remains anchored to the penetration peg during fungal invasion (Fig. S6).

We also conducted FM4-64 staining experiment as recommended by the reviewer. Although the experiment provided valuable insights, we found that the resolution was insufficient to precisely delineate the spatial relationship between Ppe1 and host membranes at the penetration peg (Author response image 3). To optimize this colocalization, we tested the localization between Ppe1-mCherry ring and rice plasma membrane marker GFP-OsPIP2 (Fig. S8). These new results provide compelling complementary evidence supporting our conclusion that Ppe1 functions extracellularly at the host-pathogen interface. We hope these additional data will help address the reviewer's concerns regarding Ppe1's localization.

Author response image 3.

FM4-64-stained rice leaf sheath inoculated with *M. oryzae* strain expressing Ppe1-GFP. Ppe1-GFP ring was positioned above the primary invasive hyphae. Scale bar = 5 μ m.



Secondly, the authors assume Ppe1 is required for host infection due to its association with the peg. However, its role in infection is minor. The majority of appressoria produced by the mutant strain penetrate host cells and elaborate invasive hyphae, and lesion sizes are only marginally reduced compared to WT (in fact, the lesion density of the 70-15 WT strain itself seems reduced compared to what would be expected from this strain). The authors did not analyze the lesions for spores to confirm that the mutant strains were non-pathogenic (non-pathogenic mutants sometimes form small pinprick-like lesions that do not sporulate). Thus, the pathogenicity phenotype of the knockout

mutant is weak, which could contribute to the inability to accurately define the molecular and cellular function of Ppe1.

We appreciate the reviewer's comments. To ensure the reliability of our findings, we conducted spray inoculation experiments with multiple independent repeats. Our results consistently demonstrated that deletion of the *PPE1* gene significantly attenuates the virulence of *M. oryzae*. Further analysis of lesion development and sporulation in the $\Delta ppe1$ mutant revealed that it retains the ability to produce conidia. To validate these observations, we generated a *PPE1* knockout in the wild-type reference strain Guy11. Similarly, we observed a significant decrease in the pathogenicity of the $\Delta ppe1$ mutants generated from the wild-type Guy11 strain compared to Guy11 in the spray assay (Fig S2). These results collectively indicate the importance of Ppe1 in the pathogenicity of *M. oryzae* to rice.

In summary, it is important that the role of Ppe1 in infection be determined.

Reviewer #2 (Public review):

The article focuses on the study of Magnaporthe oryzae, the fungal pathogen responsible for rice blast disease, which poses a significant threat to global food security. The research delves into the infection mechanisms of the pathogen, particularly the role of penetration pegs and the formation of a penetration ring in the invasion process. The study highlights the persistent localization of Ppe1 and its homologs to the penetration ring, suggesting its function as a structural feature that facilitates the transition of penetration pegs into invasive hyphae. The article provides a thorough examination of the infection process of M. oryzae, from the attachment of conidia to the development of appressoria and the formation of invasive hyphae. The discovery of the penetration ring as a structural element that aids in the invasion process is a significant contribution to the understanding of plant-pathogen interactions. The experimental methods are well-documented, allowing for reproducibility and validation of the results.

We sincerely appreciate the thoughtful and insightful evaluation of our work. Thank you for recognizing the significance of our findings regarding the penetration ring and the functional role of Ppe1 during host invasion.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

Line 48: "after appressorium- or transpressorium-mediated penetration of plant cell wall" - transpressoria do not penetrate the plant cell wall.

Thank you for your valuable suggestion. For improved clarity, we have rephrased the sentence as follows: In this study, we showed that a penetration ring is formed by penetration pegs after appressorium-mediated penetration of plant cell wall.

Line 143: "approximately 25% of the 143 appressoria formed by the $\Delta ppe1$ mutant had no penetration peg" - It is not possible to see the penetration peg by confocal microscopy.

Thank you for your valuable suggestion. We have revised the sentence as follows: In contrast, approximately 25% of the appressoria formed by the $\Delta ppe1$ mutant had no penetration.

Line 159: "inner cycle" -should be inner circle?

We gratefully acknowledge the reviewer's careful reading. The typographical error has been corrected throughout the revised manuscript.

Line 255: "These results indicate that initiation of penetration peg formation is necessary for the formation of the penetration ring." Actually, more precisely, they indicate that penetration is necessary.

We appreciate this suggestion and have revised the text to be more concise: These results indicate that penetration is necessary for the formation of the penetration ring.

Line 282: "unlike subcellular localizations of other effectors"- is this an effector if no plant targets are known?

We appreciate this suggestion and have revised the text as follows: unlike subcellular localizations of Bas4, Slp1, Pwl2, and AvrPiz-t.

Line 299: "it may function as a novel physical structure for anchoring penetration pegs on the surface of plant cytoplasm membrane after cell wall penetration" - an interaction with the plant plasma membrane was not shown and this is speculative.

We have provided new evidence to show the spatial positioning of Ppe1-mCherry ring with the rice plasma membrane (see figure S8)

Line 301: "It is also possible that this penetration ring functions as a collar or landmark that is associated with the differentiation of penetration pegs (on the surface of cytoplasm membrane) into primary invasive hyphae enveloped in the EIHM cytoplasm membrane (Figure 7)." The alternative conclusions for Ppe1 function, either interacting with host membranes or acting as a developmental landmark, need to be resolved here.

We appreciate this suggestion and have revised the text as follows: It is also possible that this penetration ring functions as a collar that is associated with the differentiation of penetration pegs into primary invasive hyphae enveloped in the EIHM (Figure 7).

Line 317: "is likely a structural feature or component for signaling the transition of penetration pegs to invasive hyphae",- if the authors think Ppe1 has these roles, why do they refer to Ppe1 as an effector?

Many thanks for these comments. We have revised this and refer to Ppe1 as a secreted protein throughout the revised manuscript.

Line 337: "After the penetration of plant cell wall, the penetration ring may not only function as a physical structure but also serve as an initial effector secretion site for the release of specific effectors to overcome plant immunity in early infection stages"- which is it? Also, no evidence is provided to suggest it is a platform for effector secretion.

We sincerely appreciate your valuable suggestion. We have revised this sentence as follows: After the penetration of plant cell wall, the penetration ring may not only function as a physical structure but also serve as a secretion site for the release of specific proteins to overcome plant immunity during the early infection stages.

Reviewer #2 (Recommendations for the authors):


(1) While the study suggests the penetration ring as a structural feature, it remains unclear whether it also serves as a secretion site for effectors. Further exploration of this

aspect would strengthen the conclusions.

We thank the reviewer for this useful suggestion. In this study, we demonstrated that Ppe1 proteins form a distinct penetration ring structure at the site where the penetration peg contacts the plant plasma membrane prior to differentiation into primary invasive hyphae (Figs. 2 and 7). Thus, we reasoned that penetration ring may function as a novel physical structure. Notably, additional Ppe family members (Ppe2, Ppe3, and Ppe5) were also found to localize to this penetration ring (Fig. 6B), suggesting that it also serves as a secretion site for releasing proteins. To test whether Ppe1 and Ppe2 label to the same site, we analyzed the colocalization between Ppe1-GFP and Ppe2-mCherry. The results showed that Ppe1-GFP and Ppe2-mCherry are well colocalized (Author response image 4). This study primarily focuses on the discovery and characterization of the penetration ring. The potential role of this structure in effector translocation will be investigated in future studies.

Author response image 4.

Ppe1 co-localizes with Ppe2 at the penetration ring in *M. oryzae*. Line graphs were generated at the directions pointed by the white arrows. Scale bar = 2 μ m.



(2) The article could benefit from a discussion on the broader implications of these findings for developing resistant crop varieties or new fungicidal strategies.

We have incorporated this discussion as suggested (lines 358-360).

(3) *What is the significance of the formation of the penetration ring in the pathogenicity of the rice blast fungus? Or, how does it assist the fungus in its infection process?*

Our findings have several significant implications. First, we believe that the discovery of the penetration ring as a novel physical structure associated with the differentiation of invasive hyphae represents a breakthrough in plant-pathogen interactions that will be of interest to fungal biologists, pathologists and plant biologists. Secondly, our study presents new role of the peg as a specialized platform for secretory protein deployment, in addition to its commonly known role as a physical penetration tool for the pathogen. Thirdly, we identify Ppe1 as a potential molecular target for controlling the devastating rice blast disease, as Ppe homologs are absent in plants and mammals. We have incorporated this discussion in the revised manuscript (lines 354-362).

<https://doi.org/10.7554/eLife.101209.2.sa0>