

# C-C chemokine receptor 4 deficiency exacerbates early atherosclerosis in mice

Reviewed Preprint

v1 • October 14, 2024

Not revised

Toru Tanaka, Naoto Sasaki , Aga Krisnanda, Hilman Zulkifli Amin, Ken Ito, Sayo Horibe, Kazuhiko Matsuo, Ken-ichi Hirata, Takashi Nakayama, Yoshiyuki Rikitake

Laboratory of Medical Pharmaceuticals, Kobe Pharmaceutical University, Kobe, Japan • Division of Cardiovascular Medicine, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kobe, Japan • Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan • Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia • Division of Chemotherapy, Faculty of Pharmacy, Kindai University, Higashi-osaka, Osaka, Japan • Department of Cardiology, Kakogawa Central City Hospital, Kakogawa, Hyogo, Japan

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

## Abstract

Chronic inflammation via dysregulation of T cell immune responses is critically involved in the pathogenesis of atherosclerotic cardiovascular disease. Improving the balance between proinflammatory T cells and anti-inflammatory regulatory T cells (Tregs) may be an attractive approach for treating atherosclerosis. Although C-C chemokine receptor 4 (CCR4) has been shown to mediate the recruitment of T cells to inflamed tissues, its role in atherosclerosis is unclear. Here, we show that genetic deletion of CCR4 in hypercholesterolemic mice accelerates the development of early atherosclerotic lesions characterized by an inflammatory plaque phenotype. This was associated with proinflammatory T helper type 1 (Th1) cell-skewed responses in peripheral lymphoid tissues, para-aortic lymph nodes, and atherosclerotic aorta. Mechanistically, CCR4 deficiency in Tregs impaired their suppressive function and migration to the atherosclerotic aorta and augmented Th1 cell-mediated immune responses through defective regulation of dendritic cell function, which accelerated aortic inflammation and atherosclerotic lesion development. Thus, we revealed a previously unrecognized role for CCR4 in controlling the early stage of atherosclerosis via Treg-dependent regulation of proinflammatory T cell responses. Our data suggest that CCR4 is an important negative regulator of atherosclerosis.

## eLife Assessment

This **important** study provides **solid** in-vivo evidence that CCR4 regulates the early inflammatory response during atherosclerotic plaque formation. The authors propose that altered T-cell response plays a role in this process, shedding light on mechanisms that may be of interest to medical biologists, biochemists, cell biologists, and immunologists. Further in vivo validation, mechanistic studies, and discussion of results in vitro suggested would be helpful to cement the significance and implications of these findings.

<https://doi.org/10.7554/eLife.101830.1.sa3>

## Introduction

Severe cardiovascular diseases, including ischemic heart disease and stroke, occur in patients with atherosclerosis and are major causes of mortality worldwide. Despite state-of-the-art intensive treatment, patients at high risk of atherosclerotic diseases have considerable residual risk, which could be partly related to vascular inflammatory responses.<sup>1</sup> Notably, recent clinical trials have provided evidence that anti-inflammatory therapies could be potentially novel approaches for preventing cardiovascular diseases in patients with the past disease history, although overall mortality has not improved.<sup>2,3</sup>

Chronic aortic inflammation via T cell-mediated immune dysregulation has been shown to be critically involved in the development of atherosclerosis.<sup>4</sup> Single-cell proteomic and transcriptomic approaches have revealed that activated and differentiated T cells are the major immune cells infiltrating human carotid artery plaques,<sup>5</sup> indicating the involvement of T cell-mediated immunoinflammatory responses in atherosclerotic plaque development in humans. A recent clinical trial provided evidence for the increased risk of atherosclerotic cardiovascular disease events in cancer patients treated with immune checkpoint inhibitors,<sup>6</sup> confirming the critical role of effector T cell (Teff) immune responses in the development of human atherosclerotic disease. T effs including T helper type 1 (Th1) cells play a proatherogenic role by producing proinflammatory cytokines including interferon (IFN)- $\gamma$ .<sup>7,8</sup> A recent translational study demonstrated that anti-CXC-motif-chemokine receptor 3 autoantibodies, which reflect Th1 cell-mediated responses, can be a novel biomarker and an inflammatory risk factor for cardiovascular morbidity and mortality beyond traditional risk factors, indicating the possible proatherogenic role of Th1 cells in humans.<sup>9</sup> On the other hand, several subsets of regulatory T cells (Tregs) including forkhead box P3 (Foxp3)-expressing Tregs play an anti-atherogenic role by suppressing Teff-mediated immunoinflammatory responses or reducing plasma atherogenic lipoprotein levels.<sup>10–12</sup> A number of approaches to dampening proatherogenic Th1 cell responses or augmenting atheroprotective Treg responses have been reported to be effective for the treatment and prevention of atherosclerosis in experimental mouse models.<sup>13</sup> Despite accumulating evidence for the critical role of the Th1 cell/Treg balance in atherosclerosis, how the balance of these T cell populations is regulated in lymphoid tissues and atherosclerotic lesions to control atherosclerosis is unclear.

Various immune cells including T cells and monocytes migrate to the aorta via interactions between chemokines and their specific chemokine receptors and are critically involved in the process of atherosclerosis.<sup>14</sup> However, few reports have identified a chemokine system that prevents atherosclerosis by improving the Th1 cell/Treg balance in lymphoid tissues and atherosclerotic lesions. C-C chemokine receptor 4 (CCR4) is expressed on several T cell subsets,

including T helper type 2 (Th2) cells, T helper type 17 (Th17) cells, skin-homing T cells, and Tregs, but not on proatherogenic Th1 cells.<sup>15</sup> CCR4 is a highly specific receptor for two CC chemokine ligands CCL17 (thymus- and activation-regulated chemokine) and CCL22 (macrophage-derived chemokine)<sup>16</sup> which were shown to be expressed in mesenteric lymph nodes (LNs) in a mouse model of inflammatory bowel disease<sup>17</sup> and lung tissues affected by allergic airway inflammation.<sup>18</sup> These chemokines play a role in guiding CCR4<sup>+</sup> T cells to the inflammatory sites with abundant expression of these chemokines.<sup>19,20</sup> Previous experimental studies have suggested that the CCL17/CCL22–CCR4 axes protect against inflammatory autoimmune diseases partly by promoting Treg accumulation in target tissues.<sup>17,18</sup> The expression of CCL17 and CCL22 was also observed in human and mouse atherosclerotic lesions.<sup>21–23</sup> Although the disruption of the CCL17/CCL22–CCR4 axes did not affect the development of advanced atherosclerotic lesions or the proportion of Tregs in peripheral lymphoid tissues in atherosclerosis-prone mice fed a high-cholesterol diet,<sup>22,24</sup> its effect on the early stage of atherosclerosis and Treg responses has not been investigated.

Here, we investigated the role of CCR4 in the development of early atherosclerotic lesions and the underlying mechanisms in CCR4-deficient (*Ccr4*<sup>−/−</sup>) mice on an apolipoprotein E-deficient (*Apoe*<sup>−/−</sup>) background fed a standard chow diet, with a particular focus on CD4<sup>+</sup> T cell immune responses.

## Results

### CCR4 is predominantly expressed on CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs, and the ligands CCL17 and CCL22 are expressed in peripheral LNs and atherosclerotic lesions

To evaluate the expression levels of CCR4 on CD4<sup>+</sup> T cells under normocholesterolemic or hypercholesterolemic conditions, we performed flow cytometric analysis of peripheral LNs, spleen, and para-aortic LNs from wild-type, *Apoe*<sup>−/−</sup>, and *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice. Notably, CCR4 expression was expressed on approximately 15–25% of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs from wild-type or *Apoe*<sup>−/−</sup> mice (Supplementary Figure 1A through 1C), while CD4<sup>+</sup>Foxp3<sup>−</sup> non-Tregs from these mice expressed CCR4 at markedly lower levels, suggesting that CCR4 is predominantly expressed on CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs and that hypercholesterolemia does not affect CCR4 expression. We also examined CCR4 expression on CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs in the atherosclerotic aorta of *Apoe*<sup>−/−</sup> mice, and consistently found that CCR4 expression in CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs was much higher than that in CD4<sup>+</sup>Foxp3<sup>−</sup> non-Tregs (Supplementary Figure 1D).

As CCL17 and CCL22, known as specific ligands for CCR4, are highly expressed by dendritic cells (DCs) in LNs,<sup>25,26</sup> we examined their expression in the peripheral LNs of wild-type, *Apoe*<sup>−/−</sup>, and *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice by immunohistochemistry. As expected, clear expression of these chemokines was observed in the peripheral LNs of wild-type, *Apoe*<sup>−/−</sup>, and *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice (data not shown). We also examined the expression of CCL17 and CCL22 in the aorta of these mice by immunohistochemistry. CCL17 expression was detected in the atherosclerotic lesions of *Apoe*<sup>−/−</sup> and *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice, and some lesional MOMA-2<sup>+</sup> macrophages expressed this chemokine (Supplementary Figure 2A). However, CCL17 expression was not detected in the aorta of wild-type mice without atherosclerotic plaques (Supplementary Figure 2A). Another ligand CCL22 was modestly expressed in atherosclerotic lesions and partially colocalized with lesional MOMA-2<sup>+</sup> macrophages in *Apoe*<sup>−/−</sup> or *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice, while its expression was not detected in the aorta of wild-type mice (Supplementary Figure 2B).

Together, these findings suggest that Tregs may migrate to peripheral LNs and atherosclerotic lesions partly via the CCL17/CCL22–CCR4 axes under hypercholesterolemia.

## CCR4 deficiency accelerates the development of early atherosclerotic lesions characterized by an inflammatory plaque phenotype


To investigate the effect of CCR4 deficiency on the development of early atherosclerosis, we analyzed the atherosclerotic lesions of 18-week-old *Apoe*<sup>-/-</sup> and *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice fed a standard chow diet. *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice developed normally without any spontaneous inflammatory disease. Notably, compared with *Apoe*<sup>-/-</sup> mice, *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice exhibited a significant increase in atherosclerotic lesion size in the aortic sinus (aortic sinus mean plaque area: control *Apoe*<sup>-/-</sup> mice:  $1.46 \pm 0.50 \times 10^5 \mu\text{m}^2$  versus *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup>:  $2.04 \pm 0.82 \times 10^5 \mu\text{m}^2$ ,  $P < 0.05$ ; **Figure 1A**). In parallel with the cross-sectional studies, we performed en face analysis of thoracoabdominal aortas, revealing no significant difference in aortic plaque burden between the 2 groups (**Figure 1B**). There were no significant differences in body weight or plasma lipid profile between the 2 groups (Supplementary Table 1).

To determine the effect of CCR4 deficiency on plaque components, we performed immunohistochemical studies of atherosclerotic lesions in the aortic sinus. Notably, compared with those of *Apoe*<sup>-/-</sup> mice, the atherosclerotic lesions of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice showed a 20% increase in macrophage accumulation (**Figure 1C**) and a marked 42% increase in CD4<sup>+</sup> T cell infiltration (**Figure 1D**). In addition, we performed immunohistochemical analysis of Foxp3<sup>+</sup> Tregs in atherosclerotic lesions using an anti-Foxp3 antibody. However, few Foxp3<sup>+</sup> Tregs were found within the plaques of *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice (data not shown). The proportion of collagen in the aortic sinus plaques of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice was significantly lower than that in the aortic sinus plaques of *Apoe*<sup>-/-</sup> mice (**Figure 1E**). These findings on atherosclerotic plaque size and components collectively suggest that CCR4 plays a role in preventing the development of early atherosclerotic lesions and inducing a less inflammatory plaque phenotype. To further evaluate aortic immunoinflammatory responses, we analyzed the mRNA expression of pro- and anti-inflammatory cytokines and transcription factors specific for Tregs or helper T cell subsets in atherosclerotic aorta by quantitative reverse transcription PCR. The mRNA expression of proinflammatory cytokines (*Il1b* and *Il6*), Th1-related *Tbx21*, and Th17-related *Rorc* was markedly upregulated in the aorta of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice, indicating augmented proatherogenic immune responses in the atherosclerotic aorta (**Figure 1F**). The mRNA expression of the Treg-specific transcription factor *Foxp3* was undetectable.

Collectively, these data suggest that CCR4 deficiency promotes the accumulation of inflammatory cells and proinflammatory immune responses in the aorta, leading to augmented development of early atherosclerotic lesions in the aortic root.

## CCR4 deficiency augments Teff immune responses in peripheral lymphoid tissues

We examined the mechanisms by which CCR4 deficiency accelerates early atherosclerosis by focusing on changes in systemic T cell responses, including those involving CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs and CD4<sup>+</sup>Foxp3<sup>-</sup> non-Tregs, in peripheral lymphoid tissues. The frequencies of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs and CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells were significantly higher in the spleen of 8- or 18-week-old *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in those of age-matched *Apoe*<sup>-/-</sup> mice (**Figure 2A** and **2B**). Similar tendency was observed for the peripheral LNs of these mice, although there was no difference in the frequency of CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells between 8-week-old *Apoe*<sup>-/-</sup> and *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice (**Figure 2A** and **2B**). The absolute numbers of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs and CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells were significantly higher in the peripheral LNs of 8- or 18-week-old *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than those of age-matched *Apoe*<sup>-/-</sup> mice (**Figure 2A** and **2B**). The absolute number of CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells was also higher in the spleen of 18-week-old *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than that of age-matched *Apoe*<sup>-/-</sup>



**Figure 1.**


### C-C chemokine receptor 4 (CCR4) deficiency accelerates the development of early atherosclerotic lesions characterized by an inflammatory plaque phenotype.

**A**, Representative photomicrographs of Oil Red O staining and quantitative analysis of atherosclerotic lesion area at 5 different levels and the average area in the aortic sinus of 18-week-old apolipoprotein E-deficient (*Apoe*<sup>-/-</sup>) mice (n=24) or CCR4-deficient mice on an *Apoe*<sup>-/-</sup> background (*Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup>; n=23). **B**, Representative photomicrographs of Oil Red O staining and quantitative analysis of atherosclerotic lesion area in the aorta of 18-week-old *Apoe*<sup>-/-</sup> (n=15) or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice (n=15). **C-E**, Representative sections and quantitative analyses of MOMA-2<sup>+</sup> macrophages (**C**), CD4<sup>+</sup> T cells (**D**), and collagen (**E**) in the aortic sinus. Arrowheads indicate the CD4<sup>+</sup> T cells. n=10 per group. **F**, mRNA expression of pro- or anti-inflammatory cytokines and helper T cell-associated transcription factors in aorta. The expression levels of the target genes were normalized so that the mean values in *Apoe*<sup>-/-</sup> mice were set to 1. n=8 to 10 per group. Eighteen-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice were used for all experiments. Black bars represent 50, 200, or 500 μm as described. Data points represent individual animals. Horizontal bars represent means. Error bars indicate s.d. \**P*<0.05, \*\**P*<0.01; Mann-Whitney *U*-test: **A** and **F** *Il1b*; 2-tailed Student's *t*-test: **C**, **D**, **E**, and **F** *Il6*, *Tbx21*, and *Rorc*.

mice (**Figure 2A** and **2B**). Similar tendency was observed for the peripheral LNs of these mice, although there was no difference in the frequency of  $CD4^+CD44^{\text{high}}CD62L^{\text{low}}$  effector memory T cells between 8-week-old  $Apoe^{-/-}$  and  $Ccr4^{-/-}Apoe^{-/-}$  mice (**Figure 2A** and **2B**). The absolute numbers of  $CD4^+Foxp3^+$  Tregs and  $CD4^+CD44^{\text{high}}CD62L^{\text{low}}$  effector memory T cells were significantly higher in the peripheral LNs of 8- or 18-week-old  $Ccr4^{-/-}Apoe^{-/-}$  mice than those of age-matched  $Apoe^{-/-}$  mice (**Figure 2A** and **2B**). The absolute number of  $CD4^+CD44^{\text{high}}CD62L^{\text{low}}$  effector memory T cells was also higher in the spleen of 18-week-old  $Ccr4^{-/-}Apoe^{-/-}$  mice than that of age-matched  $Apoe^{-/-}$  mice (**Figure 2B**). Similar results were obtained for normocholesterolemic wild-type mice (data not shown), suggesting that the expansion of Tregs and effector memory T cells in  $Ccr4^{-/-}Apoe^{-/-}$  mice is independent of hypercholesterolemia. We evaluated the proliferative capacity of  $CD4^+Foxp3^+$  Tregs and  $CD4^+Foxp3^-$  non-Tregs by analyzing Ki-67 expression using flow cytometry and found that the proportions of Ki-67-positive Tregs and non-Tregs were markedly higher in the peripheral LNs and spleen of  $Ccr4^{-/-}Apoe^{-/-}$  mice than in those of  $Apoe^{-/-}$  mice (**Figure 2C** and **2D**). To determine the effect of CCR4 deficiency on T cell development in the thymus, we performed flow cytometric analysis of thymocytes from 4-week-old  $Apoe^{-/-}$  or  $Ccr4^{-/-}Apoe^{-/-}$  mice and found no difference in the development of thymic T cells between the 2 groups (Supplementary Figure 3A). In line with a previous report in normocholesterolemic mice,<sup>27</sup> there was no difference in the frequency of thymic  $CD4^+Foxp3^+$  Tregs between the 2 groups (Supplementary Figure 3B). These data indicate that the increased numbers of  $CD4^+Foxp3^+$  Tregs and  $CD4^+Foxp3^-$  non-Tregs in peripheral LNs are due to their enhanced proliferative capacity but not to their promoted development in the thymus. We also analyzed other immune cells in the spleen of  $Apoe^{-/-}$  and  $Ccr4^{-/-}Apoe^{-/-}$  mice by flow cytometry and found no major differences in the proportions or activation-associated molecule expression between the 2 groups (Supplementary Figure 4).

To determine the activation and function of CCR4-deficient  $CD4^+Foxp3^+$  Tregs, we investigated the expression of their activation- and function-associated molecules in the peripheral LNs of 8- or 18-week-old  $Apoe^{-/-}$  or  $Ccr4^{-/-}Apoe^{-/-}$  mice by flow cytometry. Notably, CCR4 deficiency had no major effect on the expression of cytotoxic T lymphocyte-associated antigen-4 (CTLA-4) or CD103 in  $CD4^+Foxp3^+$  Tregs (**Figure 2E**). We also analyzed the mRNA expression of *Ccr4*, activation- or function-associated molecules (*Foxp3*, *Ctla4*, *Cd103*, *Tnfrsf18*, *Il10*, and *Tgfb*), and major chemokine receptors (*Ccr5*, *Ccr6*, *Ccr7*, and *Ccr8*) in splenic Tregs by quantitative reverse transcription PCR. The *Ccr4* mRNA expression in Tregs from  $Ccr4^{-/-}Apoe^{-/-}$  mice was less than the detectable levels (data not shown). In line with the data on peripheral LNs, there were no significant differences in the mRNA expression of these molecules between the 2 groups (**Figure 2F**; Supplementary Figure 5A).

In contrast to the results on  $CD4^+Foxp3^+$  Tregs, the expression of activation marker CTLA-4 in peripheral LN  $CD4^+Foxp3^-$  non-Tregs was higher in  $Ccr4^{-/-}Apoe^{-/-}$  mice than in  $Apoe^{-/-}$  mice, although the expression of another activation marker CD103 was not altered by CCR4 deficiency (**Figure 2G**). We analyzed the mRNA expression of *Ccr4*, activation-associated molecules (*Ctla4*, *Cd44*, *Cd69*, and *Cd103*), and representative chemokine receptors (*Ccr1*, *Ccr5*, *Ccr6*, *Ccr7*, *Ccr8*, *Cxcr3*, and *Cx3cr1*) in splenic non-Tregs by quantitative reverse transcription PCR. The *Ccr4* mRNA expression in non-Tregs from  $Ccr4^{-/-}Apoe^{-/-}$  mice was less than the detectable levels (data not shown). We found a marked increase in the mRNA expression of activation-associated molecules (*Ctla4*, *Cd44*, and *Cd103*) in splenic non-Tregs from  $Ccr4^{-/-}Apoe^{-/-}$  mice (**Figure 2H**). We also observed upregulated mRNA expression of various chemokine receptors including Th1-related *Ccr1*, *Cxcr3*, and *Cx3cr1*, Th2-related *Ccr8*, and Th17-related *Ccr6* in splenic non-Tregs from  $Ccr4^{-/-}Apoe^{-/-}$  mice (Supplementary Figure 5B), indicating that CCR4 deficiency may promote the activation and expansion of helper T cells and facilitate their migration to atherosclerotic lesions. There were no significant differences in the mRNA expression of other chemokine receptors (*Ccr5* and *Ccr7*) in splenic non-Tregs between the 2 groups (Supplementary Figure 5B).



**Figure 2.**

### C-C chemokine receptor 4 (CCR4) deficiency augments effector T cell immune responses in peripheral lymphoid tissues.

**A and B.** Representative flow cytometric analysis of CD4<sup>+</sup> forkhead box P3 (Foxp3)<sup>+</sup> regulatory T cells (Tregs) (**A**) and CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells (**B**) in the spleen of 8-week-old apolipoprotein E-deficient (*Apoe*<sup>-/-</sup>) mice or CCR4-deficient mice on an *Apoe*<sup>-/-</sup> background (*Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup>). The graphs represent the total numbers and proportions of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs (**A**) and CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells (**B**) in the peripheral lymph nodes (LNs) and spleen of 8- or 18-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. *n*=9 to 10 per group. **C and D.** The graphs represent the proportions of Ki-67 positive cells among CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs (**C**) and CD4<sup>+</sup>Foxp3<sup>-</sup> non-Tregs (**D**) in the peripheral LNs and spleen of 8-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice, as assessed by flow cytometry. *n*=13 per group. **E.** Expression levels of activation-associated molecules cytotoxic T lymphocyte-associated antigen-4 (CTLA-4) and CD103 were analyzed by gating on CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs in the peripheral LNs of 8- or 18-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. *n*=9 to 10 per group. **F.** mRNA expression of Treg-associated markers in splenic Tregs from 8-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. *n*=8 per group. **G.** Expression levels of activation-associated molecules CTLA-4 and CD103 were analyzed by gating on CD4<sup>+</sup>Foxp3<sup>-</sup> non-Tregs in the peripheral LNs of 8- or 18-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. *n*=9 to 10 per group. **H.** mRNA expression of activation-associated molecules in splenic non-Tregs from 8-week-old *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. *n*=8 per group. The expression levels of the target genes were normalized so that the mean values in *Apoe*<sup>-/-</sup> mice were set to 1 (**F, H**). Data points represent individual animals. Horizontal bars represent means. Error bars indicate s.d. \**P*<0.05, \*\**P*<0.01; Mann-Whitney *U*-test: **A** second (8w) from the left, **B** second (8w) and third (8w) from the left, **C** left, and **H** *Cd44* and *Cd103*; 2-tailed Student's *t*-test: **A** first, second (18w), and third from the left, **B** first, second (18w), third (18w), and fourth from the left, **C** right, **D, E, G,** and **H** *Ctla4*. MFI indicates mean fluorescence intensity.

Considering the accelerated early atherosclerosis observed in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice, these results indicate that Teff-skewed immune responses may affect atherosclerosis more strongly than the increase in Tregs in peripheral lymphoid tissues.

## CCR4 deficiency promotes proinflammatory CD4<sup>+</sup> T cell immune responses in peripheral lymphoid tissues

To determine whether CCR4 deficiency affects CD4<sup>+</sup> T cell immune responses and polarization, we examined cytokine secretion from CD4<sup>+</sup> T cells by intracellular cytokine staining. The fraction of IFN-γ-producing Th1 cells in the peripheral LNs was significantly higher in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in *Apoe*<sup>-/-</sup> mice, while there were no differences in the fractions of other CD4<sup>+</sup> T cell subsets including interleukin (IL)-4-producing Th2 cells, IL-10-producing CD4<sup>+</sup> T cells, and IL-17-producing Th17 cells between the 2 groups (**Figure 3A** [↗](#)). In line with this, the fraction of splenic Th1 cells was higher in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in *Apoe*<sup>-/-</sup> mice, although the proportion of IL-17-producing Th17 cells was also higher in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in *Apoe*<sup>-/-</sup> mice (**Figure 3B** [↗](#)).


We semiquantitatively analyzed the production of various cytokines or chemokines by splenic CD4<sup>+</sup> T cells stimulated with plate-bound anti-CD3 and anti-CD28 antibodies using a cytokine array kit. Compared with those from *Apoe*<sup>-/-</sup> mice, splenic CD4<sup>+</sup> T cells from *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice secreted more Th1-related cytokine IFN-γ, Th2-related cytokine IL-13, Th17-related cytokine IL-17, and various inflammation-related cytokines and chemokines (**Figure 3C** [↗](#)). ELISA analysis confirmed that the IFN-γ and IL-17 levels in the cell supernatants of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice were much higher than those in the cell supernatants of *Apoe*<sup>-/-</sup> mice (**Figure 3D** [↗](#)). Although the cytokine levels of Th2-related cytokine IL-4 and Treg-related anti-inflammatory cytokine IL-10 were lower than the levels detected by cytokine array analysis, ELISA analysis revealed higher IL-4 and IL-10 production in splenic CD4<sup>+</sup> T cells from *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in those from *Apoe*<sup>-/-</sup> mice (**Figure 3D** [↗](#)), which may be compensatory immune responses to the proinflammatory T cell responses caused by CCR4 deficiency.

## CCR4 deficiency promotes Th1 cell responses in para-aortic LNs and atherosclerotic aorta

Next, we investigated the mechanisms by which CCR4 deficiency accelerates early atherosclerosis by focusing on local immune responses in para-aortic LNs and atherosclerotic aorta. Consistent with the peripheral LN data, the frequency and number of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs were significantly higher in the para-aortic LNs of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in those of *Apoe*<sup>-/-</sup> mice (**Figure 4A** [↗](#)). The number of CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells was significantly higher in the para-aortic LNs of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in those of *Apoe*<sup>-/-</sup> mice, while their frequency did not differ between the 2 groups (**Figure 4B** [↗](#)).

In line with the data on peripheral lymphoid tissues, there were no significant differences in the expression of activation- or function-associated molecules or chemokine receptors in para-aortic LN Tregs between the 2 groups (**Figure 4C** [↗](#); Supplementary Figure 6A). Notably, the expression of *Cd103* and Th1-related *Tbx21* was upregulated in para-aortic LN non-Tregs from *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice, while the expression of other activation- or helper T cell-associated molecules or chemokine receptors was unaltered (**Figure 4D** [↗](#); Supplementary Figure 6B). Flow cytometric analysis of helper T cell subsets in para-aortic LNs revealed that the fractions of IFN-γ-producing Th1 cells, IL-4-producing Th2 cells, and IL-17-producing Th17 cells were significantly higher in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in *Apoe*<sup>-/-</sup> mice, suggesting augmented proinflammatory CD4<sup>+</sup> T cell immune responses by CCR4 deficiency (**Figure 4E** [↗](#)).


We explored the infiltration of T-box expressed in T cells (T-bet)-expressing Th1 cells, GATA3-expressing Th2 cells, and retinoic acid-related orphan receptor gamma t (RORγt)-expressing Th17 cells in atherosclerotic aorta by flow cytometry. Strikingly, the frequency of aortic T-bet-expressing



**Figure 3.**

### C-C chemokine receptor 4 (CCR4) deficiency promotes proinflammatory CD4 $^{+}$ T cell immune responses in peripheral lymphoid tissues.

**A and B,** The graphs represent the frequencies of interferon (IFN)- $\gamma$  $^{+}$ , interleukin (IL)-4 $^{+}$ , IL-10 $^{+}$ , and IL-17 $^{+}$  CD4 $^{+}$  T cells in the peripheral lymph nodes (LNs) (**A**) and spleen (**B**) of 8-week-old apolipoprotein E-deficient (*Apoe* $^{-/-}$ ) mice or CCR4-deficient mice on an *Apoe* $^{-/-}$  background (*Ccr4* $^{-/-}$ *Apoe* $^{-/-}$ ). *n*=10 to 11 per group. **C and D,** Purified splenic CD4 $^{+}$  T cells from 8-week-old *Apoe* $^{-/-}$  or *Ccr4* $^{-/-}$ *Apoe* $^{-/-}$  mice were stimulated with plate-bound anti-CD3 and soluble anti-CD28 antibodies *in vitro*. The levels of various cytokines and chemokines in pooled cell supernatants from 8 mice in each group were determined semiquantitatively by a cytokine array kit (**C**). Data are representative of two independent experiments. Cytokine concentrations in the cell supernatants were measured by ELISA (**D**). *n*=8 per group. Data points represent individual animals. Horizontal bars represent means. Error bars indicate s.d. \**P*<0.05, \*\**P*<0.01; Mann-Whitney *U*-test: **A, B** first from the left, and **D** second from the left; 2-tailed Student's *t*-test: **B** fourth from the left and **D** first, third, and fourth from the left.



**Figure 4.**

### C-C chemokine receptor 4 (CCR4) deficiency promotes T helper type 1 (Th1) cell responses in para-aortic lymph nodes (LNs) and atherosclerotic aorta.

**A** and **B**, Representative flow cytometric analysis of CD4<sup>+</sup> forkhead box P3 (Foxp3)<sup>+</sup> regulatory T cells (Tregs) (**A**) and CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells (**B**) in para-aortic LNs. The graphs represent the total numbers and proportions of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs (**A**) and CD4<sup>+</sup>CD44<sup>high</sup>CD62L<sup>low</sup> effector memory T cells (**B**) in para-aortic LNs. n=9 to 10 per group. **C** and **D**, mRNA expression of Treg-associated markers in Tregs (**C**) and mRNA expression of activation or helper T cell-associated molecules in non-Tregs (**D**) in para-aortic LNs. The expression levels of the target genes were normalized so that the mean values in apolipoprotein E-deficient (*Apoe*<sup>-/-</sup>) mice were set to 1. Tregs or non-Tregs purified from pooled para-aortic LNs of 9 to 10 mice were analyzed as a sample. n=4 per group. **E**, The graphs represent the frequencies of interferon (IFN)- $\gamma$ <sup>+</sup>, interleukin (IL)-4<sup>+</sup>, IL-10<sup>+</sup>, and IL-17<sup>+</sup> CD4<sup>+</sup> T cells in para-aortic LNs. n=12 per group. **F-H**, Representative flow cytometric analysis of T-box expressed in T cells (T-bet) (**F**), GATA3 (**G**), and retinoic acid-related orphan receptor gamma t (ROR $\gamma$ t) (**H**) expression in aortic CD3<sup>+</sup>CD4<sup>+</sup>CD45<sup>+</sup> T cells. The graphs represent the frequencies of T-bet<sup>+</sup> (**F**), GATA3<sup>+</sup> (**G**), and ROR $\gamma$ t<sup>+</sup> (**H**) cells among aortic CD3<sup>+</sup>CD4<sup>+</sup>CD45<sup>+</sup> T cells. n=9 per group. **I**, Representative flow cytometric analysis of Foxp3 expression in aortic CD3<sup>+</sup>CD4<sup>+</sup>CD45<sup>+</sup> T cells. The graph represents the frequency of Foxp3<sup>+</sup> Tregs among aortic CD3<sup>+</sup>CD4<sup>+</sup>CD45<sup>+</sup> T cells. n=9 per group. **J**, The graph represents the ratio of CD4<sup>+</sup>T-bet<sup>+</sup> Th1 cells to CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs (Th1 cell/Treg ratio). n=9 per group. Pooled aortic lymphoid cells from 2 mice were analyzed as a sample. Eighteen-week-old *Apoe*<sup>-/-</sup> or CCR4-deficient mice on an *Apoe*<sup>-/-</sup> background (*Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup>) were used for all experiments. Data points represent individual animals (**A**, **B**, and **E**) or individual pooled samples (**C**, **D**, **F-J**). Horizontal bars represent means. Error bars indicate s.d. \**P*<0.05, \*\**P*<0.01; Mann-Whitney *U*-test: **D** *Cd103* and **J**; 2-tailed Student's *t*-test: **A**, **B**, **D** *Tbx21*, **E**, and **F**.

Th1 cells was markedly higher in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice than in *Apoe*<sup>-/-</sup> mice, while the proportions of Th2 and Th17 subsets did not differ between the 2 groups (**Figure 4F** through **4H**). We next analyzed the accumulation of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs in the atherosclerotic aorta by flow cytometry and found no difference in the proportion of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs between the 2 groups (**Figure 4I**). Importantly, we found a marked increase in the ratio of T-bet-expressing Th1 cells to CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs (Th1 cell/Treg ratio) in the atherosclerotic aorta of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice (**Figure 4J**), indicating Th1-skewed immune responses in the atherosclerotic lesions of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice.


Collectively, these results suggest that CCR4 deficiency promotes the accumulation of proatherogenic Th1 cells in para-aortic LNs and atherosclerotic aorta and shifts the Th1 cell/Treg balance toward Th1 cell responses, leading to exacerbated aortic inflammation and early atherosclerosis.

## CCR4 expression on Tregs regulates Th1 cell responses and mediates Treg migration to the atherosclerotic aorta

Given that CCR4 plays a crucial role in attenuating immunoinflammatory responses in autoimmune or allergic disease via the modulation of Treg function,<sup>17,18</sup> we speculated that despite the expansion and unaltered expression of activation- or function-associated molecules in Tregs, these cells might be dysfunctional. To determine whether CCR4 deficiency affects the suppressive function of Tregs in hypercholesterolemia, we performed an *in vitro* suppression assay. Interestingly, CCR4 deficiency significantly impaired the suppressive function of Tregs isolated from hypercholesterolemic mice (**Figure 5A**), indicating that CCR4 expression on Tregs may be important for the regulation of proinflammatory immune responses and the development of early atherosclerosis.

Interaction with DCs is well-known as one of the core suppressive mechanisms by which Tregs control excessive immune responses. A previous report demonstrated that the cell-cell contacts between Tregs and CCL22-deficient DCs are impaired.<sup>28</sup> Tregs limit the CD80/CD86-CD28-dependent activation of T cells through CTLA-4-dependent downregulation of CD80 and CD86 expression on DCs,<sup>29</sup> which may contribute to the reduction in atherosclerosis.<sup>30</sup> Therefore, we examined the interactions between CCR4-intact or CCR4-deficient Tregs and DCs by a coculture experiment (**Figure 5B**). As expected, the upregulation of CD80 and CD86 expression on DCs mediated by conventional T cells was markedly suppressed by coculture with Tregs from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice (**Figure 5C**). Notably, the suppressive effect of CCR4-deficient Tregs was significantly attenuated compared with that of CCR4-intact Tregs (**Figure 5C**). To assess the involvement of CCR4 expression on Tregs in regulating the production of proatherogenic IFN-γ by conventional T cells, we stimulated conventional T cells with an anti-CD3 antibody and DCs in the presence or absence of CCR4-intact or CCR4-deficient Tregs and analyzed IFN-γ production by ELISA (**Figure 5B**). IFN-γ production by conventional T cells was markedly suppressed by coculture with CCR4-intact Tregs but was not significantly affected by coculture with CCR4-deficient Tregs (**Figure 5D**). Importantly, conventional T cells cocultured with CCR4-deficient Tregs produced more IFN-γ than those cocultured with CCR4-intact Tregs (**Figure 5D**). These results suggest that the augmented Th1 cell responses in the peripheral lymphoid tissues of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice are partly due to impaired Treg-dependent regulation of DC function.

As described above, CD4<sup>+</sup>Foxp3<sup>+</sup> Treg accumulation in the atherosclerotic aorta of *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice was not altered despite the marked expansion of CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs in peripheral lymphoid tissues. Based on these findings and previous reports showing the impaired migratory capacity of CCR4-deficient Tregs to inflammatory tissues,<sup>17,18</sup> we hypothesized that CCR4-deficient Tregs might have less capacity to migrate to atherosclerotic lesions. To address this issue, we performed a Treg transfer experiment using *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice on a Kaede-Tg background<sup>31</sup> which provided us with the opportunity to faithfully track transferred Tregs by monitoring the



**Figure 5.**

### C-C chemokine receptor 4 (CCR4) expression on regulatory T cells (Tregs) regulates T helper type 1 cell responses and mediates Treg migration to the aorta.

**A**, The suppressive function of Tregs was assessed by evaluating the proliferation of carboxyfluorescein diacetate succinimidyl ester (CFSE)-labeled conventional T cells (Tconv) cocultured with Tregs from apolipoprotein E-deficient (*Apoe*<sup>-/-</sup>) mice or CCR4-deficient mice on an *Apoe*<sup>-/-</sup> background (*Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup>). Data are presented as the results of triplicate wells and are representative of 2 independent experiments. Data are expressed as the mean ± s.d. **B** and **C**, CD80 and CD86 expression in live splenic dendritic cells (DCs) after 2 days of coculture with Tconv from *Apoe*<sup>-/-</sup> mice, or a mixture of Tregs from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice and Tconv from *Apoe*<sup>-/-</sup> mice in the presence of an anti-CD3 antibody. Data points represent the results of quintuplicate wells. Data are representative of 2 independent experiments. **B** and **D**, Tconv from *Apoe*<sup>-/-</sup> mice and DCs were cocultured with or without Tregs from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice in the presence of an anti-CD3 antibody. Interferon (IFN)-γ concentrations in cell supernatants were measured by ELISA. Data points represent the results of sextuplicate wells. **E**, Eighteen-week-old *Apoe*<sup>-/-</sup> mice fed a high-cholesterol diet for 10 weeks received transfer of Tregs from *Apoe*<sup>-/-</sup> Kaede-Tg or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> Kaede-Tg mice, and the accumulation of Kaede<sup>+</sup> Tregs in the peripheral lymphoid tissues and aorta was analyzed by flow cytometry 20 hours later. **F-I**, Representative flow cytometric analysis and the proportions of Kaede<sup>+</sup> Tregs among CD4<sup>+</sup> T cells in the peripheral lymph nodes (LNs) (**F**), spleen (**G**), para-aortic LNs (**H**), and aorta (**I**) of *Apoe*<sup>-/-</sup> mice that received *Apoe*<sup>-/-</sup> Kaede<sup>+</sup> Tregs or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> Kaede<sup>+</sup> Tregs. n=9 to 10 per group (**F-H**). Pooled aortic lymphoid cells from 5 mice in each group were used for analysis. The results are presented as the mean ± s.d. of 3 independent experiments. Treg migration was normalized so that the number of migrated *Apoe*<sup>-/-</sup> Kaede<sup>+</sup> Tregs was set to 1 (**I**). Data points represent individual animals (**F-H**) or individual pooled samples (**I**). Horizontal bars represent means. Error bars indicate s.d. \**P*<0.05, \*\**P*<0.01; 1-way ANOVA followed by Tukey's multiple comparisons test: **C** and **D**; 2-tailed Student's *t*-test: **A** and **I**. MFI indicates mean fluorescence intensity.

fluorescent Kaede protein (**Figure 5E** [↗](#)). There was no difference in the migratory capacity of CCR4-intact or CCR4-deficient Kaede-expressing Tregs to the peripheral LNs, spleen, or para-aortic LNs of recipient *Apoe*<sup>-/-</sup> mice (**Figure 5F** [↗](#) through **5H** [↗](#)). To promote the accumulation of T cells in atherosclerotic aorta, we fed recipient *Apoe*<sup>-/-</sup> mice with a high-cholesterol diet and analyzed the migration of transferred Tregs in the aorta. Interestingly, we found a significant reduction in the proportion of CCR4-deficient Kaede-expressing Tregs in the aorta of recipient *Apoe*<sup>-/-</sup> mice (**Figure 5I** [↗](#)), suggesting that CCR4-deficient Tregs have a reduced ability to migrate to the atherosclerotic aorta, but not to the peripheral lymphoid tissues, under hypercholesterolemia.

Taken together, these data demonstrate that CCR4 expression on Tregs plays a critical role in regulating Th1 cell responses in lymphoid tissues and mediating Treg migration to the atherosclerotic aorta under hypercholesterolemia, which cooperatively contributes to the reduction in early atherosclerosis by efficiently mitigating aortic inflammatory immune responses.

## CCR4 expression on Tregs is critical for limiting aortic inflammation and the development of atherosclerosis


To provide direct evidence for the critical role of CCR4 expression in Tregs in reducing early atherosclerosis, we injected *Apoe*<sup>-/-</sup> mice with saline or Tregs from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice and analyzed the aortic root atherosclerotic lesions of recipient *Apoe*<sup>-/-</sup> mice (**Figure 6A** [↗](#)). Although there were no significant differences in the aortic sinus mean plaque area among the 3 groups (data not shown), detailed analysis of the aortic root plaques at 5 different levels revealed significantly greater lesions in *Apoe*<sup>-/-</sup> mice that were administered CCR4-deficient Tregs than in *Apoe*<sup>-/-</sup> mice that were administered CCR4-intact Tregs (**Figure 6B** [↗](#)), suggesting that the impaired CCR4-deficient Treg function is involved in the acceleration of atherosclerosis in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. There were no significant differences in body weight or plasma lipid profile among the 3 groups (Supplementary Table 2).

To determine the effect of Treg-specific CCR4 deficiency on plaque components, we performed immunohistochemical studies of atherosclerotic lesions in the aortic sinus. Compared with those of saline-injected *Apoe*<sup>-/-</sup> mice, the atherosclerotic lesions of *Apoe*<sup>-/-</sup> mice injected with CCR4-intact Tregs showed markedly reduced accumulation of macrophages (**Figure 6C** [↗](#)) and CD4<sup>+</sup> T cells (**Figure 6D** [↗](#)), whereas there were no differences in intraplaque accumulation of these inflammatory cells between saline-injected *Apoe*<sup>-/-</sup> mice and *Apoe*<sup>-/-</sup> mice injected with CCR4-deficient Tregs (**Figure 6C** [↗](#) and **6D** [↗](#)). Notably, macrophage accumulation in the aortic sinus atherosclerotic lesions was markedly higher in *Apoe*<sup>-/-</sup> mice injected with CCR4-deficient Tregs than in *Apoe*<sup>-/-</sup> mice injected with CCR4-intact Tregs (**Figure 6C** [↗](#)). Collagen content in atherosclerotic lesions was significantly higher in *Apoe*<sup>-/-</sup> mice injected with CCR4-intact Tregs than in saline-injected *Apoe*<sup>-/-</sup> mice, while no difference in collagen was observed between between saline-injected *Apoe*<sup>-/-</sup> mice and *Apoe*<sup>-/-</sup> mice injected with CCR4-deficient Tregs (**Figure 6E** [↗](#)).

Overall, we provide evidence that CCR4 protects against early atherosclerosis partly by mediating Treg-dependent induction of a less inflammatory plaque phenotype.

## Discussion

In the present study, we demonstrated that genetic deletion of CCR4 in hypercholesterolemic *Apoe*<sup>-/-</sup> mice accelerates the development of early atherosclerotic lesions in the aortic root which exhibit an inflammatory phenotype, associated with proatherogenic Th1 cell-skewed responses in the peripheral lymphoid tissues, para-aortic LNs, and atherosclerotic aorta. Furthermore, T cell-DC coculture and Treg transfer experiments revealed that CCR4 expression on Tregs regulates the



**Figure 6.**

**C-C chemokine receptor 4 (CCR4) expression on regulatory T cells (Tregs) is critical for limiting aortic inflammation and the development of early atherosclerosis.**

**A**, Tregs purified from the peripheral lymph nodes (LNs) and spleen of apolipoprotein E-deficient (*Apoe*<sup>-/-</sup>) mice or CCR4-deficient mice on an *Apoe*<sup>-/-</sup> background (*Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup>) were intravenously transferred into 12-week-old *Apoe*<sup>-/-</sup> mice fed a standard chow diet, and atherosclerotic lesions were analyzed at 16 weeks of age. As a control without cell transfer, 12-week-old *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice were intravenously injected with saline and atherosclerotic lesions were analyzed at 16 weeks of age. **B**, Representative photomicrographs of Oil Red O staining and quantitative analysis of atherosclerotic lesion area at 5 different levels and maximal lesions in the aortic sinus of *Apoe*<sup>-/-</sup> mice injected with saline (n=19), *Apoe*<sup>-/-</sup> Tregs (n=17), or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> Tregs (n=20). **C-E**, Representative sections and quantitative analyses of MOMA-2<sup>+</sup> macrophages (**C**), CD4<sup>+</sup> T cells (**D**), and collagen (**E**) in the aortic sinus of *Apoe*<sup>-/-</sup> mice injected with saline, *Apoe*<sup>-/-</sup> Tregs, or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> Tregs. Arrowheads indicate the CD4<sup>+</sup> T cells. n=10 per group. Black bars represent 50, 200, or 500 μm as described. Data points represent individual animals. Horizontal bars represent means. Error bars indicate s.d. \*P<0.05, \*\*P<0.01; 1-way ANOVA followed by Tukey's multiple comparisons test.

development of early atherosclerosis by suppressing Th1 cell responses in lymphoid tissues and mediating Treg migration to the atherosclerotic aorta. Thus, we identified a novel role for the CCL17/CCL22–CCR4 axes in controlling early atherosclerosis via favorable modulation of the Th1 cell/Treg balance.

Emerging experimental and clinical data obtained by single-cell RNA sequencing and mass cytometry have clearly shown that CD4<sup>+</sup> or CD8<sup>+</sup> T cells are the dominant populations in human<sup>5</sup> and mouse<sup>32</sup> atherosclerotic plaques. Recent experimental and clinical evidence has demonstrated that dysregulation of the balance of proatherogenic Th1 cells and anti-atherogenic Tregs has adverse effects on atherosclerotic disease.<sup>13</sup> In the present study, we found that the formation of early atherosclerotic lesions in the aortic root was markedly accelerated by CCR4 deficiency, which was associated with the upregulation of various helper T cell immune responses in peripheral lymphoid tissues and augmented Th1 and Th17 cell-mediated responses in the atherosclerotic aorta. Given the proatherogenic nature of Th1 cells,<sup>7,8</sup> our data indicate that Th1 cell-skewed responses in peripheral lymphoid tissues, especially in atherosclerotic lesions, substantially contributed to atherosclerotic lesion formation in *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice. However, we cannot exclude the possibility that augmented immune responses of other helper T cell subsets might also contribute to accelerated atherosclerosis in *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice.

As Th2 cells and some Th17 cells, but not Th1 cells, are known to express CCR4,<sup>15</sup> the augmented Th1 cell responses observed in *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice do not seem to be direct effects of CCR4 deficiency and may involve other mechanisms. Given our recent report showing the augmented immune responses related to Th1, Th2, and Th17 cells in CD4<sup>+</sup>Foxp3<sup>+</sup> Treg-depleted atherosclerosis-prone mice,<sup>33</sup> augmentation of these helper T cell responses may be caused by dysregulated CD4<sup>+</sup>Foxp3<sup>+</sup> Treg responses in our *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice. In line with experimental studies in mouse models of inflammatory diseases showing a pivotal role for CCR4 in mediating Treg migration to inflammatory tissues,<sup>17,18</sup> we found that CCR4 expression on Tregs is critical for regulating atherosclerosis by mediating their migration to the atherosclerotic aorta. Another interesting finding is that CCR4 deficiency markedly upregulated Th1 cell responses in peripheral lymphoid tissues as well as in atherosclerotic lesions under hypercholesterolemia. T cell-DC coculture experiments clearly demonstrated that CCR4 expression on Tregs is critical for attenuating DC-dependent stimulation of Th1 cell responses by maintaining interactions between Tregs and DCs and Treg suppressive function, which is supported by the findings of a previous report showing a crucial role of the CCL22–CCR4 axis in the contact of Tregs with DCs and regulation of inflammatory responses.<sup>28</sup> Based on these findings, we propose that CCR4 deficiency in Tregs interrupts their contact with DCs in lymphoid tissues and impairs their suppressive function despite their unaltered activation status, leading to augmented Th1 cell responses and accelerated atherosclerosis. We speculate that the increased frequency of Tregs in the peripheral lymphoid tissues of *Ccr4*<sup>−/−</sup>*Apoe*<sup>−/−</sup> mice may be a compensatory effect in response to augmented Th1 cell-mediated proinflammatory responses due to dysfunctional Tregs.

T cell costimulatory signals such as the CD28–CD80/CD86<sup>10</sup> and CD27–CD70<sup>34</sup> pathways protect against atherosclerosis by systemically shifting the Th1 cell/Treg balance toward Treg responses. Several approaches involving treatment with cytokines<sup>35,36</sup>, antibodies<sup>11,36,37</sup>, an active form of vitamin D<sub>3</sub><sup>38</sup>, or ultraviolet B irradiation<sup>39,40</sup> have been reported to mitigate atherosclerosis via modulation of the Th1 cell/Treg balance in the peripheral lymphoid tissues and atherosclerotic lesions of atherosclerosis-prone mice. Despite these reports, the mechanisms that regulate the T cell balance in lymphoid tissues, particularly in atherosclerotic lesions, have not been fully elucidated. The chemokine system plays an important role in the differentiation and migration of various subsets of T cells including Tregs and has been shown to be involved in the process of atherosclerosis.<sup>14</sup> We revealed that CCR4 protects against early atherosclerosis by mediating Treg migration to the aorta and mitigating Th1 cell responses in lymphoid tissues and atherosclerotic lesions, indicating a previously unrecognized role of the chemokine system in regulating the Th1 cell/Treg balance to limit atherosclerosis.

We observed the accelerated development of early atherosclerotic lesions in *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice fed a standard chow diet. However, previous studies showed that neither hematopoietic nor systemic CCR4 deficiency affected the development of advanced atherosclerotic lesions or Treg frequency in severely hypercholesterolemic mice.<sup>22,24</sup> These findings suggest that the role of CCR4 may vary depending on the stages of atherosclerosis. Given that Treg responses affected by CCR4 deficiency in mice with early atherosclerotic lesions may be different from those in mice with advanced atherosclerotic lesions, we speculate that the difference in Treg responses may be responsible for the inconsistent findings on the role of CCR4 deficiency in atherosclerosis.

Our data show that interactions between CCR4 and CCL17/CCL22 may promote Treg-skewed responses in lymphoid tissues and atherosclerotic lesions. However, studies in hypercholesterolemic mice demonstrated that genetic deficiency of CCL17 ameliorated atherosclerotic lesion development by promoting Treg accumulation in lymphoid tissues and atherosclerotic lesions independently of the CCL17–CCR4 axis.<sup>22,24</sup> This suggests that CCL17 restrains Treg homeostasis and accelerates atherosclerosis, although the role of the CCL17–CCR4 axis in the regulation of Treg homeostasis and atherosclerosis remains elusive. A study in mouse models of myocardial injury showed that CCL17 deficiency increased Treg recruitment to the heart and attenuated myocardial inflammation and injury, which may be attributed to the cancelation of competitive inhibition of CCL22-mediated Treg chemotaxis by CCL17,<sup>41</sup> suggesting that CCL17 may impair Treg homeostasis via the inhibition of the CCL22–CCR4 axis. Given the crucial role of the CCL22–CCR4 axis in mediating the Treg suppressive function described above,<sup>28</sup> we propose that the CCL22–CCR4 axis plays a dominant role in the prevention of atherosclerosis by promoting Treg function, which may be inhibited by stimulating the CCL17–CCR4 axis. However, the specific roles of the CCL17–CCR4 and CCL22–CCR4 axes in the various stages of atherosclerosis have not been fully elucidated and further studies are needed.

This study has several limitations. Although Treg transfer experiments revealed a critical role for CCR4 in Tregs in protecting against early atherosclerosis, the use of conditional knockout mice would provide additional definitive evidence. Our data were obtained from animal experiments, and investigations using human samples will be needed to translate our findings to clinical settings.

In conclusion, we demonstrated that CCR4 protects against early atherosclerosis by favorably modulating the balance between proatherogenic Th1 cell responses and atheroprotective Treg responses in lymphoid tissues and atherosclerotic lesions. We showed that CCR4 expression on Tregs is critical for suppressing Th1 cell responses and mediating Treg migration to the atherosclerotic aorta. Our data suggest that CCR4 is an important negative regulator of atherosclerosis.

## Methods

### Animals

All mice were male on a C57BL/6 background and fed a standard chow diet or a high-cholesterol diet containing 1.25% cholesterol (CLEA Japan, Tokyo, Japan) as indicated. Wild-type mice were obtained from CLEA Japan. *Apoe*<sup>-/-</sup><sup>11</sup> and *Ccr4*<sup>-/-</sup><sup>42</sup> mice are previously described. We crossed *Ccr4*<sup>-/-</sup> mice with *Apoe*<sup>-/-</sup> mice to obtain *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice. Kaede-Tg mice (RBRC05737)<sup>31</sup> were provided by the RIKEN BRC through the National BioResource Project of the MEXT/AMED, Japan. We housed mice in cages for each strain or treatment group in a specific pathogen-free animal facility at Kobe Pharmaceutical University. Randomization and allocation concealment were performed. Littermate mice of each genotype were randomly allocated to each experimental group. During the experiments, animal/cage location was not controlled. The investigators were not blinded to the mouse genotype or treatment allocation. The criterion for exclusion was defined

as severe body weight loss and set before the study. However, during at least 2 observations per week, we did not find such symptoms, and no mice were excluded. The experimental procedures were performed in our laboratory rooms or animal facility. All animal experiments were approved by the Animal Care Committee of Kobe Pharmaceutical University (permit numbers: 2018-008, 2019-011, 2020-050, 2021-038, 2022-005, 2023-038, and 2024-014) and conformed to the National Institutes of Health Guide for the Care and Use of Laboratory Animals and the ARRIVE guidelines (Animal Research: Reporting of *In Vivo* Experiments).

## Assessment of biochemical parameters

Under anesthesia by intraperitoneal injection of medetomidine hydrochloride (0.3 mg/kg), midazolam (4 mg/kg), and butorphanol tartrate (5 mg/kg) (all from WAKO, Osaka, Japan), blood was collected by the cardiac puncture after overnight fasting, and plasma lipid profile was analyzed as described previously.<sup>40</sup> Concentrations of plasma total cholesterol, high-density lipoprotein-cholesterol, and triglycerides were determined enzymatically using an automated chemistry analyzer (Oriental Yeast Co., Ltd., Tokyo, Japan).

## Assessment of atherosclerotic lesion

The atherosclerotic lesions in the aortic root and thoracoabdominal aorta were analyzed as described previously.<sup>40</sup> For analysis of atherosclerotic lesions in the aortic root, 5 consecutive sections (10  $\mu$ m thickness), spanning 600  $\mu$ m of the aortic sinus (150  $\mu$ m interval), were collected from each mouse and stained with hematoxylin-eosin. The lesion area of the sections was quantified using the ImageJ (National Institutes of Health). Some sections were stained with Oil Red O (Sigma) for representative photomicrographs of the aortic sinus atherosclerotic lesions. For en face analysis of thoracoabdominal aortas, the aorta was opened longitudinally and stained with Oil Red O. The proportion of the lesion area was determined using the ImageJ.

## Histological analysis of atherosclerotic lesions

Immunofluorescence staining of CCL17 and CCL22 in lymphoid tissues and atherosclerotic lesions was performed on 4% paraformaldehyde-fixed cryosections of mouse aortic roots using rabbit anti-CCL17 (1:200; abcam) or goat anti-CCL22 (1:200; R&D Systems) antibodies, followed by detection with fluorescent secondary antibodies. For costaining of macrophages in the above cryosections, anti-MOMA-2 (1:400; BMA Biomedicals) and fluorescent secondary antibodies were also used. Stained sections were digitally captured using a fluorescence microscope (BZ-X810; KEYENCE, Osaka, Japan).

For the detection of macrophages or CD4<sup>+</sup> T cells, immunohistochemistry was performed on 4% paraformaldehyde-fixed cryosections of mouse aortic roots using anti-MOMA-2 or anti-CD4 (1:100; BD Biosciences) antibodies, followed by detection with biotinylated secondary antibodies and streptavidin-horseradish peroxidase. Staining with Masson's trichrome (Muto Pure Chemicals, Tokyo, Japan) was used to delineate the fibrous area. Stained sections were digitally captured using a microscope (BZ-X810; KEYENCE), and the percentage of the stained area (the stained area per total atherosclerotic lesion area) was calculated as described previously.<sup>11</sup> CD4<sup>+</sup> T cells were quantified as described previously by counting the number of positively stained cells, which was divided by the total plaque area.<sup>11</sup> The primary and secondary antibodies used are listed in Supplementary Table 3.

## Flow cytometry

For flow cytometric analysis of lymphoid tissues, peripheral LN cells and splenocytes were isolated and stained in PBS containing 2% fetal calf serum. For analysis of immune cells within the aorta, mice were anesthetized, and the aorta was perfused with cold saline. The aorta was dissected and the adventitial tissue was carefully removed. The aorta was digested with Liberase TM (Roche Diagnostics) in plane RPMI medium at 37°C for 45 min with vortexing. For the detection of CCR4

on aortic T cells, Collagenase D (Sigma-Aldrich) was used instead of Liberase TM. A cell suspension obtained by mashing the aorta through a 70- $\mu$ m strainer was stained with antibodies specific for CD3, CD4, CD45, CCR4, Foxp3, T-bet, ROR $\gamma$ t, and GATA3. The Foxp3 staining buffer set (Thermo Fisher Scientific) was used for intracellular staining of Foxp3. In all staining procedures, Fc receptors were blocked by anti-CD16/CD32 (BD Biosciences). Flow cytometric analysis was performed with FACSaria III (BD Biosciences) using FlowJo software version 10.8.1 (Tree Star). The antibodies used were listed in Supplementary Table 4.

## Intracellular cytokine staining

Immune cells from lymphoid tissues were stimulated with 20 ng/ml phorbol 12-myristate 13-acetate (Sigma) and 1 mmol/L ionomycin (Sigma) for 5 hours in the presence of Brefeldin A (Thermo Fisher Scientific). Intracellular cytokine staining was performed as described previously.<sup>40</sup>

## Cytokine assay

In cell culture experiments, RPMI 1640 medium (Sigma) supplemented with 10% fetal calf serum, 50  $\mu$ mol/L 2 $\beta$ -mercaptoethanol, and antibiotics was used. The production of several major cytokines from CD4<sup>+</sup> T cells was examined as described previously.<sup>30</sup> Splenic CD4<sup>+</sup> T cells ( $1 \times 10^5$  cells) isolated using MACS (Miltenyi Biotec) were stimulated with plate-bound anti-CD3 (10  $\mu$ g/mL, clone 145-2C11; BD Biosciences) and soluble anti-CD28 antibodies (2  $\mu$ g/mL, clone 37.51; BD Biosciences) in 96-well round-bottomed plates for 48 hours. The concentrations of IL-4, IL-10, IL-17, and IFN- $\gamma$  in culture supernatants were determined by ELISA using paired antibodies specific for corresponding cytokines (R&D Systems). The levels of multiple inflammation-associated cytokines and chemokines in culture supernatants were determined using a Mouse Cytokine Array Kit according to the manufacturer's instructions (R&D Systems).

In some experiments, CD4<sup>+</sup>CD25<sup>+</sup> Tregs and CD4<sup>+</sup>CD25<sup>-</sup> T cells were purified from the peripheral LNs and spleen of *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice using a CD4<sup>+</sup>CD25<sup>+</sup> Regulatory T Cell Isolation Kit (Miltenyi Biotec) and anti-CD4 beads (Miltenyi Biotec) according to the manufacturer's instructions. The purity of each population was >95%, as determined by flow cytometric analysis. CD11c<sup>+</sup> DCs were isolated from spleen of *Apoe*<sup>-/-</sup> mice treated with Liberase TM using MACS (Miltenyi Biotec). The purity of the CD11c<sup>+</sup> population was approximately 93%, as determined by flow cytometric analysis. CD4<sup>+</sup>CD25<sup>-</sup> T cells ( $1 \times 10^5$  cells) from *Apoe*<sup>-/-</sup> mice and splenic CD11c<sup>+</sup> DC ( $2 \times 10^4$  cells) were cocultured with or without CD4<sup>+</sup>CD25<sup>+</sup> Tregs ( $1.25 \times 10^4$  cells) from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice in the presence of soluble anti-CD3 antibody (0.5  $\mu$ g/mL) in 96-well round-bottomed plates. Culture supernatants were collected at 48 hours and analyzed by ELISA for IFN- $\gamma$  as described above.

## Treg suppression assay

For analysis of the *in vitro* suppressive function of Tregs, CD4<sup>+</sup>CD25<sup>+</sup> Tregs and CD4<sup>+</sup>CD25<sup>-</sup> T cells were purified from peripheral LNs and spleen of *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice as described above. Purified CD4<sup>+</sup>CD25<sup>+</sup> Tregs from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice were cocultured with carboxyfluorescein diacetate succinimidyl ester (Thermo Fisher Scientific)-labeled CD4<sup>+</sup>CD25<sup>-</sup> conventional T cells ( $2.5 \times 10^4$  cells) from *Apoe*<sup>-/-</sup> mice at the indicated ratios in the presence of mitomycin C (WAKO)-treated antigen-presenting cells ( $5 \times 10^4$  cells) and soluble anti-CD3 antibody (0.5  $\mu$ g/mL) in 96-well round-bottomed plates. The cocultured cells were maintained at 37°C with 5% CO<sub>2</sub> for 3 days. The proliferation of carboxyfluorescein diacetate succinimidyl ester-labeled CD4<sup>+</sup>CD25<sup>-</sup> conventional T cells was analyzed by flow cytometry.

## Flow cytometric analysis of DC phenotypic changes

CD4<sup>+</sup>CD25<sup>+</sup> Tregs and CD4<sup>+</sup>CD25<sup>-</sup> T cells were purified from peripheral LNs and spleen of *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice, and CD11c<sup>+</sup> DCs were isolated from the spleen of *Apoe*<sup>-/-</sup> mice as described above. CD4<sup>+</sup>CD25<sup>-</sup> T cells (5×10<sup>4</sup> cells) from *Apoe*<sup>-/-</sup> mice or a mixture of CD4<sup>+</sup>CD25<sup>+</sup> Tregs (5×10<sup>4</sup> cells) from *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice and CD4<sup>+</sup>CD25<sup>-</sup> T cells (5×10<sup>4</sup> cells) from *Apoe*<sup>-/-</sup> mice were cocultured with splenic CD11c<sup>+</sup> DCs (4×10<sup>4</sup> cells) in the presence of soluble anti-CD3 antibody (0.1 µg/mL) in 96-well round-bottomed plates. After 48 hours of coculture, the cells were collected, stained with antibodies specific for IAb, CD11c, CD80, CD86, and 7-amino-actinomycin D (Sigma-Aldrich), and analyzed using FACSaria III (BD Biosciences).

## Quantitative reverse transcription PCR analysis

Using TRIzol reagent (Thermo Fisher Scientific), we extracted total RNA from aorta which was perfused with cold saline and subsequently soaked in RNA later (Thermo Fisher Scientific). After the isolation of CD4<sup>+</sup>CD25<sup>+</sup> Tregs and CD4<sup>+</sup>CD25<sup>-</sup> T cells as described above, we extracted total RNA from the cells using an RNeasy Mini Kit (Qiagen). A PrimeScript RT reagent Kit (Takara, Shiga, Japan) was used for reverse transcription. Quantitative PCR analysis was conducted using a TB Green Ex Taq (Takara) and a StepOnePlus Real-Time PCR System (Thermo Fisher Scientific) according to the manufacturer's instructions. The primers used are listed in Supplementary Table 5. Amplification reactions were performed in duplicate and fluorescence curves were analyzed with the included software. GAPDH was used as an endogenous control reference.

## In vivo Treg homing assay

CD4<sup>+</sup>CD25<sup>+</sup> Tregs were purified from the peripheral LNs and spleen of *Apoe*<sup>-/-</sup> Kaede-Tg or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> Kaede-Tg mice as described above and were intravenously injected into recipient 18-week-old *Apoe*<sup>-/-</sup> mice (1×10<sup>6</sup> cells per mouse) on a high-cholesterol diet containing 1.25% cholesterol via the tail vein. At 20 hours after transfer, the Kaede<sup>+</sup> Treg proportions in the peripheral LNs, spleen, para-aortic LNs, and aorta of *Apoe*<sup>-/-</sup> mice were analyzed by flow cytometry.

## Analysis of atherosclerotic lesions in Treg-transferred mice

To clarify the role of CCR4 expression in Tregs in regulating atherosclerosis, CD4<sup>+</sup>CD25<sup>+</sup> Tregs were purified from the peripheral LNs and spleen of *Apoe*<sup>-/-</sup> or *Ccr4*<sup>-/-</sup>*Apoe*<sup>-/-</sup> mice as described above and were intravenously injected into 12-week-old recipient *Apoe*<sup>-/-</sup> mice (1×10<sup>6</sup> cells per mouse) on a standard chow diet via the tail vein. The atherosclerotic lesions were analyzed 4 weeks after transfer as described above.

## Statistical analysis

Normality was assessed by Shapiro-Wilk normality test. Two-tailed Student's *t*-test or Mann-Whitney *U*-test was used to detect significant differences between 2 groups when appropriate. One-way ANOVA followed by Tukey's multiple comparisons test or 2-way ANOVA followed by Tukey's multiple comparisons test was performed for multiple groups where appropriate. A value of *P*<0.05 was considered statistically significant. No data were excluded from the analysis. The investigators were not blinded to the data analysis. For statistical analysis, GraphPad Prism version 9.0 (GraphPad Software Inc.) was used.

## Acknowledgements

Kaede-Tg mice (RBRC05737) were provided by the RIKEN BRC through the National BioResource Project of the MEXT/AMED, Japan.

## Sources of Funding

This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers JP21K08042 and JP24K11301 (to NS) and research grants from Pfizer Japan Inc. (to YR), Astellas Pharma Inc. (to YR), Novartis Pharma K.K. (to NS), Senshin Medical Research Foundation (to NS), and ONO Medical Research Foundation (to NS).

## Disclosures

None.

## Author contributions

NS conceived the study. TT and NS designed the research, performed experiments, carried out data analyses, and wrote the manuscript. AK, HZA, and KI performed experiments, contributed to data interpretation, and reviewed the manuscript. SH, KM, KIH, TN, and YR contributed to data interpretation and reviewed and revised the manuscript.

## References

1. Weber C, Habenicht AJR, von Hundelshausen P (2023) **Novel mechanisms and therapeutic targets in atherosclerosis: inflammation and beyond** *Eur Heart J* **44**:2672–2681
2. Ridker PM *et al.* (2017) **Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease** *N Engl J Med* **377**:1119–1131
3. Tardif JC *et al.* (2019) **Efficacy and Safety of Low-Dose Colchicine after Myocardial Infarction** *N Engl J Med* **381**:2497–2505
4. Roy P, Orecchioni M, Ley K (2022) **How the immune system shapes atherosclerosis: roles of innate and adaptive immunity** *Nat Rev Immunol* **22**:251–265
5. Fernandez DM *et al.* (2019) **Single-cell immune landscape of human atherosclerotic plaques** *Nat Med* **25**:1576–1588
6. Drobni ZD *et al.* (2020) **Association Between Immune Checkpoint Inhibitors With Cardiovascular Events and Atherosclerotic Plaque** *Circulation* **142**:2299–2311
7. Gupta S, Pablo AM, Jiang X, Wang N, Tall AR, Schindler C (1997) **IFN-gamma potentiates atherosclerosis in ApoE knock-out mice** *J Clin Invest* **99**:2752–2761
8. Buono C, Binder CJ, Stavrakis G, Witztum JL, Glimcher LH, Lichtman AH (2005) **T-bet deficiency reduces atherosclerosis and alters plaque antigen-specific immune responses** *Proc Natl Acad Sci U S A* **102**:1596–1601
9. Muller FS *et al.* (2023) **Autoantibodies against the chemokine receptor 3 predict cardiovascular risk** *Eur Heart J* **44**:4935–4949
10. Ait-Oufella H *et al.* (2006) **Natural regulatory T cells control the development of atherosclerosis in mice** *Nat Med* **12**:178–180
11. Sasaki N, Yamashita T, Takeda M, Shinohara M, Nakajima K, Tawa H, Usui T, Hirata K (2009) **Oral anti-CD3 antibody treatment induces regulatory T cells and inhibits the development of atherosclerosis in mice** *Circulation* **120**:1996–2005
12. Klingenberg R *et al.* (2013) **Depletion of FOXP3+ regulatory T cells promotes hypercholesterolemia and atherosclerosis** *J Clin Invest* **123**:1323–1334
13. Tanaka T, Sasaki N, Rikitake Y (2021) **Recent Advances on the Role and Therapeutic Potential of Regulatory T Cells in Atherosclerosis** *J Clin Med* **10**
14. Noels H, Weber C, Koenen RR (2019) **Chemokines as Therapeutic Targets in Cardiovascular Disease** *Arterioscler Thromb Vasc Biol* **39**:583–592
15. Griffith JW, Sokol CL, Luster AD (2014) **Chemokines and chemokine receptors: positioning cells for host defense and immunity** *Annu Rev Immunol* **32**:659–702
16. Yoshie O (2021) **CCR4 as a Therapeutic Target for Cancer Immunotherapy** *Cancers* **13**

17. Yuan Q, Bromley SK, Means TK, Jones KJ, Hayashi F, Bhan AK, Luster AD (2007) **CCR4-dependent regulatory T cell function in inflammatory bowel disease** *J Exp Med* **204**:1327–1334
18. Faustino L, da Fonseca DM, Takenaka MC, Mirotti L, Florsheim EB, Guerreschi MG, Silva JS, Basso AS, Russo M (2013) **Regulatory T cells migrate to airways via CCR4 and attenuate the severity of airway allergic inflammation** *J Immunol* **190**:2614–2621
19. Campbell JJ *et al.* (1999) **The chemokine receptor CCR4 in vascular recognition by cutaneous but not intestinal memory T cells** *Nature* **400**:776–780
20. Montane J, Bischoff L, Soukhatcheva G, Dai DL, Hardenberg G, Levings MK, Orban PC, Kieffer TJ, Tan R, Verchere CB (2011) **Prevention of murine autoimmune diabetes by CCL22-mediated Treg recruitment to the pancreatic islets** *J Clin Invest* **121**:3024–3028
21. Greaves DR *et al.* (2001) **Linked chromosome 16q13 chemokines, macrophage-derived chemokine, fractalkine, and thymus- and activation-regulated chemokine, are expressed in human atherosclerotic lesions** *Arterioscler Thromb Vasc Biol* **21**:923–929
22. Weber C *et al.* (2011) **CCL17-expressing dendritic cells drive atherosclerosis by restraining regulatory T cell homeostasis in mice** *J Clin Invest* **121**:2898–2910
23. Kimura S, Wang KY, Yamada S, Guo X, Nabeshima A, Noguchi H, Watanabe T, Harada M, Sasaguri Y (2015) **CCL22/Macrophage-derived Chemokine Expression in Apolipoprotein E-deficient Mice and Effects of Histamine in the Setting of Atherosclerosis** *J Atheroscler Thromb* **22**:599–609
24. Döring Y *et al.* (2024) **Identification of a non-canonical chemokine-receptor pathway suppressing regulatory T cells to drive atherosclerosis** *Nat Cardiovasc Res* **3**:221–242
25. Alferink J *et al.* (2003) **Compartmentalized production of CCL17 in vivo: strong inducibility in peripheral dendritic cells contrasts selective absence from the spleen** *J Exp Med* **197**:585–599
26. Tang HL, Cyster JG (1999) **Chemokine Up-regulation and activated T cell attraction by maturing dendritic cells** *Science* **284**:819–822
27. Lee I, Wang L, Wells AD, Dorf ME, Ozkaynak E, Hancock WW (2005) **Recruitment of Foxp3+ T regulatory cells mediating allograft tolerance depends on the CCR4 chemokine receptor** *J Exp Med* **201**:1037–1044
28. Rapp M *et al.* (2019) **CCL22 controls immunity by promoting regulatory T cell communication with dendritic cells in lymph nodes** *J Exp Med* **216**:1170–1181
29. Sakaguchi S, Mikami N, Wing JB, Tanaka A, Ichiyama K, Ohkura N (2020) **Regulatory T Cells and Human Disease** *Annu Rev Immunol* **38**:541–566
30. Matsumoto T *et al.* (2016) **Overexpression of Cytotoxic T-Lymphocyte-Associated Antigen-4 Prevents Atherosclerosis in Mice** *Arterioscler Thromb Vasc Biol* **36**:1141–1151

31. Tomura M, Yoshida N, Tanaka J, Karasawa S, Miwa Y, Miyawaki A, Kanagawa O (2008) **Monitoring cellular movement in vivo with photoconvertible fluorescence protein “Kaede” transgenic mice** *Proc Natl Acad Sci U S A* **105**:10871–10876
32. Winkels H *et al.* (2018) **Atlas of the Immune Cell Repertoire in Mouse Atherosclerosis Defined by Single-Cell RNA-Sequencing and Mass Cytometry** *Circ Res* **122**:1675–1688
33. Kasahara K, Sasaki N, Amin HZ, Tanaka T, Horibe S, Yamashita T, Hirata KI, Rikitake Y (2022) **Depletion of Foxp3(+) regulatory T cells augments CD4(+) T cell immune responses in atherosclerosis-prone hypercholesterolemic mice** *Heliyon* **8**
34. Winkels H *et al.* (2017) **CD27 co-stimulation increases the abundance of regulatory T cells and reduces atherosclerosis in hyperlipidaemic mice** *Eur Heart J* **38**:3590–3599
35. Dinh TN, Kyaw TS, Kanellakis P, To K, Tipping P, Toh BH, Bobik A, Agrotis A (2012) **Cytokine therapy with interleukin-2/anti-interleukin-2 monoclonal antibody complexes expands CD4+CD25+Foxp3+ regulatory T cells and attenuates development and progression of atherosclerosis** *Circulation* **126**:1256–1266
36. Kasahara K, Sasaki N, Yamashita T, Kita T, Yodoi K, Sasaki Y, Takeda M, Hirata K (2014) **CD3 antibody and IL-2 complex combination therapy inhibits atherosclerosis by augmenting a regulatory immune response** *J Am Heart Assoc* **3**
37. Steffens S, Burger F, Pelli G, Dean Y, Elson G, Kosco-Vilbois M, Chatenoud L, Mach F (2006) **Short-term treatment with anti-CD3 antibody reduces the development and progression of atherosclerosis in mice** *Circulation* **114**:1977–1984
38. Takeda M, Yamashita T, Sasaki N, Nakajima K, Kita T, Shinohara M, Ishida T, Hirata K (2010) **Oral administration of an active form of vitamin D3 (calcitriol) decreases atherosclerosis in mice by inducing regulatory T cells and immature dendritic cells with tolerogenic functions** *Arterioscler Thromb Vasc Biol* **30**:2495–2503
39. Sasaki N *et al.* (2017) **UVB Exposure Prevents Atherosclerosis by Regulating Immunoinflammatory Responses** *Arterioscler Thromb Vasc Biol* **37**:66–74
40. Tanaka T *et al.* (2024) **Novel UV-B Phototherapy With a Light-Emitting Diode Device Prevents Atherosclerosis by Augmenting Regulatory T-Cell Responses in Mice** *J Am Heart Assoc* **13**
41. Feng G *et al.* (2022) **CCL17 Aggravates Myocardial Injury by Suppressing Recruitment of Regulatory T Cells** *Circulation* **145**:765–782
42. Matsuo K, Itoh T, Koyama A, Imamura R, Kawai S, Nishiwaki K, Oiso N, Kawada A, Yoshie O, Nakayama T (2016) **CCR4 is critically involved in effective antitumor immunity in mice bearing intradermal B16 melanoma** *Cancer Lett* **378**:16–22

## Editors

Reviewing Editor

**Carlos Escande**

Institut Pasteur de Montevideo, Montevideo, Uruguay

Senior Editor

**Tadatsugu Taniguchi**

University of Tokyo, Tokyo, Japan

**Reviewer #1 (Public review):**

Summary:

The article provides valuable information on the role of CCR4 in an inflammatory condition, namely, the arteriosclerosis plaque. The data demonstrated that in the absence of CCR4, the Th1 cells infiltrated the plaque and Tregs lost its functions. The data are clear and well-presented. Mostly importantly, the data on CCR4-specific deficiency in Regulatory T cells is more impressive.

Strengths:

The data are clear, well performed, and interesting in focusing on the plaque and compared to peripheral organs. The disease is relevant and the data could be used to understand the risk of patients under immunomodulator use.

Weaknesses:

Still, we don't know the mechanism, besides migration.

<https://doi.org/10.7554/eLife.101830.1.sa2>

**Reviewer #2 (Public review):**

Summary:

Tanaka et al. investigated the role of CCR4 in early atherosclerosis, focusing on the immune modulation elicited by this chemokine receptor under hypercholesterolemia. The study found that Ccr4 deficiency led to qualitative changes in atherosclerotic plaques, characterized by an increased inflammatory phenotype. The authors further analyzed the CD4 T cell immune response in para-aortic lymph nodes and atherosclerotic aorta, showing an increase mainly in Th1 cells and the Th1/Treg ratio in Ccr4<sup>-/-</sup>Apoe<sup>-/-</sup> mice compared to Apoe<sup>-/-</sup> mice. They then focused on Tregs, demonstrating that Ccr4 deficiency impaired their immunosuppressive function in in-vitro assays and elegantly showed that Ccr4-deficient Tregs had, as expected, impaired migration to the atherosclerotic aorta. Adoptive cell transfer of Ccr4<sup>-/-</sup> Tregs to Apoe<sup>-/-</sup> mice mimicked early atherosclerosis development in Ccr4<sup>-/-</sup>Apoe<sup>-/-</sup> mice. Therefore, this work shows that CCR4 plays an important role in early atherosclerosis but not in advanced stages.

Strengths:

Several in vivo and in vitro approaches were used to address the role of CCR4 in early atherosclerosis. Particularly, through the adoptive cell transfer of CCR4<sup>+</sup> or CCR4<sup>-</sup> Tregs, the authors aimed to directly demonstrate the role of CCR4 in Tregs' protection against early atherosclerosis.

Weaknesses:

The isolation of Tregs was inadequately controlled; they were isolated based solely on CD4 and CD25 expression. CD25 is also expressed by activated effector T cells, meaning the analyzed cells could be a pool of mainly Tregs but also include effector T cells.

The study primarily focused on Th1 and Tregs without thoroughly investigating other CD4 T cell subsets. Th17 cells are known to play an important role in atherosclerosis; non-pathogenic Th17 cells express CCR4, while pathogenic Th17 cells do not. Considering that Figure 3 shows an increased frequency of IL17-expressing CD4 T cells compared to ApoE<sup>-/-</sup> mice, and given the imprecise Treg isolation, differences in non-pathogenic Th17 cells could be contributing to the observed effects.

Furthermore, the clinical relevance of these findings is not discussed. As an initial approach, the authors could analyze public datasets to determine if certain Ccr4 single nucleotide polymorphisms correlate with a higher incidence of atherosclerosis.

<https://doi.org/10.7554/eLife.101830.1.sa1>

### Reviewer #3 (Public review):

#### Summary:

In this paper, Tanaka and colleagues address the role played by the C-C chemokine receptor 4 (CCR4) in developing early atherosclerotic plaques using ApoE-deficient mice fed with a standard chow diet as a model. Since CCR4 is expressed in several T CD4<sup>+</sup> lymphocyte subsets, the authors examined the consequences of CCR4 deficiency on the differentiation profile and traffic of T CD4<sup>+</sup> lymphocytes. By histological analysis of aortic lesions, they demonstrated that the absence of CCR4 promoted the development of early atherosclerosis, characterized by an inflammatory reaction with increased levels of macrophages and T CD4<sup>+</sup> inflammatory lymphocytes while decreased collagen content. Using flow cytometry together with mRNA expression analysis for identifying T CD4<sup>+</sup> cell subsets, the authors found that the accelerated aortic inflammation induced by CCR4 deficiency correlated with higher proliferation of T CD4<sup>+</sup> cells in lymphoid tissues, favouring the expansion of the pro-inflammatory effector Th1 cell subset, typically found in atherosclerotic lesions. Interestingly, the increased T CD4<sup>+</sup> cell response occurred despite the expansion of T CD4<sup>+</sup> Foxp3<sup>+</sup> regulatory cells (Treg), which were in higher numbers in the lymphoid tissues of CCR4-deficient mice, suggesting the absence of CCR4 interfered with the regulatory actions of Treg cells. Using in vitro and or in vivo approaches, the authors found evidence of CCR4 requirement for Treg suppressive activity and migratory capacity to inflamed aortic areas, contributing to why CCR4 deficiency induced an augmented Th1/Treg ratio in the aortic lesions. These findings might not be surprising considering the demonstrated involvement of CCR4 in driving Treg migration to inflamed tissues in immune-related pathological models and Treg-dendritic cell contact for imprinting suppressive signals. However, in previous studies using a murine model of advanced atherosclerosis, neither hematopoietic nor systemic CCR4 deficiency altered the development of the aortic lesions. The authors included a thoughtful discussion about hypothetical mechanisms explaining these contrasting results, highlighting putative differences in the role played by the CCL17/CCL22-CCR4 axis along the stages of atherosclerosis development in this murine model.

#### Major strengths and weaknesses:

The main effects of CCR4 deficiency on early atherosclerosis development and Treg functional loss are valuable and supported by collected data. In vivo studies for comparing Treg-tissue accumulation or atherosclerotic lesions in ApoE<sup>-/-</sup> mice that received Treg derived from ApoE<sup>-/-</sup> or ApoE<sup>-/-</sup>Ccr4<sup>-/-</sup> mice, strengthening results. However, an incomplete description of methods (particularly flow cytometry) and data analysis weakens some conclusions of this study. Readers should note some inconsistencies in the T CD4<sup>+</sup> response analysis in different tissues. In aortic lesions, but not in lymphoid tissues (peripheral, para-aortic, and spleen), the ratio Th1/Treg was used for evaluating the effect of CCR4 deficiency

on the profile of Th cell subsets. In lymphoid tissues, increments in the frequency of both effector Th1 and Treg were observed in CCR4-deficient Apoe<sup>-/-</sup> mice compared to CCR4-sufficient Apoe<sup>-/-</sup> mice. Therefore, it is not convincing that CCR4-deficiency shifts Th1 cell/Treg balance toward Th1 cell responses in all lymphoid tissues; this claim needs to be revised by the authors. The Treg dysfunction, caused by CCR4 deficiency, enhanced T CD4<sup>+</sup> activation and might have amplified rather than shifted, the typical biased Th1-mediated inflammatory response observed in the lymphoid tissues of hypercholesterolemic mice. A different scenario emerged in aortic lesions, where recruitment of effector Th1 cells, but not of additional effector T CD4<sup>+</sup> cell subsets expanded in lymphoid tissues, leading to a higher Th1/Treg balance. Also, effector Th17 cells seem to predominate among effector TCD45<sup>+</sup>CD3<sup>+</sup>CD4<sup>+</sup> cells in the aorta of Apoe<sup>-/-</sup> mice, and the Th1/Th17 balance appears to have increased as a consequence of CCR4 deficiency as well. Modulation of Th1/Th17 balance might be responsible for changes in the type and functional properties of recruited inflammatory cells in the aorta.

#### Study limitations:

This investigation has some limitations. Current tools for single-cell characterization have revealed the phenotypic heterogeneity and dynamics of aortic leukocytes, including T cells, which are among the principal aortic leukocytes found in mouse and human atherosclerotic lesions (doi:10.1161/CIRCRESAHA.117.312513). The flow cytometry analysis applied in this study cannot distinguish the generation of particular phenotypes within T CD4<sup>+</sup> subsets, including putative phenotypes of no-suppressive T cells expressing low levels of Foxp3, as seems could occur in other chronic inflammatory disorders (doi: 10.1038/nm.3432; doi: 10.1172/JCI79014). Limitations due to the use of a complete CCR4 knockout mouse and putative differences in CCR4-mediated mechanisms along atherosclerosis stages and in human atherosclerosis were commented on by the authors in the discussion.

#### Global Impact

This work opens the way for a deeper analysis of the contribution of CCR4 and its ligands to the activation and differentiation of T CD4<sup>+</sup> lymphocytes during atherosclerosis development, with these lymphocytes being fundamental players in the generation of pro-atherogenic and anti-atherogenic immune responses. Differences in the mechanisms mediated by the CCL17/CCL22-CCR4 axis among early and advanced atherosclerosis highlight the complex landscape to examine and validate in human samples and the need to achieve a deep knowledge for identifying genuine and safe targets capable of promoting protective anti-atherogenic immune responses.

<https://doi.org/10.7554/eLife.101830.1.sa0>