

# Dual role of FOXG1 in regulating gliogenesis in the developing neocortex via the FGF signaling pathway

Mahima Bose, Ishita Talwar, Varun Suresh, Urvi Mishra, Shiona Biswas, Anuradha Yadav, Shital T Suryavanshi, Simon Hippenmeyer, Shubha Tole 

Tata Institute of Fundamental Research, Mumbai, India • Institute of Science and Technology Austria (ISTA), Klosterneuburg, Austria

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

## eLife Assessment

This **important** study provides **convincing** evidence that developing neurons in the neocortex regulate glial cell development. The data demonstrates that the transcription factor FOXG1 negatively regulates gliogenesis by controlling the expression of a member of the FGF ligand family and by suppressing the receptor for this ligand in developing neurons. This study leads to a new understanding of the cascade of events regulating the timing of glial development in the neocortex.

<https://doi.org/10.7554/eLife.101851.1.sa3>

## Abstract

In the developing vertebrate central nervous system, neurons and glia typically arise sequentially from common progenitors. Here, we report that the transcription factor Forkhead Box G1 (*Foxg1*) regulates gliogenesis in the mouse neocortex via distinct cell-autonomous roles in progenitors and in postmitotic neurons that regulate different aspects of the gliogenic FGF signalling pathway. We demonstrate that loss of *Foxg1* in cortical progenitors at neurogenic stages causes premature astrogliogenesis. We identify a novel FOXG1 target, the pro-gliogenic FGF pathway component *Fgfr3* that is suppressed by FOXG1 cell-autonomously to maintain neurogenesis. Furthermore, FOXG1 can also suppress premature astrogliogenesis triggered by the augmentation of FGF signalling. We identify a second novel function of FOXG1 in regulating the expression of gliogenic cues in newborn neocortical upper-layer neurons. Loss of FOXG1 in postmitotic neurons non-autonomously enhances gliogenesis in the progenitors via FGF signalling. These results fit well with the model that newborn neurons secrete cues that trigger progenitors to produce the next wave of cell types, astrocytes. If FGF signalling is attenuated in *Foxg1* null progenitors, they progress to oligodendrocyte production. Therefore, loss of FOXG1 transitions the progenitor to a gliogenic state, producing either astrocytes or oligodendrocytes depending on FGF signalling levels. Our results uncover how FOXG1 integrates extrinsic signalling via the FGF

pathway to regulate the sequential generation of neurons, astrocytes, and oligodendrocytes in the cerebral cortex.

## Significance Statement

Functional brain circuitry requires the correct numbers and subtypes of neurons and glia. A fundamental feature of the vertebrate central nervous system is that common progenitors produce first neurons and then glia. Cell-intrinsic factors modulate how a progenitor responds to cell-extrinsic cues to achieve this transition. Here, we report that the transcription factor Forkhead Box G1 (*Foxg1*) regulates gliogenesis in the mouse neocortex *in vivo* by regulating the gliogenic FGF signalling pathway. Loss of *Foxg1* results in premature gliogenesis via upregulation of novel target *Fgfr3* in progenitors and dysregulation of FGF ligands in postmitotic neurons. Our findings reveal how an interplay between cell-intrinsic transcriptional regulation and cell-extrinsic cues controls the production of critical cell types in the brain.

## Introduction

A fundamental feature of the developing vertebrate central nervous system is that common progenitors produce neurons and glia sequentially. Furthermore, two broad categories of glia, astrocytes and oligodendrocytes, each subserving distinct functions, are also produced sequentially. This temporal sequence of neurogenesis, followed by astrogliogenesis and then oligogenesis, is conserved in the developing mammalian neocortex (1–3). However, the molecular mechanisms instructing a cessation of neurogenesis and initiating gliogenesis within progenitors are poorly understood, particularly in terms of how cell-intrinsic factors interact with cell-extrinsic signalling cues.

Early *in vitro* (1, 4) and *in vivo* (5) studies demonstrated that cortical progenitors first produce neurons and later glia. Quantitative MADM lineage analyses *in vivo* further extended these findings and showed that neurons, astrocytes, and oligodendrocytes arise sequentially within a clone of cells from a single progenitor (5, 6). Molecular insights into this phenomenon identified several pro-gliogenic mechanisms. These include cell-intrinsic transcription factors *Nr2f1/2*, *Zbtb20*, *Sox9*, and *Nfia* (7–10), the Notch pathway (8, 11–14); and cell-extrinsic cues such as interleukins IL6 and Cardiotrophin1 (*Ct1*), Leukemia Inhibiting Factor (*Lif*), Ciliary Neurotrophic Factor (*Cntf*), Neuropoietin (*Np*), Cardiotrophin-like Cytokine (*Clc*), Epidermal Growth Factors (*Egf*) and Fibroblast Growth Factors (*Fgf* (15–20)) which act via the Jak/Stat or Mek/Mapk signalling pathways (12, 21, 22). In these studies, the primary readout of gliogenesis was astrogliogenesis, using either morphology or molecular markers to identify glial fate. Although a few factors, such as *Sox10*, have been reported to promote oligodendrocyte precursor cell (OPC) fate (23, 24), the progenitor-level mechanisms that govern astrocyte-to-oligodendrocyte transition remain poorly understood. Furthermore, how mechanisms that maintain ongoing neurogenesis crosstalk with those that promote gliogenesis remains to be understood. Another unexplored area is how cell-intrinsic and extrinsic factors interact with or regulate each other at the level of the individual progenitor.

Forkhead family transcription factor FOXG1 plays a fundamental role in the early development of the cerebral cortex and is a well-established regulator of the sequential production of neuronal subtypes in the mammalian neocortex (25–31). *Foxg1* haploinsufficiency in humans affects a range of neurodevelopmental processes resulting in an Autism Spectrum Disorder called Foxg1

syndrome, which includes corpus callosum agenesis, microcephaly, and cognitive impairment (32). A dysregulation of *Foxg1* has also been implicated in glioblastoma pathogenesis among these individuals (29, 33).

FOXG1 is described to play a “neurogenic role” in overexpression experiments *in vitro* or cultured progenitors transplanted *in vivo*, which display decreased gliogenesis and downregulation of progliogenic pathways and markers (34–36). However, in these studies, *Foxg1* knockdown did not cause significant gliogenesis (34, 35). Furthermore, the role of FOXG1 in postmitotic neurons in regulating gliogenesis has not been examined, although these cells express *Foxg1* and are ideally positioned to signal progenitors to generate the next cell type. These are critical unanswered questions that we explored directly *in vivo*. We show that upon loss of *Foxg1*, progenitors autonomously produce glia at the expense of neurons. Using integrative analysis of the transcriptome, epigenome, and FOXG1 occupancy, we identified FGFR3, part of the established progliogenic FGF signalling pathway, as a FOXG1 target.

We show that the loss of *Foxg1* results in upregulating *Fgfr3* expression and FGF signalling in cortical progenitors. Whereas overexpression of FGF ligands induces premature gliogenesis (19), simultaneous overexpression of *Foxg1* restores neurogenesis, consistent with its established neurogenic role. Furthermore, attenuation of FGF signalling in a *Foxg1* loss-of-function (LOF) background leads to a premature production of OPCs, indicating that without FOXG1, the progenitor may progress through gliogenesis but cannot return to neurogenesis.

Independently, FOXG1 regulates cues produced by newborn postmitotic neurons that non-autonomously modulate the output of progenitors. In particular, when postmitotic neurons lack *Foxg1*, naive progenitors upregulate FGF signalling and display enhanced gliogenesis. We demonstrate that progenitors experience an enhanced level of FGF signalling due to dysregulated *Fgfl* ligand expression in postmitotic neurons upon loss of when *Foxg1*. Therefore, FOXG1 regulates both the availability of the ligand and the level of the receptor for Fgf signalling.

Our results shed light on a novel mechanism regulating the sequential generation of neurons, astrocytes, and oligodendrocytes in the cerebral cortex.

## Results

We used a *Foxg1* conditional knockout mouse line with an intrinsic reporter (*Foxg1*<sup>lox/lox</sup>; R26<sup>FRT-GFP</sup> (37)) to assess the cell-autonomous phenotypes of *Foxg1* disruption. In this line, Cre-mediated recombination of the floxed *Foxg1* allele results in the expression of flippase which then recombines a STOP-FRT-EGFP reporter inserted in the *Rosa26* locus. Since this reporter is integrated into the genome, introducing Cre recombinase into a progenitor results in GFP expression in *Foxg1*-recombined cells arising from it without dilution, serving as a label for its lineage.

### Disrupting *Foxg1* during neurogenesis causes premature gliogenesis

In wild-type mice, FOXG1 expression decreases in cortical progenitors from embryonic day (E) 15.5 to E18.5 (Supplementary Figure S1; (35)). We tested the consequences of a premature drop in FOXG1 at E15.5 when superficial layer neurons are generated in the neocortex. We introduced a plasmid encoding Cre recombinase at E15.5 using *in-utero* electroporation and examined the brains on postnatal day 14 (P14). In control brains, as expected, GFP<sup>+</sup> cells with pyramidal neuron morphologies occupied layer II/III of the neocortex, whereas cells with glial morphologies were scattered throughout the tissue. 67% of the GFP<sup>+</sup> cells also expressed neuronal marker NEUN and did not express mature astrocytic marker ALDH1L1. 16% of the GFP<sup>+</sup> cells were also ALDH1L1<sup>+</sup>,

due to the intrinsic  $R26^{FRT-GFP}$  reporter that causes EGFP expression in the entire lineage of progenitors electroporated at E15.5 (**Figure 1A–D**; Supplementary Figure S2). In contrast, Cre electroporation in  $Foxg1^{lox/lox}$  embryos results in GFP+ cells with astrocytic morphologies distributed all over the neocortex. None were NEUN+, but 56% were ALDH1L1+, confirming their astrocyte identity. We found similar results, i.e. premature gliogenesis, upon Cre electroporation at E14.5 (**Figure 1E–H**). In contrast, loss of  $Foxg1$  at E13.5 did not impede neurogenesis. The mutant cells were NEUN+ and ALDH1L1– indicating no premature gliogenesis. Consistent with the literature, there is enhanced expression of REELIN (Supplementary Figure S3; (26, 27, 38–40)).

This indicates that progenitors undergo a transition after E13.5, wherein the loss of FOXG1 from E14.5 prompts a shift from neurogenesis to gliogenesis, even if they have not yet produced the final cohorts of superficial layer neurons.

## Enhanced gliogenesis upon loss of *Foxg1* is not due to the over-proliferation of mutant astrocytes

Increased gliogenesis may arise from the over-proliferation of astrocytes and an accompanying death of neurons that have lost  $Foxg1$ . Alternatively, there could be a premature cell fate switch in the progenitors that should have produced neurons but instead produced astrocytes (**Figure 2A**). To distinguish between these scenarios, we examined the presence of KI67 in cells at one day (E16.5) or three days (E18.5) after Cre electroporation at E15.5. Similar numbers of electroporated (EGFP+) cells displayed KI67 in control and  $Foxg1^{lox/lox}$  brains, indicating that loss of  $Foxg1$  did not induce overproliferation (**Figure 2B**, B'; Supplementary Figure S4). Furthermore, no difference was observed in the expression of a cell death marker Cleaved Caspase 3 (Figure C, C'). However, consistent with a switch to gliogenesis, the  $Foxg1$  LOF cells were positive for gliogenic factor NFIA but not the neurogenic intermediate progenitor marker TBR2 (**Figure 2** D, D', E, E').


We also employed a MADM (Mosaic Analysis with Double Markers)-based strategy to distinguish between these possibilities (**Figure 2D**) (41, 42). We used the MADM-12 (M12) line (42, 43) in which the MADM GT and TG cassettes are on chromosome 12 and crossed them onto a  $Foxg1^{lox/+}$  background. In the resulting  $Foxg1$ -MADM ( $M12^{GT/TG, Foxg1}$ ) brains, a fraction of Cre-electroporated progenitors undergo recombination, resulting in progeny that is either  $Foxg1^{-/-}$ , GFP+ (green) or  $Foxg1^{+/+}$  tdTomato+ (red) cells. Since these red and green cells arise from single progenitors, the MADM system enables an accurate analysis of proliferation versus cell fate transformation upon disruption of  $Foxg1$ . A fraction of progenitors in the MADM paradigm undergoes a different recombination pattern, resulting in a yellow progeny due to the expression of both GFP and tdTomato (41, 42), which were not scored. We performed *in-utero* electroporation of Cre in  $Foxg1$ -MADM embryos in E14.5/E15.5 embryos. We analysed the brains at P14 (**Figure 2G**). Green and red cells were scored for neuronal or glial identity based on morphology (5, 20). Of the total control (red, +/+) cells, 61.9% were neurons, and 38.1% were glia. In contrast, of the total number of  $Foxg1$  mutant (green, –/–) cells, 16.7% were neurons, and 83.3% were glia (**Figure 2H**).

In summary, the results demonstrate that loss of  $Foxg1$  results in a premature cell fate switch in neurogenic progenitors that makes them gliogenic.

## Loss of *Foxg1* causes upregulation of *Fgfr3* in progenitors and enhanced FGF signalling

We investigated how FOXG1 regulates the transcriptional landscape in cortical progenitors to maintain a temporal control of neurogenesis and gliogenesis. We used the inducible  $hGFAPCreERT2$  (hGCE) driver to achieve widespread recombination in apical progenitors at






**Figure 1**

### Loss of *Foxg1* causes premature gliogenesis.

(A–D) Cre electroporation at E15.5 in control (A, *Foxg1*<sup>lox/+</sup>; *R26*<sup>FRT-GFP</sup>) and *Foxg1* LOF (B, *Foxg1*<sup>lox/lox</sup>; *R26*<sup>FRT-GFP</sup>) embryos, followed by analysis at P14. (C, D) 67.3% of GFP+ cells colocalised with NeuN in control brains and 1% in *Foxg1* LOF brains. 16% of GFP+ cells colocalised with ALDH1L1 in control brains and 56% in *Foxg1* LOF brains. *n* = 2151 (Control), 2761 (*Foxg1* LOF) cells from *N* = 3 brains (biologically independent replicates). (E–H) Cre electroporation at E14.5 in control (E, *Foxg1*<sup>lox/+</sup>; *R26*<sup>FRT-GFP</sup>) and *Foxg1* LOF (F, *Foxg1*<sup>lox/lox</sup>; *R26*<sup>FRT-GFP</sup>) embryos, followed by analysis at P14. (G, H) 77.7% of GFP+ cells colocalised with NeuN in control brains and 0% in *Foxg1* LOF brains. 20.1% of GFP+ cells colocalised with ALDH1L1 in control brains and 55.7% in *Foxg1* LOF brains. *n* = 3160 (Control), 2978 (*Foxg1* LOF) cells from *N* = 3 brains (biologically independent replicates). In each row (A, B, E, F), the boxes in the leftmost low magnification panels indicate approximate regions shown in either the NEUN or ALDH1L1 high magnification panels. Filled arrowheads depict colocalisation, and open arrowheads depict non-colocalisation of marker and electroporated cells. Statistical test: two-tailed unpaired *t*-test. \* (*p* < 0.05), \*\* (*p* < 0.01), \*\*\* (*p* < 0.001), \*\*\*\* (*p* < 0.0001). All scale bars are 50  $\mu$ m.



**Figure 2**

***Foxg1* LOF leads to premature glial fate acquisition in progenitors but no proliferation defects.**

(A) Schematic depicting the two alternative mechanisms that could result in enhanced gliogenesis upon loss of *Foxg1*: a change in cell type produced by the progenitor, i.e. “fate switch” or increase in proliferation of astrocytes accompanied by neuronal cell death. (B, B', C, C') Images of the ventricular and sub-ventricular zone (the dashed line indicates the ventricle boundary). Cre electroporation at E15.5 in control (*Foxg1*<sup>lox/+</sup>; *R26*<sup>FRT-GFP</sup>) and *Foxg1* LOF (*Foxg1*<sup>lox/lox</sup>; *R26*<sup>FRT-GFP</sup>) embryos, followed by analysis at E18.5. Proliferation marker KI67 (B, B') colocalises with similar numbers of GFP+ cells in control and *Foxg1* LOF brains. Cell death marker Cleaved Caspase 3 (C, C') does not reveal differences in colocalisation with GFP+ control and *Foxg1* LOF cells. In contrast, glial progenitor markers NFIA (D, D') display increased colocalisation with GFP+ cells in *Foxg1* LOF (51.7%) compared with controls (12.5%). Neurogenic intermediate progenitor marker TBR2 (E, E') displays decreased colocalisation with GFP+ cells in *Foxg1* LOF brains (0%) compared with controls (8%). n= 3590 (control), 2100 (mutant) cells from N=3 brains (biologically independent replicates). (G) Schematic depicting the genotype and corresponding fluorescent labels resulting from the MADM recombination events. (E–F) Cre electroporation at E14.5/E15.5 in *Foxg1*-MADM brains (*M12*<sup>GT/TG</sup>, *Foxg1*) analysed at P14. Green (*Foxg1*<sup>−/−</sup>) and red (*Foxg1*<sup>+/+</sup>) cells were scored based on neuronal (open arrowheads, E) or glial (arrowheads, E) morphology. (F) represents the number of neurons or glia as a percentage of the total population of neurons+glia of each genotype: control (red; +/+) or *Foxg1* mutant (green; −/−) neurons. n= 354 cells from N=5 brains (biologically independent replicates). Statistical test: two-tailed unpaired t-test. \*(*p*<0.05), \*\*(*p*<0.01), \*\*\*(*p*<0.001), \*\*\*\*(*p*<0.0001). All scale bars: 50µm.



embryonic ages upon induction with tamoxifen (44), (45). First, we ascertained that hGCE-driven loss of *Foxg1* at E15.5 recapitulates the premature gliogenesis phenotype we described in **Figure 1** (Supplementary Figure S5A). Also, two days after tamoxifen induction (E15.5 → E17.5), *Foxg1* LOF cells were positive for gliogenic factor NFIA, similar to that seen upon Cre electroporation (Supplementary Figure S5B, C; **Figure 2C**). To identify pathways that regulate the premature gliogenesis induced by the loss of *Foxg1*, we collected FACS-purified control and mutant cells at E17.5, 48 hours post-tamoxifen administration at E15.5, for transcriptomic analysis (Supplementary Figure S5D, D'). As expected, the loss of *Foxg1* caused the upregulation of known gliogenic factors such as *Nfia* and *Olig3* and the downregulation of neurogenic genes such as *Neurod2* and neuronal marker *Pou3f1* (**Figure 3A**).

We further filtered the list of genes upregulated upon loss of *Foxg1* using a strategy designed to narrow down candidates likely to drive gliogenesis. First, we examined a set of genes known to display FOXG1 occupancy at E15.5 (46). Second, we reasoned that cell fate transitions are driven by genes maintained in states poised for transcriptional activation or repression by displaying both epigenetic marks (47). Therefore, we analysed a dataset from E15.5 radial glial cells to identify genes that carry bivalent H3K4Me3 and H3K27Me3 marks in gene regulatory regions. Third, we obtained a list of genes enriched in cortical astrocytes (48).


The overlap of these three datasets yielded a shortlist of 19 genes representing potential FOXG1 targets that may be suppressed during neurogenesis (**Figure 3B**, C). Finally, we compared this multimodal analysis with the list of genes we discovered to be upregulated upon loss of *Foxg1* and identified four targets: *Col2a1*, *Dpysl4*, *Fgfr3*, and *Pax6* (**Figure 3D**). The presence of *Fgfr3* in this set presented the exciting possibility that FOXG1 may function to suppress the potent progliogenic FGF signalling pathway (19) in neurogenic cortical progenitors.

In summary, we identified *Fgfr3* as a novel FOXG1 target. Upregulation of *Fgfr3* may mediate the gliogenic effects of loss of *Foxg1*. Consistent with this hypothesis, the MAPK/ERK pathway, which mediates FGF signalling, was identified as a prominent upregulated pathway in a KEGG pathway analysis of the RNA-seq data (**Figure 3E**). Activation of this pathway results in the phosphorylation of ERK1/2 and its translocation to the nucleus to regulate the downstream targets of FGFR activation (49). To assess whether the loss of *Foxg1* indeed has functional consequences on Fgf signalling, we examined FGFR3 and nuclear phospho-ERK1/2 levels in control and *Foxg1* mutant cortical progenitors. Loss of *Foxg1* at E15.5 led to an increase in FGFR3 labelling and nuclear localisation of phospho-ERK1/2 by E17.5 (**Figure 3F–H**). We quantified nuclear FOXG1 levels in wild-type apical progenitors. We found a significant decrease from E15.5 to E18.5, consistent with previous findings (35), suggesting an endogenous mechanism for *Fgfr3* upregulation with the initiation of gliogenesis (Supplementary Figure S1).

These results demonstrate that FOXG1 suppresses *Fgfr3* gene expression and the MAPK/ERK pathway, and loss of *Foxg1* results in an enhancement of this pro-gliogenic pathway. The decline in nuclear FOXG1 by E18.5 suggests an endogenous mechanism for the transition of neurogenesis to gliogenesis in apical progenitors.

## ***Foxg1* overexpression cell autonomously suppresses FGF-induced astrogliogenesis**

Exogenous FGF8 is known to have potent progliogenic effects in the neocortex (19), which our results recapitulated (**Figure 4A**, B). As expected, electroporation at E15.5 of a control construct encoding EGFP did not impede neurogenesis. In wild-type mice, 100% of the electroporated GFP+ cells were NEUN+ and occupied the superficial layers of the neocortex by P7 (**Figure 4A**). Similar to the findings of (19), co-electroporation of constructs encoding *Fgf8*+*Egfp* induced premature gliogenesis, resulting in 85% of the GFP+ cells displaying astrocytic morphologies and glial marker SOX9 (**Figure 4B**). *Foxg1* overexpression alone did not affect neurogenesis but gave



**Figure 3**

### FOXG1 binds and regulates the expression of *Fgfr3*.

(A) RNA-seq analysis of FACS-purified Control and *Foxg1* LOF progenitors harvested two days after labelling at E15.5. Gliogenic factors such as *Nfia*, *Id3*, and *Olig3* are upregulated, and neuronal markers such as *Pou3f1* and *Robo4* are downregulated. (B–D) Multimodal analysis comparing FOXG1 occupancy (ChIP-seq) and bivalent epigenetic marks (H3K4Me3 and H3K27Me3) and astrocyte-enriched genes from (48) reveals a list of 19 genes common to each dataset (B). Four of these are upregulated upon loss of *Foxg1*, including the known gliogenic gene *Fgfr3* (D). FOXG1 occupies a -26kb enhancer region of *Fgfr3* (C). (E) KEGG analysis of the upregulated genes from (A) identifies the MAPK signalling pathway downstream of FGF signalling. (F) Loss of *Foxg1* from E15.5 progenitors (*hGFAPCreERT2*, tamoxifen at E15.5) causes upregulation of FGFR3 receptor by E17.5 as seen in cells near the VZ of the somatosensory cortex. Boxes (F) indicate the regions in high magnification shown in the adjacent panels (G). Dashed circles outline the ROIs identified in the DAPI channel used for intensity quantification in (H). (G; n=50 (Control and *Foxg1* LOF) ROIs from N=3 brains) and phosphorylated-ERK1/2 (H; n= 67 (Control) and 89 (*Foxg1* LOF) cells from N=3 brains (biologically independent replicates). Statistical test: Mann-Whitney test \* ( $p<0.05$ ), \*\*( $p<0.01$ ), \*\*\*( $p<0.001$ ), \*\*\*\*( $p<0.0001$ ). All scale bars: 50 $\mu$ m.



rise to neurons, some of which occupied the superficial layer and some displayed migration deficits, as previously shown ((37 [↗](#)); **Figure 4C** [↗](#)). Note that this experiment did not use the intrinsic *R26<sup>FRT-GFP</sup>* reporter. Electroporation-based introduction of GFP is expected to dilute with continued progenitor proliferation. Therefore, the control brains displayed GFP in neurons that were born at E15.5, and not in glia that arose subsequently from the same progenitors.

We tested whether FOXG1 can override FGF8-induced premature gliogenesis by co-electroporating both constructs (*Foxg1-Egfp* and *Fgf8*, **Figure 4D** [↗](#)). This led to a recovery of the neuronal fate in a cell-autonomous manner, such that GFP+ cells also displayed NEUN but not SOX9 (**Figure 4D** [↗](#)). The electroporated cells also displayed migration deficits consistent with *Foxg1* overexpression (37 [↗](#)). An interesting additional result was that while these FOXG1-EGFP+ cells themselves expressed neuronal marker NEUN, they were surrounded by SOX9+ cells consistent with the non-cell-autonomous gliogenic effects of *Fgf8* electroporation (**Figure 4D** [↗](#)).


In summary, FOXG1 is sufficient to cell-autonomously suppress the pro-gliogenic effects of FGF8, establishing it as a regulator of FGF signalling within cortical progenitors.

## Postmitotic neuron-specific loss of *Foxg1* leads to premature gliogenesis in cortical progenitors

Postmitotic neurons provide “feedback instruction” to progenitors via cues that modulate the cell types that arise subsequently (50 [↗](#), 51 [↗](#)). We examined a publicly available transcriptomic dataset from CUX2+ upper-layer neurons isolated at stages from E18.5 to P48 (52 [↗](#)) and discovered that FGF family members *Fgf9*, *Fgf10*, and *Fgf18* mRNA levels peaked in the first postnatal week and dropped thereafter, consistent with the temporal profile of astrogliogenesis in the cortex (Supplementary Figure S6A, B; (53 [↗](#))). Since *Foxg1* is expressed in postmitotic neurons (29 [↗](#), 54 [↗](#)), we tested whether it may play a role in regulating gliogenic factors secreted by these cells. We used postmitotic neuron-specific *NexCre* to disrupt *Foxg1* and examined these brains at birth (Supplementary Figure S6C, D). We found a significant increase in the levels of gliogenic factor SOX9 in the ventricular zone progenitors and an increased number of SOX9+ cells in the mantle compared to that in control brains, indicating a non-autonomous effect of neuron-specific loss of *Foxg1* on progenitors (**Figure 5A–D** [↗](#)). Consistent with enhanced gliogenesis, there was an apparent increase in astrocyte marker ALDH1L1+ cells in the entire cortical plate (Supplementary Figure S6E).

To identify potential gliogenic molecules regulated by FOXG1 in postmitotic neurons, we performed transcriptomic analysis on cortical plate tissue isolated from control and *NexCre*-driven *Foxg1* LOF brains at birth and examined genes encoding secreted factors (**Figure 5E** [↗](#), Supplementary Table 4). Among members of the FGF family, *Fgf9* transcripts decreased, *Fgf10* displayed no change, whereas *Fgf18* displayed a 2.5-fold increase (**Figure 5E** [↗](#)). This ligand has been well characterised to be preferentially expressed in neurons in the postnatal rodent brain (55 [↗](#)), displays peak expression in CUX2+ cells in the first postnatal week (Supplementary Figure S6B), and is in the same family as FGF8 which is used to induce gliogenesis (19 [↗](#)). We examined nuclear-phosphorylated p42/44-ERK1/2, indicative of FGF signalling, and found it to display increased nuclear localisation specifically in the ventricular zone of *NexCre/+*; *Foxg1<sup>lox/lox</sup>* brains (**Figure 5H** [↗](#)). This indicates a net increase in FGF signalling experienced by the progenitors.


To further confirm whether this enhanced gliogenesis upon postmitotic neuron-specific loss of *Foxg1* is indeed via an increase in FGF signalling, we used a pharmacological inhibitor NVP-BGJ398 (Infigratinib) that blocks activation of FGFR1/2/3 (56 [↗](#)). Dams bearing control and *NexCre/+*; *Foxg1<sup>lox/lox</sup>* embryos were administered this inhibitor intraperitoneally from E14.5 to E17.5, and the brains corresponding to each condition were examined at E18.5 for nuclear pERK1/2 levels. In the presence of this inhibitor, nuclear pERK1/2 levels in control VZ cells were reduced compared with untreated embryos, indicating the drug had the predicted effect of



**Figure 4**

### ***Foxg1* suppresses FGF-induced gliogenesis.**

(A–D) *In-utero* electroporations were performed in wild-type embryos at E15.5, and the brains were analysed at P7. (A) GFP electroporation labels LII/III cells that are NEUN<sup>+</sup> (arrowheads) and SOX9<sup>−</sup> (open arrowheads). (B) Overexpression of *Fgf8* leads to premature gliogenesis, and the GFP<sup>+</sup> cells are NEUN<sup>−</sup> (open arrowheads) and SOX9<sup>+</sup> (arrowheads). (C) Overexpression of *Foxg1* produced NEUN<sup>+</sup> (arrowheads) and SOX9<sup>−</sup> (open arrowheads) neurons, some of which displayed delayed migration (black asterisk), and others migrated to the cortical plate (white asterisk), as shown in (37). (D) Overexpression of *Foxg1* together with FGF8 partially rescued neuronal fate such that GFP<sup>+</sup> cells also displayed NEUN (arrowheads) but not SOX9 (open arrowheads). In A–D, the boxes in the leftmost low magnification panels indicate approximate regions shown in the adjacent high magnification panels. (E) Quantifications of GFP<sup>+</sup> cells that are also NEUN<sup>+</sup> in each condition: 98.6% (GFP); 1.8% (*Fgf8*); 98.3% (*Foxg1*); 74.1% (*Foxg1*+*Fgf8*). (F) Quantifications of GFP<sup>+</sup> cells that are also SOX9<sup>+</sup> in each condition: 0% (*Egfp*); 87.7% (*Fgf8*+*Egfp*); 0% (*Foxg1*-*Egfp*); 24.2% (*Foxg1*-*Egfp*+*Fgf8*). n= 2123 (*Egfp*), 1643 (*Fgf8*+*Egfp*), 1357 (*Foxg1*-*Egfp*), 1924 (*Foxg1*-*Egfp*+*Fgf8*) cells each from N=3 brains (biologically independent replicates). Statistical test: Two-way ANOVA with Tukey's correction. \*(*p*<0.05), \*\*(*p*<0.01), \*\*\*(*p*<0.001), \*\*\*\*(*p*<0.0001). All scale bars: 50μm.



**Figure 5**

### Postmitotic neuron *Foxg1* LOF leads to premature gliogenesis and upregulation of the MAPK pathway.

(A) SOX9 staining in the control P0 cortex identifies gliogenic progenitors at the ventricular zone (white asterisk) and scattered cells throughout the cortical plate. (B–D) *NexCre*-driven loss of *Foxg1* is specific to postmitotic neurons, as seen by GFP reporter expression (C, white bars) and causes a non-autonomous upregulation of nuclear SOX9 in the ventricular zone progenitors and an increase in the numbers of SOX9+ cells cortical plate (B; quantifications: D, E). B and C are images of the same section showing SOX9 alone (B) and together with the GFP reporter (C). (F) Transcriptomic analysis of cortical plate tissue from control and *NexCre/+; Foxg1<sup>lox/lox</sup>; R26<sup>FRT-GFP</sup>* reveals a significant upregulation of *Fgf18* upon loss of *Foxg1*. (G) *Fgf18* expression in CUX2+ upper layer cells peaks at P7, as seen in the RNA seq dataset from (52). (H) Examination at E18.5 reveals increased levels of phosphorylated p42/44-ERK1/2 (pERK1/2) within the ventricular zone of *NexCre/+; Foxg1<sup>lox/lox</sup>* brains, indicative of enhanced FGF signalling. This upregulation of pERK1/2 is reversed upon treatment with the FGF Inhibitor NVP-BGJ398 (H; Quantifications: J). (I) In sections from the same brains, levels of SOX9 within the ventricular zone (VZ) are increased upon postmitotic loss of *Foxg1*, and this is restored to baseline levels upon administration of the inhibitor. (I; quantifications: K). Quantifications of pERK1/2 in each condition: 233 (Control); 248 (*NexCre/+; Foxg1<sup>lox/lox</sup>*); 207 (Control+NVP-BGJ398); 223 (*NexCre/+; Foxg1<sup>lox/lox</sup>*+NVP-BGJ398) cells from N=2 brains (biological replicates from 2 independent experiments). Quantifications of SOX9 levels in each condition: 233 (Control); 234 (*NexCre/+; Foxg1<sup>lox/lox</sup>*); 228 (Control+NVP-BGJ398); 205 (*NexCre/+; Foxg1<sup>lox/lox</sup>*+NVP-BGJ398) cells from N=3 brains (biological replicates from 2 independent experiments). Statistical test: Mann-Whitney Test (D, E); Two-way ANOVA with Tukey's correction (J, K). \*( $p < 0.05$ ), \*\*( $p < 0.01$ ), \*\*\*( $p < 0.001$ ), \*\*\*\*( $p < 0.0001$ ). All scale bars: 50  $\mu$ m.



abrogating endogenous FGF signalling. Moreover, the increase in nuclear pERK1/2 levels in VZ progenitors in *NexCre/+; Foxg1<sup>lox/lox</sup>* was no longer seen in the presence of the inhibitor (**Figure 5** [H, J](#)). We also examined nuclear SOX9 levels in each condition and found that these paralleled the findings from the FGF inhibitor experiments (**Figure 5** [I, K](#)).

Together, these data indicate that FOXG1 regulates signals from postmitotic neurons that non-autonomously trigger a transition to gliogenesis via the FGF-ERK pathway in the progenitors residing in the ventricular zone.

## Attenuation of FGF signalling does not restore neurogenesis in *Foxg1* mutant progenitors but causes premature oligogenesis


Loss of *Foxg1* appears to promote astrogliogenesis via two distinct functions: enhancing FGFR3 expression in progenitors and increasing expression of *Fgf18* in postmitotic neurons. We, therefore, tested whether a reduction of available FGF ligands may be sufficient to restore neurogenesis in *Foxg1* LOF progenitors. We used a construct encoding soluble FGFR3c (*sFGFR3c*), which has been effectively used by other studies to sequester and limit the availability of FGFs using *in-utero* electroporation ([19](#), [57](#)).

As before (**Figure 1**), embryos were electroporated at E15.5, and the brains were harvested at P14 and examined for glial markers either common to glial precursor cells (OLIG2) or exclusive to either OPCs (PDGFRA) or astrocytes (ALDH1L1). For two of the conditions, control (Cre electroporation in *Foxg1<sup>lox/+</sup>*) and *Foxg1* LOF alone (Cre electroporation in *Foxg1<sup>lox/lox</sup>*), brains from the same set of experiments presented in **Figure 1** were used from which additional sections were examined for OLIG2 and PDGFRA (**Figure 6A and B**). In controls, the majority of GFP<sup>+</sup> cells were neurons that did not express any glial marker (**Figure 1**, 6A, 6D). Neurogenesis was significantly reduced upon loss of *Foxg1* alone, and there was a corresponding increase in both types of glia (**Figure 6B**, D). For the new experimental condition, which involves a combined loss of *Foxg1* and abrogation of FGF signalling, we co-electroporated Cre with *sFgfr3c* into *Foxg1<sup>lox/lox</sup>* embryos. Strikingly, this condition did not restore neurogenesis but instead shifted the cell fate towards OLIG2<sup>+</sup>, PDGFRA<sup>+</sup>, and ALDH1L1<sup>−</sup> OPCs (**Figure 6C**, D). Conversely, neurogenesis was enhanced at the expense of gliogenesis when Cre and *sFgfr3c* were electroporated in *Foxg1<sup>lox/+</sup>; R26<sup>FRT-GFP</sup>* (Control) embryos (Supplementary Figure S7), consistent with previous reports of *sFgfr3c* overexpression in wild-type brains ([19](#)). To verify the functional consequences of FGF signalling, we quantified nuclear phospho-ERK1/2 levels in the different conditions one day after electroporation, when the process of cell fate decision would be underway in progenitors. Loss of *Foxg1* at E15.5 led to an increase in nuclear localisation of phospho-ERK1/2 by E16.5, and this effect was abrogated by co-electroporation of *sFgfr3c* (Supplementary Figure S8).

These results indicate that neurogenesis can neither proceed normally nor be restored by abrogating FGF signalling in the absence of FOXG1. Instead, the progenitors display a premature progression to an OPC fate, which appears to be a permissive cell fate in the context of the combined loss/decrease of FOXG1 and FGF signalling, respectively.

We tested the effects of prolonged *Foxg1* expression on progenitor gliogenic potential. We performed *Foxg1* overexpression using the *Rosa26<sup>FRT-GFP</sup>* background to examine the lineage arising from E15.5 progenitors scored at P14. We co-electroporated a *Foxg1* full-length construct together with Cre in control *Foxg1<sup>lox/+</sup>; R26<sup>FRT-GFP</sup>* embryos at E15.5. This overexpression led to an increase in the neurons produced by E15.5 progenitors to 98% compared with 67% in Controls (Supplementary Figure S9). This suggests that a prolonged *Foxg1* expression extends the ability of progenitors to produce neurons at the expense of glia. This is consistent with the established role of FOXG1 as a neurogenic factor ([35](#)).






**Figure 6**

### *Foxg1-Fgf* double LOF leads to premature oligogenesis.

(A–C) Cre electroporation at E15.5 in control (A, *Foxg1*<sup>lox/+</sup>; *R26*<sup>FRT-GFP</sup>) and *Foxg1* LOF (B, *Foxg1*<sup>lox/lox</sup>; *R26*<sup>FRT-GFP</sup>) embryos, followed by analysis at P14. GFP+ cells in control brains do not colocalise with ALDH1L1, OLIG2, and PDGFRA staining (A), whereas most GFP+ cells in *Foxg1* LOF brains display these markers (B). Co-electroporation of *Cre* together with a construct encoding soluble FGFR3c (an FGF-chelator) in *Foxg1*<sup>lox/lox</sup>; *R26*<sup>FRT-GFP</sup> causes a significant increase in the co-localisation of GFP+ cells with Oligodendrocyte Precursor Cells (OPCs) markers such as OLIG2 and PDGFRA (C). In each row (A–C), the boxes in the leftmost low magnification panels indicate approximate regions from the same section or serial sections shown in the adjacent high magnification panels. A quantitative analysis reveals a drastic reduction of neurogenesis at the expense of gliogenesis (astrocytes+OPCs) upon loss of *Foxg1* and an additional increase in the percentage of OPCs with the additional abrogation of FGF signalling (D, D'). n = 4069 (Control), 3970 (*Foxg1* LOF), 3332 (*Foxg1* LOF + sFGFR3c) from N=3 brains (biologically independent replicates). Statistical test: Two-way ANOVA with Tukey's correction \* (p<0.05), \*\* (p<0.01), \*\*\* (p<0.001), \*\*\*\* (p<0.0001). All scale bars: 50µm.



**Figure 7**

**Schematic depicting the regulation of gliogenesis by FOXG1.**

In neurogenic progenitors, FGFR3 levels are suppressed by FOXG1. As time progresses, FOXG1 levels decrease within progenitors, and FGFR3 levels increase, making progenitors more sensitive to FGF signalling. Concomitantly, postmitotic neurons secrete factors, including FGF18, which is also under FOXG1 regulation. FGF signalling drives progenitors towards astrogliogenesis. Later, when both FOXG1 and FGF levels are low, the progenitors transition to oligogenesis.

In summary, the results uncover a fundamental role of FOXG1 in regulating the transition of cortical progenitors from neurogenesis to gliogenesis and also highlight that neurogenesis is not possible in the absence of this critical factor.

## Discussion

Cell type diversity arises due to genetic mechanisms that regulate how progenitors give rise to cells with unique identities. In the nervous system, neurons, astrocytes, and oligodendrocytes have distinct functions, and the production of the correct numbers of each of these three cell types must be carefully controlled. How these cells arise from common progenitors is one of the outstanding questions in developmental neuroscience. Progenitors regulate the cell fate of their progeny via cell-autonomous, i.e. intrinsic programs and cell non-autonomous, i.e. extrinsic signalling mechanisms. Understanding how these regulatory processes are orchestrated and how the progenitor integrates these effects is crucial in elucidating how these transitions in cell fate arise. Our study reveals two novel functions of transcription factor FOXG1 in controlling both the expression of FGF family ligands in cortical neurons, as well as the sensitivity of progenitors to these FGF ligands via the expression of the receptor FGFR3. Thus, FOXG1 integrates the production and receptivity to extrinsic signals so progenitors may unfold cell-intrinsic programs to produce glia.

Whereas the different stages of neurogenesis have been well studied (58 [↗](#), 59 [↗](#)), the progression from neurogenesis to gliogenesis and through the various stages of gliogenesis has been less well understood. *Foxg1* was previously identified as a regulator of sequential neurogenesis in the neocortex via controlling the timing and order of generation of specific neuronal subtypes (26 [↗](#), 29 [↗](#), 30 [↗](#), 46 [↗](#)), but its endogenous role in regulating gliogenesis was not examined. The suggestion that FOXG1 may regulate the neuron-glia cell fate switch came from *Foxg1* overexpression in E12 cortical progenitors *in vitro*, which caused decreased gliogenesis (35 [↗](#), 36 [↗](#)). However, in these studies, *Foxg1* knockdown did not result in gliogenesis. Therefore, the endogenous role of FOXG1 could not be established (35 [↗](#), 36 [↗](#)).

We discovered a novel link between FOXG1 and FGF signalling. Cortical progenitors express *Fgfr1*, *r2*, and *r3* from early stages (19 [↗](#), 60 [↗](#)), and the expression levels increase during gliogenesis (19 [↗](#)). Our work demonstrates that the timing of the cessation of neurogenesis and the subsequent emergence of glial lineages is controlled by FOXG1 in cortical progenitors *in vivo*. Furthermore, we identify a key target, *Fgfr3*, via which FOXG1 controls the response of the progenitor to extrinsic FGF signalling. Importantly, we find that progenitors at E14.5 as well as E15.5 produce glia upon loss of *Foxg1*. This indicates that *Foxg1* mutant progenitors do not simply accelerate to the immediate next stage of cell type production, i.e. layer II/III instead of layer 4, but undergo a switch to producing a different class, astroglia. A decrease in FOXG1 in apical progenitors (Supplementary Figure S1 and (35 [↗](#))) may be an endogenous mechanism for the transition of neurogenesis to gliogenesis in progenitors. Intriguingly, E13.5 progenitors do not progress to premature gliogenesis upon loss of *Foxg1*, consistent with the idea that progenitors transition through distinct competent states (61 [↗](#)), an avenue that motivates future directions of inquiry.

Postmitotic neurons are known to provide cues to progenitors to initiate the production of the next wave of cells (50 [↗](#), 51 [↗](#), 55 [↗](#)). In this context, we identify an additional function of FOXG1 in regulating the expression of FGF ligands from postmitotic neurons which results in a net increase in FGF signalling experienced by progenitors. This provides an elegant mechanism for timing the onset of astrogliogenesis after neurogenesis is complete. *In vivo*, *Foxg1* expression is dynamically downregulated in pyramidal neuronal precursors as they migrate in the intermediate zone (37 [↗](#)),

which provides a potential endogenous mechanism for the modulation of FGFs secreted by postmitotic neurons. Additional regulatory mechanisms within maturing postmitotic neurons that remain unidentified may also participate in such feedback signalling to progenitors.

FGF signalling has essential roles in the early patterning of the cortical primordium (57, 62, 63) and the guidance of major axon tracts in the brain (64). However, its importance during late embryonic and postnatal stages has yet to be fully explored. Recent evidence suggests that FGF signalling regulates neurogenesis and neuronal migration (19, 65, 66). Depletion of FGF from E15.5 progenitors leads to prolonged neurogenesis in the cortex (19). However, reducing FGF signalling in *Foxg1* mutant progenitors does not restore neurogenesis but causes a premature production of OPCs. These data indicate that FOXG1 is essential for neuronal production, and in its absence, the removal of FGF is insufficient to prolong the neurogenic programme in progenitors. Overexpression of *Foxg1*, however, does restore neurogenesis despite augmented FGF signalling, and overexpression of *Foxg1* dramatically enhances the neurogenic potential of the progenitor at the expense of gliogenesis. This establishes that FOXG1 is necessary and sufficient for neuronal fate regardless of FGF signalling levels.

A rich body of work has explored how FGF, SHH, BMP, and WNT signalling regulate the proliferation and differentiation of OPCs *in vitro* and in the spinal cord (67–70). However, there is a lack of information on how OPCs arise from gliogenic progenitors and whether there are differences in gliogenic mechanisms between the spinal cord and cerebral cortex, given that the initial molecular context of progenitors in these two structures is vastly different (71, 72).

Our data indicates that once a neocortical progenitor can no longer generate neurons due to loss of *Foxg1*, it appears to be forced into a gliogenic mode, producing astrocytes if FGF signalling is present and OPCs if FGF signalling is abrogated. Taking the results from Dinh Duong *et al.*, (19) and our study together, the emerging interpretation is that for FOXG1 mutant progenitors, FGF signalling seems to be critical for astrocyte production, but it does not appear to be required for oligogenesis. The enhanced oligogenesis we report may be the result of progenitors being unable to either return to neurogenesis due to the absence of *Foxg1* or to maintain astroglialogenesis due to a reduction of FGF signalling. *In vivo*, a decline in FGFs available to progenitors may initiate OPC production. This is consistent with a general mechanism in which the regulated secretion of the ligands from previously born cells governs the timing of generation and proliferation of later-born cells (50). In the particular context of astroglialogenesis and oligogenesis, a more in-depth analysis of the affinity of FGF-FGFR binding in progenitors and specified glial cells is required to understand the fine-tuning of these processes (73, 74).

FOXG1 modulates the cell-intrinsic response of progenitors to cell-extrinsic FGF signalling, thereby regulating the production of neurons, astroglia, and oligodendrocytes, the three major components of brain circuitry. In parallel, FOXG1 also regulates FGF ligand expression in postmitotic neurons, which triggers gliogenesis in progenitors. In humans, either gain- or loss-of-function *Foxg1* mutations cause “FOXG1 syndrome”, an Autism Spectrum Disorder (29, 33, 75) that results in a range of conditions that include microcephaly, seizures and may also increase the propensity for glioblastoma or myelination deficits (29). Our findings offer mechanistic insights into potentially novel deficits that may underlie aspects of *Foxg1* syndrome-associated dysfunction.

## Acknowledgements

We thank the animal house staff of the Tata Institute of Fundamental Research (TIFR) for their excellent support; Gord Fishell and Goichi Miyoshi for the *Foxg1* floxed mouse line; Hiroshi Kawasaki for the plasmids pCAG-FGF8 and pCAG-sFGFR3c. We thank Prof. S.K. Lee for the *Foxg1*<sup>lox/lox</sup> genotyping primers and protocol. We thank Dr. Deepak Modi and Dr. Vainav Patel for



allowing us to use the NIRRH FACS Facility and the staff of the NIRRH and TIFR FACS facilities for their assistance. We thank Denis Jabaudon for his critical comments on the manuscript and members of the Jabaudon lab for helpful discussions. This work was funded by the Department of Atomic Energy (DAE), Govt. of India (Project Identification no. RTI4003, DAE OM no. 1303/2/2019/R&D-II/DAE/2079).

## Materials and Methods

### Mice

All procedures followed the Tata Institute of Fundamental Research Animal Ethics Committee (TIFR-IAEC) guidelines. The *Foxg1*<sup>lox/lox</sup>; *R26*<sup>FRT-GFP</sup> mouse line used in this study is described in (37 [↗](#)). MADM-12 GT/GT and MADM-12 TG/TG lines are described in (42 [↗](#)). *hGFAPCreERT2* (Strain #:012849) and *Ai9* reporter mouse line (Strain #:007909) were obtained from Jackson Laboratory. The *NexCre*<sup>+</sup> mouse line was obtained from Klaus Nave, Max Planck Institute for Experimental Medicine (77 [↗](#)).

All animals were kept at an ambient temperature and humidity, with a 12h light-dark cycle and food available ad libitum. Noon of the day of the vaginal plug was designated as embryonic day 0.5 (E0.5). Both male and female animals were used for all experiments.

Primers for genotyping (expected band sizes):

*Foxg1*cKO F: CCACTCCGAACCCGCTGG,

*Foxg1*cKO R: AGGCTGTTGATGCTGAACGA, (mutant: 190 bp, WT: 156 bp);

FRT Reporter: RCE-Rosa1: CCCAAAGTCGCTCTGAGTTGTTATC,

RCE-Rosa2: GAAGGAGCGGGAGAAATGGATATG,

RCE-Cag3: CCAGGCGGGCCATTTACCGTAAG, (WT: 550bp, FRT: 350bp).

MADM 12 Cassette:

Chr 12 WT F: CACTAAGCTCCACTCGCACC,

Chr 12 WT R: TCCCTCATGATGTATCCCCT,

MADM R: TCA ATG GGC GGG GGT CGT T, (WT: 322bp, MADM Cassette: 200bp).

Cre F: ATTTGCCTGCATTACCGGTC.

Cre R: ATCAACGTTTTCTTTTCGG (Cre: 350bp)

NexCre Forward: GAGTCCTGGAATCAGTCTTTTC

NexCre Reverse: AGAATGTGGAGTAGGGTGAC

NexCre Mutant Reverse: CCGCATAACCAGTGAAACAG (WT: 770 bp, NexCre: 525 bp)

## ***In utero* electroporation**

*In utero* electroporation was performed as previously described (78). Embryos were injected with 1–2  $\mu$ L of plasmid DNA solution dissolved in nuclease-free water with 0.1% fast green with plasmid DNA into the lateral ventricle through the uterine wall using a fine glass microcapillary (Sutter capillaries #B100-75-10). Constructs pCAGG-IRES-eGFP and pCAGG-IRES-FOXP1-EGFP were gifts from Gord Fishell, Harvard Medical School. pCAGG-FGF8 and pCAGG-sFGFR3c were gifts from Hiroshi Kawasaki, Kanazawa University.

## **Tissue Preparation**

Embryos were isolated in ice-cold PBS. Embryonic brains were dissected and fixed overnight in 4% (wt/vol) paraformaldehyde at 4°C overnight and then cryoprotected by transferring to 30% (wt/vol) sucrose-PBS until sectioning. Postnatal mice were anaesthetised using Thiopentone and transcardially perfused with 4% (wt/vol) paraformaldehyde in phosphate buffer, followed by overnight fixation and then cryoprotected by transferring to 30% sucrose-PBS until sectioning. The brains were sectioned at 30  $\mu$ m and 40  $\mu$ m for MADM using a freezing microtome (Leica SM2000R).

## **FGF Inhibitor administration and analysis**

We used the FGFR inhibitor NVP-BGJ398, a selective inhibitor of FGF receptors 1, 2, and 3 for blocking FGF signalling (56). Pregnant dams were administered NVP-BGJ398 (10 mg/kg body weight in DMSO and corn oil; MedChemExpress, Cat. No.: HY-13311) or vehicle solution (corn oil) by intraperitoneal injection twice a day from E14.5 to E17.5, eight treatments in total.

## **Immunohistochemistry**

Brains were sectioned (30  $\mu$ m), mounted on Superfrost plus glass microscope slides (Cat #71869-10), and dried for 2 h at room temperature. Three washes were given for 5 minutes each (3\*5') in phosphate buffer. All antibodies except FGFR3 and PDGFR $\alpha$  required antigen retrieval at 90°C–95°C in 10 mM sodium citrate buffer (pH 6.0) for 10 min. Sections were immersed in blocking solution (5% (vol/vol) horse serum in phosphate buffer with 0.1% (vol/vol) Triton X-100 (Sigma; X100)) for 1 h at RT. Incubation with primary antibody was performed in phosphate buffer containing 0.1% (vol/vol) Triton X-100 and 2.5% (vol/vol) horse serum at 4°C overnight. For postnatal brain sections, free-floating immunohistochemistry was performed. Sections were given three washes for 5 minutes each (3\*5') in phosphate buffer and then permeabilised with phosphate buffer containing 0.3% (vol/vol) Triton X-100 for 10 minutes. Blocking was done with 5% (vol/vol) horse serum in phosphate buffer with 0.3% (vol/vol) Triton X-100 for one h at RT. This was followed by primary antibody treatment in phosphate buffer containing 0.3% (vol/vol) Triton X-100 and 2.5% (vol/vol) horse serum overnight at 4°C. The sections were washed in phosphate buffer, followed by the appropriate secondary antibody for two hours at room temperature.

This was followed by three washes for 5 minutes each (3\*5') in phosphate buffer and DAPI staining for 10 minutes, after which the sections were washed with phosphate buffer for 15 minutes (3\*5'). The slides were then mounted with Fluoroshield (Sigma cat no F6057 or F6182). Information on antibodies and their catalogue numbers are mentioned in SI Appendix Supplementary Tables 1 and 2.

## **Image acquisition and analysis**

Fluorescence images were taken using an Olympus FluoView 3000 confocal microscope with FluoView software. All the image analysis was done on Fiji-ImageJ. A nonlinear operation such as gamma correction was not performed in the figures. Brightness and contrast adjustments were performed identically for control and mutant conditions. Cell counting was performed using the Cell Counter plugin in Fiji.

Intensity quantifications for Phospho-p42/44-ERK1/2 and SOX9 were done by imaging Control and mutant samples at the same laser, gain and offset settings on the Olympus FV3000 confocal microscope. Images were stacked based on one-cell thickness stacks, *i.e.* approximately 5µm. Intensity quantification was done by drawing Regions of Interest (ROIs) around the nuclei using DAPI or SOX2, followed by intensity quantification of Phospho-p42/44-ERK1/2 or SOX9 and calculating mean intensity per unit ROI.

## Histone and FOXG1 ChIP Seq analysis

FASTQ files deposited by previously published manuscripts ([47](#)) and ([46](#)) were obtained using SRA Toolkit's fastq-dump command and aligned to mouse reference genome mm10 using Bowtie2 ([79](#)). Peak calling and annotation were performed using Homer using default options and command style “-histone” for the H3K27Me3 and H3K4Me3 datasets. For the FOXG1 ChIP-seq analysis, peaks were called using all “-style” options and only the top 50% peaks called were considered for further analysis for all the ChIP-seq datasets. Peaks were intersected using BEDTools software ([80](#)) and annotated using Homer ([81](#)) `annotatePeaks.pl` function, and genome view plots were created using IGV ([82](#)).

## Fluorescence-Assisted Cell Sorting

*Foxg1* mutant cells were obtained by administering tamoxifen (Cat #T5648) prepared in corn oil (Sigma; Cat #8267) to E15.5 *hGFAPCreERT2; Foxg1<sup>lox/lox</sup>; Ai9* dams, at 40 µg/g body weight dose and FACS-isolated using the Ai9 reporter at E17.5. We used FlashTag-labelled (Thermo Fisher Scientific SKU C34554) progenitors at E15.5 for controls and collected the cells at E17.5. Cortical tissue was dissected in HBSS without Calcium/Magnesium (Thermo Fisher Scientific Catalogue number: 14170112), and a single-cell suspension was prepared in HBSS with Calcium/Magnesium (Thermo Fisher Scientific Catalogue number: 14025092) using 0.25% Trysin (ThermoFischer Scientific Gibco Catalogue number: 15400054) and 70µm cell strainer. FACS was performed using BD Aria Fusion (BD Biosciences) with the 568 and 488 lasers using an 85µm nozzle. Singlets were selected using forward scatter and side scatter. Cells were selected for collection based on their RFP signal (Mutant) or GFP signal (Control).

## RNA seq - sample preparation and analysis

Cells obtained from FACS were stored in RNeasy lysis buffer until extraction. 400,000 cells were pooled from at least two brains to obtain a biological replicate. RNA extraction and sequencing were performed on three control replicates and four *Foxg1* mutant replicate cell suspensions. 1µg of RNA was used to obtain the cDNA library, and sequencing was performed on an Illumina platform to achieve 150 bp reads to generate 30 Million paired-end reads per sample. Fastq QC was performed as described in (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and reads > 30 Phred scores were aligned using HISAT2 ([83](#)). Feature counts were used to quantify the number of reads per transcript. Differential expression analysis was performed using DESeq2 ([84](#)). A log2FoldChange cutoff of 0.58 and P-value < 0.05 was used to identify DEGs. GO analysis was performed using Shiny GO ([85](#)). Genes described as Astrocyte-enriched were obtained from ([48](#)). Data availability: All RNA Seq samples generated for this manuscript are deposited under [GSE253919](#).

## References

1. Qian X., et al. (2000) **Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells** *Neuron* **28**:69–80
2. Hirabayashi Y., et al. (2009) **Polycomb limits the neurogenic competence of neural precursor cells to promote astrogenic fate transition** *Neuron* **63**:600–613
3. Reynolds B. A., Weiss S. (1992) **Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system** *Science* **255**:1707–1710
4. Temple S. (1989) **Division and differentiation of isolated CNS blast cells in microculture** *Nature* **340**:471–473
5. Shen Z., et al. (2021) **Distinct progenitor behavior underlying neocortical gliogenesis related to tumorigenesis** *Cell Rep* **34**
6. Gao P., et al. (2014) **Deterministic progenitor behavior and unitary production of neurons in the neocortex** *Cell* **159**:775–788
7. Naka H., Nakamura S., Shimazaki T., Okano H. (2008) **Requirement for COUP-TFI and II in the temporal specification of neural stem cells in CNS development** *Nat. Neurosci* **11**:1014–1023
8. Namihira M., et al. (2009) **Committed neuronal precursors confer astrocytic potential on residual neural precursor cells** *Dev. Cell* **16**:245–255
9. Kang P., et al. (2012) **Sox9 and NFIA coordinate a transcriptional regulatory cascade during the initiation of gliogenesis** *Neuron* **74**:79–94
10. Nagao M., Ogata T., Sawada Y., Gotoh Y. (2016) **Zbtb20 promotes astrocytogenesis during neocortical development** *Nat. Commun* **7**
11. Tanigaki K., et al. (2001) **Notch1 and Notch3 instructively restrict bFGF-responsive multipotent neural progenitor cells to an astroglial fate** *Neuron* **29**:45–55
12. Cao Q., et al. (2008) **Expression of Notch-1 receptor and its ligands Jagged-1 and Delta-1 in amoeboid microglia in postnatal rat brain and murine BV-2 cells** *Glia* **56**:1224–1237
13. Ge W., et al. (2002) **Notch signaling promotes astrogliogenesis via direct CSL-mediated glial gene activation** *J. Neurosci. Res* **69**:848–860
14. Cao Q., et al. (2010) **Nuclear factor- $\kappa$ B/p65 responds to changes in the Notch signaling pathway in murine BV-2 cells and in amoeboid microglia in postnatal rats treated with the  $\gamma$ -secretase complex blocker DAPT** *J. Neurosci. Res* **88**:2701–2714
15. Rowitch D. H., Kriegstein A. R. (2010) **Developmental genetics of vertebrate glial-cell specification** *Nature* **468**:214–222
16. Mallamaci A. (2013) **Developmental control of cortico-cerebral astrogenesis** *Int. J. Dev. Biol* **57**:689–706



17. Sloan S. A., Barres B. A. (2014) **Mechanisms of astrocyte development and their contributions to neurodevelopmental disorders** *Curr. Opin. Neurobiol* **27**:75–81
18. Takouda J., Katada S., Nakashima K. (2017) **Emerging mechanisms underlying astrogenesis in the developing mammalian brain** *Proc Jpn Acad, Ser B, Phys Biol Sci* **93**:386–398
19. Dinh Duong T. A., et al. (2019) **FGF Signaling Directs the Cell Fate Switch from Neurons to Astrocytes in the Developing Mouse Cerebral Cortex** *J. Neurosci* **39**:6081–6094
20. Zhang X., et al. (2020) **Clonal analysis of gliogenesis in the cerebral cortex reveals stochastic expansion of glia and cell autonomous responses to egfr dosage** *Cells* **9**
21. Nakashima K., Yanagisawa M., Arakawa H., Taga T. (1999) **Astrocyte differentiation mediated by LIF in cooperation with BMP2** *FEBS Lett* **457**:43–46
22. He F., et al. (2005) **A positive autoregulatory loop of Jak-STAT signaling controls the onset of astrogliogenesis** *Nat. Neurosci* **8**:616–625
23. Stolt C. C., et al. (2002) **Terminal differentiation of myelin-forming oligodendrocytes depends on the transcription factor Sox10** *Genes Dev* **16**:165–170
24. Bhattarai C., Poudel P. P., Ghosh A., Kalthur S. G. (2022) **Comparative role of SOX10 gene in the gliogenesis of central, peripheral, and enteric nervous systems** *Differentiation* **128**:13–25
25. Hanashima C., Shen L., Li S. C., Lai E. (2002) **Brain factor-1 controls the proliferation and differentiation of neocortical progenitor cells through independent mechanisms** *J. Neurosci* **22**:6526–6536
26. Toma K., Kumamoto T., Hanashima C. (2014) **The timing of upper-layer neurogenesis is conferred by sequential derepression and negative feedback from deep-layer neurons** *J. Neurosci* **34**:13259–13276
27. Hanashima C., Suzanne L., Shen Lijian Lai, Fishell G. (2004) **Foxg1 Suppresses Early Cortical Cell Fate** *Science* **303**:56–60
28. Kumamoto T., Hanashima C. (2017) **Evolutionary conservation and conversion of Foxg1 function in brain development** *Dev. Growth Differ* **59**:258–269
29. Hou P.-S., hAilín D. Ó., Vogel T., Hanashima C. (2020) **Transcription and beyond: delineating FOXG1 function in cortical development and disorders** *Front. Cell. Neurosci* **14**
30. Liu J., et al. (2022) **FOXG1 sequentially orchestrates subtype specification of postmitotic cortical projection neurons** *Sci. Adv* **8**
31. Tao W., Lai E. (1992) **Telencephalon-restricted expression of BF-1, a new member of the HNF-3/fork head gene family, in the developing rat brain** *Neuron* **8**:957–966
32. Shoichet S. A., et al. (2005) **Haploinsufficiency of novel FOXG1B variants in a patient with severe mental retardation, brain malformations and microcephaly** *Hum. Genet* **117**:536–544
33. Hettige N. C., Ernst C. (2019) **FOXG1 dose in brain development** *Front. Pediatr* **7**

34. Brancaccio M., Pivetta C., Granzotto M., Filippis C., Mallamaci A. (2010) **Emx2 and Foxg1 inhibit gliogenesis and promote neuronogenesis** *Stem Cells* **28**:1206–1218
35. Falcone C., et al. (2019) **Foxg1 antagonizes neocortical stem cell progression to astrogenesis** *Cereb. Cortex* **29**:4903–4918
36. Frisari S., et al. (2022) **Multidimensional functional profiling of human neuropathogenic FOXG1 alleles in primary cultures of murine pallial precursors** *Int. J. Mol. Sci* **23**
37. Miyoshi G., Fishell G. (2012) **Dynamic FoxG1 expression coordinates the integration of multipolar pyramidal neuron precursors into the cortical plate** *Neuron* **74**:1045–1058
38. Shen Q., et al. (2006) **The timing of cortical neurogenesis is encoded within lineages of individual progenitor cells** *Nat. Neurosci* **9**:743–751
39. Kumamoto T., Hanashima C. (2014) **Neuronal subtype specification in establishing mammalian neocortical circuits** *Neurosci. Res* **86**:37–49
40. Kumamoto T., et al. (2013) **Foxg1 coordinates the switch from nonradially to radially migrating glutamatergic subtypes in the neocortex through spatiotemporal repression** *Cell Rep* **3**:931–945
41. Zong H., Espinosa J. S., Su H. H., Muzumdar M. D., Luo L. (2005) **Mosaic analysis with double markers in mice** *Cell* **121**:479–492
42. Contreras X., et al. (2021) **A genome-wide library of MADM mice for single-cell genetic mosaic analysis** *Cell Rep* **35**
43. Hippenmeyer S., Johnson R. L., Luo L. (2013) **Mosaic analysis with double markers reveals cell-type-specific paternal growth dominance** *Cell Rep* **3**:960–967
44. Zhuo L., et al. (2001) **hGFAP-cre transgenic mice for manipulation of glial and neuronal function in vivo** *Genesis* **31**:85–94
45. Ganat Y. M., et al. (2006) **Early postnatal astroglial cells produce multilineage precursors and neural stem cells in vivo** *J. Neurosci* **26**:8609–8621
46. Cargnin F., et al. (2018) **FOXG1 Orchestrates Neocortical Organization and Cortico-Cortical Connections** *Neuron* **100**:1083–1096
47. Liu J., Wu X., Zhang H., Pfeifer G. P., Lu Q. (2017) **Dynamics of RNA Polymerase II Pausing and Bivalent Histone H3 Methylation during Neuronal Differentiation in Brain Development** *Cell Rep* **20**:1307–1318
48. Telley L., et al. (2019) **Temporal patterning of apical progenitors and their daughter neurons in the developing neocortex** *Science* **364**
49. Lavoie H., Gagnon J., Therrien M. (2020) **ERK signalling: a master regulator of cell behaviour, life and fate** *Nat. Rev. Mol. Cell Biol* **21**:607–632
50. Seuntjens E., et al. (2009) **Sip1 regulates sequential fate decisions by feedback signaling from postmitotic neurons to progenitors** *Nat. Neurosci* **12**:1373–1380

51. Barnabé-Heider F., et al. (2005) **Evidence that embryonic neurons regulate the onset of cortical gliogenesis via cardiotrophin-1** *Neuron* **48**:253–265
52. Yuan W., et al. (2022) **Temporally divergent regulatory mechanisms govern neuronal diversification and maturation in the mouse and marmoset neocortex** *Nat. Neurosci* **25**:1049–1058
53. Thompson C. L., et al. (2014) **A high-resolution spatiotemporal atlas of gene expression of the developing mouse brain** *Neuron* **83**:309–323
54. Dastidar S. G., Landrieu P. M. Z., D’Mello S. R. (2011) **FoxG1 promotes the survival of postmitotic neurons** *J. Neurosci* **31**:402–413
55. Hoshikawa M., Yonamine A., Konishi M., Itoh N. (2002) **FGF-18 is a neuron-derived glial cell growth factor expressed in the rat brain during early postnatal development** *Brain Res. Mol. Brain Res* **105**:60–66
56. Guagnano V., et al. (2011) **Discovery of 3-(2,6-dichloro-3,5-dimethoxy-phenyl)-1-{6-[4-(4-ethyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yl}-1-methyl-urea (NVP-BGJ398), a potent and selective inhibitor of the fibroblast growth factor receptor family of receptor tyrosine kinase** *J. Med. Chem* **54**:7066–7083
57. Fukuchi-Shimogori T., Grove E. A. (2001) **Neocortex patterning by the secreted signaling molecule FGF8** *Science* **294**:1071–1074
58. Huilgol D., Russ J. B., Srivas S., Huang Z. J. (2023) **The progenitor basis of cortical projection neuron diversity** *Curr. Opin. Neurobiol* **81**
59. Mukhtar T., Taylor V. (2018) **Untangling cortical complexity during development** *J. Exp. Neurosci* **12**
60. Bansal R., Lakhina V., Remedios R., Tole S. (2003) **Expression of FGF receptors 1, 2, 3 in the embryonic and postnatal mouse brain compared with Pdgfralpha, Olig2 and Plp/dm20: implications for oligodendrocyte development** *Dev. Neurosci* **25**:83–95
61. Telley L., et al. (2016) **Sequential transcriptional waves direct the differentiation of newborn neurons in the mouse neocortex** *Science* **351**:1443–1446
62. Fukuchi-Shimogori T., Grove E. A. (2003) **Emx2 patterns the neocortex by regulating FGF positional signaling** *Nat. Neurosci* **6**:825–831
63. Crossley P. H., Martinez S., Ohkubo Y., Rubenstein J. L. (2001) **Coordinate expression of Fgf8, Otx2, Bmp4, and Shh in the rostral prosencephalon during development of the telencephalic and optic vesicles** *Neuroscience* **108**:183–206
64. Tole S., Gutin G., Bhatnagar L., Remedios R., Hébert J. M. (2006) **Development of midline cell types and commissural axon tracts requires Fgfr1 in the cerebrum** *Dev. Biol* **289**:141–151
65. Kon E., et al. (2019) **N-cadherin-regulated FGFR ubiquitination and degradation control mammalian neocortical projection neuron migration** *eLife* **8**
66. Szczurkowska J., et al. (2016) **Targeted in vivo genetic manipulation of the mouse or rat brain by in utero electroporation with a triple-electrode probe** *Nat. Protoc* **11**:399–412

67. McCaughey-Chapman A., Connor B. (2023) **Cell reprogramming for oligodendrocytes: A review of protocols and their applications to disease modeling and cell-based remyelination therapies** *J. Neurosci. Res* **101**:1000–1028
68. Manzari-Tavakoli A., et al. (2022) **The Cross-Talks Among Bone Morphogenetic Protein (BMP) Signaling and Other Prominent Pathways Involved in Neural Differentiation** *Front. Mol. Neurosci* **15**
69. El Waly B., Macchi M., Cayre M., Durbec P. (2014) **Oligodendrogenesis in the normal and pathological central nervous system** *Front. Neurosci* **8**
70. Farreny M. A., et al. (2017) **FGFs are orchestra conductors of Shh-dependent oligodendroglial fate specification in the ventral spinal cord** *Neural Development* :1–19
71. Zeisel A., et al. (2018) **Molecular architecture of the mouse nervous system** *Cell* **174**:999–1014
72. Silbereis J. C., Pochareddy S., Zhu Y., Li M., Sestan N. (2016) **The cellular and molecular landscapes of the developing human central nervous system** *Neuron* **89**:248–268
73. Bansal R., Kumar M., Murray K., Morrison R. S., Pfeiffer S. E. (1996) **Regulation of FGF receptors in the oligodendrocyte lineage** *Mol. Cell. Neurosci* **7**:263–275
74. Bansal R., Magge S., Winkler S. (2003) **Specific inhibitor of FGF receptor signaling: FGF-2-mediated effects on proliferation, differentiation, and MAPK activation are inhibited by PD173074 in oligodendrocyte-lineage cells** *J. Neurosci. Res* **74**:486–493
75. Florian C., Bahi-Buisson N., Bienvenu T. (2011) **FOXG1-Related Disorders: From Clinical Description to Molecular Genetics** *Mol. Syndromol* <https://doi.org/10.1159/000327329>
76. Di Bella D. J., et al. (2021) **Molecular logic of cellular diversification in the mouse cerebral cortex** *Nature* **595**:554–559
77. Goebbels S., et al. (2006) **Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice** *Genesis* **44**:611–621
78. Pal S., et al. (2021) **An Early Cortical Progenitor-Specific Mechanism Regulates Thalamocortical Innervation** *J. Neurosci* **41**:6822–6835
79. Langmead B., Salzberg S. L. (2012) **Fast gapped-read alignment with Bowtie 2** *Nat. Methods* **9**:357–359
80. Quinlan A. R., Hall I. M. (2010) **BEDTools: a flexible suite of utilities for comparing genomic features** *Bioinformatics* **26**:841–842
81. Heinz S., et al. (2010) **Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities** *Mol. Cell* **38**:576–589
82. Robinson J. T., et al. (2011) **Integrative genomics viewer** *Nat. Biotechnol* **29**:24–26
83. Kim D., Paggi J. M., Park C., Bennett C., Salzberg S. L. (2019) **Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype** *Nat. Biotechnol* **37**:907–915



84. Love M. I., Huber W., Anders S. (2014) **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2** *Genome Biol* **15**
85. Ge S. X., Jung D., Yao R. (2020) **ShinyGO: a graphical gene-set enrichment tool for animals and plants** *Bioinformatics* **36**:2628–2629

## Editors

Reviewing Editor

**Samuel Pleasure**

University of California, San Francisco, San Francisco, United States of America

Senior Editor

**K VijayRaghavan**

National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, India

### Reviewer #1 (Public review):

Summary:

In this paper, Bose et al. investigated the role of Foxg1 transcription factor in the progenitors at late stages of cerebral cortex development.

They discover that Foxg1 is a repressor of gliogenesis and has a dual function, first as a repressor of Fgfr3 receptor in progenitors, and second as a suppressor of the Fgf ligands in young neurons.

They found that the inactivation of Foxg1 in cortical progenitors causes premature astrogliogenesis at the expense of neurogenesis. They identify Fgfr3 as a novel FOXG1 target. They show that suppression of Fgfr3 by FOXG1 in progenitors is required to maintain neurogenesis. On the other hand, they also show that FOXG1 negatively regulates the expression of Fgf gliogenic secreted factors in young neurons suppressing gliogenesis cells extrinsically.

Strengths:

The authors used time-consuming in vivo experiments utilizing several mouse strains including Foxg1-MADM in combination with RNA-Seq and ChIP to convincingly show that Foxg1 acts upstream of FGF signalling in the control of gliogenesis onset. The conclusions of this paper are mostly well supported by data.

Weaknesses:

The role of Fgf signaling in gliogenesis and Foxg1 in neurogenesis is well known. It is not clear if Fgf18 is a direct target of Foxg1.

<https://doi.org/10.7554/eLife.101851.1.sa2>

### Reviewer #2 (Public review):

Summary:

We have known for some time that neural progenitors in the cerebral cortex switch their output from cortical neurons to glia at late embryonic stages, however little is known about

how this switch is regulated at the molecular level. Bose et al present a convincing set of findings, demonstrating that the transcription factor Foxg1 plays a key role in this process, mediated through FGF signalling. Foxg1 cell-autonomously inhibits gliogenesis in progenitor cells (thereby promoting neuronal identity), and lower Foxg1 expression in postnatal neurons leads to increased expression of FGF ligand, promoting glial development from nearby progenitors.

#### Strengths:

The study is very well designed, having a systematic, thorough, and logical approach. The data is convincing. The authors make full use of a range of existing transgenic strains, published 'omics data, and elegant genetic approaches such as MADM. This combination of approaches is particularly rigorous, lending significant weight to the study. The manuscript is well-written, clear, and easy to follow.

#### Weaknesses:

It wasn't clear to me why the authors chose postnatal day 14 to examine the effects of Foxg1 deletion at E15 - this is a long time window, giving time for indirect consequences of Foxg1 deletion to influence development and thereby potentially complicating the interpretation of findings. For example, the authors show that there is no increased proliferation of astrocytes or death of neurons lacking Foxg1 shortly after cre-mediated deletion, but it remains formally possible (if perhaps unlikely) that these processes could be affected later during the time window. The rationale underlying the choice of this time point should be explained.

I don't agree with the statement in the very last sentence of the results section that "neurogenesis is not possible in the absence of [Foxg1]" as there are multiple reports in the literature demonstrating the presence of neurons in Foxg1<sup>-/-</sup> mice (eg: Xuan et al., 1995; Hanashima et al., 2002, Martynoga et al., 2005, Muzio and Mallamaci 2005). Perhaps the statement refers specifically to late-born cortical neurons. This point also arises in the discussion section.

#### Impact

This manuscript identifies a previously unknown role for Foxg1 in forebrain development and a mechanism underlying the neurogenic-to-gliogenic switch that occurs at late embryonic stages of cortex development. These findings will stimulate further research to uncover more details of how this important switch is controlled and may provide useful insight into some of the symptoms experienced by children with FOXG1 Syndrome.

<https://doi.org/10.7554/eLife.101851.1.sa1>

#### Author response:

##### **Reviewer 1:**

*The role of Fgf signaling in gliogenesis and Foxg1 in neurogenesis is well known. It is not clear if Fgf18 is a direct target of Foxg1.*

We agree with the reviewer- Fgf signaling is an established pro-gliogenic pathway (Duong et al 2019) and Foxg1 overexpression is known to promote neurogenesis in cultured neural stem cells (Branacaccio et al 2019). Our study links these two mechanisms, as the Reviewer has summarized: (a) we demonstrate that FOXG1 works via modulating Fgf signaling cell-autonomously within progenitors by regulating the levels of Fgfr3. (b) Loss of Foxg1 in postmitotic neurons results in the upregulation of Fgf ligand expression (possibly via indirect

mechanisms) and this non-cell autonomously increases Fgf signaling in progenitors. Our study is entirely performed *in vivo*.

Proposed revision: We will revise the manuscript to reflect that *Fgf18* may be an *indirect* target of FOXG1 in postmitotic neurons.

**Reviewer 2:**

*It wasn't clear to me why the authors chose postnatal day 14 to examine the effects of Foxg1 deletion at E15 - this is a long time window, giving time for indirect consequences of Foxg1 deletion to influence development and thereby potentially complicating the interpretation of findings. For example, the authors show that there is no increased proliferation of astrocytes or death of neurons lacking Foxg1 shortly after cre-mediated deletion, but it remains formally possible (if perhaps unlikely) that these processes could be affected later during the time window. The rationale underlying the choice of this time point should be explained.*

*I don't agree with the statement in the very last sentence of the results section that "neurogenesis is not possible in the absence of [Foxg1]" as there are multiple reports in the literature demonstrating the presence of neurons in Foxg1<sup>-/-</sup> mice (eg: Xuan et al., 1995; Hanashima et al., 2002, Martynoga et al., 2005, Muzio and Mallamaci 2005). Perhaps the statement refers specifically to late-born cortical neurons. This point also arises in the discussion section.*

Proposed revisions:

(a) We will revise the manuscript to explain why we chose postnatal day 14 to examine the effects of Foxg1 deletion at E15.

- We have examined the transcriptomic dysregulation after Foxg1 deletion at E17.5, which is a reasonable period to identify potential direct targets. Furthermore, FOXG1 occupies the *Fgfr3* locus in ChIP-seq performed at E15.5. Together, these support the interpretation that *Fgfr3* is a direct target of Foxg1.
- As the Reviewer notes, we have investigated the possibility of increased proliferation of astrocytes and death of neurons and found no evidence that suggests these phenomena occur in the 3 days after loss of Foxg1. Cortical neurons are postmitotic and differentiated by E18.5, the stage at which we examined CC3 staining and found no difference in cell death in control and mutants (Supplementary Figure S2C, C'). The majority of progenitors (PAX6+ve cells) that lose Foxg1 at E15.5 express the gliogenic transcription factor NFIA by E18.5 (Figure 2C, C'), but hardly any express intermediate (neurogenic) progenitor marker TBR2 (Supplementary Figure S2B, B'). It is therefore unlikely that neurons are born from Foxg1 mutant progenitors and then die at a later stage.
- The cellular consequences of loss of Foxg1 require additional time to detect e.g. it takes ~ 5 days for GFAP to be detected in astrocytes once they are born. The P14 timepoint permits the assessment of oligogenesis which begins after astroglialogenesis and therefore permits a comprehensive assessment of the lineage of E15.5 Foxg1 null progenitors.

(b) Thank you for pointing out that the last sentence of the results section implied (incorrectly) that ALL neurogenesis is not possible in the absence of Foxg1. We will modify this (and the discussion) to reflect that this applies to E14/15 progenitors and late-born cortical neurons.

<https://doi.org/10.7554/eLife.101851.1.sa0>