

Realistic mossy fiber input patterns to unipolar brush cells evoke a continuum of temporal responses comprised of components mediated by different glutamate receptors

Reviewed Preprint

v2 • January 7, 2025

Revised by authors

Reviewed Preprint

v1 • November 12, 2024

Vincent Huson, Wade G Regehr 

Department of Neurobiology, Harvard Medical School, Boston, USA

 https://en.wikipedia.org/wiki/Open_access Copyright information

eLife Assessment

This **valuable** study describes how trains of mossy fiber stimulation control cerebellar unipolar brush cell discharges. The dissection of the contributions of relevant glutamate receptors to these transformations is **convincing**. Overall, the study broadens our understanding of temporal processing in the cerebellar cortex.

<https://doi.org/10.7554/eLife.102618.2.sa3>

Abstract

Unipolar brush cells (UBCs) are excitatory interneurons in the cerebellar cortex that receive mossy fiber (MF) inputs and excite granule cells. The UBC population responds to brief burst activation of MFs with a continuum of temporal transformations, but it is not known how UBCs transform the diverse range of MF input patterns that occur *in vivo*. Here we use cell-attached recordings from UBCs in acute cerebellar slices to examine responses to MF firing patterns that are based on *in vivo* recordings. We find that MFs evoke a continuum of responses in the UBC population, mediated by three different types of glutamate receptors that each convey a specialized component. AMPARs transmit timing information for single stimuli at up to 5 spikes/s, and for very brief bursts. A combination of mGluR2/3s (inhibitory) and mGluR1s (excitatory) mediates a continuum of delayed, and broadened responses to longer bursts, and to sustained high frequency activation. Variability in the mGluR2/3 component controls the time course of the onset of firing, and variability in the mGluR1 component controls the duration of prolonged firing. We conclude that the combination of glutamate receptor types allows each UBC to simultaneously convey different aspects of MF firing. These findings establish that UBCs are highly flexible circuit elements that provide diverse temporal transformations that are well suited to contribute to specialized processing in different regions of the cerebellar cortex.

Introduction

Unipolar brush cells (UBCs) are a specialized type of excitatory interneuron in the cerebellum and cerebellar-like structures (Altman and Bayer, 1977; Floris et al., 1994; Mugnaini and Floris, 1994; Rossi et al., 1995; Meek et al., 2008). They receive a single mossy fiber (MF) input onto their large dendritic brush, and they in turn make their own mossy fiber boutons that make synapse mainly onto granule cells, but also onto other UBCs (Rossi et al., 1995; Nunzi et al., 2001; Hariani et al., 2024). UBCs transform MF firing patterns into prolonged increases or decreases in firing that lead to diverse granule cell firing patterns (Rossi et al., 1995; Kinney, Overstreet and Slater, 1997; Dino et al., 2000; Mugnaini, Sekerkova and Martina, 2011; Locatelli et al., 2013; Kennedy et al., 2014; van Dorp and De Zeeuw, 2014; Borges-Merjane and Trussell, 2015; Zampini et al., 2016; Guo et al., 2021b). In weakly electric fish, UBC firing provides a filtered signal that is suited to cancelling self-generated signals (Kennedy et al., 2014). In the rodent brain, recordings have been made from UBCs *in vivo* (Simpson et al., 2005; Barmack and Yakhnitsa, 2008; Ruigrok, Hensbroek and Simpson, 2011; Hensbroek et al., 2015; Witter and De Zeeuw, 2015), but in the absence of simultaneous recordings from connected MF-UBC pairs it is not known how UBCs transform MF firing patterns during behaviors.

Most of what is known regarding UBC temporal transformations is based on brain slice experiments in which MFs are activated with brief bursts (Rossi et al., 1995; Mugnaini, Sekerkova and Martina, 2011; Borges-Merjane and Trussell, 2015; Guo et al., 2021b). UBCs responses range from fast “ON” cells that fire for hundreds of milliseconds with virtually no delay, to intermediate UBCs that initially pause before increasing firing for seconds, to “OFF” UBCs that suppress firing for up to two seconds (Guo et al., 2021b; Huson, Newman and Regehr, 2023). Responses are mediated by a combination of different glutamate receptors. mGluR2/3s activate potassium channels to suppress firing (Russo et al., 2008; Kim et al., 2012; Borges-Merjane and Trussell, 2015), while a combination of AMPARs and mGluR1s mediate excitation (Rossi et al., 1995; Kinney, Overstreet and Slater, 1997; Kim et al., 2012; Schwartz et al., 2012; Borges-Merjane and Trussell, 2015; Guo et al., 2021b). Populations of UBCs exhibit a continuum of response durations and amplitudes, which is thought to reflect inverse expression profiles of the mGluR1 and mGluR2 signaling pathways (Guo et al., 2021b; Kozareva et al., 2021).

Little is known about how UBCs transform MF activity patterns during behavior. This issue is complicated by regional differences in MF firing properties (Eccles et al., 1971; Lisberger and Fuchs, 1978; van Kan, Gibson and Houk, 1993; Garwicz, Jorntell and Ekerot, 1998; Rancz et al., 2007; Arenz et al., 2008; Witter and De Zeeuw, 2015). Even though UBCs are present throughout the cerebellar cortex, they are most abundant in regions involved in the control of eye position and in vestibular processing (Floris et al., 1994; Dino, Willard and Mugnaini, 1999; Takacs et al., 1999; Englund et al., 2006). MFs in these UBC-rich regions often fire in brief high frequency bursts (Eccles et al., 1971; van Kan, Gibson and Houk, 1993; Garwicz, Jorntell and Ekerot, 1998; Rancz et al., 2007) that in some cases represent saccadic eye movements (Ohtsuka and Noda, 1992). MFs can also fire continuously for extended periods and change their firing rates dynamically to represent motor signals, and encode velocity, position and direction (Lisberger and Fuchs, 1978; van Kan, Gibson and Houk, 1993; Arenz et al., 2008; Barmack and Yakhnitsa, 2008; Lisberger, 2009). Brain slice experiments suggest that UBCs can perform useful computations that transform MF inputs. UBC responses to sinusoidal modulation of MF firing rates have been previously examined (Zampini et al., 2016; Balmer, Borges-Merjane and Trussell, 2021; Guo et al., 2021b). It was found that “OFF” UBCs provide a phase reversed response in which the UBC fires out of phase with the MF, while AMPAR responses provide a variety of phase transformations (Zampini et al., 2016). Such temporal transformations are thought to provide granule cell activity patterns that can be used in combination with plasticity mechanisms to modify the phase of the Purkinje cell output. However, contributions from mGluR1

were not assessed in that study. The mGluR1 component is prone to washout, and non-invasive techniques are required for reliable long-lasting recordings (Guo et al., 2021b [↗](#); Huson, Newman and Regehr, 2023 [↗](#)). As such, it is very difficult to predict how the complete UBC population will respond to the complex activity patterns observed *in vivo*, given the complex dynamics from differential activation of AMPARs, mGluR1s and mGluR2/3s.

Here we characterize UBC firing evoked by MF input patterns that are based on *in vivo* recordings of MF firing during smooth pursuit eye movements. We recorded MF-evoked increases in UBC spiking using cell-attached recordings in brain slices and pharmacologically assessed the contributions of mGluR2/3s, AMPARs, and mGluR1s. AMPARs conveyed time-locked responses to single stimuli, brief bursts, and baseline stimulation at up to 5 spikes/s (spk/s). But for bursts of 20 stimuli, excitatory responses are primarily mediated by mGluR1. An interplay of mGluR2/3s and mGluR1s dominated UBC responses to bursts of 20 stimuli or more and prolonged increases in input, creating a continuum of temporally-filtered responses. In this way a combination of glutamate receptors allows single MF to UBC connections to simultaneously convey a spike-timing component and a temporally-filtered component.

Results

Diverse MF activity evokes wide ranging UBC responses

We set out to assess the UBC response to MF input patterns as observed *in vivo* in the vestibulocerebellum. For this purpose, we utilized *in vivo* recordings of MF activity during a smooth pursuit eye movement task from the floccular complex of rhesus macaques (provided by David J. Herzfeld and Stephen G. Lisberger) and reproduced the firing patterns observed there in acute cerebellar slices (**Fig. 1a** [↗](#)). In their experiments, a head-fixed monkey was trained to track a smoothly moving visual target with minimal saccades while simultaneously recording single unit MF responses (see **Methods**). MF firing is comprised of sustained changes in firing that encode velocity and eye position, burst discharges that encode saccades, or a combination of both (Lisberger and Fuchs, 1978 [↗](#)). We selected 30 s periods of firing from 2 representative MFs recorded in separate sessions. One showed characteristic burst firing (**Fig. 1c** [↗](#), green trace) and the other showed more sustained firing that increased or decreased in accordance with the smooth pursuit trials (**Fig. 1d** [↗](#), green trace). We performed cell-attached recordings from UBCs in acute cerebellar brain slices from P30-47 mice in the presence of GABA_A, GABA_B, and glycine receptor blockers (see **Methods**) and stimulated MF inputs using a theta-glass electrode.

As a means of categorizing the UBCs (Guo et al., 2021b [↗](#); Huson, Newman and Regehr, 2023 [↗](#)), we stimulated the MF input with a standard burst of 20 stimuli at 100 spk/s (**Fig. 1b** [↗](#), green trace). These stimuli evoked diverse long-lasting changes in firing, with **Fig. 1b** [↗](#) depicting examples of a fast UBC with relatively brief increases in firing, a longer lasting mid-range UBC, a slow UBC with a long-lasting increase in firing, and a UBC that is spontaneously active and transiently stops firing (OFF). We then applied stimulation patterns from *in vivo* recordings to these same UBCs. The first representative *in vivo* MF displayed a series of bursts consisting of 2 - 23 spikes at very high frequencies (**Fig. 1c** [↗](#), green trace) in line with previous findings (Eccles et al., 1971 [↗](#); Rancz et al., 2007 [↗](#)). Activating MF inputs to the UBCs with this stimulation pattern evoked a series of responses that were qualitatively consistent with those evoked by 20 stimuli at 100 spk/s bursts. However, in the slow UBC, repeated bursts resulted in what appeared to be an increase in steady-state firing, and burst stimulation primarily induced a pause in this firing. Generally, different MF bursts evoked responses of different magnitude, which was particularly evident in the OFF cell, where even small changes in the input evoked noticeable decreases in firing.




Figure 1

UBCs perform complex temporal transformations of in-vivo MF firing patterns.

a. Scheme showing experimental paradigm. We received *in vivo* recordings of MFs in the flocculus and paraflocculus of macaques (middle) during a smooth pursuit eye movement task (left). We reproduced the *in vivo* firing patterns in MFs in acute slices of lobule X the mouse cerebellum using a theta stimulation electrode, while making cell-attached recordings of the UBC response (right).

b. MF stimulation with an artificial burst (20 stimuli at 100 spk/s; green trace, *top*) and instantaneous firing rates of the responses in 4 UBCs (fast, mid-range, slow, and OFF; individual trials in grey, mean in black). Dashed gray lines indicate 0 spk/s.

c. As in **b**, but for a MF firing pattern recorded *in vivo* during a smooth pursuit task in which the MF primarily fired in brief high frequency bursts. The numbers of spikes in each burst larger than 2 spikes are indicated in the trace.

d. As in **b**, but for a MF firing pattern recorded *in vivo* during a smooth pursuit task with characteristic long-lasting increases and decreases in firing.

Next, we stimulated MF inputs to the same four UBCs using an *in vivo* MF firing pattern associated with smooth pursuit trials (**Fig. 1d**). This firing pattern consists mossy fiber firing at 10-20 spk/s that periodically increases to 40-80 spk/s. The three excitatory UBCs responded with firing patterns that were qualitatively similar to the mossy fiber input, with irregular firing at 10-50 spk/s periodically increasing by 20-60 spk/s in response to elevations in mossy fiber firing. However, UBC responses were delayed relative to mossy fiber inputs, peaking later and remaining elevated after the MF firing rates decreased. As such, the firing of the excitatory UBCs appeared to be a low-pass filtered version of the MF input, with the time course progressively slower and the extent of filtering more pronounced for the fast, mid-range and slow cells. The slow cell increased its firing only after the elevation in MF firing. The OFF UBC completely shut down during the increase in MF firing, and only fired in between increases in firing, or in response to decreases in firing, effectively inverting the MF firing pattern. Overall, the different UBCs provided diverse temporal transformations of the *in vivo* MF input patterns.

The influence of MF burst duration on UBC responses

Given the diversity of spike numbers in MF bursts *in vivo*, we systematically examined the effects of MF burst duration on the UBC response. Previously, we found that mossy fiber activation with a burst of 20 stimuli at 100 spk/s evoked a continuum of temporal responses in a population of UBCs (Guo et al., 2021b; Huson, Newman and Regehr, 2023). We extended this approach by stimulating MFs with single stimuli, and 100 spk/s bursts of 2, 5, 10, and 20 stimuli (**Fig. 2a**, green trace). We sorted UBCs (n=70) by their responses to the 20 stimuli at 100 spk/s burst, either according to the half-widths of their increases in firing (cells #1-61), or by the pause durations following stimulation (cells #62-70) (Guo et al., 2021b) (**Fig. 2b**).

UBCs had highly diverse responses to different MF burst durations, as illustrated by four example cells (**Fig. 2a**), and by the responses of all cells summarized in a heatmap (**Fig. 2b**). Single stimuli and brief bursts evoked large rapid increases in firing in close to half of the UBCs (**Fig. 2a**, cell 20; **Fig. 2b**, most cells #1-40), and, surprisingly, peak firing rates were quite similar for single stimuli and 20 stimuli bursts for many of these UBCs. Different duration bursts evoked responses with very different time courses, as illustrated by cell #20 (**Fig. 2a**), in which a single stimulus evoked a rapid increase in firing, but a 20-stimulus burst evoked a very short-lived increase at the onset of stimulation, before briefly stopping, and then resuming after stimulation ends. Single stimuli and brief bursts evoked decreases in firing for approximately 20% of the UBCs (**Fig. 2a**, cells #55 and #67; **Fig. 2b**, most cells #55-70). Single stimuli generally had very small effects on firing for UBCs with intermediate properties, but the size of the response increased as the stimulus duration increased (**Fig. 2a**, cell #39; **Fig. 2b**, most cells #41-54).

Summaries of peak increases in firing (**Fig. 2c,d**), the number of spikes evoked by MF stimulation (**Fig. 2e**), and the half-width of firing-rate increases (**Fig. 2f**) for all burst durations revealed systematic trends in the properties of UBC responses. There were large differences in the dependence of peak firing on burst duration (**Fig. 2c**). Fast UBCs responded to all burst durations with consistently high peak responses that exceeded 100 spk/s, while intermediate and slow UBCs increased their peak response almost linearly with burst duration (**Fig. 2c**). As such, peak firing rates vary continuously across a wide range in response to 2 stimulus bursts, while in response to 20 stimulus bursts an apparent ceiling restricts the peak firing rates to a narrower range (**Fig. 2d**). The number of spikes evoked by MF stimulation increased with burst duration in almost all cells, but increases were small in fast cells, and could grow more than 100-fold in slow cells (**Fig. 2e**). The half-widths of responses also increased with input duration for almost all UBCs, with the largest increases apparent when the number of stimuli increases from one to five (**Fig. 2f**).

Overall, these results indicate that UBCs can provide long-lasting responses to bursts with a wide range of input durations. After single stimuli and short bursts only fast UBCs respond with clear increases in firing, but these responses remain consistent as burst duration increases. This




Figure 2

Diverse UBC responses to MF bursts of increasing duration.

- a.** Instantaneous firing rates of four representative UBCs in response to 100 spk/s bursts comprised of 1-20 stimuli (input pattern indicated above in green; dashed gray lines indicate 0 spk/s; cell numbers refer to the index in the summary plot **b**).
- b.** Heat maps showing the normalized instantaneous firing rates for all UBCs in response to bursts comprised of 1-20 stimuli. Responses were normalized per cell to the peak firing rate in response to 20 stimuli at 100 spk/s bursts, and sorted by the half-width for cells with clear increases in firing (cell #1-61), or sorted by pause duration for the rest (cell #62-70). Time indicates seconds since start of the MF stimulation (indicated by dotted red line). Red arrows indicate representative UBCs shown in **a**.
- c.** Summary plot showing peak increase in firing rate on a log scale, for all different bursts by the number of input spikes also on a log scale. UBCs color coded to correspond to the cell index in **b**. Cells without significant increases in firing excluded from this plot (#62-70).
- d.** Separate plots of the peak increase in firing rate in response to 2 and 20 stimuli bursts sorted by cell index as in **b**.
- e.** As in **c** but for the number of spikes evoked by the different MF burst stimulations.
- f.** As in **c** but for the half-width of the increase in firing for the different MF burst stimulations.

suggests that fast UBCs are suited to detecting the occurrence of a burst, regardless of burst duration. In contrast, slow UBCs are suited to detecting bursts of 5-20 MF spikes and scaling their responses proportionally to burst duration.

Different glutamate receptors mediate distinct components of UBC responses

Many types of glutamate receptors are present at MF to UBC synapses, and it is not clear how each type contributes to responses evoked by MF bursts of different durations. We focused our studies on AMPARs and mGluR1s that elicit excitatory currents in UBCs, and mGluR2/3s that elicit inhibitory currents (Rossi et al., 1995 [\[1\]](#); Kinney, Overstreet and Slater, 1997 [\[2\]](#); Russo et al., 2008 [\[3\]](#); Kim et al., 2012 [\[4\]](#); Schwartz et al., 2012 [\[5\]](#); Borges-Merjane and Trussell, 2015 [\[6\]](#); Guo et al., 2021b [\[7\]](#)). NMDARs are also present at UBC synapses, but we found that they do not make an appreciable contribution to responses evoked by bursts (**Fig. 3 - Supplementary Fig. 1** [\[8\]](#)), and we therefore did not explicitly examine their contributions in combination with antagonist for the other glutamate receptors.

We evoked responses in 31 UBCs with single stimuli, 2, 5, 10, and 20 stimuli at 100 spk/s while additively washing in antagonists of mGluR2/3s (LY341495; 1-5 μ M), AMPARs (NBQX; 5 μ M), and mGluR1s (JNJ16259685; 1 μ M) (**Fig. 3a** [\[9\]](#), above). Blocking mGluR2/3 eliminated the pause after MF stimulation (e.g. cell #27, **Fig. 3a** [\[9\]](#), yellow, and **Fig. 3d** [\[10\]](#), second row), in line with previous findings (Russo et al., 2008 [\[3\]](#); Kim et al., 2012 [\[4\]](#); Borges-Merjane and Trussell, 2015 [\[6\]](#)). Blocking mGluR2/3s also slightly increased peak responses (**Fig. 3b** [\[11\]](#)), and the number of evoked spikes in some cells (**Fig. 3c,e** [\[12\]](#)), especially for larger bursts, with significantly more spikes after blocking mGluR2/3s for burst of 5 stimuli or more (**Fig. 3f** [\[13\]](#)). The additional block of AMPARs eliminated increases in firing evoked by single stimuli in most cells (e.g. cell #8, **Fig. 3a-c** [\[14\]](#), red, and **Fig. 3d** [\[10\]](#), third row), but its effect diminished with increasing numbers stimuli in the burst (**Fig. 3b,c,e** [\[15\]](#)), and on average AMPARs were not the main mediator of the increase in firing for bursts of 5 stimuli or more (**Fig. 3g** [\[16\]](#)). The component of the response that was sensitive to blocking mGluR1 increased as the bursts got longer and was the main mediator of the increase in firing in almost all UBCs for 20 stimulus bursts (**Fig. 3b,c,e,h** [\[17\]](#)). In a few UBCs, some short-lived responses remained after blocking both AMPARs and mGluR1s (e.g. cell #27 **Fig. 3a,b** [\[18\]](#)), but this did not represent a significant portion of the evoked spikes (**Fig. 3c,e** [\[19\]](#)).

AMPA-mediated responses to single stimuli are more common in fast UBCs (**Fig. 3d** [\[20\]](#)). This is surprising since AMPAR subunits are expressed homogeneously across within the UBC population (Guo et al., 2021b [\[7\]](#); Kozareva et al., 2021 [\[21\]](#)) (**Fig 3 - Supplementary Fig. 2b** [\[22\]](#)). It is therefore surprising that AMPAR responses are so small or are not apparent in slower UBCs and OFF UBCs. Although mGluR2/3 inhibition obscured small AMPAR components in some cells (e.g. cell #10, **Fig. 3a** [\[9\]](#)), this was uncommon and did not account for the differences in AMPAR-mediated responses in the UBC population (**Fig. 3d** [\[20\]](#)). It is possible that differential expression of auxiliary proteins governs the amount of AMPAR current present in a UBC. TARP γ -2 controls the amplitude of slow AMPAR EPSCs in UBCs (Lu et al., 2017 [\[23\]](#)), and it may be slightly elevated in UBCs that highly express mGluR1 (**Fig 3 - Supplementary Fig. 2** [\[24\]](#)), which corresponds to fast excitatory UBCs (Borges-Merjane and Trussell, 2015 [\[6\]](#); Guo et al., 2021b [\[7\]](#); Kozareva et al., 2021 [\[21\]](#)). TARP γ -8 and TARP γ -7 have similar properties as TARP γ -2 (Jackson and Nicoll, 2011 [\[25\]](#)) and are also present in UBCs at low levels in UBCs. TARP γ -8 has an inverse expression pattern, and TARP γ -7 is expressed homogeneously, but their roles in regulating AMPARs in UBCs has not been examined. The expression of GSG1L, an AMPAR auxiliary subunit that is implicated in controlling AMPAR desensitization (Shanks et al., 2012 [\[26\]](#)), is of particular interest. GSG1L is expressed at high levels and shows a pronounced expression gradient (**Fig 3 - Supplementary Fig. 2c** [\[27\]](#)).




Figure 3

Differential glutamate receptor contributions to responses evoked by MF bursts of increasing duration.

a. Instantaneous firing rates of five representative UBCs in response to 100 spk/s bursts comprised of 1-20 stimuli (stimulus indicated by green bars). Responses shown for baseline conditions (black), and after addition of antagonists of mGluR2/3 (yellow), AMPAR (red), and mGluR1 (blue). Glutamate receptor antagonists were applied successively on top of the previous antagonist(s) as depicted in the scheme at the top. Cell numbers refer to the index in the summary plot **d**.

b. Summaries of the peak change in firing rate for the same five cells, with successive different markers for baseline and the different glutamate antagonists, and separate plots for the different MF burst stimulations.

c. As in **b** but for the number of spikes evoked by the MF stimulation.

d. Heat maps showing the normalized instantaneous firing rates for all UBCs in response to 100 spk/s burst comprised of 1-20 stimuli (columns), for baseline and after addition of glutamate receptor antagonists (rows). Responses were normalized per cell to the peak firing rate in response to the 20 stimuli at 100 spk/s burst with mGluR2/3 blocked. Cells sorted by their response to the baseline 20 stimuli at 100 spk/s input, either by the half-width of the increase in firing (cell #1-27) or by pause duration (cell #28-31). Time indicates seconds since start of MF stimulation (indicated by dotted red line). Red arrows indicate representative UBCs shown in **a-c**.

e. Summary plots of the number of spikes evoked by the MF stimulation on a log scale for all UBCs color coded to correspond to the cell index in **d**. Successive different markers indicate baseline and the different glutamate antagonists, and separate plots for the different MF burst stimulations.

f. Violin plots of the number of spikes after each burst under baseline conditions (-) and after blocking mGluR2/3 (+), normalized per cell to the number of spikes after 20 stimuli at 100 spk/s under baseline conditions. Markers indicate individual UBCs color coded by the cell index in **d**. (* $p < 0.01$, Wilcoxon signed rank test).

g. Violin plots of the percentage of the number of spikes evoked by MF stimulation that was mediated by AMPARs, estimated from the effect of blocking AMPARs on the response. Markers indicate individual UBCs color coded by the cell index in **d**. Responses smaller than 5 spikes not shown.

h. As in **g** but for the component mediated by mGluR1

In conclusion, each type of glutamate receptor makes a distinct contribution to responses to bursts. mGluR2/3s mediate pauses in firing for both short and long bursts in over half of the cells. AMPARs are important mediators of firing rate increases evoked by short bursts in fast UBCs. mGluR1s are the primary mediators of responses to medium and long duration bursts.

UBCs provide both time-locked and temporally-filtered responses

Next, we systematically characterized UBC responses evoked by stimulating MFs with input patterns that contain features that are characteristic of MF firing measured during smooth pursuit tasks. To achieve this, we exposed 31 cells to a smooth pursuit-like input pattern with prolonged baseline stimulation at 5 spk/s, interposed by 1 second steps to frequencies, ranging from 10 to 60 spk/s, at 4 second intervals (**Fig. 4a**, green, top). We interposed this stimulation protocol with burst stimulations and used the responses to 20 stimuli at 100 spk/s bursts to sort cells for display purposes (**Fig. 4a,c**, left).

UBCs showed diverse responses to this input pattern. For some UBCs, each stimulus during 5 spk/s baseline input evoked brief increases in firing, and steps to higher firing rates evoked slower, more prolonged increases in firing (**Fig. 4**, cell #4). Other fast UBCs did not respond to 5 spk/s stimulation, but steps to 10-60 spk/s evoked responses that began with a delay and persisted after the step ended (**Fig. 4**, cell #8). Along the sorted UBCs, 1 s steps evoked responses with increasingly delayed onset, and longer lasting increases in firing (**Fig. 4**, cells #14 and #17). For slow cells, steps to 10-60 spk/s silenced firing during stimulation, and persistently elevated UBC firing after the step had ended (**Fig. 4**, cells #22 and #27). Some slow UBCs also showed discrete decreases in firing during baseline 5 spk/s input (**Fig. 4a,b**, cell #22), while others did not (**Fig. 4a,b**, cell #27). These responses illustrate that UBCs provide both a temporal transformation of sustained changes in firing frequency, and discrete responses evoked by single stimuli. Single stimuli evoked changes in firing in a large percentage of cells for 5 spk/s stimulation, and responses were even larger for 1 spk/s and 2.5 spk/s (**Fig. 4 – Supplementary Fig. 1**).

As the step change of the input rate increased, the UBC population showed an increasingly wider range of responses. The number of evoked spikes that occurred during the 1 s steps increased for fast and intermediate UBCs, while slower UBCs tended to reduce their firing; with 45% of UBCs (14/31) reducing firing during the 60 spk/s step (**Fig. 4e**). Step changes more consistently evoked increases in firing in the 3 s after the step, with 87% (27/31) of UBCs firing more spikes after a 60 spk/s step (**Fig. 4f**), leaving only the slowest cells (#28-31) with net decreases in spiking. Some UBCs showed much larger responses for higher stimulus frequencies (e.g. cell #8 and #27, **Fig. 4a**), while others were strikingly consistent (e.g. cell #4 and #17, **Fig. 4a**). However, this property did not vary in alignment with the sorting of the UBCs by their half-width (**Fig. 4e,f**). In cells with clear increases in firing after step changes (cells #1-27), we estimated time to peak for the responses to 60 spk/s steps by fitting the instantaneous firing rates with a log-gaussian function (see **Methods**). We observed gradually more delayed peaks in line with the UBC sorting ranging from less than 20 ms to almost 2 s (**Fig. 4g**). Half-decay times of the step responses were also highly correlated with the half-widths of the responses to 20 stimuli at 100 spk/s bursts (for the 60 spk/s step: Spearman's $\rho = 0.873$, $p = 1.40 \times 10^{-6}$), and did not consistently change with firing frequency. Of the cells that showed increases in firing, 33% (9/27) took more than 1 s after the steps for their response amplitude to decay by half (**Fig. 4h**), causing increases in firing to gradually build during the protocol, such that the response to step to 20 spk/s late in the protocol was far larger than one early in the protocol (e.g. cell #22 and #27, **Fig. 4a**).

Overall, during smooth pursuit-like MF input, fast UBCs closely follow the input pattern, while slow UBCs produce strongly temporally-filtered responses, in line with responses to 20 stimuli at 100 spk/s bursts. However, many UBCs additionally preserve timing information with discrete changes in firing time-locked to lower rate baseline stimuli. As such, multiple streams of information appear to be simultaneously encoded by the response.




Figure 4

The UBC population displays a continuum of long-lasting responses to smooth pursuit-like input.

- a.** Instantaneous firing rates of seven representative UBCs in response to 20 stimuli at 100 spk/s burst (*left*), and smooth pursuit-like MF input (*right*) with prolonged 5 spk/s input interposed by 1 s step to 10-60 spk/s. Input pattern indicated above (green traces). Dashed gray lines indicate 0 spk/s; cell numbers refer to the index in the summary plot **c**.
- b.** The same as in **a** but on an expanded timescale, displaying only the 60 spk/s step. Dotted red lines indicate onset and offset of the 60 spk/s step.
- c.** Heat maps showing the normalized instantaneous firing rates for all UBCs in response to 20 stimuli at 100 spk/s burst (*left*) and smooth pursuit-like MF input (*right*). Responses normalized to the peak firing rate per cell separately for burst and smooth pursuit-like MF input. Cell sorted by their response to 20 stimuli at 100 spk/s burst input, either by the half-width of the increase in firing (cell #1-25) or by pause duration (cell #26-31). Dotted red lines indicate the start of the step changes in input rate, red arrows indicate representative UBCs shown in **a,b**.
- d.** The same as in **c** but on an expanded timescale, displaying only the 60 spk/s step.
- e.** Summary plot of the number of evoked spikes during the 1 s steps compared to the 1 s period preceding the step. Individual UBCs color coded to correspond to the cell index in **c**.
- f.** As in **e** but for the number of evoked spikes in the 3 s period after the 1 s steps.
- g.** Summary plot of the time to peak for log-gaussian fits of the response to the step to 60 spk/s. Cells sorted and color coded to correspond to the cell index in **c**. Cells without significant increase in firing after step changes excluded (cell #28-31).
- h.** Summary plot of the time for the increase in firing to decay by half after a step change. Cells without significant increase in firing after step changes excluded (cell #28-31).

mGluR1 is the main mediator of UBC responses to smooth pursuit-like input patterns

We used selective antagonists to determine the roles of different glutamate receptors in mediating responses evoked by the stimulus protocol in [Fig. 4a](#). NMDARs do not make an appreciable contribution to responses evoked by this stimulus ([Fig. 5 - Supplementary Fig. 1](#)), and we therefore did not explicitly examine their contributions in combination with other glutamate-receptor antagonists. We again additively washed in antagonists of mGluR2/3s, AMPARs, and mGluR1s. For a fast UBC that responded to each 5 spk/s stimulation and to 1 s increases in input rate, blocking mGluR2/3s had minor effects, while blocking AMPARs eliminated responses to single stimuli only, and blocking mGluR1 eliminated responses to 1 s steps to 10-60 spk/s ([Fig. 5a,b](#), cell #3). In another cell ([Fig. 5a,b](#), cell #7), 5 spk/s stimulation did not evoke prominent responses, but step increases in input frequency evoked delayed and long-lasting increases in firing. In this cell, blocking mGluR2/3s shortened the delay between the step increases in MF stimulation and UBC firing rate increases ([Fig. 5a,b](#), cell #7), blocking AMPARs had very little effect, and blocking mGluR1s eliminated UBC responses. In slow UBCs, suppression of mGluR2/3s also removed discrete inhibitory responses to low-rate baseline input ([Fig. 5 - Supplementary Fig. 2b,d,f-h](#)). Blocking AMPARs eliminated excitatory responses evoked by individual stimuli at 5 spk/s ([Fig. 5a-d](#)), 1 spk/s and 2.5 spk/s stimuli ([Fig. 5 - Supplementary Fig. 2](#)).

For most UBCs, 1 s 10-60 spk/s MF stimulation evoked slow changes in UBC firing that remained prominent in the presence of mGluR2/3 and AMPAR antagonists. These responses were eliminated by blocking mGluR1 ([Fig. 5a-d](#)). Plotting the change in spikes evoked by each 1 s step for the successive application of antagonists revealed that many UBCs showed clear increases in their excitatory response after suppression of mGluR2/3s ([Fig. 5e,f](#)), which were also evident in the heatmaps ([Fig. 5c,d](#)). Suppression of AMPARs only mildly reduced the number of evoked spikes for most steps ([Fig. 5g](#)), with 90% (9/10) of UBCs that showed increases in firing to the 60 spk/s step before suppression of AMPAR, still responding after ([Fig. 5e](#)). However, the number of evoked spikes was close to zero for all UBCs after suppression of mGluR1 ([Fig. 5e,h](#)).

We conclude that, during smooth pursuit-like input patterns, mGluR1 is the main mediator of excitatory responses, while mGluR2/3 is the main mediator of inhibitory responses. and, additionally, reduces excitation overall throughout stimulation. For low-rate inputs in fast UBCs, AMPARs mediate an additional excitatory response that preserves timing information of the input, while mGluR2/3s provide an analogous inhibitory response in slower UBCs.

Discussion

Our findings establish that the UBC population powerfully preprocesses MF inputs during realistic MF activation. UBCs simultaneously convey timing information, and a slow filtered version of the MF firing pattern that continuously varies across the population ([Fig. 6b](#)). The multiple types of glutamate receptors present at MF-UBC synapses underlie the ability of UBCs to create diverse, multicomponent temporal responses in granule cells ([Fig. 6c-e](#)).

The roles of AMPARs in UBC responses to MF input

We find that for many UBCs AMPARs convey timing information about MF firing by responding to single stimuli, short bursts, and individual stimuli during prolonged low-frequency MF activation ([Fig. 3](#), [Fig. 5](#), [Fig. 5 - Supplementary Fig. 2](#), [Fig. 6c](#)). AMPARs primarily mediate responses to high frequency bursts of up to 5 stimuli, but as the burst duration increases their contribution diminishes and mGluR1s are the primary driver of increases in firing for most UBCs




Figure 5

Contributions of mGluR2/3, AMPAR, and mGluR1 to responses evoked by smooth pursuit-like input.

- a.** Instantaneous firing rates of two representative UBCs in response to 20 stimuli at 100 spk/s bursts (*left*), and smooth pursuit-like MF input (*right*) as indicated by the traces at the top (green). Responses shown under baseline conditions and after successive addition of antagonists of mGluR2/3, AMPAR, and mGluR1. Dashed gray lines indicate 0 spk/s; cell numbers refer to the index in the summary plot **c**.
- b.** The same as in **a** but on an expanded timescale, displaying only the 60 spk/s step. Dotted red lines indicate onset and offset of the 60 spk/s step.
- c.** Heat maps showing the normalized instantaneous firing rates for all UBCs in response 20 stimuli at 100 spk/s bursts (*left*) and smooth pursuit-like MF input (*right*), for baseline and after addition of glutamate receptor antagonists (*rows*). Responses normalized per cell separately for burst and smooth pursuit-like MF input to the respective peak firing rates with mGluR2/3 blocked. Cells sorted by their response to the baseline 20 stimuli at 100 spk/s input, either by the half-width of the increase in firing (cell #1-9) or by the pause duration (cell #10-11). Dotted red lines indicate the start of the step changes in input rate, red arrows indicate representative UBCs shown in **a,b**.
- d.** The same as in **c** but on an expanded timescale, displaying only the 60 spk/s step.
- e.** Summary plots of the number of evoked spikes in the 1 s period during the step and the 3 s period follow it for all UBCs color coded to correspond to the cell index in **c**. Successive different markers indicate baseline and the different glutamate antagonists, and separate plots for the different step changes in input.
- f.** Violin plots of the number of spikes in the 1 s period during the step and the 3 s period after under baseline conditions (-) and after blocking mGluR2/3 (+), normalized per cell to the number of spikes associated with the step to 60 spk/s under baseline conditions. Markers indicate individual UBCs color coded by the cell index in **c**. (* $p < 0.01$, Wilcoxon signed rank test).
- g.** Violin plots of the percentage of the number of spikes evoked by MF stimulation that was mediated by AMPARs, estimated from the effect of blocking AMPARs on the response. Markers indicate individual UBCs color coded by the cell index in **d**. Responses smaller than 5 spikes not shown.
- h.** As in **g** but for the component mediated by mGluR1.




Figure 6

Schematic summarizing the range of UBC responses evoked by different stimuli.

a. MF stimuli used to evoke responses in UBCs. **b-e).** Drawn representations of instantaneous firing rates illustrate the different types of UBC responses and the contributions of different types glutamate receptors. Four "cells" are shown that range from fast "ON" to slow "OFF" cells that have the properties of classic ON and OFF cells, respectively. Two additional "cells" with intermediate properties are also shown.

b. Total responses evoked by the stimuli in **a**.

c. Contributions of the AMPAR component to responses in **b**.

d. Contributions of the mGluR1 component to responses in **b**.

e. Contributions of the mGluR2/3 component to responses in **b**.

(Fig. 3). A large fraction of the UBC population fire in response to each stimulus during 1-5 spk/s MF stimulation, and these responses are driven almost entirely by AMPARs (Fig. 4 - Supplementary Fig. 1, Fig. 5 - Supplementary Fig. 2). Although brief compared to UBC responses mediated by mGluR1s following prolonged bursts, AMPAR responses still last up to 400 ms, which is much longer than AMPA-mediated EPSCs observed at most synapses. This slow time course is consistent with the properties of AMPAR EPSCs at MF-UBC synapses that arise from the ultrastructure of UBC synapses, the resulting slow glutamate signal, and the distinctive properties of the AMPARs present in UBCs (Rossi et al., 1995; Kinney, Overstreet and Slater, 1997; van Dorp and De Zeeuw, 2014; Lu et al., 2017; Balmer, Borges-Merjane and Trussell, 2021). Several aspects of AMPAR-mediated spiking are readily explained by known properties of UBC synapses. The multiple phases of the AMPAR response evoked by 20 stimuli at 100 spk/s bursts likely reflect initial activation, desensitization during stimulation, and a large slow rebound current after stimulation as the prolonged presence of glutamate activates AMPARs as they recover from desensitization (Fig. 3a cell #8, Fig. 3d)(Kinney, Overstreet and Slater, 1997; van Dorp and De Zeeuw, 2014; Lu et al., 2017; Balmer, Borges-Merjane and Trussell, 2021). The frequency-dependent decrease in AMPAR mediated burst magnitude (compare 1 and 5 spk/s Fig. 4 - Supplementary Fig. 1, Fig. 5 - Supplementary Fig. 2) also likely arises from AMPAR desensitization (van Dorp and De Zeeuw, 2014).

Contributions of AMPARs to MF-evoked responses are most apparent in fast UBCs (Fig. 2), even though AMPAR subunits are expressed uniformly across the population (Guo et al., 2021b; Kozareva et al., 2021) (Fig. 3 - Supplementary Fig. 2b). It is therefore surprising that AMPAR responses are so small or are not apparent in slower UBCs and OFF UBCs. It is possible that differential expression of auxiliary proteins such as TARPs or GSG1L governs the amplitude of AMPAR responses (Fig. 3 - Supplementary Fig. 2c), but further experiments are required to determine if this is the case.

The roles of mGluR2/3s in MF-UBC responses

Suppression provided by mGluR2/3 receptors influences UBC responses in multiple ways (Fig. 6e). (1) It suppresses baseline firing after a MF burst in “OFF” UBCs, a role that has been extensively described previously (Knoflach and Kemp, 1998; Russo et al., 2008; Kim et al., 2012; Borges-Merjane and Trussell, 2015). (2) In most cells with a prominent pause in firing, suppression is followed by slow excitation mediated by mGluR1s. In these cells mGluR2/3s introduces a delay between the onset of MF stimulation and the response in the UBC. This is important for many UBCs that respond to 1 s MF activation with a delayed peak in firing that occurs long after the increase in MF firing rate. In this way mGluR2/3s crucially contribute to UBCs filtering of MF inputs by selectively influencing the slow rise of UBC spiking without influencing the time course of decay. (3) There is also an interplay between mGluR2/3 suppression and excitation by AMPARs and mGluR1s such that even in cells when suppression is not apparent, mGluR2/3s reduce the number of spikes evoked by stimulation. (4) In many UBCs, single stimuli lead to decreases that are mediated by mGluR2/3s.

The roles of mGluR1s in MF-UBC responses

mGluR1s mediate slow responses that gradually build during repetitive activation (Fig. 6d). Single stimuli and brief bursts build up too little glutamate to effectively activate mGluR1 and drive excitation (Batchelor, Madge and Garthwaite, 1994; Tempia et al., 1998; Brasnjo and Otis, 2001; Wadiche and Jahr, 2005). mGluR1-mediated responses increase as the total number of spikes in the burst increases. In UBCs with prominent AMPAR responses, increasing the number of spikes in a high frequency burst leads to a transition from responses being primarily mediated by AMPARs to responses being mediated by mGluR1s. In these UBCs a combination of AMPARs and mGluR1s conveys relatively consistent increases in peak firing that could not be achieved by either receptor alone (Fig. 2, 3). Such UBCs are highly effective at detecting the presence of a burst, but they are not suited to differentiating between bursts comprised of different numbers of MF

spikes. In UBCs with a small AMPAR component and a prominent mGluR1 component, the UBC peak firing rate of the UBC response is strongly related to the number of spikes in the MF input burst. Similarly, increasing the firing rate of 1 s steps during prolonged MF input, as occurs during smooth pursuit eye movements, leads to increases in the magnitude of UBC responses. Additionally, mGluR1s also participate in temporally filtering MF input by mediating broad UBC responses of different durations (Guo et al., 2021b [↗](#); Huson, Newman and Regehr, 2023 [↗](#)). However, during prolonged stimulation mGluR1s only play a crucial role in regulating the duration of UBC spiking following changes in the input rate, whereas the delay in the firing rate increase reflects the time course of mGluR2/3 suppression and is not limited by the time course of mGluR1 activation.

Functional roles of MF transformations by the UBC population

Although the manner in which UBCs contribute to cerebellar-dependent behaviors has not been established, the properties of transformations at MF-UBC synapses provide insight into several possible functions. With regard to eye movements, the cerebellum is thought to aid in adaptation for real-time error correction during target tracking, and in learning to predict target trajectory both for saccades and smooth eye movements (Shadmehr, 2017 [↗](#); Soetedjo, Kojima and Fuchs, 2019 [↗](#); Lisberger, 2021 [↗](#)).

Saccadic eye movements are represented in MF signaling by rapid bursts preceding the onset of movement. The peak firing rate of a MF burst can encode the saccade amplitude (Ohtsuka and Noda, 1992 [↗](#)). In Purkinje cells eye velocity during saccades is encoded in the average response of the population, with individual cells showing diverse firing patterns including increases and decreases that can outlast the saccade (Herzfeld et al., 2015 [↗](#)). Changes in the simple spike firing rate has been suggested to underly adaptation of saccadic eye movements (Soetedjo, Kojima and Fuchs, 2019 [↗](#)). UBC responses similarly consist of increases or decreases in firing that outlast the input, and as such, may contribute to the diversity in Purkinje cell firing rates, possibly providing a wider distribution of signals to support adaptation.

Visual feedback to the cerebellum lags eye movement by more than 100 ms (Raymond and Lisberger, 1998 [↗](#)). Therefore, to accurately track a target along its trajectory as during smooth pursuit, predictive eye movement signals are required. Temporal transformations by UBCs may be crucial to generating diverse signals to provide Purkinje cells with the appropriate substrate for learning, similar to how UBCs contribute to the diversity required for generating predictions of electrosensory feedback in weakly electric fish (Kennedy et al., 2014 [↗](#)). Furthermore, mGluR2/3-generated delays in UBC responses are suited to encode timing, which is consistent with the hypothesis that timing rather than target position predicts the trajectory (Medina, Carey and Lisberger, 2005 [↗](#)). Generally, learning a target trajectory takes place over hundreds of trials (Medina, Carey and Lisberger, 2005 [↗](#); Hall, Yang and Lisberger, 2018 [↗](#)), but short-term plasticity is also exhibited when an instructive signal given in one trial biases eye movement in the next trial (Yang and Lisberger, 2010 [↗](#)). This bias persists for 4-10 s, which is comparable to the durations of mGluR1-mediated increases in firing in slow UBCs (Guo et al., 2021b [↗](#); Huson, Newman and Regehr, 2023 [↗](#)) (Fig. 2b [↗](#)). This raises the possibility that UBCs may provide a short-term plasticity signal that could be used to bias motion.

MF signals encoding vestibular modulation are also likely transformed by UBCs to enable behavioral adaptation. Previous work has highlighted that during sinusoidal vestibular input MF signals mostly remain in-phase, while granule cells fire in a range of different phases (Arenz et al., 2008 [↗](#); Barmack and Yakhnitsa, 2008 [↗](#)). UBCs have been shown to phase shift sinusoidal inputs, enabling the cerebellum to learn to output signals at arbitrary shifts from the input which may enable learning of phase-shifted VOR (Zampini et al., 2016 [↗](#)). That study focused on AMPAR and mGluR2/3 mediated changes in firing. The combination of delayed temporally-filtered responses mediated by mGluR2/3s and mGluR1s, and the discrete changes in firing in response to single

stimuli we describe here, enables the UBC population to provide transformations that are more complex than simple phase-shifts (**Fig. 4**), and may be used for adaptations at longer time scales and to irregular input patterns.

Having separate response components also broadens the types of input UBCs can respond to, with AMPAR and mGluR2/3 responding to brief inputs, and mGluR1 and mGluR2/3 responding to long-lasting inputs. While UBCs are enriched in regions processing eye and vestibular signals, they are consistent circuit elements throughout the cerebellum (Floris et al., 1994; Dino, Willard and Mugnaini, 1999; Takacs et al., 1999; Englund et al., 2006). The separate response components may provide UBCs with the flexibility required to usefully transform MF inputs in a wide range of behaviors.

Possible Influence of Inhibition

Our experimental approach only evaluates the direct effects of MF activation on UBC firing and does not incorporate inhibition, which is expected to influence UBC firing *in vivo*. UBCs are inhibited by Golgi cells (Rousseau et al., 2012), and a subset of UBCs are also inhibited by PCs (Guo et al., 2021a). Although the potential influence of inhibition is of considerable interest, brain slice experiments are poorly suited to addressing this issue. Golgi cells are fragile and many do not survive in adult brain slices (Hull and Regehr, 2012), and many of the complex PC axon collaterals are severed before reaching UBCs (Guo et al., 2021a). In addition, it is exceedingly difficult to activate the population of PCs and Golgi cells as would occur *in vivo*. In contrast to the simplicity of a single MF input exciting a UBC, Golgi cells are directly activated by many MFs, by MF → granule cell → Golgi cell synapses, and by MF → UBC → granule cell → Golgi cell synapses (Hull and Regehr, 2022), and PCs are activated by a great many MF → granule cell → PC synapses (Witter et al., 2016), by MF → UBC → granule cell → PC synapses and granule cell → molecular layer interneuron synapses (Hull and Regehr, 2022). As a result of these complexities it will therefore be necessary to perform experiments *in vivo* to determine how Golgi cell and PC inhibition influences UBC transformations of MF inputs.

Methods

Mice

All animal procedures were carried out in accordance with the NIH and Animal Care and Use Committee (IACUC) guidelines and protocols approved by the Harvard Medical Area Standing Committee on Animals (protocol #IS00000124). C57BL/6 mice of either sex from Charles River Laboratories were used for all experiments.

Slice preparation

Juvenile (P30–P47) C57BL/6 mice of either sex were anesthetized by a peritoneal injection of 10 mg/kg ketamine/xylazine mixture and then transcardially perfused with ice-cold cutting solution (in mM): 110 choline chloride, 2.5 KCl, 1.25 NaH₂PO₄, 25 NaHCO₃, 25 glucose, 0.5 CaCl₂, 7 MgCl₂, 3.1 sodium pyruvate, and 11.6 sodium ascorbate, equilibrated with 95% O₂ and 5% CO₂. The brain was subsequently extracted, dissected, and submerged in the same solution. 220 μm parasagittal slices from the cerebellar vermis were cut on a Leica VT1200S vibratome. Slices were then incubated for 30 minutes at 34°C in artificial cerebral spinal fluid (ACSF) containing (in mM): 125 NaCl, 26 NaHCO₃, 1.25 NaH₂PO₄, 2.5 KCl, 1 MgCl₂, 1.5 CaCl₂ and 25 glucose, equilibrated with 95% O₂ and 5% CO₂. Following incubation, the slices were kept for up to 6 hours at room temperature.

Electrophysiology

Recordings were performed at $\sim 34^{\circ}\text{C}$ in ACSF (flow rate 2–4 ml / min) containing inhibitory receptor blockers picrotoxin (20 μM ; Tocris), CGP 55845 (1 μM ; Abcam), and strychnine hydrochloride (1 μM ; Abcam). Visually guided cell-attached recordings were made in lobule X of the cerebellar vermis with 3–5 M Ω patch pipettes pulled from borosilicate glass capillaries (BF150-86-10, Sutter Instrument) with a P-97 Flaming/Brown puller (Sutter Instrument), and filled with ACSF. Once a loose seal was achieved (>10 M Ω), a bipolar theta glass pipette (BT150-10, Sutter Instrument) was placed in the granular layer more than 50 μm from the patched cell, and repositioned until a mossy fiber input was stimulated. UBCs were identified based on their slightly larger soma size compared to granule cells, and their distinctive long-lasting firing upon mossy fiber stimulation. Data were collected with a Multiclamp 700B amplifier (Molecular Devices), filtered at 4 kHz (4-pole Bessel filter) online, digitized at 20 kHz with an ITC18 (Heka Instrument), and saved using mafPC3 (custom software written by M. Xu-Friedman, <https://www.xufriedman.org/mafpc>) in Igor Pro 8 (WaveMetrics Inc.) for offline analysis.

Our previous study (Guo et al., 2021b) explored issues related to the reliability of MF activation, the possibility of glutamate spillover from other synapses, and the possibility of disinaptic activation involving stimulation of MF $\rightarrow$ UBC $\rightarrow$ UBC connections (Hariani et al., 2024). We did on-cell recordings and followed that up with whole cell voltage clamp recordings from the same cell and there was good agreement with the amplitude and timing of spiking and the time course and amplitudes of the synaptic currents (Guo et al., 2021b). We also compared responses evoked by focal glutamate uncaging over the brush and MF stimulation and found that the time courses and amplitudes of the responses were remarkably similar (Guo et al., 2021b). This strongly suggests that the responses we observe do not reflect MF $\rightarrow$ UBC $\rightarrow$ UBC connections. We also showed that the responses were all-or-none: at low intensities no response was evoked, as the intensity of extracellular stimulation was increased a large response was suddenly evoked at a threshold intensity and further increases in intensity did not increase the amplitude of the response (Guo et al., 2021b).

In vivo mossy fiber firing patterns

In vivo mossy fiber recordings were provided by David J. Herzfeld and Stephen G. Lisberger. Recordings were made using either single tungsten microelectrodes (FHC, ~ 1 M Ω) or 16-channel Plexon S-probes in the ventral paraflocculus of a head-fixed rhesus monkey during a smooth pursuit eye movement task. Complete experimental details have been described previously (Herzfeld, Joshua and Lisberger, 2023). Spikes were sorted using Full-binary pursuit (Hall, Herzfeld and Lisberger, 2021), and manually curated by David J. Herzfeld. The mossy fiber traces reproduced in slice in this work were continuous 30 s recordings while the monkey was engaging in smooth pursuit eye movement trials. Briefly, smooth pursuit eye movement trials required the monkey to track a dot as it moves across a monitor at a constant speed. In each trial, the dot would appear stationary in the center of the monitor for 400–800 ms before moving at a constant speed in one of 4 cardinal directions for 650 ms. The dot would then remain stationary in an eccentric position for another 200 ms.

Pharmacology

Sequential drug wash-ins were performed by switching the bath solution to the same solution with addition of the antagonist(s), taking care to keep the flow rate consistent throughout (2 ml / min). Wash-ins were only performed when UBCs showed stable responses for at least 7.5 minutes. Cells with significant shifts in firing independent of mossy fiber stimulation during wash-ins were excluded from analysis. mGluR2/3s were blocked using LY 341495 (1–5 μM , Tocris). AMPARs were blocked using NBQX (5 μM , Abcam). mGluR1s were blocked using JNJ 16259685 (1 μM , Tocris). NMDARs were blocked using R-CPP (5 μM , Abcam).

Quantification and statistical analysis

UBC responses were characterized using instantaneous firing rates in response to 20 stimuli at 100 spk/s bursts averaged over 5–10 median filtered trials recorded before other stimulation paradigms. Baseline firing was estimated from this using the second preceding mossy fiber stimulation. Pause duration was calculated as the time between the end of the stimulation and the moment firing reached 5 spk/s or half the peak firing rate (whichever was smaller). The peak change in firing was calculated by subtracting the baseline firing rate from the peak firing rate after the end of MF stimulation. The half-width was calculated as the time between when the firing rate first exceeded and then decayed to half the peak amplitude (baseline corrected). UBCs were sorted by their half-width when the peak change in firing exceeded 8 spk/s (cell 1-61) or by their pause duration (cell 62-70).

Response parameters for different burst durations were calculated in a similar manner, with separate 20 stimuli at 100 spk/s trials recorded in proximity to other burst durations used for comparison. Additionally, the number of spikes evoked by MF stimulation was calculated by integrating the period from the start of stimulation till 2 times the UBCs half-width after the start of the response. The number of spikes was corrected for baseline firing. For smooth pursuit-like input patterns the number of spikes evoked by step changes in input were calculated in a similar manner but corrected for the average firing rate in the 1 second preceding the step change. To determine time to peak, responses to step changes were fit with a log-normal function, $A \exp\{-[(\ln(t) - \mu)/\sigma]^2\}$, where A is the amplitude, μ is the peak location, and σ is the width in log time. The half-decay of the response to step changes was calculated by finding the first point where the firing rate decreased to half the peak response, using an interpolated trace where responses to individual stimuli had been removed. For experiments with constant input at 1, 2.5, and 5 spk/s peak firing rates and number of spikes were calculated based on the average of all responses excluding the first 5, without correcting for baseline firing. For wash-in experiments all parameters were estimated based on trials 2-2.5 min after the start of the application of the antagonist, when responses to 20 stimuli at 100 spk/s bursts had reached a steady state. Significant differences in **Fig. 3f** and **Fig. 5f** were determined using non-parametric Wilcoxon signed rank tests. Correlation was determined using Spearman's correlation coefficient. All analyses were done using MATLAB R2023b (MathWorks).

Acknowledgements

We thank Stephen G. Lisberger and David J. Herzfeld for providing in vivo mossy fiber recordings and for their comments and input on the project. We thank members of the Regehr lab for comments on the manuscript. The work was supported by the NIH R35NS097284.

Data availability

All data that support the findings in this study have been deposited in Harvard Dataverse:

Huson, Vincent; Regehr, Wade, 2024, "Unipolar brush cell responses to realistic mossy fiber input", <https://doi.org/10.7910/DVN/EBL6C3>

The following previously published data sets were used

Kozareva V, Martin C, Osorno T, Rudolph S, Guo C, Vanderburg C, Nadaf N, Regev A, Regehr W, Macosko E (2021) Single Cell Portal ID SCP795. A transcriptomic atlas of mouse cerebellar cortex reveals novel cell types.

https://singlecell.broadinstitute.org/single_cell/study/SCP795/a-transcriptomic-atlas-of-the-mouse-cerebellum 




Figure 3 - Supplementary Figure 1

NMDA receptors do not significantly contribute to burst responses

a. Instantaneous firing rates of two representative UBCs in response to 100 spk/s bursts comprised of 1-20 stimuli (stimulus indicated above in green). Responses shown for baseline conditions (black), and after addition of an antagonist of NMDA receptors (pink). Cell numbers refer to the index in the summary plot **b**.

b. Heat maps showing the normalized instantaneous firing rates for all UBCs in response to 100 spk/s burst comprised of 1-20 stimuli (columns), for baseline and after addition of an NMDA receptor antagonist (rows). Responses were normalized per cell to the peak firing rate in the baseline response to the 20 stimuli at 100 spk/s burst. Cells sorted by their response to the baseline 20 stimuli at 100 spk/s input, either by the half-width of the increase in firing (cell #1-8) or by pause duration (cell #9-10). Time indicates seconds since start of MF stimulation (indicated by dotted red line). Red arrows indicate representative UBCs shown in **a**.

c. Violin plots of the number of spikes after each burst under baseline conditions (-) and after blocking NMDAR (+), normalized per cell to the number of spikes after 20 stimuli at 100 spk/s under baseline conditions. Markers indicate individual UBCs color coded by the cell index in **b**.




Figure 3 - Supplementary Figure 2

AMPA auxiliary subunits are expressed differentially in the UBC population

- a.** UMAP embedding of normalized gene expression in the UBC population for mGluR1.
- b.** Same as in **a** but showing AMPAR subunits GluA1-4.
- c.** Same as in **a** but showing auxiliary subunits TARP γ -2, γ -7, and γ -8.
- d.** Same as in **a** but showing auxiliary subunit GSG1L.




Figure 4 - Supplementary Figure 1

UBC responses to sustained 1 to 5 spk/s MF input.

a. Instantaneous firing rates of five representative UBCs in response to sustained MF input at three different rates (1, 2.5, and 5 spk/s). Dashed gray lines indicate 0 spk/s; cell numbers refer to the index in the summary plot **c**.

b. As in **a** but for the average response of all individual inputs excluding the first 5.

c. Heat maps showing the normalized instantaneous firing rates for all UBCs in response to sustained input at 1, 2.5, and 5 spk/s. Responses normalized per cell by their peak firing rate. Cells sorted by their response to 20 stimuli at 100 spk/s bursts as displayed in **Fig. 2b**. Time indicates seconds since start of sustained MF input. Red arrows indicate representative UBCs shown in **a,b**.

d. Summary plot showing the peak rate of the average response (as in **b**) for all UBCs color coded to correspond to the cell index in **c**.

e. As in **d** but for the average number of spikes fired.

f. As in **d** but for the ratio between the average steady state firing rate (200 ms after stimuli) and the peak firing rate shown in **d**.




Figure 5 - Supplementary Figure 1

NMDA receptors do not significantly contribute to responses to smooth pursuit-like input

a. Instantaneous firing rates of two representative UBCs in response to 20 stimuli at 100 spk/s burst (*left*), and smooth pursuit-like MF input (*right*) as indicated by the traces at the top (green). Responses shown under baseline conditions and after addition of an antagonist of NMDA receptors. Dashed gray lines indicate 0 spk/s; cell numbers refer to the index in the summary plot **c**.

b. The same as in **a** but on an expanded timescale, displaying only the 60 spk/s step. Dotted red lines indicate onset and offset of the 60 spk/s step.

c. Heat maps showing the normalized instantaneous firing rates for all UBCs in response 20 stimuli at 100 spk/s bursts (*left*) and smooth pursuit-like MF input (*right*), for baseline and after addition of NMDA receptor antagonist (*rows*). Responses normalized per cell to the peak firing rate during baseline conditions separately for burst and smooth pursuit-like MF input. Cells sorted by their response to the baseline 20 stimuli at 100 spk/s input, either by the half-width of the increase in firing (cell #1-8) or by the pause duration (cell #9-10). Dotted red lines indicate the start of the step changes in input rate, red arrows indicate representative UBCs shown in **a,b**.

d. The same as in **c** but on an expanded timescale, displaying only the 60 spk/s step.

e. Violin plots of the number of spikes in the 1 s period during the step and the 3 s period after under baseline conditions (-) and after blocking NMDAR (+), normalized per cell to the number of spikes associated with the step to 60 spk/s under baseline conditions. Markers indicate individual UBCs color coded by the cell index in **c**.




Figure 5 - Supplementary Figure 2

Contribution of different glutamate receptors to UBC responses evoked by sustained 1 to 5 Hz MF stimulation.

- a.** Instantaneous firing rates of two representative UBCs in response to sustained MF input at three different rates (1, 2.5, and 5 spk/s). Responses shown under baseline conditions and after successive addition of antagonists of mGluR2/3, AMPAR, and mGluR1. Dashed gray lines indicate 0 spk/s; cell numbers refer to the index in the summary plot **c**.
- b.** As in **a** but for the average response of all individual inputs excluding the first 5.
- c.** Heat maps showing the normalized instantaneous firing rates for all UBCs in response to sustained input at 1, 2.5, and 5 spk/s (*columns*), for baseline and after addition of glutamate receptor antagonists (*rows*). Responses normalized per cell to the peak firing rate after application of the mGluR2/3 antagonist. Cells sorted by their response to 20 stimuli at 100 spk/s bursts under baseline conditions as in **Fig. 3d** (with exception of **Fig. 3** cell #20 and #24).
- d.** Summary plot showing the number of spikes fired in the average response to 1, 2.5, and 5 spk/s sustained input (as in **b**) for all UBCs color coded to correspond to the cell index in **c**. Successive different markers indicate baseline and the different glutamate antagonists.

References

- Altman J, Bayer SA (1977) **Time of origin and distribution of a new cell type in the rat cerebellar cortex** *Exp Brain Res* **29**:265–274
- Arenz A, Silver RA, Schaefer AT, Margrie TW (2008) **The contribution of single synapses to sensory representation in vivo** *Science* **321**:977–980
- Balmer TS, Borges-Merjane C, Trussell LO (2021) **Incomplete removal of extracellular glutamate controls synaptic transmission and integration at a cerebellar synapse** *Elife* **10**
- Barmack NH, Yakhnitsa V (2008) **Functions of interneurons in mouse cerebellum** *J Neurosci* **28**:1140–1152
- Batchelor AM, Madge DJ, Garthwaite J (1994) **Synaptic activation of metabotropic glutamate receptors in the parallel fibre-Purkinje cell pathway in rat cerebellar slices** *Neuroscience* **63**:911–915
- Borges-Merjane C, Trussell LO (2015) **ON and OFF unipolar brush cells transform multisensory inputs to the auditory system** *Neuron* **85**:1029–1042
- Brasnjo G, Otis TS (2001) **Neuronal glutamate transporters control activation of postsynaptic metabotropic glutamate receptors and influence cerebellar long-term depression** *Neuron* **31**:607–616
- Dino MR, Willard FH, Mugnaini E (1999) **Distribution of unipolar brush cells and other calretinin immunoreactive components in the mammalian cerebellar cortex** *J Neurocytol* **28**:99–123
- Dino MR, Schuerger RJ, Liu Y, Slater NT, Mugnaini E (2000) **Unipolar brush cell: a potential feedforward excitatory interneuron of the cerebellum** *Neuroscience* **98**:625–636
- Eccles JC, Faber DS, Murphy JT, Sabah NH, Taborikova H (1971) **Afferent volleys in limb nerves influencing impulse discharges in cerebellar cortex. I. In mossy fibers and granule cells** *Exp Brain Res* **13**:15–35
- Englund C, Kowalczyk T, Daza RA, Dagan A, Lau C, Rose MF, Hevner RF (2006) **Unipolar brush cells of the cerebellum are produced in the rhombic lip and migrate through developing white matter** *J Neurosci* **26**:9184–9195
- Floris A, Dino M, Jacobowitz DM, Mugnaini E (1994) **The unipolar brush cells of the rat cerebellar cortex and cochlear nucleus are calretinin-positive: a study by light and electron microscopic immunocytochemistry** *Anat Embryol (Berl)* **189**:495–520
- Garwicz M, Jorntell H, Ekerot CF (1998) **Cutaneous receptive fields and topography of mossy fibres and climbing fibres projecting to cat cerebellar C3 zone** *J Physiol* **512**:277–293
- Guo C, Rudolph S, Neuwirth ME, Regehr WG (2021) **Purkinje cell outputs selectively inhibit a subset of unipolar brush cells in the input layer of the cerebellar cortex** *Elife* **10**

- Guo C, Huson V, Macosko EZ, Regehr WG (2021) **Graded heterogeneity of metabotropic signaling underlies a continuum of cell-intrinsic temporal responses in unipolar brush cells** *Nat Commun* **12**
- Hall NJ, Yang Y, Lisberger SG (2018) **Multiple components in direction learning in smooth pursuit eye movements of monkeys** *J Neurophysiol* **120**:2020–2035
- Hall NJ, Herzfeld DJ, Lisberger SG (2021) **Evaluation and resolution of many challenges of neural spike sorting: a new sorter** *J Neurophysiol* **126**:2065–2090
- Hariani HN, Algstam AB, Candler CT, Witteveen IF, Sidhu JK, Balmer TS (2024) **A system of feed-forward cerebellar circuits that extend and diversify sensory signaling** *Elife* **12**
- Hensbroek RA, Ruigrok TJ, van Beugen BJ, Maruta J, Simpson JI (2015) **Visuo-vestibular information processing by unipolar brush cells in the rabbit flocculus** *Cerebellum* **14**:578–583
- Herzfeld DJ, Joshua M, Lisberger SG (2023) **Rate versus synchrony codes for cerebellar control of motor behavior** *Neuron* **111**:2448–2460
- Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2015) **Encoding of action by the Purkinje cells of the cerebellum** *Nature* **526**:439–442
- Hull C, Regehr WG (2012) **Identification of an inhibitory circuit that regulates cerebellar Golgi cell activity** *Neuron* **73**:149–158
- Hull C, Regehr WG (2022) **The Cerebellar Cortex** *Annu Rev Neurosci* **45**:151–175
- Huson V, Newman L, Regehr WG (2023) **A Continuum of Response Properties across the Population of Unipolar Brush Cells in the Dorsal Cochlear Nucleus** *J Neurosci* **43**:6035–6045
- Jackson AC, Nicoll RA (2011) **The expanding social network of ionotropic glutamate receptors: TARPs and other transmembrane auxiliary subunits** *Neuron* **70**:178–199
- Kennedy A, Wayne G, Kaifosh P, Alvina K, Abbott LF, Sawtell NB (2014) **A temporal basis for predicting the sensory consequences of motor commands in an electric fish** *Nat Neurosci* **17**:416–422
- Kim JA, Sekerkova G, Mugnaini E, Martina M (2012) **Electrophysiological, morphological, and topological properties of two histochemically distinct subpopulations of cerebellar unipolar brush cells** *Cerebellum* **11**:1012–1025
- Kinney GA, Overstreet LS, Slater NT (1997) **Prolonged physiological entrapment of glutamate in the synaptic cleft of cerebellar unipolar brush cells** *J Neurophysiol* **78**:1320–1333
- Knoflach F, Kemp JA (1998) **Metabotropic glutamate group II receptors activate a G protein-coupled inwardly rectifying K⁺ current in neurones of the rat cerebellum** *J Physiol* **509**:347–354
- Kozareva V, Martin C, Osorno T, Rudolph S, Guo C, Vanderburg C, Nadaf N, Regev A, Regehr WG, Macosko E (2021) **A transcriptomic atlas of mouse cerebellar cortex comprehensively defines cell types** *Nature* **598**:214–219

- Lisberger SG (2009) **Internal models of eye movement in the floccular complex of the monkey cerebellum** *Neuroscience* **162**:763–776
- Lisberger SG (2021) **The Rules of Cerebellar Learning: Around the Ito Hypothesis** *Neuroscience* **462**:175–190
- Lisberger SG, Fuchs AF (1978) **Role of primate flocculus during rapid behavioral modification of vestibuloocular reflex. II. Mossy fiber firing patterns during horizontal head rotation and eye movement** *J Neurophysiol* **41**:764–777
- Locatelli F, Botta L, Prestori F, Masetto S, D’Angelo E (2013) **Late-onset bursts evoked by mossy fibre bundle stimulation in unipolar brush cells: evidence for the involvement of H- and TRP-currents** *J Physiol* **591**:899–918
- Lu HW, Balmer TS, Romero GE, Trussell LO (2017) **Slow AMPAR Synaptic Transmission Is Determined by Stargazin and Glutamate Transporters** *Neuron* **96**:73–80
- Medina JF, Carey MR, Lisberger SG (2005) **The representation of time for motor learning** *Neuron* **45**:157–167
- Meek J, Yang JY, Han VZ, Bell CC (2008) **Morphological analysis of the mormyrid cerebellum using immunohistochemistry, with emphasis on the unusual neuronal organization of the valvula** *J Comp Neurol* **510**:396–421
- Mugnaini E, Floris A (1994) **The unipolar brush cell: a neglected neuron of the mammalian cerebellar cortex** *J Comp Neurol* **339**:174–180
- Mugnaini E, Sekerkova G, Martina M (2011) **The unipolar brush cell: a remarkable neuron finally receiving deserved attention** *Brain Res Rev* **66**:220–245
- Nunzi MG, Birnstiel S, Bhattacharyya BJ, Slater NT, Mugnaini E (2001) **Unipolar brush cells form a glutamatergic projection system within the mouse cerebellar cortex** *J Comp Neurol* **434**:329–341
- Ohtsuka K, Noda H (1992) **Burst discharges of mossy fibers in the oculomotor vermis of macaque monkeys during saccadic eye movements** *Neurosci Res* **15**:102–114
- Rancz EA, Ishikawa T, Duguid I, Chadderton P, Mahon S, Hausser M (2007) **High-fidelity transmission of sensory information by single cerebellar mossy fibre boutons** *Nature* **450**:1245–1248
- Raymond JL, Lisberger SG (1998) **Neural learning rules for the vestibulo-ocular reflex** *J Neurosci* **18**:9112–9129
- Rossi DJ, Alford S, Mugnaini E, Slater NT (1995) **Properties of transmission at a giant glutamatergic synapse in cerebellum: the mossy fiber-unipolar brush cell synapse** *J Neurophysiol* **74**:24–42
- Rousseau CV, Dugue GP, Dumoulin A, Mugnaini E, Dieudonne S, Diana MA (2012) **Mixed inhibitory synaptic balance correlates with glutamatergic synaptic phenotype in cerebellar unipolar brush cells** *J Neurosci* **32**:4632–4644
- Ruigrok TJ, Hensbroek RA, Simpson JI (2011) **Spontaneous activity signatures of morphologically identified interneurons in the vestibulocerebellum** *J Neurosci* **31**:712–724

- Russo MJ, Yau HJ, Nunzi MG, Mugnaini E, Martina M (2008) **Dynamic metabotropic control of intrinsic firing in cerebellar unipolar brush cells** *J Neurophysiol* **100**:3351–3360
- Schwartz EJ, Rothman JS, Dugue GP, Diana M, Rousseau C, Silver RA, Dieudonne S (2012) **NMDA receptors with incomplete Mg(2)(+) block enable low-frequency transmission through the cerebellar cortex** *J Neurosci* **32**:6878–6893
- Shadmehr R (2017) **Distinct neural circuits for control of movement vs. holding still** *J Neurophysiol* **117**:1431–1460
- Shanks NF *et al.* (2012) **Differences in AMPA and kainate receptor interactomes facilitate identification of AMPA receptor auxiliary subunit GSG1L** *Cell Rep* **1**:590–598
- Simpson JI, Hulscher HC, Sabel-Goedknecht E, Ruigrok TJ (2005) **Between in and out: linking morphology and physiology of cerebellar cortical interneurons** *Prog Brain Res* **148**:329–340
- Soetedjo R, Kojima Y, Fuchs AF (2019) **How cerebellar motor learning keeps saccades accurate** *J Neurophysiol* **121**:2153–2162
- Takacs J, Markova L, Borostyankoi Z, Gorcs TJ, Hamori J (1999) **Metabotropic glutamate receptor type 1a expressing unipolar brush cells in the cerebellar cortex of different species: a comparative quantitative study** *J Neurosci Res* **55**:733–748
- Tempia F, Miniaci MC, Anchisi D, Strata P (1998) **Postsynaptic current mediated by metabotropic glutamate receptors in cerebellar Purkinje cells** *J Neurophysiol* **80**:520–528
- van Dorp S, De Zeeuw CI (2014) **Variable timing of synaptic transmission in cerebellar unipolar brush cells** *Proc Natl Acad Sci U S A* **111**:5403–5408
- van Kan PL, Gibson AR, Houk JC (1993) **Movement-related inputs to intermediate cerebellum of the monkey** *J Neurophysiol* **69**:74–94
- Wadiche JI, Jahr CE (2005) **Patterned expression of Purkinje cell glutamate transporters controls synaptic plasticity** *Nat Neurosci* **8**:1329–1334
- Witter L, De Zeeuw CI (2015) **In vivo differences in inputs and spiking between neurons in lobules VI/VII of neocerebellum and lobule X of archaocerebellum** *Cerebellum* **14**:506–515
- Witter L, Rudolph S, Pressler RT, Lahlaf SI, Regehr WG (2016) **Purkinje Cell Collaterals Enable Output Signals from the Cerebellar Cortex to Feed Back to Purkinje Cells and Interneurons** *Neuron* **91**:312–319
- Yang Y, Lisberger SG (2010) **Learning on multiple timescales in smooth pursuit eye movements** *J Neurophysiol* **104**:2850–2862
- Zampini V, Liu JK, Diana MA, Maldonado PP, Brunel N, Dieudonne S (2016) **Mechanisms and functional roles of glutamatergic synapse diversity in a cerebellar circuit** *Elife* **5**

Author information

Vincent Huson

Department of Neurobiology, Harvard Medical School, Boston, USA

Wade G Regehr

Department of Neurobiology, Harvard Medical School, Boston, USA

ORCID iD: [0000-0002-3485-8094](https://orcid.org/0000-0002-3485-8094)

For correspondence: wade_regehr@hms.harvard.edu

Editors

Reviewing Editor

Megan Carey

Champalimaud Foundation, Lisbon, Portugal

Senior Editor

Sacha Nelson

Brandeis University, Waltham, United States of America

Reviewer #1 (Public review):

In this manuscript, the authors recorded cerebellar unipolar brush cells (UBCs) in acute brain slices. They confirmed that mossy fiber (MF) inputs generate a continuum of UBC responses. Using systematic and physiological trains of MF electrical stimulation, they demonstrated that MF inputs either increased or decreased UBC firing rates (UBC ON vs. OFF) or induced complex, long-lasting modulation of their discharges. The MF influence on UBC firing was directly associated with a specific combination of metabotropic glutamate receptors, mGluR2/3 (inhibitory) and mGluR1 (excitatory). Ultimately, the amount and ratio of these two receptors controlled the time course of the effect, yielding specific temporal transformations such as phase shifts. The experiments are well-executed and properly analyzed.

Strengths:

- (1) A wide range of MF stimulation based on activity patterns observed in vivo was explored, including burst duration and frequency dependency, which could serve as a valuable foundation for explicit modeling of temporal transformations in the granule cell layer.
- (2) The pharmacological blockade of mGluR2/3, mGluR1, AMPA, and NMDA receptors helped identify the specific roles of these glutamate receptors.
- (3) The experiments convincingly demonstrate the key role of mGluR1 receptors in temporal information processing by UBCs.

Weaknesses:

- (1) This study is a follow up of previous work (Guo et al., Nat. Commun., 2021).
- (2) The MF activity used to mimic natural stimulation was previously collected from primates, whereas the recordings were conducted in mice.

Comments on revisions:

The authors included a discussion about inhibition, but I still disagree with their claim that it was not possible to study the MF-UBC connection with inhibition unblocked. This group has already conducted experiments on Golgi cell inhibition in slices.

<https://doi.org/10.7554/eLife.102618.2.sa2>

Reviewer #2 (Public review):

This study addresses the question of how UBCs transform synaptic input patterns into spiking output patterns and how different glutamate receptors contribute to their transformations. The first figure utilizes recorded patterns of mossy fiber firing during eye movements in the flocculus of rhesus monkeys obtained from another laboratory. In the first figure, these patterns are used to stimulate mossy fibers in the mouse cerebellum during extracellular recordings of UBCs in acute mouse brain slices. The remaining experiments stimulate mossy fiber inputs at different rates or burst durations, which is described as 'mossy-fiber like', although they are quite simpler than those recorded in vivo. As expected from previous work, AMPA mediates the fast responses, and mGluR1 and mGluR2/3 mediate the majority of longer-duration and delayed responses. The manuscript is well organized and the discussion contextualizes the results effectively.

Comments on revisions:

The authors have adequately addressed my concerns.

<https://doi.org/10.7554/eLife.102618.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public review):

In this manuscript, the authors recorded cerebellar unipolar brush cells (UBCs) in acute brain slices. They confirmed that mossy fiber (MF) inputs generate a continuum of UBC responses. Using systematic and physiological trains of MF electrical stimulation, they demonstrated that MF inputs either increased or decreased UBC firing rates (UBC ON vs. OFF) or induced complex, long-lasting modulation of their discharges. The MF influence on UBC firing was directly associated with a specific combination of metabotropic glutamate receptors, mGluR2/3 (inhibitory) and mGluR1 (excitatory). Ultimately, the amount and ratio of these two receptors controlled the time course of the effect, yielding specific temporal transformations such as phase shifts.

Overall, the topic is compelling, as it broadens our understanding of temporal processing in the cerebellar cortex. The experiments are well-executed and properly analyzed.

Strengths:

- (1) A wide range of MF stimulation patterns was explored, including burst duration and frequency dependency, which could serve as a valuable foundation for explicit modeling of temporal transformations in the granule cell layer.*
- (2) The pharmacological blockade of mGluR2/3, mGluR1, AMPA, and NMDA receptors helped identify the specific roles of these glutamate receptors.*
- (3) The experiments convincingly demonstrate the key role of mGluR1 receptors in temporal information processing by UBCs.*

Weaknesses:

(1) This study is largely descriptive and represents only a modest incremental advance from the previous work (Guo et al., Nat. Commun., 2021).

We feel that the present study is a major advance. It builds on (Guo et al., Nat. Commun., 2021) in which we examined the effects of bursts of 20 stimuli at 100 spk/s. In that study we found that differential expression of mGluR1 and mGluR2 let to a continuum of temporal responses in UBCs, but AMPARs make a minimal contribution for such bursts. It was not known how UBCs transform realistic mossy fiber input patterns. Here we provide a comprehensive evaluation of a wide range of input patterns that include a range of bursts comprised of 1-20 stimuli, sustained stimulation with stimulation of 1 spk/s to 60 spk/s. This more thorough assessment of UBC transformations combined with a pharmacological assessment of the contributions of different glutamate receptor subtypes provided many new insights:

- We found that UBC transformations are comprised of two different components: a slow temporally filtered component controlled by an interplay of mGluR1 and mGluR2, and a second component mediated by AMPARs that can convey spike timing information. NMDARs do not make a major contribution to UBC firing. The finding that UBCs simultaneously convey two types of signals, a slow filtered response and responses to single stimuli, has important implications for the computational potential of UBCs and fundamentally changes the way we think about UBCs.
- We found that with regard to the slow filtered component mediated by mGluR1 and mGluR2, we could extend the concept of a continuum of responses evoked by 20 stimuli at 100 spk/s (Guo et al., Nat. Commun., 2021) to a wide range of stimuli. It was not a given that this would be the case.
- The contributions of AMPARs was surprising. Even though snRNAseq data did not reveal a gradient of AMPAR expression across the population of UBCs (Guo et al., Nat. Commun., 2021), we found that there was a gradient of AMPA-mediated responses, and that the AMPA component was also most prominent in cells with a large mGluR1 component. Our finding that AMPAR accessory proteins exhibit a gradient across the population, which could account for the gradient of AMPAR responses, will prompt additional studies to test their involvement.

(2) The MF activity used to mimic natural stimulation was previously collected in primates, while the recordings were conducted in mice.

Our first task was to determine the firing properties of mossy fibers under physiological conditions in UBC rich cerebellar regions. Previous studies have estimated this in anesthetized mice using whole cell granule cell recordings (Arenz et al., 2008; Witter & De Zeeuw 2015). However, for assessing firing patterns during awake behavior, we felt that the most comprehensive data set available in a UBC rich cerebellar region was for mossy fibers involved in smooth pursuit in monkeys (David J. Herzfeld and Stephen G. Lisberger). This revealed the general features of mossy fiber firing that helped us design stimulus patterns to thoroughly probe the properties of MF to UBC transformations. The firing patterns are designed to investigate the transformations for a wide range of activity patterns and have important general implications for UBC transformations that are likely applicable to UBCs in different species that are activated in different ways.

(3) Inhibition was blocked throughout the study, reducing its physiological relevance.

The reviewer correctly brings up the very important issue of inhibition in shaping UBC responses. It is well established that UBCs are inhibited by Golgi cells (Rousseau et al., 2012), and we recently showed that some UBCs are also inhibited by PCs (Guo et al., eLife, 2021).

This will undoubtedly influence the firing of UBCs in vivo. We considered examining this issue, but felt that brain slice experiments are not well suited to this. In contrast to MF inputs that can be activated with a realistic activity pattern, it is exceedingly difficult to know how Golgi cells and Purkinje cells are activated under physiological conditions. Each UBC is activated by a single mossy fiber, but inhibition is provided by Golgi cells that are activated by many mossy fibers and granule cells, and PCs that are controlled by many granule cells and many other PCs. In addition, we found that many Golgi cells do not survive very well in slices, and the axons of many PCs are severed in brain slice. Although limitations of the slice preparation prevent us from determining the role of inhibition in shaping UBC responses, we have added a section to the discussion in which we address the important issue of inhibition and UBC responses.

Reviewer #2 (Public review):

This study addresses the question of how UBCs transform synaptic input patterns into spiking output patterns and how different glutamate receptors contribute to their transformations. The first figure utilizes recorded patterns of mossy fiber firing during eye movements in the flocculus of rhesus monkeys obtained from another laboratory. In the first figure, these patterns are used to stimulate mossy fibers in the mouse cerebellum during extracellular recordings of UBCs in acute mouse brain slices. The remaining experiments stimulate mossy fiber inputs at different rates or burst durations, which is described as 'mossy-fiber like', although they are quite simpler than those recorded in vivo. As expected from previous work, AMPA mediates the fast responses, and mGluR1 and mGluR2/3 mediate the majority of longer-duration and delayed responses. The manuscript is well organized and the discussion contextualizes the results effectively.

The authors use extracellular recordings because the washout of intracellular molecules necessary for metabotropic signaling may occur during whole-cell recordings. These cell-attached recordings do not allow one to confirm that electrical stimulation produces a postsynaptic current on every stimulus. Moreover, it is not clear that the synaptic input is monosynaptic, as UBCs synapse on one another. This leaves open the possibility that delays in firing could be due to disynaptic stimulation. Additionally, the result that AMPA-mediated responses were surprisingly small in many UBCs, despite apparent mRNA expression, suggests the possibility that spillover from other nearby synapses activated the higher affinity extrasynaptic mGluRs and that that main mossy fiber input to the UBC was not being stimulated. For these reasons, some whole-cell recordings (or perforated patch) would show that when stimulation is confirmed to be monosynaptic and reliable it can produce the same range of spiking responses seen extracellularly and that AMPA receptor-mediated currents are indeed small or absent in some UBCs.

We appreciate the reviewer's concerns regarding the reliability of mossy fiber activation, the possibility of glutamate spillover from other synapses, and the possibility of disynaptic activation involving stimulation of MFàUBCàUBC connections. We examined these issues in a previous study (Guo et al., Nat. Commun., 2021). We did on-cell recordings and followed that up with whole cell voltage clamp recordings from the same cell (Guo et al., Nat. Commun., 2021, Fig. 5), and there was good agreement with the amplitude and timing of spiking and the time course and amplitudes of the synaptic currents. We also compared responses evoked by focal glutamate uncaging over the brush and MF stimulation (Guo et al., Nat. Commun., 2021, Fig. 4). We found that the time courses and amplitudes of the responses were remarkably similar. This strongly suggests that the responses we observe do not reflect disynaptic activation (MFàUBCàUBC connections). We also showed that the responses were all-or-none: at low intensities no response was evoked, as the intensity of extracellular stimulation was increased a large response was suddenly evoked at a threshold intensity and further increases in intensity did not increase the amplitude of the response (Guo et al., Nat.

Commun., 2021, Extended data Fig. 1). We can be well above threshold and still excite the same response, and as a result we do not see stereotyped indications of an inability to stimulate during prolonged high frequency activation. We recognize the importance of these issues, so we have added a section dealing explicitly with these issues (pp. 15-16).

A discussion of whether the tested glutamate receptors affected the spontaneous firing rates of these cells would be informative as standing currents have been reported in UBCs. It is unclear whether the firing rate was normalized for each stimulation, each drug application, or each cell. It would also be informative to report whether UBCs characterized as responding with Fast, Mid-range, Slow, and OFF responses have different spontaneous firing rates or spontaneous firing patterns (regular vs irregular).

The spontaneous firing of UBCs is indeed an interesting issue that is deserving of further investigation. It is not currently known how spontaneous firing at rest is regulated in UBCs, however, in previous work we have shown that there is great diversity in the rates across the population of UBCs in the dorsal cochlear nucleus (Huson & Regehr, JNeurosci, 2023, Fig. 4). Unfortunately, during the kind of sustained high-frequency stimulation protocols (as used in this study) spontaneous firing rates tend to increase. This is likely an effect of residual receptor activation. As such, our current dataset is not suitable to performing in depth analysis of the effects of the different glutamate receptors on spontaneous firing rates. As this study aims to explore UBC responses to MF inputs we feel that specific experiments to address the issue of spontaneous firing rates are outside of the scope.

As the reviewers points out there are indeed different ways the firing rates can be normalized for display in the heatmaps, and different normalizations have been used in different figures. We have made sure that the method for normalization is clearly indicated in the figure legends for each of the heatmaps on display, specifying the protocol and drug application used for normalization.

Figure 1 shows examples of how Fast, Mid-range, Slow, and OFF UBCs respond to in vivo MF firing patterns, but lacks a summary of how the input is transformed across a population of UBCs. In panel d, it looks as if the phase of firing becomes more delayed across the examples from Fast to OFF UBCs. Quantifying this input/output relationship more thoroughly would strengthen these results.

The UBC responses to in vivo MF firing patterns are intriguing and we agree that there appears to be increasing delays for slower UBCs visible in Figure 1. However, we feel that the true in vivo MF firing patterns are too complex and irregular for rigorous interpretation. Therefore, we only tested simplified burst and smooth pursuit-like input patterns on the full population of UBCs. Here we indeed do see increasingly delayed responses as UBCs get slower (Fig. 4).

Inhibition was pharmacologically blocked in these studies. Golgi cells and other inhibitory interneurons likely contribute to how UBCs transform input signals. Speculation of how GABAergic and glycinergic synaptic inhibition may contribute additional context to help readers understand how a circuit with intact inhibition may behave.

As indicated in our response to reviewer 1, we have added a section discussing the very important issue of inhibition and UBC responses in vivo.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

(1) Including recordings without inhibition blocked would strengthen the study and provide a more comprehensive view of the transformations made by UBCs at the input stage of the cerebellar cortex.

See response to public comments.

(2) The authors claim that a continuum of temporal responses was observed in UBCs, but they also distinguish between fast, mid-range, slow, and OFF UBCs. While some UBCs fire spontaneously, others are activated by MF inputs. A more thorough classification effort would clarify the various response profiles observed under specific MF stimulation regimes. Have the authors considered using machine learning algorithms to aid in classification?

We fundamentally feel that these response properties do not conform to rigid categories. In our previous work we have shown that UBC population constitutes a continuum in terms of gene expression, and in terms of spontaneous and evoked firing patterns. While in order to answer some questions empirically it may still be useful to apply advanced algorithms to enforce separate groups to be compared, in this work we aimed to present the full range of UBC responses without introducing any additional biases that such methods would produce.

(3) A robust classification could assist in quantifying the temporal shifts observed during smooth pursuit-like MF stimulation, a critical outcome of the study.

As stated above, we prefer to present an unbiased overview of the continuous nature of the UBC population, as we believe that this is fundamentally the most accurate representation. While it is true that this prevents us from providing a quantification in the different temporal shifts, we believe that the range of shifts across the population is sufficiently large and continuously varying to be convincing (see Figure 4d).

(4) In Figure 5, contrary to what is described on page 10, Cells 10 and 11 (OFF UBCs) appear to behave differently, as mGluR1 does not seem to affect their firing rates. A specific case should be made for OFF UBCs.

Indeed, cells 10 and 11 do not show clear increases in firing and are not strongly affected by blocking of mGluR1. However, as discussed above and explored in our previous work, we feel that the range of UBC increases in firing is best described as a continuum, including the extreme where increases in firing are no longer clearly observable. As the aim in this work is to describe this continuum of responses for physiologically relevant inputs, we do not feel there is a benefit to creating a specific case for OFF UBCs here. It should be pointed out that the number of "pure" OFF UBCs completely lacking an mGluR1 component is very small.

(5) A summary diagram should be added at the end of the manuscript to highlight the key temporal features observed in this study.

This is a great suggestion and we have prepared such a summary diagram (Figure 6).

Reviewer #2 (Recommendations for the authors):

(1) Page 3- "Assed" should be "assessed"

(2) Page 19- "by integrating" is repeated twice

(3) It was not noted whether the data would be made available. It could be useful for those interested in implementing UBCs in models of the cerebellar cortex.

We agree that this data set is invaluable to those interested in implementing UBCs in models of the cerebellar cortex. We will make the dataset available as described in the text.

<https://doi.org/10.7554/eLife.102618.2.sa0>