

Microglia replacement by ER-Hoxb8 conditionally immortalized macrophages provides insight into Aicardi-Goutières Syndrome neuropathology

Reviewed Preprint

v2 • June 30, 2025

Revised by authors

Reviewed Preprint

v1 • January 6, 2025

Kelsey M Nemec, Genevieve Uy, V Sai Chaluvadi, Freddy S Purnell, Bilal Elfayoumi, Leila Byerly, Micaela L O'Reilly, Carleigh A O'Brien, William H Aisenberg, Sonia I Lombroso, Xinfeng Guo, Niklas Blank, Chet Huan Oon, Fazeela Yaqoob, Brian Temsamrit, Priyanka Rawat, Christoph A Thaiss, Will Bailis, Adam P Williamson, Qingde Wang, Mariko L Bennett, F Chris Bennett ✉

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States • Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States • Division of Neurology, Children's Hospital of Philadelphia, Philadelphia, United States • Department of Biology, School of Arts and Sciences, University of Pennsylvania, Philadelphia, United States • Epigenetics Institute, University of Pennsylvania, Perelman School of Medicine, Philadelphia, United States • Department of Biology, Bryn Mawr College, Bryn Mawr, United States • Department of Systems Pharmacology and Translational Therapeutics, University of Pennsylvania, Philadelphia, United States • Department of Surgery, University of Pittsburgh School of Medicine, Pittsburgh, United States • Department of Microbiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States • Faculty of Biology, University of Freiburg, Freiburg, Germany • Department of Pathology and Laboratory Medicine, Children's Hospital of Philadelphia, Philadelphia, United States • Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States • Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This revised study describes an **important** new model for in vivo manipulation of microglia, exploring how mutations in the *Adar1* gene within microglia contribute to Aicardi-Goutières Syndrome. The methodology is validated with **exceptional** data, supporting the authors' conclusions. The paper underscores both the advantages and limitations of using transplanted cells as a surrogate for microglia, making it a resource that is of value for biologists studying macrophages and microglia.

<https://doi.org/10.7554/eLife.102900.2.sa3>

Abstract

Microglia, the brain's resident macrophages, can be reconstituted by surrogate cells - a process termed "microglia replacement." To expand the microglia replacement toolkit, we here introduce estrogen-regulated (ER) homeobox B8 (*Hoxb8*) conditionally immortalized macrophages, a cell model for generation of immune cells from murine bone marrow, as a versatile model for microglia replacement. We find that ER-*Hoxb8* macrophages are highly comparable to primary bone marrow-derived (BMD) macrophages in vitro, and, when transplanted into a microglia-free brain, engraft the parenchyma and differentiate into microglia-like cells. Furthermore, ER-*Hoxb8* progenitors are readily transducible by virus and easily stored as stable, genetically manipulated cell lines. As a demonstration of this system's power for studying the effects of disease mutations on microglia in vivo, we created stable, *Adar1*-mutated ER-*Hoxb8* lines using CRISPR-Cas9 to study the intrinsic contribution of macrophages to Aicardi-Goutières Syndrome (AGS), an inherited interferonopathy that primarily affects the brain and immune system. We find that *Adar1* knockout elicited interferon secretion and impaired macrophage production in vitro, while preventing brain macrophage engraftment in vivo - phenotypes that can be rescued with concurrent mutation of *Ifih1* (MDA5) in vitro, but not in vivo. Lastly, we extended these findings by generating ER-*Hoxb8* progenitors from mice harboring a patient-specific *Adar1* mutation (D1113H). We demonstrated the ability of microglia-specific D1113H mutation to drive interferon production in vivo, suggesting microglia drive AGS neuropathology. In sum, we introduce the ER-*Hoxb8* approach to model microglia replacement and use it to clarify macrophage contributions to AGS.

Introduction

Microglia, the parenchymal tissue-resident macrophages of the brain and spinal cord, play critical roles in development, homeostasis, injury, and disease (Salter and Stevens, 2017; Li and Barres, 2018). When depleted of endogenous microglia, the brain parenchyma can be reconstituted by surrogate macrophages - a process termed "microglia replacement." Microglia replacement has the potential for precise and personalized delivery of therapeutic payloads or correction of dysfunction. Primary microglia are challenging to manipulate - they rapidly lose transcriptional identity ex vivo, resist viral manipulation, are minimally proliferative, and are highly sensitive to serum (Balcaitis et al., 2005; Jiang et al., 2012; Masuda et al., 2013; Su et al., 2016; Bohlen et al., 2017). These challenges limit the study of microglia replacement, underscoring the need for new, transplantable cell models. Common microglia surrogate cells include myeloid cells from the blood or bone marrow and induced pluripotent stem cell (iPSC)-derived microglia (iMG; Muffat et al., 2016; Pandya et al., 2017; Haenseler et al., 2017; Douvaras et al., 2017; Abud et al., 2017; Takata et al., 2020). Although capable of reconstituting the microglial niche (Priller et al., 2001; Bohlen et al., 2017; Bennett et al., 2018; Hasselmann et al., 2019), primary cells and iPSCs each have limitations that motivated us to consider new microglia replacement tools.

To create an alternative microglia surrogate, we turned to the estrogen-regulated (ER) homeobox B8 (*Hoxb8*) system, a method for generating and gene-editing unlimited quantities of macrophages from primary murine bone marrow (Wang et al., 2006). When overexpressed, *Hoxb8* promotes the expansion of hematopoietic progenitor cells while preventing their differentiation. When transduced with ER-*Hoxb8* virus, myeloid progenitor cells from murine bone marrow become immortalized and indefinitely expandable when in the presence of exogenous estrogen, but differentiate upon estrogen removal. The ER-*Hoxb8* approach has been used to generate so-called

“conditionally immortalized” progenitors with lymphoid and myeloid potential (Wang et al., 2006 [↗](#); Rosas et al., 2011 [↗](#); Redecke et al., 2013 [↗](#); Gurzeler et al., 2013 [↗](#); Zach et al., 2015 [↗](#); Fites et al., 2018 [↗](#); Ma et al., 2020 [↗](#); Lail et al., 2022), including macrophages.

We recently showed that ER-Hoxb8 cells can reconstitute the microglial niche (Chadarevian, Lombroso, et al., 2023 [↗](#)), supporting their potential as a new cell model for microglia replacement. Here, we aimed to deeply characterize the identity, function, and application of ER-Hoxb8s as microglia surrogates to test hypotheses about microglia-specific gene functions in health and disease.

Results

Comparison of ER-Hoxb8 to bone marrow-derived macrophages in vitro

To directly compare primary bone marrow-derived (BMD) and ER-Hoxb8 cells, we created three biologically independent lines of conditionally immortalized macrophage progenitors as described previously (Wang et al., 2006 [↗](#)). ER-Hoxb8 progenitor cells were plated, differentiated for seven days in CSF1, and compared to BMD macrophages (Stanley and Heard 1977 [↗](#); Murray et al., 2014 [↗](#); **Figure 1A** [↗](#)). We found that ER-Hoxb8 macrophages were highly similar to BMD macrophages morphologically (**Figure 1B** [↗](#)) and by CD11B/CD45 expression (**Figure 1C** [↗](#), **Supplemental Figures 1A/B** [↗](#)). To validate ER-Hoxb8 macrophage function, we tested their phagocytic capacity. In agreement with prior literature (Elhag et al., 2021 [↗](#)), ER-Hoxb8 macrophages more efficiently engulfed beads coated with 10% phosphatidylserine (PS) to mimic apoptotic cells, compared to beads coated with phosphatidylcholine (PC) alone (**Supplemental Figure 1C/D** [↗](#)).

To expand upon previous studies (Redecke et al., 2013 [↗](#); Roberts et al., 2019 [↗](#); Accarias et al., 2020 [↗](#)), we performed RNA sequencing of both cell types before and after seven days of differentiation. We generated high-quality transcriptomes (Phred Score > 35, n = 3 biological replicates) for BM progenitors, BM monocytes, seven-day differentiated BMD macrophages, ER-Hoxb8 progenitors, four-day differentiated ER-Hoxb8 cells, and seven-day differentiated ER-Hoxb8 macrophages (top 100 expressed genes in Supplemental Table 1). In agreement with previous literature (Wang et al., 2006 [↗](#)), ER-Hoxb8 progenitors express canonical macrophage genes (**Supplemental Figure 1E** [↗](#)) and, interestingly, ER-Hoxb8 cells at all time points were more similar to BMD macrophages than to BM progenitors or monocytes (**Supplemental Figure 1F** [↗](#)).

Transcriptionally, seven-day differentiated ER-Hoxb8 macrophages were highly similar to BMD macrophages as demonstrated by expression of macrophage-specific genes and lack of expression of common progenitor, lymphocyte, neutrophil, and natural killer cell-specific genes (**Figure 1D** [↗](#)). Linear regression analysis between the two cell types (**Figure 1E** [↗](#)), revealed a strong correlation ($R^2 = 0.96$) between expression levels of all 29,625 expressed genes. We identified 85 genes as differentially expressed between ER-Hoxb8 and BMD macrophages (FDR < 0.05, Log2 Fold Change ≥ 2 , counts per million (CPM) > 1; **Figure 1F** [↗](#)). Of these, 74 were upregulated in BMD macrophages, 11 were upregulated in ER-Hoxb8 macrophages, and no statistically significant Gene Ontology (GO) terms were found for either group (top ten DEGs by Log2FC in **Supplemental Figure 1G** [↗](#), all DEGs in Supplemental Table 2). These data demonstrate that ER-Hoxb8 macrophages are highly similar to primary BMD macrophages in vitro.




Figure 1

Comparison of ER-Hoxb8 to BMD macrophages in vitro.

(A) Schematic for creation of bone marrow-derived (BMD) and ER-Hoxb8 cells **(B)** Brightfield images of BMD and ER-Hoxb8 macrophages plated in the presence of 30 ng/mL mouse CSF1 and differentiated for seven days (scale bar = 100um) **(C)** Dot plot representing CD45/CD11B levels (pre-gated on live, singlet, leukocyte) by flow cytometry **(D)** Heatmap showing Log2 CPM of canonical macrophage (top) and non-macrophage (bottom) immune cell genes **(E)** Whole transcriptome comparison between BMD and ER-Hoxb8 macrophages, depicting best fit line and coefficient of determination (one dot = one gene) **(F)** Volcano plots comparing all genes or those with CPM > 1 (Log2FC >= 2, FDR < 0.05); blue = upregulated in ER-Hoxb8 macrophages, red = upregulated in BMD macrophages Panel A created with [BioRender.com](https://www.biorender.com)

ER-Hoxb8 cells engraft the brain parenchyma after intracranial transplantation and attain a microglia-like identity

To extend the utility of the ER-Hoxb8 model, we next studied their engraftment and identity following intracranial transplantation into *Csf1r*^{-/-} hosts, which lack microglia and readily permit donor cell engraftment (Bennett et al., 2018 [DOI](#); [Figure 2A](#) [DOI](#)). Using BMD cells as a primary cell control, early postnatal transplantation at days zero to five (P0-5) led to robust and comparable parenchymal engraftment by 16 days ([Figure 2B-D](#) [DOI](#)). ER-Hoxb8 macrophages engrafted at equivalent densities (171.7 cells/mm² +/- 8.0) as endogenous, wild-type (WT) microglia (160.0 cells/mm² +/- 5.5), while BMD macrophages engrafted at higher densities (264.6 cells/mm² +/- 6.9; [Figure 2E](#) [DOI](#)).

BMD macrophages become “microglia-like” after engraftment in the *Csf1r*^{-/-} brain (Bennett et al., 2018 [DOI](#)). Like BMD cells, ER-Hoxb8 cells lack expression of the microglia signature protein TMEM119 in vitro ([Supplemental Figure 2A](#) [DOI](#)). After brain engraftment, we found comparably high levels of TMEM119 in BMD and ER-Hoxb8 macrophages by flow cytometry, though both had lower TMEM119 expression than endogenous microglia, consistent with prior studies (Bennett et al., 2018 [DOI](#); Cronk et al., 2018 [DOI](#); Shemer et al., 2018 [DOI](#); [Figure 3A](#) [DOI](#)). As with WT microglia and BMD macrophages, ER-Hoxb8 macrophages likewise express P2RY12 after brain engraftment ([Supplemental Figure 2B](#) [DOI](#)). To more comprehensively assess microglial identity, we generated high-quality bulk RNA sequencing transcriptomes (Phred Score > 35; n = 6-9 biological replicates, [Supplemental Figure 2C](#) [DOI](#)) from isolated brain-engrafted BMD macrophages, ER-Hoxb8 macrophages, and endogenous microglia from age-matched WT hosts. Principal component analysis (PCA) and unsupervised hierarchical clustering revealed that brain residence induces both BMD and ER-Hoxb8 macrophages to become more similar to endogenous microglia ([Figure 3B](#) [DOI](#), [Supplemental Figure 2D](#) [DOI](#)). We saw no evidence for batch effects between harvest days, cell sorter used, or host mouse sex ([Supplemental Figure 2E](#) [DOI](#)). BMD and ER-Hoxb8 microglia-like cells (MLCs) were transcriptionally similar, by both PCA and linear regression analysis ([Figures 3B/C/D](#) [DOI](#)). ER-Hoxb8 MLCs upregulated expression of canonical microglia signature genes akin to BMD controls (*P2ry12*, *Tmem119*, *Hexb*, *Fcrls*, *Gpr34*, *Olfml3*, *P2ry13*, *Sparc*), including genes highly downregulated in vitro ([Figure 3E](#) [DOI](#)). Like BMD macrophages, ER-Hoxb8 macrophages lack *Sall1* expression in vivo compared to microglia ([Figure 3E](#) [DOI](#)). Lastly, to more completely assess the degree that ER-Hoxb8 cells become “microglia-like” after brain engraftment, we compared the top up and downregulated genes from our datasets to those of brain-engrafted yolk sac (YS) progenitors using data from Bennett et al., 2018 [DOI](#) ([Supplemental Figure 2F](#) [DOI](#), [Supplemental Table 3](#)). We find overlap between groups, which likewise provides a better understanding of genes that are robustly regulated by brain engraftment. Taken together, these data show that, similar to BMD macrophages, ER-Hoxb8 macrophages become “microglia-like” with exposure to signals in the brain microenvironment.

Despite their similarities, BMD MLCs and ER-Hoxb8 MLCs have 650 differentially expressed genes (FDR < 0.05, Log2FC >= 2, and CPM > 1 in at least six of the samples (the “n” of each group); [Supplemental Table 4](#)). We performed statistical overrepresentation tests to identify relevant GO terms ([Supplemental Table 5](#)). Of particular interest were terms relating to biological processes as they concerned immune activity and reactivity. We discovered that the 34 overrepresented genes for these GO terms ([Supplemental Table 6](#)), were unique to ER-Hoxb8 MLCs, suggesting that the brain environment differentially affects engrafted BMD and ER-Hoxb8 cells, and that ER-Hoxb8 MLCs may be more primed to activation.

To gauge how significantly these transcriptomic increases in inflammation-associated genes impact the brain environment, we stained for GFAP, a marker of reactive astrogliosis and a sensitive marker for central nervous system (CNS) perturbations. We found no increase in GFAP




Figure 2

Engraftment potential of ER-Hoxb8 compared to BMD macrophages after intracranial transplantation in *Csf1r*^{-/-} hosts.

(A) Schematic for in vivo *Csf1r*^{-/-} transplant experiments **(B)** Rendered tile stitches of *Csf1r*^{-/-} brains after intracranial injection of GFP+ bone marrow (left) or ER-Hoxb8 (right) progenitor cells **(C)** Percent of total brain area tiled by donor cells; n = five to seven biological replicates per group; each dot = one biological replicate (average area across three matched sagittal sections) **(D)** Immunostaining of cortical brain region 12-16 days post-intracranial injection (red = IBA1, green = endogenous GFP, blue = DAPI; scale bar = 100um; inset scale bar = 5um) **(E)** Cortical density calculations (cells per mm²) between groups; n = five to seven biological replicates per group; each dot = one biological replicate (average density across three regions of interest across three matched sagittal sections). All p-values calculated via one-way ANOVA with multiple comparisons; ns = not significant or p >= 0.05, ****p < 0.0001 Panel A created with [BioRender.com](https://www.biorender.com)




Figure 3

ER-Hoxb8 macrophages become microglia-like cells (MLCs) after engraftment in the *Csf1r*^{-/-} brain.

(A) Histogram of TMEM119 surface staining by flow cytometry (pre-gated on live, singlet, leukocyte, CD45⁺/CD11b⁺) for brain-engrafted cells 14 days post-intracerebral transplantation; Mg = WT Microglia, BM = BMD MLCs, Hox = ER-Hoxb8 MLCs **(B)** PCA plot comparing in vitro macrophages from [Figure 1](#) with in vivo macrophages; Mg = WT Microglia, BM = BMD MLCs, Hox = ER-Hoxb8 MLCs **(C)** Whole transcriptome comparison between WT microglia, BMD, and ER-Hoxb8 macrophages in vivo, depicting best fit line and the coefficient of determination **(D)** Comparison of microglia signature genes (Cronk, et al. *JEM* (2018)) depicting best fit line and coefficient of determination **(E)** In vitro and in vivo Log2 CPM gene expression of ten canonical microglia/myeloid genes for bone marrow (left) and ER-Hoxb8s (right) as compared to unmanipulated in vivo microglia

expression across brain regions between any groups, including in brains engrafted with ER-Hoxb8 cells compared to BMD cells (**Supplemental Figure 2G** [↗](#)).

***Adar1* mutation reduces macrophage number and induces interferon responses, effects mitigated by JAK inhibition or *Ifih1* mutation**

Having established ER-Hoxb8 macrophages as microglia-like cells, we explored their potential for study of microglia in health and disease. Building on prior work ([Gran et al., 2018](#) [↗](#); [Roberts et al., 2019](#) [↗](#); [Accarias et al., 2021](#); [Bromberger et al., 2022](#) [↗](#); [Shen et al., 2022](#) [↗](#); [Xu et al., 2022](#) [↗](#); [Möller et al., 2023](#) [↗](#)), we used CRISPR-Cas9 to knockout (KO) *Tlr4* in Cas9^{+/−} ER-Hoxb8 progenitors, using fluorescence-activated cell sorting (FACS) to establish lines of guide-transduced ER-Hoxb8 progenitors (**Supplemental Figure 3A** [↗](#)). We confirmed editing by TIDE analysis (**Supplemental Figure 3B** [↗](#), [Brinkman et al., 2014](#) [↗](#)) and protein loss by flow cytometry (**Supplemental Figures 3C/D** [↗](#)). We then verified phenotypic knockout by treating unmodified, non-targeting control (NTC) guide-transduced, and *Tlr4* guide-transduced ER-Hoxb8 macrophages with lipopolysaccharide (LPS, a *Tlr4* agonist) or R848 (a *Tlr7/8* agonist). As expected, *Tlr4* KO blunted TNF-α production in response to *Tlr4* but not *Tlr7/8* agonism (**Supplemental Figure 3E** [↗](#)).

Having validated effective gene targeting, we next leveraged strengths of the ER-Hoxb8 system to study the contribution of microglia to a monogenic CNS disease. Aicardi-Goutières Syndrome (AGS) is a rare, brain-predominant genetic interferonopathy with nine known causal genes ([Gavazzi et al., 2024](#) [↗](#)). As many cells respond to interferon, it is unclear which cells drive AGS brain pathology, particularly in cases where global gene knockout is embryonic lethal. We therefore tested the hypothesis that microglia, which produce and react strongly to interferons ([Zheng et al., 2015](#) [↗](#); [Aw et al., 2020](#); [Escoubas et al., 2024](#) [↗](#)), are a main contributor to AGS pathology. We first confirmed that the seven coding AGS-causal genes are expressed by ER-Hoxb8 cells at similar levels to those of BMD cells and true microglia (**Supplemental Figure 4A** [↗](#)). We were primarily interested in *Adar1*, an RNA editing enzyme expressed by all brain cells, including microglia ([Zhang, Chen, Sloan, et al., 2014](#); [Bennett et al., 2016](#) [↗](#)). As ADAR1 loss disrupts hematopoiesis ([Wang et al., 2004](#); [Hartner 2004](#)), hematopoietic progenitor differentiation ([XuFeng et al., 2009](#) [↗](#)), and is embryonically lethal in mice ([Wang et al., 2004](#); [Hartner et al., 2004](#) [↗](#)), the impact of ADAR1 loss in macrophages is unknown. We leveraged the ER-Hoxb8 model to study *Adar1* loss in macrophages by creating independent *Adar1* KO lines using two distinct sgRNAs targeting different exons (**Figure 4A** [↗](#), **Supplemental Figure 4B** [↗](#)).

We first explored differences during differentiation of *Adar1* KO ER-Hoxb8 cells in vitro. Despite normal numbers at three days, there were significantly fewer *Adar1* KO cells during the later stages of differentiation as compared to NTC cells (**Figures 4B/C** [↗](#), **Supplemental Figures 4C/D** [↗](#)). This suggests that either *Adar1* KO inhibits progenitor differentiation or macrophage proliferation and survival. We next performed RNA sequencing of *Adar1* KO progenitors and differentiated macrophages. *Adar1* KO progenitors were highly similar to NTC control progenitors (**Figure 4D/E/F** [↗](#); all DEGs in Supplemental Table 7), and *Adar1* KO macrophages expressed comparable levels of macrophage-identity genes to NTC (**Figure 4G** [↗](#)), suggesting that *Adar1* KO ER-Hoxb8 progenitors can generate macrophages. At the macrophage stage, however, we found 547 DEGs (**Figure 4D/E/F** [↗](#); Supplemental Table 8), remarkable for interferon-stimulated gene (ISG) upregulation and GO term enrichment for responses to virus, stress, cytokine stimulation, and innate immunity (**Figure 4E/F** [↗](#); **Supplemental Figure 4E** [↗](#), Supplemental Table 9).

To test whether *Adar1* KO-mediated interferon production underlies this macrophage phenotype, we used the Janus kinase (JAK) inhibitor Baricitinib to inhibit interferon signaling. Baricitinib, a treatment for AGS ([Vanderver et al., 2020](#) [↗](#); [Han et al., 2022](#) [↗](#); [Kanazawa et al., 2023](#) [↗](#)), caused a dose-dependent reduction in ISG expression (**Figure 4H** [↗](#)) and normalized cell counts during macrophage differentiation (**Figure 4I** [↗](#), **Supplemental Figure 4F** [↗](#)). We validated increased




Figure 4

***Adar1* mutation prevents macrophage-lineage cell expansion and causes interferon induction, rescued by JAKi or *Ifih1* mutation.**

(A) Schematic of ADAR1 locus, depicting exons, alternative start sites for p150 and p110 isoforms, and sgRNA targets (B) ER-Hoxb8 cell counts over differentiation time course (C) Immunostaining of in vitro, eight-day differentiated macrophages comparing control (NTC) and *Adar1* guide-transduced macrophages (red = CD11B, blue = DAPI; scale bar = 100um) (D) PCA plot of progenitors and macrophages in vitro (E) Volcano plots showing differentially expressed genes between *Adar1* KO and NTC progenitors and macrophages (CPM > 1, Log2FC >= 2, FDR < 0.05) (F) Heatmap showing the Log2FC (*Adar1* KO values over NTC values) for relevant interferon-stimulated genes (G) Heatmap showing Log2 CPM of canonical macrophage (top) and non-macrophage (bottom) immune cell genes (H) Heatmap showing Log2FC (*Adar1* KO over NTC expression) for interferon-stimulated genes in macrophages treated with baricitinib (I) ER-Hoxb8 cell counts over differentiation time course, comparing the effect of baricitinib on *Adar1* KO and NTC lines (dosages = 0uM, 0.00064uM, 0.16uM, 0.4uM, and 10uM); statistics calculated at the 9 day time point (J) Interferon, cytokine and chemokine production after treatment with baricitinib via cytokine bead array (K) ER-Hoxb8 cell counts over differentiation time course, comparing NTC, *Adar1* KO, and *Adar1/Ifih1* double KO (dKO) lines - dotted NTC and *Adar1* KO lines are equivalent to those shown in panel (B) (L) Interferon, cytokine and chemokine production via cytokine bead array, comparing NTC, *Adar1* KO, and *Adar1/Ifih1* dKO lines. All p-values calculated via one-way (L) or two-way (B, I, J, K) ANOVA with multiple comparisons; ns = not significant or $p \geq 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$

type I-specific interferon (IFN- β) and ISG (CXCL10, IL6, CCL5) production by *Adar1* KO ER-Hoxb8 macrophages, and their suppression by baricitinib treatment using multiplex bead array (**Figure 4J**). As interferons inhibit hematopoiesis and differentiation (Demerdash et al., 2021), we confirmed that the baricitinib-treated cells were indeed macrophages (**Figure 4G**) and that the ISG reduction improved macrophage production.

Melanoma differentiation associated protein 5 (MDA5, or *Ifih1*) is an epistatic modifier of *Adar1* such that deletion rescues some *Adar1* phenotypes (Liddicoat et al., 2015; Guo et al., 2022). We introduced a guide targeting *Ifih1* (BFP+) to *Adar1* KO cells (GFP+). We created single cell clones and validated double KO (dKO, GFP/BFP+) by TIDE analysis (**Supplemental Figure 4H**). *Ifih1* loss completely rescued the *Adar1* KO cell growth deficit during differentiation (**Figure 4K**), and normalized production of IFN- β , IL-6, and CCL5 (**Figure 4L**), but not CXCL10. As with baricitinib treatment (**Supplemental Figure 4G**), we found that multiple chemokines/cytokines are downregulated in *Adar1*-mutant cells as compared to NTC (TNF- α , IL-17, MIP-1a, MIP-1B, MIP-2, M-CSF), and are partially rescued by *Ifih1* deletion (**Supplemental Figure 4I**).

Together, these data show that *Adar1* deletion impairs macrophage but not progenitor health and is associated with interferonopathy, demonstrating the utility of the ER-HoxB8 model system for experimental isolation and study of macrophage dysfunction in a genetic disease.

***Adar1* mutation prevents ER-Hoxb8 engraftment**

We next attempted to study how *Adar1* KO macrophages affect the CNS. To our surprise, although TLR4 KO gene edited macrophages engrafted robustly ($n = 10$; **Figure 5A/E**), mice injected with *Adar1* KO cells died early and showed no engraftment in the brains of any transplanted mice that survived to endpoint ($n = 3$; **Figure 5B/C**).

Because blocking interferon signaling with baricitinib or *Ifih1* deletion correlated with better macrophage survival in vitro, we wondered if modifying our approach to limit interferon tone would permit *Adar1* KO cell engraftment. We first reduced the number of injected cells ($n = 4$) and harvested tissues earlier, between 4-8 days post-transplant ($n = 8$), and did not observe engraftment. We then tried to pre-treat donor cells and host mice with Baricitinib and likewise did not observe engraftment in surviving mice ($n = 1$; **Figure 5E**). Lastly, we transplanted *Adar1/Ifih1* dKO cells ($n = 3$), which showed partial rescue of *Adar1* KO phenotypes in vitro. Although we detected no areas of engraftment meeting the stringent criteria applied for our “percent area” quantification method, we noted diffuse patches of engrafted cells with rounded morphology and enlarged soma size, in three of three experimental replicates (**Figure 5D/E**, **Supplemental Figure 5A**), potentially consistent with a mild partial rescue of engraftment.

Csf1r^{-/-} mice have severe constitutional, skeletal, and CNS abnormalities, typically succumbing by weaning age (Dai et al., 2002). To extend our findings about *Adar1* KO macrophages in vivo we applied an inducible microglia depletion model using healthy *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} hosts (Bennett et al., 2018). After depleting endogenous microglia via subcutaneous tamoxifen injection at age P1 and P2, we intracerebrally transplanted cells into *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} brains (**Supplemental Figure 5B**). After 7-15 days in vivo, we saw engraftment of control (TLR4 KO and NTC; $n = 4$; **Supplemental Figure 5C**), but not *Adar1* KO cells ($n = 5$; **Supplemental Figure 5D**). Further mirroring the *Csf1r*^{-/-} data, we saw small, diffuse patches of engrafted *Adar1/Ifih1* dKO cells, but none reached quantifiable levels of engraftment ($n = 11$; **Supplemental Figure 5E**). Lastly, we harvested transplanted *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} hosts early, at day three, and saw a diffuse pattern of non-quantifiable but parenchymally engrafted *Adar1* KO cells ($n = 4$; **Supplemental Figure 5F**). These data suggest that *Adar1* KO cells can enter the brain parenchyma but do not persist, consistent with the reduction in cell numbers observed in vitro during macrophage differentiation.




Figure 5

***Adar1* mutation prevents ER-Hoxb8 engraftment in the *Csf1r*^{-/-} mouse, partially rescued by *Ifih1* deletion.**

(A) Representative rendering of donor cell engraftment (scale bar = 1000um) with inset microscopy of GFP+ donor cell engraftment (green = endogenous GFP, blue = DAPI; scale bar = 100um) for control cells (TLR4 KO and NTC) harvested 7-15 days post-injection (dpi), (B) *Adar1* KO (sgRNA #1) cells (harvest details in (E)), (C) *Adar1* KO (sgRNA #2) cells harvested 9-12 dpi, and (D) *Adar1/Ifih1* double KO (dKO) cells harvested 10-15 dpi (rendered dots in two right brains enlarged 5x for visualization) (E) Percent of total brain area tiled with cells between groups (numbers denote “n” per group); *Adar1* KO (#1) cells include pooled data (brains injected with 300k cells/hemisphere, harvested at 10-15 days post-injection (dpi); n = 3); brains injected with 300k cells/hemisphere, harvested at 4-8 dpi (n=8); brains injected with 50k cells/hemisphere, harvested at 13 dpi (n = 4); and brains injected with 100k cells/hemisphere pre-treated with 0.5uM Baricitinib, mice treated daily with 1 mg/kg Baricitinib, harvested at 5 dpi (n = 1)); asterisk indicates samples where engraftment is present but does not meet criteria for tiled brain area, as exemplified in (D); p-values calculated via one-way ANOVA with multiple comparisons; ns = not significant or p >= 0.05, ***p < 0.001, ****p < 0.0001

Overall, these results show that genetically modified ER-Hoxb8 cells robustly engraft in the brain parenchyma, which is prevented by *Adar1* KO, independent of interferon production.

***Adar1* D1113H mutant ER-Hoxb8 macrophages drive brain ISG expression**

Most available mouse models of AGS are either embryonic lethal (Wang et al., 2004; Hartner et al., 2004 [↗](#); Liddicoat et al., 2015 [↗](#)) or fail to produce CNS phenotypes (Morita et al., 2004 [↗](#); Ward et al., 2011 [↗](#); Hiller et al., 2012 [↗](#); Pereira-Lopes et al., 2013 [↗](#); Behrendt et al., 2013 [↗](#); Rehwinkel et al., 2013 [↗](#); Ohto et al., 2022 [↗](#)). We recently created a viable mouse model harboring a patient-derived mutation (D1113H) in the catalytic domain of ADAR1, which displays robust brain ISG expression, astrogliosis, microgliosis, and white matter calcifications (Guo et al., 2022 [↗](#)). Since AGS patient mutations are typically hypomorphic (rather than knockout), this more authentically models AGS (Rice et al., 2014 [↗](#)). Given the non-engraftment of *Adar1* KO cells, we created ER-Hoxb8 cells from *Adar1* D1113H mice to explore the impact of AGS-specific mutations in microglia. Unlike *Adar1* KO cells, D1113H ER-Hoxb8s showed normal growth and expansion during in vitro differentiation (**Figure 6A** [↗](#)), despite a similarly increased production of IFN- β , ISGs, and other cytokines, including CXCL10, CXCL1, CCL5, VEGF, MIP-1a, and MIP-1b, but not IFN- γ (**Figure 6B** [↗](#)).

We next characterized D1113H ER-Hoxb8 cells after transplantation into the *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} brain. As D1113H cells are not genetically tagged, we performed RNA in situ hybridization (ISH), using *Cre* expression (**Supplemental Figure 6A/B** [↗](#)) to distinguish endogenous microglia (*Cre*⁺) from donor cells (*Cre*⁻). Unlike *Adar* KO cells, D1113H mutant ER-Hoxb8s engrafted the parenchyma (**Figures 6C/D** [↗](#), *n* = 5 biological replicates). Notably, both D1113H homozygous mutant brains (**Figure 6C** [↗](#)) and mice engrafted with D1113H ER-Hoxb8 microglia-like cells exhibited similar and persistent upregulation in brain *Isg15*, with pockets of robust expression around engrafted Iba1⁺ MLCs, clustering around Iba1-nuclei that morphologically resemble neurons (**Figure 6D** [↗](#), **Supplemental Figure 7A** [↗](#)). To get a clearer idea of their overall engraftment potential, we transplanted D1113H mutant ER-Hoxb8s into *Csf1r*^{-/-} mice, which are unconfounded by repopulating endogenous microglia. Similar to brains transplanted with *Adar1/Ifih1* dKO cells, we found dispersed cell engraftment that did not reach quantifiable levels (**Supplemental Figure 7B** [↗](#), *n* = 3 biological replicates). Consistent with results from *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} host mice, we saw upregulation in brain *Isg15* clustering around Iba1-nuclei (**Supplemental Figure 7C** [↗](#), *n* = 3 biological replicates).

In sum, D1113H ER-Hoxb8 macrophages produce type I interferon and ISGs and induce *Isg15* production by neighboring cells after brain engraftment. These results demonstrate that *Adar1* mutation in microglia-like cells is sufficient to drive brain ISG production.

Discussion

Realizing the potential of microglia replacement requires new tools for research and therapy development. Here, we show the power of estrogen-regulated (ER) homeobox B8 (Hoxb8) conditionally immortalized myeloid cells to robustly model microglia for rapid assessment of gene-targeted perturbations on their identity and function. We find that ER-Hoxb8 macrophages are similar to bone marrow-derived (BMD) macrophages both in vitro and in vivo, where they robustly engraft the microglia-deficient brain and upregulate microglial identity genes. We then use this model to demonstrate an approach to untangle macrophage contributions to a neurological disease, establishing the impact of macrophage-specific mutations to clinically relevant phenotypes both in vitro and in vivo.




Figure 6

***Adar1* D1113H mutant ER-Hoxb8 macrophages persistently drive brain ISG expression.**

(A) In vitro ER-Hoxb8 cell counts over time (p-values calculated via two-way ANOVA with multiple comparisons) **(B)** Multiplex bead array data for interferons, cytokines, and chemokines produced via ER-Hoxb8 macrophages (p-values calculated via independent two-sample t-test) **(C)** Sagittal sections of non-transplanted (tamoxifen (tam) sham control) *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} brains (left) and *Adar1* D1113H mutant brains (right) at age P15 and P28-31; nuclei (blue, DAPI), *Isg15* (white) by RNA in situ hybridization (ISH)); scale bar = 1000um; red arrow depicts location of corresponding closeup images below, showing IBA1 (red, protein stain), Cre (green, ISH), *Isg15* (purple, ISH), and nuclei (teal, DAPI); scale bar = 20um; see **Supplemental Figure 7A** for further corresponding closeup images **(D)** Sagittal sections of *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} brains intracranially transplanted with WT ER-Hoxb8s (left) and *Adar1* D1113H ER-Hoxb8s (right) at 13 and 27 days post-injection (dpi); nuclei (blue, DAPI), *Isg15* (white, ISH); scale bar = 1000um; red arrow depicts location of corresponding closeup images below, showing IBA1 (red, protein stain), Cre (green, ISH), *Isg15* (purple, ISH), and nuclei (teal, DAPI); scale bar = 20um; see **Supplemental Figure 7A** for further corresponding closeup images. All p-values characterized by: ns = not significant or ≥ 0.05 , *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001

Microglia are difficult to target and manipulate for preclinical study, necessitating the use of robust microglial surrogates for discovery, such as the ER-Hoxb8 system described here. Powerfully programmed by the brain environment, microglia lose their unique transcriptional identity in vitro (Bohlen et al., 2017 [DOI](#); Gosselin et al., 2017 [DOI](#)) and are difficult to engineer (O'Brien et al., 2022 [DOI](#)). Common surrogates include macrophages from isolated bone marrow or immortalized cell lines derived from macrophages or microglia (Stanley and Heard, 1977 [DOI](#); Raschke et al., 1978 [DOI](#); Tsuchiya et al., 1980 [DOI](#); Righi et al., 1989 [DOI](#); Blasi et al., 1990 [DOI](#); Chanput et al., 2014 [DOI](#); Muffat et al., 2016 [DOI](#); Timmerman et al., 2018 [DOI](#)).

BMD cells are finite in number and exist as a heterogeneous population. Cell lines such as induced pluripotent stem cell-derived microglia-like cells can be expensive to generate and time-intensive to maintain (Timmerman et al., 2018 [DOI](#)). ER-Hoxb8 cells are expandable, readily transducible, and brain-engraftable, making them a robust microglia surrogate. Building on the work of others (Roberts et al., 2019 [DOI](#); Accarias et al., 2020 [DOI](#); Xu et al., 2022 [DOI](#)), we demonstrated straightforward gene editing using CRISPR-Cas9, generated stable lines using standard cell culture reagents, and found that ER-Hoxb8s are transcriptionally similar to primary BMD cells in vitro and in vivo. We used these strengths (and spared months of generating transgenic mouse lines) to readily test hypotheses about *Adar1* function in microglia to demonstrate the potential of this system. We generated *Adar1* knockout (KO) ER-Hoxb8 lines, as well as ER-Hoxb8 lines derived from mice harboring *Adar1* patient mutations, to study the effect of microglia-specific *Adar1* mutation on macrophage generation and interferon production. The ease with which ER-Hoxb8 cells can be modified, sorted, clonally expanded, banked, and transplanted was critical to these discoveries and will enable further innovation for the study of central nervous system macrophages.

In the future, the ER-Hoxb8 system will enable additional study of *Adar1* and microglia in Aicardi-Goutières Syndrome (AGS). We found that both complete *Adar1* KO, as well as an RNA-editing catalytic domain mutation in microglia-like cells, leads to increased interferon responses cell-autonomously in vitro, as well as non-cell-autonomously as demonstrated by direct transplant. This effect is blocked by *Ifih1*/MDA5 deletion, which is thought to sense abnormal RNA species formed by *Adar1* mutation and rescues *Adar1*-catalytic domain point mutation phenotypes (Liddicoat et al., 2015 [DOI](#); Guo et al., 2022 [DOI](#)). In contrast, brain engraftment is impacted by *Adar1* KO regardless of *Ifih1* genotype and not by *Adar1* catalytic domain mutation, suggesting that the full KO phenotype may be independent of RNA editing and interferon production. Mitochondrial antiviral signaling (MAVS) adaptor protein rescues the embryonic lethality of *Adar1* null mice (Mannon et al., 2014 [DOI](#)), suggesting that manipulation of other *Adar1* pathway components could reveal *Adar1*'s role in brain engraftment and macrophage maintenance. This system can allow for interrogation of alternative *Adar1* functions, as well as other AGS mutations, and enables testing of AGS-targeted therapies.

All cell models have limitations, and we highlight two of particular relevance to the study of microglia. First, like primary BMD macrophages, ER-Hoxb8 macrophages derive from definitive as opposed to primitive hematopoiesis - the authentic origin of microglia. Though very similar to microglia, ER-Hoxb8s cannot attain full microglial identity, conspicuously lacking *Sall1* expression, an important transcription factor in regulating homeostatic microglia (Buttgereit et al., 2016 [DOI](#)). Second, we observe that ER-Hoxb8 macrophages express slightly higher levels of inflammation-associated genes than primary BMD macrophages after brain engraftment. Though insufficient to elicit GFAP upregulation, a sensitive biomarker for inflammation, it remains an important consideration for experimental design. Interestingly, a recent study suggests it may be possible to address both limitations by treating yolk sac progenitors with the ER-Hoxb8 virus to generate macrophages with reduced inflammatory reactivity that may serve as more authentic microglia surrogates (Elhag et al., 2021 [DOI](#)). It also raises the exciting potential of an analogous system for study of human, patient-derived macrophages.

More broadly, the future of engineered microglia surrogates, including ER-Hoxb8 microglia-like cells, is bright. Leveraging the advantages of ER-Hoxb8 cells, one could create controllable gene editing via an inducible Cas9 or sgRNA and perform large, pooled screens in vitro and in vivo simultaneously, using progressively advanced methods for gene targeting. We envision microglia as potent neurotherapeutics (Bennett and Bennett, 2020), and toward that goal, the ER-Hoxb8 model provides an ideal milieu for the testing of therapeutic cell engineering strategies, such as payload delivery, synthetic receptors, and customizable gene circuits. Here we applied two engraftment models, utilizing both *Csf1r*^{-/-} and inducible *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} host mice. Pairing the ER-Hoxb8 system with our CSF1R inhibitor-resistant receptor transplantation models will allow even more widespread adoption of these techniques (Chadarevian, Lombroso, et al., 2023 [↗](#)). Together, the combination of accessible and modifiable microglial surrogates alongside increasingly effective transplantation models holds great potential for research and therapy development.

Materials/Methods

Mouse Models

All animal studies were performed with approval from the Children's Hospital of Pennsylvania IACUC panel in accordance with institutional and national regulations. All animals were housed in a non-barrier facility with 12-hour light/dark cycles at 23±2 degrees C in ventilated cages with no more than five animals per cage. Animals were provided water and standard chow *ad libitum*. Cages and bedding were changed weekly.

Csf1r^{-/-} (FVB.129×1-*Csf1r*^{tm1Ers}) and *Csf1r*^{+/-} littermate animals on the FVB background were a generous gift from Dr. Richard Stanley (Albert Einstein College of Medicine). *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} mice were generated by intercrossing JAX 021212 and 021160 strains. *Adar D1113H* mice (p.Asp1113His) were a generous gift from Dr. Qingde Wang (University of Pittsburgh). For experiments using GFP-expressing donor cells, we backcrossed Osb-GFP (C57BL/6-Tg(CAG-EGFP)131Osb/LeySopJ (JAX 006567)) onto FVB WT mice (JAX 001800) for 22 generations. For experiments using constitutively-expressed Cas9 donor cells, we used FVB.129(B6)-Gt(ROSA)26Sortm1.1(CAG-cas9*,-EGFP)Fezh/J mice (JAX 026558).

In Vitro Experiments

ER-Hoxb8 Conditionally Immortalized Cell Production

Immortalization of murine myeloid progenitor cells was completed as outlined in Wang et al. (2006) [↗](#). Briefly, bone marrow progenitors were isolated from the femurs and tibias of wildtype or Cas9-expressing FVB/NJ mice using a Percoll density separation (Cytiva, 17544501). Cells were cultured for three days in RPMI media (Invitrogen, 7240047) supplemented with 10 ng/mL IL3 (Peprotech, 213-13), IL6 (Peprotech, 216-16), and SCF (Peprotech, 250-03) before transduction with MSCVneo-HA-ER-Hoxb8 virus. Originally created by Dr. David Sykes (Massachusetts General Hospital), the MSCVneo-HA-ER-Hoxb8 plasmid was a generous gift from Dr. Igor Brodsky (University of Pennsylvania School of Veterinary Medicine). 24 hours after transduction, the cells were recovered and grown for 48 hours in RPMI media (Invitrogen, 7240047) supplemented with 1 mM Na-Pyruvate (Invitrogen, 11360070), 0.0005 mM beta-estradiol (Sigma Aldrich, E2758), and 10 ng/mL GM-CSF (Peprotech, 315-03). Cells were then selected for transduction with the addition of 1 mg/mL Geneticin (ThermoFisher, 10131035) for 48 hours. Cells were then expanded and passaged every 48-72 hours at 25-50,000 cells/mL for two weeks, or until all non-immortalized cells were terminally differentiated or killed.

Bone Marrow Isolations

Bone marrow harvests were completed as previously described (Bennett et al., 2016 [DOI](#)). Briefly, femurs and tibias were dissected and flushed with 1x PBS to collect whole bone marrow. Red blood cells were then lysed with ACK Lysis Buffer (Quality Biological, 118-156-101). For experiments requiring bone marrow progenitors, whole bone marrow samples were subsequently enriched for lineage-negative cells using the Miltenyi Direct Lineage Cell Depletion Kit (130-110-470) via MACS Column separation with LS Columns (Miltenyi, 130-042-401). For experiments requiring bone marrow monocytes, whole bone marrow samples were subsequently enriched for monocytes using the Miltenyi Monocyte Isolation Kit (130-100-629) via MACS Column separation with LS Columns.

Macrophage Differentiation

Bone marrow-derived macrophages were differentiated in vitro by plating isolated bone marrow-derived progenitors in petri dishes at a density of five million cells per dish in differentiation media - DMEM media (ThermoFisher, 10569010), 10% FBS (VWR, 89510-186), 1% Penn/Strep (Invitrogen, 15140122), supplemented with 30 ng/mL murine M-CSF (Peprotech, 315-02). Cells were differentiated for seven to nine days in a 37 degree C incubator with 5% CO₂, with media changes every two to three days.

ER-Hoxb8-derived macrophages were differentiated in vitro using methods adapted from Wang, et al. (2006) [DOI](#). Briefly, ER-Hoxb8 progenitors were plated in petri dishes at a density of two million cells per dish, and then differentiated in the same media and timetable as noted above.

Phagocytosis Assay

C57BL/6 WT ER-Hoxb8 cells were differentiated for 7 days as described above. The day prior to the phagocytosis assay, cells were harvested via incubation with 3 mM EDTA, resuspended in macrophage differentiation media, diluted to 25,000 cells/mL in macrophage differentiation media, and plated in separate wells of 8 Chambered Coverglass Systems (CellVis, C8-1.5H-N; 500 µl cell dilution/12,500 cells per well). The phagocytosis assay was adapted from Williamson and Vale (2018) [DOI](#) and Flannagan et al., 2014 [DOI](#). Briefly, 10% phosphatidylserine (PS) beads were used to mimic apoptotic cell corpses and phosphatidylcholine (PC) beads were used to mimic live cells. 10% PS beads were 4.07 µm diameter silica beads (Bangs Labs, SS05002) coated in a supported lipid bilayer composed of 10 mol.% 18:1 PS (DOPS – Avanti, 840035), 89.90 mol.% 16:0-18:1 PC (POPC – Avanti, 850457), 0.1 mol.% atto390 dye-labeled DOPE (Atto-Tec, AD 390-161). PC beads were 4.07 µm diameter silica beads (Bangs Labs, SS05002) coated in a supported lipid bilayer composed of 99.90 mol.% 16:0-18:1 PC (POPC – Avanti, 850457), 0.1 mol.% atto390 dye-labeled DOPE (Atto-Tec AD 390-161). Beads were diluted to 500,000 beads/mL in macrophage differentiation media and added to the macrophages via media change, replacing old media with 500ul bead dilution per chamber of the 8 Chambered Coverglass Systems. Macrophages were incubated with synthetic apoptotic cell corpse mimics or synthetic live cell mimics for 60 minutes at 37 degrees C, 5% CO₂. Cells were then fixed for 10 minutes at 37 degrees C, 5% CO₂ in 4% paraformaldehyde (Sigma, P6148) in PBS, permeabilized for 1 minute in 0.1% TritonX-100 (Sigma, T8787) in PBS, and stained using HCS CellMask Green (Invitrogen, H32714). After staining, cells were placed under Aqua-Poly/Mount for imaging (Polysciences, 18606). Macrophages were imaged via spinning disk confocal microscopy on an inverted Nikon Ti2-E microscope fitted with a CSU-W1 spinning disk unit and Hamamatsu ORCA-FusionBT camera. Images were acquired using a 40x/0.95 NA objective in .nd2 format using Nikon Elements software and converted to ome-tif stacks for analysis in Fiji/ImageJ. Beads were considered ingested if HCS CellMask Green signal was visible fully surrounding a circular area inside the cell corresponding to the 4 µm diameter of the silica beads. Experiments were replicated four times on four different days using independently prepared reagents each day. 81-100 cells were examined per replicate condition.

Viral Production and Transduction

To create retroviral supernatants, Lenti-X 293 T cells (Takara Bio, 632180) were transfected using 850 ng/mL pCL-Eco (Addgene, 12371) and 850 ng/mL desired retroviral plasmid, supplemented with 5 μ L/mL LipoD293 (SigmaGen, 504782) in media containing DMEM (ThermoFisher, 10569010), 25 mM HEPES (Invitrogen, 15630080), 10% FBS (VWR, 89510-186), and 1% Penn/Strep (Invitrogen, 15140122). Six hours post-transfection, media was replaced and viral supernatants were collected 48 and 72 hours later. Viral collections were combined and concentrated 10x using Retroviral Precipitation Solution (Alstem, VC200) according to manufacturer's instructions.

Transduction of ER-Hoxb8 progenitors was completed in 12-well plates coated with 1 μ g/mL fibronectin (Sigma-Aldrich, F1141). Cells were plated at 200,000 cells/mL and desired titer volume of virus was added. Plates were then spun for 90 minutes at 1000xg. Post-spinfection, cells were gently mixed and resuspended in an additional 2-3 mL of fresh media. 24 hours post-transduction, cells were recovered and expanded at normal growing conditions (25-50,000 cells/mL at 37 degrees C, 5% CO₂).

CRISPR Editing

Target gene guide sequences were created using the Broad Institute's CRISPick website (Mouse GRM38 reference genome, CRISPRko mechanism, SpyoCas9 enzyme, Hsu (2013) tracrRNA) and adapted to ligate into our plasmid backbone. This backbone, which was a generous gift from Dr. Will Bailis (University of Pennsylvania; [Bailis et al., 2019](#)) was adapted from the MSCV-MigR1-GFP/BFP plasmid (Addgene, 27490) to include a U6-driven gRNA scaffold for the insertion of guide sequences. The BbsI-HF restriction enzyme (NEB, R3539) was used to linearize the plasmid backbone at the gRNA scaffold site, and annealed guide oligos were ligated in. The ligated plasmids were then transformed using NEB Stable Cells (C3040), plated onto LB plates supplemented with 50 μ g/mL ampicillin, and grown for 24 hours at 30 degrees C. Correct insertion of guide sequences were confirmed using bacterial colony sanger sequencing via GeneWiz/Azenta, and full plasmid sequences were confirmed using Plasmidsaurus. Viral supernatants were then created as described above and Cas9-expressing ER-Hoxb8 macrophage progenitor cells were transduced as described above. Cells were then double-FACS sorted for transduction via fluorophore expression and expanded as normal. Successful editing was confirmed using TIDE as described in [Brinkman et al. \(2014\)](#). Briefly, gDNA was collected from both control and experimental macrophages and the cut site was amplified via PCR and sent for Sanger sequencing. Indel creation was then confirmed via the TIDE application ([Brinkman et al., 2014](#); <http://shinyapps.org/datacurators/tide/>). When an antibody was commercially available, such as with TLR4 (Biolegend, 145403, PE), loss of protein was also confirmed via flow cytometry.

TNF-a ELISA

ER-Hoxb8 progenitor cells were plated and differentiated as described in the above "Macrophage Differentiation" methodology. After seven days in vitro, cells were gently removed from the petri dishes using ice cold 1x PBS + 2 mM EDTA (ThermoFisher, 15575020). Cells were then re-plated at 50,000 cells/well in a TC-treated 96-well plate in 100 μ L fresh differentiation media. 24 hours later, LPS (Sigma, L2880-25MG) and R848 (Invivogen, TLRL-R848) were added at 100 ng/mL to respective wells. After 9.5 h, supernatants were collected and diluted to 25% initial concentration to stay within kit ranges. TNF-a concentrations were measured using the TNF alpha Mouse Uncoated ELISA Kit with Plates (ThermoFisher, 88-7324-22) according to manufacturer's instructions.

Adar1 Immunofluorescence

ER-Hoxb8 progenitor cells were plated and differentiated as described in the above “Macrophage Differentiation” methodology. NTC and *Adar1*-edited cells were plated at 5,000 cells per well in a glass-bottom 96-well plate. Cells were fixed on day eight of differentiation in 4% PFA (EMS, 15714) for 10 minutes. Cells were washed, blocked in 0.1% Tween20 (Bio-Rad, 1706531) + 5% donkey serum (Sigma, S39-100ML) for one hour, and then stained overnight with CD11B (BioLegend, 101208) and Hoechst 33342 Solution (ThermoFisher, 66249) at 4 degrees C. The following morning, cells were washed and imaged at 40x magnification Z-stacks using a BZ-X800 (Keyence) microscope.

Adar1 Cell Quantification

ER-Hoxb8 progenitor cells were plated and differentiated as described in the above “Macrophage Differentiation” methodology. NTC and *Adar1*-edited cells were plated at 5,000 cells per well in glass-bottom 96-well plates (n = 3 independently differentiated replicates per condition). Cells were plated in fresh differentiation media supplemented with nothing, vehicle (DMSO), or 10uM Baricitinib (Selleck Chemicals, S2851). Cells were plated with 125uL media on day 0, 62.5uL media was added on day 3, media was removed and replaced with 100uL media on day 6. On differentiation days 0, 3, 6, and 9, corresponding subsets of cells were stained with 1:10000 Hoechst for 30 minutes at 37 degrees C and imaged with the ImageXpress Micro Confocal System. Images were taken at 10x magnification at nine sites per well, with exposure settings consistent between wells. Cell number quantification was performed using the “Count Nuclei” pipeline in the ImageXpress software. Signal thresholds were set, masks were created over the Hoechst+ signal, and nuclei were counted (3-16 pixel width). Cell counts were averaged across the nine imaging sites and plotted as average nuclei per field per day of differentiation.

Adar1 Multiplex Arrays

ER-Hoxb8 progenitor cells were plated and differentiated as described in the above “Macrophage Differentiation” methodology. NTC and *Adar1*-edited cells (*Adar1* KO #1, *Adar1* KO #3, *Adar1*/*Ifih1* dKO) were plated at 100,000 cells per well in an untreated 24-well plate (n=4 independently differentiated replicates per condition), whereas WT and D1113H cells were plated at 25,000 cells per well in an untreated 24-well plate (n=4 independently differentiated replicates per condition). Media was changed every two days, with the baricitinib groups receiving 0uM, 0.00064uM, 0.16uM, 0.4uM, or 10uM Baricitinib (Selleck Chemicals, S2851) with each media change. On day seven of differentiation (24 h post final media change), cell supernatant was collected, spun down, and immediately frozen. Multiplexing analysis was performed using the LuminexTM 200 system by Eve Technologies Corp. (Calgary, Alberta). Assays used include the Mouse Cytokine/Chemokine 44-Plex Discovery Assay[®] Array (MD44), the Mouse Cytokine/Chemokine 32-Plex Discovery Assay[®] Array (MD32), as well as the Mouse IFN- α + IFN- β Assay (MDIFNAB). For the baricitinib groups, RNA was immediately collected from adherent cells for subsequent sequencing analyses.

In Vivo Experiments

Neonatal Transplantation

For intracranial transplantation into *Csf1r*^{-/-} mice, p0-p5 pups were injected as previously described (Bohlen et al., 2017 [\[10\]](#)) by hand using a pulled glass capillary tube (World Precision Instruments, 1B100F-4) in an electrode holder connected by silicon tubing to a syringe. One microliter containing a single-cell suspension (300,000 cells/ul unless otherwise noted) of donor cells in 1x PBS was slowly injected bilaterally into cortex, 1-2 mm anterior and 2-3 mm lateral to lambda at a depth of 0.5-1 mm. Host animals were harvested 12-16 days after injection unless otherwise noted.

Microglia/MLC Isolations

Single-cell suspensions of microglia/engrafted brain macrophages were isolated as previously described (Bennett et al., 2016 [DOI](#)). Briefly, mice were anesthetized using a cocktail of ketamine (100 mg/kg) plus xylazine (10 mg/kg) in 1x PBS and perfused with 10 mL cold PBS. Brains were dounce homogenized in 10 mL cold Medium A Buffer - 10% 10x HBSS (Fisher, 14185052), 1.5% 1M HEPES (Invitrogen, 15630080), 1.67% 30% glucose (Sigma, G7021-1KG), 86.8% ddH₂O supplemented with 2% DNase (Worthington, LS002007). Homogenate was filtered through a 70µm strainer and pelleted. Microglia were collected following a spin with isotonic percoll - 10% 10x HBSS, 90% Percoll PLUS density gradient media (GE Healthcare, 17544501). Cells were washed with Medium A Buffer and resuspended in desired media.

In Vivo Baricitinib Treatment

Baricitinib (Selleckchem, S2851) was reconstituted in DMSO (Sigma-Aldrich, D26250) and added to a diluent of 40% PEG300 (Sigma-Aldrich, 90878), 10% Tween80 (Sigma-Aldrich, P1754), and 50% water. Mice were administered 20µL daily by intraperitoneal injection at doses of 10 mg/kg, 5 mg/kg, or 1 mg/kg.

Staining Experiments

Immunofluorescence

Mice were perfused with 15 mL 1x PBS followed by 15 mL 4% PFA (EMS, 15714) and organs were then drop-fixed in 4% PFA for 16-24 hours at 4 degrees C. Organs were then cryo-protected in 30% sucrose (Neta Scientific, SIAL-S5391-25KG), embedded in OCT (Fisher, 23-730-571), cryo-sectioned at 16µm, mounted on Superfrost Plus slides (Fisher, 1255015), and frozen at -80 degrees C. Slides were then thawed at 60 degrees C, rehydrated in 1x PBS, and blocked for one hour at RT in blocking buffer - 90% 1x PBS, 9.5% donkey serum (Sigma, S39-100ML), 0.5% Triton X-100 (Sigma, T8787-50ML). After blocking, slides were incubated with primary antibodies (described below) in staining buffer (98.5% 1x PBS, 1% donkey serum, 0.5% Triton X-100) overnight at 4 degrees C. In the morning, slides were washed with 1x PBS and incubated with secondary antibodies (described below) for one hour at RT in staining buffer. Slides were washed once more and coverslipped with DAPI mounting media (EMS, 17989-60) before imaging.

Antibodies used: Rabbit Anti-Iba1 (Fujifilm, 019-19741, 1:500); Mouse Anti-GFAP (Agilent, MAB360, 1:500); Donkey anti-rabbit IgG 594 (ThermoFisher, A-21207, 1:500); Donkey anti-mouse IgG 647 (ThermoFisher, A32787, 1:500)

RNA In Situ Hybridization

In situ hybridization (ISH) was performed on mounted fixed frozen tissues using the RNAscope system per the manufacturer's protocol (Advanced Cell Diagnostics). Fluorescent detection was achieved using the RNAscope Multiplex Fluorescent Detection kit V2 (cat # 323110). Samples were probed for mouse *Isg15* (cat # 559271-C2) and *Cre* (cat # 312281-C3) and visualized with TSA Vivid dyes (cat #s 7534 and 7536). Following ISH, slides were immunostained with rabbit anti-Iba1 (cat # 019-19741, Fujifilm WAKO) and detected using fluorescently labeled donkey anti-rabbit secondary antibody (A21207, Invitrogen). Slides were coverslipped and nuclei were stained with Prolong Gold Antifade Mounting Medium with DAPI (cat # P36931, Invitrogen)

Imaging Acquisition and Processing

Slides were imaged using a BZ-X800 (Keyence) microscope. Whole-organ tile images plus z-stack images were captured and stitched/compressed using Keyence Analyzer software prior to image export. All images within experiment panels were taken with equivalent exposure settings. All images were then analyzed with FIJI (<https://imagej.net/Fiji>), equivalently adjusting only for brightness and black values (raw images available upon request).

Engraftment renderings were created using GFP (488) or Iba1 (594) and DAPI channels of 10x stitches. Background was equivalently subtracted using the “subtract background” function before manually thresholding via the “otsu” setting. Rendered dot masks were created by analyzing particles (size = 2-144 px; circularity = 0.5-1.0) and overlaying them on the corresponding DAPI channel. The boundaries of the brains were then outlined using the “polygon selections” tool, and everything outside this boundary was cleared and excluded from the final image.

16 μ m tissue sections used for RNA scope were imaged using an Andor BC43 spinning disc confocal microscope (Oxford Instruments). Sections that were directly compared were acquired using the same exposure, laser power, gain, and resolution settings. Subsequently, image analysis on raw unmanipulated images was performed using Imaris software. Representative images displayed for publication were processed equally in Imaris across all conditions to best display the data. Comprehensive notes and raw images are available upon request.

Percent Area Quantifications

For each brain sample, average total engrafted area was calculated from three matched sagittal sections (200 μ m apart starting at the medial-most point of the tissue), each using a 4x tile-scan image collected via Keyence microscope. Calculations were made in FIJI using the Iba1 and/or GFP channel tiled images. FIJI’s “polygon selections” tool and “measure” function were used to draw and calculate a total brain ROI, as well as engrafted area ROI(s). The latter were defined as regions with Iba1+ and/or GFP+ cell groups with no fewer than 50 total cells per “group” nor 50 cells/mm². Edge limits of engraftment areas were defined by the nucleus of the edge-most cell being no more than 200 μ m from nearest-neighbor cells. Percent area engrafted was then calculated by dividing total engrafted area(s) by total brain area. Olfactory bulb was omitted for consistency as it was not present in all samples.

For each brain sample, average total area of GFAP+ cells was calculated from three matched sagittal sections (250 μ m apart starting at the medial-most point of the tissue), each using a 10x tile-scan image collected via Keyence microscope. Calculations were made in FIJI using the GFAP channel tiled images. Background was equivalently subtracted from all images using the “subtract background” function. Images were then converted into 8-bit format and thresholded at 40/255 with the “intermodes” setting. ROIs were manually drawn for cortex, hippocampus, midbrain, and thalamus using FIJI’s “polygon selections” tool. Percent area covered was then calculated using the “measure” function to quantify positive GFAP signal over total ROI area.

Cell Density Quantification

For each brain sample, average cortical cell density was calculated using the same three matched sagittal sections used above to calculate area of engraftment. Here, three 20x representative images were included that spanned the cortical region of engraftment. If the region of engraftment was small, fewer images were included such that no cells overlapped and were counted more than once. Cells were counted manually using FIJI’s “Cell Counter” tool and were defined as cells that were Iba1+ and/or GFP+ and directly overlaid with a DAPI+ nucleus. Cell counts per image were averaged and divided by total area per image to obtain densities at cells/mm².

Morphological Analysis

All morphological analyses were performed using Fiji. Individual Iba1+ cells in the cortical parenchyma were randomly selected from 10x stitched images of the brain. For all images, individual cells were converted to 8-bit and thresholded equivalently. The despeckle function was used to remove noise from each sample. Each image was then converted to a binary image that was used for further analysis. Soma size was measured by drawing a region of interest (ROI) around somas and measuring the encompassed area. Primary branches were defined as branches extending directly from the soma and were counted manually. The length of each branch was measured from the center of the cell soma using the Neuroanatomy SNT plugin (Arshadi et al., 2021 [\[4\]](#)). Traces were drawn to outline branch morphology, then measured using the SNT Measurements function to obtain branch lengths.

Flow Cytometry and FACS

Once isolated into single cell suspensions, cells were washed 2x in FACS Buffer - 1x PBS, 0.5% BSA (Sigma, 126609-10GM), 0.5 mM EDTA (ThermoFisher, 15575020) - and blocked with FC Block - 0.2% CD16/CD32 (BD Biosciences, 553142) in FACS Buffer - for 10 m at RT. Cells were then stained using antibodies (described below) for 30-60 m at 4c. Cells were washed 2x more in FACS Buffer before being analyzed or sorted as described below.

Antibodies used: LIVE/DEAD fixable far red dead cell stain kit (Invitrogen, L10120, 1:1600); Anti-mouse CD45 (PE/Cyanine7, Clone 30-F11, Biolegend, 103113, 1:400); Anti-mouse/human CD11b (PerCP/Cyanine5.5, Clone M1/70, Biolegend, 101227, 1:400); Anti-mouse monoclonal TMEM119 (PE, Clone V3RT1GOsz, Invitrogen, 12-6119-82, 1:400) For flow cytometry experiments, stained cells were analyzed on a CytoFLEX LX (6 Laser) Flow Cytometer and visualized via FlowJo (10.8.1). For sorting experiments, stained cells were sorted with a 100uM nozzle on a FACSaria Fusion, FACSJazz, Aurora, or MoFlo Astrios depending on Core availability (no effects on data were seen between FACS machines). Cells were deposited into media for cell expansion, or TRIzol LS Reagent (Fisher, 10296028) for subsequent RNA extraction.

Bulk RNA-Sequencing and Analysis

RNA Extraction & Quantification

For in vitro populations, RNA was extracted using the Qiagen RNeasy Mini Kit (74104) according to manufacturer's instructions. For suspended populations, cells were collected and spun down at 200xg for 5 m, supernatant was removed, and 350ul Buffer RLT was added directly to the cell pellet. For adherent populations, supernatant was removed, plates were washed with 1x PBS, and 650ul of Buffer RLT was added directly to the dish to create cell lysates.

For in vivo populations, cells were isolated as described in the above "Microglia/MLC Isolations" methodology and FACS sorted into TRIzol LS reagent as described in the above "Flow Cytometry and FACS" methodology. Cells were briefly vortexed and 0.2 volumes of chloroform (Sigma, C2432-500ML) were added. Cells were again vortexed, incubated for 3 m at RT, and then centrifuged at 12,000xg for 15 m at 4c. The upper aqueous phase was then collected and 1 volume of fresh 70% EtOH was added. Cells were again vortexed and then run through the Qiagen RNeasy Micro Kit (74004) according to manufacturer's instructions. Isolated RNA was stored at -80.

When able, RNA Integrity Numbers (RINs) were calculated using an Agilent TapeStation with High Sensitivity RNA ScreenTape (5067-5579) and Sample Buffer (5067-5580) according to manufacturer's instructions. Cells with a RIN $\geq$ 7.5 were used for subsequent library

preparations. For in vitro datasets, the average RIN score was 9.03. Quantification was often not possible for brain-isolated cell populations due to low yield. In this case, all samples were used for subsequent library preparations, and quality was assessed later during preparation.

Library Preparation & Sequencing

Sequencing libraries were prepared in-house using either 40 ng or 8ul RNA per sample, depending on if RIN scores and concentrations were able to be calculated as discussed above, using the NEBNext Single Cell/Low Input RNA Library Prep Kit for Illumina (NEB, E6420) according to manufacturer's instructions. Quality and quantity steps were performed using an Agilent TapeStation using High Sensitivity D5000 ScreenTape (5067-5592), and High Sensitivity D5000 Reagents (5067-5593).

Bulk RNA-sequencing of completed libraries was performed by the Children's Hospital of Philadelphia's Center for Applied Genomics (CAG) Core Facility using a NovaSeq 6000 (Illumina) system with the SP Reagent Kit v1.5 (2×100 bp). Data was de-multiplexed and sent to us as FastQ files via BaseSpace (Illumina).

For sequencing of NTC and *Adar1*-edited cells with varying doses of Baricitinib, cDNA libraries were sent to Novogene for sequencing using a NovaSeq 6000 (Illumina) system with the S4 Reagent Kit (2×150 bp). Data was de-multiplexed and sent to us as FastQ files for manual download. Data sequenced by Novogene was never combined with data sequenced by CAG.

Sequencing Analysis

All pre-processing steps were run using Terminal. FastQ files were downloaded from BaseSpace (Illumina) and concatenated across lanes using the “cat” function. All files were processed using FastP (Chen et al., 2018 [DOI](#); Chen, 2023 [DOI](#)) to filter reads and trim Illumina adaptors, as well as FastQC (Andrews, 2014 [DOI](#)) to assess quality. All files were determined to be of sufficient quality for subsequent downstream processing. Reads were pseudoaligned to the mouse GRCm39 cDNA transcriptome (Ensembl release 99; Schneider et al., 2017 [DOI](#); Zerbino et al., 2018 [DOI](#)) using Kallisto (version 0.46.0; Bray et al., 2016 [DOI](#)) and run through MultiQC (Ewels et al., 2016 [DOI](#)) for a final quality check after mapping.

All post-processing steps were run using RStudio (R Core Team, 2022 [DOI](#)). Transcripts were summarized to the gene level through tximport (version 1.14.2; Soneson et al., 2015 [DOI](#)) with abundance counts calculated via TPM and then normalized using TMM via edgeR (Robinson et al., 2010 [DOI](#); McCarthy et al., 2012 [DOI](#); Chen et al., 2016 [DOI](#)). Hierarchical cluster dendrograms were created using the “dist” function (method = euclidean) and “hclust” function (method = complete) and were visualized with the plot function. PCA plots were created using the “prcomp” function. Differential gene expression analysis was performed using linear modeling via limma (Ritchie et al., 2015 [DOI](#)), and differentially expressed genes (DEGs) were decided by applying an FDR cutoff of 0.05, a counts per million (CPM) cutoff of +1, and a Log2(FC) cutoff of +/- 2. Linear models and R² values were created using the “lm” function. All output plots were visualized with ggplot2 (Wickham, 2016).

Statistical overrepresentation tests and related GO Terms were calculated using the Panther Classification System (Version 18.0). Respective lists of DEGs were manually imported into the Gene List Analysis tab, selected for mus musculus, and run through the statistical overrepresentation test option using the PANTHER GO-Slim options for biological process, molecular function, and cellular component. Tests were run using the Fisher's Exact option while calculating the false discovery rate. FDR was set to a threshold of 0.05, and terms were ranked via the hierarchy option.

Statistical Calculations


GraphPad Prism (Version 10.0.0) was used to perform statistical tests and generate p values with standard designations (ns = not significant or $p \geq 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$). All values are shown as mean $\pm$ standard error of the mean (SEM). Details regarding replicate numbers and individual statistical test used are provided in the respective figure legends.

Acknowledgements

We thank Dr. Richard Stanley (Albert Einstein College of Medicine, New York, United States) for the *Csf1r*^{-/-} (FVB.129X1-Csf1rtm1Ers) and *Csf1r*^{+/+} littermate animals on the FVB background; Dr. Qingde Wang (University of Pittsburgh, Pittsburgh, United States) for the *Adar1* D1113H mice (p.Asp1113His); Dr. David Sykes (Mass General Hospital, Boston, United States) and Dr. Igor Brodsky (University of Pennsylvania, Philadelphia, United States) for the MSCVneo-HA-ER-HoxB8 plasmid; Dr. Dan Beiting for the bulk RNA sequencing DIY Transcriptomics scripts; the Children's Hospital of Philadelphia Center for Applied Genomics for bulk RNA sequencing assistance; the Children's Hospital of Philadelphia Flow Cytometry core for flow cytometry and sorting assistance; and Kayla Peelman (Emory University, Atlanta, United States) for assistance with Illustrator. This work was supported by National Science Foundation Graduate Research Fellowship Program DGE-1845298 (KMN, SIL); NIH Training in Age Related Neurodegenerative Diseases T32 2-T32-AG-000255-26 (VSC); NIH Medical Scientist Training Program T32 GM007170 (VSC); NIH T32MH019112 (WHA); NIH T32 GM008076 (SIL); Blavatnik Family Fellowship, Blavatnik Foundation (SIL); NIH R35GM138085 (WB); Paul Allen Institute Distinguished Investigator Award (WB); NIH R15NS133939 (APW); R01AI139544 (QW); R01NS134651 (QW); NIH DP5OD036159 (MLB); NIH R01-NS-120960-01 (FCB); Klingenstein-Simons Fellowship in Neuroscience (FCB); and The Paul Allen Frontiers Group GRT-00000774 (FCB).

Additional files


Supplemental Tables [↗](#)



Supplemental Figure 1

Extended validation and comparison of ER-Hoxb8 to BMD macrophages in vitro, relating to Figure 1


(A) Gating strategy for Figure 1C (B) Flow cytometry histograms and median fluorescence intensity (MFI) of CD11B (left) and CD45 (right), relating to Figure 1C (C) Merged image of ER-Hoxb8 macrophages (green) and synthetic apoptotic cell corpse mimics (10% PS beads, purple) imaged after fixation and staining (orange denotes beads counted as ingested) (D) Quantification of phagocytosis of synthetic apoptotic cell corpse mimics (10% PS beads) as compared to synthetic live cell mimic controls (PC beads); p-value calculated via nested t-test for n=4 replicates per group, with overall distribution (81-100 cells per replicate) shown in gray; ***p < 0.001 (E) Heatmap showing Log2 CPM gene expression levels of macrophage genes across groups; progenitor and macrophage groups are as represented in Figure 1A, BM Mono = monocytes isolated from BM, 4div ER-Hoxb8 = ER-Hoxb8s collected after four days of in vitro differentiation (F) PCA plot combining bulk RNA sequencing data amongst all groups (G) Log2 CPM values of top 10 differentially expressed genes by Log2FC (CPM > 1, Log2FC >= 2, FDR < 0.05)



Supplemental Figure 2

Extended comparison of ER-Hoxb8 and BMD macrophages after intracranial transplantation in *Csf1r*^{-/-} hosts, relating to Figure 3 [↗](#).


(A) Histogram of TMEM119 surface staining by flow cytometry (pre-gated on live, singlet, leukocyte, CD45⁺/CD11b⁺) for seven-day differentiated in vitro BMD and ER-Hoxb8 macrophages **(B)** Immunostaining of cortical brain region 12-16 days post-intracranial injection (red = IBA1, green = endogenous GFP, white = P2RY12; blue = DAPI; scale bar = 100μm) **(C)** Gating strategy for in vivo TMEM119 histogram shown in **Figure 3A** [↗](#) (pre-gated on live, singlet, leukocyte, GFP) of WT microglia and intracranially transplanted (ICT) cells (BM or ER-Hoxb8s) **(D)** Unsupervised hierarchical cluster dendrogram, related to **Figure 3B** [↗](#) (distance method = euclidean; cluster method = complete) **(E)** Comparison of "batch" by PCA plots for harvest days (left), sorter used (middle), and host mouse sex (right) **(F)** Venn diagrams comparing the top 200 up and downregulated genes, as compared to unmanipulated in vivo microglia; asterisk denotes data derived from previously published dataset (Bennett et al., 2018 [↗](#)); genes were rank-ordered by absolute expression value, with only the top 1000 being considered for comparison, sorted by Log2FC > 2, FDR < 0.5, and ordered by Log2FC values **(G)** Quantification of GFAP percent area covered; n = 3 biological replicates per group; each dot = one biological replicate (average area across three matched sagittal sections); p-values calculated via one-way ANOVA with multiple comparisons; ns = not significant or p >= 0.05 Panel F created with [BioRender.com](#) [↗](#)



Supplemental Figure 3

CRISPR-Cas9 editing of ER-Hoxb8 progenitors, relating to Figures 4 and 5.


(A) Schematic for *Tlr4* knockout using Cas9^{+/-} ER-Hoxb8s and sgRNA viral transduction (B) Post-editing trace decomposition charts (via TIDE Analysis) for *Tlr4* KO macrophages (C) Histogram of TLR4 surface staining by flow cytometry (pre-gated on live, singlet, leukocyte) for eight day-differentiated in vitro BMD and ER-Hoxb8 macrophages (D) TLR4 median fluorescence intensity (MFI) normalized to FMO levels using flow cytometry data shown in C (E) TNFα production, comparing control, LPS, and R848 treated samples (eight-day differentiation, 9.5 h LPS or R848, 100 ng/mL, n = four replicates per condition); p-values calculated via two-way ANOVA with multiple comparisons; ns = not significant or p ≥ 0.05, ****p < 0.0001 Panel A created with BioRender.com



Supplemental Figure 4

Evidence for gene knockout (KO), *Adar1* sgRNA #2 results, and extended bead array data, relating to Figure 4.


(A) Heatmap of Log2 CPM expression of seven causal AGS genes in BMD and ER-Hoxb8 macrophages (B) Post-editing trace decomposition charts (via TIDE Analysis) for *Adar1* KO cells targeted with two distinct guides (C) ER-Hoxb8 cell counts over time (D) Immunostaining of in vitro, eight-day differentiated macrophages comparing control (NTC) and *Adar1* sgRNA #2-transduced macrophages (red = CD11B, blue = DAPI; scale bar = 100um) (E) Heatmap showing the Log2FC (*Adar1* KO values over NTC values) for relevant interferon-stimulated genes for both progenitors and macrophages at baseline (F) Validation of in vitro ER-Hoxb8 cell counts with vehicle or 10uM baricitinib treatment using *Adar1* sgRNA #2; statistics calculated at the 9 day time point (G) Interferon, cytokine and chemokine production for ER-Hoxb8 macrophages grown with or without baricitinib, via cytokine bead array (H) Trace decomposition charts (via TIDE Analysis) for *Adar1*/*Ifih1* double KO (dKO) cell lines (I) Interferon, cytokine and chemokine production for ER-Hoxb8 macrophages with or without subsequent *Ifih1* KO. All p-values calculated via one-way (I) or two-way (C, F, G) ANOVA with multiple comparisons; ns = not significant or $p \geq 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$



Supplemental Figure 5

Concurrent *Adar1*, *Ifih1* mutation morphology, plus engraftment in the *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} mouse, relating to Figure 5 [↗](#).


(A) Violin plot of all analyzed cortical ER-Hoxb8 cells (n=20) per animal (n=2), with yellow line denoting the median, for number of primary branches, average length of branches, longest branch length, and soma size; p-value calculated via nested t-test for n=2 animals per group; ns = not significant or p >= 0.05, **p < 0.005 (B) Schematic for in vivo *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} transplant experiments (C) Representative rendering of donor cell engraftment (scale bar = 1000μm) with inset microscopy of GFP+ donor cell engraftment (green = endogenous GFP, blue = DAPI; scale bar = 100μm) for control cells (TLR4 KO, NTC) harvested 15 days post-injection, (D) *Adar1* KO cells harvested seven days post-injection, (E) *Adar1/Ifih1* double KO (dKO) cells harvested 13-15 days post-injection, and (F) *Adar1* KO cells harvested three days post-injection Panel B created with BioRender.com [↗](#)



Supplemental Figure 6

Evidence for *Cre* expression, relating to Figure 6 [↗](#).

(A) Dual immunostaining/RNA in situ hybridization (ISH) showing IBA1 (red, protein stain), *Cre* (green, ISH), and nuclei (blue, DAPI); scale bar = 100um (B) magnified image of a repopulated microglia from the non-transplanted (tam sham control) *Cx3cr1*^{CreERT}; *Csf1r*^{fl/fl} brain shown in (A); scale bar = 10um



Supplemental Figure 7

Extended *Isg15* RNA in situ hybridization (ISH), relating to Figure 6.

(A) Dual immunostaining/RNA ISH showing IBA1 (red, protein stain), *Isg15* (purple, ISH), and nuclei (teal, DAPI); scale bar = 100um; white arrow depicts location of corresponding close up image; scale bar = 20um (B) Representative rendering of donor cell engraftment in the *Csf1r*^{-/-} brain (scale bar = 1000um) with inset microscopy of IBA1+ donor cell engraftment (red = IBA1, blue = DAPI; scale bar = 100um) for FVB Osb-GFP WT ER-Hoxb8s and *Adar1* D1113H mutant ER-Hoxb8s harvested 13-14 days post-injection (C) Dual immunostaining/RNA ISH for samples shown in (B); tile scale bar = 1000um; inset = *Isg15* (pink, ISH), nuclei (blue, DAPI); scale bar = 100um

References

- Abud Edsel M., Ramirez Ricardo N., Martinez Eric S., Healy Luke M., Nguyen Cecilia H. H., Newman Sean A., Yeromin Andriy V., et al. (2017) **iPSC-Derived Human Microglia-like Cells to Study Neurological Diseases** *Neuron* **94**:278–93 [Google Scholar](#)
- Accarias Solene, Sanchez Thibaut, Labrousse Arnaud, Ben-Neji Myriam, Boyance Aurélien, Poincloux Renaud, Maridonneau-Parini Isabelle, Le Cabec Véronique (2020) **Genetic Engineering of Hoxb8-Immortalized Hematopoietic Progenitors - a Potent Tool to Study Macrophage Tissue Migration** *Journal of Cell Science* **133**:5 <https://doi.org/10.1242/jcs.236703> | [Google Scholar](#)
- Andrews S. (2014) **FastQC a Quality-Control Tool for High-Throughput Sequence Data** <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Arshadi Cameron, Günther Ulrik, Eddison Mark, Harrington Kyle I. S., Ferreira Tiago A. (2021) **SNT: A Unifying Toolbox for Quantification of Neuronal Anatomy** *Nature Methods* **18**:374–77 [Google Scholar](#)
- Bailis Will, Shyer Justin A., Zhao Jun, Carlos Garcia Canaveras Juan, Al Khazal Fatimah J., Qu Rihao, Steach Holly R., et al. (2019) **Distinct Modes of Mitochondrial Metabolism Uncouple T Cell Differentiation and Function** *Nature* **571**:403–7 [Google Scholar](#)
- Balcaitis Stephanie, Weinstein Jonathan R., Li Sheng, Chamberlain Jeffrey S., Möller Thomas (2005) **Lentiviral Transduction of Microglial Cells** *Glia* **50**:48–55 [Google Scholar](#)
- Behrendt Rayk, Schumann Tina, Gerbaulet Alexander, Nguyen Laura A., Schubert Nadja, Alexopoulou Dimitra, Berka Ursula, et al. (2013) **Mouse SAMHD1 Has Antiretroviral Activity and Suppresses a Spontaneous Cell-Intrinsic Antiviral Response** *Cell Reports* **4**:689–96 [Google Scholar](#)
- Bennett F. Chris, Bennett Mariko L., Yaqoob Fazeela, Mulinyawe Sara B., Grant Gerald A., Gephart Melanie Hayden, Plowey Edward D., Barres Ben A. (2018) **A Combination of Ontogeny and CNS Environment Establishes Microglial Identity** *Neuron* **98**:1170–83 [Google Scholar](#)
- Bennett Mariko L., Chris Bennett F. (2020) **The Influence of Environment and Origin on Brain Resident Macrophages and Implications for Therapy** *Nature Neuroscience* **23**:157–66 [Google Scholar](#)
- Bennett Mariko L., Chris Bennett F., Liddel Shaw Shane A., Ajami Bahareh, Zamanian Jennifer L., Fernhoff Nathaniel B., Mulinyawe Sara B., et al. (2016) **New Tools for Studying Microglia in the Mouse and Human CNS** *Proceedings of the National Academy of Sciences of the United States of America* **113**:E1738–46 [Google Scholar](#)
- Blasi E., Barluzzi R., Bocchini V., Mazzolla R., Bistoni F. (1990) **Immortalization of Murine Microglial Cells by a v-Raf/v-Myc Carrying Retrovirus** *Journal of Neuroimmunology* **27**:229–37 [Google Scholar](#)

Bohlen Christopher J., Chris Bennett F., Tucker Andrew F., Collins Hannah Y., Mulinyawe Sara B., Barres Ben A. (2017) **Diverse Requirements for Microglial Survival, Specification, and Function Revealed by Defined-Medium Cultures** *Neuron* **94**:759–73 [Google Scholar](#)

Bray Nicolas L., Pimentel Harold, Melsted Páll, Pachter Lior (2016) **Near-Optimal Probabilistic RNA-Seq Quantification** *Nature Biotechnology* **34**:525–27 [Google Scholar](#)

Brinkman Eva K., Chen Tao, Amendola Mario, van Steensel Bas (2014) **Easy Quantitative Assessment of Genome Editing by Sequence Trace Decomposition** *Nucleic Acids Research* **42**:e168 [Google Scholar](#)

Bromberger Thomas, Klapproth Sarah, Sperandio Markus, Moser Markus (2022) **Humanized β 2 Integrin-Expressing Hoxb8 Cells Serve as Model to Study Integrin Activation** *Cells* **11** <https://doi.org/10.3390/cells11091532> | [Google Scholar](#)

Buttgereit Anne, Lelios Iva, Yu Xueyang, Vrohings Melissa, Krakoski Natalie R., Gautier Emmanuel L., Nishinakamura Ryuichi, Becher Burkhard, Greter Melanie (2016) **Sall1 Is a Transcriptional Regulator Defining Microglia Identity and Function** *Nature Immunology* **17**:1397–1406 [Google Scholar](#)

Chadarevian Jean Paul, Lombroso Sonia I., Peet Graham C., Hasselmann Jonathan, Tu Christina, Marzan Dave E., Capocchi Joia, et al. (2023) **Engineering an Inhibitor-Resistant Human CSF1R Variant for Microglia Replacement** *The Journal of Experimental Medicine* **220** <https://doi.org/10.1084/jem.20220857> | [Google Scholar](#)

Chanput Wasaporn, Mes Jurriaan J., Wichers Harry J. (2014) **THP-1 Cell Line: An in Vitro Cell Model for Immune Modulation Approach** *International Immunopharmacology* **23**:37–45 [Google Scholar](#)

Chen Shifu (2023) **Ultrafast One-Pass FASTQ Data Preprocessing, Quality Control, and Deduplication Using Fastp** *iMeta* **2**:e107 [Google Scholar](#)

Chen Shifu, Zhou Yanqing, Chen Yaru, Gu Jia (2018) **Fastp: An Ultra-Fast All-in-One FASTQ Preprocessor** *Bioinformatics* **34**:i884–90 [Google Scholar](#)

Chen Yunshun, Lun Aaron T. L., Smyth Gordon K. (2016) **From Reads to Genes to Pathways: Differential Expression Analysis of RNA-Seq Experiments Using Rsubread and the edgeR Quasi-Likelihood Pipeline** *F1000Research* **5**:1438 [Google Scholar](#)

Cronk James C., Filiano Anthony J., Louveau Antoine, Marin Ioana, Marsh Rachel, Ji Emily, Goldman Dylan H., et al. (2018) **Peripherally Derived Macrophages Can Engraft the Brain Independent of Irradiation and Maintain an Identity Distinct from Microglia** *The Journal of Experimental Medicine* **215**:1627–47 [Google Scholar](#)

Dai Xu-Ming, Ryan Gregory R., Hapel Andrew J., Dominguez Melissa G., Russell Robert G., Kapp Sara, Sylvestre Vonetta, Richard Stanley E. (2002) **Targeted Disruption of the Mouse Colony-Stimulating Factor 1 Receptor Gene Results in Osteopetrosis, Mononuclear Phagocyte Deficiency, Increased Primitive Progenitor Cell Frequencies, and Reproductive Defects** *Blood* **99**:111–20 [Google Scholar](#)

Demerdash Yasmin, Kain Bailee, Essers Marieke A. G., King Katherine Y. (2021) **Yin and Yang: The Dual Effects of Interferons on Hematopoiesis** *Experimental Hematology* **96**:1–12 [Google Scholar](#)

Douvaras Panagiotis, Sun Bruce, Wang Minghui, Kruglikov Ilya, Lallo Gregory, Zimmer Matthew, Terrenoire Cecile, et al. (2017) **Directed Differentiation of Human Pluripotent Stem Cells to Microglia** *Stem Cell Reports* **8**:1516–24 [Google Scholar](#)

Elhag Sara, Stremmel Christopher, Zehrer Annette, Plocke Josefine, Hennel Roman, Keuper Michaela, Knabe Clarissa, et al. (2021) **Differences in Cell-Intrinsic Inflammatory Programs of Yolk Sac and Bone Marrow Macrophages** *Cells* **10**:12 <https://doi.org/10.3390/cells10123564> | [Google Scholar](#)

Escoubas Caroline C., Dorman Leah C., Nguyen Phi T., Lagares-Linares Christian, Nakajo Haruna, Anderson Sarah R., Barron Jerika J., et al. (2024) **Type-I-Interferon-Responsive Microglia Shape Cortical Development and Behavior** *Cell* **187**:1936–54 [Google Scholar](#)

Ewels Philip, Magnusson Måns, Lundin Sverker, Käller Max (2016) **MultiQC: Summarize Analysis Results for Multiple Tools and Samples in a Single Report** *Bioinformatics* **32**:3047–48 [Google Scholar](#)

Fites J. Scott, et al (2018) **An Unappreciated Role for Neutrophil-DC Hybrids in Immunity to Invasive Fungal Infections** *PLoS Pathogens* **14**:e1007073 [Google Scholar](#)

Flannagan Ronald S., Canton Johnathan, Furuya Wendy, Glogauer Michael, Grinstein Sergio (2014) **The Phosphatidylserine Receptor TIM4 Utilizes Integrins as Coreceptors to Effect Phagocytosis** *Molecular Biology of the Cell* **25**:1511–22 [Google Scholar](#)

Gavazzi Francesco, Gonzalez Carlos Dominguez, Arnold Kaley, Swantkowski Meghan, Charlton Lauren, Modesti Nicholson, Dar Asif A., Vanderver Adeline, Bennett Mariko, Adang Laura A. (2024) **Nucleotide Metabolism, Leukodystrophies, and CNS Pathology** *Journal of Inherited Metabolic Disease* <https://doi.org/10.1002/jimd.12721> | [Google Scholar](#)

Gosselin David, Skola Dylan, Coufal Nicole G., Holtman Inge R., Schlachetzki Johannes C. M., Sajti Eniko, Jaeger Baptiste N., et al. (2017) **An Environment-Dependent Transcriptional Network Specifies Human Microglia Identity** *Science* **356** <https://doi.org/10.1126/science.aal3222> | [Google Scholar](#)

Gran Sandra, Honold Lisa, Fehler Olesja, Zenker Stefanie, Eligehausen Sarah, Kuhlmann Michael T., Geven Edwin, et al. (2018) **Imaging, Myeloid Precursor Immortalization, and Genome Editing for Defining Mechanisms of Leukocyte Recruitment in Vivo** *Theranostics* **8**:2407–23 [Google Scholar](#)

Guo Xinfeng, Steinman Richard A., Sheng Yi, Cao Guodong, Wiley Clayton A., Wang Qingde (2022) **An AGS-Associated Mutation in ADAR1 Catalytic Domain Results in Early-Onset and MDA5-Dependent Encephalopathy with IFN Pathway Activation in the Brain** *Journal of Neuroinflammation* **19**:285 [Google Scholar](#)

Gurzeler U., Rabachini T., Dahinden C. A., Salmanidis M., Brumatti G., Ekert P. G., Echeverry N., Bachmann D., Simon H. U., Kaufmann T. (2013) **In Vitro Differentiation of near-Unlimited Numbers of Functional Mouse Basophils Using Conditional Hoxb8** *Allergy* **68**:604–13 [Google Scholar](#)

Haenseler Walther, Sansom Stephen N., Buchrieser Julian, Newey Sarah E., Moore Craig S., Nicholls Francesca J., Chintawar Satyan, et al. (2017) **A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a Neuronal-Co-Culture-Specific Expression Profile and Inflammatory Response** *Stem Cell Reports* **8**:1727–42 [Google Scholar](#)

Han Velda X., Mohammad Shekeeb S., Jones Hannah F., Bandodkar Sushil, Crow Yanick J., Dale Russell C., AGS-JAKi Study Group (2022) **Cerebrospinal Fluid Neopterin as a Biomarker of Treatment Response to Janus Kinase Inhibition in Aicardi-Goutières Syndrome** *Developmental Medicine and Child Neurology* **64**:266–71 [Google Scholar](#)

Hartner Jochen C., Schmittwolf Carolin, Kispert Andreas, Müller Albrecht M., Higuchi Miyoko, Seeburg Peter H. (2004) **Liver Disintegration in the Mouse Embryo Caused by Deficiency in the RNA-Editing Enzyme ADAR1** *The Journal of Biological Chemistry* **279**:4894–4902 [Google Scholar](#)

Hasselmann Jonathan, Coburn Morgan A., England Whitney, Figueroa Velez Dario X., Shabestari Sepideh Kiani, Tu Christina H., McQuade Amanda, et al. (2019) **Development of a Chimeric Model to Study and Manipulate Human Microglia In Vivo** *Neuron* **103**:1016–33 [Google Scholar](#)

Hiller Bjoern, Achleitner Martin, Glage Silke, Naumann Ronald, Behrendt Rayk, Roers Axel (2012) **Mammalian RNase H2 Removes Ribonucleotides from DNA to Maintain Genome Integrity** *The Journal of Experimental Medicine* **209**:1419–26 [Google Scholar](#)

Jiang Xiao-Shuang, Ni Ying-Qin, Liu Tian-Jin, Zhang Meng, Ren Hui, Jiang Rui, Huang Xin, Xu Ge-Zhi (2012) **CCR2 Overexpression Promotes the Efficient Recruitment of Retinal Microglia in Vitro** *Molecular Vision* **18**:2982–92 [Google Scholar](#)

Kanazawa Nobuo, Ishii Taeko, Takita Yasushi, Nishikawa Atsushi, Nishikomori Ryuta (2023) **Efficacy and Safety of Baricitinib in Japanese Patients with Autoinflammatory Type I Interferonopathies (NNS/CANDLE, SAVI, And AGS)** *Pediatric Rheumatology Online Journal* **21**:38 [Google Scholar](#)

Li Qingyun, Barres Ben A. (2018) **Microglia and Macrophages in Brain Homeostasis and Disease** *Nature Reviews. Immunology* **18**:225–42 [Google Scholar](#)

Liddicoat Brian J., Piskol Robert, Chalk Alistair M., Ramaswami Gokul, Higuchi Miyoko, Hartner Jochen C., Li Jin Billy, Seeburg Peter H., Walkley Carl R. (2015) **RNA Editing by ADAR1 Prevents MDA5 Sensing of Endogenous dsRNA as Nonself** *Science* **349**:1115–20 [Google Scholar](#)

Ma Guangxin, Gezer Deniz, Herrmann Oliver, Feldberg Kristina, Schemionek Mirle, Jawhar Mohamad, Reiter Andreas, Brümmendorf Tim H., Koschmieder Steffen, Chatain Nicolas (2020) **LCP1 Triggers mTORC2/AKT Activity and Is Pharmacologically Targeted by Enzastaurin in Hypereosinophilia** *Molecular Carcinogenesis* **59**:87–103 [Google Scholar](#)

Mannion Niamh M., Greenwood Sam M., Young Robert, Cox Sarah, Brindle James, Read David, Nellåker Christoffer, et al. (2014) **The RNA-Editing Enzyme ADAR1 Controls Innate Immune Responses to RNA** *Cell Reports* **9**:1482–94 [Google Scholar](#)

Masuda Takahiro, Tsuda Makoto, Tozaki-Saitoh Hidetoshi, Inoue Kazuhide (2013) **Lentiviral Transduction of Cultured Microglia** *Methods in Molecular Biology* **1041**:63–67 [Google Scholar](#)

McCarthy Davis J., Chen Yunshun, Smyth Gordon K. (2012) **Differential Expression Analysis of Multifactor RNA-Seq Experiments with Respect to Biological Variation** *Nucleic Acids*

Research 40:4288–97 [Google Scholar](#)

Möller Alexander, Jauch-Speer Saskia-Larissa, Gandhi Shrey, Vogl Thomas, Roth Johannes, Fehler Olesja (2023) **The Roles of Toll-like Receptor 4, CD33, CD68, CD69, or CD147/EMMPRIN for Monocyte Activation by the DAMP S100A8/S100A9** *Frontiers in Immunology* 14:1110185 [Google Scholar](#)

Morita Masashi, Stamp Gordon, Robins Peter, Dulic Anna, Rosewell Ian, Hrivnak Geza, Daly Graham, Lindahl Tomas, Barnes Deborah E. (2004) **Gene-Targeted Mice Lacking the Trex1 (DNase III) 3'→5' DNA Exonuclease Develop Inflammatory Myocarditis** *Molecular and Cellular Biology* 24:6719–27 [Google Scholar](#)

Muffat Julien, Li Yun, Yuan Bingbing, Mitalipova Maisam, Omer Attya, Corcoran Sean, Bakiasi Grisilda, et al. (2016) **Efficient Derivation of Microglia-like Cells from Human Pluripotent Stem Cells** *Nature Medicine* 22:1358–67 [Google Scholar](#)

Murray Peter J., Allen Judith E., Biswas Subhra K., Fisher Edward A., Gilroy Derek W., Goerdts Sergij, Gordon Siamon, et al. (2014) **Macrophage Activation and Polarization: Nomenclature and Experimental Guidelines** *Immunity* 41:14–20 [Google Scholar](#)

O'Brien Carleigh A., Chris Bennett F., Bennett Mariko L. (2022) **Microglia in Antiviral Immunity of the Brain and Spinal Cord** *Seminars in Immunology* 60:101650 [Google Scholar](#)

Ohto Taisuke, Tayeh Ahmed Abu, Nishikomori Ryuta, Abe Hiroto, Hashimoto Kyota, Baba Shiro, Arias-Loza Anahi-Paula, et al. (2022) **Intracellular Virus Sensor MDA5 Mutation Develops Autoimmune Myocarditis and Nephritis** *Journal of Autoimmunity* 127:102794 [Google Scholar](#)

Pandya Hetal, Shen Michael J., Ichikawa David M., Sedlock Andrea B., Choi Yong, Johnson Kory R., Kim Gloria, et al. (2017) **Differentiation of Human and Murine Induced Pluripotent Stem Cells to Microglia-like Cells** *Nature Neuroscience* 20:753–59 [Google Scholar](#)

Pereira-Lopes Selma, Celhar Teja, Sans-Fons Gloria, Serra Maria, Fairhurst Anna-Marie, Lloberas Jorge, Celada Antonio (2013) **The Exonuclease Trex1 Restrains Macrophage Proinflammatory Activation** *Journal of Immunology* 191:6128–35 [Google Scholar](#)

Priller J., Flügel A., Wehner T., Boentert M., Haas C. A., Prinz M., Fernández-Klett F., et al. (2001) **Targeting Gene-Modified Hematopoietic Cells to the Central Nervous System: Use of Green Fluorescent Protein Uncovers Microglial Engraftment** *Nature Medicine* 7:1356–61 [Google Scholar](#)

R Core Team (2022) **R: A Language and Environment for Statistical Computing** Vienna, Austria: Foundation for Statistical Computing <https://www.R-project.org/>

Raschke W. C., Baird S., Ralph P., Nakoinz I. (1978) **Functional Macrophage Cell Lines Transformed by Abelson Leukemia Virus** *Cell* 15:261–67 [Google Scholar](#)

Redecke Vanessa, Wu Ruiqiong, Zhou Jingran, Finkelstein David, Chaturvedi Vandana, High Anthony A., Häcker Hans (2013) **Hematopoietic Progenitor Cell Lines with Myeloid and Lymphoid Potential** *Nature Methods* 10:795–803 [Google Scholar](#)

Rehwinkel Jan, Maelfait Jonathan, Bridgeman Anne, Rigby Rachel, Hayward Bruce, Liberatore Rachel A., Bieniasz Paul D., et al. (2013) **SAMHD1-Dependent Retroviral Control and Escape in Mice** *The EMBO Journal* **32**:2454–62 [Google Scholar](#)

Rice Gillian I., Del Toro Duany Yoandris, Jenkinson Emma M., Forte Gabriella Ma, Anderson Beverley H., Ariaudo Giada, Bader-Meunier Brigitte, et al. (2014) **Gain-of-Function Mutations in IFIH1 Cause a Spectrum of Human Disease Phenotypes Associated with Upregulated Type I Interferon Signaling** *Nature Genetics* **46**:503–9 [Google Scholar](#)

Righi M., Mori L., De Libero G., Sironi M., Biondi A., Mantovani A., Donini S. D., Ricciardi-Castagnoli P. (1989) **Monokine Production by Microglial Cell Clones** *European Journal of Immunology* **19**:1443–48 [Google Scholar](#)

Ritchie Matthew E., Belinda Phipson Di Wu, Hu Yifang, Law Charity W., Shi Wei, Smyth Gordon K. (2015) **Limma Powers Differential Expression Analyses for RNA-Sequencing and Microarray Studies** *Nucleic Acids Research* **43**:e47 [Google Scholar](#)

Roberts Allison W., Popov Lauren M., Mitchell Gabriel, Ching Krystal L., Licht Daniel J., Golovkine Guillaume, Barton Gregory M., Cox Jeffery S. (2019) **Cas9+ Conditionally-Immortalized Macrophages as a Tool for Bacterial Pathogenesis and beyond** *eLife* **8**:e45957 <https://doi.org/10.7554/eLife.45957> | [Google Scholar](#)

Robinson Mark D., McCarthy Davis J., Smyth Gordon K. (2010) **edgeR: A Bioconductor Package for Differential Expression Analysis of Digital Gene Expression Data** *Bioinformatics* **26**:139–40 [Google Scholar](#)

Rosas Marcela, Osorio Fabiola, Robinson Matthew J., Davies Luke C., Dierkes Nicola, Jones Simon A., Caetano Reis e Sousa, Taylor Philip R. (2011) **Hoxb8 Conditionally Immortalised Macrophage Lines Model Inflammatory Monocytic Cells with Important Similarity to Dendritic Cells** *European Journal of Immunology* **41**:356–65 [Google Scholar](#)

Salter Michael W., Stevens Beth (2017) **Microglia Emerge as Central Players in Brain Disease** *Nature Medicine* **23**:1018–27 [Google Scholar](#)

Schneider Valerie A., Graves-Lindsay Tina, Howe Kerstin, Bouk Nathan, Chen Hsiu-Chuan, Kitts Paul A., Murphy Terence D., et al. (2017) **Evaluation of GRCh38 and de Novo Haploid Genome Assemblies Demonstrates the Enduring Quality of the Reference Assembly** *Genome Research* **27**:849–64 [Google Scholar](#)

Shemer Anat, Grozovski Jonathan, Tay Tuan Leng, Tao Jenhan, Volaski Alon, Süß Patrick, Ardura-Fabregat Alberto, et al. (2018) **Engrafted Parenchymal Brain Macrophages Differ from Microglia in Transcriptome, Chromatin Landscape and Response to Challenge** *Nature Communications* **9**:5206 [Google Scholar](#)

Shen Zeyang, Li Rick Z., Prohaska Thomas A., Hoeksema Marten A., Spann Nathan J., Tao Jenhan, Fonseca Gregory J., et al. (2022) **Systematic Analysis of Naturally Occurring Insertions and Deletions That Alter Transcription Factor Spacing Identifies Tolerant and Sensitive Transcription Factor Pairs** *eLife* **11**:e70878 <https://doi.org/10.7554/eLife.70878> | [Google Scholar](#)

Soneson Charlotte, Love Michael I., Robinson Mark D. (2015) **Differential Analyses for RNA-Seq: Transcript-Level Estimates Improve Gene-Level Inferences** *F1000Research* **4**:1521 [Google Scholar](#)

Stanley E. R., Heard P. M. (1977) **Factors Regulating Macrophage Production and Growth. Purification and Some Properties of the Colony Stimulating Factor from Medium Conditioned by Mouse L Cells** *The Journal of Biological Chemistry* **252**:4305–12 [Google Scholar](#)

Su Wei, Kang John, Sopher Bryce, Gillespie James, Aloï Macarena S., Odom Guy L., Hopkins Stephanie, et al. (2016) **Recombinant Adeno-Associated Viral (rAAV) Vectors Mediate Efficient Gene Transduction in Cultured Neonatal and Adult Microglia** *Journal of Neurochemistry* **136**:49–62 [Google Scholar](#)

Takata Kazuyuki, Kozaki Tatsuya, Zhe Wei Lee Christopher, Thion Morgane Sonia, Otsuka Masayuki, Lim Shawn, Utami Kagistia Hana, et al. (2020) **Induced-Pluripotent-Stem-Cell-Derived Primitive Macrophages Provide a Platform for Modeling Tissue-Resident Macrophage Differentiation and Function** *Immunity* **52**:417–18 [Google Scholar](#)

Timmerman Raissa, Burm Saskia M., Bajramovic Jeffrey J. (2018) **An Overview of in Vitro Methods to Study Microglia** *Frontiers in Cellular Neuroscience* **12**:242 [Google Scholar](#)

Tsuchiya S., Yamabe M., Yamaguchi Y., Kobayashi Y., Konno T., Tada K. (1980) **Establishment and Characterization of a Human Acute Monocytic Leukemia Cell Line (THP-1)** *International Journal of Cancer. Journal International Du Cancer* **26**:171–76 [Google Scholar](#)

Vanderver Adeline, Adang Laura, Gavazzi Francesco, McDonald Katherine, Helman Guy, Frank David B., Jaffe Nicole, et al. (2020) **Janus Kinase Inhibition in the Aicardi-Goutières Syndrome** *The New England Journal of Medicine* **383**:986–89 [Google Scholar](#)

Wang Gang G., Calvo Katherine R., Pasillas Martina P., Sykes David B., Häcker Hans, Kamps Mark P. (2006) **Quantitative Production of Macrophages or Neutrophils Ex Vivo Using Conditional Hoxb8** *Nature Methods* **3**:287–93 [Google Scholar](#)

Ward Simone V., George Cyril X., Welch Megan J., Liou Li-Ying, Hahm Bumsuk, Lewicki Hanna, de la Torre Juan C., Samuel Charles E., Oldstone Michael B. (2011) **RNA Editing Enzyme Adenosine Deaminase Is a Restriction Factor for Controlling Measles Virus Replication That Also Is Required for Embryogenesis** *Proceedings of the National Academy of Sciences* **108**:331–36 [Google Scholar](#)

Williamson Adam P., Vale Ronald D. (2018) **Spatial Control of Draper Receptor Signaling Initiates Apoptotic Cell Engulfment** *The Journal of Cell Biology* **217**:3977–92 [Google Scholar](#)

Xu Jane Jialu, Chalk Alistair M., Nikolic Iva, Simpson Kaylene J., Smeets Monique F., Walkley Carl R. (2022) **Genome-Wide Screening Identifies Cell-Cycle Control as a Synthetic Lethal Pathway with SRSF2P95H Mutation** *Blood Advances* **6**:2092–2106 [Google Scholar](#)

XuFeng Richard, Boyer Matthew J., Shen Hongmei, Li Yanxin, Yu Hui, Gao Yindai, Yang Qiong, Wang Qingde, Cheng Tao (2009) **ADAR1 Is Required for Hematopoietic Progenitor Cell Survival via RNA Editing** *Proceedings of the National Academy of Sciences of the United States of America* **106**:17763–68 [Google Scholar](#)

Zach Frank, Mueller Alexandra, Gessner André (2015) **Production and Functional Characterization of Murine Osteoclasts Differentiated from ER-Hoxb8-Immortalized Myeloid Progenitor Cells** *PloS One* **10**:e0142211 [Google Scholar](#)

Zerbino Daniel R., Achuthan Premanand, Akanni Wasiu, Ridwan Amode M., Barrell Daniel, Bhai Jyothish, Billis Konstantinos, et al. (2018) **Ensembl 2018** *Nucleic Acids Research* **46**:D754–61 [Google Scholar](#)

Zhang Ye, Chen Kenian, Sloan Steven A., Bennett Mariko L., Scholze Anja R., O’Keeffe Sean, Phatnani Hemali P., et al. (2014) **An RNA-Sequencing Transcriptome and Splicing Database of Glia, Neurons, and Vascular Cells of the Cerebral Cortex** *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* **34**:11929–47 [Google Scholar](#)

Zheng Lian-Shun, Kaneko Naoko, Sawamoto Kazunobu (2015) **Minocycline Treatment Ameliorates Interferon-Alpha-Induced Neurogenic Defects and Depression-like Behaviors in Mice** *Frontiers in Cellular Neuroscience* **9**:5 [Google Scholar](#)

Author information

Kelsey M Nemec

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States, Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States
ORCID iD: [0000-0003-3036-254X](#)

Genevieve Uy

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States, Division of Neurology, Children’s Hospital of Philadelphia, Philadelphia, United States

V Sai Chaluvadi

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States, Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

Freddy S Purnell

Department of Biology, School of Arts and Sciences, University of Pennsylvania, Philadelphia, United States, Epigenetics Institute, University of Pennsylvania, Perelman School of Medicine, Philadelphia, United States

Bilal Elfayoumi

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

Leila Byerly

Department of Biology, Bryn Mawr College, Bryn Mawr, United States

Micaela L O’Reilly

Division of Neurology, Children’s Hospital of Philadelphia, Philadelphia, United States

Carleigh A O’Brien

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

ORCID iD: [0000-0003-4549-0640](https://orcid.org/0000-0003-4549-0640)

William H Aisenberg

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

Sonia I Lombroso

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States, Department of Systems Pharmacology and Translational Therapeutics, University of Pennsylvania, Philadelphia, United States

Xinfeng Guo

Department of Surgery, University of Pittsburgh School of Medicine, Pittsburgh, United States

Niklas Blank

Department of Microbiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States, Faculty of Biology, University of Freiburg, Freiburg, Germany

Chet Huan Oon

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

Fazeela Yaqoob

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

Brian Temsamrit

Division of Neurology, Children's Hospital of Philadelphia, Philadelphia, United States

Priyanka Rawat

Division of Neurology, Children's Hospital of Philadelphia, Philadelphia, United States

Christoph A Thaiss

Department of Microbiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States

Will Bailis

Department of Pathology and Laboratory Medicine, Children's Hospital of Philadelphia, Philadelphia, United States, Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States
ORCID iD: [0000-0001-9420-6250](https://orcid.org/0000-0001-9420-6250)

Adam P Williamson

Department of Biology, Bryn Mawr College, Bryn Mawr, United States

Qingde Wang

Department of Surgery, University of Pittsburgh School of Medicine, Pittsburgh, United States
ORCID iD: [0000-0002-2482-7972](https://orcid.org/0000-0002-2482-7972)

Mariko L Bennett*

Division of Neurology, Children's Hospital of Philadelphia, Philadelphia, United States, Department of Neurology, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, United States
ORCID iD: [0000-0003-2570-0620](https://orcid.org/0000-0003-2570-0620)

*Senior authors

F Chris Bennett*

Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States, Division of Neurology, Children's Hospital of Philadelphia, Philadelphia, United States

For correspondence: frederick.bennett@pennmedicine.upenn.edu

*Senior authors

Editors

Reviewing Editor

Florent Ginhoux

Singapore Immunology Network, Singapore, Singapore

Senior Editor

Betty Diamond

The Feinstein Institute for Medical Research, Manhasset, United States of America

Reviewer #1 (Public review):

Summary:

Aicardi-Goutières Syndrome (AGS) is a genetic disorder that primarily affects the brain and immune system through excessive interferon production. The authors sought to investigate the role of microglia in AGS by first developing bone-marrow-derived progenitors in vitro that carry the estrogen-regulated (ER) Hoxb8 cassette, allowing them to expand indefinitely in the presence of estrogen and differentiate into macrophages when estrogen is removed. When injected into the brains of Csf1r^{-/-} mice, which lack microglia, these cells engraft and resemble wild-type (WT) microglia in transcriptional and morphological characteristics, although they lack Sall1 expression. The authors then generated CRISPR-Cas9 Adar1 knockout (KO) ER-Hoxb8 macrophages, which exhibited increased production of inflammatory cytokines and upregulation of interferon-related genes. This phenotype could be rescued using a Jak-Stat inhibitor or by concurrently mutating Ifih1 (Mda5). However, these Adar1-KO macrophages fail to successfully engraft in the brain of both Csf1r^{-/-} and Cx3cr1-creERT2:Csf1r^{fl/fl} mice. To overcome this, the authors used a mouse model with a patient-specific Adar1 mutation (Adar1 D1113H) to derive ER-Hoxb8 bone marrow progenitors and macrophages. They discovered that Adar1 D1113H ER-Hoxb8 macrophages successfully engraft the brain, although at lower levels than WT-derived ER-Hoxb8 macrophages, leading to increased production of Isg15 by neighboring cells. These findings shed new light on the role of microglia in AGS pathology.

Strengths:

The authors convincingly demonstrate that ER-Hoxb8 differentiated macrophages are transcriptionally and morphologically similar to bone marrow-derived macrophages. They also show evidence that when engrafted in vivo, ER-Hoxb8 microglia are transcriptomically similar to WT microglia. Furthermore, ER-Hoxb8 macrophages engraft the Csf1r^{-/-} brain with high efficiency and rapidly (2 weeks), showing a homogenous distribution. The authors also effectively use CRISPR-Cas9 to knock out TLR4 in these cells with little to no effect on their

engraftment in vivo, confirming their potential as a model for genetic manipulation and in vivo microglia replacement.

Overall, this paper demonstrates an innovative approach to manipulating microglia using ER-Hoxb8 cells as surrogates. The authors present convincing evidence of the model's efficacy and potential for broader application in microglial research, given its ease of production and rapid brain engraftment potential in microglia-deficient mice. Using mouse-derived cells for transplantation reduces complications that can come with the use of human cell lines, highlighting the utility of this system for research in mouse models.

<https://doi.org/10.7554/eLife.102900.2.sa2>

Reviewer #2 (Public review):

Summary:

Microglia have been implicated in brain development, homeostasis, and diseases. "Microglia replacement" has gained traction in recent years, using primary microglia, bone marrow or blood-derived myeloid cells, or human iPSC-induced microglia. Here, the authors extended their previous work in the area and provide evidence to support: (1) Estrogen-regulated (ER) homeobox B8 (Hoxb8) conditionally immortalized macrophages from bone marrow can serve as stable, genetically manipulated cell lines. These cells are highly comparable to primary bone marrow-derived (BMD) macrophages in vitro, and, when transplanted into a microglia-free brain, engraft the parenchyma and differentiate into microglia-like cells (MLCs). Taking advantage of this model system, the authors created stable, Adar1-mutated ER-Hoxb8 lines using CRISPR-Cas9 to study the intrinsic contribution of macrophages to Aicardi-Goutières Syndrome (AGS) disease mechanism.

Strengths:

The studies are carefully designed and well-conducted. The imaging data and gene expression analysis are carried out at a high level of technical competences and the studies provide strong evidence that ER-Hoxb8 immortalized macrophages from bone marrow are a reasonable source for "microglia replacement" exercise. The findings are clearly presented, and the main message will be of general interest to the neuroscience and microglia communities.

<https://doi.org/10.7554/eLife.102900.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

Summary:

Aicardi-Goutières Syndrome (AGS) is a genetic disorder that primarily affects the brain and immune system through excessive interferon production. The authors sought to investigate the role of microglia in AGS by first developing bone-marrow-derived progenitors in vitro that carry the estrogen-regulated (ER) Hoxb8 cassette, allowing them to expand indefinitely in the presence of estrogen and differentiate into macrophages when estrogen is removed. When injected into the brains of Csf1r^{-/-} mice, which lack

microglia, these cells engraft and resemble wild-type (WT) microglia in transcriptional and morphological characteristics, although they lack Sall1 expression. The authors then generated CRISPR-Cas9 Adar1 knockout (KO) ER-Hoxb8 macrophages, which exhibited increased production of inflammatory cytokines and upregulation of interferon-related genes. This phenotype could be rescued using a Jak-Stat inhibitor or by concurrently mutating Ifih1 (Mda5). However, these Adar1-KO macrophages fail to successfully engraft in the brain of both Csf1r^{-/-} and Cx3cr1-creERT2:Csf1r^{fl/fl} mice. To overcome this, the authors used a mouse model with a patient-specific Adar1 mutation (Adar1 D1113H) to derive ER-Hoxb8 bone marrow progenitors and macrophages. They discovered that Adar1 D1113H ER-Hoxb8 macrophages successfully engraft the brain, although at lower levels than WT-derived ER-Hoxb8 macrophages, leading to increased production of Isg15 by neighboring cells. These findings shed new light on the role of microglia in AGS pathology.

Strengths:

The authors convincingly demonstrate that ER-Hoxb8 differentiated macrophages are transcriptionally and morphologically similar to bone marrow-derived macrophages. They also show evidence that when engrafted in vivo, ER-Hoxb8 microglia are transcriptomically similar to WT microglia. Furthermore, ER-Hoxb8 macrophages engraft the Csf1r^{-/-} brain with high efficiency and rapidly (2 weeks), showing a homogenous distribution. The authors also effectively use CRISPR-Cas9 to knock out TLR4 in these cells with little to no effect on their engraftment in vivo, confirming their potential as a model for genetic manipulation and in vivo microglia replacement.

Weaknesses:

The robust data showing the quality of this model at the transcriptomic level can be strengthened with confirmation at protein and functional levels. The authors were unable to investigate the effects of Adar1-KO using ER-Hoxb8 cells and instead had to rely on a mouse model with a patient-specific Adar1 mutation (Adar1 D1113H). Additionally, ER-Hoxb8-derived microglia do not express Sall1, a key marker of microglia, which limits their fidelity as a full microglial replacement, as has been rightfully pointed out in the discussion.

Overall, this paper demonstrates an innovative approach to manipulating microglia using ER-Hoxb8 cells as surrogates. The authors present convincing evidence of the model's efficacy and potential for broader application in microglial research, given its ease of production and rapid brain engraftment potential in microglia-deficient mice. While Adar1-KO macrophages do not engraft well, the success of TLR4-KO line highlights the model's potential for investigating other genes. Using mouse-derived cells for transplantation reduces complications that can come with the use of human cell lines, highlighting the utility of this system for research in mouse models.

Thank you for this thoughtful and balanced assessment. The major suggestion from Reviewer 1 was that confirmation of RNAseq data with protein or functional studies would add strength. We provided protein staining by IHC for IBA1 in vivo, as well as protein staining by FACS for CD11B, CD45, and TMEM119 in vitro and in vivo. For TLR4, we showed successful protein KO and blunted response to LPS (a TLR4 ligand) challenge, which we believe provides some protein and functional data to support the approach. To bolster these data, we added staining for P2RY12 on brain-engrafted ER-Hoxb8s.

Regarding the *Adar1* KO phenotypes showing non-engraftment. Because ADAR1 KO mice are embryonically lethal due to hematopoietic failure, we see the health impacts of *Adar1* KO on ER-Hoxb8s as a strength of the transplantation model, enabling the assessment of ADAR1 global function in macrophages and microglia-like cells without generation of a transgenic

mouse line. In addition, it was a surprise that the health impact occurs at the macrophage and not the progenitor stage, perhaps providing insight for future studies of ADAR1's role in hematopoiesis. Instead, we were able to show a significant impact of complete loss of *Adar1* on survival and engraftment, suggesting an important biological function of ADAR1. Macrophage-specific D1113H mutation, which affects part of the deaminase domain, shows that when the RNA deamination (but not the RNA binding) function of ADAR1 is disrupted, we find brain-wide interferonopathy. This is very exciting to our group and hopefully the community as astrocytes are thought to be a major driver of brain interferonopathy in patients with ADAR1 mutations. Instead, this suggests that disruption of brain macrophages is also a major contributor.

Reviewer #2 (Public review):

Summary:

Microglia have been implicated in brain development, homeostasis, and diseases. "Microglia replacement" has gained traction in recent years, using primary microglia, bone marrow or blood-derived myeloid cells, or human iPSC-induced microglia. Here, the authors extended their previous work in the area and provided evidence to support: (1)

Estrogen-regulated (ER) homeobox B8 (Hoxb8) conditionally immortalized macrophages from bone marrow can serve as stable, genetically manipulated cell lines. These cells are highly comparable to primary bone marrow-derived (BMD) macrophages in vitro, and, when transplanted into a microglia-free brain, engraft the parenchyma and differentiate into microglia-like cells (MLCs). Taking advantage of this model system, the authors created stable, Adar1-mutated ER-Hoxb8 lines using CRISPR-Cas9 to study the intrinsic contribution of macrophages to the Aicardi-Goutières Syndrome (AGS) disease mechanism.

Strengths:

The studies are carefully designed and well-conducted. The imaging data and gene expression analysis are carried out at a high level of technical competence and the studies provide strong evidence that ER-Hoxb8 immortalized macrophages from bone marrow are a reasonable source for "microglia replacement" exercise. The findings are clearly presented, and the main message will be of general interest to the neuroscience and microglia communities.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

This is an elegant study, demonstrating both the utility and limitations of ER-Hoxb8 technology as a surrogate model for microglia in vivo. The manuscript is well-designed and clearly written, but authors should consider the following suggestions:

(1) Validation of RNA hits at the protein level: To strengthen the comparison between ER-Hoxb8 macrophages and WT bone marrow-derived macrophages, validating several RNA hits at the protein level would be beneficial. As many of these hits are surface markers, flow cytometry could be employed for confirmation (e.g., Figure 1D, Figure 3E).

In vitro, we show protein levels by flow cytometry for CD11B (ITGAM) and CD45 (PTPRC; Figure 1C), as well as TMEM119 (Supplemental Figure 2A) and TLR4 (Supplemental Figure 3C/D). In vivo, we show TMEM119 protein levels by flow cytometry (Figure 3A), as well as their CD11B/CD45 pre-gates (Supplemental Figure 2C), plus immunostaining for IBA1 (AIF1; Figure 2D). We now provide additional data showing P2RY12 immunostaining in brain-engrafted cells (Supplemental Figure 2B).

(2) The authors should consider testing the phagocytic capacity of ER-Hoxb8-derived macrophages to further validate their functionality.

Thank you for the suggestion. We measured ER-Hoxb8 macrophage ability to engulf phosphatidylserine-coated beads that mimic apoptotic cells, compared with phosphatidylcholine-coated beads, now as new Supplemental Figure 1C/D. This agrees with existing literature showing efficient engulfment/phagocytosis by ER-Hoxb8-derived cells (Elhag et al., 2021).

(3) For Figure 3E, incorporating a wild-type (WT) microglia reference would be beneficial to establish a baseline for comparison (e.g. including WT microglia data in the graph or performing a ratio analysis against WT expression levels).

We agree - we now include bars representing our sequenced primary microglia data in Figure 3E as a comparison.

(4) Some statistical analyses may require refinement. Specifically, for Figure 4J, where the effects of Adar1 KO and Adar1 KO with Bari are compared, it would be more appropriate to use a two-way ANOVA.

Thank you for noting it. We have now done more appropriate two-way ANOVA and included the updated results in Figure 4J and the corresponding Supplemental Figure 4G. Errors in figure legend texts have also been corrected to reflect the statistical tests used.

(5) Cx3cr1-creERT2 pups injected with tamoxifen: The authors could clarify the depletion ratio in these experiments before the engraftment and assess whether the depletion is global or regional. In comparison to Csf1r-/-, where TLR4-KO ER-Hoxb8 engraft globally, in Cx3cr1-creERT2, the engraftment seems more regional (Figure 5A vs Supplementary Figure 5B); is this due to the differences in depletion efficiency?

This is an excellent question and observation, and one that we are very interested in, though that finding does not change the conclusions of this particular study. We find some region-specific differences in depletion early after tamoxifen injection, but that all brain regions are >95% depleted by P7. For instance, in a recently published manuscript (Bastos et al., 2025) we find some differences in the depletion kinetics in the genetic model. By P3, we find 90% depletion in cortex with 50-60% in thalamus and hippocampus. In other studies, we typically deliver primary monocytes, and this is the first study where we report engraftment of ER-Hoxb8 cells in the inducible model. In this sense, it is possible that depletion kinetics may regionally affect engraftment, but future studies are required to more finely assess this point with ER-Hoxb8s, as it may change how these models are used in the future.

Bastos et al., Monocytes can efficiently replace all brain macrophages and fetal liver monocytes can generate bonafide SALL1+ microglia, Immunity (2025), <https://doi.org/10.1016/j.immuni.2025.04.006>

(6) It would be helpful for the authors to clarify whether Adar1 is predominantly expressed by microglia, especially since the study aims to show its role in dampening the interferon response.

That's a wonderful point. *Adar1* is expressed by all brain cells, with highest transcript level in some neurons, astrocytes, and oligodendrocytes. It is an interferon-stimulated gene, and mutation itself leads to interferonopathy, we believe, due to poor RNA editing and detection of endogenous RNA as non-self by MDA5. We hope it can dampen the interferon response, but

in the case of mutation, *Adar1* is probably causal of interferonopathy. It is induced in microglia upon systemic inflammatory challenge (LPS). We have edited the text to highlight its expression pattern. See BrainRNAseq.org (Zhang*, Chen*, Sloan*, et al., 2014 and Bennett et al., 2016)

Reviewer #2 (Recommendations for the authors):

(1) There appears to be a morphological difference between wt and Adar1/Ifih1 double KO (dKO) cells in the engrafted brains (Figure 5). It would be good if the authors could systematically compare the morphology (e.g., soma size, number, and length of branches) of the engrafted MLCs between the wt and mutant cells.

We agree. While cells did not differ in branch number or length, engrafted dKO cells had significantly larger somas compared with controls, which we now present in Figure S5A.

(2) To fully appreciate the extent of how those engrafted ER-Hoxb8 immortalized macrophages resemble primary, engrafted yolk sac-myeloid cells, vs engrafted iPSC-induced microglia, it would be informative to provide a comparison of their RNAseq data derived from the engrafted ER-Hoxb8 immortalized macrophages with published data transcriptomic data sets (e.g. Bennett et al. Neuron 2018; Chadarevian et al. Neuron 2024; Schafer et al. Cell 2023).

Thank you for this suggestion. To address this, we provide our full dataset for additional experiments. To compare with a similar non-immortalized model, we compared top up- and down-regulated genes from our data to those of ICT yolk sac progenitor cells from our previous work (Bennett et al., 2018). We find overlap between brain-engrafted ER-Hoxb8-, bone marrow-, and yolk sac-derived cells (Supplemental Figure 2F, Supplemental Table 3).

Minor comments:

Figure 6C: red arrow showing zoom in regions are not matchable. It might be beneficial to provide bigger images with each channel for C and D as a Supplemental Figure.

We fixed this in Figure 6C to show areas of interest in the cortex for both conditions. Figure S7A shows intermediate power images to aid in interpretation.

<https://doi.org/10.7554/eLife.102900.2.sa0>