

DNA replication in primary hepatocytes without the six-subunit ORC

Reviewed Preprint

v2 • March 7, 2025

Revised by authors

Reviewed Preprint

v1 • December 31, 2024

Róża K Przanowska, Yuechuan Chen, Takayuki-Okano Uchida, Etsuko Shibata, Xiaoxiao Hao, Isaac Segura Rueda, Kate Jensen, Piotr Przanowski, Anthony Trimboli, Yoshiyuki Shibata, Gustavo Leone, Anindya Dutta 

Dept. of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, United States • Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States • Cancer Center, University of Wisconsin in Milwaukee, Milwaukee, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **valuable** descriptive manuscript builds on prior research showing that the elimination of Origin Recognition Complex (ORC) subunits does not halt DNA replication. The authors obtain **solid** data using various methods to genetically remove one or two ORC subunits from specific tissues and still observe replication. The replication appears to be primarily endoreduplication, indicating that ORC-independent replication may promote genome reduplication without mitosis. The mechanism behind this ORC-independent replication remains to be elucidated. The study and mutants described herein lay the groundwork for future research to explore how cells compensate for the absence of ORC and to develop functional approaches to investigate this process. The reviewers suggested the observations could be supported by additional experiments. This work will be of interest to those studying genome duplication and replication.

<https://doi.org/10.7554/eLife.102915.2.sa4>

Abstract

The six subunit ORC is essential for initiation of DNA replication in eukaryotes. Cancer cell-lines in culture can survive and replicate DNA replication after genetic inactivation of individual ORC subunits, ORC1, ORC2 or ORC5. In primary cells, ORC1 was dispensable in the mouse liver for endo-reduplication, but this could be explained by the ORC1 homolog, CDC6, substituting for ORC1 to restore functional ORC. Here, we have created mice with a conditional deletion of ORC2, which does not have a homolog. Although mouse embryo fibroblasts require ORC2 for proliferation, mouse hepatocytes synthesize DNA in cell culture and endo-reduplicate *in vivo* without ORC2. Mouse livers endo-reduplicate after simultaneous deletion of ORC1 and ORC2 both during normal development and after partial hepatectomy. Since endo-reduplication initiates DNA synthesis like normal S phase replication these results

unequivocally indicate that primary cells, like cancer cell lines, can load MCM2-7 and initiate replication without ORC.

Introduction

DNA replication during normal mitotic cycles or during endo-reduplication is initiated by the six subunit ORC, five of which have AAA+ATPase domains and form a ring shaped complex that binds DNA and bends it (Bell and Stillman, 1992 [↗](#), Neuwald et al., 1999 [↗](#), Dhar et al., 2001a [↗](#), Clarey et al., 2006 [↗](#), Bleichert et al., 2015 [↗](#)). In cooperation with another ATPase, CDC6, and with CDT1, ORC helps load MCM2-7 double hexamers at or near origins of replication (Costa and Diffley, 2022 [↗](#), Stillman, 2022 [↗](#), Hu and Stillman, 2023 [↗](#)). To initiate replication (or endo-reduplication), the MCM2-7 hexamer associates with additional proteins to form the functional CMG helicase that is essential for unwinding the double stranded chromosomal DNA, so that the resulting single-stranded DNAs can serve as templates for copying by DNA polymerases. In the lower eukaryotes, various ORC subunits have been consistently found to be essential for DNA replication and cell proliferation (Bell et al., 1993 [↗](#), Foss et al., 1993 [↗](#), Micklem et al., 1993 [↗](#), Loo et al., 1995 [↗](#), Semple et al., 2006 [↗](#)). In mammalian cells, knockdown of individual ORC subunits have also been reported to stop cell proliferation (Chou et al., 2021 [↗](#), Prasanth et al., 2004 [↗](#)). In humans, mutations in *Orc1*, *Orc4*, *Orc6*, *Cdt1*, and *Cdc6* cause microcephalic primordial dwarfism resembling Meier-Gorlin syndrome (Bicknell et al., 2011b [↗](#), Bicknell et al., 2011a [↗](#), Guernsey et al., 2011 [↗](#)).

There have, however, been isolated reports that suggest that under unusual circumstances, DNA replication can initiate in eukaryotes in the absence of the full ORC. First hypomorphic mutation made in the *Orc2* gene in HCT116 colon cancer cells decreased ORC2 protein levels by 90% and cells were still viable and could proliferate (Dhar et al., 2001b [↗](#)). Deletion of the *Orc1* gene in *Drosophila* permitted several rounds of replication in the resulting larvae and pupae (Park and Asano, 2008 [↗](#)). A few human cancer cell lines have been created by CRISPR-Cas9 mediated genome engineering where ORC1, ORC2 and ORC5 cannot be detected by regular immunoblots and yet the cell lines survive, proliferate and replicate DNA with the normal complement of origins of replication (Shibata et al., 2016 [↗](#), Shibata and Dutta, 2020 [↗](#)). However, the *Orc2* gene is essential for viability and in the *Orc2Δ* cell line a very minimal level (<0.1% of wild type levels) of a truncated protein can be detected that reacts with anti-ORC2 antibody and co-immunoprecipitates with ORC3 (Chou et al., 2021 [↗](#)). Given that WT cells have about 150,000 molecules of ORC2, even if this truncated protein is functional ORC2, ~150 molecules of the protein would be expected to load MCM2-7 double hexamers on at least 50,000 origins of replication. This possibility would be consistent with current models of MCM2-7 loading only if ORC was highly catalytic and one ORC hexamer was capable of loading ~667 MCM2-7 hexamers (Costa and Diffley, 2022 [↗](#), Stillman, 2022 [↗](#), Hu and Stillman, 2023 [↗](#)).

To address whether the entire ORC holocomplex is essential for DNA replication initiation in primary mammalian cells, we have reported a conditional mutation in the ORC1 gene of mouse (Okano-Uchida et al., 2018 [↗](#)). Conditional deletion of *Orc1* (using tissue-specific Cre drivers) revealed that the gene was essential for normal mitotic cell division in intestinal epithelial cells, but that endoreduplication in tissues that are known to become polyploid, like placental trophoblasts and the liver hepatocytes, was unaffected. Because the ORC1 protein is a functional ATPase with sequence homology with CDC6 (Saha et al., 1998 [↗](#), Williams et al., 1997 [↗](#)), one possibility that remained was that the mouse CDC6 protein substituted for ORC1 in the ORC holocomplex, producing enough functional ORC. Therefore, we decided to make a conditional mutation in another subunit of ORC, *Orc2*, that does not have the Walker A or B motifs that would make it a functional ATPase and that is not homologous to *Cdc6*. We designed the mutation such that even if an N terminally truncated protein is expressed from a downstream methionine from an alternately spliced mRNA the AAA+ like domain will suffer a significant deletion.

This will allow us to test whether the mouse liver survival and endoreduplication after loss of ORC1 is seen even after loss of ORC2. It will also test whether combining the conditional mutations of ORC1 and ORC2 will allow the liver cells to survive and endoreduplicate. If hepatocyte survival and endoreduplication is seen in these mice, then it would be hard to explain this by the substitution of two ORC subunits by the CDC6 protein, and we would gain support for the hypothesis that in rare circumstances even untransformed mammalian cells are capable of loading enough MCM2-7 in the absence of ORC to permit extensive DNA replication.

Results

ORC2 is essential for embryonic development and proliferation of mouse embryo fibroblasts

Mice with Lox P sites inserted flanking exons 6 and 7 of mouse *Orc2* were purchased from Cyagen Biosciences Inc. (Fig. 1A). Genotyping with primers F2 and R2 distinguished the loxP marked allele from the WT allele (Fig. 1B). In a cross of *Orc2^{f/+}* mice, there was no significant decrease in the yield of *Orc2^{f/+}* or *Orc2^{f/f}* compared to *Orc2^{+/+}* mice (Fig. 1C), suggesting that the insertion of the loxP sites in the introns of *Orc2* did not impair the function of ORC2. Recombination between the lox2 sites would delete the coding exons 6 and 7 of *Orc2*, which removes amino acids L111-V150 (Fig. 1D). If a transcript is expressed that skips exons 6 and 7, then the resulting protein mutates A110 to V110 and then throws the protein coding sequence out of frame (Fig. 1D, ΔORC2), so that the bulk of the 576 amino acid ORC2 protein, including the AAA+ like and WH domains (K230-A576) are not expressed. The AAA+ like domain and the WH domain are key elements for ORC2 assembly into ORC and for ORC function. Even in the remote possibility that a truncated protein is expressed due to alternative splicing and translation initiation from an internal methionine, the next methionine is at M301, so that half of the protein including 70 amino acids of the 239 amino acid AAA+-like domain (K230-G469) would be deleted.

The *Sox2-Cre* allele expresses active Cre during embryonic development. Crossing the *Orc2^{f/f}* mice with *Sox2-Cre* mice resulted in no *Orc2^{Δ/Δ}* embryos at E7.5 days onwards suggesting that embryonic deletion of *Orc2* is lethal (Table 1). The near expected number of *Orc2^{+/Δ}* embryos suggests that hemizygosity of *Orc2* can still support viability.

Mouse embryo fibroblasts (MEF) were obtained from *Orc2^{f/f}* E13.5 day embryos and cultured *in vitro*. 3 days after infection with an Adeno virus expressing Cre, effective recombination between the loxP sites is seen, resulting in a genotype showing that exons 6 and 7 have been deleted in most of the MEF (*Orc2^Δ*) (Fig. 1E). Consistent with this, at 5 days after adeno-Cre infection, the ORC2 protein is not detected in the MEF population (Fig. 1F). The dilution of the undeleted MEF lysate on the same blot suggests, that if any ORC2 protein is expressed in the MEF after adeno-Cre infection, it is <10% of that in the undeleted MEF. The proliferation rate of the MEF were measured by MTT assays. Even in the *Orc2^{+/+}* MEF, the infection with adeno-Cre decreased proliferation a little (the orange line compared to the blue line in Fig. 1G). However, for *Orc2^{f/f}* MEF infection with adeno-Cre impairs proliferation even further (yellow line compared to black line in Fig. 1G). Furthermore, when the MEF were cultured for 15 days, the surviving cells express significant levels of ORC2 (Fig. 1H), suggesting that the MEFs that had not undergone the Cre-mediated deletion take over the culture.

Taken together, these results suggest that Cre mediated deletion of exons 6 and 7 of *Orc2* leads to early embryonic lethality and impairs the proliferation of normal diploid MEFs in culture.

Knockout of *Orc2* in developing mouse liver makes ORC2 protein undetectable in hepatocytes and yet supports




Figure 1

Generation of *Orc2^{f/f}* mice and ORC2 KO MEFs.

- A. Scheme of introduced loxP sites in *Orc2* locus.
- B. Representative picture of genotyping of offspring coming from *Orc2^{f/+}* crossed with *Orc2^{f/+}*.
- C. The ratio of observed to expected animals coming from *Orc2^{f/+}* crossed with *Orc2^{f/+}*.
- D. Schematic of the ORC2 protein and the DeltaORC2 protein produced after deletion of exons 6 and 7. A110 is mutated to V110 and then the protein goes out of frame.
- E. Validation of *Orc2* deletion 3 days after Adeno cre transduction.
- F. Western blot of ORC2 protein 5 days after Adeno cre transduction. 10 or indicated ul of lysate loaded/lane as written on the top
- G. MTT assay of WT and *Orc2^{f/f}* MEFs without and with Adeno cre transduction.
- H. Western blot of ORC2 protein 5 and 15 days after Adeno cre transduction.

most of normal development and endoreduplication

Mice carrying one copy of the Albumin promoter driven Cre gene (*Alb-Cre*) express the Cre recombinase specifically in hepatocytes (Postic et al., 1999). When the *Orc2^{ff}* mice are crossed with mice carrying *Alb-Cre* (shortened below as *Alb*), the Cre recombinase is expected to promote the recombination mediated deletion of exons 6 and 7 of *Orc2* in the hepatocytes. The *Alb^{+/-}-Orc2^{ff}* mice were viable and fertile (Fig. 2A, 2B). Crossing such mice yielded the expected numbers of *Alb^{+/-}-Orc2^{ff}* mice, though there was a drop in the yield of *Alb^{+/-}-Orc2^{ff}* mice. One allele of *Alb-Cre* is sufficient to express enough Cre to carry out homozygous deletion of the *Orc2* allele in the hepatocytes, so that the results suggest that deletion of *Orc2* in the hepatocytes did not impair viability. The partial lethality of the *Alb^{+/-}* (HOM) mice could be due to the toxicity of high dose of Cre recombinase expressed, something that has been noted by other groups (Schmidt et al., 2000, Loonstra et al., 2001, Schmidt-Suppran and Rajewsky, 2007, Naiche and Papaioannou, 2007, Janbandhu et al., 2014). Because the *Alb-Cre* is expressed only in the hepatocytes and not in other cells in the liver, we isolated hepatocytes from these livers by growing them in culture (schematically shown in Fig. 4A) to determine whether the ORC2 protein was decreased (Fig. 2C). The ORC2 protein was significantly decreased in all five of the *Alb^{+/-}-Orc2^{ff}* mice, showing consistent and near 100% effect of the deletion in the hepatocytes. Western blots for several other ORC subunits, and CDC6 protein showed that the loss of ORC in the same hepatocytes did not significantly decrease these proteins, though MCM2 and MCM3 (two subunits of the MCM2-7 helicase component) were decreased by 50% (Fig. 2D). The body weights of the *Alb, Orc2^{ff}* mice were smaller than in the *Orc2^{ff}* animals of both sexes, though the liver size and liver size normalized to body weight was significantly smaller only in the females (Fig. 2E-G). There was some elevation of the circulating liver enzymes in the mice where *Alb-Cre* is expressed to delete the *Orc2* gene, suggesting that there is some impairment of liver function (Supp. Fig. S1).

The livers of mice can grow by proliferation (increasing the number of cells) or by hypertrophy of cells (larger cells with endoreduplicated nuclei). Histological examination of the livers revealed that the *Orc2* deletion was accompanied by the presence of fewer (~50% of WT levels), but larger nuclei and cells (Fig. 2H and 2I). The *Orc2* deleted hepatocytes had significantly larger nuclei in both males and females (Fig. 2J, K). The larger nuclei in hepatocytes with *Orc2* deletion also seen in other contexts, e.g. in the pre-resection hepatocytes with *Orc2* deletion (Fig. 5E, 0 hr point) and in the Cre expressing, *Orc1^{ff/+}, Orc2^{ff}* that have an intact *Orc1* gene but delete both alleles of *Orc2* (Supp. Fig. S3B, green bars compared to blue bars).

To determine whether the larger nuclei were generated by endoreduplication, we isolated nuclei from hepatocytes and stained them with the DNA staining dye, DRAQ5, followed by flow cytometry (Fig. 3A). Both male and female *Alb-Cre, Orc2^{ff}* hepatocytes showed a decrease in 4n DNA content and an increase in 16n DNA content (Fig. 3B-D). The expression of Cre would also delete a lox-stop-lox element upstream from EYFP and lead to the expression of EYFP. Indeed, when we gated on EYFP positive cells, the 8n and 16N DNA content increase was much more evident relative to the EYFP negative cells, while the 2N DNA containing nuclei were decreased (Fig. 3E-G). Thus *Orc2* deletion promotes endo-reduplication, a result similar to what was noted when *Orc1* was deleted in mouse livers (Okano-Uchida et al., 2018).

DNA replication of *Alb-Cre^{+/-}-Orc2^{ff}* hepatocytes *in vitro*

Primary hepatocytes isolated from 8-10 weeks old mouse liver can replicate their DNA and proliferate for a limited time *in vitro*. We isolated such hepatocytes from *Alb-cre^{-/-}* (WT) and *Alb-cre^{+/-}* livers, both from *Orc2^{ff}* mice (Fig. 4A). Genotyping reveals that over 90% of the cells from the *Alb-cre^{+/-}* livers have successfully deleted *Orc2* exons 6 and 7 (Fig. 4B), while immunoblotting shows that ORC2 protein expression is also significantly decreased in these cells (Fig. 3C). EdU labeling *in vitro* for 2 hrs, showed that the EYFP positive cells (where Cre recombinase has been active) can incorporate EdU (Fig. 4C), although the total number of nuclei




Figure 2

Development of liver in *Orc2* KO mice.

A. Scheme of $Alb^{+/-}Orc2^{ff} ROSA26^{stop-EYFP}$ crossed with $Alb^{+/-}Orc2^{ff} ROSA26^{stop-EYFP}$ (All mice are with $ROSA26^{stop-EYFP}$ and so we do not include this in the genotypes below)

B. The ratio of observed to expected animals coming from A.

C. Western blot of hepatocytes from $Orc2^{ff}$ and $Orc2^{ff} Alb-Cre$ +/- animals. Tubulin was used as loading control.

D. Quantification of the Western blots of hepatocyte lysates from $Orc2^{ff}$ (without $Alb-Cre$) mice and the same genotype but with $Alb-Cre$ to show the levels of other key replication initiation proteins in the ORC2 KO hepatocytes.

E. Average body weight of $Orc2^{ff}$ and $Alb-Orc2^{ff}$ animals.

F. Average liver weight of $Orc2^{ff}$ and $Alb-Orc2^{ff}$ animals.

G. Average liver to body weight ratio of $Orc2^{ff}$ and $Alb-Orc2^{ff}$ animals.

H. Representative H&E staining of liver tissue from $Orc2^{ff}$ (WT) and $Alb-Orc2^{ff}$ (KO) animals. Both panels at same scale.

I. Quantification of hepatocyte nuclear size in $Orc2^{ff}$ and $Alb-Orc2^{ff}$ animals.

J. Quantification of hepatocyte nuclear size in $Orc2^{ff}$ and $Alb-Orc2^{ff}$ female mice.

K. Quantification of hepatocyte nuclear size in $Orc2^{ff}$ and $Alb-Orc2^{ff}$ male mice.

*p < 0.05, **p < 0.01, two-tailed Student's t test.




Figure 3

Endoreduplication in the *Orc2* liver conditional knock-out animals.

A. Experimental design.

B-D. Quantification of nuclei ploidy on 10,000 nuclei from the livers of *Orc2*^{fl/fl} *ROSA26*^{stop-EYFP} and *Alb-Orc2*^{fl/fl} *ROSA26*^{stop-EYFP} animals.

E-G. Quantification of nuclei ploidy for EYFP low (includes negative) and high (positive) primary liver cells. *p < 0.05, **p < 0.01, ***p < 0.001, two-tailed Student's t test.

that incorporate EdU is decreased to about 30% of that seen in the ORC2 wild type hepatocytes (Fig. 4D). Thus hepatocytes in culture can continue to replicate DNA in the absence of detectable ORC2 protein.

DNA replication in *Alb-Cre^{+/-}-Orc2^{f/f}* hepatocytes during liver regeneration *in vivo*

Liver DNA synthesis peaks around 36-48 hours, and the liver regenerates to nearly 50% of its original weight within 2-3 days following partial hepatectomy. Such regeneration involves both normal mitotic DNA replication/cell division and endoreduplication accompanied by hypertrophy of the cells. We therefore tested whether deletion of the *Orc2* gene in the hepatocytes adversely affects liver regeneration after partial hepatectomy in 8-14 weeks old mice (Fig. 5A). Nearly two-thirds of the liver is removed surgically and the mice allowed to recover for 36-48 hrs before harvesting the regenerated liver.

Although the liver weight after regeneration was smaller than that of WT livers, the liver:body weight ratio was similar in the livers with *Orc2* deletion and *Orc2* WT (Fig. 5B-D). The H&E stain of the regenerated liver shows that the liver cells are larger and have larger nuclei in the liver with *Orc2* deletion (Fig. 5E). In these experiments, both before and after regeneration, the livers with the *Orc2* deletion have ~50% nuclei compared to the wild type livers (Fig. 5F). Finally, labeling the livers by injection of EdU in the mice 3-4 hrs before harvesting, shows that the EYFP positive cells (indicating activity of Cre recombinase) were proficient in synthesizing DNA and incorporating EdU (Fig. 5G, H).

Nearly 100% of the hepatocytes were positive for EYFP, suggesting that the Cre recombinase was active in the vast majority of the hepatocytes (Fig. 5G), and consistent with the complete depletion of *Orc2* protein from the hepatocytes of all mice with that genotype (Fig. 3C). Thus the 30% of the liver cells that were incorporating EdU in the Cre expressing livers (Fig. 5H) were doing so in the absence of *Orc2* protein. Also, we noted several pairs of EdU positive nuclei (marked by arrows in Fig. 5G) whose relative positions suggest that they are sisters born from the same mitosis, suggesting that some EdU positive cells can go through mitosis in the EYFP positive cells. Finally, the nuclei were significantly larger in the Cre active *Orc2^{f/f}* hepatocytes than in the hepatocytes without Cre, both before and after regeneration (Fig. 5I).

We did not explore why the EYFP protein is mostly nuclear in hepatocytes in culture (Fig. 4C) and mostly cytoplasmic in hepatocytes in the liver tissue (Fig. 5G, 7G), but speculate that differences in signaling pathways or fixation techniques between the two conditions contribute to this difference.

Viable mice with endoreduplicated hepatocyte nuclei in *Alb-Orc1^{f/f} Orc2^{f/f}* mice

We have reported that conditional deletion of *Orc1* in developing mouse livers still allowed livers to develop and induced premature endoreduplication, suggesting that significant DNA synthesis can occur in liver cells that are genetically deleted of *Orc1*. We bred the *Orc1^{f/f}* mice with *Orc2^{f/f}* mice to obtain mice where both *Orc1* and *Orc2* are floxed but *Alb-cre* is not present (Fig. 6A, B). Even though Cre was not expressed in these livers, for unknown reasons there was a decrease in the percentage of progeny when *Orc2* was floxed in the liver (with or without *Orc1* being floxed), but not when just *Orc1* was floxed (Fig. 6C). These mice were then bred with *Orc2^{f/f} Alb-cre^{+/-}* mice, and the resulting *Orc1^{f/+}, Orc2^{f/f}, Alb-cre^{+/-}* mice intercrossed to get *Orc1^{f/f}, Orc2^{f/f}, Alb-Cre^{+/-}* mice where the *Orc1* and *Orc2* alleles are conditionally deleted in hepatocytes, and the deletion is




Figure 4

***Orc2* KO primary hepatocytes are viable and can incorporate EdU *in vitro*.**

A. Experimental design

B. Genotyping and western blotting of hepatocytes

C. Representative picture of EdU, EYFP and DAPI staining on the *Orc2* WT (*Orc2^{fl/fl}*) and KO (*Orc2^{fl/fl} Alb-Cre*) primary hepatocytes.

D. The percentage of EdU positive nuclei from *Orc2* WT or *Orc2* KO primary hepatocytes.

*p < 0.05, two-tailed Student's t test.




Figure 5

The ORC2 mutant livers regenerate after partial hepatectomy.

- A. Schematic of the experiment.
- B. Body weight of the *Orc2^{fl/fl} ROSA26^{stop-EYFP}* mice without (-/-) or with *Alb-Cre* (-/+) before partial hepatectomy.
- C. Liver weight of the mice in B after liver regeneration.
- D. Regenerated Liver to pre-hepatectomy body weight ratio of the mice in B.
- E. H&E stain of *Orc2^{fl/fl} ROSA26^{stop-EYFP}* livers with intact *Orc2* (*Alb-cre* -/-, N=3) or *Orc2* knockout (*Alb-cre* +/-, N=7). Scale bar: 25 μ m.
- F. Quantitation of nuclear counts per field (76,000 μ m²). Six images were taken for each liver. 0 hr (pre-resection), 42 hr (post-regeneration).
- G. EdU incorporation of indicated livers. EYFP marks cells where Cre has been expressed. Scale bar: 25 μ m.
- H. Percent EdU+ nuclei counted in 1882 and 825 nuclei in the *Cre*- and *Cre*+ livers, respectively.
- I. Nuclear size of indicated livers. 0 hr (pre-resection), 42 hr (post-regeneration). Mean and S.D from about 40-70 nuclei, *p < 0.05, ****p < 0.0001, unpaired two-tailed Student's t test is used

accompanied by activation of EYFP expression. Immunoblotting of isolated hepatocytes showed that in four out of four mice expressing the Cre recombinase the ORC1 and ORC2 proteins were decreased significantly (**Fig. 6D** [↗](#)).

Immunoblotting of some of the other replication proteins showed no decrease of other ORC subunits, CDC6 and two of the MCM2-7 subunits (**Fig. 6E** [↗](#)). The ORC3 protein, which did not change when *Orc2* was deleted (**Fig. 2D** [↗](#)) was surprisingly elevated in **Fig. 6E** [↗](#) when *Orc1* and *Orc2* were both deleted. We do not know why this is the case, and it was not seen consistently in all four animals. Similarly, the 50% decrease of MCM2 and MCM3 that was seen in the absence of ORC2 (**Fig. 2D** [↗](#)), disappeared when both *Orc1* and *Orc2* are deleted.

The female *Alb-Orc1^{fl/fl} Orc2^{fl/fl}* mice were significantly smaller in size, with smaller livers and decreased liver to body weight ratio (**Fig. 6F** [↗](#), Supp. Fig. 2A), but the male mice were relatively less affected (**Fig. 6F** [↗](#)). We have also observed ~50% lethality of double knock-out female mice within the first month of life, but the remaining 50% survive beyond 4 months of age (Supp. Fig. 2C). H&E staining showed that the double knockout (DKO) male livers have fewer but larger cells with significantly larger nuclei (**Fig. 6G** [↗](#), **6H**). Interestingly, this phenotype is even more marked in the female liver (Supp. Fig. 2B). By flow cytometry the EYFP positive cells as evidence of Cre recombinase activity had distinctly more polyploidization than the EYFP negative cells (**Fig. 6I** [↗](#)). Taken together the results reveal that hepatocytes can synthesize DNA by endoreduplication to produce very large nuclei and very large cells so that the liver size is not hugely decreased, even in the absence of two subunits of ORC. Female mice that have a deletion in two subunits of ORC suffer more morbidity and mortality than male mice.

DNA replication in *Alb-Cre^{+/+}-Orc1^{fl/fl} Orc2^{fl/fl}* hepatocytes during liver regeneration *in vivo*

Finally, partial hepatectomy was performed in the livers of male mice with DKO of *Orc1* and *Orc2*. As can be seen even before partial hepatectomy the body weights were not significantly different from WT mice (**Fig. 7A** [↗](#)). Post regeneration, the liver weights and liver/body weight ratios were not decreased in the DKO mice compared to the WT mice (**Fig. 7B**, **C** [↗](#)). H&E sections showed that the hepatocyte nuclei were larger and hepatocyte nuclear density lower in the DKO livers both pre- hepatectomy and in the regenerated livers 36 hr post-hepatectomy (**Fig. 7D-F** [↗](#)). EdU labeling for 3-4 hrs before harvesting the livers showed that despite the DKO, the EYFP positive cells showed extensive DNA synthesis (**Fig. 7G** [↗](#)). Even though nearly 100% of the hepatocytes are EYFP positive (**Fig. 7G** [↗](#)) and ORC1 and ORC2 proteins depleted in the hepatocytes (**Fig. 6D** [↗](#)), 15 % of nuclei stained with EdU (**Fig. 7H** [↗](#)), a percentage that was higher in the DKO regenerating livers than in the WT livers, suggesting that the endoreduplication that accompanies liver regeneration can occur even after deletion of two of the six subunits of ORC.

Discussion

The six subunit ORC is important for recruiting MCM2-7, the core of the replicative DNA helicase and is essential for all forms of DNA replication in eukaryotic cells (Costa and Diffley, 2022 [↗](#), Stillman, 2022 [↗](#), Hu and Stillman, 2023 [↗](#)). Yet, it has been possible to select cancer cell-lines that have mutations in *Orc1*, *Orc2* or *Orc5* and do not express detectable levels of the proteins and yet load MCM2-7 to the chromatin and replicate their DNA as they proliferate in culture (Shibata et al., 2016 [↗](#), Shibata and Dutta, 2020 [↗](#)). This paradox suggested that there may be an alternative way to load sufficient MCM2-7 on chromatin and support DNA replication, at least in cancer cells in culture. In addition, there have been two instances where *Orc1* has been mutationally removed in *Drosophila* (Park and Asano, 2008 [↗](#)) and in mouse hepatocytes or placental trophoblasts (Okano-Uchida et al., 2018 [↗](#)), where extensive DNA replication (particularly endo-reduplication in




Figure 6

Endoreduplication in liver of *Orc1 Orc2* DKO animals.

A-B. Breeding schemes to obtain conditional double floxed animals.

C. The ratio of observed to expected animals coming from B. *Orc1* = all animals with *Orc1* f/f *ROSA26* *stop-EYFP*, *Orc2* = all animals with *Orc2* f/f *ROSA26* *stop-EYFP*, *Orc1 Orc2* = all animals with *Orc1* f/f *Orc2* f/f *ROSA26* *stop-EYFP* genotype. This was before introduction of *Alb-Cre*.

D. Immunoblot of hepatocytes from WT (*Orc1* f/f *Orc2* f/f) and DKO (*Orc1* f/f *Orc2* f/f *Alb-cre+/-*) mice to show that ORC1 and ORC2 are depleted in the DKO cells.

E. Quantitation of immunoblots to show that levels of other key initiation protein subunits are not decreased in the DKO mice hepatocytes.

F. Average body, liver weight and their ratio for WT and DKO animals.

G. Representative H&E staining of liver tissue from male WT and DKO animals.

H. Quantification of hepatocyte nuclear size in the WT and DKO animals.

I. Quantification of nuclei ploidy for EYFP low (includes negative) and high (positive) primary liver cells from DKO mice.




Figure 7

Endoreduplication in the liver of the DKO mice during liver regeneration.

A-C. Body weight pre-resection, liver weight post-regeneration and regenerated liver to body weight ratio in mice with indicated genotypes. Black bars: 4 WT males (*Orc1^{fl} Orc2^{fl} ROSA26^{stop}-EYFP* mice without *Alb-Cre*). White bars: 6 DKO males (*Orc1^{fl} Orc2^{fl} ROSA26^{stop}-EYFP* mice with *Alb-Cre* $+/-$). No significant difference between the two groups using two tailed Student t-test.

D: H&E stain of WT (N=4) or DKO mice (N=6). Scale bar: 50 μ m.

E. Quantitation of hepatocyte nuclear size post regeneration. WT: black bars. DKO: white bars. 0 hr (pre-resection). 42 hr (post-regeneration). Five-six images were taken for each liver. About 120-200 nuclei are counted.

F: Quantitation of hepatocyte nuclear density post regeneration. WT: black bars. DKO: white bars. 0 hr (pre-resection). 42 hr (post-regeneration). Five-six images were taken for each liver.

G. Micrographs of EdU, DAPI and EYFP imaging of livers with indicated genotypes post regeneration. Scale bar: 20 μ m. WT in the top row, DKO in the bottom row.

H. Quantitation of EdU positive nuclei post regeneration. WT: black bar. DKO: white bar. Five-six images were taken for each liver. * $p < 0.05$, **** $p < 0.0001$, unpaired two-tailed Student's t test were used

hepatocytes and trophoblasts) has persisted, suggesting that an alternative way of loading MCM2-7 may be available in certain unique cell-cycles that are different from the normal DNA replication that occurs during normal mitotic growth of diploid cells.

Five of the six subunits of ORC (ORC1-5) associate to form a ring-shaped structure with interactions between the WH domains and the AAA+ like domains of the five subunits (Bleichert et al., 2015 [\[1\]](#)). In mammalian cells only ORC1 and ORC4 have the intact Walker A and B motifs that are necessary for the molecules to act as AAA+ ATPases (Giordano-Coltart et al., 2005 [\[2\]](#)). Though this has never been demonstrated experimentally, it is theoretically possible that in the absence of ORC1 in flies or in mouse livers, another related AAA+ ATPase, intimately involved in MCM2-7 loading, CDC6, can substitute for ORC1 to reconstitute a functional 5 subunit ring-shaped ORC-like structure that executes its function (Takeda et al., 2005 [\[3\]](#), Bell, 2017 [\[4\]](#)). We therefore wished to test whether removal of a second subunit of ORC, ORC2, that does not have obvious ATPase activity and not much homology to CDC6, would still permit endoreduplication in mouse livers. In the cancer cells, we had removed the initiator methionine of the *Orc2* gene (Shibata et al., 2016 [\[5\]](#)). Although 99.9% of the ORC2 protein disappeared, an N terminally truncated form of ORC2 protein was expressed in the mutant cells at 0.1% of the wild type level (Chou et al., 2021 [\[6\]](#)). Thus, in this case we took the additional precaution to design a mutation such that even if a truncated protein was expressed from an internal methionine downstream from the mutation site, such a protein will be missing a significant part of its AAA+ATPase domain. Anyhow, in Western blots we do not see any evidence of such a truncated protein being expressed in the genetically altered MEFs or hepatocytes. In addition, we wondered whether the simultaneous genetic removal of two subunits of ORC, *Orc1* and *Orc2*, would successfully obliterate all forms of DNA replication, including endoreduplication.

Our results show that while *Orc2* is genetically essential for viability of early embryos *in utero* and mouse embryo fibroblasts *in vitro*, mutational inactivation of *Orc2*, or combined inactivation of *Orc1* and *Orc2*, does not significantly inhibit development of the mouse liver. Adult, viable mice are produced. Albumin expression is activated very early in hepatoblasts in the 7-8 somite stage and the albumin mRNA can be seen in the hepatic primordium emerging from the gut at E9.5 days (Gualdi et al., 1996 [\[7\]](#)). Single cell sequencing studies confirm that hepatoblasts expressing albumin are present at e11 and that undifferentiated endodermal cells (the precursors to hepatoblasts) are non-existent (Wang et al., 2020 [\[8\]](#)). ³H-thymidine incorporation can be measured in the liver as late as days 7-14 postnatally (Tilghman and Belayew, 1982 [\[9\]](#)), so that many cycles of DNA replication and cell proliferation are expected to occur in hepatoblasts and hepatocytes after the activation of Alb-Cre and subsequent homozygous knockout of *Orc2* (or *Orc1*) in the hepatocytes.

The presence of functional livers in the mutant (but viable) animals, suggested that the homozygous knockout of these genes did not have a profound effect on liver development, as would have been expected if the cells were as sensitive as MEFs in culture to the loss of the ORC holocomplex. We have been unable to find antibodies that will recognize mouse *Orc1* or *Orc2* proteins in immunohistochemistry on tissues and so decided to calculate how many cell divisions have to occur *after* the *Orc1* or *Orc2* genes are deleted in the embryonic mouse hepatocytes (see Methods and Table 2). The calculations suggest that the *Orc2* deleted livers and male DKO livers underwent at least 18 cell divisions, while the female DKO livers underwent at least 15 cell divisions since E9.5 (Table 2, bottom row).

If we allow a generous 3 cell divisions for the Cre recombinase to definitively excise the Floxed genes (consistent with what we see in MEFs in culture), this means that 15 cell divisions in the *Orc2* deleted livers and male DKO livers occurred *after* the gene(s) was/were deleted. Once a gene is deleted, each cell division decreases the corresponding protein at least by half, so that 6 cell divisions would dilute the targeted protein to <1.5% of the wild type levels. Thus even after the ORC2 (or ORC1 and ORC2 in the DKO mice) decreased to <1.5% the WT level, the *Orc2* knockout livers (male or female) and the male DKO livers could execute at least 9 more rounds of replication

and cell division. Note the amount of the relevant ORC subunit would continue to decrease by half with each further division. Also, that the levels of ORC are likely to decrease faster if the protein is actively degraded or if more cell divisions are necessary to counter any development-related apoptosis.

This suggests that mitotic DNA replication of diploid cells is not completely eliminated at least in the *Orc2*^{-/-} or male DKO livers after loss of the targeted ORC proteins. Consistent with this, we see EdU incorporation in EYFP⁺ hepatocytes, *in vitro*, which, however, could be from a terminal endoreduplication event. More tantalizing is the presence of *paired* EYFP positive EdU-positive nuclei during the regeneration after partial hepatectomy in 8-14 week old mice (**Fig. 5G** [↗](#)). Such paired nuclei are usually seen with daughter nuclei that have just been born from mitosis, suggesting that there were at least a few cells that underwent complete DNA replication and mitosis even though the hepatocytes had undergone 15 rounds of cell division after deletion of the targeted ORC gene(s).

Could the ORC deletion lead to the immediate loss of hepatoblasts (despite having inherited ORC protein from the endodermal cells) causing undifferentiated endodermal cells to persist and proliferate much longer (without activating Cre) than in normal development? We consider this unlikely, but if true it will be very unexpected, both by suggesting that deletion of ORC immediately leads to the death of the hepatoblasts (despite a healthy reserve of inherited ORC protein) and by suggesting that there is a novel feedback mechanism from the death/ORC depletion of hepatoblasts leading to the persistence and proliferation of undifferentiated endodermal cells.

The female DKO (but not the *Orc2*^{-/-}) livers seem more sensitive to the depletion of both ORC1 and ORC2 since ~12 cell divisions in the female DKO livers occur after allowing 3 cell division for the targeted genes to be definitively deleted. Thus, after the cells have reached <1.5 % of WT levels of ORC2 or ORC1 protein, they can still execute at least 6 more cell divisions. Consistent with this sex specific difference, the male mice appear to be more tolerant of deletion of *Orc2* gene or both *Orc1* and *Orc2* genes, with the livers reaching larger sizes than in the female mice. Plasma testosterone levels reach 50% of adult levels in the first day after birth before declining (to rise again at puberty at 4-5 weeks age), suggesting that enough androgens are present during the proliferative phase of liver development (Bell, 2018 [↗](#)). Sex-specific changes in gene expression have been noted to begin in the liver by 3 weeks postnatally (Conforto and Waxman, 2012 [↗](#)). Thus, it is possible that androgens directly or indirectly stimulate the division or hypertrophy of hepatocytes to ameliorate the negative effects on liver mass due to the loss of ORC. We have also shown that loss of ORC subunits in cancer cell lines causes significant changes in gene expression due to the role of ORC subunits in epigenetic regulation (Su et al., 2024 [↗](#)), so another explanation of the more severe phenotype in females could be that the epigenetic effects of ORC subunits are more important in female compared to male livers.

In contrast to diploid DNA replication in hepatocytes, we have shown that the livers with a genetic deletion of *Orc2* (or *Orc1* and *Orc2*) undergo endo-reduplication more easily during normal development. Such DNA synthesis is also clearly seen during liver regeneration after partial hepatectomy in the *Orc2*^{-/-} or *Orc1*^{-/-}, *Orc2*^{-/-} livers. Endoreduplication differs from normal replication in that although the cycles of DNA replication (S phase) are separated by a G1 like phase (G phase) they are not separated by an intervening mitosis and cytokinesis, and is commonly seen during normal development in plants and animals (Shu et al., 2018 [↗](#)). However, all evidence suggests that the biochemical mechanism of initiating DNA replication is the same in normal mitotic replication and endoreduplication. For example, cyclin E/cdk kinase activity and MCM2-7 protein association with the chromatin oscillate between G phase and S phase) during *Drosophila* endoreduplication (Lilly and Duronio, 2005 [↗](#)) (Su and O'Farrell, 1998 [↗](#)). Similarly, in the mouse, cyclin E and CDC6 promote, while geminin inhibits endoreduplication (Welch, 1992 [↗](#)) (Bermejo et al., 2002 [↗](#), Gonzalez et al., 2006 [↗](#)) (de Renty et al., 2014 [↗](#)). Thus, the persistence, and

in fact acceleration, of endoreduplication in the mouse livers in the absence of ORC2 and/or ORC1 suggests that there must be an alternate way to recruit sufficient MCM2-7 to the chromatin to support at least the three cycles of endoreduplication required to produce a 16 N nucleus from a 2 N embryonic nucleus during normal development, and to support about two cycles of endoreduplication during liver regeneration.

The acceleration and persistence of endoreduplication even when both *Orc1* and *Orc2* are genetically inactivated, makes it unlikely that a make-shift ORC-like complex is being assembled for loading the MCM2-7 proteins when two out of the five subunits in the ring are missing. Our results are virtually identical to what was observed when we conditionally deleted *Orc1* (Okano-Uchida et al., 2018 [DOI](#)). There too, endoreduplication in the liver cells not only persisted but was induced prematurely during development. One possible explanation of the greater endoreduplication in both our papers is that mitosis may be arrested earlier in development by G2 DNA damage checkpoints activated by incomplete licensing and replication of the genome in the absence of ORC. As a result, endoreduplication cycles could begin earlier in development resulting in greater endoreduplication.

Is it possible that incomplete deletion of *Orc2* or *Orc1* genes in the hepatocytes allows enough liver cells to still carry an *Orc2* or *Orc1* gene to support DNA replication? This is very unlikely, because the *in vitro* DNA replication experiment with hepatocytes shows that although the ORC2 protein is virtually undetectable in immunoblots and there is evidence of Cre activity in 100% of the hepatocytes, the number of nuclei incorporating EdU in culture are decreased to only 30% the wild type level.

Furthermore, in the partial hepatectomy experiments, nearly 100% of the hepatocytes are positive for EYFP and no ORC2 (or ORC1 protein in the DKO) is detectable in immunoblots of isolated hepatocytes, suggesting high penetrance in the expression of the Cre recombinase in the hepatocytes. Yet the liver succeeds in endoreduplicating to reach nearly normal liver size and EdU incorporation is seen in 35% (**Fig. 5H** [DOI](#)) and 15% (**Fig. 7G** [DOI](#)) of hepatocytes from the *Orc2* KO and DKO mice, respectively.

Although the liver size in the *Alb-Cre, Orc2^{ff}* mice is not significantly decreased relative to body weight, the liver function tests suggest some impairment of liver function. We cannot yet attribute this deficit of liver function to the decrease in number of cells, or to excess endoreduplication, because ORC subunits are also known to be important for epigenetic control of gene expression (Vermeulen et al., 2010 [DOI](#)). Future experiments will determine whether the liver pathology could be secondary to epigenetic dysregulation of genes important for liver function. The double knockout female mice are sicker, and here some of the explanation may lie in the smaller liver, but here again epigenetic dysregulation in the absence of ORC cannot be ruled out as a potential cause for the morbidity.

Interestingly, the deletion of *Orc2* or of *Orc1+Orc2*, in the hepatocytes does not *consistently* increase or decrease any of the other ORC subunits or the examined proteins downstream of ORC like CDC6 or two of the subunits of MCM2-7. Thus, a change in any of these proteins is unlikely to be the explanation for how the hepatocytes license enough origins to support replication during development and regeneration in the absence of the ORC holocomplex. The hepatocyte studies say that the cancer cells are not unique in their ability to bypass the requirement of two ORC subunits. Even primary cells can sometimes load enough MCM2-7 to replicate DNA in the absence of detectable amounts of the ORC holocomplex. It is also worth noting that in cancer cells in culture, we are seeing a near normal level of loading of MCM2-7 on chromatin when *Orc1*, *Orc2* or *Orc5* genes are deleted, and that 60% of the origins of replication remain at the same sites as in WT cells (Shibata et al., 2024 [DOI](#)). Although it is impossible to rule out that a very small amount of ORC

protein somehow persists in the hepatocytes (or the cancer cell lines) with these mutations and that this is sufficient to facilitate loading of enough MCM2-7 to support DNA synthesis *in vivo* or *in vitro*, this is becoming progressively unlikely.

Methods

Mice

Work involving mice adhered to the guidelines of the Institutional Animal Care and Use Committees (IACUC) at the University of Virginia (protocol number 4198), the University of Alabama at Birmingham (protocol number 22335), the Ohio State University and Medical University of South Carolina. *Orc1*^{ff} *ROSA26*^{stop-EYFP} animals were previously reported (Okano-Uchida et al., 2018 [DOI](#)). *Orc2*^{ff} mice were generated by Cyagen Biosciences Inc. Exons 6-7 (amino acids L111-V150) was selected as conditional knockout region (cKO). Mouse genomic fragments containing homology arms (HAs) and cKO region were amplified from BAC clone by using high fidelity Taq DNA polymerase and were sequentially assembled into a targeting vector together with recombination sites and selection marker, Neo cassette, flanked by SDA (self-deletion anchor) sites. The linearized vector was electroporated into C57BL/6 ES cells that were subject to G418 selection (200 µg/mL) after 24 hours. 188 G418 resistant clones were picked and amplified in 96-well plates. The PCR screening identified 29 potential targeted clones, from among which 12 were expanded and further characterized by Southern blot analysis. Eleven of the twelve expanded clones were confirmed to be correctly targeted. Targeted ES cell clone N-1F10 was injected into C57BL/6 albino embryos, which were then re-implanted into CD-1 pseudo-pregnant females. Founder animals were identified by their coat color, their germline transmission was confirmed by breeding with C57BL/6 females and subsequent genotyping of the offspring. Three male and five female heterozygous targeted mice were generated from clone N-1F10. Floxed *Orc2* mice were crossed to the *Orc1*^{ff} *ROSA26*^{stop-EYFP} animals to generate *Orc2*^{ff} *ROSA26*^{stop-EYFP} and *Orc1*^{ff} *Orc2*^{ff} *ROSA26*^{stop-EYFP} strains for usage of EYFP expression as a reporter for Cre recombinase activity and *Orc2* or *Orc1* and *Orc2* deletion. Those were further bred into tissue specific Sox2-Cre (Hayashi et al., 2002 [DOI](#)) or Alb-Cre mice to obtain *Orc2* or *Orc1* and *Orc2* knockout in all cells of the inner cell mass following implantation or in the liver, respectively.

We used two independent Alb-Cre lines to introduce the gene. At UVA/UAB we used the Alb-Cre mice from (Postic et al., 1999 [DOI](#)). The Alb-Cre mice used at University of Wisconsin are from (Schuler et al., 2004 [DOI](#)). All the mice used in this study were maintained in a mixed background (C57BL/6x129x FVB/N).

PCR

Genomic DNA from ear punches was isolated using Quick Extract DNA Extraction Solution (Lucigen., Cat# QE09050). All genotyping PCRs were carried out using MyTaq™ Red Mix (Bioline, Cat# BIO-25043) according to the manufacturer's instructions.

Orc1 genotyping was carried out with primers F1 forward (common to both alleles; 51-GCTGCTTCAGTGTGGCAATA-31), R1 reverse (specific for the WT allele; 51-CTCCAATTGTTCCCCAGCTA-3'), and R2 reverse (specific for deleted allele, 51-CACCTGTCACCTGGACCACAC-31). The PCR parameters were 95°C for 30 sec, 45× (95°C for 15 sec, 60°C for 20 sec, 72°C for 60 sec), 72°C for 5 min. PCR product was run on 2% agarose gel and WT band was detected at 439bp, transgenic band at 310bp, and deleted at 677bp.

Orc2 genotyping was carried out with primers F1 forward (common to both alleles; 51-GAGGTTGTGGCTGTAATATACGTGATC -31), and R1 reverse (common to both alleles; 51-CTGAGCCATCTAACTCCTTCCTAGC -3'), or F2 forward (common to both alleles; 51-TGGGTAGGTTTCATTCCAGTTTAGCC -31), and R2 reverse (common to both alleles; 51-

ACCTTGGTATTGGACGTCTCTATTC -3'). The PCR parameters were 95°C for 30 sec, 35× (95°C for 15 sec, 55°C for 20 sec, 72°C for 60 sec), 72°C for 5 min. To detect constitutive KO allele combination of F1, F2, and R2 was used. PCR product was run on 2% agarose gel and following bands were detected: for F1+R1 – WT at 342bp, and transgenic band at 398bp; for F2+R2 - WT at 254bp, and transgenic band at 367bp; for F1+F2+R2 - WT at 254bp, transgenic band at 367bp, and deleted at 306bp.

ROSA26 genotyping was carried out with primers F1 forward (common to both alleles; 51-AAAGTCGCTCTGAGTTGTAT-31), R1 reverse (specific for the WT allele; 51-GGAGCGGGAGAAATGGATAT-3'), and R2 reverse (specific for deleted allele, 51-GCGAAGAGTTGTCTCAACC-31). The PCR parameters were 95°C for 30 sec, 45× (95°C for 15 sec, 60°C for 20 sec, 72°C for 60 sec), 72°C for 5 min. PCR product was run on 2% agarose gel and WT band was detected at 650bp, and transgenic band at 340bp.

Sox2-Cre genotyping was carried out with primers F1 forward (common to both alleles; 51-ATGCTTCTGTCCGTTTGCCG -31) and R1 reverse (common to both alleles; 51-CCTGTTTTGCACGTTACCG - 3'), with Orc1 primers F1 and R1 for an internal control. The PCR parameters were 94°C for 3 min, 38× (94°C for 30 sec, 60°C for 30 sec, 72°C for 40 sec), 72°C for 3 min. PCR product was run on 3% agarose gel and the transgenic band was detected at 875bp with an internal control band at 439bp.

Alb-cre genotyping was carried out with primers Alb-cre-20239-F (specific for the WT allele; 5'-TGCAAACATCACATGCACAC-3'), Alb-cre-olMR5374-F (specific for the transgenic allele; 5'-GAAGCAGAAGCTTAGGAAGATGG-3') and Alb-cre-20240-R (common to both alleles; 5'-TTGGCCCCTTACCATAACTG-3'). The PCR parameters were 95°C for 30 sec, 35× (95°C for 15 sec, 55°C for 20 sec, 72°C for 60 sec), 72°C for 5 min. PCR product was run on 4% agarose gel and WT band was detected at 351bp, and transgenic band at 390bp.

To establish sex of embryos isolated for MEFs, SRY genotyping was carried out with chromosome Y specific forward (5'-TTGTCTAGAGAGCATGGAGGGCCATGT-3') and reverse primers (5'-CTCCTCTGTGACACTTTAGCCCTCCGA-3'). The PCR parameters were 95°C for 30 sec, 35× (95°C for 15 sec, 55°C for 20 sec, 72°C for 60 sec), 72°C for 5 min. PCR product was run on 2% agarose gel and Y-chromosome positive band was detected at 270bp.

MEF isolation and culture

MEFs were isolated from E12.5 *Orc2*^{+/+} or *Orc2*^{ff} embryos, transformed with SV40 large T antigen and infected with adenovirus Cre-eGFP (#VVC-U of Iowa-1174, University of Iowa). The infected MEFs were cultured for indicated days in DMEM with 10% FBS medium. To measure the cell proliferation, 24 hrs. after the Adenovirus Cre-eGFP transduction, SV40 transformed MEF cells were plated in 96 well plate. The cell viability was measured every 24 hrs using CellTiter 96® Non-Radioactive Cell Proliferation Assay (Promega, #G4100) according to the manufacturer's instructions. All experiments were conducted in triplicate and absorbance relative to that on day 1 was expressed.

Liver isolation

Control (*Orc2*^{ff} *ROSA26*^{stop-EYFP} or *Orc1*^{ff} *Orc2*^{ff} *ROSA26*^{stop-EYFP}) and experimental animals (*Alb-Orc2*^{ff} *ROSA26*^{stop-EYFP} or *Alb-Orc1*^{ff} *Orc2*^{ff} *ROSA26*^{stop-EYFP}) were euthanized using CO₂. Blood was collected was further metabolic studies. Animals were perfused with prewarmed (39°C) Hank's buffered salt solution (HBSS) containing EDTA, MgCl₂, and HEPES. The livers were dissected and weighted. Each liver was divided for following experiments: ploidy analysis (right lobe; fresh processing), EYFP flow cytometry (median lobe; fresh processing), immunoblotting and histology (left lobe; half for LN₂ flash freeze and second half into 10% formalin), and genotyping (caudate lobe; LN₂ flash freeze).

Immunoblotting

SV40 transformed MEF with or without adenovirus Cre-eGFP infection were directly lysed in 2x Laemmli Sample Buffer and sonicated. Liver was lysed in modified RIPA buffer (150 mM Sodium Chloride, 50 mM Tris-HCl, pH 7.4, 1 mM EDTA, 1 mM PMSF, 1% Triton X-100, 1% Sodium Deoxycholic Acid, 0.1% SDS), sonicated, and lysate was clarified by centrifugation. Mouse ORC2 antibody was raised against His tagged full length of ORC2 recombinant protein in Rabbit (Pacific Immunology).

Histology and analysis

All formalin-fixed paraffin-embedded (FFPE) sections and H&E staining were performed by Research Histology Cores at UVA and UAB. Nuclei size was measured using ImageJ 1.50i (Java 1.6.0_24) (56) (Schneider et al., 2012). The number of analyzed animals is annotated at each figure. The statistical method used for comparison between experimental groups was a two-tailed homoscedastic Student's t test. Statistical significance was expressed as a p value. We captured images of 5-10 fields per liver and measured nuclear size relative to scale bar and nuclear density by counting the number of nuclei per field (at a fixed scale to compare between mice). Hepatocyte nuclei can be easily distinguished from stromal nuclei by their roundness and relative de-condensation.

Metabolic measurements

Blood was collected from control *Orc2^{ff} ROSA26^{stop-EYFP}* and experimental *Alb-Orc2^{ff} ROSA26^{stop-EYFP}* animals and centrifuged at 1000g for 10 min at 4°C. To measure Alanine Transaminase (ALT) and Aspartate Aminotransferase (AST) activities EnzyChrom™ Alanine Transaminase Assay Kit (BioAssay Systems, Cat# EALT-100) and Liquid AST (SGOT) Reagent Set (Pointe Scientific, Cat# A7561150) were used respectively according to manufacturers' instructions. The number of analyzed animals is annotated at the figure. The statistical method used for comparison between experimental groups was a two-tailed homoscedastic Student's t test. Statistical significance was expressed as a p value.

Isolation of hepatocytes

Hepatocytes were isolated according to STAR protocols (Charni-Natan and Goldstein, 2020). In brief, the liver perfused with perfusion Buffer for 10 min followed by digestion buffer for 10 min was dissected out and hepatocytes were released into plating medium containing dish. A single-cell suspension was obtained by filtering through a 100-µm cell strainer. Percoll centrifugation (Cytiva #17089102) was performed to remove dead cells. Isolated live Hepatocyte were suspended in William's medium (WEM, GIBCO A1217601) with Plating Supplement (GIBCO #CM3000) and plated on collagen coated cover glass (5x10⁵ cells/6-well). After 3 hrs., the medium was exchanged for maintenance medium (William's E Medium (WEM, GIBCO A1217601) supplemented with GIBCO #CM4000).

Ploidy analysis

Nuclei Isolation Medium (NIM; 250mM Sucrose, 25mM KCl, 5mM MgCl₂, 10mM Tris-Cl, 1mM DTT, 1x Protease inhibitor) with 2% paraformaldehyde and 0.1% Triton X-100 was added to the liver pieces that were subsequently homogenized with Pestle A Dounce homogenizer (25x times). After all samples were processed, they were centrifuged at 1000g for 10 min at 4°C. Pellet was resuspended in NIM and equal volume of 50% iodixanol was added. The mixed sample was carefully layered on the top of 29% Iodixanol solution in ultracentrifuge tube and spun at 10,300rpm for 10 minutes at 4°C in ultracentrifuge. The nuclei pellet was resuspended in FACS buffer (1x PBS, 2.5% (v/v) BSA, 2mM EDTA, 2mM NaN₃) with 100 µg/mL RNase A. DRAQ5-stained liver nuclei samples were processed using Attune NxT flow cytometer

(Life Technologies). Flow cytometry data from 10,000 nuclei were analyzed with FCS express software. The bottom 40% of nuclei on the EYFP axis in the FACS profiles was considered as low EYFP and the top 20% as high EYFP. The EYFP low nuclei are mostly from non-hepatocytes (Kupffer cells, endothelial cells, bile duct cells, contaminating blood cells) and some hepatocytes that have not yet expressed sufficient levels of EYFP and they all serve as a control population. The EYFP high nuclei are exclusively from hepatocytes that have undergone the Cre-mediated recombination a sufficient time back to express high levels of EYFP and are the experimental population with ORC subunit deletion. The statistical method used for comparison between experimental groups was a two-tailed homoscedastic Student's t test. Statistical significance was expressed as a p value.

EdU incorporation in vitro

2 days after plating, Hepatocyte was labeled with 20 μ M of EdU (Lumiprobe #10540) for 2 hrs and fixed with 4% paraformaldehyde for 10 min followed by permeabilized with 0.25 % Tritonx-100 for 5 min. The fixed cells were incubated with label mix [8 μ M Sulfo-Cy3-Azide (Lumiprobe #B1330), 2 mM CuSO₄*5H₂O, 113 mM Ascorbic Acid] for 30 min. Anti-GFP antibody (Abcam #ab6556) was used to detect EYFP signal after the EdU staining.

Partial hepatectomy

The partial hepatectomy experiment was conducted following a standardized protocol (Mitchell and Willenbring, 2008 [↗](#)). Briefly, 8-14-week-old mice were utilized. After inducing anesthesia with 2% isoflurane and maintaining at 0.2%, the mice were subcutaneously injected with Buprenorphine-SR (0.6 mg/kg) and Carprofen (5 mg/kg). The abdominal wall was shaved and prepared aseptically. A 3 cm long transverse incision was made to expose the xiphoid. The left and middle lobes were tied using 4-0 thread and cut. The incision site was double-checked for bleeding following washed with 0.9% sodium chloride and sutured. Mice were placed on a warm pad for recovery. To detect DNA synthesis, 0.2mg EDU (Lumiprobe #10540) was injected via the tail vein 3-4 hours before sacrifice. At 36-48 hours post-surgery, samples from the regenerating right liver were collected and weighed. For IHC and IF staining, samples were fixed with 4% formaldehyde, dehydrated, cleared, embedded in paraffin, and sectioned at 5-8 μ m thickness. Tissue sections were deparaffinized and rehydrated, and heat antigen retrieval methods were applied in 10 mM sodium citrate buffer (pH=6.0). Permeabilization with 0.25% Triton X-100 followed EDU incorporation by Click-in reaction for 40 minutes at room temperature, protected from light. After washing and blocking, an anti-GFP antibody (Abcam, cat# ab6556, 1:500 dilution) and Alexa-488 fluorescent-conjugated secondary antibody (A11029; Life Technologies) were used to detect EYFP signaling. Finally, nuclei were stained with DAPI, and images were captured using Zeiss Confocal microscope and processed using ImageJ and GraphPad software. To improve the EYFP signal, we used frozen sections for the liver regeneration experiments in the Orc1, Orc2 double knockout mice in **Fig. 6** [↗](#).

Estimate of nuclear and cell division in hepatocytes during normal development

To estimate the minimal number of cell divisions occurring in the hepatoblasts after the appearance of albumin-driven Cre, we first estimated how many hepatocyte nuclei populate an adult liver. Given that there are 125 million liver cells/gram of tissue (Sohlenius-Sternbeck, 2006), we estimate that there are 162.5 (female) to 218 (male) million hepatocytes for the 1.3 gram (female) or 1.75 gram (male) livers (all estimates of liver weight and liver nuclear density are from **Fig. 5** [↗](#), **6** [↗](#), and **S3b**).

We next estimated the number of albumin-positive cells early in embryogenesis, from published scRNAseq results that suggest that ~1% of the embryonic cells are of hepatocyte lineage when all embryonic cells are harvested from E9.5-E13.5 embryos (Cao et al., 2019 [↗](#)). Given that there are

~200,000 cells in E9.5 embryos (Cao et al., 2019 [↗](#)), we therefore estimate that there are 2,000 albumin positive cells at the stage when albumin mRNA expression is readily detected. Thus, normal development of the liver requires at least 20 rounds of diploid cell division from the 2000 hepatoblasts/hepatocytes seen in E9.5 embryos to produce the ~200 million hepatocyte nuclei in the adult liver (Table 2, bottom row).

From the relative weights of the adult livers (row 1) and the relative densities of the hepatocyte nuclei (row 2) we can estimate the number of hepatocytes in the mutant livers relative to wild type livers (row 3). The relative deficit of hepatocytes in the mutant livers allows us to estimate how many fewer cell divisions the hepatocytes underwent in the mutant livers relative to the WT livers during development (row 4). Since WT hepatocytes undergo ~20 duplications, we thus estimate that the *Orc2*^{-/-} hepatocytes and the male DKO hepatocytes undergo at least 18 divisions from the 2000 hepatoblast-stage seen in E9.5 embryo, while the female DKO hepatocyte undergo at least 15 divisions (Table 2, bottom row). This underestimates the number of divisions because we do not take into account any apoptosis that may be occurring or any endoreduplication cycles.

Acknowledgements

This work was supported by a grant from the NIH (R01 CA60499 to A.D), Wagner fellowship from the University of Virginia (to R.K.P.), and the F99/K00 NCI Predoctoral to Postdoctoral Fellow Transition Award (F99/K00CA253732 to R.K.P.). This publication was also supported in part by the Medical College of Wisconsin Cancer Center Shared Resources. We thank the following funders for grant support: Advancing a Healthier Wisconsin Endowment (G.L., T.U., A.T.), and the Dr. Glenn R. and Nancy A. Linnerson Endowed Fund (G.L.). We thank the University of Virginia (RRid:SCR_017829) and UAB Flow Cytometry Core Facilities and Research Histology Core Facilities, that were partially supported by the NCI Grants (P30-CA044579, P30-CA013148), and thank Dr. Ulrike Lorenz from the University of Virginia for the Alb-Cre breeder, and Kody Park for help with genotyping. All images were created independently and processed using BioRender. Source: Hao, X. (2025). Created in BioRender. Retrieved from <https://BioRender.com/I15k098> [↗](#).

Additional files

Table 1. Embryonic lethality of *Orc2* KO. The *Orc2*^Δ allele was created by expressing Cre recombinase from a Sox2 promoter in the *Orc2*^{f/f} embryos. [↗](#)

Table 2. Estimate of number of hepatocyte nuclei in adult mice of indicate genotypes and thus, number of hepatocyte nuclear divisions required after E9.5 mouse embryos. [↗](#)

Supplemental Fig S1. Serum liver enzymes in *ORC2* wild type (*Orc2*^{f/f}) and *Orc2* KO (*Alb-Orc2*^{f/f}) mice. **p* < 0.05, two-tailed homoscedastic Student's *t* test. [↗](#)

Supplemental Fig S2. *Orc1 Orc2* deletion produces smaller livers with larger nuclei and more mortality in female mice. A. Representative picture of livers of female *Orc1 Orc2* WT and liver-specific double KO (dKO). B. Representative H&E staining of liver tissue from *Orc1*^{f/f} *Orc2*^{f/f} ROSA26^{stop-EYFP} (WT) and *Alb-Orc1*^{f/f} *Orc2*^{f/f} ROSA26^{stop-EYFP} (dKO) females. C. Kaplan-Meier plot for *Alb-Orc1*^{f/f} *Orc2*^{f/f} ROSA26^{stop-EYFP} (dKO) females post natally. [↗](#)

Supplemental Fig S3. From an independent breeding experiment to generate mice with *Orc1*^{-/-}; *Orc2*^{-/-} and *Orc1*^{-/-}, *Orc2*^{-/-} livers. A. Ratio of liver weight to body weight for mice at 6 weeks of age. The *Orc1* mutant, *Orc2* mutant or double mutant mice are in red. B. Enlarged nuclei seen in 6-week mouse livers in

mice expressing Alb-Cre where both alleles of one ORC subunit are floxed (underlined): $Orc1^{ff}$ or $Orc2^{ff}$.
C. Enlarged nuclei seen in 6-week mouse livers in mice expressing Alb-Cre where both alleles of two ORC subunits are floxed (underlined): $Orc1^{ff} Orc2^{ff}$. [🔗](#)

References

- Bell M. R 2018) **Comparing Postnatal Development of Gonadal Hormones and Associated Social Behaviors in Rats, Mice, and Humans** *Endocrinology* **159**:2596–2613
- Bell S. P 2017) **Rethinking origin licensing** *eLife* **6**
- Bell S. P., Kobayashi R., Stillman B 1993) **Yeast origin recognition complex functions in transcription silencing and DNA replication** *Science* **262**:1844–9
- Bell S. P., Stillman B 1992) **ATP-dependent recognition of eukaryotic origins of DNA replication by a multiprotein complex** *Nature* **357**:128–34
- Bermejo R., Vilaboa N., Cales C 2002) **Regulation of Cdc6, geminin, and CDT1 in human cells that undergo polyploidization** *Mol Biol Cell* **13**:3989–4000
- Bicknell L. S., Bongers E. M., Leitch A., Brown S., Schoots J., Harley M. E., Aftimos S., Al-Aama J. Y., Bober M., Brown P. A., van Bokhoven H., Dean J., Edrees A. Y., Feingold M., Fryer A., Hoefsloot L. H., Kau N., Knoers N. V., Mackenzie J., Opitz J. M., Sarda P., Ross A., Temple I. K., Toutain A., Wise C. A., Wright M., Jackson A. P. 2011a) **Mutations in the pre-replication complex cause Meier-Gorlin syndrome** *Nat Genet* **43**:356–9
- Bicknell L. S., Walker S., Klingseisen A., Stiff T., Leitch A., Kerzendorfer C., Martin C. A., Yeyati P., Sanna AL, Bober N., Johnson M., Wise D., Jackson C., O’driscoll A. P., Jeggo M. 2011b) **Mutations in Orc1, encoding the largest subunit of the origin recognition complex, cause microcephalic primordial dwarfism resembling Meier-Gorlin syndrome** *Nat Genet* **43**:350–5
- Bleichert F., Botchan M. R., Berger J. M 2015) **Crystal structure of the eukaryotic origin recognition complex** *Nature* **519**:321–6
- Cao J., Spielmann M., Qiu X., Huang X., Ibrahim D. M., Hill A. J., Zhang F., Mundlos S., Christiansen L., Steemers F. J., Trapnell C., Shendure J. 2019) **The single-cell transcriptional landscape of mammalian organogenesis** *Nature* **566**:496–502
- Charni-Natan M., Goldstein I. 2020) **Protocol for Primary Mouse Hepatocyte Isolation** *STAR Protoc* **1**:100086
- Chou H. C., Bhalla K., Demerdesh O. E., Klingbeil O., Hanington K., Aganezov S., Andrews P., Alsudani H., Chang K., Vakoc C. R., Schatz M. C., McCombie W. R., Stillman B. 2021) **The human origin recognition complex is essential for pre-RC assembly, mitosis, and maintenance of nuclear structure** *eLife* **10**
- Clarey M. G., Erzberger J. P., Grob P., Leschziner A. E., Berger J. M., Nogales E., Botchan M. 2006) **Nucleotide-dependent conformational changes in the DnaA-like core of the origin recognition complex** *Nat Struct Mol Biol* **13**:684–90
- Conforto T. L., Waxman D. J 2012) **Sex-specific mouse liver gene expression: genome-wide analysis of developmental changes from pre-pubertal period to young adulthood** *Biol Sex Differ* **3**:9

- Costa A., Diffley J. F. X 2022) **The Initiation of Eukaryotic DNA Replication** *Annu Rev Biochem* **91**:107–131
- de Renty C., Kaneko K. J., Depamphilis M. L. 2014) **The dual roles of geminin during trophoblast proliferation and differentiation** *Dev Biol* **387**:49–63
- Dhar S. K., Delmolino L., Dutta A 2001a) **Architecture of the human origin recognition complex** *J Biol Chem* **276**:29067–71
- Dhar S. K., Yoshida K., Machida Y., Khaira P., Chaudhuri B., Wohlschlegel J. A., Leffak M., Yates J., Dutta A 2001b) **Replication from oriP of Epstein-Barr virus requires human ORC and is inhibited by geminin** *Cell* **106**:287–96
- Foss M., McNally F. J., Laurenson P., Rine J 1993) **Origin recognition complex (ORC) in transcriptional silencing and DNA replication in *S. cerevisiae*** *Science* **262**:1838–44
- GIORDANO-Coltart J., Ying C. Y., Gautier J., Hurwitz J. 2005) **Studies of the properties of human origin recognition complex and its Walker A motif mutants** *Proc Natl Acad Sci U S A* **102**:69–74
- Gonzalez M. A., Tachibana K. E., Adams D. J., van der Weyden L., Hemberger M., Coleman N., Bradley A., Laskey R. A. 2006) **Geminin is essential to prevent endoreduplication and to form pluripotent cells during mammalian development** *Genes Dev* **20**:1880–4
- Gualdi R., Bossard P., Zheng M., Hamada Y., Coleman J. R., Zaret K. S. 1996) **Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control** *Genes Dev* **10**:1670–82
- Guernsey D. L., Matsuoka M., Jiang H., Evans S., Macgillivray C., Nightingale M., Perry S., Ferguson M., Leblanc M., Paquette J., Patry L., Rideout A. L., Thomas A., Orr A., McMaster C. R., Michaud J. L., Deal C., Langlois S., Superneau D. W., Parkash S., Ludman M., Skidmore D. L., Samuels M. E 2011) **Mutations in origin recognition complex gene ORC4 cause Meier-Gorlin syndrome** *Nat Genet* **43**:360–4
- Hayashi S., Lewis P., Pevny L., McMahon A. P 2002) **Efficient gene modulation in mouse epiblast using a Sox2Cre transgenic mouse strain** *Mech Dev* **119**:S97–S101
- Hu Y., Stillman B 2023) **Origins of DNA replication in eukaryotes** *Mol Cell* **83**:352–372
- Janbandhu V. C., Moik D., Fässler R 2014) **Cre recombinase induces DNA damage and tetraploidy in the absence of loxP sites** *Cell Cycle* **13**:462–70
- Lilly M. A., Duronio R. J. 2005) **New insights into cell cycle control from the *Drosophila* endocycle** *Oncogene* **24**:2765–75
- Loo S., Fox C. A., Rine J., Kobayashi R., Stillman B., Bell S 1995) **The origin recognition complex in silencing, cell cycle progression, and DNA replication** *Mol Biol Cell* **6**:741–56
- Loonstra A., Vooijs M., Beverloo H. B., Allak B. A., van Drunen E., Kanaar R., Berns A., Jonkers J 2001) **Growth inhibition and DNA damage induced by Cre recombinase in mammalian cells** *Proc Natl Acad Sci U S A* **98**:9209–14
- Micklem G., Rowley A., Harwood J., Nasmyth K., Diffley J. F 1993) **Yeast origin recognition complex is involved in DNA replication and transcriptional silencing** *Nature* **366**:87–9

- Mitchell C., Willenbring H 2008) **A reproducible and well-tolerated method for 2/3 partial hepatectomy in mice** *Nat Protoc* **3**:1167–70
- Naiche L. A., Papaioannou V. E 2007) **Cre activity causes widespread apoptosis and lethal anemia during embryonic development** *Genesis* **45**:768–75
- Neuwald A. F., Aravind L., Spouge J. L., Koonin E. V. 1999) **AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes** *Genome Res* **9**:27–43
- OKANO-Uchida T., Kent L. N., Ouseph M. M., Mccarty B., Frank J. J., Kladney R., Cuitino M. C., Thompson J. C., Coppola V., Asano M., Leone G 2018) **Endoreduplication of the mouse genome in the absence of ORC1** *Genes Dev* **32**:978–990
- Park S. Y., Asano M 2008) **The origin recognition complex is dispensable for endoreplication in Drosophila** *Proc Natl Acad Sci U S A* **105**:12343–8
- Postic C., Shiota M., Niswender K. D., Jetton T. L., Chen Y., Moates J. M., Shelton K. D., Lindner J., Cherrington A. D., Magnuson M. A 1999) **Dual roles for glucokinase in glucose homeostasis as determined by liver and pancreatic beta cell-specific gene knock-outs using Cre recombinase** *J Biol Chem* **274**:305–15
- Prasanth S. G., Prasanth K. V., Siddiqui K., Spector D. L., Stillman B. 2004) **Human Orc2 localizes to centrosomes, centromeres and heterochromatin during chromosome inheritance** *EMBO J* **23**:2651–63
- Saha P., Chen J., Thome K. C., Lawlis S. J., Hou Z. H., Hendricks M., Parvin J. D., Dutta A. 1998) **Human CDC6/Cdc18 associates with Orc1 and cyclin-cdk and is selectively eliminated from the nucleus at the onset of S phase** *Mol Cell Biol* **18**:2758–67
- SCHMIDT-Suppran, M. & Rajewsky, K 2007) **Vagaries of conditional gene targeting** *Nat Immunol* **8**:665–8
- Schmidt E. E., Taylor D. S., Prigge J. R., Barnett S., Capecchi M. R. 2000) **Illegitimate Cre-dependent chromosome rearrangements in transgenic mouse spermatids** *Proc Natl Acad Sci U S A* **97**:13702–7
- Schneider C. A., Rasband W. S., Eliceiri K. W. 2012) **NIH Image to ImageJ: 25 years of image analysis** *Nat Methods* **9**:671–5
- Schuler M., Dierich A., Chambon P., Metzger D 2004) **Efficient temporally controlled targeted somatic mutagenesis in hepatocytes of the mouse** *Genesis* **39**:167–72
- Semple J. W., DA-Silva L. F., Jervis E. J., AH-Kee J., AL-Attar H., Kummer L., Heikkila J. J., Pasero P., Duncker B. 2006) **An essential role for Orc6 in DNA replication through maintenance of pre-replicative complexes** *EMBO J* **25**:5150–8
- Shibata E., Dutta A 2020) **A human cancer cell line initiates DNA replication normally in the absence of ORC5 and ORC2 proteins** *J Biol Chem* **295**:16949–16959
- Shibata E., Kiran M., Shibata Y., Singh S., Kiran S., Dutta A. 2016) **Two subunits of human ORC are dispensable for DNA replication and proliferation** *eLife* **5**

- Shibata Y., Peycheva M., Shibata E., Malzl D., Pavri R., Dutta A. 2024) **Specific origin selection and excess functional MCM2-7 loading in ORC-deficient cells** *bioRxiv*
- Shu Z., Row S., Deng W. M 2018) **Endoreplication: The Good, the Bad, and the Ugly** *Trends Cell Biol* **28**:465–474
- SOHLENIUS-Sternbeck, A. K 2006) **Determination of the hepatocellularity number for human, dog, rabbit, rat and mouse livers from protein concentration measurements** *Toxicol In Vitro* **20**:1582–6
- Stillman B 2022) **The remarkable gymnastics of ORC** *eLife* **11**
- Su T. T., O'farrell P. H 1998) **Chromosome association of minichromosome maintenance proteins in Drosophila endoreplication cycles** *J Cell Biol* **140**:451–60
- Su Z., Tian M., Shibata E., Shibata Y., Yang T., Wang Z., Jin F., Zang C., Dutta A. 2024) **Regulation of epigenetics and chromosome structure by human ORC2** *bioRxiv*
- Takeda D. Y., Shibata Y., Parvin J. D., Dutta A. 2005) **Recruitment of ORC or CDC6 to DNA is sufficient to create an artificial origin of replication in mammalian cells** *Genes Dev* **19**:2827–36
- Tilghman S. M., Belayew A 1982) **Transcriptional control of the murine albumin/alpha-fetoprotein locus during development** *Proc Natl Acad Sci U S A* **79**:5254–7
- Vermeulen M., Eberl H. C., Matarese F., Marks H., Denissov S., Butter F., Lee K. K., Olsen J. V., Hyman A. A., Stunnenberg H. G., Mann M. 2010) **Quantitative interaction proteomics and genome-wide profiling of epigenetic histone marks and their readers** *Cell* **142**:967–80
- Wang X., Yang L., Wang Y. C., Xu Z. R., Feng Y., Zhang J., Wang Y., Xu C. R 2020) **Comparative analysis of cell lineage differentiation during hepatogenesis in humans and mice at the single-cell transcriptome level** *Cell Res* **30**:1109–1126
- Welch M 1992) **Management of generalized faecal peritonitis** *J R Soc Med* **85**:246
- Williams R. S., Shohet R. V., Stillman B 1997) **A human protein related to yeast Cdc6p** *Proc Natl Acad Sci U S A* **94**:142–7

Author information

Róża K Przanowska*

Dept. of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, United States

ORCID iD: [0000-0003-1278-6242](https://orcid.org/0000-0003-1278-6242)

*Co-first-authors

Yuechuan Chen*

Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States

*Co-first-authors

Takayuki-Okano Uchida

Cancer Center, University of Wisconsin in Milwaukee, Milwaukee, United States

Etsuko Shibata

Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States

Xiaoxiao Hao

Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States

Isaac Segura Rueda

Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States

Kate Jensen

Dept. of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, United States

Piotr Przanowski

Dept. of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, United States

Anthony Trimboli

Cancer Center, University of Wisconsin in Milwaukee, Milwaukee, United States

Yoshiyuki Shibata

Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States

Gustavo Leone

Cancer Center, University of Wisconsin in Milwaukee, Milwaukee, United States

Anindya Dutta

Dept. of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, United States, Dept. of Genetics, University of Alabama in Birmingham, Birmingham, United States
ORCID iD: [0000-0002-4319-0073](https://orcid.org/0000-0002-4319-0073)

For correspondence: duttaa@uab.edu

Editors

Reviewing Editor

Yamini Dalal

National Cancer Institute, Bethesda, United States of America

Senior Editor

Yamini Dalal

National Cancer Institute, Bethesda, United States of America

Reviewer #1 (Public review):

The origin recognition complex (ORC) is an essential loading factor for the replicative Mcm2-7 helicase complex. Despite ORC's critical role in DNA replication, there have been instances where the loss of specific ORC subunits has still seemingly supported DNA replication in cancer cells, endocycling hepatocytes, and *Drosophila* polyploid cells. Critically, all tested ORC subunits are essential for development and proliferation in normal cells. This presents a challenge, as conditional knockouts need to be generated, and a skeptic can always claim that there were limiting but sufficient ORC levels for helicase loading and replication in polyploid or transformed cells. That being said, the authors have consistently pushed the system to demonstrate replication in the absence or extreme depletion of ORC subunits.

Here, the authors generate conditional ORC2 mutants to counter a potential argument with prior conditional ORC1 mutants that Cdc6 may substitute for ORC1 function based on homology. They also generate a double ORC1 and ORC2 mutant, which is still capable of DNA replication in polyploid hepatocytes. While this manuscript provides significantly more support for the ability of select cells to replicate in the absence or near absence of select ORC subunits, it does not shed light on a potential mechanism.

The strengths of this manuscript are the mouse genetics and the generation of conditional alleles of ORC2 and the rigorous assessment of phenotypes resulting from limiting amounts of specific ORC subunits. It also builds on prior work with ORC1 to rule out Cdc6 complementing the loss of ORC1.

The weakness is that it is a very hard task to resolve the fundamental question of how much ORC is enough for replication in cancer cells or hepatocytes. Clearly, there is a marked reduction in specific ORC subunits that is sufficient to impact replication during development and in fibroblasts, but the devil's advocate can always claim minimal levels of ORC remaining in these specialized cells.

The significance of the work is that the authors keep improving their conditional alleles (and combining them), thus making it harder and harder (but not impossible) to invoke limiting but sufficient levels of ORC. This work lays the foundation for future functional screens to identify other factors that may modulate the response to the loss of ORC subunits.

This work will be of interest to the DNA replication, polyploidy, and genome stability communities.

<https://doi.org/10.7554/eLife.102915.2.sa3>

Reviewer #2 (Public review):

This manuscript proposes that primary hepatocytes can replicate their DNA without the six-subunit ORC. This follows previous studies that examined mice that did not express ORC1 in the liver. In this study, the authors suppressed expression of ORC2 or ORC1 plus ORC2 in the liver.

Comments:

(1) I find the conclusion of the authors somewhat hard to accept. Biochemically, ORC without the ORC1 or ORC2 subunits cannot load the MCM helicase on DNA. The question arises whether the deletion in the ORC1 and ORC2 genes by Cre is not very tight, allowing some cells to replicate their DNA and allow the liver to develop, or whether the replication of DNA proceeds via non-canonical mechanisms, such as break-induced replication. The increase in the number of polyploid cells in the mice expressing Cre supports the first mechanism,

because it is consistent with few cells retaining the capacity to replicate their DNA, at least for some time during development.

(2) Fig 1H shows that 5 days post infection, there is no visible expression of ORC2 in MEFs with the ORC2 flox allele. However, at 15 days post infection, some ORC2 is visible. The authors suggest that a small number of cells that retained expression of ORC2 were selected over the cells not expressing ORC2. Could a similar scenario also happen *in vivo*?

(3) Figs 2E-G show decreased body weight, decreased liver weight and decreased liver to body weight in mice with recombination of the ORC2 flox allele. This means that DNA replication is compromised in the ALB-ORC2f/f mice.

(4) Figs 2I-K do not report the number of hepatocytes, but the percent of hepatocytes with different nuclear sizes. I suspect that the number of hepatocytes is lower in the ALB-ORC2f/f mice than in the ORC2f/f mice. Can the authors report the actual numbers?

(5) Figs 3B-G do not report the number of nuclei, but percentages, which are plotted separately for the ORC2-f/f and ALB-ORC2-f/f mice. Can the authors report the actual numbers?

(6) Fig 5 shows the response of ORC2f/f and ALB-ORC2f/f mice after partial hepatectomy. The percent of EdU+ nuclei in the ORC2-f/f (aka ALB-CRE-/-) mice in Fig 5H seems low. Based on other publications in the field it should be about 20-30%. Why is it so low here? The very low nuclear density in the ALB-ORC2-f/f mice (Fig 5F) and the large nuclei (Fig 5I) could indicate that cells fire too few origins, proceed through S phase very slowly and fail to divide.

(7) Fig 6F shows that ALB-ORC1f/f-ORC2f/f mice have very severe phenotypes in terms of body weight and liver weight (about on third of wild-type!). Fig 6H and 6I, the actual numbers should be presented, not percentages. The fact that there are EYFP negative cells, implies that CRE was not expressed in all hepatocytes.

(8) Comparing the EdU+ cells in Fig 7G versus 5G shows very different number of EdU+ cells in the control animals. This means that one of these images is not representative. The higher fraction of EdU+ cells in the double-knockout could mean that the hepatocytes in the double-knockout take longer to complete DNA replication than the control hepatocytes. The control hepatocytes may have already completed DNA replication, which can explain why the fraction of EdU+ cells is so low in the controls. The authors may need to study mice at earlier time points after partial hepatectomy, i.e. sacrifice the mice at 30-32 hours, instead of 40-52 hours.

(9) Regarding the calculation of the number of cell divisions during development: the authors assume that all the hepatocytes in the adult liver are derived from hepatoblasts that express Alb. Is it possible to exclude the possibility that pre-hepatoblast cells that do not express Alb give rise to hepatocytes? For example, the cells that give rise to hepatoblasts may proliferate more times than normal giving rise to a higher number of hepatoblasts than in wild-type mice.

(10) My interpretation of the data is that not all hepatocytes have the ORC1 and ORC2 genes deleted (eg EYFP-negative cells) and that these cells allow some proliferation in the livers of these mice.

My comments regarding the previous version still stand, since the authors did not perform experiments to address them.

<https://doi.org/10.7554/eLife.102915.2.sa2>

Reviewer #3 (Public review):**Summary:**

The authors address the role of ORC in DNA replication and that this protein complex is not essential for DNA replication in hepatocytes. They provide evidence that ORC subunit levels are substantially reduced in cells that have been induced to delete multiple exons of the corresponding ORC gene(s) in hepatocytes. They evaluate replication both in purified isolated hepatocytes and in mice after hepatectomy. In both cases, there is clear evidence that DNA replication does not decrease at a level that corresponds with the decrease in detectable ORC subunit and that endoreduplication is the primary type of replication observed. It remains possible that small amounts of residual ORC are responsible for the replication observed, although the authors provide arguments against this possibility. The mechanisms responsible for the DNA replication observed in the absence of ORC are not examined, including why such replication would primarily be due to endoreduplication.

Strengths:

The authors clearly show that there are dramatic reductions in the amount of the targeted ORC subunits in the cells that have been targeted for deletion. They also provide clear evidence that there is replication in a subset of these cells and that it is likely due to endoreduplication. Although there is no replication in MEFs derived from cells with the deletion, there is clearly DNA replication occurring in hepatocytes (both isolated in culture and in the context of the liver). Interestingly, the cells undergoing replication exhibit enlarged cell sizes and elevated ploidy indicating endoreduplication of the genome. These findings raise the interesting possibility that endoreduplication does not require ORC while normal replication does.

Weaknesses:

There remain two significant weaknesses in this manuscript. The first is that although there is clearly robust reduction of the targeted ORC subunit, the authors cannot confirm that it is deleted in all cells. For example, the analysis in Fig. 4B would suggest that a substantial number of cells have not lost the targeted region of ORC2. In their response, the authors suggest that this is due to contaminating non-hepatocyte cells but do not provide evidence that this is the case. Although the western blots show stronger effects, this type of analysis is notorious for non-linear response curves and no standards are not provided. The second weakness is that there is no evaluation of the molecular nature of the replication observed. In response to the initial review the authors point out that a previous publication mapped Mcm2-7 loading in the absence of ORC1, ORC2 and ORC5 and saw no deficit or altered location. Unfortunately, this is not done for the mutants discussed here and this previous data supports a model that limiting residual ORC is responsible for the replication observed rather than some novel mechanism (which would be expected to alter location or amounts of loading). The manuscript provides no exploration of why "ORC-independent" replication would drive endoreduplication (which is the strongest evidence for an alternative mechanism of initiation but is unique to this experiment and not the previously mutants analyzed for Mcm2-7 loading). Most importantly, it remains true that after numerous papers from this lab and others claiming that ORC is not required for eukaryotic DNA replication, we still have no information about an alternative pathway that could explain Mcm2-7 loading in the absence of ORC. Without some insights in this area, studies such as these will remain controversial.

<https://doi.org/10.7554/eLife.102915.2.sa1>

Author response:

The following is the authors' response to the original reviews.

eLife Assessment

This descriptive manuscript builds on prior research showing that the elimination of Origin Recognition Complex (ORC) subunits does not halt DNA replication. The authors use various methods to genetically remove one or two ORC subunits from specific tissues and observe continued replication, though it may be incomplete. The replication appears to be primarily endoreduplication, indicating that ORC-independent replication may promote genome reduplication without mitosis. Despite similar findings in previous studies, the paper provides convincing genetic evidence in mice that liver cells can replicate and undergo endoreduplication even with severely depleted ORC levels. While the mechanism behind this ORC-independent replication remains unclear, the study lays the groundwork for future research to explore how cells compensate for the absence of ORC and to develop functional approaches to investigate this process. The reviewers agree that this valuable paper would be strengthened significantly if the authors could delve a bit deeper into the nature of replication initiation, potentially using an origin mapping experiment. Such an exciting contribution would help explain the nature of the proposed new type of Mcm loading, thereby increasing the impact of this study for the field at large.

We appreciate the reviewers' suggestion. Till now we know of only one paper that mapped origins of replication in regenerating mouse liver, and that was published two months back in Cell (PMID: 39293447). We want to adopt this method, but we do not need it to answer the question asked. We have mapped origins of replication in ORC-deleted cancer cell lines and compared to wild-type cells in Shibata et al., BioRxiv (PMID: 39554186) (it is under review). We report the following: Mapping of origins in cancer cell lines that are wild type or engineered to delete three of the subunits, *ORC1*, *ORC2* or *ORC5* shows that specific origins are still used and are mostly at the same sites in the genome as in wild type cells. Of the 30,197 origins in wild type cells (with ORC), only 2,466 (8%) are not used in any of the three ORC deleted cells and 18,319 (60%) are common between the four cell types. Despite the lack of ORC, excess MCM2-7 is still loaded at comparable rates in G1 phase to license reserve origins and is also repeatedly loaded in the same S phase to permit re-replication.

Citation: Specific origin selection and excess functional MCM2-7 loading in ORC-deficient cells. Yoshiyuki Shibata, Mihaela Peycheva, Etsuko Shibata, Daniel Malzl, Rushad Pavri, Anindya Dutta. bioRxiv 2024.10.30.621095; doi: <https://doi.org/10.1101/2024.10.30.621095> (PMID: 39554186)

We have now included this in the discussion.

Public Reviews:

Reviewer #1 (Public review):

The origin recognition complex (ORC) is an essential loading factor for the replicative Mcm2-7 helicase complex. Despite ORC's critical role in DNA replication, there have been instances where the loss of specific ORC subunits has still seemingly supported DNA replication in cancer cells, endocycling hepatocytes, and Drosophila polyploid cells. Critically, all tested ORC subunits are essential for development and proliferation in normal cells. This presents a challenge, as conditional knockouts need to be generated, and a skeptic can always claim that there were limiting but sufficient ORC levels for helicase loading and replication in polyploid or transformed cells. That being said, the

authors have consistently pushed the system to demonstrate replication in the absence or extreme depletion of ORC subunits.

Here, the authors generate conditional ORC2 mutants to counter a potential argument with prior conditional ORC1 mutants that Cdc6 may substitute for ORC1 function based on homology. They also generate a double ORC1 and ORC2 mutant, which is still capable of DNA replication in polyploid hepatocytes. While this manuscript provides significantly more support for the ability of select cells to replicate in the absence or near absence of select ORC subunits, it does not shed light on a potential mechanism.

The strengths of this manuscript are the mouse genetics and the generation of conditional alleles of ORC2 and the rigorous assessment of phenotypes resulting from limiting amounts of specific ORC subunits. It also builds on prior work with ORC1 to rule out Cdc6 complementing the loss of ORC1.

The weakness is that it is a very hard task to resolve the fundamental question of how much ORC is enough for replication in cancer cells or hepatocytes. Clearly, there is a marked reduction in specific ORC subunits that is sufficient to impact replication during development and in fibroblasts, but the devil's advocate can always claim minimal levels of ORC remaining in these specialized cells.

The significance of the work is that the authors keep improving their conditional alleles (and combining them), thus making it harder and harder (but not impossible) to invoke limiting but sufficient levels of ORC. This work lays the foundation for future functional screens to identify other factors that may modulate the response to the loss of ORC subunits.

This work will be of interest to the DNA replication, polyploidy, and genome stability communities.

Thank you.

Reviewer #2 (Public review):

This manuscript proposes that primary hepatocytes can replicate their DNA without the six-subunit ORC. This follows previous studies that examined mice that did not express ORC1 in the liver. In this study, the authors suppressed expression of ORC2 or ORC1 plus ORC2 in the liver.

Comments:

(1) I find the conclusion of the authors somewhat hard to accept. Biochemically, ORC without the ORC1 or ORC2 subunits cannot load the MCM helicase on DNA. The question arises whether the deletion in the ORC1 and ORC2 genes by Cre is not very tight, allowing some cells to replicate their DNA and allow the liver to develop, or whether the replication of DNA proceeds via non-canonical mechanisms, such as break-induced replication. The increase in the number of polyploid cells in the mice expressing Cre supports the first mechanism, because it is consistent with few cells retaining the capacity to replicate their DNA, at least for some time during development.

In our study, we used EYFP as a marker for Cre recombinase activity. ~98% of the hepatocytes in tissue sections and cells in culture express EYFP, suggesting that the majority of hepatocytes successfully expressed the Cre protein to delete the *ORC1* or *ORC2* genes. To assess deletion efficiency, we employed sensitive genotyping and Western blotting techniques to confirm the deletion of *ORC1* and *ORC2* in hepatocytes isolated from Alb-Cre mice. Results in Fig. 2C and Fig. 6D demonstrate the near-complete absence of ORC2 and ORC1 proteins, respectively, in these hepatocytes.

The mutant hepatocytes underwent at least 15–18 divisions during development. The inherited ORC1 or ORC2 protein present during the initial cell divisions, would be diluted to less than 1.5% of wild-type levels within six divisions, making it highly unlikely to support DNA replication, and yet we observe hepatocyte numbers that suggest there was robust cell division even after that point.

Furthermore, the EdU incorporation data confirm DNA synthesis in the absence of ORC1 and ORC2. Specifically, immunofluorescence showed that both *in vitro* and *in vivo*, EYFP-positive hepatocytes (indicating successful *ORC1* and *ORC2* deletion) incorporated EdU, demonstrating that DNA synthesis can occur without ORC1 and ORC2.

Finally, the Alb-ORC2f/f mice have 25-37.5% of the number of hepatocyte nuclei compared to WT mice (Table 2). If that many cells had an undeleted *ORC2* gene, that would have shown up in the genotyping PCR and in the Western blots.

(2) Fig 1H shows that 5 days post infection, there is no visible expression of ORC2 in MEFs with the ORC2 flox allele. However, at 15 days post infection, some ORC2 is visible. The authors suggest that a small number of cells that retained expression of ORC2 were selected over the cells not expressing ORC2. Could a similar scenario also happen *in vivo*?

This would not explain the significant incorporation of EdU in hepatocytes that are EYFP positive and do not have detectable ORC by Western blots. Also note that for MEFs we are delivering the Cre by Adenovirus infection *in vitro*, so there is a finite probability that a cell will not receive the virus, the Cre and will not delete ORC2. However, *in vivo*, the Alb-Cre will be expressed in every cell that turns on albumin. There is no escaping the expression of Cre.

(3) Figs 2E-G shows decreased body weight, decreased liver weight and decreased liver to body weight in mice with recombination of the ORC2 flox allele. This means that DNA replication is compromised in the ALB-ORC2f/f mice.

It is possible that DNA replication is partially compromised or may slow down in the absence of ORC2. However, it is important to emphasize that livers with ORC2 deletion remain capable of DNA replication, so much so that liver function and life span are near normal. Therefore, some kind of DNA replication has to serve as a compensatory mechanism in the absence of ORC2 to maintain liver function and support regeneration.

(4) Figs 2I-K do not report the number of hepatocytes, but the percent of hepatocytes with different nuclear sizes. I suspect that the number of hepatocytes is lower in the ALB-ORC2f/f mice than in the ORC2f/f mice. Can the authors report the actual numbers?

We show in Table 2 that the Alb-Orc2f/f mice have about 25-37.5% of the hepatocytes compared to the WT mice.

(5) Figs 3B-G do not report the number of nuclei, but percentages, which are plotted separately for the ORC2-f/f and ALB-ORC2-f/f mice. Can the authors report the actual numbers?

In all the FACS experiments in Fig. 3B-G we collect data for a total of 10,000 nuclei (or cells). For Fig. 3E-G we divide the 10,000 nuclei into the bottom 40% on the EYFP axis (EYFP low, which is mostly EYFP negative) as the control group, and EYFP high (top 20% on the EYFP axis) test group. We have described this in the Methods in the revision and labeled EYFP negative and positive as EYFP low and high in the Figures and Figure legends.

*(6) Fig 5 shows the response of *ORC2^{f/f}* and *ALB-ORC2^{f/f}* mice after partial hepatectomy. The percent of EdU+ nuclei in the *ORC2^{f/f}* (aka *ALB-CRE^{-/-}*) mice in Fig 5H seems low. Based on other publications in the field it should be about 20-30%. Why is it so low here? The very low nuclear density in the *ALB-ORC2^{f/f}* mice (Fig 5F) and the large nuclei (Fig 5I) could indicate that cells fire too few origins, proceed through S phase very slowly and fail to divide.*

The percentage of EdU+ nuclei in the *ORC2^{f/f}* without Alb-Cre mice is 8%, while in PMID 10623657 ~10% of wild type nuclei incorporate EdU at 42 hr post partial hepatectomy (mid-point between the 36-48 hr post hepatectomy that was used in our study). The important result here is that in the *ORC2^{f/f}* mice with Alb-Cre (+/-) we are seeing significant EdU incorporation. We have also corrected the X-axis labels in 5F, 5I, 7E and 7F to reflect that those measurements were not made at 36 hr post-resection but later (as was indicated in the schematic in Fig. 5A).

*(7) Fig 6F shows that *ALB-ORC1^{f/f}-ORC2^{f/f}* mice have very severe phenotypes in terms of body weight and liver weight (about on third of wild-type!!). Fig 6H and 6I, the actual numbers should be presented, not percentages. The fact that there are EYFP negative cells, implies that CRE was not expressed in all hepatocytes.*

The liver weight is very dependent on the body weight, and so we have to look at the liver to body weight ratio to determine if it is inordinately small, and the ratio is 70% of the WT. In females the liver and body weight are low (although in proportion to each other), which maybe is what the reviewer is talking about. However, the fact that liver weight and body weight are not as low in males, suggest that this is a gender (hormone?) specific effect and not a DNA replication defect. We had discussed this possibility. We have another paper also in BioRxiv (Su et al. doi.org/10.1101/2024.12.18.629220) that suggests that ORC subunits have significant effect on gene expression, so it is possible that that is what leads to this sexual dimorphism in phenotype. We have now added this to the discussion.

The bottom 40% of nuclei on the EYFP axis in the FACS profiles (what was labeled EYFP negative but will now be called EYFP low) contains mostly non-hepatocytes that are genuinely EYFP negative. Non-hepatocytes (bile duct cells, endothelial cells, Kupffer cells, blood cells) are a significant part of cells in the dissociated liver (as can be seen in the single cell sequencing results in PMID: 32690901). Their presence does not mean that hepatocytes are not expressing Cre. Hepatocytes are nearly 100% EYFP positive, as can be seen in the tissue sections (where the hepatocytes take up most of visual field) and in cells in culture. Also if there are EYFP negative hepatocyte nuclei in the FACS, that still does not rule out EYFP presence in the cytoplasm. The important point from the FACS is that the EYFP high nuclei (which have expressed Cre for the longest period) are polyploid relative to the EYFP low nuclei.

(8) Comparing the EdU+ cells in Fig 7G versus 5G shows very different number of EdU+ cells in the control animals. This means that one of these images is not representative. The higher fraction of EdU+ cells in the double-knockout could mean that the hepatocytes in the double-knockout take longer to complete DNA replication than the control hepatocytes. The control hepatocytes may have already completed DNA replication, which can explain why the fraction of EdU+ cells is so low in the controls. The authors may need to study mice at earlier time points after partial hepatectomy, i.e. sacrifice the mice at 30-32 hours, instead of 40-52 hours.

The apparent difference that the reviewer comments on stems from differences in nuclear density in the images in Fig. 7G and 5G (also quantitated in Fig. 7F and 5F). The quantitation

in Fig. 7H and 5H show that the % of EdU plus cells are comparable (5-8%).

(9) Regarding the calculation of the number of cell divisions during development: the authors assume that all the hepatocytes in the adult liver are derived from hepatoblasts that express Alb. Is it possible to exclude the possibility that pre-hepatoblast cells that do not express Alb give rise to hepatocytes? For example the cells that give rise to hepatoblasts may proliferate more times than normal giving rise to a higher number of hepatoblasts than in wild-type mice.

Single cell sequencing of mouse liver at e11 shows hepatoblasts expressing hepatocyte specific markers (PMID: 32690901). All the cells annotated from the single-cell seq analysis are differentiated cells arguing against the possibility that undifferentiated endodermal cells (what the reviewer probably means by pre-hepatoblasts) exist at e11. We have added this citation to the paper.

Here is a review that says the hepatoblasts expressing Albumin are present before e13. (<https://www.ncbi.nlm.nih.gov/books/NBK27068/>) says: “The differentiation of bi-potential hepatoblasts into hepatocytes or BECs begins around e13 of mouse development. Initially hepatoblasts express genes associated with both adult hepatocytes (*Hnf4a*, *Albumin*) ...” Thus, we can be certain that hepatoblasts before e13 express albumin. Our calculation of number of cell divisions in Table 2 begins from e12.

The reviewer may be suggesting that ORC deletion leads to the immediate demise of hepatoblasts (despite having inherited ORC protein from the endodermal cells) causing undifferentiated endodermal cells to persist and proliferate much longer than in normal development. We consider it unlikely, but if true it will be very unexpected, both by suggesting that deletion of ORC immediately leads to the death of the hepatoblasts (despite a healthy reserve of inherited ORC protein) and by suggesting that there is a novel feedback mechanism from the death/depletion of hepatoblasts leading to the persistence and proliferation of undifferentiated endodermal cells. We have added the reviewer’s suggestion to the discussion.

(10) My interpretation of the data is that not all hepatocytes have the ORC1 and ORC2 genes deleted (eg EYFP-negative cells) and that these cells allow some proliferation in the livers of these mice.

Please see the reply in question #1. Particularly relevant: “Finally, the *Alb-ORC2^{ff}* mice have 25-37.5% of the number of hepatocyte nuclei compared to WT mice (Table 2). If that many cells had an undeleted ORC2 gene, that would have shown up in the genotyping PCR and in the Western blots.

Reviewer #3 (Public review):

Summary:

The authors address the role of ORC in DNA replication and that this protein complex is not essential for DNA replication in hepatocytes. They provide evidence that ORC subunit levels are substantially reduced in cells that have been induced to delete multiple exons of the corresponding ORC gene(s) in hepatocytes. They evaluate replication both in purified isolated hepatocytes and in mice after hepatectomy. In both cases, there is clear evidence that DNA replication does not decrease at a level that corresponds with the decrease in detectable ORC subunit and that endoreduplication is the primary type of replication observed. It remains possible that small amounts of residual ORC are responsible for the replication observed, although the authors provide arguments

against this possibility. The mechanisms responsible for DNA replication in the absence of ORC are not examined.

Strengths:

The authors clearly show that there are dramatic reductions in the amount of the targeted ORC subunits in the cells that have been targeted for deletion. They also provide clear evidence that there is replication in a subset of these cells and that it is likely due to endoreduplication. Although there is no replication in MEFs derived from cells with the deletion, there is clearly DNA replication occurring in hepatocytes (both isolated in culture and in the context of the liver). Interestingly, the cells undergoing replication exhibit enlarged cell sizes and elevated ploidy indicating endoreduplication of the genome. These findings raise the interesting possibility that endoreduplication does not require ORC while normal replication does.

Weaknesses:

There are two significant weaknesses in this manuscript. The first is that although there is clearly robust reduction of the targeted ORC subunit, the authors cannot confirm that it is deleted in all cells. For example, the analysis in Fig. 4B would suggest that a substantial number of cells have not lost the targeted region of ORC2. Although the western blots show stronger effects, this type of analysis is notorious for non-linear response curves and no standards are provided. The second weakness is that there is no evaluation of the molecular nature of the replication observed. Are there changes in the amount or location of Mcm2-7 loading that is usually mediated by ORC? Does an associated change in Mcm2-7 loading lead to the endoreduplication observed? After numerous papers from this lab and others claiming that ORC is not required for eukaryotic DNA replication in a subset of cells, we still have no information about an alternative pathway that could explain this observation.

We do not see a significant deficit in MCM2-7 loading (amount and rate) in cancer cell lines where we have deleted *ORC1*, *ORC2* or *ORC5* genes separately in Shibata et al. bioRxiv 2024.10.30.621095; doi: <https://doi.org/10.1101/2024.10.30.621095> (PMID: 39554186). This is now cited in the discussion.

The authors frequently use the presence of a Cre-dependent eYFP expression as evidence that the *ORC1* or *ORC2* genes have been deleted. Although likely the best visual marker for this, it is not demonstrated that the presence of eYFP ensures that *ORC2* has been targeted by Cre. For example, based on the data in Fig. 4B, there seems to be a substantial percentage of *ORC2* genes that have not been targeted while the authors report that 100% of the cells express eYFP.

(1) The PCR reactions in Fig. 4B are still contaminated by DNA from non-hepatocyte cells: bile duct cells, endothelial, Kupfer cells and blood cells. Microscopy of cultured cells identifies the hepatocytes unequivocally from their morphology. <2% of the hepatocyte cells in culture in Fig. 4C are eYFP-.

Recommendations for the authors:

Reviewer #2 (Recommendations for the authors):

The authors should present the data as suggested in the review and reformulate their conclusions. If possible, mice should be examined 30-32 hours after partial hepatectomy.

Based on the Literature we chose a time that is consistent with the previous paper from us (Uchida et al., Genes & Dev).

Reviewer #3 (Recommendations for the authors):

(1) It would improve the paper to use single-cell methods (e.g. FISH) to assess the deletion of ORC subunits in the targeted cells.

This is something we will reserve for future studies.

(2) The importance of the paper would be increased dramatically by showing that the elimination of ORC changed the location of Mcm2-7 loading. This would be highly likely if the authors hypothesis that ORC is not involved is true. On the other hand, given ORC's role in origin selection, an observation that the same sites are used but less frequently would support a hypothesis that residual intact ORC is responsible for the replication observed.

Shibata et al (PMID: 39554186) has answered this question. The loss of ORC does not change the locations of origins or even the ability to specify origins. We argue that this is what is to be expected from our hypothesis, that although ORC is clearly important for MCM loading in yeast and in biochemical experiments, something unexpected is going on in human cells. Either a vanishingly small amount of ORC (undetectable by commonly used methods) can load the full complement of MCM2-7 at a rate that is comparable to wild type cells, or there is an ORC-independent mechanism of MCM2-7 loading. This is now added to the discussion.

<https://doi.org/10.7554/eLife.102915.2.sa0>