

Cellular evolution of the hypothalamic preoptic area of behaviorally divergent deer mice

Reviewed Preprint

v2 • February 27, 2025

Revised by authors

Reviewed Preprint

v1 • November 18, 2024

Jenny Chen , Phoebe R Richardson, Christopher Kirby, Sean R Eddy, Hopi E Hoekstra 

Department of Molecular & Cellular Biology, Harvard University, Cambridge, United States • Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States • Howard Hughes Medical Institute, Harvard University, Cambridge, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study identifies species- and sex-specific neuronal cell types and gene expression in the preoptic area (POA) to help understand the evolutionary divergence of social behaviors. The evidence from single-nucleus RNA sequencing and immunostaining is **compelling** and suggests that cellular differences in the POA may contribute to behavioral variations such as mating and parental care that are apparent in two closely related deer mouse species. These rich observations provide an entry point for future hypothesis-driven experiments to demonstrate a causal role for these populations in sex- or species-variable behaviors in vertebrates. These data will be a resource that is of value to behavioral neuroscientists.

<https://doi.org/10.7554/eLife.103109.2.sa4>

Abstract

Genetic variation is known to contribute to the variation of animal social behavior, but the molecular mechanisms that lead to behavioral differences are still not fully understood. Here, we investigate the cellular evolution of the hypothalamic preoptic area (POA), a brain region that plays a critical role in social behavior, across two sister species of deer mice (*Peromyscus maniculatus* and *P. polionotus*) with divergent social systems. These two species exhibit large differences in mating and parental care behavior across species and sex. Using single-nucleus RNA-sequencing, we build a cellular atlas of the POA for males and females of both *Peromyscus* species. We identify four cell types that are differentially abundant across species, two of which may account for species differences in parental care behavior based on known functions of these cell types. Our data further implicate two sex-biased cell types to be important for the evolution of sex-specific behavior. Finally, we show a remarkable reduction of sex-biased gene expression in *P. polionotus*, a monogamous species that also exhibits reduced sexual dimorphism in parental care behavior. Our POA atlas is a powerful resource to investigate how molecular neuronal traits may be evolving to give rise to innate differences in social behavior across animal species.

Introduction

Large, heritable differences in complex behavior are observed even across closely-related species and across sexes within the same species. What are the molecular substrates, encoded by genes, that facilitate this variation in innate animal behavior? Two molecular traits known to govern behavior are: 1) abundances of specific neuronal cell types, which may be controlled by developmental genes involved in neurogenesis and apoptotic pathways, and 2) abundances of proteins within neuronal cells, which may be controlled by genes involved in transcriptional and translational regulation.

Variations in both types of traits have been documented to change behavior across species and sex. In *Drosophila*, loss of song-patterning neurons underlies a species-specific loss of a type of courtship song¹ while the expansion of an olfactory neuronal population contributes to *D. sechellia*'s species-specific specialization for feeding on the noni fruit². In mammals, neuronal number differences mediating behavioral traits have largely been described across sexes rather than species. Neuronal population sizes, particularly of cell types within the preoptic area and bed nucleus of the stria terminalis, are thought to be regulated by sex-specific levels of neuroestrogens that control apoptosis, thereby leading to sex-biased cell abundances and sex-specific control of behavior^{3,4}.

Across species, gene expression level differences of neuronally-important proteins such as neurotransmitter receptors, smell and taste receptors, and protein kinases are also known to cause behavioral differences (see [5] for review). For example, the expression level of the vasopressin 1a receptor gene in the ventral pallidum contributes to the evolution of partner preference formation in monogamous voles^{6,7}. Across sexes, differential expression of sex-steroid receptors in the mammalian brain is well-known⁸, and the resulting consequences for sex-specific gene expression and control of sex-specific behavior is an area of heavy investigation^{9–12}.

Advances in single-cell and single-nucleus RNA-sequencing have enabled high-throughput characterization of both cell type abundance evolution and cell type-specific expression evolution, and recent single-cell profiling of the brain of behaviorally-divergent species have revealed intriguing insights about both modes of evolution^{13–15}. However, the majority of comparative single-cell RNA-sequencing in mammals has been performed across fairly diverged species with many phenotypic differences, such as human and mouse (~90 million years diverged¹⁶) or across primates^{17–21} such as humans and marmosets (~40 million years diverged¹⁶), lending limited insight into which of the specific behavioral difference may be affected by which of the many observed molecular changes.

Closely-related species of *Peromyscus* deer mice that exhibit variation in heritable behaviors gives us an opportunity to more precisely dissect the molecular neuronal mechanisms that can alter innate mammalian behaviors^{22,23}. In particular, *P. maniculatus* and *P. polionotus* diverged only ~2 million years ago²⁴, but lie on opposite ends of the monogamy-promiscuity spectrum^{25,26} and exhibit large, heritable differences in their parental care behavior^{25,26}. Both males and females of the monogamous *P. polionotus* are more parental than the promiscuous *P. maniculatus*^{25,26}. Further, this difference is much more distinct across males: while promiscuous *P. maniculatus* fathers exhibit little to no parental care in laboratory assays, monogamous *P. polionotus* fathers provide care to nearly the same level as *P. polionotus* mothers^{25,26}. These differences provide an opportunity to identify the molecular products that evolve to modulate social behavior, and how they do so in a sexually dimorphic manner.

Evolution of the hypothalamus has been implicated in social behavior differences across many species^{27–29}, including across these two species of *Peromyscus*²⁶. Here, we used single-nucleus RNA-sequencing (snRNA-seq) to profile the preoptic area (POA), a region of the hypothalamus known for its many cell types involved in mating and parenting behavior³⁰, amongst other innate behaviors. We profiled males and females of both *P. maniculatus* and *P. polionotus* species and, by comparing to *Mus* data, identified POA cell types conserved across rodents. We identified cell type abundance and gene expression differences across species and sex, and hypothesized how these differences may contribute to species- and sex-specific behaviors. Finally, we characterized how sex-bias is evolving across these two *Peromyscus* species and propose potential cell type abundance and gene expression changes that may be co-evolving with monogamy.

Results

A transcriptional cell atlas of *Peromyscus* POA

We performed droplet-based snRNA-seq on the POA of 6 males and 6 females from each of the *P. maniculatus* and *P. polionotus* species (Methods) (**Fig 1A**). An atlas of the POA in *Mus musculus* was previously developed that combined single-cell RNA-seq data with imaging- based spatial information and data on specific neuronal populations activated by social behaviors³¹. To create a comparable dataset, we used a dissection strategy identical to that used for creating the *Mus* POA atlas to dissect a region containing the POA and surrounding nuclei (~2.5 mm x 2.5 mm x .75 mm equivalent to *Mus* Bregma +0.5 to -0.6) (Methods, **Fig 1B**). To reduce batch effect, thereby increasing statistical power for detecting species and sex differences, we took advantage of the genetic diversity of our outbred study system and pooled nuclei from a male and female from each of the *P. maniculatus* and *P. polionotus* species (i.e. 4 samples) into each snRNA-seq run (**Fig 1C**). We collected a total of 105,647 nuclei across 6 replicates (i.e. 24 samples) (Methods, Supp **Table 1**). To computationally demultiplex the pooled data, we used bulk RNA-seq or whole genome sequencing data to call single nucleotide polymorphisms (SNPs) for each animal. We then used SNP data to assign sample information to each nucleus in the snRNA-seq dataset (Methods, Supp **Figs 1–3**, Supp Table 2). In total, we were able to confidently assign a single sample to 95,057 nuclei. The remaining 10,590 nuclei appeared to be multiplets containing two or more samples and were removed from subsequent analyses.

We combined data across all samples and clustered the data using Seurat v4.1.3³² (Methods). We first performed a low-granularity clustering using the first 30 principal components (PCs) of our data and a low resolution parameter of 0.2 to obtain a small number of clusters. This procedure identified 21 major cell classes which were assigned as neurons, microglia, astrocytes, immature and mature oligodendrocytes, endothelial, and ependymal cells using known marker genes from *Mus* hypothalamus³³ (Supp Table 3, Supp **Fig 4**). We then focused only on neuronal nuclei (52,121 nuclei), and reclustered the data at a higher granularity using the first 100 PCs of our data and a resolution parameter of 1.8 (Methods). This resulted in 53 neuronal cell clusters (**Fig 1D**). We did not find major biases in sex or species across our clusters, as shown by the even distribution of each sex and species across the UMAP (Supp **Fig 5**). Our cell clusters reflect known cell types of the POA as well as surrounding nuclei of the anterior hypothalamus, including the paraventricular (PVN), supraoptic (SON), suprachiasmatic nuclei (SCN), and more. For example, arginine vasopressin (*Avp*) and oxytocin (*Oxt*) are exclusively and highly expressed in clusters 34 and 40 (Supp **Fig 6**), respectively, and reflect the well-known populations of AVP and OXT neurons that are found in the PVN and SON. Additionally, vasoactive intestinal protein (*Vip*) is highly expressed in clusters 41 and 46 (Supp **Fig 6**), which together represent the VIP neurons of the SCN that play a major role in regulating circadian rhythms. Further inspection of these two clusters finds specific expression of neuromedin S (*Nms*) only in cluster 41 and gastric releasing




Figure 1.

Overview of *Peromyscus* hypothalamus single-cell atlas.

A) Images of *Peromyscus maniculatus* and *Peromyscus polionotus* and summary of their social behavioral differences. Image credit: ©Joel Sartore/ Photo Ark. **B**) Schematic of the preoptic area of the hypothalamus. Dotted boxes indicate the area dissected. **C**) Schematic of pooling and sample assignment strategy used for snRNA-seq. **D**) UMAP visualization of all 52,121 neuronal nuclei collected across 6 males and 6 females from each of the *P. maniculatus* and *P. polionotus* species, colored by cell cluster. **E**) UMAP visualization of all neuronal nuclei, colored by the assigned *Mus* cell type label. Only *Peromyscus* clusters for which a homologous *Mus* cell type label could be found are colored; remaining nuclei are shown in gray. **F**) Heatmap of the proportion of each *Peromyscus* cell cluster (rows) that are assigned to each *Mus* cell type label (columns). Only *Peromyscus* clusters for which a homologous *Mus* cell type label could be found are shown.




Figure 2.

Differential abundance of cell types across species and sex.

A) Schematic of procedure used to test for differential abundance of cell clusters. **B)** Barplots of log₂(fold change) across species (*P. pol* / *P. man*, left) and across sex (male / female, right) for each cell cluster. Bars are colored by -log₁₀(FDR). **C)** Boxplots of cell abundances (y-axis) of four cell types differentially abundant across species and **D)** of two cell types differentially abundant across sex. Cell abundances across samples are normalized using TMM normalization (Methods).




Figure 3.

Immunostaining and cell counts of regional populations of differentially abundant cell types.

A) Representative images of immunoreactive (ir) staining of AVP⁺ neurons in the PVN (top) and SON (bottom) of *P. maniculatus* and *P. polionotus*. Scale bar represents 200μm. **B)** Boxplots of AVP⁺ neuron number counts in the PVN (left) and SON (right) of females and males of *P. maniculatus* (*P. man*) and *P. polionotus* (*P. pol*); ***: $p < 0.001$ (one-sided Mann Whitney test). **C and D)** Same as **A** and **B**, but of OXT⁺ neurons. **E)** Representative images of ir staining of CALB1⁺ neurons in the BNST (top) and SDN-POA (bottom) of a female and male *P. maniculatus* and *P. polionotus*. Scale bar represents 200μm. **F)** Boxplots of the area of CALB1⁺ in the BNST (left) or CALB1⁺ neuron number counts in the SDN-POA (right) of females and males of *P. maniculatus* and *P. polionotus*; **: $p < 0.01$; *: $p < 0.05$; ns: not significant (one-sided Mann Whitney test).

Neuropeptide	Sex-bias	Cell Type	<i>P. maniculatus</i> fold change	<i>P. maniculatus</i> FDR	<i>P. polionotus</i> fold change	<i>P. polionotus</i> FDR
Kiss1	female	i16:Gal/Th (cluster 20)	26.0	1.2E-54	14.9	6E-49
Tac1	female	i18:Gal/Tac2 (cluster 5)	2.1	3.3E-21	1.4	1.9E-03
Gal	female	i18:Gal/Tac2 (cluster 5)	1.6	3.4E-10	1.6	3.4E-08
Tac1	female	cluster 3	1.8	2.0E-09	1.5	8.9E-04
Tac1	male	i20:Gal/Moxd1 (cluster 9)	3.5	1E-30	1.5	9.8E-03
Cck	male	i20:Gal/Moxd1 (cluster 9)	1.7	1.08E-05	2.3	7.5E-16

Table 1.

Neuropeptides conserved in sex-bias in same cell types across *P. maniculatus* and *P. polionotus*

peptide (*Grp*) only in cluster 46 (Supp Fig 6 [↗](#)), suggesting that these two clusters actually represent two subpopulations of VIP neurons, *Vip/Nms* and *Vip/Grp*, that have recently been reported as genetically and functionally distinct cell types^{34,35} [↗](#).

While some neuropeptidergic cell types were obviously identifiable in our data, they were the minority of our cell clusters. We sought to provide more biological context for the rest of our data by identifying homology between our cell clusters and cell types defined from the *Mus* POA atlas³¹ [↗](#). We used Seurat's integration workflow³⁶ [↗](#) to identify conserved gene correlation structures across the *Mus* and *Peromyscus* datasets and transfer *Mus* cell type labels onto each *Peromyscus* nucleus (Methods, Fig 1E [↗](#)). Because the *Mus* POA atlas consists of separate inhibitory and excitatory datasets, we first defined each of our nuclei as inhibitory or excitatory based on whether it expressed higher levels of inhibitory markers *Gad1* and *Gad2* or excitatory marker *Slc17a6* (Methods, Supp Fig 7). We then transferred inhibitory and excitatory *Mus* cell type labels only onto inhibitory and excitatory nuclei, respectively.

For each *Peromyscus* cluster, we calculated the proportion of nuclei within the cluster assigned to each *Mus* cell type and labeled clusters if at least 15% of the cluster was predicted to belong to the same *Mus* cell type label (Methods). This procedure identified mappings for 34 of our 53 neuronal cell clusters (Fig 1E [↗](#), 1F, Supp Fig 8). (*Mus* cell type labels were denoted by [31] as inhibitory (i) or excitatory (e), and named after marker genes identified from *Mus* POA atlas.) The cell clusters for which we were able to assign *Mus* homology include four cell types activated by relevant social behaviors in *Mus*: cluster 5 (i18: *Gal/Tac2*), which is activated by mating behaviors; cluster 17 (i23: *Crh/Nts*) which is activated by parenting behaviors; and clusters 20 (i16: *Gal/Th*) and 9 (i20: *Gal/Moxd1*), which are activated by both mating and parenting behaviors³¹ [↗](#).

We were unable to assign *Mus* homology for 19 *Peromyscus* cell clusters. These cell clusters had similar numbers of reads and genes per nuclei compared to clusters for which we did find *Mus* homology (Supp Fig 9), and therefore we do not believe lack of homology assignment is due to technical artifacts. Instead, clusters unassignable to *Mus* cell types likely fall into two categories: First, dissection and/or anatomical differences between *Peromyscus* and *Mus* may influence the presence and absence of certain cell types. For example, cluster 2 forms a clear “island” on our UMAP and is transcriptionally specified by high expression of *Pvalb*, *Gad1*, and *Gad2* (Fig 1D [↗](#), Supp Figs 6-7 [↗](#)). This cluster very likely consists of inhibitory interneurons that have been reported to reside primarily in the thalamus in *Mus*³⁷ [↗](#) but may have been inadvertently captured in our dataset due to differences in dissection or differences in spatial distribution of these interneurons between species. Second, some cell types may not be transcriptionally well-defined and homology would not be assignable based on gene expression data alone. For example, cell clusters that lie in the middle of our UMAP (e.g. clusters 0, 13, 25, 8, etc.) likely do not have strong transcriptional signatures captured by our snRNA-seq and were also less likely to be mapped to a homologous *Mus* cell type. Still, we were able to assign homology for the majority of our nuclei enabling us to better interpret cellular and molecular changes in the context of their biological functions.

Differential abundance of cell clusters across species and sex

With our *Peromyscus* POA cell atlas, we first asked if there were differences in cell abundance across species or sex (Fig 2A [↗](#)). To do so, we treated cell abundance as count data and used edgeR³⁸ [↗](#) to fit a generalized linear model that included replicate, sex, and species as covariates (Fig 2A [↗](#), Methods). Importantly, edgeR controls for differential numbers of neurons sequenced across samples (Methods). We used edgeR to test for significant coefficients (i.e. differential abundance across species or sex) and accounted for multiple hypothesis testing with a false discovery rate (FDR) correction³⁹ [↗](#). Because we would later validate our results experimentally, we used a lenient FDR cutoff of 0.3 to initially identify candidate clusters showing differential abundance. We performed this analysis on all cell clusters, regardless of whether they were mapped to a homologous *Mus* cell type or not.




Fig 4.

Differential gene expression across species.

A) Schematic of procedure used to pseudobulk gene expression counts and test for differential gene expression across species. **B)** Barplot of the number of DE genes for each cell cluster (bars) colored by *P. maniculatus* or *P. polionotus* bias. **C)** Scatter plot of cell cluster abundance (x-axis) and number of differentially expressed genes (y-axis) identified in each cell cluster (dots). Line of best fit (gray line) and 95% confidence interval (gray shading) are shown. **D)** Left: Volcano plot of $\log_2(\text{fold change})$ (x-axis) and $\log_{10}(\text{FDR})$ (y-axis) of DE genes across species. Negative values of $\log_2(\text{fold change})$ indicate *P. maniculatus* bias and positive values indicate *P. polionotus* bias. For genes DE in more than one cell type, the cell type with the highest expression level is shown. Genes are colored by gene categories listed on the right. Select outlier genes are labeled. Right: Barplot of enrichment scores (Methods) for each gene category. Dotted line at 1.0 indicates no enrichment or depletion. ***: FDR<0.001; **: FDR<0.01; *: FDR<0.05; ns: not significant.




Fig 5.

Differential gene expression across sex.

A) Barplots of the number of sex-biased genes for each cell cluster (bars) in *P. maniculatus* (top) and *P. polionotus* (bottom), colored by female- or male-bias. **B)** Barplot of the number of female- and male-biased genes aggregated across all cell clusters in *P. maniculatus* (orange), *P. polionotus* (blue), or shared across both species (gray). **C)** Scatter plots of cell cluster abundance (x-axis) and number of DE genes (y-axis) identified in each cell cluster (dots) for *P. maniculatus* (left) and *P. polionotus* (right). Line of best fit (gray line) and 95% confidence interval (gray shading) are shown in each scatter plot.




Figure 6.

Enrichment of neuropeptides in sex-biased genes.

A) Left: Volcano plot of $\log_2(\text{fold change})$ (x-axis) and $\log_{10}(\text{FDR})$ (y-axis) of sex-biased genes in *P. maniculatus*. Negative values of $\log_2(\text{fold change})$ indicate female bias and positive values indicate male bias. For genes sex-biased in more than one cell type, the cell type with the highest expression level is shown. Genes are colored by gene categories listed on the right. Select outlier genes are labeled. Right: Barplot of enrichment scores for each gene category. Dotted line at 1.0 indicates no enrichment or depletion. ***: FDR < 0.001; *: FDR < 0.05. **B)** Same as **A** but for *P. polionotus*. **D)** Venn diagrams of female- (top) and male-biased (bottom) neuropeptides, categorized by whether they are *P. maniculatus* specific, *P. polionotus* specific, or shared. Some neuropeptides (e.g. *Nts*, *Tac1*, *Oxt*, *Gnrh1*) are female-based in some cell types and male-biased in others. Neuropeptides that share sex-bias in the same cell type across species are starred and listed in more detail in [Table 1](#).

Across our 53 neuronal clusters (Supp Table 4), we found four clusters that were called differentially abundant across species and two that were called differentially abundant across sexes (**Fig 2B**, Supp Table 5). Arginine vasopressin (AVP) and oxytocin (OXT) neurons (clusters 34 and 40, respectively) were both ~1.6x more abundant (AVP FDR = 0.19; OXT FDR = 0.09) in the promiscuous *P. maniculatus* than the *P. polionotus* (**Fig 2C**). The AVP neuron number difference corroborates a previous finding that *P. maniculatus* had significantly more AVP neurons than *P. polionotus* (~1.7x) in the medial paraventricular nucleus⁴⁰. Furthermore, we previously reported based on bulk RNA-seq data that the expression level of *Avp* was nearly 3x higher in *P. maniculatus* hypothalamus compared to *P. polionotus*, and that this expression difference mediates the different amounts of parental nesting behavior performed by each species²⁶. Our data suggests this difference in *Avp* expression and parental nesting behavior may be driven, at least in part, by a neuronal number difference across species.

Two cell clusters were found to be more abundant in the monogamous *P. polionotus* compared to the promiscuous *P. maniculatus*: cluster 43 (i26:Tac1/Prok2) (3.4x higher; FDR = 0.08) and cluster 20 (i16:Gal/Th) (1.6x higher; FDR = 0.02) (**Fig 2C**). We were intrigued by the latter cluster because of its known roles in parental care. In *Mus*, i16:Gal/Th is activated by pup exposure and successful mating in both sexes³¹. This cell type likely represents a subset of the galanin-expressing neurons in the medial preoptic area (MPOA) that were previously reported to elicit pup grooming in *Mus* fathers when optogenetically activated⁴¹. In *Peromyscus*, we also observed robust and specific expression of galanin in this cell cluster (Supp Fig 10). Previously, we have shown that *P. polionotus* of both sexes perform more parental care than their *P. maniculatus* counterparts, and that this behavioral difference is heritable²⁶. Thus, we found a positive association between innate parental care behavior and number of a galanin-expressing neuronal cell cluster across *Peromyscus* species.

Finally, we identified two cell clusters to be more abundant in males than females in both *Peromyscus* species: cluster 9 (i20:Gal/Moxd1) (1.7x higher; FDR = 0.007) and cluster 5 (i18:Gal/Tac2) (1.5x higher; FDR = 0.27) (**Fig 2D**). Both cell types are activated by mating behaviors in male *Mus*³¹. i20:Gal/Moxd1 neurons are found in the sexually dimorphic nucleus of the preoptic area (SDN-POA) as well as the bed nucleus of the stria terminalis (BNST), both of which have been described to be sexually dimorphic and larger in males of many mammals including rats⁴², *Mus*⁴³, and humans^{44,45}. Here, we again found a male-bias of i20:Gal/Moxd1 in both a promiscuous and monogamous species.

Several of our differentially abundant cell types have known, highly-conserved gene markers: AVP neurons (marked by AVP), OXT neurons (marked by OXT), and i20: Gal/Moxd1 (marked by calbindin1 [CALB1])^{43,46}. We took advantage of this to experimentally validate our single-nucleus data. We additionally were interested in investigating interspecies differences in the spatial distribution of differentially abundant cell types. Therefore, we performed immunoreactive staining across the entire hypothalamus (Methods). We then counted cells within hypothalamic subregions and tested for differential abundance in the direction identified from our single-nucleus data.

AVP and OXT neurons are formed from a common developmental lineage^{47,48} and both primarily reside in the PVN and the SON. Additionally, small numbers of AVP and OXT neurons were also found by immunostaining in the anterior hypothalamus area (AHA) and BNST (Supp Fig 11). Our counts of AVP-immunoreactive (ir) neurons found 1.5x more cells in the PVN ($p = 7.0\text{e-}4$, one-sided Mann Whitney test) and 2.1x more cells in SON ($p = 6.0\text{e-}6$) (**Fig 3A**, 3B) in *P. maniculatus* compared to *P. polionotus*. Our counts of OXT-ir neurons found 3.1x more cells in the PVN ($p = 2.8\text{e-}6$) and 4.7x more cells in SON ($p = 6.2\text{e-}5$) (**Fig 3C**, 3D). Therefore both regional populations of AVP and OXT neurons appear to be contributing to their differential abundance across species.

In contrast, when we examined CALB1-ir cells (marking i20:Gal/Moxd1), we found regional differences in differential abundance. As expected, CALB1-ir cells were found in the SDN-POA and BNST. Because the cell density of CALB1-ir cells is so high in the BNST, we calculated the area of immunoreactivity in the BNST rather than counting cells individually. We note that this metric may include fibers and terminals whose signal may originate from projections from regions outside the BNST. The area of immunoreactivity in the BNST was ~2x larger in males in both *P. maniculatus* ($p = 0.004$, one-sided Mann Whitney test) and *P. polionotus* species ($p = 0.03$) (Fig 3E, 3F). However, in the SDN-POA, we found that the number of CALB1-ir cells was 1.8x more abundant in the promiscuous *P. maniculatus* males ($p = 0.03$, one-sided Mann Whitney test), but *not* significantly different across sexes in the monogamous *P. polionotus* ($p = 0.2$) (Fig 3E, 3F). i20:Gal/Moxd1 cells of the SDN-POA and BNST cannot be transcriptionally differentiated³¹, and the differential abundance observed in our scRNA-seq data was likely driven primarily by large numbers of i20:Gal/Moxd1 cells of the BNST masking the small number of i20:Gal/Moxd1 cells of the SDN-POA. However, using immunostaining to better capture spatial information, we actually find that the two subregions of i20:Gal/Moxd1 are evolving differently across species: while sex bias in the BNST has remained conserved, sex bias in the SDN-POA has become more reduced in the monogamous *P. polionotus*.

In sum, we computationally identified six neuronal cell types of the hypothalamus that have changed in abundance across species or sex, and experimentally confirmed three cell types with abundance changes. Our findings align with previous work: we identified a species difference in AVP neurons, which had been reported to be differentially abundant across our focal species⁴⁰, and sex-bias of i20:Gal/Moxd1, which is known to be male-biased across mammalian species. Our findings also identified four neuronal cell types with differential abundance across species, two of which are candidates for underlying species differences in behavior: differential abundance of AVP neurons may be partially responsible for a differential expression level of AVP across species, which we previously implicated in parental nesting behavior²⁶, and an increased abundance of a galanin-expressing cell type known to govern parental care behavior, i16:Gal/Th, was observed in the monogamous species that performs more parental care. Finally, our immunostaining found that regional populations of transcriptionally identical cell types may evolve differently across species, and suggests that regional-specificity of sex bias in i20:Gal/Moxd1 cells may be important to the evolution of sex-specific behaviors.

Differential neuronal gene expression across species and sex

We next asked what gene expression differences there were across species and sex. To do this, we pseudobulked our single-nucleus expression data and summed up gene counts across cells within the same cell cluster and sample (Fig 4A). We again used edgeR to fit a generalized linear model that included replicate, sex, and species as covariates (Methods).

Because edgeR normalizes for library size across samples, cell type abundance is controlled for in our differential expression analysis. Again, we performed this analysis on all cell clusters, regardless of whether they were mapped to a homologous *Mus* cell type or not.

Across species, we found a range of 18 to 3,401 differentially expressed (DE) genes in each cell cluster ($FDR < 0.05$), with no bias towards either species (Fig 4B, Supp Table 6). The number of DE genes in each cell cluster was highly correlated with the abundance of that cell cluster (Fig 4C). This is expected given that larger abundances is analogous to deeper sequencing of a cell population and results in more genes detected and more accurate estimates of gene counts. In total, we found 8,301 unique genes to be DE in one or more cell clusters. When compared to a background of expression-matched non-DE genes (Methods), DE genes were highly enriched in neuropeptide receptors (2.4x enrichment, $FDR = 7e-04$), and modestly enriched in neuropeptides (1.5x, $FDR = 0.005$), neurotransmitter receptors (1.3x, $FDR = 0.02$), and voltage-gated channels (1.3x,

FDR = 0.03) (**Fig 4D** [↗](#)). We did not find a significant enrichment of genes involved in neurotransmitter synthesis or transport, and only a small, though marginally significant, enrichment of transcription factors (1.1x, FDR = 0.04) (**Fig 4D** [↗](#)).

Across sexes, we performed differential expression analysis separately for each species to identify changes in sex-biased expression across species (Supp Table 7, 8). Remarkably, we observed far fewer sex-biased genes in the monogamous species compared to the promiscuous species. We found a range of 0 to 40 DE genes (FDR < 0.05) in *P. maniculatus* cell clusters and 2 to 27 DE genes (FDR < 0.05) across *P. polionotus* cell clusters (**Fig 5A** [↗](#)). In total, we found 204 female- and 194 male-biased genes in *P. maniculatus*; in contrast, only 170 (83%) female- and 70 (36%) male-biased genes were detected in *P. polionotus* (**Fig 5B** [↗](#)). Our data suggest a substantial reduction of sex-biased genes in the monogamous species, driven primarily by a reduction of genes with male-biased expression.

Despite the short evolutionary timescales between our species, we found very few sex-biased genes shared across species (**Fig 5B** [↗](#)). Only eight genes were found to be female-biased in the same cell types in both species, including three known female-specific genes associated with X-inactivation (*Xist*, *Tsix*, and *Jpx*) (Supp Table 9). Eleven genes were found to be conserved in male-bias including *Nrip1* (cluster 5, homologous to i18:Gal/Tac2), a nuclear protein that mediates estrogen signaling⁴⁹ [↗](#), and *Ecel1* (cluster 9, homologous to i20:Gal/Moxd1), a neuronal protease⁵⁰ [↗](#) also known to be male-biased in the *Mus musculus* hypothalamus⁵¹ [↗](#) (Supp Table 9).

In both *P. maniculatus* and *P. polionotus*, two cell types had the most number of sex-biased genes: cluster 5 (i18:Gal/Tac2) and cluster 9 (i20:Gal/Moxd1) (**Fig 5A** [↗](#)). When comparing the number of sex-biased genes detected to cell cluster abundance, clusters 5 and 9 were obvious outliers in the number of sex-biased genes detected (**Fig 5C** [↗](#)). The enrichment of sex-biased genes was apparent in both species, even though the two cell types displayed a large reduction in the number of male-biased genes in the monogamous *P. polionotus*.

Notably, these were the same two cell types we found to be *differentially abundant* across sexes (**Fig 2D** [↗](#)). However, we found the enrichment of sex-biased genes to be true even when we re-performed our analysis on data that was downsampled so that cell abundance was equal across sexes (Supp Fig 12A). Additionally, both clusters exhibited high expression of all four major sex-steroid hormone receptors – estrogen receptor α (*Esr1*), androgen receptor (*Ar*), prolactin receptor (*Prlr*), and progesterone receptor (*Pgr*) (Supp Fig 12B). These receptors may be playing roles regulating the observed sex differences in both cell abundance and gene expression⁸ [↗](#). When controlled for cell abundance, none of these sex-steroid hormone receptors were found to be significantly sex-biased in expression (Supp Table 7,8), and so we hypothesize the observed sex-differences may be mediated through differential levels of the sex hormones themselves. Together, our data highlights two cell types that are enriched for sex-biased gene expression in both a promiscuous and monogamous species, suggesting that these cell types may function in highly conserved, sex-specific pathways.

Finally, we performed a gene enrichment analysis on all sex-biased genes, compared to a background of expression-matched non-sex-biased genes (Methods). Despite the rapid turnover of sex-biased genes across species, we found sex-biased genes to be overwhelmingly enriched in neuropeptides in both species (29.5x enrichment, FDR < 2e-4 in *P. maniculatus*; 36.8x enrichment, FDR < 2e-4 in *P. polionotus*) (**Fig 6A** [↗](#), 6B). Out of 69 annotated neuropeptides in our dataset, 21 were found to be sex-biased in at least one cell cluster in either species (**Fig 6C** [↗](#)). We found neuropeptides were more likely to be female-biased in both species (19 female-biased vs. 9 male-biased), although this did not reach statistical significance (Fisher's exact test $p = 0.06$) (**Fig 6C** [↗](#)). We found examples of both species-specific and conserved sex-bias (**Fig 6C** [↗](#)) including 6 instances of neuropeptides sharing the same sex-bias in the same cell type across species (**Table**

1²⁶). Therefore, our data suggest that neuropeptides may be playing key roles in sex differences, but that their expression levels appear to be evolutionarily labile, potentially underlying species-specific variation in sex-specific behavior.

Our differential expression analysis identifies striking sex differences in gene expression in the POA. Overall, we found rapid turnover of sex-biased genes across closely-related species, including a significant reduction in the number of male-biased genes in the monogamous species. Conserved across species, however, we found an enrichment of sex-biased genes expressed in two cell types: cluster 9 (i20:Gal/Moxd1) and cluster 5 (i18:Gal/Tac2). These same cell types were also found to be differentially abundant across sexes in both species, further underscoring their potential significance in sex-specific biology.

Finally, our data suggest that neuropeptides may be a central class of genes mediating sex differences in the brain.

Discussion

How the evolution of neuronal traits, such as cell type abundance and neuronal gene expression, contributes to variation in innate social behaviors is poorly understood. Here, we explored this question by performing single-nucleus RNA-sequencing in the POA of males and females of a monogamous, biparental species of *Peromyscus* and a species that is promiscuous and exhibits maternal-only care. We focused on the POA as it is a region known to be important for mating and parenting behaviors, and took advantage of an existing *Mus* POA single-cell atlas to assign homology between *Peromyscus* cell clusters and functionally characterized cell types.

Differential abundance of parenting-related cell types

Of the four cell types differentially abundant across species, three have obvious ties to parenting behavior: AVP (cluster 34), OXT (cluster 40), and i16:Gal/Th (cluster 20) neurons. The differential abundance we observed in AVP neurons both supports our prior finding that variation in *Avp* levels appears to contribute to a parental nesting behavior difference across our focal species²⁶, and adds to our understanding of the evolutionary mechanisms involved.

Bendesky et al.²⁶ previously used Quantitative Trait Locus (QTL) analysis to link parental nest building to two genomic loci, one of which encompasses the *Avp* locus. With bulk RNA-sequencing, Bendesky et al. found *Avp* was 3x more highly expressed in the promiscuous, less parental species (*P. maniculatus*), and further experimentally showed that an increase in intracerebroventricular *Avp* in *Peromyscus* inhibits parental nesting behavior. The genetic linkage between the *Avp* locus and parental nesting was partially attributed to a *cis*-effect that leads to differential expression across species, though this *cis*-effect changes expression by ~1.1x and does not explain the full difference in expression across species.

In our study, we found AVP neuron count was ~2x higher in the promiscuous *P. maniculatus*, suggesting that a difference in neuron number may additionally contribute to *Avp* expression and behavioral differences. Additionally, we observed 1.8x higher expression of *Avp* within AVP neurons of *P. maniculatus* compared to *P. polionotus*, corroborating the *cis*-effect found by Bendesky, et al.²⁶. Together, the combination of expression difference and abundance difference better matches the full 3x difference observed in bulk RNA-sequencing. Our data proposes that evolution may be modulating behavior by changing expression levels through both local, *cis*-regulatory pathways and upstream pathways in neurogenesis and neurodevelopment.

We also found OXT neuron number to be $\sim 3\times$ higher in the promiscuous species. The direction of differential abundance is opposite to what may be expected given oxytocin's known roles in pair bonding and monogamy⁵². The species difference in oxytocin neurons may simply be due to the shared developmental lineage between AVP and OXT neurons, which are formed from the same precursor cells and only differentiate from each other later in embryogenesis^{47,48}. Our data suggests that genetic changes upstream of AVP/OXT differentiation have evolved across our focal species. Still, more work needs to be done to understand if the OXT abundance difference leads to behavioral changes, or if, perhaps, they have been compensated for through other changes such as in expression and/or spatial distribution of the oxytocin receptor.

Finally, we find a subtype of galanin neurons, i16:Gal/Th (cluster 20), to be more abundant in monogamous species. Galanin neurons of the MPOA were first reported to be specifically activated by parental care behavior in *Mus*⁴¹, a finding that has since been replicated in cichlids⁵³ and frogs⁵⁴. These findings suggest galanin has an evolutionarily conserved role in parenting behavior. In a study of biparental poison frogs (*Ranitomeya imitator*), frogs that performed parental behavior in a laboratory assay had significantly more galanin neurons than non-parental frogs, independent of sex⁵⁴. This study and ours find a correlation between galanin neuronal number and increased propensity for parental care behavior, both within and across species. However, further investigation is required to understand whether these observations are causally linked.

Two cell types play important roles in sex-specific biology

Across sexes, we found two cell types, i20:Gal/Moxd1 (cluster 9) and i18:Gal/Tac2 (cluster 5), that were sex-biased in cell abundance and had an enrichment of sex-biased genes across both *Peromyscus* species. i20:Gal/Moxd1 (cluster 9) is known to be more abundant in males across many mammalian species in two hypothalamic regions: BNST and SDN-POA (see Campi et al. 2013⁵⁵ for an overview). However, our immunohistochemistry experiments found the male-bias in abundance to be significantly reduced in the SDN-POA of the monogamous species, though conserved across species in the BNST.

Our findings mirror a previous study comparing sister species of polygamous and monogamous voles, which found the volume of the SDN-POA to be sexually dimorphic in the polygamous species, but not in the monogamous species⁵⁶. Another study in *P. californicus*, an independent lineage of monogamous *Peromyscus*, found male-bias in the volume of SDN-POA, but noted that the volume difference was half of what had been reported in other, promiscuous rodents⁵⁵. In contrast, the male-bias found in the volume of the BNST was consistent with fold-differences reported in other rodents⁵⁵. Together, these findings suggest that 1) sex differences in the neuronal number of the BNST and SDN-POA can be independently evolving, even within transcriptionally similar cell types, and 2) sex differences in the neuronal number of the SDN-POA, but not BNST, may be co-evolving with monogamy.

Much less is known about the second sexually dimorphic cell type, i18:Gal/Tac2 (cluster 5). In *Mus*, it is activated by mating behaviors³¹ and also reported to be enriched in estrogen receptor (*Esr1*) expression and sex-biased genes⁵⁷. Thus, this cell type appears to play conserved sex-specific roles, possibly in mating and copulatory behaviors.

Sexual dimorphism in gene expression is reduced in monogamous species

Finally, we found significantly fewer sex-biased genes in the monogamous species compared to the promiscuous species. A prevailing hypothesis, first proposed by Darwin⁵⁸, suggests that males of promiscuous species have larger variance in reproductive success compared to those of monogamous species, leading to greater intrasexual competition and more extreme sexual dimorphisms. Correlations between monogamy and reduced sexual dimorphism have been

repeatedly observed in morphological traits such as body size and animal ornamentation (e.g. coloration or plumage) across diverse taxa including primates, birds, fish, and insects^{59–61}. However, much less is known about whether the reduction of sexual dimorphism extends to molecular phenotypes such as gene expression. Work by us and others previously failed to find consistent genome-wide patterns in bulk RNA-sequencing data^{62,63}. The striking reduction we see in our snRNA-seq data highlights the importance of quantifying gene expression at the level of specific cell types to better understand evolution of gene expression and its relationship with organismal-level phenotypes.

Across our study species, there is also a reduction of sexual dimorphism in behavior, with the emergence of biparental care in the monogamous species (in contrast to maternal-only care exhibited by the promiscuous species). Whether the reduced sexual dimorphism in neuronal gene expression may be actually mediating the reduced sexual dimorphism in social behavior is an interesting avenue for future research.

Conclusion

This study begins to characterize molecular differences in a region of the brain responsible for mating and parenting behaviors across two species with divergent mating systems. Overall, we find both cell abundance and gene expression changes across sex and species, both of which contribute to differential levels of neuronally-important proteins. Though not explored here, changes in cell abundance may additionally accompany neuronal circuitry changes that affect behavior. Finally, we find intriguing differences in neuropeptidergic cell types as well as cell types expressing high levels of sex steroid hormone receptors, implicating two key classes of signaling molecules – neuropeptides and sex hormones – as important evolutionary dials for modulating species and sex-specific behavior.

We note that while our two focal species most obviously differ in mating and parenting behaviors, they also differ in other behaviors including thermoregulatory behavior and burrowing behavior²². Additionally, the POA is known to regulate behaviors other than mating and parenting behavior, including sleep and thermoregulation. Therefore, further experimental characterization is required to more precisely understand the behavioral consequences of the species differences we observed in the POA, and to reveal the precise nature of how genes can encode for the substrates needed for generating variation in innate behavior.

Methods

Animal husbandry

P. maniculatus bairdii and *P. polionotus subgriseus* animals were originally acquired from the Peromyscus Genetic Stock Center (Columbia, SC, USA). We weaned animals at 23 days of age into single-sex, same-species groups. We maintained animals on a 16 h:8 h light:dark cycle at 22°C, housed them in standard mouse cages with corncob bedding and provided them with food and water *ad libitum*. Animal husbandry and experimental procedures were approved by the Harvard University Faculty of Arts and Sciences Institutional Animal Care and Use Committee (protocol 27-15-3).

POA dissection

Virgin animals between ages 60 and 103 days were euthanized by CO₂ and brains were rapidly dissected, flash-frozen in Tissue-Tek® Optimal Cutting Temperature Compound, and stored at -70°C. The POA was dissected on a cryostat (Leica CM3050 S) at -20°C. Brains were sectioned coronally until reaching the widening of the anterior commissure. The dorsal and lateral regions

of the brain were removed so that only a ~2.5 mm x 2.5 mm region that contained the POA remained (**Fig 1B**). Five 150 µm thick sections were then made to capture the POA. Sections were placed in a 2 mL tube which was kept at -70°C until the day of nuclei extraction.

Nuclei extraction and 10x library construction

Nuclei were isolated as described in [64]. Briefly, dissected frozen tissue was placed in a single well of a 6 well plate, with 2 ml of extraction buffer. Mechanical dissociation was performed by trituration using a P1000 pipette, pipetting 1 ml of solution slowly up and down, 20 times.

Trituration was repeated 5 times, with 2 minutes of rest between each time. The entire volume of the well was then passed twice through a 26 gauge needle into the same well. The tissue solution was transferred into a 50 ml Falcon tube filled with 30 ml wash buffer and spun in a swinging-bucket centrifuge for 10 min at 600g and 4°C. Following spinning, the majority of supernatant was discarded, leaving 500 µl of tissue solution. DAPI was added (1:1,000) and debris was removed from nuclei by FACS before nuclei were loaded into a 10X Genomics 3' V3 Chip. A detailed protocol can be found at [dx.doi.org/10.17504/protocols.io.bi62khge](https://doi.org/10.17504/protocols.io.bi62khge).

Single-nucleus RNA-sequencing

Our snRNA-seq data consists of two experiments: a pilot experiment that aimed to obtain a single replicate of 10,000 nuclei which was sequenced on a NovaSeq S1 lane (rep01), and a full experiment that aimed to obtain 5 replicates of 20,000 nuclei which were sequenced on an NovaSeq S4 flowcell (rep02 - rep06). Each replicate contained pooled nuclei from a male and female from each of the *P. maniculatus* and *P. polionotus* species (**Fig 1C**). Our final dataset captured 105,647 nuclei across 6 replicates sequenced to a depth of 1,623 - 2,320 median UMI counts / cell (Supp **Table 1**). The 10x library construction and sequencing were performed by the Harvard Bauer Core Facility.

Bulk RNA-sequencing and genotyping

For our pilot study (rep01), we dissected out a portion of the hypothalamus posterior to the POA from each animal for bulk RNA-sequencing. Samples were collected in Trizol (Invitrogen), and total RNA was extracted using the RNeasy mini kit (QIAGEN). Poly(A) enrichment, directional mRNA library preparation, and paired-end sequencing of 150 base pair reads on a NovaSeq 6000 were performed by Novogene. Samples were sequenced to a depth of ~35 million read pairs. Reads were mapped to their respective genomes, *Pman_2.1.3* (GCA_003704035.1) or *Ppol_1.3.3* (GCA_003704135.2), using STAR with default parameters⁶⁵. Variants were then called using the joint genotyping procedure recommended by the Genome Analysis Toolkit (GATK)⁶⁶.

Whole genome sequencing (WGS) and genotyping

For our full experiment (rep02 - rep06), we performed WGS on animals for genotyping. We extracted genomic DNA from tail tissue using proteinase K digestion followed by the Maxwell RSC (Promega) DNA extraction. We prepared DNA libraries using the Nextera DNA Flex kit with Illumina adapters. All samples were pooled into a single library and sequenced on two NovaSeq S4 lanes with paired-end sequencing of 100 base pair reads. Reads were mapped to their respective genomes using bwa-mem with default parameters⁶⁷. The median read depth for these samples was 267,987,590 mapped reads per sample, which corresponds to ~20X sequencing coverage across the genome. Variants were called using the GATK procedure described above.

Sample demultiplexing

To demultiplex pooled scRNA-seq runs, we first used 10x Genomics CellRanger v5.0.0 with default parameters to map reads to a combined *P. maniculatus* and *P. polionotus* transcriptome. Our in-house transcriptome annotations were generated using Comparative Annotational Toolkit⁶⁸ and

GENCODE v15 annotations of *Mus musculus* (GRCm38 mm10) as reference. We used the standard Cell Ranger pipeline to filter out droplet barcodes associated with background noise⁶⁹. For each droplet that passed the background filter, we preliminarily assigned each droplet to the species for which the majority of reads mapped to (which we refer to as the “primary” species). Droplets clearly fell into two clusters: one cluster that represents singlets in which a small percentage of reads incorrectly maps to the secondary species and a second, sparser cluster of droplets that represent multiplets in which genetic material from two species were truly captured (Supp Fig 1). To delineate between the two clusters, we examined the number of reads mapping to the primary genome and binned droplets in increments of 500 reads. Within each bin, we then plotted the distribution of reads mapping to the *secondary* genome. We fit this distribution with a mixture model of two gaussians using normalmixEM⁷⁰ and identified the point at which the two distributions cross (Supp Fig 1). We then found the line of best fit through all intersection points across the bins and used this line as the threshold between singlets and multiplets (Supp Fig 1). Within each species, we next used demuxlet⁷¹ to infer sample identity based on the known sample genotype data (Supp Fig 2). Demuxlet additionally identifies multiplets when mixtures of sample genotypes are detected.

For each 10x run, we expected our detectable multiplet rate to be ~9%⁷² and our assigned detectable rate ranged from 8-13% across our replicates (Supp Table 2A). We found that our assigned singlets were evenly distributed across the four samples (Supp Fig 3A, Supp Table 2B) and the distribution of the number of genes detected per droplet was similar across samples and replicates (Supp Fig 3B). Finally, when we checked the expression level of the female-specific gene, *Xist*, we found that female-assigned cells had significantly higher *Xist* read counts than male-assigned cells and the median read counts of *Xist* across male-assigned cells was 0 (Supp Fig 3C). After sample demultiplexing, we used Cell Ranger v5.0.0 to map reads from each nucleus to the transcriptome of its assigned species and used these read counts for all subsequent analyses.

Cell type clustering

All normalization, data integration, and clustering of scRNA-seq data were performed with Seurat v4.1.3³². Within species, we normalized our scRNA-seq data using sctransform^{73,74} and then integrated data across species using Seurat’s integration workflow with default parameters. Briefly, Seurat learns a conserved gene correlation structure between two data sets using canonical correlation analysis and identifies pairwise correspondences between single nuclei across datasets in order to transform data sets into a shared space³⁶. We created UMAPs using the runUMAP function, and clustered the integrated data by using the FindNeighbors function.

Homology mapping of inhibitory and excitatory cell types

We defined each cell cluster and all nuclei within that cluster as inhibitory or excitatory based on higher expression levels of *Gad1* and *Gad2* (inhibitory) or *Slc17a6* (excitatory) (Supp Fig 7). We then mapped inhibitory and excitatory cell type labels from the *Mus* POA atlas to inhibitory and excitatory nuclei, respectively, using Seurat’s label transfer workflow with default parameters³⁶. Briefly, Seurat first identifies pairwise correspondences between single nuclei across datasets (“anchors”) and then “transfers” cell type labels by weighting the cell type identities of each anchor with their distance to the nuclei at hand. For each nuclei, this procedure calculates the probability that it belongs to each cell type label and assigns the cell type label which received the highest probability.

To find cluster-to-cell-type mappings, we calculated the proportion of each cluster that was assigned to each *Mus* cell type label (Supp Fig 8). We removed *Mus* cell type labels that lacked any notable marker genes and that indiscriminately mapped onto many of our clusters (e1:Glut, i1:Gaba, i7:Gaba, i9:Gaba, i11:Gaba, i13:Gaba, i12:Gaba, i15:Gaba, i39:Gaba) (Supp Fig 8).

We then retained only cell clusters and *Mus* cell type labels if 15% of a cluster or more were predicted to belong to the same *Mus* cell type label. The cutoff value was chosen empirically after examining clusters where homology to the *Mus* atlas that could be established by marker genes. The proportion of these clusters that mapped to *Mus* labels ranged from 17.3% (cluster 15 - i14:Avp/Cck) to 92.6% (cluster 38 - i4:Gaba/Myk), and therefore a 15% cutoff was chosen.

This resulted in 17 excitatory clusters mapping onto 11 *Mus* excitatory cell type labels and 17 inhibitory clusters mapping onto 18 *Mus* inhibitory cell type labels (Figure 1E, 1F).

Differential abundance of cell clusters

Differential abundance analysis was conducted with edgeR³⁸. After clustering our neuronal snRNA-seq data, we created a count matrix of the number of nuclei within each cluster that belonged to each sample. Cluster sizes were normalized using TMM normalization, which relies on the assumption that there are minimal differences in cluster sizes across samples⁷⁵. We used the estimateDisp function with default parameters to estimate the dispersions across cell abundance counts and calcNormFactors function with default parameters to control for the differential numbers of total neurons profiled per sample. We then fit a generalized linear model with glmQLFit function and a design that included replicate, species, and sex as covariates.

Finally, we used glmQLFTest to test for significant coefficients and corrected *p*-values with an FDR correction³⁹.

Immunohistochemistry and cell counting

We transcardially perfused mice with ice-cold 1x phosphate-buffered saline (PBS) with heparin (100 U/mL) and then with 4% paraformaldehyde in PBS. Brains were dissected out, postfixed for 24 hours at 4°C, cryopreserved in 30% sucrose, and stored at -70°C until subsequent use. To stain for protein, we sectioned brains at 30 µm. We aimed to capture the entire hypothalamus and collected sections beginning at the appearance of the corpus callosum until the appearance of the thalamus, keeping every second section (48 sections spanning ~3 mm). We blocked free floating sections for 1 hour in 10% normal goat serum (NGS) and 0.5% Triton- X in PBS. We incubated sections overnight with either rabbit anti-Avp (1:4000, ImmunoStar 20069), mouse anti-Oxt (1:2000, EMD Millipore MAB5296), or mouse anti-Calb1 antibody (1:1000, Millipore Sigma C9848) and 10% NGS in PBS. We used either donkey anti-rabbit Alexa 647 antibody (1:1000, Thermo Fisher Scientific A31573) or donkey anti-mouse Alexa 546 (1:1000, Thermo Fisher Scientific A11003) for secondary detection and mounted tissues with DAPI Fluoromount-G (SouthernBiotech, 0100-20). Slides were imaged on an AxioScan.Z1 slide scanner (Zeiss).

Following imaging, we exported images to .tif format and counted cells using a custom Fiji/ImageJ macro. Briefly, we manually outlined a region of interest (ROI) (e.g. PVN, SON, etc.). Within the ROI, we binarized the image using the “Moments” method⁷⁶ for calculating the optimal threshold. We then used the watershed segmentation algorithm to separate touching particles. Finally, we used the Analyze Particles function to count the number of particles inside the ROI.

P. maniculatus brains are ~15% larger than *P. polionotus* brains by weight which could impact species comparisons of neuron numbers. However, because we were confirming species differences in AVP and OXT neurons, which were more abundant in *P. polionotus*, we did not normalize by brain size for those analyses.

Differential expression analysis

To obtain pseudobulked data, gene counts were summed across all cells belonging to the same cell cluster and sample resulting in a matrix of 30,704 genes by 24 samples by 53 cell clusters. Differential expression analysis was then performed with edgeR³⁸, using the same procedure

described above for differential abundance analysis. For differential expression analysis across species, we fit a generalized linear model with a design that included replicate, sex, and species as covariates. For differential expression analysis across sex, we performed edgeR analysis separately for each species and fit a generalized linear model that included only replicate and sex as covariates. We accounted for multiple hypothesis testing with an FDR correction³⁹ and used an FDR cutoff of 0.05 to define significant differential expression.

Gene enrichment analysis

Gene lists were manually curated by using the Gene Ontology Resource⁷⁷ and are available in Supp Table 10. To test for enrichments of each gene category, we compared foreground genes (i.e. a significantly differentially expressed gene) to background genes that were matched in expression level to control for the fact that genes that are more highly expressed are more accurately quantified and thus more likely to be identified as differentially expressed. We created backgrounds with the following procedure: We binned all genes by their expression level in bins of $0.5 \log_{10}(\text{counts per million})$. For each gene in our foreground, we randomly chose a gene from our entire dataset from the same expression bin. If the same gene was DE in multiple cell types, we matched the highest expression level. To obtain an empirical p -value, we repeated this procedure 5,000 times and calculated the number of times a background list contained equal or higher numbers of foreground genes within a gene category, and we accounted for multiple hypothesis testing of several gene categories using an FDR correction.

We calculated the enrichment score by dividing the number of foreground genes by the mean number of background genes (across 5,000 random samples) within each gene category.

Data availability

All sequencing data has been deposited in GEO under accession number GSE272719. Analysis scripts are available at <https://github.com/jjenny/PeroPOAEvolution>.

Supplementary Figure


Reads from each droplet (dots) are aligned to a combined *P. maniculatus* (*P. man*) and *P. polionotus* (*P. pol*) transcriptome and preliminarily assigned to the species for which a majority of reads map to.

An example of droplets assigned to *P. man* is shown here. The majority of droplets are "singlets" where a small percentage of reads incorrectly map to *P. pol* genome but many droplets are "multiplets" where genetic material from *P. pol* is truly present.

Droplets are binned in increments of 500 mapped *P. man* reads. An example of droplets with 500 - 1,000 mapped *P. man* reads (black dots) are shown here.


The distribution of reads mapped to the "incorrect" genome (right histogram) is then fit with a two-gaussian mixture model. These distributions represent singlets (blue curve) and multiplets (red curve). The point where the singlet and multiplet distributions intersect is calculated. Green dot is the intersection point (y-axis) and center of 500-1,000bp window (x-axis).

The intersections of singlet and multiplet distributions are calculated for each bin (green dots). A line of best fit is calculated and used as the threshold to separate singlets from multiplets.



Supplementary Figure 1.


Schematic illustrating pipeline for singlet and multiplet assignment of pooled snRNA-seq data.



Supplementary Figure 2.

Sample assignment for pooled RNA-seq data.


Scatterplots of the number of reads that map to *P. maniculatus* (*P. man*, x-axis) and *P. polionotus* (*P. pol*, y-axis) for each droplet (dots) across six replicates. Color indicates the final sample assignment made by demuxlet (orange: *P. maniculatus* female; dark orange: *P. maniculatus* male; blue: *P. polionotus* female; dark blue: *P. polionotus* male; gray: multiplet).



Supplementary Figure 3.

Statistics of demultiplexed snRNA-seq data.


A) Barplot of the number of droplets assigned to each sample. **B)** Violin plot of the numbers of genes detected (>0 uniquely mapped read) per droplet across all samples (white dots: median). **C)** Boxplot of the read counts of female-specific gene, *Xist*, across all samples.



Supplementary Figure 4.

Clustering of snRNA-seq data into major cell classes.

A) UMAP visualization of all clustered nuclei. **B)** Dotplot of percentage of cells within each cluster expressing known marker genes (size of dots) and average expression level of marker genes (color of dots). Gene markers (x-axis) are colored by the cell class they represent (see [Supp Fig 4C](#) for key). **C, D)** UMAP visualization and dotplot of same data, grouped by cell class as inferred by expression of marker genes.




Supplementary Figure 5.

UMAP visualizations of all neuronal nuclei, separated by sex and species.

Supplementary Figure 6.

Expression level and spatial distribution of six known cell type-specific genes.


UMAP visualizations of all neuronal nuclei. Color bars indicate the expression level of *Avp* (red), *Oxt* (pink), *Vip* (purple), *Nms* (green), *Grp* (brown), and *Pvalb* (blue).




Supplementary Figure 7.

Classification of inhibitory and excitatory cell clusters.


UMAP visualization of all neuronal nuclei (gray) and expression level of inhibitory markers *Gad1* and *Gad2* (blue) and excitatory marker *Slc17a6* (orange).





Supplementary Figure 8.


Homology mapping of *Mus* cell type labels to *Peromyscus* cell clusters. Heatmap of the proportion of each *Peromyscus* cell cluster (rows) that are assigned to each *Mus* cell type label (columns) for inhibitory (**A**) and excitatory (**B**) cell clusters.



Supplementary Figure 9.

Sequencing metrics across cell clusters.


Violin plots of number of reads mapped (top) and genes found (bottom) in each cell cluster. Bolded cluster numbers are clusters where *Mus* homology could be identified.



Supplementary Figure 10.

***Galanin (Gal)* expression across cell clusters.**


A) Read count of *Gal* in each neuronal nucleus, visualized by UMAP. **B)** Dot plot of normalized expression of *Gal* (color intensity) and the percent of nuclei in which *Gal* is expressed (dot size) in each neuronal cell cluster.



Supplementary Figure 11.

AVP and OXT neuron counts across the entire hypothalamus.

A) Representative image of immunoreactive staining of AVP+ neurons and subregion definitions. AHA: Anterior hypothalamus area; BNST: Bed nucleus of the stria terminalis; PVN: Paraventricular nucleus; SON: Supraoptic nucleus. Scale bar represents 200 μ m. **B)** Boxplots of AVP+ (left) and OXT+ (right) neuron number counts in each subregion of the hypothalamus of *P. maniculatus* and *P. polionotus*.



Supplementary Figure 12.

Cell clusters 5 and 9 are enriched for sex-biased genes and sex steroid receptors.

A) Scatter plots of cell cluster abundance (x-axis) and number of DE genes (y-axis) identified in each cell cluster (dots) for *P. maniculatus* (top) and *P. polionotus* (bottom) using data downsampled so that cell abundance was equal across sexes. Line of best fit (gray line) and 95% confidence interval (gray shading) are shown in each scatter plot. **B)** Dot plot of normalized expression of sex steroid receptors *Esr1*, *Ar*, *Prlr*, and *Pgr* (color intensity) and the percent of nuclei in which they are expressed (dot size) in each neuronal cell cluster.

Acknowledgements

We thank Dhananjay Bambah-Mukuu and Harris Kaplan for advice on POA dissection; Charles Vanderburg and Naeem Nadaf for advice on single-nucleus extraction; and Doug Richardson for advice on cell counting. JC was supported by the Harvard Data Science Initiative and the National Institutes of Health (K99 GM146243-01). SRE and HEH were supported by the Howard Hughes Medical Institute.

Additional information

Author contributions

JC, SRE, and HEH conceived of the study. JC performed all experiments with contribution from CK and PRR. JC analyzed data with contributions from SRE. JC wrote the manuscript with contributions from SRE and HEH. All authors approved of the final manuscript.

Additional files

Supplementary Tables 1-10 [🔗](#)

References

1. Ye D., Walsh J. T., Junker I. P., Ding Y (2024) **Changes in the cellular makeup of motor patterning circuits drive courtship song evolution in *Drosophila*** *Curr. Biol* **34**:2319–2329
2. Auer T. O., et al. (2020) **Olfactory receptor and circuit evolution promote host specialization** *Nature* **579**:402–408
3. Amateau S. K., McCarthy M. M (2004) **Induction of PGE2 by estradiol mediates developmental masculinization of sex behavior** *Nat. Neurosci* **7**:643–650
4. Tsukahara S., Morishita M (2020) **Sexually dimorphic formation of the preoptic area and the bed nucleus of the stria terminalis by neuroestrogens** *Front. Neurosci* **14**
5. Bendesky A., Bargmann C. I (2011) **Genetic contributions to behavioural diversity at the gene- environment interface** *Nat. Rev. Genet* **12**:809–820
6. Young L. J., Winslow J. T., Nilsen R., Insel T. R (1997) **Species differences in V1a receptor gene expression in monogamous and nonmonogamous voles: Behavioral consequences** *Behav. Neurosci* **111**:599–605
7. Young L. J., Nilsen R., Waymire K. G., MacGregor G. R., Insel T. R (1999) **Increased affiliative response to vasopressin in mice expressing the V1a receptor from a monogamous vole** *Nature* **400**:766–768
8. Gegenhuber B., Tollkuhn J (2020) **Signatures of sex: Sex differences in gene expression in the vertebrate brain** *Wiley Interdiscip. Rev. Dev. Biol* **9**:e348
9. Knoedler J. R., et al. (2022) **A functional cellular framework for sex and estrous cycle-dependent gene expression and behavior** *Cell* **185**:654–671
10. Gegenhuber B., Wu M. V., Bronstein R., Tollkuhn J (2022) **Gene regulation by gonadal hormone receptors underlies brain sex differences** *Nature* **606**:153–159
11. Wei Y.-C., et al. (2018) **Medial preoptic area in mice is capable of mediating sexually dimorphic behaviors regardless of gender** *Nat. Commun* **9**
12. Karigo T., et al. (2021) **Distinct hypothalamic control of same- and opposite-sex mounting behaviour in mice** *Nature* **589**:258–263
13. Tosches M. A., et al. (2018) **Evolution of pallium, hippocampus, and cortical cell types revealed by single- cell transcriptomics in reptiles** *Science* **360**:881–888
14. Shafer M. E. R., Sawh A. N., Schier A. F (2022) **Gene family evolution underlies cell-type diversification in the hypothalamus of teleosts** *Nat Ecol Evol* **6**:63–76
15. Johnson Z. V., et al. (2023) **Cellular profiling of a recently-evolved social behavior in cichlid fishes** *Nat. Commun* **14**:4891

16. Kumar S., Stecher G., Suleski M., Hedges S. B 2017) **TimeTree: A resource for timelines, timetrees, and divergence times** *Mol. Biol. Evol* **34**:1812–1819
17. Welch J. D., et al. 2019) **Single-cell multi-omic integration compares and contrasts features of brain cell identity** *Cell* **177**:1873–1887
18. Shekhar K., et al. 2016) **Comprehensive classification of retinal bipolar neurons by single-cell transcriptomics** *Cell* **166**:1308–1323
19. Bakken T. E., et al. 2021) **Comparative cellular analysis of motor cortex in human, marmoset and mouse** *Nature* **598**:111–119
20. Khrameeva E., et al. 2020) **Single-cell-resolution transcriptome map of human, chimpanzee, bonobo, and macaque brains** *Genome Res* **30**:776–789
21. Bakken T. E., et al. 2021) **Single-cell and single-nucleus RNA-seq uncovers shared and distinct axes of variation in dorsal LGN neurons in mice, non-human primates, and humans** *eLife* **10**:e64875
22. Bedford N. L., Hoekstra H. E 2015) **The natural history of model organisms: Peromyscus mice as a model for studying natural variation** *eLife* **4**:e06813
23. Dewey M. J., Dawson W. D 2001) **Deer mice: ‘The Drosophila of North American mammalogy’** *Genesis* **29**:105–109
24. Schenk J. J., Rowe K. C., Steppan S. J 2013) **Ecological opportunity and incumbency in the diversification of repeated continental colonizations by muroid rodents** *Syst. Biol* **62**:837–864
25. Foltz D. W 1981) **Genetic evidence for long-term monogamy in a small rodent, Peromyscus polionotus** *Am. Nat.* **117**:665–675
26. Bendesky A., et al. 2017) **The genetic basis of parental care evolution in monogamous mice** *Nature* **544**:434–439
27. Xie Y., Dorsky R. I. 2017) **Development of the hypothalamus: conservation, modification and innovation** *Development* **144**:1588–1599
28. Alié A., et al. 2018) **Developmental evolution of the forebrain in cavefish, from natural variations in neuropeptides to behavior** *eLife* **7**:e32808
29. Kanda S 2019) **Evolution of the regulatory mechanisms for the hypothalamic-pituitary-gonadal axis in vertebrates-hypothesis from a comparative view** *Gen. Comp. Endocrinol* **284**
30. Mei L., Osakada T., Lin D 2023) **Hypothalamic control of innate social behaviors** *Science* **382**:399–404
31. Moffitt J. R., et al. 2018) **Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic region** *Science* **362**:eaau5324
32. Hao Y., et al. 2021) **Integrated analysis of multimodal single-cell data** *Cell* **184**:3573–3587
33. Steuernagel L., et al. 2022) **HypoMap-a unified single-cell gene expression atlas of the murine hypothalamus** *Nat Metab* **4**:1402–1419

34. Todd W. D., et al. 2020) **Suprachiasmatic VIP neurons are required for normal circadian rhythmicity and comprised of molecularly distinct subpopulations** *Nat. Commun* **11**:4410
35. Wen S., et al. 2020) **Spatiotemporal single-cell analysis of gene expression in the mouse suprachiasmatic nucleus** *Nat. Neurosci* **23**:456–467
36. Stuart T., et al. 2019) **Comprehensive integration of single-cell data** *Cell* **177**:1888–1902
37. Clemente-Perez A., et al. 2017) **Distinct thalamic reticular cell types differentially modulate normal and pathological cortical rhythms** *Cell Rep* **19**:2130–2142
38. Robinson M. D., McCarthy D. J., Smyth G. K 2010) **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data** *Bioinformatics* **26**:139–140
39. Benjamini Y., Hochberg Y 1995) **Controlling the false discovery rate: a practical and powerful approach to multiple testing** *J. R. Stat. Soc. Series B Stat. Methodol* **57**:289–300
40. Cushing B. S 2016) **Estrogen receptor alpha distribution and expression in the social neural network of monogamous and polygynous *Peromyscus*** *PLoS One* **11**:e0150373
41. Wu Z., Autry A. E., Bergan J. F., Watabe-Uchida M., Dulac C. G 2014) **Galanin neurons in the medial preoptic area govern parental behaviour** *Nature* **509**:325–330
42. Gorski R. A., Harlan R. E., Jacobson C. D., Shryne J. E., Southam A. M 1980) **Evidence for the existence of a sexually dimorphic nucleus in the preoptic area of the rat** *J. Comp. Neurol* **193**:529–539
43. Brown A. E., Mani S., Tobet S. A 1999) **The preoptic area/anterior hypothalamus of different strains of mice: sex differences and development** *Brain Res. Dev. Brain Res* **115**:171–182
44. Allen L. S., Gorski R. A 1990) **Sex difference in the bed nucleus of the stria terminalis of the human brain** *J. Comp. Neurol* **302**:697–706
45. Hofman M. A., Swaab D. F 1989) **The sexually dimorphic nucleus of the preoptic area in the human brain: a comparative morphometric study** *J. Anat* **164**:55–72
46. Tsuneoka Y., et al. 2017) **Moxd1 Is a Marker for Sexual Dimorphism in the Medial Preoptic Area, Bed Nucleus of the Stria Terminalis and Medial Amygdala** *Front. Neuroanat* **11**
47. Nakai S., et al. 1995) **The POU domain transcription factor Brn-2 is required for the determination of specific neuronal lineages in the hypothalamus of the mouse** *Genes Dev* **9**:3109–3121
48. Schonemann M. D., et al. 1995) **Development and survival of the endocrine hypothalamus and posterior pituitary gland requires the neuronal POU domain factor Brn-2** *Genes Dev* **9**:3122–3135
49. Nautiyal J., et al. 2013) **The transcriptional co-factor RIP140 regulates mammary gland development by promoting the generation of key mitogenic signals** *Development* **140**:1079–1089
50. Kaneko A., Kiryu-Seo S., Matsumoto S., Kiyama H 2017) **Damage-induced neuronal endopeptidase (DINE) enhances axonal regeneration potential of retinal ganglion cells after optic nerve injury** *Cell Death Dis* **8**:e2847

51. Xu X., et al. 2012) **Modular genetic control of sexually dimorphic behaviors** *Cell* **148**:596–607
52. Froemke R. C., Young L. J. 2021) **Oxytocin, Neural Plasticity, and Social Behavior** *Annu. Rev. Neurosci.* **44**:359–381
53. Butler J. M., Herath E. M., Rimal A., Whitlow S. M., Maruska K. P 2020) **Galanin neuron activation in feeding, parental care, and infanticide in a mouthbrooding African cichlid fish** *Horm. Behav* **126**
54. Fischer E. K., et al. 2019) **The neural basis of tadpole transport in poison frogs** *Proc. Biol. Sci* **286**
55. Campi K. L., Jameson C. E., Trainor B. C 2013) **Sexual dimorphism in the brain of the monogamous California Mouse (*Peromyscus californicus*)** *Brain Behav. Evol* **81**:236–249
56. Shapiro L. E., Leonard C. M., Sessions C. E., Dewsbury D. A., Insel T. R. 1991) **Comparative neuroanatomy of the sexually dimorphic hypothalamus in monogamous and polygamous voles** *Brain Res.* **541**:232–240
57. Kaplan H. S., et al. 2024) **Sensory input, sex, and function shape hypothalamic cell type development** *bioRxiv* <https://doi.org/10.1101/2024.01.23.576835>
58. Darwin C 1871) **The descent of man, and selection in relation to sex.** London: John Murray
59. Andersson M 1994) **Sexual Selection** Princeton: Princeton Univ. Press
60. Kvarnemo C 2018) **Why do some animals mate with one partner rather than many? A review of causes and consequences of monogamy** *Biol. Rev. Camb. Philos. Soc* **93**:1795–1812
61. Kraaijeveld K., Kraaijeveld-Smit F. J. L., Komdeur J. 2007) **The evolution of mutual ornamentation** *Anim. Behav.* **74**:657–677
62. Mishra P., Rundle H. D., Agrawal A. F 2024) **The evolution of sexual dimorphism in gene expression in response to a manipulation of mate competition** *Evolution* **78**:746–757
63. Kautt A. F., et al. 2024) **Evolution of gene expression across brain regions in behaviourally divergent deer mice** *Mol. Ecol* :e17270
64. Kamath T., et al. 2022) **Single-cell genomic profiling of human dopamine neurons identifies a population that selectively degenerates in Parkinson’s disease** *Nat. Neurosci* **25**:588–595
65. Dobin A., et al. 2013) **STAR: ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21
66. Van der Auwera G. A., et al. 2013) **From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline** *Curr. Protoc. Bioinformatics* **43**:11–11
67. Li H 2013) **Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM** *arXiv*
68. Fiddes I. T., et al. 2018) **Comparative Annotation Toolkit (CAT)—simultaneous clade and personal genome annotation** *Genome Res* **28**:1029–1038
69. Zheng G. X. Y., et al. 2017) **Massively parallel digital transcriptional profiling of single cells** *Nat. Commun* **8**:14049

- 70. Benaglia T., Chauveau D., Hunter D. R., Young D 2010) **S. mixtools: An R Package for Analyzing Mixture Models** *J. Stat. Softw* **32**:1–29
- 71. Kang H. M., et al. 2018) **Multiplexed droplet single-cell RNA-sequencing using natural genetic variation** *Nat. Biotechnol.* **36**:89–94
- 72. 10x Genomics 2022) **Chromium single cell 3' reagent kits User Guide (v3.1 Chemistry)** 10x Genomics
- 73. Hafemeister C., Satija R 2019) **Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression** *Genome Biol* **20**
- 74. Choudhary S., Satija R. 2022) **Comparison and evaluation of statistical error models for scRNA-seq** *Genome Biol.* **23**:27
- 75. Robinson M. D., Oshlack A 2010) **A scaling normalization method for differential expression analysis of RNA-seq data** *Genome Biol* **11**
- 76. Tsai W.-H 1985) **Moment-preserving thresholding: A new approach.** *Computer Vision Graphics, and Image Processing* **29**:377–393
- 77. Gene Ontology Consortium, et al. 2013) **Gene Ontology annotations and resources** *Nucleic Acids Res.* **41**:D530–5

Author information

Jenny Chen

Department of Molecular & Cellular Biology, Harvard University, Cambridge, United States,
Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States

ORCID iD: [0000-0002-6664-2597](https://orcid.org/0000-0002-6664-2597)

For correspondence: jennifer_chen@fas.harvard.edu

Phoebe R Richardson

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States

ORCID iD: [0009-0008-5650-1556](https://orcid.org/0009-0008-5650-1556)

Christopher Kirby

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States

ORCID iD: [0000-0001-7292-9044](https://orcid.org/0000-0001-7292-9044)

Sean R Eddy

Department of Molecular & Cellular Biology, Harvard University, Cambridge, United States,
Howard Hughes Medical Institute, Harvard University, Cambridge, United States

ORCID iD: [0000-0001-6676-4706](https://orcid.org/0000-0001-6676-4706)

Hopi E Hoekstra

Department of Molecular & Cellular Biology, Harvard University, Cambridge, United States,
Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States

ORCID iD: [0000-0003-1431-1769](https://orcid.org/0000-0003-1431-1769)

For correspondence: hoekstra@oeb.harvard.edu

Editors

Reviewing Editor

Kate Wassum

University of California, Los Angeles, Los Angeles, United States of America

Senior Editor

Kate Wassum

University of California, Los Angeles, Los Angeles, United States of America

Reviewer #1 (Public review):**(1) Summary of the Paper:**

This paper by Chen et al. examines the cellular composition and gene expression of the hypothalamic medial preoptic area (MPOA) in two closely related deer mouse species (*P. maniculatus* and *P. polionotus*) that exhibit distinct social behaviors. Through single-nucleus RNA sequencing (snRNA-seq), Chen et al., identify sex- and species-specific neuronal cell types that likely contribute to differences in mating and parental care. By comparing monogamous and promiscuous species, the study provides insights into how neuronal diversity and gene expression changes in the MPOA might underlie the evolution of social behaviors.

(2) Strengths of the Paper:

The paper excels in several areas. First, the data presentation is clear and well-organized, making the complex findings easy to follow. The writing is straightforward and highly accessible, which enhances the overall readability. The experimental design is innovative, particularly in how they combined samples from different species into the same dataset and then used post-hoc identification to distinguish cell types by species. This dramatically controls for potential batch effects in my opinion. Additionally, the authors contextualize their findings within the framework of previously published studies on *Mus musculus*, providing a strong comparative analysis that enhances the significance of their work.

(3) Weaknesses of the Paper:

The major limitation of the study is the absence of causal experiments linking the observed changes in MPOA cell types to species-specific social behaviors. While the study provides valuable correlational data, it lacks functional experiments that would demonstrate a direct relationship between the neuronal differences and behavior. For instance, manipulating these cell types or gene expressions *in vivo* and observing their effects on behavior would have strengthened the conclusions, although I certainly appreciate the difficulty in this, especially in non-musculus mice. Without such experiments, the study remains speculative about how these neuronal differences contribute to the evolution of social behaviors.

<https://doi.org/10.7554/eLife.103109.2.sa3>

Reviewer #2 (Public review):

Summary:

The authors report several interesting species and sex differences in cell type expression that may relate to species differences in behavior. The differential cell type abundance findings build on previously observed species/sex differences in behavior and brain anatomy. These data will be a valuable resource for behavioral neuroscientists. These findings are important but the manuscript goes too far in attributing causal influences to differences in behavior. A second important problem is that dissections used for the sequencing data include other neuropeptide-rich areas of the hypothalamus like the PVN. Although histology is included, the results into the main manuscript often do not include the mPOA making it hard to know if species/sex differences are consistent across different hypothalamic regions. The manuscript would benefit from more precise language.

Strengths:

The data are novel because cell-type atlases are available for only a few species.

The authors have clearly defined appropriate steps taken to obtain trustworthy estimations of cell type abundance. Furthermore, the criteria for each cell type assignment was described in a way for readers to easily replicate. The rigor in comparing cell abundance provides convincing evidence that these species have differences in MPOA cellular composition.

The authors have a good explanation for why 19 of the 53 neuron clusters were not classified (possible *Mus/Peromyscus* anatomical differences, some cell types don't have well-defined transcriptional profiles)

Validated findings with histology.

<https://doi.org/10.7554/eLife.103109.2.sa2>

Reviewer #3 (Public review):

Summary:

The authors performed snRNA-seq in the pre-optic area (POA), a heterogeneous brain region implicated in multiple innate behaviors, comparing two species of *Peromyscus* mice that possess strikingly different parenting behaviors. *P. polionotus* show high levels of parental care from both sexes of parent, and *P. maniculatus* show lower levels of care, predominantly displayed by dams rather than sires. The overall goal of understanding the genomic basis of behavioral variation is significant and of broad interest and comparative studies in POA in these two species is an excellent approach to tackle this question. The authors correctly point out that existing studies largely compare species that are highly divergent, such as mice and humans, which confounds the association of specific neuronal populations or gene expression patterns with distinct behaviors. They identify neuronal populations with differential abundance between species and sexes, and additionally report sex and species differences in gene expression within each transcriptomic cell type. Their cell type classification is aided by mapping their *Peromyscus* cells onto a previously existing POA single cell dataset generated in lab mice. The detection and validation of previously observed sex differences in the *Gal/Moxd1* cell type, and species differences in *Avp* expression provides additional support that their data are robust. Importantly, the authors demonstrate reduced sexual dimorphism in the POA of *P. polionotus*, compared to *P. maniculatus*, and prior knowledge in rats and mice. This finding suggests a potential neural substrate for the increased parental behavior in *P. polionotus*.

Strengths:

This is a pioneering comparative snRNA-seq study that provides a roadmap for similar approaches in non-traditional model organisms.

The authors have identified populations that may underlie sex- and species- differences in parenting behavior in rodents.

A significant strength of the manuscript is the histological validation of their most robust marker genes.

Weaknesses:

My primary concern is that the dataset is limited: 52,121 neuronal nuclei across 24 samples, which does not provide many cells per cluster to analyze comparatively across sex and species, particularly given the heterogeneity of the large region dissected, which contains adjacent regions such as the PVN and SCN.

There is no explanation for the finding that there is a female-bias in gene expression across all cell types in *P. polionotus*.

<https://doi.org/10.7554/eLife.103109.2.sa1>

Author response:

The following is the authors' response to the original reviews.

We thank the reviewers for their thoughtful comments.

Based on their suggestions we will:

(1) Use more accurate language to describe the hypothalamus regions under investigation in this study. While we aimed to primarily investigate the medial preoptic area (MPOA), our dissections and sequencing data in fact capture several regions of the anterior hypothalamus including the anteroventral periventricular (AVPV), paraventricular (PVN), supraoptic (SON), suprachiasmatic nuclei (SCN), and more. We will revise the language in our manuscript to reflect that our study in fact investigates the cellular evolution of the anterior hypothalamus across behaviorally divergent deer mice.

(2) Revise our language to clarify that while our study provides a rich dataset for generating hypotheses about which cell types may contribute to behavioral differences, it does not provide any evidence of causal relationships. We hope to investigate this further in future work.

(3) Clarify specific methodological choices for which reviewers had questions, especially about the hypothalamic regions for which we did histology to validate cell abundance differences and methodological choices related to mapping our cell clusters to *Mus* cell types.

Our responses to each reviewer's specific comments are below.

Reviewer #1:

The major limitation of the study is the absence of causal experiments linking the observed changes in MPOA cell types to species-specific social behaviors. While the study provides valuable correlational data, it lacks functional experiments that would demonstrate a direct relationship between the neuronal differences and behavior. For instance, manipulating these cell types or gene expressions in vivo and observing their

effects on behavior would have strengthened the conclusions, although I certainly appreciate the difficulty in this, especially in non-musculus mice. Without such experiments, the study remains speculative about how these neuronal differences contribute to the evolution of social behaviors.

Yes, we agree the study lacks functional experiments. We hope that the dataset is of value for generating hypotheses about how hypothalamic neuronal cell types may govern species-specific social behaviors, and for these hypotheses to be functionally tested by us and others in future work.

Reviewer #2:

Some methodology could be further explained, like the decision of a 15% cutoff value for cell type assignment per cluster, or the necessity of a multi-step analysis pipeline for gene enrichment studies.

A 15% cutoff value for cell type assignment was chosen to include all known homology correspondences between our dataset and the *Mus* atlas. For example, i14:Avp/Cck cells from the *Mus* atlas represent Avp cells from the suprachiasmatic nuclei (SCN). Though only 17.3% of cluster 15 maps to i14:Avp/Cck, we know these two clusters correspond based on the expression of *Avp* and additional SCN marker genes in cluster 15 (Supp Fig 6). We will further explain this cutoff in the revised manuscript.

Our gene enrichment study includes a multi-step analysis pipeline because we wanted to control for confounders that may be introduced because of gene expression level. Genes that are more highly expressed are more accurately quantified and thus more likely to be identified as differentially expressed. Therefore, we wanted to test for gene enrichments in our set of DE genes against a background of genes with similar expression levels. We will clarify this motivation in the revised manuscript.

The authors should exercise strong caution in making inferences about these differences being the basis of parental behavior. It is possible, given connections to relevant research, but without direct intervention, direct claims should be avoided. There should be clear distinctions of what to conclude and what to propose as possibilities for future research.

Yes, we agree that we are unable to make direct claims about neuronal differences being the basis of parental behavior. We will revise our language to be clearer about which relationships we are hypothesizing and what we propose as possibilities for future research.

Histology is not performed on all regions included in the sequencing analysis.

We apologize that our language describing the hypothalamic regions included in the sequencing analysis and those included in the histology is unclear. We aimed to dissect the medial preoptic region for the sequencing analysis, but additionally captured parts of the anterior hypothalamus including the paraventricular (PVN), supraoptic (SON), and suprachiasmatic nuclei (SCN), and more. Our histology was performed across the entire hypothalamus and includes all regions included in the sequencing data. We will revise the manuscript to more accurately describe the hypothalamic regions for which we investigated.

Reviewer #3:

My primary concern is that the dataset is limited: 52,121 neuronal nuclei across 24 samples, which does not provide many cells per cluster to analyze comparatively across sex and species, particularly given the heterogeneity of the region dissected. The

Supplementary table reports lower UMIs/genes per cell than is typically seen as well. Perhaps additional information could be obtained from the data by not restricting the analyses to cells that can be assigned to Mus types. A direct comparison of the two Peromyscus species could be valuable as would a more complete Peromyscus POA atlas.

Our dataset reports ~1,500 genes and ~1,000 UMIs per nuclei which is indeed lower than is typically reported in other single nuclei datasets. Some of this discrepancy is due to a lower quality genome and annotated transcriptome available for *Peromyscus* compared to *Mus musculus*, which results in a lower mapping rate than is typically reported in *Mus* studies. However, our dataset was sufficient to identify known peptidergic cell types (Supp Fig 6) and to map homology to *Mus* cell types for 34 (64%) of our 53 clusters. Additionally, although some of our clusters contain small numbers of cells, our differential abundance analysis accounts for the variance in cell numbers observed across samples and should be robust against any increase in variance due to small numbers. In fact, even differential abundance of very small cell clusters such as oxytocin neurons (cell type 40) was validated by histology.

We would like to clarify that all analyses were performed on all cell clusters, regardless of whether or not they could be assigned homology to a *Mus* cell type. All the cell types that we identified as differentially abundant or contained significant sex differences happened to be cell types for which homology to a *Mus* cell type could be defined. This may arise for a relatively uninteresting reason: cell types that have more distinct transcriptional signatures will be more accurately clustered, leading to more accurate identification of homology as well as more accurate measurements of differential abundance / expression. We will revise language to make this more clear in our manuscript.

In Supplement 7, it appears that most neurons can be assigned as excitatory or inhibitory, but then so many of these cells remain in the unassigned "gray blob" seen in panel 1E. Clustering of excitatory and inhibitory neurons separately, as in prior cited work in Mus POA (refs 31 and 57) may boost statistical power to detect sex and species differences in cell types. Perhaps the cells that cannot be assigned to Mus contain too few reads to be useful, in which case they should be filtered out in the QC. The technical challenges of a comparative single-cell approach are considerable, so it benefits the scientific community to provide transparency about them.

We are not certain about why we are unable to cluster and assign homology to many of our cells (i.e. cells in the unassigned “gray blob”). However, we note that even in the *Mus* atlas, many cells did not belong to obvious clusters by UMAP visualization and that several clusters lacked notable marker genes and were designated simply as “Gaba” and “Glut” clusters. Therefore, it is unsurprising that our own dataset also contains cells that lack the transcriptional signatures needed to be clustered and/or mapped to *Mus* cell types. We do know, however, that the median number of reads/nuclei is uniform across cell clusters and does not explain why some clusters could not be assigned to *Mus*. We will add this information to our revised manuscript.

We do not think that a two-stage clustering (i.e. clustering first by excitatory vs. inhibitory neurons) is expected to gain power to resolve cell types in this case. Excitatory vs. inhibitory neurons are clearly separable on our UMAP (Supp Fig 7) so that information is already being used by our clustering procedure. However, we will explore this further in our revised manuscript to see if doing so will boost statistical power.

The Calb1 dimorphism as observed by immunostaining, appears much more extensive in P. maniculatus compared to P. polionotus (Figures 3 E and F). This finding is not reflected in the counts of the i20:Gal/Moxd1 cluster. The use of Calb1 staining as a proxy for the Gal/Moxd1 cluster would be strengthened if the number of POA Calb1+ neurons that are found in each cluster was apparent. There may be additional Calb+ neurons in the cells

that are not annotated to a Mus cluster. This clarification would add support to the overall conclusion that there is reduced sexual dimorphism in P. polionotus.

From the *Mus* MPOA atlas (which includes both single-cell sequencing data and imaging-based spatial information), it is known that the i20:Gal/Moxd1 cluster comprises sexually dimorphic cells that make up both the BNST and the SDN-POA. These sexually dimorphic cells are well-studied and known to be marked by *Calb1*, which we used in immunostaining as a proxy for i20:Gal/Moxd1.

However, we would like to clarify that in our study, the immunostaining of Calb1+ neurons and the sequencing counts of the i20:Gal/Moxd1 cluster are not completely reflective of each other because our sequencing dataset only captured the ventral portion of the BNST. Therefore our i20:Gal/Moxd1 counts contain a combination of some Calb1+ BNST cells and likely all Calb1+ SDN-POA cells and is difficult to interpret on its own. Our histology, however, covers the entire hypothalamus and is more reliable for identifying sex and species differences in each region. We will clarify this in the revised manuscript.

The relationship between the sex steroid receptor expression and the sex bias in gene expression would be improved if the sex bias in sex steroid receptor expression was included in Supplementary Figure 10.

We will include this in the revised manuscript.

There is no explanation for the finding that there is a female bias in gene expression across all cell types in P. polionotus.

We also find this observation interesting but don't have a good explanation for why at this point. We plan to follow this up in future work.

<https://doi.org/10.7554/eLife.103109.2.sa0>