

Variation in albumin glycation rates in birds suggests resistance to relative hyperglycaemia rather than conformity to the pace of life syndrome hypothesis

Reviewed Preprint

v3 • April 8, 2025

Revised by authors

Reviewed Preprint

v2 • March 11, 2025

Reviewed Preprint

v1 • December 3, 2024

Adrián Moreno-Borralló , Sarahi Jaramillo-Ortiz, Christine Schaeffer-Reiss, Benoît Quintard, Benjamin Rey, Pierre Bize, Vincent A Viblanc, Thierry Boulonier, Olivier Chastel, Jorge S Gutiérrez, Jose A Masero, Fabrice Bertile, François Criscuolo

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178, Strasbourg, France • National Proteomics Infrastructure, ProFi, Strasbourg, France • Parc zoologique et botanique de Mulhouse, Mulhouse, France • Laboratoire de Biométrie et Biologie Evolutive, Villeurbanne, France • Swiss Ornithological Institute, Sempach, Switzerland • CEF, Montpellier University, CNRS, EPHE, IRD, Montpellier, France • Center of Biological Studies of Chizé (CEBC), UMR 7372 CNRS - La Rochelle University, Villiers-en-Bois, France • Ecology in the Anthropocene, Associated Unit CSIC-UEX, Faculty of Sciences, University of Extremadura, Badajoz, Spain

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study uses extensive comparative analysis to examine the relationship between plasma glucose levels, albumin glycation levels, and diet and life history, within the framework of the "pace of life syndrome" hypothesis. The evidence that glucose is positively correlated with glycation levels and lifespan is **convincing** and, although there are some limitations related to data collection, they likely make the statistically significant findings more conservative. As the first extensive comparative analysis of glycation rates, life history, and glucose levels in birds, the study has the potential to be of interest to evolutionary ecologists and the aging research community more broadly.

<https://doi.org/10.7554/eLife.103205.3.sa2>

Abstract

The pace of life syndrome hypothesis (POLS) suggests that organisms' life history, physiological and behavioural traits should co-evolve. In this framework, how glycaemia (i.e., blood glucose levels) and its reaction with proteins and other compounds (i.e. glycation) covary with life history traits remain relatively under-investigated, despite the well documented consequences of glucose and glycation on ageing, and therefore potentially on life history evolution. Birds are particularly relevant in this context given that they have the highest blood glucose levels within vertebrates and still higher mass-adjusted longevity when

compared to organisms with similar physiology as mammals. We thus performed a comparative analysis on glucose and albumin glycation rates of 88 bird species from 22 orders, in relation to life history traits (body mass, clutch mass, maximum lifespan and developmental time) and diet. Glucose levels correlated positively with albumin glycation rates in a non-linear fashion, suggesting resistance to glycation in species with higher glucose levels. Plasma glucose levels decreased with increasing body mass but, contrary to what is predicted to the POLS hypothesis, glucose levels increased with maximum lifespan before reaching a plateau. Finally, terrestrial carnivores showed higher albumin glycation compared to omnivores despite not showing higher glucose, which we discuss may be related to additional factors as differential antioxidant levels or dietary composition in terms of fibres or polyunsaturated fatty acids. These results increase our knowledge about the diversity of glycaemia and glycation patterns across birds, pointing towards the existence of glycation resistance mechanisms within comparatively high glycaemic birds.

Introduction

The pace of life (POL) hypothesis [1–3] postulates that organisms behavioural/physiological characteristics and life histories have co-evolved in answer to specific ecological conditions forming pace of life syndromes (POLS) along a fast-slow continuum (but see [4] for the more recent consideration of other axes). According to the classical approach from life history theory, slower organisms would have, e.g. large body mass, late maturation, slow growth rate, small number of large offspring and high longevity, while faster ones representing the opposite trait values (see [5–8]) within this continuum. Pace of life hypothesis added physiology to this continuum, with studies testing its predictions focusing often on metabolic rate (i.e. higher or lower metabolic rate corresponding to faster or slower pace of life, respectively; see e.g. [9–12]). However, key energy substrates related to metabolic performance, such as glucose concentrations in tissues, have been largely overlooked (but see [13–16]). Notably, glucose, a central energy source for many organisms, plays a pivotal role in metabolism and there are some indications that its circulating blood or plasma levels correlate positively with metabolic rate per gram of mass at the interspecific level [13, 17], and with whole body metabolism at the intraspecific level ([18] for the case of non-diabetic humans).

Research focusing on plasma glucose becomes highly relevant as high glycaemia can entail costs that accelerate ageing. Along with other reducing sugars, glucose can react non-enzymatically with free amino groups of proteins, lipids or nucleic acids, a reaction known as glycation [19][20]. This reaction, after several molecular reorganizations, can lead to the formation of a plethora of molecules called Advanced Glycation End-products (AGEs) [21]. AGEs can form aggregates in tissues that are well known to contribute to several age-related diseases, such as diabetes (e.g., reviews by [22–25]). AGEs are also known to promote a proinflammatory state through their action on specific receptors called RAGEs [26–27].

Remarkably, birds show circulating glucose levels much higher than other vertebrate groups, on average almost twice as high as mammals [28]. These relatively high glucose levels might support the higher metabolic rates [29, 11] and body temperatures observed in birds compared to mammals [30–31]. In addition, elevated glycaemia is thought to be an adaptation to flight, providing birds with rapid access to an easily oxidizable fuel during intense bursts of aerobic exercise (see [32–33]), such as during take-off and short-term flapping flights [34–35]. However, the presence of high glycaemia in birds is also paradoxical, given their remarkable longevity compared to their mammalian counterparts — living up to three times longer than mammals of equivalent body mass [36, 11].

Several non-excluding hypotheses have been proposed to explain how birds apparently resist the pernicious effects of a high glycaemic state. One possibility is that birds might resist protein glycation through a lower exposure of lysine residues at the surface of their proteins [37]. Another hypothesis suggests that increased protein turnover may play a role [38–39]. Additionally, birds might benefit from more effective antioxidant defences [40–42] (although within Passeriformes, [16] shows no coevolution between glycaemia and antioxidant defences) or even the presence of “glycation scavengers” (i. e. molecules that bind glucose avoiding it to react with proteins) such as polyamines which circulate at high concentration [35]. Moreover, it is generally considered that RAGEs are not present in avian tissues, suggesting possible adaptive mechanisms that would allow birds to avoid the inflammatory consequences associated with RAGE activation [43–44]. However, a putative candidate for RAGE and a case of AGE-induced inflammation in birds were identified by Wein et al. [45], warranting further investigation on the occurrence of RAGEs in birds.

To date, although protein glycation levels have been measured in birds in several species, these results were mostly descriptive and concerned a small number of species [46–54]. Additionally, many of these studies employ commercial kits that were not specifically designed for avian species, making it challenging to interpret the results [53]. These limitations restrict our ability to draw general conclusions from these studies. Here, we performed a comparative analysis on primary data (for glycation and most of glucose values, see methods and ESM6) from 88 bird species belonging to 22 orders (see ESM6) and assessed whether and how bird glycaemia and glycation rates are linked to ecological and life-history traits (including body mass). While doing this, we also checked whether glycation levels are influenced by circulating plasma glucose concentrations. Finally, as diet can influence glycaemia (see e.g. [15, 55–56] but [14] found no significant effects) and glycation, we included it in our analyses. We hypothesized that carnivorous birds would exhibit higher glycaemia and glycation levels than omnivorous or herbivorous species, as diets with low-carbohydrates, high-proteins and high-fat are associated with comparatively high glycaemia and reduced insulin sensitivity in some vertebrates (e.g.; [57–59]). Accordingly, a recent comparative study in birds showed higher blood glucose levels in carnivorous species [56]. This may be attributed to high levels of constitutive gluconeogenesis, which has been confirmed in certain raptors [60–61]. Conversely, bird species with high sugar intake, such as frugivores or nectarivores, are expected to exhibit high glycaemia, as observed in hummingbirds [49] and Passeriformes with such diet [15] (although the opposite was found in [56]). In line with POLS hypothesis, we also hypothesized that species with a slower POL should exhibit a lower glycaemia and higher glycation resistance (glycation levels for a given glycaemia) compared to species with a faster POL. Hence, we included the following four parameters in our analyses of glycaemia and glycation levels: body mass, maximum lifespan, clutch mass and developmental time. Body mass is one of the main factors underlying life history, with higher body mass associated with “slower” strategies and vice versa [62]. Previous studies have reported a negative relationship between body mass and glycaemia [63, 34, 13–14, 56] (but see [49, 64] with non-significant trends in birds, and [15] only for temperate species within Passeriformes). Therefore, we predicted that bird species that live longer, develop more slowly and invest less per reproductive event (see ESM6 for further justification of the chosen variables) should show lower plasma glucose levels and albumin glycation rates (after controlling for body mass, phylogeny, diet and glucose in the case of glycation; see methods).

Materials and methods

Species and sample collection

A total of 484 individuals from 88 species measured were included for this study (see [Table ESM6.1](#)). A detailed list with the provenance of the samples, including zoos, a laboratory population, and both designated captures and samples provided by collaborators from wild populations, is provided in [ESM3](#), with a textual description in [ESM6](#).

Blood samples were collected from the brachial or tarsal vein, or from the feet in the case of swifts, using heparin-lithium capillaries or Microvette® (Sarstedt). As samples were collected mostly by different collaborators, handling times were not always recorded and could not be adjusted for, potentially rendering the results more conservative (for a more detailed discussion on potential stress effects on glucose, see [ESM6](#)). Samples were centrifuged at 4°C, 3500 g for 10 min and plasma was aliquoted when a large volume was available. Subsequently, they were transported on dry ice to the IPHC (Institut Pluridisciplinaire Hubert Curien) in Strasbourg and stored at -80° C until analysis. Glycaemia was measured in the laboratory on the remaining plasma after taking an aliquot for glycation assessment, using a Contour Plus® glucometer (Ascensia diabetes solutions, Basel, Switzerland) and expressed in mg/dL. These of point-of-care devices have previously been assessed for its usage in birds [[65](#)–[66](#)], with several examples of its usage in recent literature related to the topic presented here (e.g. [[14](#), [67](#)–[69](#)]). We also performed an assay with this particular brand on a subset of 46 samples from this study, coming from nine species distributed across the whole range of glucose values (three species with “low” values, three with “medium” and three with “high”, from both captive and wild populations, with five individuals per species, except one including six), confirming a positive linear correlation ($R^2_{\text{marginal}}=0.66$; $R^2_{\text{conditional}}=0.84$, for a model including the species as random factor) of this device with Randox GLUC-PAP colourimetry kits (P-value<0.001, unpublished data). Due to technical issues, related to the incapability of the device to determine certain glucose values (not because of the glucose concentration, but perhaps the particular composition of the plasma samples from certain species), we could not determine glycaemia values of 95 individuals of those that we sampled and in which glycation levels were assessed (belonging to 40 species coming from different sources). For these species, if not a single individual had a glucose measurement (which was the case in 13 species), we obtained mean plasma glucose values reported for the species from the ZIMS database from Species360 (Species360 Zoological Information Management System (ZIMS) (2023), [zims.Species360.org](#)). This database provides plasma glucose data measured by colorimetric glucose kits on zoo specimens not necessarily corresponding to those in which we measured glycation values. The sample sizes for both glucose and glycation measurements, either from our measured individuals or from the ones from ZIMS, are reported on the [ESM5](#).

Glycation levels

Glycation levels for each individual were determined using liquid chromatography coupled to mass spectrometry (LC-MS), which is considered the gold standard for the assessment of protein glycation levels (see e.g. [[70](#)]) and has previously been used for birds [[52](#)–[53](#), [71](#)]. Given the relatively high time intrinsically taken by the employed methodology for analysing all the samples, linked to constraints in the access to the mass spectrometry devices, the whole set of samples of this study were analysed across several instances within a total timespan of less than 2 years (2021–2023). Only one sample from each individual was measured, given logistic limitations on the total number of samples that could be processed. Briefly, 3 µl of plasma were diluted with 22 µl of distilled water containing 0.1% of formic acid, followed by injection of 5 µl into the system. The glycation values used in the analyses represent the total percentage of glycated albumin, obtained by adding the percentages of singly and doubly glycated albumin. These percentages were calculated by dividing the areas of the peaks corresponding to each glycated molecule form (albumin plus one or two glucose) by the total albumin peak area (sum of the areas of glycated plus non-glycated molecules) observed in the spectrograms obtained from the mass spectrometry outcomes. These spectrograms represent the different intensities of signal for the components

differentiated in the sample by their TOF (Time Of Flight), which depends on the mass to charge ratio (m/z) of the ionized molecules. More detailed information regarding functioning of the method and data processing can be found in [53]. In cases where albumin glycation values dropped below the limit of detection, resulting in missing data, these individuals (10 individuals from 4 species) were excluded from the statistical analyses, as outlined below.

Data on ecology and life-history

Diets of individual species were extracted from the AVONET dataset ([72]), coming from [73], and those adapted from [74]) and life history traits from the Amniote Database [75]. After determining that AVONET diet classifications did not align with our research needs, minor changes were made after consulting the original Wilman et al. [74] database (see ESM6).

For missing data on life history traits after this stage, we extracted values, in order of preference, from the AnAge database [76] or the Birds of the World encyclopaedia [77], by calculating mean values if an interval was given, and then averaging if male and female intervals were provided separately. For European species, maximum lifespan records have always been checked against the most recent Euring database [78], and from *Chionis minor* and *Eudypates chrysolophus* (not available from the previous sources) they were extracted from the Centre d'Etudes Biologiques de Chizé, Centre National de la Recherche Scientifique (CNRS-CEBC) database. Several species from the zoo had maximum lifespan values available on the ZIMs from Species 360 (Species360 Zoological Information Management System (ZIMS) (2023), zims.Species360.org), so these were also compared with the data we had from other sources.

In the case of variables (other than maximum lifespan) for which two different sources provided different records, an average value was calculated. For the maximum lifespan value, available sources were cross-checked and the highest value was always used. In the absence of satisfactory support from another source, values for maximum lifespan indicated as being anecdotal and of poor quality have been excluded from the analyses. A table with the adequate citations for each value is provided as part of the online available data (see Data accessibility section). When the source is AVONET [72], this database is cited, and when it is Amniote Database [75], we cite the sources provided by them, so the references can be checked in [75]. A list with all the additional references not coming from any of these databases or not provided by their authors (and that are not already in the main text) is given as ESM4.

Statistical analyses

All analyses were performed in R v.4.3.2 [79]. Alpha (α) ≤ 0.05 was reported as significant, and $0.05 < \alpha \leq 0.1$ as trends. General Linear Mixed Models with a Bayesian approach (*MCMCglmm* function in R; [80]) were performed. All models were run with 6×10^6 iterations, with a thinning interval of 100 and a burn-in of 1000. The models were simplified by eliminating quadratic terms where they were not significant and selecting models with lower AIC (Akaike Information Criterion) and BIC (Bayesian Information Criterion). Gaussian distribution of the response variables was assumed (after $\log_{10}$ conversion for glucose; see ESM6 for further discussions). The priors were established assuming variance proportions of 0.2 for the G matrix and 0.8 for the R matrix, with $\nu=2$. Less informative priors with equal variance proportions for each partition (residual and random) gave similar results. Lower ν values (0.1, 0.2, 0.002) were also tested, without success (i.e. simulations aborted before the established number of iterations). For the phylogenetically controlled analyses, consensus trees (each with the species included in the model) were obtained by using the *consensus.edges* function (from the *phytools* package from R [81]) from a total of 10,000 trees downloaded from *Birdtree.org* [82] (option “Hackett All species” [83]). For such purpose, a list of species with the names adapted to the synonyms available in such website was used (see ESM5), including the change of *Leucocarbo verrucosus* for *Phalacrocorax campbelli*, as the former was not available and it was the only species from the order Suliformes in our dataset, so that neither the position in the tree nor the branch length

would be affected by this change. The consensus trees were included as *pedigree* in the models. We performed models with either glucose or glycation as dependent variables and with the diet, body mass and life history traits as predicting variables (and glucose in the models with glycation as a response to assess glycation resistance; see introduction; another set of models without glucose were performed to test if there was a covariation of glycation itself, independently of glucose levels, covaried with life-history; see [ESM6](#)). Generalized Variance Inflation Factors (GVIFs) were calculated for all the models with more than one predictor variable to assess the collinearity of them, as it may be expected for life history traits (see results on [ESM1](#)), considering values above 1.6 as slightly concerning, above 2.2 as moderately concerning and above 3.2 as severely concerning (i.e. indicating high collinearity; following [\[84\]](#)). Models testing the effects of age and sex on glucose and glycation levels and the number of exposed lysine residues on glycation were also carried out. Finally, we performed models on glucose and glycation values controlling for the taxonomic orders included in the dataset (see [ESM5](#)). The effects of phylogeny on all models was determined by calculating the ratio of the variance estimated for the “animal” variable, representing the *pedigree* (see [ESM6](#)) by the total variance (all random factors plus *units*). A thorough description of all of the models, including transformations of the variables and other details is given on the [ESM6](#).

Results

Plasma glucose and albumin glycation variability across the birds’ tree

Plasma glucose and albumin glycation values varied considerably across species ([Figure 1](#)). Considerable within species repeatability (see [ESM6](#)) was observed (Glucose: $R = 0.716$, $SE = 0.042$, $CI_{95} = [0.619, 0.785]$, $P\text{-value} < 0.0001$; Glycation: $R = 0.703$, $SE = 0.042$, $CI_{95} = [0.603, 0.768]$, $P\text{-value} < 0.0001$).

We found significant differences between some orders for both glucose and albumin glycation. [ESM1](#) shows the raw outcomes of the models, showing such significant differences to the intercept and the Credible Intervals to perform the pairwise comparisons across all groups. To explore some of the details, we can see for example how Apodiformes showed the highest average glucose values (species average model: Estimated mean=351.8 mg/dL, $CI_{95}[238.9, 514.2]$; model with individuals: Estimated mean=349.8 mg/dL, $CI_{95}[251.8, 476.7]$, $n=39$ individuals, 1 species), followed by Passeriformes (species average model: Estimated mean=349.2 mg/dL, $CI_{95}[279.6, 438.6]$; model with individuals: Estimated mean=337.6 mg/dL, $CI_{95}[274.5, 414.5]$, $n=84$ individuals, 8 species), while the Suliformes (species average model: Estimated mean=172 mg/dL, $CI_{95}[117.6, 252.9]$; model with individuals: Estimated mean=169.5 mg/dL, $CI_{95}[120.9, 239.7]$, $n=5$ individuals, 1 species), Rheiformes (species average model: Estimated mean=173.7 mg/dL, $CI_{95}[118, 254.7]$; model with individuals: Estimated mean=172 mg/dL, $CI_{95}[123.7, 239.8]$, $n=9$ individuals, 1 species) and Phoenicopteriformes (species average model: Estimated mean=176 mg/dL, $CI_{95}[130, 241.1]$; model with individuals: Estimated mean=141.2 mg/dL, $CI_{95}[100.4, 198.2]$, $n=5$ individuals, 1 species) show the lowest average values (for the species average models, the pattern is Suliformes<Rheiformes<Phoenicopteriformes, while for the individual models is Phoenicopteriformes<Suliformes<Rheiformes).

For glycation, Strigiformes (species average model: Estimated mean=25.6 %, $CI_{95}[20, 31]$; model with individuals: Estimated mean=25.3 %, $CI_{95}[19.1, 31.5]$, $n=7$ individuals, 2 species), Apodiformes (species average model: Estimated mean=25.5 %, $CI_{95}[18.6, 32.4]$; model with individuals: Estimated mean=24.8 %, $CI_{95}[17.9, 32]$, $n=40$ individuals, 1 species) and Coraciiformes (species average model: Estimated mean=24.5 %, $CI_{95}[17.9, 31.6]$; model with individuals: Estimated mean=23.9 %, $CI_{95}[15.8, 32]$, $n=2$ individuals, 1 species), all terrestrial carnivores as by our sampled species (see discussion below), had the highest average.




Figure 1.

Average plasma glucose values in mg/dL (in grey) and average albumin glycation rate as a percentage of total albumin (in blue) from all the species used in this study (some of them with glucose values coming from ZIMs database; see methods) with the orders they belong to. Glucose and glycation values are standardized in order to be compared, with the dotted lines representing half the maximum and maximum values for each variable (as indicated by the axes in their corresponding colours), from inside out. Tree from a consensus on 10,000 trees obtained from "Hackett All species" on Birdtree.org, including 88 species from 22 orders (see methods).

On the other hand, Casuariiformes (species average model: Estimated mean=10.8 %, CI₉₅[3.7, 18.1]; model with individuals: Estimated mean=11.4 %, CI₉₅[2.7, 20.6], n=1 individuals, 1 species), Phoenicopteriformes (species average model: Estimated mean=11.2 %, CI₉₅[5.6, 16.7]; model with individuals: Estimated mean=10.5 %, CI₉₅[4.6, 16.3], n=22 individuals, 2 species) and Suliformes (species average model: Estimated mean=13.8 %, CI₉₅[7, 20.7]; model with individuals: Estimated mean=13.2 %, CI₉₅[5.6, 20.5], n=5 individuals, 1 species) had the lowest average levels for the species average models, the pattern is Casuariiformes<Phoenicopteriformes<Suliformes, while for the individual models is Phoenicopteriformes<Casuariiformes<Suliformes). Graphs with raw data on species average and individual glucose and glycation values by order are shown on [ESM2](#) (Figure ESM2.3). Estimates of phylogeny effects on the residuals of the models on glucose and glycation values differ if we consider the models with intraspecific variability or without it, but not so much between the models with and without life history traits (within the previous categories). In the case of species averages, the effect of the tree on residual glucose variation is lower and the estimation is less precise than for residual glycation variation, while for the models considering individual values, it is glycation residuals what shows lower levels of tree-related variance than glucose residuals (see [Figure ESM1](#)).

Bird glycemia appears related to body mass and maximum lifespan

After controlling for intraspecific variation in our analyses, we found that variation in glucose levels were significantly explained by variations in body mass and both the linear and quadratic component of residual maximum lifespan (i.e. mass and phylogeny-adjusted maximum lifespan), but not by variations in diet ([Figure 4.A](#) with predictions from the model without life history traits and [Figure ESM2.2.A](#) with raw individual data from the same dataset), clutch mass and developmental time (see [Table 1](#) for the model including life-history trait variables). Heavier species had lower glucose levels (see [Figure 2.A](#), drawn with the estimates from the model without life history traits, which includes more species). Glucose levels increase with increasing mass-adjusted lifespan until reaching a plateau ([Figure 2.B](#)). Models that did not consider intraspecific variability show no significant effect of any of the aforementioned variables on glucose levels (see [ESM1](#)). The provenance of the samples (wild versus captive) only showed a trend to a higher glucose levels in the samples from captive individuals in the model without life history traits (Estimate = 0.058; CI₉₅[-0.008, 0.125]; P-value=0.083).

Bird albumin glycation is related to glycemia and diet

After controlling for intraspecific variability, we found that diet was affecting variation in albumin glycation rates, with terrestrial carnivorous species having higher glycation levels than omnivorous species ([Table 2](#) for complete model; see [Figure 4.B](#) with predictions of the model without life history traits and [Figure ESM2.2.B](#) with raw individual data from the same dataset). However, for the models for species averages, there was only a trend on this pattern in the one including life history traits, and no effect in the other (see [ESM1](#)). The relation between glycation and glucose levels was positive and significant in all but the model that included life history traits but not intraspecific variation (see [Table 2](#) for the outcome of the model with individual values and life history traits and [ESM1](#) for the rest; see [Figure 3](#) with estimates from the model including individual variation but no life history traits, as it contains more species and the estimates are similar). Given the logarithmic relationship between glycation and glucose (see [ESM6](#)), the slope lower than one (see [Table 2](#)) implies that birds with higher glucose levels have relatively lower albumin glycation rates for their glucose, fact that we would be referring to as higher glycation resistance. The glycation models excluding glucose levels, and therefore testing for covariates of life-history with glycation itself, without considering resistance, rendered similar results, with only the abovementioned dietary effects being significant (see [Figure ESM1](#)).




Figure 2.

Plasma glucose levels (in mg/dL) variation as a function of **A** species mean centred body mass and **B** residual maximum lifespan. Both glucose and body mass are log transformed. Maximum lifespan (in years) is given as the residues of a pGLS model with body mass (in grams), both $\log_{10}$ transformed, so the effects of body mass on longevity are factored out (see [ESM6](#)). Different bird orders, are indicated by symbols, as specified on the legends at the right side of the graphs. **Figure 2.A** uses the values and estimates from the glucose model without life history traits ($n=389$ individuals from 75 species), while **Figure 2.B** uses only the data points employed on the complete model ($n=326$ individuals of 58 species).

	Estimates	Lower 95% CI	Upper 95% CI	Sampling effort	pMCMC
Intercept (Omnivore)	2.387	2.268	2.5	59900	<0.001
Diet: Carnivore terrestrial	0.02	-0.056	0.099	59900	0.592
Diet: Aquatic predator	0.034	-0.047	0.112	59900	0.393
Diet: Herbivore	-0.053	-0.177	0.07	59900	0.393
Diet: Frugivore/granivore	-0.055	-0.239	0.135	59900	0.558
Centred Log ₁₀ Body mass	-0.061	-0.106	-0.015	59900	0.009
Maximum lifespan	0.107	-0.035	0.253	59900	0.142
Maximum lifespan ²	-0.616	-1.166	-0.095	59900	0.026
Clutch mass	-0.095	-0.265	0.069	59900	0.258
Developmental time	0.011	-0.185	0.212	59900	0.916
Provenance: Captive	0.034	-0.039	0.106	59900	0.346

Table 1.

Final glucose model with diet, body mass, life history traits and sample provenance (wild versus captive; see [ESM6](#)) as explanatory variables, including the significant quadratic effect of maximum lifespan. Posterior means, CI₉₅ and pMCMC from a phylogenetic MCMC GLMM model including n=326 individuals of 58 species. Both glucose and body mass are log₁₀ transformed and life history traits are residuals from a pGLS model of log₁₀ body mass and log₁₀ of the trait in question (see [ESM6](#)). Body mass was also centred to better explain the intercept, as 0 body mass would make no biological sense. The intercept corresponds to the Omnivore diet, being used as the reference as it is considered the most diverse and “neutral” group for this purpose. Significant predictors are indicated in bold.




Figure 3.

Individual albumin glycation rates (as a percentage of total albumin) variation as a function of individual plasma glucose values (mg/dL). **A.** Both variables $\log_{10}$ transformed, as in the model, including the line representing the predicted relationship). **B.** Both variables in a linear form, to more explicitly illustrate the phenomenon referred to as higher albumin glycation resistance in birds with higher plasma glucose levels, inferred from the faster increase in glucose than albumin glycation, i.e. the negative curvature of the relationship. Different bird orders are indicated by symbols, as specified on the legends at the right side of the graphs. The values and estimates used are from the glycation model without life history traits ($n=379$ individuals from 75 species).




Figure 4.

Outcomes of the models (estimates with interquartile ranges from the posterior distributions and whiskers representing credible intervals) on individual data on effects of diet on **(A)** plasma glucose levels and **(B)** albumin glycation in birds. Glucose levels are given in mg/dL, while glycation levels are a percentage of total plasma albumin which is found to be glycated. Terrestrial carnivores showed significantly higher glycation levels than omnivores (Estimate=21.62 %, $CI_{95}[18, 25.95]$, $p_{MCMC}=0.049$). Models without life history traits, including more individuals, are represented, but the models with life history traits do not show differences in their qualitative predictions (i.e. higher albumin glycation in terrestrial carnivores than in omnivores; see [ESM1](#)).

	Estimates	Lower 95% CI	Upper 95% CI	Sampling effort	pMCMC
Intercept	1.232	1.112	1.351	59900	<0.001
Diet: Carnivore terrestrial	0.101	0.017	0.187	59900	0.021
Diet: Aquatic predator	0.027	-0.062	0.118	59900	0.549
Diet: Herbivore	-0.019	-0.161	0.119	59936	0.781
Diet: Frugivore/granivore	0.095	-0.098	0.292	59900	0.329
Centred log ₁₀ Body mass	0.004	-0.043	0.05	58765	0.876
Log ₁₀ Glucose	0.137	0.012	0.255	59900	0.027
Maximum lifespan	0.037	-0.122	0.195	59043	0.648
Clutch mass	0.151	-0.03	0.346	59900	0.114
Developmental time	0.04	-0.188	0.266	59303	0.725

Table 2.

Final glycation model with diet, body mass, glucose and life history traits as explanatory variables. Posterior means, CI₉₅ and pMCMC from a phylogenetic MCMC GLMM model including n=316 individuals of 58 species. Glycation, glucose and body mass are log₁₀ transformed and life history traits are residuals from a linear model of log₁₀ body mass and log₁₀ of the trait in question (see **ESM6** [↗](#)). Body mass and glucose were also centred to better explain the intercept. The intercept corresponds to the Omnivore diet, being used as the reference as it is considered the most diverse and “neutral” group for this purpose. Significant predictors are indicated in bold and the credible intervals are considered for making pairwise comparisons between the groups.

Additional analyses looking at the number of exposed lysines in the albumin sequence of a species show no effect of this variable on albumin glycation rates ($\alpha=10.39$: CI_{95} [-3.713, 24.993], $\beta=0.246$: CI_{95} [-0.144, 0.629], $P_{MCMC}=0.196$; **ESM1** [↗](#)).

We observed no significant effect of age relative to maximum lifespan nor sex on either glycaemia or glycation (see **ESM1** [↗](#)). In some models, GVIF for body mass and/or clutch mass were higher than 1.6, and in one case body mass is slightly above 2.2 (2.26, see **ESM1** [↗](#)). This indicates that there may be moderate collinearity for this variables, but the fact that we used life-history trait covariates that are excluding body mass associated variation, this small effects remain mysterious, but nevertheless probably not worrying.

Discussion

Plasma glucose level and albumin glycation rate (co)variation patterns suggests resistance mechanisms

Our findings show that glucose levels vary widely among different bird groups. Interspecific differences are partly explained by the allometric relationship of glycaemia with body mass (**Figure 2.A** [↗](#)), which has already been reported in previous studies [56 [↗](#), 63 [↗](#), 34 [↗](#), 13 [↗](#), 14 [↗](#)]. Indeed, Passeriformes, Apodiformes and some Columbiformes (e.g. *Nesoenas mayeri* holds the highest value in our dataset) are found at the higher end of the glucose level continuum, in accordance with their relatively small body mass and powered flight, while groups of larger birds such as Phoenicopteriformes, Anseriformes, Rheiformes and Suliformes tend to show low glycaemia levels. This pattern is similar for glycation, with some groups of large birds (such as Phoenicopteriformes, Anseriformes and Suliformes) showing the lowest levels of glycated albumin, while small birds (such as Apodiformes) had the highest values. Nevertheless, glycation remains high in some birds in relation to their glucose levels, as in Rheiformes, or low as in Psittaciformes or Passeriformes. The case of Procellariiformes, which are typically long-lived birds, is particularly striking, with some species exhibiting some of the highest glycation levels (*Calonectris diomedea* and *Macronectes giganteus*) and others some of the lowest (*Procellaria aequinoctialis*). This suggests that, if birds are protected against the deleterious effects of high glycaemia, certain species may have evolved mechanisms to efficiently prevent proteins to be glycated at high rates while others may have evolved mechanisms to resist the consequences of protein glycation. We should also bear in mind that some taxonomic groups may be underrepresented in our study, or biases in species selection due to availability contingencies (e.g. species common in zoos or present in European countries) may exist, so further studies should target this underrepresented groups in order to confirm our predictions.

As expected for a non-enzymatic reaction, just by the law of mass action, bird albumin glycation rates increase with plasma glucose concentration. However, the logarithmic nature of the relationship, and the fact that the slope is lower than one, suggest that species with higher plasma glucose levels exhibit relatively greater resistance to glycation. This finding aligns with previous research indicating that *in vitro* glycation levels of chicken albumin increase at a slower rate than those of human albumin when glucose concentration is elevated [83 [↗](#)]. Moreover, these levels are consistently lower than those of bovine serum albumin regardless of glucose concentration and exposition time [37 [↗](#)]. As discussed in previous studies comparing chicken to bovine serum albumin [37 [↗](#)], or zebra finch to human haemoglobin [53 [↗](#)], the lower glycation rates observed in bird proteins may result from a lower number of glycatable amino acids (e.g. lysines) in their sequence and/or their lesser exposure at the protein surface.

Our analyses do not succeed in indicating a significant positive relationship between average glycation levels and the number of glycatable lysines in the albumin sequence. This may be attributed to the limited number of species employed or the weak variation in the number of

glycatable lysine residues, which are mostly ranging from 33 to 39. An interesting exception are the 18 glycatable lysine residues of flamingos (*Phoenicopterus ruber*), which also shows very low glycation levels (mean=10.1%). However, the exceptions of 44 in zebra finches and 20 in godwits (*Limosa lapponica*, used in place of *Limosa limosa*) are nevertheless associated with very similar average glycation levels of 20.2% and 20.7% respectively.

Plasma glucose relates with longevity and may be influenced by reproductive strategies

Our results were only in minor agreement with predictions from POLS theory: it holds for body mass, but not for the other life history variables tested. The only previous comparative study of glycaemia and life history traits to our knowledge [14] shows no relationship with mass-adjusted maximum lifespan in passerines. However, our study over 88 bird species on 22 orders revealed an increase in glucose with mass-adjusted longevity up to a plateau (see **Figure 2.B**). Thus, the relationship between glucose and maximum lifespan may depend on differences between bird orders or be tied to specific species particularities not explored in our study. Such species particularities might involve additional undetermined ecological factors that modify the relationship of glycaemia with longevity. Further exploration of glucose metabolism in relation with lifestyle will bring further light on species-specific life-history adaptations concerning glucose. For example, the species with the lower mass-adjusted maximum lifespan here was a cormorant (*Leucocarbo verrucosus*), which have quite low glucose values for birds.

Regarding reproductive investment (i.e. clutch mass), our results show no relationship with glycation (see **Table 2**), while previous studies reported positive relationships with glycaemia in passerines [14, 83]. Interestingly, most of the species with high clutch mass included in our study belong to the Anseriformes (Figure ESM2.1). While these species exhibit very low glycaemia and albumin glycation rates, they are also characterized by a particular reproductive strategy compared to passerines, for whom clutch mass does not imply the same in terms of parental investment. For instance, Anseriformes, unlike Passeriformes, are precocial and their offspring are not very dependent on parental care. Furthermore, they are capital breeders, accumulating the energetic resources required for reproduction in advance rather than collecting them during reproduction. These species typically have large clutch sizes and incubation is usually carried out by females, who store internal fat resources to support this main part of the parental investment. In addition, ducklings are born highly developed, reducing the amount of care required post-hatching (see e.g. [85–86]). Consequently, their dependence on glucose as a rapid energy source for reproduction may be lower, with investment in this activity likely more closely linked to lipid accumulation. This could explain why we did not detect previously reported effects of clutch mass on glucose levels.

Terrestrial carnivores show a paradoxically increased albumin glycation rate without increased plasma glucose levels

Contrary to our expectation of finding differences across dietary groups, plasma glucose did not significantly vary with species diet. This aligns with results previously reported by [14] for Passeriformes or by [55] for 160 species of vertebrates among which 48 were birds, but not with [15], that showed glycaemia to be increased with the proportion of fruits/seeds in the diet, or with [56], that showed higher (mass-adjusted) glucose levels for terrestrial carnivores and insectivores (those included in our terrestrial carnivores' category) and lower for frugivorous/nectarivorous. On the other hand, intraspecific data indicates that changes in diet composition hardly affects glycaemia in birds [87]. These studies suggest that glycaemia is tightly regulated independently of dietary composition within species, while it probably varies across species depending on the diet they are adapted to, although depending on the bird groups included in the analyses and the way of assessing it.

Although terrestrial carnivore species did not have significantly higher glycaemia levels in our study, they nonetheless demonstrated significantly higher albumin glycation rates, suggesting a potential susceptibility to protein glycation. Besides unique structural features, this phenomenon could be due to lower albumin turnover rates in terrestrial carnivores. Studies in humans have linked higher oxidative damage to albumin with lower turnover rates (reviewed in [88]), which may extend to other species and post-translational modifications of albumin, such as bird's albumin glycation. However, the contrary outcome would be anticipated if protein intake were high as seen in carnivorous species [88–89].

Interestingly, the contrast between terrestrial and aquatic predators, which did not show such high glycation rates, suggests the beneficial influence of a substantial difference in food composition or the pre-eminence of another ecological factor, yet to be determined, which may explain lower glycation levels in aquatic predators. In terms of food composition, the proportion of essential n-3 PUFA (polyunsaturated fatty acids), particularly long-chain ones such as DHA and EPA compared with n-6 PUFA and short-chain n-3 PUFA, is different in aquatic and terrestrial environments [90] and therefore in the diet of predators in each environment (e.g. [91]). For instance, low n-6 PUFA/n3-PUFA ratio have been shown to decrease insulin levels and improve insulin resistance in humans [92]. On the other hand, while some studies report that an increase in dietary n3-PUFA reduces the levels of glycated haemoglobin (HbA1c) in humans, this effect was not detected in many other studies and therefore remains inconclusive (reviewed in [93]). The detailed study of the fat composition of the diets of terrestrial and aquatic predators, in particular the ratio between different types of PUFA, merits more attention. In addition, micronutrient content, such as circulating antioxidant defences, should also be taken into consideration. Indeed, a positive relationship between oxidative stress and glycated levels of bovine serum albumin has been reported [94]. One hypothesis is that terrestrial predators have higher systemic oxidative stress levels compared to other species, which may be explained by defects in their antioxidant defences. Uric acid is one of the main non-enzymatic antioxidants in birds [95–97], and uric acid levels are especially high in carnivore species that have a rich protein diet [96, 98–101]. We should therefore take into account other important antioxidants such as vitamin E, vitamin A and carotenoids, which may be less abundant in the diets of terrestrial carnivores (e.g. [102], but see [52]). The question of whether the diet of aquatic carnivores provides a better intake of antioxidants would therefore requires a more detailed description of dietary habits. The herbivorous diet, meanwhile, despite the expected possibility contributing to lower glycaemia and glycation levels due to higher levels of PUFA compared to SFA (saturated fatty acids) (effects reviewed in [93] for humans) and higher fibre content (see [103]), did not lead to significantly lower levels of either of these two parameters. However, the relatively low sample size in that group or the presence of outliers as *Nesoenas mayeri*, with the highest glucose levels of this dataset, makes the interpretation of the obtained results somehow limited. Therefore, further research should be carried out on these species to determine if the expected pattern would emerge with a better sampling.

Finally, differences between captive and wild populations could be considered as a source of variation in glucose or glycation levels, due to a more sedentary lifestyle in captivity, with lower activity levels and a higher food intake likely to lead to increased circulating glucose levels in captive individuals. Additionally, differences in nutrient intake, such as antioxidants, specific fatty acids or amino acids, between captive and wild populations could contribute to variations in glycation levels, as we saw above. Nevertheless, we think that this factor is unlikely to significantly affect our findings regarding diet categories, because our study encompasses species from both captive and wild populations across various diet groups, particularly those exhibiting significant differences (e.g., omnivores versus terrestrial carnivores).

Conclusion and perspectives

In conclusion, the avian plasma glucose levels measured here are generally higher or in the high range of values recorded for mammalian species [28]. Our study also concludes that there is considerable variation in plasma glucose levels and albumin glycation rates among bird species, with those with the highest glucose levels showing greater resistance to glycation. The correlation between plasma glucose and life history traits are primarily influenced by its inverse association with body mass, along with non-linear, mass-independent effects of longevity. Finally, although diet does not explain plasma glucose levels in our study, terrestrial carnivores have higher albumin glycation rates than omnivores. Whether these intriguing results are explained by specific effects of certain dietary components, such as PUFAs and/or antioxidants, remains to be determined. Differences in plasma glucose levels, albumin glycation rates and glycation resistance (as glycation levels adjusting for plasma glucose concentration) across bird species do not seem consistent with predictions according to the POL hypothesis, except for body mass. Further investigation is needed to elucidate the correlation between these traits and specific life conditions, such as reproductive strategy, migration patterns, flight mode or more detailed diet composition. In addition, more in-depth exploration of glycation levels within high glycaemic groups such as the Passeriformes and other small birds, which make up a significant proportion of avian species, could provide valuable new insights. Similarly, investigating groups such as the Phoenicopteriformes or Anseriformes, which are at the other end of the glycaemic-glycation spectrum, could shed light on the origin of differences between avian orders. Furthermore, notable variations were observed between species within orders, as in the Procellariiformes, which with their high mass-adjusted longevity, and particularly given that some of the age known individuals in our dataset (indeed, two specimens of snow petrels, *Pagodroma nivea*, showed ages higher than the maximum lifespan reported in the sources we explored, determining a new record) have still relatively low levels of glycation, may suggest that some species (the ones that do show higher glycation levels) are not intrinsically resistant to glycation, but rather to its adverse consequences for health. As the main limitations from this study, the usage of individuals from both wild and captive populations, sampled at different periods of the year, and the low sample size for some species, due to logistic constraints, may have introduced noise in the values reported that should be addressed in future studies by implementing a stricter sampling protocol. Also, a more thorough and accurate report and compilation of life history traits from multiple species would allow to increase the number of species included in this kind of analyses. Future research should also focus on specific species to unravel the physiological mechanisms mediating the effects of blood glucose and protein glycation on life-history trade-offs, in particular mechanisms that may vary between taxa and contribute to characteristic adaptations in birds to mitigate the potentially negative effects of comparatively high glycaemia.

Data and code accessibility

All data used, including references for the values taken from bibliography or databases, will be available as Electronic Supplementary Material (ESM5). Code is made available as Electronic Supplementary Material (ESM7).

Acknowledgements

This research was funded by an ANR (AGEs – ANR21-CE02-0009). We thank Charles-André Bost (CEBC CNRS) for providing data on some species. We would like to thank the SYLATR Association for collecting samples of wild Passeriformes as part of the MIGROUILLE program, and Manuela Forero and Frederic Angelier (CEBC CNRS) for providing some samples of certain Procellariiformes. We thank the French Parc des Oiseaux (Villars les Dombes), the bird keepers and veterinarians at Mulhouse zoo for their contribution to the collection of blood samples and access to individual data from their captive birds. Data on seabirds from the French Southern Territories was collected within the framework of the ECONERGY (119), ECOPATH (1151) and ORNITHOECO (109) programs of the French Polar Institute (IPEV). These studies are part of the long-term Studies in Ecology and Evolution (SEE-Life) program of the CNRS. We are grateful to Mathilde Lejeune, Natacha Garcin and Camille Lemonnier for their help in collecting those samples. We are thankful to Orsolya Vincze for producing **Figure 1** [↗](#), Adrien Brown for determining the number of lysines exposed in albumin sequences, Claire Saraux for her help with the statistics and F. Stephen Dobson for reading and commenting on the manuscript. Finally, we are thankful to Ascensia Diabetes care® for their generous donation of glucometers and strips for glucose measurement.

Additional information

Ethics statement

This study followed all the legal considerations, with the ethic authorisations from the French Ministry of Secondary Education and Research, n°32475 for the zebra finches sampling, the Swiss Veterinary Office (FSVO) n° 34497 for the Alpine swifts, the Ethics Committee of the University of Extremadura (licenses112//2020 and 202//2020) and the Government of Extremadura (licenses CN0012/22/ACA and CN0063/21/ACA) for the godwits and terns from Spain, and Sampling in Terres Australes et Antarctique Françaises was approved by a Regional Animal Experimentation Ethical Committee (French Ministry of Secondary Education and Research permit APAFIS #31773-2019112519421390 v4 and APAFIS#16465–2018080111195526 v4) and by the Comité de l'Environnement Polaire and CNPN (A-2021-55). The samples from the Mulhouse zoo were taken by its licensed veterinary with capacity number: 2020-247-SPAE-162.

Authors' contributions

. Criscuolo and F. Bertile conceived the idea, directed most of the sample collection and logistics and contributed significantly to the writing. A. Moreno-Borralló contributed to the development of the questions, gathered the diet and life-history data, performed the statistical analyses, and led the writing. S. Jaramillo Ortiz and C. Schaeffer performed the mass spectrometry analyses for protein glycation measurements, and S. Jaramillo Ortiz contributed to the glucose measures and commented on the manuscript. T. Boulinier. and V. A. Viblanc coordinated the ECONERGY and ECOPATH polar programs, organised the collection of samples on subantarctic seabirds, and commented on the manuscript. Olivier Chastel collaborated collecting marine bird samples and commented on the manuscript. B. Rey contributed with the collection of samples from Parc des Oiseaux (Villars-les-Dombes, France) and commented on the manuscript. P. Bize leads the monitoring of Alpine swifts' populations from which the samples were obtained, he helped collecting them and commented on the manuscript. J. S. Gutiérrez and J. A. Masero contributed with samples from Spain and part of the statistic scripts and commented on the manuscript. B.

Quintard leads the health monitoring of Mulhouse zoo bird collection, organized and realized most of Mulhouse zoo samplings and commented on the manuscript. All authors gave their approval for publication.

Additional files

ESM1-4 & 6 

References

- [1]. Wikelski Martin, Ricklefs Robert E. (2001) **The physiology of life histories** *Trends in Ecology & Evolution* **16**:479–81 [https://doi.org/10.1016/S0169-5347\(01\)02279-0](https://doi.org/10.1016/S0169-5347(01)02279-0)
- [2]. Ricklefs Robert E., Wikelski Martin (2002) **The physiology/life-history nexus** *Trends in Ecology and Evolution* **17**:462–68 [https://doi.org/10.1016/S0169-5347\(02\)02578-8](https://doi.org/10.1016/S0169-5347(02)02578-8)
- [3]. Wikelski Martin, Spinney Laura, Schelsky Wendy, Scheuerlein Alexander, Gwinner Eberhard (2003) **Slow pace of life in tropical sedentary birds: a common-garden experiment on four stonechat populations from different latitudes** *Proceedings of the Royal Society London B: Biological Sciences* **270**:2383–88 <https://doi.org/10.1098/rspb.2003.2500>
- [4]. Stott Iain, Salguero-Gómez Roberto, Jones Owen R., Ezard Thomas H.G., Gamelon Marlène, Lachish Shelly, Lebreton Jean Dominique, Simmonds Emily G., Gaillard Jean Michel, Hodgson Dave J. (2024) **Life histories are not just fast or slow** *Trends in Ecology and Evolution* **33**:1–11 <https://doi.org/10.1016/j.tree.2024.06.001>
- [5]. Stearns Stephen C (1989) **Trade-offs in life-history evolution** *Functional Ecology* **3**:259–268 <https://doi.org/10.2307/2389364>
- [6]. Gaillard Jean-Michel, Pontier Dominique (1989) **An analysis of demographic tactics in birds and mammals** *Oikos* **56**:59–76 <https://doi.org/10.2307/3566088>
- [7]. Stearns Stephen C (1992) **The evolution of life histories** Oxford university press
- [8]. Vasilieva N. A, Batrukova M. (2022) **Pace-of-Life Syndrome (POLS): evolution of the concept** *Biology bulletin* **49**:750–62 <https://doi.org/10.1134/S1062359022070238>
- [9]. Trevelyan R., Harvey P. H., Pagel M. D. (1990) **Metabolic rates and life histories in birds** *Functional Ecology* **4**:135–141 <https://doi.org/10.2307/2389332>
- [10]. Austad Steven N., Fischer Kathleen E. (1991) **Mammalian aging, metabolism, and ecology: evidence from the bats and marsupials** *Journal of Gerontology: Biological sciences* **46**:47–53 <https://doi.org/10.1093/geronj/46.2.B47>
- [11]. Speakman John R (2005) **Body size, energy metabolism and lifespan** *Journal of Experimental Biology* **208**:1717–30 <https://doi.org/10.1242/jeb.01556>
- [12]. Magalhães João Pedro de, Costa Joana, Church George M. (2007) **An analysis of the relationship between metabolism, developmental schedules, and longevity using Phylogenetic Independent Contrasts** *The Journals of Gerontology: Series A* **62**:149–60 <https://doi.org/10.1093/gerona/62.2.149>
- [13]. Kjeld M, Ólafsson Ö. (2007) **Allometry (scaling) of blood components in mammals: connection with economy of energy?** *Canadian Journal of Zoology* **89**:890–99 <https://doi.org/10.1139/Z08-061>
- [14]. Tomášek Oldřich, Bobek Lukáš, Kauzálová Tereza, Adamkova Marie, Albrecht Tomáš (2019) **Fuel for the pace of life: baseline blood glucose concentration co-evolves with life-history**

traits in songbirds *Functional Ecology* **33**:239–49 <https://doi.org/10.1111/1365-2435.13238>

- [15]. Tomášek Oldřich, Bobek Lukáš, Kauzálová Tereza, Kauzál Ondřej, Adámková Marie, Horák Kryštof, Kumar Sampath Anandan, Manialeu Judith Pouadjeu, Munclinger Pavel, Nana Eric Djomo, Nguelefack Télesphore Benoît, Sedláček Ondřej, Albrecht Tomáš (2022) **Latitudinal but not elevational variation in blood glucose level is linked to life history across passerine birds** *Ecology Letters* **25**:2203–2216 <https://doi.org/10.1111/ele.14097>
- [16]. Vágási Csongor I., Vincze Orsolya, Adámková Marie, Kauzálová Tereza, Lendvai Ádam Z., Pátrás Laura I., Péntzes Janka, Pap Péter L., Albrecht Tomáš, Tomášek Oldřich (2024) **Songbirds avoid the oxidative stress costs of high blood glucose levels: a comparative study** *Journal of Experimental Biology* **227**:1–8 <https://doi.org/10.1242/jeb.246848>
- [17]. Umminger Bruce L (1977) **Relation of whole blood sugar concentrations in vertebrates to standard metabolic rate** *Comparative Biochemistry and Physiology A* **56**:457–460
- [18]. Guo Hai, Duolikun Dilihumai, Yao Qiaoling (2023) **Associations between basal metabolic rate and insulin resistance in non-diabetic obese adults: evidence from NHANES 2011–2018** *International Journal of Diabetes in Developing Countries* <https://doi.org/10.1007/s13410-023-01190-8>
- [19]. Maillard L. C (1912) **Action des acides amines sur les sucres; formation des melanoidies par voie methodique** *Comptes rendus de l'Académie des sciences* **154**:66–68
- [20]. Suji George, Sivakami S. (2004) **Glucose, glycation and aging** *Biogerontology* **5**:365–373 <https://doi.org/10.1007/s10522-004-3189-0>
- [21]. Cerami Anthony, Vlassara Helen, Brownlee Michael (1986) **Role of nonenzymatic glycosylation in atherogenesis** *Journal of Cellular Biochemistry* **30**:111–120 <https://doi.org/10.1002/jcb.240300203>
- [22]. Poulsen Malene W., Hedegaard Rikke V., Andersen Jeanette M., de Courten Barbora, Bügel Susanne, Nielsen John, Skibsted Leif H., Dragsted Lars O. (2013) **Advanced Glycation End-products in food and their effects on health** *Food and Chemical Toxicology* **60**:10–37 <https://doi.org/10.1016/j.fct.2013.06.052>
- [23]. Chaudhuri Jyotiska, Bains Yasmin, Guha Sanjib, Kahn Arnold, Hall David, Bose Neelanjan, Gugliucci Alejandro, Kapahi Pankaj (2018) **The role of advanced glycation end products in aging and metabolic diseases: bridging association and causality** *Cell Metabolism* **28**:337–52 <https://doi.org/10.1016/j.cmet.2018.08.014>
- [24]. Khalid Mariyam, Petroianu Georg, Adem Abdu (2022) **Advanced Glycation End products and diabetes mellitus: mechanisms and perspectives** *Biomolecules* **12**:542 <https://doi.org/10.3390/biom12040542>
- [25]. Twarda-Clapa Aleksandra, Olczak Aleksandra, Białkowska Aneta M., Koziółkiewicz Maria (2022) **Advanced Glycation End-products (AGEs): formation, chemistry, classification, receptors, and diseases related to AGEs** *Cells* **11**:1312 <https://doi.org/10.3390/cells11081312>
- [26]. Hofmann Marion A., Drury Steven, Fu Caifeng, Qu Wu, Taguchi Akihiko, Lu Yan, Avila Cecilia, Kambham Neeraja, Bierhaus Angelika, Nawroth Peter, Neurath Markus F., Slattey Timothy, Beach Dale, McClary John, Nagashima Mariko, Morser John, Stern David, Schmidt Ann Marie (1999) **RAGE mediates a novel proinflammatory axis: a central cell surface receptor for**

S100/Calgranulin polypeptides *Cell* **97**:889–901 [https://doi.org/10.1016/S0092-8674\(00\)80801-6](https://doi.org/10.1016/S0092-8674(00)80801-6)

- [27]. Schmidt Ann Marie, Du Yan Shi, Yan Shi Fang, Stern David M. (2001) **The multiligand receptor RAGE as a progression factor amplifying immune and inflammatory responses** *Journal of Clinical Investigation* **108**:949–55 <https://doi.org/10.1172/JCI200114002>
- [28]. Polakof Sergio, Mommsen Thomas P., Soengas José L. (2011) **Glucosensing and glucose homeostasis: from fish to mammals** *Comparative Biochemistry and Physiology* **160**:123–49 <https://doi.org/10.1016/j.cbpb.2011.07.006>
- [29]. Lasiewski Robert C., Dawson William R. (1967) **A re-examination of the relation between standard metabolic rate and body weight in birds** *The Condor* **69**:13–23 <https://doi.org/10.2307/1366368>
- [30]. King J. R., Farner D. S (1961) **Energy metabolism, thermoregulation and body temperature** In: Marshall A. J., editors. *Biology and Comparative Physiology of Birds* New York: Academic Press pp. 215–288
- [31]. Dawson Terence J., Hulbert A. J. (1970) **Standard metabolism, body temperature, surface areas of australian marsupials** *American Journal of Physiology* **218** <https://doi.org/10.1152/ajplegacy.1970.218.4.1233>
- [32]. Marsh Richard L., Ellerby David J., Carr Jennifer A., Henry Havalee T., Buchanan Cindy I. (2004) **Partitioning the energetics of walking and running: swinging the limbs is expensive** *Science* **303**:80–83 <https://doi.org/10.1126/science.1090704>
- [33]. Weber Jean-Michel (2011) **Metabolic fuels: regulating fluxes to select mix** *Journal Experimental Biology* **214**:286–94 <https://doi.org/10.1242/jeb.047050>
- [34]. Braun Eldon J., Sweazea Karen L. (2008) **Glucose regulation in birds** *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology* **151**:1–9 <https://doi.org/10.1016/j.cbpb.2008.05.007>
- [35]. Szwergold Benjamin S., Miller Craig B. (2014b) **Potential of birds to serve as pathology-free models of type 2 diabetes, Part 2: Do high levels of carbonyl-scavenging amino acids (e.g., taurine) and low concentrations of methylglyoxal limit the production of Advanced Glycation End-products?** *Rejuvenation Research* **17**:347–58 <https://doi.org/10.1089/rej.2014.1561>
- [36]. Lindstedt Stan L., Calder William A. (1976) **Body size and longevity in birds** *The Condor* **78**:91–94 <https://doi.org/10.2307/1366920>
- [37]. Anthony-Regnitz Claire M., Wilson Amanda E., Sweazea Karen L., Braun Eldon J. (2020) **Fewer exposed lysine residues may explain relative resistance of chicken serum albumin to in vitro protein glycation in comparison to bovine serum albumin** *Journal of Molecular Evolution* **88**:653–61 <https://doi.org/10.1007/s00239-020-09964-y>
- [38]. Muramatsu Tatsuo (1990) **Nutrition and whole-body protein turnover in the chicken in relation to mammalian species** *Nutrition Research reviews* **3**:211–228
- [39]. Makino Ryosuke, Kita Kazumi (2018) **Half-Life of Glycated Tryptophan in the Plasma of Chickens** *Journal of Poultry Science* **55**:117–19 <https://doi.org/10.2141/jpsa.0170158>

- [40]. Schweigert F. J., Uehlein-Harrell Stephanie, Hegel Giesela v., Wiesner H. (1991) **Vitamin A (retinol and retinyl esters), α -tocopherol and lipid levels in plasma of captive wild mammals and birds** *Journal of Veterinary Medicine Series A* **42**:35–42 <https://doi.org/10.1111/j.1439-0442.1991.tb00981.x>
- [41]. Ku Hung-Hai, Sohal R.S. (1993) **Comparison of mitochondrial prooxidant generation and antioxidant defenses between rat and pigeon possible basis of variation in longevity and metabolic potential** *Mechanisms of Ageing and Development* **72**:67–76 [https://doi.org/10.1016/0047-6374\(93\)90132-B](https://doi.org/10.1016/0047-6374(93)90132-B)
- [42]. Hickey Anthony J.R., Jüllig Mia, Aitken Jacqueline, Loomes Kerry, Hauber Mark E., Phillips Anthony R.J. (2012) **Birds and longevity: does flight driven aerobicity provide an oxidative sink?** *Ageing Research Reviews* **11**:242–53 <https://doi.org/10.1016/j.arr.2011.12.002>
- [43]. Eythrib Jalil (2013) **The search for the receptor for Advanced Glycation End-products in avian vasculature. Bachelor thesis** The University of Arizona
- [44]. Szwergold Benjamin S., Miller Craig B. (2014a) **Potential of birds to serve as a pathology-free model of type 2 diabetes, Part 1: Is the apparent absence of the rage gene a factor in the resistance of avian organisms to chronic hyperglycemia?** *Rejuvenation Research* **17**:54–61 <https://doi.org/10.1089/rej.2013.1498>
- [45]. Wein Yossi, Shira Enav Bar, Friedman Aharon (2020) **Increased serum levels of advanced glycation end products due to induced molting in hen layers trigger a proinflammatory response by peripheral blood leukocytes** *Poultry Science* **99**:3452–62 <https://doi.org/10.1016/j.psj.2020.04.009>
- [46]. Rendell Marc, Stephent P. M., Paulsent Robert, Valentine J. L., Rasbold Kathy, Hestorfe T. I. M., Eastberg Susan, Shintt Daniel C (1985) **An interspecies comparison of normal levels of glycosylated hemoglobin and glycosylated albumin** *Comparative Biochemistry and physiology. B, Comparative Biochemistry* **81**:819–22 [https://doi.org/10.1016/0305-0491\(85\)90072-0](https://doi.org/10.1016/0305-0491(85)90072-0)
- [47]. Miksik I., Hodny Z. (1992) **Glycated hemoglobin in mute swan (*Cygnus olor*) and rook (*Corvus frugilegus*)** *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* **103**:553–55 [https://doi.org/10.1016/0305-0491\(92\)90369-3](https://doi.org/10.1016/0305-0491(92)90369-3)
- [48]. Andersson M. S., Gustafsson L. (1995) **Glycosylated haemoglobin: a new measure of condition in birds** *Proceedings of the Royal Society B: Biological Sciences* **260**:299–303 <https://doi.org/10.1098/rspb.1995.0095>
- [49]. Beuchat Carol A., Chong Curtis R. (1998) **Hyperglycemia in hummingbirds and its consequences for hemoglobin glycation** *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology* **120**:409–16 [https://doi.org/10.1016/S1095-6433\(98\)10039-9](https://doi.org/10.1016/S1095-6433(98)10039-9)
- [50]. Ardia Daniel R (2006) **Glycated hemoglobin and albumin reflect nestling growth and condition in american kestrels** *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology* **143**:62–66 <https://doi.org/10.1016/j.cbpa.2005.10.024>
- [51]. Récapet Charlotte, Adélaïde Sibeaux, Cauchard Laure, Doligez Blandine, Bize Pierre (2016) **Selective disappearance of individuals with high levels of glycated haemoglobin in a free-living bird** *Biology Letters* **12**:20160243 <https://doi.org/10.1098/rsbl.2016.0243>

- [52]. Ingram Tana, Zuck Jessica, Borges Chad R., Redig Patrick, Sweazea Karen L. (2017) **Variations in native protein glycation and plasma antioxidants in several birds of prey** *Comparative Biochemistry and Physiology Part - B: Biochemistry and Molecular Biology* **210**:18–28 <https://doi.org/10.1016/j.cbpb.2017.05.004>
- [53]. Brun Charlotte, Hernandez-Alba Oscar, Hovasse Agnès, Criscuolo François, Schaeffer-Reiss Christine, Bertile Fabrice (2022) **Resistance to glycation in the zebra finch: mass spectrometry-based analysis and its perspectives for evolutionary studies of aging** *Experimental Gerontology* **164**:111811 <https://doi.org/10.1016/j.exger.2022.111811>
- [54]. Borger M. J (2024) **How to prime your offspring: putting behavioural ecology to the test** PhD thesis. University of Groningen <https://doi.org/10.33612/diss.849300689>
- [55]. Kapsetaki Stefania E., Basile Anthony J., Compton Zachary T., Rupp Shawn M., Duke G., Boddy Amy M., Harrison Tara M., Sweazea Karen L., Maley Carlo C. (2023) **The relationship between diet, plasma glucose, and cancer prevalence across vertebrates. Version 1** *bioRxiv* <https://doi.org/10.1101/2023.07.31.551378>
- [56]. Szarka E.Z., Lendvai Á.Z. (2024) **Trophic guilds differ in blood glucose concentrations: a phylogenetic comparative analysis in birds** *Proceedings of the Royal Society B: Biological Sciences* **291**:20232655
- [57]. Coulson Roland A., Hernandez Thomas (1983) **Alligator Metabolism studies on chemical reactions in vivo** *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* **74**:1–175 [https://doi.org/10.1016/0305-0491\(83\)90418-2](https://doi.org/10.1016/0305-0491(83)90418-2)
- [58]. Dobbs J. C (1985) **Avian carnivore: food recognition, nutrient intakes, and utilization of nitrogen.** PhD thesis, University of California-Davis, Davis *Ca*
- [59]. Verbrugghe Adronie, Hesta Myriam (2017) **Cats and carbohydrates: the carnivore fantasy?** *Veterinary Sciences* **4**:1–22 <https://doi.org/10.3390/vetsci4040055>
- [60]. Migliorini R. H., Linder J. C., Moura L., Veiga J. A. S. (1973) **Gluconeogenesis in a carnivorous bird (black vulture)** *American journal of physiology* **225**:1389–1392 <https://doi.org/10.1152/ajplegacy.1973.225.6.1389>
- [61]. Myers Merrick R., Klasing Kirk C. (1999) **Low glucokinase activity and high rates of gluconeogenesis contribute to hyperglycemia in barn owls (*Tyto alba*) after a glucose challenge** *The Journal of Nutrition* **129**:1896–1904 <https://doi.org/10.1093/jn/129.10.1896>
- [62]. Stearns Stephen C (1983) **The Influence of size and phylogeny on patterns of covariation among life-history traits in the mammals** *Oikos* **41**:173–187
- [63]. Umminger Bruce L (1975) **Body size and whole blood sugar concentrations in mammals** *Comparative Biochemistry and Physiology A* **52**:455–458
- [64]. Scanes Colin G (2016) **A re-evaluation of allometric relationships for circulating concentrations of glucose in mammals** *Food and Nutrition Sciences* **7**:240–51 <https://doi.org/10.4236/fns.2016.74026>
- [65]. Mohsenzadeh Mahdieh Sadat, Zaeemi Mahdieh, Razmyar Jamshid, Azizzadeh Mohammad (2015) **Comparison of a point-of-care glucometer and a laboratory autoanalyzer for measurement of blood glucose concentrations in domestic pigeons (*Columba livia domestica*)** *Journal of Avian Medicine and Surgery* **29**:181–86 <https://doi.org/10.1647/2014-020>

- [66]. Morales Ana, Frei Barbara, Leung Casey, Titman Rodger, Whelan Shannon, Morgan Benowitz-Fredericks Z., Elliott Kyle H. (2020) **Point-of-care blood analyzers measure the nutritional state of eighteen free-living bird species** *Comparative Biochemistry and Physiology -Part A: Molecular and Integrative Physiology* **240**:1095–6433 <https://doi.org/10.1016/j.cbpa.2019.110594>
- [67]. Downs Colleen T., Wellmann Andrea E., Brown Mark (2010) **Diel variations in plasma glucose concentrations of malachite sunbirds *Nectarinia famosa*** *Journal of Ornithology* **151**:235–39 <https://doi.org/10.1007/s10336-009-0439-6>
- [68]. Breuner Creagh W., Delehanty Brendan, Boonstra Rudy (2013) **Evaluating stress in natural populations of vertebrates: total CORT is not good enough** *Functional Ecology* **27**:24–36 <https://doi.org/10.1111/1365-2435.12016>
- [69]. McGraw Kevin J., Chou Katherine, Bridge Annika, McGraw Hannah C., McGraw Peyton R., Simpson Richard K. (2020) **Body condition and poxvirus infection predict circulating glucose levels in a colorful songbird that inhabits urban and rural environments** *Journal of Experimental Zoology* **333**:561–68 <https://doi.org/10.1002/jez.2391>
- [70]. Priego-Capote F., Ramirez-Boo M., Finamore F., Gluck F., Sanchez J.C (2014) **Quantitative analysis of glycated proteins** *Journal of Proteome Research* **13**:336–347
- [71]. Zuck Jessica, Borges Chad R., Braun Eldon J., Sweazea Karen L. (2017) **Chicken albumin exhibits natural resistance to glycation** *Comparative Biochemistry and Physiology Part - B: Biochemistry and Molecular Biology* **203**:108–114 <https://doi.org/10.1016/j.cbpb.2016.10.003>
- [72]. Tobias Joseph A., Sheard Catherine, Pigot Alex L., Devenish Adam J. M., Yang Jingyi, Sayol Ferran, Neate-Clegg Montague H. C., Alioravainen Nico, Weeks Thomas L., Barber Robert A., Walkden Patrick A., MacGregor Hannah E. A., Jones Samuel E. I., Vincent Claire, Phillips Anna G., Marples Nicola M., Montaña-Centellas Flavia A., Leandro-Silva Victor, Claramunt Santiago, Darski Bianca, Freeman Benjamin G., Bregman Tom P., Cooney Christopher R., Hughes Emma C., Capp Elliot J. R., Varley Zoë K., Friedman Nicholas R., Korntheuer Heiko, Corrales-Vargas Andrea, Trisos Christopher H., Weeks Brian C., Hanz Dagmar M., Töpfer Till, Bravo Gustavo A., Remeš Vladimír, Nowak Larissa, Carneiro Lincoln S., Amikar J., Moncada R., Matysioková Beata, Baldassarre Daniel T., Martínez-Salinas Alejandra, Wolfe Jared D., Chapman Philip M., Daly Benjamin G., Sorensen Marjorie C., Neu Alexander, Ford Michael A., Mayhew Rebekah J., Silveira Luis Fabio, Kelly David J., Annorabah Nathaniel N. D., Pollock Henry S., Grabowska-Zhang Ada M., McEntee Jay P., Carlos T Juan, Gonzalez Camila G, Meneses Marcia C, Muñoz Luke L, Powell Gabriel A, Jamie Thomas J, Matthews Oscar Johnson, Brito Guilherme R. R., Zyskowski Kristof, Crates Ross, Harvey Michael G., Zevallos Maura Jurado, Hosner Peter A., Bradfer-Lawrence Tom, Maley James M., Gary Stiles F., Lima Hevana S., Provost Kaiya L., Chibesa Moses, Mashao Mmatjie, Howard Jeffrey T., Mlamba Edson, Chua Marcus A. H., Li Bicheng, Isabel Gómez M., García Natalia C., Päckert Martin, Fuchs Jérôme, Ali Jarome R., Derryberry Elizabeth P., Carlson Monica L., Urriza Rolly C., Brzeski Kristin E., Prawiradilaga Dewi M., Rayner Matt J., Miller Eliot T., Bowie Rauri C. K., Lafontaine René-Marie, Paul Scofield R., Lou Yingqiang, Somarathna Lankani, Lepage Denis, Illif Marshall, Neuschulz Eike Lena, Templin Mathias, Matthias Dehling D., Cooper Jacob C., Pauwels Olivier S. G., Analuddin Kangkuso, Fjeldsø Jon, Seddon Nathalie, Sweet Paul R., DeClerck Fabrice A. J., Naka Luciano N., Brawn Jeffrey D., Aleixo Alexandre, Böhning-Gaese Katrin, Rahbek Carsten, Fritz Susanne A., Thomas Gavin H., Schleuning Matthias (2022) **AVONET: Morphological, ecological and geographical data for all birds** *Ecology Letters* **25**:581–97 <https://doi.org/10.1111/ele.13898>
- [73]. Pigot Alex L., Sheard Catherine, Miller Eliot T., Bregman Tom P., Freeman Benjamin G., Roll Uri, Seddon Nathalie, Trisos Christopher H., Weeks Brian C., Tobias Joseph A. (2020)

Macroevolutionary convergence connects morphological form to ecological function in birds *Nature Ecology & Evolution* **4**:230–239 <https://doi.org/10.1038/s41559-019-1070-4>

- [74]. Wilman Hamish, Rivadeneira Jonathan Belmaker Marcelo M., Simpson Jennifer, de la Rosa Carolina, Jetz Walter (2014) **EltonTraits 1.0: species-level foraging attributes of the world's birds and mammals** *Ecology* **95**:2027 <https://doi.org/10.1890/13-1917.1>
- [75]. Myhrvold N. P., Baldrige E., Chan B., Sivam D., Freeman D. L., Ernest S. K. M (2015) **An amniote life-history database to perform comparative analyses with birds, mammals, and reptiles** *Ecology* **96**:3109
- [76]. Tacutu Robi, Thornton Daniel, Johnson Emily, Budovsky Arie, Barardo Diogo, Craig Thomas, Diana Eugene, Lehmann Gilad, Toren Dmitri, Wang Jingwei, Fraifeld Vadim E., de Magalhães João P. (2018) **Human Ageing Genomic Resources: new and updated databases** *Nucleic Acids Research* **46**:D1083–D1090 <https://doi.org/10.1093/nar/gkx1042>
- [77]. Billerman S. M., Keeney B. K., Rodewald P. G., Schulenberg T. S. (2022) **Birds of the World**. Ithaca, NY: Cornell Laboratory of Ornithology <https://birdsoftheworld.org/bow/home>
- [78]. Fransson T., Kolehmainen T., Moss D., Robinson R (2023) **EURING list of longevity records for European birds**
- [79]. R: A language and environment for statistical computing (2023) **R Foundation for Statistical Computing** <https://www.R-project.org/>
- [80]. Hadfield Jarrod D. (2010) **MCMC Methods for multi-response generalized linear mixed models: the MCMCglmm R package** *Journal of Statistical Software* **33**:2
- [81]. Revell L. J (2024) **. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things)** *PeerJ* **12**:e16505 <https://doi.org/10.7717/peerj.16505>
- [82]. Jetz W., Thomas G. H., Joy J. B., Hartmann K., Mooers A. O. (2012) **The global diversity of birds in space and time** *Nature* **491**:444–448
- [83]. Hackett Shannon J., Kimball Rebecca T., Reddy Sushma, Bowie Rauri C. K., Braun Edward L., Braun Michael J., Chojnowski Jena L., Andrew Cox W., Han Kin-lan, Harshman John, Huddleston Christopher J., Marks Ben D., Miglia Kathleen J., Moore William S., Sheldon Frederick H., Steadman David W., Witt Christopher C., Yuri Tamaki (2008) **A phylogenomic study of birds reveals their evolutionary history** *Science* **320**:1763–68 <https://doi.org/10.1126/science.1157704>
- [84]. Nahhas R. W. (2024) **Introduction to Regression Methods for Public Health Using R** <https://www.bookdown.org/rwnahhas/RMPH/mlr-collinearity.html#generalized-vifs-when-at-least-one-predictor-is-categorical>
- [85]. Moreno J (1989) **Strategies of mass change in breeding birds** *Biological Journal of the Linnean Society* **37**:297–310 <https://doi.org/10.1111/j.1095-8312.1989.tb01907.x>
- [86]. Winkler D. W (2016) **Breeding biology of birds** In: Lovette Irby J., Fitzpatrick John W., editors. *Handbook of bird biology* Cornell laboratory of ornithology pp. 407–450
- [87]. Basile Anthony J., Singh Kavita C., Watson Deborah F., Sweazea Karen L. (2022) **Effect of macronutrient and micronutrient manipulation on avian blood glucose concentration: a**

systematic review *Comparative Biochemistry and Physiology, Part A* **272**:July <https://doi.org/10.1016/j.cbpa.2022.111279>

- [88]. Wada Yasuaki, Takeda Yasuhiro, Kuwahata Masashi (2018) **Potential role of amino acid/protein nutrition and exercise in serum albumin redox state** *Nutrients* **10**:1–11 <https://doi.org/10.3390/nu10010017>
- [89]. Honma Ayaka, Ogawa Chiaki, Sugahara Misaki, Fujimura Shinobu, Kita Kazumi (2017) **Influence of varying dietary protein levels on glycation of albumin, tryptophan and valine in the plasma of chickens** *Journal of Poultry Science* **54**:242–46 <https://doi.org/10.2141/jpsa.0160146>
- [90]. Colombo Stefanie M., Wacker Alexander, Parrish Christopher C., Martin J., Arts Michael T., Colombo Stefanie M., Wacker Alexander, Parrish Christopher C., Kainz Martin J., Arts Michael T. (2017) **A fundamental dichotomy in long-chain polyunsaturated fatty acid abundance between and within marine and terrestrial ecosystems** *Environmental Reviews* **25**:163–74 <https://doi.org/10.1139/er-2016-0062>
- [91]. Charles Apostolos-Manuel Koussoroplis, Alexandre Lemarchand, Berny Philippe, Bourdier Gilles (2008) **From aquatic to terrestrial food webs: decrease of the docosahexaenoic acid/linoleic acid ratio** *Lipids* **43**:461–66 <https://doi.org/10.1007/s11745-008-3166-5>
- [92]. Li Na, Yue Hao, Jia Min, Liu Wei, Qiu Bin, Hou Hanxue, Huang Fenghong, Xu Tongcheng (2019) **Effect of low-ratio n-6/n-3 pufa on blood glucose: a meta-analysis** *Food and Function* **10**:4557–65 <https://doi.org/10.1039/c9fo00323a>
- [93]. Telle-Hansen Vibeke H., Gaundal Line, Myhrstad Mari C.W. (2019) **Polyunsaturated fatty acids and glycemic control in type 2 diabetes** *Nutrients* **11**:12–15 <https://doi.org/10.3390/nu11051067>
- [94]. Bavkar Laxman N., Patil Rahul S., Rooge Sheetalnath B., Nalawade Megha L., Arvindekar Akalpita U. (2019) **Acceleration of protein glycation by oxidative stress and comparative role of antioxidant and protein glycation inhibitor** *Molecular and Cellular Biochemistry* <https://doi.org/10.1007/s11010-019-03550-7>
- [95]. Klandorf H., Probert I. L., Iqbal M. (1999) **In the defence against hyperglycaemia: an avian strategy** *World's Poultry Science Journal* **55**:251–68 <https://doi.org/10.1079/WPS19990019>
- [96]. Machín M., Simoyi M. F., Blemings K. P., Klandorf H. (2004) **Increased dietary protein elevates plasma uric acid and is associated with decreased oxidative stress in rapidly-growing broilers** *Comparative Biochemistry and Physiology Part B* **137**:383–90 <https://doi.org/10.1016/j.cbpc.2004.01.002>
- [97]. Stinefelt Beth, Leonard Stephen S., Blemings Kenneth P., Shi Xianglin, Klandorf Hillar (2005) **Free radical scavenging, dna protection, and inhibition of lipid peroxidation mediated by uric acid** *Anal of clinical and laboratory science* **35**:37–45
- [98]. Harr Kendal E (2002) **Clinical chemistry of companion avian species: a review** *Veterinary Clinical Pathology* **31** <https://doi.org/10.18637/jss.v033.i02>
- [99]. Smith Susan B., McWilliams Scott R., Guglielmo Christopher G. (2007) **Effect of diet composition on plasma metabolite profiles in a migratory songbird** *The Condor* **109**:48–58 <https://doi.org/10.1093/condor/109.1.48>

- [100]. Cohen Alan A., McGraw Kevin J., Douglas Robinson W. (2009) **Serum antioxidant levels in wild birds vary in relation to diet, season, life history strategy, and species** *Oecologia* **161**:673–83 <https://doi.org/10.1007/s00442-009-1423-9>
- [101]. Alan Rebecca R., McWilliams Scott R. (2013) **Oxidative stress, circulating antioxidants, and dietary preferences in songbirds. Comparative Biochemistry and Physiology Part B** **164**:185–93 <https://doi.org/10.1016/j.cbpb.2012.12.005>
- [102]. Schneeberger Karin, Cziráj Gábor Á., Voigt Christian C. (2014) **Frugivory is associated with low measures of plasma oxidative stress and high antioxidant concentration in free-ranging bats** *Naturwissenschaften* **101**:285–90 <https://doi.org/10.1007/s00114-014-1155-5>
- [103]. Goff H. Douglas, Repin Nikolay, Fabek Hrvoje, Khoury Dalia El, Gidley Michael J. (2018) **Dietary fibre for glycaemia control: towards a mechanistic understanding** *Bioactive Carbohydrates and Dietary Fibre* **14**:39–53 <https://doi.org/10.1016/j.bcdf.2017.07.005>

Author information

Adrián Moreno-Borralló

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178, Strasbourg, France
ORCID iD: [0000-0002-2924-1153](https://orcid.org/0000-0002-2924-1153)

For correspondence: adrian.moreno-borralló@iphc.cnrs.fr

Sarahi Jaramillo-Ortiz

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178, Strasbourg, France, National Proteomics Infrastructure, ProFi, Strasbourg, France
ORCID iD: [0000-0002-9153-4205](https://orcid.org/0000-0002-9153-4205)

Christine Schaeffer-Reiss

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178, Strasbourg, France, National Proteomics Infrastructure, ProFi, Strasbourg, France
ORCID iD: [0000-0003-0672-1979](https://orcid.org/0000-0003-0672-1979)

Benoît Quintard

Parc zoologique et botanique de Mulhouse, Mulhouse, France
ORCID iD: [0000-0001-7905-886X](https://orcid.org/0000-0001-7905-886X)

Benjamin Rey

Laboratoire de Biométrie et Biologie Evolutive, Villeurbanne, France
ORCID iD: [0000-0002-0464-5573](https://orcid.org/0000-0002-0464-5573)

Pierre Bize

Swiss Ornithological Institute, Sempach, Switzerland
ORCID iD: [0000-0002-6759-4371](https://orcid.org/0000-0002-6759-4371)

Vincent A Viblanc

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178,
Strasbourg, France

ORCID iD: [0000-0002-4953-659X](https://orcid.org/0000-0002-4953-659X)

Thierry Boulinier

CEFE, Montpellier University, CNRS, EPHE, IRD, Montpellier, France

ORCID iD: [0000-0002-5898-7667](https://orcid.org/0000-0002-5898-7667)

Olivier Chastel

Center of Biological Studies of Chizé (CEBC), UMR 7372 CNRS - La Rochelle University, Villiers-en-Bois, France

ORCID iD: [0000-0002-9435-1041](https://orcid.org/0000-0002-9435-1041)

Jorge S Gutiérrez

Ecology in the Anthropocene, Associated Unit CSIC-UEX, Faculty of Sciences, University of Extremadura, Badajoz, Spain

ORCID iD: [0000-0001-8459-3162](https://orcid.org/0000-0001-8459-3162)

Jose A Masero

Ecology in the Anthropocene, Associated Unit CSIC-UEX, Faculty of Sciences, University of Extremadura, Badajoz, Spain

ORCID iD: [0000-0001-5318-4833](https://orcid.org/0000-0001-5318-4833)

Fabrice Bertile*

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178, Strasbourg, France, National Proteomics Infrastructure, ProFi, Strasbourg, France

ORCID iD: [0000-0001-5510-4868](https://orcid.org/0000-0001-5510-4868)

*These authors share senior-authorship

François Criscuolo*

University of Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR 7178, Strasbourg, France

ORCID iD: [0000-0001-8997-8184](https://orcid.org/0000-0001-8997-8184)

*These authors share senior-authorship

Editors

Reviewing Editor

Jenny Tung

Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

Senior Editor

George Perry

Pennsylvania State University, University Park, United States of America

Reviewer #2 (Public review):**Summary:**

In this extensive comparative study, Moreno-Borralló and colleagues examine the relationships between plasma glucose levels, albumin glycation levels, diet and life-history traits across birds. Their results confirmed the expected positive relationship between plasma blood glucose level and albumin glycation rate but also provided findings that are somewhat surprising or contrast with findings of some previous studies (positive relationships between blood glucose and lifespan, or absent relationships between blood glucose and clutch mass or diet). This is the first extensive comparative analysis of glycation rates and their relationships to plasma glucose levels and life history traits in birds that is based on data collected in a single study, with blood glucose and glycation measured using unified analytical methods (except for blood glucose data for 13 species collected from a database).

Strengths:

This is an emerging topic gaining momentum in evolutionary physiology, which makes this study a timely, novel and important contribution. The study is based on a novel data set collected by the authors from 88 bird species (67 in captivity, 21 in the wild) of 22 orders, except for 13 species, for which data were collected from a database of veterinary and animal care records of zoo animals (ZIMS). This novel data set itself greatly contributes to the pool of available data on avian glycemia, as previous comparative studies either extracted data from various studies or a ZIMS database (therefore potentially containing much more noise due to different methodologies or other unstandardised factors), or only collected data from a single order, namely Passeriformes. The data further represents the first comparative avian data set on albumin glycation obtained using a unified methodology. The authors used LC-MS to determine glycation levels, which does not have problems with specificity and sensitivity that may occur with assays used in previous studies. The data analysis is thorough, and the conclusions are substantiated. Overall, this is an important study representing a substantial contribution to the emerging field evolutionary physiology focused on ecology and evolution of blood/plasma glucose levels and resistance to glycation.

Weaknesses:

Unfortunately, the authors did not record handling time (i.e., time elapsed between capture and blood sampling), which may be an important source of noise because handling-stress-induced increase in blood glucose has previously been reported. Moreover, the authors themselves demonstrate that handling stress increases variance in blood glucose levels. Both effects (elevated mean and variance) are evident in Figure ESM1.2. However, this likely makes their significant findings regarding glucose levels and their associations with lifespan or glycation rate more conservative, as highlighted by the authors.

<https://doi.org/10.7554/eLife.103205.3.sa1>

Author response:

The following is the authors' response to the previous reviews

Public Reviews:**Reviewer #2 (Public review):****Summary**

In this extensive comparative study, Moreno-Borralló and colleagues examine the relationships between plasma glucose levels, albumin glycation levels, diet and life history traits across birds. Their results confirmed the expected positive relationship between plasma blood glucose level and albumin glycation rate but also provided findings that are somewhat surprising or contrast with findings of some previous studies (positive relationships between blood glucose and lifespan, or absent relationships between blood glucose and clutch mass or diet). This is the first extensive comparative analysis of glycation rates and their relationships to plasma glucose levels and life history traits in birds that is based on data collected in a single study, with blood glucose and glycation measured using unified analytical methods (except for blood glucose data for 13 species collected from a database).

Strengths

This is an emerging topic gaining momentum in evolutionary physiology, which makes this study a timely, novel and important contribution. The study is based on a novel data set collected by the authors from 88 bird species (67 in captivity, 21 in the wild) of 22 orders, except for 13 species, for which data were collected from a database of veterinary and animal care records of zoo animals (ZIMS). This novel data set itself greatly contributes to the pool of available data on avian glycemia, as previous comparative studies either extracted data from various studies or a ZIMS database (therefore potentially containing much more noise due to different methodologies or other unstandardised factors), or only collected data from a single order, namely Passeriformes. The data further represents the first comparative avian data set on albumin glycation obtained using a unified methodology. The authors used LC-MS to determine glycation levels, which does not have problems with specificity and sensitivity that may occur with assays used in previous studies. The data analysis is thorough, and the conclusions are substantiated. Overall, this is an important study representing a substantial contribution to the emerging field evolutionary physiology focused on ecology and evolution of blood/plasma glucose levels and resistance to glycation.

Weaknesses

Unfortunately, the authors did not record handling time (i.e., time elapsed between capture and blood sampling), which may be an important source of noise because handling-stress-induced increase in blood glucose has previously been reported. Moreover, the authors themselves demonstrate that handling stress increases variance in blood glucose levels. Both effects (elevated mean and variance) are evident in Figure ESM1.2. However, this likely makes their significant findings regarding glucose levels and their associations with lifespan or glycation rate more conservative, as highlighted by the authors.

Recommendations for the authors:

Reviewer #2 (Recommendations for the authors):

I understand that your main objective regarding glycation rate and lifespan, was to analyse the species resistance to glycation with respect to lifespan, while factoring out the species-specific variation in blood glucose level. However, I still believe that the absolute glycation level (i.e., not controlled for blood glucose level) may also be important for the evolution of lifespan. Given that blood glucose is positively related to both glycation and lifespan (although with a plateau in the latter case), lifespan could possibly be positively correlated with absolute glycation levels. If significant, that would be an interesting and counterintuitive finding, which would call for an explanation, thereby potentially stimulating further research. If not significant, it would show that long-lived species do not have higher glycation levels, despite having higher blood

glucose levels, thereby strengthening your argument about higher resistance of longlived species to glycation. So, in my opinion, the inclusion of an additional model of glycation level on life-history traits, without controlling for blood glucose, is worth considering.

We include now this model as supplementary material, indicating it in several parts of the text, including some of these issues we discussed here.

Lines 230-231: Please, provide a citation for these GVIF thresholds

We include it now.

Figure 3: I think that showing both glucose and glycation rate on the linear scale, rather than log scale, would better illustrate your conclusion - the slowing rise of glycation rate with increasing glucose levels.

That is a good point, although it may also be confusing for readers to see a graph that represents the data in a different way as the models. Maybe showing both graphs (as 3.A and 3.B) can solve it?

Figure 4. I recommend stating in the caption that the whiskers do not represent interquartile ranges (a standard option in box plots) but credible intervals as mentioned in the current version of the public author response.

Sorry about that, it was missed. Now it is included. Nevertheless, interquartile ranges from the posterior distributions can still be observed here represented with the boxes. Then the whiskers are the credible intervals.

<https://doi.org/10.7554/eLife.103205.3.sa0>