

Reviewed Preprint

v1 • November 27, 2024

Not revised

For correspondence:

james.marks@vanderbilt.edu

markus.hafner@nih.gov

* Present address: Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, United States

** Present address: Department of Biological Anthropology, University of California, Los Angeles, United States

Reviewing editor: Daniel Arango, Northwestern University, United States

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0 public domain dedication](https://creativecommons.org/licenses/by/4.0/).

Determining the off-target activity of antibiotics and novel translation initiation sites in mitochondria

James Marks*, , Emma Young**, Markus Hafner 

RNA Molecular Biology Laboratory, National Institute for Arthritis and Musculoskeletal and Skin Diseases, NIH, Bethesda, United States

eLife Assessment

This **useful** study presents findings on how some antibiotics, which inhibit protein synthesis in bacteria, affect the translation in mitochondrial ribosomes. The authors provide **solid** evidence that most tested antibiotics act similarly on bacterial and mitochondrial translation. Additionally, this work shows that alternative translation initiation events might exist in two specific mt-mRNAs (MT-ND1 and MT-ND5). The conclusions of this manuscript are of broad interest to the antibiotic and the mitochondrial fields.

<https://doi.org/10.7554/eLife.103699.1.sa3>

Abstract

Mitochondria translate the 13 mitochondrially encoded proteins using a dedicated translation system that closely resembles the one found in prokaryotes. Consequently, many bacterial protein synthesis inhibitors, used as antibiotics, exhibit mitochondrial toxicity as off-target effects. However, whether these antibiotics act through the same mechanisms in mitochondria as in bacteria remains unclear. To address this, we characterized the impact of a panel of bacterial translation and elongation inhibitors on mitochondrial translation through mitoribosome profiling. We found that the mechanism of action for every antibiotic, except telithromycin, is the same in both bacteria and mitochondria. Additionally, mitoribosome profiling data showed that *MT-ND1* and *MT-ND5* have incorrectly annotated translation initiation sites and suggested several novel mitochondrial translation events. Careful study of the mechanisms of mitochondrial translation inhibition can guide development of antibiotics with increased target specificity and reduced mitochondrial toxicity.

Introduction

Mitochondria contain their own genome, which in mammals canonically encodes thirteen proteins. These proteins are essential building blocks of complexes I, III, and IV of the electron transport chain, as well as ATP synthetase. They are synthesized at the mitochondrial matrix by a dedicated translation system that is conserved from the ancestral bacteria that evolved into the present-day organelles through an endosymbiotic event¹. As a result, mitochondrial translation is wholly distinct from cytoplasmic translation and utilizes its own mitochondrial ribosomes (mitoribosomes), tRNAs, and translation factors. The RNA components of the machinery are mitochondrially encoded, while its protein components are encoded in the nucleus, translated by the cytoplasmic ribosome, and subsequently imported into the mitochondria².

Chronic impairment of mitochondrial translation, most commonly through mutations in translation factors, results in disrupted mitochondrial function and causes diseases such as MERRF (Myoclonic Epilepsy with Ragged-Red Fibers)³, MELAS (Mitochondrial Encephalopathy, Lactic Acidosis, and Stroke-like episodes)⁴, and maternally inherited deafness^{5,6}. Furthermore,

acute inhibition of mitochondrial translation caused by the off-target action of antibiotics targeting the bacterial ribosome can also cause mitochondrial toxicity^{7–10}. This toxicity can manifest as lactic acidosis, ototoxicity¹¹, nephrotoxicity¹², and peripheral neuropathy¹³, depending on the specific antibiotic and the genetic background of the recipient¹⁴. While the United States has worked to regulate antibiotics with severe side effects, they remain a global issue. For example, chloramphenicol, whose use is discontinued in the United States, causes fatal aplastic anemia at a rate of 1 in 60,000⁷ but remains widely prescribed in low-income countries due to its low production cost^{15,16}. Surveys suggest that approximately 14.3 doses of antibiotics are administered per 1,000 people every day worldwide¹⁷. The toxic effects of antibiotics can be exacerbated when taken without medical supervision. In addition, safety concerns surrounding drugs like chloramphenicol and telithromycin, which is linked to liver failure¹⁸, limits our pool of available antibiotics in the face of a rise of multidrug-resistant bacteria. Studying the off-target effect of these drugs can guide the development of safer therapies.

Multiple classes of antibiotics inhibit bacterial translation by binding functionally important sites on the ribosome and disrupting a specific process in translation, resulting in arrested translation on mRNAs at sites characteristic of the mechanism of inhibition^{19,20}. In some instances, this relationship is straightforward. For example, the pleuromutilin antibiotic retapamulin binds to the peptidyl-transferase center (PTC) of the bacterial ribosome and impedes translation initiation by interfering with positioning of initiator tRNA²¹, resulting in bacterial ribosomes accumulating at start codons²². Inhibition by other antibiotics is more complicated, only disrupting translation when specific sequences are being translated, called context-dependent translation arrest. This is exemplified by chloramphenicol, which simultaneously interacts with the nascent peptide and the peptidyl-transferase center at its binding site and disrupts accommodation of aminoacyl-tRNA, thereby acting as an elongation inhibitor²³. Due to chloramphenicol's interaction with the nascent peptide, it selectively causes ribosomes to arrest translation when an alanine, serine, or threonine is in the –1 (penultimate) position of the nascent peptide²⁴. Additionally, chloramphenicol does not interfere with the accommodation of glycine due to its minimal size. The mechanisms of these and other antibiotics have been determined by ribosome profiling in bacterial cells. However, it remains unclear whether chloramphenicol or other antibiotics inhibit mitochondrial translation by identical mechanisms.

To determine the sites of translation arrest with, we utilized mitoribosome profiling in HEK293 cells treated with a diverse panel of antibiotics^{18,25}. Ribosome profiling allows for the mapping of translating ribosomes with codon-level accuracy by identifying mRNA sequences occupied by translating ribosomes. To do this, RNase treatment degrades mRNA unprotected by the ribosome to generate ribosome-protected fragments (RPFs). Because cytoplasmic ribosomes significantly outnumber mitoribosomes, mitoribosomes containing the mitochondrial RPFs (MRPFs) are isolated through sucrose gradient fractionation, where the cytoplasmic ribosome and mitoribosome sediment in the 80S and 55S fractions, respectively (Figure 1A [↗](#)). RPFs are isolated and quantified by next-generation sequencing. Increases in MRPFs reveal the specific codons and sequence contexts prevalent at sites at which mitochondrial translation arrests due to antibiotic treatment.

Understanding the mechanisms of translation inhibitors in mitochondria can guide the development of therapeutics that are more selective to the bacterial ribosome and lays the groundwork for determining how these therapeutics disrupt mitochondrial function. In this study, we utilized mitoribosome profiling in the presence of the bacterial protein synthesis inhibitors retapamulin, tiamulin, josamycin, erythromycin, chloramphenicol, and linezolid to determine their mechanism of action. We found that the mechanism of mitochondrial inhibition is the same as bacterial inhibition for all inhibitors except telithromycin. While telithromycin selectively arrests bacterial ribosomes when either arginine (R) or lysine (K) is in the –1 position of the nascent chain and A-site, called an R/K-X-R/K motif²⁶, it instead arrests mitochondrial translation at an R/K/A-X-K motif, suggesting an underlying difference in the mechanism in the two systems. Additionally, nucleotide-level precision of mitoribosome profiling also allowed us to precisely annotate the mitochondrial gene structure. We found that *MT-ND1* and *MT-ND5* genes use




Figure 1. Mitoribosome profiling captures elongating ribosomes.

(A) Schematic representation of mitoribosome footprinting and separation from cytoplasmic ribosomes by differential centrifugation. (B) Coverage of MRPFs on *MT-CYB* from mitoribosome profiling generated by either RNase I or MNase footprinting. (C) Distribution of read lengths of MRPFs mapping to the mitochondrial genome. (D) Read phasing analysis of MRPFs, where the 5' ends of reads are grouped by their subcodon position. Each dot represents an individual mitochondrial gene. (E) Average coverage of mitochondrial genes, calculated by dividing the total number of reads mapping to an ORF and dividing by the length of the ORF. (F) Distribution of mitochondrial codons grouped by the number of mitoribosome P-sites assigned to them.

alternative translation initiation sites and identified a putative open reading frame on the *MT-RNR1* gene. This work demonstrates that mitoribosome profiling is an effective tool to study the fundamentals of mitochondrial translation and the mechanisms of translation impairment affecting human health.

Results

RNase I generates highly phased mitoribosome footprints

To optimize mitoribosome profiling method, we tested two different RNases typically used for ribosome footprinting, RNase I and MNase, followed by the isolation of the 55S mitoribosome by sucrose gradient fractionation. We mapped the sequence reads to the mitochondrial genome and its thirteen open reading frames (ORFs). For each mapped sequence read, we inferred the position of the ribosomal P-site using a 13-nucleotide (nt) offset from the 5' end, determined by the analysis of footprint distribution surrounding the initiation sites of *MT-ND4*, *MT-ND6*, and the newly reported *MT-ND5-dORF*. We observed coverage across mitochondrial ORFs, suggesting we captured translating mitoribosomes (Figure 1B). Furthermore, analysis of MRPF lengths revealed a distinct difference between RNase I and MNase digested footprint. While both libraries are enriched for reads ~30 nt in length, MNase footprints showed a bimodal distribution, with a clear secondary peak at ~22 nts (Figure 1C), which may indicate that MNase-treatment may have captured two distinct conformations of the ribosome. To determine the quality of the ribosome profiling data, we calculated read phasing, which occurs when a nuclease creates uniform MRPFs and the resulting ends of the MRPFs reflect the 3-nt periodicity of elongating ribosomes. While both datasets do show clear phasing, the read phasing is more biased toward position 1 of the codon in the RNase I-treated sample, indicating slightly more consistent digestion (Figure 1D). Nevertheless, we observed similar levels of footprint abundance per gene, with the only notable exceptions being that RNase I-digested footprints were enriched on *MT-ND3* and depleted on *MT-ND4L* and *MT-ND6* (Figure 1E). While the cause of the difference is unclear, we did not expect it to interfere when comparing an antibiotic treated sample to a control. Read coverage also appeared similar across the two methods, with 85%-87% codons having between 2-50 reads mapping to them (Figure 1F). Overall, these results indicated that both MNase and RNase I digestion capture mitoribosome footprints with only minor differences. Considering the improved read phasing, we selected RNase I for our subsequent mitoribosome profiling experiments.

Retapamulin, tiamulin, and josamycin inhibit translation initiation

First, we examined the impact of three antibiotics – retapamulin, josamycin, and tiamulin on inhibiting translation initiation on mitochondrial translation. Retapamulin and tiamulin are members of the pleuromutilin class of antibiotics. They work by binding to the peptidyl transferase center (PTC) of the bacterial ribosome and disrupting initiator tRNA and A site tRNA positioning (Figure 2A, left)^{21,22,27–29}. Josamycin is a macrolide antibiotic that binds in the nascent peptide exit tunnel near the PTC and arrests bacterial ribosomes at initiation by interfering with A-site tRNA positioning (Figure 2A, right)³⁰. We sought to understand whether these antibiotics also target translation initiation in mitochondria. To this end, we treated HEK293 cells with high concentrations of these antibiotics (100 µg/ml of tiamulin and josamycin, 10 µg/mL of retapamulin) for 30 minutes, performed mitoribosome profiling, and compared the treated experiments to a DMSO treated control.

We mapped MRPFs to the 13 mitochondrial genes. On *MT-CYB* and all other genes, we observed that after treatment with antibiotics reads accumulated at the 5' end of the transcript and MRPFs across the rest of the coding region were lost or decreased (Figure 2B). We quantified the percentage of MRPFs mapping at the 5' end relative to the total ORF, excluding polycistronic mRNAs. Treatment with translation initiation inhibitors increased the percentage of those reads from 2.4 to 5.2-fold relative to the DMSO-treated cells (Figure 2C). Additionally, we quantified the cumulative footprint abundance moving from the start to the stop codon (Figure 2D). While




Figure 2. Inhibition of translation initiation in mitochondria.

(A) (Left) Schematic representation of retapamulin (orange) or tiamulin (yellow) binding to the PTC and disrupting the first round of elongation. (Right) Josamycin (blue) disrupts accommodation of the first aminoacyl-tRNA. All three inhibit initiation in bacteria. (B) MRP coverage of *MT-CYB* treated with either retapamulin, tiamulin (note y-axis is segmented), or josamycin. (C) Average fold change increase in the percentage of reads that map to the 5' of mitochondrial transcripts. (D) Cumulative distribution of ribosome density normalized across all mitochondrial ORFs.

ribosome footprints are generally evenly spread across the open reading frame in the DMSO control, MRPFs from cells treated with the initiation inhibitors showed a sharp increase in the percentage of reads found at the 5' end. Together, these results indicate that retapamulin, tiamulin, and josamycin inhibit mitochondrial translation initiation, likely by the same mechanism as in prokaryotes.

Chloramphenicol, linezolid, and telithromycin arrest mitochondrial translation in a context-dependent manner

Chloramphenicol, linezolid, and telithromycin selectively inhibit bacterial translation when ribosomes are translating specific sequences, in a process called context-dependent translation arrest^{24,26}. Linezolid, like chloramphenicol, induces context-dependent arrest when alanine, serine, or threonine are at penultimate position of the nascent peptide (Figure 3A²⁴, left). Unlike chloramphenicol and linezolid, telithromycin binds in the nascent peptide exit tunnel near the PTC and adjacent to the nascent peptide³¹. During elongation, telithromycin causes ribosomes to arrest translation at a R/K-X-R/K motif, when the identity of both the penultimate position of the nascent peptide and the incoming aminoacyl-moiety are either arginine or lysine (Figure 3A²⁴, right).

We carried out mitoribosome profiling on HEK293 cells treated with either antibiotic to understand whether these context-dependent translation inhibitors arrest mitoribosomes in the same manner as they arrest bacterial ribosomes. Unlike the cells treated with initiation inhibitors, cells treated with chloramphenicol or linezolid carried ribosome density throughout the entirety of the ORF, suggesting that, as expected, they were not acting as inhibitors of initiation (Figure 3B²⁴). Cells treated with telithromycin did show a slight increase in MRPFs abundance at the 5' end on *MT-CYB*. However, across all genes the levels of MRPFs at the 5' end were not near the levels seen when using the initiation inhibitor retapamulin and were more similar to chloramphenicol, linezolid, and the DMSO control (Figure 3C²⁴). Cumulative distribution of MRPFs suggested that ribosome density was biased towards the 5' end of the gene for chloramphenicol and telithromycin, but not significantly for linezolid (Figure 3D²⁴). These results suggest that these three antibiotics are not acting as initiation inhibitors and that linezolid and telithromycin redistribute MRPFs to the 5' end of the ORF.

Next, we determined whether this observed redistribution of elongating ribosomes was dependent on the context of the sequence being translated. We calculated the average change of MRPFs depending on the identity of the amino acid at either the -1, P-site, or A-site positions. For both chloramphenicol and linezolid, we observed a specific increase in ribosome footprints when mitoribosomes encountered codons for alanine, serine, or threonine in the -1 position, with a 2-, 2.2-, and 1.3-fold change for chloramphenicol and a 2.3-, 1.4-, and 1.6-fold change for linezolid, respectively (Figure 3E²⁴). Additionally, we also observed context-dependent translation arrest induced by telithromycin. Unlike the R/K-X-R/K motif observed in bacteria, we found that telithromycin caused mitoribosomes to arrest at sites with a lysine positioned in the A-site, with an average 2.5-fold increase in MRPFs around those sites, and that neither residue was enriched in the -1 position. Due to the limited size of the mitochondrial genome, there are only nine instances of the R/K-X-R/K motif, nevertheless, when we observed the positions with the greatest fold change, 3 of the top 4 contained the motif with lysine in the A-site (Figure 3F²⁴), suggesting that the motif was at least partially recapitulated. To specifically look for the motif we calculated the average fold change when we fixed arginine or lysine in the A-site and varied the identity of the -1 amino acid (Figure 3G²⁴). We found that when lysine was in the A-site while either arginine, lysine, or alanine was -1 position, there was strong arrest (> 5-fold change). However, when arginine was in the A-site we did not observe a strong stalling event with any -1 position residue. Therefore, in mitochondria, telithromycin induces stalling at R/K/A-X-K motifs. We concluded that the mechanism underlying context-dependent translation arrest for telithromycin is different in mitochondria compared to bacteria.




Figure 3. Context-dependent translation arrest by elongation inhibitors.

(A) Schematic representation of chloramphenicol (purple), linezolid (cyan), and telithromycin (green) inducing context-dependent translation arrest. (B) MRPF coverage of *MT-CYB* from cells treated with either chloramphenicol, linezolid, or telithromycin. (C) Average fold change increase in the percentage of reads that map to the 5' of mitochondrial transcripts. (D) Cumulative distribution of ribosome density normalized across all mitochondrial ORFs. (E) Volcano plots of changes in MRPF fold change from cells treated with translation inhibitors. Each dot is the mean $\log_2$ -fold change grouped by the identity of the -1 amino acid (top), P-site amino acid (middle), or A-site amino acid (right). (F) Top 4 strongest stalling sites induced by telithromycin treatment. Red amino acids indicate a classical R/K-X-R/K stalling motif. (G) Fold change in ribosome density, calculated as $2^{\text{mean}(\log_2(\text{Telithromycin}/\text{Control}))}$. Bars differ in the identity of the P-site amino acid and contain either a lysine in the A-site (top) or arginine (bottom).

Translation initiation on *MT-ND1* and *MT-ND5* occurs at an alternative start codon

During the process of assigning the P-site positions of MRPFs, we observed that no ribosomes were assigned in the first 13 nucleotides of the 5' end of the transcript. This is not surprising, considering that mitochondrial transcripts often have no or very short 5' untranslated regions - typically less than 4 nts long - and we applied a 13-nt offset from the 5' end of MRPFs. It reasons that as mitoribosomes begin translation on a leaderless mRNA, the 5' end of the resulting MRPF would have a full-length region 3' of the P-site and no sequence 5' of the P-site. As the ribosomes move down the transcript, the 5' end of the MRPF would increase until it is accessible for digestion, at which point it would generate 31 nt-long MRPFs (Figure 4A [↗](#)). As accurate mapping of mitoribosomes at initiation could be a useful tool to understand mitochondrial translation, we attempted to map the location of initiating mitoribosomes at 5' end using footprint length, rather than the 5' offset. Because mitoribosome profiling with MNase generated a population of ribosomes with short MRPFs, we hypothesized that these footprints could be indicative of capturing initiating ribosomes (Figure 1C [↗](#)). We plotted the distribution of MRPFs from 20 to 34 nt length that mapped to the 5' end of the 11 leaderless or short leader ORFs and compared them to reads mapping to the whole ORF. We observed that MRPFs 22-23 nts in length with 6-fold more abundant and MRPFs >28 nts in length were depleted compared to the whole ORF (Figure 4B [↗](#)). As mitochondrial ORFs vary in the length of their 5' UTR and we expect most ribosomes to be positioned at the start codon, we tested whether leaderless ORFs were enriched in shorter reads compared to ORFs preceded by a short leader. Leaderless ORFs, such as *MT-CO2*, *MT-CO3*, *MT-ND4L* contained MRPFs mainly 20-23 nucleotides in length at the 5' end (Figure 4C [↗](#), top). *MT-ATP8*, which contains a start codon beginning at the third nt, contained footprints that were 22-24 nt long and *MT-CO1*, with a start codon at the fourth nt, contained MRPFs 23-24 nt long (Figure 4C [↗](#), middle). Not only did we observe longer MRPFs in short leader ORFs, the size of the MRPFs also appeared proportional to the location of the start codon. Together, these results indicate that the size of the footprint is directly related to the start codon position at the 5' end of mitochondrial mRNAs.

Two genes, *MT-ND1* and *MT-ND5*, deviated from this trend. The annotated start site for *MT-ND5* is a leaderless AUA codon (a start codon in mitochondria) but we observed few 20-22 nt MRPFs (Figure 4C [↗](#), bottom). Similarly, the annotated start site for *MT-ND1* is also an AUA codon that begins at the third nt and it is depleted in reads for MRPFs that are smaller than 25 nts. Instead, MRPFs derived from the 5' end of both genes are enriched in longer reads: 27 nts-long for *MT-ND5* and 28-29 nts-long for *MT-ND1*. Curiously, the third codon for both genes is an in-frame AUG and the read lengths proportionally correspond to mitoribosomes positioned at these start codons, suggesting mitoribosomes initiating translation at these sites. MRPFs generated from cells treated with initiation inhibitors and footprinted by MNase yield similarly sized reads at the 5' end of *MT-ND1* and *MT-ND5* (Figure S1 [↗](#)). Using protein mass spectrometry, we tested whether peptides matching the N-terminus of these proteins would correspond to initiation at the canonical start codons or alternative start codons. We identified a peptide that mapped specifically to the truncated version of *MT-ND1* and not the full-length peptide, indicating that translation can initiate from the alternative start codon (Figure 4D [↗](#)). While we did not identify N-terminal peptides mapping to *MT-ND5* from our mass spectrometry data, the protein mass spectrometry database MassIVE³⁰ covered the N-terminal region of *MT-ND5* (Figure 4E [↗](#)). Consistent with our mitoribosome profiling data, the majority of N-terminal fragments did not include the amino acids from the annotated start or second codon but began with methionine at codon 3. We also observed that MRPFs on *MT-ND6*, which initiates at the first of two AUGs codon immediately at the 5' end, correlated to ribosomes positioned at both AUG codons, indicating that this gene may initiate translation from the second AUG codon as well. However, we did not observe any peptides mapping to the N-terminus of *MT-ND6*. Together, this data shows that *MT-ND1* and *MT-ND5* may not initiate from their canonical AUA codons but may use alternative in-frame AUG codons. Additionally, our findings indicate that the mapping of mitochondrial translation initiation events by size, rather than 5' offset is an accurate strategy.




Figure 4. MRP sizes indicate use of alternative initiation sites.

(A) Schematic representation of changing MRP sizes as a function of the mitoribosome's position near the 3' end of mRNAs. (B) Read length measurement of reads mapping to the 5' ends of mitochondrial transcripts (red) compared to the entire ORF (black). (C) Read lengths of MRPFs mapping to the 5' ends of mitochondrial transcripts. Each row contains the nucleotide sequence of each transcript starting from the 5' end. The coloring of the box is proportional abundance of MRPFs with read lengths from 20-34 nts. Nucleotide sequence and read abundance are overlaid to compare the abundance of reads to potential ribosome start sites. Transcripts are grouped by being leaderless, having a leader, or having a potential alternative start site. Black boxes indicate annotated start codons and green boxes indicate alternative initiation sites. (D) Visual representation of the peptide coverage (green bars or highlights) from mass spectrometry data covering the N-terminus of MT-ND1 or (E) MT-ND5.

Mitoribosome profiling reveals novel translation events

During our analysis of 5' end mapping, we did not remove reads mapping to non-coding RNAs (ncRNAs) before mapping to the mitochondrial genome and performing P-site assignment. Consequently, we observed that some tRNAs and rRNAs had an abundance of reads mapping to 13 nts away from their 5' ends, similarly to what we observe on mRNAs for translation initiation. In addition, some of these ncRNAs contained a potential start codon near the 5' end. These reads could either be products of random digestion of the abundant background of ncRNAs or be genuine MRPFs. We hypothesized that if these reads were the result of mitoribosomes initiating or attempting to initiate translation at the 5' end of the ncRNAs, then we could use the same analysis of read length distribution at 5' ends of mRNAs we performed above and determine of the sizes correlated to an AUG, AUU, GUG, or UUG as potential start codons. While GUG and UUG are not used in canonical mitochondrial initiation they were included in the search because noncanonical initiation has been observed at GUG³² and both are frequently utilized in bacterial initiation. We identified sites on 4 tRNAs that contained MRPFs with a size that correlated to either AUG or UUG codons (Figure 5A [↗](#)). The predicted ORFs and thus resulting potential translation events varied widely. On *MT-TF*, the start codon is followed by an immediate stop codon. The ORF on *MT-TI* is 3 codons long and terminates with the non-canonical mitochondrial termination codon AGA. On *MT-TY*, a 7-codon ORF is terminated by a UAA codon. *MT-TS1* contains the largest open reading frame identified on tRNAs, at 20 codons and it terminates with at the U AGA sequence, as used in *MT-CO1*³³. Additionally, we identified a putative start site at the 5' end of *MT-RNR1* (Figure 5A [↗](#)) and mitoribosome profiling of cells treated with initiation inhibitors and digested with MNase show similar footprint sizes (Figure S1 [↗](#)). However, this site did not occur at a canonical AUG or AUU codon, but instead at a UUG codon. This putative start codon is followed by a 58-codon ORF and terminates at a UAG stop codon.

To determine whether translation elongation occurs at these putative ORFs, we determined the read phasing characteristic for MRPFs derived from translating ribosomes at these ORFs. However, the sites on *MT-TF*, *MT-TI*, and *MT-TY* are short and close to the 5' end, hindering phasing analysis and thus do not allow us to conclude whether our sequence reads are indicative of translation. While the ORF on *MT-TS1* is longer, MRPF density was low and we did not observe read phasing and thus is likely not translated (not shown). The reads mapping to the ORF on *MT-RNR1* had a phasing profile similar to those mapping to the 13 canonical mitochondrial genes (Figure 5B [↗](#)). In contrast, reads mapping to the remainder of *MT-RNR1* or mapping to the region of *MT-RNR2* that codes for the putative mitochondrial micropeptide Humanin, shows no or conflicting read phasing. When compared to the amino acid composition of the mitochondrially-encoded proteome, the amino acid composition of the resulting peptide shows a stronger positive correlation ($R^2=0.534$) (Figure 5C [↗](#)) to the remaining non-coding portion of *MT-RNR1* ($R^2=0.241$) (Figure 5D [↗](#)). Although mitoribosome profiling suggests that this putative ORF is translated, we did not detect any corresponding peptides in the protein mass spectrometry of isolated mitochondria. Nevertheless, our mass spectrometry dataset, or the MASSIVE database do not cover many regions of the mitochondrial proteome and thus, the absence of a spectrum is not necessarily evidence that the peptide is not translated. The read phasing analysis suggests this region of *MT-RNR1* is translated and could encode a 15th mitochondrial protein. Overall, this data shows that our mitoribosome profiling can be used as a tool to study non-canonical translation events resulting from ncRNAs.

Discussion

In this work, we sought to understand the mechanisms of bacterial protein synthesis inhibitors on their primary off-target, the mitoribosome. We found that inhibitors of bacterial translation initiation, retapamulin, tiamulin, and josamycin, also act as initiation inhibitors in mitochondria. Likewise, chloramphenicol and linezolid selectively inhibit elongation when the mitoribosome carries either an alanine, serine, or threonine in the penultimate position of the nascent peptide,




Figure 5. MRPFs on ncRNAs indicate potential novel ORFs

(A) Read lengths of MRPFs mapping to the 5' end of ncRNAs. Each row contains the nucleotide sequence beginning from the 5' end of listed ncRNAs. The coloring of the box is proportional the abundance of MRPFs with read lengths 20–34 nts. The nucleotide sequence and read MRPF abundance are overlaid to identify potential new start sites. Putative start sites are indicated by green boxes. (B) Read phasing analysis of all canonical ORFs (grey), a mutative ORF located on *MT-RNR1*, the non-coding region of *MT-RNR1*, and Humanin. (C) A correlation plot comparing the frequency of amino acid usage in canonical ORFs compared to the putative ORF on *MT-RNR1* or (D) the non-coding region of *MT-RNR1*.

matching their mechanism in bacteria. In contrast, the context-dependence of telithromycin differed when arresting mitoribosomes, wherein it preferentially arrested translation at R/K/A-X-K motif.

Context-dependent translation arrest by chloramphenicol and linezolid suggests similarities between binding of the bacterial ribosome and the mitoribosome. Structural and biochemical studies have demonstrated that binding of chloramphenicol and linezolid is stabilized by alanine, serine, or threonine in the penultimate position of nascent peptide^{34,35}. Our findings suggest a similar or identical interaction is occurring in the mitoribosome. In contrast, these inhibitors are unable to inhibit peptide bond formation in bacteria when glycine is positioned in the A-site, due to the minimal size of the side chain, allowing the inhibitor and the aminoacyl-moiety to coexist in the PTC. We did not observe a lack of mitoribosome translation arrest when glycine was positioned in the A-site. This may have technical reasons, e.g. it may be caused by an insufficient depth in sequencing, or indicate that the relatively small mitochondrial genome limits the variety of stalling and non-stalling events we can observe. But more likely it shows that chloramphenicol and linezolid are capable of arresting translation when glycine is in the A-site. Recently, mitoribosome profiling analysis showed that chloramphenicol added during lysis did not induce context-dependent translation arrest³⁶. This suggests that during lysis, there is insufficient mitochondrial translation to induce a redistribution of mitoribosomes rather than an inability of the chloramphenicol to induce context-dependent arrest.

During preparation of this manuscript, two other groups studied the effect of translation inhibitors on mitochondrial translation. One group's mitoribosome profiling in the presence of chloramphenicol and linezolid agreed with our observed context-dependent translation arrest³⁷. Their structure of linezolid in the mitochondrial peptidyl-transferase center also suggested some nuanced differences in binding between the bacterial and the mitochondrial ribosome. Another group carried out mitoribosome profiling in the presence of retapamulin³⁶. However, they observed additional ribosome density throughout the coding region and potentially some associated with putative start codons. This differs significantly from our observed accumulation at only annotated start sites and may be due to differences in mitoribosome profiling methods. Future work may reveal how each strategy contributes to our understanding of mitochondrial translation.

Inhibition of mitochondrial translation through either small molecules or mutations localized in mitochondrial translation factors can induce a wide range of pathologies^{38–40}. However, it is unclear how or whether the relationships between the mechanism of translation inhibition relates to the resulting disease. Recently the proteins mtRF-R and MTRES1 were shown to act as quality control factors, rescuing stalled mitoribosomes by removing the nascent peptide and peptidyl-tRNA⁴¹. As the mitoribosome can arrest translation through a variety of mechanisms and small molecules, it remains unclear whether these factors can rescue all stalled mitoribosomes. Further work to dissect the relationship between mechanisms of inhibition, activation of rescue factors, and stress response can complement our understanding of pathologies resulting from mitoribosomopathies and antibiotic treatment.

In this study, we also reported that the translation start codon of two mitochondrial genes, *MT-ND1* and *MT-ND5*, do not use the annotated AUA codon, but instead begin translation from a nearby in-frame AUG codon. Others have noted that haplogroups L1b and F2 contain a single nucleotide polymorphism at either position T3308C or T12338C respectively, which disrupts the annotated AUA start codons of *MT-ND1* and *MT-ND5* yet do not result in any discernable pathogenicity^{42–44}. Authors noted that the same AUG codons we identified as start codons could be used in lieu of the canonical AUA start codons. Because we observe MRPF lengths primarily correlating with the AUG codons, we suggest that these downstream start codons are the primary start site for these genes, though this does not preclude the use of both sites. Given the limited examples of translation initiation sites inherent to the mitochondrial genome, the reannotation of 2 sites is a significant revision and can guide future work to understand how start codon selection occurs in mitochondria.

Recently, mitoribosome profiling was used to identify an open reading frame in region immediately 3' of the gene *MT-ND5*, which we confirmed observing in our profiling experiments⁴⁵. However, this uses an internal initiation mechanism for polycistronic genes, similar to *MT-ND4* and *MT-ATP6*. Until now, accurate mapping of mitochondrial translation events at the 5' ends of transcripts has been unavailable. We identified several putative translation initiation events occurring on mitochondrial MT-tRNAs and *MT-RNR1*, based on the size of the reads occurring at the 5' end of the RNA and their relationship to a start codon localized in the sequence. The site of *MT-RNR1* was further supported by analysis of the resulting ORF, suggesting translation initiation and elongation occurs. Curiously, tRNA^{Tyr}, encoded by *MT-TY*, is polyadenylated during processing, which may be related to our identification of a ribosome binding event on this tRNA⁴⁶. Due to both the short ORFs, minimal read coverage, and lack of a detectable peptide we could not determine if translation elongation occurred on the mitochondrial tRNAs. Future work is needed to understand whether these are actual translation events and to determine the function of these potential novel peptides. Together our findings improve our understanding of mitochondrial translation initiation, gene expression, and the mechanisms of translation inhibition.

Materials and methods

Isolation of mitoribosomes for profiling and western blotting

Mitoribosome profiling was carried out in HEK293 cells. 15 cm plates were treated with either DMSO as a control or 100 µg/mL of protein synthesis inhibitor for 1 hour, washed with 4°C PBS, and flash frozen by floating the plate on liquid nitrogen. Cells were collected by scrapping on ice with 600 µL of lysis buffer (1.5x Lysis buffer (30 mM Tris-HCl pH 7.8, 150 mM KCl, 15 mM MgCl₂, 1.5 mM DTT, 1.5% Triton X-100, 0.15% NP40, 1 × complete phosphatase and protease inhibitors)) and lysed by passing through a 27-G needle 10 times. Lysate was centrifuged for 10 minutes at 21,000 g at 4°C and the supernatant was collected. 300 µL of lysate was footprinted with either digested with 1,5000 U of MNase, 10 µL of SUPERase-In, and 5 mM of CaCl₂ at 22°C for 1 hour and stopped by adding EGTA to a concentration of 6 mM or with 1 U/µL of RNase I at 22°C for 1 hour and stopped by adding 10 µL of SUPERase-In. The digestion was clarified by centrifugation at 21,000 g for 5 minutes. Lysates were loaded onto a 5%-30% sucrose gradient (20 mM Tris-HCl pH 7.8, 100 mM KCl, 10mM MgCl₂, 1 mM DTT) and spun for 2.5 hours at 40,000 rpm in a SW40 rotor. 55S mitoribosome fractions were collected using a BioComp Fractionator and MRPFs were isolated by SDS/hot phenol/chloroform extraction.

Mitoribosome profiling library construction

Samples were run on a 15% TBE-Urea polyacrylamide gel and footprint sizes from 15-34 nts were excised and footprints were eluted in 0.3 M NaOAc, 1 mM EDTA, and 0.25% SDS followed by purification by Oligo Clean & Concentrator. Samples were treated with Quick CIP. 3' adapters (5'-rAppNN-6 nt UMI-TGGAATTCTCGGGTGCCAAGG-L) were ligated using Rnl2(1-249)K227Q ligase for 2 hours at 16°C and pooled. Pooled samples were treated with T4 PNK for 30 minutes at 37°C. A 5' chimeric DNA-RNA adapter (GTTCAGAGTTCTACAGTCCGACGATCrNrNrN) was ligated to the samples with T4 RNL1. Reverse transcription was carried out using SuperScript IV followed by PCR amplification by Taq DNA polymerase. Libraries were size selected using a 3% agarose Pippin prep and final libraries were PCR amplified. Samples were sequenced as either singe- or paired-end 50 nt reads on either a nextseq550 or novaseq.

Mitoribosome profiling analysis

Adapter sequences were removed with cutadapt and aligned with bowtie2 to a fasta containing rRNA and tRNA sequences. To identify stalling sites, reads were mapped to a custom mitochondrial transcriptome containing all 5' UTRs, coding sequences, and 3' UTRs for all processed mRNAs. Reads were aligned with bowtie2. A 13 nt P-site offset was assigned using Plastid and mapping reads relative to the start codons of *ND4*, *ATP6*, *MT-ND5-dORF*. We

determined the relative ribosome density by normalizing the number of reads at each codon by the total number of reads mapping to its respective gene for each sample. Fold change was then calculated by dividing the relative ribosome density in a treated sample to an untreated sample. To identify putative novel translation initiation sites we mapped our footprints to the entire mitochondrial genome and visually inspected the 3' ends of mitochondrial tRNAs and rRNAs.

Mass spectrometry analysis of mitochondrial proteins

Mitochondria were isolated from 2×10^7 HEK293 cells using the Mitochondria Isolation Kit for Cultured Cells. Mitochondrial proteins were processed for trypsin/LysC digestion and the digested peptide mixture was then concentrated and desalted using C18 column. Reconstituted desalted peptides in 25 μ l of 0.1% formic acid. 12 μ l of peptides from each sample was analyzed by 110 min LC/MS/MS run. The LC/MS/MS analysis of tryptic peptides for each sample was performed sequentially with a blank run between each two sample runs using a Thermo Scientific Orbitrap Exploris 240 Mass Spectrometer and a Thermo Dionex UltiMate 3000 RSLCnano System. Peptides from trypsin digestion were loaded onto a peptide trap cartridge at a flow rate of 5 μ l/min. The trapped peptides were eluted onto a reversed-phase Easy-Spray Column PepMap RSLC, C18, 2 μ m, 100A, 75 μ m $\times$ 250 mm (Thermo Scientific) using a linear gradient of acetonitrile (3–36%) in 0.1% formic acid. The elution duration was 60 min at a flow rate of 0.3 μ l/min. Eluted peptides from the Easy-Spray column were ionized and sprayed into the mass spectrometer, using a Nano Easy-Spray Ion Source (Thermo) under the following settings: spray voltage, 1.6 kV, Capillary temperature, 275°C. Other settings were empirically determined. Raw data files were searched against human protein sequences database and client target protein sequences database using the Proteome Discoverer 2.5 software (Thermo, San Jose, CA) based on the SEQUEST algorithm. Carbamidomethylation (+57.021 Da) of cysteines was set as fixed modification, and Oxidation / +15.995 Da (M), Phospho / +79.966 Da (S, T, Y), and Deamidated / +0.984 Da (N, Q) were set as dynamic modifications. The minimum peptide length was specified to be five amino acids. The precursor mass tolerance was set to 15 ppm, whereas fragment mass tolerance was set to 0.05 Da. The maximum false peptide discovery rate was specified as 0.01. The resulting Proteome Discoverer Report contains all assembled proteins with peptides sequences and peptide spectrum match counts (PSM#) and MS1 peak area relative abundance.

Data sharing

Mitoribosome fastq files, wig files mapping to mitochondrial mRNAs, and read length files for MNase treated samples mapping to chrM are available through GEO Series GSE277563.

Author contributions

J.M. Conceived the project, designed and executed experiments, analyzed data, and wrote the manuscript. E.Y. Analyzed mitoribosome profiling data. M.H. supervised the project, advised on data interpretation and experimental design, and edited the manuscript.

Acknowledgements

We'd like to thank Nicholas Guydosh (NIDDK), Sezen Meydan (Vanderbilt University), Astrid Haase (NIDDK), the Haase lab members, and the Hafner lab members for their feedback on this work. We thank Daniel Ben Halvey (Tel Aviv University) for his assistance in carrying out ribosome profiling and Sarah Young-Baird (Uniformed Services University) for assistance with sucrose gradient fractionation. We thank Stefania Dell'Orso and Faiza Naz (NIH/NIAMS Genomic Technology Section) for sequencing experiments.

Additional information

Author ORCID iDs

Markus Hafner: <http://orcid.org/0000-0002-4336-6518>




Figure S1. MRPf lengths after treatment with initiation inhibitors correlate to alternative initiation sites.

(A) Read lengths of MNase-generated MRPf from cells treated with josamycin or (B) retapamulin.

References

1. Gray M.W., Burger G., Lang B.F (1999) Mitochondrial evolution. *Science* **283**:1476-1481 <https://doi.org/10.1126/science.283.5407.1476>
2. D'Souza A.R., Minczuk M (2018) Mitochondrial transcription and translation: overview. *Essays Biochem* **62**:309-320 <https://doi.org/10.1042/EBC20170102>
3. Enriquez J.A., Chomyn A., Attardi G (1995) MtDNA mutation in MERRF syndrome causes defective aminoacylation of tRNA(Lys) and premature translation termination. *Nat Genet* **10**:47-55 <https://doi.org/10.1038/ng0595-47>
4. Webb B.D., Diaz G.A., Prasun P (2020) Mitochondrial translation defects and human disease. *J Transl Genet Genom* **4**:71-80 <https://doi.org/10.20517/jtgg.2020.11>
5. Prezant T.R., Agopian J.V., Bohlman M.C., Bu X., Oztas S., Qiu W.Q., Arnos K.S., Cortopassi G.A., Jaber L., Rotter J.I., *et al.* (1993) Mitochondrial ribosomal RNA mutation associated with both antibiotic-induced and non-syndromic deafness. *Nat Genet* **4**:289-294 <https://doi.org/10.1038/ng0793-289>
6. Zheng J., Ji Y., Guan M.X (2012) Mitochondrial tRNA mutations associated with deafness. *Mitochondrion* **12**:406-413 <https://doi.org/10.1016/j.mito.2012.04.001>
7. Smick K.M., Condit P.K., Proctor R.L., Sutcher V (1964) Fatal Aplastic Anemia. An Epidemiological Study of Its Relationship to the Drug Chloramphenicol. *J Chronic Dis* **17**:899-914 [https://doi.org/10.1016/0021-9681\(64\)90161-4](https://doi.org/10.1016/0021-9681(64)90161-4)
8. Woodhead J.L., Yang K., Oldach D., MacLauchlin C., Fernandes P., Watkins P.B., Siler S.Q., Howell B.A (2019) Analyzing the Mechanisms Behind Macrolide Antibiotic-Induced Liver Injury Using Quantitative Systems Toxicology Modeling. *Pharm Res* **36**:48 <https://doi.org/10.1007/s11095-019-2582-y>
9. De Vriese A.S., Coster R.V., Smet J., Seneca S., Lovering A., Van Haute L.L., Vanopdenbosch L.J., Martin J.J., Groote C.C., Vandecasteele S., *et al.* (2006) Linezolid-induced inhibition of mitochondrial protein synthesis. *Clin Infect Dis* **42**:1111-1117 <https://doi.org/10.1086/501356>
10. Gupta A., Okesli-Armlovich A., Morgens D., Bassik M.C., Khosla C (2020) A genome-wide analysis of targets of macrolide antibiotics in mammalian cells. *J Biol Chem* **295**:2057-2067 <https://doi.org/10.1074/jbc.RA119.010770>
11. Fu X., Wan P., Li P., Wang J., Guo S., Zhang Y., An Y., Ye C., Liu Z., Gao J., *et al.* (2021) Mechanism and Prevention of Ototoxicity Induced by Aminoglycosides. *Front Cell Neurosci* **15**:692762 <https://doi.org/10.3389/fncel.2021.692762>
12. Morales-Alvarez M.C (2020) Nephrotoxicity of Antimicrobials and Antibiotics. *Adv Chronic Kidney Dis* **27**:31-37 <https://doi.org/10.1053/j.ackd.2019.08.001>
13. Soriano A., Miro O., Mensa J (2005) Mitochondrial toxicity associated with linezolid. *N Engl J Med* **353**:2305-2306 <https://doi.org/10.1056/NEJM200511243532123>
14. Palenzuela L., Hahn N.M., Nelson R.P., Arno J.N., Schobert C., Bethel R., Ostrowski L.A., Sharma M.R., Datta P.P., Agrawal R.K., *et al.* (2005) Does linezolid cause lactic acidosis by inhibiting mitochondrial protein synthesis?. *Clin Infect Dis* **40**:e113-116 <https://doi.org/10.1086/430441>
15. Bale B.I., Elebesunu E.E., Manikavasagar P., Agwuna F.O., Ogunkola I.O., Sow A.U., Lucero-Prisno D.E (2023) Antibiotic resistance in ocular bacterial infections: an integrative review of ophthalmic chloramphenicol. *Trop Med Health* **51**:15 <https://doi.org/10.1186/s41182-023-00496-x>
16. Lorenzo D (2019) Chloramphenicol Resurrected: A Journey from Antibiotic Resistance in Eye Infections to Biofilm and Ocular Microbiota. *Microorganisms* **7** <https://doi.org/10.3390/microorganisms7090278>
17. Browne A.J., Chipeta M.G., Haines-Woodhouse G., Kumaran E.P.A., Hamadani B.H.K., Zarea S., Henry N.J., Deshpande A., Reiner R.C., Day N.P.J., *et al.* (2021) Global antibiotic consumption and usage in humans, 2000-18: a spatial modelling study. *Lancet Planet Health* **5**:e893-e904 [https://doi.org/10.1016/S2542-5196\(21\)00280-1](https://doi.org/10.1016/S2542-5196(21)00280-1)

18. Graham D.J (2006) Telithromycin and acute liver failure. *N Engl J Med* **355**:2260-2261 <https://doi.org/10.1056/NEJMc066372>
19. Paternoga H., Crowe-McAuliffe C., Bock L.V., Koller T.O., Morici M., Beckert B., Myasnikov A.G., Grubmuller H., Novacek J., Wilson D.N (2023) Structural conservation of antibiotic interaction with ribosomes. *Nat Struct Mol Biol* **30**:1380-1392 <https://doi.org/10.1038/s41594-023-01047-y>
20. Lin J., Zhou D., Steitz T.A., Polikanov Y.S., Gagnon M.G (2018) Ribosome-Targeting Antibiotics: Modes of Action, Mechanisms of Resistance, and Implications for Drug Design. *Annu Rev Biochem* **87**:451-478 <https://doi.org/10.1146/annurev-biochem-062917-011942>
21. Yan K., Madden L., Choudhry A.E., Voigt C.S., Copeland R.A., Gontarek R.R (2006) Biochemical characterization of the interactions of the novel pleuromutilin derivative retapamulin with bacterial ribosomes. *Antimicrob Agents Chemother* **50**:3875-3881 <https://doi.org/10.1128/AAC.00184-06>
22. Meydan S., Marks J., Klepacki D., Sharma V., Baranov P.V., Firth A.E., Margus T., Kefi A., Vazquez-Laslop N., Mankin A.S (2019) Retapamulin-Assisted Ribosome Profiling Reveals the Alternative Bacterial Proteome. *Mol Cell* **74**:481-493.e486. <https://doi.org/10.1016/j.molcel.2019.02.017>
23. Choi J., Marks J., Zhang J., Chen D.H., Wang J., Vazquez-Laslop N., Mankin A.S., Puglisi J.D (2020) Dynamics of the context-specific translation arrest by chloramphenicol and linezolid. *Nat Chem Biol* **16**:310-317 <https://doi.org/10.1038/s41589-019-0423-2>
24. Marks J., Kannan K., Roncase E.J., Klepacki D., Kefi A., Orelle C., Vazquez-Laslop N., Mankin A.S (2016) Context-specific inhibition of translation by ribosomal antibiotics targeting the peptidyl transferase center. *Proc Natl Acad Sci U S A* **113**:12150-12155 <https://doi.org/10.1073/pnas.1613055113>
25. Li S.H., Nofal M., Parsons L.R., Rabinowitz J.D., Gitai Z (2021) Monitoring mammalian mitochondrial translation with MitoRiboSeq. *Nat Protoc* **16**:2802-2825 <https://doi.org/10.1038/s41596-021-00517-1>
26. Sothiselvam S., Neuner S., Rigger L., Klepacki D., Micura R., Vazquez-Laslop N., Mankin A.S (2016) Binding of Macrolide Antibiotics Leads to Ribosomal Selection against Specific Substrates Based on Their Charge and Size. *Cell Rep* **16**:1789-1799 <https://doi.org/10.1016/j.celrep.2016.07.018>
27. Schlunzen F., Pyetan E., Fucini P., Yonath A., Harms J.M (2004) Inhibition of peptide bond formation by pleuromutilins: the structure of the 50S ribosomal subunit from *Deinococcus radiodurans* in complex with tiamulin. *Mol Microbiol* **54**:1287-1294 <https://doi.org/10.1111/j.1365-2958.2004.04346.x>
28. Poulsen S.M., Karlsson M., Johansson L.B., Vester B (2001) The pleuromutilin drugs tiamulin and valnemulin bind to the RNA at the peptidyl transferase centre on the ribosome. *Mol Microbiol* **41**:1091-1099 <https://doi.org/10.1046/j.1365-2958.2001.02595.x>
29. Davidovich C., Bashan A., Auerbach-Nevo T., Yaggie R.D., Gontarek R.R., Yonath A (2007) Induced-fit tightens pleuromutilins binding to ribosomes and remote interactions enable their selectivity. *Proc Natl Acad Sci U S A* **104**:4291-4296 <https://doi.org/10.1073/pnas.0700041104>
30. Arenz S., Ramu H., Gupta P., Berninghausen O., Beckmann R., Vazquez-Laslop N., Mankin A.S., Wilson D.N (2014) Molecular basis for erythromycin-dependent ribosome stalling during translation of the ErmBL leader peptide. *Nat Commun* **5**:3501 <https://doi.org/10.1038/ncomms4501>
31. Beckert B., Leroy E.C., Sothiselvam S., Bock L.V., Svetlov M.S., Graf M., Arenz S., Abdelshahid M., Seip B., Grubmuller H., et al. (2021) Structural and mechanistic basis for translation inhibition by macrolide and ketolide antibiotics. *Nat Commun* **12**:4466 <https://doi.org/10.1038/s41467-021-24674-9>
32. Dubot A., Godinot C., Dumur V., Sablonniere B., Stojkovic T., Cuisset J.M., Vojtiskova A., Pecina P., Jesina P., Houstek J (2004) GUG is an efficient initiation codon to translate the human mitochondrial ATP6 gene. *Biochem Biophys Res Commun* **313**:687-693 <https://doi.org/10.1016/j.bbrc.2003.12.013>
33. Kruger A., Remes C., Shiriaev D.I., Liu Y., Spahr H., Wibom R., Atanassov I., Nguyen M.D., Cooperman B.S., Rorbach J (2023) Human mitochondria require mtRF1 for translation termination at non-canonical stop codons. *Nat Commun* **14**:30 <https://doi.org/10.1038/s41467-022-35684-6>
34. Tsai K., Stojkovic V., Lee D.J., Young I.D., Szal T., Klepacki D., Vazquez-Laslop N., Mankin A.S., Fraser J.S., Fujimori D.G (2022) Structural basis for context-specific inhibition of translation by oxazolidinone antibiotics. *Nat Struct Mol Biol* **29**:162-171 <https://doi.org/10.1038/s41594-022-00723-9>

35. Syroegin E.A., Flemmich L., Klepacki D., Vazquez-Laslop N., Micura R., Polikanov Y.S (2022) Structural basis for the context-specific action of the classic peptidyl transferase inhibitor chloramphenicol. *Nat Struct Mol Biol* **29**:152-161 <https://doi.org/10.1038/s41594-022-00720-y>
36. Wakigawa T., Mito M., Yamashiro H., Tomuro K., Tani H., Tomizawa K., Chujo T., Nagao A., Suzuki T., Wei F.-Y., et al. (2023) Complexity and dynamics of *in organello* translation landscape assessed by high-resolution mitochondrial ribosome profiling. *bioRxiv* <https://doi.org/10.1101/2023.07.19.549812>
37. Bibel B., Raskar T., Couvillion M., Lee M., Kleinman J.I., Takeuchi-Tomita N., Churchman L.S., Fraser J.S., Fujimori D.G. (2024) Context-specific inhibition of mitochondrial ribosomes by phenicol and oxazolidinone antibiotics. *bioRxiv* <https://doi.org/10.1101/2024.08.21.609012>
38. Brandon M.C., Lott M.T., Nguyen K.C., Spolim S., Navathe S.B., Baldi P., Wallace D.C (2005) MITOMAP: a human mitochondrial genome database--2004 update. *Nucleic Acids Res* **33**:D611-613 <https://doi.org/10.1093/nar/gki079>
39. Kalghatgi S., Spina C.S., Costello J.C., Liesa M., Morones-Ramirez J.R., Slomovic S., Molina A., Shirihai O.S., Collins J.J (2013) Bactericidal antibiotics induce mitochondrial dysfunction and oxidative damage in Mammalian cells. *Sci Transl Med* **5**:192ra185 <https://doi.org/10.1126/scitranslmed.3006055>
40. D'Achille G., Morroni G (2023) Side effects of antibiotics and perturbations of mitochondria functions. *Int Rev Cell Mol Biol* **377**:121-139 <https://doi.org/10.1016/bs.ircmb.2023.03.009>
41. Desai N., Yang H., Chandrasekaran V., Kazi R., Minczuk M., Ramakrishnan V (2020) Elongational stalling activates mitoribosome-associated quality control. *Science* **370**:1105-1110 <https://doi.org/10.1126/science.abc7782>
42. Kong Q.P., Yao Y.G., Sun C., Zhu C.L., Zhong L., Wang C.Y., Cai W.W., Xu X.M., Xu A.L., Zhang Y.P (2004) Phylogeographic analysis of mitochondrial DNA haplogroup F2 in China reveals T12338C in the initiation codon of the ND5 gene not to be pathogenic. *J Hum Genet* **49**:414-423 <https://doi.org/10.1007/s10038-004-0170-3>
43. Vilarinho L., Choraó R., Cardoso M.L., Rocha H., Nogueira C., Santorelli F.M (1999) The ND1 T3308C mutation may be a mtDNA polymorphism. Report of two Portuguese patients. *J Inherit Metab Dis* **22**:90-91 <https://doi.org/10.1023/a:1005471904710>
44. Rocha A., Flores H., Campos C., Arenas Y., Vilarinho J., Santorelli L., and Torroni F.M. (1999) About the "Pathological" role of the mtDNA T3308C mutationellipsis. *Am J Hum Genet* **65**:1457-1459 <https://doi.org/10.1086/302641>
45. Soto I., Couvillion M., Hansen K.G., McShane E., Moran J.C., Barrientos A., Churchman L.S (2022) Balanced mitochondrial and cytosolic translomes underlie the biogenesis of human respiratory complexes. *Genome Biol* **23**:170 <https://doi.org/10.1186/s13059-022-02732-9>
46. Fiedler M., Rossmannith W., Wahle E., Rammelt C (2015) Mitochondrial poly(A) polymerase is involved in tRNA repair. *Nucleic Acids Res* **43**:9937-9949 <https://doi.org/10.1093/nar/gkv891>

Peer reviews

Reviewer #1 (Public review):

Summary:

This study aimed to determine whether bacterial translation inhibitors affect mitochondria through the same mechanisms. Using mitoribosome profiling, the authors found that most antibiotics, except telithromycin, act similarly in both systems. These insights could help in the development of antibiotics with reduced mitochondrial toxicity.

They also identified potential novel mitochondrial translation events, proposing new initiation sites for MT-ND1 and MT-ND5. These insights not only challenge existing annotations but also open new avenues for research on mitochondrial function.

Strengths:

Ribosome profiling is a state-of-the-art method for monitoring the translome at very high resolution. Using mitoribosome profiling, the authors convincingly demonstrate that most of the analyzed antibiotics act in the same way on both bacterial and mitochondrial ribosomes, except for telithromycin. Additionally, the authors report possible alternative translation events, raising new questions about the mechanisms behind mitochondrial initiation and start codon recognition in mammals.

Weaknesses:

The main weaknesses of this study are:

- While the authors highlight an interesting difference in the inhibitory mechanism of telithromycin on bacterial and mitochondrial ribosomes, mechanistic explanations or hypotheses are lacking.
- The assignment of alternative start codons in MT-ND1 and MT-ND5 is very interesting but does not seem to fully align with structural data.
- The newly proposed translation events in the ncRNAs are preliminary and should be further substantiated with additional evidence or interpreted with more caution.

<https://doi.org/10.7554/eLife.103699.1.sa2>

Reviewer #2 (Public review):

In this study, the authors set out to explore how antibiotics known to inhibit bacterial protein synthesis also affect mitoribosomes in HEK cells. They achieved this through mitoribosome profiling, where RNase I and Mnase were used to generate mitoribosome-protected fragments, followed by sequencing to map the regions where translation arrest occurs. This profiling identified the codon-specific impact of antibiotics on mitochondrial translation.

The study finds that most antibiotics tested inhibit mitochondrial translation similarly to their bacterial counterparts, except telithromycin, which exhibited distinct stalling patterns. Specifically, chloramphenicol and linezolid selectively inhibited translation when certain amino acids were in the penultimate position of the nascent peptide, which aligns with their known bacterial mechanism. Telithromycin stalls translation at an R/K-X-R/K motif in bacteria, and the study demonstrated a preference for arresting at an R/K/A-X-K motif in mitochondria. Additionally, alternative translation initiation sites were identified in MT-ND1 and MT-ND5, with non-canonical start codons. Overall, the paper presents a comprehensive analysis of antibiotics in the context of mitochondrial translation toxicity, and the identification of alternative translation initiation sites will provide valuable insights for researchers in the mitochondrial translation field.

From my perspective as a structural biologist working on the human mitoribosome, I appreciate the use of mitoribosome profiling to explore off-target antibiotic effects and the discovery of alternative mitochondrial translation initiation sites. However, the description is somewhat limited by a focus on this single methodology. The authors could strengthen their discussion by incorporating structural approaches, which have contributed significantly to the field. For example, antibiotics such as paromomycin and linezolid have been modeled in the human mitoribosome (PMID: 25838379), while streptomycin has been resolved (10.7554/eLife.77460), and erythromycin was previously discussed (PMID: 24675956). The reason we can now describe off-target effects more meaningfully is due to the availability of fully modified human mitoribosome structures, including mitochondria-specific modifications and their roles in stabilizing the decoding center and binding ligands, mRNA, and tRNAs (10.1038/s41467-024-48163-x).

These and other relevant studies should be acknowledged throughout the paper to provide additional context.

<https://doi.org/10.7554/eLife.103699.1.sa1>

Reviewer #3 (Public review):

Summary:

Recently, the off-target activity of antibiotics on human mitoribosome has been paid more attention in the mitochondrial field. Hafner et al applied mitoribosome profiling to study the effect of antibiotics on protein translation in mitochondria as there are similarities between bacterial ribosome and mitoribosome. The authors conclude that some antibiotics act on mitochondrial translation initiation by the same mechanism as in bacteria. On the other hand, the authors showed that chloramphenicol, linezolid and telithromycin trap mitochondrial translation in a context-dependent manner. More interesting, during deep analysis of 5' end of ORF, the authors reported the alternative start codon for ND1 and ND5 proteins instead of previously known one. This is a novel finding in the field and it also provides another application of the technique to further study on mitochondrial translation.

Strengths:

This is the first study which applied mitoribosome profiling method to analyze multiple antibiotics treatment cells.

The mitoribosome profiling method had been optimized carefully and has been suggested to be a novel method to study translation events in mitochondria. The manuscript is constructive and written well.

Weaknesses:

This is a novel and interesting study, however, most of the conclusion comes from mitoribosome profiling analysis, as a result, the manuscript lacks the cellular biochemical data to provide more evidence and support the findings.

<https://doi.org/10.7554/eLife.103699.1.sa0>