

Evolution of lateralized gustation in nematodes

Reviewed Preprint

v1 • January 22, 2025

Not revised

Marisa Mackie, Vivian Vy Le, Heather R Carstensen, Nicole R Kushnir, Dylan L Castro, Ivan M Dimov, Kathleen T Quach, Steven J Cook, Oliver Hobert, Sreekanth H Chalasani, Ray L Hong 

Department of Biology, California State University, Northridge, United States • Molecular Neurobiology Laboratory, Salk Institute for Biological Studies, La Jolla, United States • Department of Biological Sciences, Howard Hughes Medical Institute, Columbia University, New York, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

Mackie and colleagues present a **valuable** comparison of lateralized gustation in two well-studied nematodes. The evidence they present that ASEL/R lateralization exists and is achieved by different means in *P. pacificus* compared to *C. elegans* is **solid**, but would be strengthened by the inclusion of calcium imaging data for *C. elegans* and additional verification of *P. pacificus* AFD neuron identity. This work will be of interest to evolutionary biologists, ecologists, and sensory neurobiologists interested in how small nervous systems make sense of the environment, and how evolution can take multiple paths to asymmetry within a neuron class.

<https://doi.org/10.7554/eLife.103796.1.sa3>

Abstract

Animals with small nervous systems have a limited number of sensory neurons that must encode information from a changing environment. This problem is particularly exacerbated in nematodes that populate a wide variety of distinct ecological niches but only have a few sensory neurons available to encode multiple modalities. How does sensory diversity prevail within this neuronal constraint? To identify the genetic basis for patterning different nervous systems, we demonstrate that sensory neurons in the *Pristionchus pacificus* respond to various salt sensory cues in a manner that is partially distinct from that of the distantly related nematode *C. elegans*. By visualizing neuronal activity patterns, we show that contrary to previous expectations based on its genome sequence, the salt responses of *P. pacificus* are encoded in a left/right asymmetric manner in the bilateral ASE neuron pair. Our study illustrates patterns of evolutionary stability and change in the gustatory system of nematodes.

Impact statement

Animals with small nervous systems have evolved left and right asymmetry in their neurons to process various salts at different concentrations.

Introduction

Nematodes form a vast array of ecological relationships, from specialized parasite-host dependencies to nematode-microbial interactions, each one demanding exquisitely fine-tuned sets of sensory palates that span multiple modalities (1–8). Yet, the number of sensory neurons across diverse nematode species seems to be constrained (9–11). Several free-living and parasitic nematode species examined by serial section electron microscopy have nearly identical numbers of 12–13 pairs of head sensory neurons, known as the amphid neurons (11–15). How does sensory diversity arise within this neuronal constraint? When coupled with well-described neuronal anatomy, this conserved neuron count allows for detailed comparisons at the single-cell resolution and represents an opportunity to interrogate how sensory cues are processed by anatomically similar nervous systems to produce species-specific or developmental stage-dependent behavioral outputs. To identify the genetic basis for patterning different nervous systems and to understand the processes that underlie evolutionary changes in adapting to different environments, several comparative model systems have been developed to promote comparisons to the well-studied nematode *Caenorhabditis elegans* at the genetic and cellular levels, including the predatory entomophilic nematode, *Pristionchus pacificus* (11, 16–19).

As expected from their association with insects in the wild, the olfactory preferences of *P. pacificus* are distinct from those of *C. elegans* and the human parasite *Strongyloides stercoralis* (2, 3, 20), but little is known about *P. pacificus* responses to water-soluble compounds. In *C. elegans*, the main salt receptor neuron class comprises of a bilateral pair of left and right ASE neurons (ASEL and ASER), which serve to induce an attractive locomotory response toward an increase in salt concentration (21). The gene *che-1* (chemosensory defective) encodes a transcription factor that is exclusively expressed in the ASE neurons and is required for their proper differentiation, such that a *che-1* mutant results in defective salt attraction (22–25).

One major role of *C. elegans* CHE-1 is to promote lateral asymmetry in the ASE neurons. The left and right ASE neurons asymmetrically express receptor-type guanylyl cyclases (rGCs, encoded by *gcy* genes) (26). This finding led to the realization that the ASE neurons are lateralized, such that the left and the right ASE neurons differentially respond to distinct salt ions (27–30). This observation led in turn to the identification of a complex gene regulatory network that genetically programs the distinct sensory potentials of the left and right ASE neurons (31), which includes a miRNA, *lsey-6*, at the top of this gene regulatory network (31–33). However, the *lsey-6* miRNA evolved selectively in the *Caenorhabditis* genus (34) and is absent in *P. pacificus*. Moreover, *P. pacificus* does not show an expansion of the ASEL-type and ASER-type rGCs, as *C. elegans* does (Hong et al., 2019). With these two genomic observations in mind, we had previously proposed that the ASE neurons are unlikely to be lateralized in *P. pacificus* (11).

However, we now revise this view in light of our work on mapping chemosensory responses in *P. pacificus* on the level of behavior and neuronal activity. We demonstrate that *P. pacificus* does in fact show lateralized chemosensory profiles, indicating that *P. pacificus* must have evolved independent means to establish ASE laterality. We also show that the tastant palate of *P. pacificus* is distinct from that of *C. elegans*, and that its dependence on ASE, as well as its terminal selector transcription factor *che-1*, have also diverged.

Results

P. pacificus and *C. elegans* differ in their behavioral responses to salts

Previous cross-species comparisons between *P. pacificus* and *C. elegans* indicated strong differences in olfaction preferences that reflect their divergent evolutionary histories and ecology (29). To identify the neurons that mediate gustation, we first compared the chemosensory profiles of these two species toward water-soluble ions. In this survey, we found ammonium salts to be the strongest attractants to wildtype *P. pacificus* adult hermaphrodites (NH_4Br , NH_4Cl , NH_4I), with NH_4I significantly more attractive to *P. pacificus* compared to *C. elegans* (Fig. 1). Notably, in contrast to *C. elegans* (35), we find that *P. pacificus* is only moderately and weakly attracted to NaCl and LiCl , respectively (Fig. 1). Also, *P. pacificus* is repulsed by acetate salts (NaAc , NH_4Ac), which induce attractive responses in *C. elegans* (36). We conclude that *P. pacificus* and *C. elegans* display differences in their salt preferences.

Ppa che-1 shows similarities and difference to *Cel che-1* in both expression and function

The *C. elegans che-1* mutant (*chemotaxis-defective*) was originally isolated for its inability to respond to a broad panel of salt tastants (22, 35), including those described above (29, 30, 36). *Che-1* was found to encode for a Zn finger transcription factor that is exclusively expressed in the ASE neuron pair (25), which through laser ablations had been found to be the main salt receptor neurons (37). *che-1* was found to control the entire differentiation of the ASE neurons, including the expression of putative receptors of the GCY receptor guanylyl cyclase family (23–25, 38).

To assess whether *che-1* performs a similar function in salt perception for *P. pacificus* as for *C. elegans*, we analyzed the expression and function of the *P. pacificus che-1* ortholog. In a previous paper, we reported that the 5' region of the sole *Pristionchus che-1* ortholog directs reporter expression to the *Ppa* ASE and *Ppa* ASG neuron classes (11). Re-examination of our provisional cell identifications using a newly-generated *che-1p::GFP* strain with stronger neurite expression revealed highly elaborated finger-like dendritic endings in the more anterior amphid neuron that could unambiguously be assigned to the AFD neurons (Fig. 2A–C), prompting us to reassign expression of *che-1* to ASE and AFD.

We confirmed that the *che-1* reporter transgene indicates the full expression of the endogenous *che-1* locus by tagging the endogenous *che-1* locus with an ALFA-tag (39). The *che-1::ALFA* animals showed staining in 2 pairs of head neurons whose position is consistent with being the ASE and AFD neurons (Fig. 2D–E). By crossing the *che-1* reporter transgene into a *che-1* mutant background (see below), we also found that *che-1* autoregulates its own expression (Fig. 2F), as it does in *C. elegans* (25).

Next, we examined whether *P. pacificus* salt responsiveness shows similar *che-1* dependence as in *C. elegans*. We generated two putative null alleles in the *P. pacificus che-1* homolog using CRISPR/Cas9 genome engineering, through introduction of small deletions in the first exon of the gene, thereby resulting in frameshift and premature stops (Fig. 3A). Both *Ppa che-1* alleles exhibited defects in attraction towards NH_4Cl and LiCl compared to wildtype (Fig. 3B). However, unlike in *C. elegans*, *Ppa-che-1* mutants showed no detectable difference in responses to NH_4I , NaCl , and NaAc .




Figure 1.

A comparison of chemotaxis responses to water-soluble ions between *P. pacificus* and *C. elegans*.

The J4/L4 larvae and young adult hermaphrodites from the two species responded to NH₄I, LiCl, and acetates significantly differently. ****P<0.0001, **P<0.01 significant difference found between wildtype *P. pacificus* and *C. elegans* by unpaired t-test. Each assay involves a minimum of 10 J4/adult hermaphrodites.




Figure 2.

***P. pacificus che-1* expression in ASE and AFD amphid neurons.**

(A-B) The *che-1::GFP* marker in the *che-1::RCaMP* reporter strain is expressed in the ASE and AFD amphid neurons. (C) *che-1::GFP* expression is detectable in the morphologically distinct AFD neurons with 'finger'-like dendritic endings. (D-E) Immunostaining of *CHE-1:ALFA* shows two pairs of amphid neuron cell bodies corresponding to the ASE and AFD neurons; dorsal-ventral view (n=114). (F) The loss of *che-1* results in loss *che-1::RFP* expression in the ASE (circle) while retaining AFD expression. Anterior is left and the scale bar in (C) represents 50 μ m for all panels.




Figure 3.

***Che-1* expressing amphid neurons are required for sensing water-soluble ions in *P. pacificus*.**

(A) The *che-1* locus with CRISPR/Cas9-induced mutations in Exon 1. (B) The *che-1* mutants show defects in attraction towards NH₄Br, NH₄Cl, and LiCl. (C) The *che-1p::HisC1* animals lose attraction towards NH₄Br, NH₄Cl, and NH₄I in a histamine dependent manner. (D) A schematic map of the *P. pacificus* AM7(ASE) and AM12(AFD) amphid neurons that express the *che-1p::RCaMP* used in calcium imaging. The ASE axons are the only amphid axons in *P. pacificus* to cross each other over the dorsal lateral midline contralaterally. **P<0.01, *P<0.05 Dunnett's post-hoc comparison showing alleles with significant difference to wildtype *P. pacificus* (PS312).

We considered two different possibilities for the behavioral differences of *Ppa che-1* and *Cel che-1* mutants. *P. pacificus* may use sensory neurons other than ASE to sense these cues, or alternatively, *Ppa-che-1* may not have the same fundamental impact on ASE function in *P. pacificus* as it does in *C. elegans*. To explore these different possibilities, we silenced *che-1* expressing neurons by expressing codon-optimized HisCl channel under control of the *che-1* promoter. Histamine-treated *che-1p::HisCl* animals showed complete loss of attraction to NH_4Br , NH_4Cl , and NH_4I . Histamine treatment did not significantly alter their repulsive response to NaAc (**Fig. 3C**). These findings corroborate that NaAc is sensed by neurons other than ASE (or AFD, in which the *che-1* promoter also drives HisCl). Since NH_4I sensation is affected by silencing of *che-1*(+) neurons but is unaffected in *che-1* mutants, ASE differentiation may be more greatly impacted by the silencing of ASE than by the loss of *che-1*.

The ASE neurons show left/right asymmetric responses to salt

To assess whether *Ppa* ASE neurons show the same lateralized response to salt ions as *Cel* ASE neurons, we generated transgenic *P. pacificus* lines that express RCaMP in ASE neurons and assessed calcium responses to attractive salts (**Fig. 4**, **S1**–**S2**). Specifically, we looked for changes in calcium levels immediately after the addition and removal of specific salts. When 250 mM NH_4Cl is administered, an ‘ON’ response is observed as calcium transiently increases in the left ASE neuron (ASEL). In contrast, an ‘OFF’ response was observed as calcium sharply dips in the right ASE neuron (ASER) before quickly returning to baseline when the salt was removed, which suggests hyperpolarization of this neuron (**Fig. 4A–B**) (29). However, when presented with a 10-fold lower concentration of 25 mM NH_4Cl , the ‘OFF’ response completely disappeared in ASER while the ‘ON’ response became more pronounced in ASEL. The ASER responses to 250 mM and 25 mM NaCl (**Fig. 4D**) were very similar to the ‘OFF’ response (including hyperpolarization) observed for NH_4Cl , but the ASEL responses differ between the two salts: instead of the ‘ON’ response expected of ASEL, we observed a relatively weak ‘OFF’ response without the characteristic hyperpolarization but accompanied by an attenuated ‘bump’, a profile that we classify as an ‘OFF-2’ response (**Fig. 4C**). Finally, we examined the response to NH_4I and found that it also elicited laterally asymmetric responses, but yet again distinct from the responses to both NH_4Cl and NaCl, with ASEL showing a strong ‘ON’ response, and ASER showing an ‘ON-OFF’ biphasic response to 250 mM NH_4I (**Fig. 4E–F**) (40, 41). Interestingly, the ASER exhibited an ‘OFF’-only response to 25 mM NH_4I , which was not observed for the same concentration of NH_4Cl and NaCl, and thus may reflect the higher response to NH_4I than to NH_4Cl and NaCl in the behavior assays. Altogether, *P. pacificus* ASE neurons clearly show left/right asymmetric responses to salt attractants and these asymmetric responses show similarities and differences to the *C. elegans* ASE taste neurons (see Discussion).

The AFD neurons also respond to salts in *P. pacificus*

The RCaMP line that we used to assess calcium responses in ASE is also expressed in AFD, allowing us to simultaneously examine the calcium responses in the AFD (AM12) neurons (**Fig. 5**, **S1**, **S3**–**S5**). Surprisingly, we detected a distinctly ‘ON-OFF’ biphasic response to all 3 salt types at both concentrations. Specifically, although we observed weaker or comparable responses in AFD neurons when compared to either ASE neuron’s response towards 250 mM NH_4Cl , NaCl and NH_4I (**Fig. 5A–B**, **5E–F**, **5I–J**), the AFD responses were more robust than ASER toward 25 mM NH_4Cl , NaCl, and NH_4I (**Fig. 5D**, **5H**, **5L**). Specifically, AFD neurons responded strongly to 25 mM NaCl when neither one of the ASE neurons showed a positive response (**Fig. 5G–H**). Averaging the calcium transients separately by AFD left versus right did not result in significant differences in the shape of the neuronal calcium responses, with the exception of the AFD responses to higher versus lower concentrations of NH_4I (**Fig. 54**–**55**). We have not further pursued whether these AFD responses are a reflection of a direct perception of salt or a secondary consequence of communication of salt-perceptive neurons (like ASE) to AFD.




Figure 4.

ASEL and ASER responses to different concentrations of NH₄Cl, NaCl, and NH₄I.

Average percent change in RCaMP fluorescence (dF/F) over time (seconds) of tracked ASEL (left and right) sensory neurons in *P. pacificus*. Salts were presented at 10s ("ON", left vertical line) for a duration of 20s, and then removed ("OFF", right vertical line) for the remaining 30s; the total recording time was 60s. (A-B) ASEL and ASER neuron responses to high (250 mM, red) compared to low (25 mM, blue) concentrations of NH₄Cl. (C-D) ASEL and ASER neuron responses to NaCl. (E-F) ASEL and ASER neuron responses to NH₄I. Shaded ribbons represent 95% confidence intervals. Shaded ribbons in (E-F) have been cropped to maintain consistent y-axes across the plots, allowing for easier comparison. Sample sizes are indicated (n).




Figure 5.

Combined AFD responses in comparison to ASE left or right neuron responses to NH₄Cl, NaCl, and NH₄I.

Average percent change in RCaMP fluorescence (dF/F) over time (seconds) as described in [Fig. 4](#). Averaged combined AFD (left and right neurons, orange) compared to left or right ASE (green) responses to (A-B) 250mM NH₄Cl (C-D) 25mM NH₄Cl (E-F) 250mM NaCl (G-H) 25mM NaCl (I-J) 250mM NH₄I (K-L) 25mM NH₄I. Shaded ribbons represent 95% confidence intervals. Shaded ribbons in (J-K) have been cropped to maintain consistent y-axes across the plots, allowing for easier comparison. Sample sizes are indicated (n).

A target of *che-1*, the guanylyl cyclase *gcy-22.3* is required for chemotaxis

We further explored the asymmetric salt perception by the ASE neurons, which in *C. elegans* is largely mediated through distinct receptor-type guanylyl cyclases (rGC proteins, encoded by *gcy* genes), which confer salt specificity via their extracellular domains (Ortiz 2009, Smith 2013). Our previous genome survey of *Ppa* homologs of *gcy* genes has revealed patterns that made us question whether *Ppa gcy* genes are convincing candidates for lateralized chemotactic responses. Specifically, we noted that ASER-expressed *C. elegans gcy* genes and ASEL-expressed *C. elegans gcy* genes have only expanded in the *Caenorhabditis* genus (11 [↗](#)). One outlier to this pattern is the *Cel gcy-22* gene, which is expressed in ASER, but has not expanded in *C. elegans* (11 [↗](#), 42 [↗](#)). However, this gene has duplicated several times in *P. pacificus*, resulting in 5 putative *Ppa-gcy-22* paralogs (11 [↗](#)). We fused the promoter of one of these paralogs, *Ppa-gcy-22.3*, to *gfp* and found that transgenic animals express GFP exclusively in ASER (Fig. 6A [↗](#)), identical to the *C. elegans gcy-22* ortholog (42 [↗](#)). We confirmed its expression in ASER by analyzing animals that carry both the *Ppa gcy-22p::gfp* reporter and the *Ppa che-1p::rfp* reporter, showing a unilateral overlap of these reporters in ASER (Fig. 6B [↗](#)).

To assess whether *Ppa-gcy-22.3* is a potential effector of *Ppa che-1* function, we crossed the *gcy-22.3* reporter into *che-1(ot5012)* mutants. We found expression of *gcy-22.3* was eliminated (Fig. 6C [↗](#)), leading us to conclude that *gcy-22.3* is a potential effector of *che-1* function, identical to its homolog in *C. elegans*.

To determine whether and which of the observed salt responses is mediated by the ASER-expressing *gcy-22.3*, we generated a putative *gcy-22.3 null* mutant through CRISPR/Cas9 genome editing (2 bp complex deletion that introduces frameshift)(Fig. 7A [↗](#)) and examined its response to the higher salt concentration. The ‘OFF’ response to 250 mM NH₄Cl was notably abolished in the ASER neuron in the loss-of-function *gcy-22.3* mutant, while the ‘ON’ response in the ASEL remained intact (Fig. 7B-C [↗](#)). However, the responses to 250 mM NaCl were not significantly reduced in the either the ASEL or ASER neuron in the *gcy-22.3* mutant (Fig. 7D-E [↗](#)). Furthermore, the *gcy-22.3* mutation also reduced the ‘ON’ portion of the AFD biphasic response following presentation of 250 mM NH₄Cl and NaCl (Fig. 7F-G [↗](#)). Our findings suggest GCY-22.3 is possibly a *P. pacificus* salt receptor that can affect the physiological responses to salt in the ASER gustatory neuron, as well as other gustatory neurons that do not express this receptor (AFDs).

Discussion

Our study has revealed several insights into the substrates of evolutionary changes between two distantly related nematode species, *P. pacificus* and *C. elegans*. We used intracellular calcium levels as our readout for neuronal activity with a genetically encoded calcium sensor in two pairs of *che-1*-expressing amphid sensory neurons– the first calcium imaging study in *P. pacificus*. We show that three neuron types (ASE left, ASE right and AFD neurons) each have distinct calcium responses to specific ion concentrations, revealing diversity at the single neuron level. We have identified the first laterally asymmetric marker in *P. pacificus*, *gcy-22.3p::GFP*, with its expression limited to ASER homolog (AM7). Our unexpected discovery that neuronal asymmetry is present in the ASE homologs between two distantly related nematode species, despite the lack of a *lsy-6* homolog, suggests that functional lateralization in *P. pacificus* may be mediated by a different genetic pathway compared to *C. elegans*.




Figure 6.

The laterally asymmetric expression of *gcy-22.3* is dependent on the zinc finger transcription factor CHE-1.

(A) The *gcy-22.3::GFP* marker is expressed exclusively in the right ASE neuron (ASER)(n>200). (B) The *gcy-22.3::GFP* marker co-localizes with *che-1::RFP* expression in the ASER. (C) *gcy-22.3::GFP* expression is absent in the *che-1(ot5012)* mutant (n=55). Anterior is left and dorsal is top. Scale bar in (A) represents 50 μ m for all panels.




Figure 7.

ASE and AFD responses in wildtype compared to *gcy-22.3* mutants.

Average percent change in RCaMP fluorescence (dF/F) over time (seconds) as described in [Fig. 4](#). (A) WormBase JBrowse view of the *gcy-22.3* locus with CRISPR/Cas9-induced mutations in Exon 5. (B-C) ASEL and ASER neuron responses to 250 mM NH₄Cl wildtype (gray) and *gcy-22.3* mutants (magenta). (D-E) ASEL and ASER neuron responses to 250 mM NaCl in wildtype (gray) and *gcy-22.3* mutants (magenta). AFD (combined left and right neurons) responses (F) to 250 mM NH₄Cl and (G) to 250 mM NaCl in wildtype (orange) and *gcy-22.3* mutants (magenta). Shaded ribbons represent 95% confidence intervals. Sample sizes are indicated (n). For ASEL/R comparisons, bar plots represent the difference between the minimum % dF/F value 10s pre-stimulus and maximum % dF/F value 10s post-stimulus for either the (B) ON or (C-E) OFF stimulus. For AFD comparisons, bar plots represent maximum % dF/F values 10s after the ON and OFF stimulus. *P<0.05, **P<0.01, ***P<0.001 indicate significant difference. "ns" indicate no significant difference. Comparisons between different genotypes were analyzed using unpaired t-test or Mann-Whitney test; comparisons within the same genotype were analyzed using paired t-test.

***P. pacificus* and *C. elegans* have diverged taste palates**

We have shown that while *C. elegans* is attracted to acetate salts, *P. pacificus* avoids these acetates. Previous studies have shown that ammonium acetate (NH_4Ac) is sensed both as a water-soluble compound as well as a volatile odorant and is mediated by different signaling pathways (36). Like *C. elegans*, it is therefore likely that ammonium and acetate ions involve a different set of neurons in *P. pacificus* (i.e. non-*che-1* expressing neurons), based on the finding that histamine-treated *che-1p::HisCl* animals did not significantly attenuate their repulsive response to NaAc. We find that sodium chloride is a common attractant for both *P. pacificus* and *C. elegans*, although the magnitude of response is lower than previously published results (29). This result is likely due to differences in generating the salt gradients. Nevertheless, we find that *P. pacificus* neurons have a distinct response to this salt when compared to those observed in *C. elegans* neurons. The *P. pacificus* ASEL neuron responds to a decrease in NaCl concentration (likely sodium), as evidenced by the ‘OFF-2’ response profile, whereas the *C. elegans* ASEL neuron responds to an increase in sodium concentration. However, in both nematodes, the ASER neuron responds to a decrease in chloride concentration. Additionally, the *P. pacificus* ASER neuron exhibits a unique ON-OFF response to ammonium iodide, whereas in *C. elegans*, no ON-OFF type of response is seen in the ASE neurons—the *C. elegans* ASER neuron responds only to a decrease in iodide concentration. Collectively, our findings highlight the divergence of the *P. pacificus* salt-sensory neurons compared to those observed in *C. elegans*.

***P. pacificus* ASE neurons have narrower sensitivity range**

The sensitivity range of the *P. pacificus* ASER responses is significantly less compared to the ASEL responses, as well as to *C. elegans* ASER responses. Whereas the *C. elegans* ASER has a 40-fold sensitivity range in the ‘OFF’ response to the removal of various concentrations of NaCl (1–40 mM) (29, 43–45), the *P. pacificus* ASER showed the ‘OFF’ response only to 250 mM NH_4Cl but not to a 10-fold reduction concentration of NH_4Cl (25 mM). For the ASEL in contrast, the response to 25 mM was just as strong as to 250 mM NH_4Cl (10-fold) and comparable to the 8-fold concentration range observed for *C. elegans* ASEL towards NaCl. Alternatively, the magnitude of these sensitivity differences may be also partially due to differences among calcium indicators (i.e. GCaMP and Cameleon), but multiple *P. pacificus che-1p::GCaMP* strains did not exhibit sufficient basal fluorescence to allow for image tracking and direct comparison. The narrower sensitivity range in *P. pacificus* chemosensation was also observed for attraction to volatile odors (2), which span only 10-fold, versus up to 10,000-fold in attractive odors for *C. elegans* (21).

***P. pacificus* taste neurons exhibit a unique biphasic response**

Although the left-ON and right-OFF responses are conserved in the ASE neurons in both species, the biphasic response of ASER to 250 mM NH_4I has not been observed in either of the ASE neurons towards salts. In *C. elegans*, hyperosmotic stimulus such as 1M glycerol, or high concentrations or long duration of CuSO_4 exposure both result in biphasic responses by the ASH neurons that sense noxious chemicals (41, 46). Olfactory neurons that mediate avoidance behavior such as the AWB neurons can also exhibit a biphasic response to the presence and removal of a high concentration isoamyl alcohol that normally elicits avoidance behavior (47). The neuron-specific response can also be dependent on the concentration of the chemical compound, since the ASER to 25 mM NH_4I was a weak ‘OFF’ rather than a biphasic one. In contrast to the salt-dependent response types by *P. pacificus* ASE neurons, the *P. pacificus* AFD neurons also show exclusively biphasic responses with various amplitudes. Further characterization will help determine if biphasic responses can also be found in other sensory neuron types in *P. pacificus*, specifically those neurons mediating avoidance behavior.

AFD are potentially polymodal neurons

Broadly speaking, *C. elegans* chemosensory neurons have been classically characterized as specialized neurons for dedicated modalities such as water-soluble chemicals (ASE), volatile odorants (AWA, AWB, AWC), noxious chemicals (ASH), pheromones (ADL, ADF, ASK), as well as light (ASJ) and temperature (AFD) (1, 37, 48–51). Advances in multi-neuron calcium recordings have since shown that a given odor within a certain concentration range is detected by different ensembles of the 12 amphid neuron classes, including the AFD neurons (52–54). Unexpectedly, we found that the *P. pacificus* AFD neurons exhibit a distinctive biphasic response to all three salts tested (NH₄Cl, NaCl, and NH₄I), which differ from the *C. elegans* AFD calcium responses (54, 55). Moreover, the loss of the receptor *gcy-22.3* reduced the AFD ‘ON’ response to NH₄Cl and NaCl, indicating ASER has only a limited contribution to the AFD response. The strong positive AFD response to 25 mM NaCl that is absent in both ASE neurons further supports the likelihood that AFD receives inputs, possibly neuropeptidergic, from other amphid neurons. The integration of thermosensation and chemosensation is important for memory-regulated behavior. In *C. elegans*, maximum chemotaxis indices toward NH₄Cl occurs when there is concordance between cultivation temperature and assay temperature (56). The *C. elegans* AFD neurons are also important for gustatory aversive learning in NaCl avoidance (45). Given the influence of environmental temperature on the *P. pacificus* mouth-form plasticity and the wide range of micro-climates that wild strains *P. pacificus* have been isolated from (57–59), temperature and taste preferences could be regulated at multiple genetic levels during crucial developmental decisions.

Changes in the gene regulatory architecture of sensory neuron specification

We found that the key regulator of *Cel* ASE identity, *che-1*, is also expressed in *Ppa* ASE and may play a similar role as a terminal selector in this neuron, based on its effect on ASE-mediated behavior and regulation of the *Ppa-gcy-22.3* gene. However, the stronger behavioral effect of silencing of *che-1* expressing neurons compared to a *che-1* mutant background could either indicate that *che-1* does not have as broad a role in controlling ASE differentiation in *P. pacificus* versus *C. elegans*. It is also possible that a developmental loss of ASE differentiation may result in compensatory changes in the chemosensory system during early development, as was observed in the *C. elegans* mating pheromone response by males (60).

Unexpectedly, we found that unlike in *C. elegans*, the *Ppa che-1* gene is also expressed in the AFD neurons. Since we cannot record neural activity in AFD in a *che-1* mutant (the *che-1p::RCaMP* driver fails to be expressed properly in *che-1* mutants due to autoregulation), and do not yet have molecular markers for *Ppa* AFD neurons, we cannot assess whether *che-1* affects AFD neuron differentiation.

Perhaps the most striking difference in the gene regulatory architecture of ASE neuron specification is the apparent lack of the key regulator of ASE asymmetry in *P. pacificus*, the miRNA *lsy-6*. In *C. elegans*, the expression of *lsy-6* exclusively in ASEL is prepatterned via an early embryonic Notch signal (33) and serves to downregulate the homeodomain transcription factor *cog-1* in the ASEL neuron (61). Through a network of downstream regulatory events, asymmetry of rGCs eventually becomes established (31). *cog-1* and several asymmetrically expressed downstream effectors of *cog-1*, such as the *die-1*, *lim-6* and *fozi-1* transcription factors are conserved in *P. pacificus*, but whether the function of these factors in controlling *P. pacificus* ASE laterality is conserved remains to be determined. In this context it is intriguing to note that another prominent gene regulatory pathway that is controlled by miRNAs in *C. elegans*, the heterochronic pathway (*let-7s*, *lin-4*), appears to have diverged in *P. pacificus* as well, despite the

conservation of the overall physiological readouts of this pathway (temporal patterning of cell lineage divisions)([62](#)). It is tempting to speculate that miRNA-mediated regulatory process is particularly labile.

In conclusion, our work illustrates how comparative behavioral and genetic analyses in nematodes is a powerful strategy to uncover substrates of evolutionary change in simple nervous systems.

Materials and Methods

Nematode Strains

P. pacificus and other nematode strains were maintained at ~20°C on NGM plates seeded with *E. coli* OP50 for food as described previously ([63](#)); these are derived from standard *C. elegans* culture methods ([64](#)). *P. pacificus* and other nematode strains used are listed in **Table S1**.

Chemotaxis assays

The assay for assessing response to salt gradients was adapted from *C. elegans* and *P. pacificus* chemotaxis assays ([2](#), [21](#), [42](#)). Overnight salt gradients were established on 10 cm chemotaxis plates containing 20 ml agar (5 mM KPO₄, 1 M CaCl₂, 3% Bacto-agar, 1 mM MgSO₄) by adding 10 µl of 2.5 M salt solutions for 16 hours. Then, another 4 µl of the same salt solution or water control was added to reinforce the gradient 4 hours before the assay. Just prior to the assay, 1.5 µl of 1 M sodium azide was added to both the attractive salt (A) and the control (C) spots. Young *P. pacificus* adult hermaphrodite from near-saturated cultures were washed 3x with distilled water and collected by centrifuging at 2000 rpm for 2 minutes. Approximately 200 worms were loaded onto the edge of each assay plate between the gradient sources, and at least 10 combined worms have to reach the scoring arenas to be considered. At least 12 assays constituted each experimental trial, and multiple trials were conducted and averaged for each condition. The Chemotaxis Index (CI) for each end-point assay plate is defined as (A - C)/(A + C). To conduct conditional knockdowns of neurons, 5 M histamine dihydrochloride (Sigma Aldrich H7250) stock solution in sterilized deionized water (Arrowhead CA) was filter-sterilized and top plated onto the agar plates to a final histamine concentration of 25 mM histamine approximately 10 minutes before loading the worms to commence the assay. Most assays lasted ~3.5 hours at room temperature in line with the speed of *P. pacificus* without food, with ~40% of the animals participating. When using the *Ppa-che-1p::optHisCl(csEx83)* strain, only animals expressing *Ppa-egl-20p::RFP* tail marker from the extrachromosomal array were scored. Because of *P. pacificus*' strong aversion to acetate, we could not easily assess the individual contributions of salt ions in a saturated background of ammonium acetate as conventionally practiced in *C. elegans* studies ([30](#)).

Ppa-che-1p::HisCl strain

To make the *Ppa-che-1p::optHisCl*, the *P. pacificus* codon-optimized histamine-gated chloride channel sequence used in *C. elegans* (*HisCl*)([65](#), [66](#)) was designed using (https://hallemmlab.shinyapps.io/Wild_Worm_Codon_Adapter/) and custom synthesized (Twist Bioscience), and subsequently inserted behind the *che-1* promoter (3.1 kb containing the first exon and intron) ([11](#)) to create the pH30 plasmid construct. This *Ppa-che-1p::optHisCl* plasmid (2 ng/µl) along with PS312 genomic DNA (80 ng/µl) and *Ppa-egl-20p::RFP* co-injection marker (2 ng/µl) were individually digested with HindIII and assembled as the injection mix to create *csEx83*.

Ppa-che-1p::RCaMP reporter strain

To make the *Ppa-che-1p::optRCaMP*, we generated a transgenic worm strain expressing the codon-optimized genetically-encoded calcium indicator (GECI), jRCaMP1a, in the neurons of interest (67 [↗](#), 68 [↗](#)). jRCaMP1a is an improved red GECI based on mRuby with comparable sensitivity to GCaMP6 (69 [↗](#)). The codon-optimized RCaMP sequence was custom synthesized (Twist Bioscience), and then subcloned using the plasmid pJET1.2/blunt (Thermo Fisher Scientific) to create the pMM2 plasmid construct. Using Gibson Assembly (E2611, New England Biolab), the codon-optimized RCaMP sequence was introduced downstream of the *Ppa-che-1* promoter sequence to create the pMM5 plasmid construct. Separately, we also generated a *Ppa-che-1p::optGFP* transcriptional reporter with codon-optimized *GFP* (pMM3) to enable the localization of the *che-1*-expressing neurons during video acquisition (70 [↗](#)). The *Ppa-che-1p::optRCaMP* (2 ng/μl) and *Ppa-che-1p::optGFP* constructs (1 ng/μl), along with PS312 genomic DNA (80 ng/μl) and *Ppa-egl-20p::RFP* (1.5 ng/μl) were individually digested with HindIII and assembled as the injection mix to create *csuEx93*. Despite multiple attempts, we were unable to generate an equivalent *che-1p::GCaMP* transgenic line with sufficient basal level of GCaMP expression for a comparison to the RCaMP calcium dynamics.

CRISPR mutagenesis generated mutants

CRISPR/Cas9 mutagenesis was used to generate mutations (70 [↗](#), 71 [↗](#)). crRNA and primer sequences, and induced mutations, are included in [Table S3](#) [↗](#) and [Fig. S6](#) [↗](#).

che-1 alleles (PPA01143)

Target crRNA, tracrRNA, and Cas9 nuclease were purchased from IDT Technologies (San Diego, CA). crRNA and tracrRNA were hydrated to 100 μM with IDT Duplex Buffer, and equal volumes of each (0.61 μl) were combined and incubated at 95°C for 5 minutes, then 25°C for 5 minutes. Cas9 protein (0.5 μl of 10 μg/μl) was added, then the mix was incubated at 37°C for 10 minutes. *Ppa-egl-20p::RFP* was used as a co-injection marker. To reach a final total volume of 40 μl, the Cas9-crRNA-tracrRNA complex was combined with pZH009 (*Ppa-egl-20p::RFP*) DNA to reach 50 ng/μl final concentration using nuclease-free water. F₁ progeny were screened for the presence of *Ppa-egl-20p::RFP* expression in the tail and candidate F₁'s were sequenced to identify heterozygotes (71 [↗](#)). *ot5012* has a 4 bp insertion while *ot5013* has an 8 bp complex insertion/deletion, and both mutations cause frameshift mutations and premature stop codons. Each allele was outcrossed two times to wildtype before characterization.

gcy-22.3 alleles (PPA04454)

Target crRNA, tracrRNA, and Cas9 nuclease were purchased from IDT Technologies (San Diego, CA). crRNA (RHL1400) and tracrRNA were hydrated to 100 μM with IDT Duplex Buffer, and equal volumes of each (0.61 μl) were combined and incubated at 95°C for 5 minutes, then 25°C for 5 minutes. Cas9 protein (0.5 μl of 10 μg/μl) was added, then the mix was incubated at 37°C for 10 minutes. *Ppa-egl-20p::RFP* was used as a co-injection marker. To reach a final total volume of 40 μl, the Cas9-crRNA-tracrRNA complex was combined with pZH009 (*Ppa-egl-20p::RFP*) DNA to reach 50 ng/μl final concentration using nuclease-free water. F₁ progeny were screened for the presence of *Ppa-egl-20p::RFP* expression in the tail and candidate F₁'s were sequenced to identify heterozygotes (71 [↗](#)). *csu181* has a 2 bp complex deletion, while *csu182* has a 22 bp complex deletion, both mutations cause frameshifts and premature stop codons. Each allele was outcrossed two times to wildtype before characterization.

CHE-1:ALFA C-terminal tagging and immunostaining

For induction of site-specific insertions via CRISPR/Cas-9 mediated mutagenesis, target crRNA, tracrRNA, and Cas9 nuclease were treated as described above. Single-stranded DNA repair template containing the ALFA nanobody tag (RHL1551) with 35 bp of homology arms on the 5' and 3' sides were purchased from IDT. To minimize sequence identity, the two copies of the ALFA sequence contain silent mutations. The crRNA (RHL1396) and tracrRNA were hydrated to 100 μ M with IDT Duplex Buffer, and equal volumes of each (0.61 μ l) were combined and incubated at 95°C for 5 minutes, then 25°C for 5 minutes. F₁ animals expressing the co-injection marker *egl-20::optRFP* were lysed and checked by PCR for insertions.

Non-starved healthy cultures of ALFA-tagged CHE-1 (*csu226[Ppa-che-1::2xALFA]*) (from six 6 cm plate cultures) were washed with M9 and filtered (Sartorius 84g/m2, Grade 392) and processed as previously described (39). In brief, mixed stage worms were fixed overnight at 4°C on a nutator with 500 μ l fixation buffer (4% paraformaldehyde in PBS). The worms were then incubated overnight at 37°C with 500 μ l 4% β -mercaptoethanol dissolved in 1% Triton-X100 in 0.1M Tris pH 7.4, and digested in 200 μ l of collagenase buffer with 200 units of collagenase type IV (Gibco, Thermo Fisher Scientific, Waltham, MA, USA) at 37°C for ~3.5 hours. The partially digested worms were subsequently washed three times with 500 μ l PBST. After collagenase treatment the centrifugation between the washes was done at low speed, 1000 RCF. After washes, the worms were stained in 50 μ l 1% BSA in PBST with the primary antibody (1:100 FluoTag-X2 anti-ALFA-AZdye568, N1502, NanoTag Biotechnologies, Göttingen, Germany), overnight at 4°C in a nutator. Following the washes, worms were resuspended in 50 μ l VectaShield mounting medium (H-1000, Vector Laboratories, USA) containing DAPI (Molecular Probes, Thermo Fisher Scientific) and gently mounted on freshly prepared 3% Noble agar pads. Images were acquired on a Leica DM6000 microscope.

Calcium Imaging

To conduct calcium imaging, worms were trapped and imaged within a microfluidic PDMS chip as previously described while delivering stimuli directly to the nose of an immobilized worm (46, 72). The microfluidic chip was connected to a programmable valve controller (ValveBank) that enables the user to toggle between “stimulant ON” and “stimulant OFF” states. Specifically, the ValveBank allows controlled switching of flow from the two outer buffer channels in the chip such that either the control solution (stimulant OFF) or the stimulant solution (stimulant ON) flows over the worm nose (Fig. S7). Buffer solution (for the outer two channels) consisted of M9 buffer with 0.1% Tween-20 and 1 μ g/ml fluorescein (73, 74). The worm loading solution consisted of M9 buffer with 0.1% Tween-20 and 1.5 mM tetramisole hydrochloride to immobilize the animal. Water-soluble stimulant solutions consisted of the following water-soluble compounds dissolved in nanopure/milli-Q water: ammonium chloride (NH₄Cl), ammonium iodide (NH₄I), and sodium chloride (NaCl) at concentrations of 250 mM and 25 mM. 750 mM NaCl elicited very inconsistent responses. The water-soluble stimulant solution was made by diluting the water-soluble compound in nanopure/milli-Q water with 0.1% Tween-20. The control solution consisted of 0.1% Tween-20 nanopure/milli-Q water. Worms were exposed to the control solution, stimulant solution, and then control solution using a 60-second program: 10 seconds stimulant OFF, 20 seconds stimulant ON, and 30 seconds stimulant OFF. As a negative control, animals were exposed only to the control solution, without switching channels, for the duration of the recording. Images were captured using a Zeiss Axio Observer Z1 inverted fluorescence microscope and a pco.panda 4.2 SCMOS camera. Changes in fluorescence intensity were measured in the neurons of interest while the worm was exposed to green light. Images were processed using MetaMorph software version 7.10.5.476. Images were captured at 500 ms exposure time at 2 fps because the baseline fluorescence in *Ppa-che-1p::optRCaMP* worms was too dim to capture viable data using the standard 100 ms exposure time. Baseline F₀ was measured as the average background-subtracted fluorescence from the first 9 seconds of each recording and change in fluorescence intensity was

calculated as $dF/F = (F - F_0) / F_0$, as described (74 [↗](#)). Data were analyzed and plotted using custom scripts generated in MATLAB versions R2021a–R2024a. This code is available at <https://github.com/honglabcsun/Calcium-Imaging> [↗](#).

For bar plot comparisons between wildtype and *gcy-22.3* mutants, we calculated minimum pre-stimulus and maximum post-stimulus % dF/F values using custom scripts generated in MATLAB. Min-Max data were exported as text files, manually converted and organized into an XLSX file (Microsoft Excel Office16) and imported into Prism GraphPad software (version 10) for data visualization and statistical analysis.

Nomenclature

Throughout the results section, *P. pacificus* genes will be referred to without the *Ppa*- prefix; if necessary for comparison to another species such as *C. elegans* (*Cel*-) will the *Ppa*- prefix be used.

Supplementary figures and tables

Table 1.

Comparison of ASEL/R responses between *P. pacificus* and *C. elegans*.

	<i>P. pacificus</i>		<i>C. elegans</i>	
	ASEL	ASER	ASEL	ASER
NH ₄ Cl (1 – 80mM)	ON	x	x ^{29,30}	OFF ^{29,30}
NH ₄ Cl (81 – 250mM)	ON	OFF	?	?
NaCl (1 – 80mM)	x	OFF	ON ^{29,43-45}	OFF ^{29,43-45}
NaCl (81 – 250mM)	OFF-2	OFF	ON ⁴⁴	OFF ⁴⁴
NH ₄ I (1 – 80mM)	ON	OFF	X ³⁰	OFF ³⁰
NH ₄ I (81 – 250mM)	ON	ON-OFF	?	?


“x” minimal to no response

“?” represents unknown

Figure S1.

Negative controls of ASE and AFD neuron responses to green light.

Average percent change in fluorescence (dF/F) over time (seconds) of tracked sensory neurons in *P. pacificus* (black line). Individual animals are represented by colored lines. No salt was presented; vertical lines indicate time points where salt was presented and removed for experimental samples, shown for comparison. *Ppa-che-1p::optRCaMP(csuEx93)* worms experienced the control solution for the duration of the recording and were imaged under green light. (A) ASE (n=6). (B) AFD (n=5).






Figure S2.

Individual traces and heatmaps of ASE responses to various salts.

Average percent change in fluorescence (dF/F) over time (seconds) of tracked ASEL and ASER neurons in *P. pacificus* (black line). Individual animals are represented by colored lines. For heatmaps, each row represents a single individual. Lighter colors (closer to 1) represent more positive dF/F values and darker colors (closer to 0) represent more negative dF/F values.




Figure S3.

Individual traces and heatmaps of AFD responses to various salts.

Average percent change in fluorescence (dF/F) over time (seconds) of tracked left and right AFD neurons in *P. pacificus* (black line). Individual animals are represented by colored lines. For heatmaps, each row represents a single individual. Lighter colors (closer to 1) represent more positive dF/F values and darker colors (closer to 0) represent more negative dF/F values.




Figure S4.

AFDL/R responses to different concentrations of NH₄Cl, NaCl, and NH₄I.

Average percent change in RCaMP fluorescence (dF/F) over time (seconds) as described in [Fig. 4](#). Average AFDL and AFDR neuron responses to high (250mM, red) and low (25mM, blue) concentrations of salts: (A-B) NH₄Cl, (C-D) NaCl, (E-F) NH₄I. Shaded ribbons represent 95% confidence intervals. Shaded ribbons in (D) have been cropped to maintain consistent y-axes across the plots, allowing for easier comparison. Sample sizes are indicated (n).

Figure S5.

AFDL/R responses to NH₄Cl and NaCl in wildtype and *gcy-22.3* mutants.

Average percent change in RCaMP fluorescence (dF/F) over time (seconds) as described in Fig. 4. AFDL and AFDR neuron responses to (A-B) 250mM NH₄Cl and (C-D) 250mM NaCl in wildtype (orange) compared to *gcy-22.3* mutants (magenta). Shaded ribbons represent 95% confidence intervals. Shaded ribbons in (D) have been cropped to maintain consistent y-axes across the plots. Sample sizes are indicated (n).




Figure S6

Molecular lesions

CHE-1 (PPA01143) WT	CTACGGTCTTGGGTCGGGCTCGGCCAGAA
ot5012 (4 bp insertion)	CTACGGTCTTGGGTCCTCTGGGGCTGGGCCAGAA
ot5013 (complex indel)	CTACGGTCTTGG--AACCATAGACGAACGGGTT--CCCA
CHE-1 (PPA01143) WT AA	MAYVG YD YGLG.....
ot5012 (early STOP)	MAYVG... NASDD* (frameshift +157 AA)
ot5013 (early STOP)	MAYVG... RSWTI* (frameshift +33 AA)
GCY-22.3 (PPA04454) WT	AGTGACACTGGCTCAGAAATTCGAATGGACTGATCTCACCTTTATGTTCTCTGCGGAGC
<i>csu181</i> (2 bp deletion)	AGTGACACTGGCTCAGAAATTCGAATGGACTGATCTCACCTTTT TTT--TTT GCGGAGC
GCY-22.3 Wildtype	PIILWGPPYQSTLEDSQAFPTMMSTFSARPRARAVVTLAQKFEWTDLTFFMFAERDPLV... 120
<i>csu181</i> (early STOP).	PIILWGPPYQSTLEDSQAFPTMMSTFSARPRARAVVTLAQKFEWTDLTFFFCGARSARW* 120




Figure S7.

Schematic of microfluidic apparatus to conduct calcium imaging while delivering stimuli directly to the nose of an immobilized worm. Syringes act as reservoirs for buffer solution (light blue), control solution (purple), and stimulant solution (red). Tubing inserts into the microfluidic PDMS chip. Figure panels (right) zoom in to show the microscope view of the PDMS chip: (top) fluid flow over the worm nose when stimulant is switched OFF (control solution flows over worm nose); (bottom) fluid flow over the worm nose when stimulant is switched ON (stimulant solution flows over worm nose). Created with BioRender.com

Table S1.

Nematode strains

Gene (<i>Ppa</i> -)	Allele	Strain	Genetic Lesion or Genotype
--	wildtype	PS312	<i>P. pacificus</i> California reference strain
<i>gcy-22.3</i>	<i>csu181</i>	RLH293	2 bp complex deletion causing frameshift and early STOP. 2x outcrossed.
<i>gcy-22.3p::GFP</i>	<i>csuEx90</i>	RLH334	<i>Ppa-gcy-22.3p::GFP</i> ; <i>Ppa-egl-20p::turboRFP</i> (extrachromosomal array)
<i>che-1</i>	<i>ot5012</i>	OH17879	4 bp insertion causing frameshift and STOP. 2x outcrossed.
<i>che-1</i>	<i>ot5013</i>	OH17880	8 bp complex indel causing frameshift and STOP. 2x outcrossed.
<i>che-1p::HisCl</i>	<i>csuEx83</i>	RLH336	<i>Ppa-che-1p::optHisCl</i> ; <i>Ppa-egl-20p::turboRFP</i>
<i>che-1p::RFP</i>	<i>lucEx367</i>	MLC545	<i>Ppa-che-1p::turboRFP</i> ; <i>Ppa-egl-20p::turboRFP</i>
<i>che-1::2xALFA</i>	<i>csu226</i>	RLH325	<i>Ppa-che-1::ALFA</i> C-terminal tag
<i>che-1p::RCaMP</i>	<i>csuEx93</i>	RLH335	<i>Ppa-che-1p::optRCaMP</i> ; <i>Ppa-che-1p::optGFP</i> ; <i>Ppa-egl-20p::turboRFP</i>
<i>che-1p::RCaMP</i> ; <i>gcy-22.3</i>	<i>csu183</i>	RLH300	<i>gcy-22.3(cs181)</i> ; <i>csuEx93[Ppa-che-1p::optRCaMP</i> ; <i>Ppa-che-1p::optGFP</i> ; <i>Ppa-egl-20p::turboRFP]</i>
--	wildtype	N2	<i>C. elegans</i> Bristol reference strain

Table S2.

Plasmids

Plasmid	Reporter
<i>pMM3</i>	<i>Ppa-che-1p::optGFP</i>
<i>pMM5</i>	<i>Ppa-che-1p::optRCaMP</i>
<i>pHC30</i>	<i>Ppa-che-1p::optHisCl</i>
<i>pVL2</i>	<i>Ppa-gcy-22.3p::GFP</i>

Gene (<i>Ppa</i> -)	Primer	Sequence (5'→ 3')
<i>che-1</i> (PPA01143)	SJC4	CTACGGTCTTGGGTCGGGCT (crRNA, exon 1)
<i>che-1</i>	SJC341	ATCTCTCTCCCTCATCTCCC (forward)
<i>che-1</i>	SJC342	TTCCTCCCTCTCCTGACAGC (reverse)
<i>che-1</i>	SJC343	AGACGGATGATTAGGGAAGC (forward, nested)
<i>che-1</i>	SJC344	GAAGGTGTGGGAGGGTTTCC (reverse, nested)
<i>gcy-22.3</i> (PPA04454)	RHL1553	TCTCACCTTTATGTTCTCTG (crRNA, exon 5)
2xALFA	ssDNA	TCTAGACTCGAGGAGGAGCTCAGAAGAAGACTC ACTGAGtcCCgTctcgaggaggagctcCgTCgTCgTctcac Cgag
<i>che-1</i> (C-terminal ALFA)	RHL1396	CACATGAAGACGCACAAGAA (crRNA, last exon)
<i>che-1</i>	RHL1397	AGGTCACAAGCCTTATTCGTGC (forward)
<i>che-1</i>	RHL1399	GAATCTGTTGCAGCATCTGTTGC (reverse)
<i>gcy-22.3</i>	RHL1401	CGGTCCGGCATGCAGTCAAC (forward)
<i>gcy-22.3</i>	RHL1403	GTAGTCAGGCAAAGGCTCGA (reverse)

Table S3.

Primer sequences

Acknowledgements

This research is funded by NIH SC1GM140970 to RLH, NIH R56MH096881 to SHC. OH is funded by the HHMI. MM and RLH contributed in conception, design and acquisition of work. VL, HRC, DLC, IMD, NRK, KTQ, and SJC contributed to data acquisition. RLH, OH, and SHC contributed to the analysis and writing of the work. The authors declare that they have no competing interests. We would to also thank I. Martinez and C. Igreja for technical assistance. All data needed to evaluate the conclusions in the paper are present in the paper and the Supplementary Materials.

References

1. Bargmann C., Jorgensen E. (2006) **Chemosensation in *C. elegans*** *WormBook*
2. Hong R. L., Sommer R. J. (2006) **Chemoattraction in *Pristionchus* Nematodes and Implications for Insect Recognition** *Curr Biol* **16**:2359–2365
3. Chaisson K. E., Hallem E. A. (2012) **Chemosensory behaviors of parasites** *Trends Parasitol* **28**:427–436
4. Wheeler N. J., Heimark Z. W., Airs P. M., Mann A., Bartholomay L. C., Zamanian M. (2020) **Genetic and functional diversification of chemosensory pathway receptors in mosquito-borne filarial nematodes** *Plos Biol* **18**
5. Rengarajan S., Hallem E. A. (2016) **Olfactory circuits and behaviors of nematodes** *Curr Opin Neurobiol* **41**:136–148
6. Koneru S. L., Salinas H., Flores G. E., Hong R. L. (2016) **The bacterial community of entomophilic nematodes and host beetles** *Mol. Ecol* **25**:2312–2324
7. Lo W.-S., Sommer R. J. (2022) **Vitamin B12 and predatory behavior in nematodes** *Vitam. Horm* **119**:471–489
8. Lo W.-S., Sommer R. J., Han Z. (2024) **Microbiota succession influences nematode physiology in a beetle microcosm ecosystem** *Nat. Commun* **15**
9. White J. G., Southgate E., Thomson J. N., Brenner S. (1986) **The structure of the nervous system of the nematode *Caenorhabditis elegans*** *Philosophical Transactions Royal Soc Lond B Biological Sci* **314**:1–340
10. Schafer W. (2016) **Nematode nervous systems** *Curr. Biol.* **26**:R955–R959
11. Hong R. L. *et al.* (2019) **Evolution of neuronal anatomy and circuitry in two highly divergent nematode species** *Elife* **8**
12. Ward S., Thomson N., White J. G., Brenner S. (1975) **Electron microscopical reconstruction of the anterior sensory anatomy of the nematode *Caenorhabditis elegans*** *J Comp Neurology* **160**:313–337
13. Li J., Zhu X., Ashton F. T., Gamble H. R., Schad G. A. (2001) **Sensory neuroanatomy of a passively ingested nematode parasite, *Haemonchus contortus*: amphidial neurons of the third-stage larva** *J Parasitol* **87**:65–72
14. Bumbarger D. J., Wijeratne S., Carter C., Crum J., Ellisman M. H., Baldwin J. G. (2009) **Three-dimensional reconstruction of the amphid sensilla in the microbial feeding nematode, *Acrobeles complexus* (nematoda: Rhabditida)** *J Comp Neurol* **512**:271–281
15. Zhu H., Li J., Nolan T. J., Schad G. A., Lok J. B. (2011) **Sensory neuroanatomy of *Parastrongyloides trichosuri*, a nematode parasite of mammals: Amphidial neurons of the first-stage larva** *J Comp Neurology* **519**:2493–2507

16. Loer C. M., Rivard L. (2007) **Evolution of neuronal patterning in free-living rhabditid nematodes I: Sex-specific serotonin-containing neurons** *J Comp Neurol* **502**:736–767
17. Baiocchi T., Lee G., Choe D.-H., Dillman A. R. (2017) **Host seeking parasitic nematodes use specific odors to assess host resources** *Sci Rep-uk* **7**
18. Gang S. S., Castelletto M. L., Yang E., Ruiz F., Brown T. M., Bryant A. S., Grant W. N., Hallem E. A. (2020) **Chemosensory mechanisms of host seeking and infectivity in skin-penetrating nematodes** *Proc. Natl. Acad. Sci* **117**:17913–17923
19. Bryant A. S., Ruiz F., Lee J. H., Hallem E. A. (2022) **The neural basis of heat seeking in a human-infective parasitic worm** *Curr Biol* **32**:2206–2221
20. Hallem E. A., Dillman A. R., Hong A. V., Zhang Y., Yano J. M., DeMarco S. F., Sternberg P. W. (2011) **A Sensory Code for Host Seeking in Parasitic Nematodes** *Curr Biol* **21**:377–383
21. Bargmann C. I., Horvitz H. R. (1991) **Chemosensory neurons with overlapping functions direct chemotaxis to multiple chemicals in *C. elegans*** *Neuron* **7**:729–742
22. Dusenbery, Sheridan, Russell (1975) **Chemotaxis Defective Mutants of the Nematode *Caenorhabditis elegans*** *Genetics*
23. Uchida O., Nakano H., Koga M., Ohshima Y. (2003) **The *C. elegans* che-1 gene encodes a zinc finger transcription factor required for specification of the ASE chemosensory neurons** *Development* **130**:1215–1224
24. Chang S., Johnston R. J., Hobert O. (2003) **A transcriptional regulatory cascade that controls left/right asymmetry in chemosensory neurons of *C. elegans*** *Gene Dev* **17**:2123–2137
25. Etchberger J. F., Lorch A., Sleumer M. C., Zapf R., Jones S. J., Marra M. A., Holt R. A., Moerman D. G., Hobert O. (2007) **The molecular signature and cis-regulatory architecture of a *C. elegans* gustatory neuron** *Gene Dev* **21**:1653–1674
26. Yu S., Avery L., Baude E., Garbers D. L. (1997) **Guanylyl cyclase expression in specific sensory neurons: A new family of chemosensory receptors** *Proc National Acad Sci* **94**:3384–3387
27. Pierce-Shimomura J. T., Faumont S., Gaston M. R., Pearson B. J., Lockery S. R. (2001) **The homeobox gene *lim-6* is required for distinct chemosensory representations in *C. elegans*** *Nature* **410**:694–698
28. Chang S., Johnston R. J., Frøkjær-Jensen C., Lockery S., Hobert O. (2004) **MicroRNAs act sequentially and asymmetrically to control chemosensory laterality in the nematode** *Nature* **430**:785–789
29. Suzuki H., Thiele T. R., Faumont S., Ezcurra M., Lockery S. R., Schafer W. R. (2008) **Functional asymmetry in *Caenorhabditis elegans* taste neurons and its computational role in chemotaxis** *Nature* **454**:114–117
30. Ortiz C. O. *et al.* (2009) **Lateralized Gustatory Behavior of *C. elegans* Is Controlled by Specific Receptor-Type Guanylyl Cyclases** *Curr Biol* **19**:996–1004
31. Hobert O. (2014) **Development of left/right asymmetry in the *Caenorhabditis elegans* nervous system: From zygote to postmitotic neuron** *genesis* **52**:528–543

32. Johnston R., Hobert O. (2003) **A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*** *Nature* **426**:845–849
33. Cochella L., Hobert O. (2012) **Embryonic Priming of a miRNA Locus Predetermines Postmitotic Neuronal Left/Right Asymmetry in *C. elegans*** *Cell* **151**:1229–1242
34. Ahmed R., Chang Z., Younis A. E., Langnick C., Li N., Chen W., Brattig N., Dieterich C. (2013) **Conserved miRNAs Are Candidate Post-Transcriptional Regulators of Developmental Arrest in Free-Living and Parasitic Nematodes** *Genome Biol Evol* **5**:1246–1260
35. Ward S. (1973) **Chemotaxis by the Nematode *Caenorhabditis elegans*: Identification of Attractants and Analysis of the Response by Use of Mutants** *Proc. Natl. Acad. Sci* **70**:817–821
36. Frøkjær-Jensen C., Ailion M., Lockery S. R. (2008) **Ammonium-Acetate Is Sensed by Gustatory and Olfactory Neurons in *Caenorhabditis elegans*** *Plos One* **3**
37. Bargmann C. I., Hartwig E., Horvitz H. R. (1993) **Odorant-selective genes and neurons mediate olfaction in *C. elegans*** *Cell* **74**:515–527
38. Etchberger J. F., Flowers E. B., Poole R. J., Bashllari E., Hobert O. (2009) **Cis-regulatory mechanisms of left/right asymmetric neuron-subtype specification in *C. elegans*** *Development* **136**:147–160
39. Igreja C., Loschko T., Schäfer A., Sharma R., Quiobe S. P., Aloschy E., Witte H., Sommer R. J. (2022) **Application of ALFA-Tagging in the Nematode Model Organisms *Caenorhabditis elegans* and *Pristionchus pacificus*** *Cells* **11**
40. Kato S., Xu Y., Cho C. E., Abbott L. F., Bargmann C. I. (2014) **Temporal Responses of *C. elegans* Chemosensory Neurons Are Preserved in Behavioral Dynamics** *Neuron* **81**:616–628
41. Wang W., Xu Z.-J., Wu Y.-Q., Qin L.-W., Li Z.-Y., Wu Z.-X. (2015) **Off-response in ASH neurons evoked by CuSO₄ requires the TRP channel OSM-9 in *Caenorhabditis elegans*** *Biochem. Biophys. Res. Commun* **461**:463–468
42. Ortiz C. O., Etchberger J. F., Posy S. L., Frøkjær-Jensen C., Lockery S., Honig B., Hobert O. (2006) **Searching for Neuronal Left/Right Asymmetry: Genomewide Analysis of Nematode Receptor-Type Guanylyl Cyclases** *Genetics* **173**:131–149
43. Rabinowitch I., Chatzigeorgiou M., Zhao B., Treinin M., Schafer W. R. (2014) **Rewiring neural circuits by the insertion of ectopic electrical synapses in transgenic *C. elegans*** *Nat Commun* **5**
44. Shindou T., Ochi-Shindou M., Murayama T., Saita E., Momohara Y., Wickens J. R., Maruyama I. N. (2019) **Active propagation of dendritic electrical signals in *C. elegans*** *Sci. Rep* **9**
45. Watteyne J. *et al.* (2020) **Neuromedin U signaling regulates retrieval of learned salt avoidance in a *C. elegans* gustatory circuit** *Nat. Commun* **11**
46. Chronis N., Zimmer M., Bargmann C. I. (2007) **Microfluidics for in vivo imaging of neuronal and behavioral activity in *Caenorhabditis elegans*** *Nat. Methods* **4**:727–731
47. Yoshida K., Hirotsu T., Tagawa T., Oda S., Wakabayashi T., Iino Y., Ishihara T. (2012) **Odour concentration-dependent olfactory preference change in *C. elegans*** *Nat Commun* **3**

48. Mori I., Ohshima Y. (1995) **Neural regulation of thermotaxis in *Caenorhabditis elegans*** *Nature* **376**:344–348
49. Sengupta P., Chou J. H., Bargmann C. I. (1996) **odr-10 Encodes a Seven Transmembrane Domain Olfactory Receptor Required for Responses to the Odorant Diacetyl** *Cell* **84**:899–909
50. Macosko E. Z., Pokala N., Feinberg E. H., Chalasani S. H., Butcher R. A., Clardy J., Bargmann C. I. (2009) **A hub-and-spoke circuit drives pheromone attraction and social behaviour in *C. elegans*** *Nature* **458**:1171–1175
51. Liu J. *et al.* (2010) ***C. elegans* phototransduction requires a G protein-dependent cGMP pathway and a taste receptor homolog** *Nat. Neurosci* **13**:715–722
52. Leinwand S. G., Chalasani S. H. (2013) **Neuropeptide signaling remodels chemosensory circuit composition in *Caenorhabditis elegans*** *Nat. Neurosci* **16**:1461–1467
53. Lin A. *et al.* (2023) **Functional imaging and quantification of multineuronal olfactory responses in *C. elegans*** *Sci Adv* **9**
54. Yemini E., Lin A., Nejatbakhsh A., Varol E., Sun R., Mena G. E., Samuel A. D. T., Paninski L., Venkatachalam V., Hobert O. (2021) **NeuroPAL: A Multicolor Atlas for Whole-Brain Neuronal Identification in *C. elegans*** *Cell* **184**:272–288
55. Zaslaver A., Liani I., Shtangel O., Ginzburg S., Yee L., Sternberg P. W. (2015) **Hierarchical sparse coding in the sensory system of *Caenorhabditis elegans*** *Proc. Natl. Acad. Sci* **112**:1185–1189
56. Kuhara A., Okumura M., Kimata T., Tanizawa Y., Takano R., Kimura K. D., Inada H., Matsumoto K., Mori I. (2008) **Temperature Sensing by an Olfactory Neuron in a Circuit Controlling Behavior of *C. elegans*** *Science* **320**:803–807
57. Leaver M., Kienle S., Begasse M. L., Sommer R. J., Hyman A. A. (2016) **A locus in *Pristionchus pacificus* that is responsible for the ability to give rise to fertile offspring at higher temperatures** *Biol Open* **5**:1111–1117
58. Sieriebriennikov B., Markov G. V., Witte H., Sommer R. J. (2017) **The Role of DAF-21/Hsp90 in Mouth-Form Plasticity in *Pristionchus pacificus*** *Mol Biol Evol* **34**:1644–1653
59. Leaver M., Moreno E., Kayhan M., McGaughan A., Rödelberger C., Sommer R. J., Hyman A. A. (2022) **Adaptation to environmental temperature in divergent clades of the nematode *Pristionchus pacificus*** *Evolution* **76**:1660–1673
60. White J. Q., Nicholas T. J., Gritton J., Truong L., Davidson E. R., Jorgensen E. M. (2007) **The Sensory Circuitry for Sexual Attraction in *C. elegans* Males** *Curr. Biol* **17**:1847–1857
61. Johnston R. J., Chang S., Etchberger J. F., Ortiz C. O., Hobert O. (2005) **MicroRNAs acting in a double-negative feedback loop to control a neuronal cell fate decision** *Proc National Acad Sci* **102**:12449–12454
62. Sharma D. R., Röseler W., Witte H., Werner M. S., Sommer R. J. (2024) **Developmental small RNA transcriptomics reveals divergent evolution of the conserved microRNA miR-100 and the let-7-complex in nematodes** *bioRxiv*

63. Cinkornpumin J. K., Wisidagama D. R., Rapoport V., Go J. L., Dieterich C., Wang X., Sommer R. J., Hong R. L. (2014) **A host beetle pheromone regulates development and behavior in the nematode *Pristionchus pacificus*** *Elife* **3**
64. Brenner S. (1974) **The Genetics of *Caenorhabditis elegans*** *Genetics* **77**:71–94
65. Pokala N., Liu Q., Gordus A., Bargmann C. I. (2014) **Inducible and titratable silencing of *Caenorhabditis elegans* neurons in vivo with histamine-gated chloride channels** *Proc National Acad Sci* **111**:2770–2775
66. Pokala N., Flavell S. W., elegans C. (2022) **Methods and Applications** *Methods Mol Biology* **2468**:357–373
67. Kerr R., Lev-Ram V., Baird G., Vincent P., Tsien R. Y., Schafer W. R. (2000) **Optical Imaging of Calcium Transients in Neurons and Pharyngeal Muscle of *C. elegans*** *Neuron* **26**:583–594
68. Nakai J., Ohkura M., Imoto K. (2001) **A high signal-to-noise Ca²⁺ probe composed of a single green fluorescent protein** *Nat. Biotechnol* **19**:137–141
69. Dana H. *et al.* (2016) **Sensitive red protein calcium indicators for imaging neural activity** *eLife* **5**
70. Han Z., Lo W.-S., Lightfoot J. W., Witte H., Sun S., Sommer R. J. (2020) **Improving Transgenesis Efficiency and CRISPR-Associated Tools Through Codon Optimization and Native Intron Addition in *Pristionchus* Nematodes** *Genetics* **216**
71. Nakayama K., Ishita Y., Chihara T., Okumura M. (2020) **Screening for CRISPR/Cas9-induced mutations using a co-injection marker in the nematode *Pristionchus pacificus*** *Dev Genes Evol* :1–8
72. Chalasani S. H., Chronis N., Tsunozaki M., Gray J. M., Ramot D., Goodman M. B., Bargmann C. I. (2007) **Dissecting a circuit for olfactory behaviour in *Caenorhabditis elegans*** *Nature* **450**:63–70
73. Stiernagle T. (2006) **Maintenance of *C. elegans*** *WormBook: online Rev. C elegans Biol* :1–11
74. Liu Z. *et al.* (2018) **Predator-secreted sulfolipids induce defensive responses in *C. elegans*** *Nat Commun* **9**

Author information

Marisa Mackie

Department of Biology, California State University, Northridge, United States

Vivian Vy Le

Department of Biology, California State University, Northridge, United States

Heather R Carstensen

Department of Biology, California State University, Northridge, United States

Nicole R Kushnir

Department of Biology, California State University, Northridge, United States

Dylan L Castro

Department of Biology, California State University, Northridge, United States

Ivan M Dimov

Department of Biology, California State University, Northridge, United States

Kathleen T Quach

Molecular Neurobiology Laboratory, Salk Institute for Biological Studies, La Jolla, United States

Steven J Cook*

Department of Biological Sciences, Howard Hughes Medical Institute, Columbia University, New York, United States

*Present address: Neural Coding Department Allen Institute for Brain Science, Seattle, USA

Oliver Hobert

Department of Biological Sciences, Howard Hughes Medical Institute, Columbia University, New York, United States

ORCID iD: [0000-0002-7634-2854](https://orcid.org/0000-0002-7634-2854)

Sreekanth H Chalasani

Molecular Neurobiology Laboratory, Salk Institute for Biological Studies, La Jolla, United States

Ray L Hong

Department of Biology, California State University, Northridge, United States

ORCID iD: [0000-0003-1870-8659](https://orcid.org/0000-0003-1870-8659)

For correspondence: ray.hong@csun.edu

Editors

Reviewing Editor

Albert Cardona

University of Cambridge, Cambridge, United Kingdom

Senior Editor

Albert Cardona

University of Cambridge, Cambridge, United Kingdom

Reviewer #1 (Public review):

Summary:

Mackie and colleagues compare chemosensory preferences between *C. elegans* and *P. pacificus*, and the cellular and molecular mechanisms underlying them. The nematodes have overlapping and distinct preferences for different salts. Although *P. pacificus* lacks the *lsy-6*

miRNA important for establishing asymmetry of the left/right ASE salt-sensing neurons in *C. elegans*, the authors find that *P. pacificus* ASE homologs achieve molecular (receptor expression) and functional (calcium response) asymmetry by alternative means. This work contributes an important comparison of how these two nematodes sense salts and highlights that evolution can find different ways to establish asymmetry in small nervous systems to optimize the processing of chemosensory cues in the environment.

Strengths:

The authors use clear and established methods to record the response of neurons to chemosensory cues. They were able to show clearly that ASEL/R are functionally asymmetric in *P. pacificus*, and combined with genetic perturbation establish a role for *che-1*-dependent *gcy-22.3* in the asymmetric response to NH_4Cl .

Weaknesses:

The mechanism of *lsy-6*-independent establishment of ASEL/R asymmetry in *P. pacificus* remains uncharacterized.

<https://doi.org/10.7554/eLife.103796.1.sa2>

Reviewer #2 (Public review):

Summary:

In this manuscript, Mackie et al. investigate gustatory behavior and the neural basis of gustation in the predatory nematode *Pristionchus pacificus*. First, they show that the behavioral preferences of *P. pacificus* for gustatory cues differ from those reported for *C. elegans*. Next, they investigate the molecular mechanisms of salt sensing in *P. pacificus*. They show that although the *C. elegans* transcription factor gene *che-1* is expressed specifically in the ASE neurons, the *P. pacificus* *che-1* gene is expressed in the Ppa-ASE and Ppa-AFD neurons. Moreover, *che-1* plays a less critical role in salt chemotaxis in *P. pacificus* than *C. elegans*. Chemogenetic silencing of Ppa-ASE and Ppa-AFD neurons results in more severe chemotaxis defects. The authors then use calcium imaging to show that both Ppa-ASE and Ppa-AFD neurons respond to salt stimuli. Calcium imaging experiments also reveal that the left and right Ppa-ASE neurons respond differently to salts, despite the fact that *P. pacificus* lacks *lsy-6*, a microRNA that is important for ASE left/right asymmetry in *C. elegans*. Finally, the authors show that the receptor guanylate cyclase gene *Ppa-gcy-23.3* is expressed in the right Ppa-ASE neuron (Ppa-ASER) but not the left Ppa-ASE neuron (Ppa-ASEL) and is required for some of the gustatory responses of Ppa-ASER, further confirming that the Ppa-ASE neurons are asymmetric and suggesting that Ppa-GCY-23.3 is a gustatory receptor. Overall, this work provides insight into the evolution of gustation across nematode species. It illustrates how sensory neuron response properties and molecular mechanisms of cell fate determination can evolve to mediate species-specific behaviors. However, the paper would be greatly strengthened by a direct comparison of calcium responses to gustatory cues in *C. elegans* and *P. pacificus*, since the comparison currently relies entirely on published data for *C. elegans*, where the imaging parameters likely differ. In addition, the conclusions regarding Ppa-AFD neuron function would benefit from additional confirmation of AFD neuron identity. Finally, how prior salt exposure influences gustatory behavior and neural activity in *P. pacificus* is not discussed.

Strengths:

(1) This study provides exciting new insights into how gustatory behaviors and mechanisms differ in nematode species with different lifestyles and ecological niches. The results from salt chemotaxis experiments suggest that *P. pacificus* shows distinct gustatory preferences from *C.*

C. elegans. Calcium imaging from Ppa-ASE neurons suggests that the response properties of the ASE neurons differ between the two species. In addition, an analysis of the expression and function of the transcription factor Ppa-che-1 reveals that mechanisms of ASE cell fate determination differ in *C. elegans* and *P. pacificus*, although the ASE neurons play a critical role in salt sensing in both species. Thus, the authors identify several differences in gustatory system development and function across nematode species.

(2) This is the first calcium imaging study of *P. pacificus*, and it offers some of the first insights into the evolution of gustatory neuron function across nematode species.

(3) This study addresses the mechanisms that lead to left/right asymmetry in nematodes. It reveals that the ASER and ASEL neurons differ in their response properties, but this asymmetry is achieved by molecular mechanisms that are at least partly distinct from those that operate in *C. elegans*. Notably, ASEL/R asymmetry in *P. pacificus* is achieved despite the lack of a *P. pacificus* *lsy-6* homolog.

Weaknesses:

(1) The authors observe only weak attraction of *C. elegans* to NaCl. These results raise the question of whether the weak attraction observed is the result of the prior salt environment experienced by the worms. More generally, this study does not address how prior exposure to gustatory cues shapes gustatory responses in *P. pacificus*. Is salt sensing in *P. pacificus* subject to the same type of experience-dependent modulation as salt sensing in *C. elegans*?

(2) A key finding of this paper is that the Ppa-CHE-1 transcription factor is expressed in the Ppa-AFD neurons as well as the Ppa-ASE neurons, despite the fact that Ce-CHE-1 is expressed specifically in Ce-ASE. However, additional verification of Ppa-AFD neuron identity is required. Based on the image shown in the manuscript, it is difficult to unequivocally identify the second pair of CHE-1-positive head neurons as the Ppa-AFD neurons. Ppa-AFD neuron identity could be verified by confocal imaging of the CHE-1-positive neurons, co-expression of Ppa-che-1p::GFP with a likely AFD reporter, thermotaxis assays with Ppa-che-1 mutants, and/or calcium imaging from the putative Ppa-AFD neurons.

(3) Loss of Ppa-che-1 causes a less severe phenotype than loss of Ce-che-1. However, the loss of Ppa-che-1::RFP expression in ASE but not AFD raises the question of whether there might be additional start sites in the Ppa-che-1 gene downstream of the mutation sites. It would be helpful to know whether there are multiple isoforms of Ppa-che-1, and if so, whether the exon with the introduced frameshift is present in all isoforms and results in complete loss of Ppa-CHE-1 protein.

(4) The authors show that silencing Ppa-ASE has a dramatic effect on salt chemotaxis behavior. However, these data lack control with histamine-treated wild-type animals, with the result that the phenotype of Ppa-ASE-silenced animals could result from exposure to histamine dihydrochloride. This is an especially important control in the context of salt sensing, where histamine dihydrochloride could alter behavioral responses to other salts.

(5) The calcium imaging data in the paper suggest that the Ppa-ASE and Ce-ASE neurons respond differently to salt solutions. However, to make this point, a direct comparison of calcium responses in *C. elegans* and *P. pacificus* using the same calcium indicator is required. By relying on previously published *C. elegans* data, it is difficult to know how differences in growth conditions or imaging conditions affect ASE responses. In addition, the paper would be strengthened by additional quantitative analysis of the calcium imaging data. For example, the paper states that 25 mM NH₄Cl evokes a greater response in ASEL than 250 mM NH₄Cl, but a quantitative comparison of the maximum responses to the two stimuli is not shown.

(6) It would be helpful to examine, or at least discuss, the other *P. pacificus* paralogs of Ce-gcy-22. Are they expressed in Ppa-ASER? How similar are the different paralogs? Additional discussion of the Ppa-gcy-22 gene expansion in *P. pacificus* would be especially helpful with respect to understanding the relatively minor phenotype of the Ppa-gcy-22.3 mutants.

(7) The calcium imaging data from Ppa-ASE is quite variable. It would be helpful to discuss this variability. It would also be helpful to clarify how the ASEL and ASER neurons are being conclusively identified during calcium imaging.

(8) More information about how the animals were treated prior to calcium imaging would be helpful. In particular, were they exposed to salt solutions prior to imaging? In addition, the animals are in an M9 buffer during imaging - does this affect calcium responses in Ppa-ASE and Ppa-AFD? More information about salt exposure, and how this affects neuron responses, would be very helpful.

(9) In Figure 6, the authors say that Ppa-gcy-22.3::GFP expression is absent in the Ppa-che-1(ot5012) mutant. However, based on the figure, it looks like there is some expression remaining. Is there a residual expression of Ppa-gcy-22.3::GFP in ASE or possibly ectopic expression in AFD? Does Ppa-che-1 regulate rGC expression in AFD? It would be helpful to address the role of Ppa-che-1 in AFD neuron differentiation.

<https://doi.org/10.7554/eLife.103796.1.sa1>

Author response:

We very much appreciate the reviewers' and editor's overall positive responses to our manuscript "Evolution of lateralized gustation in nematodes".

Reviewer #1:

The mechanism of lsy-6-independent establishment of ASEL/R asymmetry in P. pacificus remains uncharacterized.

We thank the reviewer for recognizing the novel contributions of our work in revealing the existence of alternative pathways for establishing neuronal lateral asymmetry despite the absence of the *lsy-6* miRNA in a divergent nematode species. We are certainly encouraged now to search for genetic factors that abolish asymmetric expression of *gcy-22.3*.

Reviewer #2:

(1) The authors observe only weak attraction of C. elegans to NaCl. These results raise the question of whether the weak attraction observed is the result of the prior salt environment experienced by the worms. More generally, this study does not address how prior exposure to gustatory cues shapes gustatory responses in P. pacificus. Is salt sensing in P. pacificus subject to the same type of experience-dependent modulation as salt sensing in C. elegans?

Proposed revision: For our live imaging experiments, we had not considered if starved *P. pacificus* animals in the presence of salt may exhibit responses different from a well-fed state. However, we will venture to address the effect of experience-dependent modulation in *P. pacificus* chemotaxis behavior using NH₄Cl.

(2) A key finding of this paper is that the Ppa-CHE-1 transcription factor is expressed in the Ppa-AFD neurons as well as the Ppa-ASE neurons, despite the fact that Ce-CHE-1 is

expressed specifically in Ce-ASE. However, additional verification of Ppa-AFD neuron identity is required. Based on the image shown in the manuscript, it is difficult to unequivocally identify the second pair of CHE-1-positive head neurons as the Ppa-AFD neurons. Ppa-AFD neuron identity could be verified by confocal imaging of the CHE-1-positive neurons, co-expression of Ppa-che-1p::GFP with a likely AFD reporter, thermotaxis assays with Ppa-che-1 mutants, and/or calcium imaging from the putative Ppa-AFD neurons.

We are happy to provide additional evidence to confirm Ppa-AFD neuron identity since the expression of Ppa-CHE-1 in non-ASE amphid neurons is one of the major differences between the two nematode species

Proposed revision: We will provide results showing the *Ppa-ttx-1::gfp* reporter expression in finger-like neuronal endings and *Ppa-TTX-1::ALFA* co-localization with *Ppa-che-1::gfp* in the putative AFD neurons and discuss the possible role of Ppa-CHE-1 in AFD differentiation. We attempted to obtain AFD markers using several reporter strains. However, *Ppa-gcy-8.1p::gfp(csuEx101)* (PPA24212) showed no expression while *Ppa-gcy-8.2p::gfp(csuEx100)* (PPA41407) showed only expression in pharyngeal cells.

(4) The authors show that silencing Ppa-ASE has a dramatic effect on salt chemotaxis behavior. However, these data lack control with histamine-treated wild-type animals, with the result that the phenotype of Ppa-ASE-silenced animals could result from exposure to histamine dihydrochloride. This is an especially important control in the context of salt sensing, where histamine dihydrochloride could alter behavioral responses to other salts.

Proposed revision: Thank you for noticing this oversight. The control for histamine-treated wild-type worms in the Ppa-ASE silencing experiments was inadvertently left out in the original submission. Because the HisCl transgene is on a randomly segregating transgene array, we have scored worms with and without the transgene expressing the co-injection marker (*Ppa-egl-20p::rfp* expressed in the tail) to show that the presence of the transgene is necessary for the knockdown of NH₄Br attraction.

We will also address most of the other more minor suggestions and clarifications sought by the reviewers.

<https://doi.org/10.7554/eLife.103796.1.sa0>