

Newly discovered and conserved role of IgM against viral infection in an early vertebrate

Reviewed Preprint

v2 • August 12, 2025

Revised by authors

Reviewed Preprint

v1 • February 17, 2025

Weiguang Kong, Xinyou Wang, Guangyi Ding, Peng Yang, Yong Shi, Chang Cai, Xinyi Yang, Gaofeng Cheng, Fumio Takizawa, Zhen Xu ✉

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China • Faculty of Marine Science and Technology, Fukui Prefectural University, Fukui, Japan

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

The manuscript is an **important** study which aims to demonstrate the conserved and crucial role of IgM in both systemic and mucosal antiviral immunity in teleost, challenging the established differential roles of IgT and IgM. The strength of the evidence is **solid** and supported by a combination of in vivo studies, viral infection models, and complementary in vitro assays. In the current version, authors validate the MoAb against IgM

<https://doi.org/10.7554/eLife.104465.2.sa2>

Abstract

IgM emerged in jawed vertebrates 500 Mya and remains the most evolutionarily conserved antibody class. However, despite extensive studies on IgM as an ancient antiviral weapon in warm-blooded vertebrates, its role and mechanisms in combating viral infections in early vertebrates remain poorly understood. Here, we found that significant virus-specific sIgM titers were generated in serum and gut mucus of teleost fish that survived infection, and fish lacking sIgM were more susceptible to viral infection. These results challenge the paradigm that IgM and IgT are specialized to systemic and mucosal immunity, respectively. More crucially, our study provides further insight into the role of sIgM in viral neutralization and firstly clarified the mechanism through which teleost sIgM blocks viral infection by directly targeting viral particles. From an evolutionary perspective, our findings indicate that sIgM in both primitive and modern vertebrates follow conserved principles in the development of specialized antiviral immunity.

Introduction

The origin of immunoglobulins (Ig) in jawed vertebrates can be traced back approximately 500 million years ago (Mya), marking a significant milestone in the evolution of the adaptive immune system in vertebrates (Parra et al., 2013; Hirano et al., 2011). Igs are highly specialized recognition glycoproteins that play a vital role in adaptive immune responses (Bilal et al., 2004). All jawed vertebrates, including mammals, birds, reptiles, amphibians, bony fish, and cartilaginous fish, express multiple classes of Igs (Parra et al., 2021; Stavnezer and Amemiya, 2004; Han et al., 2016). However, the specific isotypes vary across taxa due to evolutionary and selective pressures (Figure 1–figure supplement 1) (Stavnezer and Amemiya, 2004; Han et al., 2016). Agnathans, the most ancient vertebrate lineage, do not possess *bona fide* Ig but have variable lymphocyte receptors (VLRs) capable of mediating adaptive immune responses (Flajnik, 2018). Cartilaginous fish are the earliest vertebrates, from a phylogenetic perspective, to have evolved an adaptive immune system based on Igs. To date, three Ig isotypes have been identified in Chondrichthyes: IgM, IgW (which is homologous to IgD) (Xu et al., 2016; Edholm et al., 2010), and IgNAR (a lineage-specific isotype unique to cartilaginous fish) (Dooley, 2018). IgT is exclusively expressed in teleost fish and represents the most primitive Ig isotype with specialized in mucosal immune response (Zhang et al., 2010). Notably, while IgT has been identified in the majority of teleost species, genomic analyses reveal its absence in some species, such as medaka (*Oryzias latipes*), channel catfish (*Ictalurus punctatus*), Atlantic cod (*Gadus morhua*), and turquoise killifish (*Nothobranchius furzeri*) (Bengtén et al., 2002; Bradshaw et al., 2020; Magadán-Mompó et al., 2011; Györkei et al., 2024). The prevailing perspective is that avian and mammalian IgA evolved from a precursor of amphibian IgX, whereas mammalian IgG and IgE evolved from a precursor of amphibian IgY (Schaerlinger et al., 2008). For decades, IgD was thought to be sporadically present in mammals and absent in birds (Flajnik, 2002; Wilson et al., 1997). However, the discovery of IgD in an ostrich species (*Struthio camelus*) confirmed its presence across all classes of jawed vertebrates, fundamentally reshaping our understanding of Ig isotype evolution (Flajnik and Kasahara, 2010). Similar to IgD, IgM is considered one of the most primordial Ig isotypes throughout evolution, as it is present in all extant gnathostomes, with the potential unique exception of the coelacanth (Keyt et al., 2004). Compared to other Ig isotypes, an exceptionally conserved feature of IgM is its ability to form polymeric structures, which are particularly suited for binding to viral surface proteins (Boes, 2000). Interestingly, the assembly mechanism of IgM exhibits significant evolutionary variation across vertebrate lineages. In cartilaginous fishes and tetrapods, IgM is secreted as a J chain-linked pentamer, which may enhance multivalent antigen recognition (Hagiwara et al., 1985; Hohman et al., 2003). By contrast, teleosts have undergone J chain gene loss, resulting in the stable of tetrameric IgM formation (Bromage et al., 2004). Given the unique and intriguing features of secreted IgM (sIgM), immunologists are becoming increasingly interested in studying its functions (Blandino and Baumgarth, 2019).

Despite differences in its assembly mechanisms and the degree of polymerization of its secretory form among various vertebrate groups, IgM is known for its remarkable functional stability and conservation. It is the largest Ig classes in plasma and the first antibody produced during an antiviral humoral response (Schroeder and Cavacini, 2010). Although previous studies have suggested that IgM is involved in maintaining mucosal tissue homeostasis, it is generally considered to be a major Ig isotype that plays a key role in the antiviral systemic immune response (Ochsenbein et al., 1999; Boes et al., 1998). Indeed, IgM serves not only as a crucial effector molecule in the initial humoral defense against pathogens but may also contribute to mucosal protection during the early stages of life (Chen et al., 2020). Studies in tetrapods have unveiled the pivotal role and conserved properties of IgM in combating viral infections (Sun and Zhao, 2020). According to commonly held views, IgM is rapidly produced early in the humoral immune response against viral infections, providing fast and strong protective immunity (Ouchida et al.,




Figure 1.

Infection model of largemouth bass with LMBV.

(A) Fish were administered an intraperitoneal injection of 100 μ L ($1D \times 10^6$ TCID₅₀) of the virus solution and then sacrificed at intervals of 1, 4, 7, 14, 21, and 28 days post-infection (DPI) for tissue sampling. The timeline uses a red line to indicate fish that died or diseased symptoms of viral infection, while a green line represents fish that remained healthy, showing normal behavior and no signs of viral infection. (B) Clinical observations following infection with LMBV. The black and red arrows indicate skin ulceration and subdermal hemorrhage of infected fish, respectively. (C, D) Spleen size (C) and spleen/body weight ratio (D) following LMBV infection ($n = 9$). (E) Cumulative survival of LMBV-infected and control fish. (F) LMBV-MCP gene copies (Log₁₀) were measured via qPCR in head kidney (HK), spleen (SP), and gut ($n = 6$). (G) Immunofluorescence staining of LMBV in paraffin sections of the HK, SP, and gut from fish infected for 4 days and control fish ($n = 6$). LMBV was stained using an anti-LMBV-MCP monoclonal antibody (red). DAPI stains the cell nuclei blue. Scale bars, 50 μ m. (H) The virus particles in the HK, SP, and gut were analyzed by transmission electron microscopy (TEM). Scale bars, 500nm. (I) Cytopathic effect (CPE) of LMBV on EPC cells following exposure to supernatant of HK, SP, and gut homogenates from 4 days infected fish. Scale bars, 100 μ m. (J) Histological examination of HK, SP, and gut from LMBV-infected and control fish at 4 DPI ($n = 6$). The red arrow indicates erythrocyte infiltration. The black asterisk indicates cellular necrosis in HK and SP. The black arrow indicates epithelial cell shedding in gut. Scale bars, 50 μ m. Data are shown as mean $\pm$ CSEM. An unpaired Student's t -test was used. *** $p < 0.001$.

2012). For certain viruses predominantly controlled by B cells, such as polyomavirus and vesicular stomatitis virus (VSV), IgM provides sufficient protection against fatal infection during the acute phase (Beebe and Cooper, 1981; Nguyen et al., 2023). However, recent studies in mice revealed the presence of long-lived, antigen-specific IgM plasma cells (Bohannon et al., 2016). Neutralizing antibodies (NABs) secreted by plasma cells are the fundamental molecular armament of antiviral humoral immunity, serving as the primary mechanism through which Igs protect the body from viral infection (Ali et al., 2020). Nevertheless, there is little evidence supporting the neutralizing capacity of IgM in mucosal tissues. New research on SARS-CoV-2 provides compelling evidence that plasma from convalescent individuals also exhibits IgM-mediated neutralization activity, with neutralization titers showing a strong correlation with IgM levels (Ku et al., 2021). Although sIgM functions have been extensively studied in warm-blooded animals, less is known regarding sIgM function in primitive vertebrates. The function of IgM from teleost fish is of particular interest, considering these fish represent an early evolutionary stage in the development of vertebrate adaptive immunity (Taylor and Dimmock, 1985). Recently, we demonstrated that IgM is the predominant Ig isotype involved in the immune response to pathogens in systemic compartments (e.g., serum, spleen, and head kidney), where significant accumulations of IgM⁺ B cells were found after infection with parasitic and bacterial pathogens (Xu et al., 2016; Xu et al., 2013; Yu et al., 2018). Notably, the serum of fish that survive parasitic or bacterial infection harbors high levels of pathogen-specific IgM, coupled with negligible IgT-specific titers and no specific IgD. In contrast, IgT is the main Ig response in the mucus of the same fish. Additionally, we previously reported that induced viral-specific sIgT within the mucus is capable of neutralizing viruses, with fish lacking IgT being significantly more susceptible to viruses compared to those with IgT, suggesting that sIgT is essential for the control of mucosal pathogens (Yu et al., 2022). Interestingly, recent observations show that specific IgM levels were significantly increased in both serum and gut mucus upon *Aeromonas hydrophila* and spring viremia of carp (SVCV) infection, challenging the paradigm that pathogen-specific IgT and IgM responses are strictly compartmentalized in mucosal and systemic immunity, respectively (Mu et al., 2022; Yu et al., 2024). Previous studies have demonstrated the capacity of teleost fish to produce specific serum and mucus sIgM recognizing viral antigens in response to viruses across a wide range of teleost species (Mashoof and Criscitiello, 2016). However, the precise contribution and mechanisms through which teleost IgM provides protection against viral infection remain poorly understood. Since IgM is the most abundant Ig class in fish body fluids, we hypothesized that teleost sIgM, similar to mammalian sIgM, plays a role in viral neutralization across both primitive and modern vertebrates.

To verify the aforementioned hypothesis, we established a viral infection model by intraperitoneally injecting largemouth bass (*Micropterus salmoides*) with largemouth bass virus (LMBV), which induces typical symptoms and severe pathological changes, evoking a robust immune response in both systemic and mucosal areas. Our data revealed significant proliferation and accumulation of IgM⁺ B cells in the gut and head kidney, accompanied by increased viral-specific sIgM titers in serum and mucus, reinforcing the notion that sIgM participates in both mucosal and systemic immunity. To further validate the role of sIgM in antiviral defense, we employed a IgM depletion model in teleost fish and found that fish devoid of sIgM are more susceptible to the virus than control fish. Importantly, our study demonstrates that induced virus-specific sIgM can neutralize viruses *in vitro*, thereby revealing an effector role of sIgM in both serum and gut mucus against viral infection. More importantly, our work reveals that teleost sIgM may neutralize viruses by directly acting on viral particles to block infection. Collectively, our data provide proof-of-concept that IgM serves as an ancient weapon in the adaptive immune response against viral infections, preserved throughout evolution across all vertebrates.

Results

Establishment of LMBV infection model and immune responses in largemouth bass upon LMBV infection

To investigate the dynamic immune response of largemouth bass to viral infection, a viral infection model was established using LMBV. The fish were infected by intraperitoneal injection with LMBV (100 μ L, 1×10^6 TCID₅₀), and tissue samples were collected at 0, 1, 4, 7, 14, 21, and 28 days post-infection (DPI) (**Figure 1A**). The infected fish exhibited typical symptoms, such as skin ulceration, subdermal hemorrhage, and spleen enlargement, at 4 DPI (**Figure 1B – 1D**), reaching a cumulative mortality of 57% within 14 days (**Figure 1E**). qPCR analyses revealed the presence of high LMBV loads in the spleen (SP), head kidney (HK), and gut, especially at 4 DPI (**Figure 1F**). Immunofluorescence microscopy studies using an anti-LMBV-MCP monoclonal antibody (mAb) indicated that the virus was present in the SP, HK, and gut of the infected fish at 4 DPI (**Figure 1G**), with isotype-matched control antibodies shown in **Figure 1-figure supplement 2**. Transmission electron microscopy (TEM) images also showed the accumulation of numerous LMBV particles in the HK, SP, and gut (**Figure 1H**), indicating successful viral invasion of these tissues. Importantly, the homogenate supernatant from SP, HK, or gut of infected fish at 4 DPI resulted in a notable cytopathic effect (CPE) in cell cultures (**Figure 1I**). Moreover, histological analysis revealed that largemouth bass occurred varying degrees of cytoplasmic lesions at 4 DPI, including HK and SP cellular necrosis, intestinal epithelial cell shedding, and erythrocyte infiltration (**Figure 1J**). Collectively, these data confirmed the successful establishment of an LMBV infection model in largemouth bass.

Next, we evaluated the expression levels of 15 immune-related genes in the HK, SP, and gut of largemouth bass at 1, 4, 7, 14, 21, and 28 DPI via qPCR. Our initial findings indicated that upon viral infection, inflammation, innate and adaptive immune responses were triggered in the HK, SP, and gut tissues (**Figure 2A–C**). The innate and adaptive immune responses were most intense at 4 and 28 DPI, respectively. Consequently, these two time points were chosen for RNA-Seq analysis to further characterize the immune responses in HK and gut tissues following LMBV infection. Specifically, 1,237 and 1,388 genes were up-regulated in the HK at 4 and 28 DPI, whereas 22 and 1413 genes were up-regulated in the gut at these time points. Additionally, 1,616 and 916 genes were down-regulated in the HK, and 2,004 and 143 extensive changes in immune-related genes in the HK and gut at 4 and 28 DPI (**Figure 2D** and **2E**). Subsequent Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses of these differentially expressed genes (DEGs) indicated an antiviral immune response process occurring in the HK and gut, with enrichment in pathways such as NF-kappa B signaling, Epstein-Barr virus infection, and the intestinal immune network for IgA production (**Figure 2F** and **2G**). Furthermore, key genes involved in antiviral responses and innate immunity were up-regulated at 4 DPI, whereas genes associated with adaptive immunity were up-regulated at 28 DPI in both HK and gut tissues (**Figure 2H** and **2I**). Importantly, IgM exhibited high expression levels in systemic (HK and SP) and mucosal (gut) tissues of infected fish at 28 DPI (**Figure 2A–C, H**, and **I**), suggesting that IgM plays a role in both systemic and mucosal compartment.

IgM⁺ B cells and specific IgM responses to viral infection in largemouth bass

Here, we developed a monoclonal antibody to largemouth bass IgM and its specificity was rigorously validated through mass spectrometry (**Figure 3-figure supplement 1**), western blot and flow cytometry analysis (**Figure 3-figure supplement 2**). To further evaluate the IgM⁺ B cell responses, fish that survived the first LMBV infection were re-challenged with LMBV at 30 DPI




Figure 2.

The dynamic immune response of largemouth bass upon LMBV infection.

(A–C) Heatmaps display qPCR results for expression of selected immune genes in the HK (A), SP (B), and gut (C) of LMBV-infected and control fish ($n = 9$). Color values: $\log_2$ (mean fold change). (D) Venn diagrams visualize the overlapping genes that are either upregulated or downregulated in the HK and gut of largemouth bass at 4 or 28 DPI when compared to control fish ($n = 6$). (E) Heat map analysis of all DEGs associated with immune in the HK and gut of largemouth bass compared to control fish at 4 or 28 DPI. (F) GO enrichment analysis of the DEGs in the HK and gut of largemouth bass at 4 or 28 DPI versus control fish. (G) KEGG enrichment analysis of the DEGs in the HK and gut of largemouth bass compared to control fish at 4 or 28 DPI. (H, I) Representative adaptive immune, innate, and antiviral response genes modulated by LMBV infection in the HK (H) and gut (I) of largemouth bass at 4 or 28 DPI.

(**Figure 3A** [↗](#)). Fish were sampled at 28 DPI (28 DPI surviving group, 28DPI-S) and 42 DPI (42 DPI surviving group, 42DPI-S) after the initial infection (**Figure 3A** [↗](#)). Immunofluorescence microscopy analysis revealed a moderate accumulation in IgM⁺ B cells in the HK and gut of the 28DPI-S fish compared to the control group (Con), with a notable increase observed in the 42DPI-S fish (**Figures 3B–E** [↗](#), with isotype-matched control antibodies shown in (**Figure 3–figure supplement 3** [↗](#)). Flow cytometry analysis produced comparable results (**Figure 3–figure supplement 4A–D** [↗](#)). To determine whether the increase in IgM⁺ B cells detected in the gut of 42DPI-S fish was due to local proliferation, we evaluated the *in vivo* proliferation responses of IgM⁺ B cells using fluorescent 5-ethynyl-2'-deoxyuridine (EdU). Flow cytometry analysis revealed a significant increase in the number of EdU⁺ IgM⁺ B cells in both the HK and gut of the 28DPI-S fish compared to controls, with more pronounced proliferation observed in the 42DPI-S fish (**Figure 3F–I** [↗](#)). Along with the increases in IgM⁺ B cells, sIgM protein levels in the serum of 28DPI-S and 42DPI-S fish increased by approximately 2.3- and 3.9-fold compared to control fish, respectively (**Figure 3J** [↗](#)). In the gut mucus, the IgM concentration increased by approximately 1.6- and 3.8-fold in 28DPI-S and 42DPI-S fish, respectively (**Figure 3K** [↗](#)). We subsequently measured the LMBV-specific IgM titers in serum and gut mucus from the 28DPI-S and 42DPI-S groups by ELISA. Consistent with the sIgM protein expression levels, fish in the 42DPI-S group exhibited higher LMBV-specific IgM titers in both serum (~36,267) and gut (~230) compared to those in the serum (~7,867) and gut (~41) of fish in the 28DPI-S group (**Figure 3L** [↗](#) and **M** [↗](#)). These data support the idea that IgM is involved in both systemic and mucosal immunity against viral infection, with a greater contribution observed in the systemic compartment compared to the mucosa.

Largemouth bass surviving LMBV infection increases resistance to the reinfection

Based on the high LMBV-specific IgM titers in serum and gut mucus of 42DPI-S fish, we hypothesized that bass surviving the initial viral infection develop resistance to reinfection. To test this hypothesis, we challenged fish from both the control group (unexposed to LMBV) and the 42DPI-S survivor group (exposed to LMBV twice) with the same dose of LMBV (100 μ L, 1×10^6 TCID₅₀) via intraperitoneal injection, and monitored them daily for 30 days (**Figure 4A** [↗](#)). Following the challenge, approximately 57% of the fish in the control-challenge (CC) group died, whereas only 10% of the fish in the 42DPI-S challenge (SC) group died (**Figure 4B** [↗](#)). qPCR analyses revealed that viral loads in the HK and gut were significantly lower in the SC group compared to the CC group (**Figure 4C** [↗](#) and **D** [↗](#)). Immunofluorescence microscopy further confirmed that high viral loads were present in the HK (**Figure 4E** [↗](#) and **F** [↗](#)) and gut (**Figure 4G** [↗](#) and **H** [↗](#)) of the CC group, whereas very low viral loads were detected in the 42DPI-S group (**Figure 4E–H** [↗](#)). Pathological examination showed significant histological damage in the HK and gut of CC fish compared to control fish (**Figure 4I–L** [↗](#)). Interestingly, no visible tissue damage was observed in either the 42DPI-S or 42DPI-SC groups (**Figure 4I–L** [↗](#)). Collectively, these findings suggest that high LMBV-specific IgM titers induced by prior exposure to LMBV may confer protective resistance against viral reinfection.

IgM depletion significantly increases fish mortality

To determine the function of IgM in antiviral defense, we established an IgM⁺ B cell depletion model as previously described. We first assessed the degree and duration of IgM⁺ B cells and IgM depletion in the blood and gut, along with its effects on serum IgM and gut mucus sIgM levels. Flow cytometry analysis revealed that IgM⁺ B cells were significantly depleted in blood (49.7%–98.9%), SP (42.7%–89.8%), and gut (50.1–88.3%) from day 1 to day 7 post-depletion, with the most substantial depletion observed at day 3 compared to control fish (**Figure 5–figure supplement 1A–F** [↗](#)). Moreover, the depletion of IgM⁺ B cells also led to significant reductions in IgM protein levels in serum (49.7%–96.6%) and gut mucus (69.5%–90.1%) until day 10 post-depletion (**Figure 5–figure supplement 1G–J** [↗](#)).




Figure 3.

IgM⁺ B cells and LMBV-specific sIgM responses of largemouth bass upon LMBV infection.

(A) Diagram illustrating the infection method using LMBV through intraperitoneal injection. Fish received a 100 μ L injection of virus (1×10^6 TCID₅₀). The surviving fish from one group were sacrificed at 28 DPI, those from another group were re-injected with the same dose of virus at 30 DPI and then sacrificed 42 DPI. These groups are referred to as the 28DPI-S group and the 42DPI-S group, respectively. (B, D) Immunofluorescence staining of IgM (green) in paraffin sections of the largemouth bass HK (B) and gut (D) from control, 28DPI-S, and 42DPI-S fish. DAPI stains the cell nuclei blue. Scale bars, 20 μ m. (C, E) The number of IgM⁺ B cells the HK (C) and gut (E) from control, 28DPI-S, and 42DPI-S fish were quantified, as counted from B ($n = 12$). (F, H) Representative flow cytometry dot-plots demonstrate the proliferation of IgM⁺ B cells in the leukocytes of the HK (F) and gut (H) of control, 28DPI-S, and 42DPI-S fish. Each dot plot shows the percentage of lymphocytes that are proliferative B cells (EdU). (G, I) The percentage of EdU⁺ cells among the total IgM⁺ B cell populations in the HK (G) and gut (I) of control, 28DPI-S, and 42DPI-S fish ($n = 12$). (J, K) IgM protein concentration in serum (J) and gut mucus (K) of fish in the control, 28DPI-S, and 42DPI-S group ($n = 12$). (L, M) LMBV-specific IgM titers of serum (L) and gut mucus (M) from 28DPI-S and 42DPI-S fish ($n = 12$). Data are shown as mean $\pm$ CSEM. An unpaired Student's *t*-test was used. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.




Figure 4.

Resistance to reinfection in largemouth bass that survived LMBV infection.

(A) Strategy to obtain the control and survivor fish. Control fish (Con group) were obtained previously injection with EPC cells culture supernatant. At 42 days, fish were challenged with 100 µL of LMBV (1×10^6 TCID₅₀), and 4 days post-challenge fish (control challenge or CC group) were sacrificed for sampling. 42 DPI survivor fish (42DPI-S group) were injected twice with LMBV (100 µL, 1×10^6 TCID₅₀) at 0 and 28 days, respectively. At 42 days, fish were challenged with LMBV (100 µL, 1×10^6 TCID₅₀), and 4 days post-challenge fish (42DPI-S challenge or SC group) were sacrificed for sampling. (B) Cumulative survival rates of the Con, CC, 42DPI-S, and 42DPI-SC. (C, D) LMBV-MCP gene copies (Log₁₀) were quantified in HK (C), gut (D) of Con, CC, 42DPI-S, and 42DPI-SC using qPCR. (E, G) Immunofluorescence staining of LMBV in HK (E) and gut (G) paraffin-sections from Con, CC, 42DPI-S, and 42DPI-SC fish ($n = 6$). LMBV (red) was stained with an anti-LMBV-MCP mAb; nuclei (blue) were stained with DAPI. Scale bars, 20 µm. (F, H) Quantification of LMBV in HK (F) and gut (H) paraffin-sections from Con, CC, 42DPI-S, and 42DPI-SC fish counted from E and H. (I, K) Histological examination of HK (I) and gut (K) from Con, CC, 42DPI-S, and 42DPI-SC fish. The red arrow indicates erythrocyte infiltration. The black arrow indicates epithelial cell shedding in gut. The black asterisk indicates cellular necrosis in HK. (J, L) Pathology score of HK (J) and gut (L) from Con, CC, 42DPI-S, and 42DPI-SC fish ($n = 6$). Data are shown as mean $\pm$ CSEM. An unpaired Student's *t*-test was used. **p* < 0.05, ***p* < 0.01, and ****p* < 0.001.




Figure 5.

Depletion of IgM⁺ B cells in fish significantly enhances tissue viral load and increases mortality following LMBV infection.

(A) Schematic of the experimental strategy used to deplete bass IgM⁺ B cells. In brief, immune fish (those that survived two infections with 100 μ L of LMBV at 1×10^6 TCID₅₀ via intraperitoneal injection) were injected with anti-bass IgM mAbs (IgM-depleted immune fish) or either isotype control antibodies (non-depleted immune fish). One day after antibody injection, both non-depleted and IgM-depleted fish were infected with 100 μ L of LMBV at 1×10^6 TCID₅₀ via intraperitoneal injection. Fish from different groups were then analyzed at 2, 4, and 30 DPI for IgM titers, viral load, pathological changes, and mortality, respectively. (B, C) LMBV-specific IgM titers of serum (B) and gut mucus (C) from non-depleted and IgM⁺ B cell-depleted 42DPI-S fish at 2 DPI (*n* = 9). (D) H&E staining of largemouth bass HK, SP, and gut from non-depleted and IgM⁺ B cell-depleted 42DPI-S fish. Red arrow indicates erythrocyte infiltration. Black asterisk indicates cellular necrosis in HK, SP, and gut. Scale bars, 50 μ m. (E) Histological score of HK, SP, and gut in non-depleted and IgM⁺ B cell-depleted 42DPI-S fish at 4 DPI. (F) LMBV was detected using anti-LMBV mAb (red) in HK, SP, and gut of 42DPI-S fish, both non-depleted (left) and IgM⁺ B cell-depleted (right). Nuclei were stained with DAPI (blue). Scale bars, 20 μ m. (G) Quantification of LMBV-infected cells was performed in the HK, SP, and gut of both non-depleted and IgM⁺ B cell-depleted 42DPI-S fish, as counted from F. (H–J) LMBV gene copies (Log₁₀) were quantified by qPCR in HK (H), SP (I), and gut (J) (*n* = 9). (K) Cumulative survival of non-depleted and IgM⁺ B cell-depleted 42DPI-S fish infected with LMBV. Data are shown as mean $\pm$ CSEM. An unpaired Student's *t*-test and log-rank (Mantel-Cox) test in K were used. **p* < 0.05, ***p* < 0.01, ****p* < 0.001.

To further investigate whether IgM depletion increased susceptibility to viral infection, we compared 42DPI-S fish with and without IgM⁺ B cells depletion. Specifically, 42DPI-S fish were divided into two groups: one group received anti-bass IgM mAbs to deplete IgM⁺ B cells, whereas the other group (non-depleted 42DPI-S fish) was treated with isotype control antibodies. After 1 day of treatment, the fish were challenged with LMBV (100 μ L, 1×10^6 TCID₅₀) via intraperitoneal injection. LMBV-specific IgM titers in serum and gut mucus, viral load, and histopathological changes in fish tissues were assessed at 2 and 4 days post-challenge (**Figure 5A**). Furthermore, fish mortality was monitored daily for 30 days post-challenge (**Figure 5A**). As anticipated, LMBV-specific IgM titers were significantly lower in the IgM⁺ B cell-depleted 42DPI-S fish compared to the non-depleted group (**Figure 5B** and **C**). Hematoxylin and eosin (H&E) staining revealed pronounced histopathological changes in the HK, SP, and gut of IgM⁺ B cell-depleted 42DPI-S fish compared to the non-depleted group (**Figure 5D** and **E**). Moreover, immunofluorescence microscopy and qPCR further confirmed elevated levels of LMBV-MCP expression in the tissues of IgM⁺ B cell-depleted 42DPI-S fish (**Figure 5F-J**). Importantly, the IgM⁺ B cell-depleted 42DPI-S group exhibited significantly higher cumulative mortality compared to the non-depleted group (**Figure 5K**). These findings suggest that LMBV-specific IgM may play a role in enhancing resistance to viral infection and could be important for survival.

Neutralizing capacity of viral-specific IgM

As described above, viral-specific IgM plays an important role in protecting fish against LMBV. Therefore, we next sought to explore the specific antiviral mechanisms of IgM. To verify whether viral-specific IgM could also neutralize viruses, we first incubated the virus for 1 hour with serum or gut mucus from control and 42DPI-S fish. The role of IgM was evaluated by depleting IgM from half of the 42DPI-S serum or gut mucus to obtain IgM-depleted samples (42DPI-S-IgMDEP serum or gut mucus) (**Figure 6A**). After incubation, EPC cells were exposed to these three different LMBV-containing samples (control, 42DPI-S, and 42DPI-S-IgMDEP), and EPC cell viability, CPE, and viral loads were subsequently assessed at 2 and 4 days post-incubation (**Figure 6A**). Lower EPC viability and significantly higher CPE were observed in the control and 42DPI-S-IgMDEP groups compared to the 42DPI-S group, regardless of whether the sample was derived from serum (**Figure 6B** and **D**, **Figure 6-figure supplement 1A**) or gut mucus (**Figure 6C** and **D**, **Figure 6-figure supplement 1B**), suggesting that the protective efficacy of 42DPI-S was attributable to LMBV-specific IgM. Consistent with these findings, the expression of the LMBV-MCP gene and protein levels were notably decreased in EPC cells treated with the 42DPI-S serum (**Figure 6E, G**, and **I**) or gut mucus (**Figure 6F, H**, and **J**) compared to control fish. However, their levels were similar to control fish when IgM was depleted in the 42DPI-S samples (**Figure 6E-J**). Similarly, immunofluorescence analysis demonstrated that the number of LMBV-infected EPC cells treated with 42DPI-S samples (i.e., serum and gut mucus) decreased significantly compared to the control fish, whereas the number of infected EPC cells in the 42DPI-S-IgMDEP group increased and nearly returned to the level observed in control fish (**Figure 6K-N**). To further rule out complement or other factors, we purified specific IgM from serum and gut mucus of 42DPI-S fish for neutralization assays. A significant reduction in both LMBV-MCP gene expression and protein levels was observed in EPC cells treated with purified IgM from serum (**Figure 6-figure supplement 2A, C**, and **D**) or gut mucus (**Figure 6-figure supplement 2B, E**, and **F**). Moreover, significantly lower CPE were observed in the IgM treated group, while no CPE was observed in medium and bass IgM group (**Figure 6-figure supplement 2G**). Collectively, these findings strongly suggest that the neutralization process is a potential mechanism of IgM, serving as a key molecule in adaptive immunity against viral infection.

Viral-specific IgM acts directly on LMBV

The findings described above demonstrate that viral-specific IgM can neutralize viruses. Therefore, we next investigated the specific stage of viral infection at which viral-specific IgM exerts its anti-LMBV effect. To this end, serum or gut mucus from control and 42DPI-S fish was




Figure 6.

Viral neutralization exerted by LMBV-specific sIgM of serum and gut mucus.

(A) Schematic of the experimental strategy. Magnetic protein G beads were incubated with anti-bass IgM mAbs to produce protein G beads coated with these antibodies. IgM in 42DPI-S serum and gut mucus was removed by incubating these samples with protein G beads coated with anti-bass IgM mAbs. Thereafter, serum and gut mucus obtained from control, 42DPI-S, or 42DPI-S-IgMDEP fish, and each of these was incubated with LMBV and then added to EPC cells. EPC cells were treated with different serum, gut mucus, or medium alone, resulting in three distinct treatment groups for the EPC cells. (including: Control (control serum/gut mucus-LMBV-EPC cells), 42DPI-S (42DPI-S serum/gut mucus-LMBV-EPC cells), and 42DPI-S-IgMDEP (42DPI-S-IgMDEP serum/gut mucus-LMBV-EPC cells)) were analyzed at 2 and 4 days after addition for EPC cell viability and LMBV loads. (B, C) Cell viability in the different groups was measured using the colorimetric alamarBlue assay and expressed as relative values compared to those of EPC cells that were treated with serum (B) and gut mucus (C) ($n = 12$). (D) The cellular cytopathic effect in the different groups was assessed by the crystal violet stain. (E, F) LMBV-MCP gene copies (Log_{10}) were quantified by qPCR in EPC cells from the serum treatment group (E) and the gut mucus group (F) ($n = 12$). (G, H) The LMBV-MCP protein expression in EPC cells from the treatment with serum group (G) and gut mucus group (H) was evaluated by western blot. (I, J) Relative expression of LMBV-MCP protein in EPC cells from the serum treatment group (I) and the gut mucus group (J) was assessed using densitometric analysis of immunoblots ($n = 12$). (K, M) The LMBV-MCP protein in EPC cells from the serum treatment group (K) and the gut mucus group (M) was identified via immunofluorescence using the anti-LMBV-MCP mAb. Scale bars, 100 μm . (L, N) The number of virally infected cells, as determined from K and M ($n_{\text{IJ}} = 12$). Data are shown as mean $\pm$ CSEM. One-way analysis of ANOVA was used. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

directly added to co-incubate with the virus or EPC cells before or during infection (**Figure 7A**). To further confirm that the anti-LMBV effect is indeed carried out by IgM, we also applied IgM depletion treatment to the serum or mucus (42DPI-S-IgMDEP serum or gut mucus) as described above. The virus infection process is roughly divided into four phases: pre-infection, virus adsorption (at 4 C), virus invasion (0 to 2 hours post-infection (hpi) at 28 C), and replication (2 to 24 hpi). After 24 hpi, samples were collected to measure the levels of viral RNA (**Figure 7B** and **C**) and viral protein (**Figure 7D–G**). No significant effect was observed at any stage of viral infection in the serum or gut mucus from control fish (**Figure 7B–G**). In contrast, LMBV proliferation levels were significantly inhibited when serum or gut mucus from 42DPI-S fish were applied to viruses throughout the experiment (group 2, G2) or prior to and during adsorption (group 5, G5) (**Figure 7B–G**). It is worth mentioning that the 42DPI-S-IgMDEP serum or gut mucus no longer exhibited an inhibitory effect on the virus (**Figure 7B–G**). These results suggest that viral-specific IgM mediates neutralization through mechanisms involving directly acting on LMBV particles, causing them to lose their infectivity.

Discussion

IgM emerged over 500 Mya with the emergence of jawed fish, and homologs of this ancient Ig have been identified across all jawed vertebrate taxa (Warr et al., 1979; Hordvik et al., 1992). Recent discoveries have shed new light on IgM as an ancient antiviral weapon that plays a pivotal role in antiviral humoral immunity (Gong and Ruprecht, 2020). In mammals, IgM has been shown to neutralize viruses, such as in plasma from convalescent patients recovering from COVID-19 (Harrington et al., 2021). Depletion of IgM in these cases resulted in a loss of neutralization against SARS-CoV-2, significantly increasing susceptibility (Gasser et al., 2021). Compared to mammals, while pathogen-specific sIgM responses are highly induced in systemic and mucosal tissues of teleosts, the precise function and mechanisms of IgM in these organisms remain poorly understood. Therefore, this study fills significant gaps in our fundamental understanding of IgM function and evolution in the vertebrate lineage.

Here, we developed a model of LMBV infection by intraperitoneal injection in largemouth bass, an ancient teleost species. Following viral infection, the fish exhibited typical symptoms of LMBV, including hemorrhagic symptoms of the skin and fins and splenomegaly. Additionally, severe histological changes were observed in the HK, SP, and gut, including cellular shedding, erythrocyte infiltration, and intercellular space enlargement, consistent with symptoms reported in previous studies (Xu et al., 2023). Importantly, our findings demonstrate that the gut harbored higher LMBV loads compared to the HK and SP, indicating that in viral-infected fish, mucosal tissues mount a robust immune response alongside systemic immune organs. Previous studies have shown that several immune-related genes are significantly upregulated in teleost fish after infection with viral pathogens such as infectious hematopoietic necrosis virus (IHNV) (Kong et al., 2024; Huang et al., 2022) and viral hemorrhagic septicemia virus (VHSV) (Nuñez-Ortiz et al., 2018; Moore et al., 2017). Similarly, our findings indicate that LMBV infection significantly upregulates several immune genes in both systemic immune organs (HK and SP) and mucosal tissues (gut) of largemouth bass. These immune genes are associated with inflammatory responses (i.e., IL-6, IL-10, IL-1 β , IL-8, and TNF- α) (Shi et al., 2023), antiviral responses (i.e., IRF3, IRF7, IFN- γ , TRIM25, and DHX58) (Zou and Secombes, 2011; Ning et al., 2011), B cell activation (CD79a) (Cheng et al., 2021), B cell proliferation and antibody secretion (CD22) (Tedder et al., 2005), and Ig heavy chain genes (IgM, IgD, and IgT) (Yu et al., 2018). This is consistent with findings in other teleosts and mammals following viral infection. Furthermore, transcriptome data revealed an increase in IgM expression in both the HK and gut on day 28 post-LMBV infection. Therefore, we speculate that teleost IgM plays a role in both systemic and mucosal areas during viral infection.




Figure 7.

Time-of-addition experiments.

(A) EPC cells or LMBV was treated with serum or gut mucus from control, 42DPI-S, or 42DPI-S-IgM DEP fish at different times before and after infection. Samples were harvested at 24 hpi. G1-G8 respectively represent serum/gut mucus treatments during the different time periods. (B, C) LMBV-MCP gene copies (Log_{10}) was quantified by qPCR in EPC cells from the treatment with serum (B) or gut mucus (C) ($n = 6$). (D, F) The LMBV-MCP protein in EPC cells from the treatment with serum (D) or gut mucus (F) was detected by western blot using anti-LMBV-MCP monoclonal antibody and anti- β -actin antibody ($n = 6$). (E, G) Expression of LMBV-MCP/ β -actin in EPC cells from the treatment with serum (B) or gut mucus at different stage of LMBV infection group. Values were normalized against those for group 1 (G1, no treatment). Data are shown as mean $\pm$ SEM. One-way analysis of ANOVA was used. $**p < 0.01$ and $***p < 0.001$.

To verify this hypothesis, we further analyzed IgM⁺ B cells and IgM responses to the virus at both the cellular and protein levels. Consistent with the changes in IgM concentrations in serum and gut mucus, a large number of IgM⁺ B cells proliferated and accumulated in the HK and gut of 28DPI-S and 42DPI-S fish, indicating that these accumulated IgM⁺ B cells resulted in a significant enhancement of the sIgM response to viral infection. Similarly, numerous IgM-secreting plasma cells and a dramatic increase in sIgM concentrations have been observed in the bone marrow and serum of mammals after influenza virus infection (Skountzou et al., 2014 [↗](#); Wu et al., 2022 [↗](#)). Virus-specific IgM production has been well-documented in reptiles, birds, and mammals upon viral infection (Dascalu et al., 2024 [↗](#); Harrington et al., 2021 [↗](#); Hetzel et al., 2021 [↗](#); Neul et al., 2017 [↗](#)). While current evidence confirms the capacity of cartilaginous fish and amphibians to mount specific IgM responses against bacterial pathogens and immune antigens (Dooley and Flajnik, 2005 [↗](#); Ramsey et al., 2010 [↗](#)), the potential for viral induction of analogous IgM-mediated immunity in these species remains unresolved. In this study, we identified a significant increase in virus-specific sIgM titers in both serum and gut mucus, similar to what we previously reported in common carp (*Cyprinus carpio*) infected with SVCV (Yu et al., 2024 [↗](#)). These results suggest that sIgM contributes to immune protection at both mucosal and systemic immunity, challenging the current paradigm that IgM in fish is specialized as a systemic Ig isotype against pathogens (Sunyer et al., 2013 [↗](#)). Notably, 42DPI-S fish survived upon reinfection with a lower dose of LMBV in the SP, HK, and gut, indicating the development of protective immunity in these fish. Next, to assess the indispensable role of sIgM in LMBV resistance, we depleted IgM⁺ B cells in largemouth bass using a novel strategy recently described by Ding et al., 2023 [↗](#). Importantly, we observed that fish devoid of sIgM became more susceptible to LMBV infection compared to control fish. This was characterized by severe pathological manifestations in the SP, HK, and gut, accompanied by markedly higher viral loads and a notable increase in mortality. Studies on mammals have shown that IgM^{-/-} mice, lacking sIgM, lost their resistance to West Nile virus, even following inoculation with low viral doses (Diamond et al., 2003 [↗](#)). These data support the critical role of fish IgM in host defense against viral infection.

The main attribute determining the antiviral efficacy of specific antibodies lies in their ability to neutralize viruses (Burton, 2002 [↗](#)). Extensive studies in endotherms (birds and mammals) have demonstrated that specific IgM contributes to viral resistance by neutralizing viruses (Baumgarth et al., 2000 [↗](#); Diamond et al., 2003 [↗](#); Ku et al., 2021 [↗](#); Hagan et al., 2016 [↗](#); Singh et al., 2022 [↗](#)). In contrast, the neutralizing activity of IgM in amphibians and reptiles remains largely unexplored. Although viral infections have been shown to induce neutralizing antibodies in Chinese soft-shelled turtles (*Pelodiscus sinensis*) (Nie and Lu, 1999 [↗](#)), the specific Ig isotypes mediating this response have yet to be elucidated. In teleost fish, IgM has been shown to possess viral neutralizing activity similar to that observed in endotherms (Castro et al., 2013 [↗](#); Ye et al., 2013 [↗](#)). Furthermore, our recent work demonstrated that secretory IgT (sIgT) in rainbow trout (*Oncorhynchus mykiss*) can neutralize viruses, significantly reducing susceptibility to infection (Yu et al., 2022 [↗](#)). However, whether IgM in teleost fish possesses the antiviral neutralizing capacity necessary for fish to resist reinfection remains poorly understood. To explore this, we used IgM-depleted serum or gut mucus pre-incubated with EPC cell cultures and observed a significant reduction in their ability to protect EPC cells from viral infection. To the best of our knowledge, our study provides new insights into the role of sIgM in viral neutralization, suggesting a potential function of sIgM in combating viral infections. Similarly, viral-specific sIgM in serum plays a central role in humoral immunity, providing protection against SARS-CoV-2 infection by neutralizing viral effects and blocking viral entry into host cells (Gasser et al., 2021 [↗](#); Klingler et al., 2021 [↗](#)).

Polymeric IgM mediates neutralization due to its higher avidity and steric hindrance, inhibiting virus attachment and entry into target cells with great breadth and potency (Shen et al., 2019 [↗](#)). A growing body of evidence suggests that IgM mediates neutralization primarily by targeting the receptor-binding domain (RBD) of viruses, blocking virus binding to cell receptors, and effectively preventing infection (Warr et al., 1979 [↗](#); Zhu et al., 2023 [↗](#)). Our findings demonstrate that EPC

cells pre-incubated with serum or gut mucus from 42DPI-S fish, which contained high levels of virus-specific IgM, showed higher viability, a milder cytopathic effect (CPE), and lower LMBV loads. These data suggest that IgM-mediated neutralization can block virus entry into cells. Our findings revealed that sIgM in teleost fish neutralizes viruses by directly targeting viral particles, rendering them non-infectious and effectively preventing infection. The mechanism through which IgM neutralizes viruses may involve the direct destruction of the virus. This is consistent with recent research suggesting that physical disruption of the virus through antibody-imposed structural collisions damages its integrity, thus reducing its infectivity (Zheng et al., 2019 [↗](#)). Therefore, additional studies are needed to explore the specific antiviral mechanisms of IgM. Importantly, our findings indicate that virus-specific IgM antibodies in teleost fish constitute a primary line of defense in the systemic humoral response against viral infections, while also providing mucosal protection. Therefore, our data further support the concept that sIgM holds an important role in both mucosal and systemic immunity.

In conclusion, our findings demonstrate that teleost fish generate specific IgM responses in both systemic and mucosal tissues upon LMBV infection, providing further insight into the potential role and underlying mechanism of sIgM in the antiviral defense of teleost fish. The specific IgM response to viral infection in teleost fish is illustrated in **Figure 8** [↗](#). Upon infection, severe histological changes and robust adaptive immune responses occur in systemic (e.g., HK) and mucosal (gut) tissues. Subsequently, IgM⁺ B cells in the HK and gut were activated, underwent local proliferation, and differentiated into plasma cells to produce virus-specific IgM. Moreover, fish lacking sIgM showed increased susceptibility to viral infection, indicating the important role of sIgM in protection against viral infection. Crucially, we found that specific IgM has the capacity for viral neutralization and can directly act on viral particles to block infection. Although the definitive mechanisms of teleost sIgM in antiviral defense were not fully characterized in this study, our findings suggest that sIgM in both primitive and modern vertebrates utilize conserved mechanisms in response to viral infections. Finally, our findings demonstrated that sIgM may contribute to viral resistance in fish mucosal compartments, which has significant implications for the future design of fish vaccines aimed at enhancing mucosal sIgM responses.

Materials and methods

Animals

Largemouth bass (5–8 g) used in this study were obtained from Huangshi (Hubei, China) and maintained in a recirculating water system at 28 ± 1 °C. The fish were acclimated to laboratory conditions for 2 weeks and fed twice daily with a commercial diet. All experimental procedures were performed in compliance with the Guiding Principles for the Care and Use of Laboratory Animals and received approval from the Institute of Hydrobiology, Chinese Academy of Sciences (approval number 2019-048).

Cell lines and LMBV

The source of epithelioma papulosum cyprini (EPC) cell is the same as the previous study (Yu et al., 2024 [↗](#)). EPC cells were cultured in minimum essential medium (MEM; Gibco, USA) with 10% fetal bovine serum (FBS; Gibco), 100 µg/mL streptomycin (Gibco), and 100 U/mL penicillin (Gibco). The EPC cell monolayer cultivated in 75 cm² flasks was infected with LMBV at a concentration of 1×10^8 TCID₅₀ and incubated at a constant temperature of 28 °C. Cytopathic effects (CPEs) were monitored daily. When the proportion of CPEs reached 80% of the EPC cells, the infected cells were collected and stored at –80 °C. They were then subjected to three freeze-thaw cycles, followed by centrifugation at 4,000 g for 20 minutes at 4 °C to remove cell debris. Virus titers were determined using the Median TCID₅₀ assay (Reed et al., 1938 [↗](#)).




Figure 8.

Schematic of the proposed function and mechanism of teleost sIgM in response to LMBV infection.

In a healthy state, the gut contains few IgM⁺ B cells and lower levels of sIgM protein. Following the primary LMBV infection, the gut exhibits severe pathological changes, including erythrocyte infiltration, epithelial cell shedding, and a strong innate and adaptive immune response. Antigens (Ag) from LMBV are captured by antigen-presenting cells (APCs) and subsequently presented to naive CD4⁺ T cells. These antigen-specific CD4⁺ T cells then activate IgM⁺ B cells, prompting their proliferation. Subsequently, activated B cells may differentiate into plasma cells or memory IgM⁺ B cells. Upon secondary LMBV infection, plasma cells produce substantial quantities of LMBV-specific IgM. Critically, these virus-specific sIgM from both mucosal and systemic sources has the ability to neutralize the virus by directly binding viral particles and blocking host cell entry, thereby effectively reducing the proliferation of viruses within tissues. Consequently, the IgM-mediated neutralization confers protection against LMBV-induced tissue damage and significantly reduced mortality during secondary infection.

Infection of fish with LMBV and sample collection

Two types of infection challenges were conducted. For the first challenge (infected group), fish received 100 μL of LMBV (1×10^6 TCID₅₀) via intraperitoneal injection. Prior to sampling, fish were anesthetized using an overdose of MS-222. Tissues such as the head kidney (HK), spleen (SP), and gut were collected at 1, 4, 7, 14, 21, and 28 DPI for analysis of viral load, histological changes, and immune gene expression. Additionally, fluids such as gut mucus and serum were collected at 28 DPI (28DPI-S fish) and 42 DPI (14 days post-second infection, 42DPI-S fish) from survivor fish using the same methods as described previously (Yu et al., 2024 [DOI](#)). In the second challenge, fish that survived the primary infection were subjected to a secondary and tertiary infection with the same LMBV dose at 28 and 42 DPI, respectively. Following these infections, samples, including HK, SP, gut, gut mucus, and serum, were collected at 42 DPI (42DPI-S fish) and from survivor fish at 46 DPI (42DPI-SC fish). A mock group (control) was infected with culture medium from uninfected EPC cells.

RNA extraction and qPCR analysis

The methods for RNA extraction and reverse transcription from the HK, SP, and gut of bass are identical to those described in our previous study (Kong et al., 2024 [DOI](#)). qPCR was conducted using the qTOWER3G real-time PCR system (Analytik Jena AG, Germany) with the 2 $\times$ SYBR Green Master Mix (YEASEN, China). The cycling conditions are consistent with previous studies (Kong et al., 2024 [DOI](#)). To detect LMBV load, total DNA was extracted from the HK, SP, and gut of bass using the QIAamp DNA Mini Kit (Qiagen, Germany). Absolute quantification by RT-qPCR was performed to determine the LMBV copy numbers. A plasmid standard curve was constructed, and the Ct values of LMBV in tissues were extrapolated from the standard curve to calculate the copy number. The 2 $\times$ T5 Fast qPCR Mix (Qinke, China) was used under the following conditions: initial denaturation at 95 C for 5 minutes, followed by 40 cycles at 95 C for 10 seconds and 62 C for 40 seconds. The primers, synthesized by AuGCT Biotech (Wuhan, China), are listed in **Table supplement 1** [DOI](#).

Transmission electron microscopy

For transmission electron microscopy (TEM), the HK, SP, and gut tissues were fixed overnight in electron microscopy fixative and rinsed with 0.1 M phosphate buffer (pH 7.4). The fixed tissues were subsequently placed in 1% OsO₄ dissolved in 0.1 M phosphate buffer for 2 hours at room temperature (RT), followed by three washes with 0.1 M phosphate buffer. Subsequently, the tissues were subjected to dehydration using an ethanol gradient (50%, 60%, 70%, 80%, 90%, 95%, and 100%) followed by acetone. The tissues were subsequently embedded in resin and polymerized at 60 C for 48 hours. The resulting resin blocks were sectioned into ultrathin slices, transferred onto copper grids, and stained with 2.6% lead citrate and 2% uranyl acetate. The samples were examined using a TEM (HT-7700, Tokyo, Japan), and images were acquired at the Institute of Hydrobiology, Chinese Academy of Sciences.

Histology, light microscopy, and immunofluorescence

The fixation, embedding, sectioning, H&E staining, and light microscopy observation of the HK, SP, and gut of largemouth bass were carried out as described in previous studies (Kong et al., 2024 [DOI](#); Kong et al., 2019 [DOI](#)). To detect LMBV, sections were stained with 2 $\mu\text{g}/\text{mL}$ mouse anti-LMBV-MCP mAb overnight at 4 C, followed by 30 minutes of incubation with 2 $\mu\text{g}/\text{mL}$ Cy3-conjugated AffiniPure goat anti-mouse IgG (Servicebio, China). For the detection of IgM⁺ B cells, sections were stained with 2 $\mu\text{g}/\text{mL}$ mouse anti-IgM mAb. Afterward, the sections were incubated at RT for 40 minutes with Alexa Fluor 488-conjugated AffiniPure Goat anti-mouse IgG (Servicebio, China). Nuclei were incubated with DAPI (Invitrogen, USA). The sections were examined using an Olympus BX53 microscope, and images were acquired with CellSense Dimension software (Olympus, Japan).

RNA-seq library construction, sequencing, and data analyses

RNA-seq libraries were constructed from 24 samples, including those from control and LMBV-infected groups at 4 and 28 DPI, covering both the HK and gut tissues. RNA-seq data, generated by Illumina paired-end sequencing, were filtered to retain only reads uniquely mapped to one site. DEGs were estimated in R using the ‘edgeR’ package (version 3.12.1). Genes with expression levels 1 count per million in three or more samples were excluded from further analysis. DEGs with a false discovery rate (FDR) of ≤ 0.05 and $|\log_2(\text{fold change})| \geq 1$ were considered significant. GO enrichment analysis of DEGs was conducted using KOBAS (version 2.1.1).

Western blot analyses

Gut mucus, serum, and cells samples were analyzed by western blot as described by Yu et al., 2022 [\[1\]](#). Briefly, the samples were separated using 4%–15% SDS-PAGE Ready Gel (Thermo Fisher Scientific, USA) and subsequently transferred to Sequi-Blot polyvinylidene fluoride (PVDF) membranes (Bio-Rad, USA). The membranes were blocked using a 8% skim milk for 2 hours and then incubated with monoclonal antibody (mAb). For IgM concentration detection, the membranes were incubated with mouse anti-bass IgM mAb (clone 66, IgG1, 1 $\mu\text{g/mL}$) and then incubation with HRP goat-anti-mouse IgG (Invitrogen, USA) for 1 hour. IgM concentrations were determined by comparing the signal strength values to a standard curve generated with known amounts of purified bass IgM. For neutralizing effect detection, the membranes were incubated with mouse anti-LMBV MCP mAb (4A91E7, 1 $\mu\text{g/mL}$) followed by incubation with HRP goat-anti-mouse IgG (Invitrogen, USA) for 1 hour. The β -actin is used as a reference protein to standardize the differences between samples. Immunoblots were scanned using the GE Amersham Imager 600 (GE Healthcare, USA) with ECL solution (EpiZyme, China).

Flow cytometry analysis

Leukocytes from the HK and gut were isolated as described by Yu et al., 2022 [\[1\]](#). The percentages of IgM^+ B cells among the leukocytes were analyzed using flow cytometry. Briefly, the isolated leukocytes were stained with mouse anti-bass IgM monoclonal antibody (clone 66, IgG1, 1 $\mu\text{g/mL}$) to detect IgM^+ B cells. Stained cells were subsequently analyzed using an Alexa Fluor 488-conjugated goat anti-mouse IgG monoclonal antibody (1 $\mu\text{g/mL}$, Invitrogen, USA). The analysis of stained leukocytes was conducted with a CytoFLEX flow cytometer (Beckman Coulter, USA).

Detection of proliferation of IgM^+ B cells

To assess B cell proliferation, we adapted the methodology described by Yu et al., 2022 [\[1\]](#). Control, 28DPI-S, and 42DPI-S fish were anesthetized with MS-222 and given an intravenous injection with 200 μg of EdU (Invitrogen, USA) in 100 μL of PBS. The IgM^+ B cells were stained and analyzed by flow cytometry as described previously. EdU $^+$ cells were detected using the Click-iT EdU Alexa Fluor 647 Imaging Kit (Invitrogen, USA). All stained cells were then analyzed using a CytoFLEX flow cytometer (Beckman Coulter, USA).

Enzyme-linked immunosorbent assay (ELISA)

Specific anti-virus IgM titers were determined using ELISA. Purified LMBV (10 $\mu\text{g/mL}$, 50 μL per well) was incubated on Maxisorp microplates (Thermo Fisher Scientific, USA) overnight at 4 C. Non-specific binding sites were blocked using a 8% skim milk for 2 hours at RT. Then, each well was washed once with 200 μL of PBS-EDTA-Tween (1 $\times$ PBS, 10 mM EDTA, 0.05% Tween, pH 7.2). Subsequently, serum or gut mucus were diluted in PBS-EDTA (1 $\times$ PBS, 10 mM EDTA, pH 7.2) and added to the wells, followed by incubation for 2 hours at 4 C. The anti-bass IgM mAb (clone 66, IgG1, 2 $\mu\text{g/mL}$), diluted in PBS-EDTA-BSA (1 $\times$ PBS, 10 mM EDTA, 1% BSA, pH 7.2), was added to each well and incubated for 2 hours at RT. Bound antibodies were detected using HRP-conjugated goat anti-mouse IgG (H+L) (0.5 $\mu\text{g/mL}$) for 40 minutes at RT, followed by incubation with ABTS (2,2’-

azino-di-(3-ethylbenzthiazoline sulfonic acid), Invitrogen, USA) as the substrate. The color reaction was stopped after 15–20 minutes with 100 μ L of 2 M H_2SO_4 , and absorbance was measured at 450 nm using a microplate reader (BioTec Instruments, Inc., USA). LMBV-specific IgM titers are expressed as the reciprocal of the highest dilution of serum or gut mucus that produced an average absorbance exceeding twofold the average background.

***In vitro* depletion of IgM in serum and gut mucus**

To evaluate the neutralizing effect of specific IgM against LMBV, we first obtained IgM-depleted serum and gut mucus from 42DPI-S fish. Briefly, 200 μ L of protein G magnetic beads (Biolinkedin, China) were incubated with approximately 20 μ g of mouse anti-bass IgM monoclonal antibody (mAb, clone 66, IgG1). To eliminate complement-mediated interference, serum or mucus were all heat inactivation at 56 C preceding neutralization assays. Subsequently, the antibody-conjugated beads were then mixed with 100 μ L of 42DPI-S gut mucus and incubated for 2 hours at RT. After incubation, the beads were separated using a Magna Bind Magnet, and the clear supernatant (42DPI-S-IgMDEP) was collected. 100 μ L of LMBV (1×10^4 TCID₅₀) was pre-incubated with serum or gut mucus from control, 42DPI-S, or 42DPI-S-IgMDEP fish for 1 hour. EPC cells were then treated with each of the LMBV-containing samples. To further rule out other factors, we purified IgM from serum and gut mucus of 42DPI-S fish for neutralization assays. Briefly, anti-bass IgM mAb was coupled to CNBr-activated sepharose 4B beads and used for purification of IgM from both serum and gut mucus of 42DPI-S fish. After that, 100 μ L of LMBV (1×10^4 TCID₅₀) in MEM was incubated with PBS and purified IgM (100 μ g/mL) at 28 °C for 1 hour and then the mixtures were applied to infect EPC cells. Medium or bass IgM was added to EPC cells as controls. Two days after treatment, EPC cells were analyzed for cell viability and the load of LMBV. qPCR was performed at 2 DPI, and immunofluorescence staining was conducted at 4 DPI. Additionally, when CPEs were observed, cells were fixed using a 4% (v/v) neutral paraformaldehyde solution, subsequently washed with PBS (pH 7.2), and finally, stained with crystal violet for a duration of 15 minutes. The results of patch determination were analyzed under an optical microscope (Phenix), and images were captured using the MShot image analysis system.

***In vivo* depletion of IgM⁺ B cells**

The *in vivo* IgM⁺ B cell depletion model was adapted from previously established IgM depletion strategies (Ding et al., 2023 [DOI](#)). Largemouth bass (~3 to 5 g) were intraperitoneally injected with 300 μ g of mouse anti-bass IgM monoclonal antibody (mAb, clone 66, IgG1) or an isotype control (mouse IgG1, Abclonal, China). Fish were euthanized on days 1, 2, 3, 4, 5, 7, 10, and 14 post-mAb treatment, and leukocytes were collected from the HK, SP, blood, and gut. IgM⁺ B cells in the leukocytes were analyzed using flow cytometry. The concentration of IgM in the serum and gut mucus from these mAb-treated fish was measured by western blot as described previously.

Time-of-addition assay

EPC cells (5×10^4 cells/well) were seeded in 24-well cell culture plates and then incubated at 28 C for 12 hours. LMBV was mixed with serum or gut mucus derived from control, 42DPI-S, or 42DPI-S-IgMDEP fish, and the mixtures were incubated at 28 C for 1 hour. The LMBV mixtures were then added to the cells, and the cells were incubated at 4 C for 30 minutes to allow virus adsorption, followed by a 2-hours incubation at 28 C. Next, the cell supernatants were discarded, and the cells were rinsed to eliminate any unbound virus. Following this, DMEM or fresh serum/gut mucus obtained from control, 42DPI-S, or 42DPI-S-IgMDEP fish was introduced into each well incubated at 28 C. After 24 hours, cells were collected, and LMBV load in the cells was measured by qPCR and western blot analyses.

Cell viability assay

The activity of EPC cells was assessed using the alamarBlue™ Cell Viability Reagent (Invitrogen, USA) according to the manufacturer's instructions. EPC cells (5×10^4 cells/well) were seeded in 96-well plates and incubated at 28 C for 12 hours. LMBV was mixed with serum or gut mucus from control, 42DPI-S, or 42DPI-S-IgMDEP fish, followed by infection with LMBV. After 48 hours, 1/10th volume of alamarBlue reagent was added to the medium of LMBV-infected EPC cells in 96-well plates. Finally, the absorbance was recorded at 570 nm using a microplate reader (BMG LabTech, Germany), with background absorbance measured at 600 nm and subtracted.

Statistical analysis

Data were analyzed using GraphPad Prism (version 6.0). Statistical differences between groups were assessed using one-way ANOVA with Bonferroni correction or unpaired Student's *t*-test. Data are representative of three independent experiments (mean $\pm$ SEM). A *p*-value of less than 0.05 was considered statistically significant.

Supplemental Information




Figure 1-figure supplement 1.

Phylogeny of the Ig heavy chain isotypes in jawed vertebrates, related to Introduction.

IgM stands out as the isotype conserved among jawed vertebrates.




Figure 1-figure supplement 2.

Isotype control staining for anti-LMBV-MCP in largemouth bass head kidney (HK), spleen (SP), and gut paraffin sections, related to Figure 1 [↗](#).

Immunofluorescence staining of LMBV using isotype control antibodies for anti-LMBV-MCP in the HK, SP, and gut paraffin-sections, stained for LMBV (red); nuclei (blue) were stained with DAPI. Scale bars, 50 μ m.




Figure 3-figure supplement 1.

Validation of anti-bass IgM mAb by mass spectrometry, related to Figure 3.

(A) Affinity-purified IgM (~1 µg) from largemouth bass serum was separated by 4-15% SDS-PAGE under reducing conditions and visualized with Coomassie blue staining. The band corresponding to the IgM heavy chain was excised for in-gel tryptic digestion followed by mass analysis. (B) Quantitative mass spectrometry showed that IgM heavy-chain peptides accounted for 97.3% of the total signal. (C) Database searching of mass spectra against the NCBI non-redundant database revealed 54 peptide matches to the IgM heavy chain of largemouth bass (QLF98876.1), comprising 72.2% of the IgM heavy chain sequence. The numbered amino acid sequence with highlighted peptide matches is shown in highlight green. (D) Mass spectrum of IgM peptides.

Figure 3-figure supplement 2.

Validation and characterization of anti-bass IgM mAb, related to Figure 3.

(A, B) Serum samples were analyzed by western blot with anti-bass IgM (A) or an isotype-matched control antibody (B) under non-reducing conditions (NRD) or reducing conditions (RD). (C–F) Tissue lysates from various sources (i.e., spleen, skin, gill, and gut) were analyzed by western blot with a monoclonal antibody targeting largemouth bass IgM or an isotype-matched control antibody under NRD. (G) Flow cytometry of head kidney stained with anti-bass IgM mAb. (H, I) Gene-expression profiles of sorted IgM^+ B cells and IgM^- B cells among head kidney leukocytes.





Figure 3-figure supplement 3.

Isotype control staining for anti-IgM in largemouth bass head kidney (HK), spleen (SP), and gut paraffin sections, related to Figure 3.

Immunofluorescence staining of IgM using isotype control antibodies for anti-bass IgM in the HK, SP, and gut paraffin-sections, stained for IgM (green); nuclei (blue) were stained with DAPI (blue). Scale bars, 20 μm .






Figure 3-figure supplement 4.

Flow cytometry analysis of IgM⁺ B cells response in largemouth bass upon LMBV infection, related to Figure 3.

Head kidney (HK) and gut leukocytes of largemouth bass that injected with LMBV were isolated at 28 and 42 DPI. **(A, B)** Frequencies of IgM⁺ B cells in gated lymphocyte population of largemouth bass HK **(A)** and gut **(B)** after infection with LMBV. **(C, D)** Percentage of IgM⁺ cells from the HK **(C)** and gut **(D)** leukocytes populations in control, 28DPI-S, and 42DPI-S fish ($n = 12$). Statistical differences were evaluated by unpaired Student's t -test. Data in C and D are representative of at least three independent experiments (mean $\pm$ SEM). $**p < 0.01$ and $***p < 0.001$.




Figure 5-figure supplement 1.

Analysis of IgM⁺ B cells and IgM concentration in largemouth bass after IgM⁺ B cells depletion treatment, related to Figure 5

(A–C) Flow cytometry of blood (A), spleen (B), and gut (C) leukocytes from control (left) and IgM⁺ B cell-depleted (right) fish after 3 days depletion treatment. Numbers adjacent to outlined areas indicate the percentage of IgM⁺ B cells in the lymphocyte population. (D–F) The percentage of IgM⁺ B cells within the lymphocyte population of blood (D), spleen (E), and gut (F) in control or IgM⁺ B cell-depleted fish ($n = 6$). (G, H) The depletion effects of IgM from the serum (G) or gut mucus (H) of control or IgM⁺ B cell-depleted fish was detected by western blot using anti-bass mAb ($n = 6$). (I, J) Concentration of IgM in serum (I) and gut mucus (J) of control and IgM⁺ B cell-depleted fish as determined from G and H. Data are representative of at least three independent experiments (means $\pm$ SEM). Statistical analysis was performed by unpaired Student's *t* test. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.




Figure 6-figure supplement 1.

LMBV-specific IgM markedly decrease cytopathic effect (CPE) caused by LMBV, related to Figure 6 [↗](#).

(A) The CPE in the EPC cells after treatment with serum from the control, 42-DPI-S, and 42-DPI-S-IgMDEP fish. (B) The CPE in the EPC cells after treatment with gut mucus from the control, 42-DPI-S, and 42-DPI-S-IgMDEP fish. Scale bar = 50 μ m.




Figure 6-figure supplement 2.

Viral neutralization exerted by purified LMBV-specific sIgM of serum and gut mucus.

(**A**, **B**) LMBV-MCP gene copies (Log₁₀) were quantified by qPCR in EPC cells from the purified LMBV-specific sIgM of serum treatment group (**A**) and the gut mucus group (**B**) ($n = 6$). (**C**, **E**) The LMBV-MCP protein expression in EPC cells from the treatment with the purified LMBV-specific sIgM of serum group (**C**) and gut mucus group (**E**) was evaluated by western blot. (**D**, **F**) Relative expression of LMBV-MCP protein in EPC cells from the purified LMBV-specific sIgM of serum treatment group (**D**) and the gut mucus group (**F**) was assessed using densitometric analysis of immunoblots ($n = 6$). (**G**) The cellular cytopathic effect in the different groups was assessed by the crystal violet stain.

Gene	GenBank accession no.	Primer Sequence (5'→3')	
		Forward primer	Reverse primer
IRF-3	XM_038735465.1	TCTCATCTTTAAGGCGTGGGC	GGGGTTAGCGGTGTCGTTC
IRF-7	XM_038706685.1	GCTGCCACTGTGTTTGTC	CTGTCCTTTCCCTTATGC
TRIM25	XM_038714419.1	GTCACCAGTCACCACGCTTTC	CCACTTGGCTTATTTCCTCCC
IFN- γ	XM_046040264.1	TCCCTCTGAAGATGAACAAA	AACGCCACCCATAAACA
MX1	XM_046075478.1	TCTACGATGAAGGCCACGCTA	TCTTCCACGGTACTTTCGTTC
DHX-58	XM_038704586.1	GACTGATGCTTATCGCTCCCTA	ACGCTTTTACGGGTTTACTGA
IL-1 β	XM_038733429.1	CGTACATCCGTGCCAACAGT	ATGCTCTTAACTCCTCCT
IL-6	XM_046059156.1	GGGAGACTCGCTCTGACCTACTG	TACCTCCTCCTTGTGGCGTTGG
IL-8	XM_038704090.1	GAGCCATTTTTCCTGGTGACT	TCCTCATTGGTGCTGAAAGATC
IL-10	XM_038696252.1	CGGCACAGAAATCCCAGAGC	CAGCAGGCTCACAAAATAACATCT
TGF- β	XM_038710299.1	ACAACCTTCATCCATGGCGGT	TCGTTGATTCTTGGGCTGGG
TNF- α	XM_038723994.1	CAGTGCAATGGCAGAACCAG	TTGACCCTGAAGGACGCTTG
IgM	MN871984.1	GTTACCTTCTCCTGCTTG	GTTCCGTTCTCATAGTTTC
IgT	MZ388129.1	GAAGGTCAACAACGCTGAGTG	TGTTGCTGGTCACATCTAGTCC
IgD	MZ388128.1	AAGGGAAACAGTGCTGTGCT	TTGCCAGTGGGGTTTGACTT
β -actin	XM_038708956.1	CAACGATATCCGTCGTGGCA	ACAGCGAAACGACCAAGAGG

Table supplement 1.

Primers used in this study, related to Figure 2 [↗](#).

Data Availability

The raw RNA sequencing data have been deposited in the NCBI Sequence Read Archive under BioProject accession number PRJNA1254665. The mass spectrometry proteomics data have been deposited to the iProX platform with the dataset identifier IPX0011847000.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (32225050, U24A20461, and 32303053). We thank Yan Wang of the Institute of Hydrobiology, Chinese Academy of Sciences for the Flow Cytometry analysis procedures.

Additional information

Author Contributions

W.K. and Z.X. conceived the project and experimental design. W.K., X.W., and G.D. performed most of the experiments. W.K. and Y.S. analyzed the data, and contributed to the writing of the original manuscript. P.Y., C.C., and X.Y., contributed to immunofluorescence analysis and morphological detection. G.C. performed the RNA-seq analysis. Z.X. and T.F. contributed to the writing of the final manuscript. Z.X. conceptualized and supervised the study.

Ethics

All experimental procedures were performed in compliance with the Guiding Principles for the Care and Use of Laboratory Animals and received approval from the Institute of Hydrobiology, Chinese Academy of Sciences (approval number 2019-048).

Funding

National Natural Science Foundation of China (32225050)

References

- Ali M. G., Zhang Z., Gao Q., Pan M., Rowan E. G., Zhang J. (2020) **Recent advances in therapeutic applications of neutralizing antibodies for virus infections: an overview** *Immunologic Research* **68**:325–339 <https://doi.org/10.1007/s12026-020-09159-z> | Google Scholar
- Baumgarth N., Herman O. C., Jager G. C., Brown L. E., Herzenberg L. A., Chen J. (2000) **B-1 and B-2 cell-derived immunoglobulin M antibodies are nonredundant components of the protective response to influenza virus infection** *The Journal of Experimental Medicine* **192**:271–280 <https://doi.org/10.1084/jem.192.2.271> | Google Scholar
- Beebe D. P., Cooper N. R. (1981) **Neutralization of vesicular stomatitis virus (VSV) by human complement requires a natural IgM antibody present in human serum** *Journal of immunology* **126**:1562–1568 <https://doi.org/10.4049/jimmunol.126.4.1562> | Google Scholar
- Bengtén E., Quiniou S. M., Stuge T. B., Katagiri T., Miller N. W., Clem L. W., Warr G. W., Wilson M. (2002) **The IgH locus of the channel catfish, *Ictalurus punctatus*, contains multiple constant region gene sequences: different genes encode heavy chains of membrane and secreted IgD** *Journal of immunology* **169**:2488–2497 <https://doi.org/10.4049/jimmunol.169.5.2488> | Google Scholar
- Bilal S., Etayo A., Hordvik I. (2021) **Immunoglobulins in teleosts** *Immunogenetics* **73**:65–77 <https://doi.org/10.1007/s00251-020-01195-1> | Google Scholar
- Blandino R., Baumgarth N. (2019) **Secreted IgM: New tricks for an old molecule** *Journal of Leukocyte Biology* **106**:1021–1034 <https://doi.org/10.1002/JLB.3RI0519-161R> | Google Scholar
- Boes M. (2000) **Role of natural and immune IgM antibodies in immune responses** *Molecular immunology* **37**:1141–1149 [https://doi.org/10.1016/S0161-5890\(01\)00025-6](https://doi.org/10.1016/S0161-5890(01)00025-6) | Google Scholar
- Boes M., Prodeus A. P., Schmidt T., Carroll M. C., Chen J. (1998) **A critical role of natural immunoglobulin M in immediate defense against systemic bacterial infection** *The Journal of Experimental Medicine* **188**:2381–2386 <https://doi.org/10.1084/jem.188.12.2381> | Google Scholar
- Bohannon C., Powers R., Satyabhama L., Cui A., Tipton C., Michaeli M., ..., Jacob J. (2016) **Long-lived antigen-induced IgM plasma cells demonstrate somatic mutations and contribute to long-term protection** *Nature Communications* **7**:11826 <https://doi.org/10.1038/ncomms11826> | Google Scholar
- Bradshaw W. J., Valenzano D.R. (2020) **Extreme genomic volatility characterizes the evolution of the immunoglobulin heavy chain locus in cyprinodontiform fishes** *Proceedings of the Royal Society B-Biological Sciences* **287**:20200489 <https://doi.org/10.1098/rspb.2020.0489> | Google Scholar
- Bromage E. S., Ye J., Owens L., Kaattari I. M., Kaattari S. L. (2004) **Use of staphylococcal protein A in the analysis of teleost immunoglobulin structural diversity** *Developmental & Comparative Immunology* **28**:803–814 <https://doi.org/10.1016/j.dci.2003.12.001> | Google Scholar

- Burton D. R. (2002) **Antibodies, viruses and vaccines** *Nature Reviews Immunology* **2**:706–713 <https://doi.org/10.1038/nri891> | Google Scholar
- Castro R., Jouneau L., Pham H. P., Bouchez O., Giudicelli V., Lefranc M. P., ..., Boudinot P. (2013) **Teleost fish mount complex clonal IgM and IgT responses in spleen upon systemic viral infection** *PLoS pathogens* **9**:e1003098 <https://doi.org/10.1371/journal.ppat.1003098> | Google Scholar
- Chen K., Magri G., Grasset E. K., Cerutti A. (2020) **Rethinking mucosal antibody responses: IgM, IgG and IgD join IgA** *Nature Reviews Immunology* **20**:427–441 <https://doi.org/10.1038/s41577-019-0261-1> | Google Scholar
- Cheng G. F., Kong W. G., Zhai X., Mu Q. J., Dong Z. R., Zhan M. T., Xu Z. (2021) **Molecular cloning and expression analysis of CD79a and CD79b in rainbow trout (*Oncorhynchus mykiss*) after bacterial, parasitic, and viral infection** *Fish & Shellfish Immunology* **118**:385–395 <https://doi.org/10.1016/j.fsi.2021.09.022> | Google Scholar
- Dascalu S., Sealy J. E., Sadeyen J. R., Flammer P. G., Fiddaman S., Preston S. G., ..., Iqbal M. (2024) **Immunisation of chickens with inactivated and/or infectious H9N2 avian influenza virus leads to differential immune B-cell repertoire development** *Frontiers in immunology* **15**:1461678 <https://doi.org/10.3389/fimmu.2024.1461678> | Google Scholar
- Diamond M. S., Sitati E. M., Friend L. D., Higgs S., Shrestha B., Engle M. (2003) **A critical role for induced IgM in the protection against West Nile virus infection** *The Journal of Experimental Medicine* **198**:1853–1862 <https://doi.org/10.1084/jem.20031223> | Google Scholar
- Ding Y., Fernández-Montero A., Mani A., Casadei E., Shibasaki Y., Takizawa F., ..., Sunyer J. O. (2023) **Secretory IgM (sIgM) is an ancient master regulator of microbiota homeostasis and metabolism** *bioRxiv* :2023-02 <https://doi.org/10.1101/2023.02.26.530119> | Google Scholar
- Dooley H. (2018) **Chondrichthyes: The immune system of cartilaginous fishes** In: Cooper E., editors. *Advances in Comparative Immunology* Cham: Springer https://doi.org/10.1007/978-3-319-76768-0_18 | Google Scholar
- Dooley H., Flajnik M. F. (2005) **Shark immunity bites back: affinity maturation and memory response in the nurse shark, *Ginglymostoma cirratum*** *European journal of immunology* **35**:936–45 <https://doi.org/10.1002/eji.200425760> | Google Scholar
- Du Pasquier L., Zucchetti I., De Santis R. (2004) **Immunoglobulin superfamily receptors in protochordates: before RAG time** *Immunological Reviews* **198**:233–248 <https://doi.org/10.1016/B978-0-12-387664-5.00004-2> | Google Scholar
- Edholm E. S., Bengtén E., Stafford J. L., Sahoo M., Taylor E. B., Miller N. W., Wilson M. (2010) **Identification of two IgD+ B cell populations in channel catfish, *Ictalurus punctatus*** *The Journal of Immunology* **185**:4082–4094 <https://doi.org/10.4049/jimmunol.1000631> | Google Scholar
- Flajnik M. F. (2002) **Comparative analyses of immunoglobulin genes: surprises and portents** *Nature Reviews Immunology* **2**:688–698 <https://doi.org/10.1038/nri889> | Google Scholar
- Flajnik M. F. (2018) **A convergent immunological holy trinity of adaptive immunity in lampreys: Discovery of the variable lymphocyte receptors** *The Journal of Immunology* **201**:1331–1335 <https://doi.org/10.4049/jimmunol.1800965> | Google Scholar

- Flajnik M. F., Kasahara M. (2010) **Origin and evolution of the adaptive immune system: genetic events and selective pressures** *Nature Reviews Genetics* **11**:47–59 <https://doi.org/10.1038/nrg2703> | [Google Scholar](#)
- Gasser R., Cloutier M., Prévost J., Fink C., Ducas É., Ding S., ..., Finzi A. (2021) **Major role of IgM in the neutralizing activity of convalescent plasma against SARS-CoV-2** *Cell Reports* **34**:108790 <https://doi.org/10.1016/j.celrep.2021.108790> | [Google Scholar](#)
- Gong S., Ruprecht R. M. (2020) **Immunoglobulin M: an ancient antiviral weapon—rediscovered** *Frontiers in Immunology* **11**:1943 <https://doi.org/10.3389/fimmu.2020.01943> | [Google Scholar](#)
- Györkei Á., Johansen F. E., Qiao S. W. (2024) **Systematic characterization of immunoglobulin loci and deep sequencing of the expressed repertoire in the Atlantic cod (*Gadus morhua*)** *BMC Genomics* **25**:663 <https://doi.org/10.1186/s12864-024-10571-0> | [Google Scholar](#)
- Hagiwara K., Kobayashi K., Kajii T., Tomonga S. (1985) **J-chain-like component in 18-S immunoglobulin of the skate *Raja kenoi*, a cartilaginous fish** *Molecular Immunology* **22**:775–778 [https://doi.org/10.1016/0161-5890\(85\)90142-7](https://doi.org/10.1016/0161-5890(85)90142-7) | [Google Scholar](#)
- Hagan M., Ranadheera C., Audet J., Morin J., Leung A., Kobasa D. (2016) **Post-exposure treatment with whole inactivated H5N1 avian influenza virus protects against lethal homologous virus infection in mice** *Scientific reports* **12**:29433 <https://doi.org/10.1038/srep29433> | [Google Scholar](#)
- Han B., Yuan H., Wang T., Li B., Ma L., Yu S., ..., Zhao Y. (2016) **Multiple IgH isotypes including IgD, subclasses of IgM, and IgY are expressed in the common ancestors of modern birds** *The Journal of Immunology* **196**:5138–5147 <https://doi.org/10.4049/jimmunol.1600307> | [Google Scholar](#)
- Harrington W. E., Trakhimets O., Andrade D. V., Dambrauskas N., Raappana A., Jiang Y., ..., Sather D. N. (2021) **Rapid decline of neutralizing antibodies is associated with decay of IgM in adults recovered from mild COVID-19** *Cell Reports Medicine* **2**:100253 <https://doi.org/10.1016/j.xcrm.2021.100253> | [Google Scholar](#)
- Hetzel U., Korzyukov Y., Keller S., Szivovics L., Pesch T., Vapalahti O., ..., Hepojoki J. (2021) **Experimental Reptarenavirus infection of boa constrictor and *Python regius*** *Journal of virology* **95**:e01968–20 <https://doi.org/10.1128/JVI.01968-20> | [Google Scholar](#)
- Hirano M., Das S., Guo P., Cooper M. D. (2011) **The evolution of adaptive immunity in vertebrates** *Advances in Immunology* **109**:125–157 <https://doi.org/10.1016/B978-0-12-387664-5.00004-2> | [Google Scholar](#)
- Hohman V. S., Stewart S. E., Rumfelt L. L., Greenberg A. S., Avila D. W., Flajnik M. F., Steiner L. A. (2003) **J chain in the nurse shark: implications for function in a lower vertebrate** *The Journal of Immunology* **170**:6016–6023 <https://doi.org/10.4049/jimmunol.170.12.6016> | [Google Scholar](#)
- Hordvik I., Voie A. M., Glette J., Male R., Endresen C. (1992) **Cloning and sequence analysis of two isotypic IgM heavy chain genes from Atlantic salmon, *Salmo salar* L** *European Journal of Immunology* **22**:2957–2962 <https://doi.org/10.1002/eji.1830221130> | [Google Scholar](#)
- Huang Z., Zhan M., Cheng G., Lin R., Zhai X., Zheng H., ..., Xu Z. (2022) **IHN infection induces strong mucosal immunity and changes of microbiota in trout intestine** *Viruses*

14:1838 <https://doi.org/10.3390/v14081838> | Google Scholar

Keyt B. A., Baliga R., Sinclair A. M., Carroll S. F., Peterson M. S. (2020) **Structure, function, and therapeutic use of IgM antibodies** *Antibodies* **9**:53 <https://doi.org/10.3390/antib9040053> | Google Scholar

Klingler J., Weiss S., Itri V., Liu X., Oguntuyo K. Y., Stevens C., ..., Hioe C. E. (2021) **Role of immunoglobulin M and A antibodies in the neutralization of severe acute respiratory syndrome coronavirus 2** *The Journal of Infectious Diseases* **223**:957–970 <https://doi.org/10.1093/infdis/jiaa784> | Google Scholar

Kong W., Cheng G., Cao J., Yu J., Wang X., Xu Z. (2024) **Ocular mucosal homeostasis of teleost fish provides insight into the coevolution between microbiome and mucosal immunity** *Microbiome* **12**:10 <https://doi.org/10.1186/s40168-023-01716-6> | Google Scholar

Kong W. G., Yu Y. Y., Dong S., Huang Z. Y., Ding L. G., Cao J. F., ..., Xu Z. (2019) **Pharyngeal immunity in early vertebrates provides functional and evolutionary insight into mucosal homeostasis** *The Journal of Immunology* **203**:3054–3067 <https://doi.org/10.4049/jimmunol.1900863> | Google Scholar

Ku Z., Xie X., Hinton P. R., Liu X., Ye X., Muruato A. E., ..., An Z. (2021) **Nasal delivery of an IgM offers broad protection from SARS-CoV-2 variants** *Nature* **595**:718–723 <https://doi.org/10.1038/s41586-021-03673-2> | Google Scholar

Mashoof S., Criscitiello M. F. (2016) **Fish immunoglobulins** *Biology* **5**:45 <https://doi.org/10.3390/biology5040045> | Google Scholar

Magadán-Mompó S., Sánchez-Espinel C., Gambón-Deza F. (2011) **Immunoglobulin heavy chains in medaka (*Oryzias latipes*)** *BMC Evolutionary Biology* **11**:165 <https://doi.org/10.1186/1471-2148-11-165> | Google Scholar

Moore L. J., Jarungsriapisit J., Nilsen T. O., Stefansson S., Taranger G. L., Secombes C. J., ..., Patel S. (2017) **Immune gene profiles in Atlantic salmon (*Salmo salar* L.) post-smolts infected with SAV3 by bath-challenge show a delayed response and lower levels of gene transcription compared to injected fish** *Fish & Shellfish Immunology* **62**:320–331 <https://doi.org/10.1016/j.fsi.2017.01.034> | Google Scholar

Mu Q., Dong Z., Kong W., Wang X., Yu J., Ji W., ..., Xu Z. (2022) **Response of immunoglobulin M in gut mucosal immunity of common carp (*Cyprinus carpio*) infected with *Aeromonas hydrophila*** *Frontiers in Immunology* **13**:1037517 <https://doi.org/10.3389/fimmu.2022.1037517> | Google Scholar

Nguyen N. K., Devilder M. C., Gautreau-Rolland L., Fourgeux C., Sinha D., Poschmann J., ..., McIlroy D. (2023) **A cluster of broadly neutralizing IgG against BK polyomavirus in a repertoire dominated by IgM** *Life Science Alliance* **6**:e202201567 <https://doi.org/10.26508/lsa.202201567> | Google Scholar

Neul A., Schrödl W., Marschang R. E., Bjick T., Truyen U., von Buttlar. H., Pees M. (2017) **Immunologic responses in corn snakes (*Pantherophis guttatus*) after experimentally induced infection with ferlaviruses** *American journal of veterinary research* **78**:482–494 <https://doi.org/10.2460/ajvr.78.4.482> | Google Scholar

Nie Y. C., Lu C. P. (1999) **Antibody against testudo herpesvirus is not common in Chinese soft-shelled turtles** *Zentralbl Veterinarmed B* **46**:731–734 <https://doi.org/10.1046/j.1439-0450>

.1999.00292.x | [Google Scholar](#)

Ning S., Pagano J. S., Barber G. N. (2011) **IRF7: activation, regulation, modification and function** *Genes & Immunity* **12**:399–414 <https://doi.org/10.1038/gene.2011.21> | [Google Scholar](#)

Núñez-Ortiz N., Moore L. J., Jarungsriapisit J., Nilsen T. O., Stefansson S., Morton H. C., ..., Patel S. (2018) **Atlantic salmon post-smolts adapted for a longer time to seawater develop an effective humoral and cellular immune response against Salmonid alphavirus** *Fish & Shellfish Immunology* **82**:579–590 <https://doi.org/10.1016/j.fsi.2018.08.059> | [Google Scholar](#)

Ochsenbein A. F., Fehr T., Lutz C., Suter M., Brombacher F., Hengartner H., Zinkernagel R. M. (1999) **Control of early viral and bacterial distribution and disease by natural antibodies** *Science* **286**:2156–2159 <https://doi.org/10.1126/science.286.5447.2156> | [Google Scholar](#)

Ouchida R., Mori H., Hase K., Takatsu H., Kurosaki T., Tokuhisa T., ..., Wang J. Y. (2012) **Critical role of the IgM Fc receptor in IgM homeostasis, B-cell survival, and humoral immune responses** *Proceedings of the National Academy of Sciences* **109**:E2699–E2706 <https://doi.org/10.1073/pnas.1210706109> | [Google Scholar](#)

Parra D., Takizawa F., Sunyer J. O. (2013) **Evolution of B cell immunity** *Annual Review Of Animal Biosciences* **1**:65–97 <https://doi.org/10.1146/annurev-animal-031412-103651> | [Google Scholar](#)

Ramsey J. P., Reinert L. K., Harper L. K., Woodhams D. C., Rollins-Smith L. A. (2010) **Immune defenses against batrachochytrium dendrobatidis, a fungus linked to global amphibian declines, in the South African clawed frog, Xenopus laevis** *Infection and immunity* **78**:3981–92 <https://doi.org/10.1128/IAI.00402-10> | [Google Scholar](#)

Reed L. J., Muench H. (1938) **A simple method of estimating fifty per cent endpoints** *American Journal of Hygiene* **27**:493–497 <https://doi.org/10.1093/oxfordjournals.aje.a118408> | [Google Scholar](#)

Schaerlinger B., Bascove M., Fripiat J. P. (2008) **A new isotype of immunoglobulin heavy chain in the urodele amphibian Pleurodeles waltl predominantly expressed in larvae** *Molecular Immunology* **45**:776–786 <https://doi.org/10.1016/j.molimm.2007.06.356> | [Google Scholar](#)

Schroeder H. W., Cavacini L. (2010) **Structure and function of immunoglobulins** *Journal of Allergy and Clinical Immunology* **125**:S41–S52 <https://doi.org/10.1016/j.jaci.2009.09.046> | [Google Scholar](#)

Shen C., Zhang M., Chen Y., Zhang L., Wang G., Chen J., ..., Xia N. (2019) **An IgM antibody targeting the receptor binding site of influenza B blocks viral infection with great breadth and potency** *Theranostics* **9**:210 <https://doi.org/10.7150/thno.28434> | [Google Scholar](#)

Shi Y., Liu Y., Xie K., Dai J., Wang Y., Xie S., ..., Zhong L. (2023) **Dietary sanguinarine ameliorates growth impairment and intestinal dysfunction in Ctenopharyngodon idellus fed a high cottonseed meal diet** *Aquaculture* **576**:739864 <https://doi.org/10.1016/j.aquaculture.2023.739864> | [Google Scholar](#)

Singh T., Hwang K. K., Miller A. S., Jones R. L., Lopez C. A., Dulson S. J., ..., Bonsignori M. (2022) **A Zika virus-specific IgM elicited in pregnancy exhibits ultrapotent neutralization** *Cell* **185**:4826–4840 <https://doi.org/10.1016/j.cell.2022.10.023> | [Google Scholar](#)

Skountzou I., Satyabhama L., Stavropoulou A., Ashraf Z., Esser E. S., Vassilieva E., ..., Jacob J. (2014) **Influenza virus-specific neutralizing IgM antibodies persist for a lifetime** *Clinical and Vaccine Immunology* **21**:1481–1489 <https://doi.org/10.1128/CVI.00374-14> | [Google Scholar](#)

Stavnezer J., Amemiya C. T. (2004) **Evolution of isotype switching** *Seminars in Immunology* **16**:257–275 [Google Scholar](#)

Sun Y., Zhao Y. (2014) **Immunoglobulin genes in tetrapods** In: *Comparative immunoglobulin genetics* [Google Scholar](#)

Sunyer J. O. (2013) **Fishing for mammalian paradigms in the teleost immune system** *Nature Immunology* **14**:320–326 <https://doi.org/10.1038/ni.2549> | [Google Scholar](#)

Taylor H. P., Dimmock N. J. (1985) **Mechanisms of neutralization of influenza virus by IgM** *Journal of General Virology* **66**:903–907 <https://doi.org/10.1099/0022-1317-66-4-903> | [Google Scholar](#)

Tedder T. F., Poe J. C., Haas K. M. (2005) **CD22: a multifunctional receptor that regulates B lymphocyte survival and signal transduction** *Advances in Immunology* **88**:1–50 [https://doi.org/10.1016/S0065-2776\(05\)88001-0](https://doi.org/10.1016/S0065-2776(05)88001-0) | [Google Scholar](#)

Warr G. W., Deluca D., Griffin B. R. (1979) **Membrane immunoglobulin is present on thymic and splenic lymphocytes of the trout *Salmo gairdneri*** *The Journal of Immunology* **123**:910–917 <https://doi.org/10.4049/jimmunol.123.2.910> | [Google Scholar](#)

Wilson M., Bengtén E., Miller N. W., Clem L. W., Du Pasquier L., Warr G. W. (1997) **A novel chimeric Ig heavy chain from a teleost fish shares similarities to IgD** *Proceedings of the National Academy of Sciences* **94**:4593–4597 <https://doi.org/10.1073/pnas.94.9.4593> | [Google Scholar](#)

Wu L., Yang Y., Gao A., Li J., Ye J. (2022) **Teleost fish IgM+ plasma-like cells possess IgM-secreting, phagocytic, and antigen-presenting capacities** *Frontiers in Immunology* **13**:1016974 <https://doi.org/10.3389/fimmu.2022.1016974> | [Google Scholar](#)

Xu W., Zhang Z., Lai F., Yang J., Qin Q., Huang Y., Huang X. (2023) **Transcriptome analysis reveals the host immune response upon LMBV infection in largemouth bass (*Micropterus salmoides*)** *Fish & Shellfish Immunology* **137**:108753 <https://doi.org/10.1016/j.fsi.2023.108753> | [Google Scholar](#)

Xu Z., Parra D., Gómez D., Salinas I., Zhang Y. A., von Gersdorff Jørgensen L., ..., Sunyer J. O. (2013) **Teleost skin, an ancient mucosal surface that elicits gut-like immune responses** *Proceedings of the National Academy of Sciences* **110**:13097–13102 <https://doi.org/10.1073/pnas.1304319110> | [Google Scholar](#)

Xu Z., Takizawa F., Parra D., Gómez D., von Gersdorff Jørgensen L., LaPatra S. E., Sunyer J. O. (2016) **Mucosal immunoglobulins at respiratory surfaces mark an ancient association that predates the emergence of tetrapods** *Nature Communications* **7**:10728 <https://doi.org/10.1038/ncomms10728> | [Google Scholar](#)

Ye J., Kaattari I. M., Ma C., Kaattari S. (2013) **The teleost humoral immune response** *Fish & Shellfish Immunology* **35**:1719–1728 <https://doi.org/10.1016/j.fsi.2013.10.015> | [Google Scholar](#)

Yu J., Kong W., Wang X., Cai C., Cheng G., Ding G., Xu Z. (2024) **Mucosal immune responses of gut IgM in common carp (*Cyprinus carpio*) following infection with spring viremia of carp**

virus (SVCV) *Fish & Shellfish Immunology* **145**:109326 <https://doi.org/10.1016/j.fsi.2023.109326> | [Google Scholar](#)

Yu Y., Huang Z., Kong W., Dong F., Zhang X., Zhai X., ..., Xu Z. (2022) **Teleost swim bladder, an ancient air-filled organ that elicits mucosal immune responses** *Cell Discovery* **8**:31 <https://doi.org/10.1038/s41421-022-00393-3> | [Google Scholar](#)

Yu Y. Y., Kong W., Yin Y. X., Dong F., Huang Z. Y., Yin G. M., ..., Xu Z. (2018) **Mucosal immunoglobulins protect the olfactory organ of teleost fish against parasitic infection** *Plos Pathogens* **14**:e1007251 <https://doi.org/10.1371/journal.ppat.1007251> | [Google Scholar](#)

Zhang Y. A., Salinas I., Li J., Parra D., Bjork S., Xu Z., ..., Sunyer J. O. (2010) **IgT, a primitive immunoglobulin class specialized in mucosal immunity** *Nature Immunology* **11**:827–835 <https://doi.org/10.1038/ni.1913> | [Google Scholar](#)

Zheng Q., Jiang J., He M., Zheng Z., Yu H., Li T., ..., Xia N. (2019) **Viral neutralization by antibody-imposed physical disruption** *Proceedings of the National Academy of Sciences* **116**:26933–26940 <https://doi.org/10.1073/pnas.1916028116> | [Google Scholar](#)

Zhu W., Li J., Wu Z., Li H., Zhang Z., Zhu X., ..., Dong S. (2023) **Dual blockages of a broad and potent neutralizing IgM antibody targeting GH loop of EVCAs** *Immunology* **169**:292–308 <https://doi.org/10.1111/imm.13629> | [Google Scholar](#)

Zou J., Secombes C. J. (2011) **Teleost fish interferons and their role in immunity** *Developmental & Comparative Immunology* **35**:1376–1387 <https://doi.org/10.1016/j.dci.2011.07.001> | [Google Scholar](#)

Author information

Weiguang Kong

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Xinyou Wang

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Guangyi Ding

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Peng Yang

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Yong Shi

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Chang Cai

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Xinyi Yang

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Gaofeng Cheng

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

Fumio Takizawa

Faculty of Marine Science and Technology, Fukui Prefectural University, Fukui, Japan
ORCID iD: [0000-0002-0988-2770](https://orcid.org/0000-0002-0988-2770)

Zhen Xu

State Key Laboratory of Breeding Biotechnology and Sustainable Aquaculture, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China
ORCID iD: [0000-0001-6598-2058](https://orcid.org/0000-0001-6598-2058)

For correspondence: zhenxu@ihb.ac.cn

Editors

Reviewing Editor

Dario Riccardo Valenzano

Leibniz Institute on Aging, Jena, Germany

Senior Editor

Tadatsugu Taniguchi

The University of Tokyo, Tokyo, Japan

Reviewer #2 (Public review):

Summary:

In this manuscript, Weiguang Kong et al. investigate the role of immunoglobulin M (IgM) in antiviral defense in the teleost largemouth bass (*Micropterus salmoides*). The authors employ an in vivo IgM depletion system and viral infection models, complemented by in vitro assays, histology, and gene expression analysis. Assuming the specificity of the MoAb, their findings demonstrate that largemouth bass IgM functions in both systemic and mucosal immunity and exhibits viral neutralization capabilities by acting on viral particles.

Strengths:

The authors utilize multiple complementary methods, including an innovative teleost immunoglobulin depletion approach, to provide strong evidence for the important and conserved role of IgM in anti-viral resistance. The study also highlights the dual role of teleost IgM at both systemic and mucosal levels, challenging the established idea that IgT primarily mediates mucosal protection. Despite variability in IgM depletion levels, the authors demonstrate that fish with depleted IgM+ B cells exhibit significantly higher viral loads, more severe pathological changes, and increased mortality compared to control fish. These results

have evolutionary and practical implications, suggesting that IgM's role as an antiviral effector has been conserved across jawed vertebrates for over 500 million years. Insights into IgM's role could inform vaccine strategies targeting mucosal immunity in fish, addressing a key challenge in aquaculture.

Weaknesses:

While the authors validate the specificity of MoAb against IgM and address most of the aspects suggested by the reviewer. Some aspects are missing, mainly concerning the overstatement of the findings' novelty.

<https://doi.org/10.7554/eLife.104465.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #2 (Public review)

*In this manuscript, Weiguang Kong et al. investigate the role of immunoglobulin M (IgM) in antiviral defense in the teleost largemouth bass (*Micropterus salmoides*). The study employs an IgM depletion model, viral infection experiments, and complementary in vitro assays to explore the role of IgM in systemic and mucosal immunity. The authors conclude that IgM is crucial for both systemic and mucosal antiviral defense, highlighting its role in viral neutralization through direct interactions with viral particles. The study's findings have theoretical implications for understanding immunoglobulin function across vertebrates and practical relevance for aquaculture immunology.*

Strengths:

The manuscript applies multiple complementary approaches, including IgM depletion, viral infection models, and histological and gene expression analyses, to address an important immunological question. The study challenges established views that IgT is primarily responsible for mucosal immunity, presenting evidence for a dual role of IgM at both systemic and mucosal levels. If validated, the findings have evolutionary significance, suggesting the conserved role of IgM as an antiviral effector across jawed vertebrates for over 500 million years. The practical implications for vaccine strategies targeting mucosal immunity in fish are noteworthy, addressing a key challenge in aquaculture.

Weaknesses:

Several conceptual and technical issues undermine the strength of the evidence:

Monoclonal Antibody (MoAb) Validation: The study relies heavily on a monoclonal antibody to deplete IgM, but its specificity and functionality are not adequately validated. The epitope recognized by the antibody is not identified, and there is no evidence excluding cross-reactivity with other isotypes. Mass spectrometry, immunoprecipitation, or Western blot analysis using tissue lysates with varying immunoglobulin expression levels would strengthen the claim of IgM-specific depletion.

IgM Depletion Kinetics: The rapid depletion of IgM from serum and mucus (within one day) is unexpected and inconsistent with prior literature. Additional evidence, such as Western blot analyses comparing treated and control fish, is necessary to confirm this finding.

Novelty of Claims: The manuscript claims a novel role for IgM in viral neutralization, despite extensive prior literature demonstrating this role in fish. This overstatement

detracts from the contribution of the study and requires a more accurate contextualization of the findings.

Support for IgM's Crucial Role: The mortality data following IgM depletion do not fully support the claim that IgM is indispensable for antiviral defense. The survival of IgM-depleted fish remains high (75%) compared to non-primed controls (~50%), suggesting that other immune components may compensate for IgM loss

Presentation of IgM Depletion Model: The study describes the IgM depletion model as novel, although similar models have been previously published (e.g., Ding et al., 2023). This should be clarified to avoid overstating its novelty.

While the manuscript attempts to address an important question in teleost immunology, the current evidence is insufficient to fully support the authors' conclusions. Addressing the validation of the monoclonal antibody, re-evaluating depletion kinetics, and tempering claims of novelty would strengthen the study's impact. The findings, if rigorously validated, have important implications for understanding the evolution of vertebrate immunity and practical applications in fish health management.

This work is of interest to immunologists, evolutionary biologists, and aquaculture researchers. The methodological framework, once validated, could be valuable for studying immunoglobulin function in other non-model organisms and for developing targeted vaccine strategies. However, the current weaknesses limit its broader applicability and impact.

We would like to thank Reviewer for the helpful comments. As the reviewer suggested, we verified the specificity of anti-bass IgM MoAb using multiple well-established experimental approaches, including mass spectrometry analysis, western blot, flow cytometry, and in vivo IgM depletion models. Additionally, we included western blot analyses to further confirm the IgM depletion kinetics. Moreover, we carefully revised any overstated claims in the original manuscript and incorporated the valuable suggestions of the reviewer in the Introduction and Discussion sections to enhance the clarity and rigor of our work.

Reviewer #1 (Recommendations for the authors):

(1) Experiments and Data Validation:

Monoclonal Antibody Validation:

Provide detailed validation of the monoclonal antibody (MoAb) used for IgM depletion. Perform immunoprecipitation followed by mass spectrometry to confirm the specificity of the MoAb and identify any off-target interactions. Conduct Western blot analysis using tissue lysates with varying IgM, IgT, and IgD expression to demonstrate specificity. Include controls, such as a group treated with a control antibody of the same isotype, to confirm the depletion specificity and effects. Present data on the binding site of the MoAb and confirm it targets IgM.

We thank the reviewer for this constructive comment and have carried out a comprehensive validation of anti-bass IgM monoclonal antibody (MoAb).

Validation of anti-bass IgM MoAb by Mass Spectrometry

To validate the specificity of anti-bass IgM MoAb, target proteins were immunoprecipitated from bass serum using IgM MoAb-coupled CNBr-activated Sepharose 4B beads, followed by mass spectrometry analysis to verify exclusive IgM heavy-chain identification (Figure 3–

figure supplement 1A). Quantitative mass spectrometry verified the antibody's specificity, with IgM heavy-chain peptides representing 97.3% of total signal, indicating negligible off-target reactivity. This high target specificity was further supported by the no detectable cross-reactivity to IgT/IgD (Figure 3–figure supplement 1B). Moreover, the 72% sequence coverage (Figure 3–figure supplement 1C) and confirmed LC-MS/MS spectra of IgM peptides (Figure 3–figure supplement 1D) further validated target selectivity.

Validation of anti-bass IgM MoAb by western blot and flow cytometry

We compared the anti-bass IgM MoAb with an isotype control (mouse IgG1) under both non-reducing and reducing serum immunoblots. The western blot results showed that the developed MoAb bound specifically to IgM in largemouth bass serum. Owing to the structural diversity of fish IgM isoforms, denatured non-reducing electrophoresis typically yields multiple bands with varying molecular weights (Rombout et al., 1993; Ye et al., 2010). Immunoblot analysis revealed multiple bands with varying molecular weights under non-reducing conditions, with the main band ranging from 700 to 800 kDa and a distinct ~70 kDa band under reducing conditions (Figure 3–figure supplement 2A). Notably, the isotype control showed no detectable bands under both non-reducing and reducing conditions (Figure 3–figure supplement 2A). Additionally, we analyzed tissue lysates from various sources (i.e., Spleen, skin, gill, and gut) and observed consistently recognized bands at identical positions and sizes, whereas the isotype control showed no detectable bands (Figure 3–figure supplement 2B–F).


Next, we performed flow cytometry analysis to confirm antibody specificity. In largemouth bass head kidney leukocytes, IgM⁺ B cells accounted for 28.56% of the population, compared to only 0.41% for the isotype control (Figure 3–figure supplement 2G). Following flow sorting of negative and positive cell populations, we extracted RNA from equal cell numbers. Gene expression analysis revealed high expression of IgM and IgD in the positive population, while IgT and T cell markers were absent (Figure 3–figure supplement 2H and I). These results collectively demonstrate that the monoclonal antibody specifically targets largemouth bass IgM.

Validation of the depletion specificity and effects using an isotype-matched control antibody

Largemouth bass (~3 to 5 g) were intraperitoneally injected with 300 µg of mouse anti-bass IgM monoclonal antibody (MoAb, clone 66, IgG1) or an isotype control (mouse IgG1, Abclonal, China). The concentration of IgM in the serum and gut mucus from these MoAb-treated fish was measured by western blot. Our results indicated that anti-bass IgM treatment led to a marked reduction in IgM protein levels in serum (Author response image 1A) and gut mucus (Author response image 1B) from day 1 post-treatment, in contrast to control fish treated with an isotype-matched control antibody.

Author response image 1.

Validation of the depletion specificity and effects using an isotype-matched control antibody. (A, B) The depletion effects of IgM from the serum (A) or gut mucus (B) of control or IgM-depleted fish was detected by western blot. Iso: Isotype group; Dep: IgM-depleted group.



We fully agree with the reviewer that epitope characterization would further validate and elucidate the specificity of IgM MoAb. In the present study, we have demonstrated the antibody's IgM-specific binding through multiple classic experimental methods: (1) mass spectrometry analysis, (2) western blot analysis, (3) flow cytometry analysis, and (4) in vivo IgM depletion models. These results collectively support the conclusion that our MoAb specifically targets IgM. We feel that conformational epitope mapping requires structural biology approaches are out of the scope of this work, although future studies should address them in detail.

Kinetics of IgM Depletion:

Provide additional evidence for the observed rapid depletion of IgM from serum and mucus within one day, as this is inconsistent with previous findings. Include Western blot results to confirm IgM depletion kinetics.

Thanks for the reviewer's suggestion. Previous studies have demonstrated significant differences in the depletion efficiency and persistence of IgM⁺ B cells between warm-water and cold-water fish species. In Nile tilapia (*Oreochromis niloticus*), a warm-water species, administration of 20 µg of anti-IgM antibody resulted in a near-complete depletion of IgM⁺ B cells within 9 days (Li et al., 2023). In contrast, rainbow trout (*Oncorhynchus mykiss*), a cold-water species, required significantly higher doses (200–300 µg) to achieve similar depletion, which persisted in both blood and gut from week 1 up until week 9 post-depletion treatment (Ding et al., 2023). In this study, we investigated largemouth bass (*Micropterus salmoides*), a warm-water freshwater species. Administration of 300 µg of IgM antibody resulted in rapid IgM⁺ B cell depletion from serum and mucus within one day, indicating that the rapid depletion kinetics may be attributed to the combined effects of the elevated antibody dose and the species-specific immunological characteristics. Moreover, we provide a western blot analysis of serum and mucus after IgM depletion as shown in Figure 5–figure supplement 1G and H.

Neutralizing Capacity Assays:

Discuss the potential role of complement or other serum/mucus factors in the neutralization assays. Consider performing neutralization assays that isolate viruses, antibody, and target cells to assess the specific role of IgM.

Thanks for the reviewer's insightful suggestion regarding the potential influence of complement and other serum/mucus factors in our neutralization assays. We sincerely regret that the lack of clarity in our methodological description caused misunderstandings to the reviewer. In fact, prior to performing the virus neutralization assays, serum and mucus samples were heat-inactivated at 56 °C to eliminate potential complement interference. Now, we added the related description of heat-inactivation of serum and mucus samples in the revised manuscript (Lines 727-729). Moreover, our results showed that selective IgM depletion from high LMBV-specific IgM titer mucus and serum samples resulted in significantly increased viral loads and enhanced cytopathic effects (CPE), while no significant difference was observed compared to the control group (shown in Figure 6 of the manuscript).

To further rule out complement or other factors, we purified IgM from serum and gut mucus of 42DPI-S fish for neutralization assays. Briefly, anti-bass IgM MoAb was coupled to CNBr-activated sepharose 4B beads and used for purification of IgM from both serum and gut mucus of 42DPI-S fish. After that, 100 µL of LMBV (1×10^4 TCID₅₀) in MEM was incubated with PBS and purified IgM (100 µg/mL) at 28 °C for 1 hour and then the mixtures were applied to infect EPC cells. Medium or bass IgM was added to EPC cells as controls. We added

the new text in Materials and methods of the revised manuscript in Lines 735-741. Our result showed that a significant reduction in both LMBV-MCP gene expression and protein levels was observed in EPC cells treated with purified IgM from serum (Figure 6–figure supplement 2A, C, and D) or gut mucus (Figure 6–figure supplement 2B, E, and F). Moreover, significantly lower CPE were observed in the IgM treated group, while no CPE was observed in medium and bass IgM group (Figure 6–figure supplement 2G). Collectively, these findings strongly suggest that the neutralization process is a potential mechanism of IgM, serving as a key molecule in adaptive immunity against viral infection. Here, we have incorporated these new findings in the Results section of the revised manuscript (Lines 382-388).

IgT Depletion Model:

To fully establish the role of IgM and IgT in antiviral defense, consider including an experimental group where IgT is depleted.

Thanks for the reviewer's suggestion. The role of IgT in mucosal antiviral immunity in teleost fish has been reported in our previous studies (Yu et al, 2022). However, this study primarily investigates the antiviral function of IgM in systemic and mucosal immunity and further analyzes the mechanisms of viral neutralization. In future research, we plan to establish an IgT and IgM double-depletion/knockout model to further elucidate their specific roles in antiviral immune defense.

(2) Writing and Presentation:

Introduction:

Replace the cited review article on IgT absence with original research articles (e.g., Bradshaw et al., 2020; Györkei et al., 2024) to strengthen the context.

Thank you for your valuable suggestion. We have changed in the revised manuscript (Lines 45-50) as “Notably, while IgT has been identified in the majority of teleost species, genomic analyses reveal its absence in some species, such as medaka (*Oryzias latipes*), channel catfish (*Ictalurus punctatus*), Atlantic cod (*Gadus morhua*), and turquoise killifish (*Nothobranchius furzeri*) (Bengtén et al., 2002; Bradshaw et al., 2020; Magadán-Mompóet al., 2011; Györkei et al., 2024).”

Highlight the evolutionary contrast between the presence of the J chain in older cartilaginous fishes and amphibians and its loss in teleosts. Relevant references include Hagiwara et al., 1985, and Hohman et al., 2003.

Thank you for your valuable suggestion. We have added the relevant description in the revised manuscript (Lines 61-66) “Interestingly, the assembly mechanism of IgM exhibits significant evolutionary variation across vertebrate lineages. In cartilaginous fishes and tetrapods, IgM is secreted as a J chain-linked pentamer, which may enhance multivalent antigen recognition (Hagiwara et al., 1985; Hohman et al., 2003). By contrast, teleosts have undergone J chain gene loss, resulting in the stable of tetrameric IgM formation (Bromage et al., 2004).”

Acknowledge prior studies demonstrating the viral neutralization role of teleost IgM (e.g., Castro et al., 2021; Chinchilla et al., 2013). Avoid overstating the novelty of findings.

Thanks for the reviewer's suggestion. Here, we revised the related description: “More crucially, our study provides further insight into the role of sIgM in viral neutralization and firstly clarified the mechanism through which teleost sIgM blocks viral infection by directly targeting viral particles. From an evolutionary perspective, our findings indicate that sIgM in

both primitive and modern vertebrates follows conserved principles in the development of specialized antiviral immunity.” in the revised manuscript (Lines 20-25) and “To the best of our knowledge, our study provides new insights into the role of sIgM in viral neutralization, suggesting a potential function of sIgM in combating viral infections.” in the revised manuscript (Lines 536-538).

Clarify terms such as "primitive IgM" and avoid misleading evolutionary language (e.g., VLRs are not "candidates"; they mediate adaptive responses).

Thanks for the reviewer’s suggestion. We changed the description of the primitive IgM in the sentence of the revised manuscript as “From an evolutionary perspective, our findings indicate that sIgM in both primitive and modern vertebrates follows conserved principles in the development of specialized antiviral immunity.” in the revised manuscript (Lines 23-25) and “our findings suggest that sIgM in both primitive and modern vertebrates utilize conserved mechanisms in response to viral infections” in the revised manuscript (Lines 574-575). Moreover, we deleted the description of VLRs for “candidates” and rewrote the relevant sentence in the revised manuscript (Lines 37-39) as “Agnathans, the most ancient vertebrate lineage, do not possess bona fide Ig but have variable lymphocyte receptors (VLRs) capable of mediating adaptive immune responses (Flajnik, 2018).”

Results and Discussion:

Address inconsistencies between data and claims, such as the statement that IgM plays a "crucial role" in protection against LMBV, which is not fully supported by mortality data.

Thank you for your insightful comment. We have carefully reviewed our data and revised the language throughout the manuscript to ensure that our claims are fully consistent with the mortality data. We have changed the description of “IgM plays a crucial role in protection against LMBV” as “plays a role” (Line 119), “sIgM participates in” (Line 127), “contributes to immune protection” (Line 507) to more accurately reflect the mortality data

Revise the model in Figure 8 to reflect the concerns raised regarding proliferation data, the role of IgM in protective resistance, and the potential contributions of complement in neutralization assays.

Thank you for your insightful comment. We have added the raised concerns regarding “the viral proliferation data and the role of IgM in protective resistance” in Figure 8 (shown below). Meanwhile, we added relevant descriptions in the figure legends of the revised manuscript (Lines 587-592) as “Upon secondary LMBV infection, plasma cells produce substantial quantities of LMBV-specific IgM. Critically, these virus-specific sIgM from both mucosal and systemic sources has the ability to neutralize the virus by directly binding viral particles and blocking host cell entry, thereby effectively reducing the proliferation of viruses within tissues. Consequently, the IgM-mediated neutralization confers protection against LMBV-induced tissue damage and significantly reduced mortality during secondary infection.”

However, considering the following two reasons: (1) heat-inactivation of serum and mucus samples at 56°C prior to neutralization assays effectively abolished complement activity, and (2) purified IgM from both serum and gut mucus demonstrated comparable neutralization capacity, confirming IgM-dependent mechanisms independent of complement. Therefore, we did not add the potential function of complement in neutralization to Figure 8.

Provide a comparative analysis with other vertebrate models to strengthen the evolutionary implications of findings.

Thank you for your insightful comment. We have added comparative analyses across additional vertebrate models in the discussion of the revised manuscript to enhance the evolutionary perspective of our findings. The details are as follows:

“Virus-specific IgM production has been well-documented in reptiles, birds, and mammals upon viral infection (Dascalu et al., 2024; Harrington et al., 2021; Hetzel et al., 2021; Neul et al., 2017;). While current evidence confirms the capacity of cartilaginous fish and amphibians to mount specific IgM responses against bacterial pathogens and immune antigens (Dooley and Flajnik, 2005; Ramsey et al., 2010), the potential for viral induction of analogous IgM-mediated immunity in these species remains unresolved.” in the revised manuscript (Lines 498-504) and “Extensive studies in endotherms (birds and mammals) have demonstrated that specific IgM contributes to viral resistance by neutralizing viruses (Baumgarth et al., 2000; Diamond et al., 2013; Ku et al., 2021; Hagan et al., 2016; Singh et al., 2022). In contrast, the neutralizing activity of IgM in amphibians and reptiles remains largely unexplored. Although viral infections have been shown to induce neutralizing antibodies in Chinese soft-shelled turtles (*Pelodiscus sinensis*) (Nie and Lu, 1999), the specific Ig isotypes mediating this response have yet to be elucidated. In teleost fish, IgM has been shown to possess viral neutralizing activity similar to that observed in endotherms (Castro et al., 2013; Ye et al., 2013). Furthermore, our recent work demonstrated that secretory IgT (sIgT) in rainbow trout (*Oncorhynchus mykiss*) can neutralize viruses, significantly reducing susceptibility to infection (Yu et al., 2022). However, whether IgM in teleost fish possesses the antiviral neutralizing capacity necessary for fish to resist reinfection remains poorly understood.” in the revised manuscript (Lines 521-534)

Include a description of the Western blot procedure shown in Figures 7D and 7F in the Methods section.

Thank you for your suggestion. A detailed protocol for the western blot experiments presented in Figures 7D and 7F has been added to the Methods section (Western Blot Analysis) in the revised manuscript (Lines 684-687). The details are as follows: Gut mucus, serum, and cells samples were analyzed by western blot as described by Yu et al (2022). Briefly, the samples were separated using 4%–15% SDS-PAGE Ready Gel (Thermo Fisher Scientific, USA) and subsequently transferred to Sequi-Blot polyvinylidene fluoride (PVDF) membranes (Bio-Rad, USA). The membranes were blocked using a 8% skim milk for 2 hours and then incubated with monoclonal antibody (MoAb). For IgM concentration detection, the membranes were incubated with mouse anti-bass IgM MoAb (clone 66, IgG1, 1 µg/mL) and then incubation with HRP goat-anti-mouse IgG (Invitrogen, USA) for 1 hour. IgM concentrations were determined by comparing the signal strength values to a standard curve generated with known amounts of purified bass IgM. For neutralizing effect detection, the membranes were incubated with mouse anti-LMBV MCP MoAb (4A91E7, 1 µg/mL) followed by incubation with HRP goat-anti-mouse IgG (Invitrogen, USA) for 1 hour. The β -actin is used as a reference protein to standardize the differences between samples. Immunoblots were scanned using the GE Amersham Imager 600 (GE Healthcare, USA) with ECL solution (EpiZyme, China).

Ensure all figures are labeled appropriately (e.g., replace "Morality" with "Mortality" in Figure 5A).

Thanks for bringing this to our attention. We have corrected the label in Figure 5A (shown below) and reviewed all figures to ensure that they are appropriately labeled.

(3) Minor Corrections:

Line 117: Correct the typo "across both both."

Thanks for bringing this to our attention. We have changed “across both both” to “across both” in the revised manuscript (Line 119).

| *Line 203: Revise to "IgM plays a role (not crucial role)."*

Thank you for your valuable suggestion. We have modified the description of IgM's role from “crucial” to “plays a role” to better align with our experimental findings in the revised manuscript (Line 202).

| *Line 684: Correct the typo "given an intravenous injection with 200 µg."*

Thanks for bringing this to our attention. We have corrected the phrase to “given an intravenous injection with 200 µg” in the revised manuscript (Line 700-701).

| *Line 686: Fix the sentence fragment "previously. EdU+ cells."*

Thank you for your careful review. We have revised the sentence fragment for clarity in the revised manuscript (Lines 702-703).

| *Abstract and other sections: Adjust language to remove claims of novelty unsupported by data, particularly regarding the role of IgM in viral neutralization.*

Thank you for your constructive feedback. We have thoroughly reviewed and revised the language throughout the abstract and other sections to remove any unsupported claims of novelty, particularly regarding the role of IgM in viral neutralization in the revised manuscript (Lines 20-25).

| *(4) Technical Details:*

| *Verify data availability, including raw data and analysis scripts, in line with eLife's data policies. Include detailed descriptions of all methods, particularly those involving Western blot analysis and antibody validation.*

Thank you for your suggestion. We added the verify data availability, including raw data and analysis scripts as “The raw RNA sequencing data have been deposited in the NCBI Sequence Read Archive under BioProject accession number PRJNA1254665. The mass spectrometry proteomics data have been deposited to the iProX platform with the dataset identifier IPX0011847000.” in the revised manuscript (Lines 808-811).

| *(5) Ethical and Policy Adherence:*

| *Confirm compliance with ethical standards for animal use and antibody development. Ensure proper citation of all referenced works and accurate reporting of prior findings.*

Thank you for your valuable comment. We confirm that our study fully complies with ethical standards for animal use and antibody development. Additionally, we have carefully reviewed the manuscript to ensure that all referenced works are properly cited and that prior findings are accurately reported.

<https://doi.org/10.7554/eLife.104465.2.sa0>