

TRPV4 overactivation enhances cellular contractility and drives ocular hypertension in TGF β 2 overexpressing eyes

Reviewed Preprint

v1 • January 21, 2025

Not revised

Christopher N Rudzitis, Monika Lakk, Ayushi Singh, Sarah N Redmon, Denisa Kirdajova, Yun-Ting Tseng, Michael L De Ieso, W Daniel Stamer, Samuel Herberg, David Križaj 

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States • Department of Neurobiology, University of Utah, Salt Lake City, United States • Department of Ophthalmology and Visual Sciences, SUNY Upstate Medical University, Syracuse, United States • Department of Cell and Developmental Biology, SUNY Upstate Medical University, Syracuse, United States • Department of Ophthalmology, Duke Eye Center, Duke University, Durham, United States • Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, United States • Department of Bioengineering, University of Utah, Salt Lake City, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **fundamental** work extends our understanding of the role of TGF β 2 as a modulator of mechanosensing in the eye and identifies the TRPV4 ion channel as a common regulator of Trabecular Meshwork (TM) contractility and pathological OHT. The data and evidence are **convincing**, with some minor limitations. This work will clearly be of interest to researchers investigating the role of mechanosensors in the TM and may underpin future research into treatments that aim to lower intra ocular pressure. This work will additionally be of interest to the growing field of researchers investigating the regulation of force sensing via ion channels and their roles in health and disease, in particular the ion channel TRPV4.

<https://doi.org/10.7554/eLife.104894.1.sa3>

Abstract

The risk for developing primary open-angle glaucoma (POAG) correlates with the magnitude of ocular hypertension (OHT) and the concentration of transforming growth factor- β 2 (TGF β 2) in the aqueous humor. Effective treatment of POAG requires detailed understanding of interaction between pressure sensing mechanisms in the trabecular meshwork (TM) and biochemical risk factors. Here, we employed molecular, optical, electrophysiological and tonometric strategies to establish the role of TGF β 2 in transcription and functional expression of mechanosensitive channel isoforms alongside studies of TM contractility in biomimetic hydrogels, and intraocular pressure (IOP) regulation in a mouse model of TGF β 2-induced OHT. TGF β 2 upregulated expression of *TRPV4* and *PIEZO1* transcripts and time-dependently augmented functional TRPV4 activation. TRPV4 activation induced TM contractility whereas

pharmacological inhibition suppressed TGF β 2-induced hypercontractility and abrogated OHT in eyes overexpressing TGF β 2. *Trpv4*-deficient mice resisted TGF β 2-driven increases in IOP. Nocturnal OHT was not additive to TGF β -evoked OHT. Our study establishes the fundamental role of TGF β as a modulator of mechanosensing in nonexcitable cells, identifies TRPV4 channel as the final common mechanism for TM contractility and circadian and pathological OHT and offers insights future treatments that can lower IOP in the sizeable cohort of hypertensive glaucoma patients that resist current treatments.

Introduction

Primary open-angle glaucoma (POAG) is an irreversible blinding disease afflicting ~3.5% of the global population (1). Its incidence and severity are proportional to the amplitude and duration of ocular hypertension (OHT) (2, 3), which correlates with retinal ganglion cell dysfunction, neuroinflammation, and oxidative stress (4, 5). Biomechanical factors, glucocorticoids, and the cytokine transforming growth factor- β 2 (TGF β 2) contribute to POAG by compromising the funneling of aqueous humor (AH) from the trabecular meshwork (TM) into the canal of Schlemm (SC). IOP elevations increase the contractility of juxtacanalicular TM (JCT), a circumocular tissue formed by extracellular matrix (ECM) beams populated by mechanosensitive cells smooth muscle-like cells, by increasing its resistance to the flow of aqueous humor (AH). The molecular mechanism that links TM pressure sensing to the contractile response is not known but is likely to underpin the sensitivity to compressive, tensile, osmotic, shear and traction forces to regulate the expression of hundreds TM genes and secretion of dozens of ECM proteins (6–12).

The increase in trabecular outflow resistance induced by mechanical stress, glucocorticoids, and TGF β 2 consists of a dynamic component that can be reversed by cytoskeletal and Rho signaling blockers, and a chronic component, reflecting transdifferentiation of TM cells into fibrotic and contractile myofibroblasts (16–18). One of the clearest examples of fibrotic remodeling in glaucoma pertains to TGF β 2 signaling: (i) TM cells derived from POAG patients secrete more active TGF β 2 compared to cells isolated from healthy donors (19), (ii) the likelihood of contracting POAG is proportional to [TGF β 2]_{AH} (20–22), and (iii) ocular overexpression of TGF β 2 is sufficient to induce OHT (23, 24), presumably due to overexpression of ECM proteins and increased cellular contractility (25, 26). The cognate TGF β 1 isoform similarly induces fibrotic remodeling in fibroblasts, epithelial, and endothelial cells in the heart, kidney, skin, and lung (27–30). TGF β overexpression is thus a causal determinant of OHT that may reflect a universal fibrotic program that, however, cannot be disambiguated from the biomechanical environment. TGF β release is activated by tissue contractility and tension (31, 32), TGF β activity correlates with the distribution of mechanical stress (33), and mechanical stress may induce TGF β -dependent epithelial-mesenchymal transition (EMT; 34, 35).

Our understanding of TM mechanotransduction and its role in IOP homeostasis remains rudimentary despite its overarching clinical relevance. Strain and shear were proposed to stimulate the TM primary cilium, integrins and TRPV4, Piezo1 and TREK-1 channels (36–39) but it is not clear whether mechanosensation regulates TM contractility or is itself impacted by POAG inducers (e.g., TGF β 2 or glucocorticoids) nor is it known how it relates to chronic fibrosis. TRPV4 (Transient Receptor Potential Vanilloid isoform 4), a tetrameric channel with $P_{Ca}/P_{Na} \sim 10$ (40), is strongly expressed in rodent and human TM (36, 51) where it carries the principal component of the pressure-activated transmembrane current and stretch-evoked $[Ca^{2+}]_i$ elevations together with responsiveness to shear and swelling (7, 10, 37, 39, 41, 42). Pharmacological inhibition of the channel and deletion of the TRPV4 gene modulate pressure gradients in the brain, kidney, lung, and bladder (46–50) and mutations in the TRPV4 sequence underpin sensorimotor neuropathies, skeletal dysplasias, retinal degeneration and ocular dysfunction (43–45) while the function of TRPV4 channels in ocular hypertension remains under debate. TRPV4 activity has been implicated in IOP lowering and elevation,

respectively and linked to a diverse array of effector mechanisms that include eNOS and RhoA activation, phospholipid-cholesterol-caveolin regulation, modulation of cell-ECM contacts, primary cilia mechanosensing, polyunsaturated fatty acid release, and Piezo1 signaling (7, 37, 41, 52–55). This invites testable predictions regarding TRPV4 involvement in ocular function and POAG. If TRPV4 maintains steady-state normotension, opposes the profibrotic effects of TGF β 2 and promotes outflow via eNOS-dependent TM relaxation (7, 52), abrogation of its activation should elevate IOP. Conversely, if TRPV4 promotes OHT (37), its inhibition or deletion of the TRPV4 gene should lower IOP.

In this study, we demonstrated novel functions for the TRPV4 channel in homeostatic and pathological IOP regulation by uncovering the mechanisms through which reciprocal TRPV4-TGF β 2 interactions maintain the vicious cycle between mechanical stressors and TM contractility that underlies OHT. We found that inhibition and deletion of TRPV4 lower IOP in TGF β 2 overexpression-induced and circadian models of OHT and suppress TM contractility in TGF β 2-treated biomimetic hydrogels. The cytokine promoted upregulation of EMT-associated genes alongside increased transcription and trafficking of TRPV4, which may have increased the sensitivity of TM cells to innocuous mechanical stimuli. While physiological (nocturnal) and pathological (cytokine-induced) OHT modes both required TRPV4 activity they were not additive, indicating the involvement of a final common pathway.

Collectively, we identify TRPV4 as a fulcrum of TGF β 2-induced TM contractility and IOP regulation and a candidate target for glaucoma therapy.

Results

TGF β 2 drives overexpression of genes that encode fibrotic markers and mechanosensitive ion channels

Human TM cells respond to TGF β 2 with increased biosynthesis, deposition and degradation of ECM, altered autophagy, upregulation of F-actin stress fibers, α -smooth muscle actin (α SMA) (19, 25, 26, 56, 57), but it is unclear whether cells undergoing TGF β 2-induced fibrotic remodeling also exhibit altered capacity for sensing and transduction of mechanical stimuli. We profiled genes that encode known TM mechanochannels together with a selection of key cytoskeletal, ECM, and fibrotic markers in primary TM cells (pTM) isolated from 3–7 donors without history of visual dysfunction (Figure 1A–C). Five-day exposure of pTM cells to a physiological concentration of TGF β 2 (1 ng/mL) increased the expression of EMT-promoting transcription factor SNAI1 (*SNAI1*, $P = 0.0094$) and fibronectin (*FN1*, $P = 0.0263$), while expression of connective tissue growth factor 2 (*CCN2*, alternatively *CTGF*) was elevated in 5/5 pTM cell strains without reaching significance ($P = 0.0909$). Expression of fibroblast-specific protein 1 (*FSP1*, a calcium-binding fibroblast marker), yes-associated protein 1 (*YAP1*, a stiffness induced hippo-pathway transcription factor) and *ACTA2* (α SMA, associated with cell contractility) was not consistently impacted by TGF β 2 while transcription of myocilin (*MYOC*) decreased across 4/4 pTM strains ($P = 0.0055$) (Figure 1B). Indicative of feedback inhibition (58), TGF β 2-treatment downregulated transcript levels of transforming growth factor beta receptor 2 (*TGFBR2*, $P = 0.0219$) and upregulated expression of autoinhibitory SMAD family protein 7 (*SMAD7*, $P = 0.0461$) without affecting *SMAD2* or *SMAD3* expression. TGF β 2 thus promotes selective upregulation of ECM and fibrosis-related genes together with cell dedifferentiation and activation of autoregulatory SMAD mechanisms.

Analysis of genes encoding mechanosensitive channels implicated in outflow modulation (36, 39, 59, 60) showed a 102.5% increase in expression of *TRPV4* ($P = 0.0193$) and 78.9% increase in *PIEZO1* expression ($P = 0.0114$) across 8 replicates including 7 distinct pTM strains. (Figure 1C). Conversely, TGF β 2 exposure did not affect *TRPC1* gene expression ($P = 0.261$) and




Figure 1

TGFβ2 induces a fibrotic phenotype in pTM cells and increases expression and membrane insertion of the TRPV4 channel.

(A-B) Five-day TGFβ2 treatment (1ng/mL) significantly altered expression of TGFβ pathway effectors, cytoskeletal machinery, and canonical fibrotic markers. (C) TGFβ2 treatment significantly increased *TRPV4* and *PIEZO1* expression, but not *TREK1* and *TRPC1* expression. Mean ± SEM shown. N = 4 - 8 experiments, each gene tested in 3-7 different pTM strains (See [Table 1](#)). Two-tailed one sample t-test of TGFβ2-induced gene expression levels as a percent of control samples. (D) Isolation of membrane proteins from two separate pooled pTM samples suggests TGFβ2 treatment drives increased TRPV4 membrane insertion. N = 2 independent pooled samples, 3 pTM strains were pooled per sample. * $P < 0.05$, ** $P < 0.01$.

Location	Donor Age	Donor Sex	Experiments Used
Utah	55	M	PCR, Electrophysiology
Utah	76 (a)	F	PCR, WB, Ca ²⁺ Img.
Utah	76 (b)	F	PCR, Ca ²⁺ Img.
Utah	78	M	PCR, Ca ²⁺ Img.
Utah	64 (a)	F	PCR, WB, Ca ²⁺ Img.
Utah	64 (b)	F	PCR, Ca ²⁺ Img.
Utah	70 (a)	F	PCR, WB, Ca ²⁺ Img., Electrophysiology
Utah	70 (b)	F	PCR
Utah	53	M	Ca ²⁺ Img.
Utah	26	M	Ca ²⁺ Img., Electrophysiology
Utah	73	F	Ca ²⁺ Img.
Utah	56	M	Ca ²⁺ Img.
Utah	73	M	Ca ²⁺ Img.
Utah	80	M	WB
SUNY	39	M	Contractility
SUNY	50	F	Contractility
SUNY	56	F	Contractility

Table 1

Donor information for primary human trabecular meshwork (pTM) strains used in this study.

had variable, strain-dependent effects on transcript levels of *KCNK2* ($P = 0.293$, encoding the TREK-1 channel). Thus, TGF β 2 promotes selective transcriptional upregulation of genes that encode a subset of mechanosensitive proteins alongside fibrotic upregulation and cell dedifferentiation.

TGF β 2 exposure time-dependently augments

TRPV4-mediated current and $\Delta[\text{Ca}^{2+}]_i$

To assess the functional relevance of TGF β 2-dependent transcriptional upregulation we determined the membrane expression and functional activation of TRPV4, which mediates the pressure-activated current and calcium signaling, regulates cytoskeletal dynamics and modulates conventional outflow resistance in vitro (37, 41). Western blots showed that TGF β 2 exposure produces an increase in levels of membrane-bound TRPV4 protein (Figure 1D) in two grouped pTM membrane protein samples. While low amounts of TRPV4 were visible in the membrane fractions in control samples, TGF β 2 treatment produced an increase in the higher weight TRPV4 band, suggesting there could be isoform-specific TGF β 2-induced responses and increased TRPV4 translation leading to elevated TRPV4 trafficking, membrane insertion and/or lipid raft interaction (52).

Functional expression was assessed by tracking $[\text{Ca}^{2+}]_i$ changes in cells exposed to the selective TRPV4 agonist GSK1016790A (GSK101, 10 nM) using Fura2-AM ratiometric Ca^{2+} dye, with TGF β 2-treated and control cells tested on the same day. All pTM strains responded to GSK101 with robust $[\text{Ca}^{2+}]_i$ increases which reached peak within 5 min before the majority of responding cells gradually decreased to a steady plateau (Figure 2C). TGF β 2-treated cells exhibited a remarkable potentiation of GSK101-evoked $[\text{Ca}^{2+}]_i$ responses compared to control cells, with 5/5 cell strains showing an increase in the $\Delta\text{peak}/\text{baseline } F_{340}/F_{380}$ response equivalent to $258.4\% \pm 61.7\%$ of the control response in ($P = 0.0046$) (Figure 2A–B). The fraction of GSK101 responders and the overall time course of responses between groups were not significantly different, indicating that TRPV4 potentiation primarily affects TRPV4-expressing cells. Thus, TGF β 2 treatment promotes TRPV4 expression and functional activity.

To gain insight into the time- and dose-dependence of TGF β 2-dependent TRPV4 signaling modulation pTM cells were treated for 24 hours, at 1 ng/mL and 5 ng/mL concentrations of TGF β 2. GSK101-stimulated Ca^{2+} influx was not significantly increased by 24-hour TGF β 2 treatment at 1 ng/mL ($\Delta\text{peak}/\text{baseline } F_{340}/F_{380} = 117.0\% \pm 23.6\%$ of control) or 5 ng/mL ($\Delta\text{peak}/\text{baseline } F_{340}/F_{380} = 133.6\% \pm 34.5\%$ of control) (Figure 3; SI Appendix, Figure S1); additionally, the potentiation of both was significantly lower relative to the five-day 1 ng/mL TGF β 2 treatment ($P < 0.0011$; Figure 3A). GSK101 evoked a moderately outwardly rectifying nonselective current ($I_{\text{GSK}} - I_{\text{baseline}}$) with reversal potential at ~ 0 mV (Fig 3C). While its amplitude was variable, mean current density consistently increased in cells treated for 1 day with TGF β 2 ($n = 10$; 5 ng/mL) relative to the control group ($n = 11$). The potentiating effect of TGF β 2 on TRPV4 activity thus appears to be time-dependent but is significant after chronic exposure to relatively low-dose TGF β 2.

TGFB2-Induced TM contractility requires TRPV4 activation

The IOP-lowering effectiveness of Rho kinase inhibitors and latrunculins (57, 61–63) indicates that sustained increases in outflow resistance require tonic actin polymerization and contractility. TGF β 2 drives the TM myofibroblast contractile response (57) while the role of mechanosensation remains unknown. To ascertain whether TRPV4 upregulation (Figures 1–2) contributes to the contractile response, we seeded pTM cells into high-compliance Type I collagen hydrogels (57) (Figure 4, SI Appendix, Figure S2). Hydrogels that were incubated with TGF β 2 (5 ng/mL) showed profound increases ($P < 0.0003$) in the rate and the magnitude of contraction at all time points (Figure 4, SI Appendix Figure S2).




Figure 2

TRPV4-mediated Ca^{2+} influx is potentiated by five-day TGF β 2 treatment.

(A) Five-day TGF β 2 treatment (1 ng/mL) increased TRPV4 agonist-induced (GSK101, 10 nM) Ca^{2+} influx in pTM cells compared to serum-free media alone treated cells tested on the same day (N = 5 pTM strains, n = 3 - 5 slides/condition/day, individual data points over mean $\pm$ SEM). Two-tailed one sample t- test of TGF β 2-treated cell average GSK101 response as a percent of control samples from the same pTM strain on the same day. (B) Violin plots showing the distribution of GSK101-induced Ca^{2+} responses for each pTM strain tested in A. Thick dashed line indicates mean, while light dashed line indicates quartiles. (C) Representative traces showing TRPV4 agonist-induced Ca^{2+} influx (seen as an increase in F_{340}/F_{380}) in pTM (mean $\pm$ SEM of 4 representative cells/ group), alongside example Fura-2-loaded pTM cells before (i), during (ii), and after (iii) GSK101 application. Scale bar = 50 μm . ** $P < 0.01$




Figure 3

TGFβ2-induced TRPV4 potentiation is not seen at a shorter time period, regardless of treatment strength.

(A) TGFβ2 treatments for 24 hours at 1 ng/mL (N = 6 pTM strains, n = 3 - 5 slides/condition/day) or 5 ng/mL (N = 5 pTM strains, n = 3 - 5 slides/condition/day) did not show potentiation of GSK101-evoked TRPV4 Ca^{2+} influx (SI Appendix, Figure S1) and were significantly lower than cells treated with TGFβ2 for 5d at 1ng/mL (5d TGFβ2 results from [Figure 2A](#)). Individual data points over mean $\pm$ SEM. One-way ANOVA with Tukey's multiple comparisons test, statistics for individual 1d treatment groups compared to control groups shown in Figure S1. (B) Representative traces for GSK101 response following 24-hour TGFβ2 treatment, traces show mean $\pm$ SEM of 3-4 cells. (C) Average current density in response to GSK101 (24-hour control: n = 11 cells, 24-hour TGFβ2: n=10 cells) shows generally increased current in TGFβ2-treated cells. Data shows mean $\pm$ SEM (D - E) Violin plots of individual cell strains shown in A. Thick dashed line indicates mean, while light dashed line indicates quartiles. ** $P < 0.01$, *** $P < 0.001$




Figure 4
Effects of TRPV4 inhibition/activation on TGFβ2-induced contraction of TM cells.

(A) Representative longitudinal 24-well plate scans of collagen type I hydrogels seeded with pTM subjected to the different treatments (dashed lines outline size of contracted constructs). (B) Longitudinal quantification of hydrogel construct size compared to the control group at the 0 minute time point. (C) Detailed comparisons between groups at each experimental time point. $n = 6$ hydrogels/group. One-way ANOVA with Tukey's multiple comparisons test, data in (B,D) shows individual data points over mean $\pm$ S.D. One pTM strain shown: TGFβ2-induced contractility induction, HC-06-mediated rescue of hypercontractility, and GSK101-induced transient (15 min) contraction were consistent across (3/3) pTM strains tested (Figure S2). ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Simultaneously, treatment with the TRPV4 antagonist HC-067047 (HC-06, 5 μ M) significantly reduced the extent of TGF β 2-induced TM contractility ($P < 0.0001$). To determine whether TRPV4 activation is sufficient to induce the contractile response, the antagonist was washed out and hydrogels supplemented with GSK101 (25 nM). 15 minutes post-treatment, the constructs responded to the agonist with transient contraction (**Figure 4C**; SI Appendix, Figure S2, $P < 0.01$), with a time course mirroring GSK101-induced $[Ca^{2+}]_i$ elevations (**Figures 2–3**). The effects of TRPV4 inhibition and activation were consistent across all pTM strains tested ($N = 3$ pTM strains). TRPV4-mediated Ca^{2+} influx is therefore sufficient to induce TM contractility and necessary for pTM hypercontractility induced by TGF β 2.

TRPV4 activation is required to maintain TGF β 2-induced OHT

To test whether TRPV4 contributes to TGF β 2 induced ocular hypertension (OHT) *in vivo*, we utilized the lentiviral TGF β 2 overexpression model developed by Patil et al. (23). Adult C57BL/6J mice ($N = 5$) were intravitreally injected with lentivirus overexpressing constitutively active human TGF β 2 (LV-TGF β 2). LV-TGF β 2-injected eyes, but not the contralateral eyes injected with a lentivirus containing a scrambled transgene (LV-Ctrl), exhibited significant IOP elevations one-week post-transduction (**Figure 5A**, Week 2, $\Delta_{TGF-Ctrl} = 4.0$ mm Hg, $P = 0.0143$). By 2 weeks post-transfection, IOP in LV-TGF β 2 eyes reached 19.9 ± 4.7 mm Hg whereas IOP in LV-Control eyes remained at control levels (14.0 ± 1.2 mm Hg), with $\Delta_{TGF-Ctrl} = 5.9$ mm Hg ($P = 0.0002$). IOP remained elevated throughout the 4 weeks after the injection (Week 5, $\Delta_{TGF-Ctrl} = 4.9$ mm Hg, $P = 0.0008$). HC-06 (100 μ M) microinjection into the anterior chamber of LV-TGF β 2 and LV-Ctrl eyes lowered IOP in LV-TGF β 2 eyes to 12.2 ± 1.7 mm Hg after 24 hours ($\Delta_{postHC-preHC} = -5.8$ mm Hg) with no difference observed in IOP from LV-Ctrl eyes (12.6 ± 1.9 mm Hg, $\Delta_{post-HC-pre-HC} = -0.3$ mm Hg). LV-Ctrl eyes remained close to pre-injection levels post-HC-06 treatment (**Figure 5A–B**). IOP in LV-TGF β 2 eyes returned to hypertensive levels by 1-week post-HC-06 injection (Week 6–7, $\Delta_{TGF-Ctrl} = 3.9$ mm Hg, $P = 0.0201$). To determine the effect of the bolus injection alone, LV-TGF β 2 and LV-Ctrl eyes were reinjected with PBS 2 weeks after re-establishing the OHT baseline. The sham injection transiently reduced IOP in LV-TGF β 2 ($\Delta_{postPBS-prePBS} = -4.5$ mm Hg) and LV-Ctrl ($\Delta_{postPBS-pre-PBS} = -1.2$ mm Hg) eyes; however, LV-TGF β 2 eyes returned to hypertensive levels by 48 hours post-injection ($\Delta_{TGF-Ctrl} = 3.6$ mm Hg, $p = 0.0465$) and to pre-injection levels after 72 hours ($\Delta_{TGF-Ctrl} = 5.4$ mm Hg, $p = 0.0002$). Bolus injection was less effective than HC-06 at all time points 24 hours post-injection (Week 8–9, **Figure 5B**). These data indicate that selective pharmacological inhibition of TRPV4 effectively and reversibly blocks TGF β 2-induced OHT.

To further evaluate the TRPV4-dependence of TGF β -induced OHT we took advantage of mice with global *Trpv4* knockdown (64–66). *Trpv4*^{−/−} mice ($N = 6$) were injected with LV-TGF β 2 and LV-Ctrl vectors in contralateral eyes (**Figure 5C**). Additionally, two littermate control mice injected alongside the *Trpv4*^{−/−} animals were added to previously collected WT LV-injected cohorts measured at the same timepoints ($N = 8–15$, **Figure 5C**). Pre-LV injection, IOP levels in *Trpv4*^{−/−} animals were comparable to the WT cohort, indicating that TRPV4 activity does not regulate normotension. Similarly, IOP in LV-Ctrl-injected eyes was not significantly different between WT and *Trpv4*^{−/−} animals at any point in the experiment (peak $\Delta_{CtrlKO-CtrlWT} = -1.2$ mm Hg, SI Appendix, **Figure 5D**). By two weeks post-injection (Week 3), LV-TGF β 2-treated *Trpv4*^{−/−} eyes exhibited significantly lower IOP compared to the LV-TGF β 2 WT cohort ($\Delta_{TGFKO-TGFWT} = -3.1$ mm Hg, $P = 0.0009$, **Figure 5C**). While LV-TGF β 2 injected *Trpv4*^{−/−} eyes exhibited mild OHT, the effect was significantly reduced compared to WT eyes and IOP decreased by two weeks post-injection (**Figure 5C–D**).

TGF β 2-induced and nocturnal OHT are non-additive but require TRPV4

Mammalian IOP is modulated by the circadian rhythm, with levels elevated at night and nocturnal IOP fluctuations implicated in POAG (7, 55, 67). We measured nocturnal (9:00–10:00 PM) IOP in LV-TGF β 2 and LV-Ctrl WT eyes ($N = 4$) to determine whether nocturnal OHT is additive to




Figure 5

TRPV4 activation is necessary to maintain LV-TGFβ2-induced ocular hypertension.

(A) Intravitreal injection of LV-TGFβ2 (week 1), but not LV-Control, elevates IOP in WT mice (N = 5 eyes/group) as early as one-week post-injection. Injection of TRPV4 antagonist HC-06, but not PBS, produced multiday IOP reduction in LV-TGFβ2 treated eyes. HC-06 and PBS injections did not affect IOP in LV-Control injected eyes. Two-way ANOVA with Bonferroni post-hoc analysis (B) Direct comparison of the results of PBS and HC-06 injections in the eyes shown in A. Two-way ANOVA with Bonferroni post-hoc analysis (C) Intravitreal injection of LV-TGFβ2 in *Trpv4*^{-/-} mice (N = 6 eyes/group) resulted in only mild OHT; plotted against WT eyes at matching timepoints (3 WT cohorts including the 5 WT eyes shown in A-B, N = 8-15 eyes/group). (D) Statistical comparison of the IOP values shown in C. The IOP in LV-TGFβ2 WT eyes was significantly elevated compared to the LV-TGFβ2 *Trpv4*^{-/-} eyes from 2 weeks post-injection. LV-Control injected eyes in WT or *Trpv4*^{-/-} eyes remain close to the baseline value and are not significantly different. Two-way ANOVA with Bonferroni post-hoc analysis. (A, C) shows mean ± SEM. Data in (B, D) shows individual data points over mean ± SEM, * *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001, **** *P* < 0.0001

elevation observed during the daytime (12:00-2:00 PM, **Figure 6A**), LV-TGFβ2 injected eyes showed significant IOP elevation compared to LV-Ctrl eyes during daytime measurements (Diurnal $\Delta_{\text{TGF}\beta\text{-Ctrl}} = 7.9$ mm Hg, $P < 0.0001$) but the difference vanished at night (Nocturnal $\Delta_{\text{TGF}\beta\text{-Ctrl}} = 0.2$ mm Hg), indicating that TGFβ2-induced OHT is not additive to the circadian OHT. To determine whether physiological (nocturnal) OHT requires TRPV4 we microinjected the eyes of two animals with PBS, and two with HC-06. When IOP stably recovered after the first treatment, the treatment groups were switched. PBS injection did not affect IOP in LV-Ctrl or LV-TGFβ2 eyes at day or night (**Figure 6B-C**) except for a single LV-TGFβ2 eye exhibiting abnormally high nocturnal IOP (37 mm Hg) at the four-day timepoint. Conversely, HC-06 injection blocked LV-TGFβ2-induced IOP during the day ($P < 0.001$) and significantly lowered IOP ~5 mm Hg in both LV-Ctrl and LV-TGFβ2 eyes at night ($P < 0.01$). These data indicate that i) TRPV4 activation is necessary for OHT in the TGFβ2 overexpression mouse model (**Figures 5-6**) and the circadian IOP elevations ii) TGFβ2-evoked OHT does not affect nocturnal IOP elevation in mice, and iii) TRPV4 inhibition does not disrupt the mechanisms that maintain daytime normotensive IOP (**Figures 5-6**).

Discussion

The mechanistic framework developed in this study unifies key biochemical and biomechanical risk factors of POAG to point towards an alternative approach to mitigate fibrotic and functional dysfunction in eyes experiencing OHT. Specifically, we show that TGFβ2 drives overexpression and excessive activation of TRPV4, a Ca^{2+} -permeable channel with diverse mechanosensing functions that include mediating the principal component of the pressure-activated transmembrane current roles in TM cells and fibrotic remodeling across the body (39, 68, 69). Our central observation - that tonic TRPV4 activity is obligatory to maintain TM contractility and OHT induced by angle occlusion and TGFβ2 - identifies a potential molecular linchpin for increased resistance of the JCT TM to AH outflow.

Considering that current glaucoma treatments target secondary outflow mechanisms or are associated with side effects (such as hyperemia) (70, 71), the IOP lowering effected by TRPV4 inhibition and gene knockdown suggests a novel target within the primary outflow pathway that can be engaged without compromising the structural integrity or function of the anterior eye.

Glaucoma is a multifactorial disease with etiology that reflects the convergence of risk factors that include IOP and TGFβ2: the likelihood of POAG correlates with the amplitude of IOP and $[\text{TGF}\beta2]_{\text{AH}}$ (22, 72), and chronic increases in either $[\text{TGF}\beta2]_i$ or IOP promote fibrotic remodeling of the TM/SC and augment the AH flow resistance of the conventional pathway (17, 24, 25). TGFβ2-induced facility suppression has been historically attributed to changes in composition, crosslinking and amount of ECM (25, 26, 73, 74), activation of Hippo signaling and Rho kinase- (Rho/ROCK) mediated contractility (19, 57) and altered expression of genes encoding mitogen-activated protein kinase (MAPK), immune response, oxidative stress, and/or ECM pathways (75-77). Our discovery that TGFβ2 impacts the expression and function of TM mechanosensors and *vice versa*, that TRPV4 is required for TGFβ2-induced contractility, coalesces two key modifiable risk factors (TGFβ2 and pressure) at the level of TRPV4 signaling. Specifically, our data embed TGFβ2 and TRPV4 signaling within reciprocal feedback loops: TGFβ2 (i) induced time-dependent upregulation of TRPV4 mRNA and protein and amplified TRPV4-mediated calcium signaling, while (ii) TRPV4 was required to mediate TGFβ2-induced contractility and maintain chronic OHT in TGFβ2-treated mouse eyes. Microinjection of the selective antagonist HC-06 accordingly reduced IOP in LV-TGFβ2-treated eyes to baseline with hypotension persisting for ~4 days and reversing to pre-injection OHT by day 7. The TRPV4-dependence of TGFβ2-induced OHT and contractility was validated *in vivo* using *Trpv4*^{-/-} eyes and *in vitro* in 3D hydrogel constructs. The differential effectiveness of IOP lowering induced by gene knockdown (~50% reduction in OHT) and pharmacological inhibition (~100% reduction in OHT) may reflect compensatory upregulation of cognate mechanosensory mechanisms in *Trpv4*^{-/-} animals (55).




Figure 6

TRPV4 inhibition inhibits nocturnal IOP elevation in control and TGFβ2 overexpressing eyes.

(A) ~2 months post-LV injection daytime (12-2:00 P.M.) and nocturnal (9-10:00 P.M.) IOP compared in WT mice (N = 4 eyes/group) before drug treatment. LV-TGFβ2 eyes were elevated at daytime, but nocturnal OHT was not significantly different between LV-Ctrl and LV-TGFβ2 eyes. One-way ANOVA with Tukey's multiple comparisons test. (B-C) PBS-injected eyes did not exhibit changes in daytime or nighttime intraocular pressure; however, HC-06 injection reduced TGFβ2-induced IOP elevations during the day and LV-Ctrl and LV-TGFβ2 nocturnal IOPs (N = 4 Eyes/Group); Two-way ANOVA with Bonferroni post-hoc analysis. Figures show datapoints over mean ± SEM, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$

We've previously shown that TRPV4 channels in primary human TM cells are activated by physiological (5 – 25 mm Hg) pressure steps (39, 60) and (1 – 12%) strains (37, 41), to activate downstream outflow-relevant signaling mechanisms such as Rho kinase, F-actin, tyrosine phosphorylation of FAK, paxillin and vinculin, reorganization of membrane lipids, and ECM release (37, 41, 52). Here, we extend those observations to demonstrate that TRPV4 activation is required for TM contractility and ocular hypertension induced by TGFβ2 overexpression and circadian rhythmicity. The observation that TRPV4 activity underpins increased outflow resistance under physiological and pathological conditions resolves contradictory conclusions from prior investigations, which implicated TRPV4 signaling in ocular hypertension and hypotension, respectively (7, 36, 37, 39, 41, 52, 53, 55). TRPV4 has been proposed to lower IOP through phosphoinositide signaling in primary cilia (36), stimulate TM-resident endothelial nitric oxide synthase (eNOS) (7) and release of polyunsaturated fatty acids (PUFAs) (53) and/or activate downstream from Piezo1 mechanosensing (54). However, TRPV4-regulated Ca^{2+} influx in TM cells is unaffected by the ablation of primary cilia (37), eNOS expression in TM cells is vanishingly low (78–80), PUFAs such as arachidonic acid and EETs stimulate rather than inhibit, TRPV4 (37) and TRPV4 signaling in TM cells is unaffected by Piezo1 inhibitors and knockdown (39). Moreover, Piezo1 inhibition reduces outflow facility under *in vitro* and *in vivo* conditions (39, 81), indicating opposing homeostatic functions for Piezo1 vs. TRPV4 activation. The TRPV4-dependence of TM contractility (Figure 4) accords with reports that TRPV4 inhibition increases, and TRPV4 activation reduces outflow facility in biomimetic TM-populated scaffolds in the absence of ciliary body, Schlemm's canal, and ciliary muscle (37). Induction of the contractile response by GSK101 and its inhibition of hypercontractility by HC-06 further suggest a model whereby TRPV4 pressure transduction drives Ca^{2+} - and Rho-dependent hypercontractility and fibrosis via actin polymerization, myosin light chain phosphorylation, αSMA integration into stress fibers and reinforcement of focal ECM contacts (41, 82, 83) (Figure 7). TRPV4 channels in cells treated with TGFβ2 are likely to be constitutively active at incubator temperature, which coincides with peak TRPV4 thermoactivation (~34 – 37°C) (84, 85). The residual contractility in HC-06-treated cells may reflect TGFβ2-mediated contributions from Piezo1, TRPC, and/or TREK-1 channels and/or intracellular Ca^{2+} release (37, 60, 86, 87). Reports from heart, lung, liver, skin and articular cartilage preparations similarly implicate TRPV4 in TGFβ1-dependent fibrosis (68, 88–91) and bladder (92), heart (93, 94), and vascular (95) contractility, with conditional ablation of TRPV4 from smooth muscle cells shown to lower blood pressure (96, 97). TGFβ2-stimulated induction of *FN1*, *SNAIL1*, and *CTGF* transcripts (Figure 1) accords with RNA profiling studies which documented the cytokine's key role in TM transdifferentiation towards the contractile myofibroblast state (25, 75, 98–103) whereas the decreased expression of *TGFBR2* and increased the abundance of *SMAD7* mRNA indicate activation of autoinhibitory mechanisms in EMT-undergoing cells (104).

Treatment of TM cells with TGFβ2 concentrations comparable to those found in POAG AH (0.2–3.2 ng/ml) (20) produced 2–3-fold upregulation of TRPV4 transcripts, protein, and responsiveness to GSK101, with the time course of these effects mirroring facility reduction in human eyes treated with exogenous cytokine (105). A single exposure to 5 ng/ml TGFβ2 approximately doubled the amplitude of the GSK101-evoked current and reduced outward rectification of I_{TRPV4} (Figure 3). Difficulties with giga-ohm seal formation precluded I_{TRPV4} analyses at longer incubation times but we were able to obviate this limitation with imaging experiments, which revealed robust and reproducible time-dependent increases in the amplitude of agonist-induced Ca^{2+} signals across all 5 studied strains (Figure 2). The effects of TGFβ2 on I_{TRPV4} , membrane protein levels and $[\text{Ca}^{2+}]_{\text{GSK}}$ accord with increased expression of the TRPV4 gene, with precedents from other cell types (e.g., fibroblasts) suggesting the possibility of increased trafficking of TRPV4–PI3K complexes and/or β-arrestin 1-dependent ubiquitination (106, 107). The upregulation of TRPV4/Piezo1 transcription predicts that TGFβ2-exposed cells might exhibit exaggerated responsiveness to mechanical loading and mechanical hyperalgesia, as reported for chemotherapy (108), neuropathic pain (109, 110), cancer (111), and diabetic neuropathy (112).




Figure 7

TGFβ2-TRPV4 interactions in TM remodeling and ocular hypertension.

Chronic exposure to TGFβ2 induces upregulation of functional TRPV4 channels alongside the autoinhibitory canonical modulator SMAD7. TRPV4-mediated Ca²⁺ influx, canonical, and non-canonical TGFβ2 signaling stimulate Rho/ROCK signaling, augment cytoskeletal contractility, and stimulate ECM release to increase the flow resistance of the conventional pathway. Increased contractility drives OHT, resulting in a feedforward vicious TRPV4-dependent circle loop that maintains ocular hypertension. Schematic made using Biorender.com.

The lack of additivity between TGF β 2-induced OHT and nocturnal OHT (**Figure 6**) suggests that control of physiological and pathological hypertension converges at the level of TRPV4-Rho signaling as the final common mechanism obligatory for OHT. This conjecture is supported by the observations that TRPV4 inhibitors, ROCK inhibitors and TM-specific expression of dominant negative scAAV2.*dnRhoA* constructs lower IOP, elevated through occlusion of the iridocorneal angle, TGF β , glucocorticoids and the nocturnal cycle (55, 62, 113). Future studies of TRPV4 signaling will investigate the mechanisms that underlie the reversibility of circadian TRPV4 activation and how TGF β 2 impairs this physiological process. For example, the suprachiasmatic nucleus and the hypothalamus–pituitary–adrenal axis (114, 115) may modulate the TRPV4-Rho axis via nocturnal release of norepinephrine and melatonin (116, 117). It is worth noting that TRPV4 may directly bind to membrane proteins known to regulate conventional outflow, such as β 1 integrins (118), caveolin-1 (52), and cytoskeletal proteins (actin, actin adaptor proteins, microtubules) (119).

Our study unifies biomechanical and biochemical paradigms of fibrotic and functional remodeling in glaucoma, expands the biological significance of TGF β 2 modulation by including increased actomyosin contractility in addition to fibrotic remodeling, thereby opening a new window into the mechanisms that subserve physiological vs. pathological OHT. We propose that TGF β 2 shifts the homeostatic normotensive setpoint maintained by steady-state TRPV4, Piezo1 and TREK-1 activation (39, 60, 81) through upregulation of TRPV4 expression, which increases the cells' sensitivity to pressure and strain under normotensive conditions. In addition to hijacking the cells' contractile apparatus, TGF β 2 overexposure induces fibrosis that may facilitate the pull of stress fibers on the increasingly “stiff” ECM (19) together with increased production and secretion of ECM. The absence of structural and functional visual phenotypes in TRPV4 KO mice (55, 66, 120) predicts that small-molecule TRPV4 antagonists might lower IOP, suppress fibrosis and protect retinal neurons without compromising homeostatic IOP regulation (70). The similarities between TRPV4 expression in mouse and human TM (37) and between outflow mechanisms in mice vs. humans (121, 122) suggest that TRPV4 targeting might be explored within the clinical context.

Methods

Animals

C57BL/6J mice were from JAX laboratories (Bar Harbor, ME), *Trpv4*^{-/-} mice were a gift from Wolfgang Liedtke (Duke University) (64, 65). The animals were maintained in a pathogen-free facility with a 12-hour light/dark cycle and ad libitum access to food and water, at a temperature of ~22–23°C. Mice were 2–6 months in age prior to LV-injection; data from both male and female sexed animals were included in this study.

Human TM Culture

De-identified postmortem eyes from donors with no history of glaucoma (pTM cells) were procured from Utah Lions Eye Bank with written informed consent of the donor's families. TM cells were isolated from juxtacanalicular and corneoscleral regions as previously described (37, 39), in accordance with consensus characterization recommendations (123). pTM cells were cultured in Trabecular Meshwork Cell Medium (TMCM; Sciencell) in Collagen-I (Corning) coated culture flasks and glass coverslips at 37°C in a humidified atmosphere with 5% CO₂. Fresh media was supplied every 2–3 days. Serum free (SF) media was mixed as needed by excluding fetal bovine serum (FBS, Sciencell) from the TMCM. A list of all pTM strains used is available in **Table 1**; all cells were used between passages 2–4. Cell lines were chosen based on availability at the time of experiments.

For contractility experiments pTM cells were isolated from healthy donor corneal rims discarded after transplant surgery, as previously described (19, 57, 124), and cultured according to established protocols (123, 125). Three pTM cell strains isolated from healthy donors and validated with dexamethasone-induced myocilin expression were used for contractility experiments. pTM cells were cultured in low-glucose Dulbecco's Modified Eagle's Medium (DMEM; Gibco; Thermo Fisher Scientific) containing 10% fetal bovine serum (FBS; Atlanta Biologicals) and 1% penicillin/streptomycin/glutamine (PSG; Gibco) and maintained at 37°C in a humidified atmosphere with 5% CO₂. Fresh media was supplied every 2-3 days.

The experiments were conducted according to the tenets of the Declaration of Helsinki for the use of human tissue.

Reagents

The TRPV4 antagonist HC-067047 (HC-06) was purchased from Millipore-Sigma or Cayman Biotech and dissolved in DMSO at 20mM. The TRPV4 agonist GSK1016790A (GSK101; Cayman Biotech) was dissolved in DMSO at 1mM. Aliquots were diluted into working concentrations (10-25 nM, GSK101; 5- 100 μM, HC-06). Recombinant human TGFβ2 protein was ordered from R&D Systems and reconstituted in sterile 4 mM HCl with 0.1% BSA at 20 ug/mL.

Quantitative Real-Time PCR

Gene-specific primers were used to detect expression of target genes, as described (126). Total RNA was isolated using the Arcturus PicoPure RNA isolation kit (ThermoFisher Scientific). cDNA was generated from total RNA using qScript XLT cDNA Supermix (Quanta Biosciences). SYBR Green based real-time PCR was performed with 2X GREEN Master Mix (Apex BioResearch Products). Gapdh was used as an endogenous control to normalize fluorescence signals. Gene expression relative to GAPDH was measured using the comparative CT method ($2^{-\Delta\Delta C_T}$), All genes were assessed in 4-8 individual samples taken from 3-7 different pTM strains. The primer sequences, expected product length, and gene accession are given in **Table 2**.

Western Blot

3 SF- or TGFβ2-treated samples were pelleted and pooled together from 3 different pTM samples within the same condition. To separate membrane proteins from heavier cellular debris the pooled cell pellets were homogenized in a hypotonic lysis buffer (20mM TRIS-HCl, 3mM MgCl₂, 10mM NaCl, 10mM PMSF, 0.5 mM DTT, 20 mM NaF, 2 mM NaV, 0.5 μg/mL leupeptin) before centrifuging at 300x g for 5 minutes (4 °C). The resulting supernatant was removed and centrifuged again at >12,500 rpm for 30 minutes to pellet membrane proteins, which were then resuspended in RIPA Buffer (Santa Cruz). Proteins were separated on a 10% SDS-PAGE gel and transferred to polyvinylidene difluoride membranes (Bio- Rad). Membranes were blocked with 5% skim milk/2% BSA in TBST and incubated at 4 °C overnight with a primary antibody against TRPV4 (1:250, Alomone Labs #ACC-034) or rabbit antibody against β-tubulin (1:2000, Abcam #EPR1330). Appropriate secondary antibodies conjugated to HRP were used to visualize protein expression on an iBright CL750 imaging system (Thermo Fisher Scientific). β-Tubulin expression was used to standardize protein levels between samples.

Calcium Imaging

pTM cells were seeded onto Collagen-I (Corning) coated coverslips and cultured in TCM media (ScienCell) as described (39, 41). The cells were serum starved for 24 hours followed by serum-free TCM with or without TGFβ2 (1 or 5 ng/mL) for 24 hours or five days. The cells were loaded with 10 μM of the ratiometric indicator Fura-2 AM (K_d at RT = 225 nM (Invitrogen/ThermoFisher) for 30-60 minutes. Coverslips were placed in a RC-26G chamber

Gene	Forward	Reverse	Product Length (bp)	NCBI reference number
<i>GAPDH</i>	CTCCTGTTTCGACAGTCAGCC	GACTCCGACCTTCACCTTCC	89	NM_002046.5
<i>SMAD2</i>	GGGTTTTGAAGCCGTCTATCA GC	CCAACCACTGTAGAGGTCCATTC	149	NM_005901.6
<i>SMAD3</i>	CAAGTGGCCGCGTGTAATAA	AGTCCAGAACAGCCGAGTTG	181	NM_005902.4
<i>SMAD7</i>	CTGCTCCCATCCTGTGTGTT	CCTTGGGTATGACGGACCA	120	NM_005904.3
<i>TGFBR2</i>	AACCTCTAGGCACCTCCTC	AACCTCTAGGCACCTCCTC	100	NM_001024847.3
<i>FSP1</i>	GCTTCTTCTTCTTGGTTTGAT CCT	AAGTCCACCTCGTTGCCCT	250	NM_002961.3
<i>SNAIL1</i>	GGCTCCTTCGTCCTTCTCCTCT AC	CTGGAGATCCTTGGCCTCAGAGA G	124	NM_005985.4
<i>CCN2</i>	CCCCAGACACTGGTTTGAAG	CCCACTGCTCCTAAAGCCAC	100	NM_001901.3
<i>YAP1</i>	ACAGGGAAGTGACTTTGTAC A	GCACTGAATATTGCACCCAC	183	NM_001130145.
<i>FN1</i>	CTGAAAGACCAGCAGAGGCA	GTGTAGGGGTCAAAGCACGA	110	M10905.1
<i>SMA (ACTA2)</i>	GTCACCCACAATGTCCCAT	GGAATAGCCACGCTCAGTCA	123	NM_001141945.2
<i>MYOC</i>	CCACGTGGAGAATCGACACA	TCCAGTGGCCTAGGCAGTAT	118	NM_000261.1
<i>TRPV4</i>	TCCCATTCTTGCTGACCCAC	AGGGCTGTCTGACCTCGATA	217	NM_021625.4
<i>PIEZO1</i>	GGCCAACTTCTCACCAAGA	GGGTATTTCTTCTGTCTC	106	NM_001142864.3
<i>TREK1</i>	AGGGATTCTACTTGGCGGC	CAAGCACTGTGGGTTTCGTG	99	NM_001017424.3
<i>TRPC1</i>	TGCGTAGATGTGCTTGGGAG	CGTTCATTAGTTCTGACAACCG	107	X89066.1

Table 2

Sequences, product size, and reference numbers for PCR Primers used in this study.

platform (Warner Instrument Corp) and perfused with external saline (pH 7.4) (in mM): 80 NaCl, 4.7 KCl, 1.2 MgCl₂, 10 D-Glucose, 19.1 HEPES sodium salt, 2 CaCl₂ and osmolality adjusted to 300 mOsm using D-mannitol. External solutions were delivered via a manually controlled gravity-fed eight-line manifold system, with perfusion speed kept constant to minimize changes in shear. Epifluorescence imaging was performed using an inverted Nikon Ti microscope with a 40x 1.3 N.A. oil objective and Nikon Elements AR software. 340 nm and 380 nm excitation were delivered by a high-intensity 150W Xenon arc lamp (Lambda DG-4; Sutter Instruments), high pass-filtered at 510 nm and detected with a 12-bit Delta Evolve camera (Photometrics/Teledyne).

GSK101 (10 nM) evoked $\Delta[\text{Ca}^{2+}]_i$ was assessed as $\Delta R/R$ (dividing the difference between the peak GSK- evoked F_{340}/F_{380} signal during stimulation and baseline F_{340}/F_{380} signal by the baseline F_{340}/F_{380} signal).

Every data point represents a separate experimental day and pTM cell strain, each with 3-5 control and 3- 5 TGF β 2-treated slides tested on the same day. TGF β 2 datapoints represent the average GSK101 evoked $\Delta R/R$ across all TGF β 2 cells as a % of the average $\Delta R/R$ of control cells from the same cell strain on the same day.

Collagen hydrogel contraction assay

Rat tail collagen type I (Corning, Thermo Fisher Scientific) was prepared at a concentration of 1.5 mg/ml according to the manufacturer's instructions. Five hundred microliters of the hydrogel solution were pipetted into 24-well culture plates. Upon complete collagen polymerization, pTM cells were seeded at 1.5×10^5 cells/well atop the hydrogels and cultured in DMEM + 10% FBS + 1% PSG for 48 hours to facilitate even cell spreading. Next, constructs were cultured in serum-free DMEM + 1% PSG supplemented with: i) control (vehicle: 0.008 mM HCl + 0.0004% BSA; 0.025% DMSO), ii) TGF β 2 (5 ng/ml; R&D Systems), or iii and iv) TGF β 2 + HC067047 (5 μ M in DMSO) for 36 hours before carefully releasing the hydrogels from the walls using a sterile 10 μ l pipette tip to facilitate contraction. The next morning, fresh serum-free DMEM + 1% PSG supplemented with 0.0025% DMSO (= vehicle) was added to groups i-iii; group iv received serum-free DMEM + 1% PSG supplemented with GSK1016790A (25 nM in DMSO). Plates were longitudinally imaged at 600 dpi resolution with a CanoScan LiDE 300 flatbed scanner (Canon USA) at 0, 15, 30, 60, and 120 minutes. Hydrogel construct size was quantified using FIJI software (National Institutes of Health) ([127](https://doi.org/10.1371/journal.pone.0174311) [↗](#)).

Electrophysiology

Borosilicate patch-clamp pipettes (WPI) were pulled using a P-2000 horizontal micropipette puller (Sutter Instruments), with a resistance of 6-8 M Ω . The internal solution contained (mM): 125 K-gluconate, 10 KCl, 1.5 MgCl₂, 10 HEPES, 10 EGTA, pH 7.4. Patch clamp data was acquired with a Multiclamp 700B amplifier, pClamp 10.6 software and Digidata 1440A interface (Molecular Devices), sampled at 5kHz and analyzed with Clampfit 10.7. Current-voltage relationships were assessed using V_m steps from -100 to + 100 mV against a holding potential of -30 mV. Current density was measured as the average current during GSK101 exposure subtracted by the average current from the same cell during baseline perfusion.

IOP Measurements

A TonoLab rebound tonometer (Colonial Medical Supply) was used to measure IOP of awake mice between 12-2 P.M. IOP was determined from the mean of 10-20 tonometer readings. Nocturnal measurements were conducted between 9-10 P.M. under 2.5% isoflurane delivered by a Somnosuite isoflurane vaporizer (Kent Scientific). After animals recovered from intracameral HC-06/PBS injections, IOP was measured daily. IOP was measured every day for 4-5 consecutive days to confirm a stable return to baseline. IOP data for individual cohorts was binned into weeks of experimental time to group values for analysis.

Lentiviral Transduction

Lentiviral stock for TGFβ2 (C226,228S) was purchased from VectorBuilder Inc. (VB170816-1094fnw, pLV[Exp]-CMV> {hTGFB2[NM_003238.3](C226,228S)}) (23 [DOI](#)). Scrambled control lentivirus was purchased from SignaGen Laboratories (LM-CMV-Null-Puro). Mice were anesthetized with an intraperitoneal IP injection of ketamine/xylazine (90 mg/10 mg/ kg body weight), followed by eyedrops containing 0.5% proparacaine hydrochloride and 1% tropicamide ophthalmic solution to numb the eyes and dilate the pupils. Anesthetized mice were secured to allow stereotaxic injection of lentivirus.

Intravitreal injections were conducted by creating a guide hole with a 30-gauge needle 1-2 mm equatorial of the cornea-scleral border followed by insertion of a 12° beveled 33-gauge Hamilton syringe (Hamilton Company) secured to a stereotaxic rig (World Precision Instruments) used to insert the needle 2-3 mm into the eye. Each eye was injected with a 2μL bolus of lentivirus diluted to 1×10^6 TU/μL over the course of one minute, before the needle was quickly drawn and the pilot hole treated with erythromycin ophthalmic ointment USP (Bausch & Lomb). The efficiency of LV-TGFβ2 OHT induction in WT animals was close to 100%. No differences in observable health post-injection were detected between wild type and *Trpv4*^{-/-} animals or LV-Ctrl and LV-TGFβ2 injected animals.

Intracameral Microinjections

Mice were anesthetized and treated with eyedrops as above, before being placed on an isothermal heating pad. HC-06 (100 μM) or PBS with DMSO (0.5%) as a vehicle were injected into the anterior chamber using a blunt tip Hamilton syringe (Hamilton Company) through a guide hole made using a 30-gauge needle. At the end of each injection a small air bubble was introduced to seal the cornea and minimize fluid outflow. 0.5% Erythromycin ophthalmic ointment USP (Bausch & Lomb) was applied to the eye after the procedure. Intracameral injections were not associated with observable inflammation, corneal opacity or behavioral changes. For the nocturnal IOP experiments in **Figure 6** [DOI](#), two animals were injected with PBS while two were injected with HC-06. When OHT was stably reestablished a week post- injection, the treatment groups were switched, and experiments repeated to obtain four eyes/treatment group for **Figure 6C-D** [DOI](#).

Statistical Analysis

GraphPad Prism 9 was used for statistical analysis. Means are plotted ± SEM unless otherwise noted. One-sample t-tests were used to determine whether TGFβ2 treated groups were significantly different than untreated control groups, while one-way ANOVA or two-way ANOVA along with Tukey or Bonferroni's multiple comparisons test were used to compare multiple groups.

Data availability

Individual datapoints for in-vivo figures, and unedited/uncropped annotated western blot images, are included in the supplementary data files for this manuscript. Further information about the data presented in this manuscript is available from the corresponding authors upon reasonable request.

Acknowledgements

We thank Dr. Paloma Liton (Duke University) for the generous gift of LV-TGF β 2^{C226,228S} lentivirus stock during the pilot stages of this experiment, and Dr. Gulab Zode for the availability of the LV-TGF β 2^{C226,228S} construct on Vectorbuilder. We additionally thank Dr. Wolfgang Liedtke (Duke University and Regeneron) for *Trpv4*^{-/-} mice.

The study was supported by the National Institutes of Health (T32EY024234 to CNR and DK, R01EY022076, R01EY1031817, P30EY014800 to DK, R01EY034096 to SH, R01EY022359, R01EY005722 to WDS), Crandall Glaucoma Initiative, Stauss-Rankin Foundation, and Unrestricted Grants from Research to Prevent Blindness to Ophthalmology Departments at the University of Utah and Duke University.

Additional information

Study Approval

The animal experimental protocols were conducted in accordance with the NIH Guide for the Care and Use of Laboratory Animals and the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research and were approved by the Institutional Animal Care and Use Committee at the University of Utah.

Author Contributions

C.N.R. and D.K. designed the primary research study. C.N.R., M.L., A.S., S.N.R., D.K., Y.T.T., M.L. performed research, C.N.R., S.N.R., Y.T.T., S.H., D.K. analyzed the data, and C.N.R. and D.K. wrote the paper.

References

1. Tham Y. C., et al. (2014) **Global prevalence of glaucoma and projections of glaucoma burden through 2040: A systematic review and meta-analysis** *Ophthalmology* **121**:2081–2090
2. Gordon M. O., et al. (2002) **The Ocular Hypertension Treatment Study: Baseline factors that predict the onset of primary open-angle glaucoma** *Arch. Ophthalmol* **120**:714–720
3. Heijl A., et al. (2002) **Reduction of intraocular pressure and glaucoma progression: Results from the Early Manifest Glaucoma Trial** *Arch. Ophthalmol* **120**:1268–1279
4. Almasieh M., Wilson A. M., Morquette B., Cueva Vargas J. L., Di Polo A. (2012) **The molecular basis of retinal ganglion cell death in glaucoma** *Prog. Retin. Eye Res* **31**:152–181
5. Baudouin C., Kolko M., Melik-Parsadaniantz S., Messmer E. M. (2021) **Inflammation in Glaucoma: From the back to the front of the eye, and beyond** *Prog. Retin. Eye Res* **83**
6. Ellingsen B. A., Grant W. M. (1971) **The relationship of pressure and aqueous outflow in enucleated human eyes** *Invest. Ophthalmol* **10**:430–437
7. Patel P. D., et al. (2021) **Impaired TRPV4-eNOS signaling in trabecular meshwork elevates intraocular pressure in glaucoma** *Proc. Natl. Acad. Sci. U. S. A* **118**
8. De Kater A. W., Shahsafaei A., Epstein D. L. (1992) **Localization of smooth muscle and nonmuscle actin isoforms in the human aqueous outflow pathway** *Investig. Ophthalmol. Vis. Sci* **33**:424–429
9. Krizaj David, Samples J. R., Knepper P. A. (2020) **No cell is an island: trabecular meshwork ion channels as sensors of the ambient milieu** *Glaucoma Research and Clinical Advances: 2020 to 2022* Kugler Publications :1–10
10. Baumann J. M., et al. (2024) **TRPV4 and chloride channels mediate volume sensing in trabecular meshwork cells** *Am. J. Physiol. Physiol* <https://doi.org/10.1152/ajpcell.00295.2024>
11. Brubaker R. F. (1975) **The effect of intraocular pressure on conventional outflow resistance in the enucleated human eye** *Invest. Ophthalmol* **14**:286–292
12. 12., Karimi A., et al. (2022) **The Effect of Intraocular Pressure Load Boundary on the Biomechanics of the Human Conventional Aqueous Outflow Pathway** *Bioengineering* **9**
13. Borrás T. (2003) **Gene expression in the trabecular meshwork and the influence of intraocular pressure** *Prog. Retin. Eye Res* **22**:435–463
14. Ramos R. F., Sumida G. M., Daniel Stamer W. (2009) **Cyclic mechanical stress and trabecular meshwork cell contractility** *Investig. Ophthalmol. Vis. Sci* **50**:3826–3832
15. Luna C., Li G., Liton P. B., Epstein D. L., Gonzalez P. (2009) **Alterations in gene expression induced by cyclic mechanical stress in trabecular meshwork cells** *Mol. Vis* **15**:534–544

16. Acott T. S., Vranka J. A., Keller K. E., Raghunathan V. K., Kelley M. J. (2021) **Normal and glaucomatous outflow regulation** *Prog. Retin. Eye Res* **82**
17. Fuchshofer R., Tamm E. R. (2012) **The role of TGF- β in the pathogenesis of primary open-angle glaucoma** *Cell Tissue Res* **347**:279–290
18. Johnstone M. A. (2014) **Intraocular pressure regulation: Findings of pulse-dependent trabecular meshwork motion lead to unifying concepts of intraocular pressure homeostasis** *J. Ocul. Pharmacol. Ther* **30**:88–93
19. Li H., Raghunathan V. K., Stamer W. D., Ganapathy P. S., Herberg S. (2022) **Extracellular Matrix Stiffness and TGF β 2 Regulate YAP/TAZ Activity in Human Trabecular Meshwork Cells** *Front. Cell Dev. Biol* **10**
20. Agarwal P., Daher A. M., Agarwal R. (2015) **Aqueous humor TGF- β 2 levels in patients with open-angle glaucoma: A meta-analysis** *Mol. Vis* **21**:612–620
21. Ochiai Y., Ochiai H. (2002) **Higher concentration of transforming growth factor- β in aqueous humor of glaucomatous eyes and diabetic eyes** *Jpn. J. Ophthalmol* **46**:249–253
22. Tripathi R. C., Li J., Chan W. F. A., Tripathi B. J. (1994) **Aqueous humor in glaucomatous eyes contains an increased level of TGF- β 2** *Exp. Eye Res* **59**:723–728
23. Patil S. V., Kasetti R. B., Millar J. C., Zode G. S. (2022) **A Novel Mouse Model of TGF β 2-Induced Ocular Hypertension Using Lentiviral Gene Delivery** *Int. J. Mol. Sci* **23**
24. Shepard A. R., et al. (2010) **Adenoviral gene transfer of active human transforming growth factor- β 2 elevates intraocular pressure and reduces outflow facility in rodent eyes** *Investig. Ophthalmol. Vis. Sci* **51**:2067–2076
25. Fleenor D. L., et al. (2006) **TGF β 2-induced changes in human trabecular meshwork: Implications for intraocular pressure** *Investig. Ophthalmol. Vis. Sci* **47**:226–234
26. Montecchi-Palmer M., et al. (2017) **TGF β 2 induces the formation of cross-linked actin networks (CLANs) in human trabecular meshwork cells through the smad and non-smad dependent pathways** *Investig. Ophthalmol. Vis. Sci* **58**:1288–1295
27. Coker R. K., et al. (1997) **Transforming growth factors- β 1, - β 2, and - β 3 stimulate fibroblast procollagen production in vitro but are differentially expressed during bleomycin-induced lung fibrosis** *Am. J. Pathol* **150**:981–991
28. Santiago B., et al. (2005) **Topical application of a peptide inhibitor of transforming growth factor- β 1 ameliorates bleomycin-induced skin fibrosis** *J. Invest. Dermatol* **125**:450–455
29. Yue Y., Meng K., Pu Y., Zhang X. (2017) **Transforming growth factor beta (TGF- β) mediates cardiac fibrosis and induces diabetic cardiomyopathy** *Diabetes Res. Clin. Pract* **133**:124–130
30. Zhang Y., et al. (2020) **Overexpression of TGF- β 1 induces renal fibrosis and accelerates the decline in kidney function in polycystic kidney disease** *Am. J. Physiol. - Ren. Physiol* **319**:F1135–F1148
31. Walker M., Godin M., Pelling A. E. (2020) **Mechanical stretch sustains myofibroblast phenotype and function in microtissues through latent TGF- β 1 activation** *Integr. Biol* **12**:199–210

32. Wipff P. J., Rifkin D. B., Meister J. J., Hinz B. (2007) **Myofibroblast contraction activates latent TGF- β 1 from the extracellular matrix** *J. Cell Biol* **179**:1311–1323
33. Zhen G., et al. (2021) **Mechanical stress determines the configuration of TGF β activation in articular cartilage** *Nat. Commun* **12**
34. Cabrera-Benítez N. E., et al. (2012) **Mechanical stress induces lung fibrosis by epithelial-mesenchymal transition** *Crit. Care Med* **40**:510–517
35. Wei S. C., Yang J. (2016) **Forcing through Tumor Metastasis: The Interplay between Tissue Rigidity and Epithelial-Mesenchymal Transition** *Trends Cell Biol* **26**:111–120
36. Luo N., et al. (2014) **Primary cilia signaling mediates intraocular pressure sensation** *Proc. Natl. Acad. Sci. U. S. A* **111**:12871–12876
37. Ryskamp D. A., et al. (2016) **TRPV4 regulates calcium homeostasis, cytoskeletal remodeling, conventional outflow and intraocular pressure in the mammalian eye** *Sci. Rep* **6**
38. Yang Y. F., Sun Y. Y., Peters D. M., Keller K. E. (2022) **The Effects of Mechanical Stretch on Integrins and Filopodial-Associated Proteins in Normal and Glaucomatous Trabecular Meshwork Cells** *Front. Cell Dev. Biol* **10**
39. Yarishkin O., et al. (2021) **Piezo1 channels mediate trabecular meshwork mechanotransduction and promote aqueous fluid outflow** *J. Physiol* **599**:571–592
40. White J. P. M., et al. (2016) **TRPV4: Molecular conductor of a diverse orchestra** *Physiol. Rev* **96**:911–973
41. Lakk M., Krizaj D. (2021) **TRPV4-Rho signaling drives cytoskeletal and focal adhesion remodeling in trabecular meshwork cells** *Am. J. Physiol. - Cell Physiol* **320**:C1013–C1030
42. Katari V., Bhavnani N., Paruchuri S., Thodeti C. (2021) **TRPV4 regulates matrix stiffness-dependent activation of YAP/VEGFR2 signaling via Rho/Rho kinase/LATS1/2 pathway** *FASEB J* **35**
43. Nilius B., Voets T. (2013) **The puzzle of TRPV4 channelopathies** *EMBO Rep* **14**:152–163
44. Thibodeau M. L., et al. (2017) **Compound heterozygous TRPV4 mutations in two siblings with a complex phenotype including severe intellectual disability and neuropathy** *Am. J. Med. Genet. Part A* **173**:3087–3092
45. Klein C. J., et al. (2011) **TRPV4 mutations and cytotoxic hypercalcemia in axonal Charcot-Marie-Tooth neuropathies** *Neurology* **76**:887–894
46. Daneva Z., et al. (2021) **Caveolar peroxynitrite formation impairs endothelial TRPV4 channels and elevates pulmonary arterial pressure in pulmonary hypertension** *Proc. Natl. Acad. Sci. U. S. A* **118**
47. Pochynyuk O., Zaika O., O'Neil R. G., Mamenko M. (2013) **Novel insights into TRPV4 function in the kidney** *Pflugers Arch. Eur. J. Physiol* **465**:177–186
48. Roberts M. W. G., et al. (2020) **TRPV4 receptor as a functional sensory molecule in bladder urothelium: Stretch-independent, tissue-specific actions and pathological implications** *FASEB J* **34**:263–286

49. Shibasaki K. (2020) **TRPV4 activation by thermal and mechanical stimuli in disease progression** *Lab. Investig* **100**:218–223
50. Toft-Bertelsen T. L., et al. (2022) **Lysophosphatidic acid as a CSF lipid in posthemorrhagic hydrocephalus that drives CSF accumulation via TRPV4-induced hyperactivation of NKCC1** *Fluids Barriers CNS* **19**
51. Lapajne L., et al. (2022) **TRPV4: Cell type-specific activation, regulation and function in the vertebrate eye** *Curr. Top. Membr* **89**:189–219
52. Lakk M., et al. (2021) **Membrane cholesterol regulates TRPV4 function, cytoskeletal expression, and the cellular response to tension** *J. Lipid Res* **62**
53. Uchida T., et al. (2021) **TRPV4 is activated by mechanical stimulation to induce prostaglandins release in trabecular meshwork, lowering intraocular pressure** *PLoS One* **16**
54. Jing L., Liu K., Wang F., Su Y. (2024) **Role of mechanically-sensitive cation channels Piezo1 and TRPV4 in trabecular meshwork cell mechanotransduction** *Hum. Cell* **37**:394–407
55. Redmon S. N., et al. (2024) **TRPV4 subserves physiological and pathological elevations in intraocular pressure** *Res. Sq. [Preprint]* <https://doi.org/10.21203/rs.3.rs-4714050/v1>
56. Nettesheim A., Shim M. S., Hirt J., Liton P. B. (2019) **Transcriptome analysis reveals autophagy as regulator of TGF β /Smad-induced fibrogenesis in trabecular meshwork cells** *Sci. Rep* **9**
57. Li H., Henty-Ridilla J. L., Bernstein A. M., Ganapathy P. S., Herberg S. (2022) **TGF β 2 Regulates Human Trabecular Meshwork Cell Contractility via ERK and ROCK Pathways with Distinct Signaling Crosstalk Dependent on the Culture Substrate** *Curr. Eye Res* **47**:1165–1178
58. Yan X., Xiong X., Chen Y. G. (2018) **Feedback regulation of TGF- β signaling** *Acta Biochim. Biophys. Sin. (Shanghai)* **50**:37–50
59. Carreon T., van der Merwe E., Fellman R. L., Johnstone M., Bhattacharya S. K. (2017) **Aqueous outflow - A continuum from trabecular meshwork to episcleral veins** *Prog. Retin. Eye Res* **57**:108–133
60. Yarishkin O., et al. (2018) **TREK-1 channels regulate pressure sensitivity and calcium signaling in trabecular meshwork cells** *J. Gen. Physiol* **150**:1660–1675
61. Honjo M., et al. (2001) **Effects of protein kinase inhibitor, HA1077, on intraocular pressure and outflow facility in rabbit eyes** *Arch. Ophthalmol* **119**:1171–1178
62. Rao P. V., Pattabiraman P. P., Kopczynski C. (2017) **Role of the Rho GTPase/Rho kinase signaling pathway in pathogenesis and treatment of glaucoma: Bench to bedside research** *Exp. Eye Res* **158**:23–32
63. Ethier C. R., Read A. T., Chan D. W. H. (2006) **Effects of latrunculin-B on outflow facility and trabecular meshwork structure in human eyes** *Investig. Ophthalmol. Vis. Sci* **47**:1991–1998
64. Liedtke W., Friedman J. M. (2003) **Abnormal osmotic regulation in trpv4 $^{-/-}$ mice** *Proc. Natl. Acad. Sci. U. S. A* **100**:13698–13703

65. Ryskamp D. A., et al. (2011) **The polymodal ion channel transient receptor potential vanilloid 4 modulates calcium flux, spiking rate, and apoptosis of mouse retinal ganglion cells** *J. Neurosci* **31**:7089–7101
66. Yarishkin O., Phuong T. T. T., Lakk M., Križaj D. (2018) **TRPV4 does not regulate the distal retinal light response** *Adv. Exp. Med. Biol* **1074**:553–560
67. Ikegami K., Masubuchi S. (2022) **Suppression of trabecular meshwork phagocytosis by norepinephrine is associated with nocturnal increase in intraocular pressure in mice.** *Commun Biol* **5**
68. Rahaman S. O., et al. (2014) **TRPV4 mediates myofibroblast differentiation and pulmonary fibrosis in mice** *J. Clin. Invest* **124**:5225–5238
69. Willard V. P., et al. (2021) **Transient receptor potential vanilloid 4 as a regulator of induced pluripotent stem cell chondrogenesis** *Stem Cells* **39**:1447–1456
70. Sharif N. A. (2023) **Identifying new drugs and targets to treat rapidly elevated intraocular pressure for angle closure and secondary glaucomas to curb visual impairment and prevent blindness** *Exp. Eye Res* **232**
71. Weinreb R. N., Toris C. B., Gabelt B. T., Lindsey J. D., Kaufman P. L. (2002) **Effects of prostaglandins on the aqueous humor outflow pathways** *Surv. Ophthalmol* **47**
72. Leske M. C., et al. (2003) **Factors for glaucoma progression and the effect of treatment: The early manifest glaucoma trial** *Arch. Ophthalmol* **121**:48–56
73. Fuchshofer R., Birke M., Welge-Lussen U., Kook D., Lütjen-Drecoll E. (2005) **Transforming growth factor- β 2 modulated extracellular matrix component expression in cultured human optic nerve head astrocytes** *Investig. Ophthalmol. Vis. Sci* **46**:568–578
74. Medina-Ortiz W. E., Belmares R., Neubauer S., Wordinger R. J., Clark A. F. (2013) **Cellular fibronectin expression in human trabecular meshwork and induction by transforming growth factor- β 2** *Investig. Ophthalmol. Vis. Sci* **54**:6779–6788
75. Callaghan B., et al. (2022) **Genome-wide transcriptome profiling of human trabecular meshwork cells treated with TGF- β 2** *Sci. Rep* **12**
76. Inoue-Mochita M., et al. (2015) **P38 MAP kinase inhibitor suppresses transforming growth factor- β 2- induced type 1 collagen production in trabecular meshwork cells** *PLoS One* **10**
77. Kang M. H., Oh D. J., heon Kang J., Rhee D. J. (2013) **Regulation of SPARC by transforming growth factor β 2 in human trabecular meshwork** *Investig. Ophthalmol. Vis. Sci* **54**:2523–2532
78. Patel G., et al. (2020) **Molecular taxonomy of human ocular outflow tissues defined by single-cell transcriptomics** *Proc. Natl. Acad. Sci. U. S. A* **117**:12856–12867
79. Reina-Torres E., et al. (2021) **The vital role for nitric oxide in intraocular pressure homeostasis** *Prog. Retin. Eye Res* **83**
80. van Zyl T., et al. (2020) **Cell atlas of aqueous humor outflow pathways in eyes of humans and four model species provides insight into glaucoma pathogenesis** *Proc. Natl. Acad. Sci. U. S. A* **117**:10339–10349

81. Zhu W., et al. (2021) **The role of Piezo1 in conventional aqueous humor outflow dynamics** *iScience* **24**
82. Pattabiraman P. P., et al. (2015) **Rhoa gtpase-induced ocular hypertension in a rodent model is associated with increased fibrogenic activity in the trabecular meshwork** *Am. J. Pathol* **185**:496–512
83. Zhang M., Maddala R., Rao P. V. (2008) **Novel molecular insights into RhoA GTPase-induced resistance to aqueous humor outflow through the trabecular meshwork** *Am. J. Physiol. - Cell Physiol* **295**
84. Güler A. D., et al. (2002) **Heat-evoked activation of the ion channel, TRPV4** *J. Neurosci* **22**:6408–6414
85. Nishimoto R., et al. (2021) **Thermosensitive TRPV4 channels mediate temperature-dependent microglia movement** *Proc. Natl. Acad. Sci. U. S. A* **118**
86. Abad E., et al. (2008) **Activation of store-operated Ca²⁺ channels in trabecular meshwork cells** *Investig. Ophthalmol. Vis. Sci* **49**:677–686
87. Feger M., et al. (2017) **The production of fibroblast growth factor 23 is controlled by TGF- α and β** *Sci. Rep* **7**
88. Sharma S., et al. (2017) **TRPV4 ion channel is a novel regulator of dermal myofibroblast differentiation** *Am. J. Physiol. - Cell Physiol* **312**:C562–C572
89. Adapala R. K., et al. (2013) **TRPV4 channels mediate cardiac fibroblast differentiation by integrating mechanical and soluble signals** *J. Mol. Cell. Cardiol* **54**:45–52
90. O'Connor C. J., Leddy H. A., Benefield H. C., Liedtke W. B., Guilak F. (2014) **TRPV4-mediated mechanotransduction regulates the metabolic response of chondrocytes to dynamic loading** *Proc. Natl. Acad. Sci. U. S. A* **111**:1316–1321
91. Songa Y., et al. (2014) **TRPV4 channel inhibits TGF- β 1-induced proliferation of hepatic stellate cells** *PLoS One* **9**
92. Wu Y., Qi J., Wu C., Rong W. (2021) **Emerging roles of the TRPV4 channel in bladder physiology and dysfunction** *J. Physiol* **599**:39–47
93. Jones J. L., et al. (2019) **TRPV4 increases cardiomyocyte calcium cycling and contractility yet contributes to damage in the aged heart following hypoosmotic stress** *Cardiovasc. Res* **115**:46–56
94. Chaigne S., Barbeau S., Ducret T., Guinamard R., Benoist D. (2023) **Pathophysiological Roles of the TRPV4 Channel in the Heart** *Cells* **12**
95. Wen X., et al. (2023) **Aortic smooth muscle TRPV4 channels regulate vasoconstriction in high salt- induced hypertension** *Hypertens. Res* **46**:2356–2367
96. Chen Y. L., et al. (2022) **Novel Smooth Muscle Ca²⁺-Signaling Nanodomains in Blood Pressure Regulation** *Circulation* **146**:548–564

97. Zhu Y., et al. (2023) **Vascular Smooth Muscle TRPV4 (Transient Receptor Potential Vanilloid Family Member 4) Channels Regulate Vasoconstriction and Blood Pressure in Obesity** *Hypertension* **80**:757–770
98. Zhao X., Ramsey K. E., Stephan D. A., Russell P. (2004) **Gene and protein expression changes in human trabecular meshwork cells treated with transforming growth factor- β** *Investig. Ophthalmol. Vis. Sci* **45**:4023–4034
99. Zhavoronkov A., et al. (2016) **Pro-fibrotic pathway activation in trabecular meshwork and lamina cribrosa is the main driving force of glaucoma** *Cell Cycle* **15**:1643–1652
100. Last J. A., et al. (2011) **Elastic modulus determination of normal and glaucomatous human trabecular meshwork** *Investig. Ophthalmol. Vis. Sci* **52**:2147–2152
101. Raychaudhuri U., Millar J. C., Clark A. F. (2017) **Tissue transglutaminase elevates intraocular pressure in mice** *Investig. Ophthalmol. Vis. Sci* **58**:6197–6211
102. Read A. T., Chan D. W. H., Ethier C. R. (2007) **Actin structure in the outflow tract of normal and glaucomatous eyes** *Exp. Eye Res* **84**:214–226
103. Wang K., Read A. T., Sulchek T., Ethier C. R. (2017) **Trabecular meshwork stiffness in glaucoma** *Exp. Eye Res* **158**:3–12
104. Fuchshofer R., Stephan D. A., Russell P., Tamm E. R. (2009) **Gene expression profiling of TGF β 2- and/or BMP7-treated trabecular meshwork cells: Identification of Smad7 as a critical inhibitor of TGF- β 2 signaling** *Exp. Eye Res* **88**:1020–1032
105. Gottanka J., Chan D., Eichhorn M., Lütjen-Drecoll E., Ethier C. R. (2004) **Effects of TGF- β 2 in Perfused Human Eyes** *Investig. Ophthalmol. Vis. Sci* **45**:153–158
106. Grove L. M., et al. (2019) **Translocation of TRPV4-PI3Ky complexes to the plasma membrane drives myofibroblast transdifferentiation** *Sci. Signal* **12**
107. Shukla A. K., et al. (2010) **Arresting a transient receptor potential (TRP) channel: β -arrestin 1 mediates ubiquitination and functional down-regulation of TRPV4** *J. Biol. Chem* **285**:30115–30125
108. Matsumura Y., et al. (2014) **The prophylactic effects of a traditional Japanese medicine, goshajinkigan, on paclitaxel-induced peripheral neuropathy and its mechanism of action** *Mol. Pain* **10**
109. Zhang Y., et al. (2008) **A transient receptor potential vanilloid 4 contributes to mechanical allodynia following chronic compression of dorsal root ganglion in rats** *Neurosci. Lett* **432**:222–227
110. Alessandri-Haber N., Dina O. A., Joseph E. K., Reichling D., Levine J. D. (2006) **A transient receptor potential vanilloid 4-dependent mechanism of hyperalgesia is engaged by concerted action of inflammatory mediators** *J. Neurosci* **26**:3864–3874
111. Maqboul A., Elsadek B. (2018) **Expression profiles of TRPV1, TRPV4, TLR4 and ERK1/2 in the dorsal root ganglionic neurons of a cancer-induced neuropathy rat model** *PeerJ* **2018**

112. Cui Y. Y., et al. (2020) **Expression and functional characterization of transient receptor potential vanilloid 4 in the dorsal root ganglion and spinal cord of diabetic rats with mechanical allodynia** *Brain Res. Bull* **162**:30–39
113. Borrás T., Buie L. K., Spiga M. G., Carabana J. (2015) **Prevention of nocturnal elevation of intraocular pressure by gene transfer of dominant-negative RhoA in rats** *JAMA Ophthalmol* **133**:182–190
114. Samuels B. C., et al. (2012) **Dorsomedial/perifornical hypothalamic stimulation increases intraocular pressure, intracranial pressure, and the translaminar pressure gradient** *Investig. Ophthalmol. Vis. Sci* **53**:7328–7335
115. Ikegami K. (2024) **Circadian rhythm of intraocular pressure** *J. Physiol. Sci* **74**
116. Shen K. Z., North R. A., Surprenant A. (1992) **Potassium channels opened by noradrenaline and other transmitters in excised membrane patches of guinea-pig submucosal neurones** *J. Physiol* **445**:581–599
117. Steinberg E. A., Wafford K. A., Brickley S. G., Franks N. P., Wisden W. (2015) **The role of K2P channels in anaesthesia and sleep** *Pflugers Arch. Eur. J. Physiol* **467**:907–916
118. Matthews B. D., et al. (2010) **Ultra-rapid activation of TRPV4 ion channels by mechanical forces applied to cell surface β 1 integrins** *Integr. Biol* **2**:435–442
119. Goswami C., Kuhn J., Heppenstall P. A., Hucho T. (2010) **Importance of non-selective cation channel TRPV4 interaction with cytoskeleton and their reciprocal regulations in cultured cells** *PLoS One* **5**
120. Jo A. O., et al. (2016) **Differential volume regulation and calcium signaling in two ciliary body cell types is subserved by TRPV4 channels** *Proc. Natl. Acad. Sci. U. S. A* **113**:3885–3890
121. Boussommier-Calleja A., et al. (2012) **Pharmacologic manipulation of conventional outflow facility in ex vivo mouse eyes** *Investig. Ophthalmol. Vis. Sci* **53**:5838–5845
122. Overby D. R., et al. (2014) **The structure of the trabecular meshwork, its connections to the ciliary muscle, and the effect of pilocarpine on outflow facility in mice** *Investig. Ophthalmol. Vis. Sci* **55**:3727–3736
123. Keller K. E., et al. (2018) **Consensus recommendations for trabecular meshwork cell isolation, characterization and culture** *Exp. Eye Res* **171**:164–173
124. Bagué T., et al. (2022) **Effects of Netarsudil-Family Rho Kinase Inhibitors on Human Trabecular Meshwork Cell Contractility and Actin Remodeling Using a Bioengineered ECM Hydrogel** *Front. Ophthalmol* **2**
125. Stamer W. D., Seftor R. E. B., Williams S. K., Samaha H. A. M., Snyder R. W. (1995) **Isolation and culture of human trabecular meshwork cells by extracellular matrix digestion** *Curr. Eye Res* **14**:611–617
126. Phuong T. T. T., et al. (2017) **Calcium influx through TRPV4 channels modulates the adherens contacts between retinal microvascular endothelial cells** *J. Physiol* **595**:6869–6885
127. Schindelin J., et al. (2012) **Fiji: An open-source platform for biological-image analysis** *Nat. Methods* **9**:676–682

Author information

Christopher N Rudzitis

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States, Department of Neurobiology, University of Utah, Salt Lake City, United States

Monika Lakk

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States

Ayushi Singh

Department of Ophthalmology and Visual Sciences, SUNY Upstate Medical University, Syracuse, United States, Department of Cell and Developmental Biology, SUNY Upstate Medical University, Syracuse, United States

Sarah N Redmon

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States

Denisa Kirdajova

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States

Yun-Ting Tseng

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States

Michael L De Ieso

Department of Ophthalmology, Duke Eye Center, Duke University, Durham, United States

W Daniel Stamer

Department of Ophthalmology, Duke Eye Center, Duke University, Durham, United States

Samuel Herberg

Department of Ophthalmology and Visual Sciences, SUNY Upstate Medical University, Syracuse, United States, Department of Cell and Developmental Biology, SUNY Upstate Medical University, Syracuse, United States, Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, United States

ORCID iD: [0000-0001-7196-2010](https://orcid.org/0000-0001-7196-2010)

David Križaj

Department of Ophthalmology and Visual Sciences, University of Utah, Salt Lake City, United States, Department of Neurobiology, University of Utah, Salt Lake City, United States, Department of Bioengineering, University of Utah, Salt Lake City, United States

ORCID iD: [0000-0003-4468-3029](https://orcid.org/0000-0003-4468-3029)

For correspondence: david.krizaj@hsc.utah.edu

Editors

Reviewing Editor

Kate Poole

UNSW Sydney, Sydney, Australia

Senior Editor

Merritt Maduke

Stanford University, Stanford, United States of America

Reviewer #1 (Public review):

Summary:

This comprehensive study employed molecular, optical, electrophysiological and tonometric strategies to establish the role of TGF β 2 in transcription and functional expression of mechanosensitive channel isoforms alongside studies of TM contractility in biomimetic hydrogels, and intraocular pressure regulation in a mouse model of TGF β 2 -induced ocular hypertension. TGF β 2 upregulated expression of TRPV4 and PIEZO1 transcripts and time-dependently augmented functional TRPV4 activation. TRPV4 activation induced TM contractility whereas pharmacological inhibition suppressed TGF β 2-induced hypercontractility and abrogated ocular hypertension in eyes overexpressing TGF β 2. Trpv4 $^{-/-}$ mice resisted TGF β 2-driven increases in IOP. These data establish a fundamental role of TGF β as a modulator of mechanosensing and identifies TRPV4 channel as a common mechanism for TM contractility and pathological ocular hypertension.

Strengths:

The manuscript is very well written and details the important function of TRPV4 in TM cell function. These data provide novel therapeutic targets and potential for disease-altering therapeutics.

Weaknesses:

The experimental rigor and design of the nocturnal IOP experiments was weak with low n values and differing methods of IOP measurement (conscious versus anesthetized). The same method of IOP measurement needs to be used for all measurements to make any conclusions on the circadian patterns of IOP in each condition.

<https://doi.org/10.7554/eLife.104894.1.sa2>

Reviewer #2 (Public review):

The manuscript by Christopher N. Rudzitis et al. describes the role of TGF β 2 in the transcription and functional expression of mechanosensitive channel isoforms, alongside studies on TM contractility in biomimetic hydrogels and intraocular pressure. Overall, it is a very interesting study, nicely designed, and will contribute to the available literature on TRPV4 sensitivity to mechanical forces.

I have the following comment for the authors to address.

Figure 1A-C.

Often there is a difference between the massage and transcript data. I recommend the authors to confirm with qPCR data with another mode of protein measurements.

Does direct TRPV4 activation also induce the expression of these markers? Does inhibition of

TRPV4, after TGF- β treatment, prevent the expression of these markers? Is TRPV4 acting downstream of this response?

Figure 1D. Beta tubulin is not a membrane marker. Having staining of β tubulin in membrane fraction shows contamination from the cytoplasm.

Does the overall expression also increase?

Figure 4A: it is not very clear. I recommend including a zoom image or better resolution image.

Figure 5B and 6B.

Why there is a difference between groups in pre-injection panel. As Figure 5A, in pre-injection, there is no difference between LV-TGF β and LV-control while in 5B there is a significant difference between these groups.

Discussion section.

Line 279, . "TRPV4 channels in cells treated with TGF β 2 are likely to be constitutively active" ... needs to be discussed further.

Line 280: "The residual contractility in HC-06-treated cells may reflect TGF β 2-mediated contributions from Piezo1."

Piezo1 has a low threshold for mechanosensitivity. How do the authors discuss the observation that, in the presence of Piezo1, TRPV4 has a more prominent mechanosensory function? Is this tied to TGF β signalling?

<https://doi.org/10.7554/eLife.104894.1.sa1>

Author response:

We thank the editors and reviewers for the constructive assessment. We plan to address the comments as follows:

Reviewer #1 (Public review):

We are generating a new cohort of Lv-TGFB2 overexpressing mice in which IOP will be compared under the anesthesia conditions that are identical for diurnal and nocturnal states. Parenthetically, we used the awake (diurnal) and isoflurane (nocturnal) anesthesia to mirror the conditions in the Patel et al (2021) PNAS study.

Reviewer #2 (Public review):

We are not sure what the Reviewer means by the “difference between the message and transcript data” and are not sure whether providing evidence about the TRPV4-dependence of the expression of fibrotic genes and canonical TGF β 2 pathway genes fits within the scope of our study (which focuses on the TGF β 2-dependence of TRPV4 expression and IOP regulation). We propose to address this by including new data about the TGF β 2- and TRPV4 dependence of TRPV4 and Piezo1 expression. We could include information about the effect of TGF β 2 on fibrosis-related genes from a (submitted study) in which we used RNASeq to investigate TGF β 2 and TGF β 2 + HC067047-dependence of gene expression in TM cells on a confidential basis but not include it in the revised manuscript.

- Re: β -tubulin comment [β -tubulin associates with the plasma membrane by binding to integral membrane proteins in the plasma and organellar membranes, through palmitoylation and attachment to linker proteins and as an integral component of exocytotic vesicles (Wolff, BBA 2009; Hogerheide et al., PNAS 2017). Together with β -actin and Gapdh it is often used as a loading control to assess cellular TRPV4 protein expression (e.g., <https://www.cellsignal.com/products/primary-antibodies/trpv4-antibody/65893>; Grove et al., Science

Signaling 2019 and Moore et al., PNAS 2013). Our qPCR and RNASeq studies show that TGFB2 does not affect b-tubulin expression]

- We will provide a higher resolution image for Fig. 4A

- Will address the Fig 5A and 6A comment [We thank the Reviewer for noticing the ambiguity and revised Figure Legends to clarify that “pre-injection” in Figures 5B and 6B refers to IOP measurements before the intracameral injection of HC-06 not pre-injection of lentiviral constructs].

- We will address the issue of constitutive TRPV4 activity and Piezo1 involvement in the revised Discussion.

We hope this is sufficient information at this point but would be more than happy to provide more information if needed.

Thank you, we are very impressed by the eLife review protocols.

<https://doi.org/10.7554/eLife.104894.1.sa0>