

Parallel HIV-1 fitness landscapes shape viral dynamics in humans and macaques that develop broadly neutralizing antibodies

Reviewed Preprint

v1 • March 11, 2025

Not revised

Kai S Shimagaki, Rebecca M Lynch, John P Barton 

Department of Computational and Systems Biology, University of Pittsburgh School of Medicine, Pittsburgh, United States • Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, United States • Department of Microbiology, Immunology and Tropical Medicine, School of Medicine and Health Sciences, George Washington University, Washington, United States

 https://en.wikipedia.org/wiki/Open_access Copyright information

eLife Assessment

In this **important** quantitative study of HIV-1 evolution in humans and rhesus macaques, selection coefficients are inferred at scale over the HIV genome. Selection coefficients are similar in humans and macaques, providing **convincing** evidence that these coefficients are representative of the fitness landscapes of these viruses within hosts. This work should be of interest to the community working on quantitative evolution and fitness landscape inference, and the finding that rapid fitness gains in the HIV population predict bNAbs emergence has implications for HIV vaccine design.

<https://doi.org/10.7554/eLife.105466.1.sa3>

Abstract

Human immunodeficiency virus (HIV)-1 exhibits remarkable genetic diversity. An effective HIV-1 vaccine must therefore elicit antibodies that can neutralize many variants of the virus. While broadly neutralizing antibodies (bnAbs) have been isolated from HIV-1 infected individuals, a general understanding of the virus-antibody coevolutionary processes that lead to their development remains incomplete. We performed a quantitative study of HIV-1 evolution in humans and rhesus macaques, including individuals who developed bnAbs. We observed strong selection early in infection for mutations affecting HIV-1 envelope glycosylation and escape from autologous strain-specific antibodies, followed by weaker selection for bnAb resistance. The inferred fitness effects of HIV-1 mutations in humans and macaques were remarkably similar. Moreover, we observed a striking pattern of rapid HIV-1 fitness gains that precedes the development of bnAbs. Our work highlights strong parallels between infection in rhesus macaques and humans, and it reveals a quantitative evolutionary signature of bnAb development.

Introduction

Human immunodeficiency virus (HIV-1) rapidly mutates and proliferates in infected individuals. The immune system is a major driver of HIV-1 evolution, as the virus accumulates mutations to escape from host T cells and antibodies ^{1–3}. Due to the chronic nature of HIV-1 infection, coupled with high rates of mutation and replication, HIV-1 genetic diversity within and between infected individuals is incredibly high. Genetic diversity challenges vaccine development, as vaccine-elicited antibodies must be able to neutralize many strains of the virus to protect against infection ⁴.

However, there exist rare antibodies that are capable of neutralizing a broad range of HIV-1 viruses. These broadly neutralizing antibodies (bnAbs) have therefore been the subject of intense research ^{5–8}. Eliciting bnAbs through vaccination remains a major goal of HIV-1 vaccine design. However, the development of exceptionally broad antibody responses is rare, and such antibodies typically develop only after several years of infection ^{9–12}.

Recent years have yielded important insights into the coevolutionary process between HIV-1 and antibodies that sometimes leads to the development of bnAbs. Clinical studies have collected serial samples of HIV-1 sequences from a few individuals who developed bnAbs and characterized the resulting antibodies, their developmental stages, and binding sites ^{13–15}. The contributions of HIV-1 and its coevolution to bnAb development are complex ^{16,17}. High viral loads and viral diversity have been positively associated with bnAb development ^{16–18}. However, superinfection, which can vastly increase HIV-1 diversity, is not always associated with bnAb development ¹⁹, and it does not appear to broaden antibody responses in the absence of other factors ¹⁷.

Here, we sought to characterize the evolutionary dynamics of HIV-1 that accompany the development of bnAbs in clinical data. In particular, we inferred the landscape of selective pressures that shape the evolution of HIV-1 within hosts, reflecting the effects of the immune environment. We first analyzed data from two individuals who developed bnAbs within a few years after HIV-1 infection ^{13,14}. In both individuals, HIV-1 mutations inferred to be the most beneficial were observed early in infection. In general, mutations that provided resistance to autologous strain-specific antibodies were inferred to be more strongly selected than ones that escaped from bnAbs. We also observed clusters of beneficial mutations along the HIV-1 genome, which were associated with envelope protein (Env) structure.

To confirm the generality of these patterns in a broader sample, we studied recent data from rhesus macaques (RMs) infected with simian-human immunodeficiency viruses (SHIV) that incorporated HIV-1 Env proteins derived from the two individuals above ²⁰. This study also compared patterns of Env evolution in HIV-1 and SHIV in response to host immunity. We observed striking parallels between the inferred fitness effects of Env mutations in RMs and humans, suggesting highly similar selective pressures on the virus despite different host species and differences in individual immune responses. Furthermore, we found that RMs that developed broad, potent antibody responses could clearly be distinguished from those with narrowly focused responses using the evolutionary dynamics of the virus. Specifically, the virus population in individuals who developed greater breadth was distinguished by larger and more rapid gains in fitness than in other individuals. Collectively, these results show high similarity between SHIV evolutionary dynamics in RMs and HIV-1 in humans, and that viral fitness gain is associated with antibody breadth.

Results

Quantifying HIV-1 evolutionary dynamics

We studied HIV-1 evolution accompanying the development of bnAbs in two donors, CH505 and CH848, enrolled in the Center for HIV/AIDS Vaccine Immunology 001 acute infection cohort ²¹. CH505 developed the CD4 binding site-targeting bnAb CH103, which was first detectable 14 weeks after HIV-1 infection ¹³. CH103 maturation was found to be associated with viral escape from another antibody lineage, CH235, that ultimately developed significant breadth ^{22,23}. CH848 developed a bnAb, DH270, targeting a glycosylated site near the third variable loop (V3) of Env ¹⁴. Similar to the bnAb development process in CH505, escape from “cooperating” DH272 and DH475 lineage antibodies was observed to contribute to the maturation of DH270 (ref. ¹⁴).

To quantify HIV-1 evolutionary dynamics, we sought to infer a fitness model that best explained the changes in the genetic composition of the viral population observed in each individual over time. We assume that the effect on viral fitness of each individual mutation a at each site i is quantified by a selection coefficient $s_i(a)$, with positive coefficients $s_i(a) > 0$ denoting mutations that are beneficial for the virus and $s_i(a) < 0$ denoting deleterious ones. We further assume that the cumulative fitness effects of mutations are additive, such that the overall fitness F of a viral sequence is given by the sum of the selection coefficients for all the mutations that it bears. Following this model, we can then quantify the probability of any evolutionary history of the viral population (i.e., the distribution of viral genotypes over time) as a function of the selection coefficients (Methods). We then used MPL ^{24,25} to infer the selection coefficients that best fit the viral dynamics observed in data from CH505 and CH848 (Methods). This approach has been successfully applied to study the evolution of HIV-1 (ref. ²⁴), as well as SARS-CoV-2 (ref. ²⁶) and experimental evolution in bacteria ^{27,28}.

Broad patterns of HIV-1 selection

While the great majority of HIV-1 mutations were inferred to be neutral ($s_i(a) \sim 0$), a few mutations substantially increase viral fitness (**Fig. 1**). Strongly beneficial mutations occurred in clusters along the genome and preferentially appeared in specific regions of Env (**Fig. 2**).

To quantify patterns of selection in Env, we examined the top 2% of mutations inferred to be the most beneficial in CH505 and CH848. The fractions of nonsynonymous mutations within these subsets were 97% and 92% for CH505 and CH848, respectively. These fractions are significantly higher than chance expectations ($p = 6.7 \times 10^{-3}$ and 5.6×10^{-4} , Fisher’s exact test; Methods), supporting the model’s ability to accurately infer fitness effects in this data. For CH505, we found 10.9-fold more strongly beneficial mutations in the first variable loop (V1) than expected by chance ($p = 2.5 \times 10^{-3}$). This is consistent with the presence of V1 mutations conferring resistance to autologous strain-specific antibodies ²². Mutations in V4, a region targeted by CD8⁺ T cells ²², were also 10.0-fold enriched in this subset ($p = 9.0 \times 10^{-5}$). For CH848, mutations in V1, V3, and V5 were enriched by factors of 14.2, 6.3, and 19.1 among the top 2% most beneficial mutations ($p = 5.9 \times 10^{-10}$, 6.6×10^{-3} , and 4.8×10^{-6}). Mutations in these regions were shown to play a role in resistance to DH270 and DH475 lineage antibodies ¹⁴.

Reversions and mutations affecting N-linked glycosylation motifs were also likely to be beneficial. We define reversions as mutations where the transmitted/founder (TF) amino acid changes to match the subtype consensus sequence at the same site. Among the top 2% most beneficial mutations, reversions were enriched by factors of 19.9 and 17.8 for CH505 and CH848 viruses, respectively ($p = 2.1 \times 10^{-8}$ and 8.5×10^{-13}), consistent with past work finding strong selection for reversions ^{24,30}. For CH848, this group also includes several strongly selected mutations observed in Nef (**Fig. 1B**), a protein that plays multiple roles during HIV-1 infection ^{31,32}.

Fig. 1.

Beneficial mutations occur in clusters along the genome.

Inferred fitness effects of HIV-1 mutations in CH505 (A) and CH848 (B). The position along the radius of each circle specifies the strength of selection: mutations plotted closer to the center are more deleterious, while those closer to the edge are more beneficial. For both individuals, clusters of beneficial mutations are observed in the variable loops of Env, some of which are associated with antibody escape. For CH848, a group of strongly beneficial mutations also appears in Nef.





Fig. 2.

Visualization of the effects of HIV-1 mutations in CH505 on the Env trimer.

A, Side view of the Env trimer ²⁹, with detail views of selection for mutations in the CH103 binding site (B), CH235 binding site (C), and sites associated with escape from autologous strain-specific antibodies (D). E, Top view of the trimer, with detail views of variable loops (F-I) and the CD4 binding site (J). Generally, beneficial mutations appear more frequently near exposed regions at the top of the Env trimer, and deleterious mutations appear in more protected regions. Mutations near the V1/V2 apex can affect the binding and neutralization of antibodies targeting the CD4 binding site ²².



Mutations affecting N-linked glycosylation motifs (i.e., by adding, removing, or shifting a glycosylation motif) were enriched by factors of 4.6 ($p = 7.0 \times 10^{-3}$) and 8.7 ($p = 1.4 \times 10^{-10}$). Changes in glycosylation patterns contributed to antibody escape for both CH505 and CH848 (refs. [14](#), [22](#), [23](#)).

Selection for antibody escape

To quantify levels of selection for antibody escape, we computed selection coefficients for mutations that were observed to contribute to resistance to bnAbs as well as bnAb precursors and autologous strain-specific antibodies (**Supplementary Tables 1** [1](#)–[3](#)) [22](#), [23](#), [33](#)–[36](#)). We mapped the inferred selection coefficients to the Env protein structure [13](#), highlighting the binding sites for bnAbs and resistance mutations for strain-specific antibodies, as well as important parts of Env (**Fig. 2** [2](#), **Supplementary Fig. 1** [1](#)). We also analyzed when these resistance mutations were first observed in each individual.

Overall, we observed stronger selection for escape from autologous strain-specific antibodies and/or changes in glycosylation during the first six months of infection (**Supplementary Tables 4** [4](#)–[5](#)). This was then followed by more modest selection for escape from bnAb lineages (**Fig. 3** [3](#)). For CH505, mutations that conferred resistance to the intermediate-breadth CH235 lineage [22](#) were less beneficial than top mutations escaping from strain-specific antibodies (**Fig. 3A** [3](#)). In turn, resistance mutations for the broader CH103 lineage were less beneficial than CH235 resistance mutations (**Fig. 3A** [3](#)). CH848 is similar, with some highly beneficial mutations affecting glycosylation observed early in infection (**Fig. 3B** [3](#)). While a few mutations affecting DH270 appear strongly selected, these mutations appeared long before the DH270 lineage was detected (around 3.5 years after infection [14](#)). Thus, these mutations may have initially been selected for other reasons.

Consistent patterns of selection in SHIV evolution

Simian-human immunodeficiency viruses (SHIVs) have numerous applications in HIV/AIDS research [37](#), [38](#). Recently, Roark and collaborators studied SHIV-antibody coevolution in rhesus macaques (RMs), which they compared with patterns of HIV-1 evolution [20](#). Two of the SHIV constructs in this study included envelope sequences derived from CH505 and CH848 transmitted/founder (TF) viruses. There, it was found that 2 out of 10 RMs inoculated with SHIV.CH505 and 2 out of 6 RMs inoculated with SHIV.CH848 developed antibodies with substantial breadth.

To understand whether the patterns of HIV-1 selection observed in CH505 and CH848 are repeatable, and to search for viral factors that distinguish between individuals who develop bnAbs and those who do not, we analyzed SHIV.CH505 and SHIV.CH848 evolution in RMs [20](#). To prevent spurious inferences, we first omitted data from RMs with <3 sampling times or <4 sequences in total (Methods). After processing, we examined evolutionary data from 7 RMs inoculated with SHIV.CH505 and 6 RMs inoculated with SHIV.CH848 (**Supplementary Tables 6** [6](#)–[7](#)). We then computed selection coefficients for SHIV mutations within each RM. Reasoning that selective pressures across SHIV.CH505 and SHIV.CH848 viruses are likely to be similar, we also inferred two sets of joint selection coefficients that best describe SHIV evolution in SHIV.CH505- and SHIV.CH848-inoculated RMs, respectively (Methods).

As before, we examined the top 2% of SHIV.CH505 and SHIV.CH848 mutations that we inferred to be the most beneficial for the virus. Overall, we found consistent selection for reversions (17.9- and 14.2-fold enrichment, $p = 4.3 \times 10^{-11}$ and 1.2×10^{-11} for SHIV.CH505 and SHIV.CH848, respectively), with slightly attenuated enrichment in mutations that affect N-linked glycosylation (2.7- and 4.2-fold enrichment, $p = 5.4 \times 10^{-2}$ and 3.4×10^{-6}). However, there is a small subset of mutations that *shift* glycosylation sites by simultaneously disrupting one N-linked glycosylation




Fig. 3.

Trajectories and inferred selection coefficients for immune escape mutations and mutations affecting Env glycosylation.

A, For CH505, early, strongly selected mutations include ones that escape from CD8⁺ T cells and autologous strain-specific antibodies (ssAbs). More moderately selected bnAb resistance mutations tend to arise later. Note that mutations that affect bnAb resistance can appear in the viral population before bnAbs are generated. Open circles and error bars reflect the mean and standard deviation of inferred selection coefficients in each category. **B,** For CH848, mutations affecting glycosylation dominate the early phase of evolution, followed later by mutations affecting bnAb resistance. See [Supplementary Fig. 2](#) for a detailed view of mutation frequency trajectories by mutation type.

motif and completing another, where highly beneficial mutations occur far more often than expected by chance (158.3- and 191.1-fold enrichment, $p = 1.7 \times 10^{-4}$ and 7.9×10^{-11} for CH505 and SHIV.CH505, respectively; 118.7-fold and 90.4-fold enrichment, $p = 2.0 \times 10^{-4}$ and 2.3×10^{-7} for CH848 and SHIV.CH848).

Intuitively, one may expect that strongly beneficial SHIV mutations are more likely to be observed in samples from multiple RMs. However, we found that the number of RMs in which a mutation is observed is only weakly associated with the fitness effect of the mutation (**Supplementary Fig. 3**). While substantially deleterious SHIV mutations are rarely observed across multiple RMs, neutral and nearly neutral mutations are common. Thus, in this data set, it is not generally true that SHIV mutations observed in multiple hosts must significantly increase viral fitness.

We observed some differences between HIV-1 and SHIV in the precise locations of the most beneficial Env mutations. For example, mutations in V4 are highly enriched in CH505 due to a CD8⁺ T cell epitope in this region, but not in SHIV.CH505 (2.7-fold enrichment, $p = 0.13$). For SHIV.CH848, beneficial mutations are modestly enriched in V1 and V5 (4.7- and 4.2-fold, $p = 4.0 \times 10^{-3}$ and 6.7×10^{-2}), as in CH848, but not for V3 (0.9-fold enrichment, $p = 0.22$).

Despite some differences in the top mutations, patterns of selection over time in SHIV were very similar to those found for HIV-1. As before, highly beneficial mutations, including ones affecting glycosylation, tended to appear earlier in infection. This was followed by modestly beneficial mutations at later times, including ones involved in resistance to bnAbs in the RMs who developed antibodies with significant breadth (examples in **Fig. 4**).

Detection of SHIV mutations that increase viral load

A major goal of nonhuman primate studies with SHIV is to faithfully recover important aspects of HIV-1 infection in humans. However, due to the divergence of simian immunodeficiency viruses and HIV-1, SHIVs are not always well-adapted to replication in RMs³⁶. To combat this problem, a recent study identified six SHIV.CH505 mutations that increase viral load (VL) in RMs³⁶. These mutations result in viral kinetics that better mimic HIV-1 infection in humans.

Our analysis readily identifies the SHIV.CH505 mutations shown to increase VL. Five out of the top six SHIV.CH505 mutations with the largest average selection coefficients are associated with increased VL (**Supplementary Table 8**). The final mutation identified by Bauer et al., N130D, is ranked tenth. We also find highly beneficial mutations in SHIV.CH848 that are distinct from those in SHIV.CH505 (**Supplementary Table 9**). Highly-ranked mutations identified here may be good experimental targets for future studies aimed at increasing SHIV.CH848 replication *in vivo*.

Fitness agreement between HIV-1 and SHIV

Next, we explored the similarity in the overall viral fitness landscapes inferred for HIV-1 and SHIV, beyond just the top mutations. First, we computed the fitness of each SHIV sequence using the joint SHIV.CH505 and SHIV.CH848 selection coefficients inferred from RM data. Then, we computed fitness values for SHIV sequences using selection coefficients inferred from HIV-1 evolution in CH505 and CH848.

We observed a remarkable agreement between SHIV fitness values computed from these two sources (**Fig. 5**). For both SHIV.CH505 and SHIV.CH848, the correlation between viral fitness estimated using data from humans (i.e., CH505 and CH848) and RMs is strongly and linearly correlated (Pearson's $r = 0.96$ and 0.95 , $p < 10^{-20}$). This implies that evolutionary pressures on the envelope protein SHIV-infected RMs are highly similar to those on HIV-1 in humans with the same




Fig. 4.

Example trajectories and selection coefficients for SHIV mutations that affect viral load, bnAb recognition, or glycosylation.

A, In RM5695, infected with SHIV.CH505, mutations known to increase viral load ³⁶ and ones affecting gly-cosylation were rapidly selected. Mutations affecting resistance to broad antibodies ²⁰ arose later under moderate selection. Open circles and error bars reflect the mean and standard deviation of inferred selection coefficients in each category. **B**, Slower but qualitatively similar evolutionary patterns were observed in RM6163, infected with SHIV.CH848.

TF Env. In fact, this relationship holds even beyond the SHIV sequences observed during infection. Fitness estimates for sequences with randomly shuffled sets of mutations are also strongly correlated (**Supplementary Fig. 4** [↗](#)).

Evolutionary dynamics forecast antibody breadth

Given the similarity of HIV-1 and SHIV evolution, we sought to identify evolutionary features that distinguish between hosts who develop broad antibody responses and those who do not. **Figure 5** [↗](#) shows that SHIV sequences from hosts with bnAbs often reach higher fitness values than those in hosts with only narrow-spectrum antibodies. We hypothesized that stronger selective pressures on the virus might drive viral diversification, stimulating the development of antibody breadth. Past studies have associated higher viral loads with bnAb development and observed viral diversification around the time of bnAb emergence [13](#) [↗](#), [16](#) [↗](#), [17](#) [↗](#). Computational studies and experiments have also shown that sequential exposure to diverse antigens can induce cross-reactive antibodies [39](#) [↗](#)–[41](#) [↗](#).

To further quantify SHIV evolutionary dynamics, we computed the average fitness gain of viral populations in each RM over time. We observed a striking difference in SHIV fitness gains between RMs that developed broad antibody responses and those that did not (**Fig. 6** [↗](#)). In particular, SHIV fitness increased rapidly before the development of antibody breadth. SHIV fitness gains in RM5695, which developed exceptionally broad and potent antibodies [20](#) [↗](#), were especially rapid and dramatic. These fitness differences were not attributable to bnAb resistance mutations, which were only moderately selected and generally appeared after bnAbs developed.

One outlier in this pattern is RM6072, infected with SHIV.CH505. Antibody development in RM6072 followed a path that was remarkably similar to CH505, including a lineage of antibodies, DH650, directed toward the CD4 binding site of Env [20](#) [↗](#). However, resistance to the DH650 lineage is conferred by a strongly selected mutation that adds a glycan at site 234 (T234N, with an inferred selection coefficient of 4.5%). Broadly neutralizing antibodies similar to DH650 are able to accommodate this glycan due to shorter and/or more flexible light chains [20](#) [↗](#), [42](#) [↗](#), but DH650 cannot. Antibody evolution in RM6072 thus proceeded along a clear pathway toward bnAb development, but lacked critical mutations to achieve breadth.

Next, we quantified how different types of SHIV mutations contributed to viral fitness gains over time. We examined contributions from VL-enhancing mutations [36](#) [↗](#), antibody escape mutations [20](#) [↗](#), [36](#) [↗](#), other mutations affecting Env glyco-sylation, and reversions to subtype consensus. We found increased fitness gains across all types of mutations in RMs that developed broad antibody responses, compared to those that did not (**Supplementary Fig. 5** [↗](#)). VL-enhancing mutations, known antibody resistance mutations, and reversions typically made the largest contributions to viral fitness.

Discussion

The induction of bnAbs is a major goal of HIV-1 vaccine design [8](#) [↗](#). Both computational [39](#) [↗](#), [41](#) [↗](#), [43](#) [↗](#), [44](#) [↗](#) and experimental [40](#) [↗](#), [45](#) [↗](#), [46](#) [↗](#) studies, as well as observations from individuals who developed bnAbs [13](#) [↗](#), [14](#) [↗](#), [22](#) [↗](#), suggest that the co-evolution of antibodies and HIV-1 is important to stimulate broad antibody responses. However, a general understanding of the coevolutionary pathways by which bnAbs develop remains incomplete. Here, we quantitatively studied the evolution of HIV-1 and SHIV (featuring HIV-1-derived Env sequences) that accompanies the development of bnAbs.

Fig. 5.

Inferred fitness landscapes in HIV-1 and SHIV are highly similar.

A, Fitness of SHIV.CH505 sequences relative to the TF sequence across 7 RMs, including 2 that developed bnAbs, 1 that developed tier 2 nAbs that lacked a critical mutation for breadth, and 4 that did not develop broad antibody responses, evaluated using fitness effects of mutations using data from CH505 and using RM data. The fitness values are strongly correlated, indicating the similarity of Env fitness landscapes inferred using HIV-1 or SHIV data. Values are normalized such that the fitness gain of the TF sequence is zero. **B**, Fitness values for SHIV.CH848 sequences also show strong agreement between CH848 and SHIV.CH848 landscapes.





Fig. 6.

Rapid SHIV fitness gains precede the development of broadly neutralizing antibodies.

For both SHIV.CH505 (**A**) and SHIV.CH848 (**B**), viral fitness gains over time display distinct patterns in RM hosts that developed bnAbs versus those that did not. Notably, the differences in SHIV fitness gains between hosts with and without broad antibody responses appear *before* the development of antibody breadth, and cannot be attributed to selection for bnAb resistance mutations. RM6072 is an unusual case, exhibiting antibody development that was highly similar to CH505. Although RM6072 developed tier 2 nAbs, they lacked key mutations critical for breadth [20](#). Points and error bars show the mean and standard deviation of fitness gains across SHIV samples in each RM at each time.



Our study highlighted how different classes of mutations (e.g., mutations affecting T cell escape or Env glycosylation) affect fitness *in vivo*. In both HIV-1 and SHIV, we found strong selection for reversions to subtype consensus and some mutations that affected N-linked glycosylation motifs or resistance to autologous strain-specific antibodies. Few CD8⁺ T cell epitopes were identified in this data set, but the T cell escape mutations that we did observe were highly beneficial for the virus. Consistent with past work studying VRC26 escape in CAP256, we observed more modest selection for bnAb resistance mutations ²⁴.

Overall, we found striking similarities between Env evolution in humans and RMs. Importantly, these parallels extend beyond the observation of repetitive mutations, as the number of hosts in which a mutation was observed was only weakly associated with the mutation's fitness effect (**Supplementary Fig. 3**). Our inferred Env fitness values in humans and RMs were highly correlated, even for artificial sequences with randomly permuted mutations. This indicates that the evolutionary pressures on Env in HIV-1 and SHIV infection are very similar. Our findings reinforce SHIV as a model system that closely mirrors HIV-1 infection.

We discovered that the speed of SHIV fitness gains was clearly higher in RMs that developed broad antibody responses than in those with narrow-spectrum antibodies. Fitness gains in the viral population preceded the development of bnAbs, and they were not driven by bnAb resistance mutations. This suggests that rapid changes in the viral population are a cause rather than a consequence of antibody breadth. While our sample is limited to 13 RMs and two founder Env sequences, we find a clear separation between RMs that did or did not develop antibody breadth. Thus, the dynamics of viral fitness may serve as a quantitative signal associated with bnAb development.

As noted above, Roark and collaborators also performed a detailed comparison of HIV-1 and SHIV evolution with the same TF Env sequences ²⁰. One of their main conclusions was that most Env mutations were selected for escape from CD8⁺ T cells or antibodies. We found that many antibody resistance mutations identified by Roark et al. are also positively selected in our analysis. Mutations at sites 166 and 169 were shown to confer resistance to a V2 apex bnAb, RHA1, isolated in RM5695 (ref. ²⁰). We inferred moderately positive selection coefficients of 0.49% and 0.43% for R166K and R169K, respectively. The same mutations were found in RM6070, which also developed V2 apex bnAbs, with a selective advantage of 1.7% (**Supplementary Table 10**). Mutations conferring resistance to autologous strain-specific nAbs were identified at multiple sites by Roark and colleagues: 130, 234, 279, 281, 302, 330, and 334 in RM6072, which developed antibody responses targeting the CD4 binding site (DH650) and V3 (DH647 and DH648) regions. Mutations Y330H and N334S, which confer resistance to V3 autologous nAbs, were detected in all RMs infected with SHIV.CH505, with selective advantages of 3.0% and 4.6% in RM6072, and 1.7% and 3.2% on average across RMs, respectively. Overall, we found that mutations conferring resistance to autologous strain-specific antibodies were common and more strongly selected than bnAb resistance mutations (**Supplementary Tables 10-11**).

While we observed strong selection for strain-specific antibody resistance mutations, these results could also be affected by the effects of these mutations on viral replication independent of immune escape. In particular, many ssAb resistance mutations are also reversions to the subtype consensus sequence, which have often been observed to improve viral fitness ^{24,30}. For example, N334S, K302N, and T234N are all reversions. These are among the most beneficial mutations inferred for SHIV.CH505 (**Supplementary Table 8**). In future work, it would be interesting to attempt to fully separate the fitness effects of mutations due to antibody escape and intrinsic replication (cf. ⁴⁷).

Among other analyses, Roark et al. used LASSIE ⁴⁸ to identify putative sites under selection (**Supplementary Tables 12-13**). This method works by identifying sites where non-TF alleles reach high frequencies. We found modest overlap between the sites under selection as identified

by LASSIE and the mutations that we inferred to be the most strongly selected. For SHIV.CH505, the E640D mutation at site 640 identified by LASSIE is ranked second among 664 mutations in our analysis, and mutations at the remaining 5 sites identified by LASSIE are all within the top 20% of mutations that we infer to be the most beneficial. For SHIV.CH848, the R363Q mutation that is ranked first in our analysis appears at one of the 17 sites identified by LASSIE. Some mutations at the majority of these 17 sites fall within the top 20% most beneficial mutations in our analysis, but some are outliers. In particular, we infer both S291A/P to be somewhat deleterious, with S291P ranked 810th out of 863 mutations.

Beyond the specific context of HIV-1 and bnAb development, our study also provides insight into viral evolution. Despite differences in host biology, the inferred fitness effects of Env mutations in HIV-1 and SHIV were surprisingly similar. This is consistent with the ideas of methods that use sequence statistics across multiple individuals and hosts to predict the fitness effects of mutations [49](#)–[55](#). However, we found that the relationship between the number of individuals in which a mutation was observed and its inferred fitness effect was fairly weak. This suggests that mutational biases and/or sequence space accessibility may play significant roles in short-term viral evolution, even for highly mutable viruses such as HIV-1 and SHIV. As described above, high frequency mutations were also not necessarily highly beneficial. While the recombination rate of HIV-1 is high, correlations between mutations persist, making it difficult to unambiguously interpret frequency changes as signs of selection [24](#).

Methods

Data

We retrieved HIV-1 sequences from CH505 (703010505) and CH848 (703010848) from the HIV sequence database at Los Alamos National Laboratory (LANL) [56](#). The rhesus macaque (RM) SHIV sequences [57](#) were obtained from Gen-Bank [58](#). We then co-aligned SHIV.CH505 and SHIV.CH848 sequences with CH505 and CH848 HIV-1 sequences, respectively, using HIValign [59](#).

CH505

CH505 developed two distinct lineages of CD4 binding site (CD4bs) bnAbs, CH103 and CH235 (refs. [60](#),[61](#)). CH103 antibodies were detectable by 14 weeks after infection and further developed neutralization breadth between 41–92 weeks [62](#). IC50 values of CH235 against the TF virus were 6.5-fold lower than those of CH103 (ref. [60](#)). CH235 lineages could neutralize autologous viruses at week 30. However, viruses that acquired mutations at loop D from 53–100 weeks escaped CH235 neutralization [60](#). Although the neutralization breadth of CH235 was not as broad as that of CH103, this lineage played a critical role; escaping mutations from the CH235 lineage stimulated the development of another lineage with broader neutralization depth [60](#). Mutations in loop D enabled the virus to escape from CH235, but these sequential mutations in loop D, such as E275K, N279D, and V281S, favorably bound to the mature CH103 and continuously increased the binding affinity between mature CH103 and loop D [60](#). Gradually, CH103 matured, developing a broader neutralization breadth.

CH848

CH848 developed DH270, a bnAb that targets the glycosylated site adjacent to the third variable loop (V3). DH270 was detectable three and a half years after infection [63](#). Similar to the CH505 case, the CH848 case exhibited cooperative virus and antibody coevolution. The earlier antibody lineages, DH272 and DH475, could neutralize autologous viruses until week 51 and weeks 15–39, respectively. The virus escaped from DH272 and DH475 afterward, with escape mutations including a longer V1V2 loop. DH270 then developed, with potent and broad neutralization breadth [63](#).

Rhesus macaques

Chimeric viruses, SHIVs, were constructed by bearing the transmitted/founder (TF) Env from three HIV-1 patients, including CH505 and CH848 (ref. [57](#)). In some RMs, SHIV developed similar patterns of mutations to those observed in human donors. In our analysis, we considered RMs with SHIV sequences sampled at at least three points in time. This yielded a set of 7 RMs and 6 RMs for SHIV.CH505 and SHIV.CH848, respectively. The **Supplementary Tables 6**–**7** summarize the number of sequences, time points, and the development of bnAbs for each individual in the SHIV cases as well as HIV-1 cases.

Sequence data processing

Data quality control

To focus our analysis on functional sequences, we removed sequences with more than 200 gaps. To eliminate rare insertions or possible alignment errors, we also masked sites where gaps occurred in more than 95% of sequences within each individual host. To limit errors in virus frequencies, we only considered data from time points with four or more sequences.

Identifying reversions

A mutation is classified as a reversion if the new (mutant) nucleotide matches with the nucleotide at the same site in the HIV-1 consensus sequence from the same subtype. Here, all viruses were subtype C, so we compared with the subtype C consensus sequence as defined by LANL.

Identifying mutations that affect N-linked glycosylation

To identify mutations that affect glycosylation, we search for Env mutations that modify the N-linked glycosylation motif Asn-X-Ser/Thr, where X can be any amino acid except pro-line. We identified three types of mutations affecting glyco-sylation: “shields,” which complete a previously incomplete glycosylation motif, “holes,” which disrupt an existing glyco-sylation motif, and “shifts,” which simultaneously complete one N-linked glycosylation motif and disrupt another.

Enrichment analysis

We used fold enrichment values and Fisher’s exact test to quantify the excess or lack of mutations. For a particular subset of mutations (for example, the top $x\%$ beneficial mutations), we first computed the number of mutations in that subset that do (n_{sel}) and do not (N_{sel}) have a particular property (for example, nonsynonymous mutations in the CD4 binding site). We then computed the total number of mutations that do and do not have the property (n_{null} and N_{null} , respectively) across the entire data set. The fold enrichment value is then $\frac{n_{\text{sel}}/N_{\text{sel}}}{n_{\text{null}}/N_{\text{null}}}$. The term $\frac{n_{\text{sel}}}{N_{\text{sel}}}$ quantifies the fraction of mutations having specific properties across the selected mutations, while the denominator $\frac{n_{\text{null}}}{N_{\text{null}}}$ is the fraction of all mutations that have the property. Fisher’s exact p values are computed from the 2×2 table with n_{sel} , n_{null} , $N_{\text{sel}} - n_{\text{sel}}$ in the first row, and $N_{\text{sel}} - n_{\text{sel}}$, $N_{\text{null}} - n_{\text{null}}$, n_{null} in the second row [64](#).

Inferring fitness effects of mutations

In this section, we describe the inference framework used to infer the fitness effects of mutations (selection coefficients) from temporal genetic data.

Evolutionary model

We model viral evolution with the Wright-Fisher (WF) model, a fundamental model in population genetics ⁶⁵[\[3\]](#). In this model, a population of N individuals (viruses or infected cells, in our case) undergo discrete rounds of selection, mutation, and replication. Each individual a has a genetic sequence $g^a = (g_1^a, g_2^a, \dots, g_L^a)$. Here L is the length of the genetic sequence (number of loci). We write the number of states at each locus (i.e., nucleotides or amino acids) as q . For simplicity, we will use $q = 2$ and $g_i \in \{0, 1\}$ to describe the model below, though we use $q = 5$ and $q = 21$ for DNA and amino acid sequences, respectively, in real HIV-1 and SHIV data. In this binary representation, 0 represents a “wild-type” (WT) or TF allele, and 1 is a mutant allele.

We define the fitness of an individual with genetic sequence g by

$$F(g|s) = 1 + \sum_{i=1}^L s_i g_i. \quad (1)$$

Here s_i is a selection coefficient, quantifying the fitness effect of a mutant allele at locus i . If $s_i > 0$, the mutant allele is beneficial (enhancing replication), and if $s_i < 0$ it is deleterious (impairing replication). By convention, we set the selection coefficient for TF alleles to zero. Individuals with higher fitness values are more likely to replicate than those with lower fitness.

Mutations introduce new genotypes and drive the evolution of the population. Let us define μ^{ab} as the probability of mutation from genotype a to genotype b per replication cycle. Below, we will express this probability in terms of a mutation rate per site per round of replication. In the analysis of real data, we use asymmetric mutation rates estimated from intra-host HIV-1 data ⁶⁶[\[4\]](#).

Given these parameters, the WF model describes the dynamics of the frequencies of different genotypes in the population over time. We write the frequency of genotype a at time t as $z_a(t)$. Given that the frequency of genotypes in the population at time t is $z(t) = (z_1(t), z_2(t), \dots, z_M(t))$, where M is the total number of genotypes, the probability distribution of the frequency of genotypes in the next generation $z(t+1)$ is

$$P(z(t+1)|z(t); s, \mu, N) = N! \prod_{a=1}^M \left(\frac{p_a(z(t))^{N z_a(t+1)}}{[N z_a(t+1)]!} \right). \quad (2)$$

Here p_a is

$$p_a(z(t)) \propto F^a z_a + \sum_{b(\neq a)} [\mu^{ba} z_b(t) - \mu^{ab} z_a(t)]. \quad (3)$$

where F^a is the fitness value of genotype a , based on Eq. (1) ⁶⁷[\[5\]](#). Across K generations, the probability of an entire evolutionary trajectory, defined by the vector of genotype frequencies at each time, is then

$$P((z(t_k))_{k=0}^K | s, \mu, N) = \prod_{k=0}^{K-1} P(z(t_{k+1}) | z(t_k); s, \mu, N). \quad (4)$$

Diffusion limit

When the population size is sufficiently large, the evolution of the population defined in Eq. (2) can be reasonably well approximated by a Gaussian process, which is a solution to the Fokker-Planck (forward Kolmogorov) equation [67](#), [68](#),

$$P(z(t+\Delta t)|z(t); s, \mu, N) \sim \mathcal{N}\left(z(t) + \Delta t d(t), C(z(t))/N\right), \quad (5)$$

with the drift vector $d(t)$ and the diffusion matrix $C(z(t))/N$ such that

$$C_{ab}(z(t)) = \begin{cases} -z_a(t)z_b(t) & \text{for } a \neq b \\ z_a(t)(1-z_a(t)) & \text{for } a = b, \end{cases} \quad (6)$$

and

$$d(t) = C(z(t))s + \mu^{\text{flux}}. \quad (7)$$

Dimensional reduction

While the WF process in genotype space provides valuable insights into genotype dynamics, the mathematical expressions are sometimes challenging to interpret. To obtain more intuitive expressions, we can project the dynamics onto the space of allele frequencies,

$$x_i = \sum_a g_i^a z_a. \quad (8)$$

One can then find the drift vector d and diffusion matrix C/N in allele frequency space,

$$d_i = C_{ii}s_i + \sum_{j(\neq i)} C_{ij}s_j + \mu_i^{\text{flux}}, \quad (9)$$

and

$$C_{ij} = \begin{cases} x_{ij} - x_i x_j & \text{for } i \neq j \\ x_i(1-x_i) & \text{for } i = j. \end{cases} \quad (10)$$

Here, x_{ij} is the frequency of individuals with alleles i and j , and μ_i^{flux} is net expected change in frequency of allele i due to mutations, which is given explicitly in Eq. (14) below. The first term in Eq. (9) gives the expected change in the frequency x_i due to the direct fitness effect s_i , while the second term represents the contributions due to indirect or genetic linkage effects with other alleles j .

Maximum path likelihood

Following recent work [69](#), we employed Bayes' rule to find the selection coefficients that best explain the data. These are the coefficients that maximize the posterior distribution

$$P_{\text{posterior}}(s|(z(t_k))_k) \propto P((z(t_k))_k|s) P_{\text{prior}}(s), \quad (11)$$

which is a product of the likelihood of the evolutionary trajectory observed in the data Eq. (4) (under the diffusion limit Eq. (5)) and a prior distribution for the selection coefficients. We chose a Gaussian prior distribution with zero mean and a covariance of $I/(Ny)$, where I is the

identity matrix. This prior distribution penalizes the inferences of large selection coefficients when they are not well-supported by the data. The maximum *a posteriori* selection coefficients are then given by [69](#).

$$\hat{s} = (C_{\text{int}} + \gamma I)^{-1} [\Delta x_{\text{int}} - \Delta \mu_{\text{int}}]. \quad (12)$$

Here C_{int} , Δx_{int} , and $\Delta \mu_{\text{int}}$ represent the covariance matrix, vector of frequency changes, and mutational flux integrated over the evolution

$$\begin{aligned} C_{\text{int}} &= \sum_{k=0}^K \Delta t_k C(x(t_k)) \\ \Delta x_{\text{int}} &= x(t_K) - x(t_0) \\ \Delta \mu_{\text{int}} &= \sum_{k=0}^{K-1} \Delta t_k \mu^{\text{flux}}(t_k). \end{aligned} \quad (13)$$

The mutational flux μ^{flux} is characterized by the rates of mutations from nucleotides β to α , denoted by $\mu_{\alpha\beta}$, which are determined from longitudinal HIV-1 populations in untreated patients [66](#). The change of the α nucleotide frequency at locus i due to mutation is given by:

$$\mu_{i,\alpha}^{\text{flux}} = \sum_{\beta} (\mu_{\alpha\beta} x_{i,\beta}(t_k) - \mu_{\beta\alpha} x_{i,\alpha}(t_k)). \quad (14)$$

Inverting the integrated covariance matrix effectively reveals the underlying direct allele interactions and resolves the genetic linkage effects.

The shift in the covariance diagonal in [Eq. \(12\)](#), arising from the selection coefficients' posterior distribution, reflects the uncertainty in the selection distribution. We used $\gamma = 10$ for all simulations, but the model is robust to variation in the strength of regularization [69](#). For the mutation rates, we incorporated the transition probabilities among arbitrary DNA nucleotides, estimated from whole-genome deep sequencing of multiple untreated HIV-1 patients followed for 5 to 8 years post-infection [66](#).

Integration of covariance

When the time interval of the observation Δt is sufficiently short, the trajectory of the allele frequency would be continuous and ideally it would be a smooth curve [70](#). To accurately estimate the covariance matrix, we employ piecewise linear interpolation for frequencies. Let $\tau \in [0, 1]$, then the linear interpolation for a frequency vector can be expressed as:

$$\begin{aligned} x_i^{[k,k+1]}(\tau) &= (1 - \tau)x_i(t_k) + \tau x_i(t_{k+1}), \\ x_{ij}^{[k,k+1]}(\tau) &= (1 - \tau)x_{ij}(t_k) + \tau x_{ij}(t_{k+1}), \end{aligned} \quad (15)$$

which yields,

$$\begin{aligned} x_i^{\text{int}} &= \sum_{k=0}^{K-1} \Delta t_k \int_0^1 d\tau x_i^{[k,k+1]}(\tau) \\ C_{ij}^{\text{int}} &= \sum_{k=0}^{K-1} \Delta t_k \int_0^1 d\tau \left(x_{ij}^{[k,k+1]}(\tau) - x_i^{[k,k+1]}(\tau) x_j^{[k,k+1]}(\tau) \right) \end{aligned}$$

The explicit expression of the integrated covariance is given in refs. [69](#), [70](#).

Joint RM model

In addition to fitness models derived from SHIV data for individual RMs, we inferred a joint model under the assumption that virus evolution within each individual RM with the same TF virus is governed by a similar fitness landscape. This method improves inference accuracy from the WF process and deep mutational scanning data [71–73](#). The joint path like-likelihood for allele frequency trajectories across RMs with the same TF virus is then

$$p(((x^\alpha(t_k))_{k=0}^{K_\alpha})_{\alpha=1}^R | s, \mu, \gamma) = \left(\prod_{\alpha=1}^R \prod_{k=0}^{K-1} p_M(x^\alpha(t_{k+1}) | x^\alpha(t_k); s, \mu, N) \right) \times p(x(t_0)) p(s | \gamma). \quad (16)$$

Here, x^α is the allele frequency of the α -th individual and R is the number of replicate individuals (i.e., the number of RMs sharing the same TF virus). The initial state is $p(x^\alpha(t_0)) = p(x(t_0))$ for all α for individuals with the same TF virus. The solution of the joint path likelihood is given by

$$\hat{s} = (\overline{C}_{\text{int}} + \gamma I)^{-1} [\overline{\Delta x}_{\text{int}} - \overline{\Delta \mu}_{\text{int}}]. \quad (17)$$

Here, the overbar denotes the sum over the replicate RMs. We emphasize that the joint selection coefficients in

Eq. (17) are not the same as selection coefficients that are simply averaged across RMs with the same TF virus. The joint selection coefficients are more robust, as they are guided by the level of evidence within each individual rather than naive averaging.

Geometrical interpretation of the fitness comparison


The Pearson values we utilized to compare the fitness landscapes denoted as $F_s(g) := F(g | s)$ and $F_h(g) := F(g | h)$ can be expressed by the following simple relation:

$$r = \frac{\text{Cov}(F_s, F_h)}{\sqrt{\text{Var}(F_s)} \sqrt{\text{Var}(F_h)}} = \frac{s^\top C h}{\sqrt{s^\top C s} \sqrt{h^\top C h}}. \quad (18)$$

Here, $\text{Cov}(F_s, F_h)$ and $\text{Var}(F_s)$ represent the covariance and variance values estimated from the samples being compared, $(F_s(g^i), F_h(g^i))_i$. C is the covariance matrix defined between arbitrary loci. The last equation can be interpreted as an angle between two vectors, s and h , with a metric matrix C ; if $s = h$, the Pearson value clearly becomes 1. However, the “similarity” also depends on how these vectors are projected by C ; eigenmodes associated with larger variance of statistics will be more emphasized. The last expression readily implies an interpretation for the case of shuffled sequences; shuffling the sequences equates to diluting the covariance between loci, resulting in the metric matrix becoming a diagonal matrix. Removing the off-diagonal elements corresponds to lifting the constraints on the fitness landscape.

Acknowledgements

The work of K.S.S. and J.P.B. reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM138233.



Supplementary Fig. 1.


CH848 exhibits similar spatial patterns of selection coefficients; significantly strongly selected mutations are located at the apex, particularly on the edge of the Env protein.

A and B, Side and top views of the Env protein with inferred selection coefficient values. The apex region is enriched with moderately and strongly beneficial mutations. **C-F,** Mutations that are presumably resistant to bnAbs and autologous nAbs are highlighted. The selection values in the autologous nAbs binding regions were slightly larger than those in the bnAbs binding regions. **G-K,** Top views highlight the individual variable regions.

Supplementary Fig. 2.

Frequencies of different types of HIV-1 mutations over time.


CH505 (A) and CH848 (B) mutation frequencies over time. Mutation types are the same as in Fig. 3 in the main text, but with all mutations affecting resistance to bnAb lineage antibodies grouped together.




Supplementary Fig. 3.

Weak association between the number of RMs in which a SHIV mutation was observed and its inferred fitness effect.

Inferred selection coefficients for SHIV.CH505 (A) and SHIV.CH848 (B) mutations, sorted by the number of RMs in which the mutation was observed.






Supplementary Fig. 4.

Broad similarity between HIV-1 and SHIV fitness landscapes with the same TF Env sequence.

A, Fitness estimates for a sample of artificial Env sequences, obtained by independently shuffling observed amino acids in SHIV.CH505 sequences at each residue. This random sequence ensemble conserves single-residue frequencies, but not correlations between mutations. Even on these artificial sequences, the fitness estimates using a model trained on HIV-1 data strongly agree with the SHIV model (Pearson's $r = 0.84$, $p < 10^{-20}$). This implies that the similarity of the fitness landscapes is not confined to the specific genotypes observed in HIV-1 or SHIV evolution, but also extends to more distant sequences. **B**, Similar results also hold for fitness landscapes based on CH848 and SHIV.CH848 data (Pearson's $r = 0.74$, $p < 10^{-20}$).



Supplementary Fig. 5.

Contributions of different types of SHIV mutations to viral fitness gains over time.

SHIV.CH505 (A) and SHIV.CH848 (B) mutations were grouped into four categories to assess their contributions to SHIV fitness gains over time. For SHIV.CH505, the mutations N334S, H417R, K302N, Y330H, N279D, and N130D are classified as viral load (VL) enhancing mutations, following ref. [74](#). For both SHIV.CH505 and SHIV.CH848, we then separated out the contributions of known antibody resistance mutations, including mutations that affect N-linked glycosylation motifs. We then computed the collective fitness contributions from subsets of mutations that affect N-linked glycosylation motifs that were not known to affect resistance to specific antibodies, and reversions to the HIV-1 subtype consensus sequence. Each mutation appears in only one category in this figure, sorted in the order above. For example, a mutation that affects an N-linked glycosylation motif and which is a reversion to the subtype consensus sequence, but which has not been established to affect resistance to a specific antibody, would have its contribution to fitness counted in the glycan category.

Supplementary Table 1.

CH103 and CH235 resistance mutations.

List of mutations considered to be CH103 or CH235 resistance mutations, and supporting references. Here “X” refers to any amino acid.

Mutation	Note
T234N	Increases resistance to CD4BS Abs including CH235 (ref. ⁷⁵)
E275K	Mutation combination E275K+N279D+V281S escape from CH235 (ref. ⁶⁰)
E279D	Mutation combination E275K+N279D+V281S escape from CH235 (ref. ⁶⁰)
N279K	Mutation combination E275K+N279D+V281S escape from CH235 (ref. ⁶⁰)
N280S	Mutation combination N280S+V281A escape from CH235 (ref. ⁶⁰)
N280T	Mutation combination N280T+V281A escape from CH235 (ref. ⁶⁰)
V281A	Mutation combination V281A+V281G+V281D escape from CH235 (ref. ⁶⁰)
V281G	Mutation combination V281A+V281G+V281D escape from CH235 (ref. ⁶⁰)
V281D	Mutation combination V281A+V281G+V281D escape from CH235 (ref. ⁶⁰)
V281S	Mutation combination E275K+N279D+V281S escape from CH235 (ref. ⁶⁰)
V281S	Mutation combination E275K+N279D+V281S escape from CH235 (ref. ⁶⁰)
S365P	Complete escape from CH103 (ref. ⁶⁰)
K460E	Resistance to CH235 lineage in CH505 and RM6072 (ref. ⁵⁷)
460-465X	V5 insertion increases resistance to CH103 (ref. ⁶⁰)

Supplementary Table 2.

Strain-specific antibody resistance mutations in CH505.

List of mutations considered to be strain-specific antibody resistance mutations and supporting references. Here “X” refers to any amino acid.

Mutation	Note
132-152X	V1 insertion increases resistance to intermediate CH103 lineage ⁶⁰
X197N	Loss of an N-linked glycan leads to loss of resistance to intermediate CH235 lineages ⁷⁶
X276N	Loss of an N-linked glycan leads to loss of resistance to intermediate CH235 lineages ⁷⁶
N279A	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N279C	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N279E	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N279V	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N279S	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N279T	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N279H	Resistant to UCA and intermediate CH235 lineage ⁷⁶
N280D	Resistant to UCA and intermediate CH235 lineage ⁷⁵
X293T	Resistant to UCA and intermediate antibodies ⁷⁶
X293S	Resistant to UCA and intermediate antibodies ⁷⁶
X362N	Loss of an N-linked glycan leads to loss of resistance to intermediate CH235 lineages ⁷⁶
G458Y	Resistant to UCA of CH235 lineage ^{76,77}
G458F	Resistant to UCA of CH235 lineage ⁷⁶
G458W	Resistant to UCA of CH235 lineage ⁷⁶
G458R	Resistant to UCA of CH235 lineage ⁷⁶
G458K	Resistant to UCA of CH235 lineage ⁷⁶
G458S	Resistant to UCA of CH235 lineage ⁷⁶
G458D	Resistant to UCA and intermediate CH235 lineage ⁷⁶
G458E	Resistant to UCA and intermediate CH235 lineage ⁷⁶
X461N	Loss of an N-linked glycan leads to loss of resistance to intermediate CH235 lineages ⁷⁶
X462N	Loss of an N-linked glycan leads to loss of resistance to intermediate CH235 lineages ⁷⁶

Mutation	Note
E87K	Resistance to DH272 lineage ⁶³
E87D	Resistance to DH272 lineage ⁶³
E87G	Resistance to DH475 lineage ⁶³
D185N	Resistance to DH272 and DH475 lineages ⁶³
D185S	Resistance to DH272 and DH475 lineages ⁶³
G300N	Resistance to DH272 and DH475 lineages ⁶³
N301H	Resistant to intermediate DH270 ⁷⁸
V323X	Recognized by intermediate antibody DH270.I5 ⁷⁸
D325N	Resistance to DH272 lineage ^{63,78}
V326X	Recognized by intermediate antibody DH270.I5 ⁷⁸
P326X	Recognized by intermediate antibody DH270.I5 ⁷⁸
T326X	Recognized by intermediate antibody DH270.I3 and DH270.I5 ⁷⁸
K327X	Resistance to intermediate antibody DH270.I5 ⁷⁸
E335K	Resistance to DH272 lineage ⁶³
E344K	Resistance to DH272 and DH475 lineages ⁶³
N413T	Resistance to DH272 and DH475 lineages ⁶³
D624G	Resistance to DH272 lineage ⁶³
I818L	Resistance to DH272 and DH475 lineages ⁶³

Supplementary Table 3.

DH272, DH475, and strain-specific antibody resistance mutations in CH848.

List of mutations considered to be DH272, DH475, and strain-specific antibody resistance mutations, and supporting references. Here "X" refers to any amino acid.

Rank	Mutation (DNA)	Mutation (AA)	Date	Region	Selection coefficient (%)	Note
1	A7225G	N334S	113	Near V3 Loop	8.92	Reversion, N-linked glycan shift, resistance to DH950 antibodies ⁷⁹
2	T7066A	V281D	183	CD4bs/Loop D	5.66	Resistant to intermediate CH235 lineage antibodies ⁶⁰ , combination with N280S or N280T gives resistance to mature CH235 (ref. ⁶⁰)
3	G7776A	V518M	1093	Fusion Peptide	5.42	
4	C6709T	T162I	129	V2 Loop	5.40	N-linked glycan hole
5	G6682A	G153E	71	V1 Loop	5.24	Reversion
6	C7462A	T413N	129	V4 Loop	5.16	N-linked glycan shift
7	C7468A	T415K	71	V4 Loop	4.82	Putative CD8 ⁺ T cell escape mutation ⁶⁰
8	A7130C	K302N	183	V3 Loop	3.74	Reduces neutralization of DH950-like Abs ⁷⁹
9	G7769A	M515I	1603	Fusion Peptide	3.57	Reversion
10	A7614G	N464D	183	V5 Loop	3.46	Reversion
11	C7462T	T413I	23	V4 Loop	3.42	Putative CD8 ⁺ T cell escape mutation ⁶⁰
12	G7776T	V518L	1640	Fusion Peptide	3.37	
13	T7212C	Y330H	129	V3 Loop	3.28	Reversion, reduces neutralization of DH950-like Abs ⁷⁹
14	G6661A	S146N	129	V1 Loop	3.14	Reversion, N-linked glycan shield
15	G7453A	S410N	23	V4 Loop	3.08	Putative CD8 ⁺ T cell escape mutation ⁶²
16	G6634T	S137I	183	V1 Loop	3.00	N-linked glycan hole
17	G6634C	S137A	129	V1 Loop	2.82	Reversion, N-linked glycan hole
18	T6588A	L122M	183	Near CD4bs	2.81	
19	C7456A	T411K	29	V4 Loop	2.75	Reversion, N-linked glycan hole, and putative CD8 ⁺ T cell escape mutation ⁶²
20	A6596C	P124P	183	CD4bs	2.73	
21	A7474G	H417R	23	V4 Loop	2.63	Putative CD8 ⁺ T cell escape mutation ⁶²
22	A7601dG	N459D	1603	CD4bs	2.61	N-linked glycan shield, located in the CH235 antibody binding region ⁶²
23	A7056T	T278S	1603	Loop D	2.59	Located in the CH235 antibody binding region ⁶²
24	G6630A	A136T	113	V1 Loop	2.59	Reversion
25	A8143G	N640S	1780	Near Glycosite	2.55	

Supplementary Table 4.

Biological effects of the HIV-1 mutations inferred to be the most beneficial in CH505.

The table lists the 25 mutations in CH505 with the highest (i.e., most beneficial) selection coefficients, among over 2500 mutations in total. All of the top mutations are nonsynonymous. Mutations associated with immune evasion, changes in glycosylation, and reversions to the subtype C consensus sequence are frequently observed among these top mutations.

Supplementary Table 5.

Biological effects of the HIV-1 mutations inferred to be the most beneficial in CH848.

This table is analogous to [Supplementary Table 4](#), but for CH848. Unlike for CH505, longer sequences were available for CH848, including genes beyond Env. We observed multiple beneficial mutations in Nef and Env.

Rank	Mutation (DNA)	Mutation (AA)	Date	Region	Selection coefficient (%)	Note
1	T9043G	S83A	49	Nef	12.55	Reversion
2	A8988cT	E65D	38	Nef	8.28	
3	G8911A	E39K	38	Nef	6.68	Reversion
4	A8988bG	E65G	38	Nef	6.42	
5	A8988cC	E65D	43	Nef	5.48	
6	G6640A	S139N	358	V1 Loop	5.22	Reversion, N-linked glycan shift, presence of N-linked glycan at site 138 decreases neutralization by the earlier DH270 UCA and removes glycan at site 137 (ref. ⁷⁹)
7	G6679A	G152E	38	V1 Loop	4.83	
8	C9044G	A84G	78	Nef	4.70	Reversion
9	A8915G	H41R	38	Nef	4.48	Reversion
10	A8083T	K620M	808	gp41	4.44	
11	A8988cG	E65E	38	Nef	4.35	
12	G7123A	S300N	700	V3 Loop	4.06	Reversion, confers resistance to DH272 and DH475 (ref. ⁶³)
13	G6640T	S139V	445	V1 Loop	3.92	N-linked glycan hole
14	A7310C	K362N	780	V1 Loop	3.87	N-linked glycan shield
15	A8083C	K620T	780	gp41	3.84	
16	A8912C	E40A	32	Nef	3.83	
17	A8895G	A34A	49	Nef	3.77	Reversion
18	A8912T	E40V	43	Nef	3.74	
19	G6640C	S139T	1432	V1 Loop	3.73	N-linked glycan shield
20	A6621C	N133H	1120	V1 Loop	3.66	N-linked glycan hole
21	G8911C	E39Q	49	Nef	3.64	
22	C7066T	A281V	1651	CD4BS	3.62	Reversion
23	A6912G	N230D	194	Glycosite	3.57	N-linked glycan hole, amino acid N is sensitive to DH270.5 (ref. ⁷⁵)
24	Δ8911G	Δ39E	78	Nef	3.57	
25	A7614G	T464E	445	V5 Loop	3.39	Reversion, N-linked glycan hole

Supplementary Table 6.

CH505 and SHIV.CH505 sequence statistics.

The number of sequences and time points listed here are given after filtering. Variable sites are defined as sites where at least two different nucleotides (including deletions) were observed.

identifier	# of sequences	# of time points	# of variable sites	bnAbs detected	bnAb type
703010505	578	23	1131	Yes	CD4bs
RM5695	330	8	467	Yes	V2 Apex
RM6070	215	5	334	Yes	V2 Apex
RM6072	355	13	710	No	
RM6697	228	6	386	No	
RM6699	118	5	207	No	
RM6701	98	5	148	No	
RM6703	131	4	275	No	

Supplementary Table 7.

CH848 and SHIV.CH848 sequence statistics.

The format of this table is the same as in [Supplementary Table 6](#).

identifier	# of sequences	# of time points	# of variable sites	bnAbs detected	bnAb type
703010848	1205	31	2694	Yes	V3-glycan
RM6163	261	8	467	Yes	V3-glycan
RM6167	349	10	657	Yes	V3-glycan
RM6700	260	7	568	No	
RM6713	281	7	567	No	
RM6714	284	8	516	No	
RM6720	193	8	340	No	

Supplementary Table 8.

Biological effects of strongly selected SHIV.CH505 mutations.

Five out of the top six mutations have been shown to increase viral load in RMs [74](#). Selection coefficients and ranks are based on the joint evolutionary model across SHIV.CH505-infected RMs. These analyses focus on common mutations observed in three or more RMs.

Rank	Mutation (AA)	Region	Selection coefficient (%)	Note
1	N334S	Near V3 Loop	2.44	N-linked glycan shift, enhanced viral replication ⁷⁴ , reduced early plasma neutralization ⁵⁷
2	Y330H	V3 Loop	1.99	Improved virus entry and replication ⁷⁴
3	N279D	CD4BS	1.69	Enhanced replication ⁷⁴ , reduced early plasma neutralization, conferred resistance to strain-specific CD4bs nAbs DH650 in RM6070 (ref. ⁵⁷)
4	K462E	V5 Loop	1.40	
5	H356N	Glycosite	1.16	
6	K302N	V3 Loop	1.15	Enhanced replication and affects virus entry, conferred resistance to strain-specific V3 nAbs, DH647 and DH648 (ref. ⁷⁴)
7	G153E	V1 Loop	1.12	
8	N130D	Near V1 Loop	0.99	Reduced early plasma neutralization ⁵⁷
9	T234N	Glycosite	0.91	N-linked glycan shield, reduced early plasma neutralization, conferred resistance to strain-specific CD4bs nAbs DH650 in RM6070 (ref. ⁵⁷)
10	H417R	V4 Loop	0.87	Enhanced replication ⁷⁴ , putative CD8 ⁺ T cell escape mutation ^{57,74}

Supplementary Table 9.

Biological effects of strongly selected SHIV.CH848 mutations.

Selection coefficients and ranks are based on the joint evolutionary model across SHIV.CH848-infected RMs.

Rank	Mutation (AA)	Region	Selection coefficient (%)	Note
1	P179L	V2 Loop	1.95	
2	N413T	V4 Loop	1.70	N-linked glycan hole, loss of N413 alters sensitivity to early nAbs in RM6167 (ref. ⁵⁷)
3	N413S	V4 Loop	1.43	N-linked glycan hole, aids in antibody escape ⁵⁷
4	D137S	V1 Loop	1.29	
5	N462T	V4 Loop	1.29	N-linked glycan shift
6	S407N	V4 Loop	1.22	
7	R336E	Near V3 Loop	1.20	Mutations at site 336 observed in all RMs, R336G reduced sensitivity to plasma in RM6167 (ref. ⁵⁷)
8	P179S	V2 Loop	1.18	
9	V142fA	V1 Loop	1.15	
10	N462D	V4 Loop	1.15	N-linked glycan hole

Supplementary Table 10.

Selective advantage of mutations that confer resistance to antibodies in SHIV.CH505.

Identification of resistance mutations was performed by Roark and colleagues ⁵⁷.

Mutation	RM5695 (%)	RM6072 (%)	RM6701 (%)	RM6699 (%)	RM6697 (%)	RM6070 (%)	RM6703 (%)	Note
N130D	1.5	1.1			1.0	0.90	0.87	Confers resistance to V3 mAbs DH647 and DH648 in RM6072
R166K	0.49					1.77		Confers resistance to V2 apex bnAbs RHA1 in RM5695
R166S	0.7							Confers resistance to V2 apex bnAbs RHA1 in RM5695
R169K	0.43	-0.1				-0.04		Confers resistance to V2 apex bnAbs RHA1 in RM5695
T234N	0.36	4.5			0.34	1.1		Confers resistance to CD4bs mAbs DH650 in RM6072
N279K	1.2	0.013	-0.32	0.44	-0.19	-0.3		Confers resistance to CD4bs mAbs DH650 in RM6072
V281I	0.14	-0.03		0.59		1.4		Confers resistance to V3 mAbs DH647 and DH648 in RM6072
K302N	6.19	0.59			0.76			Confers resistance to V3 mAbs DH647 and DH648 in RM6072
Y330H	6.5	3.0	0.03	1.1	0.11	-0.56	1.4	Confers resistance to V3 mAbs DH647 and DH648 in RM6072
N334S	7.2	4.6	1.8	1.6	1.4	3.1	2.4	Confers resistance to V3 mAbs DH647 and DH648 in RM6072

Supplementary Table 11.

Selective advantage of mutations that confer resistance to antibodies in SHIV.CH848.

Identification of resistance mutations was performed by Roark and colleagues ⁵⁷.

Mutation	RM6163 (%)	RM6167 (%)	RM6700 (%)	RM713 (%)	RM6714 (%)	RM6720 (%)	Note
N332S		-0.19					Mutations at or near N332 to 334 can modulate sensitivity to neutralization
R336G	0.62	0.4	0.94	0.24	0.12	0.98	Led to a reduction in neutralization by autologous nAb in RM6167

Supplementary Table 12.

List of selected sites using LASSIE in SHIV.CH505.

This analysis was performed by Roark et al. [57](#).

HXB2 Site
N130
T234
N279
V281
N300
K302
Y330
N334
K347
H355
T413
H417
G471
G620
E649

Supplementary Table 13.

List of selected sites using LASSIE in SHIV.CH848.

This analysis was performed by Roark et al. [57](#).

HXB2 Site
N130
N289
S291
G300
R336
Q337
K340
E344
R363
A372
T411
N413
T467
H651
K658
R683

Additional information

Author contributions

All authors contributed to the research design, methods development, interpretation of results, and writing the paper. K.S.S. performed simulations and computational analyses. J.P.B. supervised the project.

Code availability

Sets of processed data and computer code used in this study are available in the GitHub repository of <https://github.com/bartonlab/paper-HIV-coevolution>. This repository contains source files that process HIV-1 and SHIV sequences, infer selection coefficients, and identify and characterize mutations. The included Jupyter note-books can be run to reproduce the figures presented here. The original HIV-1 sequences can be retrieved from the LANL database (<https://www.hiv.lanl.gov/content/index>), and SHIV sequences can be found at GenBank (<https://www.ncbi.nlm.nih.gov/>).

References

1. Wei X., et al. 2003) **Antibody neutralization and escape by hiv-1** *Nature* **422**:307–312
2. Allen T. M., et al. 2005) **Selective escape from cd8+ t-cell responses represents a major driving force of human immunodeficiency virus type 1 (hiv-1) sequence diversity and reveals constraints on hiv-1 evolution** *Journal of virology* **79**:13239–13249
3. Li B., et al. 2007) **Rapid reversion of sequence polymorphisms dominates early human immunodeficiency virus type 1 evolution** *Journal of virology* **81**:193–201
4. Altfeld M., Allen T. M. 2006) **Hitting hiv where it hurts: an alternative approach to hiv vaccine design** *TRENDS in Immunology* **27**:504–510
5. Kwong P. D., Mascola J. R., Nabel G. J. 2013) **Broadly neutralizing antibodies and the search for an hiv-1 vaccine: the end of the beginning** *Nature Reviews Immunology* **13**:693–701
6. Burton D. R., Hangartner L. 2016) **Broadly neutralizing antibodies to hiv and their role in vaccine design** *Annual review of immunology* **34**:635–659
7. Sok D., Burton D. R. 2018) **Recent progress in broadly neutralizing antibodies to hiv** *Nature immunology* **19**:1179–1188
8. Haynes B. F., et al. 2023) **Strategies for hiv-1 vaccines that induce broadly neutralizing antibodies** *Nature Reviews Immunology* **23**:142–158
9. Doria-Rose N. A., et al. 2010) **Breadth of human immunodeficiency virus-specific neutralizing activity in sera: clustering analysis and association with clinical variables** *Journal of virology* **84**:1631–1636
10. Haynes B. F., Kelsoe G., Harrison S. C., Kepler T. B. 2012) **B-cell-lineage immunogen design in vaccine development with hiv-1 as a case study** *Nature biotechnology* **30**:423–433
11. Haynes B. F., et al. 2016) **Hiv-host interactions: implications for vaccine design** *Cell host & microbe* **19**:292–303
12. Hraber P., et al. 2014) **Prevalence of broadly neutralizing antibody responses during chronic hiv-1 infection** *AIDS (London, England)* **28**:163
13. Liao H.-X., et al. 2013) **Co-evolution of a broadly neutralizing hiv-1 antibody and founder virus** *Nature* **496**:469–476
14. Bonsignori M., et al. 2017) **Staged induction of hiv-1 glycan-dependent broadly neutralizing antibodies** *Science translational medicine* **9**:eaai7514
15. Doria-Rose N. A., et al. 2014) **Developmental pathway for potent v1v2-directed hiv-neutralizing antibodies** *Nature* **509**:55–62
16. Moore P. L., Williamson C., Morris L. 2015) **Virological features associated with the development of broadly neutralizing antibodies to hiv-1** *Trends in microbiology* **23**:204–211

17. Landais E., Moore P. L. 2018) **Development of broadly neutralizing antibodies in hiv-1 infected elite neutralizers** *Retrovirology* **15**:1–14
18. Gray E. S., et al. 2011) **The neutralization breadth of hiv-1 develops incrementally over four years and is associated with cd4+ t cell decline and high viral load during acute infection** *Journal of virology* **85**:4828–4840
19. Cornelissen M., et al. 2016) **The neutralizing antibody response in an individual with triple hiv-1 infection remains directed at the first infecting subtype** *AIDS research and human retroviruses* **32**:1135–1142
20. Roark R. S., et al. 2021) **Recapitulation of hiv-1 env-antibody coevolution in macaques leading to neutralization breadth** *Science* **371**:eabd2638
21. Tomaras G. D., et al. 2008) **Initial b-cell responses to transmitted human immunodeficiency virus type 1: virion-binding immunoglobulin m (igm) and igg antibodies followed by plasma anti-gp41 antibodies with ineffective control of initial viremia** *Journal of virology* **82**:12449–12463
22. Gao F., et al. 2014) **Cooperation of b cell lineages in induction of hiv-1-broadly neutralizing antibodies** *Cell* **158**:481–491
23. Bonsignori M., et al. 2016) **Maturation pathway from germline to broad hiv-1 neutralizer of a cd4-mimic antibody** *Cell* **165**:449–463
24. Sohail M. S., Louie R. H., McKay M. R., Barton J. P. 2021) **Mpl resolves genetic linkage in fitness inference from complex evolutionary histories** *Nature biotechnology* **39**:472–479
25. Sohail M. S., Louie R. H., Hong Z., Barton J. P., McKay M. R. 2022) **Inferring epistasis from genetic time-series data** *Molecular biology and evolution* **39**:msac199
26. Lee B., et al. 2022) **Inferring effects of mutations on sars-cov-2 transmission from genomic surveillance data** *medRxiv* :2021–12
27. Li Y., Barton J. P. 2023) **Estimating linkage disequilibrium and selection from allele frequency trajectories** *Genetics* **223**:iyac189
28. Li Y., Barton J. P. 2024) **Correlated allele frequency changes reveal clonal structure and selection in temporal genetic data** *Molecular Biology and Evolution* **41**:msae060
29. Jardine J. G., et al. 2016) **Minimally mutated hiv-1 broadly neutralizing antibodies to guide reductionist vaccine design** *PLoS pathogens* **12**:e1005815
30. Zanini F., et al. 2015) **Population genomics of inpatient hiv-1 evolution** *eLife* **4**:e11282
31. Das S. R., Jameel S. 2005) **Biology of the hiv nef protein** *Indian J Med Res* **121**:315–32
32. Barton J. P., et al. 2019) **Modelling and in vitro testing of the hiv-1 nef fitness landscape** *Virus evolution* **5**:vez029
33. McCurley N. P., et al. 2017) **Hiv transmitted/founder vaccines elicit autologous tier 2 neutralizing antibodies for the cd4 binding site** *PLoS One* **12**:e0177863

34. Kong R., et al. 2019) **Antibody lineages with vaccine-induced antigen-binding hotspots develop broad hiv neutralization** *Cell* **178**:567–584
35. Saunders K. O., et al. 2022) **Stabilized hiv-1 envelope immunization induces neutralizing antibodies to the cd4bs and protects macaques against mucosal infection** *Science translational medicine* **14**:eabo5598
36. Bauer A., et al. 2023) **Adaptation of a transmitted/founder simian-human immunodeficiency virus for enhanced replication in rhesus macaques** *PLoS Pathogens* **19**:e1011059
37. Hatzioannou T., Evans D. T. 2012) **Animal models for hiv/aids research** *Nature Reviews Microbiology* **10**:852–867
38. Li H., et al. 2016) **Envelope residue 375 substitutions in simian-human immunodeficiency viruses enhance cd4 binding and replication in rhesus macaques** *Proceedings of the National Academy of Sciences* **113**:E3413–E3422
39. Wang S., et al. 2015) **Manipulating the selection forces during affinity maturation to generate cross-reactive hiv antibodies** *Cell* **160**:785–797
40. Escolano A., et al. 2016) **Sequential immunization elicits broadly neutralizing anti-hiv-1 antibodies in ig knockin mice** *Cell* **166**:1445–1458
41. Sprenger K. G., Louveau J. E., Murugan P. M., Chakraborty A. K. 2020) **Optimizing immunization protocols to elicit broadly neutralizing antibodies** *Proceedings of the National Academy of Sciences* **117**:20077–20087
42. Zhou T., et al. 2013) **Multidonor analysis reveals structural elements, genetic determinants, and maturation pathway for hiv-1 neutralization by vrc01-class antibodies** *Immunity* **39**:245–258
43. Shaffer J. S., Moore P. L., Kardar M., Chakraborty A. K. 2016) **Optimal immunization cocktails can promote induction of broadly neutralizing abs against highly mutable pathogens** *Proceedings of the National Academy of Sciences* **113**:E7039–E7048
44. Nourmohammad A., Otwinowski J., Plotkin J. B. 2016) **Host-pathogen coevolution and the emergence of broadly neutralizing antibodies in chronic infections** *PLoS genetics* **12**:e1006171
45. Dosenovic P., et al. 2015) **Immunization for hiv-1 broadly neutralizing antibodies in human ig knockin mice** *Cell* **161**:1505–1515
46. Williams W. B., et al. 2023) **Vaccine induction of heterologous hiv-1 neutralizing antibody b cell lineages in humans** *medRxiv* :2023–3
47. Gao Y., Barton J. P. 2024) **A binary trait model reveals the fitness effects of hiv-1 escape from t cell responses** *bioRxiv*
48. Hraber P., et al. 2015) **Longitudinal antigenic sequences and sites from intra-host evolution (lassie) identifies immune-selected hiv variants** *Viruses* **7**:5443–5475
49. Ferguson A. L., et al. 2013) **Translating hiv sequences into quantitative fitness landscapes predicts viral vulnerabilities for rational immunogen design** *Immunity* **38**:606–617

50. Mann J. K., et al. 2014) **The fitness landscape of hiv-1 gag: advanced modeling approaches and validation of model predictions by in vitro testing** *PLoS computational biology* **10**:e1003776
51. Łuksza M., Lässig M. 2014) **A predictive fitness model for influenza** *Nature* **507**:57–61
52. Lässig M., Mustonen V., Walczak A. M. 2017) **Predicting evolution** *Nature ecology & evolution* **1**:0077
53. Barton J. P., et al. 2016) **Relative rate and location of intra-host hiv evolution to evade cellular immunity are predictable** *Nature communications* **7**:11660
54. Louie R. H., Kaczorowski K. J., Barton J. P., Chakraborty A. K., McKay M. R. 2018) **Fitness landscape of the human immunodeficiency virus envelope protein that is targeted by antibodies** *Proceedings of the National Academy of Sciences* **115**:E564–E573
55. Hie B., Zhong E. D., Berger B., Bryson B. 2021) **Learning the language of viral evolution and escape** *Science* **371**:284–288
56. Los Alamos National Laboratory 2023) **Hiv sequence database** <https://www.hiv.lanl.gov>
57. Roark R. S., et al. 2021) **Recapitulation of hiv-1 env-antibody coevolution in macaques leading to neutralization breadth** *Science* **371**:eabd2638
58. Benson D. A., et al. 2012) **Genbank** *Nucleic acids research* **41**:D36–D42
59. Los Alamos National Laboratory 2023) **Hivalign: Hiv sequence alignment tool** <https://www.hiv.lanl.gov/content/sequence/VIRALIGN/viralign.html>
60. Gao F., et al. 2014) **Cooperation of b cell lineages in induction of hiv-1-broadly neutralizing antibodies** *Cell* **158**:481–491
61. Kreer C., et al. 2023) **Probabilities of developing hiv-1 bnab sequence features in uninfected and chronically infected individuals** *Nature Communications* **14**:7137
62. Liao H.-X., et al. 2013) **Co-evolution of a broadly neutralizing hiv-1 antibody and founder virus** *Nature* **496**:469–476
63. Bonsignori M., et al. 2017) **Staged induction of hiv-1 glycan-dependent broadly neutralizing antibodies** *Science translational medicine* **9**:eaai7514
64. Ruxton G. D., Neuhäuser M. 2010) **Good practice in testing for an association in contingency tables** *Behavioral Ecology and Sociobiology* **64**:1505–1513
65. Ewens W. J. 2004) **Mathematical population genetics: theoretical introduction** Springer
66. Zanini F., et al. 2015) **Population genomics of inpatient hiv-1 evolution** *eLife* **4**:e11282
67. Kimura M. 1964) **Diffusion models in population genetics** *Journal of Applied Probability* **1**:177–232
68. Crow J. F. 2017) **An introduction to population genetics theory** Scientific Publishers

69. Sohail M. S., Louie R. H., McKay M. R., Barton J. P. 2021) **Mpl resolves genetic linkage in fitness inference from complex evolutionary histories** *Nature biotechnology* **39**:472–479
70. Shimagaki K., Barton J. P. 2023) **Bézier interpolation improves the inference of dynamical models from data** *Physical Review E* **107**:024116
71. Sohail M. S., Louie R. H., Hong Z., Barton J. P., McKay M. R. 2022) **Inferring epistasis from genetic time-series data** *Molecular biology and evolution* **39**:msac199
72. Shimagaki K. S., Barton J. P. 2024) **Efficient epistasis inference via higher-order covariance matrix factorization** *bioRxiv* :2024–10
73. Hong Z., Shimagaki K. S., Barton J. P. 2024) **popdms infers mutation effects from deep mutational scanning data** *Bioinformatics* **40**:btae499
74. Bauer A., et al. 2023) **Adaptation of a transmitted/founder simian-human immunodeficiency virus for enhanced replication in rhesus macaques** *PLoS Pathogens* **19**:e1011059
75. Bricault C. A., et al. 2019) **Hiv-1 neutralizing antibody signatures and application to epitope-targeted vaccine design** *Cell host & microbe* **25**:59–72
76. LaBranche C. C., et al. 2019) **Neutralization-guided design of hiv-1 envelope trimers with high affinity for the unmutated common ancestor of ch235 lineage cd4bs broadly neutralizing antibodies** *PLoS pathogens* **15**:e1008026
77. Saunders K. O., et al. 2019) **Targeted selection of hiv-specific antibody mutations by engineering b cell maturation** *Science* **366**:eaay7199
78. Henderson R., et al. 2023) **Structural basis for breadth development in the hiv-1 v3-glycan targeting dh270 antibody clonal lineage** *Nature Communications* **14**:2782
79. Saunders K. O., et al. 2022) **Stabilized hiv-1 envelope immunization induces neutralizing antibodies to the cd4bs and protects macaques against mucosal infection** *Science translational medicine* **14**:eabo5598

Author information

Kai S Shimagaki

Department of Computational and Systems Biology, University of Pittsburgh School of Medicine, Pittsburgh, United States, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, United States
ORCID iD: [0000-0001-7580-4781](https://orcid.org/0000-0001-7580-4781)

Rebecca M Lynch

Department of Microbiology, Immunology and Tropical Medicine, School of Medicine and Health Sciences, George Washington University, Washington, United States

John P Barton

Department of Computational and Systems Biology, University of Pittsburgh School of Medicine, Pittsburgh, United States, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, United States

ORCID iD: [0000-0003-1467-421X](https://orcid.org/0000-0003-1467-421X)

For correspondence: jpbarton@pitt.edu

Editors

Reviewing Editor

Anne-Florence Bitbol

Ecole Polytechnique Federale de Lausanne (EPFL), Lausanne, Switzerland

Senior Editor

Joshua Schiffer

Fred Hutchinson Cancer Research Center, Seattle, United States of America

Reviewer #1 (Public review):

Summary:

The present work studies the coevolution of HIV-1 and the immune response in clinical patient data. Using the Marginal Path Likelihood (MPL) framework, they infer selection coefficients for HIV mutations from time-series data of virus sequences as they evolve in a given patient.

Strengths:

The authors analyze data from two human patients, consisting of HIV population sequence samples at various points in time during the infection. They infer selection coefficients from the observed changes in sequence abundance using MPL. Most beneficial mutations appear in viral envelop proteins. The authors also analyze SHIV samples in rhesus macaques, and find selection coefficients that are compatible with those found in the corresponding human samples.

The manuscript is well-written and organized.

Weaknesses:

The MPL method used by the authors considers only additive effects of mutations, thus ignoring epistasis.

Although the evolution of broadly neutralizing antibodies (bnAbs) is a motivating question in the introduction and discussion sections (and the title), the relevance of the analysis and results to better understanding how bnAbs arise is not clear. The only result presented in direct connection to bnAbs is Figure 6.

Questions or suggestions for further discussion:

I list here a number of points for which I believe the paper would benefit if additional discussion/results were included.

The MPL method used by the authors considers only additive effects of mutations, thus ignoring epistasis. In Sohail et al (2022) MBE 39(10), p. msac199 (<https://doi.org/10.1093>

[/molbev/msac199](#)) an extension of MPL is developed allowing one to infer epistasis. Can the authors comment on why this was not attempted here?

I presume one possible reason is that epistasis inference requires considerably more computational effort (and more data). However, since the authors find most beneficial mutations occurring in Env, perhaps restricting the analysis to Env genes only (e.g. the trimer shown in Figure 2) can lead to tractable inference of epistasis within this segment (instead of the full genome).

Do the authors find correlations in the inferred selection coefficients of the two samples CH505 and CH848? I could not find any discussion of this in the manuscript. Only correlations between Humans and RM are discussed.

<https://doi.org/10.7554/eLife.105466.1.sa2>

Reviewer #2 (Public review):

Summary

This paper combines a biological topic of interest with the demonstration of important theoretical/methodological advances. Fitness inference is the foundation of the quantitative analysis of adapting systems. It is a hard and important problem and this paper highlights a compelling approach (MPL) first presented in (1) and refined in (2), roughly summarized in equation 12.

(1) Sohail, M. S., Louie, R. H., McKay, M. R. & Barton, J. P. Mpl resolves genetic linkage in fitness inference from complex evolutionary histories. *Nature biotechnology* 39, 472-479 (2021).

(2) Shimagaki, K. & Barton, J. P. Bézier interpolation improves the inference of dynamical models from data. *Physical Review E* 107, 024116 (2023).

The authors find that positive selection shapes the variable regions of env in shared patterns across two patient donors. The patterns of positive selection are interesting in and of themselves, they confirm the intuition that hyper-variation in env is the result of immune evasion rather than a broadly neutral landscape (flatness). They show that the immune evasion patterns due to CD8 T and naive B-cell selection are shared across patients. Furthermore, they suggest that a particular evolutionary history (larger flux to high fitness states) is associated with bNAb emergence. Mimicking this evolutionary pattern in vaccine design may help us elicit bNAbs in patients in the future.

There is a lot of information to be found in the full fitness landscape of env. The enormous strength of reversion-to-consensus in the patterns is a known pattern of HIV post-infection populations but they are nicely quantified here. Agreement between SHIV and HIV evolution is shown. They find selection is larger for autologous antibodies than the bNAbs themselves (perhaps bNAbs are just too small a component of the host response to drive the bulk of selection?), and that big fitness increases precede antibody breadth in rhesus macaques, suggesting that this fitness increase is the immune challenge required to draw forth a bNAb. This is all of high interest to HIV researchers.

Strength of evidence

One limitation is, of course, that the fitness model is constant in time when the immune challenge is variable and changing. This simplification may complicate some interpretations.

Equation 12 in the methods is really a beautiful tool because it is so simple, but accounts for linkage and can be solved precisely even in the presence of detailed mutational and selection

models. However, the reliance on incomplete observations of the frequency leads to complications that must be carefully (re)addressed here.

For instance, the consistent finding of strong selection in hypervariable regions is biologically intuitive but so striking, that I worry that it might be the result of a bias for selection in high entropy regions. Mutational and covariance terms in equation 12 might be underestimated, due to finite sampling effect in highly diverse populations. Sampling effects lead to zeros in $x(t)$ when actual frequency zeros might be rare at the population sizes of HIV viral loads and mutation rates. Both mutational flux and C underestimation will bias selection upward in eq. 12. The prior papers (1) and (2) seem to show robustness to finite sampling effects, but, again, more care needs to be shown that this robustness transfers to the amino acid inference under these conditions. That synonymous sites are rarely selected for in the nucleotide level is a good sign, and it may be a matter of simply fully explaining the amino-acid level model.

Uncertainty propagates to the later parts of the paper, eg. HIV and SIV shared patterns might be the result of shared biases in the method application. However, this worry does not extend to the apples-to-apples comparison of fitness trajectories across individuals (Figures 5 and 6) which I think are robust (for these sample sizes). The timing evidence is slightly weakened by the fact that bNAb detection is different from bNAb presence and the possibility that fitness increases occurred after the bNAbs appeared remains. Still, their conclusion is plausible and fits in with the other observations which form a coherent and compelling picture.

Overall this is a convincing paper, part of a larger admirable project of accurately inferring complete fitness landscapes.

<https://doi.org/10.7554/eLife.105466.1.sa1>

Reviewer #3 (Public review):

Summary:

Shimagaki et al. investigate the virus-antibody coevolutionary processes that drive the development of broadly neutralizing antibodies (bnAbs). The study's primary goal is to characterize the evolutionary dynamics of HIV-1 within hosts that accompany the emergence of bnAbs, with a particular focus on inferring the landscape of selective pressures shaping viral evolution. To assess the generality of these evolutionary patterns, the study extends its analysis to rhesus macaques (RMs) infected with simian-human immunodeficiency viruses (SHIV) incorporating HIV-1 Env proteins derived from two human individuals.

Strengths:

A key strength of the study is its rigorous assessment of the similarity in evolutionary trajectories between humans and macaques. This cross-species comparison is particularly compelling, as it quantitatively establishes a shared pattern of viral evolution using a sophisticated inference method. The finding that similar selective pressures operate in both species adds robustness to the study's conclusions and suggests broader biological relevance.

Weaknesses:

However, the study has some limitations. The most significant weakness is that the authors do not sufficiently discuss the implications of the observed similarities. While the identification of shared evolutionary patterns (e.g., Figure 5) is intriguing, the study would benefit from a more explicit discussion of what these findings mean for instance, in the context of HIV vaccine design, immunotherapy, or fundamental viral-host interactions. Even speculative interpretations could provide valuable insights into the broader significance of these results.

A secondary, albeit less critical, limitation is the placement of methodological details in the Supplementary Information. While it is understandable that the authors focus on results in the main text - especially since the methodology is not novel and has been previously described in earlier publications - some readers might benefit from a more thorough presentation of the method within the main paper.

Conclusions:

Overall, the study presents a compelling analysis of HIV-1 evolution and its parallels in SHIV-infected macaques. While the quantitative comparison between species is a notable contribution, a deeper discussion of its broader implications would strengthen the paper's impact.

<https://doi.org/10.7554/eLife.105466.1.sa0>