

Reviewed Preprint

v1 • March 12, 2025

Not revised

Reviewed Preprint

v2 • January 12, 2026

Revised by authors

✉ For correspondence:

samuel.j.sober@emory.edu

Competing interests:

No competing interests declared

Funding:

See [page 25](#)

Reviewing editor:

Juan Alvaro Gallego, Imperial College London, United Kingdom

© 2025, Thomas et al. This article is distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Motor unit mechanisms of speed control in mouse locomotion

Kyle Thomas¹, Rhuna Gibbs², Hugo Marques³, Megan R Carey³, Samuel J Sober² ✉

¹Graduate Program in Biomedical Engineering, Emory University and Georgia Institute of Technology, Atlanta, United States • ²Department of Biology, Emory University, Atlanta, United States • ³Champalimaud Neuroscience Programme, Champalimaud Centre for the Unknown, Lisbon, Portugal

eLife Assessment

This **valuable** study characterises the activity of motor units from two of the three anatomical subdivisions ("heads") of the triceps muscle while mice walked on a treadmill at various speeds. Altogether, this is the most thorough characterisation of motor unit activity in walking mice to date, providing **solid** evidence for probabilistic recruitment of motor units that differed between the two heads.

<https://doi.org/10.7554/eLife.105829.2.sa4>

Abstract

During locomotion, the coordinated activity of dozens of muscles shapes the kinematic features of each stride, including systematic changes in limb movement across walking speed. Motor units, each of which consists of a single motor neuron and the muscle fibers it innervates, contribute to the total activation of each muscle through their recruitment and firing rate when active. However, it remains unknown how the nervous system controls locomotor speed by changing the firing of individual motor units. To address this, we combined quantitative behavioral analysis of mouse locomotion with single motor unit recordings from the lateral and long heads of the triceps brachii, which drive monoarticular extension of the elbow and biarticular movements of the elbow and shoulder, respectively. In contrast to prior studies employing bulk EMG to examine muscle activity, our recordings revealed the diversity of spike patterning across motor units as well as systematic differences in motor unit activity across muscles and locomotor speeds. First, motor unit activity differed significantly across the lateral and long heads, suggesting differential control of these two closely apposed elbow extensor muscles. Second, we found that individual units were recruited probabilistically (during only a subset of strides), showing that the highly repeatable bulk EMG signals observed across strides in fact reflect stochastically varying subsets of individual motor units. Finally, although recruitment probability and firing rate both increased at faster walking speeds, increases in recruitment were proportionally larger than rate changes, and recruitment of individual units accompanied changes in limb kinematics. Together, these results reveal how the firing of individual motor units varies systematically across muscles and walking speeds to produce flexible locomotor behavior.

Introduction

Skilled behavior depends on the nervous system's precise control of muscle activity. Motor units, which consist of a single motor neuron and all of the muscle fibers it innervates, generate the forces behind movement through their firing patterns. In locomotion, proper neural coordination of motor units within and across muscles allows for the stereotyped yet rapidly adjustable movement used for each step (Akay et al., 2014 [DOI](#); Mayer & Akay, 2018 [DOI](#); N. P. Schumann et al., 2006 [DOI](#)). In principle, the total force output of a muscle is modulated by the number of recruited motor units and the firing rate of active units (Enoka & Duchateau, 2017 [DOI](#); Heckman & Enoka, 2012 [DOI](#)), with each newly-recruited unit increasing total muscle force by activating more muscle

fibers. The firing rate and inter-spike-interval (ISI) pattern of recruited units then shape force production in concert with the biomechanics of the musculoskeletal system (Sober et al., 2018; Sponberg et al., 2011). Although studies in primates, cats, and zebrafish have shown that both the number of active motor units and motor unit firing rates increase at faster locomotor speeds (Grimby, 1984; Hoffer et al., 1981, 1987; Menelaou & McLean, 2012), the extent to which speed-dependent changes in rate and recruitment vary across muscles and species is unknown.

Mice demonstrate both physiological and biomechanical differences from other vertebrates, potentially leading to unique coordination among their motor units. Compared to cats, for example, mice have highly excitable motor units (Manuel et al., 2019; Manuel & Heckman, 2011) with muscle fibers heavily biased towards fast-twitch fibers (Burkholder et al., 1994; Mathewson et al., 2012), leading to rapid force production. Mice also locomote with greater stride frequency than larger species in order to achieve comparable speeds, requiring faster muscle activation and deactivation (Heglund & Taylor, 1988; Machado et al., 2015). The capability and need for faster force generation during dynamic behavior could implicate motor unit recruitment as a primary mechanism for modulating force output in mice (Manuel & Heckman, 2011; Dideriksen et al., 2020).

To quantify the organization of motor unit firing patterns during locomotion, we recorded mouse motor unit activity from the long head and lateral head of the triceps brachii during treadmill walking at various speeds. Both muscles extend the elbow while the long head also extends, rotates, and abducts the shoulder (Tata Ramalingasetty et al., 2021). Although bulk EMG recordings have shown that the triceps brachii is active in every step during quadrupedal locomotion (English, 1978; N. Schumann, 2002; Kirk et al., 2024), it is unknown how individual motor units are coordinated to generate this rhythmic pattern and whether motor pools from closely apposed muscles exhibit the same coordination. Using Myomatrix electrodes (Chung et al., 2023; Gilmer et al., 2024; Kirk et al., 2024) to record populations of individual motor units during locomotion, we found that units were recruited probabilistically across strides. When active, units fired in distinct locomotor phases with systematic differences in spike patterns across the long and lateral heads. At faster walking speeds, motor units increased both their recruitment probabilities and (to a lesser extent) their firing rates. Moreover, motor unit recruitment was correlated with variations in limb kinematics both within and across locomotor speeds, with recruitment of long head and lateral head units associated with different changes in limb movement. Overall, our results reveal the systematic changes in motor unit firing that regulate locomotor speed.

Results

We collected kinematic and high-resolution electromyographic (EMG) data from six mice walking on a transparent treadmill that provided simultaneous lateral and ventral views of the animal (Darmohray et al., 2019). Using DeepLabCut (Mathis et al., 2018) to track body movements during locomotion (Figure 1A), we identified the stance phase of the right forelimb, defined as the period between footstrike and liftoff of the right forepaw (Figure 1B). Additionally, computing the internal angle of the elbow joint revealed that the elbow was minimally extended approximately 50 milliseconds before the footstrike (blue squares, Figure 1C). Electrode arrays (32-electrode Myomatrix array model RF-4×8-BHS-5) were implanted in forelimb muscles (note that Figure 1D shows the EMG signal from only one of the 16 bipolar recording channels), and the resulting data were used to identify the spike times of individual motor units in the triceps brachii long and lateral heads (Table 1, Figure 1E) as described previously (Chung et al., 2023). To best capture the spike pattern given that some units begin firing prior to footstrike, we defined a stride cycle as the period between minimum elbow angles rather than consecutive footstrikes. Kinematic analysis of locomotor data at different walking speeds revealed systematic variation in the temporal (Figure 1F) and spatial (Figure 1G,H) components of limb movement, consistent with prior reports (Akay et al., 2006; Bellardita & Kiehn, 2015; Machado et al., 2015; Mendes et al., 2015).




Figure 1. High-resolution muscle recording during mouse locomotion.

(A) (Left) Anatomical landmarks (shoulder, elbow, wrist, paw) and kinematic features (elbow angle θ , paw position x) tracked via high-resolution video during treadmill walking (Darmohray et al., 2019; Mathis et al., 2018). (Right) Position of anatomical landmarks during two stride cycles with limb position captured every 15 ms. (B) Position of the right forepaw (x) relative to body center. Thick lines represent the stance phase when the paw is in contact with the floor of the treadmill. (C) Interior elbow angle (θ) during locomotion. Troughs of this measure, denoting minimum extension (blue squares), were used to define the spike window for each stride. (D) Representative single channel of electromyographic (EMG) activity recorded from the long head of the right triceps. (E) Three motor units recorded from the long head identified from the above EMG trace (see Methods). Note that the bottom-most unit is active in only a subset of stride cycles. (F) Relationship between stride duration and walking speed for all strides in an example mouse. Each dot represents a stride, with shading indicating the speed quartile within which the stride falls (see Methods). (G,H) Right forepaw position x (G) and elbow angle (H) within the walking speed quartiles. Both (G) and (H) are normalized to total stride duration beginning and ending with elbow minimum extension (blue squares) and show mean ($\pm$ SE).

Mouse	Implanted muscle (# motor units identified)
A	2 threads right lateral head, triceps brachii (3) 2 threads right long head, triceps brachii (1)
B	2 threads right lateral head, triceps brachii (0) 2 threads right long head, triceps brachii (3)
C	2 threads right lateral head, triceps brachii (0) 2 threads right long head, triceps brachii (6)
D	2 threads right lateral head, triceps brachii (4) 2 threads right biceps brachii (n/a)
E	2 threads left biceps brachii (n/a) 2 threads right long head, triceps brachii (7)
F	4 threads right lateral head triceps brachii (9)

Table 1. Motor units identified per muscle in each experimental mouse.

Each thread consisted of 8 electrode contacts used to record bipolar EMG, and numbers in parentheses indicate the number of motor units isolated using the spike sorting algorithm described in Methods. Data from biceps muscle implantation in two mice were not spike sorted and are not included in this report.

Motor units are probabilistically recruited across strides

Despite the triceps muscles as a whole being reliably activated on every step (English, 1978 [↗](#); N. Schumann, 2002 [↗](#); Kirk et al., 2024 [↗](#)), the majority of individual motor units in both the long head and lateral head were active only in a subset of strides during locomotion. Motor units in both muscles exhibited this pattern of probabilistic recruitment (defined as a unit's firing on only a fraction of strides), but with differing distributions of firing properties across the long and lateral heads (Figure 2 [↗](#)). For each motor unit, we measured the probability of a unit being recruited as the percentage of strides with at least one spike. Units demonstrated a variety of firing patterns, with some units producing 0 spikes more frequently than any non-zero spike count (Figure 2A, B [↗](#)), and units that were less likely to be recruited also had lower average spike counts (Figure –figure supplement 1 [↗](#)). A subset of units, primarily in the long head, were recruited in under 50% of the total strides and with lower spike counts (Figure 2C [↗](#)). This distribution of recruitment probabilities might reflect a functionally different subpopulation of units. However, the distribution of recruitment probabilities were not found to be significantly multimodal ($p > 0.05$ in both cases, Hartigan's dip test; Hartigan, 1985 [↗](#)). However, Hartigan's test and similar statistical methods have poor statistical power for the small sample sizes ($n = 17$ and 16 for long and lateral heads, respectively) considered here, so the failure to achieve statistical significance might reflect either the absence of a true difference or a lack of statistical resolution.

Motor unit firing patterns in the long and lateral heads of the triceps

Motor units within each muscle fired at distinct phases of the stride cycle. Units in the long head typically became active near the time of footstrike, with approximately half of the units reliably recruited prior to footstrike (Figure 3A, B [↗](#)). In contrast, units in the lateral head began spiking after the long head was already active and remained active until just prior to liftoff (Figure 3B, C [↗](#)). Furthermore, units in the long head reached their stride-dependent peak rates before the lateral head ($p < 0.01$, two-sample k-s test). These findings demonstrate that despite the synergistic (extensor) function of the long and lateral heads of the triceps at the elbow, the motor pool for the long head becomes active roughly 100 ms before the motor pool supplying the lateral head during locomotion (Figure 3C [↗](#)). This timing difference suggests distinct patterns of synaptic input onto motor neurons innervating the lateral and long heads. In contrast to the timing differences described above, motor units in the lateral and long heads displayed similar burst durations (Figure 3B [↗](#)) and peak firing rates (Figure 3D [↗](#)).

The evolution of spike patterns within each stride differed between motor unit populations in the long and lateral heads. In both muscles, motor units with longer burst durations reached higher peak firing rates (Figure 4A [↗](#)). However, the slope of this relationship was significantly higher for lateral head units ($p < 0.05$, permutation test). We also observed muscle-dependent differences in motor unit patterning when examining the inter-spike intervals (ISIs) between the first three spikes in each stride cycle. Motor units in both muscles began firing with ISIs typically below 12 ms (Figure 4B [↗](#)). Furthermore, the second ISI was generally shorter, indicating that firing rate increased throughout the first three spikes fired in the stride cycle. However, the population of motor units in the long head had a larger magnitude in the ratio ISI_1/ISI_2 (Figure 4B [↗](#), $p < 0.01$, two-sample k-s test). Together with the differences in burst timing shown in Figure 3B [↗](#), these results again suggest that the motor pools for the lateral and long heads of the triceps receive distinct patterns of synaptic input, although differences in the intrinsic physiological properties of motor neurons innervating the two muscles might also play an important role.

Motor unit mechanisms of speed control

Adjusting walking speed requires changes in the firing patterns of individual motor units, which could include speed-dependent changes in units' probability of recruitment and/or changes in firing rate. To investigate the changes in motor unit firing underlying locomotor speed control, we quantified how both recruitment probability and firing rate change across the four quartiles of




Figure 2. Motor unit spike count distributions.

(A) Example motor unit from the long head of the triceps muscle fired zero spikes on 52% of strides, but on the other 48% of strides fired 1-14 spikes. **(B)** Example motor unit from the triceps lateral head fired zero spikes in 17% of strides but 1-19 spikes on the other 83% of strides. **(C)** Percentage of strides with at least one spike (probability of recruitment) for all recorded motor units in the long (purple) and lateral (green) heads of the triceps. Symbols denote different animals and each point reflects an individual motor unit. Asterisks highlight the units shown in (A,B).




Figure 3. Motor unit firing patterns within and across muscles.

(A) Example stride with three units from the long head. (B) Mean phase ($\pm$ SE) of motor unit burst activity within each stride duration across all strides. Black dots on each bar show the mean phase of the unit's peak firing rate. Starred points refer to the examples in A. (C) Left: Mean time ($\pm$ SE) between the first spike of a unit's spike train and the right forepaw footstrike. Positive values denote the spike happening after the footstrike. Right: Mean time ($\pm$ SE) between the last spike of a unit and the liftoff. Light traces denote values for individual motor units, the heavy trace shows the mean ($\pm$ SE) across all units within a muscle. (D) Mean peak firing rate ($\pm$ SE) of each unit. Note that these measurements only include strides in which the given unit was recruited.




Figure 4. Motor unit spike patterns evolve differently in the long and lateral heads.

(A) Relationships between active duration and peak firing rate across motor pools. Type 2 regression slopes were significantly different between the lateral and long heads ($p < 0.05$, permutation test). **(B)** Motor unit inter-spike intervals (ISIs) across the first three spikes in motor unit bursts. Each data point shows the mean of the first ISI and the ratio between the first and second ISIs for a single unit. Note that by definition only strides with at least three spikes could be used for the analysis shown in panel (B). Type 2 regression slopes were not significantly different between data from the lateral and long heads ($p > 0.05$, permutation test). Data shown here are grouped across all mice; Figure 4-figure supplement 1 shows how these data are distributed across animals.

locomotor speed shown in Figure 1F-H. Motor units from the long and lateral heads of the triceps (Figure 5A,B, purple and green traces, respectively) displayed significant increases in recruitment probability as locomotor speeds increased. Figure 5C shows each motor unit's difference in recruitment probability between the slowest and fastest locomotor speed quartiles. This increase was statistically significant in 31/33 motor units in our study ($p < 0.05$, permutation test) when considered individually, and was also significant when the probabilities of all motor units were analyzed as a group ($p < 0.01$, Wilcoxon signed-rank test). Robust increases in recruitment probability across the four speed quartiles were therefore the norm in our dataset.

Quantitative analysis of motor unit activity also revealed significant speed-dependent changes in firing rate, although these were proportionally smaller than the increases in recruitment probability. Motor units in both the long and lateral heads of the triceps (Figure 5D,E, purple and green traces, respectively) often had either marginal increases or no difference in peak firing rate at faster speeds. Across all motor units in our dataset in the slowest and fastest speed quartiles (Figure 5F), we observed significant increases in peak firing rate in 22/33 individual motor units in our study ($p < 0.05$, permutation test), and also a significant speed-dependent increase in peak rate when considering all motor units together ($p < 0.01$, Wilcoxon signed-rank test). Speed-dependent increases in peak firing rate were therefore also present in our dataset, although in a smaller fraction of motor units (22/33) than changes in recruitment probability (31/33). Furthermore, the mean ($\pm$ SE) magnitude of speed-dependent increases was smaller for spike rates (mean $\text{rate}_{\text{fast}}/\text{rate}_{\text{slow}}$ of $111\% \pm 20\%$ across all motor units) than for recruitment probabilities (mean $p(\text{recruitment})_{\text{fast}}/p(\text{recruitment})_{\text{slow}}$ of $179\% \pm 3\%$ across all motor units). While fractional changes in rate and recruitment probability are not readily comparable given their different upper limits, these findings could suggest that while both recruitment and peak rate change across speed quartiles, increased recruitment probability may play a larger role in driving changes in locomotor speed.

Kinematic contributions of motor unit recruitment

We next examined whether the probabilistic recruitment of individual motor units in the triceps – an elbow extensor muscle – was correlated with stride-by-stride variations in elbow angle kinematics. To do so, we compared elbow extension ($\Delta\theta$; Figure 6A) on strides in which each individual motor unit did or did not fire at least one spike. When kinematic data are combined across all speed quartiles (Figure 6B), we found that recruitment of lateral head motor units (green symbols) is associated with greater elbow extension, whereas recruitment of long head units (purple symbols) is correlated with smaller extensions ($p < 0.001$, Wilcoxon signed-rank tests). These correlations might reflect both an influence of motor unit recruitment on limb kinematics as well as different biomechanical roles for the long and lateral heads.

Since both limb kinematics (Figure 1G,H) and recruitment probability (Figure 5) are significantly correlated with locomotor speed, the observed correlations between unit recruitment and elbow angle across all speeds (Figure 6B) does not necessarily reveal the direct influence of unit firing on limb kinematics. We therefore controlled for speed by repeating the analysis shown in Figure 6B for strides within each speed quartile. Strikingly, the correlations between motor unit recruitment and elbow angle persisted in this alternative analysis (Figure 6C; $p < 0.001$, Wilcoxon signed-rank tests), suggesting that the recruitment of individual motor units in the lateral and long heads might have significant effects on elbow angle in strides of similar speed (see Discussion). We repeated these analyses using the elbow angular velocity rather than just the angle to further identify how firing patterns related to behavior. Motor units in the lateral head had a similar effect with larger velocities correlating with motor unit recruitment across all speeds (Figure 6D,E; $p < 0.001$).

Probabilistic recruitment is correlated across motor units

Our results show that the recruitment of individual motor units is probabilistic even within a single speed quartile (Figure 5A-C) and predicts body movements (Figure 6), raising the question of whether the recruitment of individual motor units are correlated or independent.




Figure 5. Motor units alter firing rate and recruitment across walking speeds.

(A) Light traces show median of recruitment probability for individual long head motor units while the heavy trace shows mean ($\pm$ SE) across all long head motor units. (B) Recruitment probability for lateral head motor units, same plotting conventions as in (A). (C) Difference in recruitment probabilities between slowest and fastest speed quartiles for all motor units. (D) Light traces show median of peak firing rate for individual long head motor units while the heavy trace shows mean ($\pm$ SE) across all long head motor units. (E) Peak firing rates for lateral head motor units, same plotting conventions as in (D). (F) Difference in peak firing rates between slowest and fastest speed quartiles for all motor units. Across all motor units, both recruitment probabilities (C) and firing rates (F) were significantly higher at the fastest quartile than at the slowest quartile ($p < 0.01$, Wilcoxon signed-rank tests). Speed-dependent changes in recruitment vs. speed-dependent changes in firing rate for individual motor units and experimental animals are shown in Figure –figure supplement 1 and Figure –figure supplement 2, respectively.




Figure 6. Motor unit recruitment correlates with muscle-specific kinematic differences.

(A) We calculated the ranges of elbow angle ($\Delta\theta$) and elbow velocity ($\Delta\omega$) observed on strides in which each motor unit did or did not fire at least one spike (purple tick marks below EMG trace). **(B, D)** Each point represents the mean $\Delta\theta$ (B) or $\Delta\omega$ (D) observed on strides in which a single motor unit fires zero spikes (horizontal axis) vs. when the motor unit fires at least one spike (vertical axis). Note that in these panels, data for each motor unit were combined across all locomotor speeds. **(C, E)** Same analyses as before, except each motor unit contributes up to four data points, one for each of the four locomotor speed quartiles in which sufficient data were available (at least 30 strides existed in both the spiking and non-spiking conditions within a given quartile). Legends in each panel show statistical significance for a difference in kinematics tested on the motor units within each muscle (Wilcoxon signed-rank tests). Note that most of the muscle-specific differences shown in (C, E) were also present when each of the four quartiles (Figure 6-figure supplement 1) or experimental animals (Figure 6-figure supplement 2) were examined individually.

Correlated recruitment might reflect shared input onto the population of motor units innervating the muscle (De Luca, 1985 [↗](#); De Luca & Erim, 1994 [↗](#); Farina et al., 2014 [↗](#)). For example, two motor units, each with low recruitment probabilities, may still fire during the same set of strides. To assess the independence of motor unit recruitment across the recorded population, we compared each unit's empirical recruitment probability across all strides to its conditional recruitment probability during strides in which another motor unit from the same muscle was recruited (Figure 7 [↗](#)). Doing this for all motor unit pairs revealed that motor units in both muscles were biased towards greater recruitment when additional units were active ($p < 0.001$, Wilcoxon signed-rank tests for both the lateral and long heads of triceps). This finding suggests that probabilistic recruitment reflects common synaptic inputs that covary together across locomotor strides.

Discussion

Our results quantify the heterogeneity of individual motor units' firing patterns across muscles and walking speeds. Motor units were probabilistically recruited on a stride-by-stride basis with peak firing rates between 60-185 Hz when active. Motor units in the long head were recruited before the lateral head and spiking patterns evolved within each stride differently across the two muscles. Motor units in both muscles demonstrated increases in their recruitment probability and firing rate at faster walking speeds. Furthermore, motor unit recruitment was also correlated with differences in limb kinematics for strides of similar speed. As discussed below, firing patterns from motor units in the long and lateral heads likely reflect the functional and anatomical role of these two muscles, highlighting the need for high-resolution quantification of motor unit firing patterns during behavior.

Differences in motor unit activity patterns across two elbow extensors

Motor unit spike patterns differed systematically between the long and lateral heads of the triceps brachii. Motor units in the long head were consistently recruited earlier than units in the lateral head (Figure 3B,C [↗](#)). The large differences in burst timing and spike patterning across the muscle heads suggest that the motor pools for each muscle receive distinct inputs. However, differences in the intrinsic physiological properties of motor units and neuromodulatory inputs across motor pools might also make substantial contributions to the structure of motor unit spike patterns (Martínez-Silva et al., 2018 [↗](#); Miles & Sillar, 2011 [↗](#)).

The observed order of muscle activation matches past reports of bulk muscle activity in these two muscles across other quadrupedal species (Carroll & Biewener, 2009 [↗](#); Drew et al., 2008 [↗](#); Livingston & Nichols, 2014 [↗](#); Scholle et al., 2001 [↗](#)) and may reflect the biomechanical functions of each muscle. Whereas the lateral head is a monoarticular elbow extensor, the long head is biarticular, both extending the elbow along with extending and rotating the shoulder (Tata Ramalingasetty et al., 2021 [↗](#)). Although we did not measure ground reaction forces in this study, prior reports indicate that the vertical ground reaction force on the mouse forepaw reaches two peaks during locomotion (Schmitt et al., 2010 [↗](#)). The first peak, which happens soon after the footstrike, has a lower magnitude than the second peak, which occurs closer to liftoff (Clarke et al., 2001 [↗](#); Schmitt et al., 2010 [↗](#)). Studies in both rats (Sarver et al., 2010 [↗](#)) and cats (Corbee et al., 2014 [↗](#)) have demonstrated that horizontal ground reaction forces in both the medio-lateral and cranio-caudal directions are also greatest soon after footstrike, with more force variability than the vertical reaction force. Since units in the long head are most active following footstrike, this suggests that activity in the long head might be related to stabilizing the limb within each step. Our finding that recruitment of long head motor units (purple symbols, Figure 6 [↗](#)) accompanied smaller elbow extensions might therefore reflect a more complex biomechanical role for the long head, potentially relating to shoulder rotation (which was not measured in this study). This interpretation is consistent with past findings that biarticular muscles are power distributors, stabilizing the joint across multiple dimensions, while monoarticular muscles are power generators (Ryan & Gregor, 1992 [↗](#); Van Ingen Schenau et al., 1992 [↗](#), 1994). Along these lines, the




Figure 7. Motor unit recruitment probability is greater when other motor units within the muscle are recruited.

Each point reflects a motor unit's empirically measured recruitment probability, $p(MU_A)$, across all strides compared to strides when a simultaneously-recorded motor unit from the same muscle was recruited, $p(MU_A | MU_B)$. Motor units in each muscle had significantly higher recruitment when another unit was recruited in the same stride ($p < 0.001$, Wilcoxon signed-rank tests).

observed timing of lateral head motor unit activity just prior to liftoff (Figure 3B) might therefore reflect the lateral head's role of providing propulsion prior to swing, consistent with our finding that recruitment of motor units in the lateral head is correlated with both larger elbow extension and more rapid changes in angle (Figure 6).

Motor units in the lateral and long heads also differed with respect to their recruitment probabilities, with a substantial population of units in the long head (but not the lateral head) with probability of recruitment less than 50% (Figure 2C). This difference may reflect different functions of muscle fibers in different subcompartments of biarticular muscles. Prior work has established that different regions within a biarticular muscle can have different contributions across the two joints (Chanaud et al., 1991; English & Weeks, 1987; Watanabe et al., 2021). For example, different regions of the cat biceps femoris are out of phase with each other during walking, with the anterior compartment active during stance as a hip extensor and the posterior compartment active during swing as a knee flexor (Chanaud et al., 1991; English & Weeks, 1987). Additionally, the posterior compartment was only active at faster speeds. In our mouse data, functional compartments within the biarticular long head might similarly explain the differently recruited populations for motor units (Figure 2C). However, to our knowledge, no studies have investigated anatomical or functional subdivisions across subregions of the triceps long head in the mouse. Nevertheless, the group of less-frequently recruited units might contribute more to forelimb joint stability during a small number of strides affected by rare and unexpected external perturbations, whereas the other long head motor units might be recruited in a greater fraction of strides to support the weight of the body. Further examination of the anatomical microstructure of the long head, including precise characterization of the attachment points to the bone (DeWolf et al., 2024; Gilmer et al., 2024), are necessary to answer these questions.

The varied composition of fiber types in the long and lateral heads may also explain the different firing patterns across muscles. Although both muscles are heavily biased towards the fastest myosin type (type 2B), the long head has a broader composition, including a small percentage of slower isoforms as well (type 1 and 2A) (Mathewson et al., 2012). Type 2B isoforms are related to fast-twitch, fatigable units (FF) while type 1 compose slow-twitch units (S) and type 2A are intermediate between fast and slow (Bączyk et al., 2022; Schiaffino & Reggiani, 2011). While we were unable to directly quantify the unit type, the majority of units observed, particularly within the lateral head, are likely FF units given these prior histological findings. Thus, while we did not explicitly measure muscle fatigue after our recordings (up to 30 minutes of walking at 12.5-27.5 cm/sec, see Methods), it is possible that fatigue might have contributed to the observed probabilistic recruitment of later-recruited units (Martínez-Silva et al., 2018). Similarly, motor units that were recruited in nearly every stride with 10 or more spikes per stride (Figure –figure supplement 1) could represent a population with slower isoforms given their resistance to fatigue. The most prominent example of this came from the single unit in the long head that fired for over 90% of the stride phase in every stride (Figure 3B, top).

Firing rates in mouse locomotion compared to other species

The range of firing rates we observed in mice are higher than those typically observed in larger species, likely reflecting the distinctive physiology of mouse motor neurons. Motor units in the lateral and long heads of the triceps exhibited a large and overlapping range of peak firing rates ranging from 50-175 Hz (Figure 3D), in agreement with prior reports of motor unit firing rates from mouse forelimb (Kirk et al., 2024) and hindlimb (Hadzipasic et al., 2016) during locomotion. In rat hindlimb muscles during walking, motor units had mean instantaneous firing rates between 45-109 Hz (Gorassini et al., 2000). During locomotion in cats, motor units recorded from the toe and hindlimb had firing rates between 15-50 Hz (Hoffer et al., 1987; Zajac & Young, 1980). Human motor units in the short extensors of the toe fire at even lower rates (10-25 Hz) during walking (Grimby, 1984). Compared to these larger species, mice likely reach higher rates through the physiological properties of their motor neurons such as afterhyperpolarization (AHP), which influences how rapidly a neuron returns to baseline voltage after firing a spike. Although

AHP durations vary across unit types, AHP durations in mice are approximately two and three times shorter than those in cats and humans respectively (Manuel et al., 2009 [DOI](#), 2019 [DOI](#); Meehan et al., 2010 [DOI](#)). Additionally, persistent inward currents (PICs), which amplify excitatory synaptic inputs (Binder et al., 2020 [DOI](#); Heckman et al., 2005 [DOI](#)), might lead to disproportionately large gain in mouse motor neurons compared to other species (Huh et al., 2017 [DOI](#); Manuel et al., 2019 [DOI](#)). Consequently, even mice performing quiet standing have motor unit firing rates reaching up to 68 Hz (Ritter et al., 2014 [DOI](#)). Our findings (Figure 4 [DOI](#)) highlight that even with the relatively high firing rates observed in mice, there are still significant changes in firing rate and recruitment probability across the spikes within bursts (Figure 4B [DOI](#)) and across locomotor speeds (Figure 5F [DOI](#)). Future studies should more carefully examine how these rapidly changing spiking patterns derive from both the statistics of synaptic inputs and intrinsic properties of motor neurons (Manuel & Heckman, 2011 [DOI](#); Petersen & Berg, 2016 [DOI](#); Berg, 2017 [DOI](#)).

Walking speed modulation of firing rate and recruitment

To investigate the neuromuscular control of locomotor speed, we quantified speed-dependent changes in both motor unit recruitment and firing rate. We found that the majority of units were recruited more often and with larger firing rates at faster speeds (Figure 5 [DOI](#), Figure –figure supplement 1 [DOI](#)). This pattern may reflect speed-dependent differences in the common input received by populations of motor neurons with varying spiking thresholds (Henneman et al., 1965 [DOI](#)). In the case of mouse locomotion, faster speeds might reflect a larger common input, increasing the recruitment probability as more neurons, particularly those that are larger and generate more force, exceed threshold for action potentials (Farina et al., 2014 [DOI](#)). This would explain our finding that motor units are more likely to be recruited during strides in which additional motor units are also recruited (Figure 7 [DOI](#)). Although this seemed consistent for each pair of motor units we measured, this was not necessarily the case given that past studies have identified motor unit substitution rather than co-activation (Westgaard & De Luca, 1999 [DOI](#); Manning et al., 2010 [DOI](#)), potentially serving to reduce fatigue across the motor pool.

Importantly, our work only examines a subset of the movement speeds and gait patterns that mice produce. It therefore remains to be determined how rate and recruitment are reshaped as mice increase their speed up to 100 cm/s and alter coordination patterns across their limbs (trotting, bounding, etc.) (Herbin et al., 2006 [DOI](#), 2007 [DOI](#); Bellardita & Kiehn, 2015 [DOI](#); Gonçalves et al., 2022 [DOI](#)). Since a majority of observed motor units, particularly in the lateral head, were already reliably recruited at the fastest speed quartile (roughly 30-40 cm/s), further speed increases might rely on either more firing rate modulation from these active units or from recruitment of more of the motor pool. Moreover, adjustments to kinematic and kinetic strategy across speeds could result from more global changes in motor unit coordination. For example, studies in drosophila (Azevedo et al., 2020 [DOI](#)) and zebrafish (Kishore et al., 2014 [DOI](#)) have demonstrated preferential recruitment of faster motor unit subtypes during rapid movements. Future studies in mice can therefore examine faster gaits to compare how different species achieve their most rapid forms of locomotion.

Considering the force production of motor units is essential to connect our observations of firing patterns to behavioral outputs. In anesthetized mice, intracellular current injections into individual motor neurons revealed that fast motor units from the triceps surae (gastrocnemius and soleus muscles of the hindlimb) reached near tetanic force at firing rates between 60-80 Hz while slow motor units reached near tetanic forces between 30-40 Hz (Manuel & Heckman, 2011 [DOI](#)). Furthermore, motor units rapidly reached these rates once active. Despite being recorded from different muscles than the ones we examined, these earlier results are relevant to our findings given that the long and lateral heads of the triceps brachii are (similarly to the gastrocnemius) biased towards fast-twitch muscle fibers (Augusto et al., 2004 [DOI](#); Burkholder et al., 1994 [DOI](#)). Since the motor units recorded in our study had firing rates at or above the aforementioned rates immediately upon recruitment within a stride (Figure 4B [DOI](#)), it could be that each of the units identified in this study generated near-maximal force whenever active. If units are recruited with near maximal force even at slow walking speeds, generating the additional

forces needed for fast walking likely comes from recruitment of additional units. Future studies might answer this question by quantifying the force-production properties of triceps motor units during behavior, including the rapid changes in the muscle length and velocity that take place during locomotion (Edman, 1979 [↗](#); Gittings et al., 2012 [↗](#); Ting & Chiel, 2017 [↗](#)).

Although strong correlations were observed between motor unit recruitment and limb kinematics during locomotion (Figure 6 [↗](#), Figure –figure supplement 1 [↗](#)), it remains unclear whether such correlations actually reflect the causal contributions that those units make to limb movement. To resolve this ambiguity, future studies could use electrical or optical perturbations of muscle contraction levels (Kim et al., 2024 [↗](#); Lu et al., 2024 [↗](#); Srivastava et al., 2015 [↗](#), 2017 [↗](#)) to test directly how motor unit firing patterns shape locomotor movements. The short-latency effects of patterned motor unit stimulation (Srivastava et al., 2017 [↗](#)) could then reveal the sensitivity of behavior to changes in muscle spiking and the extent to which the same behaviors can be performed with many different motor commands.

Methods

Surgical implantation

All procedures described below were approved by the Emory University Institutional Animal Care and Use Committee at Emory University (IACUC protocol #201700359). Mice were anesthetized with isoflurane to implant the Myomatrix arrays. Incisions were made in the skin above the skull and above the target muscle. Forceps were used to pull the Myomatrix array through these holes so that the body of the array was entirely subcutaneous, with the Omnetics connector sitting on the skull and the array threads near the muscles. The surface of the skull was lightly scored with a scalpel and dental cement (Metabond Quick Adhesive Cement) was applied generously to fix the Omnetics connector in place and seal the opening. Myomatrix threads were then sutured (8-0 non-absorbable suture from AROSurgical) into the target muscles. Using the four threads of the customizable Myomatrix array (RF-4×8-BHS-5), we implanted a combination of muscles in each mouse, sometimes placing multiple threads within the same muscle. Threads were implanted in the triceps brachii long head and/or the triceps brachii lateral head (Table 1 [↗](#)) and confirmed through visual inspection. We did not implant in the third (medial) head of the triceps given that it would have required an additional incision, posing more risk of surgical complications. Some mice also had threads simultaneously implanted in their ipsilateral or contralateral biceps brachii, although due to limited sample size we do not present biceps EMG data in this report. Lastly, 6-0 suture was used to close the incision. Surgeries typically took under three hours and animals were mobile shortly after removal from isoflurane.

Behavioral methods and data collection

The treadmill used in this task had a transparent belt and base as described previously (Darmohray et al., 2019 [↗](#); Machado et al., 2015 [↗](#)). A 45° angled mirror below the base allowed monitoring of side and bottom views from a single camera (FLIR Grasshopper High Performance USB 3.0 Monochrome Camera) at 330 frames per second. Separate motors controlled the left and right belts, but both were run at the same speed for every experiment. The treadmill, placed within a behavior box, was dark, with the only source of light coming from infrared light.

Experiments were conducted the day following the implant surgery up to five days post-surgery. Data presented in this study came from the first day of recording, in which signal quality tended to be highest (Chung et al., 2023 [↗](#)). In each experimental session, mice were first briefly placed under anesthesia using isoflurane to attach a lightweight (1g) digitizing headstage (Intan RHD #C3313 16-Channel Bipolar-Input Recording Headstage) to the Omnetics connector on their skulls. Each recording session lasted approximately 45 minutes in total, and we waited at least 10 minutes after removal from isoflurane to ensure all animals were fully awake before recording began.

For five of six mice, we attempted to record 31 trials - each trial consisted of a single minute continuous running on the treadmill. The first three trials were at 10 cm/s, while the following trials were arranged in seven blocks of four trials each. Each block contained a trial at 12.5 cm/s, 17.5 cm/s, 22.5 cm/s, and 27.5 cm/s in a pseudo-random order presented identically across mice. This pseudo-random order of speeds as opposed to a strict ramping order ensured that we collected data across the full range while reducing potential effects of fatigue. For mice that became uncooperative before completing all trials, we ended the experiment early. Of these five mice, three mice completed all 31 trials, one completed 30, and the last completed 23 trials. For the sixth mouse, we again began with three trials at 10 cm/s, but only increased the speed in 2.5 cm/s increments for either two or three trials each up to a total of 14 trials. Mice were trained on their given running paradigm and habituated to the treadmill setup twice on the day before surgery. Each trial was initiated using custom Bonsai software (Lopes et al., 2015) and Arduino components to synchronize neural recordings with the camera and motor output.

We used DeepLabCut (Mathis et al., 2018) to track body parts of the mouse during locomotion (Figure 1A). We excluded points tracked with less than 90% confidence from DLC and interpolated those points from adjacent high-confidence points. The right elbow angle was estimated using markers from the shoulder, elbow, and wrist. We defined strides using the trough-to-trough minimums of the elbow angle, which occurred approximately 50-100 ms before the footstrike of the right paw. Each stride was also required to contain footstrike and liftoff of the right forepaw during forward movement. From here, we excluded strides with stride durations, stance durations, swing durations, body velocities, or body accelerations outside their respective 95% confidence interval. As a result, we kept about 80% of strides for each animal. Five of six mice had between 2600-3600 total strides, and the remaining mouse, which was run at the slower speed range, had just over 1000 strides included in analyses.

Electromyography (EMG)

Bipolar signals from adjacent contact pairs on the Myomatrix array were extracted at 30 kHz (Intan RHD #C3100 Recording Controller) and bandpassed between 300-7500 Hz. Using up to 16 channels of high-resolution EMG from the Myomatrix arrays, motor units were identified using Kilosort 2.5, an open-source multi-channel spike sorting algorithm (Pachitariu et al., 2016). We slightly adjusted the algorithm to better fit the assumptions behind motor unit activity, including the removal of spatial decay across channels. A full description of adjustments has been previously reported (Chung et al., 2023). A total of 33 units were identified across all animals. Each unit's isolation was verified by confirming that no more than 2% of inter-spike intervals violated a 1 ms refractory limit. Additionally, we manually reviewed cross-correlograms to ensure that each waveform was only reported as a single motor unit.

We further validated spike sorting by quantifying the stability of each unit's waveform across time (Figure -figure supplement 1). First, we calculated the median waveform of each unit across every trial to capture long-term stability of motor unit waveforms. Additionally, we calculated the median waveform through the stride binned in 50 ms increments using spiking from a single trial. This second metric captures the stability of our spike sorting during the rapid changes in joint angles that occur during the burst of an individual motor unit. In doing so, we calculated each motor unit's waveforms from the single channel in which that unit's amplitude was largest and did not attempt to remove overlapping spikes from other units before measuring the median waveform from the data. We then calculated the correlation between a unit's waveform over either trials or bins in which at least 30 spikes were present. The high correlation of a unit waveform over time, despite potential changes in the electrodes' position relative to muscle geometry over the dynamic task, provides additional confidence in both the stability of our EMG recordings and the accuracy of our spike sorting.

Data analysis

Continuous firing rates were calculated by convolving raw spike times of a motor unit with a Gaussian kernel with $\sigma = 10$ ms. This continuous result was phase-normalized across strides before calculating the mean continuous rate to identify relevant patterns such as the unit's active duration or peak firing rate. Active duration was measured between the first and last time within a stride that the smoothed curve reached the half-height from a single spike. Overall, this method allowed for quantification of firing rates even when only a single spike was present for a stride.

Joint model of rate and recruitment

We modeled the recruitment probability and firing rate based on empirical data to best characterize firing statistics within the stride. Particularly, this allowed for multiple solutions to explain why a motor unit would not spike within a stride. From the empirical data alone, strides with zero spikes would have been assumed to have no recruitment of a unit. However, to create a model of motor unit activity that includes both recruitment and rate, it must be possible that a recruited unit can have a firing rate of zero. To quantify the firing statistics that best represent all spiking and non-spiking patterns, we modeled recruitment probability and peak firing rate along the following piecewise function:

$$\text{Eq. 1: } P(Y = y) = (1 - p) + p * e^{-\lambda}, \text{ if } y = 0$$

$$\text{Eq. 2: } P(Y = y) = p * \frac{e^{-\lambda} \lambda^y}{y!}, \text{ if } y > 0$$

where y denotes the observed peak firing rate on a given stride (determined by convolving motor unit spike times with a Gaussian kernel as described above), p denotes the probability of recruitment, and λ denotes the expected peak firing rate from a Poisson distribution of outcomes. Thus, an inactive unit on a given stride may be the result of either non-recruitment or recruitment with a stochastically zero firing rate. The above equations were fit by minimizing the negative log-likelihood of the parameters given the data.

Permutation test for joint model of rate and recruitment and type 2 regression slopes

To quantify differences in firing patterns across walking speeds, we subdivided each mouse's total set of strides into speed quartiles and calculated rate (λ , Eq. 1 and 2, [Figure 5A-C](#)) and recruitment probability terms (p , Eq. 1 and 2, [Figure 5D-F](#)) for each unit in each speed quartile. Here we calculated the difference in both the rate and recruitment terms across the fastest and slowest speed quartiles ($p_{\text{fast}} - p_{\text{slow}}$ and $\lambda_{\text{fast}} - \lambda_{\text{slow}}$). To test whether these model parameters were significantly different depending on locomotor speed, we developed a null model combining strides from both the fastest and slowest speed quartiles. After pooling strides from both quartiles, we randomly distributed the pooled set of strides into two groups with sample sizes equal to the original slow and fast quartiles. We then calculated the null model parameters for each new group and found the difference between like terms. To estimate the distribution of possible differences, we bootstrapped this result using 1000 random redistributions of the pooled set of strides. Following the permutation test, the 95% confidence interval of this final distribution reflects the null hypothesis of no difference between groups. Thus, the null hypothesis can be rejected if the true difference in rate or recruitment terms exceeds this confidence interval.

We followed a similar procedure to quantify cross-muscle differences in the relationship between firing parameters. For each muscle, we estimated the slope across firing parameters for each motor unit using type 2 regression. In this case, the true difference was the difference in slopes between muscles. To test the null hypothesis that there was no difference in slopes, the null model reflected the pooled set of units from both muscles. Again, slopes were calculated for 1000 random resamplings of this pooled data to estimate the 95% confidence interval.

Supplemental Figures




Figure 1-figure supplement 1. Isolated motor units had consistent waveforms. (A,B) Example motor unit waveforms. (Left) Median waveform calculated from a random subset of strides within each trial. (Right) Median waveform calculated from spikes binned in 50ms increments of the stride. (C) (Left) Auto-correlation of each unit's median waveform between the first trial and subsequent trials. (Right) Auto-correlation of each unit's median waveform between the first 50ms of its activity and each subsequent 50ms within the stride.

Figure 2-figure supplement 1. Empirical observations of spike count distributions for all units.

Units are arranged sequentially to match the descending order presented in main text Figure 3. Units 1-17 are in the long head, while units 18-33 are in the lateral head.





Figure 4-figure supplement 1. Symbols denote different animals.

All other plotting conventions are the same as in Figure 4 in the main text.






Figure 5-figure supplement 1. Altered firing rate and recruitment across walking speed quartiles for all motor units in the long head (A) and lateral head (B).

Each point reflects the median for the model estimate of each unit across the speeds.




Figure 5-figure supplement 2. Symbols denote different animals.

All other plotting conventions are the same as in [Figure 5](#) in the main text.




Figure 6-figure supplement 1. Plots show the same analysis as main text Figure 6 for each individual speed quartile of (A) elbow angle ($\Delta\theta$) and (B) elbow velocity ($\Delta\omega$).

Speed quartiles 1 and 4 are the slowest and fastest quartiles, respectively, and p-values refer to the results of Wilcoxon signed-rank tests performed separately on data from motor units of the lateral (green) and long (purple) heads of the triceps muscle. Note that most of the muscle-specific differences shown in Figure 6 (C, E) were also present when each of the four quartiles were examined individually for each muscle.




Figure 6-figure supplement 2. Symbols denote different animals.

All other plotting conventions are the same as in Figure 6 in the main text.

Data availability

A data archive including processed data and raw EMG/kinematic data for each of six locomoting mice has been included is available at DOI: 10.5061/dryad.wdbrv162w

Acknowledgements

This work was made possible by support from NIH grants U24NS126936, R01NS109237 (SJS), NSF grant DGE-1937971 (KT), the Simons Foundation as part of the Simons-Emory International Consortium on Motor Control (SJS and MRC), the McKnight, Kavli, and Azrieli Foundations (SJS), the Portuguese Fundação para a Ciência e a Tecnologia (SFRH/BPD/119404/2016 to HGM and PTDC/MED_NEU/30890/2017 to MRC), and the European Research Council Consolidator Grant #866237 to MRC. Additionally, we thank Dr. Amanda Jacob for logistical support and William McCallum for valuable editorial comments.

Additional information

Funding

Funder	Grant reference number	Author
HHS NIH National Institute of Neurological Disorders and Stroke (NINDS)	U24NS126936	Samuel J Sober
HHS NIH National Institute of Neurological Disorders and Stroke (NINDS)	R01NS109237	Samuel J Sober
NSF National Science Foundation Graduate Research Fellowship Program (GRFP)	DGE-1937971	Kyle Thomas
Simons Foundation (SF)	Simons-Emory International Consortium on Motor Control	Samuel J Sober Megan R Carey
McKnight Foundation (McKnight)		Samuel J Sober
Kavli Foundation (Kavli)		Samuel J Sober
Azrieli Foundation (azrielifdn)		Samuel J Sober
EC European Research Council (ERC)	https://doi.org/10.3030/866237	Megan R Carey
MEC Fundação para a Ciência e a Tecnologia (FCT)	PTDC/MED_NEU/30890/2017	Megan R Carey
MEC Fundação para a Ciência e a Tecnologia (FCT)	SFRH/BPD/119404/2016	Hugo G Marques

Author ORCID iDs

Kyle Thomas: <https://orcid.org/0000-0002-2985-4718>

Hugo Marques: <https://orcid.org/0000-0002-8709-4841>

Megan R Carey: <https://orcid.org/0000-0002-4499-1657>

Samuel J Sober: <https://orcid.org/0000-0002-1140-7469>

References

- Akay T., Acharya H. J., Fouad K., Pearson K. G (2006) Behavioral and Electromyographic Characterization of Mice Lacking EphA4 Receptors. *Journal of Neurophysiology* **96**:642-651 <https://doi.org/10.1152/jn.00174.2006>
- Akay T., Tourtellotte W. G., Arber S., Jessell T. M (2014) Degradation of mouse locomotor pattern in the absence of proprioceptive sensory feedback. *Proceedings of the National Academy of Sciences of the United States of America* **111**:16877-16882 <https://doi.org/10.1073/pnas.1419045111>
- Augusto V., Padovani C. R., Campos G. E. R (2004) Skeletal muscle fiber types in C57BL/6J mice. *Braz J Morphol Sci* 89-94
- Azevedo A. W., Dickinson E. S., Gurung P., Venkatasubramanian L., Mann R. S., Tuthill J. C (2020) A size principle for recruitment of Drosophila leg motor neurons. *eLife* **9**:e56754 <https://doi.org/10.7554/eLife.56754>
- Bącznyk M., Manuel M., Roselli F., Zytynicki D. (2022) Diversity of Mammalian Motoneurons and Motor Units. In: O'Donovan M. J., Falgairolle M. (Eds). *Vertebrate Motoneurons* **28** Springer International Publishing. pp. 131-150 https://doi.org/10.1007/978-3-031-07167-6_6
- Bellardita C., Kiehn O (2015) Phenotypic Characterization of Speed-Associated Gait Changes in Mice Reveals Modular Organization of Locomotor Networks. *Current Biology* **25**:1426-1436 <https://doi.org/10.1016/j.cub.2015.04.005>
- Berg R. W (2017) Neuronal Population Activity in Spinal Motor Circuits: Greater Than the Sum of Its Parts. *Frontiers in Neural Circuits* **11** <https://doi.org/10.3389/fncir.2017.00103>
- Binder M. D., Powers R. K., Heckman C. J (2020) Nonlinear Input-Output Functions of Motoneurons. *Physiology* **35**:31-39 <https://doi.org/10.1152/physiol.00026.2019>
- Burkholder T. J., Fingado B., Baron S., Lieber R. L (1994) Relationship between muscle fiber types and sizes and muscle architectural properties in the mouse hindlimb. *Journal of Morphology* **221**:177-190 <https://doi.org/10.1002/jmor.1052210207>
- Carroll A. M., Biewener A. A (2009) Mono-versus biarticular muscle function in relation to speed and gait changes: *In vivo* analysis of the goat triceps brachii. *Journal of Experimental Biology* **212**:3349-3360 <https://doi.org/10.1242/jeb.033639>
- Chanaud C. M., Pratt C. A., Loeb G. E (1991) Functionally complex muscles of the cat hindlimb: V. The roles of histochemical fiber-type regionalization and mechanical heterogeneity in differential muscle activation. *Experimental Brain Research* **85**:300-313 <https://doi.org/10.1007/BF00229408>
- Chung B., Zia M., Thomas K. A., Michaels J. A., Jacob A., Pack A., Williams M. J., Nagapudi K., Teng L. H., Arrambide E., et al. (2023) Myomatrix arrays for high-definition muscle recording. *eLife* **12**:RP88551 <https://doi.org/10.7554/eLife.88551>
- Clarke K. A., Smart L., Still J (2001) Ground reaction force and spatiotemporal measurements of the gait of the mouse. *Behavior Research Methods, Instruments, & Computers* **33**:422-426 <https://doi.org/10.3758/BF03195396>
- Corbee R. J., Maas H., Doornenbal A., Hazewinkel H. A. W (2014) Forelimb and hindlimb ground reaction forces of walking cats: Assessment and comparison with walking dogs. *The Veterinary Journal* **202**:116-127 <https://doi.org/10.1016/j.tvjl.2014.07.001>
- Darmohray D. M., Jacobs J. R., Marques H. G., Carey M. R (2019) Spatial and Temporal Locomotor Learning in Mouse Article Spatial and Temporal Locomotor Learning in Mouse Cerebellum. *Neuron* **102**:217-231 <https://doi.org/10.1016/j.neuron.2019.01.038>
- De Luca C. J. (1985) Control properties of motor units. *Journal of Experimental Biology* **115**:125-136 <https://doi.org/10.1242/jeb.115.1.125>
- De Luca C. J., Erim Z. (1994) Common drive of motor units in regulation of muscle force. *Trends in Neurosciences* **17**:299-305 [https://doi.org/10.1016/0166-2236\(94\)90064-7](https://doi.org/10.1016/0166-2236(94)90064-7)

- DeWolf T., Schneider S., Soubiran P., Roggenbach A., Mathis M (2024) Neuro-musculoskeletal modeling reveals muscle-level neural dynamics of adaptive learning in sensorimotor cortex. *bioRxiv* <https://doi.org/10.1101/2024.09.11.612513>
- Dideriksen J. L., Del Vecchio A., Farina D (2020) Neural and muscular determinants of maximal rate of force development. *Journal of Neurophysiology* **123**:149-157 <https://doi.org/10.1152/jn.00330.2019>
- Drew T., Kalaska J., Krouchev N (2008) Muscle synergies during locomotion in the cat: A model for motor cortex control. *The Journal of Physiology* **586**:1239-1245 <https://doi.org/10.1113/jphysiol.2007.146605>
- Edman K. A (1979) The velocity of unloaded shortening and its relation to sarcomere length and isometric force in vertebrate muscle fibres. *The Journal of Physiology* **291**:143-159 <https://doi.org/10.1113/jphysiol.1979.sp012804>
- English A. W. M (1978) An Electromyographic Analysis of Forelimb Muscles During Overground Stepping in the Cat. *Journal of Experimental Biology* **76**:105-122 <https://doi.org/10.1242/jeb.76.1.105>
- English A. W. M., Weeks O. I (1987) An anatomical and functional analysis of cat biceps femoris and semitendinosus muscles. *Journal of Morphology* **191**:161-175 <https://doi.org/10.1002/jmor.1051910207>
- Enoka R. M., Duchateau J (2017) Rate coding and the control of muscle force. *Cold Spring Harbor Perspectives in Medicine* **7**:1-12 <https://doi.org/10.1101/cshperspect.a029702>
- Farina D., Negro F., Dideriksen J. L (2014) The effective neural drive to muscles is the common synaptic input to motor neurons. *The Journal of Physiology* **592**:3427-3441 <https://doi.org/10.1113/jphysiol.2014.273581>
- Gilmer J. I., Colman S. K., Cuenu G., Hutchinson J. R., Huber D., Person A. L., Al Borno M. (2024) A novel biomechanical model of the mouse forelimb predicts muscle activity in optimal control simulations of reaching movements. *bioRxiv* <https://doi.org/10.1101/2024.09.05.611289>
- Gittings W., Huang J., Vandenboom R (2012) Tetanic force potentiation of mouse fast muscle is shortening speed dependent. *Journal of Muscle Research and Cell Motility* **33**:359-368 <https://doi.org/10.1007/s10974-012-9325-6>
- Gonçalves A. I., Zavatone-Veth J. A., Carey M. R., Clark D. A (2022) Parallel locomotor control strategies in mice and flies. *Current Opinion in Neurobiology* **73**:102516 <https://doi.org/10.1016/j.conb.2022.01.001>
- Gorassini M., Eken T., Bennett D. J., Kiehn O., Hultborn H (2000) Activity of Hindlimb Motor Units During Locomotion in the Conscious Rat. *Journal of Neurophysiology* **83**:2002-2011 <https://doi.org/10.1152/jn.2000.83.4.2002>
- Grimby L (1984) Firing properties of single human motor units during locomotion. *The Journal of Physiology* **346**:195-202 <https://doi.org/10.1113/jphysiol.1984.sp015016>
- Hadzipasic M., Ni W., Nagy M., Steenrod N., McGinley M. J., Kaushal A., Thomas E., McCormick D. A., Horwich A. L (2016) Reduced high-frequency motor neuron firing, EMG fractionation, and gait variability in awake walking ALS mice. *Proceedings of the National Academy of Sciences of the United States of America* **113**:E7600-E7609 <https://doi.org/10.1073/pnas.1616832113>
- Hartigan P. M (1985) Algorithm AS 217: Computation of the Dip Statistic to Test for Unimodality. *Applied Statistics* **34**:320 <https://doi.org/10.2307/2347485>
- Heckman C. J., Enoka R. M (2012) Motor unit. *Comprehensive Physiology* **2**:2629-2682 <https://doi.org/10.1002/cphy.c100087>
- Heckman C. J., Gorassini M. A., Bennett D. J (2005) Persistent inward currents in motoneuron dendrites: Implications for motor output. *Muscle & Nerve* **31**:135-156 <https://doi.org/10.1002/mus.20261>
- Heglund N. C., Taylor C. R (1988) Speed, Stride Frequency and Energy Cost Per Stride: How Do They Change With Body Size and Gait?. *Journal of Experimental Biology* **138**:301-318 <https://doi.org/10.1242/jeb.138.1.301>

- Henneman E., Somjen G., Carpenter D. O (1965) Functional significance of cell size in spinal motoneurons. *Journal of Neurophysiology* **28**:560-580 <https://doi.org/10.1152/jn.1965.28.3.560>
- Herbin M., Gasc J.-P., Renous S (2006) How does a mouse increase its velocity? A model for investigation in the control of locomotion. *Comptes Rendus Palevol* **5**:531-540 <https://doi.org/10.1016/j.crpv.2005.12.012>
- Herbin M., Hackert R., Gasc J.-P., Renous S (2007) Gait parameters of treadmill versus overground locomotion in mouse. *Behavioural Brain Research* **181**:173-179 <https://doi.org/10.1016/j.bbr.2007.04.001>
- Hoffer J. A., O'Donovan M. J., Pratt C. A., Loeb G. E (1981) Discharge Patterns of Hindlimb Motoneurons During Normal Cat Locomotion. *Science* **213**:466-467 <https://doi.org/10.1126/science.7244644>
- Hoffer J. A., Sugano N., Loeb G. E., Marks W. B., O'Donovan M. J., Pratt C. A (1987) Cat hindlimb motoneurons during locomotion. II. Normal activity patterns. *Journal of Neurophysiology* **57**:530-553 <https://doi.org/10.1152/jn.1987.57.2.530>
- Huh S., Siripuram R., Lee R. H., Turkin V. V., O'Neill D., Hamm T. M., Heckman C. J., Manuel M (2017) PICs in motoneurons do not scale with the size of the animal: A possible mechanism for faster speed of muscle contraction in smaller species. *Journal of Neurophysiology* **118**:93-102 <https://doi.org/10.1152/jn.00045.2017>
- Kim J. J., Wyche I. S., Olson W., Lu J., Bakir M. S., Sober S. J., O'Connor D. H (2024) Myo-optogenetics: Optogenetic stimulation and electrical recording in skeletal muscles. *bioRxiv* <https://doi.org/10.1101/2024.06.21.600113>
- Kirk E. A., Hope K. T., Sober S. J., Sauerbrei B. A (2024) An output-null signature of inertial load in motor cortex. *Nature Communications* **15**:7309 <https://doi.org/10.1038/s41467-024-51750-7>
- Kishore S., Bagnall M. W., McLean D. L (2014) Systematic Shifts in the Balance of Excitation and Inhibition Coordinate the Activity of Axial Motor Pools at Different Speeds of Locomotion. *The Journal of Neuroscience* **34**:14046-14054 <https://doi.org/10.1523/JNEUROSCI.0514-14.2014>
- Livingston B. P., Nichols T. R (2014) Effects of Reinnervation of the Triceps Brachii on Joint Kinematics and Electromyographic Patterns of the Feline Forelimb during Level and Upslope Walking. *Cells Tissues Organs* **199**:405-422 <https://doi.org/10.1159/000371543>
- Lopes G., Bonacchi N., Frazão J., Neto J. P., Atallah B. V., Soares S., Moreira L., Matias S., Itskov P. M., Correia P. A., et al. (2015) Bonsai: An event-based framework for processing and controlling data streams. *Frontiers in Neuroinformatics* **9** <https://doi.org/10.3389/fninf.2015.00007>
- Lu J., Zia M., Baig D. A., Yan G., Kim J. J., Nagapudi K., Anschutz P., Oh S., O'Connor D., Sober S. J., et al. (2024) Opto-Myomatrix: μ LED integrated microelectrode arrays for optogenetic activation and electrical recording in muscle tissue. *bioRxiv* <https://doi.org/10.1101/2024.07.01.601601>
- Machado A. S., Darmohray D. M., Fayad J., Marques H. G., Carey M. R (2015) A quantitative framework for whole-body coordination reveals specific deficits in freely walking ataxic mice. *eLife* **4**:e07892 <https://doi.org/10.7554/eLife.07892>
- Manning C. D., Miller T. A., Burnham M. L., Murnaghan C. D., Calancie B., Bawa P (2010) Recovery of human motoneurons during rotation. *Experimental Brain Research* **204**:139-144 <https://doi.org/10.1007/s00221-010-2295-2>
- Manuel M., Chardon M., Tysseling V., Heckman C. J (2019) Scaling of motor output, from mouse to humans. *Physiology* **34**:5-13 <https://doi.org/10.1152/physiol.00021.2018>
- Manuel M., Heckman C. J (2011) Adult mouse motor units develop almost all of their force in the subprimary range: A new all-or-none strategy for force recruitment?. *Journal of Neuroscience* **31**:15188-15194 <https://doi.org/10.1523/JNEUROSCI.2893-11.2011>
- Manuel M., Iglesias C., Donnet M., Leroy F., Heckman C. J., Zytnicki D (2009) Fast Kinetics, High-Frequency Oscillations, and Subprimary Firing Range in Adult Mouse Spinal Motoneurons. *The Journal of Neuroscience* **29**:11246-11256 <https://doi.org/10.1523/JNEUROSCI.3260-09.2009>

- Martínez-Silva M. de L, Imhoff-Manuel R. D., Sharma A., Heckman C. J., Shneider N. A., Roselli F., Zytnicki D., Manuel M (2018) Hypoexcitability precedes denervation in the large fast-contracting motor units in two unrelated mouse models of ALS. *eLife* **7**:1-26 <https://doi.org/10.7554/eLife.30955>
- Mathewson M. A., Chapman M. A., Hentzen E. R., Fridén J., Lieber R. L (2012) Anatomical, architectural, and biochemical diversity of the murine forelimb muscles. *Journal of Anatomy* **221**:443-451 <https://doi.org/10.1111/j.1469-7580.2012.01559.x>
- Mathis A., Mamidanna P., Cury K. M., Abe T., Murthy V. N., Mathis M. W., Bethge M (2018) DeepLabCut: Markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience* **21**:1281-1289 <https://doi.org/10.1038/s41593-018-0209-y>
- Mayer W. P., Akay T (2018) Stumbling corrective reaction elicited by mechanical and electrical stimulation of the saphenous nerve in walking mice. *Journal of Experimental Biology* **221** <https://doi.org/10.1242/jeb.178095>
- Meehan C. F., Sukiasyan N., Zhang M., Nielsen J. B., Hultborn H (2010) Intrinsic Properties of Mouse Lumbar Motoneurons Revealed by Intracellular Recording In Vivo. *Journal of Neurophysiology* **103**:2599-2610 <https://doi.org/10.1152/jn.00668.2009>
- Mendes C. S., Bartos I., Márka Z., Akay T., Márka S., Mann R. S (2015) Quantification of gait parameters in freely walking rodents. *BMC Biology* **13**:1-11 <https://doi.org/10.1186/s12915-015-0154-0>
- Menelaou E., McLean D. L (2012) A Gradient in Endogenous Rhythmicity and Oscillatory Drive Matches Recruitment Order in an Axial Motor Pool. *The Journal of Neuroscience* **32**:10925-10939 <https://doi.org/10.1523/JNEUROSCI.1809-12.2012>
- Miles G. B., Sillar K. T (2011) Neuromodulation of Vertebrate Locomotor Control Networks. *Physiology* **26**:393-411 <https://doi.org/10.1152/physiol.00013.2011>
- Pachitariu M., Steinmetz N., Kadir S., Carandini M., Harris K. D. (2016) Kilosort: Realtime spike-sorting for extracellular electrophysiology with hundreds of channels. *bioRxiv* <https://doi.org/10.1101/061481>
- Petersen P. C., Berg R. W (2016) Lognormal firing rate distribution reveals prominent fluctuation-driven regime in spinal motor networks. *eLife* **5** <https://doi.org/10.7554/elife.18805>
- Ritter L. K., Tresch M. C., Heckman C. J., Manuel M., Tysseling V. M (2014) Characterization of motor units in behaving adult mice shows a wide primary range. *J Neurophysiol* **111**:543-551 <https://doi.org/10.1152/jn.00108.2014>
- Ryan M. M., Gregor R. J (1992) EMG profiles of lower extremity muscles during cycling at constant workload and cadence. *Journal of Electromyography and Kinesiology* **2**:69-80 [https://doi.org/10.1016/1050-6411\(92\)90018-E](https://doi.org/10.1016/1050-6411(92)90018-E)
- Sarver J. J., Dishowitz M. I., Kim S.-Y., Soslowsky L. J (2010) Transient decreases in forelimb gait and ground reaction forces following rotator cuff injury and repair in a rat model. *Journal of Biomechanics* **43**:778-782 <https://doi.org/10.1016/j.jbiomech.2009.10.031>
- Schiaffino S., Reggiani C (2011) Fiber Types in Mammalian Skeletal Muscles. *Physiological Reviews* **91**:1447-1531 <https://doi.org/10.1152/physrev.00031.2010>
- Schmitt D., Zumwalt A. C., Hamrick M. W (2010) The relationship between bone mechanical properties and ground reaction forces in normal and hypermuscular mice. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* **313A**:339-351 <https://doi.org/10.1002/jez.604>
- Scholle H. Ch, Schumann N. P., Biedermann F., Stegeman D. F., Graßme R., Roeleveld K., Schilling N., Fischer M. S (2001) Spatiotemporal surface EMG characteristics from rat triceps brachii muscle during treadmill locomotion indicate selective recruitment of functionally distinct muscle regions. *Experimental Brain Research* **138**:26-36 <https://doi.org/10.1007/s002210100685>
- Schumann N (2002) Multi-channel EMG of the M. triceps brachii in rats during treadmill locomotion. *Clinical Neurophysiology* **113**:1142-1151 [https://doi.org/10.1016/S1388-2457\(02\)00143-8](https://doi.org/10.1016/S1388-2457(02)00143-8)

- Schumann N. P., Biedermann F. H. W., Arnold D., Jinnah H. A., Grassme R., Fischer M. S., Scholle H. C (2006) Treadmill locomotion in normal mice—Step related multi-channel EMG profiles of thigh muscles. *Pathophysiology* **13**:245-255 <https://doi.org/10.1016/j.pathophys.2006.09.002>
- Sober S. J., Sponberg S., Nemenman I., Ting L. H (2018) Millisecond Spike Timing Codes for Motor Control. *Trends in Neurosciences* **41**:644-648 <https://doi.org/10.1016/j.tins.2018.08.010>
- Sponberg S., Spence A. J., Mullens C. H., Full R. J (2011) A single muscle's multifunctional control potential of body dynamics for postural control and running. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**:1592-1605 <https://doi.org/10.1098/rstb.2010.0367>
- Srivastava K. H., Elemans C. P. H., Sober S. J (2015) Multifunctional and Context-Dependent Control of Vocal Acoustics by Individual Muscles. *The Journal of Neuroscience* **35**:14183-14194 <https://doi.org/10.1523/JNEUROSCI.3610-14.2015>
- Srivastava K. H., Holmes C. M., Vellema M., Pack A. R., Elemans C. P. H., Nemenman I., Sober S. J (2017) Motor control by precisely timed spike patterns. *Proceedings of the National Academy of Sciences of the United States of America* **114**:1171-1176 <https://doi.org/10.1073/pnas.1611734114>
- Tata Ramalingasetty S., Danner S. M., Arreguit J., Markin S. N., Rodarie D., Kathe C., Courtine G., Rybak I. A., Ijspeert A. J. (2021) A Whole-Body Musculoskeletal Model of the Mouse. *IEEE Access* **9**:163861-163881 <https://doi.org/10.1109/ACCESS.2021.3133078>
- Ting L. H., Chiel H. J. (2017) Muscle, Biomechanics, and Implications for Neural Control. In: Hooper In S. L., Büschges A. (Eds). *Neurobiology of Motor Control* (1st) Wiley. pp. 365-416 <https://doi.org/10.1002/9781118873397.ch12>
- Van Ingen Schenau G. J., Boots P. J. M., De Groot G., Snackers R. J., Van Woensel W. W. L. M. (1992) The constrained control of force and position in multi-joint movements. *Neuroscience* **46**:197-207 [https://doi.org/10.1016/0306-4522\(92\)90019-X](https://doi.org/10.1016/0306-4522(92)90019-X)
- Van Ingen Schenau G. J., Pratt C. A., Macpherson J. M. (1994) Differential use and control of mono- and biarticular muscles. *Human Movement Science* **13**:495-517 [https://doi.org/10.1016/0167-9457\(94\)90051-5](https://doi.org/10.1016/0167-9457(94)90051-5)
- Watanabe K., Vieira T. M., Gallina A., Kouzaki M., Moritani T (2021) Novel Insights Into Biarticular Muscle Actions Gained From High-Density Electromyogram. *Exercise and Sport Sciences Reviews* **49**:179-187 <https://doi.org/10.1249/JES.0000000000000254>
- Westgaard R. H., De Luca C. J. (1999) Motor Unit Substitution in Long-Duration Contractions of the Human Trapezius Muscle. *Journal of Neurophysiology* **82**:501-504 <https://doi.org/10.1152/jn.1999.82.1.501>
- Zajac F. E., Young J. L (1980) Discharge properties of hindlimb motoneurons in decerebrate cats during locomotion induced by mesencephalic stimulation. *Journal of Neurophysiology* **43**:1221-1235 <https://doi.org/10.1152/jn.1980.43.5.1221>

Peer reviews

Reviewer #1 (Public review):

Summary:

Here, the authors have addressed the recruitment and firing patterns of motor units (MUs) from the long and lateral heads of triceps in the mouse. They used their newly developed Myomatrix arrays to record from these muscles during treadmill locomotion at different speeds, and they used template-based spike sorting (Kilosort) to extract units. Between MUs from the two heads, the authors observe differences in their firing rates, recruitment probability, phase of activation within the locomotor cycle and interspike interval patterning. Examining different walking speeds, the authors find increases in both recruitment probability and firing rates as speed increases. The authors also observed differences in the relation between recruitment and the angle of elbow extension between motor units from

each head. These differences indicate meaningful variation between motor units within and across motor pools, and may reflect the somewhat distinct joint actions of the two heads of triceps.

Strengths:

The extraction of MU spike timing for many individual units is an exciting new method that has great promise for exposing the fine detail in muscle activation and its control by the motor system. In particular, the methods developed by the authors for this purpose seem to be the only way to reliably resolve single MUs in the mouse, as the methods used previously in humans and in monkeys (e.g. Marshall et al. *Nature Neuroscience*, 2022) do not seem readily adaptable for use in rodents.

The paper provides a number of interesting observations. There are signs of interesting differences in MU activation profiles for individual muscles here, consistent with those shown by Marshall et al. It is also nice to see fine scale differences in the activation of different muscle heads, which could relate to their partially distinct functions. The mouse offers greater opportunities for understanding the control of these distinct functions, compared to the other organisms in which functional differences between heads have previously been described.

The Discussion is very thorough, providing a very nice recounting of a great deal of relevant previous results.

Weaknesses:

The findings are limited to one pair of muscle heads. While the findings are important in their own right, the lack of confirmation from analysis of other muscles acting at other joints leaves the generalization of these findings unclear.

While differences between muscle heads with somewhat distinct functions are interesting and relevant to joint control, differences between MUs for individual muscles, like those in Marshall et al., are more striking because they cannot be attributed potentially to differences in each head's function. The present manuscript does show some signs of differences for MUs within individual heads (e.g. Figure 2C), but the manuscript falls short of providing a statistical basis for the existence of distinct subpopulations.

<https://doi.org/10.7554/eLife.105829.2.sa3>

Reviewer #2 (Public review):

The present study, led by Thomas and collaborators, aims to characterise the firing activity of individual motor units in mice during locomotion. To achieve this, the team implanted small arrays of eight electrodes into two heads of the triceps and performed spike sorting using a custom implementation of Kilosort. Concurrently, they tracked the positions of the shoulder, elbow, and wrist using a single camera and a markerless motion capture algorithm (DeepLabCut). Repeated one-minute recordings were conducted in six mice across five speeds, ranging from 10 to 27.5 cm-1.

From these data, the authors demonstrate that:

- Their recording method and adapted spike-sorting algorithm enable robust decoding of motor unit activity during rapid movements.
- Identified motor units tend to be recruited during a subset of strides, with recruitment probability increasing with speed.

- Motor units within individual heads of the triceps likely receive common synaptic inputs that correlate their activity, whereas motor units from different heads exhibit distinct behaviour.

The authors conclude that these differences arise from the distinct functional roles of the muscles and the task constraints (i.e., speed).

Strengths:

- The novel combination of electrode arrays for recording intramuscular electromyographic signals from a larger muscle volume, paired with an advanced spike-sorting pipeline capable of identifying motor unit populations.

- The robustness of motor unit decoding during fast movements.

Weaknesses:

- The data do not clearly indicate which motor units were sampled from each pool, leaving uncertainty as to whether the sample is biased towards high-threshold motor units or representative of the entire pool.

- The results largely confirm the classic physiological framework of motor unit recruitment and rate coding, offering limited new insights into motor unit physiology.

I would like to thank the authors for their thorough and insightful revisions. I am particularly pleased with the inclusion of the new analyses demonstrating the robustness of motor unit decoding, as well as the improved transparency regarding spike-sorting yield for each muscle and animal. Additionally, the new analyses illustrating that recruitment within muscle heads is consistent with the presence of common synaptic inputs and orderly recruitment significantly strengthen the manuscript.

<https://doi.org/10.7554/eLife.105829.2.sa2>

Reviewer #3 (Public review):

Summary:

Using the approach of Myomatrix recording, the authors report that 1) motor units are recruited differently in the two types of muscles and 2) individual units are probabilistically recruited during the locomotion strides, whereas the population bulk EMG has a more reliable representation of the muscle. Third, the recruitment of units was proportional to walking speed.

Strengths:

The new technique provides a unique dataset, and the data analysis is convincing and well-executed.

Weaknesses:

After the revision, I no longer see any apparent weaknesses in the study.

<https://doi.org/10.7554/eLife.105829.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

Summary:

Here, the authors have addressed the recruitment and firing patterns of motor units (MUs) from the long and lateral heads of the triceps in the mouse. They used their newly developed Myomatrix arrays to record from these muscles during treadmill locomotion at different speeds, and they used template-based spike sorting (Kilosort) to extract units. Between MUs from the two heads, the authors observed differences in their firing rates, recruitment probability, phase of activation within the locomotor cycle, and interspike interval patterning. Examining different walking speeds, the authors find increases in both recruitment probability and firing rates as speed increases. The authors also observed differences in the relation between recruitment and the angle of elbow extension between motor units from each head. These differences indicate meaningful variation between motor units within and across motor pools and may reflect the somewhat distinct joint actions of the two heads of triceps.

Strengths:

The extraction of MU spike timing for many individual units is an exciting new method that has great promise for exposing the fine detail in muscle activation and its control by the motor system. In particular, the methods developed by the authors for this purpose seem to be the only way to reliably resolve single MUs in the mouse, as the methods used previously in humans and in monkeys (e.g. Marshall et al. Nature Neuroscience, 2022) do not seem readily adaptable for use in rodents.

The paper provides a number of interesting observations. There are signs of interesting differences in MU activation profiles for individual muscles here, consistent with those shown by Marshall et al. It is also nice to see fine-scale differences in the activation of different muscle heads, which could relate to their partially distinct functions. The mouse offers greater opportunities for understanding the control of these distinct functions, compared to the other organisms in which functional differences between heads have previously been described.

The Discussion is very thorough, providing a very nice recounting of a great deal of relevant previous results.

We thank the Reviewer for these comments.

Weaknesses:

The findings are limited to one pair of muscle heads. While an important initial finding, the lack of confirmation from analysis of other muscles acting at other joints leaves the general relevance of these findings unclear.

The Reviewer raises a fair point. While outside the scope of this paper, future studies should certainly address a wider range of muscles to better characterize motor unit firing patterns across different sets of effectors with varying anatomical locations. Still, the importance of results from the triceps long and lateral heads should not be understated as this paper, to our knowledge, is the first to capture the difference in firing patterns of motor units across any set of muscles in the locomoting mouse.

While differences between muscle heads with somewhat distinct functions are interesting and relevant to joint control, differences between MUs for individual muscles, like those in Marshall et al., are more striking because they cannot be attributed potentially to

differences in each head's function. The present manuscript does show some signs of differences for MUs within individual heads: in Figure 2C, we see what looks like two clusters of motor units within the long head in terms of their recruitment probability. However, a statistical basis for the existence of two distinct subpopulations is not provided, and no subsequent analysis is done to explore the potential for differences among MUs for individual heads.

We agree with the Reviewer and have revised the manuscript to better examine potential subpopulations of units within each muscle as presented in Figure 2C. We performed Hartigan's dip test on motor units within each muscle to test for multimodal distributions. For both muscles, $p > 0.05$, so we cannot reject the null hypothesis that the units in each muscle come from a multimodal distribution. However, Hartigan's test and similar statistical methods have poor statistical power for the small sample sizes ($n=17$ and 16 for long and lateral heads, respectively) considered here, so the failure to achieve statistical significance might reflect either the absence of a true difference or a lack of statistical resolution.

Still, the limited sample size warrants further data collection and analysis since the varying properties across motor units may lead to different activation patterns. Given these results, we have edited the text as follows:

“A subset of units, primarily in the long head, were recruited in under 50% of the total strides and with lower spike counts (Figure 2C). This distribution of recruitment probabilities might reflect a functionally different subpopulation of units. However, the distribution of recruitment probabilities were not found to be significantly multimodal ($p>0.05$ in both cases, Hartigan's dip test; Hartigan, 1985). However, Hartigan's test and similar statistical methods have poor statistical power for the small sample sizes ($n=17$ and 16 for long and lateral heads, respectively) considered here, so the failure to achieve statistical significance might reflect either the absence of a true difference or a lack of statistical resolution.”

The statistical foundation for some claims is lacking. In addition, the description of key statistical analysis in the Methods is too brief and very hard to understand. This leaves several claims hard to validate.

We thank the Reviewer for these comments and have clarified the text related to key statistical analyses throughout the manuscript, as described in our other responses below.

Reviewer #2 (Public review):

The present study, led by Thomas and collaborators, aims to describe the firing activity of individual motor units in mice during locomotion. To achieve this, they implanted small arrays of eight electrodes in two heads of the triceps and performed spike sorting using a custom implementation of Kilosort. Simultaneously, they tracked the positions of the shoulder, elbow, and wrist using a single camera and a markerless motion capture algorithm (DeepLabCut). Repeated one-minute recordings were conducted in six mice at five different speeds, ranging from 10 to $27.5 \text{ cm}\cdot\text{s}^{-1}$.

From these data, the authors reported that:

- (1) a significant portion of the identified motor units was not consistently recruited across strides,*
- (2) motor units identified from the lateral head of the triceps tended to be recruited later than those from the long head,*
- (3) the number of spikes per stride and peak firing rates were correlated in both muscles, and*

(4) the probability of motor unit recruitment and firing rates increased with walking speed.

The authors conclude that these differences can be attributed to the distinct functions of the muscles and the constraints of the task (i.e., speed).

Strengths:

The combination of novel electrode arrays to record intramuscular electromyographic signals from a larger muscle volume with an advanced spike sorting pipeline capable of identifying populations of motor units.

We thank the Reviewer for this comment.

Weaknesses:

(1) There is a lack of information on the number of identified motor units per muscle and per animal.

The Reviewer is correct that this information was not explicitly provided in the prior submission. We have therefore added Table 1 that quantifies the number of motor units per muscle and per animal.

(2) All identified motor units are pooled in the analyses, whereas per-animal analyses would have been valuable, as motor units within an individual likely receive common synaptic inputs. Such analyses would fully leverage the potential of identifying populations of motor units.

Please see our answer to the following point, where we address questions (2) and (3) together.

(3) The current data do not allow for determining which motor units were sampled from each pool. It remains unclear whether the sample is biased toward high-threshold motor units or representative of the full pool.

We thank the Reviewer for these comments. To clarify how motor unit responses were distributed across animals and muscle targets, we updated or added the following figures:

Figure 2C

Figure 4–figure supplement 1

Figure 5–figure supplement 2

Figure 6–figure supplement 2

These provide a more complete look at the range of activity within each motor pool, suggesting that we do measure from units with different activation thresholds within the same motor pool, rather than this variation being due to cross-animal differences. For example, Figure 2C illustrates that motor units from the same muscle and animal show a wide variety of recruitment probabilities. However, the limited number of motor units recorded from each individual animal does not allow a statistically rigorous test for examining cross-animal differences.

(4) The behavioural analysis of the animals relies solely on kinematics (2D estimates of elbow angle and stride timing). Without ground reaction forces or shoulder angle data, drawing functional conclusions from the results is challenging.

The Reviewer is correct that we did not measure muscular force generation or ground reaction forces in the present study. Although outside the scope of this study, future work

might employ buckle force transducers as used in larger animals (Biewener et al., 1988; Karabulut et al., 2020) to examine the complex interplay between neural commands, passive biomechanics, and the complex force-generating properties of muscle tissue.

Major comments:

(1) Spike sorting

The conclusions of the study rely on the accuracy and robustness of the spike sorting algorithm during a highly dynamic task. Although the pipeline was presented in a previous publication (Chung et al., 2023, eLife), a proper validation of the algorithm for identifying motor unit spikes is still lacking. This is particularly important in the present study, as the experimental conditions involve significant dynamic changes. Under such conditions, muscle geometry is altered due to variations in both fibre pennation angles and lengths.

This issue differs from electrode drift, and it is unclear whether the original implementation of Kilosort includes functions to address it. Could the authors provide more details on the various steps of their pipeline, the strategies they employed to ensure consistent tracking of motor unit action potentials despite potential changes in action potential waveforms, and the methods used for manual inspection of the spike sorting algorithm's output?

This is an excellent point and we agree that the dynamic behavior used in this investigation creates potential new challenges for spike sorting. In our analysis, Kilosort 2.5 provides key advantages in comparing unit waveforms across multiple channels and in detecting overlapping spikes. We modified this version of Kilosort to construct unit waveform templates using only the channels within the same muscle (Chung et al., 2023), as clarified in the revised Methods section (see “Electromyography (EMG)”):

“A total of 33 units were identified across all animals. Each unit’s isolation was verified by confirming that no more than 2% of inter-spike intervals violated a 1 ms refractory limit. Additionally, we manually reviewed cross-correlograms to ensure that each waveform was only reported as a single motor unit.”

The Reviewer is correct that our ability to precisely measure a unit’s activity based on its waveform will depend on the relationship between the embedded electrode and the muscle geometry, which alters over the course of the stride. As a follow-up to the original text, we have included new analyses to characterize the waveform activity throughout the experiment and stride (also in Methods):

“We further validated spike sorting by quantifying the stability of each unit’s waveform across time (Figure 1–figure supplement 1). First, we calculated the median waveform of each unit across every trial to capture long-term stability of motor unit waveforms. Additionally, we calculated the median waveform through the stride binned in 50 ms increments using spiking from a single trial. This second metric captures the stability of our spike sorting during the rapid changes in joint angles that occur during the burst of an individual motor unit. In doing so, we calculated each motor unit’s waveforms from the single channel in which that unit’s amplitude was largest and did not attempt to remove overlapping spikes from other units before measuring the median waveform from the data. We then calculated the correlation between a unit’s waveform over either trials or bins in which at least 30 spikes were present. The high correlation of a unit waveform over time, despite potential changes in the electrodes’ position relative to muscle geometry over the dynamic task, provides additional confidence in both the stability of our EMG recordings and the accuracy of our spike sorting.”

(2) Yield of the spike sorting pipeline and analyses per animal/muscle

A total of 33 motor units were identified from two heads of the triceps in six mice (17 from the long head and 16 from the lateral head). However, precise information on the yield per muscle per animal is not provided. This information is crucial to support the novelty of the study, as the authors claim in the introduction that their electrode arrays enable the identification of populations of motor units. Beyond reporting the number of identified motor units, another way to demonstrate the effectiveness of the spike sorting algorithm would be to compare the recorded EMG signals with the residual signal obtained after subtracting the action potentials of the identified motor units, using a signal-to-residual ratio.

Furthermore, motor units identified from the same muscle and the same animal are likely not independent due to common synaptic inputs. This dependence should be accounted for in the statistical analyses when comparing changes in motor unit properties across speeds and between muscles.

We thank the Reviewer for this comment. Regarding motor unit yield, as described above the newly-added Table 1 displays the yield from each animal and muscle.

Regarding spike sorting, while signal-to-residual is often an excellent metric, it is not ideal for our high-resolution EMG signals since isolated single motor units are typically superimposed on a “bulk” background consisting of the low-amplitude waveforms of other motor units. Because these smaller units typically cannot be sorted, it is challenging to estimate the “true” residual after subtracting (only) the largest motor unit, since subtracting each sorted unit’s waveform typically has a very small effect on the RMS of the total EMG signal. To further address concerns regarding spike sorting quality, we added Figure 1–figure supplement 1 that demonstrates motor units’ consistency over the experiment, highlighting that the waveform maintains its shape within each stride despite muscle/limb dynamics and other possible sources of electrical noise or artifact.

Finally, the Reviewer is correct that individual motor units in the same muscle are very likely to receive common synaptic inputs. These common inputs may reflect in sparse motor units being recruited in overlapping rather than different strides. Indeed, in the following added to the Results, we identified that motor units are recruited with higher probability when additional units are recruited.

“Probabilistic recruitment is correlated across motor units

Our results show that the recruitment of individual motor units is probabilistic even within a single speed quartile (Figure 5A-C) and predicts body movements (Figure 6), raising the question of whether the recruitment of individual motor units are correlated or independent. Correlated recruitment might reflect shared input onto the population of motor units innervating the muscle (De Luca, 1985; De Luca & Erim, 1994; Farina et al., 2014). For example, two motor units, each with low recruitment probabilities, may still fire during the same set of strides. To assess the independence of motor unit recruitment across the recorded population, we compared each unit’s empirical recruitment probability across all strides to its conditional recruitment probability during strides in which another motor unit from the same muscle was recruited (Figure 7). Doing this for all motor unit pairs revealed that motor units in both muscles were biased towards greater recruitment when additional units were active ($p < 0.001$, Wilcoxon signed-rank tests for both the lateral and long heads of triceps). This finding suggests that probabilistic recruitment reflects common synaptic inputs that covary together across locomotor strides.”

(3) Representativeness of the sample of identified motor units


However, to draw such conclusions, the authors should exclusively compare motor units from the same pool and systematically track violations of the recruitment order.

Alternatively, they could demonstrate that the motor units that are intermittently active across strides correspond to the smallest motor units, based on the assumption that these units should always be recruited due to their low activation thresholds.

One way to estimate the size of motor units identified within the same muscle would be to compare the amplitude of their action potentials, assuming that all motor units are relatively close to the electrodes (given the selectivity of the recordings) and that motoneurons innervating more muscle fibres generate larger motor unit action potentials.

We thank the Reviewer for this comment. Below, we provide more detailed analyses of the relationships between motor unit spike amplitude and the recruitment probability as well as latency (relative to stride onset) of activation.

We generated the below figures to illustrate the relationship between the amplitude of motor units and their firing properties. As suspected, units with larger-amplitude waveforms fired with lower probability and produced their first spikes later in the stride. If we were comfortable assuming that larger spike amplitudes mean higher-force units, then this would be consistent with a key prediction of the size principle (i.e. that higher-force units are recruited later). However, we are hesitant to base any conclusions on this assumption or emphasize this point with a main-text figure, since EMG signal amplitude may also vary due to the physical properties of the electrode and distance from muscle fibers. Thus it is possible that a large motor unit may have a smaller waveform amplitude relative to the rest of the motor pool.



Author response image 1. Relation between motor unit amplitude and (A) recruitment probability and (B) mean first spike time within the stride. Colored lines indicate the outcome of linear regression analyses.

Currently, the data seem to support the idea that motor units that are alternately recruited across strides have recruitment thresholds close to the level of activation or force produced during slow walking. The fact that recruitment probability monotonically increases with speed suggests that the force required to propel the mouse forward exceeds the recruitment threshold of these "large" motor units. This pattern would primarily reflect spatial recruitment following the size principle rather than flexible motor unit control.

We thank the Reviewer for this comment. We agree with this interpretation, particularly in relation to the references suggested in later comments, and have added the following text to the Discussion to better reflect this argument:

“To investigate the neuromuscular control of locomotor speed, we quantified speed-dependent changes in both motor unit recruitment and firing rate. We found that the majority of units were recruited more often and with larger firing rates at faster speeds (Figure 5, Figure 5—figure supplement 1). This result may reflect speed-dependent differences in the common input received by populations of motor neurons with varying spiking thresholds (Henneman et al., 1965). In the case of mouse locomotion, faster speeds might reflect a larger common input, increasing the recruitment probability as more neurons, particularly those that are larger and generate more force, exceed threshold for action potentials (Farina et al., 2014).”

(4) Analysis of recruitment and firing rates

The authors currently report active duration and peak firing rates based on spike trains convolved with a Gaussian kernel. Why not report the peak of the instantaneous firing rates estimated from the inverse of the inter-spike interval? This approach appears to be more aligned with previous studies conducted to describe motor unit behaviour during fast movements (e.g., Desmedt & Godaux, 1977, J Physiol; Van Cutsem et al., 1998, J Physiol; Del Vecchio et al., 2019, J Physiol).

We thank the Reviewer for this comment. In the revised Discussion (see ‘Firing rates in mouse locomotion compared to other species’) we reference several examples of previous studies that quantified spike patterns based on the instantaneous firing rate. We chose to report the peak of the smoothed firing rate because that quantification includes strides with zero spikes or only one spike, which occur regularly in our dataset (and for which ISI rate measures, which require two spikes to define an instantaneous firing rate, cannot be computed). Regardless, in the revised Figure 4B, we present an analysis that uses inter-spike intervals as suggested, which yielded similar ranges of firing rates as the primary analysis.

(5) Additional analyses of behaviour

The authors currently analyse motor unit recruitment in relation to elbow angle. It would be valuable to include a similar analysis using the angular velocity observed during each stride, re broadly, comparing stride-by-stride changes in firing rates with changes in elbow angular velocity would further strengthen the final analyses presented in the results section.

We thank the Reviewer for this comment. To address this, we have modified Figure 6 and the associated Supplemental Figures, to show relationships in unit activation with both the range of elbow extension and the range of elbow velocity for each stride. These new Supplemental Figures show that the trends shown in main text Figure 6C and 6E (which show data from all speed quartiles on the same axes) are also apparent in both the slower and faster quartiles individually, although single-quartile statistical tests (with smaller sample size than the main analysis) not reach statistical significance in all cases.

Reviewer #3 (Public review):

Summary:

Using the approach of Myomatrix recording, the authors report that:

(1) Motor units are recruited differently in the two types of muscles.

(2) Individual units are probabilistically recruited during the locomotion strides, whereas the population bulk EMG has a more reliable representation of the muscle.

(3) The recruitment of units was proportional to walking speed.

Strengths:

The new technique provides a unique data set, and the data analysis is convincing and well-performed.

We thank the Reviewer for the comment.

Weaknesses:

The implications of "probabilistical recruitment" should be explored, addressed, and analyzed further.

Comments:

One of the study's main findings (perhaps the main finding) is that the motor units are "probabilistically" recruited. The authors do not define what they mean by probabilistically recruited, nor do they present an alternative scenario to such recruitment or discuss why this would be interesting or surprising. However, on page 4, they do indicate that the recruitment of units from both muscles was only active in a subset of strides, i.e., they are not reliably active in every step.

If probabilistic means irregular spiking, this is not new. Variability in spiking has been seen numerous times, for instance in human biceps brachii motor units during isometric contractions (Pascoe, Enoka, Exp physiology 2014) and elsewhere. Perhaps the distinction the authors are seeking is between fluctuation-driven and mean-driven spiking of motor units as previously identified in spinal motor networks (see Petersen and Berg, eLife 2016, and Berg, Frontiers 2017). Here, it was shown that a prominent regime of irregular spiking is present during rhythmic motor activity, which also manifests as a positive skewness in the spike count distribution (i.e., log-normal).

We thank the Reviewer for this comment and have clarified several passages in response. The Reviewer is of course correct that irregular motor unit spiking has been described previously and may reflect motor neurons' operating in a high-sensitivity (fluctuation-driven) regime. We now cite these papers in the Discussion (see 'Firing rates in mouse locomotion compared to other species'). Additionally, the revision clarifies that "probabilistically" - as defined in our paper - refers only to the empirical observation that a motor unit spikes during only a subset of strides, either when all locomotor speeds are considered together (Figure 2) or separately (Figure 5A-C):

"Motor units in both muscles exhibited this pattern of probabilistic recruitment (defined as a unit's firing on only a fraction of strides), but with differing distributions of firing properties across the long and lateral heads (Figure 2)."

"Our findings (Figure 4) highlight that even with the relatively high firing rates observed in mice, there are still significant changes in firing rate and recruitment probability across the spikes within bursts (Figure 4B) and across locomotor speeds (Figure 5F). Future studies should more carefully examine how these rapidly changing spiking patterns derive from both the statistics of synaptic inputs and intrinsic properties of motor neurons (Manuel & Heckman, 2011; Petersen & Berg, 2016; Berg, 2017)."

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

As mentioned above, there are several issues with the statistics that need to be corrected to properly support the claims made in the paper.

The authors compare the fractions of MUs that show significant variation across locomotor speeds in their firing rate and recruitment probability. However, it is not statistically founded to compare the results of separate statistical tests based on different kinds of measurements and thus have unconstrained differences in statistical power. The comparison of the fractional changes in firing rates and recruitment across speeds that follow is helpful, though in truth, by contemporary standards, one would like to see error bars on these estimates. These could be generated using bootstrapping.

The Reviewer is correct, and we have revised the manuscript to better clarify which quantities should or should not be compared, including the following passage (see “Motor unit mechanisms of speed control” in Results):

“Speed-dependent increases in peak firing rate were therefore also present in our dataset, although in a smaller fraction of motor units (22/33) than changes in recruitment probability (31/33). Furthermore, the mean ($\pm$ SE) magnitude of speed-dependent increases was smaller for spike rates (mean $\text{rate}_{\text{fast}}/\text{rate}_{\text{slow}}$ of $111\% \pm 20\%$ across all motor units) than for recruitment probabilities (mean $p(\text{recruitment})_{\text{fast}}/p(\text{recruitment})_{\text{slow}}$ of $179\% \pm 3\%$ across all motor units). While fractional changes in rate and recruitment probability are not readily comparable given their different upper limits, these findings could suggest that while both recruitment and peak rate change across speed quartiles, increased recruitment probability may play a larger role in driving changes in locomotor speed.”

The description in the Methods of the tests for variation in firing rates and recruitment probability across speeds are extremely hard to understand - after reading many times, it is still not clear what was done, or why the method used was chosen. In the main text, the authors quote p-values and then state "bootstrap confidence intervals," which is not a statistical test that yields a p-value. While there are mathematical relationships between confidence intervals and statistical tests such that a one-to-one correspondence between them can exist, the descriptions provided fall short of specifying how they are related in the present instance. For this reason, and those described in what follows, it is not clear what the p-values represent.

Next, the authors refer to fitting a model ("a Poisson distribution") to the data to estimate firing rate and recruitment probability, that the model results agree with their actual data, and that they then bootstrapped from the model estimates to get confidence intervals and compute p-values. Why do this? Why not just do something much simpler, like use the actual spike counts, and resample from those? I understand that it is hard to distinguish between no recruitment and just no spikes given some low Poisson firing rate, but how does that challenge the ability to test if the firing rates or the number of spiking MUs changes significantly across speeds? I can come up with some reasons why I think the authors might have decided to do this, but reasoning like this really should be made explicit.

In addition, the authors would provide an unambiguous description of the model, perhaps using an equation and a description of how it was fit. For the bootstrapping, a clear description of how the resampling was done should be included. The focus on peak firing rate instead of mean (or median) firing rate should also be justified. Since peaks are noisier, I would expect the statistical power to be lower compared to using the mean or median.

We thank the Reviewer for the comments and have revised and expanded our discussion of the statistical tests employed. We expanded and clarified our description of these techniques in the updated Methods section:

“Joint model of rate and recruitment

We modeled the recruitment probability and firing rate based on empirical data to best characterize firing statistics within the stride. Particularly, this allowed for multiple solutions to explain why a motor unit would not spike within a stride. From the empirical data alone, strides with zero spikes would have been assumed to have no recruitment of a unit. However, to create a model of motor unit activity that includes both recruitment and rate, it must be possible that a recruited unit can have a firing rate of zero. To quantify the firing statistics that best represent all spiking and non-spiking patterns, we modeled recruitment probability and peak firing rate along the following piecewise function:

$$\text{Eq. 1: } P(Y = y) = (1 - p) + p * e^{-\lambda}, \text{ if } y = 0$$

$$\text{Eq. 2: } P(Y = y) = p * \frac{e^{-\lambda} \lambda^y}{y!}, \text{ if } y > 0$$

where y denotes the observed peak firing rate on a given stride (determined by convolving motor unit spike times with a Gaussian kernel as described above), p denotes the probability of recruitment, and λ denotes the expected peak firing rate from a Poisson distribution of outcomes. Thus, an inactive unit on a given stride may be the result of either non-recruitment or recruitment with a stochastically zero firing rate. The above equations were fit by minimizing the negative log-likelihood of the parameters given the data.

“Permutation test for joint model of rate and recruitment and type 2 regression slopes


To quantify differences in firing patterns across walking speeds, we subdivided each mouse’s total set of strides into speed quartiles and calculated rate (λ , Eq. 1 and 2, Fig. 5A-C) and recruitment probability terms (p , Eq. 1 and 2, Fig. 5D-F) for each unit in each speed quartile. Here we calculated the difference in both the rate and recruitment terms across the fastest and slowest speed quartiles ($p_{\text{fast}} - p_{\text{slow}}$ and $\lambda_{\text{fast}} - \lambda_{\text{slow}}$). To test whether these model parameters were significantly different depending on locomotor speed, we developed a null model combining strides from both the fastest and slowest speed quartiles. After pooling strides from both quartiles, we randomly distributed the pooled set of strides into two groups with sample sizes equal to the original slow and fast quartiles. We then calculated the null model parameters for each new group and found the difference between like terms. To estimate the distribution of possible differences, we bootstrapped this result using 1000 random redistributions of the pooled set of strides. Following the permutation test, the 95% confidence interval of this final distribution reflects the null hypothesis of no difference between groups. Thus, the null hypothesis can be rejected if the true difference in rate or recruitment terms exceeds this confidence interval.

We followed a similar procedure to quantify cross-muscle differences in the relationship between firing parameters. For each muscle, we estimated the slope across firing parameters for each motor unit using type 2 regression. In this case, the true difference was the difference in slopes between muscles. To test the null hypothesis that there was no difference in slopes, the null model reflected the pooled set of units from both muscles. Again, slopes were calculated for 1000 random resamplings of this pooled data to estimate the 95% confidence interval.”

The argument for delayed activation of the lateral head is interesting, but I am not comfortable saying the nervous system creates a delay just based on observations of the mean time of the first spike, given the potential for differential variability in spike timing across muscles and MUs. One way to make a strong case for a delay would be to show aggregate PSTHs for all the spikes from all the MUs for each of the two heads. That would distinguish between a true delay and more gradual or variable activation between the heads.

This is a good point and we agree that the claim made about the nervous system is too strong given the results. Even with Author response image 2 below that the Reviewer suggested,

there is still not enough evidence to isolate the role of the nervous system in the muscles' activation.



Author response image 2. Aggregate peristimulus time histogram (PSTH) for all motor unit spike times in the long head (top) and lateral head (bottom) within the stride.

In the ideal case, we would have more simultaneous recordings from both muscles to make a more direct claim on the delay. Still, within the current scope of the paper, to correct this and better describe the difference in timing of muscle activity, we edited the text to the following:

“These findings demonstrate that despite the synergistic (extensor) function of the long and lateral heads of the triceps at the elbow, the motor pool for the long head becomes active roughly 100 ms before the motor pool supplying the lateral head during locomotion (Figure 3C).”

The results from Marshall et al. 2022 suggest that the recruitment of some MUs is not just related to muscle force, but also the frequency of force variation - some of their MUs appear to be recruited only at certain frequencies. Figure 5C could have shown signs of this, but it does not appear to. We do not really know the force or its frequency of variation in the measurements here. I wonder whether there is additional analysis that could address whether frequency-dependent recruitment is present. It may not be addressable with the current data set, but this could be a fruitful direction to explore in the future with MU recordings from mice.

We agree that this would be a fruitful direction to explore, however the Reviewer is correct that this is not easily addressable with the dataset. As the Reviewer points out, stride frequency increases with increased speed, potentially offering the opportunity to examine how motor unit activity varies with the frequency, phase, and amplitude of locomotor movements. However, given our lack of force data (either joint torques or ground reaction forces), dissociating the frequency/phase/amplitude of skeletal kinematics from the frequency/phase/amplitude of muscle force. Marshall et al. (2022) mitigated these issues by using an isometric force-production task (Marshall et al., 2022). Therefore, while we agree that it would be a major contribution to extend such investigations to whole-body movements like locomotion, given the complexities described above we believe this is a project for the future, and beyond the scope of the present study.

Minor:

Page 5: "Units often displayed no recruitment in a greater proportion of strides than for any particular spike count when recruited (Figures 2A, B)," - I had to read this several

| times to understand it. I suggest rephrasing for clarity.

We have changed the text to read:

“Units demonstrated a variety of firing patterns, with some units producing 0 spikes more frequently than any non-zero spike count (Figure 2A, B),...”

| *Figure 3 legend: "Mean phase ({plus minus} SE) of motor unit burst duration across all strides." It is unclear what this means - durations are not usually described as having a phase. Do we mean the onset phase?*

We have changed the text to read:

“Mean phase $\pm$ SE of motor unit burst activity within each stride”

| *Page 9: "suggesting that the recruitment of individual motor units in the lateral and long heads might have significant (and opposite) effects on elbow angle in strides of similar speed (see Discussion)." I wouldn't say "opposite" here - that makes it sound like the authors are calling the long head a flexor. The authors should rephrase or clarify the sense in which they are opposite.*

This is a fair point and we agree we should not describe the muscles as ‘opposite’ when both muscles are extensors. We have removed the phrase ‘and opposite’ from the text.

| *Page 11: "in these two muscles across in other quadrupedal species" - typo.*

We have corrected this error.

| *Page 16: This reviewer cannot decipher after repeated attempts what the first two sentences of the last paragraph mean. - "Future studies might also use perturbations of muscle activity to dissociate the causal properties of each motor unit's activity from the complex correlation structure of locomotion. Despite the strong correlations observed between motor unit recruitment and limb kinematics (Fig. 6, Supplemental Fig. 3), these results might reflect covariations of both factors with locomotor speed rather than the causal properties of the recorded motor unit."*

For better clarity, we have changed the text to read:

“Although strong correlations were observed between motor unit recruitment and limb kinematics during locomotion (Figure 6, Figure 6-figure supplement 1), it remains unclear whether such correlations actually reflect the causal contributions that those units make to limb movement. To resolve this ambiguity, future studies could use electrical or optical perturbations of muscle contraction levels (Kim et al., 2024; Lu et al., 2024; Srivastava et al., 2015, 2017) to test directly how motor unit firing patterns shape locomotor movements. The short-latency effects of patterned motor unit stimulation (Srivastava et al., 2017) could then reveal the sensitivity of behavior to changes in muscle spiking and the extent to which the same behaviors can be performed with many different motor commands.”

Reviewer #2 (Recommendations for the authors):

Minor comments:

Introduction:

(1) "Although studies in primates, cats, and zebrafish have shown that both the number of active motor units and motor unit firing rates increase at faster locomotor speeds (Grimby, 1984; Hoffer et al., 1981, 1987; Marshall et al., 2022; Menelaou & McLean, 2012)." I would remove Marshall et al. (2022) as their monkeys performed pulling tasks

with the upper limb. You can alternatively remove locomotor from the sentence and replace it with contraction speed.

Thank you for the comment. While we intended to reference this specific paper to highlight the rhythmic activity in muscles, we agree that this deviates from 'locomotion' as it is referenced in the other cited papers which study body movement. We have followed the Reviewer's suggestion to remove the citation to Marshall et al.

(2) "The capability and need for faster force generation during dynamic behavior could implicate motor unit recruitment as a primary mechanism for modulating force output in mice."

The authors could add citations to this sentence, of works that showed that recruitment speed is the main determinant of the rate of force development (see for example Dideriksen et al. (2020) J Neurophysiol; J. L. Dideriksen, A. Del Vecchio, D. Farina, Neural and muscular determinants of maximal rate of force development. J Neurophysiol 123, 149-157 (2020)).

Thank you for pointing out this important reference. We have included this as a citation as recommended.

Results:

(3) "Electrode arrays (32-electrode Myomatrix array model RF-4x8-BHS-5) were implanted in the triceps brachii (note that Figure 1D shows the EMG signal from only one of the 16 bipolar recording channels), and the resulting data were used to identify the spike times of individual motor units (Figure 1E) as described previously (Chung et al., 2023)."

This sentence can be misleading for the reader as the array used by the researchers has 4 threads of 8 electrodes. Would it be possible to specify the number of electrodes implanted per head of interest? I assume 8 per head in most mice (or 4 bipolar channels), even if that's not specifically written in the manuscript.

Thank you for the suggestion. As described above, we have added Table 1, which includes all array locations, and we edited the statement referenced in the comment as follows:

"Electrode arrays (32-electrode Myomatrix array model RF-4x8-BHS-5) were implanted in forelimb muscles (note that Figure 1D shows the EMG signal from only one of the 16 bipolar recording channels), and the resulting data were used to identify the spike times of individual motor units in the triceps brachii long and lateral heads (Table 1, Figure 1E) as described previously (Chung et al., 2023)."

(4) "These findings demonstrate that despite the overlapping biomechanical functions of the long and lateral heads of the triceps, the nervous system creates a consistent, approximately 100 ms delay (Figure 3C) between the activation of the two muscles' motor neuron pools. This timing difference suggests distinct patterns of synaptic input onto motor neurons innervating the lateral and long heads."

Both muscles don't have fully overlapping biomechanical functions, as one of them also acts on the shoulder joint. Please be more specific in this sentence, saying that both muscles are synergistic at the elbow level rather than "have overlapping biomechanical functions".

We agree with the above reasoning and that our manuscript should be clearer on this point. We edited the above text in accordance with the Reviewer suggestion as follows:

"These findings demonstrate that despite the synergistic (extensor) function of the long and lateral heads of the triceps at the elbow, ..."

(5) *"Together with the differences in burst timing shown in Figure 3B, these results again suggest that the motor pools for the lateral and long heads of the triceps receive distinct patterns of synaptic input, although differences in the intrinsic physiological properties of motor neurons innervating the two muscles might also play an important role."*

It is difficult to draw such an affirmative conclusion on the synaptic inputs from the data presented by the authors. The differences in firing rates may solely arise from other factors than distinct synaptic inputs, such as the different intrinsic properties of the motoneurons or the reception of distinct neuromodulatory inputs.

To better explain our findings, we adjusted the above text in the Results (see "Motor unit firing patterns in the long and lateral heads of the triceps"):

"Together with the differences in burst timing shown in Figure 3B, these results again suggest that the motor pools for the lateral and long heads of the triceps receive distinct patterns of synaptic input, although differences in the intrinsic physiological properties of motor neurons innervating the two muscles might also play an important role."

We also included the following distinction in the Discussion (see "Differences in motor unit activity patterns across two elbow extensors") to address the other plausible mechanisms mentioned.

"The large differences in burst timing and spike patterning across the muscle heads suggest that the motor pools for each muscle receive distinct inputs. However, differences in the intrinsic physiological properties of motor units and neuromodulatory inputs across motor pools might also make substantial contributions to the structure of motor unit spike patterns (Martínez-Silva et al., 2018; Miles & Sillar, 2011)."

(6) *"We next examined whether the probabilistic recruitment of individual motor units in the triceps and elbow extensor muscle predicted stride-by-stride variations in elbow angle kinematics."*

I'm not sure that the wording is appropriate here. The analysis does not predict elbow angle variations from parameters extracted from the spiking activity. It rather compares the average elbow angle between two conditions (motor unit active or not active).

We thank the Reviewer for this comment and agree that the wording could be improved here to better reflect our analysis. To lower the strength of our claim, we replaced usage of the word 'predict' with 'correlates' in the above text and throughout the paper when discussing this result.

Methods:

(7) *"Using the four threads on the customizable Myomatrix array (RF-4x8-BHS-5), we implanted a combination of muscles in each mouse, sometimes using multiple threads within the same muscle. [...] Some mice also had threads simultaneously implanted in their ipsilateral or contralateral biceps brachii although no data from the biceps is presented in this study."*

A precise description of the localisation of the array (muscles and the number of arrays per muscle) for each animal would be appreciated.

(8) *"A total of 33 units were identified and manually verified across all animals." A precise description of the number of motor units concurrently identified per muscle and per animal would be appreciated. Moreover, please add details on the manual inspection. Does it involve the manual selection of missing spikes? What are the criteria for considering an identified motor unit as valid?*


As discussed earlier, we added Table 1 to the main text to provide the details mentioned in the above comments.

Regarding spike sorting, given the very large number of spikes recorded, we did not rely on manual adjusting mislabeled spikes. Instead, as described in the revised Methods section, we verified unit isolation by ensuring units had >98% of spikes outside of 1ms of each other. Moreover, as described above we have added new analyses (Figure 1–figure supplement 1) confirming the stability of motor unit waveforms across both the duration of individual recording sessions (roughly 30 minutes) and across the rapid changes in limb position within individual stride cycles (roughly 250 msec).

Reviewer #3 (Recommendations for the authors):

Figure 2 (and supplement) show spike count distributions with strong positive skewness, which is in accordance with the prediction of a fluctuation-driven regime. I suggest plotting these on a logarithmic x-axis (in addition to the linear axis), which should reveal a bell-shaped distribution, maybe even Gaussian, in a majority of the units.

We thank the Reviewer for the suggestion. We present the requested analysis below, which shows bell-shaped distributions for some (but not all) distributions. However, we believe that investigating why some replotted distributions are Gaussian and others are not falls beyond the scope of this paper, and likely requires a larger dataset than the one we were able to obtain.



Author response image 3. Spike count distributions for each motor unit on a logarithmic x-axis.

Why not more data? I tried to get an overview of how much data was collected.

Supplemental Figure 1 has all the isolated units, which amounts to 38 (are the colors the two muscle types?). Given there are 16 leads in each myomatrix, in two muscles, of six mice, this seems like a low yield. Could the authors comment on the reasons for this low yield?

Regarding motor unit yield, even with multiple electrodes per muscle and a robust sorting algorithm, we often isolated only a few units per muscle. This yield likely reflects two factors. First, because of the highly dynamic nature of locomotion and high levels of muscle contraction, isolating individual spikes reliably across different locomotor speeds is

inherently challenging, regardless of the algorithm being employed. Second, because the results of spike-train analyses can be highly sensitive to sorting errors, we have only included the motor units that we can sort with the highest possible confidence across thousands of strides.

Minor:

Figure captions especially Figure 6: The text is excessively long. Can the text be shortened?

We thank the Reviewer for this comment. Generally, we seek to include a description of the methods and results within the figure captions, but we concede that we can condense the information in some cases. In a number of cases, we have moved some of the descriptive text from the caption to the Methods section.

References

- Berg, R. W. (2017). Neuronal Population Activity in Spinal Motor Circuits: Greater Than the Sum of Its Parts. *Frontiers in Neural Circuits*, 11. <https://doi.org/10.3389/fncir.2017.00103> 
- Biewener, A. A., Blickhan, R., Perry, A. K., Heglund, N. C., & Taylor, C. R. (1988). Muscle Forces During Locomotion in Kangaroo Rats: Force Platform and Tendon Buckle Measurements Compared. *Journal of Experimental Biology*, 137(1), 191–205. <https://doi.org/10.1242/jeb.137.1.191> 
- Chung, B., Zia, M., Thomas, K. A., Michaels, J. A., Jacob, A., Pack, A., Williams, M. J., Nagapudi, K., Teng, L. H., Arrambide, E., Ouellette, L., Oey, N., Gibbs, R., Anschutz, P., Lu, J., Wu, Y., Kashefi, M., Oya, T., Kersten, R., ... Sober, S. J. (2023). Myomatrix arrays for high-definition muscle recording. *eLife*, 12, RP88551. <https://doi.org/10.7554/eLife.88551> 
- De Luca, C. J. (1985). Control properties of motor units. *Journal of Experimental Biology*, 115(1), 125–136. <https://doi.org/10.1242/jeb.115.1.125> 
- De Luca, C. J., & Erim, Z. (1994). Common drive of motor units in regulation of muscle force. *Trends in Neurosciences*, 17(7), 299–305. [https://doi.org/10.1016/0166-2236\(94\)90064-7](https://doi.org/10.1016/0166-2236(94)90064-7) 
- Farina, D., Negro, F., & Dideriksen, J. L. (2014). The effective neural drive to muscles is the common synaptic input to motor neurons. *The Journal of Physiology*, 592(16), 3427–3441. <https://doi.org/10.1113/jphysiol.2014.273581> 
- Hartigan, P. M. (1985). Algorithm AS 217: Computation of the Dip Statistic to Test for Unimodality. *Applied Statistics*, 34(3), 320. <https://doi.org/10.2307/2347485> 
- Henneman, E., Somjen, G., & Carpenter, D. O. (1965). FUNCTIONAL SIGNIFICANCE OF CELL SIZE IN SPINAL MOTONEURONS. *Journal of Neurophysiology*, 28(3), 560–580. <https://doi.org/10.1152/jn.1965.28.3.560> 
- Karabulut, D., Dogru, S. C., Lin, Y.-C., Pandy, M. G., Herzog, W., & Arslan, Y. Z. (2020). Direct Validation of Model-Predicted Muscle Forces in the Cat Hindlimb During Locomotion. *Journal of Biomechanical Engineering*, 142(5), 051014. <https://doi.org/10.1115/1.4045660> 
- Kim, J. J., Wyche, I. S., Olson, W., Lu, J., Bakir, M. S., Sober, S. J., & O'Connor, D. H. (2024). Myo-optogenetics: Optogenetic stimulation and electrical recording in skeletal muscles. <https://doi.org/10.1101/2024.06.21.600113> 
- Lu, J., Zia, M., Baig, D. A., Yan, G., Kim, J. J., Nagapudi, K., Anschutz, P., Oh, S., O'Connor, D., Sober, S. J., & Bakir, M. S. (2024). Opto-Myomatrix: μ LED integrated microelectrode arrays for optogenetic activation and electrical recording in muscle tissue. <https://doi.org/10.1101/2024.07.01.601601> 

- Manuel, M., & Heckman, C. J. (2011). Adult mouse motor units develop almost all of their force in the subprimary range: A new all-or-none strategy for force recruitment? *Journal of Neuroscience*, 31(42), 15188–15194. <https://doi.org/10.1523/JNEUROSCI.2893-11.2011> 
- Marshall, N. J., Glaser, J. I., Trautmann, E. M., Amematsro, E. A., Perkins, S. M., Shadlen, M. N., Abbott, L. F., Cunningham, J. P., & Churchland, M. M. (2022). Flexible neural control of motor units. *Nature Neuroscience*, 25(11), 1492–1504. <https://doi.org/10.1038/s41593-022-01165-8> 
- Martínez-Silva, M. de L., Imhoff-Manuel, R. D., Sharma, A., Heckman, C. J., Shneider, N. A., Roselli, F., Zytnicki, D., & Manuel, M. (2018). Hypoexcitability precedes denervation in the large fast-contracting motor units in two unrelated mouse models of ALS. *eLife*, 7(2007), 1–26. <https://doi.org/10.7554/eLife.30955> 
- Miles, G. B., & Sillar, K. T. (2011). Neuromodulation of Vertebrate Locomotor Control Networks. *Physiology*, 26(6), 393–411. <https://doi.org/10.1152/physiol.00013.2011> 
- Petersen, P. C., & Berg, R. W. (2016). Lognormal firing rate distribution reveals prominent fluctuation-driven regime in spinal motor networks. *eLife*, 5. <https://doi.org/10.7554/elife.18805> 
- Srivastava, K. H., Elemans, C. P. H., & Sober, S. J. (2015). Multifunctional and Context-Dependent Control of Vocal Acoustics by Individual Muscles. *The Journal of Neuroscience*, 35(42), 14183–14194. <https://doi.org/10.1523/JNEUROSCI.3610-14.2015> 
- Srivastava, K. H., Holmes, C. M., Vellema, M., Pack, A. R., Elemans, C. P. H., Nemenman, I., & Sober, S. J. (2017). Motor control by precisely timed spike patterns. *Proceedings of the National Academy of Sciences of the United States of America*, 114(5), 1171–1176. <https://doi.org/10.1073/pnas.1611734114> 
<https://doi.org/10.7554/eLife.105829.2.sa0>