

Transcriptional complexity in the insect central complex: single nuclei RNA-sequencing of adult brain neurons derived from type 2 neuroblasts

Reviewed Preprint

v2 • April 22, 2025

Revised by authors

Reviewed Preprint

v1 • February 27, 2025

Derek Epiney, Gonzalo N Morales Chaya, Noah R Dillon, Sen-Lin Lai , Chris Q Doe 

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, United States

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

This **important** study offers a molecular characterization of neurons and glia in the adult nervous system of the fruit fly *Drosophila melanogaster*. The study focuses on the progeny of a specific set of neural stem cells that contribute to the central complex, a conserved brain region that plays key roles in sensorimotor integration. The data are **convincing** and collected using validated methodology, generating an invaluable resource for future studies. The study will be of interest to developmental neurobiologists.

<https://doi.org/10.7554/eLife.105896.2.sa3>

Abstract

In both invertebrates such as *Drosophila* and vertebrates such as mouse or human, the brain contains the most diverse population of cell types of any tissue. It is generally accepted that transcriptional diversity is an early step in generating neuronal and glial diversity, followed by the establishment of a unique gene expression profile that determines morphology, connectivity, and function. In *Drosophila*, there are two types of neural stem cells, called Type 1 (T1) and Type 2 (T2) neuroblasts. In contrast to T1 neuroblasts, T2 neuroblasts generate intermediate neural progenitors (INPs) that expand the number and diversity of cell types. The diversity of T2-derived neurons contributes a large portion of the central complex (CX), a conserved brain region that plays a role in sensorimotor integration. Recent work has revealed much of the connectome of the CX, but how this connectome is assembled remains unclear. Mapping the transcriptional diversity of neurons derived from T2 neuroblasts is a necessary step in linking transcriptional profile to the assembly of the adult brain. Here we perform single nuclei RNA sequencing of T2 neuroblast-derived adult neurons and glia. We identify clusters containing all known classes of glia, clusters that are male/female enriched, and 161 neuron-specific clusters. We map neurotransmitter and neuropeptide expression and identify unique transcription factor combinatorial codes for each cluster (presumptive neuron subtype). This is a necessary step that directs functional studies to determine whether

each transcription factor combinatorial code specifies a distinct neuron type within the CX. We map several columnar neuron subtypes to distinct clusters and identify two neuronal classes (NPF+ and AstA+) that both map to two closely related clusters. Our data support the hypothesis that each transcriptional cluster represents one or a few closely related neuron subtypes.

Introduction

In all organisms, the brain has arguably the most complex cellular diversity, from human (Siletti et al., 2023 [↗](#)) to *Drosophila* (Davie et al., 2018 [↗](#); Franconville et al., 2018 [↗](#); Hulse et al., 2021 [↗](#)). Neuronal diversity is essential for the assembly and function of the adult brain, yet the “parts list” of different neuronal and glial cell types remains incomplete. In *Drosophila*, the laterally-positioned optic lobes have been well characterized for transcriptionally distinct neurons and glia (Konstantinides et al., 2018 [↗](#); Konstantinides et al., 2022 [↗](#); Özel et al., 2022 [↗](#)), as has the central brain and ventral nerve cord (Croset et al., 2018 [↗](#); Davie et al., 2018 [↗](#); Lago-Baldaia et al., 2023 [↗](#); Li et al., 2022 [↗](#); McLaughlin et al., 2021 [↗](#); Naidu et al., 2020 [↗](#); Nguyen et al., 2021 [↗](#); Sato and Suzuki, 2022 [↗](#); Shu et al., 2023 [↗](#); Velten et al., 2022 [↗](#)).

The central brain contains many diverse neurons that populate important neuropils, such as the mushroom body (MB) used for learning and memory (Sgammeglia and Sprecher, 2022 [↗](#)), or the central complex (CX) used for celestial navigation and sensory-motor integration (Fisher, 2022 [↗](#)), among other behaviors. The central brain neurons are all generated from neural stem cells, called neuroblasts (NBs). There are two types of NBs that generate central brain neurons: Type 1 (T1) and Type 2 (T2) NBs. T1 NBs undergo asymmetric divisions to self-renew and generate a ganglion mother cells (GMCs), which each produce two post-mitotic neurons (Pollington et al., 2023 [↗](#); Yu et al., 2010 [↗](#); Yu et al., 2013 [↗](#)). There are ~100 T1 NBs per larval central brain lobe that each generate 20-100 neurons and glia (Ito et al., 2013 [↗](#); Yu et al., 2013 [↗](#)). In addition, there are Type 0 (T0) NBs which generate post-mitotic neuron progeny (Baumgardt et al., 2014 [↗](#)); T0 NB lineages yet to be documented in the central brain. T2 NBs have a more complex division pattern than T1 NBs (Bello et al., 2008 [↗](#); Boone and Doe, 2008 [↗](#); Bowman et al., 2008 [↗](#)). T2 NBs undergo asymmetric division to self-renew and generate an intermediate neural progenitor (INP); each INP undergoes 4-6 divisions to self-renew and generate a GMC and its two neuron or glial progeny (Bello et al., 2008 [↗](#); Boone and Doe, 2008 [↗](#); Bowman et al., 2008 [↗](#)). Thus, each T2 NB division will generate 8-12 progeny cells. Most T2 NBs generate ~500 neurons (Yu et al., 2013 [↗](#)), have highly complex neuronal cell types based on morphology and connectivity (Hulse et al., 2020 [↗](#); Wang et al., 2014 [↗](#); Yang et al., 2013 [↗](#); Yu et al., 2013 [↗](#)), and produce the columnar neurons of the adult CX (Boyan and Williams, 2011 [↗](#); Kandimalla et al., 2023 [↗](#)). Subsequently, we will call neurons born from type 1 NBs “T1-derived” and neurons born from type 2 NBs “T2-derived.”

Although single cell RNA-seq (scRNA-seq) has been done on adult central brain neurons and glia (Croset et al., 2018 [↗](#)), the transcriptomic profile of T1 versus T2 neuronal and glial progeny has not yet been characterized. Furthermore, the T2 lineages generate diverse neuronal progeny (Hulse et al., 2020 [↗](#); Wang et al., 2014 [↗](#); Yang et al., 2013 [↗](#); Yu et al., 2013 [↗](#)) but focused transcriptional profiling of T2-derived progeny has not been reported. In this report, we generate a single nuclei (snRNA-seq) atlas of T2-derived progeny of the adult central brain. We use several complementary methods to link neuron identity to UMAP clusters. Our data will facilitate linking neuronal transcriptomes with connectomes to gain a molecular understanding of CX development, connectivity, and behavior.

Results

Generation of a transcriptomic atlas of adult central brain neurons and glia

The adult central brain (i.e. brain without optic lobes) is composed of neuronal and glial progeny derived from both T1 and T2 NBs. We first generate a transcriptomic atlas of all central brain neurons derived from both T1 and T2 NBs. The flies were of a genotype that gave permanent lineage tracing of T2 NB-derived progeny in the adult (**Figure 1A** [↗](#), A'). This allowed us to first analyze all central brain T1- and T2-derived neurons and glia (**Figure 1** [↗](#)) and then focus on neurons and glia produced specifically by T2-derived lineages.

We dissociated nuclei from 7-day-old adult fly central brains with optic lobes removed using methods previously described (Li et al., 2022 [↗](#); McLaughlin et al., 2021 [↗](#)). The cDNA libraries were then prepared from dissociated nuclei with split-pool-based barcoding for single-nuclei transcription profiling (Rosenberg et al., 2018 [↗](#)). We sequenced a total of 30,699 nuclei at 589 median genes per nuclei from the T1+T2 central brain.

To distinguish T2-derived progeny from T1-derived progeny, we expressed *worniu-gal4, asense-gal80* to drive expression of FLP recombinase (FLP) specifically in T2 NBs. This resulted in the flip out of a stop cassette and thus continuous expression of Gal4 under the ubiquitous *actin5C* enhancer. After removal of the stop cassette in T2 NBs, the *actin5c-gal4* can continuously drive expression of the reporter genes (RFP or GFP) and FLP in the T2-derived neuronal and glial progeny in adult brains (**Figure 1A** [↗](#)). We assigned 3,125 nuclei which have expression of RFP, GFP, or FLP as being derived from T2 lineages. We assigned 27,574 triple-negative nuclei as derived from the T1 lineages. We used Seurat for filtering, integrating, and clustering the atlas to provide transcriptionally unique cell clusters (Stuart et al., 2019 [↗](#)). We used uniform manifold approximation and projection (UMAP)-based dimension reduction to visualize the 152 clusters (**Figure 1B** [↗](#)). The most enriched or least expressed genes (marker genes) for each cluster are shown in Supp Table 1.

We used known markers (Supp Table 2) (Croset et al., 2018 [↗](#); Davie et al., 2018 [↗](#)) to identify distinct cell types in the central brain, including glia, mushroom body neurons, olfactory projection neurons, clock neurons, Poxn⁺ neurons, serotonergic neurons, dopaminergic neurons, octopaminergic neurons, corazonergic neurons, hemocytes, and ocelli (**Figure 1B** [↗](#); Supp Table 1). We didn't observe any cluster exclusively containing progeny from T1 or T2 NB lineages (**Figure 1C** [↗](#), Supp Figure 1). We conclude that both T1 and T2 generate transcriptionally similar cells, despite their different developmental origins. Alternatively, deeper sequencing may resolve a single cluster into two clusters with each derived from T1 or T2 lineages. We next searched for cluster-defining genes in the T2 enriched clusters (**Figure 1C** [↗](#)) and found high expression of *AstA* (Allatostatin A), *SerT* (Serotonin transporter), *Tk* (Tachykinin), and *Vmat* (Vesicular monoamine transporter) (**Figure 1D** [↗](#)). These enriched marker genes suggest that *AstA*⁺ neurons, serotonergic neurons, and *TK*⁺ neurons may be primarily derived from T2 lineages.

We explored the central brain atlas for glial cell types and their gene expression. We focused on the 3,409 glial nuclei from clusters that expressed the pan-glial marker *repo* (Campbell et al., 1994 [↗](#); Xiong et al., 1994 [↗](#)) in the T1+T2 atlas (**Figure 1E** [↗](#)). Each cluster contained a mix of T1 and T2 glial progeny (**Figure 1E** [↗](#)). We identified six known glial cell types (astrocytes, cortex, ensheathing, surface glial and the two subtypes: perineurial and subperineurial) based on canonical glial makers (**Figure 1F** [↗](#); Supp Table 3). Similar to a previous glial cell atlas (Lago-Baldaia et al., 2023 [↗](#)), we found some glial subtypes (astrocytes, ensheathing, and subperineurial) mapping to multiple clusters (**Figure 1E** [↗](#), 1F). We identified a glial cluster that expressed genes




Figure 1.

Cell atlas of central brain with single nuclei RNAseq.

(A-A') Genetics (A) to label progeny (A') derived from T2 NBs. Dashed lines show the boundary of optic lobes and central brain, and the optic lobes were removed during dissection.

(B) Central brain atlas labeled with known cell types. Abbreviations: CLK, clock neurons; CRZ, Corazonergic neurons; DOP, dopaminergic neurons; HEM, hemocytes; MBN, mushroom body neurons; OCTY, octopaminergic-tyraminergerg neurons; OC, ocelli; OL, optic lobe; OPN, olfactory projection neurons; SER: serotoninergerg neurons.

(C) Central brain atlas labeled by NB (T1 or T2) lineage. Dash line-outlined box shows the region enriched with the cells derived from T2 NBs, and the identity are shown at the bottom-right box.

(D) Dot plot of top 5 marker genes of the T2-enriched clusters.

(E) Atlas of central brain glia labeled with known cell types. **(E')** T1 and T2 derived cells colored cyan and red respectively.

(F) Dot plot of known and top marker genes of glial clusters.

associated with extracellular matrix, *vkg* and *Col4a1* (**Figure 1F**), which have previously been identified as pan surface glial markers (DeSalvo et al., 2014; Hindle and Bainton, 2014). Interestingly, the two surface glial subtypes, perineurial and subperineurial clusters, do not express these markers at cluster defining levels, and conversely the surface glia cluster 5 does not express perineurial or subperineurial specific markers (**Figure 1F**, Supp Table 4). Differential gene expression analysis for all genes between T1 and T2 glial progeny did not show differences across any glial cell types or clusters (Supp Table 5). We conclude that the adult central brain contains known glial cell types with no differences in gene expression between T1 and T2-derived glia.

Generation of a T1- and T2-specific cell atlas

We explored the diversity of T1 and T2-derived neurons by generating T1- and T2-specific cell atlases. We identified T1-derived neurons by bioinformatically *excluding* cells co-expressing T2-specific markers *FLP+/GFP+/RFP+* plus *repo+* glial clusters. We then generate a T1 neuron atlas containing 22,807 T1-derived neurons that form 114 clusters. Marker genes of each cluster are shown in Supp Table 5. We identified the neurons that are known to be generated by T1 NBs, including MB neurons and olfactory projection neurons (Croset et al., 2018; Davie et al., 2018; Li et al., 2022)(Supp Table 2). We identified other T1-derived neurons, including clock neurons, Poxn⁺ neurons, and neurons that release dopamine, serotonin, octopamine/tyramine, and other neuropeptides (Supp Figure 2, Supp Table 6) (Croset et al., 2018; Davie et al., 2018). We gathered a list of genes that represented the 10 most enriched genes from each cluster and calculated the scaled averaged expression of each gene from each cluster (Supp Figure 2). Each unique combination of enriched genes could be referred to as cluster markers. We conclude that our T1-derived nuclei contain the expected abundance of neuronal diversity seen in previous scRNA-seq atlases (Croset et al., 2018; Davie et al., 2018; Li et al., 2022).

In our whole brain atlas described above, T2-derived cells represent a minor contribution to the atlas due to their low nuclei numbers relative to the T1-derived neurons (Raji and Potter, 2021). To generate a comprehensive T2 atlas, we needed to increase the percentage of T2-derived nuclei for RNAseq. We used fluorescence-activated cell sorting (FACS) to select nuclei that are labeled by T2-specific permanent lineage tracing (**Figure 1A,B**). We then selected the nuclei *in silico* that showed expression of UAS-transgenes (FLP, GFP, or RFP, see above) to eliminate contamination of T1 nuclei during FACS. In total, we sequenced 61,118 T2 nuclei. We included 3,125 UAS-transgene labeled nuclei from the dissociated central brain without sorting from the T1+T2 atlas (see above). We used integration in Seurat to generate an atlas containing 64,243 nuclei that formed 198 clusters. T2 NBs have been estimated to produced ~5000 neurons/glia in the central brain (Ito et al., 2013; Yang et al., 2013), with ~1,800 neurons in the central complex (Hulse et al., 2020; Schlegel et al., 2024), giving our atlas ~12x coverage. We next filtered out the *repo+* glial cell clusters to generate a T2 neuron cell atlas with 50,148 non-glial nuclei forming 161 clusters (**Figure 2A**). Marker genes of each cluster are shown in Supp Table 8.

We calculated the scaled average expression of the top 10 most enriched genes from each T2 neuron cluster (Supp Table 8). Each set of genes could serve as cluster marker genes (**Figure 2B**). We conclude that each cluster within our T2 neuron atlas represents a transcriptionally unique cell type. For the remainder of the paper, we focus solely on the diversity of T2-derived neurons and glia.

T2 neuroblasts generate all major classes of fast-acting neurotransmitters

An important aspect of neuronal identity and function is fast-acting neurotransmitter expression. We determined the neuronal populations that expressed seven fast-acting neurotransmitters: glutamatergic, cholinergic, GABAergic, tyraminerbic, dopaminergic, serotonergic and octopaminergic in T2-derived neurons (**Figure 3A-G**). In neurons, we found that cholinergic




Figure 2.

Cell atlas of neurons produced by T1 or T2 NBs.

(A) Cell atlas from T2 NBs. **(B)** Heatmap of scaled average expression of top 10 markers genes from each T2 neuronal cluster.

neurons were most abundant (21%), followed by glutamatergic neurons (12%), GABAergic neurons (9%), tyraminergetic neurons (2.6%), dopaminergic neurons (1.2%), serotonergic neurons (1.2%), and octopaminergic neurons (0.7%). Additionally, 12% of neurons were co-expressing 2 or 3 neurotransmitters. The remaining 40% in our atlas were not expressing any neurotransmitter with a log normalized expression <2 (**Figure 3H**); this is similar to the ratios in the adult midbrain (Croset et al., 2018) and optic lobes (Konstantinides et al., 2022). The nuclei co-expressing two or more fast-acting neurotransmitters may reflect an authentic feature of these neurons, as seen in other systems (see Discussion). Alternatively, our double- and triple-positive neurons may be false positives (see Discussion).

T2 neuroblasts generate all major classes of glia

We next explored the complete T2 atlas for glial cell types and their gene expression. We sub-clustered from the T2 atlas for 12,315 glial nuclei from clusters that expressed the pan glial marker *repo* (**Figure 4A**) (Campbell et al., 1994; Xiong et al., 1994). Similar to our T1+T2 glial atlas, we identified glial cell types: astrocytes, cortex, ensheathing, astrocyte-like, and two surface glial subtypes, perineurial and subperineurial, based on canonical glial markers (**Figure 4B**; Supp Table 9). In line with our T1+T2 atlas and previous glia cell atlas (Lago-Baldaia et al., 2023), some subtypes mapped to several subclusters including ensheathing, astrocytes, and chiasm (**Figure 4A-B**). Interestingly, we identified chiasm glia that may represent the migratory chiasm glia generated from the DL1 lineage (Viktorin et al., 2013; see Discussion). We conclude that our T2 atlas contains the expected glial cell types.

T2 neuroblasts generate sex-biased cell types

When generating our atlas, we dissected both female and male adult brains that were processed as separate samples in order to differentiate the sex of nuclei in the snRNAseq atlas. We noticed that several cell types in the T2 glial atlas showed unequal number of male and female nuclei (**Figure 5A-B**, E-F). We identified sex-biased clusters by normalizing the number of input nuclei between male and female samples and compared the proportion of nuclei by sex within each cluster to identify sex-biased clusters (**Figure 5B,F**). We found that the T2 glial atlas contained two female enriched clusters: ensheathing/astrocyte (cluster 7) and chiasm (cluster 5), and one male enriched cluster: the astrocyte-like (cluster 3) (**Figure 5B**). We determined differential genes expressed between male and female nuclei across all glia (**Figure 5C**; Supp Table 10). We found female nuclei expressed higher levels of genes including the female-specific genes *yp1*, *yp2*, and *yp3* (**Figure 5C**) (Warren et al., 1979). Additionally, female nuclei were enriched for *dsx* (Supp Table 10). Male glial nuclei expressed higher levels of genes including the male-specific genes *lncRNA:rox1/2* and *fru* (**Figure 5C**; Supp Table 10) (Amrein and Axel, 1997; Meller et al., 1997; Ryner et al., 1996). We next looked at expression within the sex-biased clusters. We found similar results that male nuclei expressed high levels of the male-specific *lncRNA:rox1/2* and low expression of the female-specific genes *yp1*, *yp2*, and *yp3* compared to the female nuclei (**Figure 5D**; see Discussion). We conclude that male and female adult T2 glia have sex-specific differences in gene expression within the same glial cell type.

We next explored sex differences among neurons (**Figure 5E-E'**). We identified 14 neuronal clusters with disproportionate amounts of male and female nuclei (**Figure 5F**). We first tested the differential genes expressed between male and female nuclei across all T2-derived neurons (**Figure 5G**; Supp Table 11). Similar to the T2 glia sex differences, we found female nuclei expressed higher levels of the female-specific genes *yp1*, *yp2*, and *yp3* (**Figure 5G**) (Warren et al., 1979). Additionally, female nuclei were enriched for *dsx* (Supp Table 11). Male neuronal nuclei expressed higher levels of the male-specific genes *lncRNA:rox1/2* and *fru* (**Figure 5G**; Supp Table 11) (Amrein and Axel, 1997; Meller et al., 1997; Ryner et al., 1996). We next looked at expression levels within the sex-biased clusters. We found similar results that male nuclei expressed high levels of the male-specific *lncRNA:rox1/2* and low expression of the female-specific




Figure 3.

Expression of fast-acting neurotransmitters in neurons derived from T2NSC lineages.

(A-G) UMAP distribution plots demonstrate the expression of the following neurotransmitters: (A) *vesicular glutamate transporter* (VGlut, glutamatergic neurons), (B) *vesicular acetylcholine transporter* (VAcHT, cholinergic neurons), (C) *glutamic acid decarboxylase 1* (Gad1, GABAergic neurons), (D) *tyramine β hydroxylase* (Tbh, tyraminergetic neurons), (E) *tyrosine 3-monooxygenase* (Ple, dopaminergic neurons), (F) *serotonin transporter* (SerT, serotonergic neurons) (G) *tyrosine decarboxylase* (Tdc2, octopaminergic neurons). All plots have a minimum cutoff value set at 0. (H) The UpSet quantifies the number of cells in the atlas that express each neurotransmitter gene with a scaled expression > 2 (Gu, 2022 [DOI](#)).




Figure 4.

Glial cell types present in the T2-derived cell atlas.

(A) Sub-clustered of 12,315 nuclei from Repo+ T2 clusters in UMAP distribution.

(B) Dot plot of validated glial subtype markers to identify glial clusters by differential gene expression.




Figure 5.

Sex differences present in the T2-derived glia and neurons.

- (A-A'') T2 glia in UMAP distribution across samples: (A) T2 Female (3,910 nuclei), (A') T2 Male (4,385 nuclei), (A'') Female and Male mixed (4,020 nuclei).
- (B) Biased clusters for male to female ratio for number of nuclei in the T2 glia clusters.
- (C) Differential expression between male and female T2 glia.
- (D) Heatmap of top differential gene expression between male and female nuclei within glial clusters.
- (E-E'') T2 neuron in UMAP distribution across samples: (E) T2 Female (8,151 nuclei), (E') T2 Male (16,201 nuclei), (E'') Female and Male mixed (25,796 nuclei).
- (F) Biased clusters for male to female ratio for number of nuclei in the T2 neuron clusters.
- (G) Differential expression between male and female T2 neurons.
- (H) Heatmap of top differential gene expression between male and female nuclei within neuron clusters.

genes *yp1*, *yp2*, and *yp3* compared to the female nuclei (**Figure 5H** [↗](#); see Discussion). We conclude that male and female adult T2 neurons have sex-specific differences in gene expression within the same neuronal subtype.

T2 neuroblasts generate neurons expressing a diverse array of neuropeptides

Neuropeptides are often used as markers to distinguish different neuronal populations. We examined whether individual neuropeptide-encoding genes were exclusively expressed within single clusters, representing single neuronal subtypes. Among the 49 neuropeptides in *Drosophila*, we identified 13 with enriched expression in a limited number of clusters, making them suitable as cluster-defining markers (**Figure 6A** [↗](#)). For all clusters significantly expressing cluster-defining neuropeptides, we assessed their co-expression with specific fast-acting neurotransmitters (**Figure 6A** [↗](#)). We found that all neuropeptide-expressing clusters, co-express one or more neurotransmitters (see **Figure 3H** [↗](#)).

These data raise several questions. What is the relationship between neuropeptide expression and cluster identity? Can two distinct clusters express the same neuropeptide? Can two neuropeptides be co-expressed by the same cluster? Our data show that all of these patterns exist. For example, in some cases, one cluster expresses two or more neuropeptide genes (e.g. clusters 100 and 128 express NPF and AstA); conversely, we also observe one neuropeptide expressed in multiple clusters (e.g. Ms, 2 clusters; Tk, 2 clusters) (**Figure 6A** [↗](#)).

Next, we wanted to validate whether these neuropeptidergic cells indeed come from T2 lineages and target the CX. This required a method for specifically labelling of T2 peptidergic neurons, as antibody staining could be contaminated with T1-derived neurons. We developed a genetic approach to selectively visualize neuropeptide expression exclusively in neurons originating from T2 NB lineages. This technique involves driving FLP recombinase (FLP) under the control of the T2-specific driver *wor-GAL4, ase-GAL80*. The T2-specific FLP catalyzes the excision of a stop codon in the *lexAop-FRT-stop-FRT-myr::gfp* transgene, allowing existing neuropeptide-2A-LexA transgenes (Deng et al., 2019 [↗](#)) to drive GFP expression specifically in neuropeptide-positive T2-derived neurons (**Figure 6B** [↗](#)). This method revealed the morphology of specific T2-derived peptidergic neurons.

We first assayed Ms+ neurons because Wolff et al. reported that at least two FB neurons, FB4Z and FB5R express Ms and they target to two distinct FB layers (Wolff et al., 2025 [↗](#)). However, we found that type II-derived Ms-2A-LexA-expressing neurons project to multiple layers of the dorsal fan-shaped body and the entire ellipsoid body, suggesting an unknown class of Ms+ neurons targeting to EB and/or FB (**Figure 6C-C** [↗](#), Supp Movie 1). In our atlas, Ms is mainly expressed in clusters 157 and 160; however, we cannot distinguish the corresponding identity of each neuron.

Next, we assayed neurons in two clusters (100 and 128) that express the same two neuropeptides, AstA and NPF (**Figure 6D-E** [↗](#)). We wanted to determine if these clusters contain two cell types with similar gene expression, or a single cell type that co-expressed both neuropeptides. We used NPF-2A-lexA and AstA-2A-lexA to label each class of neurons. We found that they were generated at indistinguishable numbers of cells and projections: both had ~20 neurons projecting to a lateral domain of the ellipsoid body and the same two layers of the fan-shaped body (**Figure 6D-E** [↗](#); Supp Movies 2-3). Whether these T2-derived NPF and AstA neurons are indeed the same would require antibody validation.

We investigated whether specific transcription factors (TFs), or TF combinatorial codes, might correlate with the expression of cluster-defining neuropeptides, and thus provide candidates for a regulatory relationship where each cluster may have a TF code that drives transcription of a specific neuropeptides. To address this, we performed an unbiased correlation analysis of TF and




Figure 6.

Neuropeptide expression in T2-derived neurons.

(A) Dot plot showing expression of 13 neuropeptides and their co-expression with the 7 fast-acting neurotransmitters (red dashed line) across selected clusters.

(B) Genetic scheme to map neuropeptide expression in T2-derived adult neurons.

(C-E) Neuropeptide-expressing neurons labelled with GFP in three-dimensional projections. **(C'-E')** Fan-shaped body projections. **(C''-E'')** Ellipsoid body projections. nc82 counterstains (magenta) in the brain for neuropil projections.

(F) Heatmap showing the top 5 transcription factors most strongly correlated with each neuropeptide across all cells (Sigorelli, 2024). Scale bar represents 20 μ m.

neuropeptide expression patterns across all cells in our dataset to identify potential regulatory relationships. Interestingly, we found high correlations of multiple TFs to each neuropeptide (**Figure 6F**, Supp Fig 3, Supp Table 12). Our data provide a stepping-stone to the analysis of upstream TF combinatorial codes that drive neuropeptide expression.

Each neuron cluster consists of a unique combination of transcription factors

It is generally thought that neuronal diversity is generated by TF combinatorial codes that are uniquely expressed in neuronal subtypes, with an emphasis on homeodomain TFs (Reilly et al., 2020; Sagner et al., 2021). Thus, we assayed each class of TFs to determine how many were expressed in cluster-specific (i.e. neuron-specific) patterns. We found that most clusters expressed a unique TF combination (**Figure 7**). This is true for all zinc-finger TFs (**Figure 7A**) and helix-turn-helix TFs (**Figure 7B**). Nearly all clusters expressed a unique combination of homeodomain TFs (**Figure 7C**), with only a few clusters sharing a homeodomain code (**Figure 7C**). In contrast, basic domain, unspecified domain, and high mobility group TFs were more promiscuously expressed (**Figure 7D**). We conclude that zinc finger, helix-turn-helix, and homeodomain TFs are good candidates for forming unique combinatorial codes that should be tested for a role in generating neuronal diversity.

Linking neurons to UMAP clusters

We sought to link well characterized CX neurons to their transcriptomes within the T2 atlas. We chose columnar neurons due to known cell type markers and genetic access. We used four complementary approaches to identify them, described below.

1. We assigned identities of neuronal clusters based on the expression profiles of neuropeptides and neurotransmitters (**Figure 6**) characterized in the literature (Wolff et al., 2025). We further examined these neurons of their expression of selective TFs and markers with antibody staining, including *bsh*, *cut*, *DIP-y*, *Eip93F*, *Imp*, *runt*, *Syp*, *toy*, and *zfh2*. The results are summarized in Supp Table 13.
2. We used a published bulk RNAseq dataset to profile selective CX neurons and compared the expression profiles to our snRNA-seq dataset with a published algorithm (Davis et al., 2020) (**Figure 8A**). We generated a heatmap based on the coefficient of the expression profiles to predict the identity of each cluster. We predicted that E-PG neurons were in cluster 75 and 95, P-EG neurons were in cluster 102, hDeltaK neurons were in cluster 100, FB6A were in cluster 162, and FB7A were in cluster 40 (**Figure 8A**). The prediction of E-PG neurons arising from cluster 75 and 95 was validated by expression of the transcription factor *dac* (Dillon and Doe, 2024); other neuron-cluster assignments await experimental validation.
3. We assayed the enhancers in highly specific split-Gal4 lines to determine if the associated genes were expressed in the same neuron subtype as the split-Gal4 line (**Figure 8B**). For example, the stable split line SS65380 is expressed in the FB2A neurons, and the enhancer DNA used to make the split lines are associated with the *wnt10* and *lmpt* genes. These two genes show high expression in cluster 149 (**Figure 8B**), allowing us to predict that the FB2A neurons are in cluster 149. A similar approach was used to generate candidate neuron-cluster predictions for the following neurons: FB8G, FB6H, hDeltaA, hDeltaD, hDeltaH, and FB1C (**Figure 8B**). These predictions can be used to focus experimental validation on a small group of neurons; however, the percentage of validated neurons-clusters awaits experimental analysis.




Figure 7.

Combinations of transcription factor expression in T2-derived neurons.

Binarized heatmaps of positive **(A)** zinc finger **(B)** helix-turn-helix **(C)** homeodomain TF markers in T2-derived neurons. Clusters are sorted by similarity based on Jaccard index scores.

(C) Top five marker genes for clusters which had non-unique combinations of homeodomain TF expression.

(D) Percentage of clusters with unique combinations of TF expression based on TF class (zinc finger TFs = 100% (161 unique clusters), helix-turn-helix TFs = 100% (161 unique clusters), homeodomain TFs = 93.8% (150 unique clusters), basic domain TFs = 69.6% (112 unique clusters), unidentified DNA binding domain TFs = 59% (95 unique clusters), high motility group TFs = 34.2% (55 unique clusters).

4. The fourth approach was based on using known or novel markers of CX neurons together with Gal4/LexA lines that are regulated by known gene enhancers. TFs known to be expressed by PF-R, P-EN and P-FN neurons are Toy and Runt, respectively (Sullivan et al., 2019). We assigned PF-R neurons to cluster 105 based on being Toy+ Runt- and being labeled by Gal4/LexA driven by the enhancer of *rho* gene (Figure 8C, left and middle columns). We assigned P-FN to clusters 38, 46, and 123 based on being Toy-Runt+ and being labeled by split line SS00191 associated with *shakB* and *Pkc53E* genes. Lastly, we assigned P-EN neurons to clusters 66 and 130 by their expression of Toy-Runt+ and labeled by *Gψ30A-lexA* (Figure 8C, left and middle columns). This approach identified novel TF markers for P-EN, PF-R and P-FN neurons (Figure 8C, right columns), which we validated by antibody staining (Figure 8D-H) (Dillon et al., 2024). The results of cluster identity assignment are summarized in Figure 9 (Supp Table 14).

The differential expression of TFs probably contributes to the morphological or connectivity diversity found within each class of CX neurons (Hulse et al., 2021; Turner-Evans et al., 2020; Wolff and Rubin, 2018; Wolff et al., 2015). We conclude that our T2 atlas can be used to link known neuron subtypes to their transcriptome (Figure 9). This is a necessary step in determining how cluster-specific gene expression regulates neuron-specific functional properties such as morphology, connectivity, and physiology.

Discussion

We used a genetic approach to express RFP in the progeny of T2 NBs in the adult central brain. We excluded optic lobes by manual dissection, and thus our data are similar to recent scRNA-seq data from the central brain (Croset et al., 2018; Davie et al., 2018; Li et al., 2022). Similarities include neurons expressing predominantly single fast-acting neurotransmitters (discussed below). Both datasets show major classes of neurons including MB neurons, olfactory projection neurons, clock neurons, Poxn+ neurons, serotonergic neurons, dopaminergic neurons, octopaminergic neurons, corazoninergic neurons, and hemocytes. The MB neurons that appear in the T2-derived progeny may be due to off-target expression of our Gal4 driver at stages we have not assayed, or more likely, due to ambient RNA released during the dissociation and sorting steps.

Interestingly, we observed fewer distinct neuronal clusters in the T1-derived population (114 clusters; Supp Figure 2) than in the T2-derived neuronal population (161 clusters; Figure 2). This could be due to the T1-derived progeny containing a large population of transcriptionally similar Kenyon cells of the MB, whereas T2-derived clusters contain fewer, but transcriptionally more diverse, cells than T1-derived progeny. It would be interesting to know whether T1 NB lineages are generally less diverse, or whether the Kenyon cells are exceptional in their lack of transcriptional diversity, despite the presence of protein gradient in young vs. old neurons (Liu et al., 2015). Our results are consistent with lineage analysis of T2 NBs, which make distinct neurons across the temporal axis of T2 NBs, plus unique neurons across the INP temporal axis (Doe, 2017; Ito et al., 2013; Sullivan et al., 2019; Wang et al., 2014; Yang et al., 2013).

Neurotransmitters and neuropeptides

We observed that most fast-acting neurotransmitters are uniquely expressed in neuronal populations, but we also observe a smaller group of neurons that express two or three neurotransmitters. This includes excitatory and inhibitory neurons, e.g. 2.9% of all neurons express genes necessary for cholinergic and GABAergic neurotransmitters (Figure 3). Co-expression of neurotransmitters has been reported in other systems (Granger et al., 2020; Granger et al., 2023; Lozovaya et al., 2018; Meye et al., 2016; Saunders et al., 2015; Shabel et al., 2014; Spitzer, 2015; Takács et al., 2018) and it may be biologically relevant in the T2-derived neurons.




Figure 8.

Mapping T2-derived neurons to UMAP clusters.

(A) Heatmap of the coefficients (r) of expression profiles of single cell clusters (columns) to the expressions of known central complex neurons (rows) profiled with bulk RNA sequencing. The matching of the cluster to the central complex neurons is determined by the highest coefficient value (gray box) for each cluster. FB6A(1) and FB6A(2) are labeled and profiled with two different split-gal4 drivers (Wolff et al., 2025). Cluster 75 and 95 identified as E-PG were verified in Dillon et al., 2024.

(B) Scaled average expression of the enhancer genes of split-gal4 drivers (columns) in each cluster (rows). The name of central complex neurons (CX) labeled by split-gal4 drivers are listed at the top. The cluster identity was determined by the combination of positive scaled average expression of the enhancer genes (black boxes).

(C) Dotplot of the known marker genes (*toy*, *runt*) expressed in central complex neurons (PF-R, P-FN, and P-EN), *lexA*-driver enhancer gene (*rho*, *Gy30A*), split-gal4-driver (SS00191) enhancer genes (*shakB*, *Pkc53E*), and newly identified marker genes.

(D) PF-R (*R37G12-Gal4 UAS-V5*) neuronal cell bodies labeled with V5, and co-stained with marker genes boxed in (C).

(E,F) PF-N (*R16D01-Gal4 UAS-V5*) neuronal cell bodies labeled with V5, and co-stained with marker genes boxed in (C).

(G) PE-N (*R12D09-Gal4 UAS-V5*) neuronal cell bodies labeled with V5, and co-stained with marker genes boxed in (C).

(H) Percentage of marker genes in PE-N and PF-N. Scale bar: 5 μ m in all panels.

Neuropeptides are profoundly important for a vast array of behaviors (Schoofs et al., 2017), and thus our mapping of neuropeptide expression to distinct neuronal subtypes may facilitate a functional analysis linking neuron identity, neuropeptide expression, and behavior.

Glial identities

We identified six classes of glia in the T1+T2 glial atlas that reflects previous scRNAseq datasets (Figure 1E-F) (Konstantinides et al., 2018; Lago-Baldaia et al., 2023). Similar to a recent glial cell atlas (Lago-Baldaia et al., 2023), we found glial subtypes like astrocytes, ensheathing, and subperineurial glia mapped to several clusters (Figure 1E-F). It remains unclear if these clusters with the same cell type annotation represent distinct glial identities or different transcriptional states within these populations. Interestingly, we detected a small population of *vkg*+ surface glia specific to the whole atlas (T1+T2) and not the T2 glia atlas. The lack of differential gene expression between T1- and T2-derived glia suggests these cell types are similar despite different developmental origins. It will be for future work to understand how different progenitors produce similar glial fates.

Similar to the T1+T2 glial atlas, our T2 atlas captured the known glial subtypes found in other scRNAseq datasets (Figure 4) (Konstantinides et al., 2018; Lago-Baldaia et al., 2023). The T2 glia atlas contains more clusters than the T1+T2 atlas that may be due to the nearly 4x the number of glia captured (T1+T2 atlas 3,409 nuclei; T2 atlas – 12,315 nuclei).

Interestingly, we identified the astrocyte-like glia of the central brain previously described (Awasaki et al., 2008). Additionally, we captured chiasm glia that have been shown to be derived from the T2 lineage DL1 and migrate into the optic lobe (Viktorin et al., 2013). These nuclei could represent either chiasm glia that did not migrate out of the central brain or optic lobe tissue that was not completely removed during dissections. It will be interesting to investigate how these T2-derived migratory chiasm glia compare to the optic-lobe derived chiasm glia, because of their different developmental origins.

Sex differences found in the adult neurons and glia

The T2 glia and neuron atlases contained clusters with disproportionate enrichment of female and male nuclei when normalized for sample input. We found the expected differential expression of yolk protein transcripts (*yp1*, *yp2*, *yp3*) enriched in female nuclei and the long non-coding RNAs *rox1/2* and *fru* enriched in male nuclei (Amrein and Axel, 1997; Meller et al., 1997; Ryner et al., 1996). Interestingly, we found *dsx* to be enriched in both glial and neuronal female nuclei. Surprisingly, we found additional female and male-specific gene enriched in both neurons and glia including *ATPsyndelta* and undescribed, computationally predicted genes (CGs) (Figure 5). It remains to be determined if these genes are driving sex-specific differences within glial and neuronal subtypes. These genes may reflect sex-specific differences in the adult central brain and may provide insight into how behavioral circuits are linked to sex-specific behaviors. Future work should aim to characterize and test these genes for functional roles.

TF codes

Our atlas showed zinc finger TFs and homeodomain TFs as forming combinatorial codes specific for distinct cell types. This raises the possibility that these TF combinations determine neuronal functional properties, by establishing and maintaining neuronal identities. This is consistent with data from *C. elegans* (Hobert, 2021; Reilly et al., 2020), mammalian spinal cord (Briscoe et al., 2000; Sagner et al., 2021), *Drosophila* optic lobe (Holguera and Desplan, 2018; Konstantinides et al., 2018), leg motor neurons (Baek et al., 2013; Enriquez et al., 2015), and VNC (Soffers et al., 2024). Interestingly, we found more zinc finger TFs expressed in unique combinatorial codes compared to homeodomain TFs; this may be a caution to focusing too narrowly on homeodomain TFs function. We note that our results are mostly correlative and

await functional analysis of the TF combinatorial codes found in our dataset. It is also possible that the current sparse TF expression patterns are due, in part, to false negative results due to low read depths. This can be resolved by increasing read depth or by performing functional assays.

Mapping neurons to clusters

In this study, we used several complementary approaches to map identified neurons to their transcriptomic UMAP cluster. First, we used bulk-seq data from specific neurons with our atlas to match neuron to cluster. Second, we used enhancer elements from neuron-specific split Gal4 lines to link enhancer-associated genes to each hemi-driver with common expression in a cluster. Third, we used known or novel TFs and other molecular markers that are expressed by a cluster as an entry point to look for other cluster specific genes. Each of these methods provides candidate neuron-cluster association, which requires experimental validation. It is not clear which method is more robust at identifying functional neuron-cluster associations. Nevertheless, we have validated a number of neuron-cluster associations using the third approach: beginning with a single gene-cluster correlation. In the future, our data can be used in two pipelines for linking neuron subtype to transcriptome. First, start with at least one validated marker for a cluster, then search all labeled clusters for candidate TFs that co-express with the validated marker, followed by validation via RNA in situ (or antibodies). Second, we can collect neurons of interest with FACS or antibody selection (Davis et al., 2020 [↗](#)), and perform bulk RNA sequencing, followed by looking for clusters enriched for the TFs found in bulk sequencing data, and then validation via RNA in situ (or antibodies). Notably, the second pipeline requires no prior markers beyond the Gal4 or LexA line expressed in single neuronal populations. Both pipelines result in a transcriptome that can be used to identify (a) TFs for their role in neuronal specification and/or maintenance, (b) cell surface molecules that may regulate neuronal morphology and/or connectivity, or (c) functionally relevant genes encoding ion channels, neuropeptides, receptors, and signaling pathways.

Materials and methods

Key resources table

Reagent type (species of resource)	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (<i>Drosophila melanogaster</i>)	20xUAS- <i>FLPG5.PEST;worniu-gal4,asense-gal80;Act5c(FRT.CD2)gal4</i>	(Syed et al., 2017)		Type II lineage immortalization
Genetic reagent (<i>Drosophila melanogaster</i>)	20xUAS- <i>FLPG5.PEST;worniu-gal4,asense-gal80;lexAop(FRT.stop)-mCD8:GFP</i>	This work		Label type II derived lexA+ cells
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>TI{2A-lexA::GAD}AstA[2A-lexA]/TM3,Sb[1]</i>	BDSC	RRID:BDSC_84356	
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>TI{2A-lexA::GAD}Ms[2A-lexA]/TM3,Sb[1]</i>	BDSC	RRID:BDSC_84403	
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>TI{2A-lexA::GAD}NPF[2A-lexA]/TM3,Sb[1]</i>	BDSC	RRID:BDSC_84422	
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>UAS-RedStinger</i>	BDSC	RRID:BDSC_8545	
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>UAS-unc84-2xGFP</i>	(Henry et al., 2012)		
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>GMR12D09-lexA/CyO</i>	BDSC	RRID:BDSC_54419	P-EN
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>GMR16D01-lexA</i>	BDSC	RRID:BDSC_52503	P-FN
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>GMR37G12-lexA</i>	BDSC	RRID:BDSC_52765	PF-R
Genetic reagent (<i>Drosophila melanogaster</i>)	<i>13xLexAop2-IVS-myr::smGdP-V5</i>	BDSC	RRID:BDSC_62215	
Antibody, monoclonal	mouse anti-Cut 2B10	DSHB	RRID: AB_528186	2 µg/mL
Antibody, polyclonal	guinea pig anti-DIP-β	(Xu et al., 2024)		1:300
Antibody, polyclonal	guinea pig anti-E93	(Syed et al., 2017)		1:500
Antibody, polyclonal	rabbit anti-Imp	(Syed et al., 2017)		1:500
Antibody, polyclonal	rabbit anti-Lim1	C. Desplan, New York University		1:500
Antibody, polyclonal	guinea pig anti-Runt	Sullivan et al., 2019		1:1000
Antibody, polyclonal	guinea pig anti-Rx	C. Desplan, New York University		1:500
Antibody, polyclonal	rabbit anti-Syp	(Syed et al., 2017)		1:500
Antibody, polyclonal	rabbit anti-Toy	(Sullivan et al., 2019)		1:1000
Antibody, polyclonal	chicken anti-V5	Fortis Life Sciences, Waltham, MA		1 µg/mL
Antibody, polyclonal	rat anti-Zfh2	(Tran et al., 2010)		1:200
Antibody, polyclonal	DyLight405, Alexa Fluor 488, Rhodamine Red-X (RRX), or Alexa Fluor 647 conjugated donkey whole IgG	Jackson ImmunoResearch Laboratories Inc., West Grove, PA		5 µg/mL

Single nuclei isolation, library preparation, and sequencing

We first used the split-pool method (Parse Bioscience, Seattle, WA, USA) to barcode RNAs from whole brain to generate a library which includes both T1 and T2 progeny. This library, T1+T2, was used for analysis in [Figure 1](#) and [5](#). To increase the number of T2 nuclei for snRNAseq, we labeled T2 nuclei by crossing *20xUAS-FLPG5.PEST;worniu-gal4,asense-gal80; Act5c(FRT.CD2)gal4* to *UAS-RedStinger* (RFP) or *UAS-unc84-2xGFP* (GFP) flies. The adult flies were aged for 1 week at 25 °C before dissection. Equal amounts of male and female central brains (excluding optic lobes) were dissected at room temperature within 1 hour. The samples were flash-frozen in liquid nitrogen and stored separately at -80°C. Nuclei dissociated from the frozen, dissected brains was performed according to published protocols ([McLaughlin et al., 2022](#)). RFP+ or GFP+ nuclei were collected by sorting dissociated nuclei with SONY-SH800 with 100 µm chip. We performed three rounds of sorting and snRNAseq. In the first round, we pooled male and female brains together to select GFP+ nuclei and used particle-templated instant partitions to capture single nuclei to generate cDNA library (Fluent BioSciences, Watertown, MA). In the second round, RFP+ nuclei from male and female were pooled together. In the third round, RFP+ nuclei from male and female brains were collected separately. The split-pool method was then used to generate barcoded cDNA libraries from each individual nucleus from the second and third rounds. All libraries were sequenced with pair-ends reads 150bp on Illumina Novaseq 6000 (University of Oregon's Genomics and Cell Characterization Core Facility).

snRNA-seq analysis

Our bioinformatic analysis was performed using pipeline from Fluent BioSciences, Parse Bioscience, and the Seurat R package ([Hao et al., 2024](#); [Satija et al., 2015](#)). Briefly, pipelines from Fluent BioSciences and Parse Biosciences were used to perform demultiplexing, alignment, filtering, counting of barcodes and UMIs with an output being a cell-by-genes matrix of counts. We aligned our sequences to a custom reference genome by adding *flpD5* (RRID:Addgene_32132), *redstinger* (RRID:Addgene_46165), and *unc84sfGFP* (RRID:Addgene_46023) (+SV40 tail) sequences and annotations to the *Drosophila* genome release BDGP6.32.109. To further ensure that only high-quality cells were retained, we removed any cells with fewer than 200 genes or more than 2500 genes expressed and more than 5% mitochondrial RNA. For the T2 atlas, snRNA-seq data from three rounds of sorting and barcoding were integrated in Seurat with Anchor-based RPCA integration to generate an integrated dataset, and the downstream analysis was performed with the default parameters. The UMAP was generated with resolution 12. The integrated dataset was used for analyses.

Glia analyses

The T1+T2 glia atlas dataset was derived from the T1+T2 *repo*+ glial clusters and re-clustered to represent the glial subtypes. The T2 glia atlas was derived from the T2 *repo*+ glial clusters and re-clustered to represent the glial subtypes. Cell identity was determined by the validated markers for glia shown to differentially expressed between the clusters. Identities were assigned based on expression of these validated markers. Overlapping identities were assigned if multiple subtype markers were expressed within a single cluster.

Sex differences

We determined sex-biased clusters within the T2 glia and neuron atlases by identifying clusters that were disproportionate after normalizing the number of inputs for 'female' and 'male' samples respectively. The 'female and male' mixed samples were excluded from the analyses as we could not differentiate the sex origin of these nuclei. To determine differential gene expression between

females and males, we pseudo bulked nuclei by aggregating the snRNAseq for comparison between sex and clusters. Heatmaps were generated in Seurat using the Scillus package (<https://github.com/xmc811/Scillus>) to display the heatmap.

Transcription factor combinatorial analysis

We used the Seurat function FindAllMarkers to find positively differentially expressed TFs in all clusters. We then removed TFs which were not significantly differentially expressed. We gave a value of 1 to any TF that was found to be a positive marker in each cluster and a value of 0 in clusters which it was not a positive marker. Next, we found the number of unique combinations of markers for 6 classes (zinc finger, helix-turn-helix, homeodomain, basic domain, unidentified DNA binding domain, and high motility group) of TFs. The Jaccard index between each cluster was then calculated and clusters were sorted from most to least similar. The python packages Matplotlib and Seaborn were then used to generate heatmaps.

Immunohistochemistry and imaging

Standard methods were used for adult brain staining (Sullivan et al., 2019). The antibody stained brains were mounted in DPX (<https://www.janelia.org/project-team/flylight/protocols>) on poly-L-Lysine coated coverslip (Corning, Glendale, AZ) and imaged with Zeiss confocal LSM800 with software Zen.

Figure production

We used Imaris (Bitplane, Abingdon, UK) for confocal image processing, and Illustrator (Adobe, San Jose, CA) to assemble figures.

Availability of data and materials

All data and code can be found at https://github.com/dgepiney/2023_Doe_Drosophila_Central_Brain_RNAseq.

Acknowledgements

We thank Jason Carriere, and Peter Newstein for comments on the manuscript. We thank the Desplan lab (NYU) for reagents. We thank Jason Carriere (University of Oregon Genomics and Cell Characterization Core Facility) for technical assistances in cDNA library preparation and sequencing. Flies were obtained from the Bloomington Stock Center.

Additional information

Ethical Approval and Consent to participate

No vertebrate animals were used.

Consent for publication

All authors consent for this to be published

Authors' contributions

DE, NRD, GNMC and SLL performed dissections.

DE, GNMC, and SLL performed FACS, and RNAseq data integration. NRD generated **Figure 1** [↗](#), 4, 5, ST3, ST4, ST5, ST9, ST10, and ST11. GNMC generated **Figures 3** [↗](#), 6, SF3, ST12, SM1, SM2, and SM3.

DE generated **Figures 7** [↗](#), 8, and ST12.

SLL generated **Figures 1** [↗](#), 2, 8, 9, SF1, SF2, ST1, ST2, ST6, ST7, ST8, ST13, and ST14.

All authors contributed to writing the manuscript. CQD supervised the project.

Funding

<u>Funder</u>	<u>Grant reference number</u>	<u>Author</u>
Howard Hughes Medical Institute	None	CQD, SLL, NRD, DE
NIH Training Grant	5-T32-HD07348	NRD
NIH Training Grant	5-T32-GM149387	GNMC
NIH	R01-HD27056	CQD, NRD, DE, GNMC

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Additional files

Supplemental Tables 1-14 [↗](#)

References

- Amrein H., Axel R (1997) **Genes expressed in neurons of adult male *Drosophila*** *Cell* **88**:459–469
- Awasaki T., Lai S.-L., Ito K., Lee T (2008) **Organization and postembryonic development of glial cells in the adult central brain of *Drosophila*** *J. Neurosci. Off. J. Soc. Neurosci* **28**:13742–13753
- Baek M., Enriquez J., Mann R. S (2013) **Dual role for Hox genes and Hox co-factors in conferring leg motoneuron survival and identity in *Drosophila*** *Development* **140**:2027–38
- Baumgardt M., Karlsson D., Salmani B. Y., Bivik C., MacDonald R. B., Gunnar E., Thor S (2014) **Global programmed switch in neural daughter cell proliferation mode triggered by a temporal gene cascade** *Dev Cell* **30**:192–208
- Bello B. C., Izergina N., Caussinus E., Reichert H (2008) **Amplification of neural stem cell proliferation by intermediate progenitor cells in *Drosophila* brain development** *Neural Dev* **3**:5
- Boone J. Q., Doe C. Q (2008) **Identification of *Drosophila* type II neuroblast lineages containing transit amplifying ganglion mother cells** *Dev Neurobiol* **68**:1185–95
- Bowman S. K., Rolland V., Betschinger J., Kinsey K. A., Emery G., Knoblich J. A (2008) **The tumor suppressors Brat and Numb regulate transit-amplifying neuroblast lineages in *Drosophila*** *Dev Cell* **14**:535–46
- Boyan G., Williams L (2011) **Embryonic development of the insect central complex: insights from lineages in the grasshopper and *Drosophila*** *Arthropod Struct. Dev* **40**:334–348
- Briscoe J., Pierani A., Jessell T. M., Ericson J (2000) **A Homeodomain Protein Code Specifies Progenitor Cell Identity and Neuronal Fate in the Ventral Neural Tube** *Cell* **101**:435–445
- Campbell G., Goring H., Lin T., Spana E., Andersson S., Doe C. Q., Tomlinson A (1994) **RK2, a glial-specific homeodomain protein required for embryonic nerve cord condensation and viability in *Drosophila*** *Development* **120**:2957–66
- Croset V., Treiber C. D., Waddell S (2018) **Cellular diversity in the *Drosophila* midbrain revealed by single-cell transcriptomics** *eLife* **7**:e34550
- Davie K., Janssens J., Koldere D., De Waegeneer M., Pech U., Kreft L., Aibar S., Makhzami S., Christiaens V., Bravo González-Blas C., et al. (2018) **A Single-Cell Transcriptome Atlas of the Aging *Drosophila* Brain** *Cell* **174**:982–998
- Davis F. P., Nern A., Picard S., Reiser M. B., Rubin G. M., Eddy S. R., Henry G. L (2020) **A genetic, genomic, and computational resource for exploring neural circuit function** *eLife* **9**:e50901
- Deng B., Li Q., Liu X., Cao Y., Li B., Qian Y., Xu R., Mao R., Zhou E., Zhang W., et al. (2019) **Chemoconnectomics: Mapping Chemical Transmission in *Drosophila*** *Neuron* **101**:876–893

- DeSalvo M. K., Hindle S. J., Rusan Z. M., Orng S., Eddison M., Halliwill K., Bainton R. J. (2014) **The *Drosophila* surface glia transcriptome: evolutionary conserved blood-brain barrier processes** *Front. Neurosci* **8**:346
- Dillon N. R., Doe C. Q. (2024) **Castor is a temporal transcription factor that specifies early born central complex neuron identity** *Dev. Camb. Engl.* :dev.204318
- Dillon N. R., Manning L., Hirono K., Doe C. Q (2024) **Seven-up acts in neuroblasts to specify adult central complex neuron identity and initiate neuroblast decommissioning** *Dev. Camb. Engl* **151**:dev202504
- Doe C. Q (2017) **Temporal Patterning in the *Drosophila* CNS** *Annu Rev Cell Dev Biol* **33**:219–240
- Enriquez J., Venkatasubramanian L., Baek M., Peterson M., Aghayeva U., Mann R. S (2015) **Specification of individual adult motor neuron morphologies by combinatorial transcription factor codes** *Neuron* **86**:955–70
- Fisher Y. E (2022) **Flexible navigational computations in the *Drosophila* central complex** *Curr. Opin. Neurobiol* **73**:102514
- Franconville R., Beron C., Jayaraman V (2018) **Building a functional connectome of the *Drosophila* central complex** *eLife* **7**:e37017
- Granger A. J., Wang W., Robertson K., El-Rifai M., Zanello A. F., Bistrong K., Saunders A., Chow B. W., Nuñez V., Turrero García M., et al. (2020) **Cortical ChAT+ neurons co-transmit acetylcholine and GABA in a target- and brain-region-specific manner** *eLife* **9**:e57749
- Granger A. J., Mao K., Saulnier J. L., Hines M. E., Sabatini B. L (2023) **Developmental regulation of GABAergic gene expression in forebrain cholinergic neurons** *Front. Neural Circuits* **17**:1125071
- Gu Z. (2022) **Complex heatmap visualization** *iMeta* **1**:e43
- Hao Y., Stuart T., Kowalski M. H., Choudhary S., Hoffman P., Hartman A., Srivastava A., Molla G., Madad S., Fernandez-Granda C., et al. (2024) **Dictionary learning for integrative, multimodal and scalable single-cell analysis** *Nat. Biotechnol* **42**:293–304
- Henry G. L., Davis F. P., Picard S., Eddy S. R (2012) **Cell type-specific genomics of *Drosophila* neurons** *Nucleic Acids Res* **40**:9691–9704
- Hindle S. J., Bainton R. J (2014) **Barrier mechanisms in the *Drosophila* blood-brain barrier** *Front. Neurosci* **8**:414
- Hoebert O (2021) **Homeobox genes and the specification of neuronal identity** *Nat. Rev. Neurosci* **22**:627–636
- Holguera I., Desplan C (2018) **Neuronal specification in space and time** *Science* **362**:176–180
- Hulse B. K., Haberkern H., Franconville R., Turner-Evans D. B., Takemura S., Wolff T., Noorman M., Dreher M., Dan C., Parekh R., et al. (2020) **A connectome of the *Drosophila* central complex reveals network motifs suitable for flexible navigation and context-dependent action selection** *bioRxiv* :2020.12.08.413955

Hulse B. K., Haberkern H., Franconville R., Turner-Evans D., Takemura S., Wolff T., Noorman M., Dreher M., Dan C., Parekh R., et al. (2021) **A connectome of the *Drosophila* central complex reveals network motifs suitable for flexible navigation and context-dependent action selection** *eLife* **10**:e66039

Ito M., Masuda N., Shinomiya K., Endo K., Ito K (2013) **Systematic analysis of neural projections reveals clonal composition of the *Drosophila* brain** *Curr Biol* **23**:644–55

Kandimalla P., Omoto J. J., Hong E. J., Hartenstein V (2023) **Lineages to circuits: the developmental and evolutionary architecture of information channels into the central complex** *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol* **209**:679–720

Konstantinides N., Kapuralin K., Fadil C., Barboza L., Satija R., Desplan C (2018) **Phenotypic Convergence: Distinct Transcription Factors Regulate Common Terminal Features** *Cell* **174**:622–635

Konstantinides N., Holguera I., Rossi A. M., Escobar A., Dudragne L., Chen Y.-C., Tran T. N., Martínez Jaimes A. M., Özel M. N., Simon F., et al. (2022) **A complete temporal transcription factor series in the fly visual system** *Nature* **604**:316–322

Lago-Baldaia I., Cooper M., Seroka A., Trivedi C., Powell G. T., Wilson S. W., Ackerman S. D., Fernandes V. M (2023) **A *Drosophila* glial cell atlas reveals a mismatch between transcriptional and morphological diversity** *PLoS Biol* **21**:e3002328

Li H., Janssens J., De Waegeneer M., Kolluru S. S., Davie K., Gardeux V., Saelens W., David F. P. A., Brbić M., Spanier K., et al. (2022) **Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly** *Science* **375**:eabk2432

Liu Z., Yang C. P., Sugino K., Fu C. C., Liu L. Y., Yao X., Lee L. P., Lee T (2015) **Opposing intrinsic temporal gradients guide neural stem cell production of varied neuronal fates** *Science* **350**:317–20

Lozovaya N., Eftekhari S., Cloarec R., Gouty-Colomer L. A., Dufour A., Riffault B., Billon-Grand M., Pons-Bennaceur A., Oumar N., Burnashev N., et al. (2018) **GABAergic inhibition in dual-transmission cholinergic and GABAergic striatal interneurons is abolished in Parkinson disease** *Nat. Commun* **9**:1422

McLaughlin C. N., Brbić M., Xie Q., Li T., Horns F., Kolluru S. S., Kebschull J. M., Vacek D., Xie A., Li J., et al. (2021) **Single-cell transcriptomes of developing and adult olfactory receptor neurons in *Drosophila*** *eLife* **10**:e63856

McLaughlin C. N., Qi Y., Quake R.S., Luo L., Li H. (2022) **Isolation and RNA sequencing of single nuclei from *Drosophila* tissues: STAR Protocols** *STAR Protocols* **3**:101417

Meller V. H., Wu K. H., Roman G., Kuroda M. I., Davis R. L (1997) **roX1 RNA paints the X chromosome of male *Drosophila* and is regulated by the dosage compensation system** *Cell* **88**:445–457

Meye F. J., Soiza-Reilly M., Smit T., Diana M. A., Schwarz M. K., Mameli M (2016) **Shifted pallidal co-release of GABA and glutamate in habenula drives cocaine withdrawal and relapse** *Nat. Neurosci* **19**:1019–1024

Naidu V. G., Zhang Y., Lowe S., Ray A., Zhu H., Li X (2020) **Temporal progression of *Drosophila* medulla neuroblasts generates the transcription factor combination to control T1 neuron**

morphogenesis *Dev. Biol* **464**:35–44

Nguyen T. H., Viciomini R., Choudhury S. D., Coon S. L., Iben J., Brody T., Serpe M. (2021) **Single-Cell RNA Sequencing Analysis of the Drosophila Larval Ventral Cord** *Curr. Protoc.* **1**:e38

Özel M. N., Gibbs C. S., Holguera I., Soliman M., Bonneau R., Desplan C. (2022) **Coordinated control of neuronal differentiation and wiring by a sustained code of transcription factors** *bioRxiv* :2022.05.01.490216

Pollington H. Q., Seroka A. Q., Doe C. Q (2023) **From temporal patterning to neuronal connectivity in Drosophila type I neuroblast lineages** *Semin. Cell Dev. Biol* **142**:4–12

Raji J. I., Potter C. J (2021) **The number of neurons in Drosophila and mosquito brains** *PLOS One* **16**:e0250381

Reilly M. B., Cros C., Varol E., Yemini E., Hobert O (2020) **Unique homeobox codes delineate all the neuron classes of C. elegans** *Nature* **584**:595–601

Rosenberg A. B., Roco C. M., Muscat R. A., Kuchina A., Sample P., Yao Z., Graybuck L. T., Peeler D. J., Mukherjee S., Chen W., et al. (2018) **Single-cell profiling of the developing mouse brain and spinal cord with split-pool barcoding** *Science* **360**:176–182

Ryner L. C., Goodwin S. F., Castrillon D. H., Anand A., Villella A., Baker B. S., Hall J. C., Taylor B. J., Wasserman S. A (1996) **Control of male sexual behavior and sexual orientation in Drosophila by the fruitless gene** *Cell* **87**:1079–1089

Sagner A., Zhang I., Watson T., Lazaro J., Melchionda M., Briscoe J (2021) **A shared transcriptional code orchestrates temporal patterning of the central nervous system** *PLoS Biol* **19**:e3001450

Satija R., Farrell J. A., Gennert D., Schier A. F., Regev A (2015) **Spatial reconstruction of single-cell gene expression data** *Nat. Biotechnol* **33**:495–502

Sato M., Suzuki T (2022) **Cutting edge technologies expose the temporal regulation of neurogenesis in the Drosophila nervous system** *Fly (Austin)* **16**:222–232

Saunders A., Granger A. J., Sabatini B. L (2015) **Corelease of acetylcholine and GABA from cholinergic forebrain neurons** *eLife* **4**:e06412

Schlegel P., Yin Y., Bates A. S., Dorkenwald S., Eichler K., Brooks P., Han D. S., Gkantia M., dos Santos M., Munnelly E. J., et al. (2024) **Whole-brain annotation and multi-connectome cell typing of Drosophila** *Nature* **634**:139–152

Schoofs L., De Loof A., Van Hiel M. B. (2017) **Neuropeptides as Regulators of Behavior in Insects** *Annu. Rev. Entomol* **62**:35–52

Sgammeglia N., Sprecher S. G (2022) **Interplay between metabolic energy regulation and memory pathways in Drosophila** *Trends Neurosci* **45**:539–549

Shabel S. J., Proulx C. D., Piriz J., Malinow R (2014) **Mood regulation. GABA/glutamate co-release controls habenula output and is modified by antidepressant treatment** *Science* **345**:1494–1498

Shu S., Jiang M., Deng X., Yue W., Cao X., Zhang K., Wang Z., He H., Cui J., Wang Q., et al. (2023) **Heterochromatic silencing of immune-related genes in glia is required for BBB integrity and normal lifespan in drosophila** *Aging Cell* **22**:e13947

Sigorelli A. (2024) **DescTools: Tools for Descriptive Statistics** CRAN <https://doi.org/10.32614/CRAN.package.DescTools>

Siletti K., Hodge R., Mossi Albiach A., Lee K. W., Ding S.-L., Hu L., Lönnerberg P., Bakken T., Casper T., Clark M., et al. (2023) **Transcriptomic diversity of cell types across the adult human brain** *Science* **382**:eadd7046

Soffers J. H., Beck E., Sytkowski D., Maughan M., Devasi D., Zhu Y., Wilson B., Chen Y.-C. D., Erclik T., Truman J. W., et al. (2024) **A library of lineage-specific driver lines connects developing neuronal circuits to behavior in the Drosophila Ventral Nerve Cord** *BioRxiv* :2024.11.27.625713

Spitzer N. C (2015) **Neurotransmitter Switching? No Surprise.** *Neuron* **86**:1131–1144

Stuart T., Butler A., Hoffman P., Hafemeister C., Papalexi E., Mauck W. M., Hao Y., Stoeckius M., Smibert P., Satija R (2019) **Comprehensive Integration of Single-Cell Data** *Cell* **177**:1888–1902

Sullivan L. F., Warren T. L., Doe C. Q (2019) **Temporal identity establishes columnar neuron morphology, connectivity, and function in a Drosophila navigation circuit** *eLife* **8**

Syed M. H., Mark B., Doe C. Q (2017) **Steroid hormone induction of temporal gene expression in Drosophila brain neuroblasts generates neuronal and glial diversity** *eLife* **6**

Takács V. T., Cserép C., Schlingloff D., Pósfai B., Szőnyi A., Sos K. E., Környei Z., Dénes Á., Gulyás A. I., Freund T. F., et al. (2018) **Co-transmission of acetylcholine and GABA regulates hippocampal states** *Nat. Commun* **9**:2848

Tran K. D., Miller M. R., Doe C. Q (2010) **Recombineering Hunchback identifies two conserved domains required to maintain neuroblast competence and specify early-born neuronal identity** *Development* **137**:1421–30

Turner-Evans D. B., Jensen K. T., Ali S., Paterson T., Sheridan A., Ray R. P., Wolff T., Lauritzen J. S., Rubin G. M., Bock D. D., et al. (2020) **The Neuroanatomical Ultrastructure and Function of a Biological Ring Attractor** *Neuron* **108**:145–163

Velten J., Gao X., Van Nierop Y Sanchez P, Domsch K, Agarwal R, Bognar L, Paulsen M, Velten L, Lohmann I (2022) **Single-cell RNA sequencing of motoneurons identifies regulators of synaptic wiring in Drosophila embryos** *Mol. Syst. Biol* **18**:e10255

Viktorin G., Riebli N., Reichert H (2013) **A multipotent transit-amplifying neuroblast lineage in the central brain gives rise to optic lobe glial cells in Drosophila** *Dev Biol* **379**:182–94

Wang Y. C., Yang J. S., Johnston R., Ren Q., Lee Y. J., Luan H., Brody T., Odenwald W. F., Lee T (2014) **Drosophila intermediate neural progenitors produce lineage-dependent related series of diverse neurons** *Development* **141**:253–8

Warren T. G., Brennan M. D., Mahowald A. P (1979) **Two processing steps in maturation of vitellogenin polypeptides in Drosophila melanogaster** *Proc. Natl. Acad. Sci. U. S. A* **76**:2848–2852

Wolff T., Rubin G. M (2018) **Neuroarchitecture of the *Drosophila* central complex: A catalog of nodulus and asymmetrical body neurons and a revision of the protocerebral bridge catalog** *J. Comp. Neurol* **526**:2585–2611

Wolff T., Iyer N. A., Rubin G. M (2015) **Neuroarchitecture and neuroanatomy of the *Drosophila* central complex: A GAL4-based dissection of protocerebral bridge neurons and circuits** *J. Comp. Neurol* **523**:997–1037

Wolff T., Eddison M., Chen N., Nern A., Sundaramurthi P., Sitaraman D., Rubin G. M. (2025) **Cell type-specific driver lines targeting the *Drosophila* central complex and their use to investigate neuropeptide expression and sleep regulation** *BioRxiv* :2024.10.21.619448

Xiong W. C., Okano H., Patel N. H., Blendy J. A., Montell C (1994) **repo encodes a glial-specific homeo domain protein required in the *Drosophila* nervous system** *Genes Dev* **8**:981–94

Xu C., Ramos T. B., Rogers E. M., Reiser M. B., Doe C. Q (2024) **Homeodomain proteins hierarchically specify neuronal diversity and synaptic connectivity** *eLife* **12**:RP90133

Yang J. S., Awasaki T., Yu H. H., He Y., Ding P., Kao J. C., Lee T (2013) **Diverse neuronal lineages make stereotyped contributions to the *Drosophila* locomotor control center, the central complex** *J Comp Neurol* **521**:2645–62

Yu H. H., Kao C. F., He Y., Ding P., Kao J. C., Lee T (2010) **A complete developmental sequence of a *Drosophila* neuronal lineage as revealed by twin-spot MARCM** *PLOS Biol* **8**

Yu H. H., Awasaki T., Schroeder M. D., Long F., Yang J. S., He Y., Ding P., Kao J. C., Wu G. Y., Peng H., et al. (2013) **Clonal development and organization of the adult *Drosophila* central brain** *Curr Biol* **23**:633–43

Author information

Derek Epiney[#]

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, United States

ORCID iD: [0000-0002-6056-1817](https://orcid.org/0000-0002-6056-1817)

[#]equal contribution

Gonzalo N Morales Chaya[#]

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, United States

ORCID iD: [0009-0005-4263-1323](https://orcid.org/0009-0005-4263-1323)

[#]equal contribution

Noah R Dillon[#]

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, United States

ORCID iD: [0000-0002-6708-4208](https://orcid.org/0000-0002-6708-4208)

#equal contribution

Sen-Lin Lai

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, United States

ORCID iD: [0000-0002-7531-283X](https://orcid.org/0000-0002-7531-283X)

For correspondence: slai@uoregon.edu

Chris Q Doe

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, United States

ORCID iD: [0000-0001-5980-8029](https://orcid.org/0000-0001-5980-8029)

For correspondence: cdoe@uoregon.edu

Editors

Reviewing Editor

Paschalis Kratsios

University of Chicago, Chicago, United States of America

Senior Editor

Claude Desplan

New York University, New York, United States of America

Reviewer #1 (Public review):

Summary:

Epiney et al. use single-nuclei RNA sequencing (snRNA-seq) to characterize the lineage of Type-2 (T2) neuroblasts (NBs) in the adult *Drosophila* brain. To isolate cells born from T2 NBs, the authors used a genetic tool that specifically allows the permanent labeling of T2-derived cell types, which are then FAC-sorted for snRNA-seq. This effective labeling approach also allows them to compare the isolated T2 lineage cells with T1-derived cell types by a simple exclusion method. The authors begin by describing a transcriptomic atlas for all T1 and T2-derived neuronal and glia clusters, reporting that the T2-derived lineage comprises 161 neuronal clusters, in contrast to the T1 lineage which comprises 114 of them. The authors then use the expression of VAcHT, VGlut, Gad1, Tbh, Ple, SerT, and Tdc2 to show that T2 neuroblasts generate all major neuron classes of fast-acting neurotransmitters. Strikingly, they show that a subset of glia and neuronal clusters have disproportionate enrichment in males or females, suggesting that T2 neuroblasts generate sex-biased cell types. The authors then proceed to characterize neuropeptide expression across T2-derived neuronal clusters and argue that the same neuropeptide can be expressed across different cell types, while similar cell types can express distinct neuropeptides. The functional implication of both observations, however, remains to be tested. Furthermore, the authors describe combinatorial transcription factor (TF) codes that are correlated with neuropeptide expression for T2-derived neurons along with an overall TF code for all T2-derived cell types, both of which will serve as an important starting point for future investigations. Finally, the authors map well-studied neuronal types of the central complex to the clusters of their T2-derived snRNA-seq dataset. They use known marker combinations, bulk RNA-seq data and

highly specific split-GAL4 driver lines to annotate their T2-derived atlas, establishing a comprehensive transcriptomic atlas that would guide future studies in this field.

Strengths:

This study provides an in-depth transcriptomic characterization of neurons and glia derived from Type-2 neuroblast lineages. The results of this manuscript offer several future directions to investigate the mechanisms of diversifying neuronal identity. The datasets of T1-derived and T2-derived cells will pave the way for studies focused on the functional analysis of combinatorial TF codes specifying cell identity, sex-based differences in neurogenesis and gliogenesis, the relationship between neuropeptide (co)expression and cell identity, and the differential contributions of distinct progenitor populations to the same cell type.

Weaknesses:

The study presents several important observations based on the characterization of Type II neuroblast-derived lineages. However, a mechanistic insight is missing for most observations. The idea that there is a sex-specific bias to certain T2-derived neurons and glial clusters is quite interesting, however, the functional significance of this observation is not tested or discussed extensively. Finally, the authors do not show whether the combinatorial TF code is indeed necessary for neuropeptide expression or if this is just a correlation due to cell identity being defined by TFs. Functional knockdown of some candidate TFs for a subset of neuropeptide-expressing cells would have been helpful in this case.

Comments on revisions:

The authors have addressed my recommendations.

<https://doi.org/10.7554/eLife.105896.2.sa2>

Reviewer #2 (Public review):

In this manuscript, Epiney et al., present a single-nucleus sequencing analysis of *Drosophila* adult central brain neurons and glia. By employing an ingenious permanent labeling technique, they trace the progeny of T2 neuroblasts, which play a key role in the formation of the central complex. This transcriptomic dataset is poised to become a valuable resource for future research on neurogenesis, neuron morphology, and behavior.

The authors further delve into this dataset with several analyses, including the characterization of neurotransmitter expression profiles in T2-derived neurons. While some of the bioinformatic analyses are preliminary, they would benefit from additional experimental validation in future studies.

Comments on revisions:

We appreciate the authors' efforts to address some of the comments. While these revisions have improved the clarity of certain sections, some of the larger concerns remain unaddressed. Specifically, the manuscript still lacks the additional analyses that would allow for more specific conclusions, rather than the general observations currently presented. Although the revisions have certainly made the text clearer, the core issue of needing more detailed analysis to draw more concrete conclusions still stands.

<https://doi.org/10.7554/eLife.105896.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

Summary:

Epiney et al. use single-nuclei RNA sequencing (snRNA-seq) to characterize the lineage of Type-2 (T2) neuroblasts (NBs) in the adult Drosophila brain. To isolate cells born from T2 NBs, the authors used a genetic tool that specifically allows the permanent labeling of T2-derived cell types, which are then FAC-sorted for snRNA-seq. This effective labeling approach also allows them to compare the isolated T2 lineage cells with T1-derived cell types by a simple exclusion method. The authors begin by describing a transcriptomic atlas for all T1 and T2-derived neuronal and glia clusters, reporting that the T2-derived lineage comprises 161 neuronal clusters, in contrast to the T1 lineage which comprises 114 of them. The authors then use the expression of VACHT, VGlut, Gad1, Tbh, Ple, SerT, and Tdc2 to show that T2 neuroblasts generate all major neuron classes of fast-acting neurotransmitters. Strikingly, they show that a subset of glia and neuronal clusters have disproportionate enrichment in males or females, suggesting that T2 neuroblasts generate sex-biased cell types. The authors then proceed to characterize neuropeptide expression across T2-derived neuronal clusters and argue that the same neuropeptide can be expressed across different cell types, while similar cell types can express distinct neuropeptides. The functional implication of both observations, however, remains to be tested. Furthermore, the authors describe combinatorial transcription factor (TF) codes that are correlated with neuropeptide expression for T2-derived neurons along with an overall TF code for all T2-derived cell types, both of which will serve as an important starting point for future investigations. Finally, the authors map well-studied neuronal types of the central complex to the clusters of their T2-derived snRNA-seq dataset. They use known marker combinations, bulk RNA-seq data and highly specific split-GAL4 driver lines to annotate their T2-derived atlas, establishing a comprehensive transcriptomic atlas that would guide future studies in this field.

Thanks for the clear and accurate summary of our findings.

Strengths:

This study provides an in-depth transcriptomic characterization of neurons and glia derived from Type-2 neuroblast lineages. The results of this manuscript offer several future directions to investigate the mechanisms of diversifying neuronal identity. The datasets of T1-derived and T2-derived cells will pave the way for studies focused on the functional analysis of combinatorial TF codes specifying cell identity, sex-based differences in neurogenesis and gliogenesis, the relationship between neuropeptide (co)expression and cell identity, and the differential contributions of distinct progenitor populations to the same cell type.

Thank you for the positive comments.

Weaknesses:

The study presents several important observations based on the characterization of Type II neuroblast-derived lineages. However, a mechanistic insight is missing for most observations. The idea that there is a sex-specific bias to certain T2-derived neurons and

glial clusters is quite interesting, however, the functional significance of this observation is not tested or discussed extensively. Finally, the authors do not show whether the combinatorial TF code is indeed necessary for neuropeptide expression or if this is just a correlation due to cell identity being defined by TFs. Functional knockdown of some candidate TFs for a subset of neuropeptide-expressing cells would have been helpful in this case.

We agree that we do not provide mechanistic or functional insights. Our goal was to produce hypothesis generating datasets for our lab and others to use to direct functional or mechanistic studies.

Reviewer #2 (Public review):

In this manuscript, Epiney et al., present a single-nucleus sequencing analysis of Drosophila adult central brain neurons and glia. By employing an ingenious permanent labeling technique, they trace the progeny of T2 neuroblasts, which play a key role in the formation of the central complex. This transcriptomic dataset is poised to become a valuable resource for future research on neurogenesis, neuron morphology, and behavior.

Thank you for the positive comments.

The authors further delve into this dataset with several analyses, including the characterization of neurotransmitter expression profiles in T2-derived neurons. While some of the bioinformatic analyses are preliminary, they would benefit from additional experimental validation in future studies.

Thank you for the positive comments. We too hope that future research will benefit from this dataset.

Reviewer #1 (Recommendations for the authors):

Major points

(1) In Figures 1E and 4A, the T1 and T2 glia subsets reveal sub-clusters for several cell types as seen by the distribution of points on the UMAP. This observation is never validated or discussed. Do these sub-clusters represent true differences in identities or are they artifacts of the single-nucleus preparation? For Figure 1E, it is not clear whether specific sub-clusters (see Ensheathing-4 vs Ensheathing-5 and Astrocyte-2 vs. Astrocyte-6) are differentially enriched between the T1 and T2 lineages. The existence of these sub-clusters must be discussed or dismissed.

We agree that this needs to be addressed more clearly in the manuscript and have made text changes in the Results and Discussion sections to clarify. We note that a recent glial cell atlas (Lago-Baldaia et al., 2023; PMID: 37862379) of the developing fly VNC and optic lobes found sub-clusters that mapped to the same subtype annotations. Interestingly, Lago-Baldaia and colleagues found that the transcriptional diversity of glia cell types did not match the morphological diversity of glia validated *in vivo*. See text changes below.

Lines 131-133: “Similar to a previous glial cell atlas (Lago-Baldaia et al., 2023) we found some glial subtypes (astrocytes, ensheathing, and subperineurial) mapped to multiple clusters (Figure 1E, 1F).”

Lines 206-208: “In line with our T1+T2 atlas and previous glia cell atlas (Lago-Baldaia et al., 2023), some subtypes mapped to several subclusters including ensheathing, astrocytes, and chiasm (Figure 4A-B).”

Lines 397-401: “Similar to a recent glial cell atlas (Lago-Baldaia et al., 2023), we found glial subtypes like astrocytes, ensheathing, and subperineurial glia mapped to several sub-clusters (Figure 1E-F). It remains unclear if these sub-clusters with the same cell type annotation represent distinct glial identities or different transcriptional states within these populations.”

(2) The authors present evidence for sex-specific neuronal and glia subtypes and find differential expression of specific yolk proteins and long non-coding RNAs. However, whether any of these differences are driven by other canonical sex-specific genes such as Fruitless (Fru) or Double-sex (Dbx) has not been reported or discussed. The authors must re-analyze their data for these genes and claim whether they have any contribution to sex-specific sub-clusters.

Thank you for pointing this out. We have made text changes and clarifications to highlight the expression of other canonical sex-specific genes. *Fru* was enriched in male nuclei as expected. Interestingly, *dbx* was enriched in female nuclei. It remains to be determined if these genes are mechanisms that may be driving sex-specific changes.

Lines 224-226: “Additionally, female nuclei were enriched for *dbx* (Supp Table 8). Male glial nuclei expressed higher levels of genes including the male-specific genes *lncRNA:rox1/2* and *fru* (Figure 5C; Supp Table 8) (Ryner et al., 1996; Amrein and Axel, 1997; Meller et al., 1997).”

Lines 237-239: “Male nuclei expressed higher levels of genes including the male-specific genes *lncRNA:rox1/2* and *fru* (Figure 5G; Supp Table 9) (Ryner et al., 1996; Amrein and Axel, 1997; Meller et al., 1997).”

Lines 428-431: “We found the expected differential expression of yolk proteins (*yp1*, *yp2*, *yp3*) enriched in female nuclei and the long non-coding RNAs *rox1/2* and *fru* enriched in male neuronal nuclei (Ryner et al., 1996; Amrein and Axel, 1997; Meller et al., 1997; Warren et al., 1979). Interestingly, we found *dbx* to be enriched in both glial and neuronal female nuclei.”

Lines 433-435: “It remains to be determined if these genes are driving these sex-specific differences in glia and neurons.”

(3) In Figure 6C, it is unclear whether the Ms-2A-LexA-expressing neurons of clusters 157 and 160 project to two different neuropils or share projects to both neuropils. However, it is not explicitly shown in the immunostaining data whether indeed there are two populations to begin with. The authors must check for cluster 157 and 160 specific markers (such as Dh44 and ple) and test whether they appear mutually exclusively in the Ms-2A-LexA-expressing neurons. The same reasoning would apply to the data shown in Figures 6D and 6E, where the authors must test whether the NPF and AstA expressing cells are indeed neurons from clusters 100 and 128, using orthogonal cluster markers to conclude that they are similar (or the same) neurons.

We changed the focus of the paragraph to confirm that these neurons indeed come from type II and that they target the central complex. Although due to the lack of reagents we cannot test the identity of each one of these neurons, we could make meaningful interpretations of the staining to validate our ideas about neuropeptidergic cells in the central complex. We made sure to mention the limitation of our experiment to avoid any wrong conclusions.

Minor points

(1) Line 115 - "cluster that represents optic lobe neurons". How was this cluster identified?

We reexamined the most significant genes enriched in this cluster 124, and found they are *Rh2*, *ninaC*, *trpl*, and phototransduction related genes (Supplemental table 1). We reassigned the identity of this cluster as ocelli, which also express photoreceptor genes but can't be easily removed during dissection. We modified the text as follows:

"We used known markers (Croset et al., 2018; Davie et al., 2018; Supp Table 2) to identify distinct cell types in the central brain, including glia, mushroom body neurons, olfactory projection neurons, clock neurons, Poxn+ neurons, serotonergic neurons, dopaminergic neurons, octopaminergic neurons, corazonergic neurons, hemocytes, and ocelli (Figure 1B, Supp. Table 1)."

(2) As the separation in Figure 1B is not obvious, annotated cell type clusters must be re-colored instead of being labelled as the exact dots are indistinguishable. This would especially be helpful for OCTY, SER, OPN, and CLK clusters.

(3) Cluster labels in Figure 1C are barely visible and the font size must be increased for the reader. Recoloring the cluster identities and attaching a legend would again help in this case.

We recolored the atlas in Figure 1B, 1C and 1C' and increased the font size in Figure 1C'.

(4) For Figure 4A, clusters should be labelled on the UMAP along with the legend as it is difficult for the reader to match identities using Seurat colors. The same is true for the UMAPs in Figure 5A.

Yes, we agree that labeling would improve readability and have done so for UMAPs in Figure 4A and 5A-A'.

Reviewer #2 (Recommendations for the authors):

In this manuscript, Epiney et al., present a single-nucleus sequencing analysis of adult central brain neurons and glia. Through the use of an ingenious permanent labeling technique, they are able to trace the progeny of T2 neuroblasts, which contribute significantly to the formation of the central complex. This transcriptomic dataset is the first of its kind and will likely serve as a valuable resource for future studies.

The authors further explore this dataset through several analyses, including the characterization of neurotransmitter expression profiles in T2-derived neurons. However, the approach used to identify the identity of each neuron cluster could be more clearly articulated, and some of the authors' conclusions are more generalized - either already well-established or lacking sufficient support.

Detailed comments:

Abstract - "Our data support the hypothesis that each transcriptional cluster represents one or a few closely related neuron subtypes. - Is this a novel finding? If so, it would be helpful if the authors could explain why this is the case more clearly.

Our results are not generally novel, and many single cell/single nuclei RNA-seq papers have been published (more citations added to Introduction). Our work is novel in that we analyze Type 1 and Type 2 neuroblasts in the central brain.

Line 53 - In the introduction the authors should also reference other single-cell studies done in the Drosophila brain.

Done.

Line 59 - There are some typos here. The authors could also mention type zero.

Both done.

Figure 1 and Sup Table 1 - Authors show in sup table 1 the top cell markers by cluster but there is no correspondence between cluster number and identity. The authors do not say which known markers were used to give the identity to each cluster.

We have added the cell identity in the Supplemental Table 1. For the unknown cells, we left the column blank. We have also added a Supplemental Table 2 to show the markers we used to give identity to the clusters.

Supplementary Tables - For each table, more detailed information should be provided regarding what is being compared and the methods used for these comparisons.

We have added the methods we used in Seurat to generate each individual table.

Line 138 - Differential gene expression analysis between T1 and T2 glial progeny did not show differences across any glial cell types (Supp Table 4). - Was this comparison done per cluster? Is differential gene expression of top markers, which are anyway the genes that define each glial cell type, enough for this type of analysis?

Yes, we performed the differential expression analysis using all genes (i.e., not just marker defining) at a cluster-by-cluster resolution with results in Supplemental Table 4. We have edited the text to make this clarification.

Lines 139-141: "Differential gene expression analysis for all genes between T1 and T2 glial progeny did not show differences across any glial cell types or clusters (Supp Table 4)."

Line 146 - We identified T1-derived neurons by excluding cells co-expressing T2-specific. Markers FLP+/GFP+/RFP+ plus repo+ glial clusters. - Bioinformatically, correct?

Yes. We clarified the sentence as follows:

"We identified T1-derived neurons by bioinformatically excluding cells co-expressing T2-specific markers FLP+/GFP+/RFP+ plus repo+ glial clusters."

Line 156 - We found that each cluster strongly expressed a unique combination of genes. - As they are grouped by seurat in different clusters, why is this surprising?

Line 175 - "top 10 significantly enriched genes gathered from each T2 neuron cluster" - can these lists be included?

Yes they are grouped by Seurat. We toned down the sentence and refer each combination of genes as cluster markers. We modified the sentences as follows:

Each unique combination of enriched genes could be referred to as cluster markers.

Line 211- How did the authors identify sex-biased clusters? How did the authors separate the samples/cells by sex? Was it done bioinformatically by the expression of certain genes? If so, which?

We collected male and female nuclei separately. We have added text in the methods section as follows:

"Equal amounts of male and female central brains (excluding optic lobes) were dissected at room temperature within 1 hour. The samples were flash-frozen in liquid nitrogen and stored separately at -80°.

In the first round, we pooled male and female brains together to select GFP+ nuclei and used particle-templated instant partitions to capture single nuclei to generate cDNA library (Fluent BioSciences, Watertown, MA). In the second and third rounds, RFP+ nuclei from male and female brains were collected separately. The split-pool method was then used to generate barcoded cDNA libraries from each individual nucleus."

Are there sex-specific differences in genes in glia other than genes that were previously known to be sex-specific?

We report the comprehensive list of sex-specific differences in gene expression for both glia and neurons in Supp tables 8 and 9.

Line 237 - When the authors mention "We conclude that male and female adult T2 neurons have sex-specific differences in gene expression within the same neuronal subtype" does this mean that these neurons are the same in male and in female brains, but they additionally specifically express sex-specific genes?

Yes, we report that male and females contain the same neurons defined by their transcriptional profile. It remains to be seen if this sex-specific differences changes how these same neuronal subtypes function between male and females. We have added additional text in the discussion to expand on this thought.

Lines 437-441: "It remains to be determined if these genes are driving sex-specific differences within glial and neuronal subtypes. These genes may reflect sex-specific differences in the adult central brain and may provide insight into how behavioral circuits are linked to sex-specific behaviors. Future work should aim to characterize and test these genes."

Line 250 - The idea behind these sections "What is the relationship between neuropeptide expression and cluster identity?" "relation between cluster and morphology" lacks clarity. As clusters are defined based on principal component analysis, and the genes used to define a cluster are dependent on this method, there is no assumption that each cluster represents only one type of neuron or that it should include only neurons expressing the same neurotransmitter genes. Even if some clusters consist of a single neuron type, this should not be generalized to all clusters (and vice-versa).

Correct, we cannot determine from the transcriptome data whether distinct clusters will have different morphology. We have changed the focus of the question to address that we are confirming they come from type 2 and that they target the central complex while comparing to known cells that express the neuropeptide.

Line 265 - We first assayed the neuronal morphology of Ms+ neurons - why did the authors choose these neurons?

Resolved in main text: "we found that type II-derived Ms-2A-LexA-expressing neurons project to multiple layers of the dorsal fan-shaped body and the entire ellipsoid body, suggesting an unknown class of Ms+ neurons targeting to EB and/orFB".

Line 268 - "Currently we can't determine whether Ms+ neurons in clusters 157 and 160 project to different CX neuropils, or whether neurons from both clusters share projections into both neuropils. " - The purpose of this point is unclear.

Resolved in text: "we found that type II-derived Ms-2A-LexA-expressing neurons project to multiple layers of the dorsal fan-shaped body and the entire ellipsoid body, suggesting an unknown class of Ms+ neurons targeting to EB and/or FB".

Line 279 - This analysis could be more explored.

Thank you for your feedback. As the comment was somewhat broad, we were unsure of the specific revisions needed and have therefore left the text unchanged.

Line 301 - The text regarding this section, and the description and details of respective figures should be proofread to ensure clarity.

Done.

Line 386 - Alternatively, co-expression may be due to background from RNAs released during dissociation. - RNA in soup could be bioinformatically analysed.

Correct. We opted to delete this sentence since our split-pool based method does not create background RNA expression. Additionally, the analysis is performed on scaled expression >2, and any background RNA is unlikely to yield such high expression.

Discussion - Some of the conclusions are a bit too general, suggesting that the results might be meaningful, but also acknowledging the possibility of artifacts. If the authors could refine this, it would strengthen the manuscript.

We are sorry but we are uncertain what you are asking; we don't know what you want us to refine. Our apologies for the misunderstanding.

<https://doi.org/10.7554/eLife.105896.2.sa0>