

Reviewed Preprint

v1 • March 16, 2026

Not revised

Reviewed Preprint

v2 • March 30, 2026

Revised by authors

✉ For correspondence:

bfl20@cam.ac.uk

* Present address: Molecular Microbiology and Structural Biochemistry (MMSB), CNRS, UMR 5086, University of Lyon, Lyon, France

** Present address: School of Biochemistry, Faculty of Life Sciences, University of Bristol, Bristol, United Kingdom

* These authors contributed equally

Competing interests: No competing interests declared

Funding: See [page 31](#)

Reviewing editor: Lejla Zubcevic, University of Kansas Medical Center, United States

© 2026, Horne et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

A lipoprotein partner for the *Escherichia coli* outer membrane protein TolC

Jim Horne^{1,*}, Elise Kaplan^{1,*}, Ben HS Jin¹, Kieran Abbott², Victor Flores¹, Emmanouela Petsolari¹,

Jan M Gradon^{1,**}, Yvette Ntsogo¹, Andrzej Harris¹, Dingquan Yu¹, Ashraf Zarkan², Ben F Luisi¹ ✉

¹Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom • ²Department of Genetics, University of Cambridge, Cambridge, United Kingdom

eLife Assessment

In this **fundamental** work Horne et al present **compelling** evidence that YbjP is a novel binding partner of the TolC channel protein. The YbjP is characterized using cryo-EM, and its role probed using pull-down experiments, in vivo crosslinking, functional assays along with phylogenetic analysis which are all properly performed and presented and support the main conclusions. While the study does not identify a clear role for this protein, the revised manuscript offers improved clarity and contributes invaluable insight into membrane transport and antimicrobial resistance.

<https://doi.org/10.7554/eLife.110666.2.sa2>

Abstract

The outer-membrane protein TolC from *Escherichia coli* belongs to an extensive superfamily whose members are found throughout the didermal, Gram-negative bacterial lineages. The protein serves as an activated exit duct in multi-drug efflux pumps and protein secretion machinery. Many TolC homologs bear a lipid modification on the N-terminus that embeds into the inner leaflet of the outer membrane and appears to have been a conserved feature; however, the moiety is absent entirely in the *E. coli* TolC. We have discovered that the *E. coli* lipoprotein YbjP interacts extensively with the periplasmic surface of TolC and its N-terminal lipid moiety is embedded in the membrane, mimicking the intramolecular and modification-membrane interactions seen in TolC homologs. Here, we present cryo-EM structures of the MacA-MacB-TolC and AcrA-AcrB-TolC tripartite pumps complexed to YbjP. Although the association occurs spontaneously both *in vitro* and *in vivo*, the YbjP-TolC interaction is not required for efflux activity under standard laboratory conditions. YbjP may contribute stabilising the orientation and distribution of TolC in the outer membrane as well as the expression of transporters for tryptophan and cyclic peptide toxins.

Introduction

The bacterial cell envelope is a highly complex compartment and protective barrier against the external environment, ensuring survival in response to chemical threats while maintaining individual cellular identity and communication of signals within communities. In Gram-negative species, including clinical pathogens such as *Escherichia coli*, *Pseudomonas aeruginosa*, *Haemophilus influenzae*, and *Helicobacter pylori*, the envelopes are characterised by an organisation of two lipid bi-layer membranes and an interstitial partition, referred to as the periplasm, that encompasses a peptidoglycan sacculus conferring mechanical robustness (Guest and Silhavy, 2023 [↗](#); Gumbart et al., 2021 [↗](#)). The envelope is home to a diversity of dedicated machineries that selectively move molecules across the barrier, facilitated by energy transduction. Harmful compounds such as antibiotics and bactericidal agents as well as effector proteins are

displaced through this barrier by energy-dependent machines, and representative and well-characterised examples are the tripartite assemblies AcrA-AcrB-TolC and MacA-MacB-TolC of *E. coli*, which are powered respectively by proton motive force and ATP binding and hydrolysis (Du et al., 2018 [DOI](#)). The AcrA-AcrB-TolC assembly drives efflux of chemically diverse compounds, and MacA-MacB-TolC can transport macrolides, peptide toxins and host antibacterial peptides (Honeycutt et al., 2020 [DOI](#); Zhang et al., 2025 [DOI](#)). Structures of these and analogous assemblies have provided insight into how the machines operate to recognise and move complex transport substrates in a preferred direction against concentration gradients (Du et al., 2018 [DOI](#), 2018 [DOI](#); Fitzpatrick et al., 2017 [DOI](#); Glavier et al., 2020 [DOI](#); Tsutsumi et al., 2019 [DOI](#)).

The outer membrane component of the efflux assemblies, TolC, is a homotrimeric protein with a self-closing β -barrel topology that inserts into the outer membrane and a long helical structural domain that interacts with periplasmic partners such as AcrA or MacA. TolC homologs are found throughout diderm bacteria, highlighting the functional importance of this superfamily (Stubenrauch et al., 2022 [DOI](#)). The long helical portion of TolC can be subdivided into an α -helical barrel domain, which appears to be a stable structural unit (Calladine et al., 2001 [DOI](#); Huang et al., 2014 [DOI](#)), and conventional coiled-coil segments that dilate from a sealed resting state to an open-state when interacting with a periplasmic partner (Wang et al., 2017 [DOI](#)). At the junction of the two helical regions, the polypeptide meanders to form a circumferential ring known as the equatorial domain that is proposed to interact with the peptidoglycan layer of the periplasm (Shi et al., 2019 [DOI](#)).

Here, in our analysis of the MacA-MacB-TolC tripartite assembly, we observed unexpected cryo-EM density spanning between the equatorial domain of TolC and the transmembrane boundary. Using a structural topology pattern recognition query against predicted folds of *E. coli* proteins, we identified the protein as a lipoprotein of unknown function, YbjP (UniProt ID P75818). The N-terminus of YbjP trails over the surface of TolC to reach the outer membrane where its lipid moiety is embedded. Our structural comparisons, together with functional and proteomic results from YbjP deletion strains, do not support an allosteric or signalling role for the lipoprotein in TolC-mediated efflux systems but suggest that YbjP is required for expression of transporters including TnaB tryptophan permease and YojI ABC transporter that exports toxins such as microcin J25 through TolC (Baquero et al., 2024 [DOI](#); Delgado et al., 2005 [DOI](#)).

Results

Cryo-EM structure of TolC bound to a lipoprotein

Earlier cryo-EM work on the MacA-MacB-TolC assembly used samples reconstituted in detergents (Fitzpatrick et al., 2017 [DOI](#)). We explored the preparation of MacA-MacB-TolC in peptidiscs as a better mimic of the membrane environment (Carlson et al., 2018 [DOI](#)). From these novel preparations, we were able to solve the structure by cryo-EM with high-quality density for all three components of the assembly (Figure 1A [DOI](#)). In one of our preparations, additional density was observed near the equatorial domain of TolC, extending towards the outer membrane β -barrel domain (Figure 1B [DOI](#)). We refined this structure to a final resolution of 2.48 Å (Supplementary Figure S1 [DOI](#) and Table S1 [DOI](#)) and fitted the additional density with secondary structural elements (Figure 1C [DOI](#)). The incomplete candidate fold was then searched against the database of *E. coli* protein structures predicted by AlphaFold2 (Jumper et al., 2021 [DOI](#)) using the DALI topology recognition program (Holm, 2022 [DOI](#)). The result matches with the lipoprotein YbjP (P75818) which could be fitted into the cryo-EM map. The refined model is presented in Figure 1D [DOI](#). The lipoprotein globular domain is formed by a series of four consecutive α -helices (α 1- α 4), followed by four antiparallel β -strands (β 1- β 4) and a C-terminal α -helix (α 5) (Figure 1E [DOI](#)). Structural alignment of the predicted YbjP and cryo-EM refined models reveals a strikingly high degree of similarity (RMSD of 0.53 Å over 118 residues), confirming the accuracy of the prediction (Figure 1D [DOI](#), inset). A superposition of the MacAB-TolC-YbjP complex with our previously reported MacAB-TolC structure (Fitzpatrick et al., 2017 [DOI](#)) revealed that TolC and the tripartite pump are largely unaffected upon YbjP binding with an RMSD of 1.03 Å over 1258 residues for

TolC trimer (Figure 2A [↗](#)). Nonetheless, the side chains of a few residues in TolC, which mainly correspond to positively charged amino acids (R18, R24, K214, R227, R234), reorient to interact with the YbjP lipoprotein partner (Figure 2B [↗](#)).

Interactions between TolC and lipoprotein YbjP

YbjP straddles two adjacent TolC protomers, and the interface is stabilised by multiple hydrogen bonding and ionic interactions involving the linker and globular domain of YbjP and two TolC subunits (Figure 2C [↗](#)). The intermolecular contacts engage almost 20 residues in both TolC and YbjP. The lipoprotein flexible linker runs along TolC N-terminal helix from the equatorial to the transmembrane domains and is stabilised by several hydrogen bonds involving N32, R35, R24, Y98 and L17 from one TolC monomer (Figure 2C [↗](#), panel 1). This cluster is consolidated by hydrogen bonding interactions between S107, S109, T110 and N113 from YbjP loop connecting beta-strands β 1 and β 2 and R234, Q306 and Y307 from the proximal adjacent TolC monomer, maintaining this flexible loop perpendicular to the lipoprotein linker. YbjP D115 and N118 also make contacts with R18 and E314 from TolC.

Interestingly, the YbjP globular domain is strongly associated via a cluster of ionic interactions (Figure 2C [↗](#), panel 2), mediated by a trio of positive, negative, positive residues (R122, E135, R151) which each contacts a residue of opposite charge in TolC (D231, R227, E317). Finally, numerous hydrogen bonds are formed between YbjP globular domain and helices from both TolC monomers (Figure 2C [↗](#), panel 3) engaging G155, V157 and A159 backbones, and K70, N118, E135 and R151 side chains from the lipoprotein. Residue conservation of YbjP was evaluated visually with the server CONSURF, revealing that the conserved residues are located at the interface with TolC (Figure 2D [↗](#)).

Like other lipoproteins, YbjP is predicted to be processed by cleavage of the signal sequence at residue C19 (also referred as to the +1 Cys of the matured lipoprotein), then lipidated as N-palmitoyl and S-diacylglycerol cysteine. These processing and modification steps are likely to occur at the inner membrane (Guest and Silhavy, 2023 [↗](#)). Although the density is not well defined for the lipid moiety due to its flexibility, the map does reveal a portion of the hydrocarbon embedded into the inner leaflet of the outer membrane. This is consistent with mass spectrometry studies that have identified YbjP as an outer membrane protein (Molloy et al., 2000 [↗](#)). Interestingly, there is no density for the lipoprotein R3 acyl chain suggesting that the cysteine α -amino group has not been modified while *E. coli* lipoproteins are presumed to be tri-acylated to traffic via the Lol system (Grabowicz and May, 2026 [↗](#); Noland et al., 2017 [↗](#)). Further characterisation is needed to establish if YbjP is naturally di- or tri-acylated.

Validation of the TolC-YbjP interaction by in vitro pull-down and in vivo photo-crosslinking

To validate the TolC-YbjP interactions, we used an *in vitro* affinity pull-down assay with purified proteins. For this, we produced a soluble mutant of YbjP, hereafter referred to as YbjP_s, removing the lipobox conserved cysteine that carries the acyl chains, and the 8 following residues corresponding to a flexible linker joining the acylated cysteine to the protein globular domain. The protein (D28-R171) with a N-terminal His-tag was purified from the cytoplasm and immobilised on nickel resin. As shown in Figure 3A [↗](#), detergent-purified TolC remained attached to the resin only in the presence of YbjP_s, confirming the association between the two proteins and demonstrating that the lipoprotein lipid moiety is not required for the interaction to occur. We could not detect an interaction between YbjP_s and Pal (peptidoglycan associated protein) lipoprotein or between Pal and TolC using the pull-down assay.

Interactions between TolC and YbjP_s were also measured by isothermal titration calorimetry. (Figure 3B [↗](#) and Supplementary Table S3 [↗](#)). The binding event is exothermic, consistent with electrostatic interactions at the protein-protein interface. The dissociation constant is in the micromolar range ($K_D \sim 4 \mu\text{M}$). Although this value represents a weak to moderate association, the affinity is likely to be substantially higher in the presence of full-length YbjP which includes the 8-




Figure 1. TolC partner identified by cryo-EM.

(A) Cryo-EM density map (left) resolved here and model (right) for the tripartite pump MacA-MacB-TolC (PDB 5NIL). Density for the MacB dimer is less well defined. (B) Close-up view on the TolC region showing additional density in green. TolC domain organisation is indicated. (C) Modelling of secondary structural elements fitting in the additional density. (D) The refined structure of the TolC partner, YbjP. After lipoprotein maturation, the N-terminal cysteine carries a lipid modification which is not shown here. Alignment of the YbjP structure from cryo-EM and corresponding AlphaFold prediction model is shown as inset. (E) Sequence and secondary structure alignment annotation of mature YbjP. The tri-acylated cysteine (C19) conserved in lipoproteins is underlined.




Figure 2. Interactions in the TolC-YbjP complex.

(A) Superposition of TolC from the MacAB-TolC-YbjP assembly reported this paper in blue (TolC) and yellow (YbjP) (PDB 9QGY) and from the previously reported MacAB-TolC complex in green (PDB 5NIK). (B) Close-up view on the YbjP-TolC interface with TolC residues involved in the interaction shown as sticks. (C) Overview of protein-protein interactions with close-up views presented in panels 1, 2 and 3. The YbjP lipoprotein (yellow) contacts two adjacent protomers of the TolC trimer (blue). Lipoprotein acyl modification and residues involved in intermolecular contacts are shown in stick representation. (D) Residue variability analysis of YbjP highlighting the clustering of conserved residues at the TolC interface. Analysis was performed with the CONSURF server. YbjP is shown as spheres with the lipidation in yellow.




Figure 3. In vitro and in vivo validation of the TolC-YbjP interface.

(A) *In vitro* binding assay between TolC and soluble YbjP that is missing the lipid modification. TolC-FLAG was mixed with N-terminally His-tagged soluble YbjP (YbjP_s) or soluble LolB (LolB_s) prior immobilization on IMAC resin. After several washes, the elution fractions (Elu) were analysed by SDS-PAGE. Molecular masses of protein standards (M) are indicated. (B) Isothermal titration calorimetry (ITC) profile for the interaction of TolC and YbjP_s. Background-corrected heats of injection are shown together with a fitted binding curve in red. The raw thermograms corresponding to the injection of YbjP_s into a TolC-containing cell or into buffer are shown in upper and lower insets, respectively. ITC parameters are listed in [Supplementary Table S3](#). (C) *In vivo* validation of the interaction between YbjP_{FL} and TolC by photo-crosslinking. The experimental procedure was validated using another periplasmic lipoprotein, AcrA, which associates with TolC. As shown underneath, AcrA residues Q136 and Y137, proximal to TolC in the structure of the AcrABZ-TolC pump (PDB 5NG5), were replaced by pBPA. For YbjP, the two residues N113 and T110 proximal to TolC in the MacAB-TolC-YbjP complex (PDB 9QGY) and the three residues N43, N90 and H104 distal to TolC were mutated. In brief, *E. coli* C43 cells expressing YbjP_{FL,His} or AcrA_{His} mutants carrying a photoactivatable pBPA group at indicated positions were grown in the presence (+) or absence (-) of pBPA and irradiated (+) or not (-) with UV. Cells were then lysed by successive freeze-thaw cycles, solubilized with DDM and lysates purified by Ni-NTA chromatography. Eluted proteins were analysed by SDS-PAGE and immunoblotting using anti-His antibodies. Purified YbjP_s was loaded as a benchmark control. Presence of crosslinked products are indicated with a black arrow. Note that FL YbjP migrates faster than its predicted molecular weight (~19 kDa), likely due to the presence of the lipoprotein lipid chains. A double detection of YbjP_{FL,His} and endogenous TolC proteins for the N113 sample is shown in [Supplementary Figure S2](#).

residue linker connecting the lipid modification to the globular domain. This region is absent here but participates in favourable interactions with TolC and is likely to increase the affinity (Figure 2A, panel 1). Additionally, the lipid modification would localise YbjP to the membrane, reducing its configurational entropy and thereby increasing TolC binding.

To validate that YbjP and TolC interact within the cell, we engineered C-terminally His-tagged full-length YbjP (YbjP_{FL}) to contain the photo-crosslinkable amino acid para-benzophenone, pBPA (Chin et al., 2002; Young et al., 2010) at positions guided by the YbjP-TolC structure. We replaced residues at the protein-protein interface (T110, N113) or at a distance (N43, N90 and H104) as negative controls. To validate the experiment, we also constructed amber mutants of His-tagged AcrA, an inner membrane protein known to interact with TolC, and substituted Q136 and Y137, two residues proximal to TolC in the AcrABZ-TolC efflux pump structure (Wang et al., 2017). Cells were grown in a medium supplemented with pBPA, induced and then irradiated for 15 min with ultraviolet light. After cell lysis, His-tagged proteins were retrieved from the samples by affinity chromatography. Purified fractions were analyzed by SDS-PAGE and immunoblotting (Figure 3C). We observed covalent photo-adducts that migrate at higher molecular weight compatible with AcrA-TolC or YbjP-TolC adducts only for the pBPA-supplemented cells and only after illumination with UV light for the residues that are proximal to the TolC surface, but not those that are distal. We confirmed the presence of both YbjP and TolC in the N113 YbjP crosslinked product by Western blot (Supplementary Figure S2). These results are consistent with a model for YbjP lipoprotein interactions with TolC *in vivo*.

TolC-YbjP interaction in the AcrA-AcrB-AcrZ-TolC assembly

To examine whether YbjP can interact with other TolC-mediated assemblies, we supplemented the AcrABZ-TolC pump reconstituted in peptidisc with YbjP_s and resolved the cryo-EM structure of the complex at 3.17 Å resolution (Supplemental Figure S3 and Table S1). Structural comparison of the two assemblies reveals that YbjP binds at the same location on TolC (Figure 4A, B and Supplementary Figure S3) and makes the same interactions (Figure 4C). The AcrABZ-TolC-YbjP assembly was also reconstituted into peptidoglycan (Supplementary Figure S4A, B), and the position of the peptidoglycan layer between AcrA and TolC corresponds with the localization previously inferred from low-resolution cryo-ET data (Shi et al., 2019) (Supplementary Figure S4C). The reconstructed maps indicate that the YbjP may not interact with the peptidoglycan layer, but this cannot be certain due to the limited resolution of the cryo-EM map. *In vitro* characterizations indicated that YbjP can interact with peptidoglycan in pull-down assays, but this is likely to be non-specific as alanine substitutions in the domain proximal to the peptidoglycan (R80, D81, R85 and E86) had little impact on the efficiency of the pull-downs. Photo-crosslinking in substitutions of the photoreactive amino acid para-benzophenone (pBPA) at positions D77, Q166, E169 and R171 did not give evidence for peptidoglycan-linked species. The data suggest that YbjP may not form avid interactions with the peptidoglycan.

Evolutionary profile of YbjP

To explore the possible function of YbjP, we investigated its evolutionary profile. In the PFAM database, YbjP is annotated to contain one DUF3828 domain. The most well-characterized DUF3828-containing protein is Tai3, a periplasmic type VI cognate immunity protein of the T6SS amidase effector Tae3 (Dong et al., 2013; Russell et al., 2014). Another periplasmic protein YqhG contains one full and one partial DUF3828 domain and is involved in regulating type I fimbriae expression in *E. coli* by an unknown mechanism (Bessaiah et al., 2019). Structural alignment of YbjP with the Tai3 crystal structure and YqhG AlphaFold2 model indicates that these proteins share a common fold (Figure 5A).

To investigate whether YbjP is functionally homologous to Tai3 or YqhG, we reconstructed a phylogenetic history of the DUF3828-containing proteins, which could be classified into three major families based on taxonomy, gene synteny, presence of a signal peptide and a conserved QDX motif, involved in Tae3 inhibition (Supplementary Table S4). The result indicates that the function of YbjP is likely distinct from that of Tai3 or YqhG (Figure 5B). We noted that Tai3 is




Figure 4. The structural comparison of AcrABZ-TolC-YbjP and MacAB-TolC-YbjP.

(A) Cryo-EM maps and models of AcrABZ-TolC-YbjP and MacAB-TolC-YbjP reconstituted in peptidisc. (B) Superposition of TolC from the AcrABZ-TolC-YbjP and MacAB-TolC-YbjP assemblies reported here (red: TolC, green: YbjP, PDB 9TG4; blue: TolC and yellow: YbjP, PDB 9QGY). (C) Close-up views on the YbjP-TolC interface in the AcrABZ-TolC-YbjP (top) and MacAB-TolC-YbjP (bottom) structures.




Figure 5. YbjP evolution and co-evolution with TolC.

(A) Structural alignment of YbjP with other DUF3828-containing proteins: Tai3 (PDB 4HZ9, purple) and YqhG (AlphaFold2 Q46858, blue). A 90° rotation along the x-axis is shown underneath. (B) A cladogram of DUF3828-containing proteins. The reviewed proteins (Tai3, YbjP and YqhG) are marked by red arrows. (C) Surface conservation of TolC proteins from Enterobacterales containing or not containing YbjP. Results are displayed on *E. coli* TolC on the left and on *P. atrosepticum* TolC (AlphaFold model, Q6DAC5) on the right. For clarity, YbjP is displayed on the left panel but not on the right. Analysis was performed with the CONSURF server.

found in a broader range of bacterial groups (Gammaproteobacteria, Betaproteobacteria, and Bacteroidota/Chlorobiota) compared to YbjP. Similarly, YbjP is restricted to some members of the Enterobacterales Order within the class of Gammaproteobacteria, unlike TolC which is more widespread (Supplementary Figure S5 [\[4\]](#)). Notably, the TolC-YbjP interface is more conserved in Enterobacterales with YbjP than without (Figure 5C [\[4\]](#)), suggesting a scenario where YbjP evolved recently from an ancestral Tai3-like protein to become a TolC interactor.

YbjP is not required to compensate for lack of TolC lipidation

Outer membrane efflux proteins structurally homologous to TolC often carry a N-terminal acylation. This is the case for the drug exit duct OprM in *Pseudomonas* and the outer membrane efflux proteins in *E. coli* MdtP, MdtQ, and CusC, but with the notable exception of TolC. The reason for the presence or absence of acylation remains unclear, but it has been proposed that acylation has roles in anchorage to the membrane and in membrane insertion. We first postulated that YbjP could compensate for the absence of lipid modification in TolC. Indeed, YbjP linker and N-acetylation mimic the linker and acetylation arrangement seen in OprM and CusC (Figure 6A [\[4\]](#)). Similarly, AlphaFold3 predicted the interaction of YbjP with TolC but not with the other TolC structural homologues that are lipidated in *E. coli* (Figure 6B [\[4\]](#)). Consistent with this, we found that *E. coli* TolC is distinct from its lipidated structural homologues (Figure 6C [\[4\]](#)), suggesting the possibility that YbjP evolved to compensate for the absence of lipidation. However, further bioinformatic analysis reveals that while most organisms containing the *ybjP* gene possess non-lipidated TolC, non-lipidated TolC is also present in most organisms lacking *ybjP* (Figure 6D [\[4\]](#)). This indicates that YbjP may not be necessary to compensate for the lack of lipidation of TolC.

Exploring in vivo function of YbjP

To explore the physiological function of YbjP, we compared the proteomic profiles of $\Delta ybjP$, $\Delta tolC$, $\Delta tolC \Delta ybjP$ and parental strains under exponential and stationary phase conditions (Figure 7 [\[4\]](#)). Only minor differences were detected in the proteome of the $\Delta ybjP$ strain compared with the parental wild-type in both growth stages, while numerous significant changes were observed for the $\Delta tolC$ and $\Delta tolC \Delta ybjP$ strains. Levels of TolC were not impacted in the $\Delta ybjP$ strain, indicating that YbjP is not required for TolC biogenesis. Previous RNA-seq analysis highlighted an upregulation of *ybjP* in response to membrane-disrupting chemicals, including vanillin (Patrick et al., 2019 [\[4\]](#)) and EDTA (Janssens et al., 2024 [\[4\]](#)). However, we do not observe an impact on growth under standard laboratory conditions, as the minimal inhibitory concentrations for growth inhibition were unchanged in the $\Delta ybjP$ compared to the parental strains in the presence of vanillin and EDTA; nor were changes observed with benzalkonium chloride and different classes of antibiotics, under either aerobic or microaerobic conditions (Supplementary Table S5 [\[4\]](#)).

Considering that YbjP expression is regulated by the stress response sigma factor *rpoS* (Lacour and Landini, 2004 [\[4\]](#)), its physiological role is likely context-dependent and not readily appear under standard laboratory conditions. Indeed, the few proteins whose levels were influenced in the $\Delta ybjP$ strain are associated with specific contexts, including stress and changes in lifestyle (Figure 7 [\[4\]](#)). We observed the downregulation of Antigen 43 (*fla*), which is an autotransporter porin with an extensive extracellular domain that is involved in auto-aggregation and which promotes the shift from a planktonic state to sessile state. While previous work noted that YbjP overexpression influences motility (Tenorio et al., 2003 [\[4\]](#)), we did not observe any effect of the *ybjP* deleted strain on LB motility agar plates (Supplementary Table S6 [\[4\]](#)).

In addition, changes in specific transporters were noticed. The YojI ABC transporter, proposed to be involved in the efflux of the microcin J25 toxin in conjunction with TolC, was decreased in relative abundance in the $\Delta ybjP$ strain in stationary phase (Figure 7 [\[4\]](#)). However, we did not observe any change in MIC values for the $\Delta ybjP$ strain in the presence of microcin J25 during exponential phase, while the $\Delta tolC$ and $\Delta tolC \Delta ybjP$ strains did show increased sensitivity of >4,000-fold (Supplementary Table S5 [\[4\]](#)). Conversely, the TnaB low-affinity tryptophan permease level increased. We did observe a small increase in growth for the $\Delta ybjP$ strain compared to the parental strain in minimal media in the presence of tryptophan. Interestingly, the $\Delta tolC$ and $\Delta tolC$




Figure 6. YbjP lipoprotein is not required for TolC trafficking in *E. coli*.

(A) Structural alignment of *E. coli* TolC (blue), CusC (green, PDB 3PIK,) and *P. aeruginosa* OprM (red, PDB 3D5K). The YbjP- or TolC-homologue linker is shown as transparent surface and modified cysteines are indicated by arrows. Dashed lines show the extension of the lipid group (partially- or non-modelled in the structures). (B) AlphaFold3 prediction (Abramson et al., 2024) between *E. coli* outer membrane efflux proteins and YbjP. The model with the highest ipTM and pTM scores is shown out of three technical repeats. (C) CLANS (Cluster Analysis of Sequences) analysis of outer membrane efflux proteins (IPR003423) in *Pseudomonadota*. Clustering was performed in 2D until equilibrium. Outer membrane proteins in *E. coli* are shown in red, and other key reviewed proteins in black. Yellow, lipitation; blue, secreted; grey, other. (D) Distribution of the *ybjP* gene and lipitated TolC in Gammaproteobacteria. Repartitions are as follows: *ybjP* present (light blue, n=67) or absent (magenta, n=525); lipitated (yellow, n=5) or non-lipitated (dark blue, n=587) TolC.



Horne, Kaplan et al., 2026 *eLife* **15**:RP110666. <https://doi.org/10.7554/eLife.110666.2>

AybjP strains did show marked decreased growth in the absence of tryptophan (Supplementary Figure S6 [\[4\]](#)). The outer membrane pore OmpW, which plays a minor role in the efflux of quaternary cationic compounds together with the EmrE transporter (Beketskaia et al., 2014 [\[4\]](#)), was downregulated. Taken together, the changes in the proteome implicate nuanced, multiple roles of YbjP, but the phenotypic impact of the null strain is likely to be seen under environmental stress conditions.

Discussion

Our structural and functional results indicate that the lipoprotein YbjP is located to the outer membrane, where it can interact with the membrane protein TolC. The mode of interaction between TolC and YbjP resembles closely the intramolecular interactions of the lipidated N-terminus in TolC paralogs, such as OprM from *Salmonella*. The interaction is likely functionally important within some Enterobacterales. It is proposed that the TolC archetype may bear sequence or environmental features that have limited its distribution, or it is a more recently evolved form in the TolC-like protein family (Stubenrauch et al., 2022 [\[4\]](#)). It is possible that TolC and YbjP co-evolved in this lineage.

The biogenesis of TolC presents intriguing mechanistic puzzles. The insertion stage of outer membrane proteins involves the β -barrel assembly machine (BAM) complex (Voulhoux et al., 2003 [\[4\]](#); Wu et al., 2005 [\[4\]](#)), but the process is not well characterised for a non-canonical porin-like protein with an extended hydrophilic portion, such as TolC, nor are the earlier steps of translocation across the periplasm. The chaperone translocation and assembly machinery (TAM) facilitates assembly of TolC into the outer membrane in a potentially redundant pathway (Stubenrauch et al., 2022 [\[4\]](#)). Reflecting on the potential requirements suggests key aspects to consider for understanding the process. First, the protomers must be moved across the periplasm from the translocon where the nascent chains emerge. Second, this insertion would be coordinated to deliver three protomers to fold into the full trimeric architecture. Lastly, the folding does not require exogenous energy and is likely driven by hydrophobic interactions with the highly crowded and asymmetric lipid bilayer (Horne et al., 2020 [\[4\]](#)).

Mechanistic puzzles also arise when considering how lipoproteins such as OprM and YbjP arrive at the outer membrane and are inserted. Lipoproteins are processed and lipid groups are attached to a conserved cysteine residue that immediately follows the signal sequence in the inner membrane (Guest and Silhavy, 2023 [\[4\]](#)). Acylated thiols are found in components of other bacterial efflux assemblies. The structure of CusC indicates the presence of an acylated thiol on the N-terminal cysteine, and acylation is also found in CmeC, but nothing has been found to be added to the N-terminal cysteine of the MtrE structure. Post-translational lipidation is particularly essential for the secretion and localization of some membrane proteins, a process involving different acyl transferases (Aicart-Ramos et al., 2011 [\[4\]](#); Kovacs-Simon et al., 2011 [\[4\]](#); Linder and Deschenes, 2007 [\[4\]](#); Nakayama et al., 2012 [\[4\]](#)). Why proteins that are embedded within cellular membranes via a large hydrophobic structural domain need modifications such as the N-terminal lipidation appears puzzling. Akama et al. (2004) [\[4\]](#) suggested that these proteins must first be anchored to the membrane by an N-terminal lipid so that the insertion of their large hydrophobic domain may be triggered. Structural evidence indicates that this later step is critical for the correct folding of outer membrane factors (OMFs) and highlight the possible involvement of membrane-interacting components in the OMF opening process and consequently the efflux-pump function (Lei et al., 2014 [\[4\]](#)).

Translocation of lipoprotein to the outer membrane in *E. coli* requires five essential proteins (LolA-E) (Guest and Silhavy, 2023 [\[4\]](#)). The mechanism involves lipoprotein extraction from the inner membrane by the ABC transporter LolCDE, then transfer to the periplasmic carrier LolA for delivery to the outer membrane acceptor LolB, which facilitates insertion into the outer membrane. It is envisaged that this pathway is involved in the movement of OprM and its lipidated outer membrane homologs. As *E. coli* TolC is not lipidated, it is possible that YbjP acts as a chaperone that intercepts the TolC and aids its translocation across the periplasm via the Lol system. However, this proposed pathway is likely redundant or not essential, as TolC is still found

in the outer membrane in strains null for YbjP. The BAM complex impacts on TolC assembly in *E. coli* (Malinverni et al., 2006 [↗](#); Werner and Misra, 2005 [↗](#)), but surprisingly it is not involved in assembly of the TolC-like proteins OprM in *P. aeruginosa* (Hoang et al., 2011 [↗](#)). The pathway for insertion is likely to be redundant and divergent for the TolC family (Stubenrauch et al., 2022 [↗](#)). Notably, a broad screen of cell envelope protein complexes in *E. coli* did not identify physical or genetic interactions between YbjP and TolC (Babu et al., 2018 [↗](#)), but such interactions were found between YbjP and the partners of TolC in the tripartite assembly, namely the RND transporter AcrB and the periplasmic AcrA. It may be that the interactions between TolC and YbjP *in vivo* are weak and transient, as might occur for example during TolC insertion into the outer membrane.

Our results do not support an allosteric or signalling role for YbjP but suggest that the lipoprotein impacts on expression of transporters tryptophan permease TnaB and ABC family member YojI, which transports toxins such as microcin J25 through TolC (Baquero et al., 2024 [↗](#)). It is also possible that YbjP may facilitate the biogenesis of TolC during times of rapid adaptation or its distribution in the membrane. To our knowledge, YbjP is the first protein identified to interact with the lateral surface of TolC. Interestingly, the outer membrane protein SlyB has been shown to encapsulate TolC-like proteins during stress conditions and cryo-EM 2D classification images predict the binding of SlyB at the TolC-YbjP interface (Janssens et al., 2024 [↗](#)). Structural predictions from AlphaFold3 further support that SlyB binds TolC at the YbjP location (Supplementary Figure S7 [↗](#)). Given that both YbjP and TolC are upregulated in response to EDTA treatment (Janssens et al., 2024 [↗](#)), YbjP and SlyB might compete for binding to TolC under conditions of outer membrane stress. Possibly YbjP might support clustering and partitioning of transport machineries during cell division or during stress (Zhang et al., 2025 [↗](#)), which in turn can influence the efficiency of adaptation to environmental conditions.

Materials and methods

Preparation of disulphide-engineered MacAB-TolC and reconstitution into peptidisc

E. coli BL21 (DE3) strain was transformed with plasmids pET20b-hexaHis-MacA_{D271C}-MacB_{G465C} expressing the MacA_{D271C} and MacB_{G465C} mutants, allowing a disulphide link, with a C-terminal hexaHis-tag on MacB and pRSFDuet-TolC-FLAG expressing full-length TolC with a C-terminal FLAG-tag (Fitzpatrick et al., 2017 [↗](#)). Transformed cells were spread onto LB agar plates containing 50 µg mL⁻¹ carbenicillin and 50 µg mL⁻¹ kanamycin and were incubated at 37°C overnight. A single colony was used to inoculate 50 mL of LB medium containing 50 µg mL⁻¹ carbenicillin and 50 µg mL⁻¹ kanamycin. The preculture was grown at 37°C, with an orbital shaker set at 200 rpm for 4 h. When the absorbance at 600 nm reached between 0.5 and 0.7, the preculture was used to inoculate 6 L of 2YT medium containing 50 µg mL⁻¹ carbenicillin and 50 µg mL⁻¹ kanamycin in 6x2 L baffled flasks. Bacteria were grown at 37°C, 200 rpm until OD_{600nm} reached 0.5 and were subsequently induced with 0.25 mM isopropyl β-D-1-thiogalactopyranoside (IPTG). The culture was further incubated at 20°C, 200 rpm overnight. Cells were collected by centrifugation (5,000 g for 30 min at 4°C) and pellets were resuspended in a buffer containing 20 mM Tris-HCl pH 8.0 and 400 mM NaCl (Buffer A). One tablet of EDTA-free protease inhibitor per 50 mL, 5 µg mL⁻¹ lysozyme and 5 U mL⁻¹ DNase I were added to the cell suspension and incubated 1 h at 4°C. The mixture was then passed five times through a water-cooled high-pressure homogeniser (Emulsiflex), keeping the pressure below 14,500 psi throughout. The resulting lysate was centrifuged (9,000 g, 30 min, 4°C) to remove cell debris and then ultracentrifuged (175,000 g, 4 h, 4°C) to pellet membranes.

Membranes were solubilized in 50 mL of buffer 20 mM Tris pH 8.0, 400 mM NaCl with 1.5% (wt/v) β-dodecyl maltoside (β-DDM) and one tablet of EDTA-free protease inhibitor for 3 h at 4°C with gentle stirring. The solution was then purified by Ni-NTA chromatography after addition of 10 mM imidazole. The elution peak was further purified with ANTI-FLAG M2 affinity resin (0.5 mL, Sigma Aldrich), pre-equilibrated with buffer A supplemented with 0.05% wt/v β-DDM. The elution fraction was added to the resin and incubated at 4°C for 1 h with gentle agitation. The sample was then washed with buffer A containing 0.05% β-DDM and then 0.03% (wt/v) decyl maltose

neopentyl glycol (DMNG). Peptidisc (Carlson et al., 2018) at 1 mg mL⁻¹ in 400 mM Tris-HCl and additional *E. coli* lipids (100 μ L at 20 mg mL⁻¹) were added to the sample on the resin. After 16 h at 4°C, the resin was washed with buffer A containing no detergent and then eluted with 3xFLAG peptide (Sigma-Aldrich). Flow-through was collected at each step and analysed by SDS-PAGE. The presence of the full assembly was confirmed by mass photometry (Refeyn).

MacAB-TolC cryo-EM sample preparation and data collection

For single-particle cryo-EM structure solution, the samples of MacAB-TolC complex reconstituted in peptidisc (Carlson et al., 2018) were thawed and diluted to 0.5 mg mL⁻¹ in a buffer composed of 20 mM Tris pH 8.0, 100 mM KCl, 2 mM TCEP and 1 mM NaN₃. A 3- μ L aliquot of dilute protein was applied onto a Cu 300-mesh EM grid layered with R1.2/1.3 holey carbon support film (Quantifoil Micro Tools, GmbH), which had been glow-discharged immediately prior to use. The sample deposited on the EM grid was blotted of excess solution and vitrified in liquid ethane using Vitrobot Mark IV (Thermo Fisher Scientific, Inc.) robotic cryo-EM plunger.

The grids were imaged using Titan Krios G3 transmission electron microscope (Thermo Fisher Scientific Inc.) operating at accelerating voltage of 300 kV and liquid nitrogen temperature. The images were recorded in EFTEM mode using AFIS on K3 direct electron detector (Gatan, Inc.) and data acquisition software EPU (Thermo Fisher Scientific, Inc.). Exposures were for 1 s and were collected as 40-frame movies using total electron fluency (dose) of 49.75 e⁻/Å² and defocus values between -2.4 and -1.0 μ m. The movies were collected in super-resolution mode with nominal pixel size 0.415 Å, but were binned 2 $\times$ following exposure, yielding effective pixel size of 0.83 Å/px.

MacAB-TolC cryo-EM data processing

The cryo-EM data were processed using cryoSPARC 4.5 software (Structura Biotechnology, Inc.). Patch motion correction was used to correct for beam-induced motion and Patch CTF used to calculate the CTF. Circular Gaussian blob picking with a min/max diameter of 120/380 Å was used to extract an initial set of pixels in 878-px boxes Fourier-cropped to 600-px (effective pixel size 1.21 Å/px). After 2D classification these were used as input to a template-picking model and particles re-extracted at 600-px and Fourier-cropped to 400-px (effective pixel size 1.245 Å/px). This set of particles was pruned again with another round of 2D classification and then subject to *ab-initio* reconstruction. The best volume and particles from 2D classification was used for heterogeneous refinement along with 5 ‘junk’ volumes. The best output was then subjected to non-uniform refinement in *C*₁ symmetry with optimize per-particle defocus, optimize per-group CTF parameters, and fit spherical aberration options used to further improve the refinement. This yielded a 2.74 Å map with 233,835 particles. Finally, local CTF refinement, global CTF refinement, and another round of non-uniform refinement in *C*₁ was performed followed by reference-based motion correction. The unbinned output particles (233,775 particles at 600-px box size, 0.83 Å/px) were subjected to a final round of non-uniform refinement in *C*₃ symmetry yielding a final map with a GSFSC reported resolution of 2.48 Å. To facilitate model-building, Resolve from Phenix package was also run on this map. Data collection and processing parameters are shown in Supplementary Table S1.

MacAB-TolC-YbjP model building and refinement

The part of the experimental map containing an unidentified protein density adjacent to TolC trimer was visualised and traced as a poly-alanine chain using Coot 0.9 (Emsley et al., 2010). The resulting structure model was submitted to the DALI server (Holm, 2022) to query the *E. coli* species subset of the AlphaFold Database (Hierarchical AF-DB Search). The search showed strong structural homology with the uncharacterised lipoprotein YbjP (AF-P75818-F1). The AlphaFold model was fitted into the experimental map and manually adjusted in Coot 0.9, allowing the identification of the density as YbjP. Following minor structure adjustments, the model was refined using ISOLDE 1.6 plugin (Croll, 2018) for ChimeraX-1.6.1 (Pettersen et al., 2021). Molecular refinement was completed in Phenix-1.20 (Liebschner et al., 2019). Refinement statistics are listed in Supplementary Table S1.

Plasmid construction and purification of soluble YbjP

Soluble YbjP bearing a thrombin-cleavable N-terminal His-tag or a TEV-cleavable C-terminal His-tag were constructed by amplifying the protein globular domain (residues 28-171) from *E. coli* MG1655 genomic DNA. The PCR product was digested NdeI-BamHI or NdeI-XhoI prior insertion into pET28a or pET24a respectively, digested with the same enzymes. *E. coli* BL21 (DE3) cells were then transformed with pET28-YbjP₂₈₋₁₇₁ or pET24-YbjP₂₈₋₁₇₁ plasmid encoding YbjP_s. Bacteria were grown at 37°C in 2YT media supplemented with kanamycin and protein expression induced with addition at mid- to late-log phase of 0.1 mM IPTG. After 16 h at 18°C, cells were harvested by centrifugation (4,000 *g*, 10 min), and pellets resuspended in 50 mM Tris pH 7.5, 200 mM NaCl and 10% glycerol supplemented with lysozyme and DNase I. Cells were then lysed by passage through a cell disruptor (Constant Systems) at 30,200 psi. Unbroken cells and debris were removed by ultracentrifugation (107,000 *g*, 1 h, 4°C). The supernatant was supplemented with 20 mM imidazole and loaded onto a 5 mL HisTrap FF column (GE Healthcare), which was washed with 15 column volumes of the same buffer before elution with 250 mM imidazole. Protein fractions were pooled, dialysed twice against buffer containing 50 mM Tris pH 7.5 and 200 mM NaCl and concentrated using a Vivaspin filter with a 10 kDa MWCO before flash freezing and storage at -80°C.

Preparation of AcrABZ-TolC in peptidisc reconstituted with YbjP

Co-expression of the recombinant AcrABZ-TolC-3xFLAG and cellular membrane preparation was as previously described with some modifications (Du et al., 2014). The AcrABZ-TolC-3xFLAG components were expressed into *E. coli* strain C43 (Δ acrAB and Δ tolC). Cell pellets were resuspended in lysis buffer (150 mM NaCl and 50 mM Tris pH 7.5) and membranes pelleted as previously described (Du et al., 2014) and solubilized with a buffer containing 400 mM NaCl, 50 mM Tris pH 7.5 and 1.5% (wt/v) β -DDM by stirring overnight at 4°C. The soluble extract was incubated with Ni-NTA beads (Qiagen) for nickel affinity purification (2 mL for 4.5 g membrane extract) of the His-tagged AcrABZ_{6xHis}-TolC_{3xFLAG}. The beads were loaded onto a gravity flow-column and the column washed with 4 column volumes (CVs) of a buffer composed of 200 mM NaCl, 50 mM Tris pH 7.5 and 0.02% (wt/v) β -DDM. The AcrABZ_{6xHis}-TolC_{3xFLAG} protein complex was eluted with the same buffer containing 0.5 M imidazole. Eluted fractions were analysed on a 4-12% SDS-PAGE gel and fractions enriched for the pump were then pooled for a second purification step using anti-FLAG M2 affinity beads (Sigma). The fractions were incubated with 0.5 mL anti-FLAG resin for 3 h at 4°C with gentle stirring and the beads were then washed with 2 CVs of buffer (200 mM NaCl, 50 mM Tris pH 7.5 and 0.02% β -DDM) and 3 CVs of buffer 200 mM NaCl, 50 mM Tris pH 7.5 to gradually decrease the detergent of the solubilized pump to below the critical micelle concentration. For reconstitution of the AcrABZ-TolC sample, the NSPr peptidisc was used based on the protocol by Carlson et al. (2018) with some modifications. Next, 2 mg mL⁻¹ of peptidisc (Peptidisc Biotech) in 20 mM Tris-HCl pH 8.0 peptide solution was loaded to the column and incubated for 20 min at room temperature with gentle shaking. The flow-through was collected by gravity and 2 mg mL⁻¹ peptide solution containing 40 μ L of *E. coli* lipids (6 mg mL⁻¹) was loaded to the column and incubated with the beads. Finally, 0.5 mg mL⁻¹ of 3xFLAG peptide in buffer 50 mM Tris pH 7.5, 200 mM NaCl was added to the column and incubated with the beads for 15 min. The fractions were analysed on a 4-12% gradient SDS-PAGE gel and concentrated to 5-6 mg mL⁻¹ using a Vivaspin filter with a 100 kDa MWCO. Prior grid freezing, samples of AcrABZ-TolC in peptidisc were mixed with 10-fold excess of C-terminally His-tagged soluble YbjP.

AcrABZ-TolC-YbjP cryo-EM sample preparation, data collection and processing

Sample of AcrABZ-TolC in peptidisc and YbjP was applied on a Cu 300-mesh EM grid layered with R1.2/1.3 holey carbon support film (Quantifoil), which had been glow-discharged immediately prior to use. The sample deposited on the EM grid was blotted of excess solution and vitrified in liquid ethane using Vitrobot Mark IV (Thermo Fisher Scientific, Inc.) robotic cryo-EM plunger. Images were collected on a Titan Krios microscope operating at 300 kV. Electron fluency was 56 e⁻/Å² and defocus values between -2.5 and -1.0 μ m. Data were processed with cryoSPARC as detailed

in [Supplementary Figure S3](#). After patch motion correction and CTF correction, particles were selected using Blob picker and 8,603 particles were extracted prior 2D classification and best classes were used for template picking. Particles were then subjected to two rounds of 2D classification. The selected 257,681 particles were used to build three *ab initio* models. The tripartite pump was further refined by heterogenous refinement and non-uniform refinement in C_3 symmetry, yielding a final map at 3.17 Å resolution.

AcrABZ-TolC-YbjP preparation with mature peptidoglycan and cryo-EM processing

The AcrABZ-TolC-YbjP prepared in peptidisc were mixed with roughly 1 mg mL⁻¹ of peptidoglycan slurry. Samples of mature and early forms of peptidoglycan (PG) were kindly provided by Professor Waldemar Vollmer (University of Newcastle and now at University of Queensland, Brisbane, Australia), prepared from strains BW25113 ΔGLDT and C5703-1. The mixture was incubated on ice for 1 hour and then used to prepare grids on glow-discharged Ultrafoil R 1.2/1.3 using a FEI Vitrobot (IV) at settings of 95% humidity, 4°C, 3 seconds blot time, -5 force. Cryo-EM data processing and 3D-reconstructions were carried out using cryoSPARC ([Punjani et al., 2017](#)). Multiple rounds of 3D refinement were conducted including heterogeneous refinement to remove non-pump particles, along with homogeneous refinement and non-uniform refinement to improve the resolution yielding a final map of AcrABZ-TolC-PG-YbjP at 3.97 Å. The data collection and processing parameters are summarised in [Supplementary Table S2](#).

Isothermal titration calorimetry (ITC)

Experiments were performed with a ITC200 calorimeter (MicroCal, Malvern Panalytical) at 25°C. After overnight dialysis in 20 mM HEPES pH 7.5, 200 mM NaCl, 0.03% (wt/v) DDM, N-terminally His-tagged soluble YbjP at 230 μM was injected into 40 μM TolC in the same buffer. The titration program consisted of 25 injections of 1.5 μL with an initial injection of 0.4 μL every 300 s, with cell stirring rate of 750 rpm. For each condition, a control titration was performed by injecting YbjP into a cell containing buffer only. The control signal was then subtracted from the experimental titration, and the binding affinity, stoichiometry and thermodynamic parameters were obtained by nonlinear least-squares fitting of the background corrected experimental data using a single-site binding model from the MicroCal PEAQ-ITC analysis software (Malvern). ITC parameters are listed in [Supplementary Table S3](#).

IMAC (immobilised metal affinity chromatography) binding assay of TolC and YbjP

Detergent-purified FLAG-tagged TolC was mixed to N-terminally His-tagged YbjP_s for 5 min in 50 mM Tris pH 7.5, 200 mM NaCl and 0.03% (wt/v) β-DDM. Each protein was at 15 μM final concentration in a final volume of 125 μL. A control was performed with only the TolC protein, or only the YbjP, or by replacing YbjP with another periplasmic protein, soluble LolB, produced as described in [Kaplan et al. \(2018\)](#). Proteins were added to 100 μL of Ni-resin (50% slurry, profinity Biorad) in a microbatch spin column and after 5 minutes the flow-through was recovered. The resin was washed four times with 250 μL of buffer and bound proteins were eluted with the same volume of buffer containing 250 mM imidazole. Elution fractions were analysed on a gradient (4-12%) SDS-PAGE gel. As a reference the TolC flow-through fraction was also loaded.

In vivo photo-crosslinking

Plasmid constructions

Full-length YbjP was amplified from *E. coli* MG1655 genomic DNA. A Ser-Gly linker followed by a 6xHis-tag was added on the reverse primer. The PCR product was digested by NdeI-Xho and inserted into pET24a digested by the same enzymes. AcrA was cloned as described for YbjP. Amber codons were introduced at desired positions by Quikchange site-directed mutagenesis.

Culture and photo-crosslinking

E. coli C43 cells were co-transformed with pEVOL encoding an amber suppressor tyrosyl tRNA and the engineered tyrosyl tRNA synthetase (Chatterjee et al., 2014) and with YbjP_s or AcrA amber mutant for incorporation of pBPA into the TAG amber codon at desired positions. Cultures of 20 mL LB media, supplemented with 20 and 50 µg mL⁻¹ of chloramphenicol and kanamycin respectively, were inoculated with 1/100 of overnight cultures and grown at 30°C. When an optical density of 0.3 was reached, 1 mM of freshly prepared pBPA (Bachem) in NaOH was added in each culture with additional HCl to balance pH. When indicated, a control without addition of pBPA was carried out. After 30 min at 30°C, cells were induced with 0.02% (wt/v) L-arabinose and 0.5 mM IPTG and grown for 2 h at 30°C. Cells corresponding to 4 mL of culture were collected by centrifugation (4,000 g, 4°C, 4 minutes) washed with 1 mL of PBS and resuspended in 700 µL PBS. Aliquots of 150 µL of each culture were transferred in a 96-well plate and exposed under UV (365 nm) for 15 min at room temperature using a UVGL-58 handheld UV lamp.

His-tag enrichment

The next day, samples were lysed by 3 cycles of freezing in liquid nitrogen and thawing in 37°C water bath. Lysozyme at 0.5 mg mL⁻¹, DNase at 50 µg mL⁻¹ and 1% (wt/v) DDM were added. After 30 min at room temperature, samples (300 µL) were transferred into spin columns containing 100 µL of Ni-NTA resin (50% suspension, profinity Biorad) previously equilibrated with PBS (phosphate buffered saline). Proteins were allowed to bind for 5 min at room temperature. The resin was then washed 3 times with 400 µL of PBS supplemented with 0.03% DDM and bound proteins eluted with 150 µL of PBS containing 0.03% DDM and 250 µM imidazole.

Western-blot

Samples (20 µL) were analysed by SDS-PAGE and immunoblotting with mouse anti-His primary antibodies (Penta-His, Qiagen) and goat anti-mouse secondary antibodies (IRDye 800 CW, Licor) prior to imaging with an Odyssey Licor system. For the YbjP and TolC double detection (Supplementary Figure S2), proteins were simultaneously detected with mouse anti-His and rabbit anti-TolC primary antibodies and with goat anti-mouse secondary antibodies (IRDye 800 CW, Licor) and donkey anti-rabbit secondary antibodies (IRDye 680 RD, Licor).

Strains and culture conditions

E. coli BW25113 wild-type and Δ tolC were obtained from the Keio collection (Baba et al., 2006). BW25113 Δ tolC Δ ybjP was generated by λ -Red recombination from the Δ tolC strain. Due to consistent deletion of the *yecT*-*flhD* region of the Keio collection BW25113 Δ ybjP strain following FLP expression, Δ ybjP was generated by λ -Red recombination from the wild-type strain. The kanamycin-resistance cassettes were excised from these strains using pCP20 and the genomes were sequenced for confirmation.

Cells were cultured routinely in Mueller Hinton (MH) broth at 37 °C, with shaking at 200 rpm. Where specified, cells were cultured in Luria-Bertani (LB) broth (0.5 g L⁻¹ NaCl) or M9 minimal media supplemented with 0.4% glycerol. Cells containing a gentamicin-resistance version of the pKD46 plasmid were cultured in LB broth containing 10 µg mL⁻¹ gentamicin, except Δ tolC, which was cultured with 5 µg mL⁻¹ gentamicin. LB agar plates containing 50 µg mL⁻¹ kanamycin or 50 µg mL⁻¹ carbenicillin were used for routine transformant selection. Genome-integrated kanamycin-resistance cassettes were selected for on LB agar containing 30 µg mL⁻¹ kanamycin, except for Δ ybjP and Δ tolC Δ ybjP, which were plated on LB agar containing 20 µg mL⁻¹ and 10 µg mL⁻¹ kanamycin, respectively.

λ -Red recombination

λ -Red recombination (Datsenko and Wanner, 2000) was carried out using a modified version of the protocol described in Peng et al. (2022). Briefly, cultures of cells containing a gentamicin-resistance version of the pKD46 plasmid were prepared in LB broth containing gentamicin and incubated overnight at 30°C. The overnight cultures were diluted 1:200 into fresh LB broth

containing gentamicin and incubated at 30°C until the OD_{600nm} reached 0.4, at which point L-arabinose was added to a final concentration of 0.3% (w/v) and the cultures were incubated at 37°C for 1 h. The cultures were transferred to ice, then washed by four cycles of centrifugation and resuspension in ice-cold distilled water. After the final centrifugation, the distilled water was discarded, and the cells were resuspended in the residual liquid volume. The OD_{600nm} of the resuspended cells were measured and appropriate volumes of ice-cold distilled water were added to bring the cells to a final density of OD_{600nm} 50.

Amplicons of kanamycin-resistance cassettes with 50 bp homology to the gene of interest were designed as described in [Baba et al. \(2006\)](#), amplified using Q5 polymerase (NEB), and gel purified (NEB). 100 ng of each purified amplicon was added to 70 µL of the pKD46 induced electrocompetent cells and transformed by electroporation. Cells were recovered in SOC media (NEB) at 37°C for 1 h, then plated on selective agar plates. The kanamycin-resistance cassettes were excised from the strains using the plasmid pCP20, as described in [Baba et al. \(2006\)](#).

Antimicrobial susceptibility testing

Minimum inhibitory concentrations (MICs) were determined by broth microdilution. Briefly, overnight cultures in MH broth were diluted 1:1000 in fresh MH broth and incubated at 37°C until reaching an OD_{600nm} of 0.2-0.5. The cultures were then diluted to an OD_{600nm} of 0.001 in fresh MH broth and 50 µL were added to wells of flat-bottom 96-well microtiter plates already containing 50 µL of 2-fold serially diluted compounds in MH broth, to a final volume of 100 µL. The plates were sealed with a gas-permeable seal and incubated at 37°C for 18 h. Growth was determined by visual inspection and OD_{600nm} measurement using a SpectraMax iD3 microplate reader.

For antimicrobial susceptibility testing under microaerobic conditions, prior to incubation, the 96-well microtiter plates were placed in a 2.5 L anaerobic jar and Millipore Anaerocult™ C was used following the manufacturer's instructions to generate an environment containing 5-6% O₂ and 8-10% CO₂.

Bacterial growth curves

Bacterial growth curves were carried out by diluting overnight cultures to an OD_{600nm} of 0.02 in fresh media containing the relevant concentrations of supplemented solutions. 200 µL of the diluted cultures were added to wells of a flat-bottom 96-well microtiter plate and the plate was sealed with a gas-permeable seal. OD_{600nm} was read every 10 min for 24 h at 37°C with shaking using a SpectraMax iD3 microplate reader.

Swimming motility assay

LB motility agar plates were prepared by adding 25 mL of LB broth containing 0.3% agar in 94 mm diameter sterile petri dishes. Overnight cultures of the relevant strains were prepared in LB broth. A volume of 2 µL of overnight culture was injected into the centre of each motility agar plate, followed by incubation at 37°C for 24 h. The diameters of the colonies at the widest point were measured using a ruler.

Proteomics sample preparation and analysis

Cultures were prepared in triplicate under aerobic conditions in MH media for the four strains *ΔybjP*, *ΔtolC*, *ΔtolC ΔybjP* and the BW25113 parent, all prepared by the recombineering approach described above. Cultures were grown to exponential phase (0.6-0.8 OD_{600 nm}, 15 mL) and to stationary phase (4.8-5 OD_{600 nm}, 5 mL) and pelleted by centrifugation at 4,000 g, 5 min and frozen. The pellets were resuspended in 1 mL lysis buffer for set 1 (50 mM Tris-Cl pH 7.5, 100 mM NaCl, 1 mM MgCl₂, 1.4 mg mL⁻¹ hen egg white lysozyme, 1% (v/v) Triton X-100). The suspension was sonicated with a microprobe tip (Sonics vibracell) on ice for 60 sec, with 4 sec pulses, 6 sec wait time, 50 % amplitude. To the sonicate lysate, 2 units of DNase I were added to digest DNA, followed by incubation on ice for 10 min. Trichloroacetic acid (Sigma) was then added to 5 % (wt/v), then incubated on ice and collected by centrifugation in an Eppendorf centrifuge at (13,000 rpm, 5°C, 10 min). The pellet was washed with 5% TCA in water, then spun (set 1) or with 50% ethanol in 20

mM Tris-Cl pH 7.5 (set 2), and the collected pellet provided to the University of Cambridge Centre for Proteomics. Samples were digested with trypsin and analysed by quantitative label free LC-MS/MS. Signals from total unique spectral counts were normalised for each replicate and averaged.

Raw mass spectrometry data was processed with DIA-NN v2.1.1 (Demichev et al., 2020) using the reference *E. coli* proteome from UniProt (acc. no. UP000000625) along with contaminant protein sequences provided by the Cambridge Centre for Proteomics (Dawson & Smith, 2024). Precursor and fragment mass tolerances were automatically optimized within the DIA-NN search. Trypsin was set as the enzyme of choice and a maximum of one missed cleavage was allowed. Carbamidomethylation of cysteine was set as a fixed modification, one variable modification was allowed from either N-terminus acetylation or oxidation of methionine. An FDR cutoff of 1% was used at the precursor level.

Quantitative analysis was performed from the abundances at the peptide level using qfeatures and limma (Gatto and Vanderaa, 2025), adapting the code from (Hutchings et al., 2025). Briefly: peptides matching contaminant proteins, peptides with q-values higher than 0.01, ambiguously mapping peptides and peptides with a proportion of missing values greater than 0.5 across all samples were removed from the analysis. Missing values were imputed using minDet imputation for missing values not at random, and knn for missing values at random. Peptide quantitation information was aggregated at the protein level using qfeatures with the function colMedians, quantitation information was log₂ transformed and normalised. Protein-wise linear models were built with limma which were subsequently adjusted with the eBayes function. Differentially expressed proteins were considered if they had log₂ fold changes (LFC) greater than 1 or smaller than -1, statistically significant proteins were considered if their Bonferroni-Hochberg adjusted p-value was smaller than 0.05.

Phylogenetic analysis of DUF3828 domain


3327 proteins containing DUF3828 domain (PF12883) were acquired from InterPro (v103), and the domain architectures represented by *E. coli* YbjP and YqhG were selected. Taxonomy information was retrieved from UniProt on Dec 22, 2024, and the signal peptide was predicted by SignalP 6.0 (Teufel et al., 2022). After removing proteins with unidentified taxonomy, representative sequences were selected by manual inspection of *E. coli* and *Salmonella* and CD-HIT with a 95% identity threshold (Fu et al., 2012), yielding 356 sequences. The multiple sequence alignment was generated by hmmsalign using the PF12883 HMM profile and processed with TrimAl on automated mode (Capella-Gutiérrez et al., 2009). A phylogenetic tree was constructed from the resulting multiple sequence alignment using IQ-Tree2 on a default parameter (Nguyen et al., 2015). The tree was rooted using the minimal ancestor deviation method (Tria et al., 2017), yielding the same result. Annotation was performed with TreeViewer (Bianchini and Sánchez-Baracaldo, 2024) and webFlaGs (Saha et al., 2021) was used to analyze the gene synteny of DUF3828-containing proteins.

Bioinformatic analysis of outer membrane efflux protein


Outer membrane efflux proteins (IPR003423) and TolC proteins (IPR010130) were acquired from InterPro (v103) and the taxonomic information was retrieved from UniProt on Dec 22 2024. The signal peptide was predicted by SignalP6.0. Outer membrane efflux proteins of *Pseudomonadota* were selected and clustered with CLANS (Frickey and Lupas, 2004).

For the phylogenetic analysis of TolC, the method described for DUF3828 analysis was used with some modifications. Sequences of *Pseudomonadota* were collected, and the sequence showing the highest sequence similarity to *E. coli* TolC in each organism was selected. Redundant sequences were removed by CD-HIT with a 90% identity threshold. In addition, a phylogenetic tree was constructed using FastTree on -mlacc 2 -slow parameter (Price et al., 2010).

Supplementary Information




Supplementary Figure S1. Cryo-EM processing workflow for MacAB-TolC-YbjP complex. (A) Cryo-EM processing workflow. (B) Gold-standard Fourier Shell Correlation (GS-FSC) curves for final reconstruction. (C) Local resolution distribution in final map. (D) Angular distribution plot for final map, left, and Fourier Shell Correlation between half maps of MacAB-TolC-YbjP, right. These data suggest that there is not a large orientational bias in the final map.




Supplementary Figure S2. Double detection of YbjP and TolC proteins for the *in vivo* photo crosslinking.

Left, immunoblot showing purified N-terminally His-tagged soluble YbjP (YbjP_s) and His-tagged TolC detected with anti-His (green) and anti-TolC (red) antibodies. Right, immunoblot of *E. coli* C43 cells expressing His-tagged full-length YbjP (YbjP_{FL}) carrying a photoactivatable pBPA group at position N113 and endogenous, tag-free, TolC detected with anti-His (green) and anti-TolC (red) antibodies, after UV irradiation. An overlay of the two images is shown on the right. The double detection of YbjP and TolC proteins in the crosslinked sample is indicated by an arrow.




Supplementary Figure S3. Cryo-EM workflow for AcrABZ-TolC-YbjP complex.

(A) Cryo-EM processing workflow. (B) Gold-standard Fourier Shell Correlation (GS-FSC) curves for final reconstruction. (C) Local resolution distribution in the final map. (D) Angular distribution plot of the final particles.




Supplementary Figure S4. YbjP binds at the same position in the AcrABZ-TolC and MacAB-TolC tripartite pumps and may not interact with peptidoglycan.

(A) Cryo-EM map of AcrABZ-TolC-YbjP reconstituted in peptidoglycan. (B) Fourier Shell Correlation between half-maps (top) and angular distribution plot of the particles (bottom). Collection statistics for the AcrABZ-TolC-YbjP complex reconstituted in peptidoglycan are listed in [Supplementary Table S2](#). (C) Peptidoglycan location based on the cryo-ET map described by [Shi et al., 2019](#).




Supplementary Figure S5. The cladogram of TolC (IPR010130) in *Pseudomonadota*.

The sequence that showed the highest pairwise similarity to *E. coli* TolC in each organism is selected.



Supplementary Figure S6. Impact of tryptophan on growth in minimal media.

The growth curves of WT (blue), $\Delta ybjP$ (orange), $\Delta tolC$ (green), $\Delta ybjP \Delta tolC$ (red) *E. coli* were recorded in M9 media supplemented with 0.4% glycerol and indicated L-tryptophan (Trp) concentrations. Results are shown as mean $\pm$ standard deviation from two biological replicates, each with three technical replicates.



Supplementary Figure S7. Structural comparison of TolC₃-YbjP₃ and TolC₃-SlyB₁₁.

(A) AlphaFold3 structure of TolC-SlyB₁₁ with 30 palmitic acid and 30 oleic acid molecules. The model is coloured by pLDDT score. (B) PAE plot of TolC₃-SlyB₁₁. (C) Overlay of TolC₃-YbjP₃ structure and TolC₃-SlyB₁₁ AlphaFold3 structure. The lipidation in YbjP is displayed as green spheres.

	MacAB-TolC-YbjP in peptidisc	AcrABZ-TolC-YbjP in peptidisc
Organism	<i>E. coli</i>	<i>E. coli</i>
PDB ID	9QGY	9TG4
EMDB ID	EMD-53150	EMD-55890
Data collection		
Microscope	FEI Titan Krios	FEI Titan Krios
Voltage (kV)	300	300
Detector	K3	K3
Mode	counting, super-resolution	counting, super-resolution
Nominal magnification	105,000 ×	100,000 ×
Pixel size (Å/px)	0.830	1.168
Electron fluency, total (e ⁻ /Å ²)	49.75	56
Defocus range (μm)	-1.0 to -2.4	-1.0 to -2.5
Exposure (s)	1	1
Frames	40	40
Number of micrographs	5,399	13,456
Reconstruction		
Software	cryoSPARC-4.5.3	cryoSPARC-4.5.3
Final number of particle images	233,775	97,441
Point group	C ₃	C ₃
Map resolution, FSC _{0.143} (Å)	2.48	3.17
Map-sharpening B factor (Å ²)	76.6	86.3
Model composition		
Non-hydrogen atoms	64,989	107,675
Protein residues	4,198	7,048
Other	3	3
Refinement		
Software	Phenix-1.20	Phenix-1.20
Correlation coefficient, masked	0.81	0.82
Correlation coefficient, box	0.83	0.57
Model resolution, FSC _{0.5} (Å)	2.7	3.3
Validation (proteins)		
MolProbity score	1.48	1.46
Clash score	6.73	6.41
Ramachandran plot statistics		
Favoured (%)	97.46	97.45
Allowed (%)	2.54	2.54
Outliers (%)	0.00	0.01

Table S1. Cryo-EM data and refinement statistics for peptidisc-reconstituted MacAB-TolC-YbjP and AcrABZ-TolC-YbjP models.

Table S2. Cryo-EM data collection statistics for AcrABZ-TolC-YbjP in peptidoglycan.

	AcrABZ-TolC-YbjP peptidoglycan
Data collection and processing	
Magnification	100,000 ×
Voltage (kV)	300
Electron exposure (e ⁻ /Å ²)	56
Defocus range (μm)	-1 to -2.5
Pixel size (Å/px)	1.168
Number of micrographs	13,456
Symmetry	C ₁
Initial particle images (no.)	271,642
Final particle images (no.)	62,804
Map resolution, FSC _{0.143} (Å)	3.97

Table S3. Thermodynamic parameters for the TolC-YbjP_s interaction.

Results from isothermal titration calorimetry (ITC) were obtained with DDM-purified TolC-FLAG and N-terminally His-tagged YbjP_s. The mean ± spread of two independent measurements is shown in bold. Experiments were carried out at 25°C. Values of ΔG, ΔH and TΔS are in kcal mol⁻¹. An example of one of the thermograms and fit is shown in Figure 3B. N corresponds to the stoichiometry of the interaction.

K _D (μM)	N	ΔG	ΔH	-TΔS
4.1, 5.9	0.64, 0.42	-7.4, -7.1	-37.8, -14.3	30.5, 7.15
5.0 ± 1.3	0.53 ± 0.16	-7.3 ± 0.2	-26.1 ± 16.6	18.9 ± 16.5

Table S4. Criteria for classification of DUF3828-containing proteins

Family	Tai3	YbjP	Alphaproteobacteria
Taxonomy	Betaproteobacteria Gammaproteobacteria Bacteroidota/Chlorobiota group	Enterobacterales under Gammaproteobacteria	Alphaproteobacteria
Gene Synteny	Tae3	Arginine transporter Amidase	Unclear
Signal peptide	Lipoprotein	Either lipoprotein or periplasmic	Periplasmic
QDX motif	Present	Absent	Present

Table S5. Minimum inhibitory concentration (MIC) values for wild-type and indicated knockout strains of *E. coli* BW25113

. Raw values are in bold and fold-changes values for indicated knockout strains compared to wild-type *E. coli* BW25113 are in brackets, under aerobic or microaerobic (5-6% O₂, 8-10% CO₂) conditions.

Compound	Aerobic				Microaerobic			
	WT	<i>ΔybjP</i>	<i>ΔtolC</i>	<i>ΔtolC ΔybjP</i>	WT	<i>ΔybjP</i>	<i>ΔtolC</i>	<i>ΔtolC ΔybjP</i>
Oxacillin	256	256 (1)	≤ 0.5 (≥ 512)	≤ 0.5 (≥ 512)	512	512 (1)	1 (512)	2 (256)
Novobiocin	64	64 (1)	0.5 (128)	0.5 (128)	64	64 (1)	0.5 (128)	0.5 (128)
CCCP	16	16 (1)	0.25 (64)	0.25 (64)	16	16 (1)	0.25 (64)	0.25 (64)
Erythromycin	32	32 (1)	1 (32)	1 (32)	32	32 (1)	1 (32)	1 (32)
Benzalkonium chloride	8	8 (1)	0.5 (16)	0.5 (16)	32	32 (1)	8 (4)	8 (4)
Nalidixic acid	4	4 (1)	0.25 (16)	0.25 (16)	/	/	/	/
Ampicillin	16	16 (1)	16 (1)	16 (1)	/	/	/	/
Kanamycin	0.5	0.5 (1)	0.5 (1)	0.5 (1)	/	/	/	/
Colistin	0.0625	0.0625 (1)	0.0625 (1)	0.0625 (1)	/	/	/	/
Vanillin	2048	2048 (1)	2048 (1)	2048 (1)	/	/	/	/
EDTA	1024	1024 (1)	2048 (0.5)	2048 (0.5)	/	/	/	/
Microcin J25*	> 256	>256 (1)	0.03125 (>8192)	0.0625 (>4096)	/	/	/	/

*The activity of the used microcin J25 stock was significantly lower than expected, hence the stated MIC values are likely much higher than expected for a fresh active stock

Table S6. Swimming motility assay

. Colony diameters of wild-type and knockout strains of *E. coli* BW25113. Values are presented as mean ± standard deviation from at least 5 replicates.

Strain	Ring diameter (mm)	Phenotype
WT	57.2 ± 2.3	Circular spread outwards from inoculation point
<i>ΔybjP</i>	55.4 ± 3.2	Circular spread outwards from inoculation point
<i>ΔtolC</i>	29.5 ± 8.8	Uneven spread, biased in one direction
<i>ΔybjP ΔtolC</i>	29.3 ± 4.2	Uneven spread, biased in one direction

Data availability

The MacAB-TolC-YbjP and AcrABZ-TolC-YbjP atomic models and electron microscopy maps have been deposited in the Protein Data Bank (PDB) and Electron Microscopy Data Bank (EMDB) under the accession codes 9QGY, EMDB-53150, and 9TG4, EMD-55890 respectively. Proteomics data have been deposited with the PRIDE database with identification code 1-20251119-140222-7962809

Acknowledgements

JG was supported by an Amgen scholarship. Grids were prepared and cryo-EM data collected at the BIOCEM facility, Department of Biochemistry, University of Cambridge. We thank Dimitri Y. Chirgadze, Steven Hardwick, and Lee Cooper for assistance with data collection and processing at the Cryo-EM Facility. We thank Yulia Yuzenkova for providing samples of microcin J25. We thank Somenath Bakshi, Liz Sockett and Waldemar Voller for helpful discussions, and Waldemar for providing mature and early forms of peptidoglycan.

Additional information

Funding

This work is supported by an ERC Advanced Award (742210) and Wellcome Trust Investigator Awards (222451/Z/21/Z and 200873/Z/16/Z). AZ is a recipient of a Transition To Independence (TTI) fellowship from the School of Biological Sciences at the University of Cambridge, supported by funding from the Rosetrees Trust (JS16/TTI2021\1) and the Isaac Newton Trust (21.22(a)iii) and the School of Biological Sciences at the University of Cambridge.

Funding

Funder	Grant reference number	Author
Wellcome Trust (WT)	https://doi.org/10.35802/222451	Ben F Luisi
		Yvette Ntsogo
		Andrzej Harris
		Elise Kaplan
		Ben Jin
		Emmanouela
		Petsolari
		Jan Gradon
EC European Research Council (ERC)	https://doi.org/10.3030/742210	Dingquan Yu
		Yvette Ntsogo
		Ben F Luisi
		Jim Horne
cambridgeuniversity School of the Biological Sciences, University of Cambridge (School of the Biological Sciences)	Transition to Independence	Elise Kaplan
		Ashraf Zarkan
Rosetrees Trust (Rosetrees)	JS16/TTI2021\1	Ashraf Zarkan
Isaac Newton Trust	21.22(a)iii	Ashraf Zarkan

Author ORCID iDs

Jan M Gradon: <https://orcid.org/0009-0006-5043-1986>

Andrzej Harris: <https://orcid.org/0000-0003-4885-7401>

Ben F Luisi: <https://orcid.org/0000-0003-1144-9877>

References

- Abramson J**, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, *et al.* (2024) Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**:493-500 <https://doi.org/10.1038/s41586-024-07487-w> | PubMed
- Aicart-Ramos C**, Valero RA, Rodriguez-Crespo I (2011) Protein palmitoylation and subcellular trafficking. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1808**:2981-2994 <https://doi.org/10.1016/j.bbamem.2011.07.009> | PubMed
- Akama H**, Kanemaki M, Yoshimura M, Tsukihara T, Kashiwagi T, Yoneyama H, Narita S, Nakagawa A, Nakae T (2004) Crystal Structure of the Drug Discharge Outer Membrane Protein, OprM, of *Pseudomonas aeruginosa*. *Journal of Biological Chemistry* **279**:52816-52819 <https://doi.org/10.1074/jbc.C400445200> | PubMed
- Baba T**, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Molecular Systems Biology* **2**:2006-0008 <https://doi.org/10.1038/msb4100050> | PubMed
- Babu M**, Bundalovic-Torma C, Calmettes C, Phanse S, Zhang Q, Jiang Y, Minic Z, Kim S, Mehla J, Gagarinova A, *et al.* (2018) Global landscape of cell envelope protein complexes in *Escherichia coli*. *Nature biotechnology* **36**:103-112 <https://doi.org/10.1038/nbt.4024> | PubMed
- Baquero F**, Beis K, Craik DJ, Li Y, Link AJ, Rebuffat S, Salomón R, Severinov K, Zirah S, Hegemann JD (2024) The pearl jubilee of microcin J25: thirty years of research on an exceptional lasso peptide. *Natural Product Reports* **41**:469-511 <https://doi.org/10.1039/D3NP00046J> | PubMed
- Beketskaia MS**, Bay DC, Turner RJ (2014) Outer Membrane Protein OmpW Participates with Small Multidrug Resistance Protein Member EmrE in Quaternary Cationic Compound Efflux. *Journal of Bacteriology* **196**:1908-1914 <https://doi.org/10.1128/JB.01483-14> | PubMed
- Bessaiah H**, Pokharel P, Habouria H, Houle S, Dozois CM (2019) yqhG Contributes to Oxidative Stress Resistance and Virulence of Uropathogenic *Escherichia coli* and Identification of Other Genes Altering Expression of Type 1 Fimbriae. *Frontiers in Cellular and Infection Microbiology* **9**:312 <https://doi.org/10.3389/fcimb.2019.00312> | PubMed
- Bianchini G**, Sánchez-Baracaldo P (2024) TreeViewer : Flexible, modular software to visualise and manipulate phylogenetic trees. *Ecology and Evolution* **14**:e10873 <https://doi.org/10.1002/ece3.10873> | PubMed
- Calladine CR**, Sharff A, Luisi B (2001) How to untwist an α -helix: structural principles of an α -helical barrel. Edited by J. Thornton. *Journal of Molecular Biology* **305**:603-618 <https://doi.org/10.1006/jmbi.2000.4320> | PubMed
- Capella-Gutiérrez S**, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**:1972-1973 <https://doi.org/10.1093/bioinformatics/btp348> | PubMed
- Carlson ML**, Young JW, Zhao Z, Fabre L, Jun D, Jianing Li, Jun Li, Dhupar HS, Wason I, Mills AT, *et al.* (2018) The Peptidisc, a simple method for stabilizing membrane proteins in detergent-free solution. *eLife* **7**:e34085 <https://doi.org/10.7554/eLife.34085> | PubMed
- Chatterjee D**, Cooley RB, Boyd CD, Mehl RA, O'Toole GA, Sondermann H (2014) Mechanistic insight into the conserved allosteric regulation of periplasmic proteolysis by the signaling molecule cyclic-di-GMP. *eLife* **3**:e03650 <https://doi.org/10.7554/eLife.03650> | PubMed
- Chin JW**, Martin AB, King DS, Wang L, Schultz PG (2002) Addition of a photocrosslinking amino acid to the genetic code of *Escherichia coli*. *Proceedings of the National Academy of Sciences* **99**:11020-11024 <https://doi.org/10.1073/pnas.172226299> | PubMed
- Croll TI** (2018) ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps. *Acta Crystallographica. Section D. Structural Biology* **74**:519-530 <https://doi.org/10.1107/S2059798318002425> | PubMed

- Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences* **97**:6640-6645 <https://doi.org/10.1073/pnas.120163297> | PubMed
- Delgado MA, Vincent PA, Farías RN, Salomón RA (2005) YojI of *Escherichia coli* Functions as a Microcin J25 Efflux Pump. *Journal of Bacteriology* **187**:3465-3470 <https://doi.org/10.1128/JB.187.10.3465-3470.2005> | PubMed
- Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M (2020) DIA-NN: Neural networks and interference correction enable deep proteome coverage in high throughput. *Nature methods* **17**:41-44 <https://doi.org/10.1038/s41592-019-0638-x> | PubMed
- Dong C, Zhang H, Gao Z-Q, Wang W-J, She Z, Liu G-F, Shen Y-Q, Su X-D, Dong Y-H (2013) Structural insights into the inhibition of type VI effector Tae3 by its immunity protein Tai3. *Biochemical Journal* **454**:59-68 <https://doi.org/10.1042/BJ20130193> | PubMed
- Du D, Wang Z, James NR, Voss JE, Klimont E, Ohene-Agyei T, Venter H, Chiu W, Luisi BF. (2014) Structure of the AcrAB-TolC multidrug efflux pump. *Nature* **509**:512-515 <https://doi.org/10.1038/nature13205> | PubMed
- Du D, Wang-Kan X, Neuberger A, Van Veen HW, Pos KM, Piddock LJV, Luisi BF. (2018) Multidrug efflux pumps: structure, function and regulation. *Nature Reviews Microbiology* **16**:523-539 <https://doi.org/10.1038/s41579-018-0048-6> | PubMed
- Emsley P, Lohkamp B, Scott WG, Cowtan K (2010) Features and development of Coot. *Acta Crystallographica Section D: Biological Crystallography* **66**:486-501 <https://doi.org/10.1107/S0907444910007493> | PubMed
- Fitzpatrick AWP, Llabrés S, Neuberger A, Blaza JN, Bai X-C, Okada U, Murakami S, Van Veen HW, Zachariae U, Scheres SHW, et al. (2017) Structure of the MacAB-TolC ABC-type tripartite multidrug efflux pump. *Nature Microbiology* **2**:17070 <https://doi.org/10.1038/nmicrobiol.2017.70> | PubMed
- Frickey T, Lupas A (2004) CLANS: a Java application for visualizing protein families based on pairwise similarity. *Bioinformatics* **20**:3702-3704 <https://doi.org/10.1093/bioinformatics/bth444> | PubMed
- Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**:3150-3152 <https://doi.org/10.1093/bioinformatics/bts565> | PubMed
- Glavier M, Puvanendran D, Salvador D, Decossas M, Phan G, Garnier C, Frezza E, Cece Q, Schoehn G, Picard M, et al. (2020) Antibiotic export by MexB multidrug efflux transporter is allosterically controlled by a MexA-OprM chaperone-like complex. *Nature Communications* **11**:4948 <https://doi.org/10.1038/s41467-020-18770-5> | PubMed
- Grabowicz M, May KL (2026) Lipid landscaping of the bacterial cell surface. *Proceedings of the National Academy of Sciences* **123**:e2537958123 <https://doi.org/10.1073/pnas.2537958123> | PubMed
- Guest RL, Silhavy TJ (2023) Cracking outer membrane biogenesis. *Biochimica et biophysica acta. Molecular cell research* **1870**:119405 <https://doi.org/10.1016/j.bbamcr.2022.119405> | PubMed
- Gumbart JC, Ferreira JL, Hwang H, Hazel AJ, Cooper CJ, Parks JM, Smith JC, Zgurskaya HI, Beeby M. (2021) Lpp positions peptidoglycan at the AcrA-TolC interface in the AcrAB-TolC multidrug efflux pump. *Biophysical Journal* **120**:3973-3982 <https://doi.org/10.1016/j.bpj.2021.08.016> | PubMed
- Hoang HH, Nickerson NN, Lee VT, Kazimirova A, Chami M, Pugsley AP, Lory S (2011) Outer Membrane Targeting of *Pseudomonas aeruginosa* Proteins Shows Variable Dependence on the Components of Bam and Lol Machineries. *mBio* **2**:e00246-11 <https://doi.org/10.1128/mBio.00246-11> | PubMed
- Holm L (2022) Dali server: structural unification of protein families. *Nucleic Acids Research* **50**:W210-W215 <https://doi.org/10.1093/nar/gkac387> | PubMed
- Honeycutt JD, Wenner N, Li Y, Brewer SM, Massis LM, Brubaker SW, Chairatana P, Owen SV, Canals R, Hinton JCD, et al. (2020) Genetic variation in the MacAB-TolC efflux pump influences pathogenesis of invasive *Salmonella* isolates from Africa. *PLOS Pathogens* **16**:e1008763 <https://doi.org/10.1371/journal.ppat.1008763> | PubMed

- Horne JE, Brockwell DJ, Radford SE (2020) Role of the lipid bilayer in outer membrane protein folding in Gram-negative bacteria. *Journal of Biological Chemistry* **295**:10340-10367 <https://doi.org/10.1074/jbc.REV120.011473> | PubMed
- Huang P-S, Oberdorfer G, Xu C, Pei XY, Nannenga BL, Rogers JM, DiMaio F, Gonen T, Luisi B, Baker D (2014) High thermodynamic stability of parametrically designed helical bundles. *Science* **346**:481-485 <https://doi.org/10.1126/science.1257481> | PubMed
- Hutchings C, Krueger T, Crook OM, Gatto L, Lilley KS, Breckels LM (2025) An updated Bioconductor workflow for correlation profiling subcellular proteomics. *F1000Research* **14**:714 <https://doi.org/10.12688/f1000research.165543.1> | PubMed
- Janssens A, Nguyen VS, Cecil AJ, Van der Verren SE, Timmerman E, Deghelt M, Pak AJ, Collet J-F, Impens F, Remaut H. (2024) SlyB encapsulates outer membrane proteins in stress-induced lipid nanodomains. *Nature* **626**:617-625 <https://doi.org/10.1038/s41586-023-06925-5> | PubMed
- Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, et al. (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* **596**:583-589 <https://doi.org/10.1038/s41586-021-03819-2> | PubMed
- Kaplan E, Greene NP, Crow A, Koronakis V (2018) Insights into bacterial lipoprotein trafficking from a structure of LolA bound to the LolC periplasmic domain. *Proceedings of the National Academy of Sciences* **115** <https://doi.org/10.1073/pnas.1806822115> | PubMed
- Kovacs-Simon A, Titball RW, Michell SL (2011) Lipoproteins of Bacterial Pathogens. *Infection and Immunity* **79**:548-561 <https://doi.org/10.1128/IAI.00682-10> | PubMed
- Lacour S, Landini P (2004) S-Dependent Gene Expression at the Onset of Stationary Phase in *Escherichia coli*: Function of S-Dependent Genes and Identification of Their Promoter Sequences. *Journal of Bacteriology* **186** <https://doi.org/10.1128/jb.186.21.7186-7195.20>
- Lei H-T, Bolla JR, Su C-C, Yu EW (2014) Crystal structures of CusC review conformational changes accompanying folding and transmembrane channel formation. *Journal of molecular biology* **426**:403-411 <https://doi.org/10.1016/j.jmb.2013.09.042> | PubMed
- Liebschner D, Afonine PV, Baker ML, Bunkóczi G, Chen VB, Croll TI, Hintze B, Hung L-W, Jain S, McCoy AJ, et al. (2019) Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in *Phenix*. *Acta Crystallographica Section D Structural Biology* **75**:861-877 <https://doi.org/10.1107/S2059798319011471> | PubMed
- Linder ME, Deschenes RJ (2007) Palmitoylation: policing protein stability and traffic. *Nature Reviews Molecular Cell Biology* **8**:74-84 <https://doi.org/10.1038/nrm2084> | PubMed
- Malinverni JC, Werner J, Kim S, Sklar JG, Kahne D, Misra R, Silhavy TJ (2006) YfiO stabilizes the YaeT complex and is essential for outer membrane protein assembly in *Escherichia coli*. *Molecular Microbiology* **61**:151-164 <https://doi.org/10.1111/j.1365-2958.2006.05211.x> | PubMed
- Molloy MP, Herbert BR, Slade MB, Rabilloud T, Nouwens AS, Williams KL, Gooley AA. (2000) Proteomic analysis of the *Escherichia coli* outer membrane. *European Journal of Biochemistry* **267**:2871-2881 <https://doi.org/10.1046/j.1432-1327.2000.01296.x> | PubMed
- Nakayama H, Kurokawa K, Lee BL (2012) Lipoproteins in bacteria: structures and biosynthetic pathways. *The FEBS Journal* **279**:4247-4268 <https://doi.org/10.1111/febs.12041> | PubMed
- Noland CL, Kattke MD, Diao J, Gloor SL, Pantua H, Reichelt M, Katakam AK, Yan D, Kang J, Zilberleyb I, et al. (2017) Structural insights into lipoprotein N-acylation by *Escherichia coli* apolipoprotein N-acyltransferase. *Proceedings of the National Academy of Sciences* **114** <https://doi.org/10.1073/pnas.1707813114> | PubMed
- Patrick CA, Webb JP, Green J, Chaudhuri RR, Collins MO, Kelly DJ (2019) Proteomic Profiling. *Transcription Factor Modeling, and Genomics of Evolved Tolerant Strains Elucidate Mechanisms of Vanillin Toxicity in Escherichia coli*. *mSystems* **4**:e00163-19 <https://doi.org/10.1128/mSystems.00163-19>

- Peng L, Dumevi RM, Chitto M, Haarmann N, Berger P, Koudelka G, Schmidt H, Mellmann A, Dobrindt U, Berger M (2022) A Robust One-Step Recombineering System for Enterohemorrhagic *Escherichia coli*. *Microorganisms* **10**:1689 <https://doi.org/10.3390/microorganisms10091689> | PubMed
- Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE (2021) UCSF CHIMERA X : Structure visualization for researchers, educators, and developers. *Protein Science* **30**:70-82 <https://doi.org/10.1002/pro.3943> | PubMed
- Price MN, Dehal PS, Arkin AP (2010) FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments. *PLoS ONE* **5**:e9490 <https://doi.org/10.1371/journal.pone.0009490> | PubMed
- Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA (2017) cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nature Methods* **14**:290-296 <https://doi.org/10.1038/nmeth.4169> | PubMed
- Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) . limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research* **43**:e47-e47 <https://doi.org/10.1093/nar/gkv007> | PubMed
- Russell AB, Wexler AG, Harding BN, Whitney JC, Bohn AJ, Goo YA, Tran BQ, Barry NA, Zheng H, Peterson SB, et al. (2014) A Type VI Secretion-Related Pathway in *Bacteroidetes* Mediates Interbacterial Antagonism. *Cell Host & Microbe* **16**:227-236 <https://doi.org/10.1016/j.chom.2014.07.007> | PubMed
- Saha CK, Sanches Pires R, Brolin H, Delannoy M, Atkinson GC (2021) FlaGs and webFlaGs: discovering novel biology through the analysis of gene neighbourhood conservation. *Bioinformatics* **37**:1312-1314 <https://doi.org/10.1093/bioinformatics/btaa788> | PubMed
- Shi X, Chen M, Yu Z, Bell JM, Wang H, Forrester I, Villarreal H, Jakana J, Du D, Luisi BF, et al. (2019) In situ structure and assembly of the multidrug efflux pump AcrAB-TolC. *Nature Communications* **10**:2635 <https://doi.org/10.1038/s41467-019-10512-6> | PubMed
- Stubenrauch CJ, Bamert RS, Wang J, Lithgow T (2022) A noncanonical chaperone interacts with drug efflux pumps during their assembly into bacterial outer membranes. *PLOS Biology* **20**:e3001523 <https://doi.org/10.1371/journal.pbio.3001523> | PubMed
- Tenorio E, Saeki T, Fujita K, Kitakawa M, Baba T, Mori H, Isono K (2003) Systematic characterization of *Escherichia coli* genes/ORFs affecting biofilm formation. *FEMS Microbiology Letters* **225**:107-114 [https://doi.org/10.1016/S0378-1097\(03\)00507-X](https://doi.org/10.1016/S0378-1097(03)00507-X) | PubMed
- Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsigos KD, Winther O, Brunak S, Von Heijne G, Nielsen H. (2022) SignalP 6.0 predicts all five types of signal peptides using protein language models. *Nature Biotechnology* **40**:1023-1025 <https://doi.org/10.1038/s41587-021-01156-3> | PubMed
- Tria FDK, Landan G, Dagan T (2017) Phylogenetic rooting using minimal ancestor deviation. *Nature Ecology & Evolution* **1**:0193 <https://doi.org/10.1038/s41559-017-0193> | PubMed
- Tsutsumi K, Yonehara R, Ishizaka-Ikeda E, Miyazaki N, Maeda S, Iwasaki K, Nakagawa A, Yamashita E (2019) Structures of the wild-type MexAB-OprM tripartite pump reveal its complex formation and drug efflux mechanism. *Nature Communications* **10**:1520 <https://doi.org/10.1038/s41467-019-09463-9> | PubMed
- Voulhoux R, Bos MP, Geurtsen J, Mols M, Tommassen J (2003) Role of a Highly Conserved Bacterial Protein in Outer Membrane Protein Assembly. *Science* **299**:262-265 <https://doi.org/10.1126/science.1078973> | PubMed
- Wang Z, Fan G, Hryc CF, Blaza JN, Serysheva II, Schmid MF, Chiu W, Luisi BF, Du D. (2017) An allosteric transport mechanism for the AcrAB-TolC multidrug efflux pump. *eLife* **6**:e24905 <https://doi.org/10.7554/eLife.24905> | PubMed
- Werner J, Misra R (2005) YaeT (Omp85) affects the assembly of lipid-dependent and lipid-independent outer membrane proteins of *Escherichia coli*. *Molecular Microbiology* **57**:1450-1459 <https://doi.org/10.1111/j.1365-2958.2005.04775.x> | PubMed

Wu T, Malinverni J, Ruiz N, Kim S, Silhavy TJ, Kahne D (2005) Identification of a Multicomponent Complex Required for Outer Membrane Biogenesis in *Escherichia coli*. *Cell* **121**:235-245
<https://doi.org/10.1016/j.cell.2005.02.015> | PubMed

Young TS, Ahmad I, Yin JA, Schultz PG (2010) An Enhanced System for Unnatural Amino Acid Mutagenesis in *E. coli*. *Journal of Molecular Biology* **395**:361-374
<https://doi.org/10.1016/j.jmb.2009.10.030> | PubMed

Zhang W, Harper CE, Lee J, Fu B, Ramsukh M, Hernandez CJ, Chen P (2025) Transporter excess and clustering facilitate adaptor protein shuttling for bacterial efflux. *Cell Reports Physical Science* **6**:102441 <https://doi.org/10.1016/j.xcrp.2025.102441> | PubMed

Peer reviews

Reviewer #2 (Public review):

This article focuses on the study of two *E. coli* tripartite efflux pumps, both using TolC as a partner in the outer membrane, namely MacAB-TolC and AcrABZ-TolC.

By preparing MacAB-TolC in Peptidiscs rather than in detergent for cryo-EM structure determination, they visualized an extra protein localized around TolC. The resolution was sufficient to build part of the structure, and using the AlphaFold2 database and DALI topology recognition program, they identified it as the lipoprotein YbjP. This protein has an anchorage in the outer membrane, and it was suggested that it could act as a support for TolC, which is the only OMF that does not have an N-terminal extension anchored in the outer membrane, which is very puzzling for the community working in this field of research.

Authors used a large number of different approaches to evaluate the importance of YbjP (structure, genomic evolution, microbiology, photocrosslink in vivo, proteomic profile), but did not succeed in finding it a clear role so far, even if it could be important depending on environmental stress. Nevertheless, their results, obtained with extreme rigour, are of main interest for the comprehension of the complexity of such systems and deserve publication.

Comments on revisions:

Thank you for clarifying the points that puzzled me concerning the crosslink experiments. This version does not need further modifications.

<https://doi.org/10.7554/eLife.110666.2.sa1>

Author Response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

The presentation and especially main-text illustrative material seem to focus disproportionately on MacAB-TolC-YbjP complex, and the AcrABZ-TolC-YbjP is relegated to supplementary data which is somewhat confusing. There is no high-resolution side view of the AcrABZ-TolC-YbjP side-by-side to MacAB-TolC-YbjP which may be helpful to spot parallels and differences in the organisation of the two systems.

This was previously presented in Supplementary Figure S2. However, because the models were shown at a small scale, we have now included the comparison in a main manuscript (Figure 4). This figure presents AcrABZ-TolC-YbjP and MacAB-TolC-YbjP side-by-side, a structural alignment of TolC-YbjP in the two pumps, and close-up views of the interaction interface.

Supplementary Figure 2 may also be better presented in the main text, as it shows specific displacements of residues upon binding of the YbjP relative to the apo-complexes, although this can be left at the authors' discretion.

We added more text to describe the displacements of residues upon YbjP binding: 'Nonetheless, the side chains of a few residues in TolC, which mainly correspond to positively charged amino acids (R18, R24, K214, R227, R234), reorient to interact with the YbjP lipoprotein partner (Figure 2B).'

Reviewer #1 (Recommendations for the authors):

The work is of high quality and requires minimal modifications, which are mentioned as suggestions above and are mostly connected to the illustrative material.

One additional suggestion, which is connected to the earlier BioRxiv preprint, the data seen in Fig 6 of the preprint seems to have been edited out from the current version, and perhaps can be included in a revised version, as it seems to support the "rapid adaptation under stress" role for YbjP, which currently is only speculatively mentioned in p.11, line 365 of the manuscript.

We acknowledge that the BioRxiv preprint Figure 6 can support the rapid adaptation under stress role for YbjP. However, upon sequencing the $\Delta ybjP$ strain from the Keio collection used in the preprint, we identified a large deletion in the yecT-flhD region. We therefore generated a new $\Delta ybjP$ strain without the yecT-flhD deletion and repeated the experiment. However, the results with the corrected strain did not support the previous conclusion, and these data were consequently removed in the current manuscript.

Reviewer #2 (Public review):

In Figure 3C, the experiment performed with AcrA is clear and the extra band appears at the proper size. On the right panel, it is clear that the crosslink doesn't work when pBPA is placed on residues too far from TolC. Only when introduced on N113 or T110 does a band appear.

This is in accordance with an interaction in vivo. Nevertheless, $17 + 54 = 71\text{kDa}$, which is more than the two bands appearing on the gel. This difference in size migration can occur, but it is not clear when looking at Figure S3. In Figure S3a, the purified proteins are highlighted at approximately the expected size ($\approx 20\text{kDa}$ instead of 17 for YbjP and between 56 and 60kDa in two bands for TolC instead of 54kDa). On the right panel, it seems that the bands are present exactly at the same position, instead of an upper band as expected for the crosslinked YbjP-TolC (at 71kDa). It would be clearer if having the control of the same sample without illumination, revealed by anti-TolC, to see the difference.

We thank the reviewer for pointing out this discrepancy. We identified an error in the molecular weight ladder, as one band was missing. This has now been corrected: YbjP migrates just below 17 kDa, consistent with Figure 3C. In addition, we previously reported a size of 54 kDa for TolC, whereas matured TolC, after signal peptide cleavage, is actually 52 kDa.

We believe that the differences in the apparent molecular weight observed in Figures 3A, 3C and S3 (now S2) mainly result from tagging and post-translation modifications.

In Figure 3A, we used the soluble construct His-YbjP₂₈₋₁₇₁₁ (theoretical $M_w \sim 18\text{kDa}$), as also done for the controls in Figures 3C and S3 (now S2). However, for the crosslinking samples, we used full-length His-tagged YbjP, which carries a post-translational lipid modification (theoretical $M_w \sim 19\text{kDa}$, considering the protein lipidation). The presence of the lipid chains

alters the migration as this species migrates at ~15 kDa (Fig 3A). Increased hydrophobicity, due here to YbjP lipidation, could accelerate the migration (Emmanuel et al. 2025 FEBS Open Bio).

In Figure 3A, we used the TolC-FLAG whose apparent M_w is ~52 kDa, as previously reported (Fig S3, Fitzpatrick et al. 2017). In Figure S3 (now S2), we used His-tagged TolC (theoretical M_w 55 kDa) for the control, which migrates above 56 kDa. In the crosslinking samples, however, we detect tag-free, endogenous TolC, with a theoretical M_w of ~51 kDa.

In conclusion, the crosslinked complex composed of lipidated FL YbjP (~15 kDa) and endogenous TolC (~51 kDa) would be expected to migrate at ~66 kDa, which is consistent with what is observed in Figures 3C and S3 (now S2).

A second point that could be discussed further is the comparison of the structure of the pump in the presence of the peptidoglycan with the images previously obtained by tomography. It is not totally clear to me if YbjP could have been positioned in these maps.

There is density corresponding to YbjP in the map obtained in the presence of peptidoglycan. To improve clarity, we have specified the location of the peptidoglycan relative to the pumps in the revised Figure 4, and Supplementary Figure S4, together with the position of YbjP. In both figures, the lipoprotein appears distant from the peptidoglycan density.

Reviewer #2 (Recommendations for the authors):

In addition, please add explanations in the legend of Figure 3C concerning the structures.

We added the following description of the structures: ‘As shown underneath, AcrA residues Q136 and Y137, proximal to TolC in the structure of the AcrABZ-TolC pump (PDB 5NG5), were replaced by pBPA. For YbjP, the two residues N113 and T110 proximal to TolC in the MacAB-TolC-YbjP complex (PDB 9QGY) and the three residues N43, N90 and H104 distal to TolC were mutated.’

It would be clearer if having the control of the same sample without illumination, revealed by anti-TolC, to see the difference.

As the amount of crosslinked material is low, samples were enriched via His-tag purification of YbjP prior to Western blotting. In the absence of illumination (see sample N113, UV-), no crosslink would be formed, and therefore TolC would not be co-purified.

In addition, some typo errors have been noted.

Table S1 minus is missing for the defocus range for AcrABZ-TolC-YbjP.

Thank you for noting the typo. We have added the minus sign.

Table S3, please specify what is N in the legend.

N is the stoichiometry parameter, which is now specified in the table legend.

Line 237, I suppose it has to refer to Figure S6, not S5.

Thank you for noting the error. We have verified the text matches the figures here and in the entire manuscript.

Several errors are present in the legend of Figure 6.

No letters are indicated for the different panels; line 841 must be C, F and I; the indicated colors for the differentially expressed proteins do not correspond to the volcano plots.

Thank you for suggesting the improvements for the labels. We have modified the plot accordingly.

Reference Glavier 2020 has been cited as Glacier on line 72.

We have modified the writing accordingly and checked the reference.

<https://doi.org/10.7554/eLife.110666.2.sa0>