

Temperature sensitivity of the interspecific interaction strength of coastal marine fish communities

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 2

June 7, 2023

Reviewed preprint version 1

April 25, 2023 (this version)

Sent for peer review

February 14, 2023

Posted to preprint server

December 15, 2022

Masayuki Ushio , Tetsuya Sado, Takehiko Fukuchi, Sachia Sasano, Reiji Masuda, Yutaka Osada, Masaki Miya

Hakubi Center, Kyoto University, Kyoto 606-8501, Japan • Center for Ecological Research, Kyoto University, Otsu 520-2113, Japan • Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR, China • Natural History Museum and Institute, Chiba, Chiba 260-8682, Japan • Maizuru Fisheries Research Station, Kyoto University, Maizuru, Kyoto 625-0086, Japan • Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Ishigaki, Okinawa 907-0451, Japan • Graduate School of Life Sciences, Tohoku University, Sendai 980-8578, Japan

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The effects of temperature on interaction strengths are important for understanding and forecasting how global climate change impacts marine ecosystems; however, tracking and quantifying interactions of marine fish species is practically difficult especially under field conditions, and thus, how temperature influences their interaction strengths under field conditions remains poorly understood. We herein performed quantitative fish environmental DNA (eDNA) metabarcoding on 550 seawater samples that were collected twice a month from 11 coastal sites for two years in the Boso Peninsula, Japan, and analyzed eDNA monitoring data using nonlinear time-series analytical tools. We detected fish-fish interactions as information flow between eDNA time series, reconstructed interaction networks for the top 50 frequently detected species, and quantified pairwise, fluctuating interaction strengths using nonlinear time series analysis methods. Although there was a large variation, water temperature generally influenced fish-fish interaction strengths. The impact of water temperature on interspecific interaction strengths varied among fish species, suggesting that fish species identity influences the temperature effects on interactions. For example, interaction strengths that *Halichoeres tenuispinis* and *Microcanthus stringatus* received strongly increased with water temperature, while those of *Engraulis japonicus* and *Girella punctata* decreased with water temperature. An increase in water temperature induced by global climate change may change fish interactions in a complex way, which consequently influences marine community dynamics and stability. Our research demonstrates a practical research framework to study the effects of environmental variables on interaction strengths of marine communities under field conditions, and thus, would contribute to further understanding and predicting natural marine ecosystem dynamics.

eLife assessment

This study presents **important** findings regarding the quantification of dynamics in fish communities in changing ecosystems by combining a large-scale environmental DNA metabarcoding time series with novel statistical approaches. The methods are **convincing**, with controlled experiments, thorough statistical analyses, and a substantial dataset covering two years of detailed observation, which can provide sufficient power to detect fine-scale ecological interactions. This work is relevant for informing future research on assessing community stability under climate change.

Introduction

Interspecific interactions are key to understanding and predicting the dynamics of ecological communities (May 1972 [↗](#); Wootton & Emmerson 2005 [↗](#); Tang *et al.* 2014 [↗](#); Wootton & Stouffer 2015 [↗](#)). Theoretical and empirical studies have shown that various properties of interspecific interactions, namely, the number of interactions, sign (positive or negative), strength, and pair-wise correlations of interactions (e.g., prey-predator interactions), influence the dynamics, stability, and diversity of ecological communities in terrestrial and aquatic ecosystems (Mougi & Kondoh 2012 [↗](#); Tang *et al.* 2014 [↗](#); Allesina *et al.* 2015 [↗](#); Ushio *et al.* 2018a [↗](#); Ratzke *et al.* 2020 [↗](#); Ushio 2022a [↗](#)). Interaction strength is a fundamental property, and previous studies examined the relationship between interaction strengths and ecological community properties (Wootton & Emmerson 2005 [↗](#); Wootton & Stouffer 2015 [↗](#)). The dominance of weak interactions stabilizes the dynamics of a natural fish community (Ushio *et al.* 2018a [↗](#)). Interaction strengths have been reported to decrease with increases in the diversity of experimental microbial communities (Ratzke *et al.* 2020 [↗](#)), and this holds true even for more diverse ecological communities under field conditions (Ushio 2022a [↗](#)).

Environmental variables exert significant effects on interspecific interaction strengths. The effects of temperature on interaction strengths are important for understanding and forecasting the impact of the ongoing global climate change on ecosystems. The relationship between temperature and interaction strengths has been investigated in terrestrial and aquatic ecosystems for decades (Coley & Aide 1991 [↗](#); Kishi *et al.* 2005 [↗](#); Adams & Zhang 2009 [↗](#); Rall *et al.* 2010 [↗](#); Kordas *et al.* 2011 [↗](#); Kratina *et al.* 2012 [↗](#); Hein *et al.* 2014 [↗](#); Allan *et al.* 2015 [↗](#); Thakur *et al.* 2017 [↗](#)). In terrestrial ecosystems, Coley & Aide (1991) [↗](#) proposed that interactions between plants and insect herbivores were generally stronger in warmer regions, which may result in stronger negative density dependence for tree species and contribute to higher plant diversity in a tropical region (Forrister *et al.* 2019 [↗](#)). The influence of temperature on interaction strengths has also been investigated in other systems: terrestrial arthropods interactions (Rall *et al.* 2010 [↗](#)), fish-fish interactions (Hein *et al.* 2014 [↗](#); Allan *et al.* 2015 [↗](#)), and fish-prey interactions (Kishi *et al.* 2005 [↗](#)). In addition, Wieczynski *et al.* (2021) [↗](#) showed that species-level functional traits (e.g., body size and shape) may underlie the relationships between temperature and interaction strengths. Therefore, interplays among temperature, interaction strengths, species identity, and community-level function (e.g., primary production) is prevalent, and a more detailed understanding of the complex interactions among them is critical for predicting ecosystem responses to climate change.

In marine ecosystems, interspecific interactions in ecological communities play a fundamental role in system dynamics, such as population dynamics, primary production, and nutrient cycling (Hannisdal *et al.* 2017 [↗](#); Ushio *et al.* 2018a [↗](#); Penn *et al.* 2019 [↗](#)), as well as ecosystem services, including food supply (Smith *et al.* 2019 [↗](#)). Although many studies have evaluated the effects of temperature on interaction strengths of marine organisms (Kordas *et al.* 2011 [↗](#)), most of these

studies have been performed targeting relatively immobile or small organisms (Bertness & Ewanchuk 2002 [DOI](#); Chen *et al.* 2012 [DOI](#)) or under laboratory conditions (e.g., mesocosm experiments; Allan *et al.* 2015 [DOI](#)). While the previous studies have provided invaluable information, it currently remains unclear how temperature influences the strengths of interspecific interactions of larger, more mobile marine organisms, such as fish, which may exert strong top-down regulations on ecological communities, especially under field condition. This may be due to the difficulties associated with detecting and measuring interaction strengths among multiple, relatively large, mobile species under field conditions; the identification of quickly-moving fish species and the quantification of their abundance under field conditions are challenging, and the quantification of their interactions is even more difficult. Overcoming these difficulties and understanding how temperature influences the strength of interspecific interactions of marine fish communities will provide insights into how marine fish communities are assembled, how they may respond to the ongoing and future global climate change, and how the changes in fish communities may transmit to other trophic levels.

Environmental DNA (eDNA), defined here as extra-organismal DNA left behind by macro-organisms (Bohmann *et al.* 2014 [DOI](#)), has been attracting increasing attention as an indirect genetic marker for inferring the presence of species for biodiversity monitoring (Deiner *et al.* 2017 [DOI](#); Cristescu & Hebert 2018 [DOI](#)). A simple protocol for collecting eDNA samples from aquatic environments facilitates continuous biodiversity monitoring at multiple sites (Deiner *et al.* 2017 [DOI](#)), and eDNA metabarcoding (the simultaneous detection of multiple species using universal primers and a high-throughput sequencer) provides useful information on the dynamics of ecological communities (Bálint *et al.* 2018 [DOI](#); Miya *et al.* 2020 [DOI](#); Miya 2022 [DOI](#); Ushio 2022a [DOI](#)). Recent studies demonstrated that eDNA metabarcoding combined with frequent water sampling enabled the efficient monitoring of high-diversity ecological communities (Bista *et al.* 2017 [DOI](#); Djurhuus *et al.* 2020 [DOI](#); Ushio 2022a [DOI](#)). Furthermore, if eDNA metabarcoding is performed quantitatively (e.g., by including spike-in DNAs; Ushio *et al.* 2018b [DOI](#)), these data may contain information on species abundance.

More importantly, recent studies have shown that information on interspecific interactions may be embedded in multispecies eDNA time series, particularly when the data is “quantitative” (Ushio 2022a [DOI](#)). Advances in nonlinear time series analyses have enabled the quantification of interspecific interactions only from time series data. For example, transfer entropy (TE) is a method based on the information theory that quantifies information flow between two variables (Schreiber 2000 [DOI](#); Runge *et al.* 2012 [DOI](#)). Information flow may be an index of interspecific interactions when applied to ecological data. Convergent cross mapping (CCM) is a causality detection tool in empirical dynamic modeling (Sugihara *et al.* 2012 [DOI](#)), which is based on the dynamical theory. TE and CCM have more recently been understood under the information theory framework, and unified information-theoretic causality (UIC) may also quantify information flow between variables (rUIC package; Osada & Ushio 2021 [DOI](#)). In addition, a recently developed, improved version of a sequentially-weighted global linear map (S-map), called the MDR S-map, enables accurate quantifications of interaction strengths of a large interaction network even when the number of network nodes exceeds the time series length (Chang *et al.* 2021 [DOI](#)). These advanced statistical tools facilitate the detection and quantification of interspecific interactions from quantitative, multispecies eDNA time-series data.

In the present study, we collected eDNA samples twice a month from 11 coastal sites at the southern tip of the Boso Peninsula, central Japan (**Fig. 1a** [DOI](#)) for two years (a total of 550 samples). This region is located on the Pacific side of Honshu Island around 35°N and is markedly affected by the warm Kuroshio Current, cold Oyashio Current, and inner-bay water from Tokyo Bay. These geographic and oceanographic characteristics form latitudinal temperature gradients, allowing us to investigate temperature effects on interspecific interactions. We obtained quantitative eDNA metabarcoding data by combining MiFish eDNA metabarcoding data (relative abundance data) and the eDNA concentration data obtained through quantitative PCR (qPCR) for one of the most

common fish species, *Acanthopagrus schlegelii* (e.g., absolute abundance data of *A. schlegelii* eDNA was utilized as an internal spike-in DNA; see Methods). Based on quantitative, multispecies eDNA time-series data, pairwise species interactions were detected as information flow between species using a nonlinear time series analysis (Osada & Ushio 2021 [↗](#)). In addition, interaction strengths at each time point were quantified for the detected fish-fish interactions using the MDR S-map method (Chang *et al.* 2021 [↗](#)). As our eDNA time series was taken twice a month, the interactions detected should also occur at the same time scale (i.e., about two-week), which means that we focus on behavior-level interactions rather than birth-death process in the present study. In addition, the interactions we detected by the time series analysis could include various types of interactions such as competition and mutualism that could be involved in the population dynamics. We hypothesized that (1) a positive relationship exists between temperature and interaction strengths at the community level, as found in other ecosystems and many types of interactions, and (2) the relationship between the patterns of temperature and interaction strengths varies among fish species because of species-specific ecologies and the behavior of component species.

Materials and Methods

Study sites and environmental observations

We selected 11 sites (Stations [Sts.] 1–11) for seawater sampling along the shoreline of the southern tip of the Boso Peninsula (**Fig. 1a** [↗](#)). The 11 stations are located within or surrounded by rocky shores with intricate coastlines. They are arranged such that the five stations on the Pacific and Tokyo Bay sides are approximately at the same latitudinal intervals, with St. 6 in between. The northernmost stations on both sides (Sts. 1 and 11) are markedly affected by the cold Oyashio Current (Yang *et al.* 1993 [↗](#)) and inner-bay water from Tokyo Bay (Fujiwara & Yamada 2002 [↗](#)), respectively, while the remaining stations (Sts. 2–10) are markedly affected by the warm Kuroshio Current flowing northward along the Pacific coast as well as its branches (Soh 2003 [↗](#)). Before seawater sampling at each station, we measured seawater temperature (°C) and salinity (‰) using a portable water-quality meter (WQC-30, Toa DKK, Tokyo, Japan). In addition, we recorded the sampling start time, latitude/longitude, weather and sea conditions, and turbidity.

Collection of eDNA samples

Detailed information on seawater sampling and on-site filtration is provided in Supporting Information. We collected seawater samples twice a month (i.e., around the first and second quarter moons) for two years between August 2017 and August 2019 by casting a bucket fastened to a rope at the 11 sites. We filtered the collected seawater using a combination of a filter cartridge and disposable syringe until the final filtration volume reached 1,000 mL for each replicate and obtained duplicate samples at each site. We added RNAlater (Thermo Fisher Scientific, DE, USA) into the cartridge to prevent eDNA degradation. We made a filtration blank (FB) by filtering 500 mL of purified water in the same manner at the end of each day of water sampling.

Laboratory protocol

Detailed information on the laboratory protocol is provided in Supporting Information. We thoroughly sterilized the workspace and equipment in the DNA extraction, pre-PCR, and post-PCR rooms before all experiments. We used filtered pipette tips and conducted all eDNA extraction, pre-PCR, and post-PCR manipulations in three different dedicated rooms that were separate from each other to safeguard against cross-contamination.

We extracted eDNA from filter cartridges using the DNeasy Blood & Tissue kit (Qiagen, Hilden, Germany) following the methods developed by Miya *et al.* (2016) [↗](#) with slight modifications (Miya & Sado 2019a [↗](#); Minamoto *et al.* 2021 [↗](#)). After removing RNAlater from filter cartridges, they




Figure 1.

Study sites and overall dynamics of environmental DNA (eDNA) concentrations and the number of fish species detected.

(a) Study sites in the Boso Peninsula. The study sites are influenced by the Kuroshio current (red arrow; left panel) and distributed along the coastal line in the Boso Peninsula (right panel). (b) Total eDNA copy numbers estimated by quantitative eDNA metabarcoding (see Methods for detail). (c) Fish species richness detected by eDNA metabarcoding. Points and lines indicate raw values and LOESS smoothing lines, respectively. The line color indicates the sampling site. Warmer colors generally correspond to study sites with a higher mean water temperature. For the high resolution figure, see <https://bit.ly/3uUZKHE>.

were subjected to lysis using proteinase K. We purified the collected DNA extracts using the above kit following the manufacturer's protocol and the final elution volume was set to 200 μ L. An extraction blank (EB) was also created during this process.

We employed two-step PCR for paired-end library preparation using the MiSeq platform (Illumina, CA, USA). We generally followed the methods developed by [Miya et al. \(2015\)](#) and subsequently modified by [Miya & Sado \(2019b\)](#). Detailed information on library preparation is provided in Supporting Information.

We performed the first-round PCR (1st PCR) using a mixture of the following six primers: MiFish-U-forward, MiFish-U-reverse, MiFish-E-forward-v2, MiFish-E-reverse-v2, MiFish-U2-forward, and MiFish-U2-reverse (**Table 1**). The 1st PCR amplified a hypervariable region of the mitochondrial 12S rRNA gene (ca. 172 bp; hereafter called the "MiFish sequence"). We also prepared a 1st PCR blank (1B) during this process, in addition to FB and EB. After the 1st PCR, PCR products were purified, quantified, diluted to 0.1 ng/ μ L, and used as templates for the second-round PCR (2nd PCR).

We performed the 2nd PCR to append dual-indexed sequences and flow cell-binding sites for the MiSeq platform. We prepared a 2nd PCR blank (2B) during this process in addition to FB, EB, and 1B. In total, we made 195 blanks (FB = 60, EB = 50, 1B = 50, 2B = 35) to monitor contamination during the on-site filtration, subsequent DNA extraction, and 1st and 2nd PCR of 550 samples.

We pooled each of the individual paired-end libraries in an equal volume, performed electrophoresis on an agarose gel (Invitrogen, CA, USA), and excised the target amplicons (ca. 370 bp). The concentrations of the size-selected libraries were measured, diluted to 10–12.0 pM, and sequenced on the MiSeq platform using the MiSeq v2 Reagent Kit for 2 $\times$ 150 bp PE (Illumina, CA, USA) following the manufacturer's protocol.

Sequence analysis

We performed data preprocessing and analyses of raw MiSeq reads from the MiSeq run using PMiFish ver. 2.4 (<https://github.com/rogotoh/PMiFish>; [Miya et al. 2020](#)). Detailed information on sequence data processing is provided in Supporting Information. Forward (R1) and reverse (R2) reads were merged while discarding short reads (<100 bp) after tail trimming and paired reads with too many differences (>5 positions) in the aligned region (ca. 65 bp). Primer sequences were removed from merged reads and reads without the primer sequences underwent quality filtering to remove low quality reads. Preprocessed reads were dereplicated, and all singletons, doubletons, and tripletons were removed from subsequent analyses to avoid false positives ([Edgar 2010](#)). Dereplicated reads were denoised to generate amplicon sequence variants (ASVs) that removed all putatively chimeric and erroneous sequences ([Edgar 2016](#); [Callahan et al. 2017](#)).

ASVs were subjected to taxon assignments to species names with a sequence identity of >98.5% with the reference sequences (two nucleotide differences allowed) and a query coverage of $\geq$ 90%. ASVs with identical fish species names were clustered as molecular operational taxonomic units (MOTUs) following the procedure described in Supporting Information. MiFish DB ver. 43 was used for taxon assignment, comprising 7,973 species distributed across 464 families and 2,675 genera. The ASV table was rarefied to the approximate minimum read number (ca. 20,000 reads). To refine the above taxon assignments, family-level phylogenies were reproduced from MiFish sequences from MOTUs and reference sequences (contained in MiFish DB ver. 43) belonging to these families. A total of 103 family-level trees were visually inspected and taxon assignments were revised. The final list of detected species is provided in Table S1.

Primer information	Primer sequence (5'–3') ^{1,2,3,4,5}	Length
1st PCR primers		
MiFish-U-forward	<i>ACACTCTTCCCTACACGACGCTCTCCGATCT</i> NNNNNN GTCGGTAAAACTCGTGCCAGC	60
MiFish-U-reverse	<i>GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT</i> NNNNNN CATAGTGGGGTATCTAATCCCAGTTTG	67
MiFish-E-forward-v2	<i>ACACTCTTCCCTACACGACGCTCTCCGATCT</i> NNNNNN RGTGGTAAATCTCGTGCCAGC	61
MiFish-E-reverse-v2	<i>GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT</i> NNNNNN GCATAGTGGGGTATCTAATCCTAGTTTG	63
MiFish-U2-forward	<i>ACACTCTTCCCTACACGACGCTCTCCGATCT</i> NNNNNN GCCGGTAAAACTCGTGCC	57
MiFish-U2-reverse	<i>GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT</i> NNNNNN CATAGGAGGGTGTCTAATCCCCGTTTG	67
2nd PCR primers		
2nd-PCR-forward	<u>AAATGATACGGCGACACCGAGATCTACAC</u> XXXXXXX <i>ACACTCTTCCCTACACGACGCTCTTCCGATCT</i>	70
2nd-PCR-reverse	<u>CAAGCAGAAGACGGCATACGAGAT</u> XXXXXXX <i>GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT</i>	66

¹Normal characters indicate target-specific universal primers (i.e., MiFish primers).

²The six random bases (Ns) in the middle of the 1st PCR primers were appended to enhance cluster separation on flow cells.

³Italic characters indicate Illumina sequencing primers.

⁴**X** indicates index sequences to identify each sample.

⁵Underlined characters indicate P5/P7 adapter sequences for Illumina sequencing.

Table 1.

Primer sequences used in the present study

Estimation of DNA copy numbers

The nonlinear time series analytical tools used in the present study require quantitative time-series data. To estimate fish eDNA concentrations, we initially measured the eDNA concentrations of the most common fish species in the region, Japanese Black Seabream (*A. schlegelii*), using qPCR, which was detected in 504 out of 550 seawater samples (91.6%). By dividing *A. schlegelii* sequence reads by the *A. schlegelii* eDNA quantity in each sample, we estimated the number of sequence reads generated per *A. schlegelii* eDNA copy for each sample, and we applied the same conversion factor (i.e., sequence reads / *A. schlegelii* eDNA copy) to other fish species. Note that sequence reads generated per eDNA copy may vary depending on factors such as the level of PCR inhibition and PCR amplification efficiency; however, in general, this “internal spike-in DNA” method has been shown to reasonably estimate the quantity of eDNA concentrations (Ushio *et al.* 2018b; Ushio 2022a). More importantly, species-specific biases that may be introduced in the internal spike-in DNA method do not cause serious biases in the outcomes of our nonlinear time-series analysis because it standardizes time series to have a zero mean and a unit variance before analyses and also only utilizes the fluctuation patterns of time series. When we did not detect any *A. schlegelii* eDNA by qPCR, we replaced the “zero” value with the minimum eDNA copy numbers of *A. schlegelii* (0.346 copies/μl DNA). Similarly, when we did not detect any *A. schlegelii* eDNA sequence reads by metabarcoding, we replaced the “zero” value with the minimum eDNA sequence reads of *A. schlegelii* (12 reads/sample). These corrections estimated how many sequence reads were generated per eDNA copy for all samples. Details are provided in Supporting Information.

Nonlinear time series analysis to detect fish-fish interactions

We detected fish-fish interactions using a nonlinear time series analysis based on quantitative eDNA time-series data from multiple species and sites. Since the reliable detection of interspecific interactions requires sufficient information in the target time series, we arbitrarily selected the top 50 most frequently detected species and excluded other rarer fish species from the analysis. We quantified information flow between two fish species’ eDNA time series by the “unified information-theoretic causality (UIC)” method implemented in the “rUIC” v0.1.5 package (Osada & Ushio 2021) of R (R Core Team 2022). UIC tests the statistical clarity of information flow between variables in terms of TE (Schreiber 2000) computed by nearest neighbor regression based on time-delay embedding of explanatory variables (i.e., cross mapping; Sugihara *et al.* 2012). In contrast to the standard method used to measure TE, UIC quantifies information flow as follows:

$$TE = \frac{1}{T} \sum_{t=1}^T \log \left(\frac{p(y_{t+tp} | x_t, x_{t-\tau}, \dots, x_{t-(E-1)\tau}, z_t)}{p(y_{t+tp} | x_{t-\tau}, x_{t-2\tau}, \dots, x_{t-(E-1)\tau}, z_t)} \right), \# \dots [1]$$

where x , y , and z represent a potential causal variable, effect variable, conditional variable (if available), respectively. $p(A | B)$ represents conditional probability: the probability of A conditioned on B . t , tp , τ , and E represent the time index, time step, a unit of time lag, and the optimal embedding dimension, respectively. T is the total number of points in the reconstructed state space (this is equivalent to the total number of time points minus $(E-1)$). For example, if TE between y_{t+1} and $\{x_t, x_{t-\tau}, \dots, x_{t-(E-1)\tau}\}$ is measured (tp is set to 1), it tests the causal relationship between y_{t+1} and x_t . Optimal E was selected by measuring TE as follows:

$$TE = \frac{1}{T} \sum_{t=1}^T \log \left(\frac{p(x_{t+tp} | y_t, x_t, x_{t-\tau}, \dots, x_{t-(E-1)\tau})}{p(x_{t+tp} | y_t, x_{t-\tau}, \dots, x_{t-(E-1)\tau})} \right), \# \dots [2]$$

where $E_R (< E)$ is the optimal embedding dimension of lower dimensional models. Significance tests were conducted by bootstrapping data after embedding (the significance levels were set to 0.05). Eqn. [2] is a TE version of simplex projection (Sugihara & May 1990). TE measured according to Eqn. [1] gains the advantage of previous causality tests, i.e., standard TE methods (Schreiber 2000; Runge *et al.* 2012) and convergent cross mapping (CCM) (Sugihara *et al.* 2012), the algorithm of which is explained and implemented in the rUIC website (<https://github.com/yutakaos/rUIC>).

By using UIC, we quantified TE between fish eDNA time series. As environmental variables, water temperature (°C), salinity (‰), wave height (m), and tide level (cm) were considered as conditional variables. This means that the effects of the environmental variables on the fish eDNA abundance were removed in the analysis when quantifying interspecific interaction strengths. Importantly, in most cases, water temperature had statistically clear influence on fish eDNA dynamics and included as a conditional variable in the embedding. Because water temperature showed clear seasonality in the region (Fig. S1), the effect of the seasonality in detecting causation was taken into account in this analysis. We merged all eDNA time series across the 11 study sites and standardized the eDNA time series to have zero means and a unit of variance before the analysis. If TE between two fish eDNA time series was statistically clearly higher than zero, we interpreted it as a sign of an interspecific interaction. After the detections of interspecific interactions among the top 50 fish species, we visualized the interaction network (i.e., the reconstruction of fish-fish interaction network). Importantly, UIC quantifies the average information flow between two time series (see Eqn. [1]), and thus there is only one TE value for each pair of fish species, which is critically different from interaction strengths quantified by the MDR S-map (see the following section).

Nonlinear time series analysis to quantify interaction strengths

We quantified fish-fish interaction strengths for the interspecific interactions detected by UIC. For this purpose, we used an improved version of the sequential locally-weighted global linear map (S-map) (Sugihara 1994 [\[3\]](#)), called the multiview-distance, regularized S-map (MDR S-map) (Chang *et al.* 2021 [\[4\]](#)). Consider a system that has E different interacting variables, and assume that the state space at time t is given by $x(t) = \{x_1(t), x_2(t), \dots, x_E(t)\}$. For each target time point t^* , the S-map method produces a local linear model that predicts the future value $x_1(t^*+p)$ from the multivariate reconstructed state space vector $x(t^*)$. That is,

$$\hat{x}_1(t^*+p) = IS_0 + \sum_{j=1}^E IS_j x_j(t^*) \quad [3]$$

where $\hat{x}_1(t^*+p)$ is a predicted value of x_1 at time t^*+p , IS_j is a regression coefficient and interpreted as interaction strength (or called S-map coefficient), and IS_0 is an intercept of the linear model. The linear model is fit to the other vectors in the state space. However, points that are close to the target point, $x(t^*)$, are given greater weighting (i.e., locally weighted linear regression). In the standard S-map, the distances between the target point, $x(t^*)$, and other points are measured by Euclidean distance. However, Euclidean distance cannot be a good measure if the dimension of the state space is high, and in this case, it is impossible to identify nearest neighbors correctly if we rely on Euclidean distance. In the MDR S-map, the distance between the target point and other points are measured by the multiview distance (Chang *et al.* 2021 [\[4\]](#)), which can be calculated by ensembling distances measured in various low-dimensional state spaces (multiview embedding; see Ye & Sugihara 2016 [\[5\]](#)). In addition, to reduce the possibility of overestimation and to improve forecasting skill, regularization (i.e., ridge regression) is also applied (Cenci *et al.* 2019 [\[6\]](#)). Chang *et al.* (2021) [\[4\]](#) showed that the MDR S-map outperformed other S-map methods and it enables improved estimations of interaction strengths. Importantly, the MDR S-map quantifies interaction strength at each time point (i.e., the maximum possible number of interaction strength values is the number of fish-fish interactions $\times$ the number of time points – the optimal embedding dimension + 1). In the present study, we implemented the MDR S-map in our custom R package, “macam” v0.0.10 (Ushio 2022b [\[7\]](#)) and used it for computation.

Quantification of the temperature sensitivity of fish species interactions

After reconstructing fish interaction networks and quantifying interaction strengths, we analyzed the relationships among fish species interaction strengths and biotic and abiotic variables. We performed all analyses using R v4.2.1 (R Core Team 2022 [\[8\]](#)). Generalized additive mixed models

(GAMMs) were performed using the “mgcv” package of R (Wood 2004 [↗](#)). We visualized results with “ggplot2” (Wickham 2009 [↗](#)) and “cowplot” (Wilke 2017 [↗](#)). In the present study, we used the term “statistical clarity” instead of “statistical significance” to avoid misinterpretations, according to the recommendations by Dushoff et al. (2019) [↗](#).

In the first analysis, the relationships between fish-fish interaction strengths and the characteristics of the study sites were analyzed by GAMM. In the analysis, in-strength (interaction strengths that a species receives) and out-strength (interaction strengths that a species gives) were separately calculated. In GAMM, in-strength or out-strength were an explained variable, and an environmental or ecological variable was an explaining variable. Explaining variables included water temperature, species richness, total fish eDNA concentrations, salinity, tide level, and wave height. A gamma distribution was assumed as the error distribution, the “log” function was used as a link function, and study site and fish species were used as a random effect (i.e., in R, `gamm(abs(IS) ~ s(explaining_variable), family = Gamma(link = “log”), random = list(site = ~1, fish_species = ~1))`, where `s()` indicates a smoothing term). The biological assumption behind this modeling is that explaining variables, such as water temperature, may linearly or nonlinearly influence fish species interactions. Moreover, the effects of explaining variables may randomly vary depending on the study site and fish species. The effect was considered to be statistically clear at $P < 0.05$.

In the second analysis, we focused on the effects of water temperature on the interaction strengths of each fish. In the analysis, GAMM was used again. A gamma distribution was assumed as the error distribution and the “log” function was used as a link function, and the study site was used as a random effect (i.e., in R, `gamm(abs(IS) ~ s(explaining_variable), family = Gamma(link = “log”), random = list(site = ~1))`, where `s()` indicates a smoothing term). GAMM was separately performed for each fish species. We also analyzed the effects of other properties on the interaction strengths of each fish species, which are provided in the Supporting Information. Again, the effect was considered to be statistically clear at $P < 0.05$.

Code and data availability

Sequence data were deposited in DDBJ Sequence Read Archives (DRA) (DRA accession number = DRA014111). Analysis scripts are available at Github (<https://github.com/ong8181/eDNA-BosoFish-network> [↗](#)).

Results and Discussion

Taxonomic diversity and dynamics of fish eDNA

Our multiple MiSeq runs generated 71,892,685 reads in total and detected 1,130 molecular operational taxonomic units (MOTUs). We inspected their family-level phylogenies to increase the accuracy of taxonomic assignments, recognizing 856 MOTUs across 33 orders, 167 families, and 466 genera (Table S1). This taxonomic diversity was similar to that of the local fauna (948 species across 33 orders, 158 families, and 493 genera) compiled from a literature survey and museum collections (Table S2).

Sequence reads in the ASV table were converted to estimated eDNA concentrations by using the eDNA concentrations of a common fish species detected across most samples (Japanese Black Seabream, *A. schlegelii*) (i.e., an analog of the internal spike-in DNA method; see Methods and Fig. S1). The converted ASV table included 23,863 detections and we selected the top 50 most frequently detected species for our time-series analysis. The detection of these 50 species ranged between 148 and 532 with a mean of 296, and their total detection reached 14,793 (62.0%).

Fish eDNA concentrations and fish species richness showed a clear intra-annual pattern (i.e., seasonality; **Fig. 1b, c** [↗](#)), which were higher in warmer months (e.g., between July and October) than in colder months (e.g., between January and March) at all sites. Total eDNA concentrations and species richness positively correlated with water temperature (Fig. S2).

Seasonal changes in eDNA concentrations were consistent with the patterns of seasonal occurrences in tropical and subtropical fish species in the Boso Peninsula, to which they are transported by the warm Kuroshio Current, settling on the coastal waters during the warmer months and subsequently disappearing during the colder months (Senou *et al.* 2006 [↗](#); Saito 2019 [↗](#)).

Reconstruction of the fish species interaction network

We reconstructed fish species interaction networks based on the quantitative, multispecies fish eDNA time series. Regarding the 50 frequently detected fish species, we quantified pairwise information flow between fish species. **Figure 2** [↗](#) shows reconstructed fish interaction networks for the 50 frequently detected fish species in the Boso Peninsula. At the regional scale, the linear correlations among water temperature, total eDNA concentrations, interaction strengths, the number of interactions, and species richness was all statistically clear ($P < 0.05$; Fig. S3) except for the linear correlation between water temperature and mean interaction strengths ($P > 0.05$). The interaction strengths became weak as species richness increased, which is consistent with a previous study (Ratzke *et al.* 2020 [↗](#); Ushio 2022a [↗](#)), suggesting that weak interactions are key to the maintenance of species-rich communities.

Most of the statistically clear information flow may be interpreted from the viewpoint of fish-fish behavior-level interspecific interactions (Table S3 and Supporting Discussion), which is convincing considering the time resolution of our eDNA time series (but see the section “*Potential limitations of the present study*” below). The largest information flow was detected from *Pseudoblennius marmoratus* to *Pseudolabrus eoethinus* (Table S3). These fish species have overlapping habitats, and thus, they may interact. Bidirectional information flow was detected between *P. eoethinus* and *Gymnothorax kidako*. They are both carnivorous fish species and may conduct joint hunting. *Chaenogobius annularis* and *Takifugu niphobles* are often found together in the sand between reefs. *T. niphobles* frequently dive in the sand, and benthos and mysids that are dug up during the dive may be prey for *C. annularis*. Further discussions about the detected information flow are described in Supporting Information (Table S3 and Supplementary Discussion).

Interaction strengths and environmental variables

We investigated how interaction strengths (i.e., regression coefficients estimated by the MDR S-map) changed with environmental variables (e.g., water temperature) and ecological properties (e.g., species richness and total eDNA concentrations) (**Figs. 3** [↗](#) and S4). The in-strengths and out-strengths of fish species interactions were statistically clearly associated with water temperatures, species richness, and total eDNA concentrations (GAMM, $P < 0.05$; **Fig. 3** [↗](#) and Table S4) except for the effects on species richness on the in-strengths (**Fig. 3b** [↗](#)).

In-strengths of the fish-fish interactions increased with water temperature (**Fig. 3a** [↗](#)), supporting our first hypothesis while out-strengths of the interactions showed an opposite pattern (**Fig. 3d** [↗](#)), which might suggest there is a difference in the temperature dependence between the in-strengths and out-strengths of the interactions. Indeed, water temperature may influence fish physiological activity (Kishi *et al.* 2005 [↗](#); Claireaux *et al.* 2006 [↗](#); Oyugi *et al.* 2012 [↗](#)) often in a complex way, and thus, the community-level influence of water temperature may also be complex as they should arise from the individual-level influence of water temperatures on fish. Interaction strengths were also statistically clearly influenced by species richness and total eDNA concentrations (except for **Fig. 3b** [↗](#)); the interaction strengths decreased with increasing water temperature. The effects of salinity, tide and wave were less clear, although the effects were statistically clear except for the





Figure 2.

Interaction networks of the fish community in the Boso Peninsula coastal region.

The “average” interaction network reconstructed by quantifying information transfer between eDNA time series. Transfer entropy (TE) was quantified by leveraging all eDNA time series from multiple study sites to draw this network. Only information flow larger than 80% quantiles (i.e., strong interaction) was shown as interspecific interactions for visualization. The edge color indicates scaled transfer entropy values, and fish illustration colors represent their ecology (e.g., habitat and feeding behavior). Node colors and node sizes indicate the fish family and fish abundance (total eDNA copy numbers of the fish species), respectively. For the high resolution figure, see <https://bit.ly/3uUZKHE>.

In-strength



Out-strength




Figure 3.

Dependence of interaction strengths on biotic and abiotic variables (50 dominant fish species and 11 study sites were leveraged).

The panels show the overall effects of biotic and abiotic variables on interaction strengths of the 50 dominant fish species: Effects of (a, d) water temperature, (b, e) species richness, and (c, f) total eDNA copy numbers. The Y-axis indicates the effects of the variables on fish-fish interaction strengths quantified by the MDR S-map method. a–c show the effects on the species interactions that a focal species receives (i.e., In-strength), and d–f show the effects on the species interactions that a focal species gives (i.e., Out-strength). The line indicates the average effects estimated by the general additive model (GAM), and the grey region indicates 95% confidential intervals. LME and GAM indicate the statistical clarity of the linear mixed model portion and GAM portion, respectively. Detailed statistical results and raw data are shown in Table S4 and Fig. S5, respectively.

effects of tide level on the out-strength (Figs. S4 and S6). Overall, environmental and ecological variables influenced the interaction strengths statistically clearly, but large variations remained unexplained (Figs. S5–S6), suggesting that other factors (e.g., fish ecology, nutrient, and physiological status) may influence the interaction strengths.

We also investigated how temperature influenced the interaction strength of individual fish species. **Figure 4** shows how interaction strengths among fish species changed with water temperature. For visualization purpose, we show fish species of which interaction strengths changed with water temperature highly statistically clearly ($P < 0.0001$). The in-strengths of several fish species clearly increased at higher water temperatures: for example, *Halichoeres tenuispinis*, *Macrocanthus strigatus*, *Pempheris schwenkii*, *Stethojulis interrupta terina*, and *Thalassoma cupido* (**Fig. 4a**). On the other hand, in-strengths of some fish species and out-strengths decreased at higher water temperatures: for example, in-strengths of *Ditrema temminckii temminckii*, *Engraulis japonicus*, and *Girella punctata*, and out-strengths of five fish species (**Fig. 4a,b**). These results support our second hypothesis that the relationship between the patterns of temperature and interaction strengths varies among fish species. These results suggest that, although water temperature may have strong influences on fish-fish interaction strengths in general, the sign of temperature effect (i.e., positive or negative) can vary depending on fish species and environmental conditions. Previous studies showed that fish physiological activities, such as feeding rates, growth rates, and swimming speed, were influenced by water temperature (Kishi *et al.* 2005; Claireaux *et al.* 2006; Oyugi *et al.* 2012). In addition, the direction (i.e., positive or negative) of the temperature effects on fish metabolic activities may be species-specific (Oyugi *et al.* 2012), and this species specificity may underlie the species-specific effects of temperature on the interaction strength. Also, importantly, other environmental conditions may influence fish-fish interactions (Figs. S7–S8), suggesting that the interaction strengths may fluctuate through time and space as shown in previous studies (Ushio *et al.* 2018a; Ushio 2022a). Conversely, interactions detected and quantified under a controlled environment might not necessarily be observed under field conditions, and thus, our research framework that enables the detection and quantification of interactions under field conditions would play a critical role in understanding the effects of temperature on fish-fish interactions under field conditions.

Potential limitations of the present study

The present results showing that the strengths of fish-fish interactions depend on water temperature rely on several assumptions that were not fully investigated, and thus, careful interpretations and discussions are required. First, the extent to which fish eDNA concentrations accurately represent fish abundance is an open question. Previous studies demonstrated that the quantity of eDNA may be a proxy of fish abundance and/or biomass (i.e., a positive correlation between the quantity of eDNA and fish abundance/biomass) (Takahara *et al.* 2012; Thomsen *et al.* 2012; Yamamoto *et al.* 2016). Our nonlinear time series analysis used the information embedded in fluctuations in time series, and time series are always standardized before the analysis. Therefore, species-specific differences in the copy numbers of the genetic marker and eDNA release rates, which may bias the estimation of absolute abundance, do not cause serious biases in the reconstruction of the networks and estimations of interaction strengths. However, accurate estimations of the absolute abundance of fish will improve the accuracy of analyses, and the integration of eDNA concentration data, knowledge on eDNA dynamics (i.e., release and degradation rates), and the hydrodynamic modeling of ocean water flow will be a promising approach (e.g., as in Fukaya *et al.* 2021).

Another limitation is that the reconstructed interaction network was not fully validated in the present study. Although previous studies demonstrated that network reconstructions based on a time series analysis work reasonably well and show meaningful patterns (Sugihara *et al.* 2012; Ushio *et al.* 2018a; Runge *et al.* 2019; Ushio 2022a), experiment/observation-based validations of the reconstructed network have not yet been performed. Potential causal




Figure 4.

Temperature dependence of fish species interactions at the species level.

a and **b** show temperature effects on fish species interactions quantified by the MDR S-map method. Note that the MDR S-map enables quantifications of interaction strengths at each time point, and thus the number of data points is large. (**a**) Points indicate the species interactions that a focal species (indicated by the strip label and fish image) receives (i.e., In-strength). (**b**) Points indicate the species interactions that a focal species (indicated by the strip label and fish image) gives (i.e., Out-strength). For **a** and **b**, only fish species of which interactions are statistically clearly affected by water temperature are shown (to exclude fish species with relatively weak temperature effects, $P < 0.0001$ was used as a criterion here). Point color indicates the study site. Gray line is drawn by GAM (the study sites were averaged for visualization purpose).

interactions between fish species may be reasonably interpreted (Supplementary Discussion); however, developing a framework to efficiently validate “previously unknown” interactions detected by eDNA data and a nonlinear time series analysis is an important future direction.

Furthermore, although we showed that temperature effects on interaction strength are relatively clear and common in the Boso peninsula coastal region, the range of water temperatures in the present analysis was still narrow and there is currently no evidence that we can generalize these results to other regions. Nonetheless, the advantages of the eDNA method are the low cost, time, and labor needed for field sampling as well as the potential scalability of the library preparation process (Buchner *et al.* 2021 [↗](#); Ushio *et al.* 2022 [↗](#)). Therefore, frequent (Ushio 2022a [↗](#)) and large spatial-scale ecological monitoring (e.g., ANEMONE DB; <https://db.anemone.bio/> [↗](#)) is possible, which will provide a more detailed understanding of the temperature-interaction strength relationships of fish on a larger spatial scale.

Conclusions

The present study demonstrated that the strengths of fish species interactions changed with water temperature under field conditions. For several fish species, species interactions were intensified in warmer water and for some other fish species, species interactions were weakened in warmer water. This may change the correlation and distribution of interaction strengths in a community, which may consequently influence community dynamics and stability (Tang *et al.* 2014 [↗](#); Allesina *et al.* 2015 [↗](#); Ushio *et al.* 2018a [↗](#)). Therefore, a more detailed understanding of the effects of environmental conditions on species interactions under field conditions will be required to improve our capability to understand, forecast, and even manage natural ecological communities and dynamics, which is particularly important under ongoing global climate change. Developments and improvements in eDNA techniques and statistical analyses are still required; however, because eDNA analysis is potentially applicable to any type of organisms even if they are difficult to be detected using traditional methods, our framework integrating an eDNA analysis and advanced statistical analyses pave a way to understand and forecast dynamics of various ecological communities under field conditions.

Author contributions

All authors agreed to the submission. MU and MM conceived the idea and designed the research; MM, TS, and TF collected the samples, performed eDNA metabarcoding and analyzed sequence data; SS and RM performed quantitative PCR analysis; MU compiled data; MU and YO analyzed data; MU and MM wrote the first draft and completed the final manuscript with contributions from all coauthors.

Acknowledgements

We thank T. Komai, R.O. Gotoh, T. Sunobe, and K. Takiguchi for assisting with the bimonthly collection of eDNA samples. This research was supported by JSPS KAKENHI (B) Grant Numbers JP19H03291, JP22H02691, and MEXT OGAP Project Grant Number JPMXD0618068274 to MM, JSPS KAKENHI (B) Grant Number 20H03323, and the Hakubi Project in Kyoto University to MU.

References

- Adams J.M., Zhang Y. (2009) **Is there more insect folivory in warmer temperate climates? A latitudinal comparison of insect folivory in eastern North America** *J. Ecol* **97**:933–940
- Allan B.J.M., Domenici P., Munday P.L., McCormick M.I. (2015) **Feeling the heat: the effect of acute temperature changes on predator–prey interactions in coral reef fish** *Conserv. Physiol* **3**
- Allesina S., Grilli J., Barabás G., Tang S., Aljadeff J., Maritan A. (2015) **Predicting the stability of large structured food webs** *Nat. Commun* **6**
- Bálint M., Pfenninger M., Grossart H.-P., Taberlet P., Vellend M., Leibold M.A., et al. (2018) **Environmental DNA Time Series in Ecology** *Trends Ecol. Evol* **33**:945–957
- Bertness M.D., Ewanchuk P.J. (2002) **Latitudinal and climate-driven variation in the strength and nature of biological interactions in New England salt marshes** *Oecologia* **132**:392–401
- Bista I., Carvalho G.R., Walsh K., Seymour M., Hajibabaei M., Lallias D., et al. (2017) **Annual time-series analysis of aqueous eDNA reveals ecologically relevant dynamics of lake ecosystem biodiversity** *Nat. Commun* **8**
- Bohmann K., Evans A., Gilbert M.T.P., Carvalho G.R., Creer S., Knapp M., et al. (2014) **Environmental DNA for wildlife biology and biodiversity monitoring** *Trends Ecol. Evol* **29**:358–367
- Buchner D., Macher T.-H., Beermann A.J., Werner M.-T., Leese F. (2021) **Standardized high-throughput biomonitoring using DNA metabarcoding: Strategies for the adoption of automated liquid handlers** *Environ. Sci. Ecotechnology* **8**
- Callahan B.J., McMurdie P.J., Holmes S.P. (2017) **Exact sequence variants should replace operational taxonomic units in marker-gene data analysis** *Isme J*
- Cenci S., Sugihara G., Saavedra S. (2019) **Regularized S-map for inference and forecasting with noisy ecological time series** *Methods Ecol. Evol* **10**:650–660
- Chang C.-W., Miki T., Ushio M., Lu H.-P., Shiah F.-K., Hsieh C. (2021) **Reconstructing large networks with time-varying interactions** *ArXiv210204069 Q-Bio*
- Chen B., Landry M.R., Huang B., Liu H. (2012) **Does warming enhance the effect of microzooplankton grazing on marine phytoplankton in the ocean?** *Limnol. Oceanogr* **57**:519–526
- Claireaux G., Couturier C., Groison A.-L. (2006) **Effect of temperature on maximum swimming speed and cost of transport in juvenile European sea bass (*Dicentrarchus labrax*)** *J. Exp. Biol* **209**:3420–3428
- Coley P.D., Aide T.M. (1991) **A comparison of herbivory and plant defenses in temperate and tropical broad-leaved forests**

- Cristescu M.E., Hebert P.D.N. (2018) **Uses and Misuses of Environmental DNA in Biodiversity Science and Conservation** *Annu. Rev. Ecol. Evol. Syst* **49**:209–230
- Deiner K., Bik H.M., Mächler E., Seymour M., Lacoursière-Roussel A., Altermatt F., et al. (2017) **Environmental DNA metabarcoding: Transforming how we survey animal and plant communities** *Mol. Ecol* **26**:5872–5895
- Djurhuus A., Closek C.J., Kelly R.P., Pitz K.J., Michisaki R.P., Starks H.A., et al. (2020) **Environmental DNA reveals seasonal shifts and potential interactions in a marine community** *Nat. Commun* **11**
- Dushoff J., Kain M.P., Bolker B.M. (2019) **I can see clearly now: Reinterpreting statistical significance** *Methods Ecol. Evol* **10**:756–759
- Edgar R.C. (2010) **Search and clustering orders of magnitude faster than BLAST** *Bioinformatics* **26**:2460–2461
- Edgar R.C. (2016) **UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing** *bioRxiv*
- Forrister D.L., Endara M.-J., Younkin G.C., Coley P.D., Kursar T.A. (2019) **Herbivores as drivers of negative density dependence in tropical forest saplings** *Science* **363**:1213–1216
- Fujiwara T., Yamada Y. (2002) **Inflow of oceanic water into Tokyo Bay and generation of a subsurface hypoxic water mass** *J. Geophys. Res. Oceans* **107**:13–1
- Fukaya K., Murakami H., Yoon S., Minami K., Osada Y., Yamamoto S., et al. (2021) **Estimating fish population abundance by integrating quantitative data on environmental DNA and hydrodynamic modelling** *Mol. Ecol* **30**:3057–3067
- Hannisdal B., Haaga K.A., Reitan T., Diego D., Liow L.H. (2017) **Common species link global ecosystems to climate change: dynamical evidence in the planktonic fossil record** *Proc. R. Soc. B Biol. Sci* **284**
- Hein C.L., Öhlund G., Englund G. (2014) **Fish introductions reveal the temperature dependence of species interactions** *Proc. R. Soc. B Biol. Sci* **281**
- Kishi D., Murakami M., Nakano S., Maekawa K. (2005) **Water temperature determines strength of top-down control in a stream food web** *Freshw. Biol* **50**:1315–1322
- Kordas R.L., Harley C.D.G., O'Connor M.I. (2011) **Community ecology in a warming world: The influence of temperature on interspecific interactions in marine systems** *J. Exp. Mar. Biol. Ecol., Global change in marine ecosystems* **400**:218–226
- Kratina P., Greig H.S., Thompson P.L., Carvalho-Pereira T.S.A., Shurin J.B. (2012) **Warming modifies trophic cascades and eutrophication in experimental freshwater communities** *Ecology* **93**:1421–1430
- May R.M. (1972) **Will a Large Complex System be Stable?** *Nature* **238**:413–414
- Minamoto T., Miya M., Sado T., Seino S., Doi H., Kondoh M., et al. (2021) **An illustrated manual for environmental DNA research: Water sampling guidelines and experimental protocols** *Environ. DNA* **3**:8–13

- Miya M. (2022) **Environmental DNA Metabarcoding: A Novel Method for Biodiversity Monitoring of Marine Fish Communities** *Annu. Rev. Mar. Sci* **14**:161–185
- Miya M., Gotoh R.O., Sado T. (2020) **MiFish metabarcoding: a high-throughput approach for simultaneous detection of multiple fish species from environmental DNA and other samples** *Fish. Sci* **86**:939–970
- Miya M., Minamoto T., Yamanaka H., Oka S., Sato K., Yamamoto S., et al. (2016) **Use 20 of a filter cartridge for filtration of water samples and extraction of environmental DNA** *J. Vis. Exp* :e54741–e54741
- Miya M., Sado T. (2019) **DNA extraction from a filter cartridge** *In: Environmental DNA sampling and experimental manual version 2.1. eDNA Methods Standardization Committee* :31–42
- Miya M., Sado T. (2019) **Multiple species detection using MiFish primers** *In: Environmental DNA sampling and experimental manual version 2.1. eDNA Methods Standardization Committee* :55–92
- Miya M., Sato Y., Fukunaga T., Sado T., Poulsen J.Y., Sato K., et al. (2015) **MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species** *R. Soc. Open Sci* **2**
- Mougi A., Kondoh M. (2012) **Diversity of interaction types and ecological community stability** *Science* **337**:349–351
- Osada Y., Ushio M. (2021) **R package for Unified Information-theoretic Causality**
- Oyugi D.O., Cucherousset J., Robert Britton J. (2012) **Temperature-dependent feeding interactions between two invasive fishes competing through interference and exploitation** *Rev. Fish Biol. Fish* **22**:499–508
- Penn J.L., Weber T., Chang B.X., Deutsch C. (2019) **Microbial ecosystem dynamics drive fluctuating nitrogen loss in marine anoxic zones** *Proc. Natl. Acad. Sci* **116**:7220–7225
- R Core Team (2022) **R: A language and environment for statistical computing**
- Rall B.C., Vucic-Pestic O., Ehnes R.B., Emmerson M., Brose U. (2010) **Temperature, predator-prey interaction strength and population stability** *Glob. Change Biol* **16**:2145–2157
- Ratzke C., Barrere J., Gore J. (2020) **Strength of species interactions determines biodiversity and stability in microbial communities** *Nat. Ecol. Evol*
- Runge J., Bathiany S., Bollt E., Camps-Valls G., Coumou D., Deyle E., et al. (2019) **Inferring causation from time series in Earth system sciences** *Nat. Commun* **10**:1–13
- Runge J., Heitzig J., Petoukhov V., Kurths J. (2012) **Escaping the Curse of Dimensionality in Estimating Multivariate Transfer Entropy** *Phys. Rev. Lett* **108**
- Saito H., Nagai T., Saito H., Suzuki K., Takahashi M (2019) **The Kuroshio: its recognition, scientific activities and emerging issues. Kuroshio current: Physical, biogeochemical, and ecosystem dynamics** *In: Kuroshio Current: Physical, Biogeochemical, and Ecosystem Dynamics, Geophysical monograph series*
- Schreiber T. (2000) **Measuring Information Transfer** *Phys. Rev. Lett* **85**:461–464

- Senou H., Matsuura K., Shinohara G. (2006) **Checklist of Fishes in the Sagami Sea with Zoogeographical Comments on Shallow Water Fishes Occurring along the Coastlines under the Influence of the Kuroshio Current** *国立科学博物館専報* :389–542
- Smith P., Adams J., Beerling D.J., Beringer T., Calvin K.V., Fuss S., et al. (2019) **Land-management options for greenhouse gas removal and their impacts on ecosystem services and the sustainable development goals** *Annu. Rev. Environ. Resour* **44**:255–286
- Soh W. (2003) **Transport processes deduced from geochemistry and the void ratio of surface core samples, deep sea Sagami Bay, central Japan** *Prog. Oceanogr., Long-Term Monitoring of Sedimentary Processes in Central Sagami Bay Japan* **57**:109–124
- Sugihara G. (1994) **Nonlinear Forecasting for the Classification of Natural Time Series** *Philos. Trans. R. Soc. Math. Phys. Eng. Sci* **348**:477–495
- Sugihara G., May R., Ye H., Hsieh C., Deyle E., Fogarty M., et al. (2012) **Detecting causality in complex ecosystems** *Science* **338**:496–500
- Sugihara G., May R.M. (1990) **Nonlinear forecasting as a way of distinguishing chaos from measurement error in time series** *Nature* **344**:734–41
- Takahara T., Minamoto T., Yamanaka H., Doi H., Kawabata Z. (2012) **Estimation of fish biomass using environmental DNA** *PLOS ONE* **7**
- Tang S., Pawar S., Allesina S. (2014) **Correlation between interaction strengths drives stability in large ecological networks** *Ecol. Lett* **17**:1094–100
- Thakur M.P., Künne T., Griffin J.N., Eisenhauer N. (2017) **Warming magnifies predation and reduces prey coexistence in a model litter arthropod system** *Proc. R. Soc. B Biol. Sci* **284**
- Thomsen P.F., Kielgast J.O.S., Iversen L.L., Wiuf C., Rasmussen M., Gilbert M.T.P., et al. (2012) **Monitoring endangered freshwater biodiversity using environmental DNA** *Mol. Ecol* **21**:2565–2573
- Ushio M. (2022) **Interaction capacity as a potential driver of community diversity** *Proc. R. Soc. B Biol. Sci* **289**
- Ushio M. (2022) **macam: A collection of miscellaneous functions for ecology in R**
- Ushio M., Furukawa S., Murakami H., Masuda R., Nagano A.J. (2022) **An efficient early-pooling protocol for environmental DNA metabarcoding** *bioRxiv*
- Ushio M., Hsieh C., Masuda R., Deyle E.R., Ye H., Chang C.-W., et al. (2018) **Fluctuating interaction network and time-varying stability of a natural fish community** *Nature* **554**:360–363
- Ushio M., Murakami H., Masuda R., Sado T., Miya M., Sakurai S., et al. (2018) **Quantitative monitoring of multispecies fish environmental DNA using high-throughput sequencing** *Metabarcoding Metagenomics* **2**
- Wickham H. (2009) **ggplot2: Elegant Graphics for Data Analysis**

- Wieczynski D.J., Singla P., Doan A., Singleton A., Han Z.-Y., Votzke S., et al. (2021) **Linking species traits and demography to explain complex temperature responses across levels of organization** *Proc. Natl. Acad. Sci* **118**
- Wilke C.O. (2017) **cowplot: Streamlined Plot Theme and Plot Annotations for “ggplot2.”**
- Wood S.N. (2004) **Stable and efficient multiple smoothing parameter estimation for generalized additive models** *J. Am. Stat. Assoc* **99**:673–686
- Wootton J.T., Emmerson M. (2005) **Measurement of interaction strength in nature** *Annu. Rev. Ecol. Evol. Syst* **36**:419–444
- Wootton K.L., Stouffer D.B. (2015) **Many weak interactions and few strong; food-web feasibility depends on the combination of the strength of species’ interactions and their correct arrangement** *Theor. Ecol*
- Yamamoto S., Minami K., Fukaya K., Takahashi K., Sawada H., Murakami H., et al. (2016) **Environmental DNA as a ‘snapshot’ of fish distribution: A case study of Japanese jack mackerel in Maizuru bay, Sea of Japan** *PLOS ONE* **11**
- Yang S.-K., Nagata Y., Taira K., Kawabe M. (1993) **Southward intrusion of the intermediate Oyashio water along the east coast of the Boso Peninsula, Japan II. Intrusion events into Sagami Bay** *J. Oceanogr* **49**:173–191
- Ye H., Sugihara G. (2016) **Information leverage in interconnected ecosystems: Overcoming the curse of dimensionality** *Science* **353**:922–925

Article and author information

Masayuki Ushio

Hakubi Center, Kyoto University, Kyoto 606-8501, Japan, Center for Ecological Research, Kyoto University, Otsu 520-2113, Japan, Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR, China

For correspondence: ong8181@gmail.com

ORCID iD: [0000-0003-4831-7181](https://orcid.org/0000-0003-4831-7181)

Tetsuya Sado

Natural History Museum and Institute, Chiba, Chiba 260-8682, Japan

Takehiko Fukuchi

Natural History Museum and Institute, Chiba, Chiba 260-8682, Japan

Sachia Sasano

Maizuru Fisheries Research Station, Kyoto University, Maizuru, Kyoto 625-0086, Japan, Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Ishigaki, Okinawa 907-0451, Japan

Reiji Masuda

Maizuru Fisheries Research Station, Kyoto University, Maizuru, Kyoto 625-0086, Japan

ORCID iD: [0000-0002-2184-1533](https://orcid.org/0000-0002-2184-1533)

Yutaka Osada

Graduate School of Life Sciences, Tohoku University, Sendai 980-8578, Japan

Masaki Miya

Natural History Museum and Institute, Chiba, Chiba 260-8682, Japan

ORCID iD: [0000-0002-9791-9886](https://orcid.org/0000-0002-9791-9886)

Copyright

© 2023, Ushio et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Lara Urban

Helmholtz Zentrum München, Neuherberg, Germany

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #1 (Public Review):

The authors have studied the effect of temperature on the interspecific interaction strength of coastal marine fish communities, using eDNA samples. Their introduction describes the state of the art concerning the dynamics of interspecific interactions in ecological communities. This introduction is well written and highly information dense, summarizing all that the reader needs to know to further understand their study setup and execution.

The authors hypothesize that water temperature changes could have an effect on the interspecific interaction strength between marine fishes, and they studied this with a two year long, bi-weekly eDNA sampling campaign at 11 study sites in Japan with different temperature gradients. These 550 water samples were analysed for fish biodiversity through eDNA-metabarcoding using MiFish primers. By using the most abundant fish species as an internal spike in and quantifying the copy numbers from this species by qPCR, the authors were able estimate DNA copy numbers for the total dataset. From the 50 most frequently detected fish species in these samples they showed that temperature affected the interspecific interaction strength between some species. Their work provides a highly relevant approach to perform species-interaction strength analysis based on eDNA biodiversity assessments, and as such provides a research framework to study marine community dynamics by eDNA, which is highly relevant in the study of ecosystem dynamics. The models and analytical methods used are clearly described and made available, enabling application of these methods by anyone interested in applying it to their own site and species group of interest.

Strengths: The authors have a study setup that is suitable to measure the effects of temperature of the eDNA diversity, and have taken a large number of samples and all appropriate controls to be able to accurately measure and describe these dynamics. The applied internal spike in to enable relative eDNA copy number quantification is convincing.

Weaknesses: The authors aim to study the relationship between species interaction strength and ecosystem complexity, and how temperature will influence this. However, there is only limited ecological context discussed explaining their results, and a link with climate change scenario's is also limited. A further discussion of this would have strengthened the manuscript.

The authors were able to find a correlation between water temperature and interaction strengths observed. However, since water temperature is dependent on many environmental variables that are either directly or indirectly influencing ecosystem dynamics, it is hard to prove a direct correlation between the observed changes in community dynamics and the temperature alone.

- <https://doi.org/10.7554/eLife.85795.1.sa1>

Reviewer #2 (Public Review):

In this work Ushio et al. combine environmental DNA metabarcoding with novel statistical approaches to demonstrate how fish communities respond to changing sea temperatures over a seasonal cycle. These findings are important due to the need for new techniques that can better measure community stability under climate change. The eDNA metabarcoding dataset of 550 water samples over two years is, I feel, of sufficient scale to provide power to detect fine-scale ecological interactions, the experiments are well controlled, and the statistical analysis is thorough.

The major strengths of the manuscript are: (1) the magnitude of the dataset, which provides densely replicated sampling that can overcome some of the noise associated with eDNA metabarcoding data and scale up the number of data points to make unique inferences; (2) the novel method of transforming the metabarcode reads using endogenous qPCR "spike-in" data from a common reference species to obtain estimates of DNA concentration across other species; and (3) the statistical analysis of time-series and network data and translating it into interaction strengths between species provides a cross-disciplinary dimension to the work.

I feel like this kind of study showcases the power of eDNA metabarcoding to answer some really interesting questions that were previously unobtainable due to the complexities and cost of such an exercise. Notwithstanding the problems associated with PCR primer bias and PCR stochasticity, the qPCR "spike-in" method is easy to implement and will likely become a standardised technique in the field. Further studies will examine and improve on it.

Overall I found the manuscript to be clear and easy to follow for the most part. I did not identify any serious weaknesses or concerns with the study, although I am not able to comment on the more complex statistical procedures such as the "unified information-theoretic causality" method devised by the authors. The section on limitations of the study is important and acknowledges some issues with interpretation that need to be explained. The methods, while brief in parts, are clear. The code used to generate the results has been made available via a GitHub repository. The figures are clear and attractive.

- <https://doi.org/10.7554/eLife.85795.1.sa0>