

Medicine

An Unexpected Role of Neutrophils in Clearing Apoptotic Hepatocytes *In Vivo*

Reviewed Preprint

Revised by authors after peer review.

[About eLife's process](#)

Reviewed preprint version 2

June 23, 2023 (this version)

Reviewed preprint version 1

April 4, 2023

Posted to bioRxiv

February 8, 2023

Sent for peer review

February 8, 2023

Luyang Cao, Lixiang Ma, Juan Zhao, Xiangyu Wang, Xinzou Fan, Wei Li, Yawen Qi, Yingkui Tang, Jieya Liu, Shengxian Peng, Li Yang, Liangxue Zhou, Li Li, Xiaobo Hu, Yuan Ji, Yingyong Hou, Yi Zhao, Xianming Zhang, Youyang Zhao, Yong Zhao, Yuquan Wei, Asrar B. Malik, Hexige Saiyin ✉, Jingsong Xu ✉

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041 • Guangzhou Regenerative Medicine and Health Guangdong Laboratory (GRMH-GDL), Guangzhou, China, 510530 • Department of Anatomy, Histology & Embryology, Shanghai Medical College, China, 200032 • Department of Anesthesiology, Huashan Hospital, Fudan University, Shanghai, China, 200040 • Clinical Laboratory, Longhua Hospital, Shanghai University of Traditional Medicine, China, 200043 • Department of Pathology, Zhongshan Hospital Fudan University, Shanghai, China, 200043 • Department of Rheumatology and Immunology, West China Hospital, Sichuan University, China, 610041 • Program for Lung and Vascular Biology, Stanley Manne Children's Research Institute, Ann & Robert H. Lurie Children's Hospital of Chicago, and Department of Pediatrics, Division of Critical Care, Northwestern University Feinberg School of Medicine, Chicago, USA, 60611 • State Key Laboratory of Membrane Biology, Institute of Zoology, Chinese Academy of Sciences, China • Department of Pharmacology, University of Illinois, College of Medicine, Chicago, USA, 60612 • State Key Laboratory of Genetic Engineering, School of Life Sciences, Fudan University, Shanghai, China, 200433

 (https://en.wikipedia.org/wiki/Open_access)

 (<https://creativecommons.org/licenses/by/4.0/>)

Abstract

Billions of apoptotic cells are removed daily in a human adult by professional phagocytes (e.g. macrophages) and neighboring nonprofessional phagocytes (e.g. stromal cells). Despite being a type of professional phagocyte, neutrophils are thought to be excluded from apoptotic sites to avoid tissue inflammation. Here we report a fundamental and unexpected role of neutrophils as the predominant phagocyte responsible for the clearance of apoptotic hepatic cells in the steady state. In contrast to the engulfment of dead cells by macrophages, neutrophils burrowed directly into apoptotic hepatocytes, a process we term *perforocytosis*, and ingested the effete cells from the inside. The depletion of neutrophils caused defective removal of apoptotic bodies, induced tissue injury in the mouse liver and led to the generation of autoantibodies. Human autoimmune liver disease showed similar defects in the neutrophil-mediated clearance of apoptotic hepatic cells. Hence, neutrophils possess a specialized immunologically silent mechanism for the clearance of apoptotic hepatocytes through perforocytosis, and defects in this key housekeeping function of neutrophils contribute to the genesis of autoimmune liver disease.

eLife assessment

This paper reports the **fundamental** discovery of a new function of neutrophil in specifically clearing apoptotic hepatocytes by penetrating the cells rather than engulfing them without causing inflammation as a part of tissue homeostasis. This **solid** study transforms the way we think about role of neutrophil in pathogenesis of autoimmune liver disease.

Introduction

Apoptosis is a process of programmed cell death that clears aged or damaged cells to maintain internal tissue homeostasis (Hochreiter-Hufford and Ravichandran, 2013). An estimated hundred billion cells undergo apoptosis daily in a human adult (Fond and Ravichandran, 2016). These apoptotic cells must be disposed of promptly and efficiently without littering cytoplasm or causing inflammation (Franz et al., 2006). Defects in the clearance of apoptotic bodies are often linked to various inflammatory and autoimmune diseases (Nagata et al., 2010; Poon et al., 2014). It is widely believed that apoptotic bodies are cleared by professional phagocytes, such as macrophages and immature dendritic cells, or by local nonprofessional phagocytes, such as epithelial cells, endothelial cells, and fibroblasts (Lauber et al., 2004). Although the morphology of apoptotic cells is easily distinguished in histological sections (Kerr et al., 1972), these cells are infrequently observed in normal human samples due to efficient removal by phagocytes (Lauber et al., 2004; Poon et al., 2014). Hence, the phagocytes responsible for the removal of apoptotic cells in the homeostatic state remain uncertain and it is unknown whether they are tissue specific.

Neutrophils, a type of terminally differentiated and short-living phagocytic cell, represent 50-70% of the total white blood cell population in humans (Amulic et al., 2012; Kolaczowska and Kubes, 2013). However, neutrophils are not considered key players in apoptotic cell clearance (Nagata et al., 2010; Poon et al., 2014). They instead function as inflammatory cells responsible for killing bacteria and fighting infection. Blood neutrophils often swarm to the site of infection, where they release toxic mediators (e.g., reactive oxygen species and proteases) that not only kill microorganisms but also can damage tissues (Amulic et al., 2012; Kolaczowska and Kubes, 2013). To prevent inflammation, apoptotic cells have been shown to release 'keep-out' signals, such as lactoferrin, that prevent the recruitment of neutrophils to the apoptotic site (Bournazou et al., 2009). Nevertheless, neutrophils are related to multiple autoimmune diseases (Amulic et al., 2012; Kolaczowska and Kubes, 2013). For example, mild neutropenia was observed to precede and accompany the onset of type 1 diabetes, a typical autoimmune disease (Harsunen et al., 2013; Valle et al., 2013). The role of neutrophils in the clearance of apoptotic bodies and contribute to autoimmunity remain to be solved (Jorch and Kubes, 2017).

Here we report an unexpected role of neutrophils in clearing apoptotic hepatocytes under physiological conditions. We found that neutrophils, not macrophages or other cells, are responsible for apoptotic hepatocyte clearance in an immunologically silent manner. We noted that neutrophils penetrate apoptotic hepatocytes and clear them from the inside, thereby avoiding spillage of cytoplasmic content; this mechanism differs sharply from the classical engulfment of dead cells by macrophages. Defects in neutrophil-mediated apoptotic clearance have been observed in human autoimmune liver (AIL) disease. Hence, in addition

to their well-known role in combating infection, neutrophils can function as housekeepers for apoptotic clearance and thus maintain tissue homeostasis.

Results

Neutrophils burrow into apoptotic hepatocytes in human livers

We observed a large number of neutrophils in hepatocytes in human noncancerous liver tissue obtained from patients with hepatocellular carcinoma (Fig. 1A) or from patients with hepatic hemangioma (Fig. 1B). We discovered that the hepatocytes occupied by neutrophils were apoptotic, as evidenced by the condensed chromatin signature (Fig. 1A, panels i-vi, black arrowheads indicate condensed chromatin, white arrowheads point to neutrophils with a characteristic multilobed nucleus). Importantly, apoptotic hepatocytes are rarely observed in the human liver, possibly due to the rapid removal of apoptotic cells by phagocytes. In a total of 281 apoptotic hepatocytes from 32 livers, we observed that each apoptotic hepatocyte was engorged by up to 22 neutrophils (Fig. 1C, Table S1). We also confirmed apoptotic hepatocytes by TUNEL staining or Caspase-3 immunostaining (Fig. 1D). Neutrophils burrowed inside apoptotic hepatocytes were either stained with an antibody against neutrophil elastase (NE), myeloperoxidase (MPO) or recognized by their multilobed nucleus signature (Fig. 1D, S1A). The distances from burrowed neutrophils to the apoptotic hepatocyte border were analyzed by IMARIS software and recorded in Table S2. Other immune cells were rarely associated with apoptotic hepatocytes. Kupffer cells are liver-resident phagocytes that are thought to clear apoptotic bodies (Canbay et al., 2003; Eipel et al., 2007). Upon staining with an antibody against CD68, a marker of Kupffer cells, we observed few Kupffer cells invading or engulfing apoptotic hepatocytes (Fig. 1C, E, 7A and Table S3). Furthermore, we detected few CD11b⁺ or CD45RA⁺ cells associated with the apoptotic sites (Fig. S1B, Table S4, S5).




Figure 1.

Neutrophil-mediated clearance of apoptotic cells in livers.

(A) Hematoxylin staining of human noncancerous liver tissues from patients with hepatocellular carcinoma. Apoptotic hepatocytes with apparent condensed chromatin are being scavenged by several neutrophils (black arrowhead indicates condensed chromatin and white arrowheads point to neutrophils with a characteristic multilobed nucleus). There are three stages of neutrophil scavenging apoptotic hepatocytes in human samples. Stage 1 (i-iii), initial invading stage: two neutrophils start to invade an apoptotic hepatocyte with apparent condensed chromatin (outlined by the black rectangle); three other neutrophils are already inside the cytoplasm (i); at this stage, the apoptotic hepatocytes are still attached to neighboring hepatocytes and occupied by 2-16 neutrophils (ii-iii, 37 out of 241 apoptotic hepatocytes observed are at stage 1). Stage 2, phagocytosis and shrinking stage: The apoptotic hepatocytes are being phagocytosed by invading neutrophils and are partially detached from

marker of KCs (indicated by green arrowheads). KCs do not invade or phagocytose apoptotic hepatocytes (black arrowheads). Data are representative of (A, B, D, E) or from (C) a total of 281 apoptotic hepatocytes observed, mean and s.e.m. in C.

Based on the above observations that neutrophils are the predominant phagocyte associated with apoptotic hepatocytes, we hypothesized that the neutrophil-mediated clearance of apoptotic cells consists of the following three sequential steps. In the initial invading or burrowing stage, activated neutrophils identified and targeted the apoptotic hepatocytes (cells with condensed chromatin, indicated by black arrowheads) and attached to their cell membrane (Fig. 1A, panel i, outlined by black rectangle). Then, the neutrophils invaded apoptotic hepatocytes (Fig. 1A, panels ii and iii). We observed an average of 7 neutrophils entering each apoptotic hepatocyte, and we termed this process perforocytosis (from Latin *perfero* to bore) (Fig. 1A, white arrowheads point to neutrophils). The second step consisted of phagocytosis and detachment. The neutrophils within hepatocytes appeared to clear apoptotic bodies from the inside without destroying the cellular membrane or extruding the cytoplasm (two burrowed neutrophils phagocytosed apoptotic debris as shown in Fig. S1C). Following digestion by neutrophils, the apoptotic hepatocytes decreased in size, and detached from nearby hepatocytes (Fig. 1A, panels iv-vi). The third step involved the complete digestion of apoptotic hepatocytes (Fig. 1A, panels vii-ix). After the clearance of apoptotic hepatocytes, neutrophils seemed to migrate away from the cleared space, possibly to make room for new hepatocytes generated by rapid division. This ‘eating inside’ phagocytosis process differed sharply from the well-known engulfment of apoptotic cells or fragmented apoptotic bodies by other phagocytic cells, such as macrophages (Poon et al., 2014).

Visualization of neutrophil perforocytosis in mouse livers

Neutrophils within apoptotic hepatocytes were confirmed in mouse livers by intravital microscopy (Fig. 2A, B) and electron microscopy (Fig. 2C). Similar to the observations in human samples, apoptotic hepatocytes in WT mouse livers were occupied by neutrophils but not associated with macrophages (Fig. 2A, neutrophils were labeled with an i.v. injection of anti-Ly6G antibody, macrophages were labeled with anti-F4/80 antibody and apoptotic cells were labeled with Annexin V). A total of 24 apoptotic cells were observed in 8 WT livers with an average of 2 burrowed neutrophils in each mouse apoptotic hepatocyte. The burrowed neutrophils were projected and analyzed by IMARIS software and results were compared side by side with livers from MRP8cre/DTR mice (Fig. 2A, with more details in the neutrophil depletion section below). The distances of burrowed neutrophils to the border of apoptotic hepatocytes were recorded in Table S6.




Figure 2.

Neutrophil burrowing into apoptotic hepatocytes.

(A) Intravital microscopy images of liver tissues from WT and MRP8cre/DTR mice. Neutrophils are labeled with an i.v. injection of anti-Ly6G antibody (green), KCs are labeled with anti-F4/80 antibody (red), and apoptotic cells are labeled with Annexin V (blue). Neutrophils inside and KCs associated with apoptotic hepatocytes are detected and analyzed with IMARIS software as described in Methods. The distances from neutrophils to the apoptotic hepatocyte border are recorded in table S6. A total of 24 apoptotic cells were observed in the WT liver with an average of 2 burrowed neutrophils. Scale bar, 100 μm . (B) Intravital image sequences of neutrophils phagocytosing apoptotic hepatocytes in mouse livers at indicated time points. Neutrophils are labeled with an i.v. injection of anti-Ly6G antibody (green), and apoptotic cells are labeled with Annexin V (red). A total of 13 apoptotic cells with burrowed neutrophils were observed in 12 WT mouse livers. Scale bar, 20 μm . (C) Electron microscopy images of apoptotic mouse hepatocytes occupied by neutrophils. The apoptotic hepatic nucleus

(AN) is evident by distorted nuclear membrane (pointed by black arrowheads). The neutrophils are indicated by white arrowheads with a characteristic multilobed nucleus. 29 apoptotic cells with burrowed neutrophils were observed. Scale bar, 5 μm . Data are representative of three independent experiments.

By using the Cellvizio System (Confocal Miniprobes), we managed to visualize the entire process of neutrophil perforocytosis in mouse livers (Fig. 2B and Video S1, S2, apoptotic hepatocytes were labeled with Annexin V and neutrophils were labeled with anti-Ly6G antibody). Consistent with observations in human samples, Annexin V positive apoptotic hepatocytes were burrowed and cleared by Ly6G-labeled neutrophils (Fig. 2B). This eating inside phagocytic process is fast and rigorous in which neutrophils were able to completely digest apoptotic hepatocytes around 4–7 minutes (Fig. 2B, and Video S1, S2, a total of 13 apoptotic hepatocytes in 12 WT mouse livers were observed). At the end of the apoptotic clearance process, neutrophils simply left the apoptotic sites and were not labeled by Annexin V, indicating neutrophils were not apoptotic.

The selectivity of neutrophil perforocytosis of effete hepatocytes

To study neutrophil-mediated phagocytosis in live cells in vitro, we induced apoptosis of isolated human primary hepatocytes or mouse liver NCTC cells with puromycin (Fig. S2A) and then applied human neutrophils or neutrophil-like, differentiated HL60 cells (Fig. 3). Isolated human primary hepatocytes formed a hepatic plate-like structure around 7 days in the culture dishes and were further confirmed with anti-albumin antibody (Fig. 3A). Human blood neutrophils (labeled with a red membrane dye, PKH-26) were able to penetrate and burrow inside the human apoptotic hepatocytes induced by puromycin (Fig. 3B, white arrowheads point to burrowing neutrophils). These apoptotic hepatocytes had markedly

decrease in size after neutrophil burrowing as compared with cells without neutrophil burrowing (Fig. 3C, D, Video S3 shows a human neutrophil first burrowed and then started to phagocytose an apoptotic hepatocyte from inside in real time). Similar results were observed with NCTC cells (labeled with a red membrane dye, PKH-26) and HL60 cells (labeled with a green membrane dye, PKH-67). Green-labeled HL60 cells showed little response to normal NCTC cells (Fig. 3E, top row). In the presence of apoptotic NCTC cells, however, green-labeled HL60 cells polarized and penetrated dead cells (Fig. 3E, white arrowheads point to burrowing neutrophils, bottom row). Next, we quantified phagocytosis by flow cytometry using a pH sensitive dye, PHrodo (with weak fluorescent at neutral pH but high fluorescent with a drop in pH to measure engulfment, see Methods). The phagocytosis of apoptotic hepatocytes by neutrophils was markedly greater than that of normal nonapoptotic hepatocytes (Fig. 3F, G). There was little cytoplasm or DNA leakage during neutrophils phagocytosing apoptotic NCTC cells as assessed by the extracellular levels of DNA, SOD and ROS (Fig. S1D-F). To address whether neutrophils also phagocytose other apoptotic cells, we determined the ability of neutrophils to phagocytose apoptotic endothelial HUVECs or epithelial HEK293 cells. We did not observe phagocytosis of these cells compared with their nonapoptotic controls (Fig. 3G), indicating the selectivity of neutrophils in phagocytosing apoptotic hepatocytes. Upon comparing the ability of HL60 cells and macrophage U937 cells to phagocytose apoptotic hepatocytes, we found that U937 cells exhibited much lower phagocytosis of apoptotic NCTC cells than did HL60 cells (Fig. S2C). Similar results were observed with human and mouse macrophages (Fig. S2B, C), consistent with the central role of neutrophils in mediating apoptotic hepatocyte clearance.




Figure 3.

Neutrophils preferentially phagocytose apoptotic hepatocytes.

(A) Fluorescent and phase contract images of isolated and cultured human primary hepatocytes stained with anti-albumin antibody (green). Scale bar, 50µm. (B) Fluorescent and phase contract images of burrowed human neutrophils (red) inside apoptotic human primary hepatocytes from A (treated with puromycin). Scale bar, 10µm. (C, D) Cell outlines (C) and cell size quantification (D) of apoptotic human primary hepatocytes occupied with burrowed neutrophils at indicated time points. (E) Fluorescence images of PKH67 (green)-labeled HL60 cells interacting with PKH26 (red)-labeled NCTC cells at the indicated time points. HL60 cells exhibited littler responses to nonapoptotic NCTC cells (top row). In contrast, HL60 cells polarized and penetrated apoptotic NCTC cells (bottom row). Scale bar, 10 µm. (F) Flow cytometry analysis of HL60 cells (labeled with CFSE, green) incubated with nonapoptotic (first column) or apoptotic hepatocytes (second column, hepatocytes were labeled with PHrodo-red dye with weak fluorescent at neutral pH but high fluorescent with a drop in pH). The population of double fluorescent HL60 cells in this subgroup (with higher PHrodo-red fluorescent indicating engulfment) was calculated, and these cells were considered phagocytosing cells. (G) Quantification of F. HL60 cells exhibited

significantly higher phagocytosis of apoptotic NCTC cells than that of nonapoptotic controls. *, $P < 0.05$ (Student's t -test). Data are representative of (A-C, E, F) or from (D, G) three independent experiments, mean and s.e.m. in D, G.

The signals that attract neutrophils to apoptotic hepatocytes

To identify the signals that attract neutrophils to apoptotic hepatocytes, we screened for cytokine and chemokine secreted by NCTC cells before and after apoptosis. Apoptotic NCTC cells showed markedly increased secretion of the cytokines IL-1 β , IL-6, IL-8, and IL-12 compared with nonapoptotic controls (Fig. 4A-D). In contrast, GM-CSF, IFN- γ , TNF- α , IL-2 and IL-10 were not significantly changed (Fig. S3A). To address the role of the upregulated cytokines, we knocked down the receptors of IL-1 β , IL-6, IL-8, or IL-12 in HL60 cells and then examined phagocytosis ability. Knockdown of the IL-1 β receptor abolished the phagocytosis of apoptotic NCTC cells (Fig. 4E), whereas knockdown of the IL-8 receptor showed a 50% reduction in phagocytosis (Fig. 4E). Thus, neutrophil chemoattractants, IL-1 β and IL-8 secreted by apoptotic NCTC cells are key signals for attracting neutrophils and inducing subsequent phagocytosis. Consistent with above observations, we found endogenous IL-1 β associated with apoptotic hepatocytes but not with normal hepatocytes in human liver samples (Fig. 4F).




Figure 4.

Apoptotic hepatocytes release signals that attract neutrophils for phagocytosis.

(A-D) Cytokine secretion in apoptotic NCTC cells. IL-1 β (A), IL-6 (B), IL-8 (C), and IL-12 (D) levels were significantly increased in apoptotic cells compared with nonapoptotic controls, **, $P < 0.01$, ***, $P < 0.001$ (Student's t -test). (E) Phagocytosis of nonapoptotic NCTC cells by HL60 cells (ctrl) and of apoptotic NCTC cells by HL60 cells treated without (NT) or with RNAi knockdown of IL-1 β , IL-8, IL-6, IL-12 receptors, or with selectin antagonist (bimosiamose), PECAM blocking antibody. *, $P < 0.05$ (Student's t -test). (F) Images of liver tissues stained with anti-IL-1 β or anti-Cas-3. Top row, IL-1 β is only associated with apoptotic hepatocytes with condensed chromatin (indicated by black arrowheads) but not with nonapoptotic hepatocytes. Bottom row, IL-1 β and Cas-3 staining in 2 μ m consecutive liver sections. Both IL-1 β and Cas-3 signals are accumulated in the same apoptotic hepatocyte (indicated by black arrowhead). Scale bar, 20 μ m. (G) Cell surface receptors in apoptotic NCTC cells compared with nonapoptotic controls. **, $P < 0.01$ (Student's t -test). Data are from (A-E, G) or representative of three independent experiments (F), mean and s.e.m. in A-E, G.

Next, we examined the cell surface molecules in hepatocytes and found that apoptotic NCTC cells had markedly increased P-selectin, E-selectin, L-selectin and PECAM as compared with non-apoptotic controls (Fig. 4G). Inhibition of selectins (with a pan-selectin antagonist,

bimosiamose) but not PECAM (with a function blocking antibody) abrogated the neutrophil-mediated clearance of apoptotic hepatocytes (Fig. 4E), indicating selectins play a critical role during this process.

Neutrophil depletion impairs the clearance of apoptotic hepatocytes

We next examined whether neutrophil depletion with either an antibody or genetic methods (see Methods) influences the clearance of apoptotic hepatocytes. Antibody or genetic depletion yielded about 90% or 70% reduction in mouse peripheral blood neutrophils, respectively (Fig. S4A, B). We analyzed liver samples from neutrophil-depleted and control mice by intravital microscopy (Fig. 2A, genetic depletion), or immunostaining (Fig. 5, antibody depletion). Similar to human liver samples, apoptotic hepatic cells in control WT mice were occupied and phagocytosed by neutrophils (Fig. 2A, 5A, B). After neutrophil depletion, the apoptotic bodies were no longer associated with neutrophils (Fig. 2A, 5A, B). However, we observed that macrophages were associated with apoptotic hepatocytes in neutrophil-depleted samples (Fig. 2A, 5A, C), suggesting a compensatory role of macrophages in phagocytosing dead hepatocytes in the absence of neutrophils. The percentage of apoptotic hepatocytes in neutrophil-depleted samples was significantly increased compared to that in controls (0.92% VS 0.2%, $p < 0.001$, Fig. 5D). Hence, neutrophil depletion impaired the prompt clearance of apoptotic cells in the mouse liver. To rule out the effects of environmental microbes after neutrophil depletion, we also treated the neutrophil-depleted mice with antibiotic (20 mg/kg ampicillin, i.p. injected daily). We obtained similar results in neutrophil-depleted mice with or without antibiotic treatment (Fig. 5A-D). Meanwhile, we observed impaired liver function in neutrophil-depleted livers (with increased aspartate aminotransferase and alanine aminotransferase activity and increased total or direct bilirubin level) compared with nontreated controls (Fig. 6A-D, antibody depletion). Depletion of neutrophils had little effect on other tissues (e.g. kidney, Fig. S4C, D).




Figure 5.

Impaired apoptotic clearance in the neutrophil-depleted liver.

(A) Fluorescent images of liver samples from control mice (top row), or neutrophil-depleted mice (antibody depletion; without or with antibiotic, second and third rows). Neutrophils are labeled with NE immunostaining (purple), macrophages are labeled with F4/80 immunostaining (red), and apoptotic cells are labeled with TUNEL staining (green). Scale bar, 15 μ m. (B-D) Cell counts of neutrophils (B) and macrophages (C) in or associated with apoptotic hepatocytes (D) in tissue samples as described in A. *, $P < 0.05$, **, $P < 0.01$, compared to control (Student's t -test). Data are representative of (A) or from three independent experiments (B-D; mean and s.e.m. in B-D).




Figure 6.

Analysis of liver function and autoantibody generation in neutrophil-depleted mice.

(A-D) Liver function analysis of aspartate aminotransferase (AST, A), alanine aminotransferase (ALT, B), direct bilirubin (DBIL, C) and total bilirubin (TBIL, D) in ctrl or neutrophil-depleted mice (antibody depletion). *, $P < 0.05$, **, $P < 0.01$ (Student's *t*-test). (E-H) Expression of autoantibodies against smooth muscle actin (anti-α-SMA, E), antinuclear antigen (anti-ANA, G), liver-kidney microsome (anti-LKM, G), and total IgG (H) in serum of control, neutrophil-depleted (with antibody), macrophage-depleted, or neutrophil-depleted plus antibiotic treated mice. Data are from three independent experiments (A-H; mean and s.e.m. in A-H).

Defective neutrophil perforocytosis in autoimmune liver disease

Autoimmune diseases are often linked to defective clearance of apoptotic cells (Nagata et al., 2010; Poon et al., 2014). We first determined whether defective neutrophil-mediated apoptotic clearance contributes to AIL disease. In the present study, we detected an increase in autoantibodies (i.e., against antinuclear antigen, smooth muscle actin, liver-kidney microsome and total IgG antibodies) in neutrophil-depleted mice compared with controls (Fig. 6E-H, antibody depletion). This increase was not affected by the treatment of antibiotic and was not observed in macrophage-depleted mice (Fig. 6E-H, macrophages were depleted with clodronate-liposome, see Methods). Hence, neutrophil depletion not only impaired apoptotic hepatocyte clearance but also led to the generation of autoantibodies, suggesting a role of defective neutrophil-mediated removal of apoptotic bodies in AIL disease.

Next, to address whether neutrophil-mediated apoptotic clearance is impaired in AIL disease, we analyzed biopsy samples from patients diagnosed with AIL disease. In contrast to the normal human controls, a total of 22 AIL disease patient samples contained apoptotic hepatocytes that were not phagocytosed or invaded by neutrophils (Fig. 7A). More apoptotic bodies were associated with macrophages, as observed in neutrophil-depleted mouse livers (Fig. 7A, Table S7). Since the blood neutrophil count in AIL disease patients is within the normal range, we surmised potential defects in the phagocytosis ability of neutrophils from AIL disease patients. We observed markedly decreased phagocytosis of apoptotic NCTC cells by neutrophils from AIL disease patients compared to normal controls (Fig. 7B). Normal human neutrophils burrowed into apoptotic NCTC cells and demonstrated perforocytosis, while AIL disease neutrophils exhibited little response towards apoptotic NCTC cells (Fig.

7C). We further screened differential gene expression between normal human neutrophils and AIL neutrophils. We noted that the expression of IL-1 β receptor, IL1R1 and selectin binding protein, P-selectin glycoprotein ligand 1 (PSGL-1) was markedly decreased in AIL neutrophils as compared with normal human neutrophils (Fig. 7D), which further confirmed the critical roles of IL-1 β and selectins in neutrophil-mediated apoptotic clearance. The above data prove defective neutrophil-mediated apoptotic clearance in human AIL disease samples.




Figure 7.

Defective neutrophil perforocytosis in human autoimmune liver disease.

(A) Images of CD68 staining from normal or AIL disease human liver samples. Green arrowheads indicate macrophages, black arrowheads indicate apoptotic bodies and white arrowheads point to neutrophils. (B, C) Quantification (B) and images (C) of *in vitro* phagocytosis of apoptotic NCTC cells by neutrophils isolated from normal or AIL disease human patients. White arrowheads point to burrowed neutrophils. Scale bar, 10 μ m. *, $P < 0.05$ (Student's *t*-test). (D) Analysis of CD34, PSGL-1, IL1R1 and CXCR2 expression in neutrophils from normal human and AIL patients by microarray. Data are representative of (A, C) or from three independent experiments (B, D; mean and s.e.m. in B, D).

Discussion

Our finding of neutrophil burrowing into and clearing of apoptotic hepatocytes under physiological conditions reveals a fundamental mechanism for the removal of effete hepatocytes, helps to solve some mysteries surrounding the apoptotic clearance process and raises several important questions.

Neutrophils as the new scavengers for apoptotic clearance

Although billions of cells undergo apoptosis daily in our bodies, apoptotic cells are rarely observed in tissues under steady state due to the high efficiency of apoptotic removal by both professional and neighboring nonprofessional phagocytes (Poon et al., 2014). In general, professional phagocytes have much higher phagocytic efficiency and capacity than nonprofessional phagocytes, but they are greatly outnumbered by other cell types in tissues (Elliott and Ravichandran, 2016). Therefore, whether there are unidentified scavengers and/or mechanisms for the prompt removal of dead cells in distinct organs remains unclear. Compared to macrophages or their precursors, monocytes, which make up less than 10% of the total white blood cell population, neutrophils are the most abundant white blood cells (up to 70%) in humans, and thus ideally suited for the swift clearance of apoptotic hepatic cells. As macrophages were induced to phagocytose apoptotic hepatocytes following the

depletion of circulating neutrophils, our data show compensatory activation of other phagocytes under these conditions. The basis for this switch, however, is not clear.

Another important question is whether other tissues utilize a similar mechanism. As neutrophils do not phagocytose apoptotic epithelial and endothelial cells, it is possible that distinct signals are specifically released by different types of apoptotic cells. It remains largely unknown whether one or more signals are dominant in each specific tissue and how these signals are orchestrated into a complicated network to swiftly clear dying cells while avoiding tissue inflammation.

Neutrophil influx without causing tissue damage

Although apoptotic clearance has been considered an immunologically silent process that does not lead to the influx of inflammatory cells or exposure of self-antigens(Poon et al., 2014; Savill et al., 2002), it has also been suggested that this process is not completely immunologically silent (Green et al., 2009). Because neutrophils are proinflammatory cells, they have long been thought to be excluded from apoptotic sites. We observed neutrophil influx into apoptotic hepatic cells (up to 22 neutrophils in one apoptotic hepatocyte), and there was little detectable tissue injury or other inflammatory cells. One reason could be that the neutrophil attraction signals released by apoptotic hepatocytes (e.g., IL-1 β and IL-8) are sufficient for attracting neutrophils towards apoptotic cells and inducing subsequent perforocytosis, but do not elicit neutrophil inflammatory functions such as oxidative burst. Therefore, neutrophil recruitment into tissues does not always induce inflammation or tissue damage. In consistent with our observations, neutrophils can express both pro- and anti-inflammatory cytokines(Gideon et al., 2019; Mortaz et al., 2018). In contrast, neutrophil depletion caused defective removal of apoptotic bodies and induced autoantibody generation; thus neutrophils play a vital role in the genesis of AIL disease. Considering that billions of neutrophils patrol tissues under physiological circumstances without causing inflammation(Nicolas-Avila et al., 2017), we concluded that these cells not only provide immune surveillance against infection but also contribute to internal tissue homeostasis, as we report in this study.

Neutrophil burrowing to maintain tissue integrity

One intriguing question regarding apoptotic clearance is how the tissue maintains integrity while dead cells are being continually removed(Poon et al., 2014). Apoptotic cell detachment from the extracellular matrix by caspase-mediated cleavage or extrusion into the organ lumen has been proposed as the mechanism for maintaining the epithelial barrier during the clearance of dying epithelial cells(Brancolini et al., 1997; Rosenblatt et al., 2001). Our finding that burrowed neutrophils phagocytose apoptotic hepatocytes from the intracellular space provides a novel mechanism for maintaining liver integrity. The neutrophils that phagocytose cells from the inside are efficient at disposing of apoptotic bodies without extruding the cytoplasm and may help to prevent the leakage of toxic bile acids or the release of danger signals that cause tissue damage.

Neutrophil burrowing into other types of cells has also been reported previously in processes not related to apoptotic clearance(Overholtzer and Brugge, 2008). For example, neutrophils can bore into endothelial cells or megakaryocytes to temporarily form so-called cell-in-cell (or emperipolesis, entosis) structures(Overholtzer and Brugge, 2008). The apparent difference is that both the endothelial cells and megakaryocytes entered by neutrophils are viable and nonapoptotic. The purpose of forming cell-in-cell structures with neutrophils and endothelial cells or megakaryocytes is to obtain a passage out of blood

vessels or bone marrow, respectively (Overholtzer and Brugge, 2008). It will be interesting to examine whether neutrophils utilize similar invasion mechanisms during these processes.

The important apoptotic clearance function of neutrophils described in this study adds to the repertoire of other known functions of neutrophils (Amulic et al., 2012; Kolaczowska and Kubes, 2013; Wang et al., 2017). Since the failure to clear dead cells is linked to inflammatory and autoimmune diseases, the critical role of neutrophil-mediated apoptotic clearance may have implications for the pathogenesis and treatment of these diseases.

Methods

Cell culture, transfection, and isolation of human and mouse neutrophils

NCTC, HEK-293, U937 cells and HUVECs were cultured in DMEM supplemented with 10% fetal bovine serum and HL60 cells were cultured in RPMI 1640 with 10% fetal bovine serum. For experiments, HL60 cells were differentiated by adding 1.3% DMSO into the medium for 7 days (Xu et al., 2003). To establish the stable RNAi cell lines, we transfected shRNA into HEK293T cells. After virus packaging in HEK293T cells, HL60 cells were infected, screened by puromycin and sorted by flow cytometry (Liu et al., 2012).

For mouse polymorphonuclear neutrophil (PMN) isolation from bone marrow (Wang et al., 2016), mice were euthanized, and the femurs and tibias were removed and flushed by a 27-gauge needle with a 10-mL syringe filled with calcium and magnesium-free Hank's balanced salt solution (HBSS) plus 0.1% BSA. Cells were then centrifuged and resuspended in HBSS. After being filtered through a 40- μ m strainer, cells in 3 mL of HBSS were loaded onto a pre-prepared gradient solution (3 mL of NycoPrep on top and 3 mL of 72% Percoll on bottom). The samples were centrifuged at 2,400 rpm at room temperature for 20 min with the brake off. The middle layer was collected and washed once in HBSS. Sterilized distilled water (9 mL) was added for 22 s and thereafter 1 mL of 10 $\times$ PBS was added to remove red blood cells. Finally, the cells were collected and resuspended in HBSS or medium (neutrophil purity >90%). All animal experiments were performed under the protocol approved by The Institutional Animal Care and Use Committee of the State Key Laboratory of Biotherapy, West China Hospital, Sichuan University.

For the isolation of human neutrophils (Liu et al., 2015), blood was collected from healthy human donors or those with AIL disease. Erythrocytes were removed using dextran sedimentation (4.5% dextran) followed by hypotonic lysis using distilled water. Neutrophils were isolated from the resulting cell suspension using discontinuous Percoll gradient centrifugation. This procedure yielded >95% neutrophil purity and >95% viability as assessed by flow cytometry.

For the isolation of human primary hepatocytes, liver tissues from a fresh liver hemangioma surgery were suspended in DPBS and washed for 5-10 times. Surrounding connective tissues and adipose tissues were removed with a surgery knife and remaining liver tissues were further sliced into small fragments (around 1 mm²), then washed with DPBS for three times and centrifuged at 800 rpm for 3 min. Tissue samples were resuspended in 5 times volume of collagenase IV medium and incubated at 37°C for 30 min. After collagenase digestion, samples were centrifuged at 1000 rpm for 5 min, and resuspended in RPMI 1640 with antibiotics to precipitate cells, then centrifuged at 1500 rpm for 5 min. Isolated cells were cultured in RPMI 1640 medium with insulin (1: 1000), glucagon (1:1000), glucocorticoid (1:1000), HGF (10 ng/mL), EGF (10 ng/mL). Studies using human primary cells and samples

were approved by the Institutional Review Board of West China Hospital, Sichuan University and Fudan University.

Intravital microscopy, immunostaining of liver sections, 3D image reconstruction and electron microscopy

For intravital microscopy (Marques et al., 2015), freshly prepared anti-Ly6G 1A8 FITC, anti-F4/80 PE and Annexin V pacific blue were injected intravenously into mice 1 h before imaging. After anesthesia, mice were attached to a surgical stage and subjected to midline laparotomy, and all vessels in the skin were cauterized. The falciform ligament between the liver and diaphragm was subsequently cut by holding the xiphoid with a knot made of suture thread. Mice were then moved to the imaging stage, and the right lobe was exposed. The ventral side of the mouse was placed on a Plexiglas stage attached to the microscope stage of a single photon microscope (NIKON A1RMP) or Cellvizio System for the imaging of apoptotic cells (labeled with Annexin V), Kupffer cells (anti-F4/80) and neutrophils (anti-Ly-6G 1A8). Microscope acquisition settings were as follows: 405-nm laser power, 15.4%, PMT high voltage (HV), 115; 488-nm laser power, 27.9%, PMT HV, 76; 543-nm laser power, 41.4%, PMT HV, 108; pinhole size, 40 μ m; 20 $\times$ objective.

Neutrophils burrowing inside apoptotic hepatocytes were analyzed by IMARIS software (adopted from the analysis of vesicles in cells). Annexin V blue was set as the source channel to detect apoptotic hepatocyte boundary, and anti-Ly6G green was set as the source channel to detect neutrophils inside blue cells. Only neutrophils inside hepatic cells can be detected. The cells objects were calculated and displayed. The cell positions of both neutrophils and hepatocytes, and the distances from neutrophils to the hepatocyte border were recorded.

For immunostaining of liver sections, fresh mouse liver samples were fixed in freshly prepared 4% PFA and cryoprotected in 30% sucrose/PBS solution. Thick cryosections (45 μ m) were cut and washed three times with PBS. Tissue sections were incubated in blocking buffer (10% donkey serum and 0.2% Triton X-100 in PBS) for 1 h at room temperature and then in a primary antibody solution (5% donkey serum and 0.2% Triton X-100 in PBS; rat anti-Ly-6G antibody, 1:500; goat anti-E-cadherin antibody, 1:200) overnight at 4°C. Next, the sections were washed three times in PBS, incubated for 1 h in the appropriate secondary antibody solution containing DAPI for 2 h at 4°C, and then washed three times in PBS. The immunostained sections were mounted onto slides with an aqueous mounting medium. All tissue sections were scanned by a Zeiss 710 confocal microscope. The Z-stack scanned images were edited by Zen 2009 software.

To observe the liver ultrastructure with TEM, mouse livers were removed, fixed (2.5% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.4), and cut into multiple 1 mm $\times$ 1 mm strips in fluid; several strips were simultaneously cut into 1 mm $\times$ 1 mm $\times$ 1 mm blocks for TEM. Blocks were transferred to wash buffer (0.1 M cacodylate buffer, pH 7.4) to remove glutaraldehyde and postfixes for 1 h in 1% osmium tetroxide. The specimens were dehydrated in a graded series of ethanol (75%, 85%, 95%, and 100%) and embedded in Epon812. Ultrathin sections were obtained and then observed with a HITACHI H600 TEM. Apoptotic liver cells were determined based on changes in nuclear morphology (chromatin condensation and fragmentation). Neutrophils were determined by polymorphonuclear morphology.

Neutrophil depletion and macrophage depletion in mice

Two methods were used to deplete neutrophils in mice. For antibody-based depletion, wild-type mice were injected i.p. with anti-mouse Ly6G clone 1A8 (Bio X Cell) supernatant

containing 0.5 mg protein every 48 h for 1 month. Control animals were injected with isotype-matched normal control antibody. For genetic neutrophil ablation, diphtheria toxin receptor (DTR)-expressing Rosa26iDTR mice (Buch et al., 2005; Saito et al., 2001) (The Jackson Laboratory) were bred onto the *Mrp8*-Cre genetic background (The Jackson Laboratory). Cre expression in MRP-expressing cells, mainly granulocytes, will activate DTR expression in only MRP-expressing cells. At the age of 3 months, the Rosa26iDTR/*Mrp8*-Cre mice and WT control littermates (Rosa26iDTR) were treated with diphtheria toxin (DT, 20 ng/g BW, i.p., daily for 3 days) to induce *Mrp8*⁺ cell depletion.

Clodronate-liposome (YEASEN, 40337ES05) was used to deplete mouse macrophage. WT mice were injected i.v. with 100 μ l of clodronate-liposome per 10 g of body weight. 24 h later, mice were injected i.v. with 100 μ l of clodronate-liposome per 10 g of body weight again. The operation was repeated every 72 h after the second injection for 1 month. Neutrophil and macrophage depletion were confirmed by neutrophil and macrophage count in all experimental animals.

Apoptosis assay

Apoptosis of NCTC cells was induced by puromycin (Chow et al., 1995). A total of 1×10^6 NCTC cells were plated in a 35 mm glass-bottom dish (MatTek) for 10 h, and then treated with puromycin (2.5 μ g/ml) for 12 h. Apoptosis was assayed with an Alexa Fluor 488 Annexin V/Dead Cell Apoptosis Kit (Thermo Fisher) and analyzed by flow cytometry to measure the fluorescence emission at 530 nm and 575 nm (or equivalent) using 488 nm excitation.

Fluorescent labeling of cells

The fluorescent membrane stain PKH-26 (red) (Sigma-Aldrich) was used to label nonapoptotic and apoptotic NCTC and HEK293 cells and HUVECs. PKH-67 (green) was used to label HL60 cells and human neutrophils. Cells (2×10^6) were washed in serum-free DMEM or RPMI 1640 medium, and the cell pellet was resuspended in PKH-26 or PKH-67 staining solution and incubated for 5 min at room temperature. After stopping the labeling reaction by adding an equal volume of serum, the cells were washed three times with complete medium.

In vitro phagocytosis assay

Phagocytosis was quantified by flow cytometry using a pH sensitive dye, PHrodo-red (weak fluorescent at neutral pH but high fluorescent with a drop in pH) to measure engulfment. 1×10^6 human neutrophils or differentiated HL60 cells were labelled with CFSE green and then incubated with apoptotic or nonapoptotic NCTC cells (labeled with PHrodo-red, 10 μ L PHrodo-red dye mixed with 100 μ L PowerLoad concentrate, and diluted into 10 mL of HBSS, PH 7.4 for 1 h). The cells were then washed with HBSS and analyzed by flow cytometry. The forward scatter (FCS) reflects cell size, whereas the sideward scatter (SSC) reveals the degree of granularity of the cell. Apoptotic NCTC cells with burrowed neutrophils increased in cell size and granularity, causing increased diffraction of the laser beam, revealed by spreading of the dots to the upper right corner. The population of double fluorescent HL60 cells in this subgroup was calculated, and these cells were considered phagocytosing cells.

Cytokine screening and microarray

Cytokines secreted from nonapoptotic and apoptotic NCTC cells were screened with the Human Inflammatory Cytokines Multi-Analyte ELISArray Kit (QIAGEN). Nonapoptotic and

apoptotic NCTC cell supernatants were prepared in replicate serial dilutions of the Antigen Standard. Cytokine secretion was further determined with the ELISA kit.

For microarray, total RNA from normal human or AIL patient neutrophils was isolated using RNeasy Total RNA Isolation Kit (Qiagen, GmbH, Germany)/ TRIzol reagent (Life technologies, Carlsbad, CA, US) according to the manufacturer's instructions, and purified by using RNeasy Mini Kit (Qiagen, GmbH, Germany). Total RNA was checked for a RIN number to inspect RNA integration by an Agilent Bioanalyzer 2100 (Agilent technologies, Santa Clara, CA, US). RNA samples of each group were then used to generate biotinylated cRNA targets for the Agilent SurePrint Gene Expression Microarray. The biotinylated cRNA targets were then hybridized with the slides. After hybridization, slides were scanned on the Agilent Microarray Scanner (Agilent technologies, Santa Clara, CA, US). Data were extracted with Feature Extraction software 10.7 (Agilent technologies, Santa Clara, CA, US). Raw data were normalized by Quantile algorithm, R package "limma". Heatmap plots were done by a R package "pheatmap" of the target genes. GO pathway enrichment analysis was done use Fisher's exact test by R package "clusterProfiler" of the target genes. GO categories\Pathway with Fisher's exact test P values < 0.05 were selected.

Statistical analysis

All experiments were performed at least three times. Representative data are shown in the paper. An independent experiments two-tailed Student's *t*-test was used as the statistical assay for comparisons. Significant differences between samples were indicated by $P < 0.05$.

Author contributions

Conceptualization: LC, HS, JZ, ABM, JX

Methodology: LC, HS, JZ, JL, XF, WL, YQ, YT, SP, LY, LZ, LM, LL, XH, YJ, YH, YZ, XZ, YZ, YW

Investigation: LC, HS, JZ, JL, XF, WL, YQ, YT, SP, LY, LZ, LM, LL, XH, YJ, YH, YZ, XZ, YZ, JX

Visualization: LC, HS, JZ, JL, XF, WL, YQ, YT, SP, LY, LZ, LM, LL, XH, YJ, YH, YZ, XZ, YZ

Funding acquisition: LZ, YZ, YZ, YW, ABM and JX Project administration: LC, HS, JZ, JX

Supervision: LC, HS, JX Writing: LC, HS, ABM, JX

Acknowledgements

We thank Yu Fu and Ruixue Li (both from Guangzhou Regenerative Medicine and Health Guangdong Laboratory Imaging Center) for helpful imaging service. This work was supported in part by NSFC grant 32170750 and SSTEP grant 19YYJC2572.

Declaration of interests

The authors declare no competing interests.





Fig. S1

Little leakage during phagocytosis.

(A) Reconstructed 3D images of neutrophils (NE staining, red) burrowed into apoptotic hepatocytes (TUNEL, green). (B) Few CD11b⁺ or CD45RA⁺ cells around the phagocytosis site. Bar, 20 μ m. (C) Fluorescent images of human liver tissue with neutrophils scavenging apoptotic liver cells. TUNEL staining (green) shows apoptotic nucleus (white arrowheads) and apoptotic nuclear debris (white arrows). The apoptotic debris is phagocytosed by burrowed neutrophils (labeled with NE-red, analyzed and projected with IMARIS software). The burrowing neutrophils have intact nucleus (shown by DAPI staining) with negative TUNEL signals. (D-F) Extracellular levels of DNA (D), SOD (E), and ROS (F) are not significantly changed during phagocytosis of apoptotic NCTC cells by HL60 cells as compared with that of nonapoptotic control NCTC cells. Cell lysate was used as positive control. Data are representative of (A-C) or

from (D-F) 3 independent experiments (mean and s.e.m. in D-F).



C




Fig. S2.

Neutrophils selectively phagocytose apoptotic liver cells.

(A) Flow analysis of NCTC cells treated with puromycin at indicated time points in the presence of PI-red and Annexin-V-FITC. A lag time seen between PtdSer⁺ and PI⁺ is a characteristic feature of apoptotic cells, the cell population moves first from Q1 (AnnexinV⁺/PI⁻) to Q2 (AnnexinV⁺/PI⁺), then moves to Q3 (AnnexinV⁺/PI⁺). In contrast, necrotic cells immediately move from Q1 to Q3 without passing through the Q2 stage. (B) PKH67 (green)-labeled human neutrophils or macrophages interacting with PKH26 (red)-labeled NCTC, HEK293 cells and HUVECs at the indicated times. Neutrophils only burrowed inside apoptotic NCTC cells (second row, white arrowheads point to the burrowing neutrophil). Bar, 10 μ m. (C) Phagocytosis of HL60, U937, human/mouse neutrophils (PMN), or macrophages (M ϕ), towards nonapoptotic and apoptotic NCTC cells. *, $p < 0.05$, **, $p < 0.01$, (Student's t -test). Data are representative of (A, B) or from (C) three independent experiments (mean and s.e.m. in C).




Fig. S3.

Cytokines and cell surface receptors for phagocytosis.

(A) Cytokines secreted by normal nonapoptotic and apoptotic NCTC cells are not significantly changed, including IL-1 α , IL-2, IL-4, IL-10, TNF- α , IFN- γ , and GM-CSF. (B) Immunoblots of target proteins (IL-1b, IL-8, IL-6, IL-12 receptors) in non-treated (Ctrl) and RNAi-treated HL60 cells. RNAi knockdown efficiency is ranged from 70-90%. Data are from (A) or representative of (B) three independent experiments (mean and s.e.m. in A).




Fig. S4.

Neutrophil depletion impairs liver functions.

(A,B) Blood neutrophil count after antibody depletion at times indicated (A) or in WT and MRP8cre/DTR mice (B). (C,D) Analysis of creatinine (C) and blood urea nitrogen (D) in WT, MRP8Cre/DTR, and neutrophil depleted (by antibody) mice. Data are from 3 independent experiments (mean and s.e.m. in A-D).

Table 1.

The cell numbers of burrowed neutrophils (PMNs) in apoptotic hepatocytes from human liver tissues.

2D images were taken from regular tissue sections (5 μm thick), and 3D images were taken from 45 μm thick tissue sections and reconstructed by confocal microscope.

Perferocytosis stage		I	II	III
2D images	Apoptotic hepatocytes (total 241)	37	130	74
	Burrowing PMNs (total 1716)	312	931	473
	PMNs per apoptotic hepatocyte (mean $\pm$ sem)	8.4 $\pm$ 4.2	7.2 $\pm$ 3.4	6.4 $\pm$ 3.4
3D images	Apoptotic hepatocytes (total 60)	28	18	14
	Burrowing PMNs (total 820)	406	348	66
	PMNs per apoptotic hepatocyte (mean $\pm$ sem)	14.5 $\pm$ 10.0	19.3 $\pm$ 15.5	4.7 $\pm$ 2.4

Table 2.

Analysis of neutrophils burrowing inside apoptotic hepatocytes from human liver samples.

Reconstructed 3D images of neutrophils inside apoptotic hepatocytes (as shown in Figure 1D) are remodeled and analyzed with IMARIS software: anti-E-Cadherin blue staining is set as the

source channel to detect hepatic cell boundary, and anti-NE red staining is set as the source channel to detect neutrophils inside blue cells. Only neutrophils inside hepatic cells can be detected. The cell positions of both neutrophils and hepatocytes, and the distances from neutrophils to the hepatocyte border are recorded below (a total of 8 apoptotic hepatocytes are analyzed).

Apoptotic hepatocytes	Hepatocyte position X,Y,Z (μm)	PMNs inside Hepatocytes	PMN position X,Y,Z (μm)	PMN distance to hepatocyte border (μm)
#1	10.8, 10.6, 4.6	#1	13.3, 3.7, 1.9	0
		#2	10.1, 4.1, 3.9	1.9
		#3	18.6, 9.8, 3.1	1.1
		#4	8.5, 12.2, 3.5	1.5
		#5	3.2, 22.6, 7.1	1.2
		#6	8.1, 22.7, 7.7	3.8
#2	31.4, 14.7, 7.5	#1	28.1, 21.0, 0.8	0
		#2	31.2, 18.9, 3.5	1.5
		#3	30.4, 5.1, 5.1	3.1
#3	38.9, 15.1, 3.0	#1	38.9, 15.2, 3.5	1.5
#4	28.9, 24.1, 7.9	#1	34.1, 19.4, 11.3	10.1
#5	16.7, 35.4, 14.2	#1	12.6, 33.2, 7.2	5.9
		#2	15.6, 35.5, 5.6	4.4
		#3	18.9, 37.9, 5.1	3.9
		#4	13.1, 36.5, 9.9	8.6
#6	25.5, 22.8, 3.0	#1	27.5, 21.9, 2.4	1.1
		#2	25.1, 23.2, 3.4	2.2
#7	15.3, 44.1, 9.4	#1	16.0, 43.0, 8.3	3.5
		#2	13.8, 45.4, 11.0	1.2
#8	24.0, 35.4, 8.8	#1	24.0, 35.7, 9.4	8.1

Table 3.

CD68⁺ cells' distribution in human liver tissues with or without apoptotic hepatocytes (perferocytosed by PMNs).

5 μm thick tissue sections were used to stain and count CD68⁺ cells.

	Total area (μm ²)	CD68 ⁺ cells	Apoptotic hepatocytes	CD68 ⁺ cells/10 ⁴ μm ² (mean ± sem)
Liver tissue with PMN perferocytosis	579175	280	40	5.8 ± 2.2*
Liver tissue without PMN perferocytosis	403025	267	0	6.7 ± 2.8*

*, $P = 0.0962$ (student's t test), CD68⁺ cell numbers are not different in liver tissues with or without PMN perferocytosis.

Table 4.

The distribution of CD11b⁺ cells in human liver tissues with or without apoptotic hepatocytes (perferocytosed by PMNs). 45 μm thick tissue sections were used to stain and count CD11b⁺ cells.

	Total area (μm^2)	CD11b ⁺ cells	Apoptotic hepatocytes	CD11b ⁺ cells/ $10^4 \mu\text{m}^2$ (mean $\pm$ sem)
Liver tissue with PMN perferocytosis	138905	12	17	1.1 $\pm$ 1.4*
Liver tissue without PMN perferocytosis	228508	72	0	3.1 $\pm$ 3.0*

*, $P = 0.0013$ (student's t test), CD11b⁺ cell numbers are lower in liver tissues with PMN perferocytosis compared with that without PMN perferocytosis.

Table 5.

The distribution of CD45RA⁺ cells in human liver tissues with or without apoptotic hepatocytes (perferocytosed by PMNs).

45 μm thick tissue sections were used to stain and count CD45RA⁺ cells.

	Total area (μm^2)	CD45RA ⁺ cells	Apoptotic hepatocytes	CD45RA ⁺ cells/ $10^4 \mu\text{m}^2$ (mean $\pm$ sem)
Liver tissue with PMN perferocytosis	57367	7	10	1.4 $\pm$ 1.2*
Liver tissue without PMN perferocytosis	280880	54	0	1.9 $\pm$ 1.9*

*, $P = 0.309$ (student's t test), CD45RA⁺ cell numbers are not different in liver tissues with or without PMN perferocytosis.

Table 6.

Analysis of neutrophils burrowing inside apoptotic hepatocytes from mouse liver samples.

Intravital images of neutrophils inside apoptotic hepatocytes (as shown in [Figure 2A](#)) are re-modeled and analyzed with IMARIS software: Annexin V blue staining is set as the source channel to detect hepatic cell boundary, and Ly-6G green staining is set as the source channel to detect neutrophils inside blue cells. Only neutrophils inside hepatic cells are detected and calculated. The cell positions of both neutrophils and hepatocytes, and the distances from neutrophils to the hepatocyte border are recorded below (a total of 10 apoptotic hepatocytes are analyzed).

Apoptotic hepatocytes	Hepatocyte position X,Y,Z (μm)	PMNs inside hepatocytes	PMN position X,Y,Z (μm)	PMN distance to hepatocyte border (μm)
#1	33772.8, 25874.0, 3077.5	#1	33771.5, 25882.7, 3085.1	42.2
		#2	33768.9, 25877.5, 3083.2	44.1
		#3	33780.6, 25864.5, 3071.5	55.8
#2	13583.7, 17933.4, 3955.6	#1	13586.7, 17935.1, 3956.1	6.16
		#2	13581.7, 17932.3, 3955.5	6.76
#3	13550.8, 17833.9, 3950.7	#1	13551.5, 17833.9, 3950.7	11.6
#4	13493.1, 18060.3, 3947.0	#1	13491.3, 18060.9, 3946.8	15.4
#5	13539.1, 17922.4, 3946.5	#1	13539.3, 17922.8, 3946.3	16.0
#6	13504.1, 17853.5, 3946.2	#1	13504.1, 17853.3, 3946.1	16.2
#7	13440.5, 18190.0, 3941.9	#1	13440.9, 18189.2, 3941.4	20.9
#8	13368.4, 18069.1, 3938.1	#1	13369.0, 18068.6, 3938.0	24.3
#9	13034.7, 18056.6, 3909.4	#1	13034.6, 18054.8, 3909.7	24.8
		#2	13034.9, 18060.0, 3908.2	24.9
#10	13019.5, 18118.2, 3902.9	#1	13019.5, 18118.7, 3902.2	9.7

Table 7.

Apoptotic hepatocytes burrowed with PMNs or monocytes in normal human or AIL liver samples.

45 μ m thick tissue sections were used to stain and count apoptotic hepatocytes.

	Total liver sample numbers	Apoptotic hepatocytes	Apoptotic hepatocytes burrowed by PMNs	Apoptotic hepatocytes burrowed by monocytes
Normal liver samples	32	227	227	0
AIL liver samples	22	110	8	40

Video S1, S2: Mouse neutrophils burrowed and digested apoptotic hepatocytes in the mouse liver. Neutrophils were labeled with an i.v. injection of anti-Ly6G antibody (green) and apoptotic cells were labeled with Annexin V (red). Time-lapse images were acquired by the Cellvizio system. The cytoplasm of apoptotic hepatocyte with neutrophil burrowing quickly dwindled and finally disappeared. 2 out of 13 apoptotic hepatocytes are shown in video S1 and S2.

Video S3: Human neutrophils burrowed inside apoptotic hepatocytes. A human neutrophil (green, pointed by white arrow) was burrowing into an apoptotic hepatocyte (red). Time-lapse video was recorded for 60 minutes and followed by reconstructed 3D images at 20 and 60 min. At 20 min, the neutrophil partially entered the hepatic cell and did not phagocytose (neutrophil was extracted and analyzed by IMARIS software and no red dye inside the neutrophil). At 60 min, the neutrophil completely burrowed inside the hepatocyte and started to phagocytose hepatocyte (with uptake of red dye).

References

- Amulic B. , Cazalet C. , Hayes G.L. , Metzler K.D. , Zychlinsky A. (2012) **Neutrophil function: from mechanisms to disease** *Annual review of immunology* **30**:459–489
- Bournazou I. , Pound J.D. , Duffin R. , Bournazos S. , Melville L.A. , Brown S.B. , Rossi A.G. , Gregory C.D. (2009) **Apoptotic human cells inhibit migration of granulocytes via release of lactoferrin** *The Journal of clinical investigation* **119**:20–32
- Brancolini C. , Lazarevic D. , Rodriguez J. , Schneider C. (1997) **Dismantling cell-cell contacts during apoptosis is coupled to a caspase-dependent proteolytic cleavage of beta-catenin** *The Journal of cell biology* **139**:759–771
- Buch T. , Heppner F.L. , Tertilt C. , Heinen T.J. , Kremer M. , Wunderlich F.T. , Jung S. , Waisman A. (2005) **A Cre-inducible diphtheria toxin receptor mediates cell lineage ablation after toxin administration** *Nature methods* **2**:419–426

Canbay A. , Feldstein A.E. , Higuchi H. , Werneburg N. , Grambihler A. , Bronk S.F. , Gores G.J. (2003) **Kupffer cell engulfment of apoptotic bodies stimulates death ligand and cytokine expression** *Hepatology* **38**:1188–1198

Chow S.C. , Peters I. , Orrenius S. (1995) **Reevaluation of the role of de novo protein synthesis in rat thymocyte apoptosis** *Experimental cell research* **216**:149–159

Eipel C. , Hirschmann M. , Abshagen K. , Menger M.D. , Vollmar B. (2007) **Local interaction of apoptotic hepatocytes and Kupffer cells in a rat model of systemic endotoxemia** *Hepatology research: the official journal of the Japan Society of Hepatology* **37**:863–871

Elliott M.R. , Ravichandran K.S. (2016) **The Dynamics of Apoptotic Cell Clearance** *Developmental cell* **38**:147–160

Fond A.M. , Ravichandran K.S. (2016) **Clearance of Dying Cells by Phagocytes: Mechanisms and Implications for Disease Pathogenesis** *Advances in experimental medicine and biology* **930**:25–49

Franz S. , Gaip U.S. , Munoz L.E. , Sheriff A. , Beer A. , Kalden J.R. , Herrmann M. (2006) **Apoptosis and autoimmunity: when apoptotic cells break their silence** *Current rheumatology reports* **8**:245–247

Gideon H.P. , Phuah J. , Junecko B.A. , Mattila J.T. (2019) **Neutrophils express pro- and anti-inflammatory cytokines in granulomas from Mycobacterium tuberculosis-infected cynomolgus macaques** *Mucosal immunology* **12**:1370–1381

Green D.R. , Ferguson T. , Zitvogel L. , Kroemer G. (2009) **Immunogenic and tolerogenic cell death** *Nature reviews. Immunology* **9**:353–363

Harsunen M.H. , Puff R. , D'Orlando O. , Giannopoulou E. , Lachmann L. , Beyerlein A. , von Meyer A. , Ziegler A.G. (2013) **Reduced blood leukocyte and neutrophil numbers in the pathogenesis of type 1 diabetes** *Hormone and metabolic research = Hormon-und Stoffwechselforschung = Hormones et metabolisme* **45**:467–470

Hochreiter-Hufford A. , Ravichandran K.S. (2013) **Clearing the dead: apoptotic cell sensing, recognition, engulfment, and digestion** *Cold Spring Harbor perspectives in biology* **5**

Kerr J.F. , Wyllie A.H. , Currie A.R. (1972) **Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics** *British journal of cancer* **26**:239–257

Kolaczowska E. , Kubes P. (2013) **Neutrophil recruitment and function in health and inflammation** *Nature reviews. Immunology* **13**:159–175

Lauber K. , Blumenthal S.G. , Waibel M. , Wesselborg S. (2004) **Clearance of apoptotic cells: getting rid of the corpses** *Molecular cell* **14**:277–287

Liu X. , Ma B. , Malik A.B. , Tang H. , Yang T. , Sun B. , Wang G. , Minshall R.D. , Li Y. , Zhao Y. , Ye R.D. , Xu J. (2012) **Bidirectional regulation of neutrophil migration by mitogen-activated protein kinases** *Nature immunology* **13**:457–464

Liu X. , Yang T. , Suzuki K. , Tsukita S. , Ishii M. , Zhou S. , Wang G. , Cao L. , Qian F. , Taylor S. , Oh M.J. , Levitan I. , Ye R.D. , Carnegie G.K. , Zhao Y. , Malik A.B. , Xu J. (2015) **Moesin and myosin phosphatase confine neutrophil orientation in a chemotactic gradient** *The Journal of experimental medicine* **212**:267–280

Marques P.E. , Antunes M.M. , David B.A. , Pereira R.V. , Teixeira M.M. , Menezes G.B. (2015) **Imaging liver biology in vivo using conventional confocal microscopy** *Nature protocols* **10**:258–268

Mortaz E. , Alipoor S.D. , Adcock I.M. , Mumby S. , Koenderman L. (2018) **Update on Neutrophil Function in Severe Inflammation** *Frontiers in immunology* **9**

Nagata S. , Hanayama R. , Kawane K. (2010) **Autoimmunity and the clearance of dead cells** *Cell* **140**:619–630

Nicolas-Avila J.A. , Adrover J.M. , Hidalgo A. (2017) **Neutrophils in Homeostasis, Immunity, and Cancer** *Immunity* **46**:15–28

Overholtzer M. , Brugge J.S. (2008) **The cell biology of cell-in-cell structures** *Nature reviews. Molecular cell biology* **9**:796–809

Poon I.K. , Lucas C.D. , Rossi A.G. , Ravichandran K.S. (2014) **Apoptotic cell clearance: basic biology and therapeutic potential** *Nature reviews. Immunology* **14**:166–180

Rosenblatt J. , Raff M.C. , Cramer L.P. (2001) **An epithelial cell destined for apoptosis signals its neighbors to extrude it by an actin- and myosin-dependent mechanism** *Current biology: CB* **11**:1847–1857

Saito M. , Iwawaki T. , Taya C. , Yonekawa H. , Noda M. , Inui Y. , Mekada E. , Kimata Y. , Tsuru A. , Kohno K. (2001) **Diphtheria toxin receptor-mediated conditional and targeted cell ablation in transgenic mice** *Nature biotechnology* **19**:746–750

Savill J. , Dransfield I. , Gregory C. , Haslett C. (2002) **A blast from the past: clearance of apoptotic cells regulates immune responses** *Nature reviews. Immunology* **2**:965–975

Valle A. , Giamporcaro G.M. , Scavini M. , Stabilini A. , Grogan P. , Bianconi E. , Sebastiani G. , Masini M. , Maugeri N. , Porretti L. , Bonfanti R. , Meschi F. , De Pellegrin M. , Lesma A. , Rossini S. , Piemonti L. , Marchetti P. , Dotta F. , Bosi E. , Battaglia M. (2013) **Reduction of circulating neutrophils precedes and accompanies type 1 diabetes** *Diabetes* **62**:2072–2077

Wang G. , Cao L. , Liu X. , Sieracki N.A. , Di A. , Wen X. , Chen Y. , Taylor S. , Huang X. , Tiruppathi C. , Zhao Y.Y. , Song Y. , Gao X. , Jin T. , Bai C. , Malik A.B. , Xu J. (2016) **Oxidant Sensing by TRPM2 Inhibits Neutrophil Migration and Mitigates Inflammation** *Developmental cell* **38**:453–462

Wang J. , Hossain M. , Thanabalasuriar A. , Gunzer M. , Meininger C. , Kubes P. (2017) **Visualizing the function and fate of neutrophils in sterile injury and repair** *Science* **358**:111–116

Xu J. , Wang F. , Van Keymeulen A. , Herzmark P. , Straight A. , Kelly K. , Takuwa Y. , Sugimoto N. , Mitchison T. , Bourne H.R. (2003) **Divergent signals and cytoskeletal assemblies regulate self-organizing polarity in neutrophils** *Cell* **114**:201–214

Jorch S.K. , Kubes P (2017) **An emerging role for neutrophil extracellular traps in noninfectious disease** *Nat Med* **23**:279–287
<https://doi.org/10.1038/nm.4294>

Author information

Luyang Cao

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041, Guangzhou Regenerative Medicine and Health Guangdong Laboratory (GRMH-GDL), Guangzhou, China, 510530

Lixiang Ma

Department of Anatomy, Histology & Embryology, Shanghai Medical College, China, 200032

Juan Zhao

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Xiangyu Wang

Department of Anesthesiology, Huashan Hospital, Fudan University, Shanghai, China, 200040

Xinzou Fan

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Wei Li

Guangzhou Regenerative Medicine and Health Guangdong Laboratory (GRMH-GDL), Guangzhou, China, 510530

Yawen Qi

Guangzhou Regenerative Medicine and Health Guangdong Laboratory (GRMH-GDL), Guangzhou, China, 510530

Yingkui Tang

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Jieya Liu

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Shengxian Peng

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Li Yang

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Liangxue Zhou

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Li Li

Department of Anesthesiology, Huashan Hospital, Fudan University, Shanghai, China, 200040

Xiaobo Hu

Clinical Laboratory, Longhua Hospital, Shanghai University of Traditional Medicine, China, 200043

Yuan Ji

Department of Pathology, Zhongshan Hospital Fudan University, Shanghai, China, 200043

Yingyong Hou

Department of Pathology, Zhongshan Hospital Fudan University, Shanghai, China, 200043

Yi Zhao

Department of Rheumatology and Immunology, West China Hospital, Sichuan University, China, 610041

Xianming Zhang

Program for Lung and Vascular Biology, Stanley Manne Children's Research Institute, Ann & Robert H. Lurie Children's Hospital of Chicago, and Department of Pediatrics, Division of Critical Care, Northwestern University Feinberg School of Medicine, Chicago, USA, 60611

Youyang Zhao

Program for Lung and Vascular Biology, Stanley Manne Children's Research Institute, Ann & Robert H. Lurie Children's Hospital of Chicago, and Department of Pediatrics, Division of Critical Care, Northwestern University Feinberg School of Medicine, Chicago, USA, 60611

Yong Zhao

State Key Laboratory of Membrane Biology, Institute of Zoology, Chinese Academy of Sciences, China

Yuquan Wei

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

Asrar B. Malik

Department of Pharmacology, University of Illinois, College of Medicine, Chicago, USA, 60612

Hexige Saiyin

State Key Laboratory of Genetic Engineering, School of Life Sciences, Fudan University, Shanghai, China, 200433

For correspondence: saiyin@fudan.edu.cn

Jingsong Xu

Department of Neurosurgery, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, China, 610041

For correspondence: jingsong.xu@hotmail.com

Editors

Reviewing Editor

Pramod Mistry

Yale University, United States of America

Senior Editor

Mone Zaidi

Icahn School of Medicine at Mount Sinai, United States of America

Reviewer #1 (Public Review):

This study by Cao et al. demonstrates role of Neutrophil in clearing apoptotic hepatocytes by directly burrowing into the apoptotic hepatocytes and ingesting the effete cells from inside without causing inflammation. The authors applied intravital microscopy, Immunostaining and electron microscopy to visualize perforocytosis of neutrophil in hepatocytes. They also found that neutrophil depletion impairs the clearance of apoptotic hepatocytes causing impaired liver function and generation of autoantibodies, implying a role of defective neutrophil-mediated clearance of apoptotic cells in Autoimmune Liver disease. The experiments were well designed and conducted, the results were reasonably interpreted, and the manuscript was clearly written with logical inputs.

Further studies to explore the signals/mechanisms that determine why neutrophil specifically target apoptotic hepatocytes in liver would be of great clinical significance.

Reviewer #2 (Public Review):

Neutrophils are not known to be the cells responsible for removal of apoptotic cells through efferocytosis. This report provides strong evidence that neutrophils can remove apoptotic hepatocytes in vivo and in vitro. In addition, neutrophils, which are much smaller in size than hepatocytes, can burrow into apoptotic hepatocytes.

Neutrophils are the most abundant circulating leukocytes in human. They play important roles in innate immune responses to infections and tissue injuries. Although they are adept in phagocytosis of microbes, neutrophils are not known to normally conduct efferocytosis or phagocytose host cells including apoptotic cells and play a significant role in apoptotic cell removal. In this report the authors provide evidence to suggest that neutrophils are involved in removal of apoptotic hepatocytes with certain specificity (i.e., they do not remove HEK293 or HUVEC endothelial cells). Moreover, the authors also show that neutrophils can burrow

into the target cells and possibly ingest the target cells from the inside. The authors thus term this neutrophil-mediated efferocytosis process as "perforocytosis". Furthermore, evidence is provided to suggest that this neutrophil-mediated efferocytosis process keeps the number of apoptotic cells low in the livers and that defects in the processes may associate with autoimmune liver (AIL) disease phenotypes. Therefore, many of these findings are novel and the study is of important implications in our understanding of the role of neutrophils in autoimmune disease. Overall speaking, as the first report describing this novel finding, the authors have provided reasonably strong evidence for the conclusion that neutrophils burrow into apoptotic hepatocytes to perform "perforocytosis" to eliminate apoptotic hepatocytes. The importance, particularly in vivo significance, of this phenomenon needs to be further substantiated in future studies.

Author Response:

The following is the authors' response to the original reviews.

Thank you for considering our manuscript "An Unexpected Role of Neutrophils in Clearing Apoptotic Hepatocytes In Vivo". We also thank the referees for their review. We have addressed their comments in detail and added new data to buttress our conclusions.

Reviewer #1 (Public Review):

This study by Cao et al. demonstrates role of Neutrophil in clearing apoptotic hepatocytes by directly burrowing into the apoptotic hepatocytes and ingesting the effete cells from inside without causing inflammation. The authors applied intravital microscopy, Immunostaining and electron microscopy to visualize perforocytosis of neutrophil in hepatocytes. They also found that neutrophil depletion impairs the clearance of apoptotic hepatocytes causing impaired liver function and generation of autoantibodies, implying a role of defective neutrophil-mediated clearance of apoptotic cells in Autoimmune Liver disease. The experiments were well designed and conducted, the results were reasonably interpreted, and the manuscript was clearly written with logical inputs.

Thank you for your comments.

One weak point is that the signals/mechanisms that determine why neutrophil specifically target apoptotic hepatocytes in liver and no other organs or cells is not clearly understood.

We are still studying why neutrophils selectively phagocytose hepatocytes but not HUVEC or 293 cells. We have some intriguing preliminary data so far showing that apoptotic 293 cells have no significant increase of IL-1 β production as compared with their nonapoptotic controls; both apoptotic 293 cells and HUVECs do not have increased surface selectin proteins (new Fig. S3C).

Reviewer #2 (Public Review):

[...] By examination of HE-stained, noncancerous liver tissue sections from patients with hepatocellular carcinoma and hepatic hemangioma, the authors observed that cells with neutrophil nuclear morphology were inside apoptotic hepatocytes. The authors also further characterized this observation by staining the sections with neutrophil and apoptosis markers. In addition, the authors observed the same phenomena in mouse livers using intravital microscopy, which also recorded the time course of the disappearance of a neutrophil-associated apoptotic cell. The author went on further characterization of neutrophil-mediated efferocytosis of cultured hepatic cells in vitro and demonstrated the process was specific for apoptotic hepatic cells, but not HEK293 or endothelial cells. The in vitro system was then used to characterize the molecular bases for neutrophil-mediated efferocytosis of apoptotic hepatic cells. The evidence was provided to suggest that IL1b and IL-8 released from and selectins upregulated in apoptotic hepatic cells were important. Importantly, the authors used two methods to deplete the neutrophils and showed that the neutrophil depletion increased apoptotic cells in livers. Finally, the authors showed that neutrophil depletion caused defects in liver function parameters. At the end, the authors presented evidence to suggest that AIL disease may be due to defective neutrophils that fail to perform "perforocytosis."

Thank you for your comments.

Point #1. Although the evidence in its totality indicates that neutrophils burrow into apoptotic hepatocytes, the significance of this "perforocytosis" phenomenon and the circumstances under which it may occur remain to be better defined. In both neutrophil depletion models, the TUNEL-positive cells were not definitively identified rather than assuming they were hepatocytes.

Anatomically, the apoptotic hepatocytes are randomly distributed in the hepatic plate from the central vein to the portal region (please refer to the image below: hematoxylin staining of liver tissues, black arrowhead indicates perforocytosis sites).

Author response image 1.



Histologically, the structure of liver/hepatic lobe are well defined, and the cell types in the livers are easy to histologically identify based on their location, morphology and the relationship to hepatic plate and sinusoid. In addition, the hepatocytes are well known for its rich cytoplasmic components, cellular connection and prominent large round nucleus. Thus, hepatocytes are very easy to identify even without using specific molecular markers such as E-cadherin or albumin. Based on these characteristics, the TUNEL positive cells that we displayed in Fig. 5A are apoptotic hepatocytes.

Point #2. In addition, there are discrepancies in the number of neutrophils and apoptotic cells in mouse liver studies; Fig. 2a WT (many neutrophils; locations unclear) vs Fig. 5A Ctrl (a few neutrophils that appear in or near a vessel), and Fig. 2a DTR (a few apoptotic cells) vs Fig. 5A Depletion (many apoptotic cells).

In response, Fig. 2A demonstrates a larger area of the mouse liver (bar, 100 μ m), while Fig. 5A exhibits a relatively small area of the liver sample (bars, 20 μ m for Ctrl and 15 μ m for DTR). Similarly, apoptotic cells in Fig. 2A DTR need to zoom in to quantify. We apologize for the confusion, and we did quantify the apoptotic cells in Fig. 2A WT vs DTR (see the bar graph next to the images in Fig. 2A).

Point #3. Importantly, Fig 5a Ctrl, which is presumably a section from a mouse without any surgical treatment or without inflammation, the sole TUNNEL signal does not appear to be associated with neutrophils. Does this mean that "perforocytosis" primarily occurs in inflamed livers (Of note, human liver samples in Fig 1 are from patient with tumors. There should be inflammation in the livers of these patients).

In Fig 5A Ctrl, the TUNEL signal indicates apoptotic hepatocytes. The neutrophils (stained with anti-NE antibody, red) are associated with the apoptotic hepatocyte (Fig. 5A). We observed that perforocytosis primarily occurs in normal noninflamed livers.

Human liver samples in Fig 1 are from patient with tumors, hence it is possible that neutrophil burrowing is somehow associated with cancerous/inflammatory livers as the reviewer pointed out. This possibility was ruled out based on our method of sample preparation and experimental results themselves.

1. Both noncancerous and cancerous liver samples were sliced based on the anatomical appearance of normal and cancer tissues (differences were rather easy to identify, and these samples were prepared by highly experienced pathologists from the Liver Cancer Center of Zhongshan Hospital, Shanghai). Furthermore, the results were confirmed by determining whether the surrounding tissue contained microlesions characteristic of metastatic tumors. We only counted apoptotic hepatocytes in noncancerous regions having normal liver lobes and morphologically normal hepatocytes, plates, sinusoid and Kupffer cells. We also excluded hepatoma, chronic inflammatory regions, and necrotic regions.
2. We did not observe recruitment of neutrophils into apoptotic HCC cells, indicating that the clearance of apoptotic cancer cells was not mediated by neutrophils (unpublished observations).
3. It is hard for us to obtain normal human liver samples; however, we did study samples from patients with liver hemangioma characterized by aberrant vasculature in livers but with normal liver functions and the structure of hemangioma livers that we analyzed are nearly identical to a healthy liver in histology (these liver samples contained no cancerous regions and there was no apparent cirrhosis or inflammation). And here we obtained similar results (these are shown in Fig. 1B; a total of 40 apoptotic hepatocytes were examined).
4. Our data from normal mouse livers, isolated primary cells (hepatocytes and neutrophils) and cell lines (NCTC and HL60) all confirmed the central findings in this paper (Fig. 2, 3).

Point #4. The data on human AIL patient neutrophils raises more questions: how many AIL patients have been examined? Do these AIL neutrophils lack IL1, IL8 receptors, and/or selectin ligands? Are there increases in apoptotic hepatocytes in AIL patients?

In response, we have analyzed 16 AIL patient samples (see table below).

Author response table 1.

	No.	total apoptotic cells	phagocytized by PMNS	phagocytized by Monocyte
normal liver	32	227	227	0
AIL	16	110	8	40

We performed microarray assay to screen the differential gene expression of neutrophils from normal and liver autoimmune patients. We have identified that IL-1 β receptor, IL1R1 and selectin binding protein, P-selectin glycoprotein ligand 1 (PSGL-1) were all decreased in neutrophils from the AIL patients (new Fig 7D). These findings are consistent with our observations using cells and mouse models.

Point #5. Additionally, the overall numbers of apoptotic cells even in the absence of neutrophils are rare; thus, it is questionable that such rarity of apoptotic cells can cause significant AIL phenotypes.

We quantified apoptotic liver cells in percentages instead of overall numbers (Fig. 5, we were not able to precisely calculate the overall numbers, which could be large since billions of cells undergoing apoptosis daily). Depletion of neutrophils increased the percentage of apoptotic cells about 5-6-fold in livers, and we observed the generation of autoantibodies (Fig. 6).

Reviewer #1 (Recommendations For The Authors):

This study by Cao et al. was well designed and conducted, the results were reasonably interpreted, and the manuscript was clearly written with logical inputs.

It would further gain the significance of this study if authors could address the following questions:

- 1. What are the mechanisms/ signals that prevents AIL Liver neutrophils from burrowing into hepatocytes?*

We have identified that IL-1 β receptor, IL1R1 and selectin binding protein, P-selectin glycoprotein ligand 1 (PSGL-1) were all decreased in neutrophils from the AIL patients (new Fig 7D).

- 1. Have authors looked if autoantigens expressed on hepatocytes, which are often found in autoimmune liver disease trigger signaling events that activate neutrophils to burrow?*

Thank you for the comment, we have not examined autoantigens expressed in hepatocytes and plan to carry out this research as suggested.

- 1. Is perforocytosis observed in apoptotic hepatocytes induced by different agents like LPS, TNF-a , rapamycin, alcohol etc?*

We did not observe perforocytosis in LPS or TNF-a treated hepatocytes. One possible reason is that LPS or TNF-a we used induced massive necrosis instead of apoptosis. Howere, we did observe neutrophil perforocytosis in FasL-induced apoptotic hepatocytes (unpublished observations).

Reviewer #2 (Recommendations For The Authors):


In addition to the questions raised in the "Public review" section, the authors are also recommended to address the following issues:

1. Why is CD11b+ not associated with the apoptotic sites as neutrophils express CD11b

We have co-immunostained human liver samples with CD11b antibody (from Abcam: ab133357) and MPO antibody (from R&D: AF3667) and observed that tissue infiltrating neutrophils in livers have low to undetectable levels of CD11b expression (please refer the image below; white arrowheads point to neutrophils). Few CD11b+ cells in liver tissues express MPO (the CD11b+ cells are mostly macrophages, unpublished observations).

Based on these data, we conclude that CD11b is hardly expressed in neutrophils inside livers.

Author response image 2.



1. Can TUNEL signals in Fig. S1C be from apoptotic neutrophils?

In response, the fragmentation of nucleus is a hallmark of apoptosis hence TUNEL staining will uniformly label all fragmented parts of apoptotic nucleus. The nucleus of NE+ neutrophils are not labelled by TUNEL staining in Fig. S1C. The TUNEL+ nuclear fragments seen inside neutrophils are nuclear debris of apoptotic hepatocytes phagocytosed by neutrophils (Fig. S1C).

1. The Fig 2B experiment may be done with induced apoptosis so that neutrophil burrowing steps may be recorded from the very beginning and a better time course for the entire process can be assessed.

Thank you for the suggestions, we had tried many times with various conditions, yet still had no success to capture the very beginning of perforocytosis in vivo. We are continuing to work on this.

1. In "we found that U937 cells exhibited much lower phagocytosis of apoptotic NCTC cells than did HL60 cells (Fig. S2B, C)," the citation should be only S2C

Thank you for pointing this out, we have corrected this in the manuscript.

*1. Both neutrophil depletion models cause neutrophil death, which may complicate the interpretation of the liver function and AIL disease phenotypes. A neutropenic model such as *G-CSFR*^{-/-} or *Cebpe*^{-/-} mice may be used to avoid the caveat of antibody/DTR-dependent depletion models.*

Thank you for this thoughtful suggestion. We have also induced AIL phenotypes in mice by using α -Galcer. α -Galcer did not cause neutrophil death but impaired neutrophil perforocytosis and further generated AIL phenotypes in mice (unpublished observations). We plan to perform the similar experiments in *G-CSFR*^{-/-} or *Cebpe*^{-/-} mice as the reviewer suggested.

1. RNAi silencing experiments need additional controls for off-target effects

These RNAi silencing constructs were purchased from Santa Cruz Biotechnology and the off-target effects have been tested by the company. No significant off-target effects have been detected according to the manufacture report.