

ZMYM2 controls human transposable element transcription through distinct co-regulatory complexes

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

October 5, 2023 (this version)

Reviewed preprint version 1

May 2, 2023

Sent for peer review

February 25, 2023

Posted to preprint server

January 24, 2023

Danielle Owen, Elisa Aguilar-Martinez, Zongling Ji, Yaoyong Li, Andrew D. Sharrocks 

Faculty of Biology, Medicine and Health, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, UK

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

ZMYM2 is a zinc finger transcriptional regulator that plays a key role in promoting and maintaining cell identity. It has been implicated in several diseases such as congenital anomalies of the kidney where its activity is diminished and cancer where it participates in oncogenic fusion protein events. ZMYM2 is thought to function through promoting transcriptional repression and here we provide more evidence to support this designation. Here we studied ZMYM2 function in human cells and demonstrate that ZMYM2 is part of distinct chromatin-bound complexes including the established LSD1-CoREST-HDAC1 corepressor complex. We also identify new functional and physical interactions with ADNP and TRIM28/KAP1. The ZMYM2-TRIM28 complex forms in a SUMO-dependent manner and is associated with repressive chromatin. ZMYM2 and TRIM28 show strong functional similarity and co-regulate a large number of genes. However, there are no strong links between ZMYM2-TRIM28 binding events and nearby individual gene regulation. Instead, ZMYM2-TRIM28 appears to regulate genes in a more regionally defined manner within TADs where it can directly regulate co-associated retrotransposon expression. We find that different types of ZMYM2 binding complex associate with and regulate distinct subclasses of retrotransposons, with ZMYM2-ADNP complexes at SINEs and ZMYM2-TRIM28 complexes at LTR elements. We propose a model whereby ZMYM2 acts directly through retrotransposon regulation, which may then potentially affect the local chromatin environment and associated coding gene expression.

Genome browser session

UCSC browser session containing the peak tracks: http://genome.ucsc.edu/cgi-bin/hgTracks?db=hg19&position=chr1:18,078,462-18,084,961&hide=all&hgct_customText=http://bartzabel.ls.manchester.ac.uk/sharrockslab/yaoyong/ZNF198/index_file_hg19_chipSeq_ZMYM2_final.txt

Original ChIP-seq and ATAC-seq data from U2OS cells can be viewed On ArrayExpress at: E-MTAB-12292 (ADNP and TRIM28 ChIP-seq), E-MTAB-12293 (SUMO ChIP-seq) and E-MTAB-12294 (ATAC-seq)

eLife assessment

ZMYM2 is a transcriptional corepressor but little was known about how it is recruited to chromatin. This **important** study reveals that ZMYM2 homes to distinct classes of retrotransposons bound by the TRIM28 and ChAHP complexes in human cells, which is broadly relevant for the field of transcriptional regulation. Much of the evidence supporting the claims of the authors is **convincing**. Since widespread ZMYM2-mediated control of transposon activity is not apparent in RNA-seq data, further experiments are needed to demonstrate a more general role beyond the retrotransposons analysed in this study.

Introduction

The zinc finger protein ZMYM2 (otherwise known as ZNF198) was originally identified as part of an oncogenic fusion protein with the receptor tyrosine kinase FGFR1 in myeloproliferative disease (Xiao et al., 1998 [↗](#); Reiter et al., 1998 [↗](#)). In this context, the N-terminal portion of ZMYM2 promotes the oligomerisation and activation of FGFR1 (Xiao et al., 2000 [↗](#)). Further disease links have recently been uncovered where heterozygous loss of function ZMYM2 mutations lead to congenital anomalies of the kidney and urinary tract (CAKUT) and chronic kidney disease (Connaughton et al., 2020 [↗](#)). The latter observation suggests a potential developmental role and ZMYM2 has been shown to play important roles in the early stages of embryonic stem cell (ESC) differentiation and commitment. ZMYM2 promotes the transition from totipotency to pluripotency in mice (Yang et al., 2020). Similarly, in human cells, ZMYM2 plays a key role in determining ESC identity by allowing the transition to and maintenance of the primed pluripotent state (Lezmi et al., 2020 [↗](#)). Furthermore, ZMYM2 has been shown to play a key role in opposing reprogramming in human fibroblasts where ZMYM2 loss increases reprogramming efficiency (Toh et al., 2016 [↗](#); Lawrence et al., 2019 [↗](#)).

Molecularly, ZMYM2 has been shown to act as a transcriptional repressor and can bind to the LSD1-CoREST-HDAC1 corepressor complex (LCH) via its zinc fingers (Gocke and Yu, 2008 [↗](#)). It is through this repressive complex that ZMYM2 promotes pluripotency in mice (Yang et al., 2020). However, ZMYM2 has been shown to have an increasingly complex set of interaction partners such as the transcription factors TBX18 with a potential role in ureter development (Ludtke et al., 2022) and B-MYB and hence a potential link to cell cycle control (Cibis et al., 2020 [↗](#)). Proteomic screens have also revealed multiple ZMYM2 binding partners, either from using ZMYM2 itself as a bait (Connaughton et al., 2020 [↗](#)) or through detection as a component of other repressive complexes by using baits such as LSD1/KDM1A and HDAC2 (Hakimi et al., 2003 [↗](#); Shi et al., 2005 [↗](#)). The range of interaction partners solidifies ZMYM2 as a potential transcriptional repressor protein, but additional roles are hinted at such as DNA repair, exemplified by its ability to antagonise 53BP1 function at DNA double strand breaks to favour repair by homologous recombination (Lee et al., 2022 [↗](#)).

We identified ZMYM2 in a screen for multi-SUMO binding proteins (Aguilar-Martinez et al., 2015 [↗](#)) which builds on the complex interplay previously observed between ZMYM2 and SUMO. Additional work supports the non-covalent SUMO binding activity of ZMYM2 (Guzzo et al., 2014 [↗](#)) although the two studies differed in mapping of the SUMO interacting motifs, with the former mapping them to the N-terminal part of the protein and the latter mapping them to the centrally located zinc finger region. Furthermore, ZMYM2 itself is covalently modified with SUMO (Kunapuli

et al., 2006 [\[4\]](#)) suggesting a complex and important role for SUMO in determining ZMYM2 activity. Indeed, SUMO binding is needed for recruitment of ZMYM2 to chromatin (Aguilar-Martinez et al., 2015 [\[4\]](#)) and the ability of ZMYM2 to recruit SUMOylated HDAC-1 (Gocke and Yu, 2008 [\[4\]](#)).

In this study, we further investigated the molecular activities of ZMYM2 and demonstrated that it is associated with two distinct chromatin-bound complexes containing either ADNP or TRIM28. Both types of ZMYM2 chromatin binding regions are associated with different subclasses of retrotransposons. We focussed on TRIM28 and demonstrated widespread functional cooperativity with ZMYM2 in gene regulation. However, ZMYM2-TRIM28 binding regions are directly associated with ERV retrotransposable elements rather than with protein coding genes, suggesting an indirect mechanism for transcriptional control of the coding genome.

Materials and Methods

Cell culture

U2OS and HEK293T cells and derivatives were grown in DMEM supplemented with 10% fetal bovine serum. Cells were cultured at 37°C, 5% CO₂ in a humidified incubator and were passaged every 3-4 days using 0.05% Trypsin-EDTA (ThermoFisher Scientific, 25300). Stable U2OS-derived cell lines, flag-ZMYM2(WT) and SUMO3(K11R/Q90P) were created using U2OS-Flp-in cells (Aguilar-Martinez et. al. 2015).

RIME Mass Spectrometry analysis

U2OS-derived cells were grown in 15 cm dishes (6 confluent dishes per sample) and fixed using DMA for 20 minutes at room temperature, followed by 1% formaldehyde (ultra pure) addition for 5 minutes at room temperature. Cells were washed twice with cold 1X PBS. Cells were harvested in 1X PBS supplemented with protease inhibitors (cOmplete cocktail, 11873580001, Roche). Cells were centrifuged at 3,000 rpm for 10 minutes at 4°C. The supernatant was removed, the cell pellet was resuspended in 1X PBS with protease inhibitors and spun down again. The pellets were resuspended in LB1 buffer (50 mM Hepes-KOH pH 7.5, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% Igepal, 0.25% Triton X-100 supplemented with cOmplete cocktail (11873580001, Roche), 20 mM N-ethylmaleimide (NEM) and 30 mM iodoacetate (IAA)), and incubated with gently agitation for 10 minutes at 4°C. Cells were spun down at 2,000 g for 5 minutes at 4°C. Pellets were resuspended in 10 ml of LB2 buffer (10 mM Tris-HCl pH8, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, supplemented with cOmplete cocktail (11873580001, Roche), 20 mM NEM and 30 mM IAA) incubated with gently agitation for 5 minutes at 4°C. Samples were spun down at 2,000 g for 5 minutes at 4°C. Pellets were resuspended in LB3 (10 mM Tris-HCl pH 8, 100 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 0.1 % Na-Deoxycholate, 0.5% N-lauroylsarcosine, supplemented with cOmplete cocktail (11873580001, Roche), 20 mM NEM and 30 mM IAA), aliquoted and sonicated; 15 cycles of 30 seconds sonication, 30 seconds paused, in a water bath at 4°C. After sonication, triton X-100 was added to the samples. Samples were spun down at 13,000 rpm for 10 minutes at 4°C. The supernatant was removed and incubated with antibody coupled beads. Samples were incubated with gentle agitation over night at 4°C. Beads were washed 10 times with 1 ml of RIPA buffer and twice with freshly made 100 mM ammonium bicarbonate. Beads were spun down at 1,000 rpm for 1 minute to remove residual ammonium bicarbonate. To release the proteins bound to the beads, 100 ng of trypsin was added and incubated overnight at 37°C. The supernatant was carefully removed and analysed by an Orbitrap velos mass spectrometer as described previously (Aguilar-Martinez et al., 2015 [\[4\]](#)).

The data produced was searched using Mascot (Matrix Science version 2018_01), against the SwissProt (version 2018_01) with a taxonomic break of the Human (*Homo sapiens*) database. Data was validated using Scaffold (5.0.2). Proteins found in more than 50% of CRAPome samples, ribosomal proteins and proteins with zero spectral counts in any experiment were removed from

the list. Proteins showing an average of >4 spectral counts across three experiments were included in the final list and were then compared to proteins bound to ZMYM2 on mouse ESCs (Yang et al., 2020) and BioID experiment on 293 cells (Connaughton et al., 2020 [DOI](#)).

RNA interference, RT-qPCR and RNA-seq analysis

ON-TARGETplus SMART pool siRNAs from Dharmacon were used to knock-down ZMYM2, TRIM28 or ADNP (L-021348-00-0005, L-005046-00-0005 and L-012857-01-0005 respectively). Non-targeting siRNA pool (siNT, Dharmacon, D-001810-10-20) was used as a control. To carry out RNA interference (RNAi), cells were transfected with siRNA using *Lipofectamine® RNAiMAX* (Invitrogen) according to the reverse transfection method in the manufacturer's instructions. 48h after transfection, cells were transfected again with the same siRNA using forward transfection method for another 48h.

Total RNA was extracted using the RNeasy plus kit (Qiagen) following the manufacturer's protocol. To detect LTR expression, additional DNase digestion was performed using RNase-Free DNase Set (Qiagen) to completely remove genomic DNA.

For RNA-seq analysis, U2OS cells in 6-well plates were transfected twice, second transfection was done 48 h after the first one. A final concentration of 100 nM of each siRNA using *Lipofectamine® RNAiMAX* (ThermoFisher Scientific), were used following the manufacturer's instructions. Total RNA was extracted 48 h after the second transfection using an RNeasy plus kit with in-column DNaseI treatment, following the manufacturer's instructions (Qiagen). Immuno-blot and RT-qPCR for ZMYM2, TRIM28 or ADNP expression were carried out to verify knock-down of the mRNAs and proteins.

Quantitative RT-PCR was performed using 30 ng of total RNA and QuantiTect SYBR Green RT-PCR kit (Qiagen, 204243) according to the supplier's protocol. Data were analysed by Qiagen Rotor-Gene Series software. Data were normalised against the control gene *GAPDH* (primers ADS2184 ACAGTCAGCCGCATCTTCTT and ADS2185 TTGATTTTGGAGGGATCTCG) or *RPLP0* (ADS5454, ATTACACCTTCCCACTTGCT and ADS5455 CAAAGAGACCAAATCCCATATCCT). RNA samples were run in duplicate from at least three independent experiments. The primer-pairs used for RT-qPCR experiments were *ZMYM2* (ADS3573, GGTAACAACTGAGATTCGCCA and ADS3574, CCAGAACATTATTCCAGCACCT), *TRIM28* (ADS6181, GTTCAGTGGGACCTCAATGC and ADS6182, GATCATCTCCTGACCCAAAGC), *ADNP* (ADS6688, AGGCTGACAGTGTAGAGCAAG and ADS6689, GACTGCCCATTTGAGTGATTTT), *MER11A* (ADS6917, AATACACCTGGTCTCCTGC and ADS6918, AACAGGACAAGGGCAAAAGC), *LTR10A* (ADS6919, ACAACTTTCCACCAAGTCCT and ADS6920, GCAGGAGTATGAGCCAGAGT), *THEC1C-int* (ADS6543, CCAACCCGACATTTCCCTTC and ADS6544, (AGGGGCCAAGGTACATTTCA), *AluSx3* (ADS6937, TGAGGTGGGCTGATCATGAG and ADS6938, TGCAACCTCCACCTCCTAAG), *AluJr* (ADS6939, AGGCTGAGTTGGGAGGATTG and ADS6940, GCAGGATCTCATTCTGTTGCC), *MIRb* (ADS6941, AGTGCCAGCTTTGGTTTCAG and ADS6942, AGACGAGAACACTGAGGCTC).

Western blot analysis and co-immunoprecipitation assays

For immuno-blotting, the antibodies used are listed in Supplementary Table S1. Lamin-B and Tubulin were used as loading controls. Proteins were visualised using IRDye infrared dye (Li-Cor Biosciences) conjugated secondary antibodies and the signals were visualised using a LI-COR Odyssey® CLx scanner and analysed using ImageStudio v 5.2.5.

Co-immunoprecipitation was performed as previously described (Ji et al., 2012). The anti-ZMYM2, ZMYM3 and ADNP antibodies (Supplementary Table S1) were used for immunoprecipitation, with normal rabbit IgG (Millipore) as negative control. For co-immunoprecipitation analysis of endogenous TRIM28 with the EGFP-tagged ZMYM2 proteins, HEK293T cells were transfected with EGFP vector, EGFP-ZMYM2(WT) (pAS4329) or EGFP-ZMYM2(SIMmut) (pAS4330) plasmids using

Polyfect Transfection Reagent (Qiagen, Cat. No. ID: 301105) for 24 hours, cell lysates were prepared in the presence of cOmplete cocktail (11873580001, Roche) and 20 mM N-ethylmaleimide (NEM) and followed by immunoprecipitation using TRIM28 antibody.

In situ Proximity ligation assay (PLA)

U2OS cells grown on glass cover-slips were washed once in 1X PBS and then fixed and permeabilised with 3.7% para-formaldehyde plus 0.8% Triton X-100 for 15 minutes at room temperature. Cells were washed three times with 1X PBS. During all of the incubation steps cells were kept in a humidified chamber. The cells were first incubated for one hour with 1% BSA dissolved in 1X PBS, followed by one hour incubation with both primary antibodies (anti-c-myc Santa Cruz (9e10):sc-40 1:500, and anti-KAP1 Abcam, ab10483 1:500, diluted in 1% BSA-1X PBS). Cells were washed three times with 1X PBS and once with PBS⁺ (1X PBS, 0.1% Tween-20, 1% BSA). Cells were incubated with PLA probes, ligase and polymerase following Duolink-II fluorescence instructions. After the final wash, nuclei were stained by incubating with Hoechst solution (1 µg/ml) for 10 minutes at room temperature. Cells were washed twice with 1X PBS. Preparations were mounted in vectashield antifade medium (Vector) and sealed with nail varnish. Z-stack images covering the entire thickness of the samples were acquired on a Delta Vision RT (Applied Precision) restoration microscope using a 100X/1.40 Plan Apo objective. The images were collected using a Coolsnap HQ (Photometrics) camera with a Z optical spacing of 0.2 µm. Raw images were then deconvolved using the Softworx software. Deconvolved images were used to quantify interactions per cell using the single cell analysis function of Blobfinder software.

RNA-seq data analysis

RNA-seq was performed in triplicate. Libraries were generated using TruSeq stranded mRNA library kit (Illumina) and sequenced on HiSeq 4000 platform (Illumina). Reads were mapped to the human genome hg19 as described previously (Ji et al., 2021 [DOI](#)). 14.6 M-43.4 M reads were uniquely mapped for the three replicates of siZMYM2, 18.7 M-48.5 M reads were uniquely mapped for the three replicates of siTRIM28, and 22.6 M-23.1 M reads were uniquely mapped for the three replicates of siADNP (Supplementary Table S2). Differentially regulated genes between two different conditions were established using the R package edgeR (Robinson et al., 2010 [DOI](#)) with the criteria adjusted p-value < 0.01 and fold change > 1.6. The online tool Metascape (Zhou et al., 2019 [DOI](#)) was used for GO terms and pathway enrichment analysis on the DE gene sets.

ATAC-seq analysis

The ATAC-seq data in U2OS cells were generated by using the protocol essentially as described previously (Yang et al., 2019 [DOI](#)). ATAC-seq libraries were generated from U2OS cells treated with a non-targeting siRNA pool. Cells were lysed 48 h after transfection in cold lysis buffer (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl₂ and 0.1% Igepal). Nuclei were concentrated by centrifugation at 500 g for 10 minutes at 4°C. The nuclei pellet was resuspended in nuclease free water, diluted 1:10 and quantified. The equivalent volume for 50,000 nuclei was made up to 22.5 µl with nuclease free water. Nuclei were incubated for 30 minutes at 37°C, 300 rpm, with 25 µl of 2x TD buffer and 2.5 µl tagmentation enzyme (Illumina transposase FC-121-1030). DNA was purified using a Qiagen mini elute kit following the manufacturer's instructions. Library fragments were amplified using standard PCR and Nextera primers (Illumina) by adding 25 µl of 2x NEBnext master mix, 2.5 µl forward primer, 2.5 µl reverse primer and 20 µl transposased DNA. PCR was run for 5 cycles, 72°C, 5 minutes; 98°C 30 seconds; cycle of 98°C 10 seconds, 63°C 30 seconds, 72°C 1 minute, paused at 4°C. To determine the total number of cycles needed for the amplified library, 5 µl of the PCR mix were taken and added to a new mix containing 5 µl 2X master mix, 1 ml of each primer, 0.6 µl of 10X SYBR green and 2.4 µl H₂O. In a qPCR machine, 20 cycles were run, the total number of cycles for the ATAC library was chosen according to qPCR cycle that gave 1/3 of saturation. DNA was purified using Ampure XP beads following the manufacturer's instructions.

The reads in the ATAC-seq data were mapped to the human genome hg19 using Bowtie2 with the same settings as those described below for the ChIP-seq data, and only the uniquely mapped reads were used in the analysis.

Chromatin immunoprecipitation (ChIP)-qPCR and ChIP-seq assays

ChIP-qPCR was performed using primers to detect *TXNL4A*, *EMX2*, and *FAM109A*, as described previously (Aguilar-Martinez et al., 2015 [↗](#)), *MSTB* (ADS6541, GACCAGCTGACCAAAACG and ADS6542, ACTCAGGCCAAGTCTCTCTC), and *THEC1-int* (ADS6543, CCAACCCGACATTTCCTTC and ADS6544, AGGGGCCAAGGTACATTCA).

5×10^7 U2OS cells were used in each ChIP-seq experiment, which was performed essentially as described previously (Ji et al., 2012) with anti-TRIM28 antibody (Supplementary Table S1) as indicated. There was minor modification in immuno-precipitation step for the ADNP ChIP-seq experiments, which was performed overnight at 4°C by incubating the shared DNA-protein complex with 10 µg of anti-ADNP (Abcam Ab231950) antibody, followed by adding Dynabeads protein A (Invitrogen) and incubating for further 2 hours.

Immunoprecipitated DNA was purified with a PCR purification kit (Qiagen) and approximately 10 ng of DNA were processed for sequencing

ChIP-seq data processing and analysis

The data analysis of both ZMYM2(WT) and ZMYM2(SIM2mut) ChIP-seq data were described previously (Aguilar-Martinez et al., 2015 [↗](#)), including mapping the reads to human genome hg19 and obtaining 1188 peaks from the ZMYM2(WT) ChIP-seq data. We removed 8 peaks because those 8 peaks have very high signal in all the ChIP-seq data we examined and one input DNA data of U2OS cells (also in Aguilar-Martinez et al., 2015 [↗](#)), and used 1180 ZMYM2 peaks in all the analysis in the paper. The R method *kmeans* with the option 'centers=3' was used to cluster the 1180 ZMYM2 peaks into three clusters.

For TRIM28, ADNP and SUMO ChIP-seq analysis, two biological replicates were generated for each protein and sequenced using the Illumina HiSeq4000 platform. The paired-end reads were aligned to the human genome hg19 using Bowtie2 (Langmead and Salzberg, 2012) with the setting 'very sensitive' and all other default settings. The uniquely aligned reads were selected for the further analysis. Duplicate reads were removed using Picard (<http://broadinstitute.github.io/picard/> [↗](#)) and the reads mapped to the mitochondrial chromosome were also removed. The correlation coefficients between the two replicates of the read counts was between 0.85 - 0.95 for the two replicates in our ChIP-seq data. The reads in the two replicates were then pooled, and with the pooled reads from the two input replicates as control, peaks were called from the pooled ChIP-seq using MACS2 (Zhang et al., 2008) with the options "-g hs -f BAMPE". To identify overlapping peaks between two samples, one peak was considered to overlap with another peak if they overlap by at least 30% of each of their lengths. Venn diagrams showing the overlapping of peaks from different ChIP-seq datasets were created using the R package Vennrable (<https://github.com/js229/Vennerable> [↗](#)). Average tag density plots were generated as described previously (Ji et al., 2021 [↗](#)).

The genomic distribution of the ChIP-seq peaks was calculated using all the transcripts in the Ensembl human gene annotation database v75 with the human genome hg19. The genome was divided into five types of regions; promoter or TSS regions [-1 kb to 0.1 kb relative to TSS], TTS regions [-0.1 kb to 1 kb relative to TTS], exonic regions containing all exons within [TSS+0.1 kb to TTS-0.1 kb], intronic regions containing all introns within [TSS+0.1 kb to TTS-0.1 kb], and all other regions were classified as "distal intergenic" regions. A peak was linked to a region if its summit is located in the region.

The software HOMER ([Heinz et al., 2010](#)) was used with default settings for the motif enrichment analysis within a set of peaks (in 101 bp windows centred on the peak summits). The human genome annotations of LTR, SINE and LINE in hg19 contained in HOMER were also used in the analysis of their relationship to the ZMYM2 peaks.

To associate ZMYM2 or ADNP binding events with potential biological functions, we used the online tool GREAT ([McLean et al., 2010](#)) to obtain the GO terms for each set of ChIP-seq peaks.

Statistical analysis

Statistical analysis for real-time PCR results was performed using the Student t test. The error bars in all graphs represent standard deviation. The statistical test for the ATAC-seq data signal (Fig. S2E) and the distance of the ZMYM2 peaks in different clusters to the nearest TAD domains were also performed using the Student t test. Fisher Exact test was used for testing the significance of the overlaps between different sets of DE genes. The software PEGS ([Briggs et al., 2021](#)) was used to assess the statistical significance of the association between the ZMYM2 peaks and the DE genes.

Datasets

Our ChIP-seq and RNA-seq data from U2OS cells have been deposited with ArrayExpress. Accession numbers: E-MTAB-12292 (ADNP and TRIM28 ChIP-seq), E-MTAB-12293 (SUMO ChIP-seq), E-MTAB-12294 (ATAC-seq), and E-MTAB-12291 (RNAseq following ZMYM2, TRIM28 or ADNP depletion).

The following existing datasets were used: H3K18ac ChIP-seq in U2OS cells (E-MTAB-3695), ZMYM2 ChIP-seq (WT and SIM2mut) in U2OS cells (E-MTAB-2701; [Aguilar-Martinez et al., 2015](#)), ZMYM2 ChIP-seq in mouse mESC CJ7 cells (GSM3384427, GSM3384428; [Yang et al., 2020](#)), Trim28 ChIP-seq in Mouse E14 ESCs (GSM3611260; [Seah et al., 2019](#)), H3K9me3 ChIP-seq in Mouse E14 ESCs (GSM3611253, GSM3611256, GSM3611258; [Seah et al., 2019](#)), SUMO ChIP-seq in mouse ESCs (GSM2629945, GSM2629946; [Cossec et al., 2018](#)), H3K27ac and H3K9me3 in human H1 ESCs (<https://egg2.wustl.edu/roadmap/data/byFileType/alignments/consolidated/>; [Kudaje et al., 2015](#)).

Results

ADNP is a coregulatory binding partner for ZMYM2

To investigate the molecular function of ZMYM2 we used the mass spectrometry approach RIME ([Mohammed et al., 2013](#)) to identify proteins co-binding with ZMYM2 on chromatin. We performed several independent experiments using an anti-flag antibody for immunoprecipitation of ZMYM2 from a U2OS cell line containing a Flag-tagged version of ZMYM2 ([Aguilar-Martinez et al., 2015](#)) or using an antibody against endogenous ZMYM2 in U2OS cells. We identified 27 proteins in all three experiments with an average of >4 spectral counts (Supplementary Table S3). Among these, the highest scoring interactors were the transcriptional co-regulators TRIM28 and ADNP (**Fig. 1A**). ZMYM2 and ADNP are part of a larger interacting network of proteins identified in the RIME analysis (**Fig. 1B**), which includes the other two components of the ChAHP complex, CHD4 and CBX1/HP1β ([Kaaij et al., 2019](#)). Another prominent complex identified was the GTF3C complex (Supplementary Fig. S1A) which is consistent with a previous study that identified ZMYM2 from reciprocal RIME experiment using GTF3C2 as bait ([Ferrari et al., 2020](#)). A large proportion of the interactors were previously identified in two proteomic screens for ZMYM2 binding proteins from mouse ESCs ([Yang et al., 2020](#)) and/or human HEK293 cells ([Connaughten et al., 2020](#)). They are also often found as binding partners in proteomic screens for other chromatin regulators as exemplified by BEND2, whose homologue BEND3 binds to ZMYM2 in mouse testes ([Ma et al., 2022](#)). However, 10 ZMYM2 binding proteins were uniquely identified here, including

SUMO2 and SUMO3. The latter discovery is consistent with findings from screens for SUMO binding proteins where ZMYM2 was identified (Aguilar-Martinez et al., 2015; Bruninghoff et al., 2020). We validated interactions between endogenous ZMYM2 and ADNP by co-immunoprecipitation analysis (Fig. 1C) and also confirmed interactions of ADNP with the closely related ZMYM3 paralogue (Supplementary Fig. S1B). This data is consistent with the identification of ZMYM2 and ZMYM3 in other mass spectrometry datasets along with ADNP (Ostapcuk et al., 2018; Connaughten et al., 2020). We also detected the ADNP paralogue ADNP2 as a ZMYM2 interactor.

Having established a physical interaction between ZMYM2 and ADNP, we next studied their functional interplay. First, we performed two independent ChIP-seq experiments for ADNP in U2OS cells and uncovered 5,108 peaks found in both datasets. These ADNP peaks showed enrichment of binding motifs for several transcription factors with the top two motifs for HBP1 and IRF both found in over 35% of target regions (Fig. 1D). The top scoring motif for ADNP previously identified from ChIP-seq in mouse embryonic stem cells CGCCCYCTNSTG (Ostapcuk et al., 2018), was not identified in this unbiased analysis. We therefore searched for the motif in the ADNP peaks and found 775 in 5,108 peaks (15%) which was a substantially higher frequency than in a background dataset (9.5%) but well below the 63% frequency found in mouse embryonic stem cells. This motif was therefore present but at a relative low frequency and lowly enriched, suggesting an alternative chromatin recruitment mechanism. Co-binding on chromatin with ZMYM2 is inferred as there is a significant overlap in chromatin occupancy of ZMYM2 with the ADNP peaks (Fig. 1E). Example loci show clear co-binding of the two proteins (Fig. 1F).

Given this co-occupancy on chromatin we asked whether ZMYM2 and ADNP shared gene regulatory activity. We depleted each protein individually (Supplementary Fig. S1C and S3G) and performed RNA-seq to monitor transcriptional changes. Depletion of either factor alone gave roughly equivalent numbers of up and downregulated genes (Supplementary Table S4). However, there was a highly significant overlap in the genes upregulated following depletion of either factor, and also a highly significant overlap in the genes downregulated following reductions in either protein (Fig. 1G). Conversely, in contrast to these similarities, gene regulation showing reciprocal directionality following either ADNP or ZMYM2 depletion showed a very small and insignificant overlap (Supplementary Fig. S1D) and this is further emphasised by a scatter plot depicting this data (Supplementary Fig. S1E). Thus, there is a strong overlap in gene regulatory activity for ZMYM2 and ADNP, although the large numbers of genes directionally co-regulated by these two proteins (ie either positively or negatively) indicates no clear common role as either an activator or repressor.

Together these results demonstrate that ZMYM2 can function through a transcriptional regulatory complex containing ADNP to co-ordinately control gene activity.

ZMYM2 binds to molecularly distinct chromatin regions

Next, we turned to the potential interplay between ZMYM2 and TRIM28. Previous studies demonstrated that many zinc finger transcription factors can recruit TRIM28 to chromatin to repress transcription (Reviewed in Bruno et al., 2019). However this activity has previously been attributed to TRIM28 binding to the KRAB repression domain associated with many zinc finger proteins, but ZMYM2 lacks such a domain. To explore whether TRIM28 could be found in the same regions of chromatin as ZMYM2 we compared ChIP-seq data for TRIM28 in U2OS cells with our ZMYM2 binding profile. We also included ChIP-seq for SUMO2/3 and a SUMO binding defective version of ZMYM2, ZMYM2-mutSIM2 (as we previously showed that SUMO is needed for recruitment of ZMYM2 to chromatin; Aguilar-Martinez et al., 2015), markers of active chromatin including ChIP-seq data for H3K18ac (Chen et al., 2016) and ATAC-seq data. We took a ZMYM2-centric view of the data and plotted the signal for the other chromatin binding/modification events centred on the ZMYM2 binding peaks and then clustered the data (Fig. 2A and B). Three distinct clusters were revealed. Cluster 3 exhibits ZMYM2 binding but little evidence of any other




Figure 1.

RIME analysis identifies ADNP as a co-regulatory partner of ZMYM2.

(A) Summary table of the ten top scoring interactors for ZMYM2. The average number of peptides across three RIME experiments are shown and whether detected previously in ZMYM2 IP-mass spectrometry experiments is indicated (¹Yang et al., 2020; ²Connaughten et al., 2020). Core members of the ChAHP and GTF3C complexes are highlighted. (B) Depiction of interactions between ZMYM2 binding partners found in RIME experiments with known previous molecular interactions found in the STRING database (Jensen et al., 2009). (C) Co-immunoprecipitation analysis of ADNP with ZMYM2. Immunoprecipitation (IP) was performed with ZMYM2 or control IgG antibody from U2OS cells and resulting proteins detected by immunoblotting (IB) with the indicated antibodies. Molecular weight markers (kDa) and 10% input are shown. (D) *De novo* motif analysis of ADNP binding regions. The top five most significantly enriched motifs are shown, along with motif similarity to the indicated protein in brackets. (E) Venn diagram showing the overlap between ZMYM2 and ADNP binding regions. (F) UCSC genome browser of ChIP-seq data on example genomic loci showing binding of both ZMYM2 and ADNP. (G) Venn diagram showing overlaps in genes upregulated (left) or downregulated (right) following ADNP (orange) or ZMYM2 (red) depletion (fold change >1.6; Padj <0.01).

binding/modification events. However, cluster 1 is enriched for both wild-type ZMYM2 and TRIM28 binding. This cluster is relatively depleted of ATAC-seq and H3K18ac signal, suggesting that regions occupied by both ZMYM2 and TRIM28 are not areas of active chromatin (**Fig. 2A** and **B**). Indeed, when we mapped H3K9me3 signal for H1 ESCs to the ZMYM2 binding regions, stronger levels of this repressive mark were observed in cluster 1 despite the differing cell types (Supplementary Fig. S2A). We validated H3K9me3 occupancy at ZMYM2 binding sites from cluster 1 in U2OS cells (Supplementary Fig. S2B). SUMO2/3 is also strongly detected in cluster 1 and SUMO has often been associated with transcriptional repressive activity (reviewed in [Garcia-Dominguez and Reyes, 2009](#)). We also interrogated ChIP-seq data from mouse ESCs and found that a cluster of ZMYM2 binding regions (cluster B) was strongly associated with TRIM28, SUMO and H3K9me3, further validating our results from human U2OS cells (Supplementary Fig. S2C and S2D). In contrast, ZMYM2 cluster 2 shows little TRIM28 binding and higher levels of ATAC-seq and H3K18ac signal (**Fig. 2A** and **B**; Supplementary Fig. S2E), suggestive of more active chromatin. Conversely, binding of the SUMO binding defective ZMYM2(SIM2mut) is strong and SUMO occupancy is weaker suggesting that ZMYM2 activity at this cluster is independent of its SUMO binding activity. Example loci demonstrating the unique chromatin features associated with ZMYM2 binding regions in cluster 1 and cluster 2 are illustrated in **Fig. 2C**.

Next we searched for over-represented binding motifs in each ZMYM2 cluster and found numerous significantly enriched motifs which differed between clusters 1 and 2 (**Fig. 3A**). Interestingly, several of the motifs in cluster 2 are the same as in the peaks bound by ADNP (**Fig. 1D**), including motifs for HBP1 and CTCFL/BORIS. The HBP1 motif was also previously observed as a preferred binding motif for the paralogue ZMYM3 (Partridge et al., 2020). Given the similarity in binding motifs, we superimposed the ADNP ChIP-seq signal onto the ZMYM2 clusters and found highest ADNP occupancy in cluster 2 as predicted from the motif analysis (**Fig. 3B** and **C**). Cluster 3 ZMYM2 peaks showed enrichment of a different set of DNA binding motifs with ASCL2 and ETS transcription factor binding motifs figuring prominently (Supplementary Figure S2F).

Differences were also observed in genomic location distributions of each cluster with cluster 1 peaks showing a predominantly intergenic or intronic distribution, whereas cluster 2 and to a lesser extent, cluster 3 peaks were more commonly located in promoter-proximal regions (**Fig. 3D**; supplementary Fig. S2G).

Together these results reveal two classes of ZMYM2 binding region with distinct molecular features. One is SUMO-dependent and is associated with inactive chromatin and TRIM28 co-binding. The second is SUMO-independent and is associated with more active chromatin and ADNP binding. Each type of region is characterised by enrichment of distinct DNA motifs, as well as different genomic features.

ZMYM2 physically and functionally interacts with TRIM28

Next, we wanted to further understand the potential physical and functional interactions between ZMYM2 and TRIM28 implied from both RIME and chromatin co-occupancy in cluster 1 regions. First, we attempted to co-precipitate ZMYM2 and TRIM28 but were unable to detect co-binding under standard conditions (**Fig. 4A**, lanes 4-6; Supplementary Fig. S3A). However, cluster 1 is also enriched for SUMO occupancy but depleted for binding of ZMYM2 containing a mutation in a critical SUMO interacting motif (SIM). The latter mutant was previously used to demonstrate the requirement for SUMO binding for ZMYM2 recruitment to chromatin ([Aguilar-Martinez et al., 2015](#)). We therefore repeated the co-precipitation experiment in the presence of NEM (an isopeptidase inhibitor and critical to preserve protein SUMOylation) and observed ZMYM2 binding to TRIM28 (**Fig. 4A**, lanes 1-3; Supplementary Fig. S3A). Indeed, NEM addition stabilised high molecular weight SUMO conjugates and revealed the binding of ZMYM2 to SUMOylated protein(s) (Supplementary Fig. S3B). Further evidence for SUMO-dependent interactions was derived from co-IP studies where the SUMO binding defective ZMYM2(SIM2mut) showed weaker interactions with ZMYM2 than wild-type ZMYM2 (Supplementary Fig. S3C). Interactions between ZMYM2 and




Fig. 2.

ZMYM2 interactions with molecularly distinct chromatin regions.

(A) Heatmaps showing the signals of the indicated proteins or chromatin marks from ChIP-seq experiments or the ATAC-seq signal in U2OS cells plotted across a 10 kb region surrounding the centres (arrowed) of the wild-type (WT) ZMYM2 binding regions. Clustering of the data produced 3 clusters. (B) Tag density plots of the indicated ChIP-seq or ATAC-seq signals in the three clusters of ZMYM2 binding regions. (C) UCSC genome browser of the indicated ChIP-seq and ATAC-seq data on example genomic loci from clusters 1 and 2.




Fig. 3.

Characterisation of ZMYM2 chromatin binding regions.

(A) *De novo* motif analysis of cluster 1 and 2 regions. The top five most significantly enriched motifs are shown with motif similarity to the indicated protein shown in brackets. (B) Heatmaps showing the signals of the indicated proteins or chromatin marks from ChIP-seq experiments in U2OS cells plotted across a 10 kb region surrounding the centres (arrowed) of the wild-type (WT) ZMYM2 binding regions. Clustering was retained from [Fig. 2A](#) and ADNP signal superimposed on top of this. The number and percentage of ZMYM2 peaks in each cluster overlapping with ADNP2 peaks is shown on the right. (C) Tag density plot of ADNP ChIP-seq signal from U2OS cells across a 10 kb region surrounding the centres (arrowed) of the three clusters of ZMYM2 binding regions. (D) Distribution of binding regions among different genomic categories for clusters 1 and 2 and the entire genome.

TRIM28 in a cellular context were further validated using PLA (**Fig. 4B**). These findings are consistent with the SUMO-dependent enrichment of TRIM28 occupancy on chromatin in ZMYM2 cluster 1 (**Fig. 2A**) and this shared binding profile was validated by ChIP-qPCR (Supplementary Fig. S3D). This co-binding suggests that ZMYM2 might be responsible for recruiting TRIM28 to chromatin but we found no evidence for this as ZMYM2 depletion did not affect TRIM28 binding to a panel of its binding regions (Supplementary Fig. S3E). We also further validated our RIME experiments and examined whether interactions with the ZMYM2 paralog ZMYM3 could be verified. Co-immunoprecipitation detected co-binding of TRIM28 with the ZMYM2 related protein ZMYM3 (Supplementary Fig. S3F) indicating that these interactions extend to other ZMYM2 paralogs.

The physical interactions between ZMYM2 and TRIM28, and their genomic co-occupancy, suggested that they might functionally interact. We therefore depleted each one in turn (Supplementary Fig. S3G) and performed RNAseq analysis to identify their target gene repertoires. For TRIM28, roughly equal numbers of genes were up and downregulated, and a similar phenomenon was observed for ZMYM2 following depletion, albeit more skewed to downregulation in the case of ZMYM2 (**Fig. 4C**; Supplementary Table S4). However, there was a substantial and highly significant overlap between ZMYM2 and TRIM28 regulated genes irrespective of the directionality (**Fig. 4C**). Conversely, in contrast to these similarities, gene regulation showing reciprocal directionality following either TRIM28 or ZMYM2 depletion showed a very small and insignificant overlap (Supplementary Fig. S3H). This is further emphasised by a scatter plot depicting this data (Supplementary Fig. S3I). Thus, functionally there is a strong concordance between the gene regulatory events controlled via ZMYM2 and TRIM28, although the large numbers of genes directionally co-regulated by these two proteins (ie either positively or negatively) indicates no particular preference for activation or repressive activity. Gene ontology (GO) analysis revealed different pathways associated with genes up and downregulated by both ZMYM2 and TRIM28 (**Fig. 4D**). Terms such as "ossification" and "skeletal system development" were associated with the downregulated genes suggesting a disassembly of the core transcriptome associated with the cell identity (U2OS cells are derived from osteosarcomas) whereas upregulated genes contributed to more general GO terms, mainly associated with metabolism and cell responses such as "positive regulation of phosphorylation".

Together these results demonstrate that ZMYM2 and TRIM28 interact physically and functionally to control gene expression. However, these gene regulatory properties create a conundrum, as there is no obvious directionality in their effects, despite the fact that the proteins co-occupy regions of the chromatin with inactive/repressive features.

ZMYM2 regulates protein coding genes from a distance

To further explore how ZMYM2 and TRIM28 work together to control gene expression, we tried to associate ZMYM2 cluster 1 peaks with genes commonly up- or down-regulated following ZMYM2 and TRIM28 depletion by using the nearest gene model. However, very few of these peaks are associated with the co-regulated genes identified from RNA-seq analysis (Supplementary Fig. S4A). This suggested that ZMYM2 might orchestrate gene activity from a distance. We further investigated this possibility by comparing the number of genes deregulated following ZMYM2 depletion that are associated with ZMYM2 peaks across the entire ZMYM2 binding dataset. Very few differentially expressed genes are found within 20 kb of a ZMYM2 peak (**Fig. 5A**) but there is an increasing association as the window size is increased. The distance-dependent distribution of ZMYM2 binding regions is virtually the same as a randomly selected set of genes when considering genes that are downregulated following ZMYM2 depletion (**Fig. 5A**, right). However, in contrast, significantly more differentially upregulated genes are associated with ZMYM2 peaks across all of the peak-gene distance brackets compared to a control set of genes (**Fig. 5A**, left). This suggests a more direct role for ZMYM2 in transcriptional repression, albeit at a distance to target genes. Indeed when we focussed on ZMYM2 cluster 1 peaks (ie potentially ZMYM2-TRIM28 co-regulated) and the genes whose expression increased following either TRIM28 or ZMYM2




Fig. 4.

ZMYM2 interactions with TRIM28.

(A) Co-immunoprecipitation analysis of TRIM28 with ZMYM2. Immunoprecipitation (IP) was performed with ZMYM2 or control IgG antibody from U2OS cells and the resulting proteins detected by immunoblotting (IB) with the indicated antibodies. 10% input is shown (See Supplementary Fig. S3A for longer exposure). NEM was added to the extracts where indicated. (B) PLA assay of interactions between Myc-tagged ZMYM2 and endogenous TRIM28. Assays were carried out in U2OS cells transfected with a vector encoding Myc-ZMYM2 or left untransfected (UT). The addition of anti-Myc and -TRIM28 antibodies (ab) is indicated. Nuclei were stained with DAPI. Average numbers of foci per cell and numbers of cells are indicated (bottom). (C) Venn diagram showing the overlaps in downregulated (left) and upregulated (right) genes (Padj<0.05; fold change 2:1.6) from RNAseq analysis in U2OS cells treated with siRNAs against *ZMYM2* or *TRIM28*. (D) Enriched GO terms of genes commonly upregulated and downregulated by ZMYM2 and TRIM28 depletion.

depletion (ie repressed by these factors), we found a more significant distance-based association of ZMYM2 peaks with upregulated rather than downregulated genes (**Fig. 5B**). This is consistent with the observation that cluster 1 is predicted to be repressive in nature. To further explore this association, we examined the expression of all ZMYM2/TRIM28 co-regulated genes located within the TADs containing ZMYM2 binding peaks according to the three molecularly distinct clusters we identified. For this purpose we used TADs from human ESCs (Dixon et al., 2012) on the well-established assumption that the majority of these will be conserved across cell types (Stadhouders et al., 2018). TADs containing cluster 1 binding regions contained a significantly higher number of upregulated genes compared to downregulated genes compared to TADs which lacked ZMYM2 binding regions (**Fig. 5C**). Although there was a weak tendency for more upregulated genes with TADs containing cluster 2 peaks, this was insignificant. A similar trend was observed if genes were analysed based on their response to individual depletion of ZMYM2 or TRIM28, with TADs containing cluster 1 peaks being more associated with upregulated genes in both cases (Supplementary Fig. S4B). Moreover, we also tested whether a gene is more likely to be negatively regulated by ZMYM2 when located in the same TAD as a cluster 1 peak rather than being located in other TADs lacking such peaks. The observed frequency of co-localisation was significantly higher than by chance (Supplementary Fig. S4C). Although there is an association of ZMYM2 peaks with ZMYM2 regulated genes in the same TAD, we asked whether ZMYM2 might affect gene expression at a certain distance irrespective of whether the TSS of the gene is in the same TAD. To test this we took all of the ZMYM2 regions associated with genes upregulated by ZMYM2 depletion that resided in the same TAD and calculated the peak to TSS distance. Then we searched in the opposite direction for the TSS of genes at a similar distance ($\pm 25\%$) that resided in an adjacent TAD. We then asked whether these genes were upregulated by ZMYM2 depletion. 102 ZMYM2 peaks were positioned within these distance constraints with at least one gene in an adjacent TAD (716 genes in total). Of these genes, only 11 were upregulated following ZMYM2 depletion. There is therefore not a general spreading of transcriptional deregulation around ZMYM2 peaks in a distance-dependent manner. We further probed the association with TAD boundaries by comparing the distances of the ZMYM2 peaks in each of the clusters to TAD boundaries with randomly selected genomic regions. In all cases, the ZMYM2 peaks were significantly closer to a TAD boundary than expected by chance (**Fig. 5D**).

Together, these results support a role for cluster 1 ZMYM2 peaks in gene repression, and suggest a more region-specific role rather than a simple peak-to-gene association typically found with transcription factors.

ZMYM2 regulates retrotransposon elements

Several recent studies indicate a potential role for repetitive elements in controlling gene expression by demarcating TAD boundaries (reviewed in Haws et al., 2022) exemplified by the human endogenous retrovirus subfamily H (HERV-H) (Zhang et al., 2019). Zinc finger transcription factors, chiefly of the KRAB domain-containing subclass, have been associated with repetitive element repression (reviewed in Bruno et al., 2019). We therefore asked whether ZMYM2 binding regions mapped to any repetitive elements. All three clusters of ZMYM2 binding regions showed a high degree of overlap with repetitive elements, with 80% of the regions in cluster 1 showing close proximity to these elements (**Fig. 6A**). However, the distribution of retrotransposon classes differed among ZMYM2 clusters, with cluster 3 and cluster 1 in particular favouring long terminal repeats (LTRs) whereas cluster 2 regions are mainly associated with short interspersed nuclear elements (SINE). The latter is consistent with the higher levels of ADNP binding found in cluster 2 regions, as ADNP has been implicated in SINE control (Kaaij et al., 2019). We further examined the distribution of subtypes of LTR elements in clusters 1 and 3 and found that cluster 3 is dominated by endogenous retrovirus (ERV)-L elements whereas cluster 1 is dominated by ERV-1 elements (**Fig. 6B**). An example locus illustrates ZMYM2 and TRIM28 binding to a MER11A ERV-1 element (**Fig. 6C**, top). Given the association of ZMYM2 with LTRs, we next asked whether ZMYM2 represses their activity as predicted. We depleted ZMYM2 and examined the regulation of LTR expression, and found that the LTRs showed upregulation




Fig. 5.

ZMYM2 location and gene expression.

(A) Numbers of differentially expressed genes following ZMYM2 depletion (left upregulated, right downregulated) whose TSS lies within the indicated distances of a ZMYM2 binding peak (black bars). Control sets of equal numbers of randomly selected genes are shown for comparison (grey bars showing the average of 10 datasets). P-values $* = <0.05$, $** = <0.01$, $*** = <0.001$. (B) Distance-dependent association of ZMYM2 binding regions with differentially regulated genes following depletion of ZMYM2 or TRIM28. The numbers in the boxes are the number of genes among the input gene sets (x-axis) that overlap with the ZMYM2 peaks (cluster 1 peaks only) at the indicated distances to TSS (y-axis) and the colour shows $-\log_{10}$ of p-value (Hypergeometric test). (C) Relative proportion of genes commonly up- or down-regulated following ZMYM2 or TRIM28 depletion in TADs which also contain ZMYM2 peaks from the indicated clusters or have no ZMYM2 peak in the same TADs. Significance relative no regions containing no peaks is shown P-value, $*** = <0.001$; ns= non-significant. (D) Boxplots of the relative distance of ZMYM2 binding regions from each of the clusters from TAD boundaries compared to two different control sets of randomly selected regions ($n=2360$). Statistical significance of cluster 1-3 distances compared to each of the control regions is shown ($** = P\text{-value} <0.05$ in all cases; student t-test).

following ZMYM2 loss, consistent with a repressive activity (**Fig. 6D** [↗](#); Supplementary Fig. S5A). We also examined the distribution of subclasses of SINEs associated with cluster 2 ZMYM2 peaks and found equal numbers of Alu repeats and MIR elements (**Fig. 6E** [↗](#)). Importantly, depletion of either ZMYM2 (**Fig. 6F** [↗](#)) or ADNP (Supplementary Fig. S5C) both caused increased activity of several SINEs, consistent with our designation of these retroviral elements as co-regulated by ZMYM2 and ADNP from our ChIP-seq analysis (eg **Fig. 6C** [↗](#), bottom). We also examined our RNA-seq data for more widespread activation of transposable elements beyond those we have tested, but were unable to detect expression of large numbers of additional elements. This may reflect a more limited effect of ZMYM2 on a subset of elements or that our data was of insufficient depth and quality to detect their expression.

Collectively, these results show that ZMYM2 associates with LTR containing regions, and promotes their repression in combination with the corepressor protein TRIM28 whereas it works alongside ADNP to control SINE element expression. Coupled with the observation that ZMYM2 binding is associated with regional control of gene expression and that retrotransposons can participate in indirect control of gene expression by affecting genome topology, this provides a plausible mechanism through which ZMYM2 impacts widely on coding gene expression.

Discussion

ZMYM2 has been established as an important regulator of cell fate identity and commitment, particularly at the pluripotency stage. Genetic disruptions of ZMYM2 function have been linked to disease. Here, we provide further insights into the gene regulatory activities of ZMYM2 and demonstrate its physical and functional association with different corepressor complexes containing either ADNP or TRIM28 (**Fig. 6G** [↗](#)).

We identify three distinct chromatin regions where ZMYM2 binds. One has chromatin characteristics related to transcriptional repression, one appears to be associated with more active chromatin regions and a third set of regions has no distinguishing chromatin features. It remains unclear what function, if any, ZMYM2 has at the latter set of binding regions. Our main focus was on cluster 1 ZMYM2 binding regions that are also bound by a novel binding partner, TRIM28. Interestingly, ZMYM2 binding to TRIM28 is SUMO-dependent, as retention of SUMOylation is required to detect binding by co-IP and ZMYM2 binding to cluster 1 regions require the SUMO interaction motifs in ZMYM2. This is consistent with the preponderance of SUMO at cluster 1 binding regions. It is not currently clear what SUMO modified protein(s) are the direct binding target of ZMYM2, although ZMYM2 is itself SUMOylated ([Kunapuli et al., 2006](#) [↗](#)), as is TRIM28 ([Lee et al., 2007](#) [↗](#)), which can also act as a SUMO E3 ligase ([Ivanov et al., 2007](#) [↗](#)). Earlier studies may have missed the ZMYM2-TRIM28 interactions by performing ZMYM2 immunoprecipitations in the absence of SUMO stabilising agents or sample fixation as we used in our initial RIME experiments. However, ZMYM2 was identified in reciprocal co-IP experiments with TRIM28, as a glycosylated interaction partner ([Boulard et al., 2020](#) [↗](#)) suggesting that multiple post-translational modifications govern the assembly and function of this complex.

Previous studies have focussed on the role of ZMYM2 in the context of its association with the LSD1-CoREST-HDAC corepressor complex ([Hakimi et al., 2003](#) [↗](#); [Shi et al., 2005](#) [↗](#); [Gocke and Yu, 2008](#) [↗](#)). Indeed, we identify all three of these proteins in our proteomics dataset (Supplementary Table S3), further substantiating this functional interaction. Here we find a broader occurrence of ZMYM2 in chromatin binding complexes, both as part of the ChAHP complex centred on ADNP ([Ostapcuk et al., 2018](#)) and also independently in a complex with TRIM28. The latter observation is interesting, as TRIM28 has been shown to act as a widespread repressor of retrotransposons where it is recruited by KRAB domain containing zinc finger proteins (reviewed in [Bruno et al., 2019](#) [↗](#)). ZMYM2 is also a zinc finger protein but lacks the KRAB domain but still binds to TRIM28, suggesting that a broader swathe of zinc finger proteins may be involved in repetitive element




Fig. 6.

ZMYM2 functionally associates with ERV repetitive elements.

(A) Percentage of the ZMYM2 binding sites in each of the indicated clusters containing each of the indicated classes of retrotransposon elements. The genome-wide proportion of genomic regions containing each type of the retrotransposon elements is also shown. (B) Proportions of LTR subclasses in cluster 1 and cluster 3 ZMYM2 binding regions. (C) UCSC genome browser view of a MER11A ERV1 LTR element located upstream of the *HEATR6* locus, illustrating co-binding of ZMYM2 and TRIM28 (top) and a MIR SINE element located upstream of the *EIF3D* locus, illustrating co-binding of ZMYM2 and ADNP. (D) RT-qPCR analysis of expression of ZMYM2 and the indicated LTR elements following ZMYM2 depletion or control non-targeting (NT) siRNA treatment. Individual paired experiments are shown (n=4; P-values *=<0.05, ****=<0.0001). (E) Proportions of SINE subclasses in cluster 1-3 ZMYM2 binding regions. (F) RT-qPCR analysis of expression of ZMYM2 and the indicated SINE elements following ZMYM2 depletion or control non-targeting (NT) siRNA treatment. Individual paired experiments are shown (n=3; unpaired T-test P-values *=<0.05, **=<0.01, ****=<0.0001). (G) Model illustrating the two distinct complexes through which ZMYM2 functions on chromatin to control retrotransposon expression.

epigenetic repression. However, our results rule out a simple model that ZMYM2 is responsible for the recruitment of TRIM28 to chromatin. Whether KRAB-zinc finger proteins are also involved in ZMYM2 complexes and act as the relevant recruitment factor is currently unknown although we identified four such proteins in our lower confidence proteomics dataset, with ZNF202 being the most prominent consistently detected protein (Supplementary Table S3). It is also not clear what the chromatin recruitment mechanism is, although recent studies do not rule out a direct role for TRIM28 in the process as it has been shown to be able to engage directly with unmodified histone H4 tails (Bacon et al., 2020 [DOI](#)).

The identification of TRIM28 as a chromatin-associated interaction partner further supports a role for ZMYM2 in transcriptional repression, and suggests that it can do so via a variety of different means. Indeed a recent study identified ZMYM2 as an important regulator of silencing of imprinting control regions which are under the control of the zinc finger protein ZFP57 (Butz et al., 2022 [DOI](#)). In terms of global transcriptional regulation, the role of ZMYM2 is more ambivalent as we identified substantial numbers of genes that are either up- or down-regulated. However, ZMYM2 binding regions are generally more associated with genes that it represses rather than activates, suggesting that the latter may be indirectly affected. In contrast to TRIM28 co-bound regions, ZMYM2 also binds to chromatin regions occupied by ADNP, where more active characteristics are found. ZMYM2 functions in a repressive manner rather than in gene activation in the latter context and suggests that it dampens down the activity of a region rather than the complete repression which would be mediated through TRIM28 and its coregulatory partners.

The association of ZMYM2 with transposons and its ability to regulate their expression suggests several possible modes of subsequent effects on broader gene expression programmes. Retrotransposons have been shown to exhibit a multitude of different molecular mechanisms to control gene regulation (reviewed in Fueyo et al., 2022 [DOI](#)). For example, the ERV-H LTRs demarcate TADs in human pluripotent stem cells (Zhang et al., 2019), and it is possible that ZMYM2-TRIM28 complexes control ERV LTR activity which in turn influence local chromatin topology. This would help explain how ZMYM2 appears to regulate gene expression in a locally defined region rather than one ZMYM2 binding event acting to regulate expression of one specific gene. This might apply in particular to the ZMYM2-ADNP co-bound regions, as ADNP has previously been shown to play an active role in controlling chromatin domain boundary elements at SINE B2 transposable elements by blocking CTCF binding (Kaaij et al., 2019 [DOI](#)). It is also noteworthy that ZMYM2 interacts with the GTF3C complex which has previously been implicated in controlling retrotransposon transcription (Ferrari et al., 2020 [DOI](#)). Thus, ZMYM2 may contribute important regulatory activities to many different chromatin complexes associated with transposable elements. However, further extensive work will be needed to address the precise interplay between ZMYM2, retrotransposon activity, and local chromatin structure in determining ZMYM2-dependent gene regulatory outcomes.

In summary, our work adds to the growing repertoire of ZMYM2 associated chromatin regulatory proteins and complexes and provides a molecular framework for understanding the role of this protein in cell fate commitment and maintenance.

Acknowledgements

We thank Karren Palmer, Mairi Challinor and Guanhua Yan, for excellent technical assistance; Ian Donaldson, Ping Wang, Rachel Scholey and Leo Zeef in the Bioinformatics core Facility; Stacey Holden, Michal Smiga and Andy Hayes in Genomic Technologies Core Facility; staff in the Mass spectrometry and Bioimaging facilities, Nicoletta Bobola, Shen-Hsi Yang and members of our laboratory for comments on the manuscript and stimulating discussions. This work was supported by the Wellcome Trust (103857/Z/14/Z) and BBSRC (BB/V000403/1).

Author contributions

DO, EAM and ZJ performed the experiments and data analysis in this study. YL performed data analysis. ADS led the study. All authors contributed to manuscript preparation and/or critically appraised manuscript drafts.

References

- Aguilar-Martinez E, Chen X, Webber A, Mould AP, Seifert A, Hay RT, Sharrocks AD (2015) **Screen for multi-SUMO-binding proteins reveals a multi-SIM-binding mechanism for recruitment of the transcriptional regulator ZMYM2 to chromatin** *Proc Natl Acad Sci U S A* **112**:E4854–63
- Bacon CW *et al.* (2020) **KAP1 Is a Chromatin Reader that Couples Steps of RNA Polymerase II Transcription to Sustain Oncogenic Programs** *Mol Cell* **78**:1133–1151
- Boulard M, Rucli S, Edwards JR, Bestor TH (2020) **Methylation-directed glycosylation of chromatin factors represses retrotransposon promoters** *Proc Natl Acad Sci U S A* **117**:14292–14298
- Briggs P, Hunter AL, Yang SH, Sharrocks AD, Iqbal M (2021) **PEGS: An efficient tool for gene set enrichment within defined sets of genomic intervals** *F1000Res* **10**
- Brüninghoff K, Aust A, Taupitz KF, Wulff S, Dörner W, Mootz HD (2020) **Identification of SUMO Binding Proteins Enriched after Covalent Photo-Cross-Linking** *ACS Chem Biol* **15**:2406–2414
- Bruno M, Mahgoub M, Macfarlan TS (2019) **The arms race between KRAB-Zinc finger proteins and endogenous retroelements and its impact on mammals** *Annu Rev Genet* **53**:393–416
- Butz S *et al.* (2022) **DNA sequence and chromatin modifiers cooperate to confer epigenetic bistability at imprinting control regions** *Nat Genet* **54**:1702–1710
- Chen X, Ji Z, Webber A, Sharrocks AD (2016) **Genome-wide binding studies reveal DNA binding specificity mechanisms and functional interplay amongst Forkhead transcription factors** *Nucleic Acids Res* **44**:1566–78
- Cibis H, Biyanee A, Dörner W, Mootz HD, Klempnauer KH (2020) **Characterization of the zinc finger proteins ZMYM2 and ZMYM4 as novel B-MYB binding proteins** *Sci Rep* **10**
- Connaughton DM *et al.* (2020) **Mutations of the Transcriptional Corepressor ZMYM2 Cause Syndromic Urinary Tract Malformations** *Am J Hum Genet* **107**:727–742
- Cossec JC *et al.* (2018) **SUMO Safeguards Somatic and Pluripotent Cell Identities by Enforcing Distinct Chromatin States** *Cell Stem Cell* **23**:742–757
- Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B (2012) **Topological domains in mammalian genomes identified by analysis of chromatin interactions** *Nature* **485**:376–80
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) **STAR: ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21
- Ferrari R *et al.* (2020) **TFIIIC Binding to Alu Elements Controls Gene Expression via Chromatin Looping and Histone Acetylation** *Mol Cell* **77**:475–487
- Fueyo R, Judd J, Feschotte C, Wysocka J (2022) **Roles of transposable elements in the regulation of mammalian transcription** *Nat Rev Mol Cell Biol* **23**:481–497

- Garcia-Dominguez M, Reyes JC (2009) **SUMO association with repressor complexes, emerging routes for transcriptional control** *Biochim Biophys Acta* **1789**:451–9
- Gocke CB, Yu H (2008) **ZNF198 stabilizes the LSD1-CoREST-HDAC1 complex on chromatin through its MYM-type zinc fingers** *PLoS One* **3**
- Guzzo CM, Ringel A, Cox E, Uzoma I, Zhu H, Blackshaw S, Wolberger C, Matunis MJ (2014) **Characterization of the SUMO-binding activity of the myeloproliferative and mental retardation (MYM)-type zinc fingers in ZNF261 and ZNF198** *PLoS One* **9**
- Hakimi MA, Dong Y, Lane WS, Speicher DW, Shiekhatair R (2003) **A candidate X-linked mental retardation gene is a component of a new family of histone deacetylase-containing complexes** *J Biol Chem* **278**:7234–9
- Heinz S., Benner C., Spann N., Bertolino E., Lin Y.C., Laslo P., Cheng J.X., Murre C., Singh H., Glass C.K. (2010) **Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B-cell identities** *Mol. Cell* **38**:576–589
- Ivanov AV *et al.* (2007) **Rauscher FJ 3rd. PHD domain-mediated E3 ligase activity directs intramolecular sumoylation of an adjacent bromodomain required for gene silencing** *Mol Cell* **28**:823–37
- Jensen LJ *et al.* (2009) **STRING 8--a global view on proteins and their functional interactions in 630 organisms** *Nucleic Acids Res* **37**:D412–6
- Ji Z, Li Y, Liu SX, Sharrocks AD (2021) **The forkhead transcription factor FOXK2 premarks lineage-specific genes in human embryonic stem cells for activation during differentiation** *Nucleic Acids Res* **49**:1345–1363
- Kaaij LJ, Mohn F, van der Weide RH, de Wit E, Bühler M (2019) **The ChAHP Complex Counteracts Chromatin Looping at CTCF Sites that Emerged from SINE Expansions in Mouse** *Cell* **178**:1437–1451
- Kundaje A *et al.* (2015) **Integrative analysis of 111 reference human epigenomes** *Nature* **518**:317–30
- Kunapuli P, Kasyapa CS, Chin SF, Caldas C, Cowell JK (2006) **ZNF198, a zinc finger protein rearranged in myeloproliferative disease, localizes to the PML nuclear bodies and interacts with SUMO-1 and PML** *Exp Cell Res* **312**:3739–51
- Langmead, B., and Salzberg, S (2012) **Fast gapped-read alignment with Bowtie 2** *Nature Methods* **9**:357–359
- Lawrence M, Theunissen TW, Lombard P, Adams DJ, Silva JCR (2019) **ZMYM2 inhibits NANOG-mediated reprogramming** *Wellcome Open Res* **4**
- Lee YK, Thomas SN, Yang AJ, Ann DK (2007) **Doxorubicin down-regulates Kruppel-associated box domain-associated protein 1 sumoylation that relieves its transcription repression on p21WAF1/CIP1 in breast cancer MCF-7 cells** *J Biol Chem* **282**:1595–606
- Lee D, Apelt K, Lee SO, Chan HR, Luijsterburg MS, Leung JWC, Miller KM (2022) **ZMYM2 restricts 53BP1 at DNA double-strand breaks to favor BRCA1 loading and homologous recombination** *Nucleic Acids Res* **50**:3922–3943

- Lezmi E, Weissbein U, Golan-Lev T, Nissim-Rafinia M, Meshorer E, Benvenisty N (2020) **The Chromatin Regulator ZMYM2 Restricts Human Pluripotent Stem Cell Growth and Is Essential for Teratoma Formation** *Stem Cell Reports* **15**:1275–1286
- Lüdtke TH, Kleppa MJ, Rivera-Reyes R, Qasrawi F, Connaughton DM, Shril S, Hildebrandt F, Kispert A (2022) **Proteomic analysis identifies ZMYM2 as endogenous binding partner of TBX18 protein in 293 and A549 cells** *Biochem J* **479**:91–109
- McLean C.Y., Bristor D., Hiller M., Clarke S.L., Schaar B.T., Lowe C.B., Wenger A.M., Bejerano G. (2010) **GREAT improves functional interpretation of cis-regulatory regions** *Nat. Biotechnol* **28**:495–501
- Ma L, Xie D, Luo M, Lin X, Nie H, Chen J, Gao C, Duo S, Han C (2022) **Identification and characterization of BEND2 as a key regulator of meiosis during mouse spermatogenesis** *Sci Adv* **8**
- Mohammed H *et al.* (2013) **Endogenous purification reveals GREB1 as a key estrogen receptor regulatory factor** *Cell Rep* **3**:342–9
- Ostapczuk V *et al.* (2018) **Activity-dependent neuroprotective protein recruits HP1 and CHD4 to control lineage-specifying genes** *Nature* **557**:739–743
- Reiter A *et al.* (1998) **Consistent fusion of ZNF198 to the fibroblast growth factor receptor-1 in the t(8;13)(p11;q12) myeloproliferative syndrome** *Blood* **92**:1735–42
- Robinson M.D., McCarthy D.J., Smyth G.K. (2010) **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data** *Bioinformatics* **26**:139–140
- Seah MKY *et al.* (2019) **The KRAB-zinc-finger protein ZFP708 mediates epigenetic repression at RMER19B retrotransposons** *Development* **146**
- Shi YJ, Matson C, Lan F, Iwase S, Baba T, Shi Y (2005) **Regulation of LSD1 histone demethylase activity by its associated factors** *Mol Cell* **19**:857–64
- Stadhouders R *et al.* (2018) **Transcription factors orchestrate dynamic interplay between genome topology and gene regulation during cell reprogramming** *Nat Genet* **50**:238–249
- Toh CX *et al.* (2016) **RNAi Reveals Phase-Specific Global Regulators of Human Somatic Cell Reprogramming** *Cell Rep* **15**:2597–607
- Xiao S, Nalabolu SR, Aster JC, Ma J, Abruzzo L, Jaffe ES, Stone R, Weissman SM, Hudson TJ, Fletcher JA (1998) **FGFR1 is fused with a novel zinc-finger gene, ZNF198, in the t(8;13) leukaemia/lymphoma syndrome** *Nat Genet* **18**:84–7
- Xiao S, McCarthy JG, Aster JC, Fletcher JA (2000) **ZNF198-FGFR1 transforming activity depends on a novel proline-rich ZNF198 oligomerization domain** *Blood* **96**:699–704
- Yang SH, Andrabi M, Biss R, Murtuza Baker S, Iqbal M, Sharrocks AD (2019) **ZIC3 Controls the Transition from Naive to Primed Pluripotency** *Cell Rep* **27**:3215–3227
- Yang F *et al.* **DUX-miR-344-ZMYM2-mediated activation of MERVL LTRs induces a totipotent 2C-like State** :234–250

Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda SK (2019) **Metascope provides a biologist-oriented resource for the analysis of systems-level datasets** *Nat Commun* 10

Article and author information

Danielle Owen

Faculty of Biology, Medicine and Health, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, UK

Elisa Aguilar-Martinez

Faculty of Biology, Medicine and Health, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, UK

Zongling Ji

Faculty of Biology, Medicine and Health, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, UK

Yaoyong Li

Faculty of Biology, Medicine and Health, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, UK

Andrew D. Sharrocks

Faculty of Biology, Medicine and Health, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, UK

For correspondence: a.d.sharrocks@manchester.ac.uk

Copyright

© 2023, Owen et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #1 (Public Review):

Owen D et al. investigated the protein partners and molecular functions of ZMYM2, a transcriptional repressor with key roles in cell identity and mutated in several human diseases, in human U2OS cells using mass spectrometry, siRNA knockdown, ChIP-seq and RNA-seq. They tried to identify chromatin bound complexes containing ZMYM2 and

identified known and novel protein partners, including ADNP and the newly described partner TRIM28. Focusing mainly on these two proteins, they show that ZMYM2 physically interacts with ADNP or TRIM28, and co-occupies an overlapping set of genomic regions with ADNP and TRIM28. By generating a large set of knockdown and RNA-seq experiments, they show that ZMYM2 co-regulates a large number of genes with ADNP and TRIM28 in U2OS cells. Interestingly, ZMYM2-TRIM28 do not appear to repress genes directly at promoters, but the authors find that ZMYM2/TRIM28 repress LTR elements and suggest that this leads to gene deregulation at distance by affecting the chromatin environment within TADs.

A strength of the study is that, compared to previous studies of ZMYM2 protein partners, it investigates binding partners of ZMYM2 using the RIME method on chromatin. The RIME method makes it possible to identify low-affinity protein-protein interactions and proteins interactions occurring at chromatin, therefore revealing partners most relevant for gene regulation at chromatin. This allowed the identification of novel ZMYM2 partners not identified before, such as TRIM28.

The authors present solid interaction data with appropriate controls and generated an impressive amount of datasets (ChIP-seq for TRIM28 and ADNP, RNA-seq in ZMYM2, ADNP and TRIM28 knockdown cells) that are important to understand the molecular functions of ZMYM2. These datasets were generated with replicates and will be very useful for the scientific community. This study provides important novel insights into the molecular roles of ZMYM2 in human U2OS cells.

- <https://doi.org/10.7554/eLife.86669.2.sa2>

Reviewer #2 (Public Review):

In this study the authors investigate functional associations made by transcription factor ZMYM2 with chromatin regulators, and the impact of perturbing these complexes on the transcriptome of the U2OS cell line. They focus on validating two novel chromatin-templated interactions: with TRIM28/KAP1 and with ADNP, concluding that via these distinct chromatin regulators, ZMYM2 contributes to transcriptional control of LTR and SINE retrotransposons, respectively.

Strengths of the study:

-The co-localization of ZMYM2 with ADNP and TRIM28 is validated through RIME, ChIP-seq and co-IP. Since TRIM28 is a highly abundant nuclear protein, the use of multiple methods is important to add confidence in particular for the novel (SUMO-dependent interaction identified between ZMYM2 and TRIM28. That TRIM28 pulls down less of the ZMYM2-SIM mutant is reassuring.

-It is good that uniquely-mapped reads are used in the ChIP-seq analysis given the interest in repetitive elements. Likewise, though the RT-qPCR data in Fig 6 should be complemented by analysis of the RNA-seq data that the authors already have, it seems that the primers are carefully designed such that a single retrotransposon copy is amplified.

-The paper is generally written very clearly, the experiments well done and the different datasets appear to be robust.

Weaknesses of the study:

-The transcriptional response using bulk RNA-seq in ZMYM2-depleted cells remains gene-centric despite the title of the paper being about TE transcription. In fact, the only panels about TE transcription are the RT-qPCR data in Fig 6D, F. During the revision the authors said that their RNA-seq data is unfortunately too shallow to retrieve TEs. Fair enough - however, it

remains the case that the central claim is control of TE transcription by ZMYM2. Thus, without additional transcriptomic analysis we are left with only a few qPCRs, even if they are nicely done! Perhaps the title could be modified a bit in that case?

-The mechanism by which ZMYM2 and TRIM28 work together does remain a mystery. Following review the authors performed TRIM28 ChIP on ZMYM2-depleted cells, but identified no changes over three transposons. It remains unclear if H3K9me3 levels are altered.

- <https://doi.org/10.7554/eLife.86669.2.sa1>

Reviewer #3 (Public Review):

ZMYM2 is a transcriptional repressor known to bind to the post-translational modification SUMO2/3. It has been implicated in the silencing of genes and transposons in a variety of contexts, but lacking sequence-specific DNA binding, little is known about how it is targeted to specific regions. At least two reports indicate association with TRIM28 targets (Tsusaka 2020 Epigenetics & Chromatin, Graham-Paquin 2023 NAR) but no physical association with TRIM28 targets had been demonstrated. Tsusaka 2020 theorizes an indirect, potentially SUMO-independent, interaction via ATF7IP and SETDB1.

Here, Owen and colleagues show that a subset of ZMYM2-binding sites in U2OS cells are clearly TRIM28 sites, and further find that hundreds of genes are silenced by both ZMYM2 and TRIM28. They next demonstrate that ZMYM2 homes to chromatin, and interacts with TRIM28, in a SUMOylation-dependent manner, suggesting that ZMYM2 is recognizing SUMOylation on TRIM28 or a protein associated with TRIM28. ZMYM2 separately homes to SINE elements bound by the ChAHP complex in an apparently SUMOylation independent manner. Although this is not the first report to show physical interaction between ZMYM2 and ChAHP, it is the first to show that ZMYM2 homes to ChAHP-binding sites and functions as a corepressor at these sites. Finally the authors demonstrate that ZMYM2 and TRIM28 coregulate genic targets by inducing repression at LTRs within the same TADs as the genes in question.

Overall, the manuscript is well-written, convincing, and fills a significant hole in our understanding of ZMYM2's mechanistic function. The revised version of this manuscript addresses all of my previous concerns well.

- <https://doi.org/10.7554/eLife.86669.2.sa0>

Author Response

The following is the authors' response to the original reviews.

Thank you for your recent editorial decision on our manuscript. I have included a revised version of our manuscript in which we have addressed all of the required editorial and referees' comments as requested. In summary, we have added substantial amounts of new data and analysis (new Fig. 5D; Supplementary Figures S1E, S3C, S3E, S3I, S4C), amended several figures (Figures 2 and 3), added a new supplementary Table (Table S2) and we have changed the text and figure labelling/presentation in appropriate places to clarify or correct the issues raised by the reviewers.

In summary, we firmly believe that we have addressed all the outstanding issues in a positive manner and that the manuscript is now suitable for publication in eLife. I look forward to receiving your final editorial decision on this manuscript.

eLife assessment:

ZMYM2 is a transcriptional corepressor but little was known about how it is recruited to chromatin. This study reveals that ZMYM2 homes to distinct classes of retrotransposons bound by the TRIM28 and ChAHP complexes in human cells, an important finding in the field of transcriptional regulation. The evidence supporting the claims of the authors is solid, although inclusion of more functional data would have strengthened the original model proposed.

We have taken all the comments on board and provided additional new experimental data where requested and more data analysis to substantiate our claims.

Reviewer #1 (Public Review):

Owen D et al. investigated the protein partners and molecular functions of ZMYM2, a transcriptional repressor with key roles in cell identity and mutated in several human diseases, in human U2OS cells using mass spectrometry, siRNA knockdown, ChIP-seq and RNA-seq. They tried to identify chromatin bound complexes containing ZMYM2 and identified known and novel protein partners, including ADNP and the newly described partner TRIM28. Focusing mainly on these two proteins, they show that ZMYM2 physically interacts with ADNP or TRIM28, and co-occupies an overlapping set of genomic regions with ADNP and TRIM28. By generating a large set of knockdown and RNA-seq experiments, they show that ZMYM2 co-regulates a large number of genes with ADNP and TRIM28 in U2OS cells. Interestingly, ZMYM2-TRIM28 do not appear to repress genes directly at promoters, but the authors find that ZMYM2/TRIM28 repress LTR elements and suggest that this leads to gene deregulation at distance by affecting the chromatin environment within TADs.

A strength of the study is that, compared to previous studies of ZMYM2 protein partners, it investigates binding partners of ZMYM2 using the RIME method on chromatin. The RIME method makes it possible to identify low-affinity protein-protein interactions and proteins interactions occurring at chromatin, therefore revealing partners most relevant for gene regulation at chromatin. This allowed the identification of novel ZMYM2 partners not identified before, such as TRIM28. The authors present solid interaction data with appropriate controls and generated an impressive amount of datasets (ChIP-seq for TRIM28 and ADNP, RNA-seq in ZMYM2, ADNP and TRIM28 knockdown cells) that are important to understand the molecular functions of ZMYM2. These datasets were generated with replicates and will be very useful for the scientific community. This study provides important novel insights into the molecular roles of ZMYM2 in human U2OS cells.

The authors could have been more precise in the manuscript title and abstract to emphasize that these findings apply to human cells, as indeed there is no demonstration yet that the findings presented here can be transposed to mouse cells.

We have slightly changed the title and abstract to emphasise that the findings are in human cells.

The manuscript's main conceptual advance is that the authors propose a novel model of gene regulation whereby transcriptional repressors of transposable elements could regulate genes at distance by modulating the local chromatin environment within TADs. Additional experiments would be needed to strengthen this model. For example the authors could have performed TRIM28 ChIP in ZMYM2-kd cells to test if ZMYM2 favors the recruitment of TRIM28 to its genomic targets, as well as ChIP-seq of repressive

chromatin marks (such as H3K9me3) in ZMYM2-kd cells to investigate if the loss of ZMYM2 leads to reduced H3K9me3 in ERVs and over large regions surrounding the ERVs.

We have tested whether ZMYM2 is required for TRIM28 binding at several loci and find no evidence for this (new Supplementary Fig. S3E). We now discuss this in the results text and discussion where we already suggested that TRIM28 is likely recruited by KRAB-zinc finger proteins and ZMYM2 is subsequently recruited to this complex. Future extensive work is required to understand the mechanistic functions of ZMYM2 in these regions.

Reviewer #2 (Public Review):

In this study the authors investigate functional associations made by transcription factor ZMYM2 with chromatin regulators, and the impact of perturbing these complexes on the transcriptome of the U2OS cell line. They focus on validating two novel chromatin-templated interactions: with TRIM28/KAP1 and with ADNP, concluding that via these distinct chromatin regulators, ZMYM2 contributes to transcriptional control of LTR and SINE retrotransposons, respectively.

Strengths and weakness of the study:

- The co-localization of ZMYM2 with ADNP and TRIM28 is validated through RIME, ChIP-seq and co-IP. (Notably, since both RIME and ChIP-seq rely on crosslinking, and the co-IP with TRIM28 required crosslinking due to being SUMO-dependent, only the ZMYM2-ADNP co-IP experiment demonstrates an interaction in the absence of crosslinking).*

This is not correct as the co-IP experiments between endogenous ZMYM2 and TRIM28 were not performed in the presence of cross linkers. They did have NEM added, but this was to inactivate SUMO proteases rather than to cross link proteins.

- It is good that uniquely-mapped reads are used in the ChIP-seq analysis given the interest in repetitive elements. Likewise, though the RT-qPCR data in Fig5 should be complemented by analysis of the RNA-seq data that the authors already have, it seems that the primers are carefully designed such that a single retrotransposon copy is amplified.*

We re-analysed our RNA-seq data using the Tetrascripts tool and looked at TE transcription genome-wide. However very few TEs were expressed at high enough levels to get any statistically significant additional data beyond a few additional transposable elements. This likely results from the relatively low read depth we used and the lack of specific protocols being followed to preserve TE transcripts. We will return to the genome-wide effects in future studies where we plan to switch cell types and will generate more bespoke datasets (the current ones were designed for analysing effects on protein coding gene expression before we made the connection to TEs). We added additional text to the results section to indicate that we could not see widespread deregulation of subclasses of TEs but that this needs further work.

- The top-scoring interactors are highly-abundant nuclear proteins: for example, data from the contaminant repository for affinity purification mass-spec data (<https://reprint-apms.org/>) show that TRIM28 is identified in 466 / 716 AP-MS experiments with a mean spectral count of 16. While this does not indicate that the ZMYM2-TRIM28 interaction is not 'true', it would have been helpful to further dissect the interaction to strengthen this conclusion. For example, it would be nice to see the co-IP (fig 3A) repeated from the cells expressing the ZMYM2 mutant*

that is no longer competent to bind SUMO (used in the ChIP-seq data of Fig 2). Alternatively - if the model is that ZMYM2 recruits SUMOylated TRIM28 with well-characterized TRIM28 mutants that lack SUMOylation.

We are aware that TRIM28 is often present as an apparent contaminant in many mass spec studies. However we have provided co-IP, PLA and ChIP-seq data to support their co-association on chromatin. We also convincingly show that ZMYM2 and TRIM28 functionally converge on regulating the same gene expression programmes. As requested by the referee, we have added further data showing that the ZMYM2 protein that is defective in SUMO binding (ZMYM2(SIM2mut); new Supplementary Fig. S3C) shows reduced binding to TRIM28 in co-IP assays. This further strengthens the (SUMO-dependent) association between ZMYM2 and TRIM28.

- *The transcriptional response using bulk RNA-seq in ZMYM2-depleted cells is rather gene-centric despite the title of the paper being about TE transcription. In fact the only panels about TE transcription are the RT-qPCR data in Fig 5D,F. I may be missing something (and there aren't many details given about the RNA-seq experiments) but why not look at TE transcription in an unbiased way with the transcriptomic data at hand? I appreciate potential hazards of multi-mapping etc but it would be interesting to see at least some subfamily analysis (e.g. using the Tetrascripts tool). On a similar point, why not show some RNA-seq in the genome browser snapshots of the epigenomics - together with a RepeatMasker annotation track of TEs...*

See response to the same point above.

While the results broadly support the authors' conclusions, I have the overall impression that the central claim of TE transcriptional regulation by ZMYM2 could be strengthened a lot with some fairly straightforward additional experiments and analyses.

Reviewer #3 (Public Review):

ZMYM2 is a transcriptional repressor known to bind to the post-translational modification SUMO2/3. It has been implicated in the silencing of genes and transposons in a variety of contexts, but lacking sequence-specific DNA binding, little is known about how it is targeted to specific regions. At least two reports indicate association with TRIM28 targets (Tsusaka 2020 Epigenetics & Chromatin, Graham-Paquin 2022 bioRxiv) but no physical association with TRIM28 targets had been observed. Tsusaka 2020 theorizes an indirect, potentially SUMO-independent, interaction via ATF7IP and SETDB1.

Here, Owen and colleagues show that a subset of ZMYM2-binding sites in U2OS cells are clearly TRIM28 sites, and further find that hundreds of genes are silenced by both ZMYM2 and TRIM28. They next demonstrate that ZMYM2 homes to chromatin, and interacts with TRIM28, in a SUMOylation-dependent manner, suggesting that ZMYM2 is recognizing SUMOylation on TRIM28 itself. ZMYM2 separately homes to SINE elements bound by the ChAHP complex, in an apparently SUMOylation independent manner. Although this is not the first report to show physical interaction between ZMYM2 and ChAHP, it is the first to show that ZMYM2 homes to ChAHP-binding sites and functions as a corepressor at these sites.

The mode by which ZMYM2 and TRIM28 coregulate genic targets remains somewhat unclear. TRIM28/ZMYM2 bind to LTR elements, loss of these proteins results in upregulation of genes distal to (but in the same TAD as) these binding sites.

Overall, the manuscript is well-written, convincing, and fills a significant hole in our understanding of ZMYM2's mechanistic function.

We thank the referee for his/her positive evaluation of the mechanistic insights we provide. We have further added to these through addressing the specific issues raised in their “recommendations for authors”.

The reviewers appreciated the novelty of the findings, and in particular, the use of the RIME method to identify the protein partners of ZMYM2 while bound on chromatin, and multiple validation steps of these novel ZMYM2 interactors. However, they also felt that the model presented at the end of the manuscript seems preliminary and would deserve additional experiments to be really supported, the essential ones being listed below:

1 - Despite the claimed scope of the manuscript on TE regulation, their expression analysis is limited to RT-qPCR and targeted to a few families or copies. Please use the RNA-seq data generated in U2OS cells depleted for ZMYM2 to assess retrotransposon expression genome-wide, performing both family-level and copy-level analyses, and compare with TRIM28-depleted U2OS cells.

We re-analysed our RNA-seq data using the Tetrascripts tool and looked at TE transcription genome-wide. However very few TEs were expressed at high enough levels to get any statistically significant additional data beyond a few additional transposable elements. This likely results from the relatively low read depth we used and the lack of specific protocols being followed to preserve TE transcripts. We will return to the genome-wide effects in future studies where we plan to switch cell types and will generate more bespoke datasets (the current ones were designed for analysing effects on protein coding gene expression before we made the connection to TEs). We added additional text to the results section to indicate that we could not see widespread deregulation of subclasses of TEs but that this needs further work.

2 - Clarify the relationship between dysregulated genes and TAD boundaries, as this seems important to support the model of distant gene regulation by the action of ZMYM2 on local chromatin environment within TADs (see comment of Reviewer #1 and #3).

We have now provided further support for the idea that ZMYM2 functions within TADs as detailed below in response to the reviewers comments. New bioinformatics analysis has been done which is incorporated into the paper in Fig. 4D and Supplementary Fig. S4C.

3 - Perform TRIM28 ChIP-seq in ZMYM2-kd cells, to prove that ZMYM2 indeed participates to TRIM28 recruitment to TE loci. This could be complemented by H3K9me3 ChIP-seq, to see if ZMYM2 depletion reduces H3K9me3 at retrotransposons, and over the regions surrounding ERVs. This last experiment seems also important for reinforcing the distant regulation model of nearby genes through ZMYM2-mediated repression of retrotransposons.

As suggested by the referees below, we have tested whether ZMYM2 is required for TRIM28 binding at several loci and find no evidence for this (new Supplementary Fig. S3E). We now discuss this in the results text and discussion where we already suggested that TRIM28 is likely recruited by KRAB-zinc finger proteins and ZMYM2 is subsequently recruited to this complex. Future extensive work is required to understand the mechanistic functions of ZMYM2 in these regions.

Recommendations for the authors:

The reviewers appreciated the novelty of the findings, and in particular, the use of the RIME method to identify the protein partners of ZMYM2 while bound on chromatin, and multiple validation steps of these novel ZMYM2 interactors. However, they also felt that the model presented at the end of the manuscript seems preliminary and would deserve additional experiments to be really supported, the essential ones being listed below:

1 - Despite the claimed scope of the manuscript on TE regulation, their expression analysis is limited to RT-qPCR and targeted to a few families or copies. Please use the RNA-seq data generated in U2OS cells depleted for ZMYM2 to assess retrotransposon expression genome-wide, performing both family-level and copy-level analyses, and compare with TRIM28-depleted U2OS cells.

We re-analysed our RNA-seq data using the Tetrascripts tool and looked at TE transcription genome-wide. However very few TEs were expressed at high enough levels to get any statistically significant additional data beyond a few additional transposable elements. This likely results from the relatively low read depth we used and the lack of specific protocols being followed to preserve TE transcripts. We will return to the genome-wide effects in future studies where we plan to switch cell types and will generate more bespoke datasets (the current ones were designed for analysing effects on protein coding gene expression before we made the connection to TEs). We added additional text to the results section to indicate that we could not see widespread deregulation of subclasses of TEs but that this needs further work.

2 - Clarify the relationship between dysregulated genes and TAD boundaries, as this seems important to support the model of distant gene regulation by the action of ZMYM2 on local chromatin environment within TADs (see comment of Reviewer #1 and #3).

We have now provided further support for the idea that ZMYM2 functions within TADs as detailed below in response to the reviewers comments. New bioinformatics analysis has been done which is incorporated into the paper in Fig. 4D and Supplementary Fig. S4C.

3 - Perform TRIM28 ChIP-seq in ZMYM2-kd cells, to prove that ZMYM2 indeed participates to TRIM28 recruitment to TE loci. This could be complemented by H3K9me3 ChIP-seq, to see if ZMYM2 depletion reduces H3K9me3 at retrotransposons, and over the regions surrounding ERVs. This last experiment seems also important for reinforcing the distant regulation model of nearby genes through ZMYM2-mediated repression of retrotransposons.

As suggested by the referees below, we have tested whether ZMYM2 is required for TRIM28 binding at several loci and find no evidence for this (new Supplementary Fig. S3E). We now discuss this in the results text and discussion where we already suggested that TRIM28 is likely recruited by KRAB-zinc finger proteins and ZMYM2 is subsequently recruited to this complex. Future extensive work is required to understand the mechanistic functions of ZMYM2 in these regions.

Reviewer #1 (Recommendations For The Authors):

- *Figure S1D is not clear. The authors want to investigate if ADNP and ZMYM2 regulate gene expression in the same directionality. They compare the genes down in siADNP and up in siZMYM2 (or vice versa) and show very small overlaps. If I understand correctly, this shows that very few genes are regulated in opposite*

directions by ADNP and ZMYM2 and consequently that they tend to regulate genes in the same directionality. This is not what is said in the text page 19 ("with no clear common roles as either an activator or repressor") and should be clarified. Furthermore, to compare if ADNP and ZMYM2 regulate genes in the same directionality, there are better ways to represent this, for example scatter plots of log2 FC in ADNP kd vs ZMYM2 kd. Similar criticisms apply to Fig S3F.

We agree that the text could be clearer and have rewritten it as "...although the large numbers of genes directionally co-regulated by these two proteins (ie either positively or negatively) indicates no clear common role as either an activator or repressor". We have also added a scatter plot to the supplementary data (Fig. S1E) to further emphasise the common directionality of effect as suggested by the reviewer. Similarly, we changed the text and have added a scatter plot to support the conclusions on ZMYM2 and TRIM28 functional interactions (new Fig. S3I).

- The authors suggest an indirect control of genes by ZMYM2 within TADs (Fig 4C). Yet Fig 4C does not seem to address this point. Fig 4C shows that TADs with a ZMYM2/cluster 1 peak contain more upregulated than downregulated genes, but the key question should be: are upregulated genes significantly enriched in TADs containing a ZMYM2/cluster 1 peak compared to other TADs or other genomic regions?*

We have taken this suggestion on board and determined the frequency distribution of the number of TADs containing a gene upregulated (fold change >1.6; Padj <0.01) following ZMYM2 depletion. 10,000 iterations were performed by randomly selecting 216 TADs across all 3062 TADs. The observed number of TADs containing an upregulated gene (42) from 216 TADs containing a cluster 1 ZMYM2 peak is a clear outlier in this distribution (P-value = 0.0002) (see Supplementary Fig. S4C).

- A key question not addressed in the manuscript is whether ZMYM2 participates in the recruitment of TRIM28 to ERVs. I recommend performing TRIM28 ChIP in ZMYM2-kd cells.*

We have tested whether ZMYM2 is required for TRIM28 binding at several loci and find no evidence for this (new Supplementary Fig. S3E). We now discuss this in the results text and discussion where we already suggested that TRIM28 is likely recruited by KRAB-zinc finger proteins and ZMYM2 is subsequently recruited to this complex. Future extensive work is required to understand the mechanistic functions of ZMYM2 in these regions.

Reviewer #2 (Recommendations For The Authors):

Please give more details of RNA-seq analyses in the experimental section (this will be particularly important if the comment about analysing TE transcription genome-wide is acted on).

We have now expanded on the description of the RNA-seq analysis including adding in the mapping statistics to a new Supplementary table. We followed the referee's useful suggestion of looking at TE transcription genome-wide. However very few TEs were expressed at high enough levels to get any statistically significant additional data. This likely results from the relatively low read depth we used and the lack of specific protocols being followed to preserve TE transcripts. We will return to the genome-wide effects in future studies where we plan to switch cell types and will generate more bespoke datasets (the current ones were designed for analysing effects on protein coding gene expression before we made the connection to TEs).

Reviewer #3 (Recommendations For The Authors):

Major Comments:

- *The relationship of TRIM28/ZMYM2 repression of LTRs and silencing within/between TADs is interesting but underdeveloped. Upon ZMYM2 depletion, the authors observe simultaneous upregulation of genes within TADs more often than would be expected by chance, but this analysis does not distinguish "proximal to" from "in the same TAD". If a ZMYM2 binding site is X bases from a gene TSS, is it more likely to regulate that gene if it is in the same TAD? This can and should be tested bioinformatically.*

The basic question the referee is asking is whether ZMYM2 affects gene expression at a certain distance irrespective of whether the TSS of the gene is in the same TAD. We have now tested this and added text to the results section. Basically we took all of the ZMYM2 regions associated with genes upregulated by ZMYM2 depletion that resided in the same TAD and calculated the peak to TSS distance. Then we searched in the opposite direction for the TSS of genes at a similar distance ($\pm 25\%$) that resided in an adjacent TAD. We then asked whether these genes were upregulated by ZMYM2 depletion. 102 ZMYM2 peaks were positioned within these distance constraints with at least one gene in an adjacent TAD (716 genes in total). Of these genes, only 11 were upregulated following ZMYM2 depletion. There is therefore not a general spreading of deregulation around ZMYM2 peaks in a distance-dependent manner.

Furthermore, the authors note in the text and discussion that LTRs can demarkate TAD boundaries, but this is a distinct concept from the idea that they regulate genes within a TAD. Is there evidence that ZMYM2 binding sites are found at TAD boundaries?

We have provided more evidence to support the associations of ZMYM2 peaks with TADs and now show that they are closer than randomly expected to TAD boundaries (Fig. 4D). However they are clearly not all located very close to the boundaries.

- *The analysis of transposons expression was limited to qPCR of a handful of elements. Since the authors have conducted RNA-seq of U2OS cells depleted for both TRIM28 and ZMYM2, they can determine if certain classes of transposons are globally upregulated.*

We re-analysed our RNA-seq data using the Tetrascripts tool and looked at TE transcription genome-wide. However very few TEs were expressed at high enough levels to get any statistically significant additional data. This likely results from the relatively low read depth we used and the lack of specific protocols being followed to preserve TE transcripts. We will return to the genome-wide effects in future studies where we plan to switch cell types and will generate more bespoke datasets (the current ones were designed for analysing effects on protein coding gene expression before we made the connection to TEs). We added additional text to the results section to indicate that we could not see widespread deregulation of subclasses of TEs but that this needs further work.

Minor Comments:

- *Typo: "human HEK393 cells". They are HEK293 cells.*

We have corrected this error.

- *"These ADNP peaks showed enrichment of binding motifs for several transcription factors with the top two motifs for HBP1 and IRF both found in over 35% of target regions (Figure 1D)." According to Ostapczuk 2018, ADNP has its own motif (CGCCCYCTNSTG). It is intriguing that this does not appear enriched in ADNP sites in U2OS cells, this seems worthy of comment.*

This is a good point, so we did an additional search using the motif found in Ostapczuk 2018 and found this in 15% of ADNP binding regions. This value is substantially lower than the 63% seen previously. It therefore is present but is not the dominant motif. This new data and its implication regarding chromatin targeting mechanisms is now discussed in the Results section around Fig. 1D.

- *Figures S2F and S2G are central to the paper and belong in the main text.*

We have now added these to the main figures as requested (meaning that Fig.2 has now been split into two separate figures {2 and 3} as became too large for a single figure).

- *A supplementary table including libraries generated and mapping statistics should be included.*

We have now added this (new Supplementary Table S2)