

Back to the future: omnipresence of fetal influence on the human brain through the lifespan

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

May 23, 2023 (this version)

Sent for peer review

February 28, 2023

Posted to preprint server

February 10, 2023

Kristine B. Walhovd , Stine Kleppe Krogsrud, Inge K. Amlien, Øystein Sørensen, Yunpeng Wang, Anne Cecilie Sjøli Bråthen, Knut Overbye, Jonas Kransberg, Athanasia M. Mowinckel, Fredrik Magnussen, Martine Herud, Asta K. Håberg, Anders M. Fjell, Didac Vidal-Piñeiro

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway • Department of Radiology and Nuclear Medicine, Oslo University Hospital, Norway • Department of Neuromedicine and Movement Science, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Human fetal development has been associated with brain health at later stages. It is unknown whether growth in utero, as indexed by birth weight (BW), relates consistently to lifespan brain characteristics and changes, and to what extent these influences are of a genetic or environmental nature. Here we show remarkably stable and life-long positive associations between BW and cortical surface area and volume across and within developmental, aging and lifespan longitudinal samples (N = 5794, 4-82 years of age, w/ 386 monozygotic twins, followed for up to 8.3 years w/12,088 brain MRIs). In contrast, no consistent effect of BW on brain changes was observed. Partly environmental effects were indicated by analysis of twin BW discordance. In conclusion, the influence of prenatal growth on cortical topography is stable and reliable through the lifespan. This early life factor appears to influence the brain by association of brain reserve, rather than brain maintenance. Thus, fetal influences appear omnipresent in the spacetime of the human brain throughout the human lifespan. Optimizing fetal growth may increase brain reserve for life, also in aging.

eLife assessment

This **valuable** study uses multiple large neuroimaging data sets acquired at different points through the lifespan to provide **solid** evidence that birthweight (BW) is associated with robust and persistent variations in cortical anatomy, but less-substantial influences on cortical change over time. These findings, supported by robust statistical methods, illustrate the long temporal reach of early developmental influences and carry relevance for how we conceptualize, study, and potentially modify such influences more generally. The paper will be of interest to people interested in brain development and aging.

Introduction

It is established that a substantial portion of functional variation through the lifespan, including in older age, is of neurodevelopmental origin^{1,4}. Evidence converges on early life factors being important for normal individual differences in brain, mental health and cognition across the lifespan^{5–9}, as well as risk of psychiatric¹⁰ and neurodegenerative disease in older age¹¹. Obtaining reliable indicators of individual early life factors is a major challenge. In this regard, birth weight (BW) stands out as a solid available measure. BW reflects fetal and maternal genetic, but also other in utero environmental factors affecting fetal growth^{12–14}, including brain growth^{6,15–17}. By now, a series of studies have established that BW relates positively to mental health, cognitive function and brain characteristics, including neuroanatomical volumes and cortical surface area as measured in different age groups^{4–6,15,18,19}. However, it is unknown whether and how BW relates to brain characteristics through the lifespan, how consistent effects are within and across samples, whether BW is associated with lifespan brain changes, and to what extent lifespan effects of BW on the brain are of an environmental, rather than genetic nature. These questions, which are critical to understand how and when the human brain can be influenced through the lifespan, we address in the present study. On an overarching level, this study also addresses current debates in the field of lifespan cognitive neuroscience, namely: 1) whether consistent, reproducible relationships between phenotypes relevant for mental health and function and inter-individual differences in brain characteristics can be found²⁰, and 2) to what extent effects found in and ascribed to brain aging may actually reflect early life influences, rather than longitudinal changes in older age^{5,7,21}.

There are at least two different ways by which the effects of fetal growth, as indexed by birth weight, could work to produce the brain effects so far observed. 1) In line with a brain reserve model^{22,23}, higher birth weight could be associated with greater brain growth before birth. This seems likely, given that the effects are seen also in young populations^{6,15}. However, from the so far largely cross-sectional, or mixed models, several questions remain unanswered: Is this a fixed effect at the time of birth? Does higher birth weight also have carry-over effects to greater development in childhood and adolescence? In line with a brain maintenance model²⁴, is higher birth weight associated with better maintenance of brain volumes in the face of age-related changes in older adulthood? While effects are found in young populations^{5,6,15}, reduced atrophy in aging is a possible additional effect of higher BW that should be investigated, given the known relationships between birth size and brain volumes also in older age⁴. The possible effects of BW on later brain development and brain maintenance in adulthood can only be investigated by longitudinal brain imaging spanning all stages of human life. Furthermore, as BW normally reflects both genetic and prenatal environmental factors, and an environmental BW contribution to brain differences has been shown in young monozygotic (MZ) twins^{15–17}, we need to study brain effects of birth weight discordance in MZ twins in this context to disentangle possible non-genetic contributions of BW through the lifespan.

We hypothesized that there are persistent effects of BW on brain characteristics through the lifespan, and hence, that these would be consistent within and across samples of varying age and origin. We test this in a Norwegian sample covering the lifespan (LCBC)^{5,21}, the US developmental sample ABCD^{25,26}, and the older adult UK biobank (UKB)^{27,28} sample. The associations of BW and cortical surface, thickness, volume and their change were investigated vertex-wise in a total of 5794 persons (of whom 5718 with repeated scans, and 386 monozygotic twins) with 12088 longitudinal observations, 4–82 years of age at baseline, followed for up to ~ 8.3 years. Based on previous results^{5,6,15}, we hypothesized such effects to be driven primarily by positive associations between birth weight and cortical area, with lesser, if any, effects on cortical thickness. We expected positive effects on cortical volume corresponding to positive effects on cortical area. We hypothesized that effects would be stable, so that BW mainly affects the brain “intercept” and does not relate much to brain changes. That is, we hypothesize a threshold model, whereby higher BW yields greater cortical area, and hence cortical volume, to

begin with, rather than a maintenance model, whereby higher BW serves to protect against atrophy in aging. Moreover, we hypothesized that effects could not be explained solely by genetics, so that birth weight discordance in a subsample of MZ twins would also result in differences in brain characteristics through the lifespan.

Results

Cortical surfaces were reconstructed from T1-weighted anatomical MRIs by use of FreeSurfer v6.0 (LCBC and UKB), and 7.1. (ABCD) (<https://surfer.nmr.mgh.harvard.edu/>)^{29–32}, yielding maps of cortical area, thickness and volume. Vertex-wise analyses were run with spatiotemporal linear mixed effects modeling (FreeSurfer v6.0.0 ST-MLE package), to assess regional variation in the relationships between birth weight and cortical structure and its change. All analyses were run with baseline age, sex, scanner site, and time (scan interval) as covariates. For ABCD specifically, ethnicity was also included as a covariate. For consistency of multiple comparison corrections across analyses, the results were thresholded at a cluster-forming threshold of 2.0, $p < .01$, with a cluster-wise probability of $p < .025$ ($p < .05/2$ hemispheres).

The lifespan relationship of BW and cortical volume, surface area and thickness

Associations of birth weight and cortical characteristics are shown in **Figure 1** (for the right hemisphere, and in **Supplementary Figure 1** for both hemispheres). Across all cohorts, widespread positive associations were observed between BW and cortical area. These were highly consistent across lifespan (LCBC), developmental (ABCD) and aging (UKB) cohorts, and there were bilateral overlapping effects across most of the cortical mantle. As expected, BW had in general lesser effects on cortical thickness, and no significant effects on thickness were observed in the UKB. There were however some lateral positive and medial negative effects in the LCBC and ABCD cohorts. We note that corresponding effects with increased medial frontal and occipital cortical thickness have been found associated with white matter alterations (reduced FA) in young adults born preterm with very low BW compared to term-born controls³³. BW was significantly positively associated with cortical volume across much of the cortical mantle. In sum, broad, bilateral, positive associations were observed across cohorts for cortical area and volume.

Additionally controlling for education level had little effect on results (see **Supplementary Figure 2**). Information on gestational length (i.e. whether there was premature birth) was not available for all participants. Importantly, this information was lacking for the older participants, i.e. this information is not available for UKB, and since this information for the LCBC was drawn from the Medical Birth Registry of Norway (MBRN), only established in 1967, this was not available for the older part of the LCBC sample either. The majority of the LCBC sample and the ABCD sample, had information on gestational length, however (LCBC: $n = 514$; gestational length in weeks: $M = 40.0$ weeks, $SD = 1.9$, range = 25–44; ABCD: $n = 3306$; weeks premature: $M = 1.0$ weeks premature, $SD = 2.1$, range = 0.0–13.0). Controlling for gestational age in these subsamples had relatively little effect on results (see **Supplementary Figure 3**). However, as expected with reduced power in the LCBC sample, the effects in this analysis were somewhat narrower. Effects in ABCD, where almost all participants were retained for analysis, showed no sign of decrease with control for gestational length. When restricting all samples to participants with BW between 2.5 and 5.0 kg, results were also very similar (see **Supplementary Figure 4**). As expected from the widespread effects on cortical area and volume, effects were partly generic, with analyses controlling for ICV showing more restricted effects (see **Supplementary Figure 5**). However, consistent significant positive effects of BW on cortical area also when controlling for ICV were observed across all three cohorts in lateral temporal and frontal areas (see **Supplementary Figure 6**).




Figure 1.

Relationships of birthweight and cortical characteristics across LCBC, ABCD, and UKB samples.

Age, sex, time (interval since baseline) and scanner site (as well as ethnicity in the ABCD) were controlled for. Significant relationships are shown for area, thickness, and volume for each sample, from left to right: lateral view and medial view, right hemisphere.

Birth weight effects on cortical change

Significant BW x time interactions on cortical characteristics were observed in restricted and non-overlapping regions across samples, see **Figure 2** (depicting right hemisphere results, for visualization of effects in both hemispheres, see **Supplementary Figure 6**). Per direction of effect, the effect of BW differences was apparently reduced over time for area in LCBC and ABCD, whereas no interaction effects on area were significant in UKB. A mixture of positive (ABCD) and negative (LCBC, UKB) interaction effects were significant for thickness and volume.

Visualization of the interaction effects as seen in **Figure 2** and **Supplementary Figure 6**, by splitting the sample in two based on BW, did not yield convincing evidence for these interactions, as shown in Supplementary Figures 7-9. In plots of LCBC data, where number of follow-ups varied, and a select portion had longer follow-up, it appeared that the effect of BW was reduced over time in these restricted regions. However, virtually parallel trajectories for the ABCD and UKB subsamples with lower and higher BW, suggested the effect size even within the areas of significant interactions of BW and time was negligible. Since UKB and ABCD samples here consisted of samples having two time points only, whereas LCBC consisted of a mix of number of follow-ups over a longer time period, there might be sample-specific selection effects also regarding other characteristics than BW that can influence these effects in LCBC. For instance, participants who do not drop out tend to have better health, cognitive ability and education, which again may relate positively to the brain measures studied here ^{34, 35}. Thus, caution is advised in interpreting effects seen only with longer follow-up in the LCBC sample.

Consistency of spatial relationships across and within samples

Next, we assessed whether the cortical correlates of BW (beta-maps) showed a similar topographic pattern across the three independent datasets (UKB, ABCD and LCBC). The results showed that all the spatial comparisons were statistically significant ($p < 0.05$, FDR-corrected). That is, the topography of the effects of BW on cortical structure was comparable across datasets – the pairwise spatial correlation of a given cortical correlate of BW (e.g. BW effects on cortical area) was similar when estimated from two different datasets. The spatial correlations were highest for the volume measures ($r = .64-.79$), and overall also high ($r = .51-.71$) for area measures, whereas for cortical thickness, they were more moderate ($r = .24-.45$). See spatial correlations for the right hemisphere cortical volume in **Figure 3** and the full model summary in **Supplementary Table 2**. The results are qualitatively comparable when using $-\log_{10}(p)$ significance values instead of beta estimates, as shown in **Supplementary Table 2**. The same pattern of results was largely seen also for spatial correlation of the maps capturing BW-associated cortical characteristics when controlling for ICV. The correlations were then on average somewhat lower, but there were still only significant positive correlations across LCBC, ABCD, and UKB (see **Supplementary Table 2**).

In contrast, the spatial correlation of the maps capturing BW-associated cortical *change* were either unrelated ($n = 7$) or showed negative associations between cohorts ($n = 2$). The spatial correlations of birth weight on cortical change were $r = -.35-.05$ for area, $r = -.35-.08$ for volume, and $r = -.20, -.04$ for thickness. See a visual representation in **Supplementary Figure 10** and full stats in **Supplementary Table 2**.

In sum, the spatial correlation analyses imply that the different datasets show a comparable topography of BW effects across the cortical mantle – i.e. the areas more and less affected by BW were common across datasets. Thus, the BW effects on cortical structure are robust and replicable across very different datasets. In contrast, the effects of BW on cortical change are not robust across datasets, showing dissimilar topographies.




Figure 2

Interactions of BW and time on cortical characteristics across LCBC, ABCD, and UKB samples.

Age, sex, scanner site, time, and birth weight (as well as ethnicity in the ABCD) were controlled for. Significant relationships are shown, from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.




Figure 3.

Spatial correlation of birth weight effects on brain structure across datasets for cortical a) area, b) thickness, and c) volume.

Spatial correlation of birth weight effects on brain structure across datasets. For each panel, the upper triangular matrix shows Pearson's (r) pairwise spatial correlation between the different cohorts' cortical maps. Data is shown as a color-density plot. The red line represents the fitting between the two maps. The lower triangular matrix shows the significance testing. The dashed-grey line shows the empirical correlation, while the orange histogram represents the null distribution based on the spin test. The diagonal shows the effect of birth weight on cortical structure (right hemisphere shown only). Note that the β -maps are shown as a percentile red-green-blue scale, where red represents a lower (or more negative) effect of birth weight on cortical structure and vice versa. See **Supplementary Table 2** for stats. Units in the density maps represent birth weight effects as mm^2/g , mm^3/g , and mm^3/g ($10e^{-5}$) for cortical thickness, area, and volume, respectively.

Additionally, split-half analyses within datasets were performed, to investigate the replicability of significant effects ³⁶[36](#), ³⁷[37](#) of BW on cortical characteristics within samples (refer to **Figure 1** [1](#)). The results are shown in **Supplementary Figure 1** [1](#). This analysis further confirmed that the significant effects were largely replicable for volume and area, but not for thickness. The replicability of the cortical area and volume effects is thus shown across and within samples.

Effects of BW discordance on brain characteristics and changes in monozygotic twins

BW discordance analyses on twins specifically were run as described for main analyses above, with the exception that twin scans were reconstructed using FS v6.0.1. for ABCD. BW discordance was associated with cortical area, where the heavier twins had greater area in some frontal, temporal and occipitotemporal regions, with effects in the right hemisphere only surviving corrections for multiple comparisons. We note that these regions mostly overlap with regions where positive effects of BW were also seen in the bigger sample. Strikingly, the effect of BW discordance, as shown in **Figure 4** [4](#), appeared similar in size to the effect of BW itself in the MZ twin sample. However, note that this plot is merely for illustrating effects, the effect size is inflated for the BW discordance plot, since the values are derived from areas already identified as significantly related to BW. There was no association of BW discordance and cortical area changes over time.

BW discordance also had a significant negative effect on cortical thickness in restricted right frontotemporal regions, where being the lighter twin yielded greater thickness. These significant effects did not appear to overlap with regions where significant negative associations with BW were seen in the bigger sample. BW had little effect on cortical thickness in the significant region, and the effect of BW discordance in the identified regions, as shown in **Figure 5** [5](#), appeared greater than the effect of BW itself here in the MZ twin sample. However, this plot is merely for illustrating effects, it should be noted that the effect size is inflated for the BW discordance plot, since the values are derived from areas already identified as significantly related to BW.

In a very small area of the right hemisphere, there was a significant association of BW discordance and cortical thickness change, meaning the lighter twin had greater cortical thickness over time, but this effect was both regionally and quantitatively minor, as shown in **Supplementary Figure 12** [12](#). There were no significant effects of BW discordance on cortical volume or volume change over time.

Finally, to formally assess whether the cortical correlates (β -maps) of BW discordance in the twin subsample corresponded to cortical correlates of BW in the bigger samples, we did a meta-analysis of these estimates for area, thickness and volume in the UKB, ABCD and LCBC, and then assessed whether the cortical correlates of BW and BW discordance (β -maps) showed a similar topographic pattern across the datasets. The results of this meta-analysis-twin comparison showed only positive relationships, for area, $r = .23$, thickness $r = .19$, and volume $r = .22$. However, the respective uncorrected p -values were .08, .12, and .04, so the spatial comparisons would not be statistically significant ($p < 0.05$, FDR-corrected). However, the positive correlations are suggestive that the topography of the effects of BW discordance in genetically identical twins on cortical structure was to some extent comparable to effects of individual differences in BW in the bigger sample.

Effects of BW differences compared to effects of aging

We calculated the effect of 1 SD difference in BW (on average 600 g, see **Table 1** [1](#)) on cortical and brain volume across cohorts, to directly compare the early life effects on the brain to later aging changes. The effect of 1SD lower BW on cortical volume was 6708 mm³, 8466 mm³, and 5980 mm³ in LCBC, UKB and ABCD, respectively. This was equal to 1.2%, 1.6% and 1.1% lesser cortex with 600-700 grams lower BW for each sample, respectively. In the context of brain aging, this is a




Figure 4

Effects of birthweight discordance on cortical area in the sample of monozygotic (MZ) twins.

Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere. Plots are showing – for illustrative purposes – individual data points and expected trajectories for cortical area in mm (Y-axes) within the significant regions according to birth weight (BW) discordance (left panel) and BW (right panel) in kilograms (X-axes).




Figure 5

Effects of birthweight discordance on cortical thickness in the sample of monozygotic (MZ) twins.

Relationships significant corrected with cluster-forming threshold of 2.0 ($p < .01$) are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere. Plots are showing – for illustrative purposes – individual data points and expected trajectories for cortical thickness in mm (Y-axes) within the significant regions according to birth weight (BW) discordance (left panel) and BW (right panel) in kilograms (X-axes).

substantial effect. The estimated yearly cortical volume reduction from 50 to 60 years is 895 mm³ in the LCBC and 1402 mm³ in the UKB samples, respectively. Hence, the effect of 600 grams difference in BW equals 7.5 and 6 years of aging in LCBC and UKB, respectively.

Discussion

The present results indicate that BW, the earliest widely and easily obtainable congenital metric, show robust, persistent, and chiefly stable associations with brain characteristics through life. Especially, BW was associated with cortical area and volume in an age and time-invariant fashion. The robustness of this effect is quite remarkable, given the wealth of different influences individuals meet after birth, which are repeatedly assumed and reported to have major impact on the brain through the protracted human lifespan^{38, 39}. It is also in quantitative terms outstanding, compared to consistency of cortical topographies reported for other phenotypical factors²⁰. This is also special for a phenotype known to be environmentally influenced, unlike biologically hardcoded phenotypes such as sex or age, for which there are known brain-wide association studies (BWAS) patterns^{36, 40}.

Typically, other factors relating to later socioeconomic status, lifestyle, and health, get the most attention in adult and aging brain research^{38, 39}. Such factors, which are then targeted for prevention and intervention at different stages of the life-course, often do not show consistent relationships to brain characteristics⁴¹, may not actually be causal⁴¹, and may themselves be related to prenatal growth¹⁹. Another phenotype which obviously, like BW, reflects both genetic and environmental influences^{12–14}, is body mass index (BMI)⁴². Consistent BWAS patterns have been reported for BMI³⁶. BW stands out as the single chronological earliest phenotype, and besides BMI, BW appears to have the most replicable and consistent relations to cortical morphology, as shown here both across and within samples. It has been claimed that smaller than expected brain–phenotype associations and variability across population subsamples can explain widespread replication failures for brain-wide association studies (BWAS)²⁰. However, this is necessarily a question of which phenotypes are the most relevant to relate to brain characteristics. Also, the temporal order of factors needs to be considered if causal interpretations are to be made. Chronologically later factors necessarily do not cause earlier ones. While we cannot claim that BW itself causes the cortical characteristics observed in aging, the cortical variance explained by BW after one decade, and seven or eight decades of life alike, is unlikely to be explained by influences only present at some point in adulthood or aging. BW, as further discussed below, depends on genetic, as well as prenatal environmental influences^{12–14}, which likely have causal effects on early brain morphological features. Here we find that these effects are substantial also in the aging brain.

We calculated that a BW difference of one SD (about 600 g) equaled a difference in cortical volume on the order of 1.1–1.6 percent in these cohorts. This is a quite big effect, of a magnitude relevant for explaining a substantial portion of the differences typically seen between patients with neurodevelopmental or neurodegenerative diseases and healthy controls. As noted, BW differences have been reported for neurodevelopmental disorders such as ADHD⁴³, and also other neurodevelopmental disorders, such as schizophrenia or more general psychopathology, with the cortical effects detected not invariably being very large in absolute terms⁴⁴. There is a limit to the range of variation that can apply to human cortical volumes, in general, (e.g. virtually none have cortical volumes below 0.45 or above 0.65 l). In terms of sample representativity, one may assume that there can be a restriction relative to the actual range of human cortical volume variation, as the present samples specifically are largely healthy⁴⁵. Much of the differences within this limited range of variation is explained by factors here controlled for, such as age and sex. With the present effect, on the restricted range of variation, combined with big samples, it is obvious that BW differences of much less than our example magnitude (600 g) may be detectable in the cortical morphology of patients versus controls. In the context of aging and

neurodegenerative change, the estimated cortical effect of ~600 g difference in BW is of a magnitude many times the annual cortical reduction estimated to take place from e.g. 50 to 60 years in the adult cohorts, where the estimated yearly cortical volume reduction would be 895 mm³ in LCBC and 1402 mm³ in UKB. The effect of 600 grams difference in BW is equivalent to 7.5 and 6 years of aging in LCBC and UKB respectively. This is a substantial effect in terms of brain aging in imaging, and may illuminate why machine learning estimates such as “brain age”, assumed to index aging-related processes, may rather than relate to longitudinal age-changes, largely capture variance already determined at birth^{7,46, 47}. Neglecting this especially consistent and early factor is likely to lead to a substantial portion of human brain variance being either erroneously ascribed to factors only present at later life stages⁷ or left unaccounted for.

The solidity, replicability, and universality of effects as shown here for a partly environmentally influenced metric^{12, 14}, appear exceptional in human brain imaging. The within-sample replicability results are not fully comparable to other studies assessing the replicability of brain-phenotype associations due to analytical differences (e.g. sample size, multiple-comparison correction method)^{20, 36}. Still, these results too clearly show that the rate of replicability of BW associations with cortical area and volume are comparable to benchmark brain-phenotype associations such as age and BMI with brain structure³⁶. The BW-cortical volume and area associations may be among the topographically broadest and most consistent effects so far seen as stable across the lifespan of the human brain. The three cohorts studies differ on a range features known to be highly and reliably related to cortical characteristics, first and foremost age^{2,5, 36, 48, 49}, but also country of origin and representativity of the populations from which they are drawn^{26, 50}. Yet there is a comparable topography of BW effects across the samples. This is so despite the samples collectively spanning the entire human age range, within which there are always substantial age-related changes in cortical structure^{2, 5, 51–53}. The present results thus indicate that fetal growth influences an offset of brain reserve^{22, 23} and that this brain reserve effect is persistent and stable through the lifespan.

In contrast, the cortical maps capturing *change* in cortical structure associated with BW were not robust across datasets; i.e. the most positive and negative association with BW on cortical change did not overlap at all between the different cohorts. While there was evidence from ABCD that BW affected regional cortical development in the narrow age range covered, there were limited and no consistent effects of BW on cortical change across cohorts. Importantly, there was no indication whatsoever that BW could be associated with better brain maintenance²⁴ in the face of age-related changes in older adulthood. Thus, the data seem to indicate that any effect of BW on cortical *change* may be of relatively more temporary nature. The “offset effect” of BW, on the other hand, appears persistent and consistent, especially in terms of stable and widespread effects on cortical area and volume across the lifespan.

The sensitivity analyses indicate that the associations between BW and cortical characteristics are seen irrespective of not only sex and age, but also education, head size (ICV), and cases of abnormal BW. Such patterns could point to an underlying genetic pleiotropy of BW and brain characteristics. Interestingly, however, recent findings indicate that effects of exposure to environmental adversity on epigenetic programming in aging may be localized to the in-utero period⁵⁴. The effects of BW discordance in MZ twins in this context, align with other studies^{16, 17} pointing to also non-genetic, that is environmental, influences in the womb, associated with the pattern observed for cortical area effects. These analyses also account for multiple possibly confounding variables that could represent a mix of genetic-environmental effects, such as parental socioeconomic status, parity, or prenatal exposures shared between twins in the same womb such as maternal smoking or use of alcohol.

The neural basis for the observed association cannot readily be ascertained from human imaging studies tracking change^{5, 6, 15}. While the “fetal origins hypothesis”, proposing that cardiovascular disease in adulthood is related to undernourishment in utero^{55, 56} is well-

known, there has been focus on “brain-sparing” adaptations under such conditions¹⁹. However, our finding that early human development in utero appears to be associated with a persistent and stable brain reserve effect, is largely in correspondence with what is known of human nervous system development and change through the lifespan: While synaptogenesis, synaptic remodeling and myelination are known to be protracted processes long after infancy^{57–60}, numerous processes in brain development appear to be exclusively, or almost exclusively happening before birth. For instance, neurogenesis takes place almost only in fetal development⁶¹. Even if controversies remain, evidence suggests that any adult human neurogenesis must be severely restricted in location and amount^{62, 63}. Thus, human beings appear to be born with almost all cortical neurons they will have through life, and neuronal migration and differentiation are also defined early, by the place and time the neuron is born during fetal life⁶¹. Factors that affect placental function and uterine and/or umbilical blood flow on a chronic basis may lead to restricted fetal growth, including brain growth, and given the timing of brain development, it may not be surprising that effects would be stable across years. Animal studies of chronic placental insufficiency have shown effects on brain development which persist with age⁶⁴. Hence, the relationship between BW and cortical characteristics in the normal population could likely have a twofold etiology: it is likely to in part be based on normal variation in genetically determined body and brain size, but it also may be based on variations in environmental prenatal conditions, yielding differences in optimality of early brain development persisting through the lifespan.

These results indicate that there is potential to increase brain reserve throughout the lifespan, also in aging, by combating factors affecting fetal growth negatively. While it is unknown to what extent results from these specific US and European samples can be generalized to other populations, the current potential to improve prenatal factors may be especially high in low and middle-income countries, where the demographic changes will also be more marked in terms of the aging population⁶⁵. About 200 million children in developing countries are not meeting their growth potential, and improving the prenatal environment is likely important to help children reach their full potential⁶⁶ – and ultimately also to help them stay above a functional threshold into older age. Also in industrialized countries, including the US, environmental factors are associated with birth weight. The opioid epidemic is increasingly affecting pregnant women, and in-utero opioid exposure is associated with higher risk of fetal growth restriction⁶⁷. Among highly common exposures, air pollution may for instance be of relevance. Recently, local traffic congestion-pollution exposure during pregnancy, independently of a series of maternal sociodemographic characteristics, was associated with reductions in term BW in a large US sample¹⁴. Programs and policies to limit such environmental factors reducing fetal growth may thus enhance brain reserve and ultimately prevent more people from falling below a functional threshold even in advanced age.

Limitations

Some limitations should be noted. First, for most of the participants, only self-reported or parent-reported BW was available. While there was a very high correlation between registry and self-reported BW in LCBC, this is a possible source of noise. Second, pregnancy-related information of possible relevance, such as gestational age at birth, complications, method of delivery, maternal disorders, smoking, alcohol and drug intake, was not available across all participants of the different samples, and was thus not analyzed here, or, as for gestational age at birth, could only in part be controlled for. Some of these factors may be systematically related to BW, and may thus represent confounds⁶⁸. There were some premature, and very low BW participants in the samples, and these conditions are associated with known reductions in cortical volume^{69, 70}. However, the analyses controlling for gestational age, as well as on the restricted range of BW – excluding very preterm and very/extremely low BW children – and the analysis controlling for education, which may again relate to some of these factors, showed very similar results. It is unknown to what extent the BW of participants reflect their individual fetal growth potential, as a

fetus with normal BW can be growth restricted and a fetus with low BW can have appropriate growth⁷¹. We believe, however, that possible differences in such factors would likely serve to decrease consistency of results, and not lead to inflated estimates of consistency.

Conclusion

The current results show that a simple congenital marker of early developmental growth, BW, is consistently associated with lifespan brain characteristics. While some significant effects of BW on cortical change patterns were also observed, these were regionally smaller and showed no consistency across cohorts. In conclusion, while greater early human developmental growth does not appear to promote brain maintenance in aging, it does, in terms of greater cortical volume and area, relate positively to brain reserve through the lifespan. Thus, there appears to be an omnipresence of fetal factors in the spacetime of the human brain through the lifespan. This indicates a potential to increase brain reserve at all ages, including in aging, by combating factors affecting fetal growth negatively. Given the exceptional consistency and broadness of this cortical topographical effect, it should be taken into account in studies of brain research on individual differences, whether the brains studied are those of eight-or eighty-year-olds.

Materials and methods

Samples

In total, longitudinal data for 5718 persons with 12088 MRI scans from the LCBC, ABCD and UKB studies were included in the analyses. For UKB the dataset released February 2020 was used. For ABCD, the Data Release 3.0 was used (see <http://dx.doi.org/10.15154/1528313> for this NDA study). Only persons with longitudinal MRI scans were included in the main analyses, to limit the possibility that estimates of change were biased by immediate sample selection effects (i.e. those that remain for follow-up are known to have other characteristics than those who have only one time-point assessment in longitudinal studies, and this can bias effects). However, for the separate MZ twin-analyses, we also included participants with only one time-point MRI, to obtain an age-varying sample for assessment of whether non-genetic effects were found throughout the lifespan, including in adulthood and older age. Of the 386 MZ twins included, 310 had longitudinal imaging data. The twins were mostly ($n = 310$) from the developmental ABCD sample (age 10-11), whereas 64 adults were from LCBC (age 18-79 years), and 12 were from UKB (age 50-80 years). All samples consisted of community-dwelling participants.

For the most part, these were recruited by means of some type of population registry information (see Supplementary Materials, SM), but part of the LCBC cohort consisted of convenience samples, i.e. the studies were advertised broadly. Part of the LCBC sample was recruited through the Norwegian Mother, Father and Child Cohort Study (MoBa)⁷², and the Norwegian Twin Registry (NTR)⁷³. Thus, this study includes data from MoBa and NTR, and both studies are conducted by the Norwegian Institute of Public Health. LCBC participants were part of observational studies, but subsamples were part of studies including cognitive training ($n = 168$). As BW was not a criterion for assigning participants to cognitive training, these were included here. For the majority of participants, and all in UKB and ABCD, BW was collected as self-report or, for children, parent report, at the time of scan. For LCBC, BW was for the majority ($n = 526$) collected from the Medical Birth Registry (MBRN), available for those recruited through MoBa, or NTR, or if collected by consent for participants born 1967 and later, and in part by self-report in connection with scanning, or earlier self-report to the Norwegian Twin Registry (for twins recruited through this registry). MBRN is a national health registry containing information about all births in Norway. Comparative analyses in the LCBC sample for 354 persons who had available both MBRN records

and self-report/parent-report of BW, showed a very high correlation of BW as obtained from the different sources ($r = .99$). A high reliability of self-reported BW over time has also been found in broader NTR samples ⁷⁴[74](#). Demographics of the samples in the main analyses are given in **Table 1** ¹[1](#), see SM for details.

Statistical Analyses

Cortical Vertex-wise Analyses

Reconstructed cortical surfaces were smoothed with a Gaussian kernel of 15 mm full-width at half-maximum. We ran vertex-wise analyses to assess regional variation in the relationships between birth weight and cortical structure; area, thickness and volume at baseline and longitudinally. In all models we included 1) baseline age, sex and scanner site, as well as time (scan interval) as covariates. For ABCD specifically, ethnicity was also included as a covariate, as this sample is recruited to have and has, ethnic variation (see Supplemental Methods for details), whereas the other samples entered in the present analyses had little ethnic variation (i.e. in UKB, >98% of participants included in the present sample defined themselves as British/Irish/Any other white background. In LCBC, this information was unfortunately not encoded for all, but the sample was mainly of white background). In further models, we additionally included education as a covariate. General linear models were run in turn using as predictors: birth weight, the interaction term birthweight x scan interval, and the interaction term of baseline age x time (scan interval) x birth weight. When analyses were run with baseline age x scan interval x birth weight as predictor, the interaction terms of baseline age x scan interval, scan interval x birth weight, and birth weight x baseline age were included as additional covariates. Standardized values were used in analyses for age, scan interval, BW, BW discordance, and education. For consistency of multiple comparison corrections across analyses, the results were thresholded at a cluster-forming threshold of 2.0, $p < .01$, with a cluster-wise probability of $p < .025$ ($p < .05/2$ hemispheres). Finally, models were rerun only including participants with birth weights between 2.5 and 5.0 kg, to assess whether relationships were upheld also when excluding low and high birthweights. Given previous findings of broad effects of BW on cortical area and volume ⁵[5](#), ⁶[6](#), ¹⁸[18](#), we did not expect effects to be localized. Rather, we expected BW to affect gross head and brain size irrespective of sex, but we also performed supplementary analyses controlling for intracranial volume (ICV) in order to check for possible specificity of effects. Spatial correlation analyses ⁷⁵[75](#)–⁷⁷[77](#) were run on the cortical maps (for more information see SI) for analyses results using BW as predictor, from LCBC, ABCD, and UKB, to assess the overlap of BW-cortical characteristics associations in terms of topography and effect sizes. In a separate set of analyses, we restricted the sample to only monozygotic twins, and studied effects of BW discordance (number of grams BW above or below MZ twin). In these models, we included time, baseline age, sex and site as covariates.

Abbreviations

- BW
birth weight
- MRI
Magnetic Resonance Imaging

Study	N	F	Scans	Baseline age			Time since baseline			Birth weight			Education		
				M	SD	Range	M	SD	Range	M	SD	Range	M	SD	Range
LCBC	635	350	1922	19.1	20.7	4.1-81.9	2.6	2.4	0.1-8.3	3.5	0.6	0.9-6.0	17.1	2.4	9-24
ABCD	3324	1562	6648	10.0	0.6	8.9-11.1	2.0	0.1	1.4-2.8	3.2	0.7	1.0-6.7	16.0	2.6	6-21
UKB	1759	1009	3518	62.0	7.1	47-80.3	2.3	0.1	2.0-2.9	3.4	0.6	0.9-6.4	14.2	2.4	7-16

Table 1

Descriptive statistics for the longitudinal samples.

F = number of females in the sample, M = Mean, SD= Standard deviation. Numbers are given in years for baseline age, time since baseline and education, birthweight is given in kilograms. For LCBC, only 584 participants had information on education. Parental education was used in ABCD, and in LCBC when the participant was below 18 years of age, and also if no other education information was available for participants up to 21 years.

Acknowledgements

Data used for the LCBC sample were in part obtained through the Medical Birth Registry (MBRN) of Norway, the Norwegian Twin Registry (NTR) and the Norwegian Mother, Father and Child Cohort Study (MoBa). MoBa is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research. We are grateful to all the participating families in Norway who take part in this on-going cohort study. Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (<https://abcdstudy.org>), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the National Institutes of Health (NIH) or ABCD consortium investigators. The ABCD data used in this report came from DOI 10.15154/1504041 and 10.15154/1520591. DOIs can be found at <https://dx.doi.org/10.15154/1528313>. Part of the research was conducted using the UK Biobank resource under application number 32048. The ERC, RCN, EC, NIH and UK Biobank had no role in the design and conduct of this specific study. We thank all participants for contributing to all the respective data sources. This study was funded by ERC grants (771375 and 313440 to KBW; 283634 and 725025 to AMF), Research Council of Norway (RCN) Grants (to KBW, AMF and DVP), and European Commission (EC) EU Horizon 2020 Grant agreement number 732592 (Lifebrain). The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at <https://abcdstudy.org/federal-partners.html>.

Conflict of Interest Disclosures

The authors have no conflicts of interest to disclose.

Data Availability

The code used for vertex-wise ST-LME analyses and spatial correlations analyses can be accessed here <https://github.com/LCBC-UiO/paper-birthweight-brainchange-2022>. Two of the datasets used, the ABCD <https://abcdstudy.org> and UKB <https://www.ukbiobank.ac.uk> databases are open to all researchers given appropriate application, please see instructions for how to gain access here: <https://nda.nih.gov/abcd/> and <https://www.ukbiobank.ac.uk/enable-your-research>. The LCBC dataset has restricted access, requests can be made to the corresponding author, and some of the data can be made available given appropriate ethical and data protection approvals. However, the registry data on birth weight connected to this sample are not shareable by the authors, as these data are owned by the Medical Birth Registry of Norway, <https://www.fhi.no/en/hn/health-registries/medical-birth-registry-of-norway/medical-birth-registry-of-norway/>, the Norwegian Mother, Father, and Child Cohort Study <https://www.fhi.no/en/studies/moba/>, and the Norwegian Twin Registry <https://www.fhi.no/en/more/health-studies/norwegian-twin-registry/> so that any access to data must be approved by them.

Supplementary Material

Supplementary text

Supplementary Methods

Samples

Lcbc

Population, Recruitment and General Description of Study/ Procedures

Cognitively healthy, community dwelling participants across the lifespan were drawn from studies coordinated by the Research Group for Lifespan Changes in Brain and Cognition (LCBC www.oslobrains.no ⁷⁴), approved by a Norwegian Regional Committee for Medical and Health Research Ethics (REK South-East). Written informed consent was obtained from all adult participants and from parents or other legal guardians for participants below age of majority. The samples were recruited in part by newspaper and web page adds, and in part by population registry-based research studies. Part of the developmental sample was recruited through the population registry-based study MoBa, the Norwegian Mother, Father and Child Cohort Study at The Norwegian Institute of Public Health <https://www.fhi.no/en/studies/moba/> ⁷⁸. The Norwegian Mother, Father and Child Cohort Study (MoBa) is a population-based pregnancy cohort study conducted by the Norwegian Institute of Public Health. Participants were recruited from all over Norway from 1999-2008. The establishment of MoBa and initial data collection was based on a license from the Norwegian Data Protection Agency and approval from The Regional Committees for Medical and Health Research Ethics. The MoBa cohort is currently regulated by the Norwegian Health Registry Act. The current study was approved by The Regional Committees for Medical and Health Research Ethics (REK sør-øst C 2010/2359). Part of the adult sample was recruited through the Norwegian Twin Registry <https://www.fhi.no/en/more/health-studies/norwegian-twin-registry/> ⁷³. The current NTR sub-study was approved by The Regional Committees for Medical and Health Research Ethics (REK South-East C 2018/94). Both MoBa and NTR are linked to The Medical Birth Registry (MBRN), which is a national health registry containing information about all births in Norway. By individual consent, information on BW was also obtained from the MBRN also for some LCBC participants not being part of MoBa or NTR. Approval for these studies was given by The Regional Committees for Medical and Health Research Ethics (REK Sør-Øst B 2017/653). Most participants, including all children, were recruited for observational studies, while some adults were recruited to enter into cognitive training studies after baseline assessment.

Inclusion/Exclusion Criteria/ Screening

Adult participants were screened using a standardized health interview prior to inclusion in the study. Participants with a history of self- or parent-reported neurological or psychiatric conditions, including clinically significant stroke, serious head injury, untreated hypertension, diabetes, and use of psychoactive drugs within the last two years, were excluded. Further, participants reporting worries concerning their cognitive status, including memory function, were excluded. Participants above 40 years scored >26 on the Mini Mental State Examination ⁷⁹. Additionally, a generalized additive mixed model regressing area/thickness/volume nonlinear on age with random intercept per participant, was fitted independently at each of 163842 vertices. 60 observations were defined as outliers and excluded, as the absolute value of their residual was large than four times the residual standard error at more than 6000 vertices.

Variables Used for Birth weight and Education

When possible, by participant consent, birth weight was obtained from the records of the Medical Birth Registry of Norway (MBRN, for participants born 1967 when the Registry was started, or after). MBRN records of birth weight was obtained for at total 526 of the participants. For 15 participants, birth weight records based on historical self-report to the Norwegian Twin Registry were obtained. Otherwise, self-report, or for children, parental report, of birth weight in connection with participation in the MRI scanning studies were used. For analyses controlling for gestational length, only information from the MBRN was used. For education, if multiple values were reported across timepoints, the highest was chosen. Education was recorded as total years of education to the highest obtained degree, for adults, and for participants < 20 years of age, average of parental education was used. However, for some participants in the age range up to 21.3 years at baseline scan, parental education was used, if no report of own education existed. Age was recorded in years and months at the time of the baseline MRI scan.

MRI Scanning and Processing

Brain imaging data were collected across 2 sites on 1.5 T and 3T scanners (Siemens Avanto, Skyra and Prisma (Siemens Corp., Erlanger, Germany, see **Supplementary Table 1** [↗](#)). At baseline 345 participants were scanned at Avanto 1 at Oslo University Hospital, Oslo, and 105 were scanned at Avanto 2 at St. Olav's Hospital, Trondheim, 74 were scanned at Prisma and 111 at Skyra, Oslo University hospital. All were followed up longitudinally at least once at the same scanner. To the extent that scanners were switched for longer follow-up, double scanning (at old and new scanner) was implemented to the extent possible, and both scans were from that time point was entered into analysis. 369 participants did not switch scanner during the study, 266 switched scanner at least once, of which 106 participants were scanned with both scanners at the same time point. MRI data were processed using FreeSurfer 6.0.

Abcd

Population, Recruitment and General Description of Study/ Procedures

The primary aim of the Adolescent Brain Cognitive Development (ABCD) study <https://abcdstudy.org> [↗](#) is to track human brain development from childhood through adolescence to determine biological and environmental factors that impact or alter developmental trajectories ²⁵[↗](#). ABCD has recruited > 10 000 9-10 years olds across 21 US sites with harmonized measures and procedures, including imaging acquisition <https://abcdstudy.org/scientists-workgroups.html> [↗](#). A goal of the ABCD study is that its sample should reflect, as best as possible, the sociodemographic variation of the US population ²⁶[↗](#). Of relevance to the present analyses, children were ineligible to participate if they had any MRI contraindications, or prematurity at birth <28 weeks ⁸[↗](#), ⁸⁰[↗](#). For ABCD, the birth weight values were extracted from the dataset release 2.0.1 at baseline (consisting of a total of 11875 participants), and the remainder of data were extracted from release 3.0.

Inclusion/Exclusion Criteria for present analyses

All participants who had birth weight reported and had underwent MRI scanning at more than one timepoint with FreeSurfer Quality Control OK were included in the main analyses. Additionally, a generalized additive mixed model regressing area/thickness/volume nonlinear on age with random intercept per participant, was fitted independently at each of 163842 vertices. 223 observations were defined as outliers and excluded, as the absolute value of their residual was large than four times the residual standard error at more than 6000 vertices.

Variables used for Birth Weight and Education

We used the fields `devhx_2_birth_wt_lbs_p` and `devhx_2b_birth_wt_oz_p`. These fields were converted to grams, and added together. If `devhx_2b_birth_wt_oz_p` was missing, it is set to 0, so that the only contribution came from `devhx_2_birth_wt_lbs_p`. If `devhx_2_birth_wt_lbs_p` was missing, the final birth weight value was set to NA, and hence the participant was excluded from analysis. In cases where birth weight was reported multiple times for one participant, and values deviated, participants were excluded from analysis if the discrepancy was greater than 10%, else the mean reported number was entered. For the analyses controlling for gestational length, we used the variable `weeks_premature`. Education was entered as the maximum education of the parents in years.

MRI Scanning and Processing

Brain imaging data were collected across 21 sites on 3T scanners (Siemens Prisma (Siemens Corp., Erlanger, Germany), GE Discovery MR750 (GE Healthcare, Chicago, IL), and Philips Achieva (Philips, Amsterdam, the Netherlands). Acquisition parameters are listed in ²⁵ and at https://abcdstudy.org/images/Protocol_Imaging_Sequences.pdf. Images were processed using the FreeSurfer 7.1.0 software package for the analysis of the main longitudinal sample, whereas for the MZ twin analysis (done collectively with UKB and LCBC MZ samples), the FreeSurfer 6.0 software package was used.

Ukb

Population, Recruitment and General Description of Study/Procedures

UK Biobank (UKB) (<https://www.ukbiobank.ac.uk/about-biobank-uk/>) is a major national and international health resource with the aim of improving the prevention, diagnosis and treatment of a wide range of illnesses. UK Biobank recruited ~500,000 people aged between 40-69 years in 2006-2010 from across the country to take part in this project ⁸¹. Potential participants were identified through National Health Service (NHS) registers according to being aged 40-69 and living within a reasonable travelling distance of an assessment centre. Assessment centres (22 in total) are located in accessible and convenient locations with a large surrounding population. Participants have undergone measures and provided samples and detailed information about themselves and agreed to have their health followed. Age was calculated from year and month of birth (day of month is missing, and was set to 1 for all subjects) to date of assessment.

Inclusion/Exclusion Criteria/Screening

Participants are excluded from scanning in the UKB according to fairly standard MRI safety/quality criteria, such as exclusions for metal implants, recent surgery, or health conditions directly problematic for MRI scanning, (e.g. problems hearing, breathing or extreme claustrophobia) ⁸². A generalized additive mixed model regressing area/thickness/volume nonlinear on age with random intercept per participant, was fitted independently at each of 163842 vertices. 40 observations were defined as outliers and excluded, as the absolute value of their residual was large than four times the residual standard error at more than 6000 vertices.

Variables Used for Birth weight and Education

Birth weight was extracted from UKB data field 20022. Units are in kilogram. If multiple birth weights were reported, we removed participants if the discrepancy between reported birth weights were greater than 10%, and we took the mean of the reported birth weights Education: For the Biobank participant's generation, the UK school system provided free universal compulsory education between the ages of 5 and 15 to 16 years. Based on the UKB education data field 6138 (1 College or University degree; 2 A levels/AS levels or equivalent; 3 O levels/GCSEs or equivalent; 4 CSEs or equivalent; 5 NVQ or HND or HNC or equivalent; 6 Other professional

qualifications eg: nursing, teaching; -7 None of the above; -3 Prefer not to answer, education was recoded to years using the following dictionary: edu_ukb_to_years = (1: 16, 2: 13, 3: 11, 4: 11, 5: 11, 6: 12, -7: 10, -3: np.NaN). If multiple education values were reported, the highest education value was chosen.

MRI scanning and processing

Imaging data were collected and processed by the UK Biobank (<https://www.ukbiobank.ac.uk>) as described in ²⁷. Imaging data were collected using 3.0 T Siemens Skyra (32-channel head coil). Anatomical T1-weighted magnetization-prepared rapid gradient echo (MPRAGE) images were obtained in the sagittal plane at 1mm isotropic resolution, and T2 weighted FLAIR images were acquired at 1.05 x 1 x 1mm resolution in the sagittal plane. Images were processed by the UK Biobank using the FreeSurfer 6.0 software package.

Specific information on image acquisition in the different samples

T1 weighted structural scans were acquired at Siemens, Philips and GE scanners at the various sites. The sequence parameters for each cohort are given in **Supplementary Table 1**.

Supplementary Statistical methods

Assessment of consistency of effects across and within samples

We assessed the spatial relationship between birth weight cortical correlates (Beta-maps) for cortical area, volume, and thickness across the different datasets (LCBC, UKB, ABCD) using Pearson's correlations. Permutation-based significance testing ($n = 10,000$, $p < 0.05$, two-tailed, FDR-corrected) was performed using non-0 parametric spatial permutation models, i.e. spin tests as implemented by ⁸³, ⁸⁴. Briefly, spin-tests generate spatially-constrained null distributions by applying random rotations to spherical projections of the brain. For each permutation, the original values at each coordinate are replaced with those of the closest rotated coordinate. Rotations are generated in one hemisphere and mirrored in the other.

Within-sample replicability was assessed in two different ways, an exploratory and a confirmatory analysis³⁶. In the exploratory analysis, we assessed within-sample replicability by conducting the same vertex-wise cortical analysis in different subsamples. For each dataset and cortical measure, we assessed the effects of birth weight on cortical structure $|N| = 500$ times using 50% of the original sample (participants' split half). Beyond sample selection, all parameters remained identical as described in the *main* analysis (refer to **Figure 1**). Replicability – of a given vertex – was determined by the proportion of multiple comparison-corrected results across the different subsamples. In the confirmatory analysis, we assessed the proportion of significant vertices obtained in each *exploratory* (train) analysis that was also deemed significant – and in the same direction – in the remaining (test; 50%) subsample ($p < 0.05$ uncorrected). Subsamples without significant results were not considered (no results only in cortical thickness analyses). This is a criterion often followed for determining replicability³⁷. Replicability analyses were performed on *fsaverage4* for computational reasons.

Supplementary Results


The degree of within-sample replicability was high for both area and volume but not for cortical thickness (see **Supplementary Figure 11**) in the three datasets (LCBC, ABCD, and UKB). For area and volume, some vertices were fully replicable, i.e., significant results were found in this vertex

Sample	Scanner	Tesla	Sequence parameters
LCBC	Avanto	1.5	TR: 2400 ms, TE: 3.61 ms, TI: 1000 ms, flip angle:
	Siemens		8°, slice thickness: 1.2 mm, FoV: 240 × 240 mm, 160 slices, iPat = 2
	Avanto	1.5	TR: 2400 ms, TE = 3.79 ms, TI = 1000 ms, flip
	Siemens		angle = 8, slice thickness: 1.2 mm, FoV: 240 x 240 mm, 160 slices
	Skyra	3.0	TR: 2300 ms, TE: 2.98 ms, TI: 850 ms, flip angle:
	Siemens		8°, slice thickness: 1 mm, FoV: 256 × 256 mm, 176 slices
	Prisma	3.0	TR: 2400 ms, TE: 2.22 ms, TI: 1000 ms, flip angle:
	Siemens		8°, slice thickness: 0.8 mm, FoV: 240 × 256 mm, 208 slices, iPat = 2
ABCD	Prisma	3.0	TR: 2500 ms, TE: 2.88 ms, TI: 1060 ms, flip
	Siemens		angle: 8°, slice thickness: 1 mm, FoV: 256 × 256 mm, 176 slices, parallel imaging = 2
	Achieva/ dStream/ Ingenia Phillips	3.0	TR: 6.31 ms, TE: 2.9 ms, TI: 1060 ms, flip angle: 8°, slice thickness: 1 mm, FoV: 256 × 240 mm, 225 slices, parallel imaging = 1.5 x 2.2
	MR750/ DV25-26 GE	3.0	TR: 2500 ms, TE: 2 ms, TI: 1060 ms, flip angle: 8°, slice thickness: 1 mm, FoV: 256 × 256 mm, 208 slices, parallel imaging = 2x
UKB	Skyra	3.0	TR: 2000 ms, TI: 880 ms, slice thickness: 1 mm,
	Siemens		FoV: 208 × 256 mm, 256 slices, iPAT=2

Supplementary Table 1

MR acquisition parameters


TR: Repetition time, TE: Echo time, TI: Inversion time, FoV: Field of View, iPat: in-plane acceleration, GRAPPA: GRAPPA acceleration factor. *Customized



Supplementary Figure 1.

Relationships of birthweight and cortical area across LCBC, ABCD, and UKB samples when controlling for age, sex, time (interval since baseline) and scanner site (as well as ethnicity in the ABCD).

Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.




Supplementary Figure 2.

Relationships of birthweight and cortical characteristics across LCBC, ABCD, and UKB samples when controlling for education, age, sex, time (interval since baseline) and scanner site (as well as ethnicity in the ABCD).

Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.


Associations of birth weight and cortical characteristics across cohorts when controlling for gestational length



Supplementary Figure 3.

Relationships of birthweight and cortical characteristics across LCBC and ABCD samples when controlling for gestational length in weeks (LCBC) or weeks born prematurely (ABCD), age, sex, time (interval since baseline) and scanner site (as well as ethnicity in the ABCD).


Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.



Supplementary Figure 4.

Relationships of birthweight and cortical characteristics across LCBC, ABCD, and UKB when restricting the samples to participants with birth weights between 2.5 and 5.0 kg, controlling for age, sex, time (interval since baseline) and scanner site (as well as ethnicity in the ABCD).


(Compare to [Figure 1](#).) Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.



Supplementary Figure 5.

Relationships of birthweight and cortical characteristics across LCBC, ABCD, and UKB samples when controlling for intracranial volume (ICV), age, sex, time (interval since baseline) and scanner site (as well as ethnicity in the ABCD).


Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.



Supplementary Figure 6


Interactions of BW and time on cortical characteristics across LCBC, ABCD, and UKB samples when controlling for age, sex, scanner site, time, birth weight, and the interaction of baseline age and time (as well as ethnicity in the ABCD).

Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere.




Supplementary Figure 7.

Plots showing individual data points and expected trajectories for cortical area within the significant regions (refer to **Figures 1** and **2**) of each sample split in two based on BW (higher BW in red color= upper half, lower BW in blue color= lower half of BW distribution) are shown. Y-axis: cortical area: mm2, thickness, X-axis: time in years.




Supplementary Figure 8.

Plots showing individual data points and expected trajectories for cortical thickness within the significant regions (refer to **Figures 1** and **2**) of each sample split in two based on BW (higher BW in red color= upper half, lower BW in blue color= lower half of BW distribution) are shown. Y-axis: cortical thickness: mm, X-axis: time in years.



Supplementary Figure 9.

Plots showing individual data points and expected trajectories for cortical volume within the significant regions (refer to [Figures 1](#) and [2](#)) of each sample split in two based on BW (higher BW in red color= upper half, lower BW in blue color= lower half of BW distribution) are shown. Y-axis: cortical thickness: mm, X-axis: time in years.



Supplementary Figure 10.

Spatial correlation of birth weight effects on brain structure change across datasets.


For each panel, the upper triangular matrix shows Pearson's (r) pairwise spatial correlation between the different cohorts' cortical maps. Data is shown as a color-density plot. The red line represents the fitting between the two maps. The lower triangular matrix shows the significance testing. The dashed-grey line shows the empirical correlation, while the orange histogram represents the null distribution based on the *spin test*. The diagonal shows the effect of birth weight on cortical structure (right hemisphere shown only). Note eta-maps are shown as a percentile red-green-blue scale, where red β represents a lower (or more negative) effect of birth weight on cortical structure and vice versa. See **Supplementary Table 2** for stats. The different panels show the spatial correlation of birth weight effects on cortical a) area, b) thickness, and c) volume. Units in the density maps represent birth weight effects as mm/g, mm²/g, and mm³/g (10e-5) for cortical thickness, area, and volume, respectively.

Cortical measure	Compared Datasets	Birth Weight		Birth weight, ICV-corrected (-log10(p))		Birth Weight, (-log10(p))		Birth Weight x Time	
		β	p	β	p	β	p	β	p
Area	LCBC - UKB	.51	<.001	.24	.010	.35	<.001	-.20	.073
	LCBC - ABCD	.71	<.001	.61	<.001	.60	<.001	-.05	.770
	UKB - ABCD	.63	<.001	.47	<.001	.54	<.001	-.35	<.001
Thick- ness	LCBC - UKB	.24	.011	.25	.006	.23	.005	-.04	.800
	LCBC -ABCD	.45	<.001	.45	<.001	.43	<.001	-.11	.430
	UKB-ABCD	.44	<.001	.44	<.001	.37	<.001	-.20	.150
Volume	LCBC-UKB	.64	<.001	.38	<.001	.45	<.001	.08	.520
	LCBC-ABCD	.79	<.001	.70	<.001	.66	<.001	-.34	.005
	UKB-ABCD	.73	<.001	.56	<.001	.60	<.001	.03	.760

Supplementary Table 2.

Spatial correlation of birth weight effects on brain structure across datasets. Pearson's correlation and significance for pairwise spatial correlations between the different cohorts' birth weight, with and without correction for ICV, and birth weight x time (years) -related cortical maps; i.e. effects of birth weight on cortical structure and change in cortical structure. Significance was assessed using the spin test. Spatial correlations were assessed using the Beta maps (and -log10(p) for birth weight). See [Figure 3](#) and [Supplementary Figure 9](#) and 10 for a visual representation. P-values are FDR-corrected (n = 9).


regardless of sample variability. *Exploratory* replicability [median (interquartile range); LCBC, ABCD, UKB] was .46 (.56), .85 (.47), .89 (.43) for area, .36 (.47), .50 (.65), .79 (.68) for volume, and .008 (.04), .00 (.00), .002 (.02) for thickness. *Exploratory* replicability indexes for each voxel the proportion of significant results found using different $|N| = 500$ subsamples. The confirmatory replicability analyses also showed good replicability for area and volume but not for cortical thickness. Confirmatory replicability indexes the proportion of voxels (per analysis) that was significant both in the exploratory subsample ($p < 0.01$, FWE-corrected) and in the test subsample ($p < 0.05$). The proportion of vertices that were significant both in the *exploratory* and *test* subsamples was .74 (.17), .91 (.08), .93 (.08) for area, .68 (.19), .84 (.12), .92 (.09) for volume, and .28 (.28), .00 (.00), .58 (.18) for thickness. These results are not fully comparable to other studies assessing the replicability of brain-phenotype associations due to analytical differences (e.g. sample size, multiple-comparison correction method)²⁰,³⁶, yet clearly show that the rate of replicability of BW associations with cortical area and volume are comparable to benchmark brain-phenotype associations such as body-mass index and age⁶⁸. Lower levels of replicability in the LCBC subsample are likely attributed to higher sample variability (e.g. increased age span). Kinship may lead to inflated patterns of replicability within the ABCD cohort.



Supplementary Figure 11.

The degree of within-sample replicability of birth weight effects on cortical structure for LCBC, ABCD, and UKB.

Brain images show the exploratory replicability analyses; i.e. the proportion of results (refer to Figure 1) that are significant across different subsamples (50% of the original sample; $|N| = 500$ subsamples). The overlay is thresholded at replicability (p) = .1. For visual purposes data has been up-sampled and smoothed (FWHM = 5) on fsaverage space. Left-side violin plots show the exploratory replicability; i.e., the proportion of significant results across the different subsamples (unit = voxel). Right-side plots show the confirmatory replicability; i.e. for each sample, the proportion of results ($p < 0.01$; FWE-corrected) that also pass significance criteria ($p < 0.05$) in a test subsample ($|N| = 500$ subsamples) (unit = analysis). Embedded boxplots display median, interquartile range, and range.



Supplementary Figure 12.

Interaction effects of birth weight discordance and time on cortical thickness in the sample of monozygotic (MZ) twins.

Significant relationships are shown from left to right: lateral view, right and left hemisphere, and medial view, right and left hemisphere. Plots showing individual data points and expected trajectories for cortical thickness in mm (Y-axes) within the significant regions according to BW discordance (left panel) and BW (right panel) in kilograms (X-axes) are shown.

References

1. Kovacs G. G., et al. (2014) **Linking pathways in the developing and aging brain with neurodegeneration** *Neuroscience* **269**:152–172 <https://doi.org/10.1016/j.neuroscience.2014.03.045>
2. Fjell A. M., et al. (2015) **Development and aging of cortical thickness correspond to genetic organization patterns** *Proceedings of the National Academy of Sciences of the United States of America* **112**:15462–15467 <https://doi.org/10.1073/pnas.1508831112>
3. Bale T. L (2015) **Epigenetic and transgenerational reprogramming of brain development** *Nature reviews. Neuroscience* **16**:332–344 <https://doi.org/10.1038/nrn3818>
4. Muller M., et al. (2014) **Birth size and brain function 75 years later** *Pediatrics* **134**:761–770 <https://doi.org/10.1542/peds.2014-1108>
5. Walhovd K. B., et al. (2016) **Neurodevelopmental origins of lifespan changes in brain and cognition** *Proceedings of the National Academy of Sciences of the United States of America* **113**:9357–9362 <https://doi.org/10.1073/pnas.1524259113>
6. Walhovd K. B., et al. (2012) **Long-term influence of normal variation in neonatal characteristics on human brain development** *Proceedings of the National Academy of Sciences of the United States of America* **109**:20089–20094 <https://doi.org/10.1073/pnas.1208180109>
7. Vidal-Pineiro D., et al. (2021) **Individual variations in ‘brain age’ relate to early-life factors more than to longitudinal brain change** *Elife* **10** <https://doi.org/10.7554/eLife.69995>
8. Dooley N., Clarke M., Cotter D., Cannon M (2022) **Birth Weight and Childhood Psychopathology in the ABCD Cohort: Association is Strongest for Attention Problems and is Moderated by Sex** *Res Child Adolesc Psychopathol* **50**:563–575 <https://doi.org/10.1007/s10802-021-00859-0>
9. Elliott M. L., et al. (2021) **Disparities in the pace of biological aging among midlife adults of the same chronological age have implications for future frailty risk and policy** *Nat Aging* **1**:295–308 <https://doi.org/10.1038/s43587-021-00044-4>
10. Anderson P. J, et al. (2021) **Psychiatric disorders in individuals born very preterm / very low-birth weight: An individual participant data (IPD) meta-analysis** *Eclinicalmedicine* **42** <https://doi.org/10.1016/j.eclinm.2021.101216>
11. Tuovinen S., et al. (2013) **Maternal hypertensive disorders in pregnancy and self-reported cognitive impairment of the offspring 70 years later: the Helsinki Birth Cohort Study** *American journal of obstetrics and gynecology* **208**:e201–209 <https://doi.org/10.1016/j.ajog.2012.12.017>
12. Beaumont R. N., et al. (2018) **Genome-wide association study of offspring birth weight in 86 577 women identifies five novel loci and highlights maternal genetic effects that are independent of fetal genetics** *Human molecular genetics* **27**:742–756 <https://doi.org/10.1093/hmg/ddx429>

13. Li X., et al. (2021) **Association of low birth weight with cardiometabolic diseases in Swedish twins: a population-based cohort study** *BMJ Open* **11** <https://doi.org/10.1136/bmjopen-2020-048030>
14. Willis M. D., et al. (2022) **A population-based cohort study of traffic congestion and infant growth using connected vehicle data** *Sci Adv* **8** <https://doi.org/10.1126/sciadv.abp8281>
15. Raznahan A., Greenstein D., Lee N. R., Clasen L. S., Giedd J. N (2012) **Prenatal growth in humans and postnatal brain maturation into late adolescence** *Proceedings of the National Academy of Sciences of the United States of America* **109**:11366–11371 <https://doi.org/10.1073/pnas.1203350109>
16. Casey K. F., et al. (2017) **Birth weight discordance, DNA methylation, and cortical morphology of adolescent monozygotic twins** *Human brain mapping* **38**:2037–2050 <https://doi.org/10.1002/hbm.23503>
17. Halevy T., et al. (2021) **Twin discordance: a study of volumetric fetal brain MRI and neurodevelopmental outcome** *Eur Radiol* **31**:6676–6685 <https://doi.org/10.1007/s00330-021-07773-5>
18. Wheeler E., et al. (2021) **Birth weight is associated with brain tissue volumes seven decades later but not with MRI markers of brain ageing** *Neuroimage Clin* **31** <https://doi.org/10.1016/j.nicl.2021.102776>
19. Eriksson J. G., Forsen T., Tuomilehto J., Osmond C., Barker D. J (2000) **Early growth, adult income, and risk of stroke** *Stroke* **31**:869–874 <https://doi.org/10.1161/01.str.31.4.869>
20. Marek S., et al. (2022) **Reproducible brain-wide association studies require thousands of individuals** *Nature* **603**:654–660 <https://doi.org/10.1038/s41586-022-04492-9>
21. Walhovd K. B., et al. (2020) **Genetic risk for Alzheimer disease predicts hippocampal volume through the human lifespan** *Neurol Genet* **6** <https://doi.org/10.1212/NXG.0000000000000506>
22. Katzman R., et al. (1988) **Clinical, pathological, and neurochemical changes in dementia: a subgroup with preserved mental status and numerous neocortical plaques** *Annals of neurology* **23**:138–144 <https://doi.org/10.1002/ana.410230206>
23. Schofield P. W., Logroscino G., Andrews H. F., Albert S., Stern Y (1997) **An association between head circumference and Alzheimer's disease in a population-based study of aging and dementia** *Neurology* **49**:30–37 <https://doi.org/10.1212/wnl.49.1.30>
24. Nyberg L., Lovden M., Riklund K., Lindenberger U., Backman L. (2012) **Memory aging and brain maintenance** *Trends in cognitive sciences* **16**:292–305 <https://doi.org/10.1016/j.tics.2012.04.005>
25. Casey B. J., et al. (2018) **The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites** *Developmental cognitive neuroscience* **32**:43–54 <https://doi.org/10.1016/j.dcn.2018.03.001>
26. Garavan H., et al. (2018) **Recruiting the ABCD sample: Design considerations and procedures** *Developmental cognitive neuroscience* **32**:16–22 <https://doi.org/10.1016/j.dcn.2018.04.004>

27. Alfaro-Almagro F., et al. (2018) **Image processing and Quality Control for the first 10,000 brain imaging datasets from UK Biobank** *NeuroImage* **166**:400–424 <https://doi.org/10.1016/j.neuroimage.2017.10.034>
28. Nobis L., et al. (2019) **Hippocampal volume across age: Nomograms derived from over 19,700 people in UK Biobank** *Neuroimage Clin* **23** <https://doi.org/10.1016/j.nicl.2019.101904>
29. Dale A. M., Fischl B., Sereno M. I. (1999) **Cortical surface-based analysis. I. Segmentation and surface reconstruction** *NeuroImage* **9**:179–194 <https://doi.org/10.1006/nimg.1998.0395>
30. Fischl B., et al. (2002) **Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain** *Neuron* **33**:341–355
31. Reuter M., Schmansky N. J., Rosas H. D., Fischl B (2012) **Within-subject template estimation for unbiased longitudinal image analysis** *NeuroImage* **61**:1402–1418 <https://doi.org/10.1016/j.neuroimage.2012.02.084>
32. Jovicich J., et al. (2013) **Brain morphometry reproducibility in multi-center 3T MRI studies: a comparison of cross-sectional and longitudinal segmentations** *Neuroimage* **83**:472–484 <https://doi.org/10.1016/j.neuroimage.2013.05.007>
33. Rimol L. M., et al. (2019) **Reduced white matter fractional anisotropy mediates cortical thickening in adults born preterm with very low birthweight** *NeuroImage* **188**:217–227 <https://doi.org/10.1016/j.neuroimage.2018.11.050>
34. Nyberg L., et al. (2010) **Longitudinal evidence for diminished frontal cortex function in aging** *Proceedings of the National Academy of Sciences of the United States of America* **107**:22682–22686 <https://doi.org/10.1073/pnas.1012651108>
35. Wolke D., et al. (2009) **Selective drop-out in longitudinal studies and non-biased prediction of behaviour disorders** *Br J Psychiatry* **195**:249–256 <https://doi.org/10.1192/bjp.bp.108.053751>
36. Masouleh Kharabian, Eickhoff S., Hoffstaedter S. B., Genon F. (2019) **Empirical examination of the replicability of associations between brain structure and psychological variables** *Elife* **8** <https://doi.org/10.7554/eLife.43464>
37. Open Science, C. PSYCHOLOGY (2015) **Estimating the reproducibility of psychological science** *Science* **349** <https://doi.org/10.1126/science.aac4716>
38. Livingston G., et al. (2020) **Dementia prevention, intervention, and care: 2020 report of the Lancet Commission** *Lancet* **396**:413–446 [https://doi.org/10.1016/S0140-6736\(20\)30367-6](https://doi.org/10.1016/S0140-6736(20)30367-6)
39. Livingston G, et al. (2017) **Dementia prevention, intervention, and care** *Lancet* **390**:2673–2734 [https://doi.org/10.1016/S0140-6736\(17\)31363-6](https://doi.org/10.1016/S0140-6736(17)31363-6)
40. Liu S., Seidlitz J., Blumenthal J. D., Clasen L. S., Raznahan A (2020) **Integrative structural, functional, and transcriptomic analyses of sex-biased brain organization in humans** *Proceedings of the National Academy of Sciences of the United States of America* **117**:18788–18798 <https://doi.org/10.1073/pnas.1919091117>
41. Walhovd K. B., et al. (2022) **Education and Income Show Heterogeneous Relationships to Lifespan Brain and Cognitive Differences Across European and US Cohorts** *Cerebral cortex* **32**:839–854 <https://doi.org/10.1093/cercor/bhab248>

42. Couto Alves A., et al. (2019) **GWAS on longitudinal growth traits reveals different genetic factors influencing infant, child, and adult BMI** *Sci Adv* **5** <https://doi.org/10.1126/sciadv.aaw3095>
43. Momany A. M., Kamradt J. M., Nikolas M. A (2018) **A Meta-Analysis of the Association Between Birth Weight and Attention Deficit Hyperactivity Disorder** *J Abnorm Child Psych* **46**:1409–1426 <https://doi.org/10.1007/s10802-017-0371-9>
44. Patel Y., et al. (2022) **Virtual Ontogeny of Cortical Growth Preceding Mental Illness** *Biological psychiatry* **92**:299–313 <https://doi.org/10.1016/j.biopsych.2022.02.959>
45. Carretta T. R., Ree M. J (2022) **Correction for range restriction: Lessons from 20 research scenarios** *Mil Psychol* **34**:551–569 <https://doi.org/10.1080/08995605.2021.2022067>
46. Franke K., Gaser C., Roseboom T. J., Schwab M., de Rooij S. R (2018) **Premature brain aging in humans exposed to maternal nutrient restriction during early gestation** *NeuroImage* **173**:460–471 <https://doi.org/10.1016/j.neuroimage.2017.10.047>
47. Wagen A. Z., et al. (2022) **Life course, genetic, and neuropathological associations with brain age in the 1946 British Birth Cohort: a population-based study** *Lancet Healthy Longev* **3**:e607–e616 [https://doi.org/10.1016/S2666-7568\(22\)00167-2](https://doi.org/10.1016/S2666-7568(22)00167-2)
48. Kharabian Masouleh S., et al. (2022) **Empirical facts from search for replicable associations between cortical thickness and psychometric variables in healthy adults** *Sci Rep* **12** <https://doi.org/10.1038/s41598-022-17556-7>
49. Schnack H (2019) **Assessing reproducibility in association studies** *Elife* **8** <https://doi.org/10.7554/eLife.46757>
50. Stamatakis E., et al. (2021) **Is Cohort Representativeness Passe? Poststratified Associations of Lifestyle Risk Factors with Mortality in the UK Biobank** *Epidemiology* **32**:179–188 <https://doi.org/10.1097/EDE.0000000000001316>
51. Mills K. L., et al. (2016) **Structural brain development between childhood and adulthood: Convergence across four longitudinal samples** *NeuroImage* **141**:273–281 <https://doi.org/10.1016/j.neuroimage.2016.07.044>
52. Storsve A. B., et al. (2014) **Differential longitudinal changes in cortical thickness, surface area and volume across the adult life span: regions of accelerating and decelerating change** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **34**:8488–8498 <https://doi.org/10.1523/JNEUROSCI.0391-14.2014>
53. Tamnes C. K., et al. (2010) **Brain maturation in adolescence and young adulthood: regional age-related changes in cortical thickness and white matter volume and microstructure** *Cerebral cortex* **20**:534–548 <https://doi.org/10.1093/cercor/bhp118>
54. Schmitz L. L., Duque V (2022) **In utero exposure to the Great Depression is reflected in late-life epigenetic aging signatures** *Proceedings of the National Academy of Sciences of the United States of America* **119** <https://doi.org/10.1073/pnas.2208530119>
55. Barker D. J (1990) **The fetal and infant origins of adult disease** *Bmj* **301** <https://doi.org/10.1136/bmj.301.6761.1111>

56. Barker D. J (1995) **Fetal origins of coronary heart disease** *Bmj* **311**:171–174 <https://doi.org/10.1136/bmj.311.6998.171>
57. Greenough W. T., Black J. E., Wallace C. S. (1987) **Experience and brain development** *Child development* **58**:539–559
58. Huttenlocher P. R (1979) **Synaptic density in human frontal cortex – developmental changes and effects of aging** *Brain research* **163**:195–205
59. Petanjek Z., et al. (2011) **Extraordinary neoteny of synaptic spines in the human prefrontal cortex** *Proceedings of the National Academy of Sciences of the United States of America* **108**:13281–13286 <https://doi.org/10.1073/pnas.1105108108>
60. Yeung M. S., et al. (2014) **Dynamics of oligodendrocyte generation and myelination in the human brain** *Cell* **159**:766–774 <https://doi.org/10.1016/j.cell.2014.10.011>
61. Rakic P (1988) **Specification of cerebral cortical areas** *Science* **241**:170–176
62. Bhardwaj R. D., et al. (2006) **Neocortical neurogenesis in humans is restricted to development** *Proceedings of the National Academy of Sciences of the United States of America* **103**:12564–12568 <https://doi.org/10.1073/pnas.0605177103>
63. Paredes M. F., Sorrells S. F., Garcia-Verdugo J. M., Alvarez-Buylla A (2016) **Brain size and limits to adult neurogenesis** *The Journal of comparative neurology* **524**:646–664 <https://doi.org/10.1002/cne.23896>
64. Rehn A. E., et al. (2004) **An animal model of chronic placental insufficiency: relevance to neurodevelopmental disorders including schizophrenia** *Neuroscience* **129**:381–391 <https://doi.org/10.1016/j.neuroscience.2004.07.047>
65. WHO (2019) **Risk reduction of cognitive decline and dementia: WHO guidelines** *World Health Organization*
66. McGovern M. E (2019) **How much does birth weight matter for child health in developing countries? Estimates from siblings and twins** *Health Econ* **28**:3–22 <https://doi.org/10.1002/hec.3823>
67. Azuine R. E., et al. (2019) **Prenatal Risk Factors and Perinatal and Postnatal Outcomes Associated With Maternal Opioid Exposure in an Urban, Low-Income Multiethnic US Population.** *JAMA Netw Open* **2** <https://doi.org/10.1001/jamanetworkopen.2019.6405>
68. Knickmeyer R. C., et al. (2017) **Impact of Demographic and Obstetric Factors on Infant Brain Volumes: A Population Neuroscience Study** *Cerebral cortex* **27**:5616–5625 <https://doi.org/10.1093/cercor/bhw331>
69. Thompson D. K., et al. (2007) **Perinatal risk factors altering regional brain structure in the preterm infant** *Brain : a journal of neurology* **130**:667–677 <https://doi.org/10.1093/brain/awl277>
70. Thompson D. K., et al. (2020) **Tracking regional brain growth up to age 13 in children born term and very preterm** *Nat Commun* **11** <https://doi.org/10.1038/s41467-020-14334-9>

71. Zhang J., Merialdi M., Platt L. D., Kramer M. S (2010) **Defining normal and abnormal fetal growth: promises and challenges** *American journal of obstetrics and gynecology* **202**:522–528 <https://doi.org/10.1016/j.ajog.2009.10.889>
72. Magnus P., et al. (2016) **Cohort Profile Update: The Norwegian Mother and Child Cohort Study (MoBa)** *International journal of epidemiology* **45**:382–388 <https://doi.org/10.1093/ije/dyw029>
73. Nilsen T., Brandt I., Harris J. R (2019) **The Norwegian Twin Registry** *Twin research and human genetics : the official journal of the International Society for Twin Studies* **22**:647–650 <https://doi.org/10.1017/thg.2019.59>
74. Nilsen T. S., Kutschke J., Brandt I., Harris J. R (2017) **Validity of Self-Reported Birth Weight: Results from a Norwegian Twin Sample** *Twin research and human genetics : the official journal of the International Society for Twin Studies* **20**:406–413 <https://doi.org/10.1017/thg.2017.44>
75. Burt J. B., Helmer M., Shinn M., Anticevic A., Murray J. D (2020) **Generative modeling of brain maps with spatial autocorrelation** *NeuroImage* **220** <https://doi.org/10.1016/j.neuroimage.2020.117038>
76. Markello R. D., Misic B (2021) **Comparing spatial null models for brain maps** *NeuroImage* **236** <https://doi.org/10.1016/j.neuroimage.2021.118052>
77. Viladomat J., Mazumder R., McInturff A., McCauley D. J., Hastie T (2014) **Assessing the significance of global and local correlations under spatial autocorrelation: a nonparametric approach** *Biometrics* **70**:409–418 <https://doi.org/10.1111/biom.12139>
78. Magnus P., et al. (2006) **Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa)** *International journal of epidemiology* **35**:1146–1150 <https://doi.org/10.1093/ije/dyl170>
79. Folstein M. F., Folstein S. E., McHugh P. R (1975) **Mini-mental state". A practical method for grading the cognitive state of patients for the clinician** *J Psychiatr Res* **12**:189–198
80. Palmer C. E., et al. (2021) **A Comprehensive Overview of the Physical Health of the Adolescent Brain Cognitive Development Study Cohort at Baseline** *Front Pediatr* **9** <https://doi.org/10.3389/fped.2021.734184>
81. Guggenheim J. A., Williams C., Consortium U. B. E. V (2015) **Role of Educational Exposure in the Association Between Myopia and Birth Order** *Jama Ophthalmol* **133**:1408–1414 <https://doi.org/10.1001/jamaophthalmol.2015.3556>
82. Miller K. L., et al. (2016) **Multimodal population brain imaging in the UK Biobank prospective epidemiological study** *Nature neuroscience* **19**:1523–1536 <https://doi.org/10.1038/nn.4393>
83. Alexander-Bloch A. F., et al. (2018) **On testing for spatial correspondence between maps of human brain structure and function** *NeuroImage* **178**:540–551 <https://doi.org/10.1016/j.neuroimage.2018.05.070>
84. Lefevre J., et al. (2018) **SPANOL (SPECTral ANALysis of Lobes): A Spectral Clustering Framework for Individual and Group Parcellation of Cortical Surfaces in Lobes** *Front Neurosci* **12** <https://doi.org/10.3389/fnins.2018.00354>

Article and author information

Kristine B. Walhovd

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway, Department of Radiology and Nuclear Medicine, Oslo University Hospital, Norway

For correspondence: k.b.walhovd@psykologi.uio.no

ORCID iD: [0000-0003-1918-1123](https://orcid.org/0000-0003-1918-1123)

Stine Kleppe Krogsrud

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0001-8402-3688](https://orcid.org/0000-0001-8402-3688)

Inge K. Amlie

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0002-8508-9088](https://orcid.org/0000-0002-8508-9088)

Øystein Sørensen

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0003-0724-3542](https://orcid.org/0000-0003-0724-3542)

Yunpeng Wang

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0001-9831-1090](https://orcid.org/0000-0001-9831-1090)

Anne Cecilie Sjøli Bråthen

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

Knut Overbye

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0003-0809-960X](https://orcid.org/0000-0003-0809-960X)

Jonas Kransberg

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

Athanasia M. Mowinckel

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0002-5756-0223](https://orcid.org/0000-0002-5756-0223)

Fredrik Magnussen

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

Martine Herud

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

Asta K. Håberg

Department of Neuromedicine and Movement Science, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology

ORCID iD: [0000-0002-9007-1202](https://orcid.org/0000-0002-9007-1202)

Anders M. Fjell

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway, Department of Radiology and Nuclear Medicine, Oslo University Hospital, Norway

ORCID iD: [0000-0003-2502-8774](https://orcid.org/0000-0003-2502-8774)

Didac Vidal-Piñeiro

Center for Lifespan Changes in Brain and Cognition, University of Oslo, Norway

ORCID iD: [0000-0001-9997-9156](https://orcid.org/0000-0001-9997-9156)

Copyright

© 2023, Walhovd et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Alex Fornito

Monash University, Clayton, Australia

Senior Editor

Christian Büchel

University Medical Center Hamburg-Eppendorf, Hamburg, Germany

Reviewer #1 (Public Review):

This manuscript uses 3 large neuroimaging datasets - which together span childhood to late adulthood - to model the relationship between birthweight (BW) and cortical anatomy over time. The authors separately consider BW associations with the "height" of cortical anatomy trajectories (intercept effects) vs. BW associations with trajectory shape. The authors also distinguish between BW associations with cortical surface area (SA) and cortical thickness (CT), which together determine cortical volume (CV). Prior studies have firmly established robust positive associations between BW and cortical SA, but this study adds evidence for the protracted lifespan persistence of these associations, and the degree to which BW associations with cortical change over time are much weaker.

The study has several strengths including: clear motivation of this work in the Introduction and contextualization of the results in Discussion; use of three large neuroimaging datasets; inclusion of sensible sensitivity analyses; disambiguation of SA and CT findings; and use of formal spatial analysis to quantify the reproducibility of effects across cohorts.

The primary way in which this work seeks to extend beyond established findings is to determine if BW is associated with differences in cortical change over time. The results presented clearly establish that such BW-change associations are much more localized and less consistent across cohorts than BW-intercept associations. However, the evidential basis for this statement is partly limited by the nature of the neuroimaging cohorts used and the specific approaches taken to statistical modeling. Interpretation of findings for both BW-change and BW-intercept associations would also be assisted by greater clarity regarding the specification of statistical models, and the provision of effect-size maps.

Moreover, several factors complicate interpretation of the BW effects on cortical change - which are arguably the main way in which this work could extend on established knowledge of BW associations with brain anatomy. Under the study design presented, inferences regarding age-varying BW effects come from two main sources ... age effects which are quantitatively modeled within each sample, and qualitative differences in age effects between samples. Any inferences from the latter source of evidence are weakened by the fact that (i) no direct statistical comparisons are conducted between samples (beyond the spin tests), and (ii) the composition of samples with regard to age span covered (e.g. 2 in ABCD vs longer in UKB and longest in LCBC) and density of longitudinal data makes it hard to know if between-samples differences in age*BW effects are about biology or methodology. Inferences about age*BW effects from models within each sample are also limited by the fact that (i) some samples (ABCD) have very narrow age ranges precluding detection of age-related effects, and (ii) the modeling strategy used does not allow for non-linear interactions between age and BW or linear interactions that occur in the context of e.g. non-linear BW effects. For this last concern, it would be helpful to know that there is no evidence in the data for such non-linear effects

The tests for spatial consistency between BW effects are a valuable aspect of the manuscript and provide a solid quantitative test for the main effects of BW. For the reasons detailed above however, I think that the more variable (and sometimes negative) correlations in age*BW maps are harder to interpret. One could argue for example that bivariate spline models of age*BW interactions on a lifespan dataset assembled from different COMBAT-aligned cohorts would provide a more solid basis for inference regarding the degree to which BW effects on cortical anatomy vary with age

Overall, this work provides a valuable new data point in our understanding of the profound and protracted influences that prenatal developmental features can have on postnatal outcomes.

- <https://doi.org/10.7554/eLife.86812.1.sa2>

Reviewer #2 (Public Review):

This study focuses on the association between weight at birth and area, volume and thickness of the cerebral cortex measured at timepoints throughout the lifespan. Overall, the study is well designed, and supported by evidence from a large sample drawn from three geographically distinct cohorts with robust analytical and statistical methods.

The authors test three hypotheses: (1) that higher birth weight is associated with greater cortical area in later life; (2) that associations are robust across samples and age; and (3) that associations are stable across the lifespan. Analyses are performed separately in three cohorts: ABCD, UKBB and LCBC and the pattern of associations compared by means of spatial correlations. They find that BW is positively associated with cortical area (and, as a consequence, cortical volume) across most of the cortex, with effect sizes being greatest in frontal and temporal regions. These associations remain largely unchanged when accounting for age, sex, length of gestation and (in one cohort) ethnicity. Variations due to MRI scanner and site are accounted for statistically. Measures are taken to determine within sample replicability through split-half analyses.

The authors conclude that BW, as a marker of early development, is consistently associated with brain characteristics throughout the lifespan, acting as an 'intercept' and promoting brain reserve, i.e.: the capacity of the brain to withstand aging effects. Indeed, the authors calculate that 600g lower BW results in reductions in cortical volume akin to 6-7 years of aging in middle to later life. This is perhaps a startling statistic but one that is not entirely supported by the data presented.

A key piece of information lacking from this study is the functional importance of the reported associations. That lower BW is associated with lower cortical volume and that cortical volume decreases with age is perhaps not surprising - the same could be said for height - one cannot conclude that the same processes underpin the two factors without examining the functional consequences of BW-related volume reductions in older age. The notion of 'brain reserve' indicates a protective effect. If this is the case, one might expect to see a mediating effect of BW on age-related cognitive effects. Without this data, it is difficult to reach the authors' conclusions that decreased birthweight has the same effect as 7 years of aging in later life.

In addition, it is not clear to what degree the association between BW and cortical area/volume is simply reflecting overall somatic growth: brain mass scales with body height, and birth weight and length are associated with adult height. While the specificity of the associations between cortical area/volume and BW are not fully tested, the effects are significantly diminished when controlling for a related measure of somatic growth: intracranial volume (Fig S5). In this context, additional commentary on the specificity of the reported BW-brain associations (or lack thereof) would be helpful.

Finally, the authors use linear models to model brain area, thickness and volume as a function of age. The authors' previous studies have demonstrated nonlinear trajectories of cortical thickness in the LCBC cohort across most of the cortex. A stronger rationale (e.g.: theoretical or model selection) supporting the use of GLM in this study would be more compelling.

- <https://doi.org/10.7554/eLife.86812.1.sa1>

Author Response

First, we would like to thank eLife and the reviewers for the positive assessment of our manuscript and for providing the medium to expand the scientific dialogue on the present study. We greatly appreciate the reviewers' assessment and will revise the manuscript considering their input. Please see our provisional response below.

Reviewer #1's main concern is centered on the evidential strength of the study's conclusion that age-specific effects of birth weight on brain structure are more localized and less consistent across cohorts than age-uniform, stable effects. More specifically, the reviewer points out the evidence (or lack of such) for age-specific effects. The reviewer specifies four methodological concerns: that #1 no direct statistical comparisons are conducted between samples (beyond the spin-tests) and that #2 the differential composition of samples in terms of age distribution leads to the possibility that lack of results is explained by methodological differences. Further, he/she adds that #3 some datasets have a narrow age range precluding detection of age-related effects and #4 that the modeling strategy does not allow for non-linear interaction between age and BW suggesting the use of spline models instead in a _mega_ analytical fashion. Reviewer #1 also asks for greater clarity regarding the statistical models and the provision of effect-size maps.

As a response to the reviewer's comments, we will submit a revised version of the manuscript with a revised description of the statistical models and submit effect-size maps in an accompanying repository. We will tackle the first two concerns by performing additional statistical analyses. The planned analyses will include across-sample reliability and within-sample reliability for the *time*birth weight* analyses. These analyses will address the concerns that the lack of age-specific effects is due to sample differences rather than a lack of biological effects (#2) and provide further explicit statistical comparisons across samples (#1).

We do not believe concern #3 is critically problematic since *time*birth weight* refers to a within-subject contrast, e.g. longitudinal-only based contrast. Birth weight, even when self-reported is a highly reliable measure and the sample sizes are relatively large ($n = 635, 1759$, and 3324 unique individuals). Note that the smaller dataset does have longer follow-up times and more observations per participant increasing the reliability of estimations in individual change. Structural MRI measures have very high reliability (often $ICC > .95$). Clearly, longitudinal brain change is less reliable, yet the present sample size and the high reliability of birth weight should provide enough statistical power to capture even small time-varying effects of birth weight on brain structure.

We agree that some - if not most - brain structures follow non-linear trajectories throughout life (#4). In the present study, age regressors are used only for accounting for variance in the data rather than capturing any effect of interest. Rather, it is the *time*birth weight* regressor that captures age-varying changes in brain structure. *Time* reflects within-subject follow-up time. In any case, the inclusion of non-linear regressors is superfluous in ABCD given the small age range and will not regress out additional variance. Likewise, published data on UKB has shown that most age trajectories of cortical data follow grossly linear fits (age range in the UKB: 45 – 85 years). In LCBC, however, we recognize that non-linear modeling may be able to capture additional variance since it is a lifespan dataset. This may provide better estimates for *birth weight* and *time*birth weight* though it is unlikely that it has a big effect on the *slope*. In any case, #4 is a valid concern and we will repeat the main analysis using «spline models to fit age We will not follow the reviewer's suggestion for a *mega-analytical* approach. It is a valid approach, but it is best suited to other configurations of data than those used here where we used longitudinal data only. The use of cross-sectional data for deriving differential age trajectories is also not without problems.

We see Reviewer #2's concerns mainly being: #1 The translation of birth-weight effects on brain volume to years of aging is inadequate and not fully supported by the data. #2 Lack of data regarding the functional relevance of brain weight effects on brain structure; suggesting the inclusion of cognitive data and (birth weight – brain structure – cognition) mediation analyses. #3 The degree to which the association between birth weight and cortical area/volume is explained by overall somatic growth. #4 The lack of non-linear regressors for fitting age.

#1 We will modify the section on translation of birth weight effects into years of aging. While it is often used - see for example the literature on brain age - and is relatable, it can be easily misinterpreted and does not reflect any objective measure of effect size. #2 The degree to which birth weight relates to cognition throughout the lifespan through individual variations in brain structure is a key question in the field for which currently we only have indirect evidence. Unfortunately, the study of this triple relationship is out of the scope of the current study. This, we agree with the reviewer, warrants further research. We disagree, however, in that mediation analyses are able to provide a satisfactory answer to this question. Also, we believe that the conclusions of the present study are not affected by the omission of cognitive data beyond what has been commented on in #1. The relationship between birth weight, structure, and cognition will be further discussed in the revised version of the manuscript.

#3 Indeed, part of birth weight effects on brain structure can be explained by overall somatic growth. This study does not provide - and is not designed to provide - a specific mechanism for which birth weight is associated with brain structure. Yet, research has also shown that differences in birth weight partially reflect factors associated with variations in neurodevelopmental processes in gestation. Mechanisms are thought to be partially genetic (including genetically influenced potential for somatic growth) and partially environmental prenatal influences. The amount of variance explained by such mechanisms is unknown and may differ across samples (e.g. population samples vs. monozygotic twins). In our view, the association is no less meaningful if mostly attributed to somatic growth if it is associated with

partly modifiable variance in brain growth. The revised manuscript will include further considerations on the possible mechanisms explaining the association between birth weight and cortical structure throughout the lifespan.

Concern #4 is equivalent to Reviewer #1, concern #4. In short, the age regressor is used only to account for variance. Linear modeling in the ABCD and UKB datasets has been shown to fit relatively well the different cortical regions due to the samples' narrow age range. In LCBC, however, we recognize that non-linear modeling may be able to capture additional variance since it is a lifespan dataset. This may provide better estimates for *birth weight* and *time*birth weight* though it is unlikely that has a big effect on the *slope* of the effects. We will repeat the main analysis using «spline models to fit age.