

Effort cost of harvest affects decisions and movement vigor of marmosets during foraging

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

November 3, 2023 (this version)

Reviewed preprint version 1

July 5, 2023

Sent for peer review

March 14, 2023

Posted to bioRxiv

February 6, 2023

Paul Hage , In Kyu Jang, Vivian Looi, Mohammad Amin Fakharian, Simon P. Orozco, Jay S. Pi, Ehsan Sedaghat-Nejad, Reza Shadmehr 

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Our decisions are guided by how we perceive the value of an option, but this evaluation also affects how we move to acquire that option. Why should economic variables such as reward and effort alter the vigor of our movements? In theory, both the option that we choose and the vigor with which we move contribute to a measure of fitness in which the objective is to maximize rewards minus efforts, divided by time. To explore this idea, we engaged marmosets in a foraging task in which on each trial they decided whether to work by making saccades to visual targets, thus accumulating food, or to harvest by licking what they had earned. We varied the effort cost of harvest by moving the food tube with respect to the mouth. Theory predicted that the subjects should respond to the increased effort costs by choosing to work longer, stockpiling food before commencing harvest, but reduce their movement vigor to conserve energy. Indeed, in response to an increased effort cost of harvest, marmosets extended their work duration, but slowed their movements. These changes in decisions and movements coincided with changes in pupil size. As the effort cost of harvest declined, work duration decreased, the pupils dilated, and the vigor of licks and saccades increased. Thus, when acquisition of reward became effortful, the pupils constricted, the decisions exhibited delayed gratification, and the movements displayed reduced vigor.

Significance statement

Our results suggest that as the brainstem neuromodulatory circuits that control pupil size respond to effort costs, they alter computations in the brain regions that control decisions, encouraging work and delaying gratification, and the brain regions that control movements, reducing vigor and suppressing energy expenditure. This coordinated response suggests that decisions and actions are part of a single control policy that aims to maximize a variable relevant to fitness: the capture rate.

eLife assessment

This **important** study unravels the interaction between effort cost, pupil-indexed brain state, and movement (saccadic) vigor during foraging decisions in marmoset monkeys. Based on a normative computational model, the authors derive the prediction that anticipated effort should affect both decisions and movement vigor during foraging; and then provide **solid** behavioural and pupillometric evidence for this prediction in a foraging task. This paper will be of interest to decision and motor neuroscience as well as to all researchers studying animal behavior.

Introduction

During foraging, animals work to locate a food cache and then spend effort harvesting what they have found. As they forage, their decisions appear to maximize a measure that is relevant to fitness: the sum of rewards acquired, minus efforts expended, divided by time, termed the capture rate (1–3). For example, a crow will spend effort extracting a clam from a sandy beach, but if the clam is small, it will abandon it because the additional time and effort required to extract the small reward –dropping it repeatedly from a height onto rocks – can be better spent finding a bigger prize (4). In other words, if going to the bank entails waiting in a long line, one should go infrequently, but make each transaction a large amount.

Intriguingly, reward expectation not only affects decisions, it also affects movements: we not only prefer the less effortful option, we also move vigorously to obtain it (5, 6). This modulation of movement vigor can be justified if we consider that movements require expenditure of time and energy, which discount the value of the promised reward (7, 8). Thus, from a theoretical framework, it seems rational that the brain should have a mechanism to coordinate control of decisions with control of movements so that both contribute to maximizing a measure of fitness (9).

To study this coordination, we designed a task in which marmosets decided how long to work before they harvested their food. On a given trial, they made a sequence of saccades to visual targets and received an increment of food as their reward. However, the increment was small, and its harvest was effortful, requiring inserting their tongue inside a small tube. Theory predicted that in order to maximize the capture rate, harvest should commence only when there was sufficient reward accumulated to justify the effort required for its extraction. Indeed, the subjects chose to work in order to stockpile food, and only then initiated their harvest.

On some days the effort cost of harvest was low: the tube was placed close to the mouth. On other days the same amount of work, i.e., saccade trials, produced food that had a higher effort cost: the tube was located farther away. The theory made two interesting predictions: as the effort cost increased, the subjects should choose to work more trials, thus delaying their harvest so to stow more food, but reduce their movement vigor, thus saving energy. Indeed, when marmosets encountered an increased effort cost, they extended their work period, stockpiling food, but reduced their vigor, slowing their saccades during the work period, and slowing their licks during the harvest period.

What might be a neural basis for this coordinated response of the decision-making and the motor-control circuits? During the work and the harvest periods, momentary changes in pupil size closely tracked the changes in vigor: pupil dilation accompanied increases in vigor, while pupil constriction accompanied decreases in vigor. Remarkably, this was true regardless of whether the

movement that was being performed was a saccade, or a lick. Moreover, in response to the increased effort cost, the pupils exhibited a global change, constricting during both the work and the harvest periods.

If we view the changes in pupil size as a proxy for activity in the brainstem noradrenergic circuits (10 [↗](#)), our results suggest that as these circuits respond to effort costs (11 [↗](#)), they alter computations in the brain regions that control decisions, delaying gratification and encouraging work, and the brain regions that control movements, promoting sloth and conserving energy.

Results

We tracked the eyes and the tongue of head-fixed marmosets as they performed visually guided saccades in exchange for food (**Fig. 1A** [↗](#)). Each successful trial consisted of 3 visually guided saccades, at the end of which we delivered an increment of food (a slurry mixture of apple sauce and monkey chow). Because the reward amount was small (0.015–0.02 mL), the subjects rarely harvested following a single successful trial. Rather, they worked for a few trials, allowing the food to accumulate, then initiated their harvest by licking (**Fig. 1B** [↗](#)). The key variables were how many trials they chose to work before starting harvest, and how vigorously they moved their eyes and tongue during the work and the harvest periods.

Over the course of 2.5 years, we recorded 56 sessions in subject M (29 months) and 56 sessions in subject R (23 months). A typical work period lasted about 10 seconds, during which the subjects attempted ~8 trials, and succeeded in 4–5 trials (**Fig. 1D** [↗](#)) (a successful trial was when all three saccades were within 1.25° of the center of each target). The work period ended when the subject decided to stop tracking the targets and instead initiated harvest, which lasted about 6 seconds, resulting in 16–18 licks. Subject M completed an average of 909.5 ± 61 successful trials per session (mean $\pm$ SEM), producing an average of 241 ± 13.9 work-harvest pairs, and subject R completed an average of 1431 ± 65 successful trials, producing an average of 263 ± 8.9 work-harvest pairs.

We delivered food via either the left or the right tube for 50–300 consecutive trials, and then switched tubes. We tracked the motion of the tongue using DeepLabCut (12 [↗](#)), as shown for a typical session in **Fig. 1B** [↗](#). The licks required precision because the tube was just large enough (4.4 mm diameter) to allow the tongue to penetrate. As a result, about 30% of the of the reward-seeking licks were successful and contacted food ($30 \pm 1.6\%$ for subject M, $28 \pm 2.5\%$ for subject R), as shown in Supporting Information [Video 1](#). Example of licks that failed to contact food are shown in Supporting Information [Video 2](#), [Video 3](#), and [Video 4](#).

Theory and predictions

During the decision-making part of the task, the brain explicitly determined how long to work before initiating harvest. During the work and the harvest periods, the brain implicitly controlled the vigor of movements. We imagined that these two forms of behavior were not independent, but rather coordinated via a control policy that maximized a single utility: the sum of rewards acquired, minus efforts expended, divided by time, termed the capture rate. We chose this formulation because it presents a normative approach that ecologists have used to understand the decisions that animals make regarding how far to travel for food, what mode of travel to use, and how long to stay before moving on to another patch (4 [↗](#), 13 [↗](#), 14 [↗](#)).




Figure 1.

Elements of a foraging task. **A.** During the work period, marmosets made a sequence of saccades to visual targets. A trial consisted of 3 consecutive saccades, at the end of which the subject was rewarded by a small increment of food. We tracked the eyes, the tongue, and the food. **B.** An example of two consecutive work-harvest periods, showing reward-relevant saccades (eye velocity) and tongue endpoint displacement with respect to the mouth. **C.** Data for two sessions, one where the tube was placed close to the mouth (orange trace), and one where it was placed farther away (red trace). Two types of licks are shown: inner-tube licks, and outer-tube licks. Depending on food location, both types of licks can contact the food. Data on the right two panels show endpoint displacement and velocity of the tongue during inner-tube licks. Error bars are SEM. **D.** During the work period, the subjects attempted about 8 trials on average, succeeding in 4-5 trials before starting harvest, and then licked about 18 times to extract the food.

During a work period, our subjects decided to complete a number of saccade trials n_s , a fraction β_s of which were successful, earning food increment α , but expended effort c_s that consumed time T_s for each trial. They then stopped working and initiated harvest, producing a number of licks n_l , a fraction β_l of which succeeded, thus expending effort c_l and consuming time T_l for each lick. These actions produced the following capture rate:

$$J = \frac{\alpha\beta_s n_s (1 - \frac{1}{1 + \beta_l n_l}) - n_l c_l - n_s^2 c_s}{n_l T_l + n_s T_s} \quad (1)$$

In the numerator of Eq. (1), the first term represents the fact that the food cache increased linearly with successful trials and was then consumed gradually with successful licks. The second term represents the effort expenditure of licking, and the third term represents the effort expenditure of working. Notably, the effort expenditure of work, $n_s^2 c_s$, grows faster than linearly as a function of trials. This nonlinearity is essential to reflect the idea that following a long work period, the capture rate must be more negative than following a short work period (i.e., more work trials produce a greater reduction in utility).

A control policy describes how long to work and harvest, and an optimal policy produces periods of working and harvesting, n_s^*, n_l^* , that maximize Eq. (1). A closed-form solution for the optimal policy can be obtained (*Mathematica* notebook simulations.nb) and **Fig. 2A** provides an example. As the work period concludes and the harvest period begins ($n_l = 0$), the capture rate is negative. This reflects the fact that the subject has performed a few trials and stockpiled food, thus expended effort but has not been rewarded yet. The capture rate rises when licking commences. Critically, the peak capture rate is not an increasing function of the work period. Rather, there is an optimal work period (n_s^* , red trace, **Fig. 2A**) associated with a given effort cost of licking c_l . If we now move the tube away from the mouth, that is, increase the effort cost of licking c_l , the peak of the capture rate shifts and the optimal work period changes: the proper response to an increased effort cost of licking is to work longer, stowing more food before commencing harvest.

Notably, the higher cost of licking inevitably reduces the maximum capture rate (**Fig. 2A**). This should impact movement vigor: animals tend to respond to a reduced capture rate by slowing their movements (9), which can be viewed as an effective way to save energy (8). To incorporate vigor into the capture rate, we tried to define the effort cost of a single lick c_l in terms of its energetic cost, a relationship that is currently unknown. Fortunately, other movements provide a clue: the energetic cost of reaching (8, 15), as well as the energetic cost of walking (16, 17), are both concave upward functions of the movement's duration. That is, from an energetic standpoint, there is a reach speed, and a walking speed, that minimizes the cost of each type of movement. We generalized these empirical observations to licking and assumed that the energetic cost of a single lick was a concave upward function of its duration:

$$c_l(T_l) = \frac{d^2}{T_l} + kT_l \quad (2)$$

In Eq. (2), the lick is aimed at a tube located at distance d , and has a duration T_l . The parameter k describes the rate with which the cost grows as a function of duration. For example, the lick duration that minimizes the energetic cost is $d/\sqrt{k}$. Thus, for an energetically optimal lick, duration grows linearly with tube distance (**Fig. 2B**). However, our objective is not to minimize the cost of licking, but to maximize the capture rate. To do so, we insert Eq. (2) into Eq. (1) and find the optimal policy (n_s^*, n_l^*, T_l^*), which now depends on the distance of the food tube to the mouth (*Mathematica* notebook simulations.nb).

The theory predicts that to maximize the capture rate (Eq. 1), the response to an increased effort cost of harvest (i.e., tube distance) should be as follows: n_s^* should increase (**Fig. 2C**), n_l^* should decrease (**Fig. 2C**), and T_l^* should increase (**Fig. 2D**). Notably, the rate of increase in T_l^* as a function of tube distance is faster than linear, while from an energetic point of view (Eq. 2),




Figure 2.

Theoretical results of an optimal control policy. **A.** Capture rate (Eq. 1) is plotted during the harvest period as a function of lick number following various number of work trials. When the effort cost of licking is low (left plot), the optimal work period is (red trace). When the effort cost is higher (right plot), it is best to work longer before initiating harvest. **B.** The metabolic cost of licking (Eq. 2) is minimized when a lick has a specific duration. Tube distance varied from 0.1 to 0.3. Optimal duration that minimizes lick cost grows linearly with tube distance. **C.** Optimal number of work trials and licks as a function food tube distance. As the effort cost of harvest increases, one should respond by working longer, delaying harvest. **D.** Optimal lick duration as a function of food tube distance. The lick duration that maximizes the capture rate is smaller than the one that minimizes the lick metabolic cost (Fig. 2B). That is, it is worthwhile moving vigorously to acquire reward. However, grows faster than linearly as a function of tube distance. Thus, as the tube moves farther, it is best to reduce lick vigor. Hunger, modeled as increased value of reward, should promote work and increase vigor, while effort cost of harvest (tube distance) should promote work but reduce vigor. Parameter values for all simulations: , , , (low food value, less hunger), (high food value, hungry).

increase in distance should produce a linear increase in lick duration. Thus, as the harvest becomes more effortful, the subject should work longer to stockpile food, but move slower to save energy.

To test our theory further, we thought it useful to have a way to alter decisions in one direction (say work longer) but change movement vigor in the opposite direction (move faster). In theory, this is possible: if the subject is hungry (darker lines in [Fig. 2C](#) and [2D](#)), i.e., the reward is more valuable, then they should again work longer before initiating harvest. Paradoxically, they should also move faster.

In summary, if decisions and actions are coordinated via a policy that aims to maximize the capture rate, then in response to an increased cost of harvest, one should work longer, but move with reduced vigor. In response to an increased reward value, as in hunger, one should also work longer, but now move with increased vigor.

Increased effort cost of harvest promoted work but reduced saccade vigor

To vary the effort cost of harvest, we altered the tube distance to the mouth (but kept it constant during each session). Varying tube distance affected the decisions of the subjects: when the tube was placed farther, they chose to work longer before starting harvest ([Fig. 3A](#), left subplot): they attempted more trials during each work period (ANOVA, subject M: $F(2,7908)=41.5$ $p=5.2\times 10^{-25}$, subject R: $F(2,10948)=88.2$ $p=7\times 10^{-50}$), and produced more successful trials per work period (ANOVA, subject M: $F(2,7908)=63$ $p=2.8\times 10^{-24}$, subject R: $F(2,10948)=163$ $p<10^{-50}$). This policy of delayed gratification was present throughout the recording session ([Fig. 3A](#), middle plot). That is, when the harvest required more effort, the subjects worked longer to stockpile more food before initiating their harvest ([Fig. 3A](#), right plot, effect of tube distance on food cached: subject M: $F(2,9566)=176$ $p<10^{-50}$, subject R: $F(2,8907)=204$ $p<10^{-50}$).

During the work period the subjects made saccades to visual targets and accumulated their food. They also made saccades that were not toward visual targets and thus were not eligible for reward. For each animal we computed the relationship between peak saccade velocity and saccade amplitude across all sessions and then calculated the vigor of each saccade: defined as the ratio of the actual peak velocity with respect to the expected peak velocity for that amplitude ([18](#), [19](#)). For example, a saccade that exhibited a vigor of 1.10 had a peak velocity that was 10% greater than the average peak velocity of the saccades of that amplitude for that subject. As expected, the reward-relevant saccades, i.e., saccades made to visual targets (primary, corrective, and center saccades) were more vigorous than other saccades ([Fig. 3B](#), 2-way ANOVA, effect of saccade type, subject M: $F(1,391459)=7248$ $p<10^{-50}$, subject R: $F(1,355839)=13641$ $p<10^{-50}$).

As a work period began, the reward-relevant saccades exhibited high vigor, but then trial-by-trial, this vigor declined, reaching a low vigor value just before the work period ended ([Fig. 3C](#) vigor). Remarkably, on days in which the tube was placed farther, saccade vigor was lower (RMANOVA, effect of tube distance, subject M: $F(2,59033)=224$ $p<10^{-50}$, subject R: $F(2,50103)=75.51$ $p=1.8\times 10^{-33}$). Thus, increasing the effort cost of extracting food during the harvest period reduced saccade vigor during the work period.

By definition, a more vigorous saccade had a greater peak velocity. This might imply that high vigor saccades should suffer from inaccuracy due to signal dependent noise ([20](#)). However, we observed the opposite tendency: as saccade vigor increased, both the magnitude and the variance of the endpoint error decreased ([Fig. 3C](#), 2-way ANOVA, effect of vigor on error magnitude, subject M: $F(8,59046)=480$ $p<10^{-50}$, subject R: $F(8,50184)=252$ $p<10^{-50}$, effect of vigor on error variance, subject M: $F(8,2673)=18200$ $p<10^{-50}$, subject R: $F(8,2673)=4170$ $p<10^{-50}$). That is, reducing the effort costs of harvest not only promoted vigor, it also facilitated accuracy ([21](#)).




Figure 3.

As the effort cost of harvest increased, subjects chose to work more trials, but slowed their movements. **A**. Left: the number of trials attempted and succeeded per work period as a function of tube distance. Middle: successful trials per work period as a function of time during the recording session. Tube distance is with respect to a marker on the nose. Right: food available in the tube at the start of the harvest. **B**. Peak saccade velocity as a function of amplitude for reward-relevant and other saccades. **C**. Vigor of reward relevant saccades as a function of trial number during the work period. Saccade vigor was greater when the tube was closer. Pupil size is quantified during the same work periods. Accuracy is quantified as the magnitude of the saccade's endpoint error vector (with respect to the target) and the variance of that error vector (determinant of the variance-covariance matrix), plotted as a function of the vigor of the saccade (bin size=0.05 vigor units). Error bars are SEM.

Cognitive signals such as effort and reward are associated with changes in pupil size (10 [↗](#)), as well as transient activation of brainstem neuromodulatory circuits in locus coeruleus (11 [↗](#)). We wondered if the changes in tube position altered the output of these neuromodulatory circuits, as inferred via pupil size. For each reward-relevant saccade, we measured the pupil size during a ± 250 ms window centered on saccade onset, and then normalized this measure based on the distribution of pupil sizes that we had measured during the entire recording for that session in that subject, resulting in a z-score.

At the onset of each work period the pupils were dilated, but as the subjects performed more trials, the pupils constricted, exhibiting a trial-by-trial reduction that paralleled the changes in saccade vigor (**Fig. 3C** [↗](#)). Notably, the effort cost of harvest affected pupil size: during the work period the pupils were more dilated if the tube was placed closer to the mouth (**Fig. 3C** [↗](#), RMANOVA, effect of tube distance, subject M: $F(2,60502)=20$ $p=2\times 10^{-9}$, subject R: $F(2,50431)=23.8$ $p=4.9\times 10^{-11}$). That is, when the effort cost of harvest was lower, the pupils dilated, and the saccades were invigorated.

In summary, when we increased the effort cost of harvest, both the movements and the decisions changed: the pupils constricted and the movements slowed, but they chose to work more trials before initiating harvest.

Increased effort cost of harvest reduced lick vigor

The work period ended when the subject chose to stop tracking the target and initiated harvest via a licking bout. As in saccades, we defined lick vigor via the ratio of the actual peak velocity of the lick with respect to the expected velocity for that lick amplitude. As amplitude increased, lick peak velocity increased during both protraction and retraction (**Fig. 4A** [↗](#)). Some of the licks were reward seeking and directed toward the tube, while others were grooming licks, cleaning the tongue and the area around the mouth (Supporting Information [Video 5](#)). Reward-seeking licks were more vigorous than grooming licks (2-way ANOVA, effect of lick type, protraction, subject M: $F(1,272233)=66$ $p=4.5\times 10^{-16}$, subject R: $F(1,229052)=698$ $p<10^{-50}$), and retraction was more vigorous than protraction (reward seeking licks, retraction vs. protraction, subject M: $t(241145)=532$ $p<10^{-50}$, subject R: $t(213674)=665$ $p<10^{-50}$).

As the harvest began, the first lick was very low vigor, but lick after lick, the movements gathered velocity, reaching peak vigor by the 3rd or the 4th lick (**Fig. 4B** [↗](#)). As the harvest continued, lick vigor gradually declined. Like saccades, licks had a lower vigor in sessions in which the tube was placed farther from the mouth (RMANOVA, effect of tube distance, subject M: $F(2,59033)=222.5$ $p<10^{-50}$, subject R: $F(2,133502)=224$ $p<10^{-50}$), and this pattern was present during the entire recording session (**Fig. 4C** [↗](#), left subplot). Thus, an increased effort cost of harvest promoted sloth: reduced vigor of saccades during the work period, and reduced vigor of licks during the harvest period.

For each reward seeking lick, we measured pupil size during a ± 250 ms window centered on the moment of peak tongue displacement. During licking, the pupil size changed with a pattern that closely paralleled lick vigor: as the harvest began, pupil size was small, but it rapidly increased during the early licks, then gradually declined as the harvest continued (**Fig. 4B** [↗](#), right subplot). Importantly, the pupils were more dilated in sessions in which the tube was closer to the mouth (**Fig. 4C** [↗](#), right subplot, effect of tube distance, subject M: $F(2,166742)=583$ $p<10^{-50}$, subject R: $F(2,130493)=118$ $p<10^{-50}$). As a result, when the effort cost of reward increased, the pupils constricted, and the vigor of both saccades and licks decreased.

While the theory predicted that moving the tube farther would result in a longer work period and reduced movement vigor, it also predicted that the subjects would reduce their harvest duration (reduced licks, **Fig. 2C** [↗](#)). That is, it predicted that the subjects would work longer, stowing more




Figure 4.

As the effort cost of harvest increased, lick vigor declined, and the pupils constricted. **A.** Peak speed of reward-seeking and grooming licks during protraction and retraction as a function of lick amplitude. **B.** Vigor of reward seeking licks (protraction) and pupil size as a function of lick number during harvest at various tube distances. **C.** Lick vigor and pupil size as a function of time during the entire recording session. Line colors depict tube distance as in part B. **D.** Average lick vigor and pupil size during a harvest as a function of number of trials successfully completed in the previous work period. Lick vigor and pupil size were greater when more food had been stored. **E.** Following a successful lick (contact with food), the next lick was more vigorous, and pupils dilated. Following a failed lick, the next lick was slowed, and pupils were less dilated. **F.** We observed no consistent effect of lick vigor on lick accuracy across subjects or across tube distances. Error bars are SEM.

food, but leave more of it behind. This last prediction did not agree with our data (see Discussion). For Subject R, the number of licks were approximately the same across the various tube distances, and for Subject M the number of licks increased with tube distance (**Supplementary Fig. 1** [↗](#)).

In summary, within a harvest period, lick vigor rapidly increased and then gradually declined. Simultaneous with the changes in vigor, the pupils rapidly dilated and then gradually constricted. In sessions where the tube was placed farther from the mouth, the licks had lower vigor, and the pupils were more constricted.

Expectation of greater reward increased lick vigor

As the subject worked, they accumulated food, thus increasing the magnitude of the available reward. To check whether reward magnitude affected movement vigor, for each tube distance we computed the average lick vigor during the harvest as a function of the number of trials completed in the preceding work period. We found that when the work period had included many completed trials, then the movements in the ensuing harvest period were more vigorous (**Fig. 4D** [↗](#), 2-way ANOVA, effect of trials, subject M: $F(4, 164242)=353$ $p<10^{-50}$, subject R: $F(4,123411)=152$ $p<10^{-50}$). Thus, the licks were invigorated by the amount of food that awaited harvest.

Because the tube was small, many of the licks missed their goal and failed to contact the food. The success or failure of a lick affected both the vigor of the subsequent lick, and the change in the size of the pupil. Following a successful lick there was a large increase in lick vigor (**Fig. 4E** [↗](#), subject M: $t(85182)=40$ $p<10^{-50}$, subject R: $t(81378)=104$ $p<10^{-50}$), and a large increase in pupil size (subject M: $F(84969)=57$ $p<10^{-50}$, subject R: $t(80318)=94$ $p<10^{-50}$). In contrast, following a failed lick the subjects either reduced or did not increase their lick vigor (**Fig. 4E** [↗](#), subject M: $t(114159)=0.88$ $p=0.37$, subject R: $t(97164)=-44$ $p<10^{-50}$). This failure also produced a smaller increase in pupil size (comparison to successful lick, two sample t-test, subject M: $t(198722)=14.8$ $p=4.3\times 10^{-50}$, subject R: $t(176044)=53$ $p<10^{-50}$). Thus, a single successful lick led to acquisition of reward, which then was followed by a relatively large increase in pupil size, and an invigorated subsequent lick.

For saccades we had found that increased vigor was associated with greater accuracy. To quantify the relationship between lick vigor and accuracy, for each tube distance we labeled each reward seeking lick as being high or low vigor. For subject M, high vigor licks tended to be more successful, but this was not the case for subject R (**Fig. 4F** [↗](#)). Moreover, tube distance did not produce a consistent effect on lick success.

In summary, the subject licked more vigorously following a long work period in which they had accumulated more reward. Moreover, when a lick was successful in acquiring reward, they increased the vigor of the subsequent lick.

Hunger promoted work and increased vigor

Our theory predicted that it should be possible to change decisions in one direction (say work longer), while altering movement vigor in the opposite direction (move faster). An increase in the subjective value of reward, as might occur when the subject is hungry, should have two effects: increase the number of trials that the subject chooses to perform before commencing harvest, and increase movement vigor.

We did not explicitly manipulate the weight of the subjects. Indeed, to maintain their health, we strived to keep their weights constant during the roughly 2.5-year period of these experiments. However, there was natural variability, which allowed us to test the predictions of the theory.

We found that when their weight was lower than average, the subjects chose to work a greater number of trials before commencing harvest (**Fig. 5A**, two sample t-test, subject M: $t(11052)=7.9$ $p=3.4\times 10^{-25}$, subject R: $t(12549)=10.1$ $p=9.3\times 10^{-24}$). This result was similar to the effect that we had seen when the effort cost of harvest was increased. However, the theory had predicted that the effect on vigor should be in the opposite direction: if hunger increased reward valuation, then one should speed the movements and hasten food acquisition. Notably, weight did not have a consistent effect on saccade vigor across the two subjects (**Fig. 5A**), yet during the harvest, both subjects licked with greater vigor when their weight was lower (**Fig. 5A**, subject M: $t(219752)=88$ $p<10^{-50}$, subject R: $t(205163)=22$ $p<10^{-50}$).

Thus, while both the effort cost of reward and hunger promoted greater work, effort promoted sloth while hunger promoted lick vigor.

Pupil size variations strongly correlated with changes in decisions and movements

Finally, we considered the data across both the work and the harvest periods and asked how well movement vigor tracked pupil size. The results demonstrated that in both the work and the harvest periods, for both saccades and licks, an increase in pupil size was associated with an increase in vigor (**Fig. 5B**, reward-relevant saccades, subject M: $r=0.989$ $p=7.7\times 10^{-9}$, subject R: $r=0.97$ $p=7.1\times 10^{-7}$, reward-seeking protraction licks, subject M: $r=0.969$ $p=9.8\times 10^{-7}$, subject R: $r=0.989$ $p=6.3\times 10^{-9}$). Moreover, when the pupil was dilated, the work periods tended to be shorter (**Fig. 5C**, subject M: $r=-0.90$ $p=0.00014$, subject R: $r=-0.97$ $p=6.3\times 10^{-7}$), while harvest durations tended to be longer (**Fig. 5C**, subject M: $r=0.894$ $p=0.00021$, subject R: $r=0.935$ $p=2.4\times 10^{-5}$). Thus, pupil dilation was associated with choosing to work less, while moving faster.

Discussion

What we choose to do is the purview of the decision-making circuits of our brain, while the implicit vigor with which we perform that action is the concern of the motor-control circuits. From a theoretical perspective (9), our brain should coordinate these two forms of behavior because both the act that we select, and its vigor, dictate expenditure of time and energy, contributing to a capture rate that affects longevity and fecundity (22). Does the brain coordinate decisions and movements to maximize a capture rate? If so, how might the brain accomplish this coordination?

Here, we designed a foraging task in which marmosets worked by making saccades, accumulating food for each successful trial, then stopped working and harvested their cache by licking. On every trial they decided whether to work, or to harvest. Their decision was carried out by the motor system, producing either a visually guided saccade, or a lick, each exhibiting a particular vigor. The theory predicted that to maximize the capture rate, the appropriate response to an increased effort cost of harvest was to do two things: work longer to cache more food but reduce vigor to conserve energy.

We varied the effort costs by moving the food tube with respect to the mouth. This changed the effort cost of harvest but not the effort cost of work. The subjects responded by altering how they worked as well as how they harvested. When the harvest was more effortful, they performed more saccade trials to stockpile food. They also slowed their movements, reducing saccade velocity during the work period, and reducing lick velocity during the harvest period. Notably, the most vigorous saccades were also the most accurate: as saccade vigor increased, so did endpoint accuracy.




Figure 5.

Relatively low body weight, potentially reflecting a greater valuation of reward, coincided with longer work periods and greater vigor. Pupil size correlated with both vigor and decisions. **A.** Trials successfully completed during a work period as a function of normalized body weight at the start of the session. **B.** Left: saccade vigor as a function of trial number for low and high body weights. Right: pupil size during the same saccades. **C.** Left: lick vigor as a function of lick number during the harvest period. Right: pupil size during the same licks. **D.** Saccade vigor during the work period, and lick vigor during the harvest period, as a function of pupil size. **E.** Work duration and harvest duration as a function of pupil size. Error bars are SEM.

The theory made a second prediction: as the value of reward increased, the subjects should again choose to work a longer period before initiating harvest, but unlike the effort costs, now respond by moving more vigorously. We did not directly manipulate the subjective value of reward, but rather relied on the natural fluctuations in body weight and assumed that when their weight was low, the subjects were hungrier for reward. Indeed, when their weight was low, the subjects again chose to work longer, but now elevated their vigor during the harvest period.

Finally, we quantified the effect of reward magnitude on vigor. Within a session, lick vigor increased robustly as a function of the number of trials completed in the preceding work period. Thus, the licks were invigorated by the amount of food that awaited harvest.

Notably, some of the predictions of the theory did not agree with the experimental data. An increased effort cost did not accompany a reduction in the duration of harvest, and hunger did not increase saccade vigor robustly. Indeed, earlier experiments have shown that if the effort cost of harvest increases, animals who expend the effort will then linger longer to harvest more of the reward that they have earned (2 [↗](#)). This mismatch between observed behavior and theory highlights some of the limitations of our formulation. For example, our capture rate reflected a single work-harvest period, rather than a long sequence. Moreover, the capture rate did not consider the fact that the food tube had finite capacity, beyond which the food would fall and be wasted. This constraint would discourage a policy of working more but harvesting less. Finally, if we assume that a reduced body weight is a proxy for increased subjective value of reward, it is notable that we observed a robust effect on vigor of licks, but not saccades. A more realistic capture rate formulation awaits simulations, possibly one that describes capture rate not as the ratio of two sums (sum of gains and losses with respect to sum of time), but rather the expected value of the ratio of each gain and loss with respect to time (23 [↗](#), 24 [↗](#)).

A shortcoming of our model is that we did not include a link between lick vigor and its probability of success. As a result, when we moved the food tube away, the model did not consider the possibility that maintaining lick accuracy may involve reduced vigor. The reason for this is that we searched for but could not find a consistent relationship, across subjects or effort conditions, between protraction speed of the tongue and its success probability. Thus, we cannot exclude this alternate hypothesis. However, the most interesting aspect of our results was that when we increased tube distance, making harvest more effortful, there was not only a reduction in lick vigor, but also a reduction in saccade vigor. That is, the decisions and actions during the work period responded to the increased effort cost of reward during the harvest period.

What might be a neural basis for this coordination of decisions and movements? A clue was the fact that the pupils were more constricted in sessions in which the effort cost of harvest was greater. This global change in pupil size accompanied delayed harvest and reduced vigor across sessions, but surprisingly, even within a session, transient changes in pupil size accompanied changes in vigor. During the work period the trial-to-trial reduction in saccade vigor accompanied trial-to-trial constriction of the pupil, and within a harvest period, the rapid rise and then the gradual fall in lick vigor paralleled rapid dilatation followed by gradual constriction of the pupil.

Pupil dilation is a proxy for activity in the brainstem neuromodulatory system (25 [↗](#)) and is a measure of arousal (26 [↗](#)). Control of pupil size is dependent on spiking of norepinephrine neurons in locus coeruleus (LC-NE): an increase in the activity of these neurons produces pupil dilation (27 [↗](#), 28 [↗](#)). Some of these neurons show a transient change in their activity when acquisition of reward requires expenditure of either physical (11 [↗](#)) or mental effort (29 [↗](#)), even when there is no concomitant movement to be made. It is possible that in the present task, as the effort cost of harvest increased, LC-NE neurons decreased their activity, producing pupil constriction. If so, the reduced NE release may have had two simultaneous effects: encourage work and promote delayed gratification in brain regions that control decisions, discourage energy expenditure and promote sloth in brain regions that control movements. Thus, the idea that

emerges is that the response of NE to economic variables, as inferred via changes in pupil size, might act as a bridge to coordinate the computations in the decision-making circuits with the computations in the motor-control circuits, aiming to implement a consistent control policy that improves the capture rate.

In addition to NE, the basal ganglia, and in particular the neurotransmitter dopamine, are likely the key contributors to the coordination of decisions with actions (30, 31). When the effort price of a preferred food increases, animals choose to work longer, pressing a lever a greater number of times (32, 33). This desire to expend effort to acquire a valuable reward is reduced if dopamine is blocked in the ventral striatum (34–36). Hunger activates circuits in the hypothalamic nuclei, disinhibiting dopamine release in response to food cues (37). Dopamine concentrations in the striatum drop when the effort price of a food reward increases (38), and dopamine release before onset of a movement tends to invigorate that movement (39). Thus, the presence of dopamine may not only alter decisions by encouraging expenditure of effort, but also modify movements by promoting vigor.

Experiments of Hayden et al. (40) and Barack et al. (41) suggest that the decision of when to stop work and commence harvest may rely on computations that are carried out in the cingulate cortex. They found that as monkeys deliberated between the choice of staying and acquiring diminishing rewards, or leaving and incurring a travel cost, these neurons encoded a decision-variable that reflected the value of leaving the patch. The prediction that emerges from our work is that the rate of rise of these decision variables may be modulated by the presence of NE.

From a motor control perspective, a surprising aspect of our results was that an increase in saccade vigor accompanied an improvement in endpoint accuracy. In our earlier work we found that during reaching, reward increased vigor without reducing accuracy (42). Thus, the brain has the means to increase movement vigor and improve its accuracy. How is this achieved?

We found that the high vigor saccades were produced when the pupils were dilated, implying an increased release of NE. In songbirds, increased NE release acts on the basal ganglia to suppress activity of spiny neurons, and this reduced activity in the basal ganglia accompanies reduced variance in the songs that the animal sings (43). Thus, NE may play a critical role in control of movement variability. For saccades, control of endpoint accuracy depends on the coordinated activity of Purkinje cells in the oculomotor region of the cerebellar vermis (44, 45). LC projects to the cerebellum, and stimulation of LC neurons increases the sensitivity of Purkinje cells to their inputs (46).

Is movement vigor increased following increased NE inputs from LC to the basal ganglia, and accuracy improved following increased NE inputs from LC to the cerebellum? Does decision-making shift toward greater work and delayed gratification following reduced NE inputs from LC to the frontal lobe? These are some of the questions that await future experiments.

Methods

Behavioral and neurophysiological data were collected from two marmosets (*Callithrix Jacchus*, male and female, 350–390 g, subjects R and M, 6 years old). The neurophysiological data focused on the cerebellum and are described elsewhere (44, 47, 48). Here, our focus is on the behavioral data.

The marmosets were born and raised in a colony that Prof. Xiaoqin Wang has maintained at the Johns Hopkins School of Medicine since 1996. The procedures on the marmosets were evaluated and approved by the Johns Hopkins University Animal Care and Use Committee in compliance with the guidelines of the United States National Institutes of Health.

Data acquisition

Following recovery from head-post implantation surgery, the animals were trained to make saccades to visual targets and rewarded with a mixture of apple sauce and lab diet (47). They were placed in a monkey chair and head-fixed while we presented visual targets on an LCD screen (Curved MSI 32" 144 Hz - model AG32CQ) and tracked both eyes at 1000 Hz using an EyeLink-1000 system (SR Research, USA). Timing of target presentation on the video screen was measured using a photo diode. Tongue movements were tracked with a 522 frame per second Sony IMX287 FLIR camera, with frames captured at 100 Hz.

Each trial began with a saccade to the center target followed by fixation for 200 ms, after which a primary target (0.5×0.5 deg square) appeared at one of 8 randomly selected directions at a distance of 5-6.5 deg. Onset of the primary target coincided with presentation of a tone. As the animal made a saccade to the primary target, that target was erased, and a secondary target was presented at a distance of 2-2.5 deg, also at one of 8 randomly selected directions. The subject was rewarded if following the primary saccade it made a corrective saccade to the secondary target, landed within 1.5 deg radius of the target center, and maintained fixation for at least 200 ms. Onset of reward coincided with presentation of another distinct tone. Following an additional 150-250 ms period (uniform random distribution), the secondary target was erased, and the center target was displayed, indicating the onset of the next trial. Thus, a successful trial comprised of a sequence of 3 saccades: center, primary, and corrective, after which the subject received a small increment of food (0.015 mL).

The food was provided in two small tubes (4.4 mm diameter), one to the left and the other to the right of the animal (Fig. 1A). A successful trial produced a food increment in one of the tubes and would continue to do so for 50-300 consecutive trials, then switch to the other tube. Because the food increment was small, the subjects naturally chose to work for a few consecutive trials, tracking the visual targets and allowing the food to accumulate, then stopped tracking and harvested the food via a licking bout. The licking bout typically included a sequence of 15-40 licks. The subjects did not work while harvesting. As a result, the behavior consisted of a work period (targeted saccades), followed by a harvest period (targeted licking), repeated hundreds of times per session.

The critical variables were the number of trials that the subject chose to perform before initiating harvest, the vigor of their saccades during the work period, and the vigor of their licks during the harvest period.

Data analysis

All saccades, regardless of whether they were instructed by presentation of a visual target or not, were identified using a velocity threshold. Saccades to primary, secondary, and central targets were labeled as reward-relevant saccades, while all remaining saccades were labeled as task irrelevant.

We analyzed tongue movements using DeepLabCut (12). Our network was trained on 89 video recordings of the subjects with 15-25 frames extracted and labeled from each recording. The network was built on the ResNet-152 pre-trained model, and then trained over 1.03×10^5 iterations with a batch size of 8, using a GeForce GTX 1080Ti graphics processing unit (49). A Kalman filter was further applied to improve quality and smoothness of the tracking, and the output was analyzed in MATLAB to quantify varying lick events and kinematics.

We tracked the tongue tip and the edge of the food in the tube, along with control locations (nose position and tube edges). We tracked all licks, regardless of whether they were aimed toward the tube (reward seeking), or not (grooming). Reward seeking licks were further differentiated based

on whether they aimed to enter the tube (inner-tube licks), hit the outer edge of the tube (outer-edge licks), or fell below the tube (under tube). If any of these licks successfully contacted the tube, we labeled that lick as a success (otherwise, a failed lick).

Pupil area was measured during a ± 250 ms period centered at the onset of each reward-relevant saccade, and the onset of each lick. We then normalized the pupil measurements by representing it as a z-score with respect to the mean value for that session.

Saccade and tongue vigor

We relied on previous work to define vigor of a movement (5, 9, 18, 19). Briefly, if the amplitude of a movement is x and the peak speed of that movement is v , then for each subject the relationship between the two variables can be described as:

$$v = \alpha \left(1 - \frac{1}{1 + \beta x} \right) \quad (3)$$

In the above expression, $\alpha, \beta \geq 0$ and are subject-specific parameters. For a movement with amplitude x , its vigor was defined as the ratio of the actual peak speed with respect to the expected value of its peak speed, i.e., $\frac{v}{v_{\text{exp}}(x)}$. Expected value was computed by fitting Eq. (3) to all the data acquired across all sessions. When vigor is greater than 1, the movement had a peak velocity that was higher than the mean value associated with that amplitude.

Model formulation

We chose a formulation of utility (Eq. 1) based on a normative approach that ecologists have used to understand the decisions that animals make regarding how far to travel for food, what mode of travel to use, and how long to stay before moving on to another reward opportunity (4, 13, 14). In a typical formulation of the theory, the numerator represents the reward gained (in units of energy), minus the effort expended (also in units of energy), while the denominator represents the amount of time spent during that behavior. We represented this idea in Eq. (1) with saccades that produced reward accumulation, and licks that produced reward consumption. Thus, the utility that we aim to maximize is the rate of energy gained.

The specific functions that we used to represent the energy gained through reward acquisition, and the energy expended through effort expenditure, came either from experiment design, or from the measurements we have made in other experiments. We modeled reward accumulation as a linear rise in energy stored because successful saccades produced a linear increase in the food cache. We modeled harvesting of the food as a hyperbolic function of the number of licks to represent the fact that as the licking bout began, each successful lick depleted the food, and thus the first few licks produced a greater amount of food consumption than the last few licks. We modeled the effort cost of licking as a linear function of the number of licks.

A critical assumption that we made is that energy expended performing the saccade trials (which grew faster than linearly as a function of the number of trials attempted), grew faster than the time spent attempting those same trials (which grew linearly with the number of trials). This assumption is based on the heuristic that the average rate of energy lost following a large number of attempted trials is greater than the average rate of energy lost following a small number of attempted trials.

The model's simplicity provided closed-form solutions across all parameter values, allowing us to make predictions without having to fit the model to the measured data. For example, for all parameter values that produce a real solution (as opposed to imaginary), the optimal number of saccade trials increases with the square root of the cost of licking. Thus, the basic prediction of the

model is that to maximize the capture rate, regardless of parameter values, an increase in the effort required for harvest should be met with a greater willingness to work. The closed-form solutions are presented in the supplementary document (simulations.nb).

Other models of utility

In composing our utility (Eq. 1), we chose to combine reward and effort additively. This is in contrast to other approaches in which effort discounts reward multiplicatively (50–52). Our reasoning is that multiplicative interactions have the limitation that they are incompatible with the observation that reward invigorates movements.

To compare additive and multiplicative approaches, let us consider an arbitrary function u that specifies how effort varies with movement duration T . Typically, this is a U-shaped function that describes energy expenditure as a function of movement duration, as in (8). In the case of multiplicative interaction between reward and effort, we can consider the following representation of utility:

$$J = \frac{\alpha}{1+T} U^{-1}(T) \quad (4)$$

In the above formulation, reward α is discounted hyperbolically with time, and an increase in reward increases the utility of the action. The optimum movement vigor has the duration T^* that maximizes this utility. Notably, because increasing reward merely scales this utility, it has no effect on vigor. Thus, a utility in which reward is multiplied by a function of effort generally fails to predict dependence of movement vigor on reward.

Simulations

The optimal policy specifies the decisions and movements that for the effort cost defined in Eq. (2), maximizes the capture rate defined in Eq. (1). This policy selects the number of saccade trials n_s to perform during the work period, the number of licks n_l to perform during the harvest period, and the vigor of each lick, represented by the average duration of a lick T_l . To compute the optimal policy, we found the derivative of the capture rate with respect to each policy variable n_s , n_l , and T_l , then set each derivative equal to zero, producing three simultaneous nonlinear equations. In all three cases, we were able to solve for the relevant control variable analytically (see Mathematica notebook *simulations.nb* for the derivations). We found that if the solution was a real number, then regardless of parameter values, an increase in d (distance of the tube to the mouth), the optimal policy produced an increase in n_s^* , decrease in n_l^* , and increase in T_l^* . Thus, the results illustrated in Fig. 2 are robust to changes in parameter values.

To generate the plots in Fig. 2A, we used the following parameter values: $\alpha = 20$, $\beta_s = 0.5$, $\beta_L = 0.3$, $c_s = 0.5$, $T_s = 1$, $T_L = 0.2$, $c_L = 0.5$ (low effort), $c_L = 2.5$ (high effort). For the plots in Fig. 2C and 2D, we used the same parameter values, but c_L was defined via Eq. (2). Thus, tube distance d varied, and T_L was unknown and was solved for. In Eq. (2), $k_L = 1$. In the simulations, to describe state of hunger, we set $\alpha = 20$ for a sated state and $\alpha = 25$ for a hungry state.

Statistical analysis

Hypothesis testing was performed using functions provided by the *MATLAB Statistics and Machine Learning Toolbox*, version R2021b. For t-tests, across the one-sample, paired-sample, and two-sample conditions, p values were computed using the *ttest* and *ttest2* functions with data that was combined across sessions, separated by condition. For ANOVA tests, in the one-way condition, p values were computed using a nonparametric Kruskal-Wallis test, using the *kruskalwallis* function. In the 2-way condition, the *anovan* function was used to compute p values, accounting for an unbalanced design resulting from a varied number of samples across conditions. In both cases, like in the t-tests, data was combined across sessions, separated by condition. In the

repeated measures condition, each session was treated as a subject with multiple repeated measures representing a given variable (i.e., lick vigor per lick in a harvest period). To fit a repeated measures model, the *fitrm* function was used, then analyzed using the *ranova* function. In all cases of repeated measures ANOVA, compound symmetry assumptions were tested using the Mauchly sphericity test with the *maulchy* function. In cases where the assumption was violated (Mauchly test $p < 0.05$), epsilon adjustments were used, with the *epsilon* function, to compute corrected p values (for $\epsilon > .75$, use Huynh-Feldt p value and for $\epsilon < .75$, use Greenhouse-Geisser p values). For correlation analyses, Pearson's correlation coefficient, r , and corresponding p values were computed using the *corrcoef* function.

Acknowledgements

The work was supported by grants from the NIH (R01-EB028156, R01-NS078311, R37-NS128416), and the Office of Naval Research (N00014-15-1-2312).

Supporting Information


Video 1. Example of a successful inner-tube lick.

Video 2. Example of an under-tube lick that failed to contact food. Note the corrective sub-movements, as has been observed in mice (53 [↗](#)).

Video 3. Example of an outer-tube lick that failed to contact food.

Video 4. Example of a lick that hit the outer edge of the tube and failed to contact food.

Video 5. Example of a grooming lick.



Supplementary Fig. 1.

Number of licks per harvest as a function of tube distance.

References

1. Charnov E. L. (1976) **Optimal foraging, the marginal value theorem** *Theor.Popul.Biol* **9**:129–136
2. Cowie R. J. (1977) **Optimal foraging in great tits (*Parus major*)** *Nature* **268**:137–139
3. Shadmehr R., Ahmed A. A. (2020) **Vigor: neuroeconomics of movement control**
4. Richardson H., Verbeek N. A. M. (1986) **Diet selection and optimization by northwestern crows feeding on Japanese littleneck clams** *Ecology* **67**:1219–1226
5. Yoon T., Jaleel A., Ahmed A. A., Shadmehr R. (2020) **Saccade vigor and the subjective economic value of visual stimuli** *J. Neurophysiol* **123**:2161–2172
6. Korbisch C. C., Apuan D. R., Shadmehr R., Ahmed A. A. (2022) **Saccade vigor reflects the rise of decision variables during deliberation** *Curr. Biol* <https://doi.org/10.1016/j.cub.2022.10.053>
7. Shadmehr R., Orban de Xivry J. J., Xu-Wilson M., Shih T. Y. (2010) **Temporal discounting of reward and the cost of time in motor control** *J.Neurosci* **30**:10507–10516
8. Shadmehr R., Huang H. J., Ahmed A. A. (2016) **A representation of effort in decision-making and motor control** *Curr.Biol* **26**:1929–1934
9. Yoon T., Geary R. B., Ahmed A. A., Shadmehr R. (2018) **Control of movement vigor and decision making during foraging** *Proc.Natl.Acad.Sci.U.S.A* **115**:E10476–E10485
10. Joshi S., Gold J. I. (2020) **Pupil Size as a Window on Neural Substrates of Cognition** *Trends Cogn. Sci* **24**:466–480
11. Bornert P., Bouret S. (2021) **Locus coeruleus neurons encode the subjective difficulty of triggering and executing actions** *PLOS Biol* **19**
12. Mathis A., et al. (2018) **DeepLabCut: markerless pose estimation of user-defined body parts with deep learning** *Nat. Neurosci* **21**:1281–1289
13. Stephens D. W., Krebs J. R. (1986) **Foraging Theory**
14. Bautista L. M., Tinbergen J., Kacelnik A. (2001) **To walk or to fly? How birds choose among foraging modes** *Proc.Natl.Acad.Sci.U.S.A* **98**:1089–1094
15. Huang H. J., Ahmed A. A. (2014) **Older adults learn less, but still reduce metabolic cost, during motor adaptation** *J.Neurophysiol* **111**:135–144
16. Ralston H. J. (1958) **Energy-speed relation and optimal speed during level walking** *Int.Z.Angew.Physiol* **17**:277–283
17. Bastien G. J., Willems P. A., Schepens B., Heglund N. C. (2005) **Effect of load and speed on the energetic cost of human walking** *Eur.J.Appl.Physiol* **94**:76–83

18. Reppert T. R., Lempert K. M., Glimcher P. W., Shadmehr R. (2015) **Modulation of Saccade Vigor during Value-Based Decision Making** *J.Neurosci* **35**:15369–15378
19. Reppert T. R., et al. (2018) **Movement vigor as a trait-like attribute of individuality** *J.Neurophysiol* **120**:741–757
20. Harris C. M., Wolpert D. M. (1998) **Signal-dependent noise determines motor planning** *Nature* **394**:780–784
21. Wang C., Xiao Y., Burdet E., Gordon J., Schweighofer N. (2016) **The duration of reaching movement is longer than predicted by minimum variance** *J.Neurophysiol* **116**:2342–2345
22. Lemon W. C. (1991) **Fitness consequences of foraging behaviour in the zebra finch** *Nature* **352**:153–155
23. Bateson M., Kacelnik A. (1995) **Preferences for fixed and variable food sources: variability in amount and delay** *J.Exp.Analysis.Behav* **63**:313–329
24. Bateson M., Kacelnik A. (1996) **Rate currencies and the foraging starling: the fallacy of the averages revisited** *Behav. Ecol* **7**:341–352
25. Vazey E. M., Moorman D. E., Aston-Jones G. (2018) **Phasic locus coeruleus activity regulates cortical encoding of salience information** *Proc. Natl. Acad. Sci.* **115**:E9439–E9448
26. Mathot S. (2018) **Pupillometry: Psychology, Physiology, and Function** *J. Cogn.* **1**:1–23
27. Joshi S., Li Y., Kalwani R. M., Gold J. I. (2016) **Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex** *Neuron* **89**:221–234
28. Breton-Provencher V., Sur M. (2019) **Active control of arousal by a locus coeruleus GABAergic circuit** *Nat. Neurosci* **22**:218–228
29. Contadini-Wright C., Magami K., Mehta N., Chait M. (2023) **Pupil Dilation and Microsaccades Provide Complementary Insights into the Dynamics of Arousal and Instantaneous Attention during Effortful Listening** *J. Neurosci* **43**:4856–4866
30. Thura D., Cisek P. (2017) **The Basal Ganglia Do Not Select Reach Targets but Control the Urgency of Commitment** *Neuron* **95**:1160–1170
31. Herz D. M., et al. (2022) **Dynamic control of decision and movement speed in the human basal ganglia** *Nat. Commun* **13**
32. Salamone J. D., et al. (1991) **Haloperidol and nucleus accumbens dopamine depletion suppress lever pressing for food but increase free food consumption in a novel food choice procedure** *Psychopharmacol. Berl* **104**:515–521
33. Aberman J. E., Salamone J. D. (1999) **Nucleus accumbens dopamine depletions make rats more sensitive to high ratio requirements but do not impair primary food reinforcement** *Neuroscience* **92**:545–552
34. Koch M., Schmid A., Schnitzler H. U. (2000) **Role of nucleus accumbens dopamine D1 and D2 receptors in instrumental and Pavlovian paradigms of conditioned reward** *Psychopharmacol. Berl* **152**:67–73

35. Farrar A. M., et al. (2010) **Nucleus accumbens and effort-related functions: behavioral and neural markers of the interactions between adenosine A2A and dopamine D2 receptors** *Neuroscience* **166**:1056–1067
36. Yohn S. E., et al. (2015) **The role of dopamine D1 receptor transmission in effort-related choice behavior: Effects of D1 agonists** *Pharmacol.Biochem.Behav* **135**:217–226
37. Cassidy R. M., Tong Q. (2017) **Hunger and Satiety Gauge Reward Sensitivity** *Front. Endocrinol* **8**
38. Schelp S. A., et al. (2017) **A transient dopamine signal encodes subjective value and causally influences demand in an economic context** *Proc.Natl.Acad.Sci.U.S.A* **114**:E11303–E11312
39. da Silva J. A., Tecuapetla F., Paixao V., Costa R. M. (2018) **Dopamine neuron activity before action initiation gates and invigorates future movements** *Nature* **554**:244–248
40. Hayden B. Y., Pearson J. M., Platt M. L. (2011) **Neuronal basis of sequential foraging decisions in a patchy environment** *Nat.Neurosci* **14**:933–939
41. Barack D. L., Chang S. W. C., Platt M. L. (2017) **Posterior Cingulate Neurons Dynamically Signal Decisions to Disengage during Foraging** *Neuron* **96**:339–347
42. Summerside E. M., Shadmehr R., Ahmed A. A. (2018) **Vigor of reaching movements: reward discounts the cost of effort** *J.Neurophys* **119**:2347–2357
43. Singh Alvarado J., et al. (2021) **Neural dynamics underlying birdsong practice and performance** *Nature* **599**:635–639
44. Sedaghat-Nejad E., Pi J. S., Hage P., Fakharian M. A., Shadmehr R. (2022) **Synchronous spiking of cerebellar Purkinje cells during control of movements** *Proc. Natl. Acad. Sci.* **119**
45. Barash S., et al. (1999) **Saccadic dysmetria and adaptation after lesions of the cerebellar cortex** *J.Neurosci* **19**:10931–10939
46. Moises H. C., Waterhouse B. D., Woodward D. J. (1981) **Locus coeruleus stimulation potentiates Purkinje cell responses to afferent input: The climbing fiber system** *Brain Res* **222**:43–64
47. Sedaghat-Nejad E., et al. (2019) **Behavioral training of marmosets and electrophysiological recording from the cerebellum** *J.Neurophysiol* **122**:1502–1517
48. Muller S. Z., et al. (2023) **Complex spikes perturb movements, revealing the sensorimotor map of Purkinje cells** *bioRxiv* :2023–4
49. He K., Zhang X., Ren S., Sun J. (2016) **K. He, X. Zhang, S. Ren, J. Sun, Deep Residual Learning for Image Recognition in (2016), pp. 770–778. Deep Residual Learning for Image Recognition :770–778**
50. Sugiawaka H., Okouchi H. (2004) **Reformative self-control and discounting of reward value by delay or effort** *Jpn. Psychol. Res* **46**:1–9
51. Prevost C., Pessiglione M., Metereau E., Clery-Melin M. L., Dreher J. C. (2010) **Separate valuation subsystems for delay and effort decision costs** *J.Neurosci* **30**:14080–14090

52. Klein-Flugge M. C., Kennerley S. W., Saraiva A. C., Penny W. D., Bestmann S. (2015) **Behavioral modeling of human choices reveals dissociable effects of physical effort and temporal delay on reward devaluation** *PLoS.Comput.Biol* **11**
53. Bollu T., et al. (2021) **Cortex-dependent corrections as the tongue reaches for and misses targets** *Nature* :1–6

Article and author information

Paul Hage

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

For correspondence: paulhage@jhu.edu

ORCID iD: [0000-0002-7006-9877](https://orcid.org/0000-0002-7006-9877)

In Kyu Jang

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

ORCID iD: [0000-0002-1832-3743](https://orcid.org/0000-0002-1832-3743)

Vivian Looi

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

ORCID iD: [0000-0002-9392-4178](https://orcid.org/0000-0002-9392-4178)

Mohammad Amin Fakharian

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

ORCID iD: [0000-0002-7741-8689](https://orcid.org/0000-0002-7741-8689)

Simon P. Orozco

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

ORCID iD: [0000-0001-6092-7228](https://orcid.org/0000-0001-6092-7228)

Jay S. Pi

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

ORCID iD: [0000-0003-4036-2877](https://orcid.org/0000-0003-4036-2877)

Ehsan Sedaghat-Nejad

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

ORCID iD: [0000-0001-6732-1296](https://orcid.org/0000-0001-6732-1296)

Reza Shadmehr

Laboratory for Computational Motor Control, Dept. of Biomedical Engineering Johns Hopkins School of Medicine, Baltimore, Maryland USA

For correspondence: shadmehr@jhu.edu

ORCID iD: [0000-0002-7686-2569](https://orcid.org/0000-0002-7686-2569)

Copyright

© 2023, Hage et al.

This article is distributed under the terms of the [Creative Commons Attribution License \(https://creativecommons.org/licenses/by/4.0/\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Tobias Donner

University Medical Center Hamburg-Eppendorf, Germany

Senior Editor

Joshua Gold

University of Pennsylvania, United States of America

Reviewer #1 (Public Review):

The manuscript by Hage et al. presents interesting results from a foraging behavior in Marmosets that explores the interactions of saccade and lick vigor with pupil dilation and performance as well as a marginal value theory and foraging theory-inspired value-based decision-making model thereof. The results are generally robust and carefully presented and analyses, particularly of vigor, are carefully executed.

Reviewer #2 (Public Review):

Hage et al examine how the foraging behavior of marmoset monkeys in a laboratory setting systematically takes into account the reward value and anticipated effort cost associated with the acquisition and consumption of food. In an interesting comprehensive framework, the authors study how experimental and natural variation of these factors affect both the decisions and actions necessary to gather and accumulate food, as well as the actions necessary to consume the food.

The manuscript proposes a computational model of how the monkeys may guide all these aspects of behavior, by maximizing a food capture rate that trades off the food that can be gathered with the effort and duration of the underlying actions. They use this model to derive qualitative predictions for how monkeys should react to an increase in the effort associated with food consumption: Monkeys should work longer before deciding to consume the accumulated food, but should move more slowly. The model also predicts that monkeys should show a different reaction to an increase in reward value of the food, also working longer but moving faster. The authors test these predictions in an interesting experimental setup that requires monkeys to collect small increments of food rewards for successful eye movements to targets. The monkeys can decide freely when to interrupt work and consume the accumulated food, and the authors measure the speed of the eye movements involved in the food acquisition as well as the tongue movements involved in the food consumption.

By and large, the behavioral findings fall in line with the qualitative model predictions: When the effort involved in food consumption increases, monkeys collect more food before deciding to consume it, and they move slower both during food acquisition and food consumption. In a second test, the authors approximate the effects of reward value of the food at stake, by comparing monkey behavior during different days with natural variations

in body weight. These quasi-experimental increases in the reward value of food also lead to longer work times before consumption, but to faster movements during food consumption. Finally, the authors show that these effects correlate with pupil size, with pupils dilating more for low-effort foraging actions with increased saccade speed and decreased work duration. The authors conclude that the effort associated with anticipated actions can lead to changes in global brain state that simultaneously affect decisions and action vigor.

The paper proposes an interesting model for how one unified action policy may simultaneously affect multiple types of decisions and movements involved in foraging. The methods employed to measure behavior and test these predictions are generally sound, and the paper is well written.

Author Response

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

The manuscript by Hage et al. presents interesting results from a foraging behavior in Marmosets that explores the interactions of saccade and lick vigor with pupil dilation and performance as well as a marginal value theory and foraging theory-inspired value-based decision-making model thereof. The results are generally robust and carefully presented and analyses, particularly of vigor, are carefully executed.

The authors constructed a model that makes two predictions: "In summary, this simple theory made two sets of predictions: in response to an increased cost of harvest, one should work longer, but move with reduced vigor. In response to an increased reward value, as in hunger, one should also work longer, but now move with increased vigor." Their behavioral data meets these predictions. It is not clear if the model was designed and tweaked in order to make those predictions and match the data, or derived from principles. Furthermore, it is not clear what other models would make similar predictions. It would help to assess what is predicted by other simple models, as well as different functional forms for the effort costs in their model.

We chose this formulation of utility (Eq. 1) because it is a normative approach that ecologists have used to understand the decisions that animals make regarding how far to travel for food, what mode of travel to use, and how long to stay before moving on to another reward opportunity (Richardson and Verbeek 1986; Stephens and Krebs 1986; Bautista et al. 2001). In a typical formulation of the theory, the numerator represents the reward gained (in units of energy), minus the effort expended (also in units of energy). The denominator represents the amount of time spent during that behavior. We represented this idea in Eq. (1) with saccades that produced reward accumulation, and licks that produced reward consumption. Thus, the utility that we are trying to maximize is the rate of energy gained.

The specific functions that we used to represent the energy acquired through reward acquisition, and the energy expended through effort expenditure, came a priori either from experiment design, or from the measurements we have made in other experiments. We modeled reward accumulation as a linear rise in energy stored because successful saccades produced a linear increase in the food cache. We modeled consumption of the food as a hyperbolic function of the number of licks to represent the fact that as the licking bout began, each successful lick depleted the food, and thus the first few licks produced a greater amount of food consumption than the last few licks. We modeled the effort cost of licking to grow linearly with the number of licks.

A critical assumption that we made is that energy spent performing the saccade trials (which grew faster than linearly as a function of the number of trials attempted), grew faster than the time spent attempting those same trials (which grew linearly with the number of trials). This assumption is based on the heuristic that the average rate of energy lost following a large number of attempted trials is greater than the average rate of energy lost following a small number of attempted trials.

Sensitivity to parameter values: The model's simplicity provides closed-form solutions across all parameter values, allowing one to make predictions without having to fit the model to the measured data. For example, for all parameter values that produce a real solution (as opposed to imaginary), the optimal number of saccade trials increases with the square root of the cost of licking. Thus, the basic prediction of the model is that in order to maximize the capture rate, an increase in the effort that it takes to harvest the reward should produce a greater willingness to work longer, caching more food. The closed-form solutions are presented in the Mathematica supplementary document.

Other models of utility: In composing our utility (Eq. 1), we chose to combine reward and effort additively. This is in contrast to other approaches in which effort discounts reward multiplicatively (47–49). Here, let us show that multiplicative interactions may have the limitation that they are incompatible with the observation that reward invigorates movements. To compare additive and multiplicative approaches, let us consider an arbitrary function $U(T)$ that specifies how effort varies with movement duration. Typically, this is a U-shaped function that describes energy expenditure as a function of movement duration, as in Shadmehr et al. (2016). In the case of multiplicative interaction between reward and effort, we can consider the following representation of utility:

$$J = \frac{\alpha}{1+\tau} U^{-1}(T)$$

In the above formulation, reward α is discounted hyperbolically with time, and an increase in reward increases the utility of the action. The optimum movement vigor has the duration T^* that maximizes this utility. Notably, because increasing reward merely scales this utility, it has no effect on vigor. Thus, a utility in which reward is multiplied by a function of effort generally fails to predict dependence of movement vigor on reward.

Line 37 page 6; the link of pupil to NE/LC is tenuous. Other modulators systems and circuits may be equally important and should be mentioned (e.g. Reimer, Jacob, Matthew J. McGinley, Yang Liu, Charles Rodenkirch, Qi Wang, David A. McCormick, and Andreas S. Tolias. "Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex." Nature communications 7, no. 1 (2016): 13289.)

Reimer et al. (2016) used two-photon microscopy to measure activity of ACh and NE projections in layer 1 of mouse visual cortex while tracking pupil diameter fluctuations. During stillness, elevated pupil diameter was followed by cholinergic and noradrenergic axonal activity. Notably, NE activity levels were larger and with shorter latency than ACh. In primates, Joshi et al., (2016) recorded from LC during a fixation task. Using spike-triggered averaging, they found that following a spike in an LC neuron, there was pupil dilation at 200–300 ms latency. Moreover, microstimulation in LC produced pupil dilation at 500ms latency. More recently, Breton-Provencher and Sur (2019) provided causal evidence that LC activity drives pupil size. They optogenetically activated (1s) or silenced (5 sec) locus coeruleus noradrenergic neurons and found strong increase in pupil size or modest decrease: increase had a slow time scale of 1 second or more, similar slow timescale for decrease. The LC-NA neurons are surrounded by GABA-ergic neurons. Stimulation of the GABA-ergic neurons produced mild, slow constriction. They identified GABA-ergic and NA neurons by photo-

tagging and then tried to identify them via spike shape and found that “spike shape of some GABA neurons were not well separated from NA neurons, demonstrating the difficulty of cell-type identification based on spike shape alone.” They noted that a subset of GABAergic neurons received coincident inputs with the NA neurons. When the GABA neurons were excited, the gain of the pupil response to an auditory tone was diminished, producing an increase as a function of tone intensity that had a lower gain. Thus, LC-NA neurons causally drive pupil size, and the GABA neurons that surround them control the gain of the response of LC-NA neurons to arousal stimuli.

Line 35 page 6-page 7 line 10 emphasizes a cognitive interpretation of the pupil dilations that is emphasized, in relation to effort costs. But there are also more concomitant vigorous movements. Could all of their pupil results be explained by motor correlates? This should be tested and ruled out before making cognitive interpretations.

Pupil dilation is a proxy for activity in the brainstem neuromodulatory system (Vazey et al., 2018) and is a measure of arousal (Mathot, 2018). Control of pupil size is dependent on spiking of norepinephrine neurons in locus coeruleus (LC-NE): an increase in the activity of these neurons produces pupil dilation (Joshi et al., 2016; Breton-Provencher and Sur, 2019). Some of these neurons show a transient change in their activity when acquisition of reward requires expenditure of physical effort (Bornert and Bouret, 2021). However, the link between effort costs and pupil size appears to go beyond motor control, as a recent paper found that pupil size increases during effortful speech perception (Contadini-Wright et al., 2023). Thus, although in our work increases in pupil size were always associated with increased movement vigor, the results from other studies suggest that economic variables such as cognitive effort in tasks in which there is no concomitant movement also drive an increase in pupil size.

Page 7, line 37-42: How would the model need to be modified in order to account for this discrepancy with the data? Ideally, this would be tested.

We comment on potential modifications that can be made to the model that may account for the discrepancy referred to by the reviewer in the discussion section: “Notably, some of the predictions of the theory did not agree with the experimental data. An increased effort cost did not accompany a reduction in the duration of harvest, and hunger did not increase saccade vigor robustly. Indeed, earlier experiments have shown that if the effort cost of harvest increases, animals who expend the effort will then linger longer to harvest more of the reward that they have earned (2). This mismatch between observed behavior and theory highlights some of the limitations of our formulation. For example, our capture rate reflected a single work-harvest period, rather than a long sequence. Moreover, the capture rate did not consider the fact that the food tube had finite capacity, beyond which the food would fall and be wasted. This constraint would discourage a policy of working more but harvesting less. Finally, if we assume that a reduced body weight is a proxy for increased subjective value of reward, it is notable that we observed a robust effect on vigor of licks, but not saccades. A more realistic capture rate formulation awaits simulations, possibly one that describes capture rate not as the ratio of two sums (sum of gains and losses with respect to sum of time), but rather the expected value of the ratio of each gain and loss with respect to time (Bateson et al., 1995 & 1996).”

Page 9, line 2-11: In this section, it would help to also consider 'baseline' pupil size (in between trials). This would give a signal that is not 'contaminated' by movements, and may reflect control state. Relatedly, changes in control state may impact and confound the movement-related dilation magnitudes due to e.g. floor and ceiling effects on pupil size, which has a strong tendency for reversion to the mean.

The experiment design included little or no between-trial periods because during the trials the subjects worked (performed saccades to accumulate reward), while after completing a few trials they stopped working and started harvesting through licking. Because primates make saccades during their entire wake state, it is probably not possible to find a significant period in which the subjects do not make any movements. We selected a window of 500 ms around each lick in the harvest period, and each saccade during the work period, and computed the average pupil size per movement, which includes data from both before and after movements. We then computed a within-session z-score by normalizing these measures by the average pupil size acquired for that day.

The hunger-related and reward-size related analyses are both heavily confounded since they were not manipulated directly and could co-vary with many latent factors. For example, why might a given Marmoset be lower weight on a given day? Could it affect sleep, stress, activity, or other factors during the preceding 24 hours? If so, could these other variables be driving the results that are interpreted as 'hunger?' Relatedly, since the reward size is determined by the animals behavior on each trial (how much they worked), factors (internal brain state, external noises, etc.) that alter how much they worked will influence the subsequent reward size. Therefore interpretations about reward expectancy are confounded. Both of these issues should be discussed and manipulations of them (different feeding schedules and reward size-work functions proposed, respectively).

Weight of the subjects was measured prior to the start of the experiment on each day. The natural fluctuations are typically the result of factors such as time of the experiment and corresponding weight measurement (AM vs PM) relative to the time of feeding on the previous day, day of the week of the experiment (following a weekend vs. during the week), and volume of food given during the previous day. Animals were maintained at 90% of their baseline weight during food restriction, and fluctuations typically occurred within that range (Sedaghat-Nejad et al., 2019). We used weight as a proxy for hunger, and thus value of reward, and the resulting analyses yielded results consistent with predictions made by our model, as seen in Fig. 5. Critically, other factors that may co-vary with lower weights, like those mentioned by the reviewer (sleep conditions, stress levels, and activity levels) often lead to very poor task performance by the subjects. In sharp contrast, the model predicted increased work period, and increased movement vigor for high reward value, both of which we observed when the subject's weight was low. Thus, a low relative weight did not seem to impair performance, but rather act as a motivating factor. Subjects were closely monitored for well characterized stress-related behaviors and impaired attentive states by experimenters, veterinarian staff, and caretaker staff, and, in the event of abnormalities, were removed from food restriction and experimentation until behavior stabilized.

Effect of reward size: As you noted, we did not manipulate reward size directly. Rather, because our emphasis was on quantifying the effect of effort, the subjects received the same increment of reward per each completed trial, but on some sessions this reward was easy to harvest, while in other sessions the reward required greater effort to harvest. Because the reward amount accumulated during the work period, some harvests encountered a small amount of reward, while other harvests encountered a large amount of reward. Indeed, the amount of reward available for harvest depended linearly on the number of successful saccade trials completed during the work period. We found that the vigor of licks grew with the reward magnitude.

A major issue is a lack of alternative models. The authors seem to have constructed a particular model designed to capture the behavioral patterns they observed in the data. The model fails in some instances, as they point out. Even more importantly, there are no results or discussion about how other plausible models could or couldn't fit the data. The

To model behavior, we chose a formulation of utility that represented a normative approach that ecologists have used to understand the decisions that animals make regarding how far to travel for food, what mode of travel to use, and how long to stay before moving on to another patch. In the model, the objective of decisions and actions is to maximize the sum of reward acquired, minus the efforts expended, divided by time. This is termed the capture rate. However, there are other models to consider, and thus we added a new section titled Model formulation and Other models of utility.

Reviewer #2 (Public Review):

The model proposed in the paper takes a very specific functional form that is neither motivated by the previous literature nor particularly useful for indexing the behavioral tendencies of individual monkeys (or of the same monkey in different contexts). For example, while it is clear that the saccade effort cost will need to outgrow the increase in the utility of the accumulated food for the monkey to start feeding, it is unclear why this needs to be modeled with a fixed quadratic exponent on the number of saccades? Similarly, why do licks deplete the food stash with the specific rate hard-coded in the model?

We added a section titled Model formulation and Other models of utility to better explain the rationale behind the model.

We chose this formulation of utility (Eq. 1) because it is a normative approach that ecologists have used to understand the decisions that animals make regarding how far to travel for food, what mode of travel to use, and how long to stay before moving on to another reward opportunity (Richardson and Verbeek, 1986; Stephens and Krebs, 1986; Bautista et al., 2001). In a typical formulation of the theory, the numerator represents the reward gained (in units of energy), minus the effort expended (also in units of energy), while the denominator represents the amount of time spent during that behavior. We represented this idea in Eq. (1) with saccades that produced reward accumulation, and licks that produced reward consumption. Thus, the utility that we aim to maximize is the rate of energy gained.

The specific functions that we used to represent the energy gained through reward acquisition, and the energy expended through effort expenditure, came either from experiment design, or from the measurements we have made in other experiments. We modeled reward accumulation as a linear rise in energy stored because successful saccades produced a linear increase in the food cache. We modeled consumption of the food as a hyperbolic function of the number of licks to represent the fact that as the licking bout began, each successful lick depleted the food, and thus the first few licks produced a greater amount of food consumption than the last few licks. We modeled the effort cost of licking to grow linearly with the number of licks.

A critical assumption that we made is that energy expended performing the saccade trials (which grew faster than linearly as a function of the number of trials attempted), grew faster than the time spent attempting those same trials (which grew linearly with the number of trials). This assumption is based on the heuristic that the average rate of energy lost following a large number of attempted trials is greater than the average rate of energy lost following a small number of attempted trials. A quadratic function is one example of such a function, which has the advantage of providing closed form solutions for the optimal policy.

The model's simplicity provided closed-form solutions across all parameter values, allowing us to make predictions without having to fit the model to the measured data. Critically, for all

parameter values that produce a real solution (as opposed to imaginary), the optimal number of saccade trials increases with the square root of the cost of licking. Thus, the basic prediction of the model is that to maximize the capture rate, regardless of parameter values, an increase in the effort required for harvest should be met with a greater willingness to work. The closed-form solutions are presented in the supplementary document (simulations.nb).

Finally, the proportion of successful saccades and lick events is assumed to be fixed, even though it very likely to be directly influenced by movement speed (speed- accuracy trade-off), which is also contained in the model. It would strongly increase the plausibility and potential impact of the model if the authors could clearly state where these hard-coded model terms come from. Ideally, they would formulate the model in more general terms and also consider other functional forms, as briefly suggested in the discussion. This latter point would be particularly important since not all model predictions were actually borne out in the data.

Thank you for this excellent suggestion. Regarding saccades, contrary to the speed accuracy trade-off hypothesis, we found that faster saccades were also more accurate (Fig. 3C). Thus, increased pupil size was not only associated with more vigorous saccades, but also more accurate saccades. Importantly, these vigor-related changes in accuracy were too small to affect the probability of reward: the reward area for the saccades was much larger (1.5 deg) than the endpoint accuracy changes that was produced due to changes in the food tube distance. For example, on average saccade vigor changed from 0.95 to 1.05 when the food tube distance changed from 12 mm to 8 mm. These changes in vigor would produce a fraction of degree reduction in endpoint error (Fig. 3C).

Regarding licks, we added new data to the manuscript to assess the relationship between vigor of the licks and endpoint accuracy. We saw no consistent relationship, across subjects or effort conditions, between protraction speed and the outcome of a lick, that is, if the lick was successful in making it inside the tube. On average, in subject R we observed an improvement in lick accuracy with increased vigor, and in subject M we saw no change (Fig. 4F). Thus, we used the average success rate of licks, which was roughly 30% for both subjects.

The authors derive qualitative predictions, by simulating their model with apparently arbitrary parameters. They then test these qualitative predictions with conventional statistics (e.g., t-tests of whether monkeys lick more for high vs low effort trials). The reader wonders why the authors chose this route, instead of formulating their model with flexible parameters and then fitting these to data. This would allow them (and future researchers) to test their model not just qualitatively but also quantitatively, and to compare the plausibility of different functional forms. The authors certainly have enough data and power to do this, given the vast number of sessions the monkey completed.

The model's simplicity provides closed-form solutions across all parameter values, allowing one to make predictions without having to fit the model to the measured data. For example, for all parameter values that produce a real solution (as opposed to imaginary), the optimal number of saccade trials increases with the square root of the cost of licking. Thus, the basic prediction of the model is that to maximize the capture rate, an increase in the effort that it takes to harvest the reward should produce a greater willingness to work longer, caching more food. The closed-form solutions are presented in the Mathematica supplementary document.

The effort manipulation chosen by the authors (distance of food tube) goes hand in hand with a greater need for precision since the monkey's tongue needs to hit an opening of

similar size, but now located at a greater distance. This raises the question of whether the monkeys moved slower to enhance its chance of collecting the food (in line with a speed-accuracy trade off). The manuscript would benefit from an explicit test of this possibility, for example by reporting whether for each of the two conditions, the speed of tongue movements on a trial-by-trial basis predicts the probability of food collection? At the very least, the manuscript should explicitly discuss this issue and how it affects the certainty with which effects of tube distance can be linked to anticipated effort cost alone.

Thank you for the excellent point. We looked for but found no consistent relationship, across subjects or effort conditions, between protraction speed of the tongue and the success probability of a lick (probability of insertion into the tube). Regardless, we agree with you that it is an excellent alternate hypothesis that reductions in lick vigor that accompanied increased distance of the tube may be due to a desire to maintain accuracy, and not a reflection of increased effort cost of reward. To incorporate this idea into the model, we would need a measure of speed-accuracy for the licks, something that we do not have but hope to develop in the future.

However, perhaps the most interesting aspect of our results is that when we increased tube distance, making reward more effortful, there was not only a reduction in lick vigor, but also a reduction in saccade vigor. That is, the decisions and actions during the work period responded to the increased effort cost of reward during the harvest period. These changes accompanied dilation of the pupil, both in the work period and in the harvest period. We now include a paragraph regarding this in the Discussion.

The manuscript measures pupil dilation in a time period ranging from -250ms before to 250 ms after saccade onset. However, the pupil changes strongly during saccade execution relative to the preceding baseline, leaving doubts as to whether the aggregated measure blurs several interesting and potentially different effects. It would be more conclusive if the manuscript could report the analyses of pupil size separately for a period prior to saccade onset and during/after the saccade.

Our goal was to test for general correlations between the state of the pupil and both movement vigor and decisions. We chose a window of 500 ms around saccade onset, as referred to by the reviewer, as it allowed us a large enough time window to measure pupil size outside of the movement itself (~30 ms duration), to accurately capture the state of the animal around initiation and end of a saccade. Critically, pupil tracking during a saccade itself, when using infrared eye tracking techniques, can be prone to slight measurement error in certain cases due to tracking jitter. Thus, averaging across this window, following processing of the signal, results in a more accurate measure of pupil size.