

Genetics and Genomics

Histone variants shape chromatin states in *Arabidopsis*

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

July 4, 2023 (this version)

Reviewed preprint version 1

May 18, 2023

Sent for peer review

March 22, 2023

Posted to bioRxiv

March 10, 2023

Bhagyshree Jamge, Zdravko J. Lorković, Elin Axelsson, Akihisa Osakabe, Vikas Shukla, Ramesh Yelagandula, Svetlana Akimcheva, Annika Luisa Kuehn, Frédéric Berger ✉

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria
• Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical University of Vienna, A-1030, Vienna, Austria • Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan • PRESTO, Japan Science and Technology Agency, Honcho, Kawaguchi, Saitama, 332-0012, Japan • Institute of Molecular Biotechnology, IMBA, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

 (https://en.wikipedia.org/wiki/Open_access)

 (<https://creativecommons.org/licenses/by/4.0/>)

Abstract

Summary

How different intrinsic sequence variation or regulatory modifications of histones regulate nucleosome interactions with transcription remain unclear. By contrast with H3 and H2B variants, H2A variants occupy specific domains of chromatin in *Arabidopsis thaliana*. Broad domains of chromatin are affected by the loss of remodelers that affect the deposition or the exchange of H2A variants. Notably, the chromatin remodeler DECREASED IN DNA METHYLATION (DDM1) is required to maintain enrichment in all markers of constitutive heterochromatin including DNA methylation, H3K9me1/2 and the variant H2A.W. To test the importance of histone variants in the organization of chromatin we investigated how histone variants and histone modifications assemble in the *Arabidopsis thaliana* genome and showed that a limited number of chromatin states divide euchromatin and heterochromatin into several subdomains. We found that histone variants are as significant as histone modifications in determining the composition of chromatin states. Particularly strong associations were observed between H2A variants and specific combinations of histone modifications. To study the role of H2A variants in organizing chromatin states we determined the role the chromatin remodeler DECREASED IN DNA METHYLATION (DDM1) in the organization of chromatin states. We showed that the loss of DDM1 prevented the exchange of the histone variant H2A.Z to H2A.W in constitutive heterochromatin, resulting in significant effects on the definition and distribution of chromatin states in and outside of heterochromatin. We thus propose that dynamic exchanges of histone variants control the organization of histone modifications into chromatin states, acting as molecular landmarks.

eLife assessment

This study presents an **important** description on the dynamics of histone variant exchange controlling the organization of the chromatin state of the Arabidopsis genome, combining the analysis of histone variants, histone modification, and chromatin states. The evidence supporting the claims of the authors is **compelling**. This work will be of great interest to those in the field of epigenetics and chromatin biology.

Introduction

Eukaryotic genomes are packaged into chromatin, a structure defined by repeating units of ~147 bp of DNA wrapped around a protein complex known as the nucleosome (Luger et al., 1997). Nucleosomes contain two copies each of the core histones H2A, H2B, H3, and H4. Histone variants have arisen through the divergence of their intrinsically disordered loop and tail regions and dozens of histone variants in the H2A, H2B, and H3 families have been identified in eukaryotes (Loppin and Berger, 2020; Talbert and Henikoff, 2021). These variants regulate the properties of nucleosomes (Chakravarthy et al., 2004; Horikoshi et al., 2013; Koyama and Kurumizaka, 2018; Tachiwana et al., 2012) and affect transcription (Rudnizky et al., 2016; Subramanian et al., 2015; Weber and Henikoff, 2014; Wollmann et al., 2012; Wollmann et al., 2017). Animals and plants have evolved unique H2A variants associated with transcriptional repression (Loppin and Berger, 2020; Talbert and Henikoff, 2021) and with gametes (Osakabe and Molaro, 2023). In vascular plants, the H2A.W variants are distinguished by a C-terminal KSPK motif (Lei et al., 2021; Schmucker et al., 2021; Yelagandula et al., 2014). In synergy with H3K9 methylation, H2A.W favors heterochromatic silencing by directly altering the interaction between H2A.W and DNA (Bourguet et al., 2021; Osakabe et al., 2021; Schmucker et al., 2021; Yelagandula et al., 2014). The roles of the recently recognized H2B variants (Jiang et al., 2020; Raman et al., 2022) in transcription remain unknown. However, H2B variants might differ in their capacity to be ubiquitinated, an important modification for RNA polymerase II elongation (Feng and Shen, 2014; Kim et al., 2009; Minsky et al., 2008). Histone variants thus represent complex, diverse nucleosome components that could regulate and organize the functional activities of chromatin.

Histones are subject to a wide range of post-translational modifications (Talbert and Henikoff, 2021; Tessarz and Kouzarides, 2014). Applying a Hidden Markov Model (ChromHMM) (Ernst and Kellis, 2012, 2017) to genomic profiles of histone modifications revealed chromatin states that reflect the combination of post-transcriptional histone modifications present in mammals, *Drosophila*, *Arabidopsis*, and rice (Ernst and Kellis, 2012, 2017; Kharchenko et al., 2011; Liu et al., 2018; Roudier et al., 2011; Sequeira-Mendes et al., 2014). Chromatin states distinguish the major domains of chromatin within the genome, comprising euchromatin (with active genes), facultative heterochromatin (with repressed genes), and constitutive heterochromatin (with transposons and repeats) in animals and plants. They also highlight specific features such as promoters and enhancers.

Compared with the wealth of data on the relationship between histone modifications and transcription, the role of histone variants in this process remains to be explored. In *Arabidopsis*, there is a clear correlation between transcript levels and enrichment of the variant H3.3 (Stroud et al., 2012; Wollmann et al., 2012; Wollmann et al., 2017) and H2A and H2A.X (Yelagandula et al., 2014) on gene bodies marked by H3K36me2 (Leng et al., 2020). In

contrast, H2A.Z is enriched on repressed genes marked by H3K27me3 (Carter et al., 2018), and the histone variant H2A.W is present on transposons and repeats marked by H3K9me2 (Osakabe et al., 2021; Yelagandula et al., 2014). Although it was suggested that histone variants might have a global role in defining chromatin states (Hake and Allis, 2006; Henikoff et al., 2004; Weber and Henikoff, 2014) the complex, intimate associations between histone variants and modifications have not been fully characterized in plants or mammals.

Here, we analyzed how all thirteen histone variants expressed in vegetative tissues associate with twelve prominent histone modifications to form chromatin states in the model flowering plant *Arabidopsis thaliana*. Our findings indicate that H2A variants are major factors that differentiate euchromatin, facultative heterochromatin, and constitutive heterochromatin. This hypothesis is supported by in silico analyses and the mis-assembly of chromatin states caused by the deregulation of the exchange of H2A variants.

Results

Co-occurrence of H3 modifications and histone variants in nucleosomes

In *Arabidopsis*, homotypic nucleosomes containing a single type of H2A variant are prevalent (Osakabe et al., 2018). However, it was not determined whether H3 variants also assemble in a homotypic manner and if H2A variants preferably assemble with a specific H3 variant or histone modification. To answer these questions, we immunoprecipitated mononucleosomes (**Figure 1 figure supplement 1A-C**) from transgenic *Arabidopsis* lines expressing HA-tagged *HTR13* (H3.1) and *HTR5* (H3.3) under the control of their native promoters in the respective mutant backgrounds (Jiang & Berger, 2017). Because tagged H3 ran slower than endogenous H3 on SDS-PAGE (**Figure 1 figure supplement 1B**), this approach allowed us to analyze the composition of H3.1 and H3.3 nucleosomes by mass spectrometry (MS) and immunoblotting. Spectral counts of peptides covering regions around lysine 27 (K27), where H3.1 and H3.3 can be distinguished by MS analysis, revealed that ~60% of H3.1 and ~42% of H3.3 nucleosomes contained both H3 variants, (**Figure 1A**). We analyzed H3.1 and H3.3 nucleosomes for the presence of H2A variants by western blotting and found that neither H3.1 nor H3.3 was preferentially associated with a specific H2A variant (**Figure 1B**). This was further confirmed by MS analysis of nucleosomes pulled down with H2A variant-specific antibodies (**Figure 1 figure supplement 1D**). Thus, by contrast with H2A variants, H3 variants do not necessarily form homotypic nucleosomes, despite the presence of H3 variant-specific deposition mechanisms (Loppin and Berger, 2020; Nie et al., 2014; Probst, 2022), and they do not associate with a specific H2A variant.




Figure 1.

Biochemical analysis of the association between histone variants and histone marks.

(A) Histone H3.1 and H3.3 form homotypic and heterotypic nucleosomes. Spectral counts of H3.1- and H3.3-specific peptides in the respective immunoprecipitations (T – transgenic, E – endogenous H3.1 and H3.3). **(B)** H2A variants do not preferentially associate with H3.1- or H3.3- containing nucleosomes. HA-tagged H3.1 and H3.3 mononucleosomes were immunoprecipitated with HA agarose and analyzed for the presence of H2A variants by immunoblotting. **(C)** Histone H3 marks are present on both H3.1 and H3.3. HA-tagged H3.1 and H3.3 mononucleosomes were immunoprecipitated with HA agarose and analyzed for the presence of H3 marks by immunoblotting. Arrows indicate transgenic (T) and endogenous (E) H3. **(D)** MS analysis of cumulative H3K27, H3K36, and H3K37 modifications on H3.1 and H3.3. All measured spectra corresponding to H3.1 and H3.3 peptides from both IPs were used for analysis. **(E)** Relative abundance of H3K27, H3K36, and H3K37 modifications on H3.1 variant analyzed separately from MS data of H3.1 and H3.3 purified nucleosomes (left panel). Relative abundance of H3K27, H3K36, and H3K37 modifications on H3.3 variant analyzed separately from MS data of H3.1 and H3.3 purified nucleosomes (right panel). **(F)** Co-occurrence of H2A variants and H3

marks. Mononucleosomes were immunoprecipitated with the indicated antibodies and analyzed for the presence of H2A variants and H3 marks by western blotting.

Next, we explored the potential link between H3 modifications and H3 variants. We found that all the methylation and acetylation marks surveyed were present on transgenic (T) and endogenous (E) H3 in both H3.1 and H3.3 immunoprecipitations (**Figure 1C**), although there was some degree of preferential association of H3K4me, H3K36me and H3 acetylation with H3.3 and of H3K27me1 with H3.1 (**Figure 1C**). H3.1 and H3.3 modifications were also analyzed by MS by using information from both transgenic and endogenous H3 in respective immunoprecipitations. This approach unambiguously distinguishes the modifications of peptides covering lysines 27 (H3K27) and 36 (H3K36) originating from the endogenous or the transgenic copies of H3.1 and H3.3 (**Figure 1 figure supplement 1E**). We put our focus on modifications at these two positions, because they are diagnostic of constitutive heterochromatin (H3K27me1), facultative heterochromatin (H3K27me3) and actively

transcribed euchromatin (H3K36me). In contrast to previous reports (Johnson *et al.*, 2004; Zhang *et al.*, 2007) we identified similar levels of H3K27me2 and H3K27me3 marks on both H3.1 and H3.3 whereas the constitutive heterochromatin mark H3K27me1 was higher on H3.1 (**Figure 1D and E; Figure 1figure supplement 1E-G**). High levels of H3K27me1 on H3.3 were unexpected considering specificity of H3K27 methyltransferases ATXR5/6 for H3.1 (Jacob *et al.*, 2014) and are the likely product of demethylation of H3K27me3 by the JUMONJI H3K27 demethylases (Gan *et al.*, 2015; Liu *et al.*, 2010) or intermediates of methylation by the Polycomb Repressive Complex 2. Acetylation and all three methylation states at H3K36 that are associated with active chromatin were likewise found on both H3.1 and H3.3 but with higher enrichment on H3.3 (**Figure 1D and E; Figure 1figure supplement 1E-G**). This is consistent with western blot analyses (**Figure 1C**) and with the established role of H3.3 as a replacement variant during transcription (Delaney and Almouzni, 2023). The levels of H3K27 and H3K36 modifications on H3.1 and H3.3 displayed the same trends irrespective of whether they were precipitated with either H3.1 or H3.3 nucleosomes, contrasting with the predominant enrichment of H3K37 modifications on H3.3 (**Figure 1E**). The H3 region covering K4, K9, K14, K18, and K23 residues cannot be distinguished between H3.1 and H3.3 by either bottom-up MS or western blotting. Therefore, we analyzed H3K9me1 and H3K9me2, which mark constitutive heterochromatin in plants, only on transgenic H3 and found that both marks were similarly distributed among H3.1 and H3.3 (**Figure 1figure supplement 1H**) as reported previously (Johnson *et al.*, 2004). Also, acetylation of H3K9, H3K14, H3K18, H3K23 and combinations thereof were very similar in H3.1 and H3.3 immunoprecipitations (**Figure 1figure supplement 1I-K**). Modifications on H3K4 were not analyzed by MS because our experimental setup produced too short peptides to be captured by nanoLC. As the total numbers of measured spectra covering H3 modifications were nearly identical for H3.1 and H3.3 (**Figure 1figure supplement 1F**) our MS together with western blot data suggest that the vast majority of H3 modifications are not H3 variant specific to the notable exception of H3K36 and H3K37 modifications that are associated preferentially but not uniquely with H3.3. This preference for H3.3 might be related to the presence of substitutions at position 41 specific that distinguishes H3.1 from H3.3 in plants (Borg *et al.*, 2021a; Probst, 2022). These findings suggested that in the heterotypic nucleosomes the same modifications are found on either H3 variants present irrespective of their identity.

Unlike H3 variants, homotypic nucleosomes containing either H2A.W, H2A.Z, H2A, or H2A.X showed marked enrichment of specific histone modifications (**Figure 1F**). Histone modifications in constitutive heterochromatin (H3K9me1, H3K9me2, and H3K27me1) were primarily associated with H2A.W consistent with their synergistic impact on silencing transposons (Bourguet *et al.*, 2022). The modification H3K27me3 (facultative heterochromatin) was detected only in H2A.Z nucleosomes, and H3K36me3 (marking euchromatin) was primarily detected in H2A, H2A.X, and H2A.Z nucleosomes. Other modifications, such as H3K4me1 and H3K4me3, displayed complex patterns of association with H2A variants, among themselves, and with other H3 modifications (**Figure 1F**). Surprisingly, H3K4me1 and H3K4me3 co-precipitated high levels of H3K27me1, H3K27me3, and H3K36me3 and low levels of H3K9me1 and H3K9me2 (**Figure 1F**) showing that repressive and active H3 marks co-exist on the same nucleosomes or potentially on the same H3 tail. Similar complex interplay between H3 marks has also been observed in mouse ES cells (Schwammle *et al.*, 2016). Therefore, H2A but not H3 variants form homotypic nucleosomes that preferentially carry specific histone modifications associated with either the transcriptional status of protein-coding genes or transposons.

Associations of histone variants with the chromatin states

We performed ChIP-seq using Arabidopsis seedlings to identify the combinations of all histone variants present in somatic cells and their associations with twelve prominent histone modifications. We included the most abundant isoforms of H2A.W (H2A.W.6 and H2A.W.7), H2A.Z (H2A.Z.9 and H2A.Z.11), and H2A (H2A.2 and H2A.13) in our analysis. We used specific polyclonal antibodies to detect each H2A variant (see [Figure 2 figure supplement 1](#) for the demonstration of antibodies against H2A.Z.11 and H2A.2 and (Osakabe et al., 2021; Osakabe et al., 2018). The algorithm ChromHMM (Ernst and Kellis, 2017) was used to define chromatin states in Arabidopsis (see Methods for details). We chose to analyze the 26 chromatin states model, which proved to be most parsimonious ([Figure 2 figure supplement 2A](#); see Methods). Chromatin states were clustered based on the emission probability for each modification and histone variant ([Figure 2A](#)) and were distinguished based on distinct combination of enrichment amongst the 27 chromatin components analyzed ([Figure 2A](#)). Chromatin states occupied different proportions of the genome ([Figure 2 figure supplement 2B](#)) and showed distinct relative abundance on transposons, repeats, and elements of protein-coding genes ([Figure 2B](#)). For each chromatin state, we measured the average level of transcriptional activity ([Figure 2C](#)), the degree of enrichment in CG methylation ([Figure 2D](#)), the degree of enrichment in CHG and CHH methylation ([Figure 2 figure supplement 2C and D](#)), the level of accessibility by DNase I-seq ([Figure 2E](#)), the nucleosome density determined by MNase-seq ([Figure 2 figure supplement 2E](#)), and the length in base pairs ([Figure 2 figure supplement 2F](#)). Chromatin states H1–H6 showed low accessibility, the highest levels of DNA methylation, and primarily occupied transcriptionally inactive transposons and repeats, thus representing constitutive heterochromatin. The states F1–F6 were associated with low transcriptional activity and were present over genes and pseudogenes, indicating that they represented facultative heterochromatin. As defined by state occupancy, constitutive and facultative heterochromatin composed 17% and 20% of the genome respectively ([Figure 2 figure supplement 2B](#)), corresponding to previous estimates (Roudier et al., 2011). The states E1–E11 occupied expressed genes and thus comprised euchromatin. Three states (I1–I3) showed the lowest nucleosome density and highest accessibility and occupied a large fraction of non-coding RNAs ([Figure 2 figure supplement 2G](#)) and a quarter of untranslated regions of genes (3'UTR and 5'UTR) ([Figure 2B](#)). Hence, the states I1–I3 were classified as intergenic. We conclude that specific groups of chromatin states define constitutive heterochromatin (H1–6), facultative heterochromatin (F1–6), euchromatin (E1–11), and intergenic regions (I1–3).




Figure 2.

Histone variants define chromatin states in *Arabidopsis thaliana*. **(A)** Bubble plot showing the emission probabilities for histone modifications/variants across the 26 chromatin states. (The size of the bubble represents the emission probability ranging from 0 to 1). The colors are ascribed for each type of chromatin. **(B)** Stacked bar plot showing the overlap between annotated genomic features and chromatin states. **(C)** Box plot showing the expression of protein-coding genes overlapping with each chromatin state in Transcripts per Million (TPM). **(D)** Box plot showing levels of CG methylation across chromatin states. **(E)** Box plot comparing DNase I-seq read coverage across chromatin states representing chromatin accessibility. **(F)** Heatmap showing the Jaccard similarity index between the states generated using the whole model and states using a subset of marks, i.e. excluding a set of marks and variants as indicated on the X-axis. The comparison

with 9-state model (Sequeira-Mendes et al., 2014) did not include CG content, DNA methylation, H4K5ac and H3K4me2 which were not used in the 26-state model.

Overall, the chromatin states recapitulate the preferential associations between H2A variants and histone modifications observed in mononucleosomes (Figure 1; Figure 1 figure supplement 1). H2A.W and H3K9me1 are the determining marks of all six heterochromatic states. H2A.Z, with the polycomb histone modifications H2AK121ub and H3K27me3, are the hallmark of facultative heterochromatin states. H2A and H2A.X associated with H2Bub and H3K36me3 mark euchromatic states. Contrasting with the strong association between H2A variants and the three major domains of chromatin, much less prominent associations are observed between histone modifications and either H3 or H2B variants.

To examine the importance of H2A variants in the definition of chromatin states, we compared the 26 chromatin states defined in our study with the 9 states defined in a previous study that did not include the comprehensive set of histone variants present in *Arabidopsis* chromatin (Sequeira-Mendes et al., 2014) (Figure 2 figure supplement 2H). The blocks of heterochromatic states H1–H6 corresponded to the previously identified states 8 and 9, which were also defined as heterochromatin in a previous study (Sequeira-Mendes et al., 2014). States F1–F6 tended to map to the previously defined facultative heterochromatin states 5 and 6, although state F6 was split among the three states 2, 4, and 6. Similarly, although there was a broad correspondence between euchromatin states E1–E11 with states 1, 3, and 7, there were noticeable differences. Overall, several newly defined states were not associated with the corresponding types of chromatin domains described in previous

studies. Previously defined states 2, 4, and 6 contained elements belonging to multiple newly defined chromatin states. To test the dissimilarity between the 9-state and the 26-state models we calculated a 26-state model based on the chromatin marks used to define the 9-state model. The difference between the resulting matrices was measured by the Jaccard index (Figure 2F). This showed an overall coherence between the two models, with only mis-assignment of chromatin identity by the 9-state model of chromatin in intergenic regions, H1, H5, H6, and a lower diversity of euchromatin states. Altogether, the addition of the comprehensive set of histone variants in the 26-state model provides a more refined and coherent classification of elements of chromatin than the 9-state model defined primarily based on chromatin modifications, suggesting the importance of histone variants in the definition of chromatin states.


To evaluate the contribution of individual components in defining the chromatin states, we recalculated chromatin states after excluding either H2B variants, H3 variants, H2A variants, H3 modifications (no H3 PTMs) or all histone variants (no H2A/H3/H2B) from the dataset used to learn the 26-state model. The difference between the resulting matrices of chromatin states was measured by the Jaccard index (Figure 2F). Removing H2B variants did not affect most chromatin states, except for H1 and I2, which showed high emissions probabilities for H2B variants (Figure 2A). In contrast, H3 variants showed larger contributions in defining specific states, and removing all H2A variants caused the loss of several states marked by a Jaccard index < 0.7 (Figure 2F). Eventually, removing all histone variants had a stronger impact than removing all H3 modifications from the model. In summary, our computational and biochemical analyses led to conclude that, among histone variants, H2A variants have the strongest effect on defining the chromatin states by their association with histone modifications.

Perturbation of H2A variant dynamics affects the genomic localization of chromatin states

We considered the idea that the dynamics of exchange of H2A variants would affect the chromatin states. To test this idea, we studied the impact of the loss of the chromatin remodeler DDM1, which binds and controls the deposition of H2A.W (Osakabe et al., 2021), on the definition of chromatin states. In the *ddm1* mutant, TEs show a loss of DNA methylation, H3K9 methylation, and linker histone H1 as well as an increase in H3K27me3 (Jeddeloh et al., 1998; Kakutani et al., 1999; Osakabe et al., 2021; Rougee et al., 2021). This suggested that the loss of H2A.W deposition had a profound impact on chromatin states.

In our previous study we used leaves of five-week-old plants to show the impact of *ddm1* on the profiles of H2A.W.6, H2A.X, H1, H3K9me2, H3K36me3 and H3K27me3 (Osakabe et al., 2021). This study showed that DDM1 causes the deposition of H2A.W.6 to heterochromatin and we thus extended this investigation to the two other marks of heterochromatin H3K9me1, H2A.W.7 and H2A.Z.9 and H2A.Z.11 in leaves (11 profiles in total) of *ddm1* and wild type. Because these profiles were acquired on leaves while seedlings were used to define the 26-state model in the wild type, we first considered whether the development stage could affect the definition of chromatin states in the wild type. To test this idea, we built a concatenated chromHMM model based on the profiles of the same set of histone variants and modifications from 10-day-old seedlings and leaves of five-week-old plants (Figure 3 figure supplement 1A). Although this model had only 15 chromatin states we observed an association between H2A variants and histone modifications comparable to the model describing the 26-state model (Figure 2A). The absence of most typical euchromatic histone modifications in the model likely explained the strong reduction of the number of states of euchromatin compared with the 26-state model (Figure 3 figure supplement 1A). The complexity of the different heterochromatin states was preserved in the concatenated

model, supporting that the set of H2A variants and histone modifications used to study *ddm1* is sufficient to represent the complexity of the heterochromatin affected directly by *ddm1*. The composition of the chromatin states did not vary significantly between seedlings and leaves and each state occupied a similar proportion of the genome in seedling or leaves to the exception of state 5 present primarily in leaves and state 13 only present in seedlings (Figure 3figure supplement 1A, right column with green bars). However, as expected by the dissimilar transcriptomes of these two developmental stages euchromatic states occupied distinct genes in seedlings and leaves (Figure 3figure supplement 1B). This indicated that neither tissue type nor developmental stage have a dramatic effect on the definition of chromatin states in the wild type.



with H2A.W and H2A.Z. Coomassie stained 15% SDS-PAGE gel showing input protein samples (top panel) used for *in vitro* pull-down (bottom panel) with His₆-tagged DDM1 and histone dimers. The lane Δ CTH2A.W shows that the deletion of the C-terminal tail of H2A.W does not influence binding to DDM1. (H) Summary of the pull-down assays to identify regions in DDM1 binding to H2A.W and H2A.Z. Blue and purple boxes indicate the H2A.W binding regions in DDM1 identified by previous work (Osakabe et al., 2021). Original gel pictures are shown in Figure 3 figure supplement 2A-C.

Figure 3.

DDM1 loss of function disrupts chromatin states in *Arabidopsis thaliana*. (A) Heatmap showing the emission probability for each mark/variant across the 16 chromatin states of the concatenated wild type and *ddm1* mutant model. The bar plot on the left represents the proportion of the genome covered by each state in wild type (green) and in *ddm1* (red). (B) Bar plot showing the Jaccard indices between the state assignments in wild type and *ddm1* mutant. (C) Bar plot showing the state assignment overlap between the wild type and *ddm1* for each chromatin state. The red vertical line represents the genome wide overlap (62.2%). (D) Bar plot showing the log₂ fold changes of proportion of genome covered by each state across the *ddm1* genome compared to wild type. (E) Stacked bar plot showing the overlap between annotated genomic features and chromatin states from the concatenated model in wild type. (F) Stacked bar plot showing the overlap between annotated genomic features and chromatin states from the concatenated model in *ddm1* mutant. (G) DDM1 interaction

To directly compare chromatin states between *ddm1* and wild type, we built a concatenated chromHMM model (Ernst and Kellis, 2017). The concatenated model created a common set of 16 chromatin states that are shared between wild type and *ddm1* (Figure 3A) and identified the percentage of the genome occupied by each chromatin states for each genotype individually (see Methods) (Figure 3A, and bar plot on the left). The states were classified as constitutive heterochromatin (hI-hIII), facultative heterochromatin (fI-fVI), euchromatin (eI-eIII), and intergenic (iI) (Figure 3A), based on their emission probability and genomic overlap with states from those groups in the 26-state model (see Methods). The presence of chromatin profiles from *ddm1* caused three additional mixed states in the 16-state model, which represent regions covered by facultative and constitutive heterochromatin (fhI) or by a combination of intergenic and constitutive heterochromatin (ihI,ihII) in the 26-state model. Because several euchromatic histone PTMs were not included in the new model, only three states represented euchromatin (eI-eIII) compared with the 11 euchromatic states in the 26-state model. In the 16-state concatenated model with *ddm1* the complexity of facultative (fI-fVI) was preserved but the complexity of heterochromatin states was reduced to three (hI-hIII) compared with the six heterochromatin states in the wild type 26-state model. Overall, *ddm1* caused a perturbation of the chromatin states associated with constitutive heterochromatin and did not significantly affect the group similarity for facultative heterochromatin and euchromatin states between the 16- and 26- state models, with genomic overlaps between 68 and 100% for the non-mixed states (Figure 3 figure supplement 1C).

To compare the chromatin state assignments between *ddm1* and wild type genomes, we measured the Jaccard index (Figure 3B; Figure 3 figure supplement 1D) and overlap (Figure 3C) of each state between the wild type and *ddm1*. The strongest differences, in terms of genomic coverage, between *ddm1* and wild type were observed in constitutive and facultative heterochromatin (Figure 3D). Accordingly, the regions marked by heterochromatin (red states h) became covered by states typical of facultative heterochromatin (blue states f) or intergenic states (gray states iI,ihII) (Figure 3 figure supplement 1C and 1E). To a lesser extent, regions covered by facultative state fVI and euchromatic state eIII were converted to intergenic state iI. Hence, the loss of DDM1 broadly affected the association of chromatin states with regions covered with constitutive heterochromatin in the wild type. Accordingly, chromatin states were changed over TEs, including TE fragments and TE genes (assigned to TE families in TAIR10 annotation) but did not change over protein coding genes (Figure 3 figure supplement 1F). Overall, the constitutive heterochromatin present over TEs in wild type was replaced in *ddm1* by chromatin states found in intergenic, facultative heterochromatin, and euchromatin in wild type (Figure 3E and F; Figure 3 figure supplement 1E and F). In addition, we observed that *ddm1* affected not only constitutive heterochromatin but also regions of euchromatin and facultative heterochromatin that adopted chromatin states distinct from these found in the wild type. suggesting indirect effects of the disruption of constitutive heterochromatin caused directly by the loss of DDM1 (Figure 3 figure supplement 1F). We thus concluded that loss of DDM1 causes a profound perturbation of both the definition of chromatin states and their distribution.

The H2A.W-binding domains of DDM1 bind H2A.Z

The conversion of chromatin states occupied by H2A.W in wild type to chromatin states occupied by other H2A variants in *ddm1* suggested that DDM1 could also control the dynamics of other H2A variants. We have previously shown that DDM1 specifically binds H2A.W but neither H2A.X nor H1 linker histones (Osakabe et al., 2021). To determine if DDM1 binds to H2A or H2A.Z in addition to H2A.W, we performed a Ni-NTA agarose pulldown assay with recombinant histone heterodimers composed of H2B and one of the

four H2A variants. We found that His₆-tagged DDM1 co-precipitated heterodimers containing H2A.W and H2A.Z but not H2A or H2A.X (**Figure 3G**). This demonstrated that DDM1 bound H2A.W and H2A.Z but not H2A and H2A.X. Notably, the C-terminal tail of H2A.W was dispensable for DDM1 binding (**Figure 3G**). To test whether the conserved motifs involved in H2A.W binding also bind H2A.Z, we prepared a series of DDM1 fragments fused with either His₆- or GST-tag. Our pulldown assays showed that fragments containing residues 100-123 or 639-673 were able to bind H2A.W and H2A.Z, whereas some additional regions of DDM1 showed weak binding only to H2A.Z (**Figure 3H**; **Figure 3figure supplement 2A-C**). Thus, DDM1 uses the same conserved sites to bind H2A.Z and H2A.W.

Exchange of chromatin states defined by H2A.W and H2A.Z is not directly associated with transcription

H2A.W evolved in land plants independently from macroH2A in animals (Osakabe and Molaro, 2023). macroH2A is also associated with heterochromatin although not as strictly as H2A.W (Sun and Bernstein, 2019). The deposition of macroH2A depends at least in part on the DDM1 murine ortholog LSH (Ni et al., 2020) and it was proposed that LSH binds and exchanges H2A to macroH2A (Ni and Muegge, 2021). Finding that DDM1 uses the same sites to bind both H2A.Z and H2A.W suggested that DDM1 controls the deposition of H2A.W and H2A.Z. To test this hypothesis, we surveyed the enrichment of H2A variants and histone PTMs over TEs in *ddm1* and compared this with wild type. To prevent carry-over mutations from generations of self-fertilized *ddm1* homozygous mutants we used leaves from *ddm1* homozygous mutant plants segregated from heterozygous *ddm1* mutants. We observed an overall enrichment of H2A.Z over TE genes in *ddm1*, supporting the idea that DDM1 promotes the removal of H2A.Z at TE genes where DDM1 deposits H2A.W in the wild type (**Figure 4A**). Remarkably, non-transcribed TE genes also accumulated H2A.Z (**Figure 4A**) supporting the conclusion that the replacement of H2A.W by H2A.Z observed in *ddm1* mutants is not governed by transcription. Transcribed TE genes showed enrichment in H2A.Z at their TSS (**Figure 4A**) but also H2A.X on the gene body (**Figure 4 figure supplement 1A**). Transcription of TEs in *ddm1* did not affect the degree of enrichment for H3K27me₃ of facultative heterochromatin but with increasing levels of transcription, there was increased enrichment of H3K36me₃, resulting in a chromatin profile typical of expressed protein-coding genes (**Figure 4 figure supplement 1B**). Accordingly, strongly expressed TE genes showed an accumulation of euchromatic states, in contrast with non-expressed TE genes that became covered with states typical of facultative heterochromatin, irrespective of the type of constitutive heterochromatin observed on these TE genes in wild type (**Figure 4B and C**; **Figure 4 figure supplement 2A**). The change of chromatin states did not correlate with the length of the TE gene (**Figure 4 figure supplement 2B**) or the homogeneity of the chromatin landscape (**Figure 4 figure supplement 2C**). We conclude that at TE genes, DDM1 prevents the replacement of H2A.W by H2A.Z in a transcription-independent manner. Whether the replacement of H2A.W by H2A.Z in *ddm1* is caused by a direct exchange comparable to the exchange of H2A by H2A.Z catalyzed by SWR1 (Ranjan et al., 2020) or is the result of a cascade of activities of chaperones and remodelers is unclear. Yet, the alteration in the dynamics of H2A.W and H2A.Z in *ddm1* perturbs the allocation of chromatin states at constitutive heterochromatin and in other regions of chromatin. In some TEs, the new chromatin state acquired in *ddm1* permits transcription, which is then accompanied by a further exchange of H2A.Z to H2A and H2A.X and further loss of marks of constitutive heterochromatin and H1 and gain of marks of euchromatin (**Figure 4 figure supplement 1**).




Figure 4.

Impact of expression on chromatin states over TE genes in *ddm1*. **(A)** Enrichment profiles of H2A.W.6 and H2A.Z.9 over TE genes in *ddm1*. TE genes were grouped by expression in *ddm1* mutant. Out of the 3901 TE genes in the Arabidopsis genome annotation, 497 were excluded because they showed expression in wild type, 2116 TE genes showed no expression in *ddm1* (non-expressed TE genes) while 1288 TE genes were expressed. Because many of these TE genes showed very low expression levels, we divided the expressed TEs into 4 quartiles (322 TE genes each) based on their expression values where the 1st quartile contains TE genes with lowest expression and the 4th quartile contains TE genes with highest expression. Given that the TE genes in 1st and 2nd quartile showed nominal expression values, we placed only TE genes in 3rd and 4th quartile (644 TE genes) in the category of expressed TEs. n represents the number of TE genes in each group. **(B)** Stacked bar plots of the proportion of states in wild type (top panel) and in *ddm1* (bottom panel) overlapping TE genes grouped by expression in *ddm1*. **(C)** Box plot showing the expression of TE genes overlapping the 16 concatenated model states in *ddm1*.

Discussion

We report that histone modifications and histone variants combine into chromatin states that subdivide the intergenic space, the non-protein-coding space, and the protein-coding genic space of the genome into biologically significant subunits. These chromatin states summarize the prevalent organization of chromatin across cell types and stages of vegetative development from seedlings to mature leaves. Our data suggest that minor variations of chromatin states occur between vegetative cell types. We predict that highly specialized cell types such as gametes (Borg et al., 2020; Borg et al., 2021b) with specific histone variants and unusually high or low levels of chromatin regulators harbor specific chromatin states that are not reflected in our study. Calculation of chromatin states and biochemical analyses demonstrate the strongest associations between nucleosomes homotypic for H2A variants and histone modifications while nucleosomes are primarily heterotypic for H3 variants with a weaker association with histone modifications.

The prominent role of H2A variants in the organization of chromatin states is supported by systematic evaluation of individual contribution of histone variants, H3 marks and combination thereof (Figure 2F) and the impact of the loss of the remodeler DDM1, which deposits H2A.W to maintain TE silencing (Osakabe et al., 2021). We further show that the DDM1 binding sites for H2A.W also bind H2A.Z, but not H2A or H2A.X. In *ddm1*, we observe a net replacement of H2A.W by H2A.Z. The effect can be interpreted in several manners. DDM1 could exchange directly H2A.Z to H2A.W. This model would support the idea of the convergent evolution of DDM1 and its orthologs LSH and HELLS in mammals with LSH catalyzing the exchange between replication dependent H2A and macroH2A (Berger et al., 2023; Ni and Muegge, 2021). An alternative model is that the replacement of H2A.W by H2A.Z is indirect in *ddm1*. For example, H2A.Z might be deposited because of the loss of DNA methylation in *ddm1*, which is consistent with the well-established mutually exclusive patterns of H2A.Z and DNA methylation (Zilberman et al., 2008). It is also possible that the binding of DDM1 to H2A.Z is relevant in a different context that is not illustrated by our study. Whichever mechanism is involved it is unlikely that on its own, the loss of H2A.W in *ddm1* causes the deposition of H2A.Z on constitutive heterochromatin because this is not observed in mutants lacking H2A.W (Bourguet et al., 2021). The resulting enrichment of H2A.Z in *ddm1* is accompanied by a corresponding exchange of constitutive to facultative chromatin states over TEs. Notably, this does not require transcription, as TEs that remain silent in *ddm1* are likewise occupied by a facultative heterochromatin marked by H2A.Z and H3K27me3. When TEs are strongly expressed in *ddm1*, they also acquire euchromatin states marked by H2A and H2A.X over their gene body, while H2A.Z remains confined to the transcription start site. Hence, H2A variants appear to be important in shaping the deposition of histone modifications and defining chromatin states.

Overall, our observations suggest that chromatin writers and erasers operate with different affinities and efficacies based on the H2A variants in the nucleosome, an idea that is supported by previous reports. The inhibition of the H2B ubiquitin ligase by the H2A.Z tail (Surface et al., 2016) is sufficient to explain the anticorrelation between H2Bub and H2A.Z over gene bodies. H2A.Z is the major substrate for H2AK121Ub deposited by PRC1, as indicated by their co-occurrence within chromatin states as well as previous biochemical analyses (Gomez-Zambrano et al., 2019). PRC2 strongly prefers arrays of H2A.Z as a substrate (Wang et al., 2018), thus supporting the link between H2A.Z and H3K27me3 (Carter et al., 2018). We propose to expand these examples to the more general idea that the nature of the H2A variant controls the activity of the machinery that either deposits or erases the histone modifications. Such a model would explain the importance of histone variants in the establishment of chromatin states.

How are chromatin states maintained? In mammalian cells, there is a broad maintenance of the patterns of H3 modifications by recycling H3 and H4 at the replication fork (Reveron-Gomez et al., 2018). As plant cells divide, H3.1 sustains the deposition of H3K27me3 (Jiang and Berger, 2017) and H3K27me1 (Jacob et al., 2014) thus providing a positive feedback loop to sustain the polycomb repressive state and constitutive heterochromatin in dividing cells, respectively. Recent data in mammalian cells show that the recycling of H2A-H2B after replication provides another positive feedback loop to maintain the patterns of H2A variants (Flury et al., 2023). Such replication-dependent maintenance mechanisms no longer operate in non-dividing cells, providing opportunities for changing the allocation of chromatin states. This is illustrated by several studies in Arabidopsis. The positive feedback loops that maintain histone modifications become inactive and reversion to active chromatin states becomes possible by the deposition of H3.3 and eviction of H2A.Z by INO80 in response to temperature (Zhao et al., 2023) or to light (Willige et al., 2023). Mechanisms coupling H3.3 and H2A.Z dynamics are likely general for the regulation of responsive genes that are covered with facultative chromatin states (Long et al., 2023). Conversely on expressed genes,

acetylated histones assist in the docking of the chromatin remodeler SWI2/SNF2-related 1 (SWR1), which deposits H2A.Z at the transcription start sites (TSS) of expressed genes (Aslam et al., 2019; Bieluszewski et al., 2022).

Although H3.1 and H3.3 do not strongly differentiate the three main classes of chromatin, they likely refine their definition. A relative enrichment of H3.1 with constitutive heterochromatin modifications (Johnson et al., 2004) was previously noted and confirmed in our study. Accordingly, the preferred substrate for the deposition of the heterochromatic mark H3K27me1 is H3.1 (Jacob et al., 2014). We also observed a relationship between H3.3 and H3K36me1/2, pointing to a possible preference of H3.3 for the deposition of H3K36me or for the demethylation of H3K36me3. In addition, it is becoming apparent that the dynamics of H3.3 and H2A.Z are coordinated in Arabidopsis (Zhao et al., 2023) and mammals (Yang et al., 2022). We thus propose that as yet unknown mechanisms link the deposition of H2A and H3 variants, resulting in specific types of nucleosomes that are sufficient to orchestrate the deposition of PTMs in distinct chromatin states. These will be read, interpreted, and further modified by silencing mechanisms in heterochromatin and the transcriptional machinery in euchromatin, leading to a changing chromatin landscape responding to the activity of the cell and its responses to the environment.

Methods

Generation of antibodies, isolation of nuclei, MNase digestion, immunoprecipitation, SDS-PAGE and Western blotting

Antibodies against H2A.Z.11 (KGLVAAKTMAANKDKC) and H2A.2 (CPKKAGSSKPTTEED) peptides were raised in rabbits (Eurogentec) and purified by a peptide affinity column. Purified IgG fractions were tested for specificity on nuclear extracts from WT and knock out lines or with overexpressed histone variants (Figure 2 figure supplement 1). Specificities of custom-made polyclonal antibodies against Arabidopsis H2A.Z.9, H2A.X, H2A.W.6, H2A.13, H2A.W.7, H2Bs, and linker histone H1 were validated in previous publications (Jiang et al., 2020; Lorkovic et al., 2017; Osakabe et al., 2018; Yelagandula et al., 2014).

For MNase digestion followed by immunoprecipitation, nuclei were isolated from 4 grams of 2-3 weeks old leaves and the procedure as described in (Lorkovic et al., 2017) was followed. Isolated nuclei were washed once in 1 ml of N buffer (15 mM Tris-HCl pH 7.5, 60 mM KCl, 15 mM NaCl, 5 mM MgCl₂, 1 mM CaCl₂, 250 mM sucrose, 1 mM DTT, 10 mM β-glycerophosphate) supplemented with protease inhibitors (Roche). After spinning for 5 min at 1,800 × g at 4°C nuclei were re-suspended in N buffer to a volume of 1 ml. Twenty microliters of MNase (0.1 u/μl) (SigmaAldrich) were added to each tube and incubated for 15 min at 37°C and during the incubation nuclei were mixed 4 times by inverting the tubes. MNase digestion was stopped on ice by addition of 110 μl of MNase stop solution (100 mM EDTA, 100 mM EGTA). Nuclei were lysed by addition of 110 μl of 5 M NaCl (final concentration of 500 mM NaCl). The suspension was mixed by inverting the tubes and they were then kept on ice for 15 min. Extracts were cleared by centrifugation for 10 min at 20,000 × g at 4°C. Supernatants were collected and centrifuged again as above. For each immunoprecipitation extract, an equivalent of 4 g of leaf material was used, usually in a volume of 1 ml. To control MNase digestion efficiency, 100 μl of the extract were kept for DNA extraction. Antibodies, including non-specific IgG from rabbit, were bound to protein A magnetic beads (GE Healthcare) and then incubated with MNase extracts overnight at 4°C. Beads were washed two times with N buffer without sucrose, containing 300 mM NaCl, followed by three washes with N buffer

containing 500 mM NaCl. Beads were re-suspended in 150 μ l of 1 $\times$ Laemmli loading buffer in 0.2 $\times$ PBS.

Proteins were resolved on 15% SDS-PAGE, transferred to a Protran nitrocellulose membrane (GE Healthcare) and analyzed by Western blotting using standard procedures. The blots were developed with an enhanced chemiluminescence kit (Thermo Fisher Scientific) and signals acquired by a ChemiDoc instrument (BioRad). All primary histone variant-specific and H3 marks-specific antibodies were used at 1 μ g/ml dilution. H3 specific antibody was used at 1:5,000 dilution. Rat anti-HA antibody (Roche 3F10) was used at 1:2,000 dilution. Secondary antibodies were goat anti-rabbit IgG (BioRad) and goat-anti rat IgG (SigmaAldrich), both at a 1:10,000 dilution.

Mass spectrometry

For mass spectrometry immunoprecipitated nucleosomes were resolved on 4–20% gradient gels (Serva) and silver-stained. Histone H3 bands were excised, reduced, alkylated, in-gel trypsin or LysC digested, and processed for MS. The nano HPLC system used was a Dionex UltiMate 3000 HPLC RSLC nano system (Thermo Fisher Scientific, Amsterdam, Netherlands) coupled to a Q Exactive mass spectrometer (Thermo Fisher Scientific, Bremen, Germany), equipped with a Proxeon nanospray source (Thermo Fisher Scientific, Odense, Denmark). Peptides were loaded onto a trap column (Thermo Fisher Scientific, Amsterdam, Netherlands, PepMap C18, 5 mm 3 300 mm ID, 5 mm particles, 100 Å pore size) at a flow rate of 25 ml/min using 0.1% TFA as the mobile phase. After 10 min, the trap column was switched in line with the analytical column (Thermo Fisher Scientific, Amsterdam, Netherlands, PepMap C18, 500mm 3 75 mm ID, 2 mm, 100 Å). Peptides were eluted using a flow rate of 230 nl/min and a binary 2-h gradient. The gradient starts with the mobile phases: 98% solution A (water/formic acid, 99.9/0.1, v/v) and 2% solution B (water/acetonitrile/formic acid, 19.92/80/0.08, v/v/v), increases to 35% of solution B over the next 120 min, followed by a gradient in 5 min to 90% of solution B, stays there for 5 min and decreases in 5 min back to the gradient 98% of solution A and 2% of solution B for equilibration at 30°C. The Q Exactive HF mass spectrometer was operated in data-dependent mode, using a full scan (m/z range 380–1500, nominal resolution of 60 000, target value 1E6) followed by MS/MS scans of the 10 most abundant ions. MS/MS spectra were acquired using a normalized collision energy of 27%, an isolation width of 1.4 m/z , and a resolution of 30,000, and the target value was set to 1E5. Precursor ions selected for fragmentation (exclude charge state 1, 7, 8, >8) were put on a dynamic exclusion list for 40 s. Additionally, the minimum AGC target was set to 5E3 and intensity threshold was calculated to be 4.8E4. The peptide match feature was set to preferred, and the exclude isotopes feature was enabled. For peptide identification, the RAW files were loaded into Proteome Discoverer (version 2.1.0.81, Thermo Scientific). The resulting MS/MS spectra were searched against *Arabidopsis thaliana* histone H3 sequences (7 sequences; 951 residues) using MS Amanda v2.1.5.8733 (Dorfer *et al.*, 2014). The following search parameters were used: Beta-methylthiolation on cysteine was set as a fixed modification, oxidation on methionine, deamidation of asparagine and glutamine, acetylation on lysine, phosphorylation on serine, threonine, and tyrosine, methylation and di-methylation on lysine and threonine, tri-methylation on lysine, and ubiquitinylation on lysine were set as variable modifications. The peptide mass tolerance was set to ± 5 ppm, and the fragment mass tolerance was set to ± 15 ppm. The maximal number of missed cleavages was set to 2. The result was filtered to 1% FDR at the peptide level using the Percolator algorithm integrated in Thermo Proteome Discoverer. The localization of the phosphorylation sites within the peptides was performed with the tool ptmRS, which is based on the tool phosphoRS (Taus *et al.*, 2011).

Peptides diagnostic for H3.1 and H3.3 covering positions K27 and K36 (see [Figure 1 figure supplement 1E](#)) were used for the analysis of modifications. All peptides covering these two

positions were selected in H3.1 and H3.3 immunoprecipitation samples and analyzed for the presence of modifications with threshold for modification probability set to 95% or higher. Relative modification levels were expressed as the number of modified peptides (**Figure 1figure supplement 1G**) divided by the total number of peptides (**Figure 1figure supplement 1F**) that were measured for each lysine position resulting in total modification levels for H3.1 and H3.3 (see **Figure 1D**). We also analyzed the same data by splitting H3.1 and H3.3 specific peptides in each immunoprecipitation and obtained highly similar trends for H3.1 and H3.3 irrespective of whether they were precipitated with H3.1 or H3.3 (**Figure 1E**). Histone acetylation was analyzed by selecting all peptides covering indicated positions and expressed as relative acetylation levels in each immunoprecipitation without differentiating H3.1 and H3.3 variants (**Figure 1figure supplement 1I and 1J**). We also analyzed H3K9, H3K14 and H3K18 acetylation from peptides derived from transgenic copy alone because these data reflect modification levels of each H3 variant and obtained highly similar levels (**Figure 1figure supplement 1K**) as without differentiation between H3.1 and H3.3.

Purification of recombinant *Arabidopsis* DDM1 and its truncations

His₆-tagged DDM1 and its truncations (DDM1(1-440), DDM1(1-196), DDM1(441-764)), GST-tagged DDM1 truncations, DDM1(1-24), DDM1(1-55), DDM1(1-74), DDM1(1-99), DDM1(1-123), DDM1(1-152), DDM1(1-172), DDM1(1-196), were expressed and purified as described previously (Osakabe et al., 2021).

Purification and reconstitution of recombinant *Arabidopsis* histone heterodimers

To purify and reconstitute *Arabidopsis* histone dimers, recombinant *Arabidopsis* histones H2A.13, H2A.X.3, H2A.Z.9, H2A.W.6, H2A.W.6 ΔCT (aa 1-128) and H2B.9 were expressed and purified as previously described (Osakabe et al., 2013; Tachiwana et al., 2010; Tanaka et al., 2004). The H2A-H2B heterodimers were reconstituted and purified as previously described (Osakabe et al., 2013).

Pull-down assay

The pull-down assay using Ni-NTA and GS4B beads for His₆-tagged DDM1 or truncations and GST-tagged DDM1 truncations were performed as described previously (Osakabe et al., 2021). Proteins precipitated with beads were analyzed by 15% SDS-PAGE stained with Coomassie Brilliant Blue.

Plant material for ChIP-seq

Col-0 wild type (WT) *Arabidopsis thaliana* seeds were stratified at 4°C in the dark for three days. Seeds were germinated on ½ MS sterilized plates in the growth chamber under long day (LD) conditions (21°C; 16 h light/8 h dark). After 10 days seedling tissue was freshly harvested. For ChIP-seq from leaf tissue, Col-0 wild type (WT) and seeds from *ddm1*^{+/+} plants were stratified at 4°C in the dark for three days. To prevent carry-over mutations from generations of *ddm1* homozygous mutants we used leaves from *ddm1* homozygous mutant plants segregated from heterozygous *ddm1*^{+/+} mutants. Plants were grown on soil in the growth chamber under long day (LD) conditions (21°C; 16 h light/8 h dark) and leaf tissue was harvested 5 weeks after germination.

Chromatin immunoprecipitation (ChIP)

For ChIP nuclei isolation was performed using 10-day-old seedlings (WT Col-0) or leaves (WT Col-0, *ddm1*). The procedure for nuclei isolation and chromatin immunoprecipitation was performed as described in (Osakabe et al., 2021). Freshly harvested tissues (0.3 g of tissue was used for each immunoprecipitation) were fixed with 1% formaldehyde for 15 min and the cross-linking reaction was stopped by the addition of 125 mM glycine. Crosslinked tissues were frozen and ground with a mortar and pestle in liquid nitrogen to obtain a fine powder. Ground tissues were resuspended in M1 buffer (10 mM sodium phosphate pH 7.0, 100 mM NaCl, 1 M hexylene glycol, 10 mM 2-mercaptoethanol, and protease inhibitor (Roche)), and the suspension was filtered through miracloth. Nuclei were precipitated by centrifugation and washed six times with M2 buffer (10 mM sodium phosphate pH 7.0, 100 mM NaCl, 1 M hexylene glycol, 10 mM MgCl₂, 0.5% Triton X-100, 10 mM 2-mercaptoethanol, and protease inhibitor), and then further washed once with M3 buffer (10 mM sodium phosphate pH 7.0, 100 mM NaCl, 10 mM 2-mercaptoethanol, and protease inhibitor). Nuclei pellets were rinsed and resuspended in a sonication buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 0.1% SDS, and protease inhibitor) and sonicated with a Covaris E220 High Performance Focused Ultrasonicator for 15 min at 4°C (Duty factor 5.0; PIP 140.0; Cycles per Burst 200) in 1 ml Covaris milliTUBE. After chromatin shearing, the debris were removed by centrifugation and the solutions containing chromatin fragments were diluted with three times the volume of ChIP dilution buffer (16.7 mM Tris-HCl pH 8.0, 167 mM NaCl, 1.2 mM EDTA, 1.1% Triton X-100, 0.01% SDS, and protease inhibitor). After dilution, protein A/G magnetic beads (50 µl for one gram of tissue; Thermo Fisher Scientific) were added to sheared chromatin and incubated at 4°C for 1 hour with rotation. Pre-cleared samples were collected and incubated with 5 µg of in house prepared anti-H2A.X.3/5, anti-H2A.13, anti-H2A.2, anti-H2A.Z.9, anti-H2A.Z.11, anti-H2A.W.6, anti-H2A.W.7, anti-H1, and anti-H3 (Abcam, ab1791), anti-H3K36me3 (Abcam, ab9050), anti-H3K27me3 (Millipore, 07-449), anti-H3K4me3 (Abcam, ab8580), anti-H3K4me1 (Abcam, ab8895), anti-H3K27me1 (Millipore 17-643), anti-H4K20me1 (Abcam, ab9051), anti-H3K9me1 (Abcam, ab8896) or anti-H3K9me2 (Abcam, ab1220) antibodies at 4°C overnight with rotation. After incubation, samples were mixed with 30 µl of protein A/G magnetic beads, incubated at 4°C for 3 hours with rotation, washed twice with low salt buffer (20 mM Tris-HCl pH 8.0, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, and 0.1% SDS), once with high salt buffer (20 mM Tris-HCl pH 8.0, 500 mM NaCl, 2 mM EDTA, 1% Triton X-100, and 0.1% SDS), once with LiCl buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 0.25 M LiCl, 1% IGEPAL CA-630, and 0.1% deoxycholic acid), and twice with TE buffer (10 mM Tris-HCl pH 8.0 and 1 mM EDTA). Immunoprecipitated DNA was eluted by adding 500 µl of elution buffer (1% SDS and 0.1 M NaHCO₃), incubated at 65°C for 15 min, and mixed with 51 µl of reverse crosslink buffer (40 mM Tris-HCl pH 8.0, 0.2 M NaCl, 10 mM EDTA, 0.04 mg/ml proteinase K (Thermo Fisher Scientific)). The reaction mixture was then incubated at 45°C for 3 hours and subsequently at 65°C for 16 hours. After crosslink reversal, DNA was treated with 10 µg of RNaseA (Thermo Fisher Scientific), incubated at room temperature for 30 min, and purified with a MinElute PCR purification kit (Qiagen).

ChIP-seq library prep and data processing

For ChIP-seq, libraries were prepared either with Nugen Ovation Ultralow V2 DNA-Seq library prep kit (NuGen) or NEBNext Ultra II DNA library prep kit for Illumina (New England Biolabs) following the manufacturer's instructions. The libraries were sequenced with an Illumina HiSeq 2000 to generate single-end 50 bp reads. For alignment and quality check of sequenced samples, bedtools v2.27.1 (Quinlan, 2014) was used to convert the raw BAM files to fastq. FastQC v0.11.8 (<https://qubeshub.org/resources/fastqc>) was used to generate quality reports for all sequencing data. Reads were trimmed using trim_galore v0.6.5

(DOI:10.5281/zenodo.5127898.) (*trim_galore* —*dont_gzip* —*stringency 1* —*fastqc* —*length 5* \$) and then aligned to the TAIR10 *Arabidopsis* genome using Bowtie2 v2.4.1 (Langmead and Salzberg, 2012) with default settings, Picard v2.22.8 (broadinstitute.github.io/picard/) was used to remove duplicated reads. Deeptools v3.1.2 (Ramirez *et al.*, 2016) was used to examine correlations between the ChIP samples. The bamCompare function from Deeptools was used to normalize ChIP samples to Input or H3 and to generate log2 ratio (ChIP/ (Input or H3)) bigwig files.

ChIP-seq data processing for published data

Publicly available ChIP-seq data for H2B variants was downloaded from GEO GSE151166 (Jiang *et al.*, 2020). ChIP-seq data for H3 variants was downloaded from GEO GSE34840 (Stroud *et al.*, 2012). ChIP-seq data for H3K14Ac, H3K9Ac was downloaded from GEO GSE89768 (Kim *et al.*, 2016). ChIP-seq data for H2AK121Ub and H3K27me3 was downloaded from GEO GSE89357 (Zhou *et al.*, 2017). Raw data was downloaded and processed as described above. Quality control metrics for all samples are included in source file 1.

Defining chromatin states

Aligned ChIP-seq BAM files for chromatin features were generated as described above. Aligned BAM files were then converted to BED format using bedtools bamtobed v2.27.1 (Quinlan, 2014). These genome wide BED files were then used for learning chromatin states by the multivariate Hidden Markov Models. For the extensive wild type model, we used the BinarizeBed and LearnModel programs from ChromHMM (Ernst and Kellis, 2017) with default settings and window size of 200 bp. We generated models from 2 to 50 chromatin states. To compare emission parameters of models with different numbers of states, the command CompareModel from ChromHMM was used. Thus, we selected a reference model with the largest number of states (50) and compared it with other models with lower number states (from 49 to 2). This resulted in a correlation matrix between each state of the reference model and any state of the other models being compared (Figure 2 figure supplement 2). This comparison revealed that the correlation dropped for models including less than 13 states, suggesting that some of the biologically relevant states were not resolved. We thus concluded that at least 13 states should be used. However, ChromHMM (Ernst & Kellis, 2017) does not establish the optimal state number as it does not consider genomic features which are associated with specific biological function. Therefore, we analyzed the association between chromatin states with genomic features in all models ranging from 13 to 30 states. As a result, for example, we observed that models with more than 13 states could resolve biologically meaningful heterochromatic states (H2 to H6) (H2 to H4 associated with pericentromeric repeats/CMT2 associated repeats; H5 and H6 associated with chromosome arm repeats/RdDM repeats) (as shown in Jamge *et al.*, 2022, Figure 3). Further, only with a 26-state model we observed a clearly defined novel state H1 with H2B enrichment, largely deprived of TE genes and located in the closest proximity to centromeres. Increasing the number of states to 27 and above gave rise to an additional chromatin state that could no longer be associated with any distinct genomic feature. Hence, we decided that, with the set of chromatin components included, a 26-state model is optimal for our analysis.

For the mixed seedling and leaf data, and for the mixed wild type and *ddm1* mutant data we used concatenated ChromHMM models (Ernst and Kellis, 2017). Those models use the data from both tissues/genotypes to build a common model. To this end we used the BinarizeBed and LearnModel with the default settings and window size of 200 bp but with the tissue or genotype as additional information. The number of states in the final model was decided on as described for the extensive model.

Analysis of sub-epigenome models

The ChromHMM command EvalSubset was used to compare models where histone variants or modifications were excluded from the model. Five models were generated and evaluated against the full model: no H2B variants, no H3 variants, no H2A variants, no histone modifications, and no variants (H2A/H3/H2B). The diagonals of the resulting confusion matrices (representing the Jaccard indices) were extracted and visualized using the R package ComplexHeatmap v2.10.0 (Gu et al., 2016).

Analysis of chromatin states

The emission matrices from the ChromHMM models were read into R 4.1.2 (<https://www.R-project.org/>) and plotted using the R packages ggplot2 from tidyverse (doi:10.21105/joss.01686) and package ComplexHeatmap v2.10.0 (Gu et al., 2016). The state assignment files from ChromHMM for the different models and tissues/genotypes were read in and analyzed using packages from tidyverse and valr v0.6.6 (Riemondy et al., 2017). To compare the wild type and *ddm1* assignments from the concatenated model, separate files were joined. To this end, for each (200bp) bin of the genome, the information about which state had been assigned in wild type and in *ddm1* respectively was combined. The state assignment datasets were then overlapped, using the function bed_intersect from valr, with regions for genomic features defined in the file Araport11_GFF3_genes_transposons.Jun2016.gff downloaded from (https://www.arabidopsis.org/download_files/Genes/Araport11_genome_release/archived/Araport11_GFF3_genes_transposons.Jun2016.gff.gz). The transcriptome, methylome and DNase I datasets were integrated to generate plots using R and the ggplot2 package from tidyverse. All box plots show the data excluding outliers.

Comparison of states from extensive and concatenated models

To compare the states from the two tissue model with the chromatin types defined in the extensive model, the states assigned to seedling by the two tissue model were overlapped with the chromatin types from the extensive model. For the comparison of the wild type and *ddm1* mutant model states with the chromatin types from the extensive model, the states assigned to the wildtype were used. The genomic overlaps were calculated using bed_intersect from valr v0.6.6 (Riemondy et al., 2017). States overlapping one chromatin type with >66% and all others with < 20% were assigned the largest overlapping chromatin type. In other cases, the state was classified as mixed (Figure 3 figure supplement 1B and 3C).

Analysis of chromatin state changes in *ddm1* mutant

The Jaccard index for each state was calculated as the total length of regions assigned to that state in both wild type and mutant divided by the combined length of all regions belonging to that state in at least one of the two genotypes. The overlap was calculated as the total length of regions assigned to that state in both wild type and mutant divided by the total length of regions assigned to that state in wild type. The size fold change was calculated as the total length of regions assigned to that state in *ddm1* mutant divided by the total length of all regions assigned to the state in the wild type.

Or, formally, if $B_{w,m}$ is the total number of bins that are assigned to state w in the wild type and state m in *ddm1*, then the (JI) Jaccard index, (O) overlap and (FC) size change for states can be calculated as:

$$JI_s = \frac{B_{s,s}}{B_{s,\cdot} + B_{\cdot,s} - B_{s,s}}, O_s = \frac{B_{s,s}}{B_{s,\cdot}}, FC_s = \log_2 \left(\frac{B_{\cdot,s}}{B_{s,\cdot}} \right)$$

The plots were generated using R and the ggplot2 package from tidyverse. The metaplots were generated using Deeptools v3.1.2 (Ramirez et al., 2016). Using the bamCompare function of Deeptools v3.1.2 bigWig files were generated by normalizing ChIP samples with input/H3. These bigWig files were used for plotting the metaplot heatmap (Figure 4A) and profile plots (Figure 4 figure supplement 1A and 1B) using deeptools v3.1.2 plotHeatmap and plotProfile functions respectively.

RNA-seq of WT seedlings

Total RNA was extracted with Spectrum plant total RNA kit (Sigma Aldrich) from 10-day seedlings of WT Col-0. A DNA-free DNA removal kit was used to remove contaminated DNA (Thermo Fisher Scientific). RNA-seq poly-A libraries were generated with NEBNext UltraII directional RNA library prep kit for Illumina (New England Biolabs) following the manufacturer's instructions. The libraries were sequenced with Illumina HiSeq 2500 to generate single-end 50 bp reads. Samples were prepared from three independent biological replicates. The RNA-seq data was processed as described in (Osakabe et al., 2021).

Methylation data

BS-seq data for WT Col-0 was downloaded from GSE39901 (Stroud et al., 2013). BS-seq reads were processed using the nf-core pipeline (github.com/nf-core/methylseq) as described in (doi: <https://doi.org/10.1101/2022.12.04.519028>). Cutadapt v2.10 doi: <https://doi.org/10.14806/ej.17.1.200> was used to trim the adaptors (default parameters). Trimmed reads were then mapped to TAIR10 (Col-0) assembly using bismark v0.2.1 (Krueger and Andrews, 2011) allowing mismatches to 0.5. Methylation calls were performed using methylpy v1.3.7 (Schultz et al., 2015) on the aligned bam files. Methylation call bed files were used to calculate average methylation over chromatin state bed regions.

DNase I-seq

We downloaded processed DNaseI – seq bigwig files data from GEO series GSE53322 for WT Col-0 (GSM1289358) (Sullivan et al., 2014). Bedtools map v2.3 (Quinlan, 2014) was used to calculate the averaged signal over bed regions in each chromatin state. Box plots were generated in R v3.5 using ggplot2 to compare the accessibility across chromatin states.

RNA-seq data analysis of wild type and ddm1 mutant leaves

We used the wild type and *ddm1* mutant RNA-seq data from GSE150435. The data was processed as described in (Osakabe et al., 2021). Before grouping the TE genes into five groups we first excluded all TE genes showing any expression in wild type (tpm>0). TE genes without expression (tpm<0.1) in *ddm1* mutant were put in the “no expression” group. The remaining TE genes were divided into 4 quartiles based on the tpm values in the *ddm1* mutant.

Data availability

The genome-wide sequencing (ChIP-seq) generated for this study as well as published datasets (ChIP-seq, RNA-seq) utilized to support the findings in this study have been deposited on NCBI's Gene Expression Omnibus (GEO) with the accession number of subseries GSE226469, GSE231398, GSE150434 and GSE150433. Super series associated with this study and published data are GSE231408 and GSE150436. Source data for all the main figures as well as supplementary figures have been provided with this manuscript. All other data supporting the conclusions of the study will be available from the corresponding author upon reasonable request.

Code availability

All the code used to analyze the genome-wide sequencing data presented in this study, as described in the Methods are available upon request. See also https://github.com/Gregor-Mendel-Institute/jamge_states_2023

Acknowledgements

FB acknowledges support from the next generation sequencing and PlantS facilities at the Vienna BioCenter Core Facilities (VBCF), the BioOptics facility, the Proteomics facility, and Molecular Biology Services from the Institute for Molecular Pathology (IMP), the Molecular Biology Services at the GMI. We thank all members of the Berger laboratory for their technical help. We thank J. Matthew Watson for insightful discussions and critical reading of the manuscript. This work was supported by the Japan Society for the Promotion of Science (JSPS) Overseas Research Fellowships [to A.O.], the Austrian Science Fund (FWF): M2539-B21 [to A.O.], P32054, P30802, and P33380 [to F.B.], and DK1238 [to B.J.], the Austrian Academy of Sciences [to F.B., Z.J.L, E.A., S.A., R.P. and R.Y.], JSPS Grant-in-Aid for Research Activity Start-up (JP21K20628), JSPS Grant-in-Aid for Transformative Research Areas (JP22H05172 and JP22H05178), and the Japan Science and Technology Agency (JST) PRESTO (JPMJPR20K3) [to A.O.].

Author contributions

Z.J.L. performed all the biochemical analysis of the composition of mononucleosomes. A.O. and A.L.K. performed biochemical experiments on DDM1 binding to H2A variants. S.A. prepared biological material. B.J., A.O. and Z.J.L. performed the ChIPseq and RNAseq. R.Y. contributed the initial set of ChIP-seq data. B.J., E.A. and V.S. performed the genome-wide analysis. F.B. conceived, designed, and supervised the project. The manuscript was drafted by F.B., E.A. and Z.J.L., with comments from all co-authors.

Competing interests

All authors discussed the results and have no conflict of interest.

References

Aslam M. , Fakher B. , Jakada B.H. , Cao S. , Qin Y (2019) **SWR1 Chromatin Remodeling Complex: A Key Transcriptional Regulator in Plants** *Cells* **8**

Berger F. , Muegge K. , Richards E.J (2023) **Seminars in cell and development biology on histone variants remodelers of H2A variants associated with heterochromatin** *Semin Cell Dev Biol* **135**:93–101

Bieluszewski T. , Sura W. , Dziegielewski W. , Bieluszewska A. , Lachance C. , Kabza M. , Szymanska-Lejman M. , Abram M. , Wlodzimierz P. , De Winne N. , et al. (2022) **NuA4 and H2A.Z control environmental responses and autotrophic growth in Arabidopsis** *Nat Commun* **13**

Borg M. , Jacob Y. , Susaki D. , LeBlanc C. , Buendia D. , Axelsson E. , Kawashima T. , Voigt P. , Boavida L. , Becker J. , et al. (2020) **Targeted reprogramming of H3K27me3 resets epigenetic memory in plant paternal chromatin** *Nat Cell Biol* **22**:621–629

Borg M. , Jiang D. , Berger F (2021) **Histone variants take center stage in shaping the epigenome** *Curr Opin Plant Biol* **61**

Borg M. , Papareddy R.K. , Dombey R. , Axelsson E. , Nodine M.D. , Twell D. , Berger F (2021) **Epigenetic reprogramming rewires transcription during the alternation of generations in Arabidopsis** *Elife* **10**

Bourguet P. , Picard C.L. , Yelagandula R. , Pelissier T. , Lorkovic Z.J. , Feng S. , Pouch-Pelissier M.N. , Schmucker A. , Jacobsen S.E. , Berger F. , et al. (2021) **The histone variant H2A.W and linker histone H1 co-regulate heterochromatin accessibility and DNA methylation** *Nat Commun* **12**

Carter B. , Bishop B. , Ho K.K. , Huang R. , Jia W. , Zhang H. , Pascuzzi P.E. , Deal R.B. , Ogas J (2018) **The Chromatin Remodelers PKL and PIE1 Act in an Epigenetic Pathway That Determines H3K27me3 Homeostasis in Arabidopsis** *Plant Cell* **30**:1337–1352

Chakravarthy S. , Bao Y. , Roberts V.A. , Tremethick D. , Luger K (2004) **Structural characterization of histone H2A variants** *Cold Spring Harb Symp Quant Biol* **69**:227–234

Delaney K. , Almouzni G. (2023) **Transcription-coupled H3.3 recycling: A link with chromatin states** *Semin Cell Dev Biol* **135**:13–23

Ernst J. , Kellis M (2012) **ChromHMM: automating chromatin-state discovery and characterization** *Nat Methods* **9**:215–216

Ernst J. , Kellis M (2017) **Chromatin-state discovery and genome annotation with ChromHMM** *Nat Protoc* **12**:2478–2492

Feng J. , Shen W.H (2014) **Dynamic regulation and function of histone monoubiquitination in plants** *Front Plant Sci* **5**

Flury V. , Reveron-Gomez N. , Alcaraz N. , Stewart-Morgan K.R. , Wenger A. , Klose R.J. , Groth A (2023) **Recycling of modified H2A-H2B provides short-term memory of chromatin states** *Cell* **186**:1050–1065

Gan E.S. , Xu Y. , Ito T (2015) **Dynamics of H3K27me3 methylation and demethylation in plant development** *Plant Signal Behav* **10**

Gomez-Zambrano A. , Merini W. , Calonje M (2019) **The repressive role of Arabidopsis H2A.Z in transcriptional regulation depends on AtBMI1 activity** *Nat Commun* **10**

Gu Z. , Eils R. , Schlesner M (2016) **Complex heatmaps reveal patterns and correlations in multidimensional genomic data** *Bioinformatics* **32**:2847–2849

Hake S.B. , Allis C.D (2006) **Histone H3 variants and their potential role in indexing mammalian genomes: the “H3 barcode hypothesis”** *Proc Natl Acad Sci U S A* **103**:6428–6435

Henikoff S. , McKittrick E. , Ahmad K. (2004) **Epigenetics, histone H3 variants, and the inheritance of chromatin states** *Cold Spring Harb Symp Quant Biol* **69**:235–243

Horikoshi N. , Sato K. , Shimada K. , Arimura Y. , Osakabe A. , Tachiwana H. , Hayashi-Takanaka Y. , Iwasaki W. , Kagawa W. , Harata M. , et al. (2013) **Structural polymorphism in the L1 loop regions of human H2A.Z.1 and H2A.Z.2** *Acta Crystallogr D Biol Crystallogr* **69**:2431–2439

Jacob Y. , Bergamin E. , Donoghue M.T. , Mongeon V. , LeBlanc C. , Voigt P. , Underwood C.J. , Brunzelle J.S. , Michaels S.D. , Reinberg D. , et al. (2014) **Selective methylation of histone H3 variant H3.1 regulates heterochromatin replication** *Science* **343**:1249–1253

Jeddeloh J.A. , Bender J. , Richards E.J (1998) **The DNA methylation locus DDM1 is required for maintenance of gene silencing in Arabidopsis** *Genes Dev* **12**:1714–1725

Jiang D. , Berger F (2017) **DNA replication-coupled histone modification maintains Polycomb gene silencing in plants** *Science* **357**:1146–1149

Jiang D. , Borg M. , Lorkovic Z.J. , Montgomery S.A. , Osakabe A. , Yelagandula R. , Axelsson E. , Berger F (2020) **The evolution and functional divergence of the histone H2B family in plants** *PLoS Genet* **16**

Johnson L. , Mollah S. , Garcia B.A. , Muratore T.L. , Shabanowitz J. , Hunt D.F. , Jacobsen S.E (2004) **Mass spectrometry analysis of Arabidopsis histone H3 reveals distinct combinations of post-translational modifications** *Nucleic Acids Res* **32**:6511–6518

Kakutani T. , Munakata K. , Richards E.J. , Hirochika H (1999) **Meiotically and mitotically stable inheritance of DNA hypomethylation induced by ddm1 mutation of Arabidopsis thaliana** *Genetics* **151**:831–838

Kharchenko P.V. , Alekseyenko A.A. , Schwartz Y.B. , Minoda A. , Riddle N.C. , Ernst J. , Sabo P.J. , Larschan E. , Gorchakov A.A. , Gu T. , et al. (2011) **Comprehensive analysis of the chromatin landscape in *Drosophila melanogaster*** *Nature* **471**:480–485

Kim J. , Guermah M. , McGinty R.K. , Lee J.S. , Tang Z. , Milne T.A. , Shilatifard A. , Muir T.W. , Roeder R.G (2009) **RAD6-Mediated transcription-coupled H2B ubiquitylation directly stimulates H3K4 methylation in human cells** *Cell* **137**:459–471

Kim Y.J. , Wang R. , Gao L. , Li D. , Xu C. , Mang H. , Jeon J. , Chen X. , Zhong X. , Kwak J.M. , et al. (2016) **POWERDRESS and HDA9 interact and promote histone H3 deacetylation at specific genomic sites in *Arabidopsis*** *Proc Natl Acad Sci U S A* **113**:14858–14863

Koyama M. , Kurumizaka H (2018) **Structural diversity of the nucleosome** *J Biochem* **163**:85–95

Krueger F. , Andrews S.R (2011) **Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications** *Bioinformatics* **27**:1571–1572

Langmead B. , Salzberg S.L (2012) **Fast gapped-read alignment with Bowtie 2** *Nat Methods* **9**:357–359

Lei B. , Capella M. , Montgomery S.A. , Borg M. , Osakabe A. , Goiser M. , Muhammad A. , Braun S. , Berger F (2021) **A Synthetic Approach to Reconstruct the Evolutionary and Functional Innovations of the Plant Histone Variant H2A.W** *Curr Biol* **31**:182–191

Leng X. , Thomas Q. , Rasmussen S.H. , Marquardt S (2020) **A G(enomic)P(ositioning)S(ystem) for Plant RNAPII Transcription** *Trends Plant Sci* **25**:744–764

Liu C. , Lu F. , Cui X. , Cao X (2010) **Histone methylation in higher plants** *Annu Rev Plant Biol* **61**:395–420

Liu Y. , Tian T. , Zhang K. , You Q. , Yan H. , Zhao N. , Yi X. , Xu W. , Su Z (2018) **PCSD: a plant chromatin state database** *Nucleic Acids Res* **46**:D1157–D1167

Long J. , Carter B. , Johnson E.T. , Ogas J (2023) **Contribution of the histone variant H2A.Z to expression of responsive genes in plants** *Semin Cell Dev Biol* **135**:85–92

Loppin B. , Berger F (2020) **Histone Variants: The Nexus of Developmental Decisions and Epigenetic Memory** *Annu Rev Genet* **54**:121–149

Lorkovic Z.J. , Park C. , Goiser M. , Jiang D. , Kurzbauer M.T. , Schlogelhofer P. , Berger F (2017) **Compartmentalization of DNA Damage Response between Heterochromatin and Euchromatin Is Mediated by Distinct H2A Histone Variants** *Curr Biol* **27**:1192–1199

Luger K. , Mader A.W. , Richmond R.K. , Sargent D.F. , Richmond T.J (1997) **Crystal structure of the nucleosome core particle at 2.8 Å resolution** *Nature* **389**:251–260

Minsky N. , Shema E. , Field Y. , Schuster M. , Segal E. , Oren M (2008) **Monoubiquitinated H2B is associated with the transcribed region of highly expressed genes in human cells** *Nat Cell Biol* **10**:483–488

Ni K. , Muegge K (2021) **LSH catalyzes ATP-driven exchange of histone variants macroH2A1 and macroH2A2** *Nucleic Acids Res* **49**:8024–8036

Ni K. , Ren J. , Xu X. , He Y. , Finney R. , Braun S.M.G. , Hathaway N.A. , Crabtree G.R. , Muegge K (2020) **LSH mediates gene repression through macroH2A deposition** *Nat Commun* **11**

Nie X. , Wang H. , Li J. , Holec S. , Berger F (2014) **The HIRA complex that deposits the histone H3.3 is conserved in Arabidopsis and facilitates transcriptional dynamics** *Biol Open* **3**:794–802

Osakabe A. , Jamge B. , Axelsson E. , Montgomery S.A. , Akimcheva S. , Kuehn A.L. , Pisupati R. , Lorkovic Z.J. , Yelagandula R. , Kakutani T. , et al. (2021) **The chromatin remodeler DDM1 prevents transposon mobility through deposition of histone variant H2A.W** *Nat Cell Biol* **23**:391–400

Osakabe A. , Lorkovic Z.J. , Kobayashi W. , Tachiwana H. , Yelagandula R. , Kurumizaka H. , Berger F (2018) **Histone H2A variants confer specific properties to nucleosomes and impact on chromatin accessibility** *Nucleic Acids Res* **46**:7675–7685

Osakabe A. , Molaro A (2023) **Histone renegades: Unusual H2A histone variants in plants and animals** *Semin Cell Dev Biol* **135**:35–42

Osakabe A. , Tachiwana H. , Takaku M. , Hori T. , Obuse C. , Kimura H. , Fukagawa T. , Kurumizaka H (2013) **Vertebrate Spt2 is a novel nucleolar histone chaperone that assists in ribosomal DNA transcription** *J Cell Sci* **126**:1323–1332

Probst A.V (2022) **Deposition and eviction of histone variants define functional chromatin states in plants** *Curr Opin Plant Biol* **69**

Quinlan A.R (2014) **BEDTools: The Swiss-Army Tool for Genome Feature Analysis** *Curr Protoc Bioinformatics* **47**

Raman P. , Rominger M.C. , Young J.M. , Molaro A. , Tsukiyama T. , Malik H.S (2022) **Novel Classes and Evolutionary Turnover of Histone H2B Variants in the Mammalian Germline** *Mol Biol Evol* **39**

Ramirez F. , Ryan D.P. , Gruning B. , Bhardwaj V. , Kilpert F. , Richter A.S. , Heyne S. , Dundar F. , Manke T (2016) **deepTools2: a next generation web server for deep-sequencing data analysis** *Nucleic Acids Res* **44**:W160–165

Ranjan A. , Nguyen V.Q. , Liu S. , Wisniewski J. , Kim J.M. , Tang X. , Mizuguchi G. , Elalaoui E. , Nickels T.J. , Jou V. , et al. (2020) **Live-cell single particle imaging reveals the role of RNA polymerase II in histone H2A.Z eviction** *Elife* **9**

Reveron-Gomez N. , Gonzalez-Aguilera C. , Stewart-Morgan K.R. , Petryk N. , Flury V. , Graziano S. , Johansen J.V. , Jakobsen J.S. , Alabert C. , Groth A (2018) **Accurate Recycling of Parental Histones Reproduces the Histone Modification Landscape during DNA Replication** *Mol Cell* **72**:239–249

Riemondy K.A. , Sheridan R.M. , Gillen A. , Yu Y. , Bennett C.G. , Hesselberth J.R (2017) **valr: Reproducible genome interval analysis in R** *F1000Res* **6**

Roudier F. , Ahmed I. , Berard C. , Sarazin A. , Mary-Huard T. , Cortijo S. , Bouyer D. , Caillieux E. , Duvernois-Berthet E. , Al-Shikhley L. , et al. (2011) **Integrative epigenomic mapping defines four main chromatin states in Arabidopsis** *EMBO J* **30**:1928–1938

Rougee M. , Quadrana L. , Zervudacki J. , Hure V. , Colot V. , Navarro L. , Deleris A (2021) **Polycomb mutant partially suppresses DNA hypomethylation-associated phenotypes in Arabidopsis** *Life Sci Alliance* **4**

Rudnizky S. , Bavly A. , Malik O. , Pnueli L. , Melamed P. , Kaplan A (2016) **H2A.Z controls the stability and mobility of nucleosomes to regulate expression of the LH genes** *Nat Commun* **7**

Schmucker A. , Lei B. , Lorkovic Z.J. , Capella M. , Braun S. , Bourguet P. , Mathieu O. , Mechtler K. , Berger F (2021) **Crosstalk between H2A variant-specific modifications impacts vital cell functions** *PLoS Genet* **17**

Schultz M.D. , He Y. , Whitaker J.W. , Hariharan M. , Mukamel E.A. , Leung D. , Rajagopal N. , Nery J.R. , Urich M.A. , Chen H. , et al. (2015) **Human body epigenome maps reveal noncanonical DNA methylation variation** *Nature* **523**:212–216

Schwammle V. , Sidoli S. , Ruminowicz C. , Wu X. , Lee C.F. , Helin K. , Jensen O.N (2016) **Systems Level Analysis of Histone H3 Post-translational Modifications (PTMs) Reveals Features of PTM Crosstalk in Chromatin Regulation** *Mol Cell Proteomics* **15**:2715–2729

Sequeira-Mendes J. , Aragues I. , Peiro R. , Mendez-Giraldez R. , Zhang X. , Jacobsen S.E. , Bastolla U. , Gutierrez C (2014) **The Functional Topography of the Arabidopsis Genome Is Organized in a Reduced Number of Linear Motifs of Chromatin States** *Plant Cell* **26**:2351–2366

Stroud H. , Greenberg M.V. , Feng S. , Bernatavichute Y.V. , Jacobsen S.E (2013) **Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis methylome** *Cell* **152**:352–364

Stroud H. , Otero S. , Desvoyes B. , Ramirez-Parra E. , Jacobsen S.E. , Gutierrez C (2012) **Genome-wide analysis of histone H3.1 and H3.3 variants in Arabidopsis thaliana** *Proc Natl Acad Sci U S A* **109**:5370–5375

Subramanian V. , Fields P.A. , Boyer L.A (2015) **H2A Z: a molecular rheostat for transcriptional control.** *F*

Sullivan A.M. , Arsovski A.A. , Lempe J. , Bubbs K.L. , Weirauch M.T. , Sabo P.J. , Sandstrom R. , Thurman R.E. , Neph S. , Reynolds A.P. , et al. (2014) **Mapping and dynamics of regulatory DNA and transcription factor networks in *A. thaliana*** *Cell Rep* **8**:2015–2030

Sun Z. , Bernstein E (2019) **Histone variant macroH2A: from chromatin deposition to molecular function** *Essays Biochem* **63**:59–74

Surface L.E. , Fields P.A. , Subramanian V. , Behmer R. , Udeshi N. , Peach S.E. , Carr S.A. , Jaffe J.D. , Boyer L.A (2016) **H2A.Z.1 Monoubiquitylation Antagonizes BRD2 to Maintain Poised Chromatin in ESCs** *Cell Rep* **14**:1142–1155

Tachiwana H. , Kagawa W. , Kurumizaka H (2012) **Comparison between the CENP-A and histone H3 structures in nucleosomes** *Nucleus* **3**:6–11

Tachiwana H. , Kagawa W. , Osakabe A. , Kawaguchi K. , Shiga T. , Hayashi-Takanaka Y. , Kimura H. , Kurumizaka H (2010) **Structural basis of instability of the nucleosome containing a testis-specific histone variant, human H3T** *Proc Natl Acad Sci U S A* **107**:10454–10459

Talbert P.B. , Henikoff S (2021) **Histone variants at a glance** *J Cell Sci* **134**

Tanaka Y. , Tawaramoto-Sasanuma M. , Kawaguchi S. , Ohta T. , Yoda K. , Kurumizaka H. , Yokoyama S (2004) **Expression and purification of recombinant human histones** *Methods* **33**:3–11

Taus T. , Kocher T. , Pichler P. , Paschke C. , Schmidt A. , Henrich C. , Mechtler K (2011) **Universal and confident phosphorylation site localization using phosphoRS** *J Proteome Res* **10**:5354–5362

Tessarz P. , Kouzarides T (2014) **Histone core modifications regulating nucleosome structure and dynamics** *Nat Rev Mol Cell Biol* **15**:703–708

Wang Y. , Long H. , Yu J. , Dong L. , Wassef M. , Zhuo B. , Li X. , Zhao J. , Wang M. , Liu C. , et al. (2018) **Histone variants H2A.Z and H3.3 coordinately regulate PRC2-dependent H3K27me3 deposition and gene expression regulation in mES cells** *BMC Biol* **16**

Weber C.M. , Henikoff S (2014) **Histone variants: dynamic punctuation in transcription** *Genes Dev* **28**:672–682

Willige B.C. , Zander M. , Yoo C.Y. , Phan A. , Garza R.M. , Wanamaker S.A. , He Y. , Nery J.R. , Chen H. , Chen M. , et al. (2023) **Author Correction: PHYTOCHROME-INTERACTING FACTORS trigger environmentally responsive chromatin dynamics in plants** *Nat Genet* **55**

Wollmann H. , Holec S. , Alden K. , Clarke N.D. , Jacques P.E. , Berger F (2012) **Dynamic deposition of histone variant H3.3 accompanies developmental remodeling of the *Arabidopsis* transcriptome** *PLoS Genet* **8**

Wollmann H. , Stroud H. , Yelagandula R. , Tarutani Y. , Jiang D. , Jing L. , Jamge B. , Takeuchi H. , Holec S. , Nie X. , et al. (2017) **The histone H3 variant H3.3 regulates gene body DNA methylation in *Arabidopsis thaliana*** *Genome Biol* **18**

Yang Y. , Zhang L. , Xiong C. , Chen J. , Wang L. , Wen Z. , Yu J. , Chen P. , Xu Y. , Jin J. , et al. (2022) **HIRA complex presets transcriptional potential through coordinating depositions of the histone variants H3.3 and H2A.Z on the poised genes in mESCs** *Nucleic Acids Res* **50**:191–206

Yelagandula R. , Stroud H. , Holec S. , Zhou K. , Feng S. , Zhong X. , Muthurajan U.M. , Nie X. , Kawashima T. , Groth M. , et al. (2014) **The histone variant H2A.W defines heterochromatin and promotes chromatin condensation in *Arabidopsis*** *Cell* **158**:98–109

Zhang K. , Sridhar V.V. , Zhu J. , Kapoor A. , Zhu J.K (2007) **Distinctive core histone post-translational modification patterns in *Arabidopsis thaliana*** *PLoS One* **2**

Zhao F. , Xue M. , Zhang H. , Li H. , Zhao T. , Jiang D (2023) **Coordinated histone variant H2A.Z eviction and H3.3 deposition control plant thermomorphogenesis**

Zhou Y. , Romero-Campero F.J. , Gomez-Zambrano A. , Turck F. , Calonje M (2017) **H2A monoubiquitination in *Arabidopsis thaliana* is generally independent of LHP1 and PRC2 activity** *Genome Biol* **18**

Zilberman D. , Coleman-Derr D. , Ballinger T. , Henikoff S (2008) **Histone H2A.Z and DNA methylation are mutually antagonistic chromatin marks** *Nature* **456**:125–129

Author information

Bhagyshree Jamge

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria, Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical University of Vienna, A-1030, Vienna, Austria

Zdravko J. Lorković

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

Elin Axelsson

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

Akihisa Osakabe

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria, Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan, PRESTO, Japan Science and Technology Agency, Honcho, Kawaguchi, Saitama, 332-0012, Japan

Vikas Shukla

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria, Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical University of Vienna, A-1030, Vienna, Austria

Ramesh Yelagandula

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria, Institute of Molecular Biotechnology, IMBA, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

Svetlana Akimcheva

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

Annika Luisa Kuehn

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

Frédéric Berger

Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Dr. Bohr-Gasse 3, 1030 Vienna, Austria

For correspondence: Frederic.berger@gmi.oeaw.ac.at

Editors

Reviewing Editor

Paula Casati

Center of Photosynthetic And Biochemical Studies (CEFOBI), Argentina

Senior Editor

Jürgen Kleine-Vehn

University of Freiburg, Germany

Reviewer #1 (Public Review):

Jamge et al. sought to identify the relationships between histone variants and histone modifications in Arabidopsis by systematic genomic profiling of 13 histone variants and 12 histone modifications to define a set of "chromatin states". They find that H2A variants are key factors defining the major chromatin types (euchromatin, facultative heterochromatin, and constitutive heterochromatin) and that loss of the DDM1 chromatin remodeler leads to loss of typical constitutive heterochromatin and replacement of this state with features common to genes in euchromatin and facultative heterochromatin. This study deepens our understanding of how histone variants shape the Arabidopsis epigenome and provides a wealth of data for other researchers to explore.

Strengths:

1. The manuscript provides convincing evidence supporting the claims that: A) Arabidopsis nucleosomes are homotypic for H2A variants and heterotypic for H3 variants, B) that H3 variants are not associated with specific H2A variants, and C) H2A variants are strongly

associated with specific histone post-translational modifications (PTMs) while H3 variants show no such strong associations with specific PTMs. These are important findings that contrast with previous observations in animal systems and suggest differences in plant and animal chromatin dynamics.

2. The authors also performed comprehensive epigenomic profiling of all H2A, H2B, and H3 variants and 12 histone PTMs to produce a Hidden Markov Model-based chromatin state map. These studies revealed that histone H2A variants are as important as histone PTMs in defining the various chromatin states, which is unexpected and of high significance.

3. The authors show that in *ddm1* mutants, normally heterochromatic transposable element (TE) genes lose H2A.W and gain H2A.Z, along with the facultative heterochromatin and euchromatin signatures associated with H2A.Z at silent and expressed genes, respectively.

Reviewer #2 (Public Review):

Jamge et al. set out to delineate the relationship between histone variants, histone modifications and chromatin states in *Arabidopsis* seedlings and leaves. A strength of the study is its use of multiple types of data: the authors present mass-spec, immunoblotting and ChIP-seq from histone variants and histone modifications. They confirm the association between certain marks and variants, in particular for H2A, and nicely describe the loss of constitutive heterochromatin in the *ddm1* mutant.

Overall, this study nicely illustrates that, in *Arabidopsis*, histone variants (and H2A variants in particular) display specificity in modifications and genomic locations, and correlate with some chromatin sub-states. This encourages future work in epigenomics to consider histone variants with as much attention as histone modifications.

Reviewer #3 (Public Review):

How chromatin state is defined is an important question in the epigenetics field. Here, Jamge et al. proposed that the dynamics of histone variant exchange control the organization of histone modifications into chromatin states. They found 1) there is a tight association between H2A variants and histone modifications; 2) H2A variants are major factors that differentiate euchromatin, facultative heterochromatin, and constitutive heterochromatin; 3) the mutation in *DDM1*, a remodeler of H2A variants, causes the mis-assembly of chromatin states in TE region. The topic of this paper is of general interest and the results are novel.

Overall, the paper is well-written and the results are clearly presented. The biochemical analysis part is solid.

Reviewer #4 (Public Review):

This work aims at analyzing the impact of histone variants and histone modifications on chromatin states of the *Arabidopsis* genome. Authors claim that histone variants are as significant as histone modifications in determining chromatin states. They also study the effect of mutations in the *DDM1* gene on the exchange of H2A.Z to H2A.W, which convert the silent state of transposons into a chromatin state normally found on protein coding genes.

This is an interesting and well done study on the organization of the *Arabidopsis* genome in different chromatin states, adding to the previous reports on this issue.

Author Response

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

Jamge et al. sought to identify the relationships between histone variants and histone modifications in Arabidopsis by systematic genomic profiling of 13 histone variants and 12 histone modifications to define a set of "chromatin states". They find that H2A variants are key factors defining the major chromatin types (euchromatin, facultative heterochromatin, and constitutive heterochromatin) and that loss of the DDM1 chromatin remodeler leads to loss of typical constitutive heterochromatin and replacement of this state with features common to genes in euchromatin and facultative heterochromatin. This study deepens our understanding of how histone variants shape the Arabidopsis epigenome and provides a wealth of data for other researchers to explore.

Strengths:

1. The manuscript provides convincing evidence supporting the claims that: A) Arabidopsis nucleosomes are homotypic for H2A variants and heterotypic for H3 variants, B) that H3 variants are not associated with specific H2A variants, and C) H2A variants are strongly associated with specific histone post-translational modifications (PTMs) while H3 variants show no such strong associations with specific PTMs. These are important findings that contrast with previous observations in animal systems and suggest differences in plant and animal chromatin dynamics.

1. The authors also performed comprehensive epigenomic profiling of all H2A, H2B, and H3 variants and 12 histone PTMs to produce a Hidden Markov Model-based chromatin state map. These studies revealed that histone H2A variants are as important as histone PTMs in defining the various chromatin states, which is unexpected and of high significance.

1. The authors show that in *ddm1* mutants, normally heterochromatic transposable element (TE) genes lose H2A.W and gain H2A.Z, along with the facultative heterochromatin and euchromatin signatures associated with H2A.Z at silent and expressed genes, respectively.

Weaknesses:

1. Following up on the finding that H2A.Z replaces H2A.W at TE genes in *ddm1* mutants, the authors provide in vitro evidence that DDM1 binds to H2A.Z-H2B dimers. These results are taken together to conclude that DDM1 normally removes H2A.Z-H2B dimers from nucleosomes at TE genes and replaces them with H2A.W-H2B dimers. However, the evidence for this model is circumstantial and such a model raises a variety of other questions that are not addressed by the authors.

The Reviewer raises a series of interesting questions. We proposed that DDM1 exchanges H2A.Z to H2A.W because it is the simplest model and also because LSH - the mammalian ortholog of DDM1 exchanges H2A to macroH2A. However we do stress in the revised manuscript that this is a model and other possible models that could involve chaperones and additional remodelers are possible. Addressing why the loss of DDM1 results in a net exchange of H2A.W to H2A.Z is not the purpose of this study. Here we use the perturbation caused by *ddm1* as a means to address the importance of the dynamics exchange of H2A variants in setting up the chromatin states. We do observe that perturbing this dynamic exchange causes an important perturbation of chromatin states. This further supports our

main conclusion: H2A variants dynamics are one important factor that organizes chromatin states.

For example: if DDM1 does remove H2A.Z from TE genes, how does H2A.Z normally come to occupy these sites, given that they are highly DNA methylated and that H2A.Z is known to anticorrelate with DNA methylation in plants and animals?

The anticorrelation between H2A.Z and DNA methylation is observed at steady state. The exchange of H2A.Z to H2A.W that results from the action of DDM1 would indeed remove unwanted H2A.Z from regions occupied by DNA methylation as suggested by the Reviewer.

Given that H2A.Z does not accumulate in TEs in h2a.w mutants, how would H2A.X and H2A instead become enriched at these sites if DDM1 cannot bind these forms of H2A?

This is a valid question: We envisage that H2A.X and H2A are deposited by remodelers and chaperones other than DDM1 in the h2a.w mutant.

Given that there are no apparent regions with common sequence between H2A.Z and H2A.W variants that are not also shared with other H2A classes, how would DDM1 selectively bind to H2A.W-H2B and H2A.Z-H2B dimers to the exclusion of H2A(X)-H2B dimers?

It was shown by the Muegge Lab both in vitro and in vivo that LSH - the mammalian ortholog of DDM1 binds to macroH2A and H2A, and these two H2A variants do not share similar specific region. Yet it remains to determine which region of H2A.Z and H2A.W binds to DDM1, which does not fit in the scope of this study.

Reviewer #2 (Public Review):

Jamge et al. set out to delineate the relationship between histone variants, histone modifications and chromatin states in Arabidopsis seedlings and leaves. A strength of the study is its use of multiple types of data: the authors present mass-spec, immunoblotting and ChIPseq from histone variants and histone modifications. They confirm the association between certain marks and variants, in particular for H2A, and nicely describe the loss of constitutive heterochromatin in the ddm1 mutant.

The support for some of the conclusions is weak. The title of the discussion, "histone variants drive the overall organization of chromatin states" implies a causation which wasn't investigated, and overstates the finding that some broad chromatin states can be further subdivided when one considers histone variants (adding variables to the model).

We have removed subtitles in the discussion and have taken care to avoid over simplified statements.

Adding variables to a ChromHMM model naturally increases the complexity of the models that can be built, however it is difficult to objectively define which level of complexity is optimal. The differences between states may be subtle to the point that they may be considered redundant. The authors claim that the sub-states they define are biologically important, but provide little evidence to support this claim. It is not obvious whether the 26 states model is much more useful than a 9-states model. Removing variables naturally affects the definition of states that depend on these variables, but it is also hard to define the biological significance of that change. This sensitivity analysis is thus not very developed.

We agree that adding more input tracks/ data will increase the complexity.

But we would like to mention the differences of this study and the 9-state model,

1. We have included the histone variants which have been previously missed in chromatin state definition.
2. The previous 9-state model used data from different tissue types. In this study all the data generated and analyzed is from seedlings.
3. Increasing the number of states allowed us to resolve heterochromatin states compared to 9-state model which was previously missed. (BioRxiv)
4. The biological relevance of the 26 states model is analyzed and described in depth (States BioRxiv paper).

In addition we have now updated the Figure 2F to include a more direct comparison of marks used in both models. And we have expanded the description in the methods section and our reasoning behind using 26 state model to be analyzed in depth.

There are issues with the logical sequence of arguments in Fig1 and Fig3. Fig1A shows that nucleosomes often contain both H3.1 and H3.3. Therefore pulling-down H3.1-containing nucleosomes also pulls down H3.3 and whether specific H2A variants associated with H3.1 cannot be answered in this way (Fig1B).

We thank the Reviewer for pointing this out. If 60% of nucleosomes are homotypic and if they would associate with a specific H2A variant this would be clearly visible on WB as a much stronger band. Also, the MS data presented in Figure 1 figure supplement 1D clearly show that all H2A variants associate with both H3.1 and H3.3. We have included in the revised version more detailed explanation to clarify this point.

The same issue likely carries to the investigation of the association with H3 modifications in Fig1C and 1D, since the H3.1-HA pull-down also pulls down endogenous H3.1 (so presumably the rest of the nucleosome, with H3.3, as well).

We disagree on this point. The H3 band corresponding to the transgene copy is either H3.1 or H3.3, so all signals on upper band (T) in Figure 1C are associated with either H3.1 (H3.1 IP) or H3.3 (H3.3 IP), thus unambiguously showing that all modifications we analyzed are present on both H3.1 and H3.3. Furthermore, data shown in Figure 1D and E, where we analyzed modifications on K27 and K36 which are in the H3 region that can be distinguished between H3.1 and H3.3 by MS clearly demonstrate that these modifications are present on both H3.1 and H3.3. In order to make this clearer, we also extended the description of this part in the Results section to emphasize this.

In Fig3, the conclusion that it is the loss of H2A.Z → H2A.W exchange in the ddm1 mutant that causes loss of constitutive heterochromatin is rushed. The fact that the h2a.w mutant does not recapitulate the loss of constitutive heterochromatin seen in ddm1 argues against this interpretation.

We agree that at first the minimal impact of the loss of H2A.W alone is surprising. However, we point to the preprint <https://www.biorxiv.org/content/10.1101/2022.05.31.493688v1>. There it is shown that the joint loss of H2A.W and H3K9 methylation (also observed in ddm1) affects silencing of a large range of transposons that also lose silencing in ddm1.

It's also difficult to conclude about the importance of dynamic exchanges when the ddm1 mutation has been present for generations and the chromatin landscape has fully readapted. Further work is needed to support the authors' hypothesis.

We apologize that the Reviewer could not find the information regarding the origin of ddm1 mutant material. We did not use a mutant where ddm1 mutations were kept for generations.

We were in fact very careful on this point and used leaves from ddm1 first homozygous plants segregated from heterozygous ddm1 kept heterozygous.

The study also relies on a large number of custom (polyclonal) antibodies with no public validation data. Lack of specificity, a common issue with antibodies, would muddle the interpretation of the data.

We added information about validation of custom made antibodies into Methods: "Specificities of custom made polyclonal antibodies against Arabidopsis H2A.Z.9, H2A.X, H2A.W.6, H2A.13, H2A.W.7, H2Bs, and linker histone H1 were validated in previous publications (Yelagandula et al., 2014; Lorkovic et al., 2017; Jiang et al., 2020; Osakabe et al., 2021)." For H2A.2 and H2A.Z.11 antibodies we provide validation data as Figure 2 figure supplement 1.

Overall, this study nicely illustrates that, in Arabidopsis, histone variants (and H2A variants in particular) display specificity in modifications and genomic locations, and correlate with some chromatin sub-states. This encourages future work in epigenomics to consider histone variants with as much attention as histone modifications.

Reviewer #3 (Public Review):

How chromatin state is defined is an important question in the epigenetics field. Here, Jamge et al. proposed that the dynamics of histone variant exchange control the organization of histone modifications into chromatin states. They found 1) there is a tight association between H2A variants and histone modifications; 2) H2A variants are major factors that differentiate euchromatin, facultative heterochromatin, and constitutive heterochromatin; 3) the mutation in DDM1, a remodeler of H2A variants, causes the mis-assembly of chromatin states in TE region. The topic of this paper is of general interest and results are novel.

Overall, the paper is well-written and results are clearly presented. The biochemical analysis part is solid.

Reviewer #4 (Public Review):

This work aims at analyzing the impact of histone variants and histone modifications on chromatin states of the Arabidopsis genome. Authors claim that histone variants are as significant as histone modifications in determining chromatin states. They also study the effect of mutations in the DDM1 gene on the exchange of H2A.Z to H2A.W, which convert the silent state of transposons into a chromatin state normally found on protein coding genes.

This is an interesting and well done study on the organization of the Arabidopsis genome in different chromatin states, adding to the previous reports on this issue.

Reviewer #1 (Recommendations For The Authors):

- 1. The rationale for switching from using 10-day old seedlings for chromatin profiling to using mature leaves in Figure 3 and beyond is not explained and introduces additional complexity into the analyses. The reasoning should be clearly explained in the text, and there are several additional suggestions or questions related to this that should be addressed:*

This was done for practical reasons. We had already obtained some profiles of marks in ddm1 mutants and extended the dataset using the same stage of development because this tied this study with our previous study. Using different stages of development provides an additional benefit. The same chromatin states are observed in 10 day old seedlings and leaves of older plants. Constitutive heterochromatin is occupied by the same chromatin states and

logically euchromatin is positioned on different genes as expected by the distinct pattern of gene expression at the two stages of development.

A) In the 16-state model (Figure 3A), euchromatin states were not well defined compared to the 26-state model. Why did the authors not profile these marks also, and could this explain why ddm1 mutants did not show a significant effect on euchromatin states in this model?

We apologize for the lack of detailed explanation: In our previous study we used leaves of five weeks ld plants to show the impact of ddm1 on the profiles of H2A.W.6, H2A.X, H1, H3K9me2, H3K36me3 and H3K27me3 in leaves (Jamge, Osakabe et al., 2021). This study showed that DDM1 causes the deposition of H2A.W.6 to heterochromatin and we thus used leaves to extend this investigation to the two other marks of heterochromatin (constitutive or facultative) H3K9me1, H2A.W.7 and H2A.Z.9 and H2A.Z.11.

B) The authors state that the tissue types do not impact the definition of chromatin states. However, there is a clear difference in the portion of the genome occupied by each chromatin state between leaf and seedling (states 1, 5, 8, 13, and 14; Figure S3A).

We had missed a comment on supFig3B and have now provided more explanation: "Although the composition of the chromatin states did not vary significantly between seedlings and leaves, each state occupied a similar proportion of the genome in seedling or leaves to the exception of state 5 present primarily in leaves and state 13 only present in seedlings (Figure 3 figure supplement 3A, right column with green bars) and the euchromatin states occupied different genes (Figure 3 figure supplement 3B) as expected by the dissimilar transcriptomes of these two developmental stages."

1. The naming of supplemental figures throughout the text is confusing as the legends refer to them as "Figure SX" but they are called out in the text as "Figure X figure supplement XA-B". The eLifeconvention is "Figure X figure supplement XA-B".

This was changed.

1. In Figure 4, Panel D is mislabeled as C in the figure, and C is lacking a label.

1. Please remove the word "the" from the title.

This was done

Reviewer #2 (Recommendations For The Authors):

Fig1D legend should also mention K37.

This was corrected.

Fig2F legend should say "no H3 modifications" rather than "no histone modifications" This was corrected.

Fig4 labels C/D do not correspond to the legend. D is missing and C should go to the ddm1 stacked barplot.

This was corrected.

H3 variants analysis: Taking the relative abundance of H3.1 and H3.3 (and transgenes) into account would be useful to interpret the results of the nucleosome composition results. If they are at equivalent amounts, the null hypothesis of independent association would give 50% heterotypic nucleosomes and 50% homotypic.

This is a valid comment. In an ideal system the last statement would be correct, but this does not take into account chromatin dynamics associated with replication, transcription, etc. Also, total amounts of H3.1 and H3.3 in tissue we used for the experiment is not known. It could possibly be inferred from RNAseq data, but if this would reflect real amounts of the protein is highly questionable. In Arabidopsis there are 5 H3.1 genes and 3 H3.3 genes. Nevertheless, we recalculated data for H3.1 and H3.3 and this has been updated in the main text (~60% of H3.1 and ~42% of H3.3 immunoprecipitated nucleosomes contained both H3 variants). Thus, from the available data these numbers are the best we can get.

p. 5 bottom paragraph. Repetition.

This was corrected

p12. The reference to LSH is dropped in without making clear how it is relevant. Expand on mechanism to suggest similar DDM1 mechanism?

This section was expanded to provide more background in the interpretation of the results.

p13. inversion between H2A.W and H2A.Z in "the loss of DDM1 prevents the replacement of H2A.W by H2A.Z".

This was corrected

p13. make it clear that the last sentence of the results is a working model, not a fully backed up conclusion.

Alternative models are mentioned in this section and in the discussion in the revised version.

p14 middle paragraph. Not clear what "in silico simulation" refers to. Simply chromatin-state classification with ChromHMM?

This refers to the Jacard index calculation in Fig. 2F that models the impact of the loss of H2A variants (or other elements of chromatin) on the definition of chromatin states by ChromHMM. This is now clarified.

p14 bottom paragraph: the H2A.Z tail repression of ubiquitin ligase but its being the favoured substrate for H2AK121Ub is apparently contradictory. Can this be explained?

This refers to H2B Ubiquitination and is now clarified

p15. Correlation between variants and modifications/chromatin states does not necessarily mean causation.

We agree and have improved the revised version in this respect.

p15 "forward feedback loop" is ambiguous (is it a feed-forward loop? A feedback loop?), just use "positive feedback loop".

This was corrected.

p23 top "(Ingouff et al)" doesn't seem properly formatted.

This reference did not belong there and has been removed.

Data availability: GSE226469 is not public. The manuscript also mentions availability of source data for all the main figures, but I could not find it. It would be great to make the code publicly available too.

All the data and code will be public upon posting the revised version of the manuscript.

Reviewer #3 (Recommendations For The Authors):

My major concern is authors only used DDM1 as an example to show that the exchange of the histone variant contributes to definition and distribution of chromatin state on transposons (i.e., constitutive heterochromatin regions associated with H2A.W). Readers may wonder whether similar mechanisms also work at the euchromatin region. This point should be clearly discussed and mentioned in the Results (for example, cite recent work on INO80).

We discuss the impact of other remodelers in the Discussion in the revised version. We hope that the reviewer will understand that doing a study on the impact of other remodelers on chromatin states which would require dozens of new ChIP profiles and is clearly beyond the scope of revising a manuscript.

Minor:

- 1. Fig. 2A and 2B, what does color mean? I guess the color code is referred to chromatin states (Fig. 2F).*

We have clarified on Figure 2A the attribution of a specific color to each chromatin state. This same color is used also in other panels of Figures 2 and S2.

- 1. Supplemental Figures: All the figure panels should be on the same page.*

We rearranged supplemental figures so that each figure fits on one page. In places where this was not possible, we created additional supplemental figures.

- 1. "We observed that increasing state numbers from 26 to 27 gave rise to biologically redundant states.": Where are the data? Fig S2A? This figure is hard to understand.*

In the updated manuscript, we have described the legend and the methods for FigS2A in more detail.

Reviewer #4 (Recommendations For The Authors):

A general concern refers to the text that frequently falls into excessive oversimplifications and/or overstatements, with the danger of being misleading for the reader. This needs to be thoroughly revised.

We added more careful statements and proposed alternative models when it was possible.

Specific comments.

- 1. Fig 1A. Authors found the ~40% of nucleosomes contained both H3.1 and H3.3. This is a significant finding that deserves a more detailed comment.*

We now provide a more detailed description of IP and MS data presented in Figure 1. This should also help to avoid oversimplifications and/or overstatements as criticized in a general comment.

- 1. Fig 1C. "H3. And H3.3 bore the same sets and comparable levels of methylation and acetylation...". Too general statement, please specify. Is this also the case for H3K9me2?*

Others?

We did describe this part into more detail to emphasize more precisely what Figure 1 shows. We also included data on K9me into Figure 1 figure supplement 1H.

1. Fig 1D. Could you confirm the high level of H3K27me1 on H3.3?

H3K27me1 data are shown both by WB (Figure 1C) and Mass spectrometry (Figure 1D and E). We also provide a possible explanation for high levels of this mark on H3.3 by taking into account the fact that H3K27me1 is also produced by demethylation of H3K27me3 by JMJ demethylases.

1. All WB in Fig 1. They need to be quantified and normalized (plus statistical analysis) in order to provide strong support to the conclusions.

The conclusion of all WB are supported by quantified Mass spectrometry data and many WB were even repeatedly shown in Figure 1F (for example IPs for H2A variants and a large set of H3 marks used for WBs) with the same results. Also, association of H3K4me3 and H3K36me3 with H2A variants was analyzed in both ways (Figure 1F); IPs of variants and WBs of variants and marks and IPs of marks and WBs of marks and variants. For most of the data we do not have more than two repeats, so statistical analysis may not be possible.

Nevertheless, we are convinced that our major conclusions from data presented in Figure 1 and Supporting figure 1 (these are: that H3 variants form both homotypic and heterotypic nucleosomes, that H3 marks do not preferentially associate with H3 variants but some of them do so with H2A variants and that H3 modifications show very complex pattern of associations with each other) are fully valid as they were drawn from two orthogonal approaches and further supported by the chromatin states identified.

1. Fig. 2A. Authors focus on "the most parsimonious model" based on 26 chromatin states. This needs to be justified in a more explicit manner. It is surprising that this number emerges for an analysis of 27 independent variants and marks. What are the differences in the conclusions when other number of states are used? See also below (reduced number of number derived from the "concatenated model").

Why 26 states were chosen is now explained in great details in the method section. Since to the exception of H2A variants that are invariably homotypic, nucleosomes can be heterotypic for all other histone variants and histone modifications, the random combination of the 27 marks in one nucleosome representing one states is 4 H2A (without the subtypes) x 4H3 x 2H1 x 2(power16) (for each mark) which is well above the circa 26 states observed. This shows that our probabilistic model reduces the potential complexity of a theoretical random association in a remarkable manner.

1. As a summary, it would be very helpful to generated a table (or similar) where is proposed chromatin state is ascribed to functional genomic elements.

This aspect of the work is presented in a preprint where the biological association with the chromatin is described in details. See Jamge et al 2002, <https://www.biorxiv.org/content/10.1101/2022.06.02.494419v1>

1. Fig 2F (and S2B). A comprehensive comparison a various approaches should include others and estimate the Jaccard similarity index: (1) the same of marks and variants used in the Sequeira-Mendes et al paper, and (2) the subset of marks and variants added in this study. In this way, a direct evaluation of the contributions could be more properly made.

We thank the reviewer for this suggestion and have now included a new column with the combination of marks and variants as used in Sequeira-Mendes et al., 2014 (see Figure 2F). These data clearly demonstrate that adding histone variants significantly contribute to the definition of chromatin states.

1. Fig. 3. Explain in more detail the concatenated model used here. Does the reduction in the number of chromatin states mean that the other do not add new information?

ChromHMM concatenated model allows to identify common definition of chromatin state in multiple tissue types. Here multiple cell types are concatenated leading to a shared definition of chromatin states, but specific to each cell type.

In our paper we used the concatenated model to identify common chromatin states in two different genotypes (WT and ddm1). The data for WT and ddm1 was obtained from leaves. As we had a limited number of ChIP-seq profiles in the leaves dataset The complexity of the concatenated model was also reduced compared to the extensive 26 chromatin state model. We chose to analyze 16-states in the concatenated model because this was the minimal number of states that gave rise to a similar complexity of heterochromatic states.

1. The ddm1 mutant. The text in page 14 is a bit confusing. It seems that H2A.Z is deposited on TEs and the exchanged by the H2A.W.

We have provided additional alternative models that could explain our observations.

1. Page 15: link between H2A.Z and H3K27me3. Gomez-Zambrano et al (2018, cited in the text, found that only a relatively small subset of (putative) targets are common to H2A.Z and H3K27me3. How do authors reconcile this with their statement supporting a link between both of them?

We refer to Gomez-Zambranao et al to illustrate the link between H2A.Z and H2AK121ub so we do not understand this comment. The strong link between H2A.Z and H3K27me3 is shown without ambiguity by our work and also Carter et al., 2018.