

Binding to nucleosome poises human SIRT6 for histone H3 deacetylation

Ekaterina Smirnova, Emmanuelle Bignon, Patrick Schultz, Gabor Papai , Adam Ben-Shem 

Department of Integrated Structural Biology, IGBMC, Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), 67400 Illkirch, France; Université de Strasbourg, IGBMC UMR 7104-UMR-S 1258, 67400 Illkirch, France; CNRS, UMR 7104, 67400 Illkirch, France; Inserm, UMR-S 1258, 67400 Illkirch, France; Equipe labellisée Ligue Contre le Cancer • Université de Lorraine and CNRS, UMR 7019 LPCT, F-54000 Nancy, France

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 3

December 7, 2023

Reviewed preprint version 2

October 5, 2023 (this version)

Reviewed preprint version 1

June 2, 2023

Sent for peer review

March 28, 2023

Posted to preprint server

March 20, 2023

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Sirtuin 6 (SIRT6) is an NAD^+ dependent histone H3 deacetylase that is prominently found associated with chromatin, attenuates transcriptionally active promoters and regulates DNA repair, metabolic homeostasis and lifespan. Unlike other sirtuins, it has low affinity to free histone tail but demonstrates strong binding to nucleosomes. It is poorly understood how SIRT6 docking on nucleosomes stimulates its histone deacetylation activity. Here we present the structure of human SIRT6 bound to a nucleosome determined by cryogenic electron microscopy. The zinc finger domain of SIRT6 associates tightly with the acidic patch of the nucleosome through multiple arginine anchors. The Rossmann fold domain binds to the terminus of the looser DNA half of the nucleosome, detaching two turns of the DNA from the histone octamer and placing the NAD^+ binding pocket close to the DNA exit site. This domain shows flexibility with respect to the fixed zinc finger and moves with, but also relative to, the unwrapped DNA terminus. We apply molecular dynamics simulations of the histone tails in the nucleosome to show that in this mode of interaction, the active site of SIRT6 is perfectly poised to catalyze deacetylation of the H3 histone tail and that the partial unwrapping of the DNA allows even lysines close to the H3 core to reach the enzyme.

eLife assessment

This manuscript provides a **useful** reconstruction of the structure of the sirtuin-class histone deacetylase Sirt6 bound to a nucleosome based on cryo-EM observations, and additional characterization of the flexibility of the histone tails in the complex based on molecular dynamics simulations. Similar structures have recently been published, but this work provides **solid** support for the conclusions of those papers and also includes some novel insights into the potential dynamics of Sirt6 bound to a nucleosome that help explain its substrate specificity.

Introduction

The state of lysine acetylation in histones, as well as in several other proteins, is determined by the balancing act of acetyltransferases and deacetylases. Sirtuin is an evolutionarily conserved family of lysine deacetylases that comprises seven members in humans and plays an important role in regulating numerous physiological and pathological processes including aging, cancer, inflammation and metabolism (Cen et al., 2011; Vitiello et al., 2017; Watroba et al., 2017). Unlike other lysine deacetylases sirtuins carry out their enzymatic reaction by transferring the acetyl group to an essential NAD^+ cofactor before acetylating the target protein (Imai et al., 2000).

Human Sirtuin 6 (SIRT6) is a multitasking enzyme implicated in diverse pathways. Through its histone H3 deacetylation at gene promoters, it is a transcription repressor attenuating inter-alia ribosome biogenesis (Sebastian et al., 2012), glycolysis (Taylor et al., 2022) and the expression of various transcription factors, including nuclear factor κB (NF- κB) (Kawahara et al., 2009) and hypoxia-inducible factor-1 α (HIF-1 α) (Zhong et al., 2010). Its H3K9 deacetylation of telomeric chromatin maintains telomere integrity (Michishita et al., 2008). In addition, via its second enzymatic activity, ADP-ribosylation of PARP1 (Mao et al., 2011) and by stabilizing DNA-PK on chromatin (McCord et al., 2009), SIRT6 also fosters DNA double-strand break repair.

These activities position SIRT6 as a key regulator of longevity as indeed shown by several lines of evidence. In mice, knockout of SIRT6 leads to a shortened lifespan and a premature aging-like phenotype (Kanfi et al., 2012). Similar observations were made in monkeys deficient for the enzyme (Zhang et al., 2018). Conversely, overexpression of SIRT6 extends lifespan and diminishes frailty in mice (Roichman et al., 2021). Furthermore, a study with several rodent species found a positive correlation between SIRT6 ability to promote DNA double-strand break repair and longevity -in other words, more active SIRT6 leads to longer lifespan (Tian et al., 2019).

The development of cancer can also be regulated by SIRT6. However, whereas in some malignancies SIRT6 plays a role as a tumor suppressor (Marquardt et al., 2013; Min et al., 2012; Sebastian et al., 2012), in others it acts as an oncogene (Bauer et al., 2012; Khongkow et al., 2013; Liu et al., 2013). The molecular mechanisms underlying SIRT6 effects on longevity or on the two contradictory roles in cancer development are not well understood, limiting the search for medically useful drugs that target the enzyme.

SIRT6 is a 37 KDa protein composed of two globular parts, a Rossmann fold domain that harbors the NAD^+ binding site and a zinc finger domain (Pan et al., 2011). A striking difference between SIRT6 and other members of the sirtuin family is the presence of a splayed zinc finger and the lack of a stabilizing helix bundle linking the two globular domains. This absence probably endows SIRT6 with a unique flexibility between the domains (Supplemental Figure 1).

These structural differences are manifested in function. Unlike other sirtuins, SIRT6 has very low activity on acetylated free H3 histones or histone peptides (Jiang et al., 2013). This seemed to contradict findings concerning the function of SIRT6, in particular data demonstrating its activity *in vivo* as H3 deacetylase (Michishita et al., 2008) as well as the fact that SIRT6 is nuclear and found mainly associated with chromatin (Michishita et al., 2005). The conundrum was settled when it was revealed that nucleosomes, the *in vivo* cognate substrate of SIRT6, dramatically enhance the enzyme activity towards acetylated H3 histones (Gil et al., 2013). Further biochemical analysis showed that SIRT6 binds nucleosomes with high affinity via multiple interacting sites (Liu et al., 2020).

However, a mechanistic understanding of SIRT6 binding to nucleosomes and how this association promotes higher enzyme efficacy is currently lacking.

Here we present the structure of human SIRT6 bound to the nucleosome. We identify interactions between the zinc finger domain and the acidic patch of the nucleosome as well as between the Rossmann fold and the DNA terminus, positioning the NAD⁺ binding pocket close to the nucleosome DNA exit site and partially detaching the DNA from the histone octamer. We use molecular dynamics simulations of the nucleosome to show that in this position the enzyme active site is perfectly poised to bind the H3 tail for deacetylation and that due to the unwrapping of the DNA even lysines close to the core of H3 can reach the enzyme. Furthermore, we find conformational changes, facilitated by the flexibility in SIRT6 between the zinc finger and the Rossmann fold, that could serve to fine-tune the position of the active site with respect to the H3 tail in order to trap its target acetylated residues.

Results

Overall structure

To investigate how nucleosomes stimulate the activity of SIRT6 we prepared nucleosome with the Widom-601 positioning sequence and reconstituted its complex with full-length recombinant human SIRT6 that was overexpressed in and purified from *E. coli*. The complex was stabilized via mild crosslinking before its deposition on cryo-EM grids.

We obtained a single particle cryogenic electron microscopy structure showing the nucleosome at 2.9 Å overall resolution and a protein density bound to its disc-like surface (**Fig. 1a** [↗](#) and Supplemental Fig. 2 and 3). SIRT6 could be fitted into this density with its zinc finger domain (resolved at 3.0 Å resolution), associated with the acidic patch of the nucleosome and its Rossmann fold domain (4.0-6.0 Å resolution) interacting with the DNA terminus at superhelix locations (SHL) 6 and 7 (**Fig. 1b** [↗](#) and Supplemental Fig. 4). The active site of SIRT6 is thus positioned just above where the H3 tail enters the nucleosome disc between the two DNA gyres, pseudosymmetric orientations of the Widom-601 DNA (Supplemental Fig. 5a). We find that SIRT6 is bound predominantly at the “right” TA-poor arm of the Widom-601 sequence where DNA interactions with the histone octamer are looser (Ngo et al., 2015 [↗](#)). We detect however a small population with a second SIRT6 bound at the “left” TA-rich tight arm (Supplemental Fig. 2).

Conformational heterogeneity analyses in cryoDRGN (Zhong et al., 2021 [↗](#)) reveal two principle large movements within the structure (**Fig. 2a,c** [↗](#)). First, we find that where the SIRT6 Rossmann fold binds, DNA is pulled away and detached from the histone octamer (**Fig. 2b** [↗](#)). Particles greatly vary in the extent of this shift that therefore probably represents a continuous movement of the DNA (compare **Fig. 2b1** [↗](#) with **2b2** [↗](#)). The Rossmann fold moves with and is attached to the DNA whereas the zinc finger domain remains fixed at the acidic patch of the nucleosome. In the full extent of the DNA movement, two turns from SHL 6 and 7 are detached from the histone octamer. This alters the path of the terminus DNA by 37° and breaks DNA interactions with histones H2A, H2B, and H3 (Supplemental Fig. 5b). Such a large change in DNA trajectory is reminiscent of the chromatin remodelling ATPases or pioneer transcription factors binding to nucleosome but was not observed in other histone modifiers (Dodonova et al., 2020 [↗](#); Farnung et al., 2017 [↗](#)). Importantly, the DNA terminus without bound SIRT6 or free nucleosomes in our dataset did not show any deviation from the canonical conformations (Supplemental Fig. 5c). In addition, our MD simulations indicate higher flexibility of the DNA termini when SIRT6 is bound to the nucleosome (Supplemental Fig. 6). All together these findings strongly suggest that the DNA movement we observed is coupled to SIRT6 interactions. Second, in particles where the SIRT6-bound DNA remains in its canonical position, we observe shifts by up to 20 Å in the position of the Rossmann fold domain of SIRT6, partially loosening its association with the DNA (**Fig. 2d** [↗](#)). We show that this shift could be involved in fine tuning the position of the active site relative to the H3 tail.




Figure 1

Structure of human SIRT6 in complex with the nucleosome

a, Front (top) and side (bottom) views of a composite cryo-EM reconstruction of human SIRT6-nucleosome. Maps from focused refinements of SIRT6 (magenta – Rossmann fold and Zn-finger domains labelled) and the nucleosome (H2A – yellow; H2B – orange; H3 – blue; H4 – green; DNA – light and dim grey). **b**, Corresponding views of the atomic model of the complex.




Figure 2

Flexibility of the SIRT6 Rossmann fold

a, UMAP projection of the latent embeddings of a subset of SIRT6-nucleosome particles representing the concerted movement of the Rossmann fold and the DNA terminus. **b**, Structural representation of the two endpoints of the latent embeddings shown in panel **a**. **c**, Overlay of the DNA ends and the SIRT6 Rossmann fold of two endpoints shown in panel **b**. Arrows show their displacements between the two endpoints. **d**, UMAP projection of the latent embeddings of another subset of SIRT6-nucleosome particles representing the movement of the Rossmann fold with respect to the nucleosomal DNA. **e**, Structural representation of the two endpoints of the latent embeddings shown in panel **d**. **f**, Overlay of the DNA ends and the SIRT6 Rossmann fold of two endpoints shown in panel **e**. Arrow show the displacement of the Rossmann fold between the two endpoints.

These conformational changes are made possible by a flexible hinge region between the two domains of SIRT6, which is unique to this member of the sirtuins family.

Zinc finger domain binds the acidic patch

An important feature of the nucleosome surface is a cluster of acidic residues (E56, E61, E64, D90, E91, E92 of H2A and E102, E110 of H2B) that forms a negatively charged ‘acidic patch’ and takes the shape of a narrow groove. It serves as a docking site for many nucleosome-binding proteins. Four side chains of the SIRT6 zinc-finger domain extend into different sections of the acidic patch groove and interact with the negatively charged residues. **R175** packs against H2B L103 and serves as a classical “arginine anchor motif” (McGinty and Tan, 2016 [\[1\]](#)), binding to the acidic triad of E61, D90 and E92 from H2A. SIRT6 presents two other arginines, **R172**, to contact E56 and Q24 of H2A as well as Q44 and E110 of H2B and **R178** to contact H2A E91 and E102 from H2B. Finally, **K170** interacts with E64 and N68 from H2A (Fig. 3a [\[2\]](#)).

Substituting these four positively charged residues to alanines (SIRT6-4A) significantly weakened the interaction of SIRT6 with the nucleosomes as shown by electrophoretic gel-shift assay (Supplemental Fig. 7a). To test also the activity of SIRT6-4A we first acetylated the H3 tails of the nucleosomes with purified SAGA complex. The mutant protein showed strongly diminished activity in deacetylating H3K9Ac, further corroborating the role of the four basic residues R175, R172, R178 and K170, in binding to the nucleosome (Supplemental Fig. 7b).

All four SIRT6 positive residues are located in a loop that makes a sharp turn, due to a conserved glycine at position 173, connecting two beta strands and penetrates the acidic patch. Interestingly, this loop appears to be flexible in crystal structures of SIRT6 but rendered stable by its association with the acidic patch. Binding to different sections of the acidic patch maintains a strong hold of the zinc finger at the surface of the nucleosome and orients the Rossmann fold domain towards the DNA terminus. This strong hold is maintained throughout the movements of the Rossmann fold domain (Fig. 2 [\[3\]](#)).

It is important to note that the four positive residues of SIRT6 that bind the nucleosome acidic patch are conserved among vertebrates but in contrast are not present in other human sirtuins (Fig. 3b,c [\[4\]](#)). This probably reflects the fact that SIRT6 is the only sirtuin predominantly associated with chromatin.

Rossmann fold domain binds nucleosome DNA exit site

Due to its flexibility with respect to the fixed zinc finger domain, the resolution of the Rossmann fold domain is not sufficiently high to distinguish most individual side chains. However secondary structure elements, mostly helices, are clearly visible and guided the precise fitting of the crystal structure of this domain.

We find that three highly conserved arginines from the Rossmann fold domain (R220, R231, R248) bind the phosphate backbone at SHL 6 and 7 of the nucleosomal DNA (Fig. 3d [\[5\]](#)). These interactions are mostly maintained throughout the movement of the DNA terminus with the Rossmann fold domain bound. However, by comparison to dedicated DNA binding proteins such as transcription factors or TBP, the number of observed DNA-protein interactions is low in SIRT6. A biochemical study suggests that the intrinsically disordered C-terminal domain of SIRT6 is necessary for establishing tight interaction of SIRT6 with the nucleosome (Liu et al., 2020 [\[6\]](#)) via its ability to bind DNA. Our maps do show an additional isolated protein density that we could not assign with confidence to SIRT6 but associates with the DNA terminus Supplemental Fig. 8a). We propose however that the three arginines are required to correctly position the Rossmann fold, and hence the catalytic site, with respect to the H3 tail. Indeed, applying SAGA-acetylated nucleosomes as substrates, we find that a SIRT6 mutant, where the three arginines are substituted




Figure 3

Binding of SIRT6 to the nucleosome

a, Close up view on zinc-finger interactions with the acidic patch. Color code as in Fig 1. **b**, Protein sequence alignment of human sirtuins. Red boxes and asterisks depict the residues of SIRT6 interacting with the acidic patch of the nucleosome. **c**, Protein sequence alignment of SIRT6 from different species highlighting the same amino acids as in panel b. Organisms: mm *Mus musculus*, rn *Rattus norvegicus*, bt *Bos taurus*, hs *Homo sapiens* and cc *Castor canadensis*. **d**, Depiction of the 3 arginines of SIRT6 (magenta) interacting with the DNA (grey). **e**, H2A c-terminal tail (yellow) interacts with SIRT6(magenta). **f**, SIRT6 binding to H2A or H2AZ containing nucleosomes. Bars show the fraction of residual nucleosomes that did not shift with bound SIRT6 in an electron-mobility shift assay (Supplemental Fig. 9). Bars represent mean $\pm$ SD of three biological replicates (shown as dots). ** indicates a statistically significant difference between the fraction of residual H2A and H2A.Z containing nucleosomes ($P=0.0016$ in paired t-test).

to alanines (SIRT6-3A), has severely diminished H3K9Ac deacetylase activity (Supplemental Fig. 7b). These findings confirm the importance of the three lysines R220, R231 and R248 to the enzymatic function.

Unlike other sirtuins, SIRT6 binds NAD^+ in the absence of an acetylated substrate. This is attributed to the fact that crystal structures of SIRT6 describe a rigid helix ($\alpha 3$) where other sirtuins contain a flexible cofactor binding loop (Pan et al., 2011 [↗](#)). In our map, we do not observe helix $\alpha 3$ (Supplemental Figure 8b) and we speculate that in the context of the nucleosome, this helix assumes a flexible loop conformation that like in other sirtuins will bind the cofactor only after stabilization by the acetyl lysine substrate.

In addition, we find that the N-terminal part of the Rossmann fold domain associates with the C-terminal tail of histone H2A at residue K119 (**Fig. 3e** [↗](#)). This observation suggests that monoubiquitination of H2A at this position, a major histone modification in mammalian cells, could interfere with the binding of SIRT6 or regulate its activity. Furthermore, we used electrophoretic mobility shift assay to follow the binding of SIRT6 to nucleosomes harboring the histone variant H2A.Z which deviates considerably from the canonical H2A in its C-terminal tail but not in other residues that contact SIRT6. Surprisingly, we find that binding of SIRT6 to nucleosomes is mildly (but statistically significant) stronger when H2A.Z is incorporated instead of H2A (**Fig. 3f** [↗](#) and Supplemental Fig. 9). This finding could be related to SIRT6's role in repressing gene expression because H2A.Z is enriched in nucleosomes that flank the transcription start site.

SIRT6 is poised to deacetylate the H3 tail

The classical model of the nucleosome posits an elongated and disordered histone H3 N-terminal tail that freely roams and extends away from the core. However recent NMR studies and molecular dynamics predictions show that the tail actually collapses onto the nucleosome core, driven by robust interactions with DNA (Ikebe et al., 2016 [↗](#); Stutzer et al., 2016 [↗](#)). The tail adopts multiple structurally heterogeneous, but energetically similar, states. It dynamically transitions between these conformations that all interact with the last superhelix location of nucleosomal DNA (Morrison et al., 2018 [↗](#)). We used a published set of 15 predicted H3-tail conformations resulting from end-states of molecular dynamics simulations of the whole nucleosome (Morrison et al., 2018 [↗](#)), to analyze whether the position of SIRT6 on the nucleosome serves the enzyme's role in trapping residues of histone H3 tail. The nucleosome from each molecular dynamics simulation was superposed on our model of SIRT6 bound to a nucleosome with the DNA terminus at the canonical position. To represent the full range of movement of the Rossmann fold domain relative to the DNA we used two endpoints of this movement namely fully bound to DNA and partially dissociated from it (**Fig. 4a** [↗](#) and **b** [↗](#), respectively). We then examined the position of the SIRT6 active site with respect to the predicted paths of the histone H3 tail.

We find that in several H3 conformations, the tail is located in close proximity ($<15\text{\AA}$) to the SIRT6 active site. In some of these cases, H3K9 or H3K18, the main target residues of SIRT6 (Wang et al., 2016 [↗](#)), are practically situated in the active site. Furthermore, when analyzing the two endpoints of the Rossmann fold domain shift with respect to the DNA, we observe that in some H3 conformations, this shift draws the active site closer to the target residues (**Fig. 4c,d** [↗](#)). We conclude that the high-affinity binding of SIRT6 to the nucleosome poises the active site to trapping its main target residues. In addition, the flexibility of the Rossmann fold domain serves as a fine-tuning mechanism that allows for short-range scanning of the tail in order to locate the acetylated residues. We note also that except H3 no other histone is present at the vicinity of the SIRT6 active site. This gives rise to selectivity of SIRT6 towards the H3 tail.

To shed light on possible benefits offered by SIRT6-mediated DNA unwrapping we conducted molecular dynamic simulations of a nucleosome with bound SIRT6. Strikingly, new H3 tail conformations appear in the simulations that are absent from a control run with nucleosome alone. These pathways have the H3 tail protruding out of the histone core through the space




Figure 4

SIRT6 poised to deacetylate lysine residues of H3

a, Side View of SIRT6 bound to the nucleosome with histone H3 K9 residue (red sphere) closest to the SIRT6 active site (blue spheres) from a set of molecular dynamics simulations. **b**, Side View of SIRT6 bound to the nucleosome with histone H3 K18 residue (orange sphere) closest to the SIRT6 active site from a set of molecular dynamics simulations. **c**, All H3K9 positions (red spheres) in close proximity (<15 Å) to the SIRT6 active site (blue) taken from a set of 15 molecular dynamics simulations and depicted on the surface view of SIRT6 bound to nucleosome. **d**, All H3K18 positions (orange spheres) in close proximity (<15 Å) to the SIRT6 active site (blue) taken from a set of 15 molecular dynamics simulations and depicted on the surface view of SIRT6 bound to nucleosome. **e**, Molecular dynamics simulations show that H3 c-terminal tail (blue) can protrude towards SIRT6 in a space formed between the histone octamer and the DNA. H3K27 is shown in red.

between the DNA and the octamer that is formed via partial unwrapping of the DNA by SIRT6. Importantly, in these additional conformations H3 tail does not need to circle around the DNA to reach SIRT6 and therefore lysines that are closer to the core of H3 (e.g., K27) could also be accessible to the enzyme (**Fig. 4e**). To lend support to this claim we tested whether SIRT6 can deacetylate residue H3K27 that was first acetylated by SAGA (Supplemental Fig. 7c). We find that indeed SIRT6 could efficiently deacetylate H3K27Ac, although at a somewhat slower rate than H3K9Ac. We conclude that partial DNA unwrapping by SIRT6 allows H3-tail conformations that make lysines that are close to the core of H3 accessible to the enzyme.

Discussion

Most histone-modifying enzymes, including sirtuin family members such as SIRT1, show significant activity on isolated target peptides and form multiple interactions with the residue to be modified as well as with flanking amino acids to provide selectivity. It is therefore striking that the measured turnover rate for SIRT6 on isolated H3K9 peptides is 1000 slower than for other sirtuins. Moreover, unlike SIRT1, the activity of SIRT6 is dramatically stimulated by binding to its cognate nucleosome substrate.

We solved the structure of SIRT6 bound to a nucleosome and used heterogeneity analysis to uncover the main conformational changes in the complex. Examining these findings in the context of molecular dynamics studies of the nucleosome that produce putative paths of histone tails, shed light on the unique properties of SIRT6.

We find that multiple interactions of SIRT6 with the nucleosome position its active site at the putative path of relatively long-lived conformations of H3 so that residues K9 and K18 of the tail spend significant time very close to, or even at, the enzyme active site. In addition, SIRT6 moves with respect to the nucleosomal DNA allowing it to fine-tune the position of the active site to trap the target residue. Hence it is the binding of SIRT6 to the nucleosome and an intrinsic scanning capability that account for its much higher enzymatic efficiency on nucleosomes. Importantly, the position of SIRT6 also ensures high selectivity because the space surrounding the active site is devoid of any other tails apart from H3.

The recognition mode of nucleosomal H3 employed by SIRT6 is reminiscent of that used by the PRC1 complex in the specific ubiquitylation of H2A Lys119 (McGinty et al., 2014). Similar to SIRT6, PRC1 requires a nucleosome substrate, showing no activity on an H2A-H2B dimer, and multiple interactions with the nucleosome position an active site in close proximity to its target residue. However, the flexibility and scanning capacity we observed for SIRT6 is lacking in PRC1 perhaps echoing the fact that its target residue is rather stable in the nucleosome compared to the H3 tail.

It is interesting to note that the flexibility of SIRT6 that is at the root of its scanning mechanism was suggested to be, at least in part, responsible for its nearly undetected activity on isolated peptides (Pan et al., 2011). It is only when it is constrained to the proximity of the target residues, through binding to the nucleosome, that this capacity facilitates the function of the enzyme.

A second major conformational change observed in our 3D reconstructions is the detachment of the DNA terminus, with a bound Rossmann fold domain, from the octamer. Our molecular dynamics simulations suggest that the space formed between DNA and octamer allows for additional conformations of the H3 tail that protrudes towards SIRT6. These new pathways do not require circling around the DNA gyre and therefore lysines that are closer to the core of H3 (e.g.,

K27) can reach in this way the active site of SIRT6. We note that K56 which lies at the core of H3 within a stable helix was also reported to be a target for SIRT6. It is therefore possible that the enzyme employs additional recognition modes of its targets.

Because the position of the SIRT6 active site seems to be well suited for modifying the H3 tail, it stands to reason that other H3 modifying enzymes, such as the GCN5 HAT module of SAGA and ATAC, would place their active site in a position similar to the one of SIRT6 and possibly also displace the DNA from the histone octamer. Gcn5 deviates from SIRT6 in having significant affinity for the H3 tail and therefore might pose a less stringent requirement for precise positioning of the active site or for specific conformations of the H3 tail. Thus, we speculate the Gcn5 recognition mode will be similar to SIRT6 but will also include some sequence specific binding. While this manuscript was under review two studies describing the structure of SIRT6-nucleosome appeared in press (Chio et al., 2023 [DOI](#); Wang et al., 2023 [DOI](#)). The conclusion of these papers regarding the position of SIRT6 on the nucleosome and the unwinding of DNA by the enzyme are similar to our findings. We however dissected in addition the movements of SIRT6 on the nucleosome and analyzed via molecular dynamics the conformations of the H3 tail with respect to the SIRT6 active site. Our results point to the importance of the flexibility between the globular domains of SIRT6 and also explain how SIRT6 can access lysines that are much closer to the histone core than H3K9.

Methods

Human SIRT6 purification

Plasmid (AddGene #28271) was transformed into *E. coli* strain BL21-RIL. Cells in 1L TB (with 50 µg/ml Kanamycin and 25 µg/ml Chloramphenicol) grew at 37C degrees till OD₂₆₀ reached 0.8 when they were rapidly cooled on ice to 18C. Over-expression was induced by the addition 0.5 mM IPTG and cells continued to grow at 18C over-night. All subsequent steps were performed at 0-4C and solutions were always supplemented with a protease inhibitor cocktail (PMSF, Peptstatin, E-64). Cells were harvested by centrifugation at 8250g for 20 minutes, suspended in buffer A200 (20 mM Hepes pH 7.5, 5% glycerol v/v, 2 mM beta-mercaptoethanol and 200 mM NaCl) and treated with lysosyme (1 mg/ml) for 30 minutes on ice. Salt concentration was increased to 0.5M and cells were then lysed by sonication (6*10sec). Debris were pelleted by centrifugation 45,000g*20minutes. The supernatant was incubated with 0.5 ml of Ni beads (1 ml slurry, Roche), equilibrated in buffer A500, for 1h. The solution was poured into a gravity column (CliniScience). Beads were then washed by 10 ml of buffer A500 with 5mM imidazole before elution by 2ml of buffer B300 (20 mM Hepes pH 7.2, 5% glycerol v/v, 2 mM beta-mercaptoethanol and 300mM NaCl) with 150mM imidazole. The eluted protein was diluted with two volumes of buffer D (20 mM Hepes pH 7.0, 5% glycerol v/v, 2 mM beta-mercaptoethanol), loaded on a 1ml Heparin column and eluted with 0.1M-1M salt gradient in buffer D. Fraction were analyzed by SDS-PAGE and those containing SIRT6 were pooled and dialyzed against buffer E (20 mM Hepes pH 7.5, 2% glycerol v/v, 0.5 mM TCEP and 150 mM NaCl. Human SIRT6 was then concentrated to 7.8 mg/ml, flash frozen in N₂, and kept in aliquotes at -80C.

Cloning of SIRT6 mutants

SIRT6 mutants were created as follows: the region of AddGene plasmid #28271 between SacI and HindIII sites were synthesized by SynBio company with changes in DNA sequence leading to K170A, R172A, R175A, R178A, R220A, R231A and R248A mutations in amino acid sequence. A NheI restriction site was also introduced between the first four and last three mutations without changes in amino acid sequence in order to facilitate cloning. A plasmid expressing SIRT6-7A that incorporates all seven mutations was created via ligation of the synthetic DNA fragment into the original expression plasmid between the SacI and HindIII sites. Fragments of the original expression plasmid corresponding to the regions between HindIII and NheI or NheI and SacI were

amplified with PCR and were cloned separately through corresponding restriction sites into the plasmid expressing SIRT6-7A to yield plasmids expressing SIRT6 with the first four mutations (SIRT6-4A) or the last three mutations (SIRT6-3A), respectively.

Octamer reconstitution and nucleosome DNA

Octamers were reconstituted from individual *Xenopus leavis* (canonical) or human (H2A.Z) histones expressed as inclusion bodies according to the standard protocol (Dyer et al., 2004 [↗](#); Luger et al., 1999 [↗](#)). Widom 601 145bp DNA was produced using a plasmid harboring 16 copies of this sequence as described by Dyer et al. (Dyer et al., 2004 [↗](#)).

Nucleosome core particle reconstitution

Nucleosomes with 145bp Widom 601 positioning sequence were prepared according to NEB Dilution Assembly Protocol (E5350) (<https://international.neb.com/protocols/2012/06/02/dilution-assembly-protocol-e5350> [↗](#)) with some modifications as follows: 2.75 μ M 145 bp Widom601 DNA was mixed with 2.75 μ M canonical or H2AZ histone octamers in a solution containing 2M NaCl, 1mM EDTA, 5mM beta-mercaptoethanol. The solution was incubated for 30 min at RT and then underwent serial dilutions down to 1.48M, 1M, 0.6M, 0.25M NaCl with buffer LowSalt (10 mM Hepes-KOH pH 8.0, 2.5 mM beta-mercaptoethanol). After each dilution the solution was kept at RT for 30 min. In order to reduce the final NaCl concentration, nucleosomes were concentrated in 0.5 mL 100KDa cutoff Amicon up to 100 μ l, then diluted 5 times with buffer LowSalt. This step was repeated one more time. Finally, nucleosomes were concentrated to 3-4 μ M and analyzed in a 5% native 0.2xTBE polyacrylamide gel to ascertain the quality of the sample and absence of free DNA.

Nucleosome acetylation

Canonical W601 nucleosomes (final concentration 1.8 μ M) were mixed with the holo-SAGA complex (final 18 nM) purified from *Pichia pastoris* cells according to (Papai et al., 2020 [↗](#)) in 20 mM Hepes-KOH pH 7.5, 10% v/v glycerol, 50 mM NaCl, 2 mM $MgCl_2$, 5 mM beta-mercaptoethanol, 0.005% Tween-20, and 0.05 mM Acetyl-CoA. Reaction was incubated on ice for 16 hours. On the next day SAGA was removed via incubation of the mixture with Streptavidin Sepharose beads. In order to reduce the remaining Acetyl-CoA concentration, nucleosomes were concentrated in 0.5 mL 100KDa cutoff Amicon up to 100 μ l, then diluted 5 times with buffer 10 mM Hepes-KOH pH 8.0, 2.5 mM beta-mercaptoethanol. This step was repeated four more time. Finally, the nucleosomes were concentrated to 3.5 μ M.

Binding and gel-shift assays

For nucleosome-binding assays, 100 nM of recombinant nucleosomes (harboring either H2A or H2AZ histone) were incubated in binding buffer (10 mM Hepes, 50 mM NaCl, 0.5 mM TCEP, 6% glycerol, and pH 7.5) with SIRT6 at 1:2 molar ratio. The reactions were performed on ice for 1h in LoBind tubes (Eppendorf) to minimize protein loss. The reactions were then resolved in a 0.2x TBE 5% gel. The gel was stained with Ethidium Bromide. Bands were quantified with ImageJ.

Deacetylation assay

Non-acetylated or SAGA-acetylated canonical W601 nucleosomes were mixed with SIRT6 (wt, SIRT6-4A or SIRT6-3A) at 100 nM and 40 nM, respectively, in 50 mM Hepes-KOH pH 7.5, 1 mM DTT, 0.2 mg/mL BSA, 1 mM NAD^+ . Reactions were set on ice and then incubated at 37 °C for 1 or 2 hours. Reactions were quenched by boiling with 5X Laemmli buffer. The proteins were separated on 15% SDS-PAGE and acetylation was analyzed by Western blot using the following antibodies: For H3K27Ac CellSignaling #8173S diluted 1:1000; For H3K9Ac Abcam Ab4441 1:2500; For H3 CellSignaling #14269S 1:1000.

Grid preparation

SIRT6 was first diluted in buffer (20mM Hepes-K pH 7.5, 150mM NaCl, 0.5 mM TCEP, 2% Trehalose). Nucleosome and SIRT6 were then mixed at a 1:2 molar ratio. Final conditions included 1.7 μ M Nucleosome and 3.4 μ M SIRT6 in buffer (20mM Hepes-K pH 7.5, 18mM NaCl, 0.5mM TCEP). After 1h on ice glutaraldehyde and the detergent dodecyl-maltoside were added at a final concentration of 0.05% v/v and 0.0125% w/v, respectively. The sample was incubated with the cross-linker for an additional 30-60 minutes. 2nm carbon foil was floated on 3.5/1 300 mesh Quantifoil grids. 3 μ l of sample was deposited onto such grids, blotted 5s with blot force 5 after 60s incubation in a Vitrobot IV (Thermo Fisher) at 6 °C using 100 % humidity and flash-frozen in liquid ethane.

cryo-EM data acquisition

Two different datasets were collected. For the first dataset images were acquired on a Cs-corrected Titan Krios G1 (Thermo Fisher) microscope operating at 300 kV in nanoprobe mode using serialEM for automated data collection. Movie frames were recorded on a Gatan K3 direct electron detector after a Quantum LS 967 energy filter using a 20 eV slit width in zero-loss mode. Images were acquired hardware-binned at a nominal magnification of 130,000, which yielded a pixel size of 0.862 Å. Forty movie frames were recorded at a dose of 1.12 and 1.30 e^- per Å² per frame. The second image dataset was acquired on a Titan Krios G4 microscope operating at 300 kV in nanoprobe mode using EPU for automated data collection. Movie frames were recorded on a Flacon 4i direct electron detector after a Selectris X energy filter using a 10 eV slit width in zero-loss mode. Images were acquired at nominal magnification of 270,000, which yielded a pixel size of 0.458 Å.

Image analysis

For the first dataset movie frames were aligned, dose-weighted in Warp (Tegunov and Cramer, 2019) together with CTF estimation and particles picking. For the second dataset frame alignment, dose-weighting and CTF estimation was done in cryoSPARC (Punjani et al., 2017), picking was done with crYOLO (Wagner et al., 2019). These datasets were analyzed in cryoSPARC and RELION 3.1 (Zivanov et al., 2018) according to standard protocols. Briefly, images were subjected to reference-free 2D classification in cryoSPARC to remove images with contaminations. Ab-initio reconstruction was performed on the selected images followed by heterogeneous and homogeneous refinements in cryoSPARC. Structural variability and flexibility were explored with Relion 3.1 using 3D classification without alignment with different T values and cryoDRGN (Zhong et al., 2021) according to standard protocols. Sharpening of the maps was done with DeepEmhancer (Sanchez-Garcia et al., 2021).

Model building and refinement

A composite map was created from the part corresponding to the SIRT6 Rossmann-fold from structure of the first dataset and the SIRT6 Zn-finger plus nucleosome from the structure of the second dataset. Crystal structure of Widom 601 sequence containing nucleosome (PDB: 3LZ0) and human SIRT6 (PDB: 5×16) were rigid-body fitted into this composite cryo-EM structure in Chimera (Pettersen et al., 2004). Flexible-fittings were performed with Isolde (Croll, 2018) in ChimeraX (Goddard et al., 2018) into the cryoDRGN structures. All models were refined in real-space with PHENIX (Liebschner et al., 2019) and Isolde.

Molecular Dynamics simulations

All molecular dynamics (MD) simulations were performed with the AMBER20 suit of programs, using the ff14SB (Maier et al., 2015) force field with bsc1 DNA corrections (Ivani et al., 2016) and CUFIX ions corrections (Yoo and Aksimentiev, 2018) that improve the description of disordered region conformations. Parameters for the zinc finger were taken from Macchiagodena et al. (Macchiagodena et al., 2020).

The histone tails were added to the cryo-EM structure based on the work described by Armeev et al (Armeev et al., 2021 [↗](#)), and protonation states of the protein residues were checked with the propka3 software (Olsson et al., 2011 [↗](#)). The system was soaked in a truncated octahedral TIP3P water box of 20Å buffer and 0.15 mM of NaCl ions were added to simulate physiological conditions, resulting in a total of 330,000 atoms.

The minimization of the energy was carried out in 5 steps, each one alternated with a 20ps NVT equilibration step at 100K to ensure a very smooth optimization of the starting geometry. Each minimization run included 5,000 steps in steepest descent and 5,000 steps in conjugate gradient, with decreasing restraints on the protein and nucleic atoms from 20kcal/mol to 5kcal/mol that were kept in the associated equilibration step. The last minimization step was released of any constraints and followed by a final NVT equilibration run at 300K. The time step was then switched from 2 to 4fs as allowed by the use of the SHAKE bond length constraints (Kräutler et al., 2001 [↗](#)) and the Hydrogen Mass Repartitioning algorithm (Hopkins et al., 2015 [↗](#)). The latter was applied on the topology using the parmed tool of Amber20. A 40ns equilibration run in NTP was then carried out, followed by a 5 μs production. Coordinates were outputted every 1ns during the production run. The temperature was kept constant using the Langevin thermostat with a 2ps⁻¹ collision frequency, and the Berendsen barostat was used to maintain the pressure at 1 atm in the NTP runs. Electrostatic interactions were treated using the Particle Mesh Ewald approach (Darden et al., 1993 [↗](#)) with an 8Å cut-off. Three replicates were performed starting with different velocities at the unconstrained 300K NVT equilibration step, resulting in a total of 15 μs of sampling.

A cluster analysis was performed on the MD ensembles based on the DNA and protein RMSD. Conformations were extracted every 5 frames of the trajectories to avoid memory overload and were sorted using the hierarchical agglomerative and average linkage algorithms. 10 clusters were generated by replicate (Supplemental Table 2).

The DNA flexibility analysis was performed on each MD ensemble by using a Principal Component Analysis-based machine learning script quantifying the contribution of each residue to the overall dynamics of the system. Based on the Scikit-learn58 library (Fleetwood et al., 2020 [↗](#)), it calculates a covariance matrix from the internal coordinates of the MD ensemble. In this matrix, the eigenmodes showing the highest amplitude are the most implicated in the main fluctuations of the system, and the contribution of each residue is normalized based on these values.

The control system was generated by removing SIRT6 from the initial structure and was submitted to the same protocol.

Data availability

The cryo-EM maps have been deposited in the Electron Microscopy Data Bank (EMDB) under accession codes EMD-16842 (High-resolution on the nucleosome), EMD-16843 (SIRT6-nucleosome refined for SIRT6) and EMD-16845 (composite map). The model coordinates for SIRT6-nucleosome were deposited in the Protein Data Bank (PDB) under the accession code 8OF4.

All MD input files are publicly available on Github: <https://github.com/emmanuellebignon/NCP-SIRT6> [↗](#)

Acknowledgements

We acknowledge support from the Institut National de la Santé et de la Recherche Médicale (INSERM), the Centre National pour la Recherche Scientifique (CNRS), the Ligue contre le Cancer. We acknowledge the use of resources of the French Infrastructure for Integrated Structural Biology FRISBI ANR-10-INBS-05 and of Instruct-ERIC. E.B. thanks GENCI (HPC resources of IDRIS, allocation A0120713412) for computational resources.

References

- Armeev G.A., Kniazeva A.S., Komarova G.A., Kirpichnikov M.P., Shaytan A.K. (2021) **Histone dynamics mediate DNA unwrapping and sliding in nucleosomes** *Nat Commun* **12**
- Bauer I. *et al.* (2012) **The NAD⁺-dependent histone deacetylase SIRT6 promotes cytokine production and migration in pancreatic cancer cells by regulating Ca²⁺ responses** *J Biol Chem* **287**:40924–40937
- Cen Y., Youn D.Y., Sauve A.A. (2011) **Advances in characterization of human sirtuin isoforms: chemistries, targets and therapeutic applications** *Curr Med Chem* **18**:1919–1935
- Chio U.S., Rechiche O., Bryll A.R., Zhu J., Leith E.M., Feldman J.L., Peterson C.L., Tan S., Armache J.P. (2023) **Cryo-EM structure of the human Sirtuin 6-nucleosome complex** *Sci Adv* **9**
- Croll T.I. (2018) **ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps** *Acta Crystallogr D Struct Biol* **74**:519–530
- Darden T., York D., Pedersen L. (1993) **Particle mesh Ewald: AnN-log(N) method for Ewald sums in large systems** *The Journal of Chemical Physics* **98**:10089–10092
- Dodonova S.O., Zhu F., Dienemann C., Taipale J., Cramer P. (2020) **Nucleosome-bound SOX2 and SOX11 structures elucidate pioneer factor function** *Nature* **580**:669–672
- Dyer P.N., Edayathumangalam R.S., White C.L., Bao Y., Chakravarthy S., Muthurajan U.M., Luger K. (2004) **Reconstitution of nucleosome core particles from recombinant histones and DNA** *Methods Enzymol* **375**:23–44
- Farnung L., Vos S.M., Wigge C., Cramer P. (2017) **Nucleosome-Chd1 structure and implications for chromatin remodelling** *Nature* **550**:539–542
- Fleetwood O., Kasimova M.A., Westerlund A.M., Delemotte L. (2020) **Molecular Insights from Conformational Ensembles via Machine Learning** *Biophysical Journal* **118**:765–780
- Gil R., Barth S., Kanfi Y., Cohen H.Y. (2013) **SIRT6 exhibits nucleosome-dependent deacetylase activity** *Nucleic Acids Res* **41**:8537–8545
- Goddard T.D., Huang C.C., Meng E.C., Pettersen E.F., Couch G.S., Morris J.H., Ferrin T.E. (2018) **UCSF ChimeraX: Meeting modern challenges in visualization and analysis** *Protein Sci* **27**:14–25
- Hopkins C.W., Le Grand S., Walker R.C., Roitberg A.E. (2015) **Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning** *J Chem Theory Comput* **11**:1864–1874
- Ikebe J., Sakuraba S., Kono H. (2016) **H3 Histone Tail Conformation within the Nucleosome and the Impact of K14 Acetylation Studied Using Enhanced Sampling Simulation** *PLoS Comput Biol* **12**
- Imai S., Armstrong C.M., Kaeberlein M., Guarente L. (2000) **Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase** *Nature* **403**:795–800

- Ivani I. *et al.* (2016) **Parmbcs1: a refined force field for DNA simulations** *Nat Methods* **13**:55–58
- Jiang H. *et al.* (2013) **SIRT6 regulates TNF-alpha secretion through hydrolysis of long-chain fatty acyl lysine** *Nature* **496**:110–113
- Kanfi Y., Naiman S., Amir G., Peshti V., Zinman G., Nahum L., Bar-Joseph Z., Cohen H.Y. (2012) **The sirtuin SIRT6 regulates lifespan in male mice** *Nature* **483**:218–221
- Kawahara T.L. *et al.* (2009) **SIRT6 links histone H3 lysine 9 deacetylation to NF-kappaB-dependent gene expression and organismal life span** *Cell* **136**:62–74
- Khongkow M. *et al.* (2013) **SIRT6 modulates paclitaxel and epirubicin resistance and survival in breast cancer** *Carcinogenesis* **34**:1476–1486
- Kräutler V., van Gunsteren W.F., Hünenberger P.H. (2001) **A fast SHAKE algorithm to solve distance constraint equations for small molecules in molecular dynamics simulations** *Journal of Computational Chemistry* **22**:501–508
- Liebschner D. *et al.* (2019) **Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix** *Acta Crystallogr D Struct Biol* **75**:861–877
- Liu W.H., Zheng J., Feldman J.L., Klein M.A., Kuznetsov V.I., Peterson C.L., Griffin P.R., Denu J.M. (2020) **Multivalent interactions drive nucleosome binding and efficient chromatin deacetylation by SIRT6** *Nat Commun* **11**
- Liu Y., Xie Q.R., Wang B., Shao J., Zhang T., Liu T., Huang G., Xia W. (2013) **Inhibition of SIRT6 in prostate cancer reduces cell viability and increases sensitivity to chemotherapeutics** *Protein Cell* **4**:702–710
- Luger K., Rechsteiner T.J., Richmond T.J. (1999) **Expression and purification of recombinant histones and nucleosome reconstitution** *Methods Mol Biol* **119**:1–16
- Macchiagodena M., Pagliai M., Andreini C., Rosato A., Procacci P. (2020) **Upgraded AMBER Force Field for Zinc-Binding Residues and Ligands for Predicting Structural Properties and Binding Affinities in Zinc-Proteins** *ACS Omega* **5**:15301–15310
- Maier J.A., Martinez C., Kasavajhala K., Wickstrom L., Hauser K.E., Simmerling C. (2015) **ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB** *J Chem Theory Comput* **11**:3696–3713
- Mao Z., Hine C., Tian X., Van Meter M., Au M., Vaidya A., Seluanov A., Gorbunova V. (2011) **SIRT6 promotes DNA repair under stress by activating PARP1** *Science* **332**:1443–1446
- Marquardt J.U. *et al.* (2013) **Sirtuin-6-dependent genetic and epigenetic alterations are associated with poor clinical outcome in hepatocellular carcinoma patients** *Hepatology* **58**:1054–1064
- McCord R.A. *et al.* (2009) **SIRT6 stabilizes DNA-dependent protein kinase at chromatin for DNA double-strand break repair** *Aging (Albany NY)* **1**:109–121
- McGinty R.K., Henrici R.C., Tan S. (2014) **Crystal structure of the PRC1 ubiquitylation module bound to the nucleosome** *Nature* **514**:591–596

- McGinty R.K., Tan S. (2016) **Recognition of the nucleosome by chromatin factors and enzymes** *Curr Opin Struct Biol* **37**:54–61
- Michishita E. *et al.* (2008) **SIRT6 is a histone H3 lysine 9 deacetylase that modulates telomeric chromatin** *Nature* **452**:492–496
- Michishita E., Park J.Y., Burneskis J.M., Barrett J.C., Horikawa I. (2005) **Evolutionarily conserved and nonconserved cellular localizations and functions of human SIRT proteins** *Mol Biol Cell* **16**:4623–4635
- Min L. *et al.* (2012) **Liver cancer initiation is controlled by AP-1 through SIRT6-dependent inhibition of survivin** *Nat Cell Biol* **14**:1203–1211
- Morrison E.A., Bowerman S., Sylvers K.L., Wereszczynski J., Musselman C.A. (2018) **The conformation of the histone H3 tail inhibits association of the BPTF PHD finger with the nucleosome** *Elife* **7**
- Ngo T.T., Zhang Q., Zhou R., Yodh J.G., Ha T. (2015) **Asymmetric unwrapping of nucleosomes under tension directed by DNA local flexibility** *Cell* **160**:1135–1144
- Olsson M.H., Sondergaard C.R., Rostkowski M., Jensen J.H. (2011) **PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions** *J Chem Theory Comput* **7**:525–537
- Pan P.W., Feldman J.L., Devries M.K., Dong A., Edwards A.M., Denu J.M. (2011) **Structure and biochemical functions of SIRT6** *J Biol Chem* **286**:14575–14587
- Papai G., Frechard A., Kolesnikova O., Crucifix C., Schultz P., Ben-Shem A. (2020) **Structure of SAGA and mechanism of TBP deposition on gene promoters** *Nature* **577**:711–716
- Pettersen E.F., Goddard T.D., Huang C.C., Couch G.S., Greenblatt D.M., Meng E.C., Ferrin T.E. (2004) **UCSF Chimera--a visualization system for exploratory research and analysis** *J Comput Chem* **25**:1605–1612
- Punjani A., Rubinstein J.L., Fleet D.J., Brubaker M.A. (2017) **cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination** *Nat Methods* **14**:290–296
- Roichman A. *et al.* (2021) **Restoration of energy homeostasis by SIRT6 extends healthy lifespan** *Nat Commun* **12**
- Sanchez-Garcia R., Gomez-Blanco J., Cuervo A., Carazo J.M., Sorzano C.O.S., Vargas J. (2021) **DeepEMhancer: a deep learning solution for cryo-EM volume post-processing** *Commun Biol* **4**
- Sebastian C. *et al.* (2012) **The histone deacetylase SIRT6 is a tumor suppressor that controls cancer metabolism** *Cell* **151**:1185–1199
- Stutzer A., Liokatis S., Kiesel A., Schwarzer D., Sprangers R., Soding J., Selenko P., Fischle W. (2016) **Modulations of DNA Contacts by Linker Histones and Post-translational Modifications Determine the Mobility and Modifiability of Nucleosomal H3 Tails** *Mol Cell* **61**:247–259
- Taylor J.R. *et al.* (2022) **Sirt6 regulates lifespan in Drosophila melanogaster** *Proc Natl Acad Sci U S A* **119**

- Tegunov D., Cramer P. (2019) **Real-time cryo-electron microscopy data preprocessing with Warp** *Nat Methods* **16**:1146–1152
- Tian X. *et al.* (2019) **SIRT6 Is Responsible for More Efficient DNA Double-Strand Break Repair in Long-Lived Species** *Cell* **177**:622–638
- Vitiello M., Zullo A., Servillo L., Mancini F.P., Borriello A., Giovane A., Della Ragione F., D’Onofrio N., Balestrieri M.L. (2017) **Multiple pathways of SIRT6 at the crossroads in the control of longevity, cancer, and cardiovascular diseases** *Ageing Res Rev* **35**:301–311
- Wagner T. *et al.* (2019) **SPHIRE-crYOLO is a fast and accurate fully automated particle picker for cryo-EM** *Commun Biol* **2**
- Wang W.W., Zeng Y., Wu B., Deiters A., Liu W.R. (2016) **A Chemical Biology Approach to Reveal Sirt6-targeted Histone H3 Sites in Nucleosomes** *ACS Chem Biol* **11**:1973–1981
- Wang Z.A., Markert J.W., Whedon S.D., Yapa Abeywardana M., Lee K., Jiang H., Suarez C., Lin H., Farnung L., Cole P.A. (2023) **Structural Basis of Sirtuin 6-Catalyzed Nucleosome Deacetylation** *J Am Chem Soc* **145**:6811–6822
- Watroba M., Dudek I., Skoda M., Stangret A., Rzedkiewicz P., Szukiewicz D. (2017) **Sirtuins, epigenetics and longevity** *Ageing Res Rev* **40**:11–19
- Yoo J., Aksimentiev A. (2018) **New tricks for old dogs: improving the accuracy of biomolecular force fields by pair-specific corrections to non-bonded interactions** *Phys Chem Chem Phys* **20**:8432–8449
- Zhang W. *et al.* (2018) **SIRT6 deficiency results in developmental retardation in cynomolgus monkeys** *Nature* **560**:661–665
- Zhong E.D., Bepler T., Berger B., Davis J.H. (2021) **CryoDRGN: reconstruction of heterogeneous cryo-EM structures using neural networks** *Nat Methods* **18**:176–185
- Zhong L. *et al.* (2010) **The histone deacetylase Sirt6 regulates glucose homeostasis via Hif1alpha** *Cell* **140**:280–293
- Zivanov J., Nakane T., Forsberg B.O., Kimanius D., Hagen W.J., Lindahl E., Scheres S.H. (2018) **New tools for automated high-resolution cryo-EM structure determination in RELION-3** *Elife* **7**

Article and author information

Ekaterina Smirnova

Department of Integrated Structural Biology, IGBMC, Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), 67400 Illkirch, France; Université de Strasbourg, IGBMC UMR 7104-UMR-S 1258, 67400 Illkirch, France; CNRS, UMR 7104, 67400 Illkirch, France; Inserm, UMR-S 1258, 67400 Illkirch, France; Equipe labellisée Ligue Contre le Cancer

Emmanuelle Bignon

Université de Lorraine and CNRS, UMR 7019 LPCT, F-54000 Nancy, France
ORCID iD: [0000-0001-9475-5049](https://orcid.org/0000-0001-9475-5049)

Patrick Schultz

Department of Integrated Structural Biology, IGBMC, Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), 67400 Illkirch, France; Université de Strasbourg, IGBMC UMR 7104-UMR-S 1258, 67400 Illkirch, France; CNRS, UMR 7104, 67400 Illkirch, France; Inserm, UMR-S 1258, 67400 Illkirch, France; Equipe labellisée Ligue Contre le Cancer
ORCID iD: [0000-0002-7310-6186](https://orcid.org/0000-0002-7310-6186)

Gabor Papai

Department of Integrated Structural Biology, IGBMC, Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), 67400 Illkirch, France; Université de Strasbourg, IGBMC UMR 7104-UMR-S 1258, 67400 Illkirch, France; CNRS, UMR 7104, 67400 Illkirch, France; Inserm, UMR-S 1258, 67400 Illkirch, France; Equipe labellisée Ligue Contre le Cancer
For correspondence: papai@igbmc.fr
ORCID iD: [0000-0002-2779-8679](https://orcid.org/0000-0002-2779-8679)

Adam Ben-Shem

Department of Integrated Structural Biology, IGBMC, Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), 67400 Illkirch, France; Université de Strasbourg, IGBMC UMR 7104-UMR-S 1258, 67400 Illkirch, France; CNRS, UMR 7104, 67400 Illkirch, France; Inserm, UMR-S 1258, 67400 Illkirch, France; Equipe labellisée Ligue Contre le Cancer
For correspondence: adam@igbmc.fr
ORCID iD: [0000-0003-0297-8662](https://orcid.org/0000-0003-0297-8662)

Copyright

© 2023, Smirnova et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Tim Formosa

University of Utah School of Medicine, Salt Lake City, United States of America

Senior Editor

Merritt Maduke

Stanford University, Stanford, United States of America

Joint Public Review:

Smirnova et al. present a cryo-EM structure of human SIRT6 bound to a nucleosome as well as the results from molecular dynamics simulations. The results show that the combined conformational flexibilities of SIRT6 and the N-terminal tail of histone H3 limit the residues with access to the active site, partially explaining the substrate specificity of this sirtuin-class histone deacetylase. Two other groups have recently published cryo-EM structures of SIRT6:nucleosome complexes; this manuscript confirms and complements these previous findings, with the addition of some novel insights into the role of structural flexibility in substrate selection.

This manuscript is the third in a recent series of reports of cryo-EM structures of Sirt6:nucleosome complexes. The main conclusions of the three studies are similar, but this manuscript from Smirnova et al. includes additional molecular dynamics analysis of the histone tails. These studies suggest that part of the specificity for sites on the H3 tail is the result of only this tail having significant access to the active site. The results are partially validated by showing that H3-K27Ac is sometimes found near the active site in the simulations, and is a weak substrate for the deacetylase in vitro. All of the structures show Sirt6 contacting the acidic patch of H2A-H2B, partial displacement of the H2A C-terminal tail, and displacement of the DNA at the entry-exit site to "unclamp" the H3 N-terminal tail. This manuscript provides additional support for the conclusions drawn in the first two published structures, adds molecular dynamics simulations that provide further insight and includes a biochemical assay that helps to resolve an apparent conflict regarding the deacetylation of H3-K27Ac from the other two papers.

- <https://doi.org/10.7554/eLife.87989.2.sa0>

Author Response

The following is the authors' response to the original reviews.

We are grateful to the reviewers for their remarks which significantly improved the paper. Following these remarks we completed the analysis and validation of our cryo-EM data and performed several biochemical tests to support our conclusions, lending credibility to the paper. Please find our detailed answers below each recommendation of the reviewers.

Major recommendations

1. *Errors and omissions in the presentation make the manuscript difficult to access.*

a) *The text should be edited for grammatical errors more carefully*

- We corrected the grammatical errors.

b) *Figures should be labeled to allow the reader to follow the logic of the presentation and identify the features being discussed. Identification through the color coding (the identity of the histones, the location of zinc fingers, the active site, and so on) would be helpful.*

- We labeled the Rossman fold and Zn-finger domains in Figure 1 and described the histone color codes. The active site of SIRT6 is depicted in Figure 4.

1. *The recent publications from the Farnung/Cole and Peterson/Tan/Armache labs need to be cited and the results from Smirnova et al. compared and contrasted with those publications explicitly.*

- We added the following paragraph to the discussion section:
 "While this manuscript was under review two studies describing the structure of SIRT6-NCP appeared in press (Wang et al., 2023 ; Chio et al., 2023). The conclusion of these papers regarding the position of SIRT6 on the nucleosome and the unwinding of DNA by the enzyme are similar to our findings. We however dissected in addition the movements of SIRT6 on the nucleosome and analyzed via molecular dynamics the conformations of the H3 tail with respect to the SIRT6 active site. Our results point to

the importance of the flexibility between the globular domains of SIRT6 and also explain how SIRT6 can access lysines that are much closer to the histone core than H3K9.”

a) Notably, the Peterson/Tan/Armache labs suggest that H3K27 cannot be deacetylated by SIRT6 whereas the Farnung/Cole labs show deacetylation of H3K27 by SIRT6. Do the results of the Smirnova et al. structure help to resolve this situation?

- We performed deacetylation tests of H3K27Ac nucleosomes and show that SIRT6 deacetylate H3K27Ac albeit at somewhat lower efficiency than H3K9Ac. Our molecular dynamics simulations explain how H3K27, which is close to the histone core, can still be reached by SIRT6 active site. We added the following text to the paper: “To lend support to this claim we tested whether SIRT6 can deacetylate residue H3K27 that was first acetylated by SAGA (Supplemental Fig. 7c). We find that indeed SIRT6 could efficiently deactylate H3K27Ac, although at a somewhat slower rate than H3K9Ac. We conclude that partial DNA unwrapping by SIRT6 allows H3-tail conformations that make lysines that are close to the core of H3 accessible to the enzyme.”

b) The Farnung/Cole labs have visualized an intermediate state of deacetylation. How does this compare to the structure presented in this manuscript? Addressing these points would facilitate further research and discussion in the community.

- We believe the resolution of the SIRT6 Rossmann fold precludes addressing these points.

c) Can the authors exclude the possibility that the additional density observed in Supplemental Figure 6 is not coming from the H3 tail, as observed in the two other structures?

- One density is the continuation of the H2A histone tail. We strongly believe that this density corresponds to this tail. The other density indeed can originate from the H3 tail. Therefore, we didn’t model anything inside it.

d) It would be useful to comment on how much flexibility has been observed in the other structures for the SIRT6 interaction with the acidic patch, and also how other acidic-patch binding proteins compare with the results here.

- We refrain from estimating the flexibility observed in the other structures as no such analysis is provided by these papers. Regarding the interaction with the acidic patch we mention that R175 packs against H2B L103 and serves as a classical “arginine anchor motif” and refer the reader to a review on the topic.

e) Does the presence or absence of NAD⁺ affect the comparisons among the structures?

- NAD⁺ binding might affect the fine structure of the active site although NAD⁺ was not observed in crystal stuctures of SIRT6 in its presence. The resolution of this part precludes further addressing this issue.

1. The lack of biochemical validation of conclusions should be acknowledged and the reasoning behind this choice discussed.

- We added experiments to validate our conclusions with biochemical tests. We produced nucleosomes with acetylated histone H3 by employing purified SAGA acetyltransferase complex. We isolated SIRT6 where the four residues implicated in interactions with the acidic patch are mutated to alanines (SIRT6-4A). We show that this mutant has very weak interaction with the nucleosome and much lower H3K9Ac deacetylation activity than WT. Similarly SIRT6-3A with mutations in the residues we suggest involved in binding to nucleosomal DNA also shows weak activity and binding to the nucleosome. We added Supplement Figure 7 that depicts the results of these experiments and embedded reference to these results in the appropriate sections of the text. Furthermore, we also show that SIRT6 is active in deacetylating H3K27Ac. This supports our molecular dynamics simulations showing that when SIRT6 binds the nucleosome, H3 tail can assume conformations where H3K27 is accessible by the enzyme's active site. These results also appear in Supplement Figure 7.

1. *The authors nicely analyze and discuss the conformational flexibility of SIRT6 binding. This is an interesting finding, but Fig. 2 does not adequately convey this flexibility.*

- We now considerably improved Figure 2. We added panels c and f which depict clearly the movements we observe.

1. *The authors need to explain why two cryo-EM datasets were collected but were not merged, and the labeling of the datasets in the Supplemental Table appear to be switched.*

- The two datasets were collected with two very different pixel spacing therefore merging the two was possible only in Relion. This process, however, did not improve the resolution of the SIRT6's Rossmann fold domain. We thank the reviewer to notice the discrepancy in the text and the Supplemental Table 1, it was corrected.

1. *Supplemental Figure 4 should be expanded to show additional representative densities with the respective fit of the model. This will allow the reader to better judge the quality of the data. At least the acidic patch interaction, the DNA-SIRT6 interactions, and the H2A should be shown in this context.*

- To illustrate the high-resolution features of the structure as well as the key regions we added Supplemental Figure 4.

1. *Standard elements of data analysis and validation should be included (angular distribution plots for cryo-EM reconstructions, a 3D FSC sphericity plot, a Q-score and EMRinger score for the cryo-EM data and atomic model, a model-to-map FSC curve). In general, model building is poorly described as it is unclear which maps (or to what degree different maps) were used for this process. This should be clarified in the methods section and in the Supplemental Table 1.*

- The model validation and data analysis details were added to Supplemental Figures 2 and 3 as well as in Supplemental Table 1.

1. The provided maps also do not fully recapitulate the path of the H2A tail. The various density maps and PDB provided for this review do not support the final modeled residues of H2A between residues #118/119-123. This affects the validity of figure 3E and the discussion of the proximity of the potential substrates to the active site. The authors should clarify how they inferred that this is the H2A tail rather than the loosely bound SIRT6 Nterminal loop (whose stability could be altered by the presence or absence of NAD+) as suggested by overlaying the relevant crystal structures.

- We added a panel to Supplemental Figure 4 (d) depicting the density where the H2A tail was modelled.

1. The authors should explain how the data produced an asymmetrically oriented complex with a single SIRT6 molecule bound to one face. Were complexes with two SIRT6 molecules excluded? Is supplementary figure 4A the basis for the orientation and is this sufficient for this purpose?

- Complexes with two SIRT6 molecules were present but only at around 1.5 percent of the whole dataset. These images were excluded from the refinement (shown in Supplementary Figure 2). The DNA orientation is depicted in Supplementary Figure 5A. The resolution obtained at the dyad (~2.5Å) allowed us to distinguish purine and pyrimidine bases. The Widom 601 sequence is asymmetric and the densities clearly show that there is only one orientation of the DNA observed with respect to SIRT6.

1. The authors should clarify how supplemental figure 4B supports the conclusion that DNA is unwrapped. The density is not readily visible and docking of a simple DNA model in the ZN-focused map does not clearly rule out the possibility that this density comes from the H3 N-terminal tail.

- We added to this figure the cryo-EM densities used to model the DNA path and the orientation of SIRT6. This image is now Supplemental Figure 5c.

Minor recommendations

1. The scale bar is missing for the 2D classes shown in Supplemental Figure 2.

- We added the scale bar to the image depicting the 2D classes in Supplemental Figure 2.

1. Masked classifications should be shown in the classification tree (Supplemental Figure 2 +3) with the masks shown as a transparent volume.

- We now show the mask used for the 3D classifications of the SIRT6's Rossman fold domain in Supplemental Figure 2.

1. Supplemental Figure 3 should show the indicated 3D classifications in the classification tree.

- We added the 3D classifications in Supplemental Figure 3.

1. *The authors should consider applying local CTF refinement and particle polishing to improve their resolution.*

- We did local and global CTF refinements. Polishing didn't improve the resolution as movie frame alignment was done outside of Relion.

1. *The descriptions of the Widome 601 sequence orientation should be less ambiguous, perhaps mentioning the AT-rich and AT-poor arms instead of left and right arms.*

- We corrected the text as required.

1. *The statement "Such a large change in DNA trajectory is reminiscent of the chromatin-remodeler ATPases or pioneer transcription factors binding to nucleosome but was not observed in other histone modifiers" requires a citation.*

- We added appropriate references.

1. *The authors should provide a supplemental figure of the nucleosome-SIRT6 and PRC1-nucleosome structure comparison to complement the discussion section.*

- We refer the reader to the paper describing the PRC1-nucleosome structure.