

Thalamic regulation of ocular dominance plasticity in adult visual cortex

Reviewed Preprint

Revised by authors after peer review.

[About eLife's process](#)**Reviewed preprint version 2**

September 12, 2023 (this version)

Reviewed preprint version 1

June 22, 2023

Sent for peer review

April 10, 2023

Posted to bioRxiv

March 25, 2023

Yi Qin, Mehran Ahmadiou, Samuel Suhai, Paul Neering, Leander de Kraker, J. Alexander Heimel, Christiaan N. Levelt 

Molecular Visual Plasticity Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands • University of Strasbourg, Strasbourg, France • Circuits, Structure and Function Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands • Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive Research, VU University Amsterdam, de Boelelaan 1085, 1081 HV, the Netherlands

 https://en.wikipedia.org/wiki/Open_access

 <https://creativecommons.org/licenses/by/4.0/>

Abstract

Summary

Experience-dependent plasticity in the adult visual system is generally thought of as a cortical process. However, several recent studies have shown that perceptual learning or monocular deprivation can also induce plasticity in the adult dorsolateral geniculate nucleus (dLGN) of the thalamus. How plasticity in the thalamus and cortex interact in the adult visual system is ill understood. To assess the influence of thalamic plasticity on plasticity in primary visual cortex (V1), we made use of our previous finding that during the critical period, ocular dominance (OD) plasticity occurs in dLGN and requires thalamic synaptic inhibition. Using multielectrode recordings we find that this is also true in adult mice, and that in the absence of thalamic inhibition and plasticity, OD plasticity in adult V1 is absent. To study the influence of V1 on thalamic plasticity we silenced V1 and show that during the critical period, but not in adulthood, the OD shift in dLGN is partially caused by feedback from V1. We conclude that during adulthood, the thalamus plays an unexpectedly dominant role in experience-dependent plasticity in V1. Our findings highlight the importance of considering the thalamus as a potential source of plasticity in learning events that are typically thought of as cortical processes.

eLife assessment

This is an **important** study that demonstrates that plasticity of ocular dominance of binocular neurons in the visual thalamus persists in adulthood. The evidence supporting the authors' conclusion is **convincing**, and the findings are an **important** contribution to a growing body of work identifying plasticity in the adult visual system. This work will interest those in the field of ocular dominance plasticity in the visual system as well as scientists investigating the function of synaptic plasticity in the brain.

Introduction

Experience-dependent plasticity in the adult visual system is largely thought of as a cortical process (Gilbert et al., 2009 [↗](#); Gilbert and Wiesel, 1992 [↗](#)). However, several recent studies have demonstrated that plasticity also occurs in the adult dorsal lateral geniculate nucleus (dLGN) of the thalamus of human subjects during perceptual learning (Yu et al., 2016 [↗](#)) and mice upon monocular deprivation (MD) (Huh et al., 2020 [↗](#); Jaepel et al., 2017 [↗](#)). How plasticity in adult dLGN is regulated and whether plasticity in dLGN and V1 influence each other has barely been studied.

In this study, we addressed these questions using ocular dominance (OD) plasticity in mice as a model. OD is the property that neurons preferentially respond to visual stimuli presented to one eye versus the other (Wiesel and Hubel, 1963a [↗](#)). Visual experience affects OD and a period of MD results in an OD shift in V1 neurons due to weakened responses to the deprived eye and strengthened responses to the non-deprived eye (Hensch and Stryker, 1996 [↗](#); Wiesel and Hubel, 1963a [↗](#)). OD plasticity is most prominent during a critical period of development (Gordon and Stryker, 1996 [↗](#); Wiesel and Hubel, 1963a [↗](#)) but can also be induced in young adult mice (Heimel et al., 2007 [↗](#); Hofer et al., 2006 [↗](#); Lehmann and Löwel, 2008 [↗](#); Sato and Stryker, 2008 [↗](#); Sawtell et al., 2003 [↗](#)). This requires a longer period of deprivation, however, and the shift is smaller and less persistent and is mediated predominantly by strengthening of responses to the non-deprived eye (Frenkel and Bear, 2004 [↗](#); Heimel et al., 2007 [↗](#); Hofer et al., 2006 [↗](#); Lehmann and Löwel, 2008 [↗](#); Sato and Stryker, 2008 [↗](#)).

Previously, we demonstrated that during the critical period, extensive OD plasticity can be induced in dLGN and that this requires synaptic inhibition in the thalamus (Sommeijer et al., 2017 [↗](#)). Multielectrode recordings revealed that OD plasticity in dLGN is strongly reduced in mice in which thalamic synaptic inhibition is inactivated by deleting the gene encoding the GABA receptor alpha1 subunit (*Gabra1*) selectively in the dorsal thalamus (*Gabra1^{fl-hom} x Olig3-cre⁺* mice, hereafter referred to as “KO mice”). Interestingly, OD plasticity induced by long-term monocular deprivation (MD) is also reduced in V1 of these mice due to the absence of ipsilateral eye response strengthening (Sommeijer et al., 2017 [↗](#)), suggesting that during the critical period, thalamic plasticity contributes to plasticity in V1. Here, we investigated how dLGN and V1 influence each other during OD plasticity in adulthood. We find that in adult mice lacking thalamic synaptic inhibition, OD plasticity is absent in both dLGN and V1. Silencing V1 of adult wild-type (WT) mice does not affect the OD shift in dLGN, showing that it does not depend on feedback from V1. In contrast, we find that during the critical period, the OD shift in dLGN is partially depends on activity in V1. Together, our findings show that thalamocortical interactions underlying OD plasticity change with age and suggest that the thalamus may be an important source of plasticity in adult learning events that have generally been considered cortical processes.

Visual responses in adult dLGN of WT and KO mice

Previous work showed that during the critical period, visual responses of dLGN neurons in $Gabra1^{fl-hom} \times olig3-cre^+$ KO mice were less sustained due to the lack of thalamic synaptic inhibition, while average response strengths and basic receptive field properties seemed surprisingly unaffected (Sommeijer et al., 2017 [↗](#)).

To assess whether this situation remained similar in adulthood, we measured visual responses in dLGN using a 16-channel silicon probe in anesthetized KO mice and $Gabra1^{fl-hom} \times olig3-cre^-$ (WT) siblings. Recordings were performed in the ipsilateral projection zone of dLGN (Fig. 1A [↗](#)). Receptive field sizes and positions were determined by presenting white squares (5 deg) at random positions on a black background (Fig. 1B [↗](#)). We only included recordings from channels with receptive fields corresponding to the central 30° of the visual field. We observed no differences in receptive field sizes in KO and WT mice (Fig. 1C [↗](#)). As receptive field sizes in dLGN are known to become smaller between eye opening and critical period onset (Tschetter et al., 2018 [↗](#)), this observation suggests that dLGN develops surprisingly normally in the absence of synaptic inhibition. To investigate this further, we analyzed the numbers, densities and sizes of inhibitory and cholinergic boutons, which are also known to increase during the same developmental window (Bickford et al., 2010 [↗](#), Sommeijer et al. 2017 [↗](#), Sokhadze et al., 2018 [↗](#)). Again, in adult WT and KO mice we observed no differences (Suppl. Fig. 1 [↗](#)).

To record visual responses of dLGN neurons to the contra- or ipsilateral eye separately, the other eye was closed and visual stimuli (full screen, full contrast black/white reversals, at 3s intervals) were presented (Fig. 1B [↗](#)). We selected single units from non-deprived WT and KO mice and assessed their responses to the contra- and ipsilateral eye. Examples of monocular and binocular single units are shown in Figure 1D [↗](#). In WT mice, 51% of recorded neurons were binocular and in KO mice 59%. To assess whether the temporal profile of visual responses differed in adult WT and KO mice, we compared the areas under the curve (AUC) of the peristimulus time histogram (PSTH) during different time bins (Fig. 1E [↗](#)). This revealed that in KO mice, visual responses attenuated faster than in wild-type siblings: peak responses were stronger, but weaker responses were observed during the phase of prolonged firing (Fig. 1E [↗](#)). Thus, while average response strength in KO mice was similar to that in WT mice, the attenuation index was increased. These results show that like the situation during the critical period (Sommeijer et al., 2017 [↗](#)), visual responses in dLGN neurons in adult KO and WT mice mostly differ in their temporal profile, while average response strengths and receptive fields sizes are hardly changed.

OD plasticity in dLGN is reduced in adult mice lacking thalamic synaptic inhibition

We then continued experiments to assess OD plasticity in the dLGN of WT and KO mice. We monocularly deprived adult WT and KO mice for 7 days, long enough to induce an OD shift in adult V1 (Frenkel and Bear, 2004 [↗](#); Heimel et al., 2007 [↗](#); Hofer et al., 2006 [↗](#); Lehmann and Löwel, 2008 [↗](#); Sato and Stryker, 2008 [↗](#)), by suturing one eye closed. We recorded responses to the ipsi- and contralateral eye in dLGN neurons (Fig. 2A [↗](#)) and calculated the OD index (ODI) of all units recorded in monocularly deprived and non-deprived KO and WT mice and averaged them to obtain an OD score (Fig. 2B [↗](#)). We found that after one week of MD, a significant OD shift occurred in dLGN of adult WT mice. This was predominantly caused by a significant increase in the responses to the non-deprived ipsilateral eye (Fig. 2C [↗](#)). In KO mice, no OD shift could be induced in dLGN (Fig. 2B [↗](#)) and no significant changes were observed in the responses to the ipsi- or contralateral eye (Fig. 2D). Together, these results show that also in adulthood, OD plasticity in dLGN depends on thalamic synaptic inhibition.




Fig. 1

| Visual responses of dLGN neurons in mice lacking thalamic Gabra1

A, Recording electrodes were placed in the ipsilateral projection zone of dLGN (see green fluorescent trace of actual electrode penetration in dLGN). All receptive field centers of multi-units recorded in wild-type (WT, blue) and knockout (KO, green) mice ($n=61$ units from 13 non-deprived or monocularly deprived (MD) mice and $n=80$ units from 18 NO MD or MD mice). Nose position is at 0 degrees horizontally and vertically. The black dashed lines indicate -30 degree and $+30$ degree horizontal angles. **B**, Experimental setup to measure receptive field (RF) and single eye responses. **C**, RF sizes of multi-units in NO MD and MD (shaded area) KO and WT mice do not differ (two-way ANOVA, interaction of genotype with MD: $P=0.07$; Tukey's post-hoc test; WT NO MD vs KO NO MD: $P=0.19$; WT MD vs KO MD: $P=0.11$). **D**, Examples of dLGN neuron responses to full screen OFF-ON flash stimuli in WT (blue) and KO (green) mice. Colored and black lines indicate responses of contra- and ipsilateral eyes, respectively. Waveforms of each unit responding to the contra- or ipsilateral eye are shown in the upper right corner. WT: left panel is a monocular unit, right panels are binocular units. KO: left two panels are monocular units, right panel is a binocular unit (Zeta-test). **E**, Left, average responses of contralateral eye in WT (blue) and KO (green) mice. KO mice show higher peak (90-110ms) and lower prolonged responses (200-230ms). (Repeated measure two-way ANOVA, interaction of genotype with time, $P=0.0001$; Post-hoc, Fisher's LSD test). Right, attenuation index of visual responses in WT and KO mice.




Fig. 2

| Loss of OD plasticity in dLGN of mice lacking thalamic synaptic inhibition

A, Illustration of the experiment design. In experiments, four groups of animals were used: deprived (MD) or non-deprived (NO MD) wild-type and knockout mice. Mice in the MD group had the eyelids of the eye contralateral to the recording side sutured 7 for days. **B**, 7 days of MD reduces the ODI in WT mice but not in KO animals (interaction of genotype with MD: two-way ANOVA, $P=0.046$, Tukey's post-hoc test; WT NO MD vs. WT MD, $P=0.040$; WT NO MD, $n=40$ units, 7 mice; WT MD, $n=22$ units, 6 mice; KO NO MD, $n=45$ units, 9 mice; KO MD, $n=34$ units, 9 mice). **C**, In WT mice, responses to the ipsilateral eye are significantly increased after 7-d MD. Responses to the contralateral eye are unchanged (Mann-Whitney; contralateral, NO MD vs. MD, $P=0.29$; ipsilateral, NO MD vs. MD, $P=0.032$). **D**, In KO mice, MD causes no significant changes in responses to either the contralateral or the ipsilateral eye (Mann-Whitney; contralateral, NO MD vs. MD, $P=0.73$; ipsilateral, NO MD vs. MD, $P=0.59$).

OD plasticity in adult V1 is reduced in mice lacking thalamic synaptic inhibition

During the critical period, OD plasticity in V1 is partially deficient in KO mice. Brief MD induces a normal OD shift, but longer MD does not cause the OD shift to strengthen further (Sommeijer et al., 2017). This suggests that the critical period opens normally in V1 of KO mice, but that the second, homeostatic phase of the OD shift depends on thalamic inhibition and plasticity. Residual OD plasticity in adult V1 has various similarities with the second phase of OD plasticity during the critical period and also requires long term MD. We were therefore investigated whether OD plasticity in V1 was deficient in adult KO mice.

To address this question, we monocularly deprived WT and KO mice for 7 days, after which we recorded visual responses from single units in V1 of these mice and of normally sighted siblings (Fig. 3A). Again, we only included channels with receptive fields within the central 30° of the visual field to ascertain we recorded from binocular V1. Like in dLGN, receptive field sizes did not differ between KO and WT mice and were not affected by a week of MD (Fig. 3B).

Visual responses to the contra- or ipsilateral eye were recorded in the same way as for dLGN, using the same visual stimuli (examples shown in Fig. 3C). Like in dLGN, visual responses in V1 were more attenuated in KO mice than in WT mice (Fig. 3D). Next, we calculated the OD index from all single units in the four groups (Fig. 3E). A clear OD shift was induced in V1 of monocularly deprived WT mice. As expected, the OD shift involved an increase in open, ipsilateral eye responses (Fig. 3F). We also found a significant decrease of deprived, contralateral eye responses (Fig. 3F). While several studies have provided evidence that a loss of contralateral eye responses contributes less to adult OD plasticity than during the critical period (Frenkel and Bear, 2004; Kalogeraki et al., 2017; Sato and Stryker, 2008), others have shown that it still occurs in adulthood (Rose et al., 2016). Possibly, the OD shift and the loss of deprived eye responses are more pronounced in our recordings due to them being limited to the center of the visual field or the use of flash stimuli instead of moving gratings. In adult KO mice, the OD shift after 7 days of MD was negligible and significantly smaller than in WT mice (Fig. 3G). The contra- and ipsilateral eye responses in V1 of non-deprived KO mice were of comparable strength as those observed in WT mice, despite the lack of synaptic inhibition in the thalamus. After 7 days of monocular deprivation, no significant strengthening of ipsilateral eye responses or weakening of deprived eye responses occurred (Fig. 3F). We conclude that in adult mice, absence of synaptic inhibition in the thalamus reduces OD plasticity in V1.

Effect of feedback from V1 to dLGN responses in the presence or absence of thalamic synaptic inhibition

These results so far show that OD plasticity in dLGN affects the OD shift in V1. OD plasticity in V1 may also influence the OD shift in dLGN. Apart from the retinal input dLGN relay cells receive, they are also strongly innervated by excitatory feedback connections from layer 6 cells in V1. Additionally, dLGN neurons receive bisynaptic inhibitory feedback from V1 via the thalamic reticular nucleus (TRN) and local interneurons (Fig. 4A). Depending on whether excitatory or inhibitory feedback dominates, responses of dLGN relay cells to the ipsi- or contralateral eye in dLGN may be strengthened or inhibited by V1 feedback (Denman and Contreras, 2015; Howarth et al., 2014; Jaepel et al., 2017; Kirchgeßner et al., 2020; Olsen et al., 2012). These feedback inputs from V1 can thus influence the OD of relay cells in dLGN.

To investigate how dLGN responses were influenced by V1 feedback and how synaptic thalamic inhibition affected this, we silenced V1 of WT and KO mice with the GABA-receptor agonist muscimol while recording from dLGN. Muscimol injections effectively silenced V1 (Fig. 4A). On average, silencing V1 did not alter responses to the contra- or ipsilateral eye in individual units in dLGN of adult WT mice (Fig. 4B-D). In KO mice, V1 silencing also did not significantly affect




Fig. 3

| Reduced OD plasticity in adult V1 lacking thalamic OD plasticity

A, Recording electrodes are located in binocular V1. All receptive field centers of multi-units recorded in WT (blue) and KO (green) mice ($n=112$ units from 13 NO MD or MD mice and $n=138$ units from 18 NO MD or MD mice). Nose position is at 0 degrees horizontally and vertically. The black dashed lines indicate -30 degree and $+30$ degree horizontal angles. **B**, RF sizes of units in KO and KO mice do not differ (interaction of genotype with MD: two-way ANOVA, $P=0.07$). **C**, Two examples of single unit responses in V1 of a WT (blue) and KO (green) mouse to the contra- and ipsilateral eyes to ON and OFF visual stimuli. Each stimulus lasted 3s. Colored and black lines indicate contra- and ipsilateral eye responses, respectively. **D**, Attenuation index of contralateral eye responses in V1 of WT and KO mice. **E**, 7 days of MD reduces the ODI in WT but not KO mice (interaction of genotype with MD: two-way ANOVA, $P<0.001$, Tukey's post-hoc test; WT NO MD vs. WT MD, $P<0.001$; WT NO MD, $n=71$ units, 7 mice; WT MD, $n=42$, 6 mice; KO NO MD, $n=63$ units, 9 mice; KO MD, $n=78$ units, 9 mice). **F**, In WT mice, responses to the contralateral eye are significantly reduced after 7-d MD, while those to the ipsilateral eye are significant increased (Mann-Whitney; contralateral, NO MD vs. MD, $P=0.0043$; ipsilateral, NO MD vs. MD, $P=0.0062$). **G**, In KO mice, MD causes no significant changes in responses to either the contralateral or the ipsilateral eye (Mann-Whitney; contralateral, NO MD vs. MD, $P=0.17$; ipsilateral, NO MD vs. MD, $P=0.66$).




Fig. 4

| Effect of feedback from V1 to dLGN responses

A, Left, illustration of corticothalamic-thalamocortical feedback network. dLGN is innervated by V1 and receives glutamatergic feedback. All these projections send excitatory collaterals to the thalamic reticular nucleus (TRN) which sends inhibitory inputs to dLGN. By muscimol injection in V1, corticothalamic projections are silenced. Right, V1 is effectively silenced by muscimol injection (Wilcoxon signed rank, $P < 0.001$, $n = 31$ mice). **B**, Examples of dLGN responses before and after muscimol injection in V1 of non-deprived WT mice. Waveforms of each unit are shown in upper right corner. Left and right panels correspond to contralateral and ipsilateral eye responses respectively. Dark and light lines represent responses before and after muscimol injection respectively. **C & D**, Silencing V1 feedback has no significant effect on contralateral (**C**) or ipsilateral (**D**) responses in WT mice (Wilcoxon signed rank; contralateral, WT NO MD vs. WT NO MD with muscimol, $P = 0.62$; ipsilateral, WT NO MD vs. WT NO MD with muscimol, $P = 0.94$, $n = 40$ units, 7 mice). **E**, Examples of dLGN responses before and after muscimol injection in V1 of non-deprived KO mice. Waveforms of each unit are shown in upper right corner. Left and right panels correspond to contralateral and ipsilateral eye responses respectively. Dark and light lines represent responses before and after muscimol injection respectively. **F & G**, There is no significant effect of V1 silencing on contralateral (**F**) or ipsilateral (**G**) eye responses in KO mice, but a trend towards decreased ipsilateral eye responses is present (Wilcoxon signed rank; contralateral, KO NO MD vs. KO NO MD with muscimol, $P = 0.19$; ipsilateral, KO NO MD vs. KO NO MD with muscimol, $P = 0.059$, $n = 45$ units, 9 mice).

responses to the contralateral eye (**Fig. 4E-G**). Responses to the ipsilateral eye showed a trend towards weakening after silencing V1, but this did not reach significance ($P=0.094$). Thus, also in the absence of synaptic inhibition in the thalamus, V1 feedback has relatively little influence on dLGN responses to the contra- or ipsilateral eye.

Feedback from V1 does not affect the OD shift in adult dLGN

Finally, we investigated whether feedback from V1 influenced the OD shift in dLGN of adult WT and KO mice. In non-deprived mice, the OD index did not change after silencing V1 of WT mice (**Fig. 5A**), as expected considering that contra- and ipsilateral eye response strengths were not affected by V1 feedback (**Fig. 4C, D**). Similarly, the OD index in non-deprived KO mice did not change upon silencing V1 (**Fig. 5B**), which was also expected based on the small changes in contra- and lateral eye responses that we observed (**Fig. 4F, G**)

Despite the considerable OD shift we observed in V1 of adult WT mice, silencing V1 did not affect the OD measured in dLGN (**Fig. 5C**). Also, average responses to the two eyes were not altered in the absence of cortical feedback (**Fig. 5D, E**). This confirms that the OD shift in adult dLGN is not inherited from V1 (Jaepel et al., 2017) and supports the idea that dLGN plasticity involves the plasticity of retinogeniculate afferents. Interestingly, when we repeated this experiment in (C57Bl/6Jrj) WT mice during the critical period, we found that silencing V1 also did not affect ipsi- or contralateral eye responses in non-deprived mice (**Suppl. Fig. 2A, B**), but selectively reduced ipsilateral eye responses in monocularly deprived mice (**Fig. 5G-H**). Consequently, silencing V1 significantly reduced the OD shift in these animals (**Fig. 5I**). The change in ODI caused by V1 silencing in mice during the critical period was significantly larger than that in adult WT ($P=0.034$). Thus, during the critical period, corticothalamic connections strengthen the OD shift in dLGN, while they do not in adulthood.

In monocularly deprived adult KO mice (**Fig. 5I-K**), silencing V1 did not affect responses to the contralateral eye, but significantly reduced those to the ipsilateral eye (**Fig. 5J, K**), similarly to what we observed in WT mice during the critical period. However, despite this effect of V1 silencing, the average ODI in monocularly deprived KO mice was not significantly altered by it (**Fig. 5I**). We conclude that feedback from V1 does not affect OD in the dLGN of adult mice, independently of whether they are monocularly deprived or lack synaptic inhibition in the thalamus. During the critical period, however, V1 silencing does reduce the OD shift observed in dLGN.

Discussion

In summary, we show that OD plasticity in dLGN is reduced in adult mice lacking thalamic synaptic inhibition. In these mice, OD plasticity in V1 is absent, suggesting that it requires thalamic plasticity. We do not find evidence that feedback from V1 affects the thalamic OD shift in adult mice. This differs from the situation during the critical period, in which the OD shift in dLGN is partially inherited from V1.

In line with previous work (Jaepel et al., 2017), we find that in dLGN of adult mice, 7 days of MD causes a strong OD shift. The main substrate of this plasticity in adult dLGN appears to be the retinogeniculate synapse, as silencing V1, the other main source of visual input to dLGN, does not affect the OD shift. Recent work has shown that while many relay cells in dLGN receive binocular inputs (Rompani et al., 2017), synapses from the non-dominant eye are often silent and dominated by NMDA receptors (Bauer et al., 2021). Because silent synapses contribute strongly to OD plasticity in V1, it is interesting to speculate that OD plasticity in dLGN also involves the




Fig. 5

| The OD shift in dLGN is independent from V1 feedback in adult mice but not in critical period mice

A & B, Muscimol injection in V1 has no effect on the ODI in dLGN of adult non-deprived WT (**A**) and KO (**B**) mice (Wilcoxon signed rank; WT NO MD vs. WT NO MD with muscimol, $P=0.86$, $n=40$ units, 7 mice; KO NO MD vs. KO NO MD with muscimol, $P=0.45$, 45 units, 9 mice). **C, D & E**, Muscimol injection in V1 has no significant effect on the ODI (**C**), or contralateral (**D**) or ipsilateral (**E**) eye responses in dLGN of monocularly deprived WT mice (Wilcoxon signed rank; ODI, WT MD vs. WT MD with muscimol, $P=0.89$; contralateral, WT MD vs. WT MD with muscimol, $P=0.21$; ipsilateral, WT MD vs. WT MD with muscimol, $P=0.10$, $n=22$ units, 6 mice). **F, G & H**, During the critical period, V1 silencing has a significant influence on the ODI (**F**) and ipsilateral eye responses (**H**) in dLGN of monocularly deprived WT mice, but not on contralateral eye responses in these mice (**G**) (Wilcoxon signed rank; ODI, WT MD vs. WT MD with muscimol, $P<0.001$; contralateral, WT MD vs. WT MD with muscimol, $P=0.46$; ipsilateral, WT MD vs. WT MD with muscimol, $P=0.003$, $n=41$ units, 10 mice). **I, J & K**, Muscimol injection in V1 has no significant influence on the ODI and contralateral eye responses in KO MD mice, but significantly modulates dLGN ipsilateral eye responses (Wilcoxon signed rank; ODI, KO MD vs. KO MD with muscimol, $P=0.13$; contralateral, KO MD vs. KO MD with muscimol, $P=0.97$; ipsilateral, KO MD vs. KO MD with muscimol, $P=0.004$, $n=34$ units, 9 mice).

unsilencing and strengthening of these synapses (Huang et al., 2015 [DOI](#); Yusifov et al., 2021 [DOI](#)). Indeed, the OD shift in adult mice involves many (monocular) contralateral eye-selective neurons to become binocular (Jaepel et al., 2017 [DOI](#)).

In apparent contradiction to our findings and the work by Jaepel et al., a recent study (Li et al., 2022 [DOI](#)) did not observe an OD shift in dLGN of adult mice upon 6 days of monocular deprivation. A possible explanation for this apparent discrepancy is that Li et al only included binocular dLGN neurons in the analysis. This approach will exclude the monocular, contralateral-eye selective neurons in non-deprived animals from the analysis, thus decreasing the measured OD shift.

We find that in adult KO mice, OD plasticity in dLGN is absent. As we have previously shown that OD plasticity is also absent in dLGN of KO mice during the critical period, it is possible that maturation of dLGN is halted in a pre-critical period like state in adult KO mice, in analogy to the situation in GAD65 deficient mice in which reduced GABA release interferes with the onset of the critical period of cortical OD plasticity (Fagiolini and Hensch 2000). However, we did not find evidence for halted thalamic development. Receptive fields of dLGN relay cells become smaller between eye opening and critical period onset (Tschetter et al., 2018 [DOI](#)), and we find that in WT and KO mice receptive field sizes are the same. Furthermore, there is a substantial increase in inhibitory and cholinergic boutons during this developmental stage (Bickford et al., 2010 [DOI](#); Sokhadze et al., 2018 [DOI](#); Sommeijer et al., 2017 [DOI](#)), but again, we find that adult WT and KO mice are not different in this respect. The primary difference between WT and KO mice in adult dLGN thus appears to be the lack of synaptic inhibition. We do not know how this lack of synaptic inhibition in adult dLGN interferes with OD plasticity, but a possible explanation may lie in altered temporal profile of the visual responses it causes, which could interfere with spike timing-dependent plasticity.

How does inactivation of synaptic inhibition and OD plasticity in dLGN interfere with the OD shift in V1? We believe that it is unlikely that OD plasticity in V1 is deficient due to halted development of V1 in a pre-critical period like state. We have previously shown that brief MD induces an OD shift in V1 of these mice (Sommeijer et al., 2017 [DOI](#)), suggesting that the critical period starts normally in KO mice. The previously observed plasticity deficit during the critical period was limited to the second stage of the OD shift in V1, which requires long-term monocular deprivation. It thus appears that both during the critical period and in adulthood, the late phase of OD plasticity in V1 induced by long-term monocular deprivation requires thalamic plasticity or inhibition. The most straight-forward explanation for the cortical plasticity deficit is that MD-induced changes in dLGN relay cell responses to inputs from the two eyes contribute to the OD shift in V1. Additionally, the strengthening of responses to the non-deprived eye in dLGN neurons may provide their axons with a competitive advantage during OD plasticity in V1, further enhancing the OD shift in V1. Finally, it is possible that OD plasticity in V1 is also affected by deficits in spike dependent plasticity due to the more attenuated nature of dLGN responses that we observe in KO mice. Together, our results indicate that thalamic inhibition and plasticity play a crucial role in OD plasticity in adult V1, regardless of a possible developmental contribution to the plasticity deficit in dLGN.

The study by Bauer et al (Bauer et al., 2021 [DOI](#)) showed that binocularity in mouse dLGN may be lower than suggested by the current study and earlier work (Howarth et al., 2014 [DOI](#); Li et al., 2022 [DOI](#); Sommeijer et al., 2017 [DOI](#)) that involved multi-electrode recordings in dLGN. Although this difference may be caused by technical limitations of single unit recordings or calcium imaging, we think it is most likely explained by the fact that studies employing electrophysiological recording in dLGN targeted the frontal ipsilateral projection zone of dLGN, which is its most binocular region (Bauer et al., 2021 [DOI](#)). Recording in this region is essential when studying OD plasticity or binocularity in dLGN, but will strongly bias towards binocularly-responding relay cells. When

using two-photon imaging of dLGN boutons in V1 (Bauer et al., 2021 [DOI](#); Huh et al., 2020 [DOI](#); Jaepel et al., 2017 [DOI](#)), neurons from other parts of dLGN including the monocular shell- and caudal regions are also sampled.

It is unknown whether adult thalamic OD plasticity also occurs in species in which retinal inputs from the two eyes are organized in more strictly separated layers in dLGN, such as cats or primates. Studies in cats consistently found that upon MD or squint during the critical period, the layers responding to the affected eye were thinner (Hickey et al., 1977 [DOI](#); Wiesel and Hubel, 1963b [DOI](#)) and that the neuronal responses in these layers were slower or weaker (Eysel et al., 1979 [DOI](#); Ikeda and Wright, 1976 [DOI](#); Sestokas and Lehmkuhle, 1986 [DOI](#); Wiesel and Hubel, 1963b [DOI](#)). Also in human amblyopes it was noted that dLGN responses to the amblyopic eye were weaker (Hess et al., 2009 [DOI](#)). So far, studies on dLGN plasticity by prolonged visual deprivation in adulthood are missing, though it was noted that in human subjects that suffered from glaucoma, the dLGN layers representing the affected eye were thinner (Yücel et al., 2001 [DOI](#)). Future research will need to establish whether plasticity in dLGN in humans contributes to amblyopia, and whether it can be enhanced to treat the disorder. That enhancement of thalamic plasticity is in principle possible is shown by experiments in mice, demonstrating that inactivation of the nogo-66 receptor in thalamus allows recovery of reduced acuity in adult mice that were monocularly deprived during development (Stephany et al., 2018 [DOI](#)). Despite extensive monosynaptic excitatory feedback and bisynaptic inhibitory feedback (through TRN) from V1, we found that feedback does not affect the average strength of response to ipsi- or contralateral eye stimulation and thus neither the measured OD shift in dLGN of adult mice. The absence of V1 on the OD shift in adult mice was also reported previously (Jaepel et al., 2017 [DOI](#)). However in that study, silencing V1 with muscimol reduced the imaged calcium responses in boutons of dLGN axons projecting to V1, suggesting that dLGN neurons became less responsive to inputs from both eyes. A possible explanation for this apparent discrepancy may be that muscimol also has a direct effect on thalamocortical inputs (Liu et al., 2007 [DOI](#); Wang et al., 2019 [DOI](#); Yamauchi et al., 2000 [DOI](#)), which may reduce calcium responses in synaptic boutons of dLGN neurons without actually reducing their spiking activity at the soma.

Studies involving optogenetic stimulation of layer 6 neurons find that feedback from V1 suppresses dLGN responses (Denman and Contreras, 2015 [DOI](#); Kirchgeßner et al., 2020 [DOI](#); Olsen et al., 2012 [DOI](#)), though this differs per cell and changes with stimulus strength and frequency (Kirchgeßner et al., 2020 [DOI](#)). When V1 feedback is silenced, however, the average strength of dLGN responses is not reduced in most studies (Denman and Contreras, 2015 [DOI](#); Howarth et al., 2014 [DOI](#); Kirchgeßner et al., 2020 [DOI](#)). This suggests that broad optogenetic stimulation of layer 6 predominantly recruits inhibitory feedback, while visual stimulation provides either more balanced or more limited excitatory and inhibitory feedback to dLGN. In line with these previous studies, we find that silencing V1 does not significantly alter dLGN responses to either eye in WT mice. If the lack of effect of V1 silencing is due to balanced excitatory and inhibitory feedback, one would expect that in KO mice lacking thalamic synaptic inhibition, silencing V1 would cause a reduction of dLGN responses. However, such an effect in KO mice was only observed in dLGN responses to the ipsilateral eye, and reached significance only in monocularly deprived mice. This suggests that V1 feedback affects dLGN responses to the ipsilateral eye more strongly than those to the contralateral eye. This may be the result of V1 receiving input from the ipsilateral eye through dLGN but also through callosal V1 inputs (Cerri et al., 2010 [DOI](#)).

We found that during the critical period, the influence of V1 feedback on the OD shift in dLGN was much stronger. It is possible that inhibitory innervation of dLGN is still weaker during the critical period than in adulthood, causing excitatory feedback from V1 to dominate like in KO mice.

Additionally, a stronger OD shift occurs in V1 during the critical period, adding to the strength of the cortical feedback representing the ipsilateral eye. Interestingly, an excitotoxic lesion of V1 was found to alter OD in dLGN during development and affect OD plasticity in dLGN at various ages (Li

et al., 2022 [↗](#)). This suggests that continuous crosstalk between thalamus and cortex during development guides plasticity, possibly optimizing thalamocortical and corticothalamic connections. The continued absence of corticothalamic feedback is likely to have a much larger impact on dLGN plasticity than the acute silencing we performed.

During the critical period, an experience-dependent phase of retinogeniculate refinement takes place, probably optimizing direction-selective inputs from the retina (Hooks and Chen, 2020 [↗](#); Rompani et al., 2017 [↗](#); Thompson et al., 2017 [↗](#)). This experience-dependent refinement, like OD plasticity in dLGN, also depends on feedback from V1 (Thompson et al., 2017 [↗](#), 2016 [↗](#)). It is thus possible that refinement of binocular inputs and direction-selective inputs in dLGN are one and the same process.

We conclude that dLGN retains a high level of plasticity in adulthood and has considerable influence on cortical plasticity. This plasticity may not be restricted to binocular responses, but could also be relevant for other forms of perceptual learning (Yu et al., 2016 [↗](#)). The findings stress that a thalamic involvement needs to be considered in amblyopia and learning disabilities. Additionally, the results may help understanding brain disorders that are thought to involve dysfunctional thalamocortical circuits, ranging from attention deficit disorder (Wells et al., 2016) to schizophrenia (Benoit et al., 2022; Pinault, 2011; Pratt et al., 2017). Future experiments focusing on changes in thalamic responses and their interaction with the cortex may provide exciting new insights in how the brain learns.

Methods

Animals

All mice used to assess OD plasticity in adulthood were bred from homozygous conditional Gabra1-deficient mice (Gabra1^{fl-hom}) (Vicini et al., 2001 [↗](#)) crossed with homozygous Gabra1-deficient, heterozygous Olig3-cre knockin mice (Vue et al., 2009 [↗](#)) (Gabra1^{fl-hom} Olig3-cre⁺). Before cross-breeding the lines, Gabra1^{fl-hom} mice had been backcrossed to C57Bl/6J mice (Janvier) for at least 6 generations. Olig3-cre⁺ mice were crossed to C57Bl/6J mice for at least 2 generations, but should be considered mixed background. All animals were tested for unintended germline recombination of the Gabra1^{fl} locus, and such mice were excluded from breeding or experiments. In our experiments, we used 4 groups of animals: monocularly deprived or non-deprived Gabra1^{fl-hom} Olig3-cre⁺ mice and monocularly deprived or non-deprived Gabra1^{fl-hom} Olig3-cre⁺ siblings (P45-P90). The experimenter was blind to the genotype of the mice until the end of the experiment. Mice used for OD plasticity experiments during the critical period were C57Bl/6J mice. All mice were housed in a 12h/12h dark/light cycle. Both male and female mice were used. Mice housing conditions were according to Dutch law. All experiments were approved by the institutional animal care and use committee of the Royal Netherlands Academy of Arts and Sciences.

Immunohistochemistry

Age-matched mice were anesthetized with 0.1 ml/g body weight Nembutal (Janssen) and perfused with 4% paraformaldehyde (PFA) in PBS (~50 ml per mouse) and post-fixed for 2–6 h. Post fixation time was consistent between compared groups. Sections from dLGN of 50 µm were made by using a vibratome (Leica VT1000S). Mouse-α-GAD67 (1:500, Chemicon, MAB5406) was used to label inhibitory boutons and and guinea pig-α-VaChT (1:500, SySy 139105) to label cholinergic boutons. Primary antibodies were visualized using Fluor 594 Goat-α-mouse (1:1000, Invitrogen, A11032 and Alexa Fluor 488 Goat-α-Guinea pig (1:1000, Invitrogen, A11073). Free-floating sections were briefly washed in PBS followed by 1–2 h blocking in PBS containing 5% normal goat serum and 0.1% Triton X-100 at room temperature. Primary antibody incubation was performed overnight at 4 °C in fresh blocking solution. Next, the sections were washed three times for 10 min in PBS with 0.1%

Tween-20 (PBST) followed by secondary antibody incubation in fresh blocking solution for 1.5–2 h at RT. After washing three times for 10 min in TBST the sections were mounted on glass slides using Mowiol (Calbiochem/ MerckMillipore), glass covered and stored at 4°C.

Confocal microscopy and data analysis

Sections were imaged using a confocal microscopy (Leica SP5) with constant gain and laser power across compared samples. Care was taken that no signal clipping was present. For quantification of GAD67 and VAcHT puncta, and images were taken with a 40x objective (2048 x 2048 resolution). Background fluorescence was subtracted with ImageJ. VAcHT and GAD67 puncta were quantified using the ‘SynQuant’ ImageJ plugin (Wang et al., 2020), creating ROIs corresponding to synaptic puncta. For each image we calculated the average size of identified puncta, the number of puncta per unit of area and the percentage of image area identified as part of punctum.

Monocular deprivation

The eyelids of the eye contralateral to the recording side were sutured for MD. The surgery was performed under isoflurane anesthesia (5% induction, 1.5–2% maintenance in 0.7 l/min O₂). The eye was rinsed with saline. The eyelids were sutured together with 7.0 Ethilon thread. Eyes were checked for infection in the following days and reopened during recording. Only mice with healthy eye conditions were included.

Electrophysiology recordings, visual stimulation and V1 silencing

Mice were anesthetized by intraperitoneal injection of urethane (Sigma; 20% solution in saline, 1.2g/kg body weight), supplemented by intraperitoneal injection of chlorprothixene (Sigma; 2.0mg/ml in saline, 8mg/kg body weight), followed by subcutaneous injection of atropine (Sigma; 1mg/ml, 6mg/kg body weight), and head-fixed by ear- and bite bars. The temperature was measured with a rectal probe and maintained at 36.5 °C. The craniotomies for dLGN (2.0 mm lateral, 2.5 mm posterior from bregma) and V1 (2.95 mm lateral, 0.45 mm anterior from lambda) recording windows were made using a dental drill. During recordings of V1 or dLGN responses to input from one eye, the other eye was covered with a double layer of black cloth and black tape.

Using a linear silicon microelectrode (A1×16–5mm–25–177–A16, 16 channels spaced 50 µm apart, Neuronexus), extracellular recordings from V1 and dLGN were performed separately. Visual stimuli were projected by a gamma-corrected projector (PLUS U2-X1130 DLP) on a back-projection screen (Macada Innovision, covering a 60X42 cm area) positioned 17.5 cm in front of the mouse. The visual stimuli were programmed using the MATLAB (MathWorks) scripts package Psychophysics Toolbox 3 (Brainard, 1997 [link](#)). V1 was first recorded at a depth of approximately 800 µm from the cortical surface. Receptive field position was checked by showing white squares (5 deg) at random positions on a black background. If the receptive field was not within 30° from the center, we relocated the electrode and checked again. ODI was measured by presenting alternating white and gray full-screen stimuli to each eye in turn. Each stimulus lasted 3s. Both white and gray screens were presented with 100 repetitions. When the V1 recording was finished, we relocated the electrode to dLGN at a depth of 2700–3000 µm from bregma. The receptive field and ODI measurement procedures were repeated in dLGN. We then silenced V1 by injecting muscimol (Sigma; 10 mM; ~150 µL per mouse), a selective agonist for GABA_A receptors, in V1 and measured the ODI again in dLGN. After recording, we moved electrode back to V1 to verify that muscimol had silenced V1.

The extracellular signals were amplified and bandpass filtered at 500 Hz–10 kHz and digitized at 24 kHz using a Tucker-Davis Technologies RX5 Pentusa base station. The spike detection was done by a voltage thresholder at 3x s.d. online per recording, or offline using the open-source sorting

package KiloSort (Pachitariu et al., 2016 [↗](#)). Spikes were sorted and clustered by either principle component analysis-based custom-written MATLAB scripts or integrated template matching-based KiloSort scripts.

Analysis of electrophysiological data

Data analysis was done using custom-made MATLAB scripts (<http://github.com/heimel/inVivoTools> [↗](#)). For each 3 s stimulus related activity, we treated the last 500ms of the previous trial as baseline. Therefore, we defined the visual responses as the difference between the first 500ms of the stimulus and the mean of the last 500ms activities of the previous stimulus. The peak visual responses of stimuli were considered as the maximum firing rates in first 300ms of visual related responses. The visual responses were calculated as average responses of 300ms. ODI was calculated as $(R_{contra} - R_{ipsi}) / (R_{contra} + R_{ipsi})$, where the R_{contra} is the average firing rate of the unit when contralateral eye was open and ipsilateral eye was covered; R_{ipsi} is opposite. For receptive field mapping, we computed the spike – triggered average of the random sparse squares stimulus. The peak rate threshold was set to 5 Hz when the patch was within the receptive field. The actual position and size of visual field were computed and corrected for the actual distance between stimuli and animal.

To assess the monocular and binocular categorical responsiveness, we performed Zeta-test (<https://github.com/JorritMontijn/ZETA> [↗](#)) (Montijn et al., 2021 [↗](#)) over all trials on all units that we recorded. ZETA (Zenith of Event-based Time-locked Anomalies)-test is a more powerful and sensitive tool than t-test or ANOVA to detect whether a cell is responsive to stimulation in a statistically robust way and avoids binning and parameter selection together. The significant on- and off responses were calculated separately. Therefore, for each unit, four responses were tested: contra on, contra off, ipsi on and ipsi off. For each unit, if the contra on or off response passed the ZETA-test ($P < 0.05$), but neither the ipsi on nor the off response did, we categorized this unit as pure contra. The same approach, but then reversely, was used to determine pure ipsilateral eye-responsive units. If a contralateral eye on or off response and an ipsilateral eye on or off response passed the ZETA-test, we considered this unit as binocular.


Statistics

For testing the interaction between genotype with MD on OD plasticity and RF size in V1 and dLGN, we used a two-way ANOVA test with post-hoc Tukey-Kramer tests. Differences in the temporal profile of the visual responses in WT vs KO mice was determined using a repeated measure two-way ANOVA with post-hoc Fisher's LSD test. Quantitation of immunohistochemical analyses and difference in OD shift upon cortical silencing were performed using Student's t-test. All other tests were done with non-parametric tests. Statistical analyses of the response changes of V1 and dLGN units in WT and KO mice, were done by non-parametric Mann-Whitney U tests. For testing the significance of the effect of silencing V1 on adult and critical period dLGN responses, on the OD in adult deprived and non-deprived KO mice and WT mice, on the OD in critical period deprived WT mice, and the effects of muscimol injection on V1 responses, Wilcoxon signed rank tests were used.

Acknowledgements


The authors thank Emma Ruimschotel for technical assistance, Robin Haak and Huub Terra for help with spike sorting, Huub Terra for providing feedback on the manuscript and the Mechatronics department and Animal facilities of the NIN for their services. **Figures 1A** [↗](#) & B, 2A, 3A and 4A were created with *BioRender.com* (<http://biorender.com/>) [↗](#) (2022). This project has

Supplementary data



Supplementary Figure 1.

A, Examples of immunohistochemical staining for GAD67 in dLGN of adult WT and KO mice. **B**, Quantification of the density, number, and sizes of GAD67 puncta (putative inhibitory boutons). No differences were detected between WT and KO mice (T-test; puncta density, $P=0.85$; puncta number, $P=0.82$; puncta size, $P=0.80$. $n=3$ mice per group, 3 slices were imaged per mouse). **C**, Examples of immunohistochemical staining for VACHT puncta (putative cholinergic boutons) in dLGN of adult WT and KO mice. **D**, Quantification of the density, number, and sizes of VACHT. No differences were detected between WT and KO mice (T-test; puncta density, $P=0.29$; puncta number, $P=0.35$; puncta size, $P=0.22$. $n=3$ mice per group, 3 slices were imaged per mouse).



Supplementary Figure 2.

A, MD induces ocular dominance plasticity in dLGN of WT mice during the critical period (Mann-Whitney; WT NO MD vs. WT MD, $P=0.0015$; WT NO MD, $n=36$, 10 mice; WT MD, $n=41$, 10 mice). **B**, During the critical period, V1 feedback has no effect on OD in non-deprived WT mice (Wilcoxon signed rank; ODI, WT NO MD vs. WT MD with muscimol, $P=0.20$).

References

- Bauer J, Weiler S, Fernholz MHP, Laubender D, Scheuss V, Hübener M, Bonhoeffer T, Rose T (2021) **Limited functional convergence of eye-specific inputs in the retinogeniculate pathway of the mouse** *Neuron* **109**:2457–2468 <https://doi.org/10.1016/j.NEURON.2021.05.036>
- Bickford ME, Slusarczyk A, Dilger EK, Krahe TE, Kucuk C, Guido W (2010) **Synaptic development of the mouse dorsal lateral geniculate nucleus** *J Comp Neurol* **518**:622–635 <https://doi.org/10.1002/CNE.22223>
- Brainard DH (1997) **The Psychophysics Toolbox** *Spat Vis* **10**:433–436 <https://doi.org/10.1163/156856897X00357>
- Cerri C, Restani L, Caleo M (2010) **Callosal contribution to ocular dominance in rat primary visual cortex** *Eur J Neurosci* **32**:1163–1169 <https://doi.org/10.1111/j.1460-9568.2010.07363.x>
- Denman DJ, Contreras D (2015) **Complex Effects on In Vivo Visual Responses by Specific Projections from Mouse Cortical Layer 6 to Dorsal Lateral Geniculate Nucleus** *J Neurosci* **35**:9265–9280 <https://doi.org/10.1523/JNEUROSCI.0027-15.2015>
- Eysel UT, Grüsser OJ, Hoffmann KP (1979) **Monocular deprivation and the signal transmission by X- and Y-neurons of the cat lateral geniculate nucleus** *Exp brain Res* **34**:521–539 <https://doi.org/10.1007/BF00239147>
- Frenkel MY, Bear MF (2004) **How monocular deprivation shifts ocular dominance in visual cortex of young mice** *Neuron* **44**:917–923 <https://doi.org/10.1016/j.NEURON.2004.12.003>
- Gilbert CD, Li W, Piech V (2009) **Perceptual learning and adult cortical plasticity** *J Physiol* **587**:2743–2751 <https://doi.org/10.1113/JPHYSIOL.2009.171488>
- Gilbert CD, Wiesel TN (1992) **Receptive field dynamics in adult primary visual cortex** *Nature* **356**:150–152 <https://doi.org/10.1038/356150A0>
- Gordon A, Stryker P (1996) **Experience-Dependent Plasticity of Binocular Responses in the Primary Visual Cortex of the Mouse** *J Neurosci* **76**:3274–3286
- Heimel JA, Hartman RJ, Hermans JM, Levitt CN (2007) **Screening mouse vision with intrinsic signal optical imaging** *Eur J Neurosci* **25**:795–804 <https://doi.org/10.1111/j.1460-9568.2007.05333.x>
- Hensch TK, Stryker MP (1996) **Ocular dominance plasticity under metabotropic glutamate receptor blockade** *Science* **272**:554–557 <https://doi.org/10.1126/SCIENCE.272.5261.554>
- Hess RF, Thompson B, Gole G, Mullen KT (2009) **Deficient responses from the lateral geniculate nucleus in humans with amblyopia** *Eur J Neurosci* **29**:1064–1070 <https://doi.org/10.1111/j.1460-9568.2009.06650.x>
- Hickey TL, Spear PD, Kratz KE (1977) **Quantitative studies of cell size in the cat's dorsal lateral geniculate nucleus following visual deprivation** *J Comp Neurol* **172**:265–281 <https://doi.org/10.1002/CNE.901720206>

- Hofer SB, Mrsic-Flogel TD, Bonhoeffer T, Hübener M (2006) **Prior experience enhances plasticity in adult visual cortex** *Nat Neurosci* **9**:127–132 <https://doi.org/10.1038/nn1610>
- Hooks BM, Chen C (2020) **Circuitry Underlying Experience-Dependent Plasticity in the Mouse Visual System** *Neuron* **106**:21–36 <https://doi.org/10.1016/j.NEURON.2020.01.031>
- Howarth M, Walmsley L, Brown TM (2014) **Binocular integration in the mouse lateral geniculate nuclei** *Curr Biol* **24**:1241–1247 <https://doi.org/10.1016/j.CUB.2014.04.014>
- Huang X, Stodieck SK, Goetze B, Cui L, Wong MH, Wenzel C, Hosang L, Dong Y, Löwel S, Schlüter OM (2015) **Progressive maturation of silent synapses governs the duration of a critical period** *Proc Natl Acad Sci* **112**:E3131–E3140 <https://doi.org/10.1073/pnas.1506488112>
- Huh CYL, Abdelaal K, Salinas KJ, Gu D, Zeitoun J, Figueroa Velez DX, Peach JP, Fowlkes CC, Gandhi SP (2020) **Long-term Monocular Deprivation during Juvenile Critical Period Disrupts Binocular Integration in Mouse Visual Thalamus** *J Neurosci* **40**:585–604 <https://doi.org/10.1523/JNEUROSCI.1626-19.2019>
- Ikeda H, Wright MJ (1976) **Properties of LGN cells in kittens reared with convergent squint: a neurophysiological demonstration of amblyopia** *Exp brain Res* **25**:63–77 <https://doi.org/10.1007/BF00237326>
- Jaepel J, Hübener M, Bonhoeffer T, Rose T (2017) **Lateral geniculate neurons projecting to primary visual cortex show ocular dominance plasticity in adult mice** *Nat Neurosci* **20**:1708–1714 <https://doi.org/10.1038/s41593-017-0021-0>
- Kalogeraki E, Pielecka-Fortuna J, Löwel S (2017) **Environmental enrichment accelerates ocular dominance plasticity in mouse visual cortex whereas transfer to standard cages resulted in a rapid loss of increased plasticity** *PLoS One* **12** <https://doi.org/10.1371/journal.pone.0186999>
- Kirchgessner MA, Franklin AD, Callaway EM (2020) **Context-dependent and dynamic functional influence of corticothalamic pathways to first- And higher-order visual thalamus** *Proc Natl Acad Sci U S A* **117**:13066–13077 <https://doi.org/10.1073/PNAS.2002080117/-/DCSUPPLEMENTAL>
- Lehmann K, Löwel S (2008) **Age-Dependent Ocular Dominance Plasticity in Adult Mice** *PLoS One* **3** <https://doi.org/10.1371/journal.pone.0003120>
- Leresche N, Lambert RC (2018) **GABA receptors and T-type Ca²⁺ channels crosstalk in thalamic networks** *Neuropharmacology* **136**:37–45 <https://doi.org/10.1016/j.NEUROPHARM.2017.06.006>
- Li N, Liu Q, Zhang Y, Yang Z, Shi X, Gu Y (2022) **Cortical feedback modulates distinct critical period development in mouse visual thalamus** *iScience* **26** <https://doi.org/10.1016/j.ISCI.2022.105752>
- Liu BH, Wu GK, Arbuckle R, Tao HW, Zhang LI (2007) **Defining cortical frequency tuning with recurrent excitatory circuitry** *Nat Neurosci* **10**:1594–1600 <https://doi.org/10.1038/NN2012>
- Montijn JS, Seignette K, Howlett MH, Cazemier JL, Kamermans M, Levelt CN, Heimel JA (2021) **A parameter-free statistical test for neuronal responsiveness** *Elife* **10** <https://doi.org/10.7554/eLife.71969>

- Olsen SR, Bortone DS, Adesnik H, Scanziani M (2012) **Gain control by layer six in cortical circuits of vision** *Nature* **483**:47–54 <https://doi.org/10.1038/NATURE10835>
- Pachitariu M, Steinmetz N, Kadir S, Carandini M D. HK. (2016) **Kilosort: realtime spike-sorting for extracellular electrophysiology with hundreds of channels** *bioRxiv* <https://doi.org/10.1101/061481>
- Rompani SB, Müllner FE, Wanner A, Zhang C, Roth CN, Yonehara K, Roska B (2017) **Different Modes of Visual Integration in the Lateral Geniculate Nucleus Revealed by Single-Cell-Initiated Transsynaptic Tracing** *Neuron* **93**:767–776 <https://doi.org/10.1016/J.NEURON.2017.01.028>
- Rose T, Jaepel J, Hübener M, Bonhoeffer T (2016) **Cell-specific restoration of stimulus preference after monocular deprivation in the visual cortex** *Science* **352**:1319–1322 <https://doi.org/10.1126/SCIENCE.AAD3358>
- Sato M, Stryker MP (2008) **Distinctive Features of Adult Ocular Dominance Plasticity** *J Neurosci* **28**:10278–10286 <https://doi.org/10.1523/JNEUROSCI.2451-08.2008>
- Sawtell NB, Frenkel MY, Philpot BD, Nakazawa K, Tonegawa S, Bear MF (2003) **NMDA receptor-dependent ocular dominance plasticity in adult visual cortex** *Neuron* **38**:977–985 [https://doi.org/10.1016/S0896-6273\(03\)00323-4](https://doi.org/10.1016/S0896-6273(03)00323-4)
- Sestokas A, Lehmkuhle S (1986) **The effects of monocular deprivation on the visual latency of geniculate X- and Y-cells in the cat** *Brain Res* **395**:93–95 [https://doi.org/10.1016/S0006-8993\(86\)80013-0](https://doi.org/10.1016/S0006-8993(86)80013-0)
- Sokhadze G, Seabrook TA, Guido W (2018) **The absence of retinal input disrupts the development of cholinergic brainstem projections in the mouse dorsal lateral geniculate nucleus** *Neural Dev* **13** <https://doi.org/10.1186/S13064-018-0124-7>
- Sommeijer J-PP, Ahmadlou M, Saiepour MH, Seignette K, Min R, Heimel JA, Levelt CN (2017) **Thalamic inhibition regulates critical-period plasticity in visual cortex and thalamus** *Nat Neurosci* **1** <https://doi.org/10.1038/s41593-017-0002-3>
- Stephany C-É, Ma X, Dorton HM, Wu J, Solomon AM, Frantz MG, Qiu S, McGee AW (2018) **Distinct Circuits for Recovery of Eye Dominance and Acuity in Murine Amblyopia** *Curr Biol* **28**:1914–1923 <https://doi.org/10.1016/j.cub.2018.04.055>
- Thompson A, Gribizis A, Chen C, Crair MC (2017) **Activity-dependent development of visual receptive fields** *Curr Opin Neurobiol* **42**:136–143 <https://doi.org/10.1016/J.CONB.2016.12.007>
- Thompson AD, Picard N, Min L, Fagiolini M, Chen C (2016) **Cortical Feedback Regulates Feedforward Retinogeniculate Refinement** *Neuron* **91**:1021–1033 <https://doi.org/10.1016/J.NEURON.2016.07.040>
- Tschetter WW, Govindaiah G, Etherington IM, Guido W, Niell CM (2018) **Refinement of Spatial Receptive Fields in the Developing Mouse Lateral Geniculate Nucleus Is Coordinated with Excitatory and Inhibitory Remodeling** *J Neurosci* **38**:4531–4542 <https://doi.org/10.1523/JNEUROSCI.2857-17.2018>

- Vicini S, Ferguson C, Prybylowski K, Kralic J, Morrow AL, Homanics GE (2001) **GABA(A) receptor alpha1 subunit deletion prevents developmental changes of inhibitory synaptic currents in cerebellar neurons** *J Neurosci* **21**:3009–3016 <https://doi.org/10.1523/JNEUROSCI.21-09-03009.2001>
- Vue TY, Bluske K, Alishahi A, Yang LL, Koyano-Nakagawa N, Novitsch B, Nakagawa Y (2009) **Sonic hedgehog signaling controls thalamic progenitor identity and nuclei specification in mice** *J Neurosci* **29**:4484–4497 <https://doi.org/10.1523/JNEUROSCI.0656-09.2009>
- Wang L, Kloc M, Maher E, Erisir A, Maffei A (2019) **Presynaptic GABAA Receptors Modulate Thalamocortical Inputs in Layer 4 of Rat V1** *Cereb Cortex* **29**:921–936 <https://doi.org/10.1093/CERCOR/BHX364>
- Wiesel TN, Hubel DH (1963) **SINGLE-CELL RESPONSES IN STRIATE CORTEX OF KITTENS DEPRIVED OF VISION IN ONE EYE** *J Neurophysiol* **26**:1003–1017 <https://doi.org/10.1152/JN.1963.26.6.1003>
- Wiesel TN, Hubel DH (1963) **EFFECTS OF VISUAL DEPRIVATION ON MORPHOLOGY AND PHYSIOLOGY OF CELLS IN THE CATS LATERAL GENICULATE BODY** *J Neurophysiol* **26**:978–993 <https://doi.org/10.1152/JN.1963.26.6.978>
- Yamauchi T, Hori T, Takahashi T (2000) **Presynaptic inhibition by muscimol through GABAB receptors** *Eur J Neurosci* **12**:3433–3436 <https://doi.org/10.1046/J.1460-9568.2000.00248.X>
- Yu Q, Zhang P, Qiu J, Fang F (2016) **Perceptual Learning of Contrast Detection in the Human Lateral Geniculate Nucleus** *Curr Biol* **26**:3176–3182 <https://doi.org/10.1016/J.CUB.2016.09.034>
- Yücel YH, Zhang Q, Weinreb RN, Kaufman PL, Gupta N (2001) **Atrophy of relay neurons in magno- and parvocellular layers in the lateral geniculate nucleus in experimental glaucoma** *Investig Ophthalmol Vis Sci* **42**:3216–3222
- Yusifov R, Tippmann A, Staiger JF, Schlüter OM, Löwel S (2021) **Spine dynamics of PSD-95-deficient neurons in the visual cortex link silent synapses to structural cortical plasticity** *Proc Natl Acad Sci U S A* **118** <https://doi.org/10.1073/PNAS.2022701118>

Author information

Yi Qin

Molecular Visual Plasticity Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands, University of Strasbourg, Strasbourg, France

Mehran Ahmadi

Circuits, Structure and Function Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands
ORCID ID: [0000-0003-4688-2064](https://orcid.org/0000-0003-4688-2064)

Samuel Suhai

Molecular Visual Plasticity Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands

Paul Neering

Molecular Visual Plasticity Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands

Leander de Kraker

Molecular Visual Plasticity Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands
ORCID iD: [0000-0001-5598-3727](https://orcid.org/0000-0001-5598-3727)

J. Alexander Heimel

Circuits, Structure and Function Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands2
ORCID iD: [0000-0002-5291-4184](https://orcid.org/0000-0002-5291-4184)

Christiaan N. Levelt

Molecular Visual Plasticity Group, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Meibergdreef 47, 1105 BA Amsterdam, the Netherlands, Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive Research, VU University Amsterdam, de Boelelaan 1085, 1081 HV, the Netherlands
For correspondence: c.levelt@nin.knaw.nl
ORCID iD: [0000-0002-1813-6243](https://orcid.org/0000-0002-1813-6243)

Editors

Reviewing Editor

Mario Penzo

National Institute of Mental Health, United States of America

Senior Editor

Joshua Gold

University of Pennsylvania, United States of America

Reviewer #1 (Public Review):

Qin et al., demonstrate, convincingly, that plasticity of ocular dominance of binocular neurons in the visual thalamus persists in adulthood. The adult plasticity is similar to that described in critical period juveniles in that it is absent in transgenic mice with the deletion of the GABA $\alpha 1$ receptor in thalamus, which also blocks ocular dominance plasticity in primary visual cortex. However, the adult plasticity is not dependent on feedback from primary visual cortex, an important difference from juveniles. These findings are an important contribution to a growing body of work identifying plasticity in the adult visual system, and identifies the visual thalamus as a potential target for therapies to reverse adult amblyopia.

Reviewer #2 (Public Review):

In this work, the authors found in the mouse line of GABA $\alpha 1$ subunit KO in thalamic neurons, which was previously reported lacking ocular dominance (OD) plasticity in juvenile V1 and dLGN (Sommeijer et al., 2017), the adult V1 and dLGN OD plasticity was also missing.

Through muscimol inhibiting the V1 feedback, thalamic OD plasticity was unaffected in both WT and KO adult mice. However, during the critical period, the thalamic OD plasticity was dependent on V1 feedback in WT mice.

Strengths:

1. The experiments were well designed. The authors used both MD and No MD controls with both WT and KO mice. The authors used in vivo SU recording, which is broadly accepted as the major method for evaluating OD plasticity.
2. The data analysis was solid. The authors used proper statistical tests for non-parametric data set.

Weaknesses:

1. In my previous review I pointed out that an alternative interpretation of the results is that the lack of OD plasticity in adult V1 and dLGN was caused by an early blockade of the development of the inhibitory circuit in dLGN, which causes life-long deficits in the functional connection of dLGN. The best way to rule out this possibility is by using conditional KO mice that dLGN synaptic inhibition was only interfered in adulthood. In response to my concern, the authors replied with a long text of reasoning why the current results are solid enough and the proposed experiment was unnecessary. I agree with most of the explanation that the current conclusion is solid, but I still think that the cKO experiment will be a good supplement to the current study, and if we do see a similar result in the cKO mice, the conclusion that the adult perturbation of thalamic inhibitory circuit interfere with the OD plasticity will be more convincing. However, I do understand that repeating the experiments again in another mouse line will be difficult and time-consuming, so the authors could choose if they want to perform the experiment or not.
2. Now the discussion part is very long and complex. Rearranging the discussion with sub-sections will make it easy to read.

Author Response

The following is the authors' response to the original reviews.

Reviewer #1 (Recommendations For The Authors):

Please describe the criteria for binocularity of dLGN neurons, and what % of recorded neurons meet this criteria. Do all the example neurons in figure 1D meet the criteria for binocular neurons?

We now include criteria for binocularity of dLGN neurons in the methods section on page 24, and mention the percentage of binocular neurons that we detected. We also indicate which of the example neurons in figure 1D are monocular or binocular according to these criteria. We would like to stress that these percentages are not representative for the level of binocularity in dLGN as a whole, as our recordings were limited to the frontal ipsilateral projection zone of dLGN, which is its most binocular region, and only units with a receptive field within 300 from the center were included in the analysis. We mention this in the discussion on page 23.

Fig 1: Please perform statistical comparison of data presented in Figure 1c by genotype, as in other figures.

We conducted post-hoc Tukey's tests exclusively when a significant interaction between phenotype and genotype was detected in the two-way ANOVA (as seen in Figs. 2B and 3E). This decision was made because interpreting a significant post-hoc test becomes uncertain when there is no interaction, which is evident in Fig. 1C. In that case, the posthoc Tukey's test yielded a p-value of 0.044 for the difference in RF size between KO NOMD and KO MD, while all other comparisons were not significant (WT NO-MD vs WT MD: $P=0.15$, WT NO-MD vs KO NO-MD: $p=0.99$, WT MD vs KO MD: $p=0.21$). However, since there was no significant interaction between genotype and phenotype, we cannot conclude that there is an effect in KO mice that is absent in WT mice. In Fig. 3B, all posthoc Tukey's tests resulted in P-values greater than 0.05.

Fig 1e: There is no justification for splitting the data into two time epochs before and after 150 msec. A repeated measures anova of smaller time bins across the full time course would be more effective/appropriate here.

The reviewer is correct. We have now performed a repeated measures ANOVA.

Fig 2: GABA $\alpha 1R$ KO results in a loss/absence of OD plasticity, not a reduction

We agree. We have changed the wording.

Fig 3: Please be specific about the location of V1 recordings. Was layer-specificity determined?

The location of V1 recordings is mentioned in the methods section under "Electrophysiology recordings, visual stimulation and V1 silencing", page 23. We have assessed OD per depth, but found that we do not have sufficient units to draw any conclusions about differences in plasticity per layer.

Why is feedback from V1 more influential in dLGN OD plasticity in KO?

We believe this is because the reduced thalamic inhibition causes the excitation/inhibition ratio to shift in favor of excitation. We discuss this more extensively on page 19 of the discussion.

Fig 4: Inclusion of a GABA αR antagonist protects thalamic axons from muscimol silencing (Liu BH, Wu GK, Arbuckle R, Tao HW, Zhang LI. Defining cortical frequency tuning with recurrent excitatory circuitry. Nat. Neurosci. 2007;10:1594-600.)

We now mention the possible direct influence of muscimol on thalamic axons in the discussion on page 19 and cite the suggested article.

The observation that feedback from primary visual cortex does not contribute to adult visual thalamus plasticity is interesting and important. The authors should expand on their discussion of this observation to include changes in cortical circuitry that may help to explain this observation.

We have expanded this part of the discussion on page 20.

The authors should describe the pathway by which inhibition enables plasticity in dLGN.

We discuss this more extensively on page 17 in the updated manuscript.

Reviewer #2:

1. The current work was basically a follow-up of a previous study in juvenile mice, and the results were also very similar to the juvenile results (Sommeijer et al., 2017). One possible interpretation of the results is that the lack of OD plasticity in adult V1 and dLGN was caused by an early blockade of the development of the inhibitory circuit in dLGN, which retains the dLGN in an immature stage till adulthood. The authors indeed claimed in the discussion that the 2-day OD shift is intact in juvenile dLGN and V1 in KO mice, and provided evidence in supplementary figure that GABAergic and cholinergic synapse amount are similar between WT and KO mice. However, the 7-day OD shift is indeed defected in juvenile V1 and dLGN in KO mice (Sommeijer et al., 2017), and it is possible that this early functional deficit didn't induce a structural remodeling in adulthood. To better support the author's claim that the lack of adult V1 OD plasticity is specifically due to reduced dLGN synaptic inhibition, the author should generate conditional KO mice that dLGN synaptic inhibition was only interfered in adulthood.

In order to address this criticism it is important to discuss the plasticity deficits in dLGN and V1 separately.

Concerning V1 plasticity: We have previously shown that brief MD induces an OD shift in V1 of mice lacking thalamic synaptic inhibition in dLGN. OD plasticity induced by brief MD is a hallmark of critical period plasticity in V1, and it thus seems highly unlikely that critical period onset in V1 is defective or that development of V1 is halted in an immature state that does not support OD plasticity in thalamus-specific GABRA1 deficient mice.

The observed plasticity deficit during the critical period was limited to the second stage of the OD shift in V1, which requires long-term monocular deprivation. The straightforward explanation for this result and our current findings is that both during the critical period and in adulthood, the second stage of OD plasticity in V1 induced by long-term monocular deprivation requires thalamic plasticity or inhibition. The proposed alternative, that lack of thalamic synaptic inhibition during development results in a possible lack of structural change in V1 that would cause a lifelong deficiency selectively affecting OD plasticity induced by long-term monocular deprivation, requires many more assumptions.

Concerning dLGN plasticity: The simplest explanation for the observed lack of OD plasticity in dLGN is that it is a direct consequence of the absence of synaptic inhibition in the KO mice. However, an alternative explanation could indeed be that dLGN is kept in an immature (pre-critical period-like) state due to the developmental absence of synaptic inhibition. This situation would be analogous to that in V1 of GAD65 deficient mice (which have reduced GABA release), in which OD plasticity cannot be induced by brief monocular deprivation during the critical period or in adulthood (Fagiolini and Hensch, 2000). Because this deficit can be reversed by treating the mice with benzodiazepines (allosteric modulators of GABA receptors) at any age, it is thought that development of V1 in GAD65 mice is halted in a pre-critical period-like state until inhibition is strengthened. We cannot exclude that something similar occurs in dLGN of mice lacking thalamic synaptic inhibition, although we did not observe any changes in hallmarks of dLGN maturity, such as reduced receptive field size, and increased cholinergic and inhibitory bouton densities.

However, if the analogy with the developmental deficit in V1 of GAD65 deficient mice is valid, the reduced plasticity is still likely to be a direct consequence of reduced inhibition. In GAD65 deficient mice, long term monocular deprivation during the critical period causes a full OD shift, showing that no additional deficits (besides reduced inhibition) limit OD plasticity in V1 of these mice (Fagiolini and Hensch 2000). And, as already mentioned, increasing inhibition rescues OD plasticity in GAD65 KO mice. Thus, the immature state of V1 in these mice is

probably nothing more than a situation in which inhibition tone is too low to support efficient OD plasticity. In dLGN, knocking out GABRA1 at a later age could therefore also create a situation in which inhibition is too low to support thalamic OD plasticity, which is not different from the situation in which the gene is inactivated at birth. Only if lack of synaptic inhibition in thalamus affects another, unknown developmental process that is of importance later in life to support OD plasticity in dLGN, the proposed experiment would result in a different outcome. We are not convinced that this scenario is likely enough to justify repeating most of this study, but now using mice in which GABRA1 is inactivated in dLGN through bilateral AAV-cre injections.

Independently of the exact cause of the plasticity deficit in dLGN, our results make clear that a cortical plasticity deficit in adulthood can have a thalamic origin, which we believe is an important insight that is highly relevant.

We have included part of these arguments in the discussion on page 17.

1. The authors found that in juveniles, dLGN OD shift is dependent on V1 feedback, but not in adults. However, a recent work showed that the effects of V1 silencing on dLGN OD plasticity could differ with various starting points and duration of the V1 silencing and MD (Li et al., 2023). Could the authors provide more details of the MD and V1 silencing for an in-depth discussion?

We discuss some of the findings of the Li et al paper on pages 16 and 20 of the manuscript now.