

Immunology and Inflammation

Mouse gingival single-cell transcriptomic atlas: An activated fibroblast subpopulation guides oral barrier immunity in periodontitis

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

2 June 2023 (this version)

Sent for peer review

20 April 2023

Posted to bioRxiv

14 April 2023

Takeru Kondo, Annie Gleason, Hiroko Okawa, Akishige Hokugo, Ichiro Nishimura 

Weintraub Center for Reconstructive Biotechnology, UCLA School of Dentistry, Los Angeles, CA 90095, USA •

Division of Molecular and Regenerative Prosthodontics, Tohoku University Graduate School of Dentistry, Sendai,

Miyagi 980-8575, Japan • UCLA Bruin in Genomics Summer Program • Regenerative Bioengineering and Repair

Laboratory, Division of Plastic and Reconstructive Surgery, Department of Surgery, David Geffen School of

Medicine at UCLA, Los Angeles, CA 90095, USA

 (https://en.wikipedia.org/wiki/Open_access)

 (<https://creativecommons.org/licenses/by/4.0/>)

Abstract

Chronic cardiovascular and metabolic diseases have been linked with oral inflammation in the tooth-supporting gingiva. Therefore, elucidating the mechanisms underlying development of gingival inflammation may hold critical insight into the pathogenesis of these debilitating non-communicable diseases. Here, we report a unique fibroblast subpopulation activated to guide oral inflammation (AG fibroblasts), identified in a single-cell RNA sequencing-based gingival cell atlas constructed from the mouse ligature-induced periodontitis model. Collagen-XIV-positive AG fibroblasts localized beneath gingival epithelium express chemokine ligands and Toll-like receptor-related molecules upon ligature placement, which were linked to receptors expressed by neutrophils and lymphocytes, including innate lymphoid cells (ILCs). We further identify ILCs as the primary source of proinflammatory interleukin-17 cytokines and show that cervical alveolar bone resorption is absent in Rag2^{-/-}γc^{-/-}, but not Rag2^{-/-}, mice suggesting ILC3s mediate the human periodontitis-like phenotype. We therefore propose AG fibroblasts function as a previously unrecognized surveillant to orchestrate chronic gingival inflammation in periodontitis.

eLife assessment

The findings of this article provide **valuable** information on the changes of cell clusters induced by chronic periodontitis. The observation of a new fibroblast subpopulation, which was named as AG fibroblasts, was quite interesting, but needs further evidence. The strength of evidence presented is **incomplete**.

Introduction

The oral mucosa must constantly respond to a diverse array of oral microbes, as well as to dietary and airborne antigens, making it one of the most active barrier tissues in the human

body^{(1), (2)}. Consequently, oral barrier immunity acts as a crucial surveillance system, which must defend from pathogenic organisms, while ensuring sustained tolerance of commensal microbes and harmless antigens, without excessive inflammation⁽³⁾. Within this system, the junction between the gingiva and the tooth surface represents a weak link, wherein the initiation of localized inflammation often leads to chronic gingivitis and periodontitis. Once this gingival inflammation progresses, adjacent alveolar bone supporting the cervical area of the dentition is subjected to a localized, but often severe, osteoclastic bone resorption, resulting in tooth loss and disruption of the maxillofacial structure⁽⁴⁾. Globally, periodontitis ranks among the most significant contributors to poor health and decreased quality of life, imposing substantial economic and healthcare burdens, particularly in low and middle-income countries⁽⁸⁾. It is therefore acutely important to elucidate the molecular mechanisms underlying the initiation of gingival inflammation, in order to facilitate the development of effective preventive and therapeutic measures for this debilitating condition.

Critically, periodontitis is not only the most frequent cause of tooth loss in adults⁽⁵⁾, it is also associated with a number of severe non-communicable diseases, such as metabolic and cardiovascular diseases. A recent systematic review revealed that type 2 diabetes mellitus patients had significantly worse periodontal status with more tooth loss, and the presence of severe periodontitis increased the incidence of type 2 diabetes mellitus by 53%⁽⁶⁾. Chronic inflammatory conditions are also known to increase the risk of cardiovascular diseases, including valvular heart disease. Recently, a possible association has been recognized between periodontal inflammation—distinct from transient bacteremia due to dental procedures—and atherosclerotic cardiovascular disease⁽⁷⁾. Although a direct causal relationship between these non-communicable diseases and periodontitis has not been established, it is postulated that periodontitis triggers an inflammatory cascade that begins in oral barrier tissues and subsequently spreads throughout the body. Thus, elucidating the mechanisms underlying periodontitis development may further provide insight on the signaling pathways that induce and sustain systemic inflammation.

Clinical characterization of periodontitis patients and observations made during treatment have not yielded a comprehensive understanding of the mechanisms that promote periodontal disease. Rather, our current model for the pathological framework of periodontitis has been reconstructed from animal models, harvested immune cell analyses, and oral microbial associations^{(4), (9)}. Of particular importance has been the development of a ligature-induced periodontitis model in mice, combined with targeted gene mutations⁽¹⁰⁾. This model has shown that fully developed periodontal inflammation is closely associated with the abundant recruitment and excessive activation of neutrophils⁽¹¹⁾ and pathological induction of interleukin (IL)-17-secreting proinflammatory effector CD4⁺ T helper (Th)17 cells⁽¹²⁾. Based on our current understanding of osteoimmunology, it was hypothesized that these pathological immune cells activate osteoclasts, leading to induction of periodontal alveolar bone resorption⁽¹³⁾⁽¹⁴⁾. However, the precise cascade of events leading to this pathological outcome has not been fully deciphered.

Investigation of the molecular mechanisms underlying chronic inflammation in immunological disorders is an intense area of research. Prior studies have shown that chemokines are involved in the trafficking and activation of inflammatory cells during both homeostasis and disease-associated inflammation⁽¹⁵⁾. These small proteins, which typically contain disulfide cysteine–cysteine (CC) and cysteine–X–cysteine (CXC) molecular signatures, act as ligands for cellular receptors that modulate inflammatory signaling pathways⁽¹⁶⁾. Such chemokine–receptor networks direct the migration of inflammatory cells, potentially amplifying the resulting tissue damage. However, in addition to immune cells, barrier tissues are also composed of epithelial cells and fibroblastic stromal cells, and increasing evidence suggests that stromal cells are involved in innate immune cell regulation^{(17)–(20)}. In particular, we and others have found that oral fibroblasts secrete cytokines and chemokines

in response to microbial stimuli or a proinflammatory environment, suggesting a possible mechanism by which they may regulate immune cell function^{(21), (22)}. Therefore, a comprehensive elucidation of chemokine–receptor signaling networks will likely need to include all types of gingival cells.

In the present study, to better understand the pathways contributing to chronic oral inflammation, we determined the gingival single-cell transcriptomic atlas of mice with ligature-induced periodontitis and identified a novel subpopulation of fibroblasts activated to guide chronic inflammation (AG fibroblasts). Our findings suggest that chemokines and ligands derived from AG fibroblasts bind to and activate receptors involved in neutrophil and lymphocyte activation. We further show that *Cd4*⁺ innate lymphoid cells (ILCs) produce proinflammatory IL-17 cytokines in the inflamed gingiva, and provide evidence that type 3 ILCs (ILC3s) are responsible for cervical alveolar bone resorption in mice—a pathological feature consistent with the human periodontitis phenotype. Thus, the results from this study identify the AG fibroblast–neutrophil–ILC3 axis as a previously unrecognized mechanism contributing to pathological inflammation in periodontitis.

Results

Alterations in major cell type proportions during periodontitis development

Periodontal disease was induced in C57BL/6J mice by placing a ligature around the maxillary left second molar⁽²³⁾; gingival defect developed after Day 3 following ligature placement (**Figure 1A** and **B**). Three-dimensional (3D) reconstruction of micro-computed tomography (microCT) images (**Figure 1C**) further revealed a reduction in alveolar bone height starting from Day 3, which progressively increased on Day 7 (**Figure 1D—figure supplement 1**). Overall, the observed pattern of periodontal tissue degradation was consistent with that reported in previous studies^{(24), (25)}.




Figure 1.

Changes in proportions of major cell types during periodontitis development in mice. **(A)** A ligature (5.0 silk suture) was placed around the maxillary second molar (M2) of wild-type (WT) mice. Representative intra-oral photographs of the maxilla on Day 0, prior to ligature placement (healthy gingiva), and on Days 1, 3, and 7 after ligature placement. **(B)** The gingival defect area was measured and normalized to the circumferential area of the maxillary first molar (M1) ($n = 6$). Gingival defects appeared on Day 3. **(C)** Representative micro-computed tomography (microCT) images of the maxilla taken from the lateral view. **(D)** Alveolar bone loss was determined from the total distance between the cemento-enamel junction (CEJ) and the alveolar bone crest (ABC) of the buccal or palatal bone at six sites in the ligated side ($n = 6$). Alveolar bone loss was apparent on Day 7. **(E)** Single-cell RNA sequencing (scRNA-seq) t -distributed stochastic neighbor embedding (t -SNE) projection plots showing the major cell types present within gingival tissue during periodontitis development on Day 0, 1, 4, and 7. Colors indicate cell type, as follows: green, epithelial cells; blue, fibroblasts; pink, endothelial cells; yellow, B cells; red, T cells; and purple, myeloid cells. **(F)** Proportion plots showing the relative amounts of each major cell type on Days 0, 1, 4, and 7. Significance was determined by analysis of variance (ANOVA), with Tukey's multiple-comparison test (**B**, **D**). Data are presented as mean values $\pm$ standard deviation (SD); $p < 0.05$ was considered statistically significant.

and purple, myeloid cells. **(F)** Proportion plots showing the relative amounts of each major cell type on Days 0, 1, 4, and 7. Significance was determined by analysis of variance (ANOVA), with Tukey's multiple-comparison test (**B**, **D**). Data are presented as mean values $\pm$ standard deviation (SD); $p < 0.05$ was considered statistically significant.

Left-side palatal gingiva tissue was harvested from mice on Day 0 (*i.e.*, healthy gingiva without ligature placement) and on Days 1, 4, and 7 after ligature placement, and gingival cells were dissociated for single-cell RNA sequencing (scRNA-seq)^{(22), (26)}. On Days 0 and 1, the major cell types identified included epithelial cells expressing cadherin 1 (*Cdh1*)⁽²⁷⁾ and type XVII collagen (*Col17a1*)⁽²⁸⁾, fibroblasts expressing type I collagen (*Col1a1*)⁽²⁹⁾ and lumican (*Lum*)⁽³⁰⁾, B cells expressing membrane spanning 4 domains A1 (*Ms4a1*)⁽³¹⁾ and cluster of differentiation 79A (*Cd79a*)⁽³²⁾, T cells expressing epsilon subunit of T cell receptor complex (*Cd3e*)⁽³³⁾ and cluster of differentiation 5 (*Cd5*)⁽³⁴⁾, and myeloid cells expressing lysozyme 2 (*Lyz2*)⁽³⁵⁾ and integrin subunit alpha M (*Itgam*)⁽³⁶⁾ (**Figure 1E** and **F—figure supplement 2**). On Days 4 and 7, an additional endothelial cell fraction expressing selectin P (*Selp*)⁽³⁷⁾ and selectin E (*Sele*)⁽³⁸⁾ emerged (**Figure 1E** and **F—figure supplement 2**), suggesting increased inflammatory neovascularization with the progression of periodontal inflammation. The proportion of B cells was increased on Day 1, and the proportion of myeloid cells increased progressively from Day 4 to 7 (**Figure 1F**). In addition, the proportion of fibroblasts was increased on Day 7 (**Figure 1F**).

Myeloid cell composition and activity during periodontitis development

The infiltration of proinflammatory neutrophils into the gingiva has been extensively characterized in this mouse model of periodontitis^{(11), (39)}. Here, we found that macrophages expressing cluster of differentiation 86 (*Cd86*)⁽⁴⁰⁾ and integrin subunit alpha X (*Itgax*)⁽⁴¹⁾ were predominant on Day 0 in the myeloid cell fraction from healthy gingiva (**Figure 2A** and **B—figure supplement 3**). In contrast, the proportion of neutrophils expressing CXC motif chemokine receptor (*Cxcr2*)⁽⁴²⁾ and G0/G1 switch gene 2 (*G0s2*)⁽⁴³⁾ increased after ligature placement and during periodontitis development from Day 1 to 7 (**Figure 2A** and **B—figure supplement 3**), suggesting continuous neutrophil infiltration into the early and established gingival lesion. Moreover, after ligature placement, gingival neutrophils upregulated the expression of triggering receptor expressed on myeloid cells 1 (*Trem1*), indicating that these cells are activated and participating in the amplification of inflammatory signals^{(44), (45)} (**Figure 2C**). Strikingly, the *Trem1*-expressing activated neutrophils also show upregulation of matrix metalloproteinase 9 (*Mmp9*; **Figure 2C**)—a protein associated with extracellular matrix degeneration within gingival tissue⁽⁴⁶⁾. We further detected expression of tumor necrosis factor alpha (*Tnfa*) and transforming growth factor beta 1 (*Tgfb1*) in both macrophages and neutrophils on Day 0 and after ligature placement (**Figure 2D**). Collectively, these myeloid cell behaviors are consistent with those reported in prior studies on periodontitis development, thus validating our scRNA-seq data.




Figure 2.

Myeloid cell composition and activity in gingival tissue during periodontitis development. **(A)** t-SNE projection plots showing myeloid cell subpopulations in gingival tissue during periodontitis development on Days 0, 1, 4, and 7. Colors indicate cell type, as follows: green, macrophages and red, neutrophils. **(B)** Proportion plots showing the relative amounts of neutrophils and macrophages on Days 0, 1, 4, and 7. **(C)** Violin plots showing *Trem1* and *Mmp9* expression levels in myeloid cells on Days 0, 1, 4, and 7; both genes are upregulated in neutrophils after ligature placement. **(D)** Violin plots showing *Tgfb1* and *Tnfa* expression in myeloid cells on Days 0, 1, 4, and 7; no obvious induction is observed in response to ligature placement. Dot plots depicting expression levels of the C motif chemokine ligand (CCL) genes *Ccl2*, *Ccl3*, *Ccl4*, *Ccl6*, *Ccl9* **(E)**, and the CXC motif chemokine ligand (CXCL) genes *Cxcl2*, *Cxcl3*, *Cxcl4*, and *Cxcl6* **(F)**. Chemokine expression in myeloid cells was unrelated to progression of gingival inflammation from Day 1 to 7.

Myeloid cells are also known to stimulate other immune cells through the expression of CC motif chemokine ligands (CCLs) and CXC motif chemokine ligands (CXCLs), many of which

are associated with periodontitis development⁽⁴⁷⁾. Our scRNA-seq analysis revealed that macrophages expressed *Ccl2*, *Ccl9*, *Cxcl4*, and *Cxcl6*, and neutrophils expressed *Ccl3*, *Ccl4*, *Ccl6*, *Cxcl2*, and *Cxcl3* throughout periodontitis development (Figure 2E and F). These data suggest that chemokines and cytokines produced by macrophages and neutrophils in inflamed tissue may amplify and polarize the immune response toward chronic gingival inflammation.

Fibroblasts activated to guide leukocyte migration in periodontitis development

We previously identified two distinct subpopulations of gingival fibroblasts, differentiated by expression of type XIV collagen (*Col14a1*)⁽²²⁾, and these were also detected in our current scRNA-seq data from Day 0 to 7 (Figure 3A). Gene ontology (GO) enrichment analysis of the *Col14a1*-expressing fibroblast subpopulation revealed expression of major gene clusters related to immune regulation, including “Regulation of leukocyte migration” (Figure 3B). This immune regulatory phenotype appears to be unique to *Col14a1*-expressing fibroblasts, which are therefore referred to as ‘fibroblasts activated to guide leukocyte migration’ or AG fibroblasts. The other subpopulation of *Col14a1*-expressing fibroblasts appeared to keep typical fibroblastic features and is thus referred to as KT fibroblasts (Figure 3C). An additional fibroblast subpopulation expressing smooth muscle aortic actin 2 (*Acta2*) was detected on Day 7, and these were identified as myofibroblasts⁽⁴⁸⁾ (MF) (Figure 3A). We found that the proportions of KT and AG fibroblasts were equal on Day 0 and on Days 1 and 4 after ligature placement (Figure 3E and F). However, on Day 7, the proportion of AG fibroblasts decreased, and the MF fraction emerged (Figure 3E and F).




Figure 3.

Fibroblasts activated to guide leukocyte migration (AG fibroblasts) are one of three fibroblast subpopulations in gingival tissue during periodontitis development. (A) Violin plots showing gene expression levels of type I collagen (*Col1a1*), type XIV collagen (*Col14a1*), and smooth muscle aortic actin 2 (*Acta2*) in gingival fibroblast subpopulations during periodontitis development. AG, AG fibroblasts; KT, ‘keep typical’ (KT) fibroblasts; MF, myofibroblasts. Gene ontology (GO) enrichment analysis of the biological functions of AG fibroblasts and KT fibroblasts on Day 0 without ligature placement (B) and on Day 1 (C) and Day 7 (D) after ligature placement. (E) t-SNE projection plots showing fibroblast subpopulations in gingival tissue during periodontitis development. Colors indicate cell type, as follows: blue, KT fibroblasts; red, AG fibroblasts; and yellow, MFs. (F) Proportion plots showing the relative amounts of each fibroblast subpopulation on Days 0, 1, 4, and 7.

AG fibroblasts and immune surveillance in periodontitis development

To further characterize AG fibroblasts and their “Regulation of leukocyte migration” phenotype, we analyzed expression of CCL and CXCL chemokines within fibroblast subpopulations in our scRNA-seq dataset⁽¹³⁾. Results show that AG fibroblasts activated expression of *Ccl8*, *Ccl11*, *Ccl19*, *Cxcl1*, *Cxcl11*, and *Cxcl12* immediately after ligature placement on Days 1 and 4 (Figure 4A). Chemokine expression levels then decreased on Day 7, suggesting that AG fibroblasts might initiate early leukocyte migration into gingival tissue. Given that activation of Toll-like receptors (TLR) is known to increase chemokine expression, we further assessed expression of TLRs in fibroblast subpopulations. We found that AG fibroblasts displayed temporal activation of *Tlr2*, *Tlr3*, and *Tlr4* expression on Days 1 and 4 (Figure 4B). Similarly, the TLR downstream genes *Myd88*, *Irak1*, *Map3k7*, and *Rela* were also expressed on Days 1 and 4 (Figure 4B). These data suggest that AG fibroblasts may guide the establishment of an early inflammatory environment within the gingiva and thereby promote periodontitis pathogenesis.




Figure 4.

AG fibroblasts and immune surveillance in periodontitis development. (A) Dot plots depicting expression levels of the CCL genes *Ccl8*, *Ccl11*, *Ccl19*, *Cxcl1*, *Cxcl11*, and *Cxcl12* in gingival fibroblast subpopulations during periodontitis development. (B) Dot plots depicting expression levels of the Toll-like receptor and related genes *Tlr2*, *Tlr3*, *Tlr4*, *Myd88*, *Irak1*, *Map3k7*, and *Rela* in gingival fibroblast subpopulations during periodontitis development. Upregulation of chemokines and TLR-related molecules is predominantly observed in the AG fibroblast subpopulation. (C) Hematoxylin and eosin (HE) staining and immunohistochemical (IHC) staining for COL14A1 and CXCL12 in periodontal tissue on Day 1; scale bars, 100 μm (HE) and 20 μm, (IHC). Yellow arrows indicate COL14A1- and CXCL12-positive cells in the connective tissue papillae and periodontal ligament (PDL).

To validate the presence of AG fibroblasts, Day 1 gingiva and periodontal tissue histological sections were subjected to immunohistochemistry with antibodies against COL14A1 and CXCL12 (Figure 4C). We detected COL14A1- and CXCL12-positive AG fibroblasts localized near gingival epithelial cells in the connective tissue papillae and free gingiva, as well as in the cervical zone of the periodontal ligament (PDL) space (Figure 4C). We note that this localization pattern of AG fibroblasts appears to be highly suitable for early immune surveillance during periodontitis pathogenesis.

Role of AG fibroblasts in myeloid cell activation

The interaction between chemokine ligands and their receptors has been extensively studied. Here, to evaluate the interaction between chemokine ligands strongly expressed by AG fibroblasts and chemokine receptors in innate immune cells, we matched AG fibroblast-expressed chemokines to expression of their putative receptors in myeloid cells. Results show that AG fibroblasts may regulate macrophages via the expression *Ccl8* and *Ccl11*, which encode chemokines that can interact with CC chemokine receptors (CCRs) encoded by *Ccr2* and *Ccr5* in macrophages. Similarly, gene expression signatures suggest that AG fibroblast-mediated neutrophil regulation may occur through CCL8–CCR1, CXCL1–CXCR2, and CXCL12–CXCR4 interactions (Figure 5A and B). Notably, all chemokine–receptor pairs are expressed throughout periodontitis development, although expression of factors mediating the interaction between AG fibroblasts and macrophages was decreased on Day 7.




Figure 5.

Role of AG fibroblasts in myeloid cell activation. Interaction between chemokine ligands expressed by AG fibroblasts and their putative chemokine receptors expressed by myeloid cells during periodontitis development. Dot plots depicting expression levels of the CC chemokine receptor (CCR) and the CXC chemokine receptor (CXCR) genes *Ccr1*, *Ccr2*, *Ccr5*, *Ccr7*, *Cxcr2*, *Cxcr3*, and *Cxcr4* in myeloid cell subpopulations on Day 1 (A) and Day 7 (B) following ligature placement. NicheNet ligand–target matrix indicating the regulatory potential between active ligands expressed in fibroblasts and target genes expressed in myeloid cells from the p-EMT program on Day 1 (C) and Day 7 (D). (E) Dot plot depicting expression levels of active ligand genes from panel (C) in fibroblast subpopulations on Day 1. (F) Dot plots depicting expression levels of active ligand genes from panel (D) in fibroblast subpopulations on Day 7. (G) Dot plot depicting expression levels of target genes from panel (C) in myeloid cell subpopulations on Day 1. (H) Dot plot depicting expression levels of target genes from panel (D) in myeloid cell

subpopulations on Day 7. Results suggest a strong intercellular communication network from AG fibroblasts to neutrophils.

Intercellular communication may occur via other ligand–receptor interactions, which induce downstream target gene expression in recipient cells. Here, we performed NicheNet analysis to identify potential interactions between gingival fibroblasts and myeloid cells, revealing a trend toward increasing ligand–target interactions between fibroblasts and myeloid cells from Day 1 to Day 7 (Figure 5C and D). On Day 1, ligand expression was


present primarily in AG fibroblasts (Figure 5E), whereas on Day 7, MF fibroblasts also exhibited prominent ligand expression (Figure 5F). Analysis of target gene expression revealed that both macrophages and neutrophils expressed receptors capable of interacting with AG fibroblast ligands on Day 1 (Figure 5G). In contrast, on Day 7, receptor expression was almost exclusively present in neutrophils, suggesting these cells are primarily targeted by MFs and AG fibroblasts at later stages of periodontitis development (Figure 5H).

T cells and ILCs in periodontitis development

We next analyzed expression of T cell-related genes in our scRNA-seq dataset and found that the Day 0 healthy gingiva exclusively contained Th cells expressing *Cd4*⁽⁴⁹⁾. However, after ligature placement, cytotoxic T (Tc) cells expressing *Cd8*⁽⁴⁹⁾ emerged, and on Day 7, *Cd4*⁺*Zbtb16*⁺ ILCs and *Cd4*⁺*Foxp3*⁺ regulatory T (Treg) cells⁽⁵⁰⁾ were also detected (Figure 6A and B). The ILCs expressed *Nfil3*, a basic leucine zipper (bZIP) transcription factor required for ILC development⁽⁵¹⁾ (Figure 6C). However, to our surprise, expression of *Rory*, *Il17a*, and *Il17f* was only detected in ILCs (Figure 6C), indicating that this gingival subpopulation is predominantly composed of ILC3s.

Figure 6.

T cell subpopulations in periodontitis development. (A) Proportion plots showing the relative amounts of T cell subpopulations in gingival tissue during periodontitis development. Treg, T regulatory cells; ILC, innate lymphoid cells; Th, T helper cells; Tc, cytotoxic T cells. (B) Violin plots showing expression levels of the T cell marker genes *Cd8* (Tc), *Cd4* (Th), *Zbtb16* (ILC), and *Foxp3* (Treg) on Day 7 following ligature placement. (C) Violin plots showing expression levels of *Nfil3*, *Rory*, *Il17a*, *Il17f*, *Tbx21*, and *Gata3* on Day 7 following ligature placement. These gene signatures indicate that gingival ILCs primarily comprise type 3 ILCs (ILC3s). (D) Violin plots showing expression levels of the macrophage-colony stimulating factor (M-CSF)-encoding gene, *Csf1* (M-CSF), in each major cell type. (E) Violin plots showing expression levels of *Csf1* in fibroblast subpopulations, myeloid cell subpopulations, and T cell subpopulations. (F) Violin plots showing expression levels of the nuclear factor kappa-B ligand (RANKL)-encoding gene, *Tnfsf11*, in each major cell type. (G) Violin plots showing expression levels of *Tnfsf11* in fibroblast subpopulations, myeloid cell subpopulations, and T cell subpopulations.



ILC3s and Th17 cells share similar regulatory functions. However, the role of ILC3s in osteoimmunology has not been fully deciphered. Macrophage-colony stimulating factor (M-CSF), encoded by *Csf1*, and receptor activator of nuclear factor kappa-B ligand (RANKL), encoded by *Tnfsf11*, are known to play critical roles in osteoclast induction. We found that in gingival tissue, AG fibroblasts and neutrophils are the primary cellular sources of *Csf1*; ILC3s also express this gene, albeit at lower levels (Figure 6E). Expression of *Tnfsf11* was detected in AG fibroblasts and ILC3s (Figure 6G). These observations suggest that ILC3s may function as inducers of inflammatory osteoimmune responses, similar to Th17 cells, leading to osteoclastogenesis in periodontitis downstream of AG fibroblast activation.

ILC3s are critical for cervical alveolar bone resorption in the mouse periodontitis model

We next examined the role of ILC3s in periodontitis pathogenesis by measuring ligature-induced gingival defects and alveolar bone resorption in *Rag2*^{-/-} mice, which lack functional B, Th, and Tc cells, and *Rag2*-IL-2 receptor common gamma (*Il2rg*) double-knockout mice (*Rag2*^{-/-}*γc*^{-/-}), lacking all lymphocytes, including ILCs. After ligature placement, gingival defects developed similarly in wild-type (WT), *Rag2*^{-/-}, and *Rag2*^{-/-}*γc*^{-/-} mice (Figure 7A and B). However, we found that alveolar bone loss was decreased in *Rag2*^{-/-} mice and nearly eliminated in *Rag2*^{-/-}*γc*^{-/-} mice (Figure 7C and D—figure supplement 4A). MicroCT image analysis further indicated a better preservation of alveolar bone structure in *Rag2*^{-/-}*γc*^{-/-} mice, relative to the other groups (Figure 7C—figure supplement 4B). Histologically, osteoclastic resorption lacunae were observed on the alveolar bone surface at the cervical PDL and tooth apex PDL zones only in WT mice (Figure 7E and F). In addition, we detected a significant decrease in the number of tartrate-resistant acid phosphatase (TRAP)-positive osteoclasts in the cervical PDL zone of *Rag2*^{-/-}*γc*^{-/-} mice and in the apical PDL zone of both *Rag2*^{-/-} and *Rag2*^{-/-}*γc*^{-/-} mice (Figure 7G and H). Collectively, these data suggest that ILC3s, not Th17 cells, are responsible for cervical alveolar bone resorption in the mouse periodontitis model, a pathological phenotype consistent with human disease.




Figure 7.

ILC3s are critical for cervical alveolar bone resorption in the mouse periodontitis model. (A) Representative intra-oral photographs of maxilla from WT, *Rag2*^{-/-}, and *Rag2*^{-/-}*γc*^{-/-} mice taken on Day 7 following ligature placement. (B) The gingival defect area was measured and normalized to the circumferential area of M1 (*n* = 5). (C) Representative microCT images of the maxilla taken from the lateral view for the ligated side and from the contralateral view for the unligated side. (D) Alveolar bone loss was determined from the total distance between the CEJ and the ABC of the buccal bone or palatal bone at six sites in the ligated side (*n* = 6). (E) HE staining of the periodontal tissue on Day 7. gCT, gingival connective tissue; Bone, alveolar bone; PDL, periodontal ligament; scale bars, 100 μm. (F) Tartrate-resistant acid phosphatase (TRAP) staining of periodontal tissue from WT mice on Day 7; scale bar, 100 μm. Total number of TRAP-positive cells in a 0.01 mm²-area of the buccal and palatal bone in the cervical PDL site (G) and apical PDL site (H) (*n* = 6). Significance was determined by ANOVA, with Tukey's multiple-comparison test (D, G, H). Data are presented as mean values ± SD; *p* < 0.05 was considered significant.

The role of AG fibroblasts and neutrophils in ILC3 development in periodontitis

Based on our present data, we hypothesize that ILC3s within the gingival tissue play a pathological role in alveolar bone resorption in our mouse model. We therefore aimed to identify the cells that promote ILC3 development in mice. Similar to Th17 cells, ILC3 development was shown to be triggered by IL-6 and IL-23a^{(52), (53)}. A survey of our scRNA-seq data identified epithelial cells, fibroblasts, and myeloid cells as the source of *Il6* in gingival tissue (Figure 8A). On Days 1 and 4 following ligature placement, AG fibroblasts primarily expressed *Il6*, whereas neutrophils became the predominant source on Day 7. We further found that gingival epithelial cells comprise at least four different subpopulations, plus an additional group displaying an epithelial–mesenchymal transition (EMT) phenotype on Day 7; *Il6* was expressed by several of these epithelial subpopulations, including the EMT subgroup (Figure 8A—figure supplement 5A–D). *Il23a* was also detected in AG fibroblasts and myeloid cells (Figure 8B), with expression present in all epithelial cell subsets at various points in periodontitis development (Figure 8B—figure supplement 5D).




Figure 8.

The role of AG fibroblasts and neutrophils in ILC3 development in periodontitis. Violin plots showing expression levels of the genes encoding interleukin (IL)-6 (*Il6*) (A) and IL-23 (*Il23a*) (B) in each major cell type, fibroblast subpopulations, and myeloid cell subpopulations during periodontitis development. (C) Interaction between chemokine ligands strongly expressed by AG fibroblasts and their putative chemokine receptors expressed by T cells, including ILCs. Dot plots depict gene expression levels of *Ccr1*, *Ccr2*, *Ccr5*, *Ccr7*, *Ccr8*, *Cxcr2*, *Cxcr3*, and *Cxcr4* in T cell subpopulations on Day 7 following ligature placement. (D) Interaction between chemokine ligands strongly expressed by neutrophils and their putative chemokine receptors expressed by T cells. Dot plots depicting gene expression levels of *Ccr1*, *Ccr4*, and *Ccr5* in T cell subpopulations on Day 7 following ligature placement. (E) NicheNet ligand–target matrix denoting the regulatory potential between active ligands in fibroblasts and target genes in T cells from the p-EMT program on Day 7 following ligature placement. (F) NicheNet ligand–target matrix denoting the regulatory potential between active ligands in myeloid cells and target genes in T cells from the p-EMT program on Day 7 following ligature placement. (G) Dot plot depicting expression levels of active ligand genes from panel (E) in fibroblast subpopulations. (H) Dot plot depicting expression levels of active ligand genes from panel (F) in myeloid cell subpopulations. (I) Dot plot depicting expression levels of target genes from panel (E) in T cell subpopulations. (J) Dot plot depicting expression levels of target genes from panel (F) in T cell subpopulations.

Lastly, we evaluated potential interactions between chemokine ligands expressed by AG fibroblasts and neutrophils and chemokine receptors in innate immune cells. Our data suggest the presence of a chemokine–receptor association between ILC3s and both AG fibroblasts (**Figure 8C**) and neutrophils (**Figure 8D**), although interactions with other innate immune cells are also possible. NicheNet analysis further identified potential ligand–target gene associations between lymphocytes, including ILC3s, and both fibroblasts (**Figure 8E**) and myeloid cells (**Figure 8F**). Ligand expression was more prominent in AG fibroblasts than in other fibroblast subpopulations (**Figure 8G**) and elevated in neutrophils relative to macrophages (**Figure 8H**). Additionally, target gene expression was detected in ILC3s, although it was not specific to these cells (**Figure 8I, J**). Thus, in total, our data suggest a regulatory role for a newly identified AG fibroblast subpopulation in the gingiva, which appears to orchestrate chronic gingival inflammation, at least in the early stages, and to promote alveolar bone resorption via stimulation of neutrophils and ILC3s.

Discussion

In this study, we used scRNA-seq to generate a gingival cell transcriptomic atlas across the time course of periodontal inflammation development, leading to pathological alveolar bone resorption, in a mouse ligature-induced model of disease. Our results suggest the presence of a unique and previously uncharacterized subpopulation of gingival fibroblasts, referred to as AG fibroblasts, which show upregulated expression of TLRs and chemokines immediately after ligature placement (**Figure 3** and **4**). Previous studies have proposed that TLR-expressing gingival fibroblasts regulate innate immune responses in the oral barrier tissue^{(54), (55)}; however, the precise mechanisms underlying this fibroblast-mediated regulation remained poorly understood. Here, we found that not all gingival fibroblasts acquired an immune-sensing capability, but rather, AG fibroblasts represent a distinct fibroblast subpopulation, capable of upregulating TLRs and potentially responding to microbial and tissue damage signals. Consistent with this possibility, the temporal expression of TLRs was found to be associated with increased expression of CC and CXC chemokine ligands.

The signature gene of AG fibroblasts is *Col14a1*, which encodes a member of the Fibril-Associated Collagen with Interrupted Triple-helices (FACIT) superfamily⁽⁵⁶⁾. To our knowledge, our present findings represent the first observation of COL14A1 in the gingiva (**Figure 4**). However, in a previous study, we detected COL14A1-positive AG fibroblasts immediately under the gingival epithelium and in the cervical region of PDL, making them ideally situated to mediate immune surveillance for oral stimuli and microbial signals⁽²²⁾. Given the strategic localization within the gingiva and the temporal expression of TLRs and chemokines after ligature placement, we hypothesize that AG fibroblasts play a previously unrecognized role in innate immune regulation during periodontitis development.

Neutrophils are well-established as crucial players in periodontitis progression. One study found that gingival neutrophils in chronic periodontitis patients are longer-lived than those from healthy patients⁽⁵⁷⁾. Moreover, it was shown that dysregulated neutrophils can induce destruction of periodontal connective tissue through the production of toxic molecules, such as reactive oxygen species, leading to initiation of bone resorption^{(58), (59)}. Critically, neutrophil function in periodontitis is dependent on trafficking to the affected gingiva, and accordingly, we found that the proportion of neutrophils increased after ligature placement (**Figure 2**). Neutrophils were previously reported to express a relatively limited number of chemokine receptors, including CXCR2, CXCR4, and CCR1^{(60), (61)}. Consistent with prior studies, we detected expression of these chemokine receptors in gingival neutrophils and

linked them to ligands expressed by AG fibroblasts—CXCL1, CXCL12, and CCL8, respectively (Figure 5). In addition to chemokines, neutrophils can respond to various chemoattractants that modulate and finetune various cellular behaviors, such as migration direction, adhesion strength, and functional heterogeneity⁽⁶²⁾. Here, we observed evidence for substantial intracellular communication between AG fibroblasts and neutrophils through ligand–target gene interactions (Figure 5). Although validation of each putative ligand–receptor interaction is beyond the scope of this study, our data suggest that AG fibroblasts may play an important role in neutrophil trafficking in both early and established periodontitis lesions.

A primary pathological consequence of periodontitis is uncontrolled alveolar bone resorption, leading to tooth loss, which is thought to be mediated by osteoimmune pathways involving IL-17–expressing Th17 cells⁽⁶³⁾. In contrast, analysis of our scRNA-seq revealed expression of *Rory*, *Il17a*, and *Il17f* not in CD4⁺ Th cells, but rather, in lymphocytes with ILC characteristics (Figure 6). ILCs share phenotypic and functional features with CD4⁺ T cells, although they lack antigen-specific T cell receptors⁽⁵¹⁾. In particular, the gingival ILCs detected in our study strongly expressed *Rory*, *Il17a*, and *Il17f*, but not *Tbx21* and *Gata3*, indicating they are ILC3s, which are functionally similar to Th17 cells. Notably, recent clinical studies have reported the presence of ILCs in gingiva from human periodontitis patients^{(64), (65)} and leptin receptor–deficient mice⁽⁶⁶⁾. However, the specific role of these cells in alveolar bone loss had not been previously established.

In the present study, we explored the role of ILC3s in periodontitis-associated bone loss using Rag2^{−/−} and Rag2^{−/−}γc^{−/−} mice and observed differential patterns of alveolar bone resorption following ligature placement in these animals. Rag2^{−/−} mutation prevents V(D)J recombination required for generating immunoglobulin and T cell receptors, resulting in the production of functionally immature B and T cells, including Th17 cells. However, ILCs do not undergo genomic receptor rearrangements and, thus, are unaffected by Rag2^{−/−} mutation. In contrast, Rag2^{−/−}γc^{−/−} mice have the Rag2^{−/−} mutation in combination with the Il2rg^{−/−} mutation, which disables common γ chain cytokines (γc). Therefore, in addition to non-functional B and T cells, these animals also have defective γc-dependent ILCs. Here, we found that Rag2^{−/−} mice contained a reduced number of osteoclasts in the apical PDL area, indicative of reduced bone resorption in this region, whereas a significant loss of bone resorption was present in both the cervical and apical PDL areas of Rag2^{−/−}γc^{−/−} mice (Figure 7). Human periodontitis induces alveolar bone at the cervical PDL zone relative to the gingival legion. In contrast, apical alveolar bone resorption is observed in clinical cases of root canal infection (apical periodontitis). Therefore, our findings suggest that ILCs, which are differentially impacted by Rag2^{−/−}γc^{−/−} vs. Rag2^{−/−} mutation, may be involved in human periodontitis-like alveolar bone resorption near the site of gingival inflammation. It must be noted that Rag2^{−/−} and Rag2^{−/−}γc^{−/−} mice developed gingival defects similar to WT mice. However, this observation may be explained by results from our previous study, which suggest that early gingival degradation in the mouse model is mediated by cathepsin K (CTSK) secreted from gingival fibroblasts⁽²²⁾, rather than by activated lymphocytes.

Differentiation of ILC3s and Th17 cells is mediated by similar environmental cues, with IL-6 and IL-23a playing critical roles in both pathways^{(67), (68)}. Our present scRNA-seq data (Figure 8) suggest that AG fibroblasts and neutrophils are the primary cellular sources of *Il6* during the early and later stages, respectively, of periodontitis development. In contrast, *Il23a* was found to be expressed by a variety of cell types, such as AG fibroblasts, myeloid cells, and multiple subsets of epithelial cells, including those with an EMT phenotype⁽⁶⁹⁾. Further, CC and CXC chemokine–receptor associations between ILC3s and both AG fibroblasts and neutrophils appeared to be non-specific. Therefore, our data suggest that rather than a specific trigger, the collective gingival environment, which includes AG fibroblasts, might contribute to ILC3 differentiation.

In conclusion, based on our present findings, we propose that a previously unrecognized AG fibroblast subpopulation in the gingiva can facilitate immune surveillance and participate in the pathological regulation of innate immune cells, such as proinflammatory neutrophils, within oral barrier tissue (Figure 9). Moreover, we hypothesize that ILC3s in the inflamed gingiva play a critical role in pathological alveolar bone resorption in the mouse model of periodontitis and, potentially, in human disease. Thus, the newly proposed AG fibroblast–neutrophil–ILC3 axis may hold valuable clues for unraveling the pathological mechanisms underlying periodontitis development. Moreover, these findings also provide a basis for investigation of new preventive and therapeutic strategies to contain oral barrier inflammation and potentially sever the link between periodontitis and debilitating non-communicative metabolic and cardiovascular diseases.




Figure 9.

Schematic overview of the newly proposed AG fibroblast–neutrophil–ILC3 axis. We propose that periodontal inflammation is initiated by the activation of AG fibroblasts, which secrete chemokines and cytokines that recruit neutrophils to sites of tissue damage. Activated neutrophils and AG fibroblasts, in turn, activate ILC3s, leading to production of proinflammatory IL-17 cytokines. Ultimately, cervical alveolar bone resorption is facilitated by a localized osteoclastogenic environment, induced by activated ILC3s, together with AG fibroblasts, neutrophils, myofibroblasts, and gingival epithelial cells, including those with an epithelial-mesenchymal transition (EMT) phenotype.

Methods

Animal care

All protocols for animal experiments were reviewed and approved by the University of California Los Angeles (UCLA) Animal Research Committee (ARC# 2003-009) and followed the Public Health Service Policy for the Humane Care and Use of Laboratory Animals and the UCLA Animal Care and Use Training Manual guidelines. C57BL/6J WT, Rag2^{-/-}, and Rag2^{-/-}yc^{-/-} mice were purchased from the Jackson Laboratory (Bar Harbor, ME, USA). Animals had free access to regular rodent diet and water and were maintained in standard housing conditions with 12-h light/dark cycles in the Division of Laboratory Animal Medicine at UCLA.

Evaluation of gingival defect and alveolar bone resorption in a ligature-induced mouse model of periodontitis

A silk thread was gently tied around the left maxillary second molar of 8- to 12-week-old female WT, Rag2^{-/-}, and Rag2^{-/-}yc^{-/-} mice under general inhalation anesthesia with isoflurane (Henry Schein, Melville, NY, USA). On Days 1, 3, and 7 after ligature placement, WT mice were randomly chosen and euthanized by 100% CO₂ inhalation. WT mice without ligature

placement were used as Day 0 pre-periodontitis control. Rag2^{-/-} and Rag2^{-/-}γc^{-/-} mice were euthanized on Day 7. The palatal tissue was digitally photographed, and maxillae were harvested and fixed in 10% buffered formalin (Thermo Fisher Scientific, Waltham, MA, USA). The gingival defect area was measured from digital photographs, using the ImageJ Java-based image-processing program (NIH, Bethesda, MD, USA), and normalized to the circumferential area of the maxillary first molar. Fixed maxillae were subjected to microCT imaging at an energy level of 60 kV and 166 μA, and 3D images were reconstructed from microCT scans (Skyscan 1275: Bruker, Billerica, MA, USA). Alveolar bone loss was assessed at three sites (mesiobuccal cusp, distobuccal cusp, and distal cusp) of the first molar, two sites (mesiobuccal cusp and distobuccal cusp) of the second molar, and one site (buccal cusp) of the third molar by measuring the distance from the cemento-enamel junction (CEJ) to the alveolar bone crest (ABC) on the buccal and palatal side of the alveolar bone. Total bone loss was calculated from the six-site total CEJ–ABC distance. The bone volume/total volume (BV/TV) ratio, bone surface, trabecular number (Tb.N), and trabecular thickness (Tb.Th) in the buccal side of alveolar bone of the second molar were determined using the proprietary analysis program (CTan: Bruker, Billerica, MA, USA). Statistical analysis to assess differences among multiple experimental groups was performed using two-way analysis of variance (ANOVA) with Tukey's multiple-comparison test; $p < 0.05$ was considered to be statistically significant.

Single cell dissociation from mouse maxillary gingiva

On Days 1, 4, and 7 after ligature placement, mice were euthanized by 100% CO₂ inhalation. And maxillary gingival tissues ($n = 4$ per group) were harvested from freshly isolated mouse maxillae.

Collagenase II treatment

The tissues were cut into 1-mm pieces and placed immediately into digestion buffer, containing 1-mg/ml collagenase II (Life Technologies, Thermo Fisher Scientific, Waltham, MA, USA), 10-units/ml DNase I (Sigma–Aldrich, St. Louis, MO, USA), and 1% bovine serum albumin (BSA; Sigma–Aldrich) in Dulbecco's Modified Eagle Medium (DMEM; Life Technologies). The tissues were incubated in digestion buffer for 20 min at 37 °C on a shaker at 150 rpm and then passed through a 70-μm cell strainer. The collected cells were pelleted at 1,500 rpm for 10 min at 4 °C and resuspended in phosphate-buffered saline (PBS; Life Technologies), supplemented with 0.04% BSA (Cell suspension A).

Trypsin treatment

Immediately following collagenase II treatment, tissues were incubated in 0.25% trypsin (Life Technologies) and 10-units/ml DNase I for 30 min at 37 °C on a 150-rpm shaker. Trypsin was neutralized with fetal bovine serum (Life Technologies), and the tissues were passed through a 70-μm cell strainer and washed with DMEM. The collected cells were then pelleted at 1,500 rpm for 10 min at 4 °C and resuspended in PBS with 0.04% BSA (Cell suspension B). Cell suspension A and Cell suspension B were combined in one tube. An equal number of combined cell suspensions A and B from four animals per group were combined for scRNA-seq analysis (10X Genomics, San Francisco, CA, USA).

Cell clustering and Identification

Cell Ranger was used to align reads, generate feature–barcode matrices, and perform clustering and gene expression analyses on the scRNA-seq data, and the output from this program was analyzed using the R-program Seurat (<https://satijalab.org/seurat/>). Cells with

<2,400 genes detected or >1% mitochondrial gene expression were filtered out as low-quality cells. Individual gene counts for each cell were divided by the total gene counts for that cell and multiplied by a scale factor of 10,000; natural-log transformation was then applied to the counts. The FindVariableFeatures function was used to select 2,000 variable genes with default parameters, and the ScaleData function was used to scale and center the counts in the dataset. Principal component analysis and Uniform Manifold Approximation and Projection dimensional reduction were performed on variably expressed genes. The cluster markers were found using the FindAllMarkers function, and cell types were manually annotated based on the cluster markers. Cell types were assigned based on expression of cell marker genes, and gene expression within different cell types was displayed using dot plots and violin plots.

Histological analysis

Fixed maxillae were decalcified in 10% EDTA (Sigma–Aldrich) for 3 weeks and then embedded in paraffin. Histological cross-sections were stained with hematoxylin and eosin (HE) and evaluated on a light microscope. Adjacent paraffin sections (4 μ m) were subjected to a heat-induced epitope retrieval procedure and then immunohistochemically stained with polyclonal antibodies to COL14A1 (#PA5-49916, Thermo Fisher Scientific) and CXCL12 (#PA5-30603, Thermo Fisher Scientific) at a 1:100 dilution, followed by secondary antibody application, diaminobenzidine staining, and methylene blue counterstaining.

Using maxillary cross-sections of WT, Rag2^{-/-}, and Rag2^{-/-} γ c^{-/-} mice, osteoclasts were evaluated by TRAP staining using a commercially available kit (Acid Phosphatase TRAP kit, Sigma–Aldrich), according to the manufacturer's instructions. TRAP-positive cells were counted in the 0.01-mm⁽²⁾ area. Statistical analysis to assess the differences among multiple experimental groups was performed using two-way ANOVA with Tukey's multiple-comparison test; $p < 0.05$ was considered to be statistically significant.

Functional annotation and pathway enrichment analysis

Annotation and visualization of GO terms were performed by Metascape (<http://metascape.org/gp/index.html#/main/step1>). The top-100 differentially expressed genes in each population were input and filtered with the term “immune”. Filtered genes were then input, and only “biological process” gene sets were retrieved from the GO database.

Ligand–target matrix prediction

NicheNet (v.1.0.0, <https://github.com/saeyslab/nichenetr>) was used to predict interactions between cell types. In brief, the integrated Seurat object containing each cell subpopulation was input into the NicheNet Seurat wrapper. Sender cells and receiver cells were determined, and interactions between active ligands expressed by sender cells and target receptors expressed by receiver cells were predicted based on information in signaling and ligand–receptor databases.

Acknowledgements

This study was supported by NIH grants R01DE022550, R44DE025524, and C06RR014529 and by SINTX Technologies.

Author contributions

I.N. conceptualized the project and obtained funding. T.K. and A.H. performed animal studies.

H.O. collected microCT data from the animal model. T.K. and A.G. performed scRNA-seq and data analysis. T.K. and I.N. drafted the manuscript. All authors gave final approval and agreed to be accountable for all aspects of the work.

Competing interests

I.N. is a consultant for FUJI FILM Corp and BioVinc LLC and received a research fund from SINTX Technologies, Inc. I.N. and A.H. received a research fund from Maruho Co. Ltd. The other authors declare no competing interests.

Data availability statement

We specify where the data supporting the results of this study can be found and provide hyperlinks to publicly archived datasets analyzed or generated during the study, where applicable. scRNA-seq data are found in NIH Gene Expression Omnibus (GSE228635).

Supplemental Information




Figure supplement 1.

Alveolar bone loss in the ligature-induced mouse periodontitis model. Alveolar bone loss was assessed at the mesiobuccal cusp (M1-1), distobuccal cusp (M1-2), and distal cusp (M1-3) of the first molar, the mesiobuccal cusp (M2-1) and distobuccal cusp (M2-2) of the second molar, and the buccal cusp (M3) of the third molar by measuring the distance from the cementoenamel junction (CEJ) to the alveolar bone crest (ABC) on the buccal or palatal side of the alveolar bone ($n = 6$). Significance was determined by analysis of variance (ANOVA), with Tukey's multiple-comparison test. Data are presented as mean values $\pm$ standard deviation (SD); $p < 0.05$ was considered significant.




Figure supplement 2.

Identification of major cell types in mouse gingival tissue during periodontitis development by single-cell RNA sequencing (scRNA-seq). Violin plots showing expression levels of cell-type marker genes in each major cell type: epithelial cells, cadherin 1 (*Cdh1*) and type XVII collagen (*Col17a1*); fibroblasts, type I collagen (*Col1a1*) and lumican (*Lum*); endothelial cells, selectin P (*Selp*) and selectin E (*Sele*); B cells, membrane spanning 4 domains A1 (*Ms4a1*) and cluster of

differentiation 79A (*Cd79a*); T cells, epsilon subunit of T cell receptor complex (*Cd3e*) and cluster of differentiation 5 (*Cd5*); and myeloid cells, lysozyme 2 (*Lyz2*) and integrin subunit alpha M (*Itgam*).




Figure supplement 3.

Identification of myeloid cell subpopulations in mouse gingival tissue during periodontitis development by scRNA-seq. Violin plots showing expression levels of macrophage and neutrophil marker genes in myeloid cell subpopulations: macrophages, cluster of differentiation 86 (*Cd86*) and integrin subunit alpha X (*Itgax*); and neutrophils, CXC motif chemokine receptor 2 (*Cxcr2*) and G0/G1 switch gene 2 (*G0s2*).




Figure supplement 4.

Effects of innate lymphoid cell (ILC) deletion on alveolar bone loss in the mouse periodontitis model. (A) Alveolar bone loss was assessed at six sites by measuring the distance from the CEJ to the ABC on the buccal or palatal side of the alveolar bone of wild-type (WT), *Rag2*^{-/-}, and *Rag2*^{-/-}*yc*^{-/-} mice on Day 7 following ligature placement (*n* = 6). (B) Bone volume/total volume (BV/TV), bone surface, trabecular number (Tb.N), and trabecular thickness (Tb.Th) in the buccal side of alveolar bone of the second molar were measured on Day 7 (*n* = 6). Significance was determined by ANOVA, with Tukey's multiple-comparison test. Data are presented as mean values ± SD; *p* < 0.05 was considered significant.




Figure supplement 5.

Epithelial cell subpopulations involved in periodontitis development in mice. (A) *t*-distributed stochastic neighbor embedding (*t*-SNE) projection plots showing epithelial cell subpopulations in gingival tissue on Days 1, 4, and 7 following ligature placement. Colors indicate cell type, as follows: dark green, epithelial cell population 1 (Epi 1); sea green, epithelial cell population 2 (Epi 2); pale green, epithelial cell population 3 (Epi 3); light green, epithelial cell population 4 (Epi 4); and light blue, epithelial cell population expressing epithelial-mesenchymal transition (EMT) genes. (B) Heatmap showing expression of the top-10 differentially expressed genes in Epi 1, Epi 2, Epi 3, Epi4, and EMT. (C) Violin plots showing gene expression levels of keratin 5 (*Krt5*), keratin 14 (*Krt14*), cadherin 11 (*Cdh11*), *Col1a1*, tenascin C (*Tnc*), and smooth muscle aortic actin 2 (*Acta2*) in epithelial cell subpopulations. (D) Violin plots showing gene expression levels of interleukin 6 (*Il6*) and interleukin 23A (*Il23a*) in epithelial cell subpopulations.

References

- Abusleme L , et al. (1970) **The subgingival microbiome in health and periodontitis and its relationship with community biomass and inflammation** *ISME J* 7:1016–1025
- Belkaid Y , Harrison OJ. (1970) **Homeostatic Immunity and the Microbiota** *Immunity* 46:562–576
- Moutsopoulos NM , Moutsopoulos HM (1970) **The oral mucosa: A barrier site participating in tissue-specific and systemic immunity** *Oral Dis* 24:22–25
- Lamont RJ , Koo H , Hajishengallis G (1970) **The oral microbiota: dynamic communities and host interactions** *Nat Rev Microbiol* 16:745–759
- Helal O , Gostemeyer G , Krois J , Fawzy El Sayed K , Graetz C , Schwendicke F. (1970) **Predictors for tooth loss in periodontitis patients: Systematic review and meta-analysis** *J Clin Periodontol* 46:699–712
- Wu CZ , et al. (2020) **Epidemiologic relationship between periodontitis and type 2 diabetes mellitus** *BMC Oral Health* 20

7. Naderi S , Merchant AT (1970) **The Association Between Periodontitis and Cardiovascular Disease: an Update** *Curr Atheroscler Rep* **22**
8. Peres MA , et al. (1970) **Oral diseases: a global public health challenge** *Lancet* **394**:249–260
9. Hajishengallis G , Chavakis T (1970) **Local and systemic mechanisms linking periodontal disease and inflammatory comorbidities** *Nat Rev Immunol* **21**:426–440
10. Lin P , et al. (1970) **Application of Ligature-Induced Periodontitis in Mice to Explore the Molecular Mechanism of Periodontal Disease** *Int J Mol Sci* **22**
11. Hajishengallis G (1970) **New developments in neutrophil biology and periodontitis** *Periodontol* **82**:78–92
12. Hasiakos S , Gwack Y , Kang M , Nishimura I (1970) **Calcium Signaling in T Cells and Chronic Inflammatory Disorders of the Oral Cavity** *J Dent Res* **100**:693–699
13. Cekici A , Kantarci A , Hasturk H , Van Dyke TE (1970) **Inflammatory and immune pathways in the pathogenesis of periodontal disease** *Periodontol* **64**:57–80
14. Ikeuchi T , Moutsopoulos NM (1970) **Osteoimmunology in periodontitis; a paradigm for Th17/IL-17 inflammatory bone loss** *Bone* **163**
15. Sokol CL , Luster AD (1970) **The chemokine system in innate immunity** *Cold Spring Harb Perspect Biol* **7**
16. Hughes CE , Nibbs RJB (1970) **A guide to chemokines and their receptors** *FEBS J* **285**:2944–2971
17. Hodzic Z , Schill EM , Bolock AM , Good M (1970) **IL-33 and the intestine: The good, the bad, and the inflammatory** *Cytokine* **100**:1–10
18. Ina K , et al. (1970) **Intestinal fibroblast-derived IL-10 increases survival of mucosal T cells by inhibiting growth factor deprivation and Fas-mediated apoptosis** *J Immunol* **175**:2000–2009
19. Pinchuk IV , et al. (1970) **PD-1 ligand expression by human colonic myofibroblasts/fibroblasts regulates CD4+ T-cell activity** *Gastroenterology* **135**:1228–1237
20. Cheng HW , et al. (1970) **CCL19-producing fibroblastic stromal cells restrain lung carcinoma growth by promoting local antitumor T-cell responses** *J Allergy Clin Immunol* **142**:1257–1271
21. Williams DW , et al. (1970) **Human oral mucosa cell atlas reveals a stromal-neutrophil axis regulating tissue immunity** *Cell* **184**:4090–4104

22. Kondo T , et al. (1970) **Oral microbial extracellular DNA initiates periodontitis through gingival degradation by fibroblast-derived cathepsin K in mice** *Commun Biol* **5**
23. Abe T , Hajishengallis G (1970) **Optimization of the ligature-induced periodontitis model in mice** *J Immunol Methods* **394**:49–54
24. Tamura H , Maekawa T , Hiyoshi T , Terao Y (1970) **Analysis of Experimental Ligature-Induced Periodontitis Model in Mice** *Methods Mol Biol* :237–250
25. Marchesan J , et al. (1970) **An experimental murine model to study periodontitis** *Nat Protoc* **13**:2247–2267
26. Okawa H , et al. (1970) **Mechanism of bisphosphonate-related osteonecrosis of the jaw (BRONJ) revealed by targeted removal of legacy bisphosphonate from jawbone using competing inert hydroxymethylene diphosphonate** *Elife* **11**
27. Groeger S , Meyle J (1970) **Oral Mucosal Epithelial Cells** *Front Immunol* **10**
28. Kamaguchi M , et al. (1970) **High Expression of Collagen XVII Compensates for its Depletion Induced by Pemphigoid IgG in the Oral Mucosa** *J Invest Dermatol* **138**:1707–1715
29. Takahashi A , et al. (1970) **Autocrine regulation of mesenchymal progenitor cell fates orchestrates tooth eruption** *Proc Natl Acad Sci U S A* **116**:575–580
30. Lallier TE , Spencer A , Fowler MM (1970) **Transcript profiling of periodontal fibroblasts and osteoblasts** *J Periodontol* **76**:1044–1055
31. Bruno KF , Silva JA , Silva TA , Batista AC , Alencar AH , Estrela C (1970) **Characterization of inflammatory cell infiltrate in human dental pulpitis** *Int Endod J* **43**:1013–1021
32. Sparger EE , et al. (1970) **Investigation of immune cell markers in feline oral squamous cell carcinoma** *Vet Immunol Immunopathol* **202**:52–62
33. Alcover A , Alarcon B , Di Bartolo V (1970) **Cell Biology of T Cell Receptor Expression and Regulation** *Annu Rev Immunol* **36**:103–125
34. Fujihashi K , et al. (1970) **Immunoregulatory function of CD3+, CD4-, and CD8- T cells. Gamma delta T cell receptor-positive T cells from nude mice abrogate oral tolerance** *J Immunol* **143**:3415–3422
35. Cross M , Mangelsdorf I , Wedel A , Renkawitz R (1970) **Mouse lysozyme M gene: isolation, characterization, and expression studies** *Proc Natl Acad Sci U S A* **85**:6232–6236
36. Okubo M , et al. (1970) **M2-polarized macrophages contribute to neovasculogenesis, leading to relapse of oral cancer following radiation** *Sci Rep* **6**

37. Gotsch U , Jager U , Dominis M , Vestweber D (1970) **Expression of P-selectin on endothelial cells is upregulated by LPS and TNF-alpha in vivo** *Cell Adhes Commun* **2**:7–14
38. Komatsu T , et al. (1970) **E-selectin mediates Porphyromonas gingivalis adherence to human endothelial cells** *Infect Immun* **80**:2570–2576
39. Silva LM , Brenchley L , Moutsopoulos NM (1970) **Primary immunodeficiencies reveal the essential role of tissue neutrophils in periodontitis** *Immunol Rev* **287**:226–235
40. Lam RS , et al. (1970) **Macrophage depletion abates Porphyromonas gingivalis-induced alveolar bone resorption in mice** *J Immunol* **193**:2349–2362
41. Agarbati S , et al. (1970) **Prognostic Relevance of Macrophage Phenotypes in High-grade Oral Tongue Squamous Cell Carcinomas** *Appl Immunohistochem Mol Morphol* **29**:359–365
42. Hashim A , Alsam A , Payne MA , Aduse-Opoku J , Curtis MA , Joseph S (1970) **Loss of Neutrophil Homing to the Periodontal Tissues Modulates the Composition and Disease Potential of the Oral Microbiota** *Infect Immun* **89**
43. Zhang F , Zhu YL , Deng WL , Zhu J , Zhang J (1970) **Activation of G0S2 is coordinated by recruitment of PML/RARalpha and C/EBPepsilon to its promoter during ATRA-induced APL differentiation** *J Leukoc Biol* **101**:655–664
44. Bouchon A , Dietrich J , Colonna M (1970) **Cutting edge: inflammatory responses can be triggered by TREM-1, a novel receptor expressed on neutrophils and monocytes** *J Immunol* **164**:4991–4995
45. Dopheide JF , et al. (1970) **Critical limb ischaemia is characterised by an increased production of whole blood reactive oxygen species and expression of TREM-1 on neutrophils** *Atherosclerosis* **229**:396–403
46. Corotti MV , et al. (1970) **Immunolocalization of matrix metalloproteinases-2 and -9 during apical periodontitis development** *Arch Oral Biol* **54**:764–771
47. Souto GR , Queiroz CM , Costa FO , Mesquita RA (1970) **Relationship between chemokines and dendritic cells in human chronic periodontitis** *J Periodontol* **85**:1416–1423
48. Sasaki H , et al. (1970) **Neuronal PAS Domain 2 (Npas2)-Deficient Fibroblasts Accelerate Skin Wound Healing and Dermal Collagen Reconstruction** *Anat Rec (Hoboken)* **303**:1630–1641
49. Mahanonda R , et al. (1970) **Memory T cell subsets in healthy gingiva and periodontitis tissues** *J Periodontol* **89**:1121–1130
50. Wei L , Xu M , Xiong H (1970) **An update of knowledge on the regulatory role of Treg cells in apical periodontitis** *Oral Dis* **27**:1356–1365

51. Eberl G , Colonna M , Di Santo JP , McKenzie AN. (1970) **Innate lymphoid cells. Innate lymphoid cells: a new paradigm in immunology** *Science* **348**
52. Bielecki P , et al. (1970) **Skin-resident innate lymphoid cells converge on a pathogenic effector state** *Nature* **592**:128–132
53. Powell N , et al. (1970) **Interleukin 6 Increases Production of Cytokines by Colonic Innate Lymphoid Cells in Mice and Patients With Chronic Intestinal Inflammation** *Gastroenterology* **149**:456–467
54. Qian SJ , et al. (1970) **Single-Cell RNA Sequencing Identifies New Inflammation-Promoting Cell Subsets in Asian Patients With Chronic Periodontitis** *Front Immunol* **12**
55. Naruishi K (1970) **Biological Roles of Fibroblasts in Periodontal Diseases** *Cells* **11**
56. Agarwal P , et al. (1970) **Collagen XII and XIV, new partners of cartilage oligomeric matrix protein in the skin extracellular matrix suprastructure** *J Biol Chem* **287**:22549–22559
57. Lakschevitz FS , Aboodi GM , Glogauer M (1970) **Oral neutrophil transcriptome changes result in a pro-survival phenotype in periodontal diseases** *PLoS One* **8**
58. Chapple IL , Matthews JB (1970) **The role of reactive oxygen and antioxidant species in periodontal tissue destruction** *Periodontol* **43**:160–232
59. Ryder MI (1970) **Comparison of neutrophil functions in aggressive and chronic periodontitis** *Periodontol* **53**:124–137
60. Sabroe I , Jones EC , Whyte MK , Dower SK (1970) **Regulation of human neutrophil chemokine receptor expression and function by activation of Toll-like receptors 2 and 4** *Immunology* **115**:90–98
61. Huston J , Muhm JR. (1970) **Solitary pulmonary nodules: evaluation with a CT reference phantom** *Radiology* **170**:653–656
62. Metzemaekers M , Gouwy M , Proost P (1970) **Neutrophil chemoattractant receptors in health and disease: double-edged swords** *Cell Mol Immunol* **17**:433–450
63. Deng J , Lu C , Zhao Q , Chen K , Ma S , Li Z (1970) **The Th17/Treg cell balance: crosstalk among the immune system, bone and microbes in periodontitis** *J Periodontal Res*
64. Brown JL , Campbell L , Malcolm J , Adrados Planell A , Butcher JP , Culshaw S (1970) **Enrichment of Innate Lymphoid Cell Populations in Gingival Tissue** *J Dent Res* **97**:1399–1405
65. Dutzan N , Konkel JE , Greenwell-Wild T , Moutsopoulos NM (1970) **Characterization of the human immune cell network at the gingival barrier** *Mucosal Immunol* **9**:1163–1172

66. Wang Q , et al. (1970) **Single-Cell Transcriptomic Atlas of Gingival Mucosa in Type 2 Diabetes** *J Dent Res* **101**:1654–1664
67. Klose CS , et al. (1970) **A T-bet gradient controls the fate and function of CCR6-RORgammat+ innate lymphoid cells** *Nature* **494**:261–265
68. Sawa S , et al. (1970) **Lineage relationship analysis of RORgammat+ innate lymphoid cells** *Science* **330**:665–669
69. Wadie KW , Bashir MH , Abbass MMS (1970) **Epithelial-mesenchymal transition in gingival tissues from chronic periodontitis patients: A case-control study** *Dent Med Probl* **58**:311–319

Author information

Takeru Kondo

Weintraub Center for Reconstructive Biotechnology, UCLA School of Dentistry, Los Angeles, CA 90095, USA, Division of Molecular and Regenerative Prosthodontics, Tohoku University Graduate School of Dentistry, Sendai, Miyagi 980-8575, Japan

Annie Gleason

Weintraub Center for Reconstructive Biotechnology, UCLA School of Dentistry, Los Angeles, CA 90095, USA, UCLA Bruin in Genomics Summer Program

Hiroko Okawa

Weintraub Center for Reconstructive Biotechnology, UCLA School of Dentistry, Los Angeles, CA 90095, USA, Division of Molecular and Regenerative Prosthodontics, Tohoku University Graduate School of Dentistry, Sendai, Miyagi 980-8575, Japan

Akishige Hokugo

Weintraub Center for Reconstructive Biotechnology, UCLA School of Dentistry, Los Angeles, CA 90095, USA, Regenerative Bioengineering and Repair Laboratory, Division of Plastic and Reconstructive Surgery, Department of Surgery, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095, USA

Ichiro Nishimura

Weintraub Center for Reconstructive Biotechnology, UCLA School of Dentistry, Los Angeles, CA 90095, USA

For correspondence: inishimura@dentistry.ucla.edu

Editors

Reviewing Editor

Bian Zhuan

Wuhan University, China

Reviewer #1 (Public Review):

In this article, the authors found a distinct fibroblast subpopulation named AG fibroblasts, which are capable of regulating myeloid cells, T cells and ILCs, and proposed that AG fibroblasts function as a previously unrecognized surveillant to orchestrate chronic gingival inflammation in periodontitis. Generally speaking, this article is innovative and interesting, however, there are some problems that need to be addressed to improve the quality of the manuscript.

Results:

1. It is recommended to add HE staining and immunohistochemistry staining to observe the inflammation, tissue damage, and repair status from 0 to 7 days, so that readers can understand cell phenotype changes corresponding to the periodontitis stage. The observation index can include inflammation and vascular related indicators.
2. Figure 1A-1D can be placed in the supplementary figure.
3. I suggest the authors to put the detection of the existence of AG fibroblasts before exploring its relationship with other types of cells.
4. The layout of the picture should be closely related to the topic of the article. It is recommended to readjust the layout of the picture. Figure 1 should be the detection of AG cells and their proportion changes from 0 to 7 days. In other figures, the authors can separately describe the proportion changes of myeloid cells, T cells and ILCs, and explored the association between AG fibroblasts and these cell types.

Methods:

It is recommended to separately list the statistical methods section. The statistical method used in the article should be one-way ANOVA.

Reviewer #2 (Public Review):

This study proposed the AG fibroblast-neutrophil-ILC3 axis as a mechanism contributing to pathological inflammation in periodontitis. However, the immune response in the vivo is very complex. It is difficult to determine which is the cause and which is the result. This study explores the relevant issue from one dimension, which is of great significance for a deeper understanding of the pathogenesis of periodontitis. It should be fully discussed.

1. Many host cells participate in immune responses, such as gingival epithelial cells. AG fibroblast is not the only cell involved in the immune response, and the weight of its role needs to be clarified. So the expression in the conclusion should be appropriate.
2. This study cannot directly answer the issue of the relationship between periodontitis and systemic diseases.

Author Response:

We thank all reviewers for constructive critiques. We plan to perform new experiments and revise our manuscript accordingly. The text and Figures are currently undergoing the revision process. Below highlights our revision plan.

eLife assessment

The findings of this article provide valuable information on the changes of cell clusters induced by chronic periodontitis. The observation of a new fibroblast subpopulation, which was named as AG fibroblasts, was quite interesting, but needs further evidence. The strength of evidence presented is incomplete.

RESPONSE: We discovered a new subpopulation of gingival fibroblasts, named AG fibroblasts, using non-biased single cell RNA sequencing (scRNA-seq) of mouse gingival samples undergoing the development of ligature-induced periodontitis. AG fibroblasts exhibited a unique gene expression profile: [1] constitutive expression of type XIV collagen; and [2] ligature-induced upregulation of chemokines such as CXCL12. As a biomedical data science experiment, we validated the scRNA-seq observation using immunohistochemical experiment, which showed the presence of type XIV collagen-positive and CXCL12-positive gingival fibroblasts localized immediately under the gingival epithelium and the coronal region of periodontal ligament.

We agree that the functional/pathological role of AG fibroblasts must be further explored. We have hypothesized that AG fibroblasts initially sense the pathological stress including oral microbial stimuli and secrete inflammatory signals through chemokine expression. To address this hypothesis, in this revision, we plan to analyze a separate scRNA-seq data for AG fibroblast gene expression profile derived from mouse gingival tissues that have been stimulated by Toll-Like Receptor 9 (TLR9) ligand (unmethylated CpG oligonucleotide) and TLR2/4 ligand (LPS). This approach mimics the initial pathological stress applied to gingival tissue. The new insight of AG fibroblasts will be presented in the revision.

Reviewer #1 (Public Review):

In this article, the authors found a distinct fibroblast subpopulation named AG fibroblasts, which are capable of regulating myeloid cells, T cells and ILCs, and proposed that AG fibroblasts function as a previously unrecognized surveillant to orchestrate chronic gingival inflammation in periodontitis. Generally speaking, this article is innovative and interesting, however, there are some problems that need to be addressed to improve the quality of the manuscript.

RESPONSE: We appreciate this comment. As suggested, we further investigated the surveillant function of AG fibroblasts by reanalyzing the scRNA-seq data for stress sensing receptors such as Toll-Like Receptors (TLR). Therefore, we analyzed AG fibroblast gene expression profile when the putative ligands to TLR2/4 and TLR9 are applied to mouse gingival tissue instead of ligature placement. We believe that this first step analysis should warrant to dissect further the function of AG fibroblasts in the future.

Results:

1. *It is recommended to add HE staining and immunohistochemistry staining to observe the inflammation, tissue damage, and repair status from 0 to 7 days, so that readers can understand cell phenotype changes corresponding to the periodontitis stage. The observation index can include inflammation and vascular related indicators.*

RESPONSE: As recommended, representative histological figures will be included. We will further perform new immunohistochemistry experiment of mouse gingival tissue (D0, D1, D4, D7). We plan to highlight the infiltration of CD45+ immune cells. We also plan to highlight the progressive degeneration of gingival collagen fiber by picrosirius red staining.

1. *Figure 1A-1D can be placed in the supplementary figure.*

RESPONSE: Combining the new data above, Figure 1 will be revised as suggested.

1. *I suggest the authors to put the detection of the existence of AG fibroblasts before exploring its relationship with other types of cells.*
1. *The layout of the picture should be closely related to the topic of the article. It is recommended to readjust the layout of the picture. Figure 1 should be the detection of AG cells and their proportion changes from 0 to 7 days. In other figures, the authors can separately describe the proportion changes of myeloid cells, T cells and ILCs, and explored the association between AG fibroblasts and these cell types.*

RESPONSE: As suggested, the presentation order of Figures and text will be revised to bring the information about AG fibroblasts first. The chemokine-receptor analysis is moved below.

Methods:

It is recommended to separately list the statistical methods section. The statistical method used in the article should be one-way ANOVA.

RESPONSE: A separate statistical method section is created. As pointed out, we used one-way ANOVA with post-hoc Tukey test (when multiple groups were compared).

Reviewer #2 (Public Review):

This study proposed the AG fibroblast-neutrophil-ILC3 axis as a mechanism contributing to pathological inflammation in periodontitis. However, the immune response in the vivo is very complex. It is difficult to determine which is the cause and which is the result. This study explores the relevant issue from one dimension, which is of great significance for a deeper understanding of the pathogenesis of periodontitis. It should be fully discussed.

RESPONSE: We agree with this comment. We expanded the current understanding of oral immune signal communication in Discussion and highlight how AG fibroblast may fit to it.

1. *Many host cells participate in immune responses, such as gingival epithelial cells. AG fibroblast is not the only cell involved in the immune response, and the weight of its role needs to be clarified. So the expression in the conclusion should be appropriate.*

RESPONSE: Following this critique, we revised INTRODUCTION, DISCUSSION and CONCLUSION, to highlight how AG fibroblasts function within a comprehensive immune response network.

1. *This study cannot directly answer the issue of the relationship between periodontitis and systemic diseases.*

RESPONSE: We agree with this critique. We either deleted or de-emphasized the relationship between periodontitis and systemic diseases throughout the text.