

Neuroscience

Retinal input integration in excitatory and inhibitory neurons in the mouse superior colliculus *in vivo*

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

2 June 2023 (this version)

Posted to bioRxiv

8 April 2023

Sent for peer review

7 April 2023

Carolin Gehr, Jérémie Sibille, Jens Kremkow ✉

Neuroscience Research Center, Charité-Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin • Bernstein Center for Computational Neuroscience Berlin, Philippstraße 13, 10115 Berlin • Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Philippstraße 13, 10115 Berlin • Einstein Center for Neurosciences Berlin, Charitéplatz 1, 10117 Berlin

 (https://en.wikipedia.org/wiki/Open_access)

 (<https://creativecommons.org/licenses/by/4.0/>)

Abstract

The superior colliculus (SC) is a midbrain structure that contains one of the highest densities of inhibitory neurons in the brain and, together with the thalamocortical visual system, it plays a key role in visually guided behaviors. The SC receives direct inputs from retinal ganglion cells (RGCs) but whether excitatory and inhibitory SC neurons differentially integrate retinal activity *in vivo* is still largely unknown. We recently established an extracellular recording approach using high-density electrodes to measure the activity of RGCs simultaneously with their postsynaptic SC targets *in vivo*, that allows addressing how SC neurons integrate RGC activity. Here, we employ this method to study the functional properties and dynamics that govern retinocollicular signaling in a cell-type specific manner by identifying GABAergic SC neurons using optotagging in anesthetized VGAT-ChR2 mice. We measured 305 monosynaptically connected RGC-SC pairs, out of which approximately one third of retinal afferents connect onto inhibitory SC neurons. We show that both excitatory and inhibitory SC neurons receive comparable strong RGC inputs, with functionally similar RGC-SC pairs showing stronger connections. Our results demonstrate that similar wiring rules apply for RGCs innervation of both excitatory and inhibitory SC neurons, which is unlike the cell-type specific connectivity in the thalamocortical system. Contrasting the similar RGC-SC connection strength, we observed that RGC activity contributed more to the activity of postsynaptic excitatory SC neurons than to the activity of postsynaptic inhibitory SC neurons. This implies that the excitatory SC neurons are more specifically coupled to RGC afferent inputs, while inhibitory SC neurons may integrate additional inputs from other sources. Taken together, our study deepens the understanding of cell-type specific retinocollicular functional connectivity and emphasizes that the two major brain areas for visual processing, the visual cortex and the superior colliculus, differently integrate sensory afferent inputs.

eLife assessment

This **valuable** study contributes to understanding how retinal activity shapes the response properties of excitatory and inhibitory neurons in a major visual target, the superior colliculus. The evidence supporting the claim is **solid**: the work is technically excellent but the interpretation was limited without more detailed knowledge of cell types involved and the lack of loss or gain of function manipulations. This work will be of interest to visual neuroscientists.

Introduction

The mouse superior colliculus (SC) is a midbrain structure that receives direct inputs from retinal ganglion cells (RGCs) (Ellis et al., 2016; Kremkow and Alonso, 2018). Together with the visual cortex (Glickfeld et al., 2013; Lashley, 1931; Niell and Scanziani, 2021; Petruno et al., 2013), the SC plays a key role in visually guided behaviors (Evans et al., 2018; Hoy et al., 2019; Shang et al., 2019; 2015; Wei et al., 2015). Intriguingly, the visual SC layers contain one of the highest densities of inhibitory (GABAergic) neurons in the brain (Kaneda et al., 2008; Kasai and Isa, 2016) and motion processing (Barchini et al., 2018; Gale and Murphy, 2016), but also in the regulation of wakefulness (Zhang et al., 2019). However, how GABAergic SC neurons are recruited by their retinal inputs *in vivo* remains largely unknown (Shi et al., 2017). An understanding how RGC inputs are integrated by SC INs and SC excitatory neurons (EXNs) is crucial for reaching a mechanistic understanding of the computations within the SC microcircuit.

In sensory systems, the divergence of long-range afferent axons onto GABAergic and non-GABAergic neurons ensures a balance between excitation and inhibition (E/I) (Miller, 2016). For example, in sensory cortices, thalamic afferents differentially activate excitatory and local inhibitory neurons (Cruikshank et al., 2007; Ji et al., 2016) with a stronger drive onto inhibitory neurons compared to excitatory neurons (Bruno and Simons, 2002; Cruikshank et al., 2007; Gabernet et al., 2005; Gibson et al., 1999; Inoue and Imoto, 2006; Swadlow, 2003). This stronger drive on inhibitory neurons establishes an effective feedforward inhibition (Agmon and Connors, 1992; Bereshpolova et al., 2020; Gabernet et al., 2005; Kremkow et al., 2010a; 2010b; Sun et al., 2006) that balances the excitation from thalamic afferents (Isaacson and Scanziani, 2011) and contributes to sharpening stimulus feature selectivity (Lee et al., 2012). Moreover, this wiring motif can partly be linked to the response properties of cortical inhibitory neurons as they typically show higher firing rates, broader tuning, low selectivity and high sensitivity (Alonso and Swadlow, 2005; Bruno and Simons, 2002; Gibson et al., 1999; Kremkow et al., 2016; Porter et al., 2001; Swadlow et al., 2002) to facilitate feedforward inhibition in response to thalamic input. Despite the importance of this afferent wiring motif for sensory processing in the thalamocortical visual circuit, it is currently unknown whether the retinocollicular pathway follows similar or different principles.

We recently demonstrated that SC neurons can receive strong input from RGC axons *in vivo* (Sibille et al., 2022a). Moreover, around one third of retinocollicular synapses connect onto inhibitory neurons that form intrinsic connections within the SC (Whyland et al., 2020). These findings support the notion that an effective feedforward inhibition might also be at play in the retinocollicular system. But whether afferent inputs are differentially integrated by excitatory and inhibitory SC neurons, similar to the thalamocortical system, is unknown. However, our previous study in the retinocollicular system lacks the cell-type specificity, and

it remains an open question whether the strong drive is specific to inhibitory SC neurons or is a general property of retinocollicular connections, and thus cell-type independent. Therefore, whether there are any differences in connection strengths between RGCs that project to excitatory and inhibitory SC neurons is different remains elusive. In this study, we investigate how excitatory and inhibitory neurons in the SC integrate retinal inputs *in vivo*.

We aim to answer these questions by characterizing the properties of the long-range synaptic connections between RGC-SC pairs employing a set of measurements: 1. The connection efficacy is used to study how strongly excitatory and inhibitory SC neurons are driven by their RGC input. 2. The functional similarity is analyzed between connected RGC-SC pairs to examine whether comparable wiring motifs govern retinocollicular signaling in excitatory and inhibitory neurons. 3. Short-term dynamics of retinal activity are studied to estimate whether the retinal spiking pattern shows paired-spike enhancement on retinocollicular synapses. Finally, 4. the connection contribution is employed to investigate how tightly the SC activity is coupled to retinal spiking.

This study intends to generate a more detailed picture of cell-type specific retinocollicular innervation and to fill a gap concerning the different long-range functional connectivity motifs that exist along the visual pathways.

Results

Recording excitatory and inhibitory neurons simultaneously with RGC axons in the mouse SC *in vivo*

To investigate the integration of retinal input in excitatory and inhibitory neurons in the SC *in vivo*, we recorded extracellular neural activity in the visual layers of SC in VGAT-ChR2 mice using high-density Neuropixels probes (Jun et al., 2017). We employed a tangential insertion approach to maximize the extent of visually driven channels in the SC (Fig. 1A) (Sibille et al., 2022a). Notably, we recently demonstrated that this approach allows us to not only measure spiking activity from SC neurons, but also to record electrical signals from retinal ganglion cell axons (RGC) that terminate at the SC (Sibille et al., 2022a). The somatic action potentials of SC neurons can be distinguished from the axonal action potentials of RGCs based on their spatiotemporal waveforms profile (Sibille et al., 2022a). RGC axonal waveforms are triphasic with a larger spatial spread (Fig. 1D left) while somatic SC neurons show a characteristic biphasic waveform profile (Fig. 1D, middle and right).




Fig. 1

Simultaneous extracellular recordings of RGC axons and SC neurons combined with optotagging identifies GABAergic neurons in VGAT-ChR2 mice.

A, Recording configuration for tangential electrode insertion and optotagging in the visual layers of mouse SC. The optogenetic fiber is inserted perpendicularly to the Neuropixels probe to activate GABAergic cells in VGAT-ChR2 mice. RGC afferents (orange) project onto GABAergic (blue) and non-GABAergic (black) neurons in SC. **B**, Multi-unit response (MUA) to optogenetic stimulation along the 384 recording sites during the presentation of a black screen. The large spatial extent of the optogenetic activation is evident. Gray vertical bar = light artifacts induced by the LED stimulation. **C**, Visually evoked MUA during the presentation of a sparse noise stimulus along the recording sites. Receptive field contours of recording sites with high signal-to-noise ratio. The color code reflects the location within the SC. Scale bar represents 10 degree. **D**, Spatiotemporal waveform profiles. Single channel waveforms identified at the peak channel (top) and multi-channel waveforms (bottom) for RGC axon (left, orange) and inhibitory (middle, blue) and excitatory (right, gray) SC neurons. Receptive fields (RFs) indicate visually responsive neurons. **E**, Identification of SC cell-types via optotagging. Raster plots and peri-stimulus time histograms (PSTHs) for single-neuron responses to blue light pulses (100ms) presented under baseline conditions. Excitatory SC neurons (EXNs, bottom) and RGCs (top) do not respond to the LED pulse, while GABAergic SC neurons respond to the light pulse with an increase in spiking response (middle). Optogenetic stimulation period is highlighted in blue. The colored scale bars on the right represent 20 Hz firing rate. **F**, PSTHs and raster plots for different cell types shown in E in response to a natural movie stimulus (10s, 30 trials). Note the high firing rate in RGCs. **G**, Top: Proportion of identified GABAergic (INs, blue), non-GABAergic (EXNs, gray) SC neurons and retinal axons (RGCs, orange) populations. Note that around one third of the captured SC neurons are GABAergic ($n = 326$ RGCs, $n = 468$ EXNs, $n = 212$ INs). Bottom: Mean firing rates in response to a natural movie stimulus presented for 10s, 30 trials. Two-sided Wilcoxon rank-sum test.

To study cell-type specific differences, we combined the Neuropixels recordings with an optogenetic identification strategy (optotagging) to identify GABAergic SC neurons in VGAT-ChR2-EYFP (Zhao et al., 2011) mice (Fig. 1A, B; data from $n = 11$ experiments and $n = 9$ mice included in the analysis, if not stated otherwise). In these transgenic mice, GABAergic

neurons specifically express Channelrhodopsin-2 and can be identified upon blue light stimulation. To photo-activate a large proportion of GABAergic SC neurons using short light pulses (100ms pulses, 2Hz, 2.7 mW) we inserted the optic fiber perpendicularly to the Neuropixels probe (Fig. 1A, right, see Material and Methods). Using this recording and light stimulation configuration we could activate GABAergic SC neurons across an average of ~ 1.5 mm of SC tissue (light-activated range = 1.50 ± 0.41 mm, Fig. 1B) which also allowed us to record SC neurons along continuous retinotopy (Fig. 1C, right) in a cell-type specific manner (Fig. 1G). We used the visually evoked activity and continuous retinotopy to demarcate the boundaries of the SC (Fig. 1C). Moreover, for the characterization of excitatory and inhibitory classes of SC neurons, we limited our analysis to single units that were localized within the range of optogenetic stimulation (Fig. 1B/C, see Material and Methods). Cell types were characterized using a custom-written graphic-user-interface (GUI, see Material and Methods) to label units that were activated by the LED pulse as inhibitory neurons (INs, Fig. 1E, middle). These neurons were distinguished based on their reliable, short-latency responses to optogenetic stimulation during spontaneous (black screen) and visually evoked conditions (checkerboard stimulus, see Material and Methods). SC neurons that were not modulated by the LED pulses, but located within the light-activated range, were labeled as excitatory (EXNs, Fig. 1E bottom). As expected, retinal axons were not activated by the LED pulse (Fig. 1E, top), except for a few cases (4/326 RGCs) which were excluded from the connectivity analysis in this study. In total, we recorded 326 RGC axons and 680 SC neurons. Among the recorded SC neurons one third (31.2%) were GABAergic (Fig. 1G, $n = 468$ EXNs, $n = 212$ INs).

Having sorted the SC neurons into the cell classes using optogenetic stimulation (EXNs and optogenetically identified INs) we compared different spike waveform features to test whether SC neurons show differences in their waveforms. Characterizing the spike waveforms is commonly used in the cortex or hippocampus to differentiate between glutamatergic and GABAergic neurons (Lee et al., 2016; Moore and Wehr, 2013; Niell and Stryker, 2008). However, the separation based on differences in waveform features was not an adequate measure to separate SC EXNs from SC INs (Supplementary Fig. 1). This result is in line with findings by Essig et al. who did not observe a cell type specific difference between waveform features in the SC (Essig et al., 2021). Similar findings have been shown in the inferior colliculus (Ono et al., 2017), a neighboring midbrain structure, suggesting that waveform classification analyses used in the cortex are not suitable for distinguishing neurons in the midbrain.

To further characterize the response properties of the different cell classes (RGC, SC INs, SC EXNs), we measured their firing rates (FR) in response to a natural movie stimulus provided in (Froudarakis et al. (2014)) (Froudarakis et al., 2014) (see Material and Methods). We found that RGCs showed the highest FR (mean firing rate RGC = 11.3 ± 10.5 spikes/s, $n = 326$ EXNs; Fig. 1G, bottom). On average, the mean FR in SC INs (mean firing rate IN = 5.9 ± 6.0 spikes/s, $n = 212$ INs) was 1.8 times higher compared to that of SC EXNs (mean firing rate EXN = 3.2 ± 4.9 spikes/s, $n = 468$ INs, SC EXN vs SC IN $p = 3.1567 \times 10^{-15}$, SC EXN vs RGC $p = 4.8362 \times 10^{-40}$, SC IN vs RGC $p = 2.7388 \times 10^{-8}$, two-sided Wilcoxon rank-sum test). Our results are in line with previous studies by Reid (1998, 2003) showing that RGCs show high mean firing rates (Kara and Reid, 2003; Usrey et al., 1998). Moreover, from other midbrain areas it is also known that GABAergic neurons show higher FRs compared to glutamatergic cells (Ono et al., 2017).

In summary, by combining Neuropixels recordings to measure the activity of RGC axons and SC neurons with optogenetic identification of GABAergic SC neurons, it becomes possible to measure monosynaptic retinocollicular connectivity *in vivo* and retinal innervation strength in a cell-type specific manner.

Retinal input integration in excitatory and inhibitory SC neurons

To study the monosynaptic retinocollicular connectivity *in vivo* and retinal innervation strength, we identified putative monosynaptically connected RGC-SC pairs employing established cross-correlation analysis (CCG) methods (Alonso et al., 2001; Bereshpolova et al., 2020; Sibille et al., 2022a). A significant transient and a short-latency peak in the spike train CCG is a hallmark of monosynaptic connectivity (Fig. 2A) and identifies connected RGC-SC pairs. We measured up to 98 monosynaptic connections in individual recordings, depending on the number of measured RGC axons (Fig. 2C). In total, we identified 305 connected RGC-SC pairs, of which 29.8% were retinal axons that connected onto GABAergic SC neurons ($n = 214$ connected RGC-SC EXN pairs, $n = 91$ connected RGC-SC IN pairs, Fig. 2B/D). This high yield in monosynaptically connected pairs was due to the close proximity of the simultaneously recorded RGC axons and SC neurons on the high-density electrode (Fig. 2D, bottom) (Sibille et al., 2022a). For most of the connected RGC-SC pairs the peak channels on the electrode were on average 50-60 μm apart on the probe and hence within SC (distance RGC-EXN: median = 40.0 μm , first quartile = 25.61 μm , third quartile = 62.1 μm ; distance RGC-IN: median = 51.22 μm , first quartile = 32.0 μm , third quartile = 83.08 μm). We found a significant difference in peak channel distance between RGC-SC EXN and RGC-SC IN pairs with longer distances between RGC axons and inhibitory SC neurons ($p = 0.03287$, two-sided Wilcoxon rank-sum test), however, the effect size was small (Cohen's $d = 0.0915$). The maximum distance for retinal axons and inhibitory SC neurons was 240 μm , while the peak channels of one RGC-SC EXN pair were 960 μm apart. The latter pair might include a SC wide field neuron as these cells are excitatory and their dendritic arbor widths have been reported to spread up to 900 μm (Gale and Murphy, 2014).




Fig. 2

Retinal innervation is similarly strong to excitatory and inhibitory SC neurons.

A, Monosynaptically connected RGC-SC EXN (gray) and RGC-SC IN (blue) pairs are identified via cross-correlation analysis (CCG). **B**, CCGs of connected RGC-SC EXN and RGC-SC IN pairs sorted by their peak latency ($n = 214$ RGC-SC EXN, $n = 91$ RGC-SC IN, $n = 11$ recordings). **C**, Connectivity matrix from a single recording. Gray marks indicate connections onto excitatory SC neurons, blue marks indicate connections onto inhibitory SC neurons. RGC axons and SC neurons are sorted by their peak channel along the electrode. **D**, Distribution of peak channel distances between RGC axons and connected SC neurons ($p = 0.0328$, two-sided Wilcoxon rank-sum test).

Inset shows pie chart of identified RGC-SC IN and RGC-SC EXN pairs. **E**, Elicited SC spiking in response to firing of a presynaptically connected retinal ganglion cell (RGC). Raster plot shows SC firing to 1000 randomly selected RGC spikes. Both SC cell types show robust activation upon RGC spiking (top) but also weaker connections can be found (bottom). **F**, Synaptic efficacy as a measure for connection strength for RGC-SC EXN and RGC-SC IN connected pairs ($p = 0.053$, $n = 214$ RGC-SC EXN, $n = 91$ RGC-SC IN). **G**, Peak latency responses for RGC-SC EXN and RGC-SC IN pairs triggered on retinal spiking ($p = 0.321$). Two-sided Wilcoxon rank-sum test.

To assess the connection strength between RGCs and excitatory and inhibitory SC neurons, we calculated the connection efficacy (Usrey et al., 1999). The efficacy measures the connection strength as the probability of an RGC spike triggering an action potential in the postsynaptic SC neuron (see Material and Methods). We found highly diverse connectivity patterns for RGC-SC EXC and RGC-SC IN pairs. RGC activity evoked robust firing in both SC cell types (Fig. 2E, top), however, pairs with weak connection strength were also observed as recently described in (Sibille et al. (2022a)) (Fig. 2E, bottom). Across the population, both SC cell types received similarly strong input from the retina (efficacy RGC-SC IN: median = 4.54 %, Q1 = 2.2 %, Q3 = 7.96 %, maximum = 62.05 %, $n = 91$ connected pairs, Fig. 2F) compared to RGC-SC EXNs (efficacy RGC-SC EXN: median = 3.5 %, Q1 = 1.96 %, Q3 = 5.89 %, maximum = 41.12, $n = 214$ connected pairs). While we observed a tendency for slightly stronger RGC-SC IN connections, this difference was not statistically significant ($p = 0.0534$, two-sided Wilcoxon rank-sum test). In addition, we examined whether excitatory and inhibitory cells show differences in their peak latencies in response to presynaptic retinal spiking. However, we did not find a significant difference in peak latencies between RGC-SC EXN and RGC-SC IN pairs (Fig. 2G, peak latency for RGC-SC EXN = 1.4 ± 0.3 ms, $n = 214$ connected pairs; peak latency for RGC-SC IN = 1.4 ± 0.4 ms, $n = 91$ connected pairs, $p = 0.321$, two-sided Wilcoxon rank-sum test).

Here, we set out to explore how RGC spikes drive the activity in postsynaptic excitatory and inhibitory SC neurons. We did not find a cell-type specific difference in connection strength and therefore conclude that inhibitory and excitatory SC neurons are innervated comparably strong and are both reliably driven by their retinal inputs.

Functional similarity of retinocollicular connections

In sensory cortices, inhibitory neurons are less selective and receive a diverse and nonspecific set of thalamic inputs compared to excitatory neurons that receive more specific thalamic inputs (Alonso and Swadlow, 2005; Bruno and Simons, 2002). However, the selectivity and functional similarity between RGCs and their excitatory and inhibitory target populations in SC remains largely unknown (Shi et al., 2017). Despite both SC populations showing a similar innervation strength from retinal axons, inhibitory neurons could sample from a more functionally diverse population of RGCs. To test this, we characterized the functional similarity between the RGCs and their connected postsynaptic SC neurons. To this end, we calculated the correlation coefficients between the trial-averaged visually evoked activity of connected RGC-SC pairs in response to light and dark sparse noise stimuli (r_{SL} & r_{SD}) as well as the natural movie stimulus (r_{NM}) (Fig. 3A/B, RGC-SC with high signal-to-noise in their receptive, see Material and Methods). To quantify the overall functional similarity by a single similarity value, we averaged the three correlation measurements (similarity = $(r_{SL} + r_{SD} + r_{NM})/3$). A similarity value of 1 corresponds to perfectly correlated visually driven responses, while a value of 0 signifies uncorrelated responses. In our data, the RGC-SC pairs span a wide range of functional similarity, however, we found no significant difference between RGC-SC EXN and RGC-SC IN pairs (similarity for RGC-SC EXN = 0.45 ± 0.15 and RGC-

SC IN = 0.40 ± 0.17 , $n = 85$ RGC-SC EXN pairs, $n = 29$ RGC-SC IN pairs, $p = 0.245$, two-sided Wilcoxon rank-sum test, Fig. 3C). In a next step, we correlated the connection efficacy measure to the similarity index. The synaptic efficacy was positively correlated with the functional similarity of the connected pairs, with a stronger correlation for RGC-SC IN pairs (RGC-SC EXN $r = 0.29$, $p = 0.007$; RGC-SC IN $r = 0.471$, $p = 0.01$, Fig. 3D).




Fig. 3

Characterization of functional similarity between retinocollicular connected pairs.

A, Spatiotemporal receptive fields (STRF) evoked by a dark sparse noise stimulus for RGC-SC EXN and RGC-SC IN connected pairs. The functional similarity of the RGC axon and the postsynaptic SC neuron is characterized by the correlation coefficient r_{SD} . **B**, Visually evoked activity in response to a natural movie stimulus for the connected pairs shown in **A** together with their correlation coefficient values r_{NM} . **C**, The overall functional similarity between the presynaptic RGC and the postsynaptic SC neurons is reflected in the similarity index calculated from the averaged correlation coefficients $(r_{SD} + r_{SL} + r_{NM})/3$ ($p = 0.245$, two-sided Wilcoxon rank-sum test, $n = 85$ RGC-SC EXN pairs, $n = 29$ RGC-SC IN pairs). **D**, Relationship between similarity index and connection efficacy (RGC-SC EXN $r = 0.29$, $p = 0.007$; RGC-SC IN $r = 0.471$, $p = 0.01$; Pearson correlation coefficient test; $n = 85$ RGC-SC EXN pairs, $n = 29$ RGC-SC IN pairs).

Overall, we observed that functionally similar pairs are more strongly connected. However, we also observed cases of relatively strong connected RGC-SC pairs (~10%) with low similarity (Fig. 3D), suggesting that some SC neurons receive convergent input from a functionally more diverse pool of RGC afferents (Sibille et al., 2022a). Our findings delineate that both RGC-SC EXN and RGC-SC IN connected pairs are organized, while functionally similar properties between the pre- and postsynaptic neuron are reflected in a stronger connection.

Paired-spike interactions of the retinocollicular connections

We showed that retinal spikes are efficient in driving SC activity for both excitatory and inhibitory neurons. As the timing of presynaptic activity has been assessed as a parameter

that modulates postsynaptic firing (Zucker, 1989) and connection strength (Kara and Reid, 2003; Usrey et al., 1998), we wanted to know whether the temporal pattern of retinal activity has different effects on the synaptic efficacy of excitatory and inhibitory SC neurons. To that end, we studied the short-term dynamics (facilitation or depression) of the measured connections and performed a paired-spike analysis approach as described by Usrey and colleagues in 1998 (Usrey et al., 1998). A pair of retinal spikes (1st, 2nd RGC) was defined by its temporal firing patterns: inter-spike interval (ISI) and dead time. We selected RGC spikes for which a 1st RGC followed the 2nd RGC by a specific ISI (minimum ISI = 5ms, maximum ISI = 30ms). The dead time ensured that the 1st RGC was preceded by a minimum period without activity (30ms) (Fig. 4A).




Fig. 4

Paired-spike dynamics: Second retinal spikes are more efficient in driving SC response.

A, Schematic illustrating the temporal dynamics between two successive RGC spikes. Pairs of RGC spikes with a minimum inter-spike interval (ISI) of 5 ms and maximum ISI of 30 ms were included if there were no preceding spikes before the 1st RGC for a dead time of at least 30ms. **B**, Cross-correlograms (CCG) of example RGC-SC EXN and RGC-SC IN pairs calculated from spike trains selected for 1st and 2nd RGCs and the corresponding raster plots to 1000 trials triggered on RGC spikes. **C**, Scatter plot of efficacies for 1st versus 2nd retinal spikes. The majority of connected pairs (185/214 RGC-SC EXN pairs; 73/91 RGC-SC IN pairs) showed paired-spike enhancement in response to 2nd RGC spikes. **D**, Paired-spike ratio (PSR) for RGC-SC EXN and RGC-SC IN connected pairs. The paired pulse enhancement is stronger in SC-EXNs ($n = 214$ RGC-SC EXN; $n = 91$ RGC-SC IN, $p = 2.581 \times 10^{-5}$, two-sided Wilcoxon rank-sum test). **E**, Correlation of 1st RGC efficacy with the PSR for connected pairs ($r = 0.3314$ for $n = 214$ RGC-SC EXN pairs and $r = 0.4767$ for $n = 91$ RGC-SC IN pairs; Pearson correlation coefficient test).

We found that overall 2nd retinal spikes were more efficient in driving SC activity, and thus facilitating. This was true for both RGC-SC EXN (efficacy 1st RGC: median = 1.8 %, Q1 = 0.77 %, Q3 = 4.47 %; efficacy 2nd RGC: median = 2.66 %, Q1 = 1.16 %, Q3 = 6.07 %, $p_{1^{st} \text{ vs } 2^{nd}} = 3.13 \times 10^{-25}$, $n = 214$ pairs) and RGC-SC IN connected pairs (efficacy 1st RGC: median = 3.66 %, Q1 = 1.6, Q3 = 8.61; efficacy 2nd RGC: median = 4.33, Q1 = 2.03, Q3 = 8.25, $p_{1^{st} \text{ vs } 2^{nd}} = 1.68 \times 10^{-6}$, $n = 91$ pairs, Wilcoxon signed-rank test) (Fig. 4B, C). This effect is known as paired-spike enhancement and has been reported earlier in retinogeniculate (Mastrorade, 1987; Usrey et al., 1998) and geniculocortical (Usrey et al., 2000) connections. The majority of retinocollicular connected pairs showed paired-spike enhancement where 2nd retinal spikes were more effective in driving SC response (86.45% (185/214) RGC-SC EXN pairs, 80.22 % (73/91) RGC-SC IN pairs). The effect was more pronounced in RGC-SC EXN pairs where the 2nd retinal spike was 1.7 times as effective in driving responses in the postsynaptic SC neuron while for RGC-SC IN pairs, the 2nd retinal spike was only 1.3 times more efficacious in driving SC spiking. To quantify this observation, we calculated the paired-spike ratio (PSR

= efficacy 2nd/efficacy 1st) which confirmed a higher facilitation effect for RGCs connecting into excitatory neurons (PSR RGC-SC EXN = 1.711 ± 1.14 ; RGC-SC IN: 1.281 ± 0.414 , $p = 0.00002582$, Wilcoxon rank-sum, $n = 305$ pairs, Fig. 4D).

Previous work in the thalamocortical system has shown that weaker synaptic connections can be more strongly modulated compared to strong synaptic connections (Ferrarese et al., 2018). To test the notion that also in the retinocollicular system, connected pairs with lower efficacy in response to 1st RGCs show greater enhancement in response to the 2nd RGCs, we correlated the efficacy of the 1st RGC to the PSR (Fig. 4E). We found a negative correlation of efficacy to 1st RGC and PSR for both RGC-SC EXN ($r = 0.3314$, $p = 7.0148 \times 10^{-7}$, $n = 214$ pairs) and RGC-SC IN pairs ($r = 0.4767$, $p = 1.7714 \times 10^{-6}$, $n = 214$ pairs). However, we found pairs that show low efficacy to 1st RGC and low PSR. Taken together, we observed paired-spike enhancement in RGCs connecting onto both inhibitory and excitatory SC neurons with a stronger facilitation effect for retinal afferents connecting onto excitatory, in comparison to inhibitory, SC neurons.

Inhibitory SC neurons are less coupled to the retina compared to excitatory SC neurons

Our data so far reveals that excitatory and inhibitory SC neurons receive similarly strong drive from their retinal afferents. However, we observed a higher mean firing rate in SC INs compared to SC EXNs, on average (Fig. 1G), which is intriguing given the similarity in the RGC drive on both cell types. A possible explanation could be that the activity of SC INs is less coupled to the RGC inputs as compared to that of SC EXNs, i.e. SC INs spikes are less often driven by their RGC inputs. To examine how strongly the activity of the two populations of SC neurons is coupled to the activity of individual RGC inputs, we estimated the synaptic coupling strength from the connection contribution (Levick et al., 1972; Sibille et al., 2022a; Usrey et al., 1999) of the measured retinocollicular pairs. The connection contribution reflects the fraction of postsynaptic SC spikes that are preceded by a presynaptic retinal spike within a short time window (Fig. 5A). High contribution values indicate that SC neurons are primarily driven by individual RGC afferent inputs, while low contribution values point towards an integration of multiple inputs.




Fig. 5

Connection contribution is higher in connected RGC-SC EXN pairs.

A, Example CCGs of monosynaptically connected RGC-SC EXN (top) and RGC-SC IN (bottom) pairs. Connected pairs were identified by their peaks in the CCGs (left). Raster plots of RGC spiking activity triggered on 1000 SC spikes (right). **B**, Population data on contribution measures for RGC-SC EXN and RGC-SC IN pairs ($n = 114$ RGC-SC EXN pairs, $n = 91$ RGC-SC IN pairs, $p = 0.0003$, two-sided Wilcoxon rank-sum test). **C**, Relationship between functional similarity index and connection contribution (Pearson correlation coefficient test; $n = 85$

RGC-SC EXN pairs, $n = 29$ RGC-SC IN pairs). **D**, Correlation of mean firing rate (to natural movie stimulus) and contribution for RGC-SC EXN and RGC-SC IN pairs (Pearson correlation coefficient, two-sided Wilcoxon rank-sum test, $n = 214$ RGC-EXN and $n = 91$ RGC-IN connected pairs; $n = 11$ penetrations from 9 mice).

Interestingly, we found that the contribution of individual retinal spikes on SC activity was higher for excitatory compared to inhibitory SC neurons (Fig. 5B) (contribution RGC-SC EXN: median = 18.27% Q1 = 10.79%, Q3 = 29.69%, $n = 214$; contribution RGC-SC IN: median = 12.2%, Q1 = 6.61%, Q3 = 23.02%, $n = 91$ connected pairs, $p = 0.00034694$, two-sided Wilcoxon rank-sum test). This opens up the possibility that SC EXNs might be more specifically linked to the retinal input compared to SC INs. To assess the relationship between the coupling strength and the functional similarity, we correlated both measures and found a positive correlation suggesting that stronger coupled pairs show higher similarity (Fig. 5C). This correlation was higher for RGC-SC EXN pairs (RGC-SC EXN $r = 0.502$, $p = 1.0 \times 10^{-6}$, $n = 85$ pairs) compared to RGC-SC IN ($r = 0.481$, $p = 0.0081839$, $n = 29$ pairs, Pearson correlation coefficient test). Moreover, we observed a diversity of firing rates in both SC EXNs and SC INs (Fig. 1G) which suggests that low contribution values are related to the mean firing rates of the SC neurons. And indeed, when we correlated the contribution to the mean firing rate, we found a negative correlation for both RGC-SC EXN and RGC-SC IN pairs (RGC-SC EXN $r = -0.16985$, $p = 0.01284$; RGC-SC IN $r = -0.40215$, $p = 0.00008$, Pearson correlation coefficient, $n = 214$ RGC-EXN; $n = 91$ RGC-IN connected pairs) (Fig. 5D).

Taken together, the cell-type specific difference in retinal coupling strength suggests that retinal input contributes a substantial part to the spiking activity of excitatory SC neurons, while inhibitory SC neurons seem to also integrate inputs from other sources. This finding denotes that excitatory SC neurons are more tightly linked to the retinal input compared to inhibitory SC neurons.

Discussion

In this study, we elucidate how inhibitory and excitatory SC neurons integrate visual information from the retina *in vivo*. By combining the tangential Neuropixels recording approach with optotagging techniques, we were able to measure RGC-SC connected pairs in a cell-type specific manner and study their functional organization. Our findings allow the following conclusions: (1) The retinal drive is similarly strong for RGCs connecting onto GABAergic and non-GABAergic SC neurons. (2) The connection strength between the RGCs and both SC cell classes is positively correlated with their functional similarity, while still allowing for the presence of strong connections with low functional similarity. (3) Paired-spike enhancement in RGC-SC connections was a signature for both excitatory and inhibitory SC neurons. Finally, (4) the contribution of a retinal spike to overall SC firing is higher in excitatory SC neurons compared to inhibitory SC neurons.

We studied retinocollicular connectivity strength and aimed to answer the question whether excitatory and inhibitory SC populations use similar or different coding schemes to integrate visual information compared to the thalamocortical visual system. Connectivity patterns and circuit motifs have been extensively studied in the thalamocortical system, yet our understanding of the encoding mechanisms that govern retinocollicular signaling remains rudimentary albeit recent studies suggested that feedforward inhibition might be at play (Villalobos et al., 2018; Whyland et al., 2020). While cortical inhibitory neurons are more strongly activated by thalamic axons compared to excitatory neurons (Bruno and Simons, 2002; Cruikshank et al., 2007; Ji et al., 2016), our results now show that in the SC, the afferent retinal drive is comparably strong on excitatory and inhibitory neurons. This finding is intriguing and suggests that the two major visual brain structures in the vertebrate brain, the visual cortex and the SC, integrate afferent sensory information differentially.

Several parameters for example the degree of convergence (Kara and Reid, 2003), response properties (Alonso et al., 2001), synaptic mechanisms (Cruikshank et al., 2007) or spiking

patterns (Usrey et al., 1998) can have meaningful effects on modulating the input strength on the target cell. SC neurons receive input from a small number (~6) of RGCs (Chandrasekaran et al., 2007). In contrast, cortical neurons receive numerous (approximately one order of magnitude more) thalamic inputs that are usually weak (Bruno and Sakmann, 2006; Jin et al., 2011; Lien and Scanziani, 2018). Overall, we observed strong retinal drive onto both SC populations indicated by high efficacy values (Fig. 2). One hypothesis is that convergence might be low in retinocollicular pairs as recently suggested by Sibille et al. (2022); however, despite the limited input from RGCs, these inputs may be sufficiently strong to drive the SC. This is in contrast to thalamocortical inputs, where individually efficacy values are low, but the thalamic drive is boosted by synchronously active thalamic inputs (Alonso et al., 1996; Bruno and Sakmann, 2006). Our recording method allows for the simultaneous recording of multiple RGC afferents from SC neurons making it possible to evaluate how the synchrony among RGC afferents modulates the firing of postsynaptic SC neurons in future studies.

The difference in connection strength between cortical excitatory and inhibitory neurons could be partly attributed to their differences in response properties. Cortical inhibitory neurons display higher sensitivity, receive more convergent thalamic input, have higher firing rates and sample more randomly while excitatory neurons receive highly functional specific thalamic input (Kyriazi et al., 1994; Reid and Alonso, 1995; Simons and Carvell, 1989). While we observed higher firing rates in inhibitory SC neurons (Fig. 1G), the efficacy of the RGC-SC connections was comparable between both types of neurons (Fig. 2F), suggesting that the sensitivity of both neuron types is not markedly disparate. In general, visual response properties between excitatory and inhibitory neurons in the SC may not differ to such a great extent (Inayat et al., 2015; Kasai and Isa, 2016; Shi et al., 2017), (but see (Barchini et al., 2018); (Li and Meister, 2022)), and experiments in the inferior colliculus show that GABAergic and glutamatergic cells share similar response properties (Ono et al., 2017). These observations imply that in the SC and other parts of the midbrain, the cell-type specific difference in terms of selectivity might not be as strongly separated as in the cortex. Future studies are required to deeper investigate the differences in response properties of the subclasses of SC neurons and whether these properties are correlated to the RGC-SC connection strength.

In our results, we observed an overall trend that functionally similar RGC-SC pairs show stronger connections which was true for both excitatory and inhibitory pairs (Fig. 3). However, we also find pairs with considerably strong retinocollicular connectivity (efficacy values >10%) albeit the functional similarity between these connected cells was low (Fig. 3D). This observation suggests that SC neurons may either show high projection-specificity by integrating inputs from a functional homogeneous population of RGCs (Li and Meister, 2022; Shi et al., 2017; Sibille et al., 2022a) or receive diverse inputs from multiple functional RGC types (Li and Meister, 2022; Sibille et al., 2022a). Moreover, a high degree of projection-specificity was recently shown to be maintained even disynaptically towards brain regions downstream of the SC (Reinhard et al., 2019), suggesting that at least one important role of the RGC-SC projection is to preserve the representation of the functionally diverse retinal channels (Baden et al., 2016). In our study, we investigated the functional similarity between connected RGC-SC pairs in response to sparse noise and natural movie stimuli. And while we observed a diversity among the similarities between RGCs and connected SC neurons, supporting the idea that SC neurons may follow diverse wiring motifs (Sibille et al., 2022a), we observed no clear difference between inhibitory and excitatory SC neurons (Fig. 3C). To further address the influence of functional properties on wiring mechanisms, it would be interesting to target the pool of different RGC types and to study how strongly they recruit SC subpopulations and modulate postsynaptic responses.

Temporal dynamics of presynaptic activity patterns are known to play a role in modulating synaptic strength (Usrey et al., 1998; Zucker, 1989). To study how the temporal spiking

pattern of the RGC activity shapes RGC-SC connectivity, we analyzed the paired-spike dynamics. Across the population, we found a facilitating effect where responses to 2nd retinal spikes were more efficient in driving SC activity for both SC neuron populations. However, the paired-spike enhancement was more pronounced for excitatory than inhibitory neurons (Fig. 4). Moreover, our observation highlights differences in the encoding strategies used by different regions of visual processing. While retinogeniculate connections show paired-spike enhancement (Usrey et al., 1998), thalamocortical and corticotectal synapses express mostly depressing dynamics (Bereshpolova et al., 2006; Swadlow et al., 2002; Swadlow and Gusev, 2001), but see (Usrey et al., 2000). Our data now reveals that retinocollicular connections facilitate, suggesting that the temporal integration of RGC inputs by SC neurons may be similar to the visual thalamus.

In our study, we used cross-correlograms that were calculated from spike trains throughout the full recording period, to be applied on different connectivity strength measures. However, paired-spike dynamics are known to be modulated by the local temporal structure of inputs, for example the visual stimulus presented (Usrey et al., 1998). In addition, facilitating and depressing effects can change with the target population (Bereshpolova et al., 2006). Further studies are required to investigate whether the facilitating effect observed in our study changes upon targeting defined collicular subpopulations that may have differences in firing rates or during different stimulus conditions (e.g. spontaneous and stimulus evoked). In this study, retinocollicular connectivity was measured in anesthetized conditions. For future studies it would be interesting to examine connectivity dynamics and innervation strength in awake conditions as it has been shown that retinocollicular synapses are modulated by the animals' level of arousal (Schröder et al., 2020).

Despite our observation that retinal drive is similarly strong onto both SC cell types, the contribution of retinal activity to individual SC neurons firing was higher in excitatory SC neurons. This opens up the possibility that excitatory SC neurons may be more specifically linked to retinal inputs, while GABAergic SC cells might receive excitation from other sources aside from the retina. This finding raises the question on what other sources could drive spiking in inhibitory SC neurons?

1) RGCs provide the main input to the superficial SC (Ellis et al., 2016) with limited convergence (Sibille et al., 2022a). Nonetheless, the convergence may be higher for RGCs connecting onto inhibitory SC neurons which could explain the lower contribution and higher mean firing rates we found in inhibitory neurons. However, it has to be considered that our *in vivo* recording technique is limited in capturing the total pool of converging RGC inputs on a neuron (Sibille et al., 2022a). Viral retrograde transsynaptic tracing methods (Muellner and Roska, 2023; Rompani et al., 2017), could be employed to study the full presynaptic RGC pool of single SC neurons.

2) In addition to retinal inputs, the SC receives extensive projections from the visual cortex (V1) that could act as a potential candidate to excite inhibitory neurons. And while indeed GABAergic horizontal cells have been shown to receive V1 input (Whyland et al., 2020), the majority (93%) of corticotectal terminals target the dendrites of non-GABAergic neurons in the SC (Masterson et al., 2019). Therefore, more work is needed and studying V1 excitation to defined inhibitory subpopulations will be an important topic to be investigated by the use of tracing studies and will shed light into this complexity.

3) Intracollicular connections can provide a source of recurrent excitation on SC neurons, amplifying the retinal drive (Shi et al., 2017). However, the cell-type specificity of the intracollicular amplification is still debated. For example, while GAD2-positive horizontal cells get excitatory inputs from stellate cells (Gale and Murphy, 2018), intracollicular connectivity originating from neighboring excitatory cell types such as narrow-field (NF)

and wide-field (WF) cells has shown to be rare (Gale and Murphy, 2018). Next-generation *in vitro* multi-neuron patch-clamp recordings (Campagnola et al., 2022) could reveal important new insights into the cell-type specific (Ayupe et al., 2023) intracollicular wiring.


4) In addition, collicular neurons in the visual layers also get inputs from deeper SC layers. However, excitatory connections from deep SC to the upper layers are rather uncommon (Gale and Murphy, 2018; Ghitani et al., 2014). Despite these studies have highlighted different sources of excitation onto inhibitory SC neurons, further studies are required that elucidate the role and strength of input that deep SC layers or external sources provide to drive inhibitory SC populations.

In this study, we recorded populations of well-isolated single units including RGCs as well as GABAergic and non-GABAergic SC neurons in VGAT-ChR2 mice *in vivo*. Our approach using VGAT-ChR2 mice to target all types of inhibitory SC neurons that express the vesicular GABA transporter VGAT enables a first impression on how inhibitory neurons sculpt retinocollicular signaling. Our inhibitory population is made up of 30% of the recorded SC neurons and connected pairs. These observations are in line with previous findings showing that in the upper SC layers, a substantial part of neurons (30%) are GABAergic (Ranney Mize, 1992; Whyland et al., 2020) and that around one third of the postsynaptic targets of retinocollicular terminals are GABAergic (Boka et al., 2006; Whyland et al., 2020). Yet, in spite of this high density of GABAergic neurons in the SC, they comprise of a high diversity of subpopulations that have been shown to be involved in distinct visual processing (Barchini et al., 2018; Kasai and Isa, 2016) and visually guided behaviors for example prey capture (Hoy et al., 2019) or saccadic suppression (Phongphanphane et al., 2011). An understanding of the connectivity and encoding principles of these diverse subpopulations provides the foundation for a deeper understanding of how inhibition shapes functional circuits and their underlying behavior.

In the last decade, the diversity of SC cell types has been mainly characterized based on morphology and response properties (Franceschi and Solomon, 2018; Ito et al., 2017). Advanced techniques have started to shed light on this diversity by providing cell-type characterizations on the molecular level (Barchini et al., 2018; Gale and Murphy, 2018; 2014; Li and Meister, 2022; Villalobos et al., 2018). Three categories of potential GABAergic interneurons have been classified in the SC by the use of Cre-lines (Whyland et al., 2020) that can be further subdivided into intrinsic interneurons and potential projection neurons. RNA-sequencing has become an approach to characterize defined subtypes of neuronal populations. A recent study by (Tsai et al., 2022) has mapped retinocollicular connectivity by targeting specific RGC-SC circuits using a novel approach (Trans-Seq) to classify subsets of SC cell-types that are innervated by genetically defined RGC types (Tsai et al., 2022). They identified five inhibitory clusters of retinorecipient SC cells that allow to specifically map retinocollicular connectivity. In addition, 12 distinct inhibitory subtypes have been recently identified using single-nucleus RNA sequencing where Pax7 was shown to be exclusively expressed in inhibitory neurons (Ayupe et al., 2023). Thus, we are at an enthralling stage where information on SC cell-types finally becomes available based on gene expression patterns, alike to the cortical circuit (Tremblay et al., 2016). However, while these novel techniques shed light on the diversity of inhibitory subpopulations and allow to map cell-type specific connectivity, these methods cannot provide insights into the strength of the RGC-SC connections. Hence, a key step forward would be to leverage the tracing and transcriptomics studies and combine these methods with our high-density recording approach to enable a finer characterization of the collicular interneuron diversity and the complexity of their connectivity patterns *in vivo*.

Taken together, our study targeted a heterogeneous population of GABAergic neurons in VGAT-ChR2 mice and our results reveal that excitatory and inhibitory neurons in the two

main brain areas for visual processing, the visual cortex and the superior colliculus, integrate sensory afferent inputs in different ways.



Supplementary Fig. 1

Spike waveform features analysis for GABAergic and non-GABAergic neuron populations in the SC.

A, Top: Illustration of features extracted from single-channel waveforms (magenta circles indicate the trough and peak). Middle and bottom: Single-channel mean waveforms for non-GABAergic (gray; $n = 468$) and GABAergic (blue, $n = 212$, $n = 9$ mice) SC populations (mean $\pm$ SEM). **B-D**, Distribution of waveform features extracted from single-channel waveforms for inhibitory and excitatory SC populations. Peak-to-trough duration represents the time between trough and peak ($p = 4.05 \times 10^{-10}$). Peak-to-trough ratio represents the ratio between amplitudes of peak and trough ($p = 4.45 \times 10^{-9}$). Amplitude is the absolute difference between trough and peak ($p = 1.38 \times 10^{-9}$). **E**, Distribution of waveform spread along the probe extracted from multichannel waveforms for GABAergic and non-GABAergic SC populations ($p = 1.58 \times 10^{-4}$, $n = 468$ EXNs, $n = 212$ INs, Two-sided Wilcoxon rank-sum test).

Acknowledgements

We thank J. Poulet for providing us the VGAT-ChR2 mice; KL. Teh, T. Lupashina, J. Kosubek-Langer and P. Norton for comments on the manuscript. We thank the Neuropixels community for their support. This work was supported by the DFG Emmy-Noether grants KR 4062/4-1 and KR 4062/4-2 (JK).

Author contributions

C.G. and J.K. conceived and designed the study; C.G. collected the data; C.G., J.S., and J.K. analyzed the data and C.G. and J.K. wrote the manuscript with inputs from all authors.

Declaration of interests

The authors declare no competing interests.

Materials and methods

Experimental Design Animals

All experiments were carried out in accordance with the guidelines given by Landesamt für Gesundheit und Soziales (LAGeSo - G 0142/18) Berlin and were approved by the authority. Adult VGAT-ChR2 (Slc32a1-COP4*H134R/EYFP, The Jackson Laboratory, stock no. 014548, n=9) transgenic mice of either sex were used to activate inhibitory neurons (Zhao et al., 2011). Mice were aged 7-9 months old on the day of recordings.

Surgical procedures

All experiments were conducted in VGAT-ChR2 mice of both sexes. Mice were induced with 2.0-2.5% isoflurane (CP-Pharma) in an induction chamber and transferred to the surgery. Once anesthetized, the surgery was performed in a stereotactic frame (Narishige) with a closed-loop temperature controller (FHC DC) for monitoring the animal's body temperature and maintaining it at 37°C. The isoflurane level was gradually lowered during surgery and maintained at 0.8-1.5% while ensuring a complete absence of vibrissa twitching or responses to tactile stimulation. During surgery, the eyes were protected with eye ointment (Vidisic, Bausch+Lomb). Mice were head-fixed in the frame, and the skull was exposed. The head was aligned along the anterior-posterior axis and marks were made for the craniotomy at 600-1500 µm ML and 0-1500 µm AP from Lambda (Paxinos and Franklin, Nixdorf 2007 stereotactic atlas) using a micromanipulator (Luigs and Neumann). For optotagging experiments, an additional mark for the optic fiber placement was made at 3500 µm AP, 500-1200 µm ML to Bregma. A headpost was placed on the skull and implanted with dental cement (Paladur). Dental cement was also used to build a recording chamber to provide a bath for grounding. During this step, a silver wire (AG-10W, Science Products) was attached to the dental cement chamber for grounding. It is important to keep the dental cement chamber low enough on the anterior and posterior sides to allow for the shallow angle probe and fiber implantations, see Sibille et al (2022) for a detailed description of the method (Sibille et al., 2022b). Craniotomies were made at the marked positions using a dental drill (Johnson-Promident). To allow for the probe insertion at a horizontal angle between 15° and 20°, a small part of the skull at the posterior part of the craniotomy was slightly thinned using the drill to ensure a smooth insertion (Sibille et al., 2022b).

Anesthetized extracellular recordings and optogenetic stimulation

Electrophysiological recordings in SC were performed in 9 anesthetized mice of both sexes (4 female, 5 male) using Neuropixels 1.0 probes (Jun et al., 2017). Data were acquired at a sampling rate of 30 kHz using the Open Ephys acquisition software and GUI (Siegle et al., 2017) (www.open-ephys.org(<http://www.open-ephys.org/>)) and the PXIe hardware acquisition system (National Instrument NI-PXIe-1071). The signal was amplified and stored in both the local field potential band (LFP, high pass-filtered 0-300 Hz) and the action potential band (AP, 300 Hz to 3 kHz). The tangential recordings along the anterior-posterior

axis in SC were performed as described in Sibille et al. (2022) (Sibille et al., 2022a). The fiber for optogenetic stimulation was inserted in the SC prior to the Neuropixels probe implantation to decrease mechanical pressure and reduce drift of the electrode probe. The angled optogenetic fiber was zeroed at the brain surface and inserted from the anterior in a 60-70° angle towards the AP axis (see Fig. 1A) and slowly lowered using a manual micromanipulator until reaching an insertion of 150-200 µm. Following the fiber implantation, the Neuropixels probe was inserted tangentially in a 15 to 20° angle in the superior colliculus (SC) using a micromanipulator (NewScale). The probe was zeroed at Lambda and inserted 500-1200 µm ML and 2800-3800 µm DV. After the Neuropixels probe was implanted, the optic fiber was lowered further slowly (to a maximum insertion depth of ~250 µm) and LED pulses were applied to test for optogenetic activation of cells. After correct placement, the probe and fiber were allowed to settle for 15-20 minutes before data acquisition was started. In addition, confirmation of probe placement within the SC was done via receptive field (RF) mapping to check the number of channels with RFs. RFs were estimated from the multi-unit activity (MUA) using spike-triggered average (STA) (Sibille et al., 2022b)) (Fig. 1C).

Optotagging to identify GABAergic neurons in VGAT-ChR2 mice

We identified (“optotagged”) populations of single GABAergic SC neurons via Channelrhodopsin-2 (ChR2) activation in VGAT-ChR2 mice. The optic fiber was calibrated prior to each experiment to ensure a stable light output (2.5mW) measured at the fiber tip. Before the electrode probe insertion (see above), the optic fiber (NA .66, 200 µm core diameter, with a bare fiber tip, PlexBright, Plexon) coupled to a blue (470 nm) LED module (PlexBright, Plexon) was inserted using a 60-70° angle tilted towards the AP axis (Fig. 1A). Blue light pulses were used to evoke spiking activity in VGAT-expressing neurons and to verify the proper alignment of the optic fiber and the Neuropixels probe. If no optogenetically evoked neuronal activity was observed, either the fiber was lowered to a maximum of 250 µm or the probe position was re-adjusted to ensure both the activation and recording of GABAergic SC neurons. Once the Neuropixels electrode probe and optic fiber were properly aligned, optogenetic stimulation using either a square wave or Gaussian function stimulus pulse was applied to identify light-activated channels (Fig. 1A, B). As the stimulation using a square wave pulse led to the induction of light artefacts in the spiking response (Jun et al., 2017), we manually removed these artefacts in a post-processing step by interpolating the raw values in each channel during the onset and offset of the square pulse in 2 out of 11 experiments. In order to decrease light artefacts induced by the sharp light onset in the square wave pulse, we changed the pulse to using a Gaussian function stimulus pulse to gradually increase light intensity in the remaining 8 experiments. The light pulses were applied during spontaneous conditions (presentation of a black background, 100 ms pulse duration, 2 Hz, 200 trials, 2.7-3.5 mW measured at the fiber tip) and conditions with visually evoked activity (checkerboard stimulus, 1000 ms pulse duration, 30 trials ON, 30 trials OFF, 2.7-3.5 mW measured at the fiber tip, n = 10/11 experiments). Both stimulation protocols were shown twice within the recording to ensure stable responses throughout the recording session. GABAergic neurons were identified by their increased short-latency responses to blue light stimulation pulses. When we compared the visually evoked responses to those induced by ChR2 activation we found that they varied in their temporal activation profile (Fig. 1B, C). The response to the visual sparse noise stimulus starts ~80ms after the stimulus onset (Fig. 1C) while responses induced by ChR2 activation had a shorter latency following light onset (Fig. 1C).

Visual stimulation

Visual stimuli were generated in Python using the PsychoPy2 toolbox (Peirce et al., 2019). The visual stimuli were displayed on a calibrated screen (Dell, refresh rate = 120 Hz, mean luminance = 120 cd/m²). For receptive field mapping, we presented the sparse noise visual stimulus on a grid of 24x 14 squares. The sparse noise targets were either dark (on light background) or light (on dark background) and 10 deg in size with [n_targets per frame, n_trials per position] = [2,20] and presented for 100 ms. A set of standardized different visual stimuli was presented including a natural movie (30 trials, randomized with other stimuli, 10s) taken from (Froudarakis et al. (2014)) (Froudarakis et al., 2014) and a full-field chirp stimulus (Baden et al., 2016). The frametimes of the visual stimuli were marked by stimulus-locked synchronizing events (TTLs).

Histology

For histological reconstruction of the electrode track, the probe was removed, coated with fluorescent dye (DiI, Abcam-ab145311) and re-inserted in the same location. In a few experiments we performed multiple recordings and only stained the last insertion. Subsequently, the animal was terminated with an excess of isoflurane (> 4%). Cardiac perfusion was performed with phosphate buffer saline (PBS) followed by 4% paraformaldehyde (PFA) in PBS. Brains were post-fixed overnight in 4% PFA, and stored in PBS until slicing (100 µm) on a vibratome (Leica VT1000S). The brain slices were mounted using a mounting medium containing DAPI (DAPI-Fluoromount-G, Biozol Cat. 0100-20). Slices were imaged on a fluorescence microscope using a 2.5 x objective for post hoc visualization of the electrode probe track.

Data analysis and statistical analyses

Data analysis was performed in Python 3.0 (www.anaconda.com(<http://www.anaconda.com/>)) using custom-written scripts. MATLAB 2019b (www.mathworks.com(<http://www.mathworks.com/>)) was used for spike sorting, see below. Statistical tests were performed using the two-sided Wilcoxon rank-sum test for unpaired samples and the Wilcoxon signed-rank test (two-sided) for paired samples using the scipy.stats module in Python, unless stated otherwise. For correlation analyses we used Pearson correlation implemented via the linregress function. Population results are indicated as mean ± standard deviation if not stated otherwise. The number of neurons and animals used in each analysis is reported in the Results section or figure legends.

Multi-unit activity extraction

For MUA analysis, common average referencing was applied to the bandpass filtered AP data (Butterworth filter order 2, 0.3 to 3 kHz), where each event is extracted using custom-written Python scripts. Spike detection was performed for each channel independently at a threshold of 4 standard deviations in the AP band (double side detection). The scripts for performing the MUA analysis are available on the GitHub repository (https://github.com/KremkowLab/tangential_recording)

Offline Spike sorting and criteria for unit classification

KiloSort2 and 3 (Pachitariu et al., 2016) (<https://github.com/MouseLand/Kilosort>) were used for automatic detection and clustering of spikes followed by manual curation using phy2

(Rossant et al., 2016) (<https://github.com/cortex-lab/phy>). Double-counted spikes were removed for each cluster (within ± 0.16 ms) (Siegle et al., 2021). Furthermore, clusters with between-unit overlapping spikes that show above chance zero-lag peaks in the cross-correlograms (CCG, peak windows ± 0.5 ms) were re-evaluated individually in phy to be either refined or removed. Inter-spike-interval (ISI) violations were calculated as the ratio of the spikes within the refractory period (± 1.5 ms) to the total number of spikes. Units with ISI $> 0.05\%$ were removed. We also checked for stable firing of clusters throughout the recording. Furthermore, we excluded units from further analysis that were lying outside the optogenetic stimulation range (see below, Identification of Inhibitory neurons, see Fig. 1B) and outside the SC borders (from STA RFs on MUA, see Fig. 1C).

Waveform classification to identify RGC axons and SC neurons

A waveform classification approach was applied to distinguish action potentials (APs) from retinal ganglion cell (RGC) axons and somatic APs from SC neurons (Sibille et al., 2022a). In brief, we calculated the multi-channel waveform (MCW) which reflects the spatiotemporal profile of the AP signals. Afferent axons and somatic signals could be classified based on their distinct waveform which allowed us to classify clusters into afferent axon AP vs somatic AP. In essence, due to the SC electrode probe implantation along the anterior-posterior (AP) axis, the electrode captures signals from retinal axons that innervate the SC along the electrode channels. This propagation of RGC axonal action potentials along the axonal path is visible in the spatiotemporal profile of their multi-channel waveforms (Fig. 1D, left).

Waveform classification approach to separate GABAergic and non-GABAergic SC neurons

We performed standard waveform analyses to attempt to separate GABAergic SC neurons based on waveform features (Jia et al., 2019; Quirk et al., 2009) (Supplementary Fig. 1). Briefly, we detected the negative (trough) and positive peak (peak) from single-channel waveforms and calculated their peak-to-trough duration. The amplitude was defined as the absolute difference between trough and peak. The peak-to-trough ratio was defined as the ratio between amplitudes of peak and trough.

Post hoc separation of GABAergic and non-GABAergic SC neurons

To study retinal input integration in a cell-type specific manner, we used a method to separate the neural responses of GABAergic neurons from that of excitatory neurons in the SC. First, we identified the minimum and maximum recording sites on the probe that were photo-activated. These borders were defined via the minimum and maximum peak channel of light-induced units that are activated on the probe. SC neurons that did not fall inside this optogenetic stimulation range (identified by their peak channel) were excluded from further analysis. SC neurons were separated based on their short-latency increase in responses to blue light pulses during two different protocols: (1) baseline activity (black background, 100 ms pulse duration, 2 Hz pulse frequency) and (2) during a checkerboard stimulus (1000 ms pulse duration, 30 trials LED ON, 30 trials LED OFF). This classification was done using a custom-written graphical-user-interface (GUI) approach where units were labeled as either “light-responsive”, “non light-responsive”. The labeling step was conducted by three independent observers. A unit was included if 2 out of 3 observers chose the same label.

Units with unstable firing throughout the recording period were excluded from further analysis.

Identification of synaptically connected RGC-SC pairs

Monosynaptic connections between RGC axons and SC neurons were detected using established methods (Bereshpolova et al., 2020; Reid and Alonso, 1995; Denman and Contreras, 2015; Usrey et al., 1998) based on statistically significant peaks at synaptic delays. Briefly, we calculated a jittered (timescale jitter 5 ms) version of each spike train by randomizing all spike times within a consecutive 10-15 ms window (Smith and Kohn, 2008). We then calculated the cross-correlation between a pair of neurons both for the original (raw CCG) and the jittered spike train (jittered CCG). Subtracting the jittered CCG from the raw CCG results in a jitter-corrected version of the CCG which was used for monosynaptic pairs detection. Connected pairs were identified using cross-correlation analysis and significant peaks (peak detection window from + 0.5 to 3.5 ms) had to extend over the threshold (baseline + 4x standard deviation) for at least 5 consecutive time bins (0.1 ms resolution) (as described previously, in Sibille et al. (2022) (Sibille et al., 2022a). The cross-correlations were calculated using the Python `pycorrelate` package (<https://github.com/tritemio/pycorrelate>). Please note that for CCG analysis, spike trains were correlated over the full recording duration, except for periods with optogenetic stimulation. Latencies to peak response were calculated from the CCGs as the maximum time within the response window.

Analysis of retinocollicular connection strength

Synaptic efficacy and contribution measures of connected RGC-SC pairs were estimated from spike trains during the entire recording session (except for optotagging periods which were excluded) using standard approaches (Bereshpolova et al., 2020; Reid and Alonso, 1995; Swadlow and Gusev, 2002; Usrey et al., 1998). Briefly, efficacy was estimated from the baseline-corrected CCGs by dividing the area of the CCG peak (baseline window: > -1 to < 0.4 ms, peak window: > 0.4 to < 3.5 ms) by the total number of presynaptic spikes. Thus, an efficacy measure of 1 (100%) would reflect that for each presynaptic retinal spike, a postsynaptic spike could be detected. To estimate the connection contribution, we counted the number of SC spikes that were preceded by a retinal afferent spike, integrated over a time window between -3 to 0.5 ms, and divided this number by the total number of SC spikes. A contribution of 1 (100%) indicates that all spikes of an SC neuron are preceded by retinal afferent spikes.

Paired-spike dynamics analysis

For estimation of paired-spike dynamics, baseline-corrected CCGs were re-calculated triggered on 1st and 2nd RGCs using the same baseline and peak windows as mentioned above. We identified pairs of retinal spikes (1st and 2nd RGC) for each RGC-SC connected pair (Usrey et al., 1998). Briefly, a pair of retinal spikes was defined by a specific inter-spike interval (ISI) where the second RGC spike followed the 1st RGC by a ISI of 5 to 30 ms. The second criterion for classifying a pair of retinal spikes was a certain dead time where the 1st retinal spike was preceded by a minimum period of silence (dead time: 30ms). The dead time criterion was applied to ensure a comparable level of activity immediately preceding all first spikes. The preceding window (> -1 to < 0.4 ms) was used as a baseline window. Paired-spike enhancement was calculated as the ratio of efficacy for the 2nd RGC to the efficacy for the 1st RGC.

Functional similarity analysis of retinocollicular connected pairs

To characterize the functional similarity between connected RGCs and the target neurons in the SC, we estimated the functional similarity index. To do so, we calculated the correlation coefficients from responses during the light and dark sparse noise (r_{SL} and r_{SD}) as well as during the natural movie stimulus (r_{NM}). The correlation of activity during sparse noise stimuli (r_{SL} and r_{SD}) was estimated from the spatiotemporal receptive fields (STRFs) which were calculated from the STA. The similarity during the natural movie stimulus (r_{NM}) was calculated from natural movie PSTHs. We limited this analysis step to neurons with high signal-to-noise ratio (SNR) in the visually evoked activity ($SNR > 8$ for the STRFs). The overall similarity index was then calculated from the averaged correlation coefficients: functional similarity index = $(r_{SD} + r_{SL} + r_{NM})/3$.

Data availability

The datasets that support the findings of this study are available from the corresponding author on request.

Code availability

Custom Python code is available from the corresponding author on request.

References

- Agmon A , Connors BW (1992) **Correlation between intrinsic firing patterns and thalamocortical synaptic responses of neurons in mouse barrel cortex** *J Neurosci Off J Soc Neurosci* **12**:319–329
<https://doi.org/10.1523/JNEUROSCI.12-01-00319.1992>
- Alonso J-M , Swadlow HA (2005) **Thalamocortical specificity and the synthesis of sensory cortical receptive fields** *J Neurophysiol* **94**:26–32
<https://doi.org/10.1152/jn.01281.2004>
- Alonso J-M , Usrey WM , Reid RC (2001) **Rules of Connectivity between Geniculate Cells and Simple Cells in Cat Primary Visual Cortex** *J Neurosci* **21**:4002–4015
<https://doi.org/10.1523/JNEUROSCI.21-11-04002.2001>
- Alonso JM , Usrey WM , Reid RC (1996) **Precisely correlated firing in cells of the lateral geniculate nucleus** *Nature* **383**:815–819
<https://doi.org/10.1038/383815a0>
- Ayupe AC , Choi JS , Beckedorff F , McCartan R , Levay K , Park KK (2023) **Single-Nucleus RNA Sequencing of Developing and Mature Superior Colliculus Identifies Neuronal Diversity and Candidate Mediators of Circuit Assembly** *BioRxiv Prepr Serv Biol*
<https://doi.org/10.1101/2023.02.01.526254>

Baden T , Berens P , Franke K , Román Rosón M , Bethge M , Euler T (2016) **The functional diversity of retinal ganglion cells in the mouse** *Nature* **529**:345–350
<https://doi.org/10.1038/nature16468>

Barchini J , Shi X , Chen H , Cang J (2018) **Bidirectional encoding of motion contrast in the mouse superior colliculus** *eLife* **7**
<https://doi.org/10.7554/eLife.35261>

Bereshpolova Y , Hei X , Alonso J-M , Swadlow HA (2020) **Three rules govern thalamocortical connectivity of fast-spike inhibitory interneurons in the visual cortex** *eLife* **9**
<https://doi.org/10.7554/eLife.60102>

Bereshpolova Y , Stoelzel CR , Gusev AG , Bezdudnaya T , Swadlow HA (2006) **The impact of a corticotectal impulse on the awake superior colliculus** *J Neurosci Off J Soc Neurosci* **26**:2250–2259
<https://doi.org/10.1523/JNEUROSCI.4402-05.2006>

Boka K , Chomsung R , Li J , Bickford ME (2006) **Comparison of the ultrastructure of cortical and retinal terminals in the rat superior colliculus** *Anat Rec A Discov Mol Cell Evol Biol* **288**:850–858
<https://doi.org/10.1002/ar.a.20359>

Bruno RM , Sakmann B (2006) **Cortex is driven by weak but synchronously active thalamocortical synapses** *Science* **312**:1622–1627
<https://doi.org/10.1126/science.1124593>

Bruno RM , Simons DJ (2002) **Feedforward Mechanisms of Excitatory and Inhibitory Cortical Receptive Fields** *J Neurosci* **22**:10966–10975
<https://doi.org/10.1523/JNEUROSCI.22-24-10966.2002>

Campagnola L , Seeman SC , Chartrand T , Kim L , Hoggarth A , Gamlin C , Ito S , Trinh J , Davoudian P , Radaelli C , Kim M-H , Hage T , Braun T , Alfiler L , Andrade J , Bohn P , Dalley R , Henry A , Kebede S , Alice M , Sandman D , Williams G , Larsen R , Teeter C , Daigle TL , Berry K , Dotson N , Enstrom R , Gorham M , Hupp M , Dingman Lee S , Ngo K , Nicovich PR , Potekhina L , Ransford S , Gary A , Goldy J , McMillen D , Pham T , Tieu M , Siverts L , Walker M , Farrell C , Schroedter M , Slaughterbeck C , Cobb C , Ellenbogen R , Gwinn RP , Keene CD , Ko AL , Ojemann JG , Silbergeld DL , Carey D , Casper T , Crichton K , Clark M , Dee N , Ellingwood L , Gloe J , Kroll M , Sulc J , Tung H , Wadhwani K , Brouner K , Egdorf T , Maxwell M , McGraw M , Pom CA , Ruiz A , Bomben J , Feng D , Hejazinia N , Shi S , Szafer A , Wakeman W , Phillips J , Bernard A , Esposito L , D’Orazi FD , Sunkin S , Smith K , Tasic B , Arkhipov A , Sorensen S , Lein E , Koch C , Murphy G , Zeng H , Jarsky T (2022) **Local connectivity and synaptic dynamics in mouse and human neocortex** *Science* **375**
<https://doi.org/10.1126/science.abj5861>

Chandrasekaran AR , Shah RD , Crair MC (2007) **Developmental Homeostasis of Mouse Retinocollicular Synapses** *J Neurosci* **27**:1746–1755
<https://doi.org/10.1523/JNEUROSCI.4383-06.2007>

Reid R Clay , Alonso J-M (1995) **Specificity of monosynaptic connections from thalamus to visual cortex** *Nature* **378**:281–284
<https://doi.org/10.1038/378281a0>

Cruikshank SJ , Lewis TJ , Connors BW (2007) **Synaptic basis for intense thalamocortical activation of feedforward inhibitory cells in neocortex** *Nat Neurosci* **10**:462–468
<https://doi.org/10.1038/nn1861>

Denman DJ , Contreras D (2015) **Complex Effects on In Vivo Visual Responses by Specific Projections from Mouse Cortical Layer 6 to Dorsal Lateral Geniculate Nucleus** *J Neurosci* **35**:9265–9280
<https://doi.org/10.1523/JNEUROSCI.0027-15.2015>

Ellis EM , Gauvain G , Sivyer B , Murphy GJ (2016) **Shared and distinct retinal input to the mouse superior colliculus and dorsal lateral geniculate nucleus** *J Neurophysiol* **116**:602–610
<https://doi.org/10.1152/jn.00227.2016>

Essig J , Hunt JB , Felsen G (2021) **Inhibitory neurons in the superior colliculus mediate selection of spatially-directed movements** *Commun Biol* **4**
<https://doi.org/10.1038/s42003-021-02248-1>

Evans DA , Stempel AV , Vale R , Ruehle S , Lefler Y , Branco T (2018) **A synaptic threshold mechanism for computing escape decisions** *Nature* **558**:590–594
<https://doi.org/10.1038/s41586-018-0244-6>

Ferrarese L , Jouhanneau J-S , Remme MWH , Kremkow J , Katona G , Rózsa B , Schreiber S , Poulet JFA (2018) **Dendrite-Specific Amplification of Weak Synaptic Input during Network Activity In Vivo** *Cell Rep* **24**:3455–3465
<https://doi.org/10.1016/j.celrep.2018.08.088>

Franceschi G , Solomon SG (2018) **Visual response properties of neurons in the superficial layers of the superior colliculus of awake mouse** *J Physiol* **596**:6307–6332
<https://doi.org/10.1113/JP276964>

Froudarakis E , Berens P , Ecker AS , Cotton RJ , Sinz FH , Yatsenko D , Saggau P , Bethge M , Tolias AS (2014) **Population code in mouse V1 facilitates readout of natural scenes through increased sparseness** *Nat Neurosci* **17**:851–857
<https://doi.org/10.1038/nn.3707>

Gabernet L , Jadhav SP , Feldman DE , Carandini M , Scanziani M (2005) **Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition** *Neuron* **48**:315–327
<https://doi.org/10.1016/j.neuron.2005.09.022>

Gale SD , Murphy GJ (2018) **Distinct cell types in the superficial superior colliculus project to the dorsal lateral geniculate and lateral posterior thalamic nuclei** *J Neurophysiol* **120**:1286–1292
<https://doi.org/10.1152/jn.00248.2018>

Gale SD , Murphy GJ (2016) **Active Dendritic Properties and Local Inhibitory Input Enable Selectivity for Object Motion in Mouse Superior Colliculus Neurons** *J Neurosci* **36**:9111–9123

<https://doi.org/10.1523/JNEUROSCI.0645-16.2016>

Gale SD , Murphy GJ (2014) **Distinct representation and distribution of visual information by specific cell types in mouse superficial superior colliculus** *J Neurosci Off J Soc Neurosci* **34**:13458–13471

<https://doi.org/10.1523/JNEUROSCI.2768-14.2014>

Ghitani N , Bayguinov PO , Vokoun CR , McMahon S , Jackson MB , Basso MA (2014) **Excitatory Synaptic Feedback from the Motor Layer to the Sensory Layers of the Superior Colliculus** *J Neurosci* **34**:6822–6833

<https://doi.org/10.1523/JNEUROSCI.3137-13.2014>

Gibson JR , Beierlein M , Connors BW (1999) **Two networks of electrically coupled inhibitory neurons in neocortex** *Nature* **402**:75–79

<https://doi.org/10.1038/47035>

Glickfeld LL , Histed MH , Maunsell JHR (2013) **Mouse Primary Visual Cortex Is Used to Detect Both Orientation and Contrast Changes** *J Neurosci* **33**:19416–19422

<https://doi.org/10.1523/JNEUROSCI.3560-13.2013>

Hoy JL , Bishop HI , Niell CM (2019) **Defined Cell Types in Superior Colliculus Make Distinct Contributions to Prey Capture Behavior in the Mouse** *Curr Biol* **29**:4130–4138

<https://doi.org/10.1016/j.cub.2019.10.017>

Inayat S , Barchini J , Chen H , Feng L , Liu X , Cang J (2015) **Neurons in the Most Superficial Lamina of the Mouse Superior Colliculus Are Highly Selective for Stimulus Direction** *J Neurosci* **35**:7992–8003

<https://doi.org/10.1523/JNEUROSCI.0173-15.2015>

Inoue T , Imoto K (2006) **Feedforward inhibitory connections from multiple thalamic cells to multiple regular-spiking cells in layer 4 of the somatosensory cortex** *J Neurophysiol* **96**:1746–1754

<https://doi.org/10.1152/jn.00301.2006>

Isaacson JS , Scanziani M (2011) **How Inhibition Shapes Cortical Activity** *Neuron* **72**:231–243

<https://doi.org/10.1016/j.neuron.2011.09.027>

Ito S , Feldheim DA , Litke AM (2017) **Segregation of Visual Response Properties in the Mouse Superior Colliculus and Their Modulation during Locomotion** *J Neurosci* **37**:8428–8443

<https://doi.org/10.1523/JNEUROSCI.3689-16.2017>

Ji X , Zingg B , Mesik L , Xiao Z , Zhang LI , Tao HW (2016) **Thalamocortical Innervation Pattern in Mouse Auditory and Visual Cortex: Laminar and Cell-Type Specificity** *Cereb Cortex* **26**:2612–2625

<https://doi.org/10.1093/cercor/bhv099>

Jia X, Siegle JH, Bennett C, Gale SD, Denman DJ, Koch C, Olsen SR (2019) **High-density extracellular probes reveal dendritic backpropagation and facilitate neuron classification** *J Neurophysiol* **121**:1831–1847
<https://doi.org/10.1152/jn.00680.2018>

Jin J, Wang Y, Swadlow HA, Alonso JM (2011) **Population receptive fields of ON and OFF thalamic inputs to an orientation column in visual cortex** *Nat Neurosci* **14**:232–238
<https://doi.org/10.1038/nn.2729>

Jun JJ, Steinmetz NA, Siegle JH, Denman DJ, Bauza M, Barbarits B, Lee AK, Anastassiou CA, Andrei A, Aydın Ç, Barbic M, Blanche TJ, Bonin V, Couto J, Dutta B, Gratiy SL, Gutnisky DA, Häusser M, Karsh B, Ledochowitsch P, Lopez CM, Mitelut C, Musa S, Okun M, Pachitariu M, Putzeys J, Rich PD, Rossant C, Sun W, Svoboda K, Carandini M, Harris KD, Koch C, O’Keefe J, Harris TD (2017) **Fully integrated silicon probes for high-density recording of neural activity** *Nature* **551**:232–236
<https://doi.org/10.1038/nature24636>

Kaneda K, Isa K, Yanagawa Y, Isa T (2008) **Nigral Inhibition of GABAergic Neurons in Mouse Superior Colliculus** *J Neurosci* **28**:11071–11078
<https://doi.org/10.1523/JNEUROSCI.3263-08.2008>

Kara P, Reid RC (2003) **Efficacy of Retinal Spikes in Driving Cortical Responses** *J Neurosci* **23**:8547–8557
<https://doi.org/10.1523/JNEUROSCI.23-24-08547.2003>

Kasai M, Isa T (2016) **Imaging population dynamics of surround suppression in the superior colliculus** *Eur J Neurosci* **44**:2543–2556
<https://doi.org/10.1111/ejn.13371>

Kremkow J, Aertsen A, Kumar A (2010) **Gating of Signal Propagation in Spiking Neural Networks by Balanced and Correlated Excitation and Inhibition** *J Neurosci* **30**:15760–15768
<https://doi.org/10.1523/JNEUROSCI.3874-10.2010>

Kremkow J, Alonso J-M (2018) **Thalamocortical Circuits and Functional Architecture** *Annu Rev Vis Sci* **4**:263–285
<https://doi.org/10.1146/annurev-vision-091517-034122>

Kremkow J, Perrinet LU, Masson GS, Aertsen A (2010) **Functional consequences of correlated excitatory and inhibitory conductances in cortical networks** *J Comput Neurosci* **28**:579–594
<https://doi.org/10.1007/s10827-010-0240-9>

Kremkow J, Perrinet LU, Monier C, Alonso J-M, Aertsen A, Frégnac Y, Masson GS (2016) **Push-Pull Receptive Field Organization and Synaptic Depression: Mechanisms for Reliably Encoding Naturalistic Stimuli in V1** *Front Neural Circuits* **10**

Kyriazi HT, Carvell GE, Simons DJ (1994) **OFF response transformations in the whisker/barrel system** *J Neurophysiol* **72**:392–401
<https://doi.org/10.1152/jn.1994.72.1.392>

Lashley KS (1931) **The mechanism of vision IV. The cerebral areas necessary for pattern vision in the rat** *J Comp Neurol* **53**:419–478
<https://doi.org/10.1002/cne.900530304>

Lee CM , Osman AF , Volgushev M , Escabí MA , Read HL (2016) **Neural spike-timing patterns vary with sound shape and periodicity in three auditory cortical fields** *J Neurophysiol* **115**:1886–1904
<https://doi.org/10.1152/jn.00784.2015>

Lee S-H , Kwan AC , Zhang S , Phoumthippavong V , Flannery JG , Masmanidis SC , Taniguchi H , Huang ZJ , Zhang F , Boyden ES , Deisseroth K , Dan Y (2012) **Activation of specific interneurons improves V1 feature selectivity and visual perception** *Nature* **488**:379–383
<https://doi.org/10.1038/nature11312>

Levick WR , Cleland BG , Dubin MW (1972) **Lateral geniculate neurons of cat: retinal inputs and physiology** *Invest Ophthalmol* **11**:302–311

Li Y , Meister M (2022) **Functional Cell Types in the Mouse Superior Colliculus (preprint)** *Neuroscience*
<https://doi.org/10.1101/2022.04.01.486789>

Lien AD , Scanziani M (2018) **Cortical direction selectivity emerges at convergence of thalamic synapses** *Nature* **558**:80–86
<https://doi.org/10.1038/s41586-018-0148-5>

Masterson SP , Zhou N , Akers BK , Dang W , Bickford ME (2019) **Ultrastructural and optogenetic dissection of V1 corticotectal terminal synaptic properties** *J Comp Neurol* **527**:833–842
<https://doi.org/10.1002/cne.24538>

Mastronarde DN (1987) **Two classes of single-input X-cells in cat lateral geniculate nucleus II. Retinal inputs and the generation of receptive-field properties.** *J Neurophysiol* **57**:381–413
<https://doi.org/10.1152/jn.1987.57.2.381>

Miller KD (2016) **Canonical computations of cerebral cortex** *Curr Opin Neurobiol* **37**:75–84
<https://doi.org/10.1016/j.conb.2016.01.008>

Moore AK , Wehr M (2013) **Parvalbumin-expressing inhibitory interneurons in auditory cortex are well-tuned for frequency** *J Neurosci Off J Soc Neurosci* **33**:13713–13723
<https://doi.org/10.1523/JNEUROSCI.0663-13.2013>

Muellner FE , Roska B (2023) **Individual thalamic inhibitory interneurons are functionally specialized towards distinct visual features (preprint)** *Neuroscience*
<https://doi.org/10.1101/2023.03.22.533751>

Niell CM , Scanziani M (2021) **How Cortical Circuits Implement Cortical Computations: Mouse Visual Cortex as a Model** *Annu Rev Neurosci* **44**:517–546
<https://doi.org/10.1146/annurev-neuro-102320-085825>

- Niell CM , Stryker MP (2008) **Highly Selective Receptive Fields in Mouse Visual Cortex** *J Neurosci* **28**:7520–7536
<https://doi.org/10.1523/JNEUROSCI.0623-08.2008>
- Ono M , Bishop DC , Oliver DL (2017) **Identified GABAergic and Glutamatergic Neurons in the Mouse Inferior Colliculus Share Similar Response Properties** *J Neurosci* **37**:8952–8964
<https://doi.org/10.1523/JNEUROSCI.0745-17.2017>
- Pachitariu M , Steinmetz N , Kadir S , Carandini M , Kenneth D. H (2016) **Kilosort: realtime spike-sorting for extracellular electrophysiology with hundreds of channels (preprint)** *Neuroscience*
<https://doi.org/10.1101/061481>
- Peirce J , Gray JR , Simpson S , MacAskill M , Höchenberger R , Sogo H , Kastman E , Lindeløv JK (2019) **PsychoPy2: Experiments in behavior made easy** *Behav Res Methods* **51**:195–203
<https://doi.org/10.3758/s13428-018-01193-y>
- Petruno SK , Clark RE , Reinagel P (2013) **Evidence that primary visual cortex is required for image, orientation, and motion discrimination by rats** *PloS One* **8**
<https://doi.org/10.1371/journal.pone.0056543>
- Phongphanphanee P , Mizuno F , Lee PH , Yanagawa Y , Isa T , Hall WC (2011) **A Circuit Model for Saccadic Suppression in the Superior Colliculus** *J Neurosci* **31**:1949–1954
<https://doi.org/10.1523/JNEUROSCI.2305-10.2011>
- Porter JT , Johnson CK , Agmon A (2001) **Diverse Types of Interneurons Generate Thalamus-Evoked Feedforward Inhibition in the Mouse Barrel Cortex** *J Neurosci* **21**:2699–2710
<https://doi.org/10.1523/JNEUROSCI.21-08-02699.2001>
- Quirk MC , Sosulski DL , Feierstein CE , Uchida N , Mainen ZF (2009) **A Defined Network of Fast-Spiking Interneurons in Orbitofrontal Cortex: Responses to Behavioral Contingencies and Ketamine Administration** *Front Syst Neurosci* **3**
<https://doi.org/10.3389/neuro.06.013.2009>
- Mize R Ranney (1992) **Chapter 11 The organization of GABAergic neurons in the mammalian superior colliculus** *Progress in Brain Research Elsevier* :219–248
[https://doi.org/10.1016/S0079-6123\(08\)63616-X](https://doi.org/10.1016/S0079-6123(08)63616-X)
- Reid RC , Alonso JM (1995) **Specificity of monosynaptic connections from thalamus to visual cortex** *Nature* **378**:281–284
<https://doi.org/10.1038/378281a0>
- Reinhard K , Li C , Do Q , Burke EG , Heynderickx S , Farrow K (2019) **A projection specific logic to sampling visual inputs in mouse superior colliculus** *eLife* **8**
<https://doi.org/10.7554/eLife.50697>

Rompani SB , Müllner FE , Wanner A , Zhang C , Roth CN , Yonehara K , Roska B (2017) **Different Modes of Visual Integration in the Lateral Geniculate Nucleus Revealed by Single-Cell-Initiated Transsynaptic Tracing** *Neuron* **93**:767–776
<https://doi.org/10.1016/j.neuron.2017.01.028>

Rossant C , Kadir SN , Goodman DFM , Schulman J , Hunter MLD , Saleem AB , Grosmark A , Belluscio M , Denfield GH , Ecker AS , Tolias AS , Solomon S , Buzsaki G , Carandini M , Harris KD (2016) **Spike sorting for large, dense electrode arrays** *Nat Neurosci* **19**:634–641
<https://doi.org/10.1038/nn.4268>

Schröder S , Steinmetz NA , Krumin M , Pachitariu M , Rizzi M , Lagnado L , Harris KD , Carandini M. (2020) **Arousal Modulates Retinal Output** *Neuron* **107**:487–495
<https://doi.org/10.1016/j.neuron.2020.04.026>

Shang C , Liu A , Li D , Xie Z , Chen Z , Huang M , Li Y , Wang Y , Shen WL , Cao P (2019) **A subcortical excitatory circuit for sensory-triggered predatory hunting in mice** *Nat Neurosci* **22**:909–920
<https://doi.org/10.1038/s41593-019-0405-4>

Shang C , Liu Z , Chen Z , Shi Y , Wang Q , Liu S , Li D , Cao P (2015) **BRAIN CIRCUITS. A parvalbumin-positive excitatory visual pathway to trigger fear responses in mice** *Science* **348**:1472–1477
<https://doi.org/10.1126/science.aaa8694>

Shi X , Barchini J , Ledesma HA , Koren D , Jin Y , Liu X , Wei W , Cang J (2017) **Retinal origin of direction selectivity in the superior colliculus** *Nat Neurosci* **20**:550–558
<https://doi.org/10.1038/nn.4498>

Sibille J , Gehr C , Benichov JI , Balasubramanian H , Teh KL , Lupashina T , Vallentin D , Kremkow J (2022) **High-density electrode recordings reveal strong and specific connections between retinal ganglion cells and midbrain neurons** *Nat Commun* **13**
<https://doi.org/10.1038/s41467-022-32775-2>

Sibille J , Gehr C , Teh KL , Kremkow J (2022) **Tangential high-density electrode insertions allow to simultaneously measure neuronal activity across an extended region of the visual field in mouse superior colliculus** *J Neurosci Methods* **376**
<https://doi.org/10.1016/j.jneumeth.2022.109622>

Siegle JH, Jia X, Durand S, Gale S, Bennett C, Graddis N, Heller G, Ramirez TK, Choi H, Luviano JA, Groblewski PA, Ahmed R, Arkhipov A, Bernard A, Billeh YN, Brown D, Buice MA, Cain N, Caldejon S, Casal L, Cho A, Chvilicek M, Cox TC, Dai K, Denman DJ, de Vries SEJ, Dietzman R, Esposito L, Farrell C, Feng D, Galbraith J, Garrett M, Gelfand EC, Hancock N, Harris JA, Howard R, Hu B, Hytnen R, Iyer R, Jessett E, Johnson K, Kato I, Kiggins J, Lambert S, Lecoq J, Ledochowitsch P, Lee JH, Leon A, Li Y, Liang E, Long F, Mace K, Melchior J, Millman D, Mollenkopf T, Nayan C, Ng L, Ngo K, Nguyen T, Nicovich PR, North K, Ocker GK, Ollerenshaw D, Oliver M, Pachitariu M, Perkins J, Reding M, Reid D, Robertson M, Ronellenfitch K, Seid S, Slaughterbeck C, Stoecklin M, Sullivan D, Sutton B, Swapp J, Thompson C, Turner K, Wakeman W, Whitesell JD, Williams D, Williford A, Young R, Zeng H, Naylor S, Phillips JW, Reid RC, Mihalas S, Olsen SR, Koch C (2021) **Survey of spiking in the mouse visual system reveals functional hierarchy** *Nature*
<https://doi.org/10.1038/s41586-020-03171-x>

Siegle JH, López AC, Patel YA, Abramov K, Ohayon S, Voigts J (2017) **Open Ephys: an open-source, plugin-based platform for multichannel electrophysiology** *J Neural Eng* **14**
<https://doi.org/10.1088/1741-2552/aa5eea>

Simons DJ, Carvell GE (1989) **Thalamocortical response transformation in the rat vibrissa/barrel system** *J Neurophysiol* **61**:311–330
<https://doi.org/10.1152/jn.1989.61.2.311>

Smith MA, Kohn A (2008) **Spatial and Temporal Scales of Neuronal Correlation in Primary Visual Cortex** *J Neurosci* **28**:12591–12603
<https://doi.org/10.1523/JNEUROSCI.2929-08.2008>

Sun Q-Q, Huguenard JR, Prince DA (2006) **Barrel cortex microcircuits: thalamocortical feedforward inhibition in spiny stellate cells is mediated by a small number of fast-spiking interneurons** *J Neurosci Off J Soc Neurosci* **26**:1219–1230
<https://doi.org/10.1523/JNEUROSCI.4727-04.2006>

Swadlow HA (2003) **Fast-spike Interneurons and Feedforward Inhibition in Awake Sensory Neocortex** *Cereb Cortex* **13**:25–32
<https://doi.org/10.1093/cercor/13.1.25>

Swadlow HA, Gusev AG (2002) **Receptive-field construction in cortical inhibitory interneurons** *Nat Neurosci* **5**:403–404
<https://doi.org/10.1038/nn847>

Swadlow HA, Gusev AG (2001) **The impact of “bursting” thalamic impulses at a neocortical synapse** *Nat Neurosci* **4**:402–408
<https://doi.org/10.1038/86054>

Swadlow HA, Gusev AG, Bezdudnaya T (2002) **Activation of a Cortical Column by a Thalamocortical Impulse** *J Neurosci* **22**:7766–7773
<https://doi.org/10.1523/JNEUROSCI.22-17-07766.2002>

Tremblay R, Lee S, Rudy B (2016) **GABAergic interneurons in the neocortex: From cellular properties to circuits** *Neuron* **91**:260–292
<https://doi.org/10.1016/j.neuron.2016.06.033>

Tsai NY , Fei Wang , Toma K , Yin C , Takatoh J , Pai EL , Wu K , Matcham AC , Yin L , Dang EJ , Marciano DK , Rubenstein JL , Fan Wang , Ullian EM , Duan X (2022) **Trans-Seq maps a selective mammalian retinotectal synapse instructed by Nephronectin** *Nat Neurosci* **25**:659–674

<https://doi.org/10.1038/s41593-022-01068-8>

Usrey WM , Alonso J-M , Reid RC (2000) **Synaptic Interactions between Thalamic Inputs to Simple Cells in Cat Visual Cortex** *J Neurosci* **20**:5461–5467

<https://doi.org/10.1523/JNEUROSCI.20-14-05461.2000>

Usrey WM , Reppas JB , Reid RC (1999) **Specificity and Strength of Retinogeniculate Connections** *J Neurophysiol* **82**:3527–3540

<https://doi.org/10.1152/jn.1999.82.6.3527>

Usrey WM , Reppas JB , Reid RC (1998) **Paired-spike interactions and synaptic efficacy of retinal inputs to the thalamus** *Nature* **395**:384–387

<https://doi.org/10.1038/26487>

Villalobos CA , Wu Q , Lee PH , May PJ , Basso MA (2018) **Parvalbumin and GABA Microcircuits in the Mouse Superior Colliculus** *Front Neural Circuits* **12**

<https://doi.org/10.3389/fncir.2018.00035>

Wei P , Liu N , Zhang Z , Liu X , Tang Y , He X , Wu B , Zhou Z , Liu Y , Li J , Zhang Y , Zhou X , Xu L , Chen L , Bi G , Hu X , Xu F , Wang L (2015) **Processing of visually evoked innate fear by a non-canonical thalamic pathway** *Nat Commun* **6**

<https://doi.org/10.1038/ncomms7756>

Whyland KL , Slusarczyk AS , Bickford ME (2020) **GABAergic cell types in the superficial layers of the mouse superior colliculus** *J Comp Neurol* **528**:308–320

<https://doi.org/10.1002/cne.24754>

Zhang Z , Liu W-Y , Diao Y-P , Xu W , Zhong Y-H , Zhang J-Y , Lazarus M , Liu Y-Y , Qu W-M , Huang Z-L (2019) **Superior Colliculus GABAergic Neurons Are Essential for Acute Dark Induction of Wakefulness in Mice** *Curr Biol CB* **29**:637–644

<https://doi.org/10.1016/j.cub.2018.12.031>

Zhao S , Ting JT , Atallah HE , Qiu L , Tan J , Gloss B , Augustine GJ , Deisseroth K , Luo M , Graybiel AM , Feng G (2011) **Cell type-specific channelrhodopsin-2 transgenic mice for optogenetic dissection of neural circuitry function** *Nat Methods* **8**:745–752

<https://doi.org/10.1038/nmeth.1668>

Zucker RS (1989) **Short-term synaptic plasticity** *Annu Rev Neurosci* **12**:13–31

<https://doi.org/10.1146/annurev.ne.12.030189.000305>

Author information

Carolin Gehr

Neuroscience Research Center, Charité-Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Bernstein Center for Computational Neuroscience Berlin, Philippstraße 13, 10115 Berlin, Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Philippstraße 13, 10115 Berlin, Einstein Center for Neurosciences Berlin, Charitéplatz 1, 10117 Berlin
ORCID iD: [0000-0003-0892-506X](https://orcid.org/0000-0003-0892-506X)

Jérémie Sibille

Neuroscience Research Center, Charité-Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Bernstein Center for Computational Neuroscience Berlin, Philippstraße 13, 10115 Berlin, Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Philippstraße 13, 10115 Berlin, Einstein Center for Neurosciences Berlin, Charitéplatz 1, 10117 Berlin

Jens Kremkow

Neuroscience Research Center, Charité-Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Bernstein Center for Computational Neuroscience Berlin, Philippstraße 13, 10115 Berlin, Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Philippstraße 13, 10115 Berlin, Einstein Center for Neurosciences Berlin, Charitéplatz 1, 10117 Berlin

For correspondence: jens.kremkow@charite.de

ORCID iD: [0000-0001-7077-4528](https://orcid.org/0000-0001-7077-4528)

Editors

Reviewing Editor

Marla Feller

University of California, Berkeley, United States of America

Senior Editor

Sacha Nelson

Brandeis University, United States of America

Reviewer #1 (Public Review):

Gehr and colleagues used an elegant method, using neuropixels probes, to study retinal input integration by mouse superior collicular cells in vivo. Compared to a previous report of the same group, they opto-tagged inhibitory neurons and defined the differential integration onto each group. Through these experiments, the author concluded that overall, there is no clear difference between the retina connectivity to excitatory and inhibitory superior colliculus neurons. The exception to that rule is that excitatory neurons might be driven slightly stronger than inhibitory ones. Technically, this work is performed at a high level, and the plots are beautifully conceived, but I have doubts if the interpretation given by the authors is solid. I will elaborate below.

Some thoughts about the interpretation of the results.

My main concern is the "survivor bias" of this work, which can lead to skewed conclusions. From the data set acquired, 305 connections were measured, 1/3 inhibitory and 2/3 excitatory. These connections arise from 83 RGC onto 124 RGC (I'm interpreting the axis of Fig.2 C). Here it is worth mentioning that different RGC types have different axonal diameters (Perge et al., 2009). Here the diameter is also related to the way cells relay information (max frequencies, for example). It is possible that thicker axons are easier to measure, given the larger potential changes would likely occur, and thus, selectively being picked up by the neuropixels probe. If this is the case, we would have a clear case of "survivor bias", which should be tested and discussed. One way to determine if the response properties of axonal termini are from an unbiased sample is to make a rough functional characterization as generally performed (see Baden et al. 2006). This is fundamental since all other conclusions are based on unbiased sampling.

One aspect that is not clear to me is to measure of connectivity strength in Figure 2. Here it seems that connectivity strength is directly correlated with the baseline firing rate of the SC neuron (see example plots). If this is a general case, the synaptic strength can be assumed but would only differ in strength due to the excitability of the postsynaptic cell. This should be tested by plotting the correlation coefficient analysis against the baseline firing rate.

My third concern is the assessment of functional similarity in Fig. 3. It is not clear to me why the similarity value was taken by the arithmetic mean. For example, even if the responses are identical for one connected pair that exclusively responds either to the ON or OFF sparse noise, the maximal value can only be 0.67. Perhaps I misunderstood something. Secondly, correlations in natural(istic) movies can differ dramatically depending on the frame rate that the movie was acquired and the way it is displayed to the animal. What looks natural to us will elicit several artifacts at a retinal level, e.g., due to big jumps between frames (no direction-selective response) or overall little modulation (large spatial correlations). I would rather opt for uniform stimuli, as suggested previously. Of course, these are also approximations but can be easily reproduced by different labs and are not subjected to the intricacies of the detailed naturalistic stimulus used.

Fourth. It is important to control the proportion of inhibitory cells activated optogenetically across the recording probe. Currently, it is not possible to assess if there are false negatives. One way of controlling for this would be to show that the number of inhibitory interneurons doesn't vary across the probe.

Fifth. In Fig. 4, the ISI had a minimal bound of 5 ms. Why? This would cap the firing rate at 200Hz, but we know that RGC in explants can fire at higher frequencies for evoked responses. I would set a lower bound since it should come naturally from the after-depolarization block. Another aspect that remains unclear is to what extent the paired-spike ratio depends on the baseline firing rate. This would change the interpretation from the particular synaptic connection to the intrinsic properties of the cell and is plausible since the baseline firing rate varies tremendously. One related analysis would be to plot the change of PSR depending on the ISI. It would be intuitive to make a scatter plot for all paired spikes of all recorded neurons (separated into inhibitory and excitatory) of ISI vs. PSR.

Panel 4E is confusing to me. Here what is plotted is efficacy 1st against PSR (which is efficacy 2nd/efficacy 1st). Given that you have a linear relation between efficacy 1st and efficacy 2nd (panel 4C), you are essentially re-plotting the same information, which should necessarily have a hyperbolic relationship: $f(x) = y/x$. Thus, fitting this with a linear function makes no sense and it has to be decaying if efficacy 2nd > efficacy 1st as shown in 4C.

Finally, in Figure 5, the perspective is inverted, and the spike correlations are seen from the perspective of SC neurons. Here it would also be good to plot the cumulative histograms and not look at the averages. Regarding the similarity index and use of natural stats, please see

my previous comments. Also, would it be possible to plot the contribution v/s the firing rate with the baseline firing rate with no stimulation or full-field stimulation? This is important since naturalistic movies have too many correlations and dependencies that make this plot difficult to interpret.

Overall, the paper only speaks from excitatory and inhibitory differences in the introduction and results. However, it is known that there are three clear morphologically distinct classes of excitatory neurons (wide-field, narrow-field, and stellate). This topic is touched in the discussion but not directly in the context of these results. Smaller cells might likely be driven much stronger. Wide-field cells would likely not be driven by one RGC input only and will probably integrate from many more cells than 6.

Reviewer #2 (Public Review):

This study follows up on a previous study by the group (Sibille et al Nature Communications 2022) in which high density Neuropixel probes were inserted tangentially through the superficial layers of the superior colliculus (SC) to record the activity of retinocollicular axons and postsynaptic collicular neurons in anesthetized mice. By correlating spike patterns, connected pairs could be identified which allowed the authors to demonstrate that functionally similar retinal axon-SC neuron pairs were strongly connected.

In the current study, the authors use similar techniques in vGAT-ChR2 mice and add a fiber optic to identify light-activated GABAergic and non-light-activated nonGABAergic neurons. Using their previously verified techniques to identify connected pairs, within regions of optogenetic activation they identified 214 connected pairs of retinal axons and nonGABAergic neurons and 91 pairs of connected retinal axons and GABAergic neurons. The main conclusion is that retinal activity contributed more to the activity of postsynaptic nonGABAergic SC neurons than to the activity of postsynaptic GABAergic SC neurons.

The study is very well done. The figures are well laid out and clearly establish the conclusions. My main comments are related to the comparison to other circuits and further questions that might be addressed in the SC.

It is stated several times that the superior colliculus and the visual cortex are the two major brain areas for visual processing and these areas are compared throughout the manuscript. However, since both the dorsal lateral geniculate nucleus (dLGN) and SC include similar synaptic motifs, including triadic arrangements of retinal boutons with GABAergic and nonGABAergic neurons, it might be more relevant to compare and contrast retinal convergence and other features in these structures.

The GABAergic and nonGABAergic neurons showed a wide range of firing rates. It might be interesting to sort the cells by firing rates to see if they exhibit different properties. For example, since the SC contains both GABAergic interneurons and projection neurons it would be interesting to examine whether GABAergic neurons with higher firing rates exhibit narrower spikes, similar to cortical fast spiking interneurons. Similarly, it might be of interest to sort the neurons by their receptive field sizes since this is associated with different SC neuron types.

The recording techniques allowed for the identification of the distance between connected retinocollicular fibers and postsynaptic neurons. It might also be interesting to compare the properties of connected pairs recorded at dorsal versus ventral locations since neurons with different genetic identities and response properties are located in different dorsal/ventral locations (e.g. Liu et al. Neuron 2023). Also, regarding the strength of connections, previous

electron microscopy studies have shown that the retinocollicular terminals differ in density and size in the dorsal/ventral dimension (e.g Carter et al JCN 1991).

Was optogenetic activation of GABAergic neurons ever paired with visual activation? It would be interesting to examine the receptive fields of the nonGABAergic neurons before and after activation of the GABAergic neurons (as in Gale and Murphy J Neurosci 2016).

Reviewer #3 (Public Review):

This study performs in vivo recordings of neurons in the mouse superior colliculus and their afferents from the retina, retinal ganglion cells (RGCs). Building on a preparation they previously published, this study adds the use of optogenetic identification of inhibitory neurons (aka optotagging) to compare RGC connectivity to excitatory and inhibitory neurons in SC. Using this approach, the authors characterize connection probability, strength, and response correlation between RGCs and their target neurons in SC, finding several differences from what is observed in the retina-thalamus-visual cortex pathway. As such, this may be a useful dataset for efforts to understand retinocollicular connectivity and computations.