

The involvement of the human prefrontal cortex in the emergence of visual awareness

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

August 10, 2023 (this version)

Posted to bioRxiv

June 9, 2023

Sent for peer review

May 30, 2023

Zepeng Fang, Yuanyuan Dang, Zhipei Ling, Yongzheng Han, Hulin Zhao , Xin Xu , Mingsha Zhang 

State Key Laboratory of Cognitive Neuroscience and Learning and IDG/McGovern Institute for Brain Research, Division of Psychology, Beijing Normal University, Beijing 100875, China • Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China • Department of Anesthesiology, Peking University Third Hospital, Beijing, China

 https://en.wikipedia.org/wiki/Open_access

 <https://creativecommons.org/licenses/by/4.0/>

Abstract

Exploring the neural mechanisms of awareness is a fundamental task of cognitive neuroscience. There is an ongoing dispute regarding the role of the prefrontal cortex (PFC) in the emergence of awareness, which is partially raised by the confound between report-and awareness-related activity. To address this problem, we designed a visual awareness task that can minimize report-related confounding. Our results show that saccadic latency is significantly shorter in the aware trials than in the unaware trials. Local field potential (LFP) data from 6 patients consistently show early (200-300 ms) awareness-related activity in the PFC, including event-related potential and high-gamma activity. Moreover, the awareness state can be reliably decoded by the neural activity in the PFC since the early stage, and the neural pattern is dynamically changed rather than being stable during the representation of awareness. Furthermore, the enhancement of dynamic functional connectivity, through the phase modulation at low frequency, between the PFC and other brain regions in the early stage of the awareness trials may explain the mechanism of conscious access. And the spatiotemporal overlap between the awareness-related activity and the inter-regional connectivity in PFC suggested that conscious access and phenomenal awareness may be closely coupled. These results indicate that the PFC is critically involved in the emergence of awareness.

eLife assessment

This paper reports **valuable** results regarding the potential role and time course of the prefrontal cortex in conscious perception. Although the sample size is small, the results are clear and **convincing**, and strengths include the use of several complementary analysis methods. The behavioral test includes subject report so the results do not allow for distinguishing between theories of consciousness; nevertheless, results do advance our understanding of the contribution of prefrontal cortex to conscious perception.

A conscious state means that a person has some kind of subjective experiences, such as seeing a painting, hearing a sound, thinking about a problem, or feeling an emotion ¹. The neural mechanism of consciousness is one of the most fundamental, exciting and challenging pursuits in the 21st-century brain science ². Among the various aspects of consciousness, visual awareness, as a core of consciousness, has attracted enormous attention in consciousness research. Visual awareness seems quite simple: we see an object and seem to immediately know its shape, color and other properties. However, the neural processes behind the emergence of visual awareness are quite sophisticated and largely unknown ³.

The typical method to study the neural mechanism of visual awareness is to present external stimuli with similar physical properties but to elicit different conscious experiences, such as using near perceptual threshold stimuli, visual masking, or attentional blinks ^{4,5}. Then, the aware and unaware states are determined through the subjective reports of participants, and the awareness-related neural activity is dissociated by the brain activation difference between the two states. Using this approach, some studies have found that activities in broad brain regions are correlated with visual awareness. For example, an fMRI study of lexical visual masking found that only consciously perceived and reported words can induce activation of higher-order cortical regions such as the frontal and parietal lobes ⁶. Another study using a visual masking task found changes in the amplitude of intracranial electroencephalography (iEEG) signals and high-gamma power in occipital-temporal regions, which indicated that these regions played an important role in the emergence of awareness ⁷.

However, there is still much debate about the origin of conscious awareness in the brain. Some hypotheses (global neuronal workspace theory, higher-order theory, etc.) argued that visual awareness arose from the frontoparietal network, in which the prefrontal cortex played a key role, and they argued that the earliest biomarker of awareness in EEG is the P3b wave onset approximately 300 ms after stimulus onset ^{8,9}. For example, based on the electrophysiological study of a visual awareness-guided saccade selection paradigm in macaques, by comparing the action potential firing characteristics of neurons in the dorsolateral prefrontal cortex (dlPFC), V4 and V1, only dlPFC neurons were closely associated with visual awareness states. Therefore, the authors argued that stimuli must elicit a threshold level of prefrontal activity for conscious detection to occur ¹⁰. In contrast, other hypotheses (integrated information theory, recurrent processing theory, etc.) proposed that the "hot zone" of the posterior cortex of the brain, including the parietal lobe, occipital lobe, and temporal lobe, is sufficient to generate visual awareness, and its biomarkers on EEG are the relatively earlier visual awareness negativity (VAN) at approximately 200 ms after stimulus onset ^{11,12-13}, whereas the PFC is not necessary for the emergence of conscious experience.

This controversy is partially caused by the confounding effect of report-related activity, such as motor planning and execution, on awareness-related activity. The reason is that the study of awareness usually requires subjective reports to determine different awareness states, while the prefrontal cortex is closely related to such report-related functions ¹⁴. To solve this problem, some researchers use the no-report paradigm to rule out report-related impacts. For instance, a functional magnetic resonance imaging (fMRI) study employing human binocular rivalry paradigms found that when subjects need to manually report their awareness states, the frontal, parietal, and occipital lobes are all involved in the process of binocular rivalry, but when a report is not needed, rivalry-related activation disappears in the frontal lobe but remains in the occipital and parietal lobes ¹⁵. However, the no-report paradigm may overestimate unconscious processing because it cannot directly measure the awareness state but infers the awareness state through other relevant physiological indicators, such as optokinetic nystagmus and pupil size ¹⁶.

Therefore, neither the typical report paradigm nor the no-report paradigm is ideal for visual awareness research; thus, it is difficult to solve the problem of whether the PFC participates in conscious perception.

In addition, when studying the role of the PFC in visual awareness, the noninvasive technologies mostly used in previous studies, i.e., Functional magnetic resonance imaging (fMRI) and M/EEG have limitations. Although fMRI research can detect the whole brain activation pattern, it is difficult to describe the dynamic changes and sequences of different brain regions in the rapid process of visual awareness due to its limited time resolution (approximately several seconds). For example, even though a recent fMRI study using the no-reported paradigm found that there are still visual awareness-related activities in the PFC ¹⁷, it remains unclear when these activities begin and whether PFC activity occurs earlier or later than other brain regions. At the same time, although EEG and MEG have a very high temporal resolution, their low spatial resolution and low signal-to-noise ratio limit their interpretation of awareness-related activities, especially high-frequency activities (highly correlated with local population neuronal activities). For example, although increasing evidence shows that the earliest biomarker of visual awareness in EEG is more likely to be the VAN, which starts at approximately 180-300 ms (may be delayed to 300-460 ms for low-visible stimulus) and peaks at the parieto-occipital region ¹⁸⁻²⁰, rather than the P3b wave onset at approximately 300-500 ms which peaks in the central frontoparietal region, it is still unclear whether there are local visual awareness-related activities in the PFC during the VAN period. In addition, a few recent studies have suggested that awareness may be related to the activities of brain networks, including the PFC, rather than that of a single brain region. However, most of these brain network studies used noninvasive research methods ^{17,21}, and due to the limitations mentioned above, it is difficult to detect the dynamic changes in brain networks in a more precise spatial-temporal profile during the generation of visual awareness.

To address the questions mentioned above, we first designed a novel visual awareness task (**Figure 1**). The contrast of the cue stimulus (grating) was roughly divided into three levels: well above the visual perceptual threshold, near the visual perceptual threshold and zero contrast (no stimulus). During each experiment, in nearly 80% of trials, the contrast of the grating was close to the subject's visual perceptual threshold so that the subjects could sometimes see the grating and sometimes could not. Subjects were required to choose a saccade direction according to the visual awareness state (seeing the grating or not seeing the grating) and the color of the fixation point. Notably, subjects were unable to choose the saccade direction or effectively prepare for saccades until the color of the fixation point was changed. Moreover, the subjects needed to choose the saccade direction, both in the awareness and unawareness states, according to the color of the fixation point so that the report behaviors were matched between the two awareness states. Such a paradigm can effectively dissociate awareness-related activity from report-related activity in terms of time (the fixation point changes color 650 ms after the grating onset) and report behavior, thereby being report-independent of the awareness states ²², thus minimizing the impact of report-related factors on visual awareness. In addition, saccade parameters (such as reaction time, etc.) under different awareness states can be compared to explore the influence of visual awareness on saccade behavior. Then, we applied the above behavioral paradigm to 6 clinical patients with implanted electrodes (stereoelectroencephalography, sEEG) in prefrontal and other cortices and 10 healthy subjects. The behavior and local field potential (LFP) data of 6 patients were recorded while they performed the task, and we also collected the behavioral data of 10 healthy subjects for comparison with the patients' behavioral data.




Figure 1.

Schematic diagram of the visual awareness task.

A trial started when a fixation point ($0.5^\circ \times 0.5^\circ$, white cross) appeared at the center of the screen (radius of eye position check window is 4° , the dotted circle). After the subject fixated on the fixation point for 600 ms, a cue stimulus (Gabor grating, $2 \times 2^\circ$ circle) was presented for 50 ms at a fixed position (7°) on the left (or right, see Methods) side of the screen for all participants. In 70% of the trials, the grating contrast was maintained near the subject's perceptual threshold by a staircase method; in 10% of the trials, the stimulus contrast was well above the threshold; and in the other 20% of the trials, the stimulus contrast was 0, namely, no stimulus appeared. After another 600 ms delay, the color of the fixation point turned red or green, and two saccade targets ($1 \times 1^\circ$, white square) appeared at fixed positions (10°) on the left and right sides of the screen. If the grating was seen, the green fixation point was required to make a saccade to the right target, while the red fixation point was required to make a saccade to the left target. If the grating was not seen, the rule of saccadic direction was inverted.

Behavioral Results

To detect the effect of cue stimulus (grating) contrast on the emergence of visual awareness, we first analyzed the proportion of participants being aware of the grating as a function of its contrast. **Figure 2A** shows exemplified sessions from a patient and healthy subject, and **Figure 2B** shows the population results of the two participant groups. The results showed that with increasing grating contrast, the ratio of reporting ‘awareness’ of the grating gradually increased for all participants (including patients and healthy subjects), showing a classical psychometric curve. Such results showed that the grating contrast level had a direct impact on the emergence of visual awareness and thus is a reliable way to study visual awareness. In addition, in trials with no grating and grating contrast well above the perceptual threshold, subjects showed high accuracy (patients $94.75\% \pm 2.65$; healthy subjects, $96.80\% \pm 0.71$, mean $\pm$ SEM), indicating that the subjects understood and performed the task well following the task rules.

To explore whether visual awareness affects behavioral performance, we divided the trials into 4 groups according to the level of grating contrast and the participants’ reported awareness state, i.e., high-contrast aware (HA), near-threshold-contrast aware (NA), near-threshold-contrast unaware (NU), and low-contrast unaware (LU) (**Figure 2A**, also see Methods). We compared the saccadic parameters between the awareness and unawareness states in near-threshold contrast levels (NA vs. NU), as well as in high and low contrast levels (HA vs. LU), as a control. While saccadic reaction time showed a significant difference between the two awareness states (**Figure 2B**), the other parameters did not. **Figure 2B** shows the mean saccadic reaction time of patients and healthy subjects under different awareness states and contrast levels. The results showed that the average saccadic reaction time was significantly shorter under the awareness state than under the unawareness state at both the near-threshold contrast level and the high-low contrast level ($p < 0.05$, Wilcoxon signed rank test), and it was also explicit and consistent at the individual level. The results showed that the saccadic reaction time in the aware trials was systematically shorter than that in the unaware trials. Such results demonstrate that visual awareness significantly affects the speed of information processing in the brain.

Local Field Potential Results

While the patients were performing our visual awareness task, we recorded the local field potential (LFP) from 901 recording sites (**Figure 2D**, projected on Montreal Neurological Institute (MNI) brain template ICBM152), in which 245 sites were located in the PFC. The table in **Figure 2D** shows a summary of the proportions of recording sites in different lobes and structures.

Early visual awareness-related iERP activity in the PFC

While looking through the event-related response of the LFP data, a striking phenomenon in all patients was that the amplitude of LFP showed remarkably different patterns between awareness and unawareness states. We show the grand average (left) and single-trial data (right) of LFP activity of recording sites in the PFC from each patient in **Figure 3A**, which represents the typical awareness-related neural activity of the PFC. There was a vigorous visual evoked response under the HA condition (light red line) compared to the LU (light blue line) condition. However, in the near-threshold condition, although the contrast of the grating was similar (see Results below), the LFP amplitude of the NA (red line) and NU (blue line) trials started to differ significantly at approximately 200-300 ms (thick black line in panels, $p < 0.01$ corrected, by independent t-test, details see Methods). This difference is also very explicit and consistent at the single-trial (>180 trials for NA and NU conditions in individual participants, see Methods) and single-subject level.




Figure 2.

Behavioral performance and electrode location

A. Psychometric detection curve in a single session. Left panels from one patient and right from one healthy subject. Each black point in the graph represents the aware percent in a contrast level, and the black curve represents the fitted psychometric function. The contrast level resulted in an awareness percentage greater than 25%, and less than 75% was defined as near-threshold, whereas an awareness percentage less than 25% was low and greater than 75% was high.

B. Psychometric detection curves for all participants. The same as Panel A except the contrast is aligned to the individual subject's perceptual threshold, i.e., the contrast 0 represents each subject's perceptual threshold. (N, number of subjects; R^2 , coefficient of determination).

C. Saccadic reaction time in the aware and unaware trials under high+low-and near-threshold-level contrasts of gratings. Bars show the mean values of saccadic reaction time. In the near-threshold condition, the mean reaction times of patients and healthy subjects in the aware trials were 676.55 ± 42.38 and 556.16 ± 28.45 ms, respectively. The mean reaction times in the unaware trials were 892.57 ± 43.28 and 702.30 ± 30.93 ms, respectively, and the p values were 0.03 and 9.77×10^{-4} , respectively, Wilcoxon signed-rank test. In the high+low condition, the mean reaction times for patients and healthy subjects under the aware trials were 577.78 ± 55.28 and 524.08 ± 31.06 ms, and the mean reaction times of the unaware trials were 898.64 ± 57.21 and 687.59 ± 49.07 ms, respectively, with p values of 0.002 and 9.77×10^{-4} , respectively, Wilcoxon signed-rank test. Each black dot represents one participant. The gray solid line represents the paired dots. Error bars represent the standard error of the mean (SEM).

D. Left, right and top views of all recording sites projected on an MNI brain template. Each color represents a participant.





Figure 2. (continued)

The divergence onset time (DOT, see Methods) of different recording sites in the PFC is consistently within the period of 200–300 ms after grating onset, i.e., the time window of VAN in scalp EEG mentioned above.

As proposed above, the different activities between NA and NU are regarded as awareness-related activities. We found 89 sites with awareness-related iERP activity in the PFC (323 for all recording sites). **Figure 3B** shows the spatial-temporal dynamics of visual awareness-related activities in all recording sites during the 0–650 ms period (for a clearer display, projected to the cortex, see the Supplementary Materials for the complete video). Visual awareness-related activities clearly appeared in the PFC during 200–300 ms, and these early activities were mainly concentrated in the middle lateral prefrontal cortex (LPFC) and then spread to other brain areas of the PFC. We further calculated the start time of awareness-related activity, i.e., DOT, of all recording sites. **Figure 3C** shows the spatial distribution of all recording sites DOT (standardized and convenient for visual display), and it can be found that the DOT of the PFC is not explicitly later than those in posterior brain regions, including the posterior parietal lobe, temporal lobe, occipital lobe, etc., and the earliest region in PFC is still located in the middle lateral prefrontal cortex.

Considering the possibility that the brain region with earlier awareness-related activity might be more important for visual awareness, we tried to detect awareness-related activity in the PFC during the ‘early’ period, i.e., the VAN time window mentioned above. We focused on the sites in PFC with earlier DOTs. Because of the low-visible stimulus used in this experiment, the onset time of VAN may be delayed to 300–350 ms, and we took 350 ms as the cutoff value and divided DOT into early ($n=43$, 48.31%) and late ($n=46$, 51.69%) phases. We focused on the early DOT sites and calculated the normalized average iERP response of the early DOT sites (**Figure 3D**). At the population level, visual awareness-related activities in the PFC were also significant and started at approximately 200–300 ms, which is located in the VAN time window. Importantly, there were sites with DOT < 350 ms from each patient and these sites from different patients were uniformly concentrated in the middle lateral prefrontal cortex (**Figure 3D**, right). That is, the early visual awareness-related activities in the prefrontal cortex are robust and consistent not only at the group level but also at the individual level.

Early visual awareness-related high-gamma activity in the PFC

Although the above iERP results indicate that there are early visual awareness-related neural activities in the prefrontal cortex, it is difficult to explain whether this response is the result of local processing in the PFC or transmission from other brain regions. Therefore, we calculated the LFP response in the high-gamma (HG) band, which is more representative of the local processing, of the prefrontal cortex. **Figure 4A** shows the exemplified high-gamma activity in the prefrontal cortex from each patient at the grand average (left), spectrogram (middle), and single-trial level (right). It is similar to the iERP results (**Figure 3**) except the data are the HG power and there was a vigorous visual evoked power increase under the HA condition compared to the LU condition. However, in the near-threshold condition, while the contrast of the grating was similar, the high-gamma power of the NA and NU trials started to differ significantly after ~ 230 ms (thick black line in panels, $p < 0.01$ corrected by independent t-test; for details, see Methods). This difference is also very explicit and consistent at the single-trial and single-subject levels. The HG activity divergence onset time (DOT) of different recording sites in the PFC was also consistent within the early stage, which was similar to the above iERP result.

We found 31 sites with awareness-related HG activity in the PFC (74 for all recording sites). **Figure 4B** shows the spatial-temporal dynamics of visual awareness-related HG activities in all recording sites during the 0–650 ms period (for a clearer display, projected to the cortex, see the Supplementary Materials for the complete video). Visual awareness-related HG activities also clearly appeared in the PFC during 200–300 ms, while they were rarely seen in lower-level visual areas (occipital and temporal lobe), and these early activities were mainly concentrated in the




Figure 3.

Visual awareness-related iERP activities in the PFC.

A. LFP activity of example sites from each patient. Each panel represents one example site from one patient. In each panel, the left plot shows the grand average of LFP activity in different conditions. The pink line represents data under HA conditions, the red line represents data under NA conditions, the blue line represents data under NU conditions, and the light blue line represents data under LU conditions. The shaded area of the curve represents the SEM. The two black dotted lines at 0 ms and 650 ms represent the time when the grating and fixation point change color (the appearance of saccade targets), and the black thick solid line area represents the period when the LFP amplitude is significantly different under NA and NU conditions ($p < 0.01$ corrected, independent sample t-test). The right figure in each panel shows the single-trial data in the NA (upper) and NU (lower) conditions. The color represents the voltage.

B. The spatial-temporal distribution of awareness-related ERP activities after the appearance of the grating. Each brain image showed a significant ($p < 0.01$ corrected, see Methods) difference in local field potential in NA and NU trials at all visual awareness-related sites at a specific time point (lower left corner) after the grating appeared. The color represents the standardized voltage difference (see Methods).

C. Spatial distribution of divergence onset time. The color represents the normalized DOT value.

D. Population results of early phase ERP response in the prefrontal cortex. Different lines represent different conditions, as shown in Panel A. The right table shows the number of recording sites for different patients. The lower right is the location of these sites. The dots with different colors represent different subjects, as shown in [Figure 2D](#).




Figure 3. (continued)




Figure 4.

Visual awareness-related high-gamma activities in the PFC.

A. High-gamma activity of example sites from each patient. Each panel represents one example site from one patient. In each panel, the left figure shows the grand average of high-gamma power in different conditions. The pink line represents data under HA conditions, the red line represents data under NA conditions, the blue line represents data under NU conditions, and the light blue line represents data under LU conditions. The shaded area of the curve represents the SEM. The two black dotted lines at 0 ms and 650 ms represent the time when the grating and fixation point change color (the appearance of saccade targets), and the black thick solid line area represents the period when the high-gamma power is significantly different under NA and NU conditions ($p < 0.01$ corrected, independent sample t-test). The right figure in each panel shows the single-trial data in the NA (upper) and NU (lower) conditions. The color represents the normalized power.

B. The spatial-temporal distribution of awareness-related high-gamma activities after the appearance of the grating. Each brain image showed a significant ($p < 0.01$ corrected, see Methods) difference in high-gamma activity in NA and NU trials at all visual awareness-related sites at a specific time point (lower left corner) after the grating appeared. The color represents the standardized power difference.

C. Spatial distribution of divergence onset time for high-gamma activity. The color represents the normalized DOT value.

D. Population results of early phase high-gamma response in the prefrontal cortex. Different lines represent different conditions, as shown in Panel A. The middle panel shows the spectrogram in NA (upper) and NU (lower) trials. The right table shows the number of recording sites for different patients. The lower right is the location of these sites. The dots with different colors represent different subjects, as shown in [Figure 2D](#).




Figure 4. (continued)

middle lateral prefrontal cortex (LPFC) and then spread to other brain areas of the PFC. We further calculated the start time of awareness-related HG activity, i.e., DOT, of all recording sites. **Figure 4C** [↗](#) shows the spatial distribution of all recording sites DOT (standardized and convenient for visual display), and it can be found that the DOT of the PFC is not explicitly later than those in posterior brain regions, and the earliest region is still located in the middle lateral prefrontal cortex.

We continue to take 350 ms as the cutoff value and divide the HG DOT into early ($n=18$, 58.06%) and late (13, 41.94%). We focused on the early DOT sites and calculated the normalized mean response (**Figure 4D** [↗](#)) and spectrum of the early DOT sites. At the population level, early visual awareness-related HG activities of the prefrontal sites still exist, and the start time is approximately 200 ms, which is within the VAN time window. Importantly, there are sites with HG DOT<350 ms for each patient except one, and these sites are also uniformly concentrated in the middle lateral prefrontal cortex. In other words, this effect is significant and consistent at both the group and individual levels. It is worth noting that in the PFC recording sites, the early sites with a high-gamma visual response (7.35%) were explicitly less than those with ERP visual response (17.55%), which may indicate that in the process of visual awareness generation, there may also be a variety of other bands activity for information interaction between multiple brain regions.

Decoding awareness state through ERP and HG activity in the PFC

In addition to the univariate analysis above, we also adopted the machine learning decoding method based on multivariate pattern analysis (MVPA) to test the ability of broadband and high-gamma activities in the PFC to predict the awareness state at the single-trial level. We applied a linear discriminant analysis (LDA) to classify aware versus unaware trials based on broadband and high-gamma LFP activities in the PFC.

Figure 5A [↗](#) shows the decoding performance through the ERP activity in the PFC. We found that the ERP activity of the PFC can accurately predict the state of visual awareness at the single-trial level in the delayed period after grating onset, and this result is also relatively consistent at the individual level. Moreover, this decoding accuracy has begun to be much higher than the chance level in 200-300 ms, consistent with our above univariate analysis results. Furthermore, we applied the temporal generalization method to test the predictive ability across time. The results (**Figure 5B** [↗](#)) show that the decoding performance of population ERP is very limited in temporal generalization and cannot be well generalized over a long period. Importantly, this phenomenon is also very consistent at the individual level, which indicates that the representation of visual awareness via broadband activities in the PFC is relatively dynamic rather than static. Moreover, we also applied temporal generalization across the control conditions, i.e., train on NA/NU trials but test on HA/LU trials to test the predictive ability across conditions. The classifier trained on the NA/NU trials can be well generalized to the HA/LU trials, and the generalization across time is also very consistent. These results further indicated the robustness of the above decoding results.

Figure 5C-D [↗](#) shows the results of a similar analysis of HG activity in the prefrontal cortex. **Figure 5C** [↗](#) shows the decoding performance through HG activity in the PFC. We found that the HG activity of the PFC can accurately predict the state of visual awareness at the single-trial level in the delayed period, and this result is also relatively consistent at the individual level. Moreover, this decoding accuracy began to be significantly higher than chance at approximately 200-300 ms, which is also consistent with our above univariate analysis results. Furthermore, we applied the temporal generalization method to the HG activity. Although the group results still show that the decoding performance of HG activity is limited in time generalization, the generalization results are not consistent at the individual level, which may be attributed to the variability of electrode location in different patients. This may suggest that the representation of visual awareness-related




Figure 5.

Decoding the awareness state through broadband and HG activities.

A. Decoding the awareness state through broadband LFP activity. The population result (left) is the average result of six individual subjects (right) after decoding analysis.

B. Time generalization of broadband activity decoding. The left panel represents generalization across time. The right panel represents generalization across different conditions, which means training on NA&NU data and testing on HA&LU data. In each panel, the population result (left) is the average result of six subjects (right) after decoding analysis.

C. The awareness state is decoded by HG activity. Same as A, except that the HG magnitude is used for decoding.

D. Time generalization of HG activity decoding. Same as B, except that the HG magnitude is used for decoding.

information by population neurons at different regions in the PFC is inconsistent, i.e., some are relatively stable, and others are dynamic. Similarly, we applied temporal generalization across the HA/LU conditions, and the result was also very consistent.

Dynamic functional connectivity changes in the PFC associated with visual awareness

We further explored the dynamic functional connectivity between the prefrontal cortex and other brain regions in the generation of visual awareness. First, we calculated the time-across phase-locking value (PLV) ²³ in NA and NU conditions at the sensor level (baseline removed, see Methods for details). **Figure 6A-B** shows the example results of two exemplified patients (see SM for other patients). We found that in the low-frequency band (1-8 Hz), overall, there was a significant difference in the PLV between the NA and NU trials. In the first 150 ms after the appearance of the grating, there was no significant difference in the PLV between these two conditions. However, in the period of approximately 200-300 ms, the PLV between many recoding sites in the NA trials began to increase explicitly and lasted until after 600 ms. In contrast, there was no such phenomenon in the NU trials. Importantly, this phenomenon is strikingly explicit and consistent in each patient.

Furthermore, we grouped the recording sites by brain regions according to the DKT brain atlas ²⁴ and averaged the PLV between different sites and different subjects according to brain regions (**Figure 6C**). The functional connectivity between the prefrontal cortex and subcortical structures, such as the thalamus, and the posterior cortex, such as the posterior parietal cortex, was significantly enhanced in the NA trials compared with the PLV in NU trials. In addition, the functional connectivity between the brain regions within the PFC was also significantly enhanced. For clearer visualization, we show the dynamics of PLV in the NA and NU trials on a topographic map (**Figure 6D**).

Interestingly, we found that at the single-subject level, the sites in the prefrontal cortex that showed early awareness-related PLV activity coincided with the above sites with early awareness-related HG activity. **Figure 6E** shows the averaged PLV between these sites ($n=18$) with early visual awareness-related HG activity in the PFC and other sites in the NA and NU trials. The averaged PLV also showed significant visual awareness-related activities. Moreover, at the population level, the start time of awareness-related PLV activity at these sites also started at approximately 200 ms, which is very similar to that of HG activity.

The different neural activity between NA and NU trials is not caused by the grating contrast

Since there may be a slight difference in the grating contrast under the NA and NU trials, the aforementioned different activity between the NA and NU conditions may be caused by the different contrast of grating. To rule out this possibility, we compared the distribution of grating contrast in the NA and NU conditions for each patient (**Figure 7A**). Overall, we find that the distributions of grating contrasts largely overlap with each other (red and blue curves). The Kolmogorov–Smirnov test resulted in no significant difference ($p>0.05$) between the two contrast distributions in all patients except patient 2. Although intuitively the trials with higher contrast contributed more to the NA condition and the trials with lower contrast contributed more to the NU condition, the contrast difference between the NA and NU conditions were not significantly different perhaps because the grating contrast was designed to be near the perceptual threshold (the ‘aware’ percent is 50%) in most trials. Moreover, for patient 2, the difference between the mean grating contrast in the NA and NU trials was also subtle (0.15%), much less than the difference between NA and HA (17.35%) or NU and LU (0.84%). However, it is clear in the results above that the different activities between NA and NU were much stronger than those between the same awareness states, i.e., NA vs. HA or NU vs. LU. Therefore, the difference in neural activity between aware and unaware trials cannot be explained by the contrast differences in the grating.




Figure 6.

Functional connectivity analysis results.

A-B. Phase-locking value (PLV) changes at the sensor level (NxN) in two example patients. From left to right, the changes in PLV between each recording site at the five time points of 0/150/300/450/600 ms after the appearance of the grating in the near-threshold aware (upper) and near-threshold unaware (lower) trials are displayed (baseline removed, see the Methods). The color represents PLV.

C. The population results of functional connectivity analysis averaged according to brain regions. From left to right, PLV between different brain regions at the five time points of 0/150/300/450/600 ms after the appearance of the grating in the near-threshold aware (upper) and near-threshold unaware (lower) trials (baseline removed, see the Methods). The color represents the PLV.

D. For the same data as C, only the strongest 15% PLV is displayed on the brain template for clear visualization.

E. PLV under different conditions at early-phase HG awareness-related sites in the PFC. The red line represents the average PLV in the NA trials, and the blue line represents the average PLV in the NU trials. The faded line represents the PLV of each site.




Figure 6. (continued)




Figure 6. (continued)

Moreover, we generalized the classifier trained under the NA/NU condition to the NU/LU condition. If the decoding result under the NA/NU condition is based on the difference in the grating contrast rather than on the awareness state, then this classifier should also be able to generalize across the NU/LU condition because the difference in the grating contrast between the NU/LU conditions is greater than that under the NA/NU condition, while the awareness state remains the same (both being unaware of the grating). **Figure 7B** shows the results of generalization decoding results of ERP (left) and HG (right) data. The classifier trained between NA/NU cannot be well generalized to NU/LU conditions for broadband or HG data. Therefore, the above decoding results also cannot be attributed to the difference in grating contrast.

Discussion

We employed a novel visual awareness task and found that, first, the saccadic reaction time of all subjects was shorter when they were aware of grating than when they were not aware. Second, there are ERP and HG activities related to visual awareness in the prefrontal LFP, and these activities begin at the early stage (200-300 ms). Then, we found that the awareness state can be reliably decoded from the broadband and HG activities of the PFC at approximately 200-300 ms after stimulus onset. Finally, the functional connectivity between the PFC and other brain regions in the low-frequency band (1-8 Hz) also appeared awareness-related changes, and the initial time and location of this change are consistent with the above HG results. The results above are not caused by the difference in the physical properties of external stimuli. Therefore, we propose that the prefrontal cortex also plays a key role in the early stage of visual awareness generation even if motor effects are excluded.

Visual awareness states affect saccadic reaction time

Many factors affect saccadic reaction time, including the characteristics of external stimuli and the internal state of the brain. The physical properties of external visual stimuli have a direct impact on the performance of saccades; for example, saccadic reaction time decreases with increasing visual stimulus contrast^{25,26}. A widely accepted view is that the greater contrast of visual stimuli will automatically attract more bottom-up attention and speed up information processing in the brain, thereby shortening the saccadic reaction time^{27,28}. Other studies found that endogenous factors, such as top-down attention, can also significantly shorten the saccadic reaction time^{28,29}.

Consciousness, as a high-level cognitive function of the brain, should have some similar effects as other cognitive functions on behavior (for example, saccadic reaction time). With this question in mind, we carefully searched the literature about the relationship between consciousness and behavior; surprisingly, we failed to find any relevant literature. One study briefly reported that the average reaction time (button press) for word semantic classification in the awareness state was approximately 300 ms faster than in the unawareness state, but the statistical test did not reach a significant level³⁰. In this study, we found that the saccadic reaction time of all subjects in the aware trials was significantly smaller than that in the unaware trials (**Figure 2C**). More critically, the grating contrast near the threshold was not significantly different between the aware and unaware trials (**Figure 7A**); thus, such results cannot be caused by the contrast difference in the grating. Here, for the first time, we showed that awareness states significantly affect saccadic reaction time, which supports our previous hypothesis.




Figure 7.

Grating contrast in NA and NU conditions

A. Distribution of grating contrast in the NA and NU conditions for each patient. Each panel represents one patient. The red and blue lines represent the distribution of grating contrast in NA and NU conditions, respectively.

B. Temporal generalization of broadband and HG activity across the NA&NU condition and NU&LU condition. The left panel represents the generalization of ERP activity across NA&NU and NU&LU conditions, which means trained on NA&NU data and tested on NU&LU data. The right panel is the same analysis except for high-gamma activity.

Visual awareness-related activities appeared in the PFC at an early stage

The long debate about the neural mechanism of consciousness focuses on the location, i.e., ‘front’ vs. ‘back’, and the time, i.e., ‘early’ vs. ‘late’, of its origin in the brain ¹⁴. Concerning the dispute of location, our results show that the prefrontal cortex still displays visual awareness-related activities even after eliminating the influence of the confounding variables related to subjective reports such as motion preparation, which indicates that the prefrontal cortex does participate in the information processing of visual awareness. Regarding the time dispute, our results show that at 200-300 ms after stimulus onset, the VAN time window, the prefrontal cortex has activities related to visual awareness, and these activities include not only ERP activities but also HG activities, which are highly related to the firing of local population neurons ⁷. Notably, the recording sites with early ERP and HG visual awareness-related activities are mostly concentrated in the middle lateral prefrontal cortex. This shows that the prefrontal cortex, especially the middle lateral prefrontal cortex, started visual awareness-related activities, including local activities, at a relatively early stage.

Some recent studies utilizing the no-report paradigm supported our results. For example, a recent electrophysiological study of rhesus monkeys found that even if monkeys do not need to report their conscious content under the binocular competition paradigm, there were also visual awareness-related spiking activities in the prefrontal cortex ³¹. In addition, some previous human intracranial studies also reported ERP, HG, and single-neuron activities related to visual perception in the higher visual areas of the temporal lobe, such as the superior temporal gyrus ^{7,32}. However, due to the limitations of the experimental paradigm, these studies cannot completely exclude the impact of subjective reports. Moreover, most of the electrodes in previous studies were located in the temporal lobe, and electrodes in the prefrontal cortex are rare. As our data show that the proportion of sites with earlier ERP and HG responses in the prefrontal cortex is low and relatively concentrated, it is not hard to explain that earlier visual awareness-related activities in the prefrontal cortex were not found in previous iEEG studies, especially in the HG band. Our results, on the one hand, support that the prefrontal cortex may play an important role in the emergence of visual awareness and thus support the ‘front of the brain’ theories of consciousness in the origin brain regions. However, our data also support the prediction of the early emergence of consciousness proposed by the ‘back of the brain’ theories. This means that our result also challenges some predictions of these mainstream theories.

Awareness state can be effectively decoded from PFC activities from an early stage

Compared with traditional univariate analysis, MVPA-based machine learning decoding may be more sensitive to the information contained in brain activities ³³, so it has gradually become a powerful tool for the study of neural correlates of consciousness (NCCs). Therefore, we used linear discriminant analysis (LDA) to decode the awareness states at the single-trial level through broadband and HG activities in the prefrontal cortex. Our decoding results show that whether through broadband or HG activities, the decoding performance began well above the chance level at approximately 200-300 ms. This shows that the prefrontal cortex has begun to encode visual awareness-related information in 200-300 ms, which is consistent with the above univariate analysis results. In addition, a recent electrophysiological study of rhesus monkeys using the no-report paradigm also found that even without the requirement of explicit reporting, awareness-related neurons can still be found in the lateral prefrontal cortex of rhesus monkeys, and conscious content can be accurately decoded by the MVPA method ³¹. Another human fMRI study also found similar results ¹⁷. These results supported our experimental results.

In addition, we also applied the temporal generalization method during decoding. Temporal generalization analysis shows how different brain activity patterns change and transform over time³⁴. Our results show that the representation pattern of neural activity in the prefrontal cortex for visual awareness-related information is dynamic, not static, and this result is very consistent at the individual level and across the control conditions. Such a result may be consistent with the nature of conscious perception. The process of conscious perception (visual awareness) includes a series of processes from phenomenal consciousness ('qualia') to conscious access. Therefore, at the level of large, high-order brain regions, the representation of this complex process in the brain should be continuously transmitted and transformed, that is, dynamically changed.

Evaluate the role of the prefrontal cortex in visual awareness at the brain network level

There is growing evidence showing that conscious experience may be represented at the brain network level^{17,21}. Thanks to the simultaneously recorded LFP data in multiple brain regions with a high signal-to-noise ratio and high spatial-temporal resolution, we can analyze the dynamic functional connectivity changes under the aware and unaware conditions at the millisecond scale. Since the signals in the low-frequency band are more involved in the information interaction between different brain regions and the activities in the high-frequency band are more representative of the local neural activities, it is easy to understand the functional connectivity changes in the low-frequency band between relatively distant brain regions. Some previous studies have found that the phase synchronization of neural activities in beta and gamma bands is related to visual awareness, but because most of these studies are based on fMRI, it is difficult to describe the dynamic changes in such brain networks on a more precise time scale. The only human iEEG study³⁰ reported that the phase synchronization of the beta band in the aware condition also occurred relatively late (>300 ms). Moreover, most of these studies did not rule out the influence of report-related cognitive processing, and these advanced cognitive processing processes are often associated with changes in the functional connectivity of multiple brain regions. To the best of our knowledge, we are the first to report dynamic changes in functional connectivity on the millisecond time scale in the low-frequency band (1-8 Hz) associated with visual awareness, and this change is not caused by reported-related confounding factors, such as motion preparation. Moreover, the early sites of PLV and HG overlapped each other to a large extent. Such converging results may further suggest that the middle lateral prefrontal cortex, the brain regions where these sites are located, plays a crucial role in the emergence of visual awareness.

Our results may contribute to some theoretical concepts related to consciousness

Compared with the unaware trials, the high-frequency activity in the PFC and the functional connectivity between many brain regions in the aware trials increased. These results may confirm some theoretical concepts, such as 'ignition' and conscious access. Global neuronal workspace theory (GNWT) emphasizes ignition and conscious access^{5,8}. Ignition means nonlinear brain activation to small stimulus changes (such as perceptual threshold stimulation). For the near-threshold stimulus used in the current study, while the sensory information in the brain passes through a certain threshold in a specific brain area, such as the lateral prefrontal cortex, the population of neurons discharges (HG activity of local field potential), that is, "ignition". This is consistent with many previous human intracranial studies^{7,32}. However, due to clinical limitations, these studies are largely confined to the temporal lobe. To the best of our knowledge, our study detected 'ignition' in the human PFC for the first time.

Conscious access means that conscious contents can be shared by many other local information processors in the brain ^{5,8}. For example, when you see the grating, you can either report it orally or draw it down. Such a function may require the participation and coordination of many related brain regions or networks, but there is no clear biomarker for how these brain regions coordinate. The explicit increase in the PLV, specifically appearing in the aware trials, in the low-frequency band found in the current study may provide potential biological meaning for conscious access; that is, many brain regions interact with each other through phase synchronization in the low-frequency band to accomplish information sharing of conscious content.

Interestingly, the beginning time of prefrontal visual awareness-related activities found in ERP and HG activities is very consistent with the onset time of dynamic functional connectivity enhancement, both of which start at 200-300 ms. In addition, the dynamic functional connectivity of the sites with early HG visual awareness-related activity is usually the earliest to enhance compared to other sites, and the two have good consistency. And this consistency suggested that conscious access and phenomenal awareness may be closely coupled, occurring initially in the lateral PFC at the early stage, and indicate a potential process of conscious perception: subtle change in the external stimulus or internal brain state - population neuron discharge in specific brain areas (ignition, phenomenal consciousness appeared) - phase synchronization of various brain areas induced by specific brain areas (conscious access). Further research is needed to regulate the activity of the middle lateral prefrontal cortex in the high-gamma band or the phase synchronization in the low-frequency band to explore whether the prefrontal cortex plays a causal role in the process of conscious perception.

Methods

Data acquisition

Six adult patients with drug-resistant epilepsy (6 males, 32.33±4.75 years old, mean±SEM) and 10 healthy subjects (4 males, 6 females, age 27.80 ± 3.92, mean ± SEM) participated in this study. All subjects had normal or corrected-to-normal vision. Electrophysiological signals (LFPs) were obtained from 6 patients. Stereotaxic EEG depth electrodes (SINOVATION MEDICAL TECHNOLOGY CO., LTD. Ltd., Beijing, China) containing 8 to 20 sites were implanted in patients in the Department of Neurosurgery, PLA General Hospital. Each site was 0.8 mm in diameter and 2 mm in length, with 1.5 mm spacing between adjacent sites. A few electrodes were segmented, and the distance between the two segments was 10 mm. Electrode placement was based on clinical requirements only. Recordings were referenced to a site in white matter. The sEEG signal was sampled at a rate of 1 kHz, filtered between 1 and 250 Hz, and notched at 50 Hz (NEURACLE Technology Co., Ltd., Beijing, China). Stimulus-triggered electrical pulses were recorded simultaneously with sEEG signals to precisely synchronize the stimulus with electrophysiological signals.

Ten healthy subjects performed the experiment in a quiet dark room. Stimuli were presented on a 27-inch LED screen (BenQ, refresh rate of 120 Hz, resolution of 1920 x 1080), and eye movement data were recorded by an infrared eye tracker (EyeLink 1000 Desktop; SR Research) with a sampling rate of 1 kHz. Patients performed the experiment in a quiet dark environment. The stimuli were presented on a 24-inch LED screen (Admiral Overseas Corporation, refresh rate of 144 Hz, resolution of 1920 x 1080), and eye position data were obtained by an infrared eye tracker (Jsmz EM2000C, Beijing Jasmine Science & Technology, Co. Ltd), sampling at 1 kHz. The experimental paradigm was presented by MATLAB (The MathWorks) and the toolbox Psychtoolbox-3 (PTB-3; Brainard and Pelli). Patients were regular recipients of standard medication for epilepsy treatment during the experimental period.

Electrode localization

Electrode locations were determined by registering postoperative computed tomography (CT) scans and preoperative T1 MRI using a standardized mutual information method in SPM12³⁵. Cortical reconstruction and parcellation (Desikan-Killiany Atlas) were conducted by FreeSurfer²⁴. To demonstrate the MNI coordinates of recording sites, a nonlinear surface registration was performed using a nonlinear registration in SPM12, aligning to the MNI ICBM152 template.

Experimental task

A trial started when a fixation point ($0.5^\circ \times 0.5^\circ$, white cross) appeared at the center of the screen (radius of eye position check window is 4° , the dotted circle). After the subject fixated on the fixation point for 600 ms, a cue stimulus (Gabor grating, $2 \times 2^\circ$ circle) was presented for 50 ms at a fixed position (7°) on the left (or right) side of the screen for all participants. The side where the grating appears (left or right) is opposite to the hemisphere where the patients' electrodes are implanted (left or right). For example, if the patient's electrodes are implanted on the right side, the grating would appear on the left side of the screen. For patients 1, 5, and 6, the electrodes were implanted in both hemispheres, and the grating is set on the right side. In 70% of the trials, the grating contrast (weber contrast, see¹⁰) was maintained near the subject's perceptual threshold by a 1 up/1 down staircase method, and the step was either 0.39% or 0.78% for each participant. In 10% of the trials, the stimulus contrast was well above the threshold, and in the other 20% of the trials, the stimulus contrast was 0, namely, no stimulus appeared. After another 600 ms delay, the color of the fixation point turned red or green, and two saccade targets ($1 \times 1^\circ$, white square) appeared at fixed positions (10°) on the left and right sides of the screen. If the grating was seen, the green fixation point was required to make a saccade to the right target, while the red fixation point was required to make a saccade to the left target. If the grating was not seen, the rule of saccadic direction was inverted. Gratings with high contrast (well above the perceptual threshold) and zero contrast (grating absent) served as control conditions to evaluate the understanding and performance of the task. Before starting to collect data, subjects accepted 1-2 training sessions, and the contrast perceptual threshold was determined for each subject. In the formal experiments, each session consisted of 180 trials, and the intertrial interval (ITI) was 800 ms.

Data analysis

Each healthy subject completed 2 sessions, and each patient completed 5-7 sessions. Each session contained 180 trials in total. We excluded trials in which subjects broke fixation during the fixation period (eye position exceeded $4 \times 4^\circ$ check window by more than 40 ms) or the saccadic latency exceeded max response duration (2000 ms for most patients, 5000 ms for patient 6 according to the patient's performance) after the target appeared. For all subjects, more than 85% of the trials were available for further analysis.

In each session, according to the subjects' reports of being aware and unaware of the existence of grating, we divided the trials into awareness and unawareness states. According to the percentage of subjects reporting 'awareness', we further divided grating contrast into 3 levels: high contrast (aware percentage $> 75\%$), near-threshold contrast ($25\% \leq \text{aware percentage} \leq 75\%$), and low contrast (aware ratio $< 25\%$). Due to the few trials of low-contrast-aware and high-contrast-unaware trials, we next analyzed only trials under four conditions of awareness: low-contrast-unaware (LU), near-threshold-unaware (NU), near-threshold-aware (NA) and high contrast-awareness (HA). In total, under LU/NU/NA/HA conditions, there were 243/353/343/160 trials (22.1/32.1/31.2/15.6%, averaged across patients). Since we were more concerned with the differences in subjects' behavior and electrophysiological responses during the physical stimulus, we focused on the

results under NA and NU conditions in the subsequent analysis. The results under LU and HA conditions were classified as the control group and were only used to verify and check the results during calculation.

Behavioral data analysis

Psychophysical curve fitting uses the following formula:

$$AP = a + \frac{b}{1 + \exp(-d * (SC - c))}$$

AP, ratio of reported awareness; SC, stimulus contrast; a-d, individual fitting parameters;

For population psychometric curve fitting, adjacent contrast levels were averaged, resulting in 8 contrast levels in each session.

The saccadic latency was calculated according to a custom program. Eye position and saccade velocity were smoothed with 10 ms sliding windows, and each step was 1 ms. Saccade onset time was calculated as the time when the saccade velocity first exceeded 30°/s and lasted more than 20 ms. Saccadic latency was defined as the time interval between the onset of the saccadic target and the onset of the saccade.

LFP data analysis

Data preprocessing

Data preprocessing was performed by the software package EEGLAB ³⁶ in MATLAB. Data were filtered between 1-250 Hz and notch-filtered at 50, 100, 150, 200, and 250 Hz with an FIR filter. Epochs were extracted from -490 ms before grating onset to 1299 ms after grating onset. Bad recording sites were discarded for analysis based on visual inspection and power spectral density (PSD) analysis. each recording site was applied a bipolar reference, i.e., rereferenced to its direct neighbor.

Spectral analyses

Time-frequency decomposition in high-gamma bands (60-150 Hz) was achieved by Morlet wavelets in the Brainstorm toolbox ³⁷. Before computation, the mean ERP in each condition was removed. For visualization, we used a prestimulus baseline correction over -200 to 0 ms, and the results in **Figure 4** show the HG activity increases or decreases relative to this baseline. To avoid windowing artifacts, we only reported results of -200 to 800 ms.

Quantitative definition of visual awareness-related activity in recording sites

We first identified visual awareness-related sites. For all recording sites, we compared the broadband amplitude (for ERP analysis) or high-gamma magnitude (for HG analysis) of LFP between the NA and NU conditions within 650 ms after grating onset by statistical tests for each time point. We defined the site where there was a significant difference ($p < 0.01$ FDR corrected for time points and channels, independent t-test) in the LFP between NA and NU conditions and persisted for more than 20 ms as an awareness-related site. We also defined the start time of divergence as the divergence onset time (DOT) and defined the mean of the LFP amplitude difference in the first divergence period as the divergence amplitude (DA).

Normalized DOT:

$$T_d = T_{\text{target}} - \text{DOT}$$

$$\text{normDOT} = \frac{T_d - T_{d\min}}{T_{d\max} - T_{d\min}}$$

DOT, divergence onset time; T_{target} , time from target onset to grating onset, 650 ms; T_d , time from DOT to T_{target} ; $T_{d\min}$, the minimum value of T_d in all awareness-related sites within one patient; $T_{d\max}$, the maximum value of T_d of all awareness-related sites within one patient; normDOT, normalized DOT value, a larger normDOT value means a shorter DOT;

For population analysis of recording sites, we conducted baseline normalization for each site.

Baseline normalization for ERP amplitude:

$$x_{\text{norm}} = \frac{x - \mu}{\mu} * 100$$

x_{norm} , normalized value; x , LFP amplitude; μ , mean LFP amplitude in the baseline period (-200 to 0 ms).

Baseline normalization for HG magnitude:

$$x_{\text{norm}} = \frac{x - \mu}{\sigma} * 100$$

x_{norm} , normalized value; x , high-gamma magnitude or power; μ , mean of high-gamma magnitude or power in the baseline period (-200 to 0 ms); σ , standard error of high-gamma magnitude or power in the baseline period (-200 to 0 ms).

Baseline normalization for spectrogram:

$$x_{\text{norm}} = 10 * \log_{10}(x/\mu)$$

x_{norm} , normalized value; x , power value; μ , mean power in the baseline period (-200 to 0 ms).

Topographic display of ERP and HG data

To visualize the topography of the ERP and HG activities, we used the Brainstorm³⁷ package of MATLAB to map the results on the ICBM152 cortical surface. It is a simple 3D interpolation with Shepard's method (weights decreasing with the square of the distance), similar to what is done when projecting subject sources to an anatomy template, where vertices of the cortical surface that are further than 15 mm from the center of the sEEG contact are ignored.

Decoding Analysis

We implemented a machine learning framework for trial-by-trial classification using broadband amplitude and high-gamma magnitude of LFP activities in the PFC via the MVPA-Light toolbox³⁸. We applied the LDA classification and temporal generalization on each participant

separately, and all trials from each class were undersampled to balance the unequal trial number between different conditions. Then, the trials were randomly assigned into 10 folds, and all trials within each fold were subaveraged across every 4 trials to increase the signal-to-noise ratio. Decoding was then followed with a leave-one-out cross-validation procedure on the subaveraged trials. This procedure was repeated 10 times so that each fold in the dataset was used exactly once for testing, and at least once for training, but never at the same time. And the fold assignment was repeated 5 times, which resulted in total 50 iterations for the decoding analysis of each patient.

Functional Connectivity Analysis

The time-across phase-locking value (PLV) was calculated in the *brainstorm* toolbox. Time-frequency transformation was conducted via the Hilbert transform method. The time-across PLV measures if the phase difference between two signals is consistent across trials with respect to the event that defines the trials (i.e., the output is a mean across trials per time point)³⁹. PLV values range between 0 (for totally random) and 1 (for perfect phase locking). For clear demonstration here, the mean PLV in the baseline period (-200 to 0 ms) of each pair of recording sites was subtracted. The group analysis of PLV was performed according to^{40,41}. The topographic display of the connectivity was conducted by the *BrainNet Viewer* toolbox⁴².

Data and software availability

Electrophysiological data were analyzed using MATLAB in conjunction with the toolboxes mentioned above. Data and customized MATLAB analysis scripts are available upon reasonable request from Ming-sha Zhang (mingsha.zhang@bnu.edu.cn).

Acknowledgements

We thank the participants for volunteering to take part in the study. This study was funded by STI2030-Major Projects 2021ZD0204300 and National Natural Science Foundation of China (32061143004, 32030045 and 31871078).

Supplementary Information

SI Figures and Videos

Video S1. The spatial-temporal dynamics of visual awareness-related activities (event-related potential (ERP)) in all recording sites during the 180-650 ms period (for a clearer display, projected to the cortex). The curves in the upper left panel represent the significant ($p < 0.01$ corrected, see Methods) difference (normalized) in averaged amplitude of local field potential in NA and NU trials at all visual awareness-related ERP sites. The brain image showed the difference projected on the surface at a specific time point after the grating appeared (top, left and right views were showed at upper right, lower left and lower right panels, respectively). The color represents the normalized voltage difference (see Methods).

Video S2. The spatial-temporal dynamics of visual awareness-related activities (high-gamma (HG) activity) in all recording sites during the 180-650 ms period (for a clearer display, projected to the cortex). The same as Video S1 but for high-gamma activity. The curves in the upper left panel represent the significant ($p < 0.01$ corrected, see Methods) difference (normalized) in averaged magnitude of high-gamma activity in NA and NU trials at all visual awareness-related HG sites. The brain image showed the difference projected on the surface at a specific time point after the grating appeared (top, left and right views were showed at upper right, lower left and lower right panels, respectively). The color represents the normalized magnitude difference (see Methods).




Figure S1.

Functional connectivity analysis results of Patient 3-6.

A-D. Phase-Locking Value (PLV) changes at sensor level (NxN) in four patients P3-P6. From left to right, the changes of PLV between each recording site in the five time points of 0/150/300/450/600ms after the appearance of the grating in the near-threshold aware (upper) and near-threshold unaware (lower) trials are displayed respectively (baseline removed, see the Method). The color represents PLV.




Figure S1. (continued)

References

1. Koch C., Massimini M., Boly M., Tononi G (2016) **Neural correlates of consciousness: progress and problems** *Nat Rev Neurosci* **17**:307–321 <https://doi.org/10.1038/nrn.2016.22>
2. Mashour G.A (2018) **The controversial correlates of consciousness** *Science* **360**:493–494 <https://doi.org/10.1126/science.aat5616>
3. Dehaene S., Lau H., Kouider S (2017) **What is consciousness, and could machines have it?** *Science* **358**:486–492 <https://doi.org/10.1126/science.aan8871>
4. Kim C.Y., Blake R (2005) **Psychophysical magic: rendering the visible 'invisible'** *Trends Cogn Sci* **9**:381–388 <https://doi.org/10.1016/j.tics.2005.06.012>
5. Dehaene S., Changeux J.P (2011) **Experimental and theoretical approaches to conscious processing** *Neuron* **70**:200–227 <https://doi.org/10.1016/j.neuron.2011.03.018>
6. Dehaene S., Naccache L., Cohen L., Bihan D.L., Mangin J.F., Poline J.B., Riviere D (2001) **Cerebral mechanisms of word masking and unconscious repetition priming** *Nat Neurosci* **4**:752–758 <https://doi.org/10.1038/89551>
7. Fisch L., Privman E., Ramot M., Harel M., Nir Y., Kipervasser S., Andelman F., Neufeld M.Y., Kramer U., Fried I., Malach R (2009) **Neural "ignition": enhanced activation linked to perceptual awareness in human ventral stream visual cortex** *Neuron* **64**:562–574 <https://doi.org/10.1016/j.neuron.2009.11.001>
8. Mashour G.A., Roelfsema P., Changeux J.P., Dehaene S (2020) **Conscious Processing and the Global Neuronal Workspace Hypothesis** *Neuron* **105**:776–798 <https://doi.org/10.1016/j.neuron.2020.01.026>
9. Brown R., Lau H., LeDoux J.E (2019) **Understanding the Higher-Order Approach to Consciousness** *Trends Cogn Sci* **23**:754–768 <https://doi.org/10.1016/j.tics.2019.06.009>
10. van Vugt B., Dagnino B., Vartak D., Safaai H., Panzeri S., Dehaene S., Roelfsema P.R. (2018) **The threshold for conscious report: Signal loss and response bias in visual and frontal cortex** *Science* **360**:537–542
11. Koch C (2018) **What Is Consciousness?** *Nature* **557**:S8–S12 <https://doi.org/10.1038/d41586-018-05097-x>
12. Lamme V.A.F (2018) **Challenges for theories of consciousness: seeing or knowing, the missing ingredient and how to deal with panpsychism** *Philos Trans R Soc Lond B Biol Sci* **373** <https://doi.org/10.1098/rstb.2017.0344>
13. Dembski C., Koch C., Pitts M (2021) **Perceptual awareness negativity: a physiological correlate of sensory consciousness** *Trends in Cognitive Sciences* **25**:660–670 <https://doi.org/10.1016/j.tics.2021.05.009>
14. Seth A.K., Bayne T (2022) **Theories of consciousness** *Nat Rev Neurosci* **23**:439–452 <https://doi.org/10.1038/s41583-022-00587-4>

15. Frassle S., Sommer J., Jansen A., Naber M., Einhauser W (2014) **Binocular rivalry: frontal activity relates to introspection and action but not to perception** *J Neurosci* **34**:1738–1747 <https://doi.org/10.1523/JNEUROSCI.4403-13.2014>
16. Tsuchiya N., Wilke M., Frassle S., Lamme V.A.F (2015) **No-Report Paradigms: Extracting the True Neural Correlates of Consciousness** *Trends Cogn Sci* **19**:757–770 <https://doi.org/10.1016/j.tics.2015.10.002>
17. Kronemer S.I., Aksen M., Ding J.Z., Ryu J.H., Xin Q.L., Ding Z.X., Prince J.S., Kwon H., Khalaf A., Forman S., et al. (2022) **Human visual consciousness involves large scale cortical and subcortical networks independent of task report and eye movement activity** *Nat Commun* **13** <https://doi.org/10.1038/s41467-022-35117-4>
18. Northoff G., Lamme V (2020) **Neural signs and mechanisms of consciousness: Is there a potential convergence of theories of consciousness in sight?** *Neurosci Biobehav Rev* **118**:568–587 <https://doi.org/10.1016/j.neubiorev.2020.07.019>
19. Koivisto M., Revonsuo A (2010) **Event-related brain potential correlates of visual awareness** *Neurosci Biobehav Rev* **34**:922–934 <https://doi.org/10.1016/j.neubiorev.2009.12.002>
20. Koivisto M., Lahteenmaki M., Sorensen T.A., Vangkilde S., Overgaard M., Revonsuo A (2008) **The earliest electrophysiological correlate of visual awareness?** *Brain Cogn* **66**:91–103 <https://doi.org/10.1016/j.bandc.2007.05.010>
21. Huang Z., Mashour G.A., Hudetz A.G (2023) **Functional geometry of the cortex encodes dimensions of consciousness** *Nat Commun* **14** <https://doi.org/10.1038/s41467-022-35764>
22. Merten K., Nieder A (2012) **Active encoding of decisions about stimulus absence in primate prefrontal cortex neurons** *P Natl Acad Sci USA* **109**:6289–6294
23. Tass P., Rosenblum M.G., Weule J., Kurths J., Pikovsky A., Volkman J., Schnitzler A., Freund H.J (1998) **Detection of $\mathit{n}:\mathit{m}$ Phase Locking from Noisy Data: Application to Magnetoencephalography** *Physical Review Letters* **81**:3291–3294 <https://doi.org/10.1103/PhysRevLett.81.3291>
24. Fischl B (2012) **FreeSurfer** *Neuroimage* **62**:774–781 <https://doi.org/10.1016/j.neuroimage.2012.01.021>
25. Allik J., Kreegipuu K (1998) **Multiple visual latency** *Psychol. Sci* **9**:135–138 <https://doi.org/10.1111/1467-9280.00025>
26. Deplancke A., Madelain L., Chauvin A., Cardoso-Leite P., Gorea A., Coello Y (2010) **Influence of near threshold visual distractors on perceptual detection and reaching movements** *J Neurophysiol* **104**:2249–2256 <https://doi.org/10.1152/jn.01123.2009>
27. Posner M.I., Petersen S.E (1990) **The Attention System of the Human Brain** *Annual Review of Neuroscience* **13**:25–42 <https://doi.org/10.1146/annurev.ne.13.030190.000325>
28. Corbetta M., Shulman G.L (2002) **Control of goal-directed and stimulus-driven attention in the brain** *Nat Rev Neurosci* **3**:201–215 <https://doi.org/10.1038/nrn755>
29. Posner M.I (1980) **Orienting of Attention** *Q J Exp Psychol* **32**:3–25 <https://doi.org/10.1080/00335558008248231>

30. Gaillard R., Dehaene S., Adam C., Clemenceau S., Hasboun D., Baulac M., Cohen L., Naccache L (2009) **Converging intracranial markers of conscious access** *PLoS Biol* **7** <https://doi.org/10.1371/journal.pbio.1000061>
31. Kapoor V., Dwarakanath A., Safavi S., Werner J., Besserve M., Panagiotaropoulos T.I., Logothetis N.K (2022) **Decoding internally generated transitions of conscious contents in the prefrontal cortex without subjective reports** *Nat Commun* **13** <https://doi.org/10.1038/s41467-022-28897-2>
32. Reber T.P., Faber J., Niediek J., Bostrom J., Elger C.E., Mormann F (2017) **Single-Neuron Correlates of Conscious Perception in the Human Medial Temporal Lobe** *Curr Biol* **27** <https://doi.org/10.1016/j.cub.2017.08.025>
33. Grootswagers T., Wardle S.G., Carlson T.A (2017) **Decoding Dynamic Brain Patterns from Evoked Responses: A Tutorial on Multivariate Pattern Analysis Applied to Time Series Neuroimaging Data** *J Cogn Neurosci* **29**:677–697 https://doi.org/10.1162/jocn_a_01068
34. King J.R., Dehaene S (2014) **Characterizing the dynamics of mental representations: the temporal generalization method** *Trends Cogn Sci* **18**:203–210 <https://doi.org/10.1016/j.tics.2014.01.002>
35. Ashburner J., Friston K. (1997) **Spatial transformation of images.** *spm course notes*
36. Delorme A., Makeig S (2004) **EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics** *Journal of Neuroscience Methods* **134**:9–21
37. Tadel F., Baillet S., Mosher J.C., Pantazis D., Leahy R.M (2011) **Brainstorm: a user-friendly application for MEG/EEG analysis** *Comput Intell Neurosci* **2011** <https://doi.org/10.1155/2011/879716>
38. Guggenmos M., Sterzer P., Cichy R.M (2018) **Multivariate pattern analysis for MEG: A comparison of dissimilarity measures** *Neuroimage* **173**:434–447 <https://doi.org/10.1016/j.neuroimage.2018.02.044>
39. Mercier M.R., Dubarry A.S., Tadel F., Avanzini P., Axmacher N., Cellier D., Vecchio M.D., Hamilton L.S., Hermes D., Kahana M.J., et al. (2022) **Advances in human intracranial electroencephalography research, guidelines and good practices** *Neuroimage* **260** <https://doi.org/10.1016/j.neuroimage.2022.119438>
40. Betzel R.F., Medaglia J.D., Kahn A.E., Soffer J., Schonhaut D.R., Bassett D.S (2019) **Structural, geometric and genetic factors predict interregional brain connectivity patterns probed by electrocorticography** *Nat Biomed Eng* **3**:902–916 <https://doi.org/10.1038/s41551-019-0404-5>
41. Sadaghiani S., Brookes M.J., Baillet S (2022) **Connectomics of human electrophysiology** *Neuroimage* **247** <https://doi.org/10.1016/j.neuroimage.2021.118788>
42. Xia M., Wang J., He Y (2013) **BrainNet Viewer: a network visualization tool for human brain connectomics** *PLoS One* **8** <https://doi.org/10.1371/journal.pone.0068910>

Author information

Zepeng Fang

State Key Laboratory of Cognitive Neuroscience and Learning and IDG/McGovern Institute for Brain Research, Division of Psychology, Beijing Normal University, Beijing 100875, China

Yuanyuan Dang

Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China

Zhipei Ling

Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China

Yongzheng Han

Department of Anesthesiology, Peking University Third Hospital, Beijing, China

Hulin Zhao

Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China

For correspondence: zhaohulin_90@sohu.com

Xin Xu

Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China

For correspondence: xuxinmm@hotmail.com

Mingsha Zhang

State Key Laboratory of Cognitive Neuroscience and Learning and IDG/McGovern Institute for Brain Research, Division of Psychology, Beijing Normal University, Beijing 100875, China

For correspondence: mingsha.zhang@bnu.edu.cn

Editors

Reviewing Editor

Marius Peelen

Radboud University, Netherlands

Senior Editor

Floris de Lange

Donders Institute for Brain, Cognition and Behaviour, Netherlands

Reviewer #1 (Public Review):

This is a clear and rigorous study of intracranial EEG signals in the prefrontal cortex during a visual awareness task. The results are convincing and worthwhile, and strengths include the use of several complementary analysis methods and clear results. The only methodological weakness is the relatively small sample size of only 6 participants compared to other studies in the field. Interpretation weaknesses that can easily be addressed are claims that their task removes the confound of report (it does not), and claims of primacy in showing early prefrontal cortical involvement in visual perception using intracranial EEG (several studies already have shown this). Also the shorter reaction times for perceived vs not perceived stimuli (confident vs not confident responses) has been described many times previously and is not a new result.

Reviewer #2 (Public Review):

The authors attempt to address a long-standing controversy in the study of the neural correlates of visual awareness, namely whether neurons in prefrontal cortex are necessarily involved in conscious perception. Several leading theories of consciousness propose a necessary role for (at least some sub-regions of) PFC in basic perceptual awareness (e.g., global neuronal workspace theory, higher order theories), while several other leading theories posit that much of the previously reported PFC contributions to perceptual awareness may have been confounded by task-based cognition that co-varied between the aware and unaware reports (e.g., recurrent processing theory, integrated information theory). By employing intracranial EEG in human patients and a threshold detection task on low-contrast visual stimuli, the authors assessed the timing and location of neural populations in PFC that are differentially activated by stimuli that are consciously perceived vs. not perceived. Overall, the reported results support the view that certain regions of PFC do contribute to visual awareness, but at time-points earlier than traditionally predicted by GNWT and HOTs.

Major strengths of this paper include the straightforward visual threshold detection task including the careful calibration of the stimuli and the separate set of healthy control subjects used for validation of the behavioral and eye tracking results, the high quality of the neural data in six epilepsy patients, the clear patterns of differential high gamma activity and temporal generalization of decoding for seen versus unseen stimuli, and the authors' interpretation of these results within the larger research literature on this topic. This study appears to have been carefully conducted, the data were analyzed appropriately, and the overall conclusions seem warranted given the main patterns of results.

Weaknesses include the saccadic reaction time results and the potential flaws in the design of the reporting task. This is not a "no report" paradigm, rather, it's a paradigm aimed at balancing the post-perceptual cognitive and motor requirements between the seen and unseen trials. On each trial, subjects/patients either perceived the stimulus or not, and had to briefly maintain this "yes/no" judgment until a fixation cross changed color, and the color change indicated how to respond (saccade to the left or right). Differences in saccadic RTs (measured from the time of the fixation color change to moving the eyes to the left or right response square) were evident between the seen and unseen trials (faster for seen). If the authors' design achieved what they claim on page 3, "the report behaviors were matched between the two awareness states", then shouldn't we expect no differences in saccadic RTs between the aware and unaware conditions? The fact that there were such differences may indicate differences in post-perceptual cognition during the time between the stimulus and the response cue. Alternatively, the RT difference could reflect task-strategies used by subjects/patients to remember the response mapping rules between the perception and the color cue (e.g., if the YES+GREEN=RIGHT and YES+RED=LEFT rules were held in memory, while the NO mappings were inferred secondarily rather than being actively held in memory). This saccadic RT result should be better explained in the context of the goals of this particular reporting-task.

Nevertheless, the current results do help advance our understanding of the contribution of PFC to visual awareness. These results, when situated within the larger context of the rapidly developing literature on this topic (using "no report" paradigms), e.g., the recent studies by Vishne et al. (2023) Cell Reports and the Cogitate consortium (2023) bioRxiv, provide converging evidence that some sub-regions of PFC contribute to visual awareness, but at latencies earlier than originally predicted by proponents of, especially, global neuronal workspace theory.

Reviewer #3 (Public Review):

The authors report a study in which they use intracranial recordings to dissociate subjectively aware and subjectively unaware stimuli, focusing mainly on prefrontal cortex. Although this paper reports some interesting findings (the videos are very nice and informative!) the interpretation of the data is unfortunately problematic for several reasons. I will detail my main comments below. If the authors address these comments well, I believe the paper may provide an interesting contribution to further specifying the neural mechanisms important for conscious access (in line with Gaillard et al., Plos Biology 2009).

The main problem with the interpretation of the data is that the authors have NOT used a so-called "no-report paradigm". The idea of no report paradigms is that subjects passively view a certain stimulus without the instruction to "do something with it", e.g., detect the stimulus, immediately or later in time. Because of the confusion of this term, specifically being related to the "act of reporting", some have argued we should use the term no-cognition paradigm instead (Block, TiCS, 2019, see also Pitts et al., Phil Trans B 2018). The crucial aspect is that, in these types of paradigms, the critical stimulus should be task-irrelevant and thus not be associated with any task (immediately or later). Because in this experiment subjects were instructed to detect the gratings when cued 600 ms later in time, the stimuli are task relevant, they have to be reported about later and therefore trigger all kinds of (known and potentially unknown) cognitive processes at the moment the stimuli are detected in real-time (so stimulus-locked). You could argue that the setup of this delayed response task excludes some very specific report related processes (e.g., the preparation of an eye-movement), which is good, however this is usually not considered the main issue. For example when comparing masked versus unmasked stimuli (Gaillard et al., 2009 Plos Biology), these conditions usually also both contain responses but these response related processes are "averaged out" in the specific contrasts (unmasked > masked). In this paper, RT differences between conditions (that are present in this dataset) are taken care of by using this delayed response in this paper, which is a nice feature for that and is not the case for the above example set-up.

Given the task instructions, and this being merely a delayed-response task, it is to be expected that prefrontal cortex shows stronger activity for subjectively aware versus subjectively unaware stimuli. Unfortunately, given the nature of this task, the novelty of the findings is severely reduced. The authors cannot claim that prefrontal cortex is associated with "visual awareness", or what people have called phenomenal consciousness (this is the goal of using no-cognition paradigms). The only conclusion that can be drawn is that prefrontal cortex activity is associated with accessing sensory input: and hence conscious access. This less novel observation has been shown many times before and there is also little disagreement about this issue between different theories of consciousness (e.g., global workspace theory and local recurrency theories both agree on this).

The best solution at this point seems to rewrite the paper entirely in light of this. My advice would be to state in the introduction that the authors investigate conscious access using iEEG and then not refer too much to no-cognition paradigm or maybe highlight some different strategies about using task-irrelevant stimuli (see Canales-Johnson et al., Plos Biology 2023; Hesse et al., eLife 2020; Hatamimajoumerd et al Curr Bio 2022; Alilovic et al., Plos Biology 2023; Pitts et al., Frontiers 2014; Dwarakanth et al., Neuron 2023 and more). Obviously, the authors should then also not claim that their results solve debates about theories regarding visual awareness (in the "no-cognition" sense, or phenomenal consciousness), for example in relation to the debate about the "front or the back of the brain", because the data do not inform that discussion. Basically, the authors can just discuss their results in detail (related to timing, frequency, synchronization etc) and relate the different signatures that they have observed to conscious access.

I think the authors have to discuss the Gaillard et al PLOS Biology 2009 paper in much more detail. Gaillard et al also report a study related to conscious access contrasting unmasked and masked stimuli using iEEG. In this paper they also report ERP, time frequency and phase synchronization results (and even Granger causality). Because of the similarities in approach, I think it would be important to directly compare the results presented in that paper with results presented here and highlight the commonalities and discrepancies in the Discussion.

In the Gaillard paper they report a figure plotting the percentage of significant frontal electrodes across time (figure 4A) in which it can be seen that significant electrodes emerge after approximately 250 ms in PFC as well. It would be great if the authors could make a similar figure to compare results. In the current paper there are much more frontal electrode contacts than in the Gaillard paper, so that is interesting in itself.

In my opinion, some of the most interesting results are not highlighted: the findings that subjectively unaware stimuli show increased activations in the prefrontal cortex as compared to stimulus absent trials (e.g., Figure 4D). Previous work has shown PFC activations to masked stimuli (e.g., van Gaal et al., J Neuroscience 2008, 2010; Lau and Passingham J Neurosci 2007) as well as PFC activations to subjectively unaware stimuli (e.g., King, Pescetelli, and Dehaene, Neuron 2016) and this is a very nice illustration of that with methods having more detailed spatial precision. Although potentially interesting, I wonder about the objective detection performance of the stimuli in this task. So please report objective detection performance for the patients and the healthy subjects, using signal detection theoretic d' . This gives the reader an idea of how good subjects were in detecting the presence/absence of the gratings. Likely, this reveals far above chance detection performance and in that case I would interpret these findings as "PFC activation to stimuli indicated as subjectively unaware" and not unconscious stimuli. See Stein et al., Plos Biology 2021 for a direct comparison of subjectively and objectively unaware stimuli.

In Figure 7 of the paper the authors want to make the case that the contrast does not differ between subjectively aware stimuli and subjectively unaware stimuli. However so far they've done the majority of their analyses across subjects, and for this analysis the authors only performed within-subject tests, which is not a fair comparison imo. Because several P values are very close to significance I anticipate that a test across subjects will clearly show that the contrast level of the subjectively aware stimuli is higher than of the subjectively unaware stimuli, at the group level. A solution to this would be to subselect trials from one condition (NA) to match the contrast of the other condition (NU), and thereby create two conditions that are matched in contrast levels of the stimuli included. Then do all the analyses on the matched conditions.

Related, Figure 7B is confusing and the results are puzzling. Why is there such a strong below chance decoding on the diagonal? (also even before stimulus onset) Please clarify the goal and approach of this analysis and also discuss/explain better what they mean.

I was somewhat surprised by several statements in the paper and it felt that the authors may not be aware of several intricacies in the field of consciousness. For example a statement like the following "Consciousness, as a high-level cognitive function of the brain, should have some similar effects as other cognitive functions on behavior (for example, saccadic reaction time). With this question in mind, we carefully searched the literature about the relationship between consciousness and behavior; surprisingly, we failed to find any relevant literature." This is rather problematic for at least two reasons. First, not everyone would agree that consciousness is a high-level cognitive function and second there are many papers arguing for a certain relationship between consciousness and behavior (Dehaene and Naccache, 2001 Cognition; van Gaal et al., 2012, Frontiers in Neuroscience; Block 1995, BBS; Lamme, Frontiers in Psychology, 2020; Seth, 2008 and many more). Further, the explanation for the reaction time differences in this specific case is likely related to the fact that subjects' confidence in

that decision is much higher in the aware trials than in the unaware trials, hence the speeded response for the first. This is a phenomenon that is often observed if one explores the "confidence literature". Although the authors have not measured confidence I would not make too much out of this RT difference.

I would be interested in a lateralized analysis, in which the authors compare the PFC responses and connectivity profiles using PLV as a factor of stimulus location (thus comparing electrodes contralateral to the presented stimulus and electrodes ipsilateral to the presented stimulus). If possible this may give interesting insights in the mechanism of global ignition (global broadcasting), supposing that for contralateral electrodes information does not have to cross from one hemisphere to another, whereas for ipsilateral electrodes that is the case (which may take time). Gaillard et al refer to this issue as well in their paper, and this issue is sometimes discussed regarding to Global workspace theory. This would add novelty to the findings of the paper in my opinion.