

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

November 7, 2023 (this version)

Posted to medRxiv

July 27, 2023

Sent for peer review

June 23, 2023

The impact of colectomy and chemotherapy on risk of type 2 diabetes onset in patients with colorectal cancer: Nationwide cohort study in Denmark

Caroline Elisabeth Krag, Maria Saur Svane, Sten Madsbad, Susanne Boel Graversen, Jesper Frank Christensen, Thorkild IA Sørensen, Louise Lang Lehrskov , Tinne Laurberg

Department of Endocrinology, Copenhagen University Hospital Hvidovre, 2650 Hvidovre, Denmark • Department of Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, 2200 Copenhagen, Denmark • Department of Surgical Gastroenterology, Copenhagen University Hospital Hvidovre, 2650 Hvidovre, Denmark • Steno Diabetes Center Aarhus, Aarhus University Hospital, 8200 Aarhus, Denmark • Centre for Physical Activity Research, Copenhagen University Hospital - Rigshospitalet, Copenhagen, 2100 Copenhagen, Denmark • Department of Sports Science and Clinical Biomechanics, Faculty of Health Sciences, University of Southern Denmark, 5000 Odense, Denmark • Digestive Disease Center, Bispebjerg Hospital, 2400 Copenhagen, Denmark • Novo Nordisk Foundation Center for Basic Metabolic Research and Department of Public Health, University of Copenhagen, 2200 Copenhagen, Denmark

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Background

Comorbidity with type 2 diabetes (T2D) results in worsening of cancer-specific and overall prognosis in colorectal cancer (CRC) patients. The treatment of CRC *per se* may be diabetogenic. We assessed the impact of different types of surgical cancer resections and oncological treatment on risk of T2D development in CRC patients.

Method

We developed a population-based cohort study including all Danish CRC patients, who had undergone CRC surgery between 2001-2018. Using nationwide register data, we identified and followed patients from date of surgery and until new-onset of T2D, death or end of follow-up.

Results

In total, 46,373 CRC patients were included and divided into six groups according to type of surgical resection: 10,566 Right-No-Chemo (23%), 4,645 Right-Chemo (10%), 10,151 Left-No-

Chemo (22%), 5,257 Left-Chemo (11%), 9,618 Rectal-No-Chemo (21%) and 6,136 Rectal-Chemo (13%). During 245,466 person-years of follow-up 2,556 patients developed T2D. The incidence rate (IR) of T2D was highest in the Left-Chemo group 11.3 (95%CI: 10.4-12.2) per 1,000 person-years and lowest in the Rectal-No-Chemo group 9.6 (95%CI: 8.8-10.4). Between-group unadjusted hazard ratio (HR) of developing T2D was similar and non-significant. In the adjusted analysis, Rectal-No-Chemo was associated with lower T2D risk (HR 0.86 [95%CI 0.75-0.98]) compared to Right-No-Chemo.

For all six groups, an increased level of BMI resulted in a nearly twofold increased risk of developing T2D

Conclusion

This study suggests postoperative T2D screening should be prioritized in CRC survivors with overweight/obesity regardless of type of colorectal cancer treatment applied.

Funding

The Novo Nordisk Foundation (NNF17SA0031406); TrygFonden (101390; 20045; 125132).

Impact statement

Colon cancer treatment per se can affect type 2 diabetes risk, but bmi at time of surgery holds the strongest association.

eLife assessment

This study presents **valuable** findings on diabetogenic risk from colorectal cancer (CRC) treatment. The authors claim that postoperative screening for type 2 diabetes should be prioritized in CRC survivors with overweight/obesity, irrespective of the oncological treatment received. The evidence supporting the claims is **solid** but requires confirmation in different populations. These results have theoretical or practical implications and will be of interest to endocrinologists, oncologists, general practitioners, gastrointestinal surgeons, and policymakers working on CRC and diabetes.

Introduction

Colorectal cancer (CRC) is the third most common cancer worldwide,¹ and the incidence rate is still increasing.² The same applies to the global incidence rate of type 2 diabetes (T2D). The association between CRC and T2D is well-established, as patients with T2D have approximately 30% increased risk of developing CRC,³ and T2D is known to negatively influence the prognosis of CRC.⁴ Furthermore, some studies even show an increased T2D risk in CRC survivors.^{5, 6} Fortunately, survival after a CRC diagnosis is steadily improving due to early detection, diagnostic precision, and more advanced cancer treatment; and today, more than 70% of non-metastatic CRC

patients are alive after 5 years.⁷ Accordingly, it becomes increasingly important to identify CRC patients at risk of developing long-term health problems like T2D in order to initiate preventive strategies or treatment.⁶ The underlying mechanisms behind the increased T2D risk in CRC survivors are poorly understood. However, shared life-style related risk factors such as obesity, physical inactivity, and unhealthy diet may play a role.^{8–10}

Another explanation may be treatment-induced hyperglycemia. Prior studies showed that treatment with adjuvant chemotherapy can induce hyperglycemia and development of T2D in CRC patients.^{6, 11, 12} Moreover, glucocorticoid, administered to alleviate side effects of chemotherapy, may impair glucose homeostasis and thus increase the risk of T2D.^{13, 14} In addition to the drug-induced metabolic disturbances, a recent study revealed that the type of CRC surgery *per se* is associated with diverging risk of T2D.¹⁵ A previous nationwide Danish study revealed that patients, who had undergone left-sided colon resection for CRC and other colonic illnesses had an increased risk of developing T2D, whereas this risk was unaltered in patients who had undergone right-sided colonic resections compared with a control group.¹⁵ However, this study lacks essential information on the role of potential modifying factors like BMI, treatment with chemotherapy etc. Along the same line, in a population of non-CRC patients, Lee et al. published that patients who had undergone right-sided or transversal colonic resections had a reduced risk of developing T2D compared with a matched non-colectomy control group.¹⁶ These findings raise an intriguing question of whether the risk of developing T2D in CRC survivors may be casually affected by the type of CRC surgery. Possible explanations of this side-specific risk modification include removal of different enteroendocrine cell types resulting in gut changes that is important for insulin secretion and action and for appetite regulation as well as changes in the microbiome, but causality remains to be investigated further.^{17–19}

In the present study, we tested the hypothesis whether different types of surgical cancer resection and oncological treatment have an impact on risk of developing diabetes in CRC survivors. For this purpose, we used a nationwide register to compare the frequency of diabetes development in CRC patients who had undergone right-sided colectomy, left-sided colectomy or rectal resection with and without chemotherapy, when adjusting for potential modifying factors like BMI etc.

Methods and material

Study population

The study was based on information from Danish nationwide health registers. All data are recorded with reference to a civil registration number, which is a unique personal identification number assigned to all Danish residents. This number permits accurate linkage of recorded information at the personal level.²⁰

The study population was identified through the Danish Colorectal Cancer Group (DCCG) database, which contains prospectively collected and validated data on all patients with CRC since 2001, including data on patient's characteristics, diagnostics and treatment.²¹ The degree of patient inclusion in the database is high (>95%) and thus adequately represents the population of Danish CRC patients undergoing anti-cancer resections.²¹

We included all CRC cancer patients undergoing cancer surgery between 1st of May 2001 until 31st of December 2018 (**Figure 1**). Patients with unspecified surgery or total colectomy were excluded as these procedures are not standard treatment for CRC. Finally, individuals with a preoperatively diagnosed T2D or Type 1 diabetes (T1D) diagnosis, as well as those who were diagnosed with T2D within the first 2 weeks postoperatively were excluded, as the last group probably represents patients with preoperatively unknown pre-existing prediabetes or diabetes.




Figure 1

Flowchart of the study

T2D: New-onset of type 2 diabetes

Right: Right-sided colonic resections; Left: Left-sided colonic resections; Rectal: Rectal resections

No-chemo: No chemotherapy; Chemo: Chemotherapy

Exposure

Patients were subdivided into six groups according to the surgical procedure and whether the patients had received chemotherapy or not. The group “right-sided colonic resection” (Right) included patients with cancer in cecum, ascending colon, or the oral part of colon transversum, the group “left-sided colonic resection” (Left) included patients with cancer in the anal part of colon transversum, descending colon, and sigmoid colon, and the group “rectal resection” (Rectal) included patients with rectal cancer. Information regarding chemotherapy was limited to whether the patient had received chemotherapy (Chemo) or not (No-Chemo).

Based on the combination of surgical procedures and chemotherapy, the study population was divided into six groups: Right-No-Chemo, Right-Chemo, Left-No-Chemo, Left-Chemo, Rectal-No-Chemo, Rectal-Chemo.

Radiation therapy is only part of standard treatment for rectal cancer, and therefore this oncological modality was restricted to Rectal patients (Rectal-No-Radiation vs. Rectal-Radiation).

Finally, a sub analysis was made comparing all colonic resected patients (All Colon) with all rectal resected (All Rectal)

Outcome

The primary outcome was first-time diagnosis of T2D after different types of CRC cancer surgery and chemotherapy. Patients were followed from date of surgery until time to T2D diagnosis, emigration, death, or the end of follow up (31st of December 2018), whichever came first. Individuals with T2D were identified from data in the Danish National Patient Register (comprising information on diagnoses during all hospital contacts in Denmark), the National Prescription Registry (comprising information on all prescriptions filled at any pharmacy in Denmark), and the Danish National Health Service Register (comprising information on podiatrist services) as described by Carstensen et al. [22](#). Individuals were classified as having T1D if they had received prescriptions for insulin combined with a diagnosis of type 1 from a medical hospital department. Otherwise, diabetes was classified as type 2.

Statistical analysis

Descriptive data was presented as means with standard deviations (SD) for continuous variables and proportions (n, %) for categorical variables. Crude incidence rates (number of T2D diagnoses by person-time at risk) were calculated for the six groups (type of surgery ± chemo), and characterised by preoperative variables (sex, age, body mass index (BMI), smoking, alcohol status, American Society of Anesthesiologists physical status classification score (ASA), and treatment with radiotherapy in Rectal cancer patients). Unadjusted and adjusted Cox regression analyses were used to calculate hazard ratios (HRs) for developing T2D after different types of surgery with and without chemotherapy (using Right-No-Chemo as a reference group), after radiation therapy among Rectal resected (using Rectal-No-Radiation as reference), and after overall type of surgery (All Colon (ref) vs All Rectal). The adjusted analyses were performed stepwise by first adjusting for year of surgery, age at surgery, sex and BMI (Model 1), then further adjusting ASA, smoking and alcohol consumption (Model 2a), and lastly, further adjusting for chemotherapy (Model 2b) in the analyses on radiation therapy and overall type of surgery. Finally, the impact of BMI on the risk of developing of T2D was investigated within each of the six groups. BMI was categorized into 4 subgroups (18.5-24.9, 25.0-29.9, 30-34.9 and 35-39.9), and underweight (BMI<18.5) or severely obese (BMI ≥ 40) subjects were excluded from these analyses due to low numbers. For each group (type of surgery ± chemotherapy), the HR for developing T2D depending on BMI subgroups was calculated by using Cox regression analysis adjusted for age, sex, year of surgery, and ASA score using normal weight (BMI:18.5-24.9) as the reference group. A sensitivity analyses was performed restricting the inclusion period of surgery patients from 2001-2018 to 2010-2018, as baseline information was complete from 2010.

All analyses were conducted using STATA version 16 (STATA Corp., College Station, Tex., USA). P-value of $P < 0.05$ was considered statistically significant. Results were presented with 95% confidence interval (CI) when relevant.

Results

Baseline characteristics

A total of 46,373 CRC patients without preoperatively recorded diabetes were found eligible for the study (**Figure 1**) and were divided into six groups: 10,566 Right-No-Chemo (23%), 4,645 Right-Chemo (10%), 10,151 Left-No-Chemo (22%), 5,257 Left-Chemo (11%), 9,618 Rectal-No-Chemo (21%) and 6,136 Rectal-Chemo (13%).

Compared with the Right groups, the Left and Rectal groups were younger, more often men, and less often classified as ASA>II (**Table 1**). Those that had received chemotherapy were generally younger, less often classified as ASA>II and as expected more often stage III/IV cancers than No-Chemo.

Incidence rates of diabetes

During the 245,466 person-years of follow up, 2,556 cases of T2D were diagnosed (10.4 per 1000 person-years) (**Table 2**). The incidence rate (IR) of T2D was almost similar between the six groups with the highest IR observed in the Left-No-Chemo group (IR 11.3 (95%CI: 10.4-12.2) per 1,000 person-years), and the lowest IR observed in the Rectal-No-Chemo group (IR 9.6 (95%CI: 8.8-10.4) per 1,000 person-years).

Across all groups, the T2D IR was consistently higher in males than in females, in patients with BMI ≥ 25 compared with BMI of 20-24.9, and in ASA > I compared with ASA score I (**Table 2**).

Risk of developing T2D after different types of colorectal cancer surgery with and without chemotherapy

When compared to the Right-No-Chemo group, the other groups had similar and non-significant unadjusted hazard ratio (HR) of developing T2D (**Table 3**). In the adjusted analysis Rectal-No-Chemo was associated with a lower risk of T2D (HR 0.86 [95%CI 0.75-0.98]) compared to Right-No-Chemo. In the sub analysis comparing All Colon resected with All Rectal resected, the unadjusted and adjusted HR for developing T2D were significant lower for Rectal resected.

Risk of developing T2D after radiation therapy among rectal resected patients

Radiation therapy in rectal resected had no impact on the incidence rate of T2D in neither the Rectal-No-Chemo group nor the Rectal-Chemo group (**Table 1**), and the unadjusted/adjusted HR of developing T2D was non-significant when comparing Rectal-No-Radiation patients with Rectal-Radiation patients (**Table 3**).

The impact of BMI on development of T2D

BMI had a strong association with T2D regardless of surgical resection and treatment with chemotherapy (**Figure 2**). The risk of developing T2D increased near twofold every time the BMI group increased one level, for example in the Right-No-Chemo group: HR 2.13 in BMI 25-30, HR 3.63 in BMI 30-35, HR 8.01 in BMI 35-40, compared with the reference BMI of 18.5-25. A similar

	Right-sided resection (Right) (N=15,211)		Left-sided resection (Left) (N=15,408)		Rectum resection (Rectal) (N=15,754)	
	<i>Right-No-Chemo</i>	<i>Right-Chemo</i>	<i>Left-No-Chemo</i>	<i>Left-Chemo</i>	<i>Rectal-No-Chemo</i>	<i>Rectal-Chemo</i>
All patients, n (%)	10,566 (23%)	4,645 (10%)	10,151 (22%)	5,257 (11%)	9,618 (21%)	6,136 (13%)
Sex, n (%)						
Male	4,338 (41%)	2,126 (46%)	5,450 (54%)	2,834 (54%)	5,743 (60%)	3,745 (61%)
Female	6,228 (59%)	2,519 (54%)	4,701 (46%)	2,423 (46%)	3,875 (40%)	2,391 (39%)
Age, year, mean (SD)	75.2 (10.0)	66.9 (10.1)	72.1 (10.7)	64.8 (10.1)	70.0 (10.7)	64.3 (10.1)
BMI, mean (SD)	25.0 (4.7)	25.4 (4.7)	25.7 (4.6)	25.8 (4.5)	25.6 (4.4)	25.6 (4.4)
Smoking, n (%)						
Never	3,132 (30%)	1,533 (33%)	3,030 (30%)	1,804 (34%)	2,828 (29%)	2,014 (33%)
Current	1,671 (16%)	791 (17%)	1,486 (15%)	821 (16%)	1,679 (17%)	1,305 (21%)
Former	3,376 (32%)	1,465 (32%)	3,415 (34%)	1,700 (32%)	3,421 (36%)	2,021 (33%)
Missing	2,387 (22%)	856 (18%)	2,220 (21%)	932 (18%)	1,690 (18%)	796 (13%)
Alcohol^a, n (%)						
1-14	4,819 (46%)	2,342 (51%)	4,803 (47%)	2,771 (53%)	4,886 (51%)	3,399 (55%)
14-21	478 (5%)	254 (5%)	618 (6%)	333 (6%)	711 (7%)	494 (8%)
>21	433 (4%)	256 (6%)	593 (6%)	323 (6%)	671 (7%)	412 (7%)
None	2,269 (21%)	889 (19%)	1,722 (17%)	872 (17%)	1,446 (15%)	1,006 (16%)
Missing	2,567 (24%)	895 (19%)	2,415 (24%)	958 (18%)	1,904 (20%)	825 (13%)
ASA score^b, n (%)						
Healthy (=I)	1,633 (15%)	1,331 (29%)	2,185 (22%)	1,735 (33%)	2,541 (26%)	2,082 (34%)
Mild (=II)	5,457 (52%)	2,596 (56%)	5,299 (51%)	2,857 (54%)	5,265 (55%)	3,415 (56%)
Sick (=>II)	3,160 (30%)	613 (13%)	2,442 (24%)	548 (11%)	1,653 (17%)	557 (9%)
Missing	316 (3%)	105 (2%)	295 (3%)	117 (2%)	159 (2%)	82 (1%)
Screening detected tumor, n (%)	770 (7%)	299 (6%)	1,077 (11%)	457 (9%)	664 (7%)	394 (6%)
UICC stage^c, n (%)						
I/II	7,002 (66%)	674 (15%)	7,339 (72%)	745 (14%)	5,940 (62%)	420 (7%)
III/IV	3,330 (32%)	3,504 (75%)	2,506 (25%)	3,989 (76%)	2,246 (23%)	2,471 (40%)
Unknown	234 (2%)	467 (10%)	306 (3%)	523 (10%)	1,432 (15%)	3,245 (53%)
Radiation, n (%)	.	.	.	.	1,157 (12%)	2,983 (49%)

Chemo: Chemotherapy
^a Alcohol items per week
^b The American Society of Anesthesiologist score: I: Healthy, but with CRC, III: Mild systemic disease without substantial functional limitations, >II: Severe systemic disease, includes ASA stage III, IV, V, VI
^c UICC stage, Union of International Cancer Control, Stage I: T1 or T2, Stage II: T3 or T4, Stage III: N1 or N2, Stage IV: Disseminated disease at time of diagnosis

Table 1.

Baseline characteristics

	Right-sided resection (Right) (N=15,211)		Left-sided resection (Left) (N=15,408)		Rectum resection (Rectal) (N=15,754)	
	<i>Right-No-Chemo</i>	<i>Right-Chemo</i>	<i>Left-No-Chemo</i>	<i>Left-Chemo</i>	<i>Rectal-No-Chemo</i>	<i>Rectal-Chemo</i>
All patients, n	10,566	4,645	10,151	5,257	9,618	6,136
Development of T2D, n	495	235	609	323	560	334
Person-years	48,039	22,317	53,935	29,301	58,448	33,422
Incidence rates	10.3 (9.4-11.2)	10.5 (9.3-12.0)	11.3 (10.4-12.2)	11.0 (9.9-12.3)	9.6 (8.8-10.4)	10.0 (9.0-11.1)
Sex						
Male	13 (11-15)	14 (12-16)	14 (12-15)	14 (12-16)	11 (10-12)	12 (11-14)
Female	9 (8-10)	8 (7-10)	9 (8-10)	8 (7-10)	8 (7-9)	7 (6-8)
Age						
<50	3 (1-8)	6 (3-11)	7 (5-12)	8 (5-12)	5 (3-8)	6 (4-10)
50-64.9	12 (10-14)	13 (11-16)	11 (9-12)	10 (8-12)	10 (8-12)	10 (9-12)
65-74.9	12 (11-14)	11 (9-14)	12 (11-14)	14 (12-17)	14 (12-17)	11 (10-13)
≥75	9 (8-10)	6 (4-9)	11 (10-13)	8 (6-12)	8 (6-12)	8 (5-11)
BMI subgroups						
<18.5	3 (2-8)	2 (0.2-14)	4 (2-10)	6 (2-20)	5 (2-12)	8 (3-18)
18.5-24.9	6 (5-7)	5 (4-7)	6 (5-8)	4 (3-5)	4 (4-5)	4 (3-6)
25-29.9	13 (11-16)	11 (9-14)	13 (11-15)	12 (10-15)	11 (9-12)	12 (10-14)
30-34.9	23 (18-28)	30 (23-40)	21 (17-25)	24 (19-30)	25 (21-30)	24 (19-30)
35-39.9	51 (38-70)	53 (34-84)	32 (23-45)	34 (23-52)	30 (21-44)	31 (20-48)
>40	55 (31-97)	61 (26-128)	51 (31-84)	80 (47-138)	38 (21-71)	49 (22-109)
Missing	7 (6-9)	9 (6-12)	11 (9-12)	12 (10-16)	9 (8-12)	10 (8-14)
Smoking						
Never	9 (8-11)	11 (9-14)	9 (8-11)	8 (7-10)	8 (7-10)	9 (7-11)
Current	13 (10-15)	9 (7-13)	11 (9-14)	12 (10-16)	10 (8-12)	10 (7-12)
Former	11 (10-13)	12 (10-15)	13 (11-15)	13 (11-15)	11 (9-12)	11 (9-13)
Missing	8 (7-10)	9 (6-12)	11 (10-14)	12 (9-15)	9 (8-11)	11 (8-14)
Alcohol^a						
1-14	11 (9-12)	10 (8-12)	10 (9-12)	10 (9-12)	9 (8-10)	10 (8-11)
14-21	9 (6-14)	7 (3-14)	9 (6-13)	10 (7-16)	8 (6-11)	9 (6-14)
>21	10 (7-16)	16 (11-25)	14 (11-19)	12 (8-18)	11 (9-15)	13 (9-18)
None	12 (10-14)	14 (10-18)	13 (11-16)	12 (9-16)	13 (11-15)	10 (7-13)
Missing	9 (7-10)	10 (7-13)	12 (10-14)	12 (10-16)	10 (9-10)	10 (8-13)
ASA score^b						
Healthy (=I)	7 (6-9)	5 (3-7)	7 (6-9)	7 (5-9)	5 (4-6)	6 (4-7)
Mild (=II)	10 (9-12)	12 (11-15)	12 (11-13)	13 (11-15)	11 (10-12)	13 (11-14)
Sick (=>II)	14 (12-17)	17 (13-24)	17 (12-17)	18 (14-25)	15 (12-18)	12 (8-17)
Missing	6 (3-11)	15 (7-31)	13 (8-20)	11 (5-23)	9 (5-17)	10 (4-25)
Radiation						
Yes	.	.	.	.	11 (9-13)	11 (9-13)
No	.	.	.	.	9 (9-10)	10 (9-12)

Chemo: Chemotherapy
^a Alcohol items per week
^b The American Society of Anesthesiologist score: I: Healthy, but with CRC, III: Mild systemic disease without substantial functional limitations, >II: Severe systemic disease, includes ASA stage III, IV, V, VI

Table 2.

Absolute incidence rates of type 2 diabetes (T2D) per 1000 person-years (95% CI) among colorectal cancer patients treated with different types of colorectal cancer surgery with and without chemotherapy.

	Unadjusted		Model 1		Model 2 ^(a/b)	
	HR (95%CI)	<i>p-value</i>	HR (95%CI)	<i>p-value</i>	HR (95%CI)	<i>p-value</i>
Surgery and chemotherapy						
Right-No-Chemo	ref					
Right-Chemo	1.02 (0.87;1.19)	0.799	0.99 (0.83;1.17)	0.880	1.01 (0.85;1.20) ^a	0.884
Left-No-Chemo	1.10 (0.98;1.24)	0.112	0.92 (0.81;1.05)	0.228	0.94 (0.82;1.07) ^a	0.333
Left-Chemo	1.07 (0.93;1.23)	0.339	0.93 (0.79;1.09)	0.366	0.97 (0.83;1.14) ^a	0.708
Rectal-No-Chemo	0.94 (0.83;1.06)	0.317	0.82 (0.72;0.94)	0.004	0.86 (0.75;0.98) ^a	0.028
Rectal-Chemo	0.96 (0.84;1.11)	0.610	0.84 (0.72;0.99)	0.031	0.89 (0.76;1.03) ^a	0.126
Surgery and radiation therapy						
Rectal-No-Radiation	ref					
Rectal-Radiation	1.07 (0.93;1.24)	0.336	0.98 (0.85;1.13)	0.804	0.97 (0.84;1.12) ^b	0.691
Surgery						
All Colon	ref					
All Rectal	0.90 (0.83;0.98)	0.013	0.87 (0.79;0.95)	0.002	0.89 (0.82;0.98) ^b	0.015
Right: Right-sided colonic resections; Left: Left-sided colonic resections; Rectal: Rectal resections No-Chemo: No chemotherapy; Chemo: Chemotherapy No-radiation: No radiotherapy; Radiation: Radiation therapy All Colon: All right- and left-sided colonic resections with and without chemotherapy All Rectal: All rectal resections with and without chemotherapy, and with and without radiation therapy Model 1: adjusted for year of surgery, age at surgery, sex, BMI Model 2a: adjusted like Model 1 and further adjusted for performance status before surgery assessed by American Society of Anesthesiologist physical scale, smoking and alcohol Model 2b: adjusted like Model 2a and further adjusted for chemotherapy						

Table 3.

Likelihood of developing type 2 diabetes after different types of colorectal cancer surgery with and without oncological treatment, unadjusted and adjusted analysis

pattern was observed within each of the other five groups, although a less pronounced impact of BMI was observed in the Left-No-Chemo group: HR 1.85 in 25-30, 2.93 in 30-35, 4.21 in 35-40 BMI group.

Discussion

This national cohort study demonstrated that rectal cancer resected patients had a slightly lower risk of developing T2D when compared to colon cancer resected patients. Treatment with chemotherapy had no impact on T2D risk in any of the resection groups, and radiation therapy didn't affect T2D risk either.

Within each of the six groups (different types of surgery with and without chemotherapy), BMI was strongly associated to developing T2D, the risk increased almost twofold every time the BMI group increased one level.

Only few studies have examined the risk of developing T2D after different kinds of colonic- and rectal surgical procedures, and in contrast to the present study the following studies were not able to adjust for several confounders incl. preoperative BMI. Jensen et al. found a higher risk of T2D after the left-sided compared with the right-sided colonic resections, but, in contrast to the current study, patients operated for both non-malignant as well as malignant diseases were included.¹⁵ In support of side-specific effect of colonic resections, Wu et al. investigated a population of non-CRC patients and showed that right-sided and transversal colonic resections were associated with a reduced risk of T2D development compared with patients without colectomy.¹⁶ In contrast to both the abovementioned studies the most recent epidemiological study found an increased T2D risk after colectomy compared with small bowel resection in non-cancer patients, but when stratifying by resection type, they found no difference in T2D risk.²³

Obesity is one of the predominant factors in the development of T2D²⁴⁻²⁶, and the linear increase in new onset T2D with increased BMI reported in this study is well described in the literature.²⁷ It is important to emphasize that only pre-operative BMI was included in the adjustments as post-surgery BMI information was not available and accordingly, weight trajectories after surgery could not be included in the analyses. It's still unknown how body weight and body composition changes after different types of CRC resection, and thus it is not clear whether changes in weight in any groups play a role in diabetes development.

Like our findings, Jensen et al. and Wu et al. found a tendency towards a decreased risk of T2D after rectal resection, although these findings were not statistically significant in the previous studies.^{15, 16} Further studies are needed to elucidate the potential mechanism behind this finding.

We found no impact of treatment with chemotherapy on T2D risk, which was a bit surprising. Prior studies indicated that treatment with chemotherapy may induce hyperglycemia, and especially the combined use of chemotherapy and glucocorticoid, administered to alleviate side-effects of chemotherapy, may have synergistic effects on later diabetes risk.²⁸

In the study period of the present study, different types of chemotherapy have been part of the standard treatment of colon and rectum cancer respectively, and knowledge regarding the specific effects of each drug on glucose homeostasis is sparse.²⁹ In most cases CRC patients included in the study, will however have been treated with the chemotherapy 5-Fluorouracil.¹² In a recent study a significant increase in glycemic variability, a precursor of T2D development, was found in non-diabetic early-stage colon cancer patients during 5-Fluorouracil and Capecitabine chemotherapy treatment.³⁰ Moreover, Feng et al. showed a 10% increased risk of developing T2D in CRC patients during and after treatment with adjuvant 5-Fluorouracil. They argued that 5-




Figure 2

Likelihood of developing type 2 diabetes depending on BMI subgroups (using normal weight, BMI:18.5-25 as reference) after different types of colorectal cancer surgery with and without chemotherapy.

$\diamond$ Hazard ratios were adjusted for age, sex, year of surgery and ASA score (performance status before surgery assessed by American Society of Anesthesiologist physical scale).

Flourouracil could induce toxicity in B-cells leading to hyperglycaemia.¹² Similarly, Lee et al revealed an impairment in glucose homeostasis in both breast cancer and CRC patients treated with chemotherapy and glucocorticoid following treatment.¹³ Nonetheless 65% of the newly diagnosed T2D cases remitted 6 months after discontinuation of chemotherapy, suggesting that chemotherapy is a reversible diabetogenic factor.¹³ Thus, in summary, treatment with chemotherapy may primarily induce transient effects on glucose homeostasis in CRC patients. In contrast to the lack of influence of chemotherapy treatment, it was not surprising that radiation therapy had no impact on diabetes risk in our study.

Due to the strong negative prognostic effects of comorbidity with T2D on CRC outcomes, identifying CRC patients at increased risk of developing T2D is of great clinical importance. Our study didn't support a role of cancer surgery with left-sided colonic resection or treatment with chemotherapy in T2D development in CRC survivors. However, in line with existing literature we found BMI to be strongly associated with T2D development.

A core strength of the present study is the use of a nationwide database, allowing inclusion of large numbers of CRC patients treated with different types of CRC resections, providing valid real-life data regarding with close to complete coverage and long-term follow-up and a register-based approach to identify individuals with T2D. We were able to adjust for several confounders such as BMI, alcohol consumption and smoking, which has been a limitation in previous studies.¹⁵¹⁶

The limitation of the present study includes that information on BMI and other risk factors for developing T2D were only available preoperatively, eliminating the possibility of exploring an impact of BMI trajectories after cancer treatment until the T2D diagnosis.³¹ Secondly, information on dose, length of treatment and type of chemotherapy were not available. Thirdly, information on physical activity, diet, medication, and comorbidities, apart from the ASA score, were not available. In addition, our findings are limited by the register-based data collection method. Clinical well-designed mechanistic investigations are warranted to establish whether left-sided CRC resection *per se* drives metabolic alterations leading to an increased risk of T2D.

In summary, we report a slightly increased risk of developing T2D after CRC treatment with colonic resection compared with rectal resection, whereas chemotherapy had no impact on new onset T2D. Within each surgical subgroup, BMI at time of surgery was strongly associated to developing T2D. This study suggests that postoperative diabetes screening should be prioritized in CRC survivors with overweight/obesity regardless of type of colorectal cancer treatment applied.

Data Availability

All data produced in the present study are available upon reasonable request to the authors

Acknowledgements

The authors thank the DCCG for access to data and Danish Cancer Society for support.

Funding

This work was supported by the Danish Diabetes Association and the Danish Diabetes Academy, which is funded by the Novo Nordisk Foundation, grant number NNF17SA0031406.

Ethics approval

According to Danish law and the Committee on Health Research Ethics in the Central Denmark Region, the study required no ethical approval because it was based on secondary use of register data for research purposes. This committee also waived patient consent for the use of register data. The study was approved by the Central Region Denmark (file no. 1-16-02-304-19).

Patient consent for publication

Not applicable

Contributors

Conceptualization, TL and LLL. Methodology, TL, LLL, CK, TIAS, MSS and JFC. Access to and verified the raw data and formal analysis TL and SBG. Resources, TL, LLL. Writing - Original draft CK, TL, LLL, MSS and SM. Writing – Review and editing, all authors contributed.

All authors approved the final version for publication, and LLL had final responsibility for the decision to submit.

Declaration of interests

The authors disclose no conflicts.

Data sharing

Anonymized data will be made available upon reasonable request for academic use and within the limitations of the informed consent. Requests should be made to last author tinlaurb@rm.dk. Due to the Danish law it is not possible to make national health registers available, but on request we would like to collaborate and discuss reasonable access to data.

References

1. Araghi M, Soerjomataram I, Jenkins M, et al. (2019) **Global trends in colorectal cancer mortality: projections to the year 2035** *Int J Cancer* **144**:2992–3000 <https://doi.org/10.1002/ijc.32055>
2. Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F (2017) **Global patterns and trends in colorectal cancer incidence and mortality** *Gut* **66**:683–691 <https://doi.org/10.1136/gutjnl-2015-310912>
3. González N, Prieto I, del Puerto-Nevado L, et al. (2017) **Update on the Relationship between Diabetes and Colorectal Cancer: Epidemiology, Potential Molecular Mechanisms and Therapeutic Implications**
4. Mills KT, Bellows CF, Hoffman AE, Kelly TN, Gagliardi G (2013) **Diabetes mellitus and colorectal cancer prognosis: A meta-analysis** *Dis Colon Rectum* **56**:1304–1319 <https://doi.org/10.1097/DCR.0b013e3182a479f9>
5. Singh S, Earle CC, Bae SJ, et al. (2016) **Incidence of Diabetes in Colorectal Cancer Survivors** *Natl Cancer Inst* **108** <https://doi.org/10.1093/jnci/djv402>
6. Xiao Y, Wang H, Tang Y, et al. (2021) **Increased risk of diabetes in cancer survivors: a pooled analysis of 13 population-based cohort studies** *ESMO Open* **6** <https://doi.org/10.1016/j.esmoop.2021.100218>
7. Colorectal D Nordcan (2022) **5-Year Age-Standardised Relative Survival (%), Males and Females**
8. Nano J, Dhana K, Asllanaj E, et al. (2020) **Trajectories of BMI Before Diagnosis of Type 2 Diabetes: The Rotterdam Study** *Obesity* **28**:1149–1156 <https://doi.org/10.1002/oby.22802>
9. Maddatu J, Anderson-Baucum E, Evans-Molina C (2017) **Smoking and the risk of type 2 diabetes** *Translational Research* **184**:101–107 <https://doi.org/10.1016/j.trsl.2017.02.004>
10. Lega IC, Lipscombe LL. (2020) **Review: Diabetes, Obesity, and Cancer-Pathophysiology and Clinical Implications** *Endocr Rev* **41** <https://doi.org/10.1210/endrev/bnz014>
11. Jo A, Scarton L, O'Neal LTJ, et al. (2021) **New onset of type 2 diabetes as a complication after cancer diagnosis: A systematic review** *Cancer Med* **10**:439–446 <https://doi.org/10.1002/cam4.3666>
12. Feng JP, Yuan XL, Li M, et al. (2013) **Secondary diabetes associated with 5-fluorouracil-based chemotherapy regimens in non-diabetic patients with colorectal cancer: Results from a single-centre cohort study** *Colorectal Disease* **15**:27–33 <https://doi.org/10.1111/j.1463-1318.2012.03097.x>
13. Lee EK, Koo B, Hwangbo Y, et al. (2021) **Incidence and disease course of new-onset diabetes mellitus in breast and colorectal cancer patients undergoing chemotherapy: A prospective multicenter cohort study** *Diabetes Res Clin Pract* **174** <https://doi.org/10.1016/j.diabres.2021.108751>

14. Jeong Y, Han HS, Lee HD, et al. (2016) **A pilot study evaluating steroid-induced diabetes after antiemetic dexamethasone therapy in chemotherapy-treated cancer patients** *Cancer Res Treat* **48**:1429–1437 <https://doi.org/10.4143/crt.2015.464>
15. Jensen AB, Sørensen TI, Pedersen O, Jess T, Brunak S, Allin KH (2018) **Increase in clinically recorded type 2 diabetes after colectomy** *Published online* <https://doi.org/10.7554/eLife.37420.001>
16. Wu CC, Lee CH, Hsu TW, et al. (2021) **Is colectomy associated with the risk of type 2 diabetes in patients without colorectal cancer? A population-based cohort study** *J Clin Med* **10** <https://doi.org/10.3390/jcm10225313>
17. Holst JJ, Gribble F, Horowitz M, Rayner CK (2016) **Roles of the gut in glucose homeostasis** *Diabetes Care* **39**:884–892 <https://doi.org/10.2337/dc16-0351>
18. Lin XH, Jiang JK, Luo JC, et al. (2019) **The long term microbiota and metabolic status in patients with colorectal cancer after curative colon surgery** *PLoS One* **14** <https://doi.org/10.1371/journal.pone.0218436>
19. Hansen C Palnæs, Andreassen JJ, Holst JJ (1997) **The release of gastric inhibitory peptide, glucagon-like peptide-I, and insulin after oral glucose test in colectomized subjects** *Scand J Gastroenterol* **32**:473–477 <https://doi.org/10.3109/00365529709025084>
20. Pedersen CB, Gøtzsche H, Møller JØ, Mortensen PB **Med Bull D**
21. Klein MF, Gögenur I, Ingeholm P, et al. (2020) **Validation of the Danish Colorectal Cancer Group (DCCG.dk) database – on behalf of the Danish Colorectal Cancer Group** *Colorectal Disease. Published online* <https://doi.org/10.1111/codi.15352>
22. Carstensen B, Kristensen JK, Marcussen MM, Borch-Johnsen K (2011) **The national diabetes register** *Scand J Public Health* **39**:58–61 <https://doi.org/10.1177/1403494811404278>
23. Allin KH, Agrawal M, Iversen AT, Antonsen J, Villumsen M, Jess T (2022) **The Risk of Type 2 Diabetes in Patients With Inflammatory Bowel Disease After Bowel Resections: A Nationwide Cohort Study** *Gastro Hep Advances* **1**:777–784 <https://doi.org/10.1016/j.gastha.2022.06.007>
24. Wang Y, Rimm EB, Stampfer MJ, Willett WC, Hu FB (2005) **Comparison of Abdominal Adiposity and Overall Obesity in Predicting Risk of Type 2 Diabetes among Men 1-3**
25. Hu Y, Bhupathiraju SN, de Koning L, Hu FB (2014) **Duration of obesity and overweight and risk of type 2 diabetes among US women** *Obesity* **22**:2267–2273 <https://doi.org/10.1002/oby.20851>
26. Lee DH, Keum NN, Hu FB, et al. (2018) **Comparison of the association of predicted fat mass, body mass index, and other obesity indicators with type 2 diabetes risk: two large prospective studies in US men and women** *Eur J Epidemiol* **33**:1113–1123 <https://doi.org/10.1007/s10654-018-0433-5>
27. Karin A Easson, Jon E, Martin A, et al. (2022) **Body mass index in adolescence, risk of type 2 diabetes and associated complications: A nationwide cohort study of men** *Published online* <https://doi.org/10.1016/j>

28. Hwangbo Y, Kang D, Kang M, et al. (2018) **Incidence of diabetes after cancer development a Korean national cohort study** *JAMA Oncol* **4**:1099–1105 <https://doi.org/10.1001/jamaoncol.2018.1684>
29. Ariaans G, de Jong S, Gietema JA, Lefrandt JD, de Vries EGE, Jalving M (2015) **Cancer-drug induced insulin resistance: Innocent bystander or unusual suspect** *Cancer Treat Rev* **41**:376–384 <https://doi.org/10.1016/j.ctrv.2015.02.007>
30. Mandolfo NR, Berger AM, Struwe LA, et al. (2022) **Glycemic Variability in Patients With Stage II–III Colon Cancer Treated With Surgery and Adjuvant Chemotherapy** *Oncol Nurs Forum* **49**:571–584 <https://doi.org/10.1188/22.ONF.571-584>
31. Winkels RM, Snetselaar T, Adriaans A, et al. (2016) **Changes in body weight in patients with colorectal cancer treated with surgery and adjuvant chemotherapy: An observational study** *Cancer Treat Res Commun* **9**:111–115 <https://doi.org/10.1016/j.ctarc.2016.09.002>

Article and author information

Caroline Elisabeth Krag

Department of Endocrinology, Copenhagen University Hospital Hvidovre, 2650 Hvidovre, Denmark, Department of Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, 2200 Copenhagen, Denmark

Maria Saur Svane

Department of Endocrinology, Copenhagen University Hospital Hvidovre, 2650 Hvidovre, Denmark, Department of Surgical Gastroenterology, Copenhagen University Hospital Hvidovre, 2650 Hvidovre, Denmark

Sten Madsbad

Department of Endocrinology, Copenhagen University Hospital Hvidovre, 2650 Hvidovre, Denmark

Susanne Boel Graversen

Steno Diabetes Center Aarhus, Aarhus University Hospital, 8200 Aarhus, Denmark

Jesper Frank Christensen

Centre for Physical Activity Research, Copenhagen University Hospital - Rigshospitalet, Copenhagen, 2100 Copenhagen, Denmark, Department of Sports Science and Clinical Biomechanics, Faculty of Health Sciences, University of Southern Denmark, 5000 Odense, Denmark, Digestive Disease Center, Bispebjerg Hospital, 2400 Copenhagen, Denmark

Thorkild IA Sørensen

Novo Nordisk Foundation Center for Basic Metabolic Research and Department of Public Health, University of Copenhagen, 2200 Copenhagen, Denmark

Louise Lang Lehrskov

Centre for Physical Activity Research, Copenhagen University Hospital - Rigshospitalet, Copenhagen, 2100 Copenhagen, Denmark

For correspondence: louise.lang.lehrskov.01@regionh.dk

Tinne Laurberg

Steno Diabetes Center Aarhus, Aarhus University Hospital, 8200 Aarhus, Denmark

Copyright

© 2023, Krag et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Rui Medeiros

Instituto Português de Oncologia-Porto, Portugal

Senior Editor

Eduardo Franco

McGill University, Canada

Reviewer #1 (Public Review):

Summary:

In this study, the authors set out to determine whether colorectal cancer surgery site (right, left, rectal) and chemotherapy impact the subsequent risk of developing T2DM in the Danish national health register.

Strengths:

- The research question is conceptually interesting
- The Danish national health register is a comprehensive health database
- The data analysis was thorough and appropriate
- The findings are interesting, and a little surprising that there was no impact of chemotherapy on the development of T2DM

Weaknesses:

- This is not a weakness as such, but in the discussion, I would consider adding some brief comment on the international generalizability of the findings - e.g. demographic make up of the Danish population health register and background rates of DM and obesity in this population with CRC compared to countries on other continents.
- A little more information would be helpful regarding how T2DM was diagnosed in the registry. If someone did develop transient hyperglycemia requiring DM medications during chemotherapy, would the investigators have been able to identify these people? Would they have been classified as T2DM based on filling a prescription for DM meds for a period of time? Also, did the authors have information regarding time to development of T2DM after surgery?
- In the adjusted Models, the authors did not adjust for cancer stage, even though cancer stage appears to be very different between the chemo and no chemo groups. It would be interesting to know if it affects the results if the model adjusted for cancer stage
- It would be worthwhile to report if mortality rates were different between the groups during follow up, and if the authors investigated whether perhaps differences in mortality rates led to specific groups living longer, and therefore having more time to develop DM

Overall, the authors achieved their aims, and the conclusions are supported by their results as reported.

The results are unlikely to significantly change patient treatment or T2DM screening in this population. With some additional information, as described above, the results would be of interest to the community.

Reviewer #2 (Public Review):

Summary: The study showed the impact of cancer treatment on new onset of diabetes among patients with colorectal cancer using the national database. Findings reported that individuals with rectal cancer without chemotherapy were less likely to develop diabetes but among other groups, treatment didn't show any impact on the development of diabetes. BMI still played a significant role in developing diabetes regardless of treatment types.

Strengths:

One of the strengths of this study is innovative findings about the prognosis of colorectal cancer treatment stratified by treatment types. Especially, as it examined the impact of treatment on the risk of new chronic disease after diagnosis, it became significant evidence that suggests practical insights in developing a proper monitoring system for patients with colorectal cancer and their outcomes after treatment and diagnosis. It is imperative for providers to guide patients and caregivers to prevent adverse outcomes like new onset of chronic disease based on BMI and types of treatment. The next strength is the national database. As the study used the national database, the generalizability is validated.

Weaknesses: Even though the study attempted to examine the impact of each treatment option, the dosage of chemotherapy and the types of chemotherapy were not able to be examined due to the data source.