

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

September 5, 2023 (this version)

Posted to bioRxiv

August 20, 2023

Sent for peer review

July 11, 2023

SUMOylation of Bonus, the *Drosophila* homolog of Transcription Intermediary Factor 1, safeguards germline identity by recruiting repressive chromatin complexes to silence tissue-specific genes

Baira Godneeva, Maria Ninova, Katalin Fejes Tóth, Alexei A. Aravin 

California Institute of Technology, Division of Biology and Biological Engineering Pasadena, CA 91125, USA •
Institute of Molecular Genetics, National Research Center "Kurchatov Institute", Moscow 123182, Russia •
University of California, Riverside, Riverside, CA 92521, USA

 https://en.wikipedia.org/wiki/Open_access

 <https://creativecommons.org/licenses/by/4.0/>

Abstract

The conserved family of Transcription Intermediary Factors (TIF1) proteins consists of key transcriptional regulators that control transcription of target genes by modulating chromatin state. Unlike mammals that have four TIF1 members, *Drosophila* only encodes one member of the family, Bonus. Bonus has been implicated in embryonic development and organogenesis and shown to regulate several signaling pathways, however, its targets and mechanism of action remained poorly understood. We found that knockdown of Bonus in early oogenesis results in severe defects in ovarian development and in ectopic expression of genes that are normally repressed in the germline, demonstrating its essential function in the ovary. Recruitment of Bonus to chromatin leads to silencing associated with accumulation of the repressive H3K9me3 mark. We show that Bonus associates with the histone methyltransferase SetDB1 and the chromatin remodeler NuRD and depletion of either component releases Bonus-induced repression. We further established that Bonus is SUMOylated at a single site at its N-terminus that is conserved among insects and this modification is indispensable for Bonus's repressive activity. SUMOylation influences Bonus's subnuclear localization, its association with chromatin and interaction with SetDB1. Finally, we showed that Bonus SUMOylation is mediated by the SUMO E3-ligase Su(var)2-10, revealing that although SUMOylation of TIF1 proteins is conserved between insects and mammals, both the mechanism and specific site of modification is different in the two taxa. Together, our work identified Bonus as a regulator of tissue-specific gene expression and revealed the importance of SUMOylation as a regulator of complex formation in the context of transcriptional repression.

eLife assessment

This **important** study advances our understanding of the functions and regulation of the *Drosophila* transcriptional regulator Bonus, an ortholog of mammalian TIF1 family members. **Solid** evidence, leveraging both in vivo and in vitro studies, reveals the molecular requirements for Bonus' function as a transcriptional repressor, including a key role for sumoylation.

Introduction

Epigenetic regulation of gene expression is an essential mechanism that guides cell differentiation during development. The post-translational modifications of chromatin proteins act in combination with various chromatin-remodeling proteins to mediate changes in transcriptional activities and chromatin structure (reviewed in [Berger, 2007](#) [↗](#); [Kouzarides, 2007](#) [↗](#)). TRIM/RBCC is an ancient protein family characterized by the presence of an N-terminal RING finger domain closely followed by one or two B-boxes and a coiled coil domain. Additional protein domains found at their C termini have been used to classify TRIM proteins into subfamilies. The Transcriptional Intermediary Factor 1 (TIF1) proteins present in Bilaterian species contain PHD and Bromo domains at their C-terminus and belong to Subfamily E according to [Marín, 2012](#) [↗](#) or structural class E according to [Ozato et al., 2008](#) [↗](#). In vertebrates this subfamily contains four proteins: TIF1 α /TRIM24, TIF1 β /TRIM28, TIF1 γ /TRIM33, and TIF1 δ /TRIM66, while only one protein, Bonus (Bon), is present in *Drosophila*, making it an attractive model to understand the conserved functions of TIF1 proteins.

Mammalian TIF1 proteins are chromatin-associated factors that have been shown to play an essential role in transcription, cell differentiation, cell fate decisions, DNA repair and mitosis ([Bai et al., 2010](#) [↗](#); [Cammis et al., 2004](#) [↗](#), [2000](#) [↗](#); [Kulkarni et al., 2013](#) [↗](#); [Le Douarin et al., 1996](#) [↗](#); [Nielsen et al., 1999](#) [↗](#); [Sedgwick et al., 2013](#) [↗](#)). TIF1 proteins modulate the transcription of target genes by binding to co-regulators in the genome and controlling the chromatin state ([Khetchooumian et al., 2004](#) [↗](#); [Nielsen et al., 1999](#) [↗](#); [Schultz et al., 2002](#) [↗](#), [2001](#) [↗](#); [Venturini et al., 1999](#) [↗](#)). One of the best characterized TIF1 proteins, KAP-1 (TIF1 β), is the universal cofactor for the large family of Kruppel-associated box zinc finger proteins (KRAB-ZFPs) composing one of the best-studied gene silencing systems in vertebrates ([Friedman et al., 1996](#) [↗](#)). Diverse KRAB-ZFPs recognize specific DNA sequences with the majority targeting endogenous retroviruses, ensuring their repression. After target recognition by KRAB-ZFPs, KAP-1 suppresses target transcription with the help of the H3K9-specific histone methyltransferase SetDB1, the H3K9me3 reader HP1 and the NuRD histone deacetylase complex ([Schultz et al., 2002](#) [↗](#), [2001](#) [↗](#)).

The only member of the TIF1 subfamily in *Drosophila*, Bon was shown to be important in the development of several organs and somatic tissues during embryogenesis and metamorphosis, including the nervous system and the eye ([Allton et al., 2009](#) [↗](#); [Beckstead et al., 2001](#) [↗](#); [Ito et al., 2012](#) [↗](#); [Kimura et al., 2005](#) [↗](#); [Salzberg et al., 1997](#) [↗](#); [Zhao et al., 2023](#) [↗](#)). Bon has been shown to regulate the function of different signaling pathways to drive developmental fate decisions, such as the ecdysone pathway in the ovary ([Beckstead et al., 2001](#) [↗](#)) and the Hippo pathway in the eye ([Zhao et al., 2023](#) [↗](#)). Bon can act as both an Enhancer and a Suppressor of position-effect variegation ([Beckstead et al., 2005](#) [↗](#)), suggesting that it might play different roles that depend on specific interactors.

Many TRIM proteins from different subfamilies, including the mammalian TIF1 γ /TRIM33, act as ubiquitin ligases, suggesting that this was the ancient function of the family. On the other hand, several members, including the mammalian KAP-1 was shown to be active as E3 SUMO-ligases. Furthermore, SUMOylation plays an essential role in KAP-1 function: KAP-1 is SUMOylated through its own activity and SUMOylation is required for its repressive function by facilitating recruitment of the SetDB1 histone methyltransferase (Ivanov et al., 2007 [↗](#); Lee et al., 2007 [↗](#); Li et al., 2007 [↗](#); Mascle et al., 2007 [↗](#)). SUMO (small ubiquitin like modifier) is a small protein that is covalently conjugated to lysine residues of substrates that can modify and enhance protein-protein interactions (Gareau and Lima, 2010 [↗](#); Jentsch and Psakhye, 2013 [↗](#); Martin et al., 2007 [↗](#)). SUMOylation has been implicated in facilitating formation of protein complexes and condensates, especially in the nucleus, in different contexts including DNA repair, transcriptional repression and formation of subnuclear structures and chromatin domains (reviewed in Garvin and Morris, 2017 [↗](#); Gill, 2005 [↗](#); Verger et al., 2003 [↗](#)). The SUMO conjugation cascade involves the E1-activating enzyme, the E2-conjugating enzyme, and multiple E3-ligases that interact with E2 and facilitate the transfer of SUMO to the final substrates (Gill, 2004 [↗](#), p. 20; Johnson and Gupta, 2001 [↗](#)).

Here, we show that depletion of Bon in the female germline results in defective oogenesis and female infertility. We found that Bon controls oogenesis through repression of ectopic gene expression indicating that it serves as a guardian of cell type identity. Mechanistically, we found that Bon induces transcriptional repression through interaction with the dNuRD chromatin remodeler and the SetDB1 histone methyltransferase. We show that Bon is SUMOylated at a single site at its N-terminus and that this modification is essential for Bon-induced transcriptional silencing. Furthermore, this modification is important for Bon subnuclear localization and chromatin association as well as its interaction with SetDB1. The N-terminal SUMOylation site is conserved in insect species, but not in mammalian KAP-1 where several SUMOylation sites were reported at the C-terminal portion of the protein. Finally, we established that Bon SUMOylation depends on a distinct SUMO E3-ligase, Su(var)2-10, in contrast to mammalian KAP-1 that auto-SUMOylates itself. Our results identify Bon as a regulator of tissue-specific gene expression and highlight the universal function of SUMOylation as a regulator of complex formation in the context of transcriptional repression. On the other hand, our work suggest that SUMOylation of *Drosophila* Bon and mammalian KAP-1 has evolved independently and through distinct mechanisms revealing a remarkable case of parallel evolution in insects and vertebrates.

Results

***bon* knockdown in the female germline interferes with germline stem cells function and leads to arrested oogenesis and sterility**

According to FlyAtlas, the *bon* gene encodes a nuclear protein that is expressed throughout development with high level of expression in several tissues including the brain, gut, and ovaries (FlyAtlas [Chintapalli et al., 2007 [↗](#)]). Immunostaining with antibodies against Bon revealed that it is expressed in both the germline and somatic cells at all stages of oogenesis, starting from the germarium which contains germline stem cells (GSCs) to late-stage egg chambers where GSC-derived nurse cells support maturing oocytes (Fig. 1A [↗](#)). While *bon* was shown to be required for metamorphosis and the development of the nervous system (Beckstead et al., 2001 [↗](#); Ito et al., 2012 [↗](#)), its function in the germline remained unknown. To gain insights into the germline functions of Bon, we generated transgenic flies expressing short hairpin RNAs (shRNAs) against *bon* under control of the UAS/Gal4 system and performed germline-specific RNAi knockdown (GLKD). Using the *maternal tubulin-Gal4* (*MT-Gal4*) driver, we found by RT-qPCR that two distinct shRNAs targeting *bon* led to 75% and 88% reduction in ovarian Bon expression, respectively (Fig. 1B [↗](#)). Because the *MT-Gal4* driver is active in the germline, but not in follicular cells, the actual knockdown efficiency of *bon* in germ cells is even higher than what we detected from whole

ovarian lysates. Indeed, immunofluorescence confirmed that Bon protein had been efficiently depleted from germline cells (**Fig. 1C**). For all subsequent experiments, we used the shRNA that resulted in higher knockdown efficiency.

To analyze the role of Bon throughout the developmental progression of the germline we combined the *bon* shRNA construct with three different germline Gal4 drivers using different stage-specific promoters: *bam-Gal4*, which is expressed from cystoblasts to eight-cell cysts; *MT-Gal4*, which drives expression in germ cells starting in stage 2 of oogenesis, and *nos-Gal4* driver, which induces expression in two distinct stages, in GSCs and at late stages of oogenesis (Extended Data Fig. 1A) (Chen and McKearin, 2003; Doren et al., 1998; McKearin and Ohlstein, 1995). Bon GLKD driven by either *MT-Gal4*, *bam-Gal4* or the double driver (*MT+bam*) did not result in significant changes in ovarian morphology compared to controls (**Fig. 1D**). Furthermore, such females laid eggs and were fertile. Thus, germline depletion of Bon starting at the cystoblast stage does not lead to morphological or obvious functional defects in oogenesis. In contrast, silencing of Bon beginning in the GSCs by expressing the shBon using *nos-Gal4* driver induces visible morphological changes with 34% of flies having only rudimentary ovaries lacking late stages of oogenesis and another ~39% having one of the two ovaries rudimentary (**Fig. 1E**). An even stronger phenotype was observed upon GLKD using a double *nos+bam* driver which drives expression at all stages of oogenesis (**Fig. 1D, F**). 100% of such females displayed rudimentary ovaries and were completely sterile (**Fig. 1D, E**). Consistent with this, immunostaining for the germ cell marker Vasa demonstrates that depletion of Bon results in partial loss of germ cells and arrested oogenesis as morphological defects were accompanied by loss of vasa-positive cells from the egg chambers (**Fig. 1F**). Additionally, we proved the importance of Bon in the *Drosophila* ovarian germline by using CRISPR/Cas9 mediated mutagenesis. Transgenic flies from the Heidelberg CRISPR Fly Design Library (Port et al., 2020) expressing sgRNAs targeting *bon* were crossed to *nos-Gal4;UAS-Cas9* to achieve germline-specific knockout of *bon*. Almost 65% of the female offspring with *nos-Cas9;sgRNA-bon* were sterile and had defects in ovarian morphology, another 23% had one rudimentary ovary and only 12% showed normal phenotype (Extended Data Fig. 1B). These results indicate that Bon function in the early stages of oogenesis, particularly in GSCs, is essential for proper oogenesis.

To further analyze the role of Bon in the maintenance of GSCs and early oogenesis we used immunofluorescence against the cytoskeletal protein α -spectrin, which marks the spectrosome, a spherical intracellular organelle present in GSCs and cystoblasts. At later stages, spectrosomes become fusomes, branched structures that are localized in cytoplasmic bridges connecting differentiating germ cells in the growing cysts. Thus, fusome formation is a hallmark of normal oogenesis progression. In ovarioles of control flies, we observed a normal germarium organization with 2-3 spectrosome-containing GSCs, and branched fusomes in germ cells at later stages. In contrast, germ cells with normal fusomes were absent upon depletion of Bon using *nos-Gal4*. Instead, the germarium of Bon-depleted flies harbored several cells containing spherical spectrosomes, a hallmark of GSCs or cystoblast-like undifferentiated germ cells (Extended Data Fig. 1C, D). Overall, our results indicate that loss of Bon in early germ cells interferes with maintenance of GSCs and arrests their further differentiation.

Loss of Bonus triggers the ectopic expression of tissue-specific genes in the ovary

To investigate the effect of Bon depletion on gene expression in the female germline, we performed transcriptome profiling using RNA sequencing (RNA-seq) analysis. We tested the effects of loss of Bon in both early and later stages of oogenesis using the *nos-Gal4* or *MT-Gal4* driver to drive *bon* shRNA expression, respectively. RNA-seq libraries were prepared in triplicates and compared to respective control libraries. As knockdown using the *nos-Gal4* driver causes early arrest of oogenesis and rudimentary ovaries, while later-stage knockdown with the *MT-gal4* driver yields normal ovaries, we used different controls depending on the driver to assure that ovary size




Figure 1.

Germline expression of *Bonus* is required for oogenesis.

A) *Bon* is expressed throughout oogenesis. Stacked confocal image of wild-type Oregon-R flies stained for *Bon*. **B**) Bar graph shows the relative expression of *Bon* (normalized to *rp49* level) in control and *Bon*-depleted ovaries (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **C**) Confocal images of egg chambers from wild-type Oregon-R flies (control) and flies expressing *MT-Gal4*-driven shRNA against *Bon* stained for *Bon* (scale bar: 20 μ m). **D**) *Bon* depletion leads to rudimentary ovaries. Phase contrast images of dissected ovaries from flies of indicated genotypes. Wild-type Oregon-R flies were used as control. **E**) Top: Phase contrast image of dissected ovaries with different phenotypes from flies with *Bon* GLKD driven by *nos-Gal4*. Bottom: graph showing the percentage of normal, hypomorphic and rudimentary ovary phenotypes of indicated genotypes (n = 85, 175, 52, 71, 64, 58, respectively). **F**) Confocal images of whole ovaries from wild-type Oregon-R flies (control) and flies with *Bon* GLKD driven by *bam+nos* double driver stained for Vasa (red) and DAPI (blue) (scale bar: 30 μ m).

and cell composition of the Bon GLKD and control are similar. For *nos-Gal4* we used ovaries from young (0-1 day old) flies that lack later stages of oogenesis and compared them to their age-matched siblings that lack the shRNA, and for *MT-Gal4* we used 1-2-old flies that express either shRNA against *bon* or the *white* gene, which is not expressed in the germline. Thus, in both cases, GLKD and control flies had the same age and similar ovary size. As the mammalian homolog of Bon, KAP-1, plays a central role in repression of many transposable elements (TEs) through its function as co-repressor for multiple KRAB-ZFPs that recognize TE sequences, we analyzed expression of both host genes and TEs.

Most TE families were not affected by Bon depletion using either driver. Using the *nos-Gal4*-driven shRNA, only 6 out of 207 (~3%) TE families present in the *Drosophila* genome significantly increased their expression more than 2-fold ($\log_2FC > 1$, and $qval < 0.05$, LRT test, Sleuth) (Extended Data Fig. 2A) and none showed strong (>6-fold) upregulation. Similarly, depletion of Bon at later stages of oogenesis also did not lead to strong (>6-fold) change in transposon expression (Extended Data Fig. 2A). This phenotype is in stark contrast to the significant activation of many TE families when the main TE repression pathway in the ovary – the piRNA pathway – is abolished, suggesting that Bon is not involved in the piRNA pathway and that oogenesis defects observed upon Bon depletion likely have a different molecular basis.

In contrast to TEs, the protein-coding transcriptome was severely disrupted upon Bon depletion - differential gene expression analysis using Sleuth revealed many genes with altered steady state RNA levels upon Bon GLKD at either stage. As expected, Bon was one of the most strongly downregulated genes, showing ~7-fold reduction and confirming the efficiency of KD and the validity of the RNA-seq data (Fig. 2A). Early Bon GLKD resulted in 694 differentially expressed genes ($qval < 0.05$, LRT test) (Fig. 2A), of which 464 (~67%) and 28 (~4%) genes, respectively, increased and decreased their expression more than two-fold, while late Bon GLKD using *MT-Gal4* revealed 1769 genes that were differentially expressed ($qval < 0.05$, LRT test), with 231 genes (~13.6%) showing more than two-fold increase, while 72 genes (~4.2%) showed more than a two-fold decrease in mRNA level (Extended Data Fig. 2B). Interestingly, the sets of genes that change their expression upon Bon GLKD at the early and late stages of oogenesis are quite different: only 51 genes were derepressed at both stages of oogenesis, while the remaining genes that changed their expression were unique for one or the other stage (Extended Data Fig. 2C). Overall, our results indicate that Bon plays an important role in regulation of gene expression during oogenesis with distinct targets at different stages.

To characterize Bon targets in oogenesis, we performed gene ontology (GO) analysis of genes strongly upregulated upon *nos-Gal4*-driven Bon GLKD ($n=464$). GO analysis identified enrichment of genes from 27 biological process in the set of Bon-repressed genes (Fig. 2B). These included terms such as mesoderm development, myofibril assembly, sarcomere organization, hemolymph coagulation, motor neuron axon guidance, and visceral muscle development, suggesting that Bon loss leads to the ectopic ovarian activation of genes normally expressed in other tissues. To comprehensively explore the specific expression patterns of the 464 Bon-repressed genes, we used modENCODE RNA-seq data from different tissues. This analysis revealed that many genes that are derepressed in the ovary upon Bon GLKD are normally expressed in other tissues and have no (55%) or low (33%) expression in the ovary of wild-type flies. Instead, many of these genes are predominantly expressed in the head (50%), digestive system (38%), and central nervous system (28%) of wild-type flies (Fig. 2C).

We used RT-qPCR and *in situ* hybridization chain reaction (HCR) for selected upregulated genes including *rbp6*, *CG34353* and *ple*, which are highly expressed in the head, and *pst*, which is highly expressed in the gut, to confirm that germline depletion of Bon triggers their ectopic activation. No signal for these genes was detected in wild-type ovaries, while abundant *rbp6*, *CG34353* and *pst* transcripts were identified in germ cells upon *MT-Gal4>Bon GLKD* (Fig. 2D, E, F, Extended Data Fig. 2D, E, F). Surprisingly, depletion of Bon in germ cells caused the appearance of *ple* transcripts




Figure 2.

Bonus functions as a repressor of tissue-specific genes in ovary.

A) Bon GLKD leads to misexpression of tissue-specific genes in the ovary. Volcano plot shows fold changes in genes expression upon Bon GLKD driven by *nos-Gal4* in the ovary as determined by RNA-seq ($n=3$). Siblings that lack shRNA against Bon produced in the same cross were used as a control. Genes that change significantly ($\log_2FC>1$, $qval<0.05$, LRT test, sleuth (Pimentel et al., 2017)) are highlighted. Genes *bon*, *pst*, *Rbp6*, *ple* are labeled. Genes with infinite fold change values (zero counts in control ovaries) are not shown. **B)** Bon represses genes with diverse functions. Bubble plot shows the analysis of GO enrichment at the level of biological processes (BP) for genes that are de-repressed upon Bon GLKD driven by *nos-Gal4* ($\log_2FC>1$, $qval<0.05$, LRT test, sleuth (Pimentel et al., 2017)). Only GO terms above the established cut-off criteria ($p\text{-value}<0.01$ and >3 genes per group) are shown. BP are ranked by fold enrichment values. The most significant processes are highlighted in purple, and the less significant in yellow according to $\log_{10}(FDR)$ values. The bubbles size reflects the number of genes, assigned to the GO BP terms. **C)** Normal expression level of de-regulated genes upon Bon GLKD in the tissues where they are normally expressed indicates Bon-mediated silencing of genes normally expressed in the head and digestive system. The graph shows the percentage of de-repressed genes upon Bon GLKD driven by *nos-Gal4* ($\log_2FC>1$, $qval<0.05$, LRT test, sleuth (Pimentel et al., 2017)) with given expression level in the indicated enriched tissues. Expression levels according RPKM values from modENCODE anatomy RNA-seq dataset are no expression (0-0), very low (1-3), low (4-10), moderate (11-25), moderate high (26-50), high (51-100), very high (101-1000), extremely high (>1000). **D)** GLKD of Bon leads to *ple* expression in follicular cells. Confocal images of egg chambers show RNA *in situ* HCR detecting *ple* and *bonus* mRNAs in flies with *MT-Gal4>Bon GLKD* and control siblings from the same cross that lack Bon shRNA (scale bar: 20 μm). **E)** Bon represses *rbp6* in the germline. Confocal images of egg chambers show RNA *in situ* HCR detecting *rbp6* and *bonus* mRNAs in flies with *MT-Gal4>Bon GLKD* and control siblings from the same cross that lack Bon shRNA (scale bar: 20 μm). **F)** Bar graph shows the relative expression of *ple* and *rbp6* (normalized to rp49 level) in control and Bon-depleted ovaries (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.).

in somatic follicular cells that surround germline cells, suggesting that Bon depletion causes activation of *ple* indirectly, through a process that involves signaling between the adjacent germline and follicular cells (**Fig. 2D** [↗](#)). Overall, our results indicate that Bon is required for repression of tissue-specific genes in the ovary.

Recruitment of Bonus to a genomic locus induces transcriptional repression associated with accumulation of the H3K9me3 mark

Transcriptome profiling upon Bon germline depletion demonstrated global changes in steady-state RNA levels of hundreds of genes. As exemplified by the activation of the *ple* gene in the somatic follicular cells some of these effects might be indirect and even mediated by intercellular signaling. To test the ability of Bon to directly induce transcriptional silencing, we took advantage of a tethering approach in which Bon is recruited to a reporter locus via binding to nascent transcripts (**Fig. 3A** [↗](#)). Tethering was achieved through fusion of Bon to the λ N RNA-binding domain that has high affinity for BoxB RNA hairpins encoded in the 3'UTR region of the reporter gene (De Gregorio, 1999 [↗](#)). λ N-eGFP-Bon and the reporter were co-expressed in the germline using the *MT-Gal4* driver; recruitment of λ N-eGFP was used as a control.

RT-qPCR showed that tethering of Bon triggers ~22-fold reporter repression (**Fig. 3B** [↗](#)). Similar results were obtained with a different reporter in another genomic location, indicating that recruitment of Bon induces strong repression regardless of the genomic locus (Extended Data Fig. 3A). ChIP-qPCR analysis revealed that Bon recruitment results in a strong increase in the repressive H3K9 trimethylation (H3K9me3) chromatin mark, at the reporter locus (**Fig. 3C** [↗](#)), suggesting that repression induced by Bon is mediated, at least in part, by the deposition of H3K9me3.

We also examined changes in H3K9me3 enrichment on genes upregulated upon Bon depletion (Bon GLKD driven by *nos-Gal4*). Global ChIP-seq analysis revealed that many Bon-dependent genes show low or no H3K9me3 signal in control ovaries and no change upon Bon depletion, hence likely represent secondary targets (**Fig. 3D** [↗](#)). However, several Bon-regulated genes are enriched in H3K9me3 mark in wild-type ovaries including in the proximity of the transcription start site (TSS) and show prominent loss of H3K9me3 upon Bon depletion (**Fig. 3D** [↗](#)). For example, gene *CG1572*, which was activated two-fold upon Bon GLKD, showed almost a two-fold decrease in H3K9me3 level upstream of its TSS (**Fig. 3E** [↗](#)). Independent ChIP-qPCR analysis of few Bon-regulated genes such as *CG3191* and *Spn88Eb* also showed a slight decrease in the repressive mark upon Bon depletion (**Fig. 3F** [↗](#)).

Altogether, these data indicate that Bon recruitment to genomic targets induces transcriptional repression associated with accumulation of the H3K9 trimethylation mark. However, repression of many Bon-regulated genes might be indirect and/or independent of H3K9me3.

Bonus interacts with dNuRD complex components Mi-2 and Rpd3, as well as the histone methyltransferase SetDB1

Mammalian KAP-1 was shown to associate with the NuRD histone deacetylase and chromatin remodeling complex and with the H3K9me3 writer SetDB1, and their interactions are important for its function in transcriptional repression (Schultz et al., 2002 [↗](#), 2001 [↗](#)). To study whether dNuRD and SetDB1 are required for Bon's ability to trigger transcriptional repression in the *Drosophila* germline, we tested reporter expression upon Bon tethering and concomitant knockdown of SetDB1 and dNuRD components. GLKD of either Mi-2 or SetDB1, but not Rpd3, inhibited silencing, indicating that Mi-2 and SetDB1 act downstream of Bon to induce repression (**Fig. 4A** [↗](#)).




Figure 3.

Bonus induces transcriptional silencing.

A) Schematics of the reporter construct in flies that allows Bon recruitment to nascent reporter transcript in flies. λ N-GFP-Bonus and the mKate reporter encoding 4BoxB hairpins are co-expressed in germline cells of the ovary (driven by *MT-Gal4*). **B)** Bon tethering leads to transcriptional silencing of the reporter. Bar plot shows reporter expression (normalized to rp49 level) upon tethering of λ N-GFP-Bonus or λ N-GFP control ovaries (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **C)** Bon tethering leads to H3K9me3 accumulation. Bar plot shows H3K9me3 enrichment upon tethering of λ N-GFP-Bonus or λ N-GFP control ovaries (ChIP-qPCR, dots correspond to 2 independent biological replicates; error bars indicate st. dev.). **D)** Heatmap shows H3K9me3 distribution across Bon targets in control and *nos-Gal4>Bon GLKD* ovaries (input-normalized log2 values). **E)** RNA-seq and ChIP-seq tracks show counts per million (CPM)-normalized coverage for *CG1572* in control and *nos-Gal4>Bon GLKD* ovaries. The gene structure is depicted at the top; arrow indicates the direction of transcription. The ChIP (blue) and input (gray) signals are overlaid. Numbers show the CPM values of the exonic regions (RNA-seq) or the normalized ChIP/input signal (ChIP-seq) in a manually selected genomic location. **F)** Bon depletion results in a slight decrease in H3K9me3 over some Bon target genes. Bar graph shows H3K9me3 levels at the genes *CG3191* and *Spn88Eb* in control and Bon-depleted ovaries (ChIP-qPCR, dots correspond to 2 independent biological replicates; error bars indicate st. dev.).




Figure 4.

Bonus interacts with Mi-2, Rpd3 and SetDB1.

A) Reporter silencing by Bon depends on Mi-2 and SetDB1. Bar plot showing the reporter expression (normalized to rp49 level) upon tethering of control λ N-GFP or λ N-GFP-Bonus in ovaries with Rpd3, Mi-2, SetDB1 GLKD, and control *white* GLKD (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **B-C**) Bon interacts with SetDB1 and Rpd3. Western blot analysis of immunoprecipitation experiment using GFP nanotrap beads from S2 cells co-expressing GFP-Bonus and Flag-tagged SetDB1 (**B**) and Flag-tagged Rpd3 (**C**). Lysates not expressing GFP-Bonus were used as negative control. **D**) Bon interacts with the C-terminus of Mi-2. Top: schematic illustration of full-length *Drosophila* Mi-2 and its truncated versions as defined by the amino acids: C-terminal truncated Mi-2 (1-1680) and N-terminal truncated Mi-2 (1681-1982). Bottom: Western blot analysis of immunoprecipitation experiment using GFP nanotrap beads from S2 cells co-expressing GFP-Bonus and Flag-tagged Mi2 fragments. Lysate not expressing GFP-Bonus was used as negative control.

To explore physical interactions of Bon with components of the dNuRD and SetDB1 complexes, we employed co-immunoprecipitation assay using tagged proteins in S2 cells. In *Drosophila*, the dNuRD complex mediates chromatin remodeling and histone deacetylation through dMi-2 and Rpd3 (HDAC1 homolog), respectively (Bouazoune and Brehm, 2006 [DOI](#); Brehm, 2000 [DOI](#); Kunert and Brehm, 2009 [DOI](#); Rubertis et al., 1996 [DOI](#); Tong et al., 1998 [DOI](#)). We found that both components of dNuRD, Mi-2 and Rpd3, as well as SetDB1 co-purify with Bon (Fig. 4B, C, D [DOI](#), Extended Data Fig. 3B). The interaction between Bon and Mi-2 is mediated by the C-terminus of Mi-2 (Fig. 4D [DOI](#)), similar to interaction between Mi-2 α /CHD3 and KAP-1 in mammals (Schultz et al., 2001 [DOI](#)). In *Drosophila* Mi-2 is found in two distinct complexes, the canonical dNuRD complex and the dMec complex that contains the zinc-finger protein Mep-1 (Kunert et al., 2009 [DOI](#)). We did not detect an interaction between Bon and Mep-1 (Extended Data Fig. 3C), indicating that Bon interacts with Mi-2 in the context of the dNuRD complex but not dMec. Overall, our results indicate that the interactions between Bon and the NuRD and SetDB1 chromatin remodeler and modifying complexes are essential for its repressor activity and evolutionarily conserved among members of the TIF1 protein family between insects and mammals.

Bonus is SUMOylated at a single site close to its N-terminus

Self-SUMOylation of mammalian KAP-1 is essential for its repressive function (Ivanov et al., 2007 [DOI](#)). In our analysis of immunopurified Bon by Western blotting, we noticed a band of higher molecular weight, indicative of a post-translationally modified form. To explore if Bon is SUMOylated, we co-expressed tagged SUMO and Bon in S2 cells followed by immunoprecipitation of Bon under stringent washing conditions to remove non-covalently bound proteins in the presence of N-ethylmaleimide (NEM), an inhibitor of SUMO-specific deconjugating enzymes. Western blot revealed the presence of unmodified and single SUMO-modified forms of Bon (Extended Data Fig. 4A, top). To explore if Bon is SUMOylated in fly ovaries, we immunoprecipitated Bon from ovarian extracts of flies that express Flag-tagged SUMO in the germline. As in S2 cells, we observed unmodified and single SUMO-modified Bon forms, indicating that a fraction of the Bon protein pool is SUMOylated in both S2 cells and ovaries (Extended Data Fig. 4A, bottom).

To find potential SUMOylation sites in Bon we used the SUMOplotTM Analysis Program which yielded three high-scoring predicted residues: two sites, lysine K9 (LKND) and K20 (IKQE), are conforming to the canonical consensus site for SUMOylation, Ψ KxD/E, whereas the third one, K763, resides in a noncanonical motif, LKSP (Fig. 5A [DOI](#)). Unlike wild-type Bon, the triple mutant with all three lysine residues substituted to arginine was not SUMOylated when expressed in either S2 cells or fly ovaries (Fig. 5B [DOI](#), Extended Data Fig. 4B). To further narrow the modification site, we created individual point mutants and checked their SUMOylation. The single K20R mutation completely abolished Bon SUMOylation (Fig. 5C [DOI](#)). Together with the characteristic shift in modified Bon migration on SDS-PAGE gel, these results indicate that the bulk of SUMO-modified Bon carries a single SUMO moiety at the K20 residue. To explore whether this SUMOylation site is conserved in other *Drosophila* species, we performed an alignment of predicted Bon homologs from the genomes of 12 fully sequenced species using ClustalW. We found that the consensus SUMOylation site at the K20 position of *D. melanogaster* Bon is conserved in all analyzed *Drosophila* species (Fig. 5D [DOI](#), Extended Data Fig. 4C). Furthermore, analysis revealed conservation of the position of SUMOylated lysine residue in distantly related insects including the buff-tailed bumblebee *Bombus terrestris*, the honeybee *Apis mellifera*, yellow fever mosquito *Aedes aegypti*, and diamondback moth *Plutella xylostella* (Fig. 5D [DOI](#)), suggesting that SUMOylation at the Bon N-terminus is conserved across many insects. On larger evolutionary distances, although SUMOylation is conserved between the mammalian KAP-1 and the *Drosophila* Bon, its position is different: while mammalian KAP-1 is SUMOylated at multiple sites that all reside in its C-terminus, Bon is SUMOylated at a single site close to its N-terminus.




Figure 5.

Bonus is SUMOylated.

A) Schematic representation of putative SUMOylation sites within Bon. SUMOylation consensus sites are shown and boxed. Canonical consensus sites are in bold. Putative SUMOylated lysines were mutated to arginines individually (K9R, K20R, K763R) or in combination (3KR). **B**) Bon is SUMOylated at specific residues. Western blot analysis shows the SUMOylation levels of GFP-tagged Bon and SUMO-deficient triple mutant 3KR expressed in fly ovaries. SUMOylated form of Bon was detected only in wild-type GFP-Bonus (WT). Total protein lysates from flies co-expressing Flag-SUMO and λN-GFP-Bonus or λN-GFP-Bonus[3KR] were immunopurified using anti-GFP nanotrap beads. Flies not expressing λN-GFP-tagged protein were used as a negative control. **C**) Bon is predominantly SUMOylated at K20. Western blot analysis shows the associated SUMOylation levels of GFP-tagged Bon and SUMO-deficient triple mutant 3KR and single mutated K9R, K20R expressed in S2 cells. Single mutation K9R reduced, while the K20R mutation and triple 3KR mutation completely abolished Bon SUMOylation. Total protein lysates from S2 cells co-expressing HA-SUMO and GFP-Bonus or GFP-Bonus[3KR], GFP-Bonus[K9R], GFP-Bonus[K20R] were immunopurified using anti-GFP nanotrap beads. Lysate not expressing GFP-tagged protein was used as a negative control. **D**) SUMOylation site of Bon is conserved in insects. Sequence alignment of the Bon protein sequence from 12 *Drosophila* species and other insects shows conserve action of canonical SUMOylation consensus at K20 (boxed and indicated by the arrowhead). **E**) SUMO-deficient Bon mislocalizes into nuclear foci. Confocal images of egg chambers show the localization of *MT-Gal4*-driven λN-GFP-tagged Bonus and SUMO-deficient triple mutant λN-GFP-Bonus[3KR] flies. Images on the right panel show isolated nurse cell nuclei (scale bar: 20μm). **F**) Chromatin association of Bon depends on its SUMOylation. Western blot analysis shows the fractionation of cytoplasmic (cyto), nuclear (nuclei) and chromatin compartments of *MT-Gal4*-driven λN-GFP-tagged Bonus (WT) and SUMO-deficient triple mutant λN-GFP-Bonus (3KR) fly ovaries. Lamin and Histone H3 were used as markers for nuclear and chromatin fractions. **G**) Bon-mediated reporter repression depends on Bon SUMOylation. Bar plot shows the reporter expression (normalized to rp49 level) upon tethering of λN-GFP-Bonus, SUMO-deficient triple mutant λN-GFP-Bonus[3KR] or λN-GFP control ovaries (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **H**) Bar plot shows the reporter expression (normalized to rp49 level) upon tethering of control λN-GFP or λN-GFP-Bonus in ovaries with SUMO GLKD, and control *white* GLKD (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **I**) Western blot analysis shows the SUMO-dependent interaction between Bon and SetDB1. Total protein lysates from S2 cells co-expressing Flag-SetDB1 and GFP-Bonus (WT) or triple mutant GFP-Bonus[3KR] (3KR) were immunopurified using anti-GFP nanotrap beads. Lysate from cells not expressing GFP-tagged protein was used as a negative control.

We compared the subcellular localization of wild-type and SUMO-deficient Bon in the germline. Wild-type Bon is localized in nurse cell nuclei and overall nuclear localization was not affected by lack of SUMOylation (**Fig. 5E**). Wild-type Bon shows enrichment in some nuclear regions, but is generally distributed throughout the nucleus. ~37% of Bon colocalized with DAPI-dense chromatin regions (Extended Data Fig. 4D), however, Bon showed poor (8%) overlap with HP1 protein, the mark of gene-poor and repeat-rich heterochromatin (Nakayama et al., 2001; Rea et al., 2000), indicating that Bon is not localized to the bulk of heterochromatin (Extended Data Fig. 4D, bottom). Remarkably, the lack of SUMOylation affected Bon distribution within the nucleus: mutant Bon was localized in more discrete nuclear foci compared to the wild-type protein (**Fig. 5E**). To explore whether SUMOylation affects Bon localization on chromatin we separated the chromatin fraction and probed Bon presence by Western blot. SUMO-deficient mutant exhibited significantly reduced association with the chromatin fraction (**Fig. 5F**), indicating that SUMOylation is essential for Bon's subnuclear localization and chromatin association.

To test the functional role of Bon SUMOylation, we explored the ability of SUMO-deficient Bon to induce transcriptional repression in the tethering assay. Unlike wild-type Bon, tethering of SUMO-deficient protein did not trigger repression of the reporter (**Fig. 5G**). Furthermore, knockdown of *smt3*, the single gene encoding SUMO in the *Drosophila* genome, resulted in a partial release of the reporter silencing caused by tethering of wild-type Bon (**Fig. 5H**). Thus, SUMOylation seems to be essential for the ability of Bon to induce transcriptional repression. Next, we tested if SUMOylation affects Bon interaction with SetDB1 and dNuRD. Immunoprecipitation assays showed that SUMOylation is dispensable for Bon interaction with Mi-2 and Rpd3 (Extended Data Fig. 4E, F), however, it decreased its interaction with SetDB1 (**Fig. 5I**). Overall, our results revealed that SUMOylation of Bon at a single site modulates its subnuclear localization and is essential for Bon's ability to induce transcriptional silencing and interact with SetDB1.

Bonus SUMOylation depends on the SUMO E3-ligase Su(var)2-10

To test if Bon acts as a SUMO E3-ligase and can promote self-SUMOylation similar to mammalian KAP-1, we explored its interaction with Ubc9, the only SUMO E2-conjugating enzyme in *Drosophila*. As E3-ligases facilitate transfer of SUMO from the E2 enzyme to the final substrates they form complexes with E2 that are readily detected by co-immunoprecipitation (Melchior, 2003; Pichler et al., 2002; Reverter and Lima, 2005). Co-immunoprecipitation of tagged Bon and Ubc9 did not reveal an interaction (Extended Data Fig. 5A), suggesting that Bon does not act as an E3-ligase. Interestingly, K20R but not K9R Bon point mutant coimmunoprecipitates with Ubc9 (**Fig. 6A**, Extended Data Fig. 5B), suggesting that inability to transfer SUMO from Ubc9 to Bon stabilizes the transiently formed complex between these proteins and the E3 SUMO-ligase potentially involved in the process.

Recently, Su(var)2-10 protein was identified as a Bon interactor in S2 cells (Zhao et al., 2023). We showed that Su(var)2-10 is an E3 SUMO-ligase that is required for suppressing tissue-inappropriate gene expression in the *Drosophila* germline, a function similar to the one we observed for Bon (Ninova et al., 2020a; Ninova et al., 2020b). To explore a possible interaction between Bon and Su(var)2-10, we first used co-immunoprecipitation, which confirmed binding of Bon and Su(var)2-10 (**Fig. 6B**). Second, we tested Bon SUMOylation upon germline depletion of Su(var)2-10. Remarkably, knockdown of Su(var)2-10 led to a complete loss of Bon SUMOylation, indicating that Bon modification strongly depends on Su(var)2-10 (**Fig. 6C**). Finally, germline depletion of Su(var)2-10 resulted in the derepression and loss of H3K9me3 signal on the reporter silenced by recruitment of Bon (**Fig. 6D, E**). Combined, these results show that Bon modification and its repressive function depend on the E3 SUMO-ligase Su(var)2-10. Altogether, our results indicate that Su(var)2-10 promotes Bon SUMOylation and is required for Bon-induced H3K9me3 deposition and transcriptional silencing.




Figure 6.

SUMO E3-ligase Su(var)2-10 interacts with Bonus and regulates its SUMOylation.

A) Western blot analysis shows the interaction between Bon and SUMO E2-conjugating enzyme Ubc9. Total protein lysates from S2 cells co-expressing Flag-Ubc9 and GFP-Bonus (WT), SUMO-deficient triple mutant 3KR or single mutated K9R, K20R were immunoprecipitated using anti-GFP nanotrap beads. Lysate from cells not expressing GFP-tagged proteins was used as a negative control. **B)** Bon interacts with Su(var)2-10. Western blot analysis of immunoprecipitation experiment using GFP nanotrap beads from S2 cells co-expressing GFP-Bonus and Flag-tagged Su(var)2-10. Lysate expressing only Flag-Su(var)2-10 was used as a negative control. **C)** Western blot analysis shows the loss of SUMOylated Bon in fly ovaries upon Su(var)2-10 depletion. Total protein lysates from flies co-expressing *MT-Gal4*-driven Flag-SUMO and λN-GFP-Bonus and shRNAs against Su(var)2-10 (Sv210) or control *white* (ctrl) were immunoprecipitated using anti-GFP nanotrap beads. Ovarian lysates from flies expressing only Flag-SUMO, only expressing λN-GFP-Bonus, or lacking Su(var)2-10 shRNA were used as controls. **D)** Reporter repression by Bon depends on Su(var)2-10. Bar plot shows reporter expression (normalized to rp49 level) upon tethering of control λN-GFP or λN-GFP-Bonus in ovaries with Su(var)2-10 GLKD (shSv210), and control *white* GLKD (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **E)** Bon H3K9me3 depositing requires Su(var)2-10. Bar plot shows H3K9me3 enrichment upon tethering of control λN-GFP or λN-GFP-Bonus in ovaries with Su(var)2-10 GLKD (shSv210), and control *white* GLKD (ChIP-qPCR, dots correspond to 2 independent biological replicates; error bars indicate st. dev.).

Bon, the only member of the TIF1 protein family in *Drosophila*, is ubiquitously expressed throughout development and previous studies have demonstrated its role in the development of several organs and tissues, particularly the nervous system (Allton et al., 2009; Beckstead et al., 2001; 2005; Ito et al., 2012; Kimura et al., 2005; Salzberg et al., 1997). We have found that depletion of Bon in the female germline leads to defective oogenesis resulting in rudimentary ovaries, loss of germline stem cells and sterility (Fig. 1D, E, F). Germline knockdown of Bon resulted in the misexpression of hundreds of genes, which are normally restricted to other tissues, such as the nervous system and the gut (Fig. 2A, 2B). Interestingly, Bon affects a diverse set of targets at different stages of oogenesis, as depletion of Bon using different germline drivers led to the ectopic expression of only partially overlapping sets of genes (Extended Data Fig. 2C). As Bon acts as a strong transcriptional repressor when recruited to a genomic locus (Fig. 3B), genes upregulated upon Bon depletion might be its direct targets. However, the finding that only a fraction of these loci have Bon-dependent H3K9me3 mark suggests that many of these genes are regulated by Bon indirectly, possibly through repression of other transcriptional regulators. Indirect regulation is further confirmed by the finding that germline depletion of Bon leads to enhanced expression of the *ple* gene in follicular cells surrounding the germline, where Bon levels were not perturbed (Fig. 2D). Unfortunately, several attempts to map direct targets of Bon on chromatin using ChIP-seq turned out to be unsuccessful. Previous studies support the idea that Bon controls the expression of other developmental regulators that affect cell fate decisions. For example, a recent study has demonstrated that Bon together with the Hippo pathway is involved in cell fate decisions during eye development (Zhao et al., 2023). In the ovary, the Hippo pathway acts downstream of Hedgehog signaling to regulate follicle stem cells maintenance (Hsu et al., 2017). Bon has also been shown to control genes in the ecdysone response pathway (Beckstead et al., 2001), which in turn regulates multiple steps during oogenesis and controls the development of the *Drosophila* ovary (Gancz et al., 2011; Hodin and Riddiford, 1998; König et al., 2011).

The interesting unresolved question is how Bon identifies its genomic targets. In mammals, KAP-1 is recruited to chromatin through interaction with the large and diverse family of KRAB domain-containing C2H2-zinc finger transcription factors (KRAB-ZFPs), which recognize their target DNA sequences – primarily various endogenous retroviruses and other types of transposable elements – via their zinc finger domains (Friedman et al., 1996; Margolin et al., 1994; Pengue et al., 1994; Vissing et al., 1995; Witzgall et al., 1994). However, KRAB-ZFPs appeared during vertebrate evolution and are absent in insects. In agreement with this, we found that depletion of Bon did not activate the expression of transposable elements and instead affects host genes (Fig. 2A, Extended Data Fig. 2A, B). A different class of ZFPs, ZAD (Zinc-finger-associated domain)-zinc finger proteins have expanded during insect evolution (Chung et al., 2007, 2002). It will be interesting to explore whether ZAD-ZFPs interact with and recruit Bon to its genomic targets, which if true, would represent a remarkable case of parallel evolution.

Our results reveal both similarities and differences between molecular mechanisms and functions of *Drosophila* Bon and mammalian TIF1 members, such as KAP-1. Similar to KAP-1, Bon induces transcriptional repression associated with accumulation of the H3K9me3 repressive histone mark (Fig. 3B, C) (Khetchoumian et al., 2004; Nielsen et al., 1999; Schultz et al., 2002, 2001; Venturini et al., 1999). Furthermore, both proteins interact with the histone methyltransferase SetDB1 and a member of the NuRD histone deacetylase complex and these interactions are important for their repressive functions (Fig. 4B, C, D, Extended Data Fig. 3B) (Ito et al., 2012; Schultz et al., 2002, 2001; Zhao et al., 2023). Finally, the repressive function of both KAP-1 and Bon requires their post-translational modification by SUMO (Fig. 5G-H) (Ivanov et al., 2007; Lee et al., 2007; Li et al., 2007; Mascle et al., 2007). SUMOylation is a highly dynamic, reversible process, and SUMOylation of even a small fraction of a given protein was shown to drastically influence a protein's cellular function (reviewed in Geiss-Friedlander and

Melchior, 2007 [\[1\]](#); Hay, 2005 [\[2\]](#)). We found that SUMO-deficient Bon loses its association with chromatin and mislocalizes into discrete nuclear foci (**Fig. 5E**, **5F**). SUMOylation has been implicated in the assembly of functional nuclear condensates, such as PML bodies (Ishov et al., 1999 [\[3\]](#); Muller, 1998 [\[4\]](#); Shen et al., 2006 [\[5\]](#); Zhong et al., 2000 [\[6\]](#)), yet Bon is more dispersed and associates with chromatin in its SUMOylated form and concentrates into foci in its unmodified form, suggesting that these foci might represent inactive Bon, which due to its lack of SUMOylation fails to form complexes with its partners such as SetDB1. Indeed, SUMOylation is known to enhance protein-protein interactions and promote efficient assembly of protein complexes during heterochromatin formation and transcriptional silencing (Gareau and Lima, 2010 [\[7\]](#); Gill, 2005 [\[8\]](#), 2004 [\[9\]](#); Jentsch and Psakhye, 2013 [\[10\]](#); Martin et al., 2007 [\[11\]](#); Shio and Eisenman, 2003 [\[12\]](#)).

Despite the universal role of SUMOylation in the repressive function of KAP-1 and Bon, there are crucial differences that suggests parallel and independent evolution in vertebrates and insects rather than conservation of an ancient mechanism. First, SUMOylation sites are not conserved between two groups (but conserved within each group) and in fact are located in different regions in Bon and KAP-1. Bon is SUMOylated at a single residue, K20, close to its N terminus (**Fig. 5C**) and this site seems to be conserved in other insects. In KAP-1 several SUMOylation sites are located in C-terminal bromodomain, the most prominent ones being K779 and K804 (Ivanov et al., 2007 [\[13\]](#)). Other mammalian TIF1 proteins are also SUMOylated at C-terminal region. SUMOylation of human TIF1 α at lysine residues K723 and K741 was proposed to play a role in regulating genes involved in cell adhesion pathways (Appikonda et al., 2018 [\[14\]](#)), and multiple SUMO-conjugations at lysines 776, 793, 796 and 839 of human TIF1 γ were reported to be required for the transcriptional repression of TGF β signaling (Fattet et al., 2013 [\[15\]](#)).

The second important difference is the molecular mechanism of SUMOylation of TIF1 members in insects and vertebrates. The PHD domain of mammalian KAP-1 functions as a SUMO E3-ligase to induce modification of the adjacent bromodomain (Ivanov et al., 2007 [\[13\]](#)). Importantly, evolutionary analysis of the TRIM/RBCC family suggest that the E3 SUMO-ligase activity has developed recently and only in a few specific members of this family and does not represent an ancient function of this protein family (Marín, 2012 [\[16\]](#)). In agreement with the evolutionary analysis, the lack of stable interaction between Bon and the E2-conjugating enzyme Ubc9 suggests that Bon does not have E3-ligase function (**Fig. 6A**, Extended Data Fig. 5A, 5B). Instead, we found that Bon is SUMOylated by the distinct SUMO E3-ligase Su(var)2-10, which belong to the PIAS family of SUMO ligases conserved among Metazoa (**Fig. 6B, C**). Importantly, Su(var)2-10 is required for the repressive function of Bon. Thus, both the specific sites and the molecular mechanisms of SUMOylation of TIF1 proteins are different in insects and vertebrates, suggesting that TIF1 SUMOylation developed independently during evolution in these two groups.

In conclusion, our study reveals an essential function of *Drosophila* TIF1 factor, Bon, in repression of tissue-specific genes in the germline and suggests that Bon SUMOylation at a single site by the SUMO E3-ligase Su(var)2-10 is critical for its role as a transcriptional repressor.

Acknowledgements

We thank members of the Aravin and Fejes Toth labs for discussion. We thank Peiwei Chen for suggesting some of the experiments. We appreciate the help of Anastasiya Grebin with the experiments. We are grateful to Julius Brennecke, Gregory Hannon, Albert Courey, the Bloomington Stock Center, and the Vienna *Drosophila* Resource Center for providing fly stocks, Hugo Bellen for providing antibodies. We thank Igor Antoshechkin (Millard and Muriel Jacobs Genetics and Genomics Laboratory, Caltech) for the help with sequencing, Giada Spigolon (Biological Imaging Facility, Caltech) for the help with microscopy, and Grace Shin for the help with HCR experiments. This work was supported by grants from the National Institutes of Health (R01 GM097363 to AA and R01 GM110217 to KFT) and by the HHMI Faculty Scholar Award to AAA.

Author contributions

Conceptualization: B.G, M.N., A.A.A. and K.F.T. Investigation: B.G. Methodology: B.G., M.N., K.F.T. and A.A.A. Formal analysis: B.G., M.N., A.A.A. and K.F.T. Visualization: B.G. Software: B.G. and M.N. Funding acquisition: K.F.T. and A.A.A. Writing – original draft: B.G. Writing – review & editing: B.G., M.N., K.F.T. and A.A.A.

Supplementary figures

Extended Data Figure 1.

A) Schematics of a *Drosophila* ovariole and the expression pattern of different Gal4 driver lines during developmental stages of oogenesis. **B)** Table shows the count of ovaries with indicated phenotypes after *nos-Gal4*; *UAS-Cas9* knockout of *bon*. **C)** Confocal images of germarium from flies with Bon GLKD driven by *nos-Gal4* and control siblings from the same cross that lack Bon shRNA stained for α -spectrin (green), vasa (red) and DAPI (blue). The numbers on the left represent different germarium categories observed: I: 2-3 GSC and no fusomes, II: >4 GSC, III: empty germarium (scale bar: 20 μ m). **D)** Graph showing the percentage of different germarium phenotypes for the indicated genotypes (n= 23 for control flies and n =36 for flies with Bon GLKD driven by *nos-Gal4*).

Extended Data Figure 2.

A) Scatterplots display changes in transposons expression upon Bon GLKD driven by *MT-Gal4* (left) or *nos-Gal4* (right) as determined by RNA-seq data (log2-transformed TPM values, n =3). Flies with *white* GLKD driven by *MT-Gal4* (left) or siblings from the same cross that lack Bon shRNA (right) were used as a control. **B)** Volcano plot shows fold changes in genes expression upon Bon GLKD driven by *MT-Gal4* in the ovary as determined by RNA-seq data (n =3). Flies with *white* GLKD driven by *MT-Gal4* were used as a control. Genes that change significantly (qval< 0.05, LRT test, sleuth (Pimentel et al., 2017 [DOI](#))) and >2-fold are highlighted, and *bon*, *Rbp6*, *ple* are labeled. Genes with infinite fold change values (zero counts in control ovaries) are not shown. **C)** Venn diagram of the number of significantly derepressed genes (qval< 0.05, LRT test, log2FC>1) upon Bon GLKD driven by *MT-Gal4* and *nos-Gal4*. **D)** Bar graph shows the relative expression of *CG34353* and *pst* (normalized to rp49 level) in control and Bon-depleted ovaries (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **E-F)** Confocal images of egg chambers show RNA *in situ* HCR detecting *bonus* and *CG34353* (**E**) and *pst* (**F**) mRNAs in flies with *MT-Gal4*>*Bon GLKD* and control siblings from the same cross that lack Bon shRNA (scale bar: 20 μ m).

Extended Data Figure 3.

A) Bar plot shows the Luciferase reporter expression (normalized to rp49 level) in ovaries upon tethering of λ N-GFP-Bonus or λ N-GFP control ovaries (RT-qPCR, dots correspond to 3 independent biological replicates; error bars indicate st. dev.). **B-C)** Western blot analysis of

immunoprecipitation experiment using GFP nanotrap beads from S2 cells co-expressing GFP-Bonus and Flag-tagged Mi-2 (**B**) or Flag-tagged Mep-1 (**C**). Lysates not expressing the GFP-Bonus were used as negative control.

Extended Data Figure 4.

A) Western blot analysis of GFP-Bonus SUMOylation in S2 cells (top) and ovaries (bottom). Total protein lysates from S2 cells co-expressing HA-SUMO and GFP-Bonus or from ovaries of flies co-expressing Flag-SUMO and λ N-GFP-Bonus were used to immunopurify Bon using anti-GFP nanotrap beads. S2 cells or flies not expressing GFP-tagged protein were used as negative control. **B)** Phylogenetic relationships between the 12 *Drosophila* species and some other insects (using iTOL). **C)** Western blot analysis of SUMOylation of GFP-tagged Bon and SUMO-deficient triple mutant 3KR expressed in S2 cells. Triple 3KR mutation completely abolished Bon SUMOylation. Total protein lysates from S2 cells co-expressing HA-SUMO and GFP-Bonus or GFP-Bonus[3KR] were immunopurified using anti-GFP nanotrap beads. Cell lysates not expressing GFP-tagged proteins were used as a negative control. **D)** Confocal images of egg chambers from wild-type Oregon-R flies stained for Bon (red), DAPI (blue) and HP1 (green) (scale bar: 20 μ m). Bottom images show isolated nurse cell nuclei. Bottom right represent fluorescence intensity analysis of overlayed signal (along the yellow line) performed by ImageJ software and expressed as arbitrary fluorescence units (AFU). **E-F)** Western blot analysis shows the SUMO-independent interaction between Bon and Mi-2 or Rpd3. Total protein lysates from S2 cells co-expressing GFP-Bonus (WT) or triple mutant GFP-Bonus[3KR] (3KR) and full-length or truncated Flag-Mi2 (**E**), and Flag-Rpd3 (**F**) were immunopurified using anti-GFP nanotrap beads. Lysates from cells not expressing GFP-tagged proteins were used as a negative control.

Extended Data Figure 5.

A, B, Western blot analysis of immunoprecipitation experiment using GFP nanotrap beads from S2 cells co-expressing Flag-tagged Ubc9 and GFP-Bonus (**A**) or Flag-tagged Ubc9 and GFP-Bonus or SUMO-deficient triple mutant 3KR GFP-Bonus (**B**). Lysates not expressing the GFP-tagged proteins were used as negative control.

Methods

Drosophila fly stocks

All fly stocks and crosses were raised at 24°C. 1-2 day old females were put on yeast for 1 day prior to dissection. Females from crosses with GLKD at early stages of oogenesis and respective control were 0-1 days old and were dissected right away. The following stocks were used: stocks with shRNAs targeting *Su(var)2-10* (shSv210, BDSC #32956), *Rpd3* (shRpd3, BDSC #33725), *Mi-2* (shMi-2, BDSC #35398) and *white* (shWhite, BDSC #33623) and *nos-Gal4;UAS-Cas9* (BDSC #54593) were obtained from the Bloomington Drosophila Stock Center. The fly line expressing sgRNAs targeting *bon* (VDRC #341851) was obtained from the Vienna Drosophila Resource Center. Fly lines UASp-mKate2-4xBoxB-K10polyA, UASp- λ N-GFP-eGFP control, UASp- λ N-GFP-Su(var)-10, shSmt3 were described previously (Chen et al., 2016 [\[1\]](#); Ninova et al., 2020a [\[2\]](#)). shSetDB1 was a gift from Julius Brennecke, the luciferase 8BoxB reporter and wild-type Oregon-R fly lines were a gift from Gregory Hannon, UASp-Flag-SUMO was a gift from Albert Courey. To obtain the shBonus fly lines, the short hairpin sequences (**Table 1** [\[3\]](#)) was ligated into the pValium20 vector (Ni et al., 2011 [\[4\]](#)) and then integrated into the attP2 landing site (BDSC #8622). To generate the UASp- λ N-GFP-Bonus and UASp- λ N-GFP-Bonus[3KR] fly lines, full length cDNA sequences of wild-type Bon or triple

mutant Bon, respectively, were cloned in vectors containing a miniwhite marker followed by the UASp promoter sequence, and λ N-GFP. Transgenic flies carrying these constructs were generated by phiC31 transformation by BestGene Inc and were integrated into the attP40 landing site ($y^1 w^{67c23}; P\{CaryP\}attP40$). The expression of constructs was driven by maternal alpha-tubulin67C-Gal4 (*MT-Gal4*) (BDSC #7063), *nos-Gal4* (BDSC #4937), or *bam-Gal4* (BDSC #80579) drivers.

Immunofluorescent microscopy and image processing

Seven to fifteen pairs of ovaries from *Drosophila* lines expressing UASP- λ N-GFP-Bonus and UASP- λ N-GFP-Bonus[3KR] under the control of the *MT-Gal4* driver were dissected in ice-cold PBS and then fixed in PBST solution (PBS, 0.1% Tween-20) supplemented with 4% formaldehyde for 20 min at room temperature with end-to-end rotation. Samples were washed three times 10 min with PBST and mounted in SlowFade® Gold antifade Mountant with DAPI. Seven to fifteen pairs of ovaries from *Drosophila* Oregon-R flies and lines with GLKD of Bon under the control of the *nos-Gal4* or *bam+nos* drivers were fixed in PBST supplemented with 4% formaldehyde for 20 min at room temperature with rotation and then washed three times 10 min with PBST. Fixed ovaries were incubated for 30 min with PBX (PBS, 0.1% Tween-20, 0.3% Triton X-100), and blocked in 5% normal goat serum (NGS) in PBX for 1 hour at room temperature. Samples were incubated with primary antibody in 3% NGS in PBX overnight at 4°C with rotation, followed by three washes in PBX solution for 10 min, and an overnight incubation with secondary antibody in 3% NGS in PBX at 4°C with rotation in the dark. After three washes in PBX, SlowFade® Gold antifade Mountant with DAPI was added to the samples. Confocal images were acquired with a Zeiss LSM 800 using a 63× oil immersion objective and were processed using Fiji. Primary antibodies to Vasa (rat, DSHB), to α -spectrin (mouse, 3A9 DSHB) and to Bon (a gift from Hugo Bellen) were used. Secondary antibodies were anti-mouse Alexa Fluor488, anti-rat Alexa Fluor546 (Invitrogen) and anti-guinea pig Cy3 (Jackson ImmunoResearch Inc.).

RNA *in situ* hybridization chain reaction (HCR)

For RNA *in situ* HCR, probes, amplifiers, and buffers were purchased from Molecular Instruments (molecularinstruments.org) (<http://molecularinstruments.org/>) [↗](#) for *bon* (unique identifier: 4165/E324), *ple* (unique identifier: 4324/E516), *Rbp6* (unique identifier: 4408/E662), *pst* (unique identifier: 4408/E660) and *CG34353* (unique identifier: 4408/E658) transcripts. RNA *in situ* HCR v3.0 was performed according to manufacturer's instructions for generic samples in solution.

S2 cell line

Drosophila S2 cells (DGRC catalog #006) were cultured at 25°C in Schneider's *Drosophila* Medium supplemented with 10% heat-inactivated FBS and 1X Penicillin-Streptomycin.

Protein co-immunoprecipitation from S2 cells

S2 cells were transfected with plasmids encoding HA-, GFP- and FLAG-tagged proteins under the control of the Actin promoter using TransIT-LT1 reagent (Mirus). 24–40 h after transfection, cells were collected and resuspended in lysis buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 0.2% NP-40, 0.2% Triton-X100, 5% glycerol, 20 mM N-Ethylmaleimide (NEM) (Sigma) and Complete Protease Inhibitor Cocktail (Roche)). The cell lysate was incubated on ice for 30 min, centrifuged and the supernatant collected. The supernatant was incubated with magnetic agarose GFP-Trap beads (Chromotek) for 2 h at 4°C with end-to-end rotation. Beads were washed three-four times for 10 min with wash buffer (20 mM Tris-HCl pH 7.4, 0.1% NP40, 150 mM NaCl) and boiled in 2xLaemmli buffer for 5 min at 95°C. The eluate was used for western blot analysis. For detection of SUMO-modified Bon, cells were lysed in RIPA-like buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% NP-40, 0.5% Sodium deoxycholate, 0.1% SDS, 20 mM NEM, cOmplete Protease Inhibitor Cocktail), and washed in high salt wash buffer (20 mM Tris-HCl pH 7.4, 500 mM NaCl, 1% NP40, 0.5% Sodium deoxycholate, 0.5% SDS, 20 mM NEM, cOmplete Protease Inhibitor Cocktail).

Table 1.

Purpose	Name	Sequence (5' to 3')
TRiP Short Hairpin	shBonus	CACACTCGTTTCCGAAATCAA
Oligos for RT-qPCR and ChIP-qPCR	bon-for	ACTTCTGGGTCTGACTGGCGAAG
	bon-rev	TCAACGCACCACGACGTGG
	rp49-for	CCGCTTCAAGGGACAGTATCTG
	rp49-rev	ATCTCGCCGCAGTAAACGC
	ple-for	AGCAGACCAAACAAACCGTCCTC
	ple-rev	CACATCTTGCTCGGAGGGAAT
	rbp6-for	CGCACCTCCCTATCTATCTATCT
	rbp6-rev	GTGTTCTTTTTTGCATTTTCTGTGCTT
	CG34353-for	TACCGTTTTTTTGGCACCTACGC
	CG34353-rev	GATGGTGCTAGAATCTTGCCC
	pst-for	GCAGCAAGCTATTTCGATCGCAG
	pst-rev	CTGCCAATCCCACAGTGCCAC
	mKate_reporter-for	TCAGAGGGGTGAACTTCCCA
	mKate_reporter-rev	CTCCCAGCCGAGTGTTTTCT
	luc_reporter-for	TGATTATGTCCGGCTACGTGAAT
	luc_reporter-rev	GGTCCACGATGAAGAAGTGCTC
	CG3191-for	CGAAGGAGAGGTGCAGATATTC
	CG3191-rev	GCTGGAGGAACTTGATGTAGG
	Spn88Eb-for	CGAGCGTGAGCAGGAAAT
	Spn88Eb-rev	GACTTGTATATGGTGCGGTAGG

Protein co-immunoprecipitation from fly ovaries

For detection of SUMO-modified Bon, 70-90 pairs of dissected ovaries co-expressing Flag-tagged SUMO and λ N-GFP-Bonus or λ N-GFP-Bonus[3KR] under the control of the *MT-Gal4* driver were lysed and dounced in 500 μ l lysis buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 0.4% NP40, 10% glycerol, 20 mM NEM and cOmplete Protease Inhibitor Cocktail (Roche)). The ovary lysate was incubated on ice for 30 min, then centrifuged and the supernatant collected. The supernatant was incubated with magnetic agarose GFP-Trap beads (Chromotek) for 2-3 h at 4°C with end-to-end rotation. Beads were washed 4 times $\geq$ 10 min at 4°C with ovary high salt wash buffer (20 mM Tris-HCl pH 7.4, 500 mM NaCl, 1% NP40, 0.5% Sodium deoxycholate, 0.5% SDS, 20 mM NEM, cOmplete Protease Inhibitor Cocktail). Washed beads were further in 2xLaemmli buffer for 5 min at 95°C, and the eluate was analyzed by western blotting.

Ovary fractionation

For whole ovaries and subcellular compartment extraction, 30-40 pairs of fly ovaries expressing λ N-GFP-Bonus or λ N-GFP-Bonus[3KR] were lysed and dounced in ice-cold “AT” buffer (15 mM HEPES-NaOH pH 7.6, 10 mM NaCl, 5 mM MgOAc, 3mM CaCl₂, 300 mM sucrose, 0.1% Triton X-100, 1 mM DTT, cOmplete Protease Inhibitor Cocktail (Roche)). A small fraction of lysate was saved as whole cell lysate for western blot analysis. For subcellular fractionation 2 volumes of buffer “B” (15 mM HEPES-NaOH pH 7.6, 10 mM NaCl, 5 mM MgOAc, 3mM CaCl₂, 1 M sucrose, 1 mM DTT, cOmplete Protease Inhibitor Cocktail (Roche)) were added to the lysate. The lysate was incubated for 5 min on ice and then centrifuged at 5900g for 15 min at 4°C. The supernatant was transferred to a new tube, centrifuged at 19000g for 10 min 4°C, and saved as cytoplasmic fraction for western blot analysis. The cell pellet was resuspended in “E2” buffer (10 mM Tris-HCl pH 7.5, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA and cOmplete Protease Inhibitor Cocktail (Roche)) and centrifuged at 1500g for 2 min at 4°C. The supernatant was transferred to a new tube, centrifuged at 19000g for 10 min 4°C, and saved as nuclear fraction for western blot analysis. The pellet was once washed in E2 buffer, then resuspended in E2 buffer and incubated for 10 min at 4°C, followed by centrifugation at 1500g for 2 min at 4°C. For chromatin extraction, the pellet was resuspended in “E3” buffer (500 mM Tris-HCl pH 7.5, 500 mM NaCl, cOmplete Protease Inhibitor Cocktail (Roche)) and then sonicated for 5 min at high setting with 30sON/30sOFF in a Bioruptor sonicator (Diagenode) and centrifuged for 5 min at 19000g at 4°C. The supernatant was saved as chromatin fraction for western blot analysis. Protein concentration of all fractions was measured and further all saved fractions were boiled with 1x final concentration of Laemmli buffer for 5 min at 95°C and analyzed by western blotting.

Western blotting

Proteins were separated by SDS-PAGE gel electrophoresis and transferred to a 0.45- μ m nitrocellulose membrane (Bio-Rad) according to standard procedures. The membrane was blocked with 5% milk or with 0.2% I-block (Invitrogen) in PBST (PBS, 0.1% Tween-20) for 1 hour. The membrane was incubated with primary antibodies for 2 hours at room temperature or overnight at 4°C, followed by 3x washes for 5 min in PBST and incubation with secondary antibodies for 1 hour at room temperature. The membrane was washed three times for 5 min with PBST and then imaged with Odyssey system (Li-Cor). When primary antibody was HRP-conjugated, the membrane was washed 3x 5 min with PBST, incubated with the HRP substrate, and X-ray film developed on an X-Ray Film Processor (Konica Minolta). The following antibodies were used: HRP-conjugated anti-FLAG (Sigma, A8592), mouse anti-FLAG (Sigma, F1804), rabbit polyclonal anti-GFP (Chen et al., 2016 [DOI](#)), IRDye anti-rabbit and anti-mouse secondary antibodies (Li-Cor, #925-68070 and #925-32211).

RNA extraction and RT-qPCR

For RNA extraction 10-20 pairs of dissected ovaries were homogenized in TRIzol (Invitrogen), RNA extracted, isopropanol precipitated, and treated with DNaseI (Invitrogen) according to the manufacturer's instructions. Reverse transcription was performed using random hexamer oligonucleotides with Superscript III Reverse Transcriptase (Invitrogen). qPCR was performed on a Mastercycler[®]ep realplex PCR machine (Eppendorf). 3 biological replicates per genotype were used for all RT-qPCR experiments. Target expression was normalized to rp49 mRNA expression. The data were visualized using Python 3 via JupyterLab. Primers used for qPCR analysis are listed in [Table 1](#).

RNA-seq and RNA-seq analysis

For RNA-seq libraries, total RNA was extracted from fly ovaries using TRIzol reagent. PolyA+ selection was performed using an NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB, #E7490) for total RNA from lines with Bon GLKD driven by *nos-Gal4* (and matched siblings that lack the shRNA as control). Total RNA from line with Bon GLKD driven by *MT-Gal4* (and control flies that express an shRNA against the *white* gene) was depleted of ribosomal RNA with the Zymo-Seq RiboFree Total RNA Library Kit (Zymo Research, #R3000). RNA-seq libraries were made using the NEBNext Ultra II Directional RNA Library Prep kit for Illumina (NEB, #E7760) according to the manufacturer's instructions. Libraries were sequenced on the Illumina HiSeq 2500 platform. To quantify expression level of protein-coding genes and TEs, RNA-seq libraries were pseudoaligned to the *D. melanogaster* transcriptome (RefSeq, dm6) and transposon consensus (from RepBase, [Jurka et al., 2005](#)), using kallisto ([Bray et al., 2016](#)). Differential expression analysis was done with sleuth using the gene analysis option ([Pimentel et al., 2017](#)). The average TPM between three biological replicates in knockdown versus control were calculated for fold changes in gene expression. For RNA-seq coverage tracks reads first were aligned to the *D. melanogaster* genome (dm6) using bowtie1 (v.1.2.2) allowing 2 mismatches and single mapping position. Tracks were generated using deepTools (v.3.5.1) bamCoverage function with 10 bp bin sizes.

Gene ontology (GO) biological-process term enrichment analysis was performed on the genes that were significantly de-repressed upon Bon GLKD driven by *nos-Gal4* ($\log_2\text{FC} > 1$, $q\text{-val} < 0.05$, LRT test, sleuth ([Pimentel et al., 2017](#))), using DAVID Bioinformatics Resources and all *Drosophila* genes that were not filtered out by sleuth as background. The enriched GO terms associated with 2 or less submitted genes were excluded. A significant threshold was applied using a multiple testing correction (Fisher's Exact test p-value < 0.01). The data visualization was performed using the "ggplot2" R package.

ChIP-qPCR and ChIP-seq

ChIP experiments were performed in two biological replicates as previously described ([Le Thomas et al., 2014](#)). In brief, 80-120 pairs of dissected ovaries were crosslinked with 1% formaldehyde in PBS for 10 min at room temperature, then quenched with Glycine (final concentration 25 mM). Frozen ovaries were dounced in RIPA buffer and then sonicated (Bioruptor sonicator) to a desired fragment sizes of 200–800 bp. Lysates were centrifuged at 19000g, and supernatants collected. The supernatants were first precleared for 2 h at 4°C using Protein G Dynabeads (Invitrogen). Precleared samples were immunoprecipitated with anti-H3K9me3 (abcam, ab8898) antibodies for 3-5 h at 4°C, then 50 μl Protein G Dynabeads were added, and samples were further incubated overnight at 4°C. Beads were washed 3x 10 minutes in LiCL buffer, followed by proteinase K treatment for 2 h at 55°C and then overnight at 65°C. DNA was extracted by standard phenol-chloroform extraction. ChIP-qPCR was performed on the Mastercycler[®]ep realplex PCR machine (Eppendorf). All ChIPs were normalized to respective inputs and to control region rp49. The data were visualized using Python 3 via JupyterLab. Primers used for qPCR analysis are listed in [Table 1](#).

ChIP-seq libraries were prepared using NEBNext Ultra DNA Library Prep Kit Illumina (NEB) and sequenced on the Illumina HiSeq 2500 platform (PE 50bp). After removal of the adaptors, reads with a minimal length of 18 nucleotides were aligned to the *D. melanogaster* genome (dm6) using bowtie1 (v.1.2.2) allowing 2 mismatches and single mapping position. Genome coverage tracks were generated using deepTools (v.3.5.1) bamCoverage function with 10 bp bin sizes. ChIP signal was normalized to input counts by calculating cpm (counts per million) using the deepTools bamCompare function with 50 bp bin sizes (ChIP/Input). Heatmaps were generated with deepTools plotHeatmap using normalized (ChIP/Input) BigWig files.

Computational analysis

Tissue specificity identification

To classify upregulated genes upon Bon GLKD driven by *nos-Gal4* ($\log_2FC > 1$, $qval < 0.05$, LRT test, sleuth (Pimentel et al., 2017 [DOI](#))), according to the tissues they are normally expressed in, we used RPKM values from the modENCODE anatomy RNA-Seq dataset. The expression levels according to RPKM values from modENCODE anatomy RNA-seq dataset are no expression (0-0), very low (1-3), low (4-10), moderate (11-25), moderate high (26-50), high (51-100), very high (101-1000), extremely high (>1000). The analysis of enrichment in each tissue was calculated as the number of genes expressed at a certain expression level to the total number of provided genes. The data were visualized using Python 3 via JupyterLab.

Phylogenetic analysis of Bon

The *D. melanogaster* Bon protein sequence was used to BLAST against the National Center for Biotechnology Information (NCBI) nonredundant protein database with the Position-Specific Iterated BLAST (PSI-BLAST) program. Orthologs of Bon in other *Drosophila* species and some insects were identified based on high sequence similarity. Multiple sequence alignment was performed using the ClustalW program. The species distribution of the orthologs was visualized using SeaView v5.0.5. Phylogenetic tree was built with ClustalW and iTOL v6 software.

References

- Allton K., Jain A.K., Herz H.-M., Tsai W.-W., Jung S.Y., Qin J., Bergmann A., Johnson R.L., Barton M.C. (2009) **Trim24 targets endogenous p53 for degradation** *Proc. Natl. Acad. Sci. U.S.A.* **106**:11612–11616 <https://doi.org/10.1073/pnas.0813177106>
- Appikonda S., Thakkar K.N., Shah P.K., Dent S.Y.R., Andersen J.N., Barton M.C (2018) **Cross-talk between chromatin acetylation and SUMOylation of tripartite motif-containing protein 24 (TRIM24) impacts cell adhesion** *Journal of Biological Chemistry* **293**:7476–7485 <https://doi.org/10.1074/jbc.RA118.002233>
- Bai X., Kim J., Yang Z., Jurynek M.J., Akie T.E., Lee J., LeBlanc J., Sessa A., Jiang H., DiBiase A., Zhou Y., Grunwald D.J., Lin S., Cantor A.B., Orkin S.H., Zon L.I (2010) **TIF1y Controls Erythroid Cell Fate by Regulating Transcription Elongation** *Cell* **142**:133–143 <https://doi.org/10.1016/j.cell.2010.05.028>
- Beckstead R., Ortiz J.A., Sanchez C., Prokopenko S.N., Chambon P., Losson R., Bellen H.J (2001) **Bonus, a Drosophila homolog of TIF1 proteins, interacts with nuclear receptors and can inhibit betaFTZ-F1-dependent transcription** *Mol Cell* **7**:753–765 [https://doi.org/10.1016/s1097-2765\(01\)00220-9](https://doi.org/10.1016/s1097-2765(01)00220-9)
- Beckstead R.B., Ner S.S., Hales K.G., Grigliatti T.A., Baker B.S., Bellen H.J (2005) **Bonus, a Drosophila TIF1 homolog, is a chromatin-associated protein that acts as a modifier of position-effect variegation** *Genetics* **169**:783–794 <https://doi.org/10.1534/genetics.104.037085>
- Berger S.L (2007) **The complex language of chromatin regulation during transcription** *Nature* **447**:407–412 <https://doi.org/10.1038/nature05915>
- Bouazoune K., Brehm A (2006) **ATP-dependent chromatin remodeling complexes in Drosophila** *Chromosome Res* **14**:433–449 <https://doi.org/10.1007/s10577-006-1067-0>
- Bray N.L., Pimentel H., Melsted P., Pachter L (2016) **Near-optimal probabilistic RNA-seq quantification** *Nat Biotechnol* **34**:525–527 <https://doi.org/10.1038/nbt.3519>
- Brehm A (2000) **dMi-2 and ISWI chromatin remodelling factors have distinct nucleosome binding and mobilization properties** *The EMBO Journal* **19**:4332–4341 <https://doi.org/10.1093/emboj/19.16.4332>
- Cammas F., Herzog M., Lerouge T., Chambon P., Losson R (2004) **Association of the transcriptional corepressor TIF1beta with heterochromatin protein 1 (HP1): an essential role for progression through differentiation** *Genes Dev* **18**:2147–2160 <https://doi.org/10.1101/gad.302904>
- Cammas F., Mark M., Dollé P., Dierich A., Chambon P., Losson R (2000) **Mice lacking the transcriptional corepressor TIF1β are defective in early postimplantation development** *Development* **127**:2955–2963 <https://doi.org/10.1242/dev.127.13.2955>
- Chen D., McKearin D.M (2003) **A discrete transcriptional silencer in the bam gene determines asymmetric division of the Drosophila germline stem cell** *Development* **130**:1159–1170 <https://doi.org/10.1242/dev.00325>

- Chen Y.-C.A., Stuwe E., Luo Y., Ninova M., Le Thomas A., Rozhavska E., Li S., Vempati S., Laver J.D., Patel D.J., Smibert C.A., Lipshitz H.D., Fejes Toth K., Aravin A.A. (2016) **Cutoff Suppresses RNA Polymerase II Termination to Ensure Expression of piRNA Precursors** *Molecular Cell* **63**:97–109 <https://doi.org/10.1016/j.molcel.2016.05.010>
- Chintapalli V.R., Wang J., Dow J.A.T (2007) **Using FlyAtlas to identify better Drosophila melanogaster models of human disease** *Nat Genet* **39**:715–720 <https://doi.org/10.1038/ng2049>
- Chung H.-R., Löhr U., Jäckle H (2007) **Lineage-specific expansion of the Zinc Finger Associated Domain ZAD** *Molecular Biology and Evolution* **24**:1934–1943 <https://doi.org/10.1093/molbev/msm121>
- Chung H.-R., Schäfer U., Jäckle H., Böhm S (2002) **Genomic expansion and clustering of ZAD-containing C2H2 zinc-finger genes in Drosophila** *EMBO Rep* **3**:1158–1162 <https://doi.org/10.1093/embo-reports/kvf243>
- De Gregorio E. (1999) **Translation driven by an eIF4G core domain in vivo** *The EMBO Journal* **18**:4865–4874 <https://doi.org/10.1093/emboj/18.17.4865>
- Doren M.V., Williamson A.L., Lehmann R (1998) **Regulation of zygotic gene expression in Drosophila primordial germ cells** *Current Biology* **8**:243–246 [https://doi.org/10.1016/S0960-9822\(98\)70091-0](https://doi.org/10.1016/S0960-9822(98)70091-0)
- Fattat L., Ay A.-S., Bonneau B., Jallades L., Mikaelian I., Treilleux I., Gillet G., Hesling C., Rimokh R (2013) **TIF1y requires sumoylation to exert its repressive activity on TGFβ signaling** *Journal of Cell Science jcs* **126**:748 <https://doi.org/10.1242/jcs.126748>
- Friedman J.R., Fredericks W.J., Jensen D.E., Speicher D.W., Huang X.P., Neilson E.G., Rauscher F.J (1996) **KAP-1, a novel corepressor for the highly conserved KRAB repression domain** *Genes Dev* **10**:2067–2078 <https://doi.org/10.1101/gad.10.16.2067>
- Gancz D., Lengil T., Gilboa L (2011) **Coordinated Regulation of Niche and Stem Cell Precursors by Hormonal Signaling** *PLoS Biol* **9** <https://doi.org/10.1371/journal.pbio.1001202>
- Gareau J.R., Lima C.D (2010) **The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition** *Nat Rev Mol Cell Biol* **11**:861–871 <https://doi.org/10.1038/nrm3011>
- Garvin A.J., Morris J.R (2017) **SUMO, a small, but powerful, regulator of double-strand break repair** *Phil. Trans. R. Soc. B* **372** <https://doi.org/10.1098/rstb.2016.0281>
- Geiss-Friedlander R., Melchior F (2007) **Concepts in sumoylation: a decade on** *Nat Rev Mol Cell Biol* **8**:947–956 <https://doi.org/10.1038/nrm2293>
- Gill G (2005) **Something about SUMO inhibits transcription** *Curr Opin Genet Dev* **15**:536–541 <https://doi.org/10.1016/j.gde.2005.07.004>
- Gill G (2004) **SUMO and ubiquitin in the nucleus: different functions, similar mechanisms?** *Genes Dev* **18**:2046–2059 <https://doi.org/10.1101/gad.1214604>
- Hay R.T (2005) **SUMO** *Molecular Cell* **18**:1–12 <https://doi.org/10.1016/j.molcel.2005.03.012>

- Hodin J., Riddiford L.M (1998) **The ecdysone receptor and ultraspiracle regulate the timing and progression of ovarian morphogenesis during *Drosophila* metamorphosis** *Development Genes and Evolution* **208**:304–317 <https://doi.org/10.1007/s004270050186>
- Hsu T.-H., Yang C.-Y., Yeh T.-H., Huang Y.-C., Wang T.-W., Yu J.-Y (2017) **The Hippo pathway acts downstream of the Hedgehog signaling to regulate follicle stem cell maintenance in the *Drosophila* ovary** *Sci Rep* **7** <https://doi.org/10.1038/s41598-017-04052-6>
- Ishov A.M., Sotnikov A.G., Negorev D., Vladimirova O.V., Neff N., Kamitani T., Yeh E.T.H., Strauss J.F., Maul G.G (1999) **Pml Is Critical for Nd10 Formation and Recruits the Pml-Interacting Protein Daxx to This Nuclear Structure When Modified by Sumo-1** *Journal of Cell Biology* **147**:221–234 <https://doi.org/10.1083/jcb.147.2.221>
- Ito H., Sato K., Koganezawa M., Ote M., Matsumoto K., Hama C., Yamamoto D (2012) **Fruitless Recruits Two Antagonistic Chromatin Factors to Establish Single-Neuron Sexual Dimorphism** *Cell* **149**:1327–1338 <https://doi.org/10.1016/j.cell.2012.04.025>
- Ivanov A.V., Peng H., Yurchenko V., Yap K.L., Negorev D.G., Schultz D.C., Psulkowski E., Fredericks W.J., White D.E., Maul G.G., Sadofsky M.J., Zhou M.-M., Rauscher F.J (2007) **PHD domain-mediated E3 ligase activity directs intramolecular sumoylation of an adjacent bromodomain required for gene silencing** *Mol Cell* **28**:823–837 <https://doi.org/10.1016/j.molcel.2007.11.012>
- Jentsch S., Psakhye I (2013) **Control of Nuclear Activities by Substrate-Selective and Protein-Group SUMOylation** *Annu. Rev. Genet* **47**:167–186 <https://doi.org/10.1146/annurev-genet-111212-133453>
- Johnson E.S., Gupta A.A (2001) **An E3-like factor that promotes SUMO conjugation to the yeast septins** *Cell* **106**:735–744 [https://doi.org/10.1016/s0092-8674\(01\)00491-3](https://doi.org/10.1016/s0092-8674(01)00491-3)
- Jurka J., Kapitonov V.V., Pavlicek A., Klonowski P., Kohany O., Walichiewicz J (2005) **Repbase Update, a database of eukaryotic repetitive elements** *Cytogenet Genome Res* **110**:462–467 <https://doi.org/10.1159/000084979>
- Khetchoumian K., Teletin M., Mark M., Lerouge T., Cerviño M., Oulad-Abdelghani M., Chambon P., Losson R (2004) **TIF1δ, a Novel HP1-interacting Member of the Transcriptional Intermediary Factor 1 (TIF1) Family Expressed by Elongating Spermatids** *Journal of Biological Chemistry* **279**:48329–48341 <https://doi.org/10.1074/jbc.M404779200>
- Kimura K.-I., Ote M., Tazawa T., Yamamoto D (2005) **Fruitless specifies sexually dimorphic neural circuitry in the *Drosophila* brain** *Nature* **438**:229–233 <https://doi.org/10.1038/nature04229>
- König A., Yatsenko A.S., Weiss M., Shcherbata H.R (2011) **Ecdysteroids affect *Drosophila* ovarian stem cell niche formation and early germline differentiation: Steroids in stem cells and their niches** *The EMBO Journal* **30**:1549–1562 <https://doi.org/10.1038/emboj.2011.73>
- Kouzarides T (2007) **Chromatin Modifications and Their Function** *Cell* **128**:693–705 <https://doi.org/10.1016/j.cell.2007.02.005>

- Kulkarni A., Oza J., Yao M., Sohail H., Ginjala V., Tomas-Loba A., Horejsi Z., Tan A.R., Boulton S.J., Ganesan S (2013) **Tripartite Motif-containing 33 (TRIM33) Protein Functions in the Poly(ADP-ribose) Polymerase (PARP)-dependent DNA Damage Response through Interaction with Amplified in Liver Cancer 1 (ALC1) Protein** *Journal of Biological Chemistry* **288**:32357–32369 <https://doi.org/10.1074/jbc.M113.459164>
- Kunert N., Brehm A (2009) **Novel Mi-2 related ATP-dependent chromatin remodelers** *Epigenetics* **4**:209–211 <https://doi.org/10.4161/epi.8933>
- Kunert N., Wagner E., Murawska M., Klinker H., Kremmer E., Brehm A (2009) **dMec: a novel Mi-2 chromatin remodelling complex involved in transcriptional repression** *EMBO J* **28**:533–544 <https://doi.org/10.1038/emboj.2009.3>
- Le Douarin B., Nielsen A.L., Garnier J.M., Ichinose H., Jeanmougin F., Losson R., Chambon P. (1996) **A possible involvement of TIF1 alpha and TIF1 beta in the epigenetic control of transcription by nuclear receptors** *EMBO J* **15**:6701–6715
- Le Thomas A., Stuwe E., Li S., Du J., Marinov G., Rozhkov N., Chen Y.-C.A., Luo Y., Sachidanandam R., Toth K.F., Patel D., Aravin A.A. (2014) **Transgenerationally inherited piRNAs trigger piRNA biogenesis by changing the chromatin of piRNA clusters and inducing precursor processing** *Genes Dev* **28**:1667–1680 <https://doi.org/10.1101/gad.245514.114>
- Lee S.-J., Lee J.-R., Hah H.-S., Kim Y.-H., Ahn J.-H., Bae C.-D., Yang J.-M., Hahn M.-J (2007) **PIAS1 interacts with the KRAB zinc finger protein, ZNF133, via zinc finger motifs and regulates its transcriptional activity** *Exp Mol Med* **39**:450–457 <https://doi.org/10.1038/emm.2007.49>
- Li X., Lee Y.-K., Jeng J.-C., Yen Y., Schultz D.C., Shih H.-M., Ann D.K (2007) **Role for KAP1 serine 824 phosphorylation and sumoylation/desumoylation switch in regulating KAP1-mediated transcriptional repression** *J Biol Chem* **282**:36177–36189 <https://doi.org/10.1074/jbc.M706912200>
- Margolin J.F., Friedman J.R., Meyer W.K., Vissing H., Thiesen H.J., Rauscher F.J. (1994) **Krüppel-associated boxes are potent transcriptional repression domains** *Proc. Natl. Acad. Sci. U.S.A.* **91**:4509–4513 <https://doi.org/10.1073/pnas.91.10.4509>
- Marín I (2012) **Origin and Diversification of TRIM Ubiquitin Ligases** *PLoS ONE* **7** <https://doi.org/10.1371/journal.pone.0050030>
- Martin S., Wilkinson K.A., Nishimune A., Henley J.M (2007) **Emerging extranuclear roles of protein SUMOylation in neuronal function and dysfunction** *Nat Rev Neurosci* **8**:948–959 <https://doi.org/10.1038/nrn2276>
- Masclé X.H., Germain-Desprez D., Huynh P., Estephan P., Aubry M (2007) **Sumoylation of the Transcriptional Intermediary Factor 1β (TIF1β), the Co-repressor of the KRAB Multifinger Proteins, Is Required for Its Transcriptional Activity and Is Modulated by the KRAB Domain** *Journal of Biological Chemistry* **282**:10190–10202 <https://doi.org/10.1074/jbc.M611429200>
- McKearin D., Ohlstein B (1995) **A role for the Drosophila bag-of-marbles protein in the differentiation of cystoblasts from germline stem cells** *Development* **121**:2937–2947 <https://doi.org/10.1242/dev.121.9.2937>

- Melchior F (2003) **SUMO: ligases, isopeptidases and nuclear pores** *Trends in Biochemical Sciences* **28**:612–618 <https://doi.org/10.1016/j.tibs.2003.09.002>
- Muller S (1998) **Conjugation with the ubiquitin-related modifier SUMO-1 regulates the partitioning of PML within the nucleus** *The EMBO Journal* **17**:61–70 <https://doi.org/10.1093/emboj/17.1.61>
- Nakayama J., Rice J.C., Strahl B.D., Allis C.D., Grewal S.I (2001) **Role of histone H3 lysine 9 methylation in epigenetic control of heterochromatin assembly** *Science* **292**:110–113 <https://doi.org/10.1126/science.1060118>
- Ni J.-Q., Zhou R., Czech B., Liu L.-P., Holderbaum L., Yang-Zhou D., Shim H.-S., Tao R., Handler D., Karpowicz P., Binari R., Booker M., Brennecke J., Perkins L.A., Hannon G.J., Perrimon N (2011) **A genome-scale shRNA resource for transgenic RNAi in Drosophila** *Nat Methods* **8**:405–407 <https://doi.org/10.1038/nmeth.1592>
- Nielsen A.L., Ortiz J.A., You J., Oulad-Abdelghani M., Khechumian R., Gansmuller A., Chambon P., Losson R (1999) **Interaction with members of the heterochromatin protein 1 (HP1) family and histone deacetylation are differentially involved in transcriptional silencing by members of the TIF1 family** *EMBO J* **18**:6385–6395 <https://doi.org/10.1093/emboj/18.22.6385>
- Ninova M., Chen Y.-C.A., Godneeva B., Rogers A.K., Luo Y., Fejes Tóth K., Aravin A.A (2020) **Su(var)2-10 and the SUMO Pathway Link piRNA-Guided Target Recognition to Chromatin Silencing** *Molecular Cell* **77**:556–570 <https://doi.org/10.1016/j.molcel.2019.11.012>
- Ninova M., Godneeva B., Chen Y.-C.A., Luo Y., Prakash S.J., Jankovics F., Erdélyi M., Aravin A.A., Fejes Tóth K (2020) **The SUMO Ligase Su(var)2-10 Controls Hetero- and Euchromatic Gene Expression via Establishing H3K9 Trimethylation and Negative Feedback Regulation** *Molecular Cell* **77**:571–585 <https://doi.org/10.1016/j.molcel.2019.09.033>
- Ozato K., Shin D.-M., Chang T.-H., Morse H.C (2008) **TRIM family proteins and their emerging roles in innate immunity** *Nat Rev Immunol* **8**:849–860 <https://doi.org/10.1038/nri2413>
- Pengue G., Calabró V., Bartoli P.C., Pagliuca A., Lania L (1994) **Repression of transcriptional activity at a distance by the evolutionary conserved KRAB domain present in a subfamily of zinc finger proteins** *Nucl Acids Res* **22**:2908–2914 <https://doi.org/10.1093/nar/22.15.2908>
- Pichler A., Gast A., Seeler J.S., Dejean A., Melchior F (2002) **The Nucleoporin RanBP2 Has SUMO1 E3 Ligase Activity** *Cell* **108**:109–120 [https://doi.org/10.1016/S0092-8674\(01\)00633-X](https://doi.org/10.1016/S0092-8674(01)00633-X)
- Pimentel H., Bray N.L., Puente S., Melsted P., Pachter L (2017) **Differential analysis of RNA-seq incorporating quantification uncertainty** *Nat Methods* **14**:687–690 <https://doi.org/10.1038/nmeth.4324>
- Port F., Strein C., Stricker M., Rauscher B., Heigwer F., Zhou J., Beyersdörffer C., Frei J., Hess A., Kern K., Lange L., Langner N., Malamud R., Pavlović B., Räddecke K., Schmitt L., Voos L., Valentini E., Boutros M (2020) **A large-scale resource for tissue-specific CRISPR mutagenesis in Drosophila** *eLife* **9** <https://doi.org/10.7554/eLife.53865>
- Rea S., Eisenhaber F., O’Carroll D., Strahl B.D., Sun Z.W., Schmid M., Opravil S., Mechtler K., Ponting C.P., Allis C.D., Jenuwein T (2000) **Regulation of chromatin structure by site-specific histone H3 methyltransferases** *Nature* **406**:593–599 <https://doi.org/10.1038/35020506>

- Reverter D., Lima C.D (2005) **Insights into E3 ligase activity revealed by a SUMO–RanGAP1–Ubc9–Nup358 complex** *Nature* **435**:687–692 <https://doi.org/10.1038/nature03588>
- Rubertis F.D., Kadosh D., Henchoz S., Pauli D., Reuter G., Struhl K., Spierer P (1996) **The histone deacetylase RPD3 counteracts genomic silencing in Drosophila and yeast** *Nature* **384**:589–591 <https://doi.org/10.1038/384589a0>
- Salzberg A., Prokopenko S.N., He Y., Tsai P., Pál M., Maróy P., Glover D.M., Deák P., Bellen H.J (1997) **P-element insertion alleles of essential genes on the third chromosome of Drosophila melanogaster: mutations affecting embryonic PNS development** *Genetics* **147**:1723–1741
- Schultz D.C., Ayyanathan K., Negorev D., Maul G.G., Rauscher F.J (2002) **SETDB1: a novel KAP-1-associated histone H3, lysine 9-specific methyltransferase that contributes to HP1-mediated silencing of euchromatic genes by KRAB zinc-finger proteins** *Genes Dev* **16**:919–932 <https://doi.org/10.1101/gad.973302>
- Schultz D.C., Friedman J.R., Rauscher F.J (2001) **Targeting histone deacetylase complexes via KRAB-zinc finger proteins: the PHD and bromodomains of KAP-1 form a cooperative unit that recruits a novel isoform of the Mi-2alpha subunit of NuRD** *Genes Dev* **15**:428–443 <https://doi.org/10.1101/gad.869501>
- Sedgwick G.G., Townsend K., Martin A., Shimwell N.J., Grand R.J.A., Stewart G.S., Nilsson J., Turnell A.S (2013) **Transcriptional intermediary factor 1y binds to the anaphase-promoting complex/cyclosome and promotes mitosis** *Oncogene* **32**:4622–4633 <https://doi.org/10.1038/onc.2012.501>
- Shen T.H., Lin H.-K., Scaglioni P.P., Yung T.M., Pandolfi P.P (2006) **The Mechanisms of PML–Nuclear Body Formation** *Molecular Cell* **24**:331–339 <https://doi.org/10.1016/j.molcel.2006.09.013>
- Shiio Y., Eisenman R.N. (2003) **Histone sumoylation is associated with transcriptional repression** *Proc. Natl. Acad. Sci. U.S.A.* **100**:13225–13230 <https://doi.org/10.1073/pnas.1735528100>
- Tong J.K., Hassig C.A., Schnitzler G.R., Kingston R.E., Schreiber S.L (1998) **Chromatin deacetylation by an ATP-dependent nucleosome remodelling complex** *Nature* **395**:917–921 <https://doi.org/10.1038/27699>
- Venturini L., You J., Stadler M., Galien R., Lallemand V., Koken M.H., Mattei M.G., Ganser A., Chambon P., Losson R., de Thé H. (1999) **TIF1gamma, a novel member of the transcriptional intermediary factor 1 family** *Oncogene* **18**:1209–1217 <https://doi.org/10.1038/sj.onc.1202655>
- Verger A., Perdomo J., Crossley M (2003) **Modification with SUMO: A role in transcriptional regulation** *EMBO Rep* **4**:137–142 <https://doi.org/10.1038/sj.embor.embor738>
- Vissing H., Meyer W.K.-H., Aagaard L., Tommerup N., Thiesen H.-J (1995) **Repression of transcriptional activity by heterologous KRAB domains present in zinc finger proteins** *FEBS Letters* **369**:153–157 [https://doi.org/10.1016/0014-5793\(95\)00728-R](https://doi.org/10.1016/0014-5793(95)00728-R)
- Witzgall R., O’Leary E., Leaf A., Onaldi D., Bonventre J.V (1994) **The Krüppel-associated box-A (KRAB-A) domain of zinc finger proteins mediates transcriptional repression** *PNAS* **91**:4514–4518 <https://doi.org/10.1073/pnas.91.10.4514>

Zhao H., Moberg K.H., Veraksa A (2023) **Hippo pathway and Bonus control developmental cell fate decisions in the Drosophila eye** *Developmental Cell* 51534580723000473
<https://doi.org/10.1016/j.devcel.2023.02.005>

Zhong S., Salomoni P., Pandolfi P.P (2000) **The transcriptional role of PML and the nuclear body** *Nat Cell Biol* 2:E85–E90 <https://doi.org/10.1038/35010583>

Author information

Baira Godneeva

California Institute of Technology, Division of Biology and Biological Engineering Pasadena, CA 91125, USA, Institute of Molecular Genetics, National Research Center “Kurchatov Institute”, Moscow 123182, Russia
ORCID iD: [0009-0004-1662-8844](https://orcid.org/0009-0004-1662-8844)

Maria Ninova

University of California, Riverside, Riverside, CA 92521, USA

Katalin Fejes Tóth

California Institute of Technology, Division of Biology and Biological Engineering Pasadena, CA 91125, USA

Alexei A. Aravin

California Institute of Technology, Division of Biology and Biological Engineering Pasadena, CA 91125, USA
For correspondence: aaa@caltech.edu

Editors

Reviewing Editor

Claude Desplan

New York University, United States of America

Senior Editor

Claude Desplan

New York University, United States of America

Reviewer #1 (Public Review):

Summary:

This important study from Godneeva et al. establishes a Drosophila model system for understanding how the activity of Tif1 proteins is modified by SUMO. The authors nicely show that Bonus, like homologous mammalian Tif1 proteins, is a repressor, and that it interacts with other co-repressors Mi-2/NuRD and setdb1 in Drosophila ovaries and S2 cells. They also show that Bonus is SUMOylated by Su(var)2-10 on at least one lysine at its N-terminus to promote its interaction with setdb1. By combining nice biochemistry with an elegant reporter gene approach, they show that SUMOylation is important for Bonus interaction with setdb1, and that this SUMO-dependent interaction triggers high levels of H3K9me3 deposition and gene silencing. While there are still major questions of how SUMO molecularly promotes this process, this study is a valuable first step that opens the door for interesting future experimentation.

Major Point:

The RNAseq and ChIPseq data is not available. This is critical for the review of the paper and would help the readers and reviewers interpret the Bonus mutant phenotype and its mechanism of repressing genes.

1. The author's conclusion that Bonus SUMOylation is "essential for its chromatin localization" is not supported by the data. Figure 5F shows less 3KR mutant in the chromatin fraction but there is still significant signal.
2. The author's conclusion that Bonus is SUMOylated at a single site close to its N-terminus is not necessarily true. In several SUMO and Bonus blots throughout the paper (5B, 6C, S4A), there are >2 differentially migrating species that could represent more than one SUMO added to Bonus. While the single K20R mutation eliminates all of these species in Fig 5C, it is possible that K20R SUMOylation is required for additional SUMOylation events on other residues. One way to determine if Bonus is SUMOylated on multiple sites is to add recombinant SUMO protease to the extract and see if multiple higher molecular weight bands collapse into a single migrating species (implying multiple SUMOs) or multiple migrating species (implying something else is altering gel migration).
3. The authors state that most upregulated genes in BonusGLKD are not highly enriched in H3K9me3. The heatmap in figure 3D is not an ideal presentation of this argument. The authors should show an example of what the signal on a highly enriched gene looks like for comparison. The authors also argue that because most upregulated genes in BonusGLKD are not highly enriched in H3K9me3, they must be indirectly repressed. Another possibility is that bonus-mediated H3K9me3 is only important (and present) during early nurse cell differentiation and is later lost and dispensable during the rapid endocycles. After bonus establishes repression though H3K9me3, it might be maintained through bonus-Mi2/Nurd, something else, or nothing at all. The authors could discuss this possibility or perform H3K9me3 ChIP during cyst formation and early nurse cell differentiation rather than in whole ovaries, which are enriched for later stages.
4. The BonusGLKD RNAseq analysis is underwhelming. The conclusion that "Bonus represses tissue-specific genes" has limited value. Every gene that is not expressed in ovaries is "tissue-specific." What subset of tissue-specific genes does Bonus repress? What common features do these genes have and how do they compare to other sets of tissue-specific genes, such as those reportedly repressed by setdb1, Polycomb proteins, small ovary, l(3)mbt, and stonewall (among others in female germ cells). Comparing these available data sets could help the authors understand the mechanism of Bonus repression and how BonusGLKD leads to sterility. The authors could also further analyze the differences between nos-Gal4 and MT-Gal4 to better understand why nos-but not MT-driven knockdown is sterile.

Main Study Limitations:

1. It is unclear which genes are directly vs indirectly regulated by bonus, which makes it difficult to understand Bonus's repressive mechanism. Several lines of experiments could help resolve this issue. 1) Bonus ChIPseq, which the authors mentioned was difficult. 2) RNAseq of BonusGLKD rescued with KR3 mutation. This would help separate SUMO/setdb1-dependent regulation from Mi-2 dependent regulation. Similarly, comparing differentially expressed genes in Su(var)2-10GLKD, setdb1GLKD, 3KR rescue, and MI-2 GLKD could identify overlapping targets and help refine how bonus represses subsets of genes through these different corepressors.

2. The paper falls short in discussing how SUMO might promote repression. This is important when considering the conservation (of lack thereof) of SUMOylation sites in Tif1 proteins in distantly related animals. One piece of data that was not discussed is the apparent localization of SUMOylated bonus in the cytoplasmic fraction of the blot in Figure 5F. Su(var)2-10 is mostly a nuclear protein, so is bonus SUMOylated in the nucleus and then exported to the cytoplasm? Also, setdb1 is a nuclear protein, so it is unlikely that the SUMOylated bonus directly interacts with setdb1 on target genes. Together with Fig 5E (unSUMOylatable Bonus aggregates in the nucleus), one could make a model where SUMO solubilizes bonus (perhaps by disassembling aggregates) and indirectly allows it to associate with setdb1 and chromatin. It is also important to note that in Figure 5I, the K3R mutation appears to lessen but not eliminate Bonus interaction with setdb1. This data again disfavors a model where SUMO establishes an interaction interface between setdb1 and Bonus. To determine which form of Bonus interacts with setdb1, the authors could perform a setdb1 pulldown and monitor the SUMOylation state of coIPed Bonus through mobility shift. If mostly unSUMOylated bonus interacts with setdb1, and SUMO indirectly promotes Bonus interaction with setdb1 (perhaps by disassembling Bonus aggregates), then the precise locations of Bonus SUMOylation sites could more easily shift during evolution, disfavoring the author's convergent evolution hypothesis.

Reviewer #2 (Public Review):

Summary:

The authors analyze the functions and regulation of Bon, the sole *Drosophila* ortholog of the TIF1 family of mammalian transcriptional regulators. Bon has been implicated in several developmental programs, however the molecular details of its regulation have not been well understood. Here, the authors reveal the requirement of Bon in oogenesis, thus establishing a previously unknown biological function for this protein. Furthermore, careful molecular analysis convincingly established the role of Bon in transcriptional repression. This repressor function requires interactions with the NuRD complex and histone methyltransferase SetDB1, as well as sumoylation of Bon by the E3 SUMO ligase Su(var)2-10. Overall, this work represents a significant advance in our understanding of the functions and regulation of Bon and, more generally, the TIF1 family. Since Bon is the only TIF1 family member in *Drosophila*, the regulatory mechanisms delineated in this study may represent the prototypical and important modes of regulation of this protein family. The presented data are rigorous and convincing. As discussed below, this study can be strengthened by a demonstration of a direct association of Bon with its target genes, and by analysis of the biological consequences of the K20R mutation.

Strengths:

1. This study identified the requirement for Bon in oogenesis, a previously unknown function for this protein.
2. Identified Bon target genes that are normally repressed in the ovary, and showed that the repression mechanism involves the repressive histone modification mark H3K9me3 deposition on at least some targets.
3. Showed that Bon physically interacts with the components of the NuRD complex and SetDB1. These protein complexes are likely mediating Bon-dependent repression.
4. Identified Bon sumoylation site (K20) that is conserved in insects. This site is required for repression in a tethering transcriptional reporter assay, and SUMO itself is required for repression and interaction with SetDB1. Interestingly, the K20-mutant Bon is mislocalized in the nucleus in distinct puncta.
5. Showed that Su(var)2-10 is a SUMO E3 ligase for Bon and that Su(var)2-10 is required for Bon-mediated repression.

Weaknesses:

The study would be strengthened by demonstrating a direct recruitment of Bon to the target genes identified by RNA-seq. Given that the global ChIP-seq was not successful, a few possibilities could be explored. First, Bon ChIP-qPCR could be performed on the individual targets that were functionally confirmed (e.g. *rbp6*, *pst*). Second, a global Bon ChIP-seq has been reported in PMID: 21430782 - these data could be used to see if Bon is associated with specific targets identified in this study. In addition, it would be interesting to see if there is any overlap with the repressed target genes identified in Bon overexpression conditions in PMID: 36868234.

The second area where the manuscript can be improved is to analyze the biological function of the K20R mutant Bonus protein. The molecular data suggest that this residue is important for function, and it would be important to confirm this *in vivo*.