

# Dual-color optical activation and suppression of neurons with high temporal precision

Reviewed Preprint

v2 • February 24, 2025

Revised by authors

Reviewed Preprint

v1 • September 19, 2023

Noémie Mermet-Joret, Andrea Moreno, Agnieszka Zbela, Milad Nazari, Bárður Eyjólfsson Ellendersen, Raquel Comaposada-Baro, Nathalie Krauth, Anne von Philipsborn, Andreas Toft Sørensen, Joaquin Piriz, John Y Lin , Sadegh Nabavi 

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark • Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark • Center for Proteins in Memory - PROMEMO, Danish National Research Foundation, University of Tasmania, Hobart, Australia • Tasmanian School of Medicine, College of Health and Medicine, University of Tasmania, Hobart, Australia • Department of Neuroscience, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark • Instituto de Fisiología Biología Molecular y Neurociencias (IFIBYNE), Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

## eLife Assessment

This study develops **useful** tools for distinct optogenetic control of neuronal activity by red or blue light. The basic characterization of the activation of a red-shifted channelrhodopsin paired with a blue-light sensitive anion channel engineered to obtain desired inhibitory current kinetics is **solid**. However, evidence for their practical use under simultaneous multi-color or high frequency stimulation in cells are missing.

<https://doi.org/10.7554/eLife.90327.2.sa2>

## Abstract

A well-known phenomenon in the optogenetic toolbox is that all light-gated ion channels, including red-shifted channelrhodopsins (ChRs), are activated by blue light, whereas blue-shifted ChRs are minimally responsive to longer wave-lengths. Here, we took advantage of this feature to create a system which allows high-frequency activation of neurons with pulses of red light, while permitting the suppression of action potentials with millisecond precision by blue light. We achieved this by pairing an ultrafast red-shifted ChR with a blue light-sensitive anion channel of appropriately matching kinetics. This required screening several anion-selective ChRs, followed by a model-based mutagenesis strategy to optimize their kinetics and light spectra. Slice electrophysiology in the hippocampus as well as behavioral inspection of vibrissa movement demonstrate a minimal excitation from blue light. Of significant potential value, in contrast to existing tools, the system we introduce here allows high frequency optogenetic excitation of neurons with red light, while blue light suppression of action potentials is confined within the duration of the light pulse.

## Introduction

Despite the short history of optogenetics, optical manipulation of neuronal activity has become an indispensable tool in the studies for functional analysis of brain circuits. The increasing needs for a more precise temporal and spatial manipulation of the neurons along with the untapped potentials of optogenetics have been driving forces in expanding the repertoire of the optogenetic toolbox. Main development among these expansions is the discovery of opsins of diverse excitation spectra, with the current excitatory opsins showing light sensitivity ranging from near UV to near far-red spectra (Yizhar et al. 2011 [↗](#); Deisseroth and Hegemann 2017 [↗](#); Marshel et al. 2019 [↗](#)).

A major concern in the use of opsins with longer excitation spectra is that all opsins, in addition to their own excitation spectral peak, are activated by blue light, a problem known in the field as cross-talk or bleed-through. Attempts to create blue light-insensitive opsins will likely be unsuccessful due to the intrinsic blue absorption by the retinal molecule, the non-encoded chromophore, which captures light energy in channelrhodopsins (Waddell, Schaffer, and Becker 1973 [↗](#); Wang et al. 2012 [↗](#)).

There have been several standouts attempts to overcome the problem of cross-talk. One uses a variant of blue-shifted ChR with fast kinetics in combination with a slow kinetics red-shifted ChR (R-ChR) (Klapoetke et al. 2014 [↗](#)). The reasoning is that short pulses of blue light will activate only fast ChR. Since it requires a tight regulation of the protein expression as well as the light intensity, this method has not been adopted for *in vivo* application. Another approach is to expose axons expressing an R-ChR to prolonged photostimulation, during which axons become unresponsive to light pulses and cease to fire (Hooks et al. 2015 [↗](#); Faress et al. 2024 [↗](#)). This method is limited to axonal regions as prolonged optical stimulation of the soma induces trains of action potentials. Additionally, prolonged stimulation of the axons of neurons with slow spike-frequency adaptation, such as fast-spiking interneurons, may result in repetitive activity rather than depolarization block (Hooks et al. 2015 [↗](#)).

A potential solution to the problem of cross-talk is to co-express a blue-shifted anion channelrhodopsin (B-ACR) along with the R-ChR to ‘suppress’ the excitation of R-ChR by blue light. The intention is to have a neuron co-expressing such opsins selectively produce action potentials to red light and not blue light. A recent study achieved this, by pairing GtACR2 and Chrimson (Vierock et al. 2021 [↗](#)). Due to its relatively slow kinetics, this system is more suitable for optical manipulation of lower frequency stimulation. Additionally, optical activation of these opsins, which have a long opening lifetime, may elevate the concentration of intracellular ions with unintended physiological complications (see Discussion).

For these reasons, we created a system with fast closing kinetics. We chose vfChrimson currently the fastest R-ChR to gain precise temporal control of red-light excitation (Mager et al. 2018 [↗](#)). To generate a B-ACR with vfChrimson matching kinetics we used a model-based mutagenesis approach to modify the kinetic properties of ZipACR, an ultrafast blue light-activated chloride channel (Govorunova et al. 2017 [↗](#)). It must be noted that due to the high intracellular chloride concentration in the axonal terminal (Turecek and Trussell 2001 [↗](#); Messier et al. 2018 [↗](#)) the use of a chloride channel-based expression system must be restricted to the somato-dendritic regions (Mahn et al. 2016 [↗](#); Messier et al. 2018 [↗](#)).

We describe several variants of ZipACR, focusing on two, I151T and I151V, which display appropriate kinetics, light spectra, and photocurrent magnitudes for the purposes of our co-expressing system. We then paired the optimized anion channels with a membrane-trafficking optimized vfChrimson (IvfChr) in a single expression unit. The new system, Zip-IvfChr, when

expressed in neurons, can drive time-locked high-frequency action potentials (APs) in response to red pulses (635 nm). On the other hand, in response to blue pulses (470 nm), Zip-Ivf-Chr produces no action potentials.

As expected, blue light pulses transiently suppress APs in Zip-IvfChr-expressing neurons, but the suppression is fully reversed within 5 ms after termination of light pulses due to the use of fast ZipACR mutants. When tested in the facial motor nucleus in the brainstem, Zip-IvfChr activation by red but not blue light triggered vibrissa movement of large amplitudes. Due to its high temporal precision of activation and suppression, this tool could be further developed for independent optical excitation of distinct neural populations.

## Results

### The strategy to reduce blue light-mediated excitation by red-shifted channelrhodopsins


We initially tested our strategy using existing variants of excitatory and inhibitory channelrhodopsin. The channelrhodopsin Chrimson and its variants have the most red-shifted spectra (Klapoetke et al., 2014), and their response to 470 nm has a slower on-rate compared to the 590 nm light (Figure 1-figure supplement 1A and B). Despite this, 470 nm light is equally effective in inducing APs (Figure 1-figure supplement 1C). To overcome this limitation, we decided to take advantage of blue light-gated inhibitory chloride channels which effectively block action potential upon exposure to blue light (Govorunova et al. 2015, 2017). By co-expressing the B-ACR variant GtACR2, along with a Chrimson variant, photostimulation of targeted neurons by red light would be expected to only activate R-ChR, which causes photoactivation of these neurons. Blue light, however, activates R-ChR as well as GtACR2. This allows the chloride channel to induce shunting inhibition of the neurons which in turn would be expected to prevent firing of the neurons by blue light (Fig. 1A).

### Proof of concept of the co-expression strategy in *Drosophila*

As the first step to test the potential validity of this approach, we took advantage of the well-studied male-specific P1 neurons in the fly central brain that elicit courtship singing upon activation (von Philipsborn et al. 2011; Inagaki et al. 2014; Ellendersen and von Philipsborn 2017; Mohammad et al. 2017) (Fig. 1B 1-B2). When Chrimson was co-expressed with an inactive control protein (tdTomato), illumination at 470 nm and 590 nm wavelengths led to similar amounts of courtship song ( $110.1 \pm 12.8$  and  $78.5 \pm 11.6$  pulses per 10s,  $n = 16-17$ , respectively), demonstrating that blue light at the chosen intensity can efficiently activate Chrimson (Fig 1B3). When Chrimson was co-expressed with the blue-sensitive anion channelrhodopsin GtACR2, 590 nm light illumination reliably elicited male courtship song ( $95.2 \pm 9.14$  pulses per 10s); while 470 nm light illumination did not evoke any detectable pulse song ( $n = 17-18$ ). Neither 590 nm nor 470 nm light pulses exposure induced courtship song in flies expressing only GtACR2 ( $n = 5$ ).

### Identifying a suitable pair of ultrafast light-gated anion channel and a red-shifted Chrimson variant

Although GtACR2 proved to be effective in protocols with pulses of long duration, its slow current decay (Fig. 2A, see GtACR2,  $161 \pm 24$  ms,  $n = 3$ ) limits its application to optical stimulations of lower frequencies (Govorunova et al. 2015, 2017). For this reason, we searched for a pair of blue-shifted anion channelrhodopsin and red-shifted cation channelrhodopsin with fast and



**Figure 1.**

### Reducing blue-light mediated excitation in red-shifted channelrhodopsins- Proof of concept.

**(A)** Schematic representation of the approach. A red-shifted channelrhodopsin (R-ChR) is co-expressed with a blue-shifted anion channel (B-ACR). When red light is ON, there is an overall excitation of the cell due to the more dominant R-ChR response. When blue-light is ON, the shunting inhibitory effect of B-ACR reduces the excitation induced by R-ChR to blue light. This panel was created using [BioRender.com](#). **(B)** Validation of the approach in flies expressing Chrimson, Chrimson and GtACR2, or GtACR2 alone. **(B1)** Experimental setup to record the courtship song of solitary male flies. This panel was created using [BioRender.com](#). **(B2)** Top, Example of a reconstructed neuronal arborization of P1 neurons, Bottom, example of a typical courtship song induced by LED pulses. **(B3)** Courtship song production of the solitary male flies during 10-sec of constant illumination with 590 nm (amber) or 470 nm (blue) light. Each dot represents an individual drosophila (mean  $\pm$  SEM).

matching kinetics. For a fast R-ChR, we evaluated the recently published vfChrimson, the fastest red-shifted channelrhodopsin available at this time, which can induce APs up to the intrinsic limit of the neurons (Mager et al. 2018 [↗](#)).

Previously, we found that Chrimson and its variants traffic poorly to the membrane and developed a strategy to improve the trafficking (C-Chrimson-ts, (Bonaventura et al. 2019 [↗](#)). We adapted this modification to improve the trafficking of vfChrimson (referred to as IvfChr, **Figure 2 - figure supplement 1** [↗](#)). In our measurements in HEK 293 cells, IvfChr has a channel off-rate consistent with previously reported (Mager et al. 2018 [↗](#)) (off-rate time constant of  $5.6 \pm 0.3$  ms,  $n = 10$  compared to C-Chrimson-ts  $21.5 \pm 1.2$ ,  $n = 10$ ; mean  $\pm$  SEM, **Fig. 2A** [↗](#), **Figure 2 - figure supplement 2** [↗](#)). When comparing C-ChrimsonR-ts and IvfChr photocurrents, we observe 40% reduction of photocurrent of IvfChr ( $0.086 \pm 0.055$  pA/ pF/A.U,  $n = 25$ ) compared to C-ChrimsonR-ts  $0.145 \pm 0.126$  pA/pF/A.U.,  $n = 25$ ; mean  $\pm$  SD, **Figure 2 - figure supplement 2B** [↗](#)) when the peak photocurrent amplitude was adjusted to membrane capacitance and membrane fluorescence.


To counteract the IvfChr excitation, we estimated that desired properties of the blue light activated anion ChR should have the off-rate time constant of  $\sim 15$ -30 ms, minimal response to light of wavelength  $> 590$  nm and a strong photocurrent. We identified ZipACR, RsACR\_995, GcACR\_457, RIACR\_477 and GtACR2 as possible templates for further development based on published data (Govorunova et al. 2017 [↗](#)). GtACR2, RsACR\_995 and RIACR\_477 are spectrally ideal with negligible response to 590 nm light (**Figure 3 - figure supplement 1A** [↗](#)), but their kinetics are slow (mean off-rate of  $161.0 \pm 24.1$ ,  $122.7 \pm 36.3$  and  $95.8 \pm 49.4$  ms, respectively,  $n = 3$ , **Fig. 2A** [↗](#)) which could have undesired long lasting inhibitory effects after light termination. GcACR\_457 has very small photocurrents and cannot be characterized accurately, possibly due to poor membrane trafficking. ZipACR, although spectrally not ideal ( $\sim 515$  nm; (Govorunova et al. 2017 [↗](#)) has a kinetic close to the desired range ( $9.5 \pm 3.0$  ms,  $n = 9$ ) and it has strong photocurrents (**Fig. 2A** [↗](#)).

To test the inhibitory property of ZipACR, we injected 40 Hz somatic current in organotypic hippocampal neurons expressing the channel. Consistent with the original report, individual pulses of 470 nm light resulted in the time-locked suppression of APs (**Fig. 2B** [↗](#)). Next, we examined whether ZipACR can block blue light-mediated APs in neurons expressing vfChrimson. Recordings from the granule cells in acute hippocampal slices co-expressing ZipACR and vfChrimson showed that, in 8 out of ten cases 470 nm light induced significantly fewer APs than 635 nm light (**Fig. 2C1** [↗](#)), in line with our prediction. However, we consistently observed the appearance of individual APs time-locked to the blue light offset (**Fig. 2C1-C2** [↗](#)). We concluded that due to the fast-closing rate, the photoinhibitory current generated by ZipACR decays too early to suppress the remaining excitatory response of vfChrimson. Therefore, we decided to generate variants of ZipACR with slightly slower off-kinetics.

## Optimizing ZipACR off-kinetics

It is technically easier to generate mutations with slower kinetics and blue-shifted spectra than generating variants with faster on- and off-kinetics. Thus, we decided to use ZipACR as a template for further mutagenesis instead of further modifying GtACR2. To identify relevant mutation sites in ZipACR, we generated a homology model of ZipACR with the crystal structure of GtACR1 (Govorunova et al., 2017 [↗](#)) and identified the putative residues forming the retinal binding pocket in ZipACR (**Fig. 3A** [↗](#)). Multiple alignments were performed with the kinetically slower GtACR1 and GtACR2 to identify possible replacement of these residues that would result in slower kinetics and/or small blue-shift in ZipACR without disrupting the channel function (**Fig. 3A2** [↗](#)). We tested these variants in HEK 293 cells for photocurrent ratio 590 nm / 470 nm LED stimulating light and measured the channel off-rate kinetics and photocurrent amplitude (**Fig. 3B** [↗](#)).


Of these candidates, we identified I151V and I151T at the 4th transmembrane domain of ZipACR that resulted in a small increase in channel off-rate time constant ( $15.5 \pm 8.4$  ms for I151V and  $20.2 \pm 6.4$  ms for I151T, mean  $\pm$  SD, **Fig. 3B** [↗](#)) compared to the original ZipACR ( $9.5 \pm 2.9$  ms, mean  $\pm$



**Figure 2.**

### Comparison of channel properties of light-gated chloride channels and Chrimson variants.


(A) Representative photocurrents of the Chrimson and ACRs variants in whole-cell patch-clamp recordings from HEK 293 cells with 590 nm and 470 nm illumination (top black bars, 10 mW/mm<sup>2</sup>). The plot shows the off-time constant (top) following 1 sec of 470 nm illumination, and the peak photocurrent (bottom) at -60 mV (mean  $\pm$  SD,  $n = 3$ -13 cells). (B) Typical response of CA1 cells from hippocampal organotypic slices expressing ZipACR to 40 Hz, 5 ms pulses (top black trace) and to an illumination protocol using 40 Hz of 10 ms light pulses at 470 nm (4 mW/mm<sup>2</sup>) (bottom blue trace). (C) Recordings from dentate gyrus granule cells in acute slices expressing ZipACR and vfChrimson. (C1) Representative responses to 500 ms illumination (bottom black bar) at 470 nm and 635 nm (10 mW/mm<sup>2</sup>) (blue and red traces respectively). Note, 635 nm illumination evoked action potential firing while 470 nm illumination only evoked a single action potential time-locked to the end of the light pulse (arrowhead). Inset, example of an action potential upon 470 nm light termination and plot showing the number of light-evoked spikes at 635 nm (red bar) and 470 nm (blue bar) illumination.  $n = 10$  cells, paired  $t$ -test  $**p < 0.01$ ; (C2) Example of light-induced firings by 635 nm and 470 nm illumination (red and blue traces respectively, 5 ms pulses (black squares) at 10 Hz). The arrowheads show consistent APs time-locked to the end of the blue light pulse. Inset, example of a voltage-clamp recording of currents evoked by 5 ms pulses of 635 nm and 470 nm light (red and blue traces respectively). Light-induced currents with 470 nm illumination were initially outward but turned inward gradually and peaked immediately following the light offset. Note that the late inward current matches the red light induced current, reflecting a slower kinetics for vfChrimson compare to ZipACR.



**Figure 3.**

### Characterization of the optimized variants of the ultrafast anion channelrhodopsin ZipACR.

**(A)** Structure-based design of ZipACR variants. **(A1)** The retinal binding pocket of GtACR1 and the putative retinal binding pocket of ZipACR homology model. **(A2)** The alignment of the transmembrane domains (TM) of GtACR1, GtACR2 and ZipACR. The residues surrounding the retinal binding pocket of GtACR1 and the corresponding residues from GtACR2 and ZipACR are highlighted in red. \*marks the residues targeted for mutation. The lysine that forms the Schiff base with retinal is indicated in green. The numbers on the left correspond to the number of the first residue in each line. **(B)** Basic properties of the ZipACR variants compared to IvfChr. Off-time constant (top) following 1 sec of 470 nm illumination, peak photocurrent (middle) and 590/470 ratio (bottom). The data have been obtained from whole-cell recordings in HEK 293 cells. Each dot corresponds to one cell (mean  $\pm$  SD,  $n = 2-12$ ). **(C)** On-time constant (**C1**) and normalized photocurrent (**C2**) of the selected variants Zip(151T) and Zip(151V) as compared to the original ZipACR and IvfChr, with 470 nm or 590 nm illumination of various light intensities (in  $\text{mW}/\text{mm}^2$ ). For the photocurrent, the values have been measured at the maximum response. The data points are the mean value  $\pm$  SEM ( $n = 6-7$ ). **(D1)** The action spectra of ZipACR, Zip(151V) and Zip(151T) as measured from 650 nm to 400 nm **(D2)** For the action spectra, the maximum photocurrent amplitudes measured at each wavelength were normalized to the peak values obtained from the same cell across the spectrum. The data points are the mean value  $\pm$  SEM ( $n = 6-7$ ). **(E)** Validation of the efficiency of the Zip(151T) variant in slices. **(E1)** The response of dentate gyrus granule cells expressing Zip(151T) variant at 470 nm ( $4\text{mW}/\text{mm}^2$ ) and 635 nm ( $7\text{mW}/\text{mm}^2$ ) light pulses delivered through a 63x objective. This panel was created using *BioRender.com*. **(E2)** Representative traces of the firing induced by current injection at 40 Hz. Overlapping 470 nm but not 635 nm pulses block action potentials. The plot on the right shows the efficiency (in %) of Zip(151T) in blocking individual action potentials at 470 and 635 nm light pulses (mean  $\pm$  SD,  $n = 6-10$  cells). The traces of the two outliers are presented in **Figure 3 - figure supplement 3A**.



**Figure 3.** (continued)

SD). We also generated double and triple mutants combining the I151V or I151T with Y170F, Y177F or V229L mutations but most of these double or triple mutants appeared to have smaller photocurrents and off-rate kinetics that could not be fitted well with single exponential back to zero baseline. Based on these results, we concluded the variants I151V (Zip(151V)) and I151T (Zip(151T)) are the best candidates for the pairing and proceeded for further characterizations.


Neither Zip(151V) nor Zip(151T) mutations changed the ion permeability of ZipACR as measured by channel reversal potential in potassium gluconate and cesium chloride based intracellular recording solutions (**Figure 3 - figure supplement 1B** and **C**). In terms of light sensitivity, it is desirable that the ZipACR variant would respond to 470 nm light faster and stronger and to red-shifted light slower and weaker than IvfChr at the same intensity (**Figure 3 - figure supplement 2**). With both Zip(151V) and Zip(151T), the channel on-rate kinetics have the desirable properties as described above to 470 nm and 590 nm light, although the Zip(151T) on-rate starts to approach the on-rate of IvfChr at light intensities  $>5 \text{ mW/mm}^2$  at 590 nm (**Fig. 3C1**). With 470 nm light, IvfChr does not appear to reach saturating response even at  $20 \text{ mW/mm}^2$  (**Fig. 3C2**). With 590 nm light, IvfChr reaches saturating response at  $10 \text{ mW/mm}^2$  whereas none of the ZipACR variants appears to reach saturating response even at  $20 \text{ mW/mm}^2$  (**Fig. 3C2**). Spectrally, Zip(151T) shows a wider red-shifted action spectra than ZipACR whereas Zip(151V) may have a slightly narrower blue-shifted action spectra than ZipACR (**Fig. 3D**).

Since Zip(151T) is a more red-shifted variant of the two (**Fig. 3B,D**), we asked whether red light pulses may induce photoinhibitory currents that suppress APs. We expressed Zip(151T) in the granule cells of the dentate gyrus in hippocampal slices using AAV vector and tested its inhibitory effect upon 470 nm and 635 nm light pulses. We injected somatic current at 40 Hz, while alternately delivering overlapping 5 ms pulses of light. Except in two instances, blue light pulses were consistently effective in blocking APs, while red light had no effect on APs induction (**Fig. 3E**,  $n = 10$  for 470 and  $n = 6$  for 635, the two outliers are presented in **Figure 3 - figure supplement 3A**). It has been reported that at resting or hyperpolarized membrane potentials ZipACR may produce light-evoked APs (Kato et al. 2018). We did not observe such a phenomenon under our experimental setup (**Figure 3 - figure supplement 3B**).

## Validation of the co-expression strategy in brain slices

Since co-injection of two viruses results in variability in the ratio of expression in different neurons, we used the bicistronic 2A approach (Supplementary material 1). We favored this system to the internal ribosomal entry sites (IRES) (Pelletier and Sonenberg 1988; Douin et al. 2004) and tandem fusion strategies (Kleinlogel et al. 2011). As observed by us and reported by other groups, IRES produces an unbalanced expression level with the second protein at much lower level (Mizuguchi et al. 2000). The testing of vfChrimson in the recently published BiPOLES tandem fusion system with GtACR2 (Vierock et al. 2021) resulted in high levels of intracellular accumulation of the tandem opsins in HEK293 cells (**Figure 2 - figure supplement 1**).

Therefore, we generated constructs containing IvfChr and Zip(151T/V) (ZipT-IvfChr and ZipV-IvfChr) using the bicistronic 2A cassette with KV2.1 soma targeting sequence for both ZipACR mutant and IvfChr (**Fig. 4A**, **Figure 2 - figure supplement 1** and Supplementary material 1). This ensures an equimolar translation of the two proteins from one mRNA and soma enrichment of the opsins in neurons. In comparison to BiPOLES with vfChrimson, the vfChrimson in our bicistronic 2A cassette led to  $\sim 2.7 \times$  higher photocurrent ( $-7.57 \pm 5.37 \text{ pA/pF}$  in ZipV-IvfChr compared to  $-2.72 \pm 1.33 \text{ pA/pF}$  in BiPOLES,  $n = 14$ ) when tested in identical conditions, consistent with the improved membrane trafficking observed with IvfChr (**Figure 2 - figure supplement 1A** and **B**). The photocurrent amplitudes of the IvfChr and ZipV-IvfChr constructs in HEK 293 cells are comparable after adjustment to cell size and membrane fluorescence (**Figure 2 - figure supplement 1D**). We did not test the ZipT-IvfChr variant as the red-shifted response of Zip(151T) makes it more difficult to isolate the IvfChr response to 590 nm light available on our LED system.



**Figure 4.**

### Zip(151T) variant co-expressed with the red ChR2 IvfChr prevents blue-light induced action potentials, in-vitro and in-vivo.

(A) The expression cassette for IvfChr and bicistronic ZipT-IvfChr used in rAAV vector. The selected ZipACR variant Zip(151T) is co-expressed with the red-sensitive IvfChr by using the 2A self-cleaving peptide (T2A). All the opsins are soma-targeted with a Kv2.1 peptide. (B) Comparison of the high-frequency red and blue light-induced spiking (B1), and spike probability (B2,  $n = 6-13$  for each opsin), of DG cells expressing IvfChr or the ZipT-IvfChr, with 10 and 20 Hz of light pulses (4-7 mW/mm<sup>2</sup>). Holding potential = -60 mV. (C) Blue and red lightdriven spike fidelity at various light intensities in DG cells ( $n = 4-10$  cells for each opsin; 10 ms pulse width, single pulse). (D) Experimental design. ZipT-IvfChr or IvfChr are expressed in the facial nucleus (FN) of the brainstem (top) and light-induced whisker movements are recorded. Example of soma-targeted expression of our ZipT-IvfChr construct in FN (bottom). This panel was created using *BioRender.com*. (E) Averaged Z-scores of the angle of the whiskers under 450 nm and 638 nm illumination (20 mW/mm<sup>2</sup>) of the FN motoneurons. ZipT-IvfChr does not trigger whiskers movement under blue-light ( $n=4$  animals per condition). (F) Ratio of the AUC of the responses to 450 nm and 638 nm illumination for ZipT-IvfChr and IvfChr ( $n=4$  animals per condition, \*\* $p < 0.01$ , t-test). (G) Distribution of the AUC for three individual trials from each animal at 450 nm and 638 nm illumination ( $n=4$  animals per condition).

After characterization in HEK293 cells, we examined the power at which responses to 470 nm and 635 nm lights induce APs in neurons expressing ZipT-IvfChr, ZipV-IvfChr, or IvfChr. We expressed AAV constructs in the granule cells of the dentate gyrus in the hippocampal slices (as described in [Fig. 3E1](#)). We recorded cells under the current-clamp mode and injected current through the patch pipette to maintain the membrane potentials at  $-60$  mV. To examine the efficiency of 635 nm and 470 nm lights in inducing APs, we delivered 10 Hz and 20 Hz light pulses of 5 ms duration ([Fig. 4B](#)). As expected, in IvfChr expressing neurons, both 470 nm and 635 nm lights induced APs ( $n=6$ ). In neurons expressing ZipT-IvfChr ([Fig. 4B](#)), 635 nm pulses were equally effective in driving APs. On the other hand, 470 nm light only induced modest depolarization, which did not produce the threshold for AP induction ( $n = 5-12$  for each opsin). Similarly, co-illumination with 470 nm and 635 nm also failed to generate AP, showing blue light effectively nullifies the depolarizing property of red light ([Figure 4 - figure supplement 1A](#)).

In IvfChr expressing neurons 470 nm and 635 nm lights induced APs with light powers as low as  $1 \text{ mW/mm}^2$  ( $n = 10$ , [Fig. 4C](#)). In contrast, ZipT-IvfChr expressing neurons produced APs only to 635 nm light stimulation ([Figure 4 - figure supplement 1B](#)), whereas 470 nm light powers as high as  $10 \text{ mW/mm}^2$  failed to induce APs ( $n = 7-8$ , [Fig. 4C](#)). As for ZipV-IvfChr, we occasionally observed blue light-induced APs ([Figure 4 - figure supplement 1B](#)), which may be explained by its relatively faster closure rate ([Fig. 3B](#)). In neurons expressing IvfChr, we noticed a reduction in probability of AP upon successive optical stimulation specifically to 635 nm pulses ([Fig. 4B](#)). Consistent with this, we observed a reduction in response in HEK 293 cells expressing IvfChr ([Figure 4 - figure supplement 2](#)). This phenomenon, likely caused by the desensitization of the channel, is consistent with previous reports with other red-shifted channelrhodopsins ([Lin et al. 2013](#)).

## Validation of the co-expression strategy in vivo

To evaluate the system *in vivo*, we compared the performance of ZipT-IvfChr and IvfChr in driving vibrissae movements by optically activating motoneurons of the facial nucleus (FN) expressing either of the two constructs ([Lin et al. 2013](#); [Sreenivasan et al. 2015](#); [Gharraei et al. 2020](#)) ([Fig. 4D](#)). Four to six weeks after virus injection, a 200- $\mu\text{m}$  diameter optic fiber was placed above the FN of head-fixed anesthetized mice and motoneurons were stimulated at 0.33 Hz using 100ms pulses of either 450 nm or 638 nm light. In mice expressing IvfChr, whiskers movements of large amplitude were elicited with both red and blue pulses ([Fig. 4E-G](#), the Area Under the Curve (AUC) of  $2.9 \pm 0.09$  and  $1.95 \pm 0.13$ , respectively, with a ratio  $\text{AUC}_{450}/\text{AUC}_{638} = 0.68 \pm 0.05$ ,  $n=4$ ) (Supplementary material 2). Pulses of red light elicited strong whisking (AUC of  $1.8 \pm 0.29$ ,  $n=4$ ) in mice expressing ZipT-IvfChr, which was comparable to the IvfChr group. On the other hand, 450 nm light pulses failed to reliably elicit whiskers movement, as represented by a significant decrease of the AUC under blue light illumination ( $0.25 \pm 0.07$ , ratio  $\text{AUC}_{450}/\text{AUC}_{638} = 0.15 \pm 0.04$ ,  $n=4$ ,  $p = 0.0012$ , t-test) (Supplementary material 2). Overall, we confirmed that *in vivo*, unlike motoneurons expressing IvfChr, motoneurons expressing ZipT-IvfChr are largely non-responsive to blue light, while preserving the red shifted property of an R-ChR.

## Fast recovery of blue-light induced suppression of action potentials in vitro

We expected that blue-light pulses in ZipV-IvfChr- and ZipT-IvfChr-expressing cells would have an overall inhibitory effect on membrane potential due to the activation of ZipV/T but this effect is short in duration due to the fast kinetics of ZipACR variants. To test this, we stimulated the neurons by somatic current injection at 40 Hz, while alternately delivering perfectly overlapping 5 ms pulses of lights ([Fig. 5A](#)). Light pulses of 470 nm blocked APs, while 635 nm light pulses had virtually no effect. Importantly, we asked how fast a neuron recovers from the blue-light inhibition associated with the expression of ZipV-IvfChr and ZipT-IvfChr. We induced APs through somatic current injection at different time points from the offset of a light pulse ([Fig. 5B1](#)). In

neurons expressing ZipT-IvfChr, current injection evokes APs with high probability after 5 ms, which is consistent with our goal of minimizing the unintended disruption of membrane excitability after the light stimulation (**Fig. 5B** [↗](#)). The recovery time for ZipV-IvfChr was immediate, with 100% success rate immediately following the light offset (**Fig. 5B2** [↗](#)).

## Evaluation and modification of recently developed light-gated potassium channelrhodopsins


A main drawback of using chloride channels is that their opening in the axonal terminals causes depolarization and neurotransmitter release (Turecek and Trussell 2001 [↗](#); Mahn et al. 2016 [↗](#)). This is due to the high concentration of chloride in the terminals. Since potassium channels are hyperpolarizing in all neuronal compartments, we evaluated the recently published potassium selective channelrhodopsins HcKCR1, HcKCR2 (Govorunova et al. 2022 [↗](#)), WiChR and B1ChR2 (Vierock et al. 2022 [↗](#)) for possible pairing with IvfChrimson in next version of our tool. We observed significant light-induced membrane currents from HEK293 cells in all the examined potassium-selective channelrhodopsins, except for B1ChR2 (**Figure 5 - figure supplement 1A** [↗](#)). This is consistent with previously reported data in mammalian cells (Vierock et al., 2022 [↗](#)). Therefore, B1ChR2 was not tested any further.

The channel reversal potentials, as measured by a ramp protocol, were  $-58.0 \pm 8.7$  mV ( $n = 9$ ),  $-50.7 \pm 8.0$  mV ( $n = 14$ ) and  $-65.2 \pm 6.5$  mV ( $n = 7$ ) for HcKCR1, HcKCR2 and WiChR, respectively (**Figure 5 - figure supplement 1B** [↗](#) and **C** [↗](#)) which were more hyperpolarized than the reversal potential for ZipACR measured previously in similar conditions ( $-39.0$  mV). In terms of the channel kinetics, HcKCR1, HcKCR2 and WiChR have the off-rate time constants (measured at holding potential of  $-10$  mV) of  $49.7 \pm 11.3$  ms ( $n = 9$ ),  $58.6 \pm 47.8$  ms ( $n = 14$ ) and  $229.3 \pm 53.8$  ms ( $n = 9$ ) (**Figure 5 - figure supplement 1D** [↗](#)) which is much slower than the channel kinetics of ZipACR.

As an observation, the off-rate for the HcKCRs often do not fit a single exponential function well and the value measured here underestimate the speed of the channel (see the traces in **Figure 5 - figure supplement 1E1** [↗](#)). HcKCR1 has spectral peak at 530 nm as previously reported (Govorunova et al. 2022 [↗](#)) and still responds strongly to 590 nm light ( $73.0 \pm 6.4$  % of 530 nm response,  $n = 10$ ) whereas HcKCR2 and WiChR have spectral peaks at 470 nm and have small responses to 590 nm light ( $9.0 \pm 3.0$  % of 470 nm response,  $n = 8$  for HcKCR2 and  $27.6 \pm 7.2$  % of 470 nm response,  $n = 6$ ; for WiChR; **Figure 5 - figure supplement 1E2** [↗](#)). In light of these results, none of the wild-type K-ChRs appear to be optimal for the pairing and further mutagenesis, and engineering in future study would be required to optimize the pairing.

## Discussion

Here, we describe a dual-color expression system by which pulses of red-light drive time-locked high frequency APs, while blue light allows precise suppression of AP confined within the duration of the pulse. We created this system by pairing a B-ACR with the ultrafast red-shifted ChR, IvfChr. This requires the B-ACR to have a significantly larger photocurrent compared to R-ChR as well as a faster channel opening and slightly slower channel closing. As we discuss later, we considered anion channels with slow decay of inhibition are not suited for many experimental manipulations of long duration. For this reason, we chose ZipACR, currently the fastest B-ACR, as the template for further modifications. Based on our homology modeling and multiple alignments with the kinetically slower B-ACRs, GtACR1 and GtACR2, we identified potential residue replacements within the putative retinal binding pocket which would result in slightly slower kinetics without impairing the channel function.



**Figure 5.**

### The blue-light induced suppression of action potentials is confined within the duration of the light pulse.

**(A)** Optical inhibition of spiking in DG cells expressing the ZipT-IvfChr and ZipV-IvfChr variants. Representative traces of the firing induced by current injection at 40 Hz. Overlapping 470 nm but not 635 nm light pulses block action potentials. Holding potential = -60 mV. Plots on the right shows the efficiency (in %) of ZipT-IvfChr in blocking individual action potentials at 470 and 635 nm ( $n = 9-10$ ). **(B)** Temporal recovery of neuronal excitability following light-induced inhibition by the ZipACR variants. **(B1)** Experimental paradigm. DG granule cells expressing ZipT-IvfChr or ZipV-IvfChr have been stimulated by a somatic current injection (5 ms pulse) overlapping or preceded by a 470 nm pulse (5 ms, 4 mW/mm<sup>2</sup>) at 0, 5, 10, 15 or 20 ms intervals. The traces in current clamp (bottom) are representative of the responses to current injections. This panel was created using [BioRender.com](https://www.biorender.com). **(B2)** The spike probability shows a fast recovery of ZipT-IvfChr (<5ms) and an immediate recovery of ZipV-IvfChr, from the off-set of the 470 nm light ( $n = 7$  for ZipT-IvfChr,  $n = 6$  for ZipV-IvfChr, mean  $\pm$  SD).

To our surprise, many mutations of ZipACR around the alltrans retinal do not alter the channel properties or compromise channel conductance very strongly. This is contrary to our past experiences with channelrhodopsins where conductance is highly sensitive to mutations around the retinal binding pocket (data not shown). ZipACR double mutants appear to have more compromised conductance or altered properties. Among the tested positions, the I151 residue appears to have significant influences in kinetics and spectral properties. Indeed, we found that two variants, Zip(151T) and Zip(151V), have properties optimal for our purpose. More specifically, the channels off-rate kinetics have the desirable property of being slightly but sufficiently slower than IvfChr without compromising the opening rate, photocurrent and light spectrum. Thus, when paired with IvfChr, they prevent AP induction by blue light, without interfering with red light-induced high frequency neural spiking.

Of the two variants, we observed Zip(151T), and consequently ZipT-IvfChr, to be more consistent in blocking blue light-induced APs, which could be explained by its slightly slower closing rate. Although Zip(151T) is the more red-shifted of the two, within the tested range of light intensities and expression, only rarely did we observe a block of APs at 635 nm. However, we do not recommend the use of orange light pulses with ZipT-IvfChr, as we observed a significant photocurrent in this wavelength ([Fig. 3D1](#) and [D2](#); [Figure 3 - figure supplement 2](#)). Comparing the ZipV-IvfChr and the ZipT-IvfChr constructs, we would recommend users to use the ZipT-IvfChr for most general experiments if fast light source with wavelength > 600 nm is available given the reduced probability of neuronal spiking at the end of blue light pulse. The ZipV-IvfChr system is faster in kinetics and is suitable in situation where kinetics is critical but there is some tolerance in spiking at the end of blue light pulse in some neurons.

A recent study ([Vierock et al. 2021](#)) proposed an R-ChR expression system by relying on the pairing between GtACR2 and Chrimson, as we did in our early stage work with *Drosophila*. For experimental reasons linked to controlling neuronal physiology, the priority we set was to create a system with fast closing kinetics to minimize any physiological disruption after the termination of light illumination. All channelrhodopsins have the potentials to alter osmotic balance, ion gradient and pH of small neuronal compartments and this is exacerbated by long light pulses and prolonged channel opening. This can be problematic in long-term neuroplasticity experiments. Cationic channelrhodopsins are known to conduct calcium ions, long opening can evoke significant elevation of intracellular calcium ([Lin et al. 2013](#)) and potentially activate second messenger pathways.

The alteration of pH, osmotic balance and ion gradient (either directly or secondary to elevated transporter activity) applies to both cationic and anionic channelrhodopsins and can have unpredictable biological consequences. Hence, channelrhodopsins with rapid kinetics such as the ones used in our system and brief stimulation light pulses are more ideal for experiments where neuronal or synaptic plasticity are of importance. We were able to demonstrate that neurons expressing ZipT-IvfChr can recover from this inhibition within 5 ms after light termination and ZipV-IvfChr recovers almost instantaneously. If we model our results to a B-ACR with slower kinetics such as GtACR2 used in BiPOLES, we could expect the inhibitory effects of blue light to last up to 10-20 ms. This could disrupt normal neuronal physiology and signal processing that occurs in millisecond time scale, as well as alter intracellular ion concentrations which may lead to short-term and long-term alteration in physiology.

From an experimental viewpoint, a fast photostimulation system has the flexibility to perform a wide range of optical manipulations. Ultimately, ZipT-IvfChr and ZipV-IvfChr will allow dual optical activation of segregated groups of neurons when co-expressed with a fast blue-shifted channelrhodopsin such as oChIEF in the same brain region. In these experiments, it is critical to minimize the blue-light mediated inhibition due to possible disruption of responses resulting from local monosynaptic connection that may have important modulatory effects on postsynaptic cells. Spike-timing dependent experiments requiring high temporal precision would be one of the

experiments that would benefit from our system. Blue-light activated cationic channelrhodopsin such as oChIEF can be expressed presynaptically to control neurotransmitter release and ZipV/T-IvfChr can be expressed postsynaptically to control spiking to induce long-term potentiation or long-term depression.

In cortical experiments, blue cationic channelrhodopsin and ZipV/T-IvfChr can be expressed in projection neurons and interneurons to study the effects of excitation/inhibition balance in cortical processing of information or decision making. In the striatum, the independent control of neurons of direct and indirect pathways would benefit from the dual wavelength manipulation. Many of these experiments would require the use of retrograde AAV, cell-type specific enhancer/promoter or transgenic animals with defined driver line. Although ZipV/T-IvfChr can theoretically be used with single-photon jRCaMP imaging, this would require blue light illumination with duration greater than several milliseconds which will lead to prolonged inhibition of ZipV/T-IvfChr-expressing.

The main limitation of our current approach is that there is limited control of the IvfChr and ZipV/T expression. Although the 2A approach should achieve better translation ratio compared to the use of Internal Ribosome Entry Site (IRES), the levels of respective proteins would also depend on the amount of protein degradation and the fraction of channelrhodopsins that failed to fold or incorporate the retinal chromophore correctly. This can vary with different neurons in different regions. Therefore, similar to other optogenetic systems, it is important for the users of our system to validate the approach in their cells of interest before commencing behavioral experiments.

There can also be situations where dual virus delivery would be preferable compared to bicistronic expression cassette. As shown in our testing, the tethered-tandem expression system used in the BiPOLES reduces the photocurrent of vfChrimson compared to our IvfChr, possibly associated with hindered membrane trafficking. Given the reduced conductance of vfChrimson compared to Chrimson, a BiPOLES based tethered-tandem expression system is also not ideal. The development of an improved tunable dual expression system would therefore be beneficial for the neuroscientific community. Ultimately, the co-expression strategy remains a temporary solution to the spectral cross-talk of opsins and the development of a true spectrally narrow channelrhodopsin would still be the preferred solution. However, whether this is achievable due to the chemical properties of the retinal chromophore is questionable.

Before concluding, we must note that our approach, which acts based on chloride flux, is unlikely to be suitable for axonal terminal inactivation, where the intracellular chloride concentration is high (Mahn et al. 2016 [↗](#), 2018 [↗](#)). Potassium selective channelrhodopsins, having a hyperpolarized reversal potential in all cellular compartments, may appear better positioned for our purpose. However, as we show in our characterizations, none of the wild-type variants would be optimal for the pairing kinetically. WiChR is kinetically too slow but is the most potent inhibitor. HcKCR1 has strong desensitization, is kinetically too slow and spectrally too red-shifted. HcKCR2 is the least potassium selective but is spectrally and kinetically closest to the ideal candidate of the three. Overall, further engineering and modification would still be required to identify the ideal mutants for the pairing similar to what we have done with ZipACR in this study.

At present, for axonal or terminal stimulation, the prior published approaches utilizing differential kinetics and light sensitivity of the opsins or optical manipulation combined with synaptic release properties would still be the preferred options (Klapoetke et al. 2014 [↗](#); Hooks et al. 2015 [↗](#)). Specifically, we refer our readers to a recent work, where we demonstrate effective independent dual optical activation of converging axons, electrophysiologically as well as behaviorally (Faress et al. 2024 [↗](#)). Thus, the system introduced here complements these studies that are unsuitable for somatic photoactivation. This permits researchers to choose complementary methods based on their experimental needs. Future engineering of a faster and

blue-shifted potassium selective channelrhodopsin would provide and unlock a more comprehensive approach suitable for both somatic and axonal stimulation (Govorunova et al. 2022 [↗](#); Vierock et al. 2022 [↗](#)).

## Materials and Methods

### Drosophila optogenetic activation experiments

Flies (*Drosophila melanogaster*) were kept on standard food medium (water, cornmeal, oatmeal, sucrose, yeast, agar, acetic acid, preservative methyl-4-hydroxybenzoate) at 25°C, 60% humidity, and a 12h/12h light-dark cycle. Male flies were collected after eclosion and aged for 7 days in isolation and protected from light on standard food, followed by 3 days on food containing 400  $\mu$ M all-trans retinal (Sigma). The following genotypes were assayed: co-expression of Chrimson and GtACR2: w-; NP2631.GAL4/ UAS-CsChrimson.mVenus. attp40; tubP-FRT-stop-FRT-GAL80, fruFLP/ UAS-GtACR2. attp2; co-expression of Chrimson and tdTomato: w-; NP2631. GAL4/ UAS-CsChrimson.mVenus.attp40; tubP-FRT-stop-FRT-GAL80, fruFLP/ UAS-myr-tdTomato. attp2; co-expression of GtACR2 and tdTomato: w-; NP2631. GAL4/ UAS-myr-tdTomato.attp2; tubP-FRT-stop-FRT-GAL80, fru-FLP/ UAS-GtACR2.attp2.

For optogenetic activation experiments, we used a previously described multi-channel array of electret condenser microphones (CMP-5247TF-K, CUI Inc), amplified with a custom-made circuit board and digitized with a multifunction data acquisition device (NI USB-6259 MASS Term, National Instruments)(“Cellular and Behavioral Functions of Fruitless Isoforms in *Drosophila* Courtship” 2014) (von Philipsborn et al. 2014 [↗](#)). Single male flies were placed in recording chambers equipped with microphones at the bottom and light diodes (Amber 591 nm: Cree 5-mm C503B-AAS/AAN-015 or Blue 470 nm: Cree 5-mm C503B-BAS-CY0C0461) were placed 1 cm above the song recording chambers at a 60° angle, illuminating the whole chamber (10-sec of constant light). Tentative courtship song was detected by a MATLAB script and corrected manually, using a custom made user interface for visualization and annotation of song oscillograms (von Philipsborn et al., 2014 [↗](#)).

### Cell culture, transfection and stable cell line generation

HEK 293 cells (Invitrogen) were grown in DMEM (1g/L glucose) supplemented with 8% FBS and 1% Pen/Strep. For recordings, cells were plated at low density on glass coverslips and kept in DMEM media supplemented with 4% FBS and 1% Pen/Strep. Transfection was done with Xtremegene 9 (Sigma-Aldrich) according to the manufacturer’s instructions. Recordings of the cells were performed between 38-50 hours after transfection.

For generation of recombinant lentivirus for stable HEK 293 cell lines expressing IvfChr and ChrimsonR-tdTomato cells, HEK 293 cells were co-transfected with psPAX2 and pMD2.G (courtesy of Trono laboratory) and the media was transferred to a new dish with untreated cells 2 and 3 days after transfection. The cells receiving the media were kept as stable cell lines of respective constructs.

### Structure alignment and modeling

Homology modeling of ZipACR structure was performed with Modeller within the UCSF Chimera to the published GtACR1 structure (PDB #6CSM). This was further aligned to the GtACR2 alignment from (Govorunova et al. 2017 [↗](#)). Residues surrounding the retinal binding pocket of GtACR1 and the modeled ZipACR were identified and those residues that were different in ZipACR were changed to the corresponding residues of GtACR2 or GtACR1.

## Molecular biology

For GcACR\_457, RsACR\_995 and RlACR\_477, the cDNA were in the original pcDNA3.1 vector as EYFP fusion as available on Addgene (#103137, 103772, 103771). The GtACR2 construct was synthesized (IDT DNA) and modified with a N-terminal signalling peptide from the commercial pDisplay vector (Invitrogen) and a trafficking signal prior to tdTomato or citrine fusion and inserted in the pcDNA3.1+ vector (Invitrogen) for expression. For the ZipACR constructs, the original EYFP fusion from Addgene (#88844) was retained and the construct was either inserted into the pcDNA3.1+ vector or a second generation lentiviral vector pLenti with CMV promoter. No functional or fluorescent differences in the expressed protein were observed between expression from pcDNA3.1+ and pLenti vector in HEK 293 cells and the results were combined for comparison and analysis.

Site-directed mutagenesis of the ZipACR, Chrimson and GtACR2 were performed with overlap extension PCR using Phusion DNA Polymerase (Life Technologies) and ligated into the pcDNA3.1+ or pLenti expression vector.

For ChrimsonR and vfChrimson, additional improvements for membrane trafficking were used as previously described (Bonaventura et al. 2019). In brief, the N-terminal leading sequence from ChR1/oChIEF/ReaChR was transplanted to the transmembrane A-G of ChrimsonR and vfChrimson. A trafficking sequence was added after the transmembrane G before fusion to tdTomato. The modified ChrimsonR/vf-Chrimson-tdTomato constructs were inserted into the 2nd generation lentiviral pLenti vector with CMV promoter and used to make stable ChrimsonR/vfChrimson-tdTomato HEK 293 cell lines. The improved vf-Chrimson is named IvfChr.

For the comparison of IvfChr-citrine and Zip(151V)-Kv2.1-2A-IvfChr-citrine-Kv2.1 (ZipV-IvfChr) photocurrents, the two cDNAs were inserted into pcDNA3.1+ for expression. The IvfChr in both designs contains the modified N-terminal sequence from ChR1/oChIEF/ReaChR but the trafficking sequence was only used in the IvfChr-citrine design and not the Kv2.1 design.

For the expression of HcKCR1, HcKCR2 and WiChR, the mammalian codon-optimized sequences of the 3 potassium-selective ChRs were synthesized as gBlocks (IDT DNA) based on the published sequences and ligated into pcDNA3 or pcDNA3.1H(+) vectors 5' in-frame of coding sequences of EYFP or citrine using BamHI and NotI sites. Trafficking sequences were placed between potassium-selective ChRs and fluorescent protein. N-terminal signaling peptide from IvfChr was placed in some constructs tested. Membrane currents were similar in the presence and absence of N-terminal signaling peptides and the results were combined.

## rAAV production

rAAV2-retro containing vfChrimson\_EYFP\_Kv2.1, IvfChr\_Citrine\_Kv2.1, Zip(151T)-Kv2.1\_IRES\_DsRed, Zip(151T)-Kv2.1-2A-IvChrimson-citrine-Kv2.1 (ZipT-IvfChr), and Zip(151V)-Kv2.1-2A-IvfChr-citrine-Kv2.1 (ZipV-IvfChr) were produced according to the protocol at <https://www.salk.edu/science/core-facilities/viral-vector-core/resources/>. In brief, HEK 293A cells (Life Technologies, Carlsbad, CA) were grown to 90% confluence and transfected with vectors containing opsin proteins, and the helper plasmids XX6-80 and rAAV2-retro (gift from A. Y. Karpova, Howard Hughes Medical Institute Janelia Farm Research Campus). rAAV2-retro were released and purified according to protocol at [https://www.takarabio.com/documents/User%20Manual/6232/6232\\_e.v1509Da.pdf](https://www.takarabio.com/documents/User%20Manual/6232/6232_e.v1509Da.pdf).

Alternatively, virus vectors were produced at the University of Copenhagen, using triple-transfection protocol and purified with an iodixanol density gradient column following (Challis et al., 2019) protocol. In brief, 420 million HEK293t cells were seeded in DMEM 1965 (Glutamax, 10% FBS, 1% P/S). The day after, cells were transfected with 400 µg DNA in a 1:4:2 ratio (pAAV: rep/cap:

pHelper), using linear PEI polyethylenimine (PEI, MW 25000). 18 hrs post-transfection, the media was exchanged to low serum DMEM 1965 (Glutamax, 1% FBS, 1% P/S) and incubated for 72 hrs before harvesting AAV vector from both media and cells using PEG8000. The AAVs were purified from an iodixanol density gradient column following 15 hrs ultracentrifugation (rotor SW28, 28000 rpm, 4 °C). Purity was confirmed with SDS-PAGE, and titer (vg/ mL) was determined with Quant-iT PicoGreen dsDNA Assay Kit (Invitrogen).

## Electrophysiological characterization and imaging of HEK 293 cells

Whole-cell patch-clamping was performed with Axopatch 200A patch-clamp amplifier with Digidata 1322A data acquisition board and pCLAMP10 software on an Olympus IX73 inverted microscope. Cells were clamped at -60mV except in experiments with the ramp protocol and experiments comparing the 590 nm light-induced photocurrents between IvfChr-citrine and ZipV-IvfChr expressing cells. In the experiment comparing the IvfChr-citrine photocurrent to the 2A configuration, cells were clamped at -40mV to reduce the influence of ZipACR response with IvfChr photocurrent. Series resistance for the recorded cells were mostly under 15 MΩ and series resistance were compensated up to 70% during recordings where possible. For comparison of IvfChr and vfChrimson photocurrent in ZipV-IvfChr and BiPOLES-vfChrimson, the potassium gluconate based intracellular solution was used and cells were voltage-clamped at -40 mV (close to the reversal potential of gtACR2 and ZipACR) when stimulated with 590 nm light.

The extracellular solution was composed of (in mM): 145 NaCl, 3 KCl, 2 CaCl<sub>2</sub>, 1 MgCl<sub>2</sub>, 20 D-glucose and 10 HEPES (pH 7.35). For most experiments, the intracellular solution contained (in mM): 110 CsCl, 15 TEACl, 10 HEPES, 5 Cs<sub>2</sub>EGTA, 1 MgCl<sub>2</sub> and 0.1mM CaCl<sub>2</sub> (pH 7.25). In experiments comparing the 590 nm light-induced photocurrents between IvfChr-citrine and ZipV-IvfChr expressing cells and some ramp experiments, a K-gluconate based intracellular solution containing: 115 K-gluconate, 10 K<sub>2</sub>EGTA, 10 HEPES, 5 NaCl, 10 KCl, 2 MgATP and 0.1 Tris-GTP (pH 7.25), was used. For potassium-selective ChR recordings, only the K-gluconate-based intracellular solution was used.

For imaging the expression pattern of channelrhodopsin-expressing HEK 293 cells, cells on coverslips were imaged with an Olympus BX51 equipped with a 40x water immersion objective and a Hamamatsu Orca Flash 4.0LT sCMOS camera and X-cite LED110 light source. Images were acquired using Micromanager 1.4 with no gain and no electron multiplication and exposure time of 250 ms. Citrine and EYFP expression were imaged with a GFP filter cube (Semrock, FF02-470/30 excitation, FF01-520/35 emission and FF495-Di03 dichroic) and tdTomato was imaged with a TRITC filter cube (Semrock, FF01-543/22 excitation, FF01-593/40 emission and FF562-Di03 dichroic). For co-relating photocurrent amplitude and membrane expression, pre-recording images of the cells were captured with an Olympus IX73 microscope with a 40x air objective equipped with Photometrics Cascade II 512 EM-CCD camera and excitation by the Thorlabs 4 channel LED (LED4D118 with 405, 470, 530 and 590 nm channels and only 470 nm and 530 nm light were used for imaging with LED driving currents of 150 and 80 mA, respectively). An identical filter set was used as in the Olympus BX51 microscope. Images were acquired using Micromanager 1.4 with no gain and no electron multiplication and exposure time of 250 ms.

For most channelrhodopsin stimulation experiments 10 mW/ mm<sup>2</sup> of 470 nm and 590 nm stimulation lights from Thorlabs 4 channel LED light source (LED4D118) was delivered to the specimen plane. The light intensity was controlled with a Thorlabs DC4104 4 channel controller electronically by the separate analogue outputs of a HEKA ITC-18 DAQ board under the control of WinWCP 5.5 software. In the spectral response experiment, a Sutter VF-5 system with a smart shutter in the neutral density mode was used to achieve cross spectrum stimulation at the same photon flux at  $1.586 \times 10^{15}$  photons / s / mm<sup>2</sup> (25.1 mW/mm<sup>2</sup> at 400 nm and 15.4 mW/mm<sup>2</sup> at 650 nm) between 400-650 nm (25 nm increments). The VF-5 has the following filters and bandwidth settings: 451-398 nm with 15 nm bandwidth, 503-446 nm with 15 nm bandwidth, 564-498 nm with 14 nm bandwidth, 632-557 nm with 14 nm bandwidth, 703-627 nm with 13 nm bandwidth. A

second Sutter shutter was placed in the light path before the light guide to control the light output. For experiments with ACRs, a 15 sec interval was used between stimulation episodes. For ChromsonR/IvfChr stimulation, stimulation light pulses either have 15 sec intervals between stimulation episodes (no preconditioning) or a 15 sec interval – 1 sec 405 nm light pulse – 15 sec interval stimulation pattern for reconditioning of the protein with 405 nm before the subsequent stimulation. The stimulation light was guided into the fluorescent excitation light path of the Olympus IX73 microscope as with imaging but reflected with a Semrock FF685-DiO2 dichroic mirror to the objective.

Light intensity (in mW/mm<sup>2</sup>) was measured with a Thorlabs slide based light detector (S170C) at the specimen plane with illumination area defined by the microscope aperture and manually calculated for photon flux or illumination intensity.

For measuring the membrane expression of vfChrimson in various modified constructs, BiPOLES and 2A based cassettes, a membrane localized mCherry (encoded by pcDNA3-mCherry-CaaX) was co-expressed and live cell imaging were done on an Olympus Fluoview FV3000 confocal microscope with 40 × 0.8 NA water immersion objective. The citrine fluorescence was acquired with a 488 nm laser excitation and 500-550 emission. mCherry images were acquired with a 594 nm laser excitation and 600-700 nm emission. For the analysis, single in-focus z-planes where the membrane boundaries were clearly defined were used. A straight line was drawn across the cell between 2 membrane boundaries devoid of the nucleus. The pixel intensities of the citrine and mCherry channels were measured to generate the Pearson Correlation coefficient.

## Organotypic slices and virus infection

All procedures involving animals were approved by the Danish Animal Experiment Inspectorate. Hippocampi were isolated from postnatal P5-6 wild-type Sprague-Dawley rats (Janvier). The pups were decapitated and the brain removed and kept in an icy low sodium ACSF solution composed (in mM) of 1 CaCl<sub>2</sub>, 4 KCl, 1 MgCl<sub>2</sub>, 26 NaHCO<sub>3</sub>, 10 D-Glucose, 234 Sucrose, and 0.1% Phenol Red Solution, in milliQ H<sub>2</sub>O, and bubbled with a mixture of CO<sub>2</sub> (5%) and O<sub>2</sub> (95%). From that moment, all the steps were performed in a laminar flow tissue culture hood using sterile equipment and in total asepsis. The brain was poured into a petri dish filled with the low sodium ACSF solution for hippocampus dissection under microscope guidance (Stereomicroscope, Olympus x1 objective). After extraction, the hippocampi were sliced at 400 μm thickness using an automatic tissue chopper (McILWAIN) and moved to a dish with pre-heated culture medium containing MEM Eagle medium 78.8% (Gibco), 20% heat-inactivated horse serum (Gibco), 1 mM L-glutamine, 1 mM CaCl<sub>2</sub>, 2mM MgSO<sub>4</sub>, 170 nM insulin, 0.0012% ascorbic acid, 12.9 mM D-glucose, 5.2 mM NaHCO<sub>3</sub>, 300 mM HEPES (Sigma), pH = 7.2-7.3, osmolality adjusted to 315–325. The slices with intact DG and CA regions were then transferred onto air-fluid interface-style Milli-cell culture inserts (Millipore) in 6-well culture plates (ThermoFisherScientific) with 800 μL of sterile medium added below each insert. The slices were kept in a sterile incubator at 37 °C degrees with 5% CO<sub>2</sub> (ThermoScientific, Steri-cycle i-160). The medium was replaced by a pre-warmed medium (37 °C) three times a week.

After two to three days of culture, the slices were microinjected in DG (or in CA1 when specified) along the characteristic horseshoe pattern with a pulled glass pipette containing the following viruses: retro-IvfChr (titer: 2.5×10<sup>12</sup> particles/mL), retro-Zip(151T) (titer: 9×10<sup>12</sup> particles/mL), retro-ZipT-IvfChr (titer: 2.9×10<sup>12</sup> particles/mL), or retro-ZipV-IvfChr (titer: 1.5×10<sup>12</sup> particles/mL). The injections were done under a microscope (using a Picospitzer III (Parker) connected to a Pulse Pal (10 ms pulse, every 0.5-sec). In the process of evaluating the R-ChR expression system *in vivo*, we noticed variability in the performance between batches from different virus sources. Although we adjusted for the nominal titers given by the providers, some batches produced toxicity while others fared well (data not shown). Therefore, the titers should not be taken as face value, and each batch must be tested in the region of interest before proceeding with the experiment. This concern applicable to any viral expression system has an added complication here. Higher titers, in addition to toxicity, may lead to the leakage of the soma-targeted channels expression in the

terminals, which may cause unintended consequences. Recent developments in engineering soma-targeted motifs and peptides must be considered in future works (“Precision Calcium Imaging of Dense Neural Populations via a Cell-Body-Targeted Calcium Indicator” 2020) (Shemesh et al. 2020 [DOI](#)).

## Whole-cell electrophysiology and light delivery in vitro

Two to three weeks post-infection, the organotypic slices were transferred into the recording chamber and continuously perfused with Artificial Cerebrospinal Fluid containing (in mM): 119 NaCl, 2.5 KCl, 26 NaHCO<sub>3</sub>, 1 NaH<sub>2</sub>PO<sub>4</sub>, supplemented with 11 D-glucose, 4 CaCl<sub>2</sub>, 4 MgCl<sub>2</sub>. Additionally, APV (50μM), NBQX (10μM) and picrotoxin (100μM) were added to block excitatory and inhibitory fast transmission. The solution was adjusted to pH 7.4 (osmolarity ~330 mOsm) and bubbled with a mixture of CO<sub>2</sub> (5%) and O<sub>2</sub> (95%), at room temperature. The chamber was mounted on an upright microscope (Scientifica) linked to a digital camera (QImaging Exi Aqua) and the cells were visualized using 60X water-immersion objective (Olympus, LumiPlan). Acquisitions were performed in whole-cell configuration using Clampex 10.6 connected to a Multiclamp 700B amplifier via a Digidata 1550A digitizer (all from Molecular Devices). Voltage-clamp data were low-pass filtered at 200 Hz and digitized at 10 kHz and the whole-cell capacitance was compensated. Patch pipettes (2–4 MΩ of resistance) were filled with an internal solution containing (in mM): 153 K-gluconate, 10 HEPES, 4.5 NaCl, 9 KCl, 0.6 EGTA, 2 MgATP, and 0.3 NaGTP. The pH and osmolarity of the internal solution were close to physiological conditions (pH 7.4, osmolarity 297 mOsm). The access resistance of the cells in our sample was ~25 MOhm. For voltage-clamp experiments, the Dentate Gyrus granule cells (otherwise specified when done in CA1 pyramidal cells) were clamped at -60mV. For current-clamp experiments, the cells were maintained at -60mV (except for the protocol where the potential was brought from -70 to -40mV) by constant current injection (holding current ± 100pA).

The recorded neurons were illuminated with 470 nm and 635 nm (nominal wavelengths) light with a coolLED pE-4000 system connected to the Digidata via a single TTL. The light irradiance was controlled via the control pod and was adjusted to 4mW/mm<sup>2</sup> for 470 nm and 7 mW/mm<sup>2</sup> for 635 nm (otherwise specified). Light irradiance was measured with a Thorlabs digital optical power meter (PM100D) at the specimen plane with illumination area defined by the microscope aperture. Between the recordings, wavelengths were switched from 470 nm to 635 nm and vice versa. At the end of the recordings, the location of the recorded cell was confirmed by inspection at 10x magnification. Data analysis was performed using Clampfit 10.6 (Molecular device).

After recordings, slices were kept in 4% formalin, and later mounted under coverslips with Fluoromount (Sigma-Aldrich). Pictures were taken with an Apotome.2 (Zeiss) and ZEN software at 10X and 40X magnifications.

## Acute slices

Experimental procedures were approved by the Animal Care and Use Committee of the University of Buenos Aires (CIC-UAL). Briefly, 4-week-old C57 mice (n = 4) were co-injected with AAV-8/2-hSyn1-ChrimsonVF\_EYFP\_Kv2.1 and AAV-8/2-hSyn1-ZipACR\_Kv2.1\_IRES\_DsRed2 within the dentate gyrus of the hippocampus (250nl of a 1:1 mixture, coordinates from Bregma: 1 mm mediolateral, -2 mm anteroposterior, -1.5 dorsoventral). After 3 weeks of expression, the animals were sacrificed and the brain removed and cut in a solution composed of (in mM): 234 sucrose, 11 glucose, 26 NaHCO<sub>3</sub>, 2.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 10 MgSO<sub>4</sub>, and 0.5 CaCl<sub>2</sub> (equilibrated with 95% O<sub>2</sub>–5% CO<sub>2</sub>). The slices were maintained at room temperature before being transferred in a recording chamber mounted on a microscope (Nikon) connected to a Mightex Illumination system for 470 nm and 635 nm light delivery (at 10mW/mm<sup>2</sup>). For current clamp experiments the amount of current injected was corrected in the inter-sweep interval to keep it close to -60 mV. In voltage clamp experiments holding potential was -60 mV.

Recordings were done in a solution of composition (in mM): 126 NaCl, 26 NaHCO<sub>3</sub>, 2.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 2 MgSO<sub>4</sub>, 2 CaCl<sub>2</sub> and 10 Glucose (pH 7.4).

Patch-pipettes (2–4 MΩ of resistance) were filled with a K-glu-conate based internal solution (in mM, 130 K-gluconate, 5 KCl, 10 HEPES, 0.6 EGTA, 2.5 MgCl<sub>2</sub>·6H<sub>2</sub>O, 10 Phospho-creatine, 4 ATP-Mg, 0.4 GTP-Na<sub>3</sub>).

## In vivo expression of ZipT-IvfChr and IvfChr in the facial nucleus

For expression of ZipT-IvfChr and IvfChr in the facial motor nucleus of the brainstem (FN), 8 to 10 weeks old c57Bl6/j mice (males) from Janvier (France) were anesthetized with Isoflurane (5% for induction, 2% for maintenance) and head-fixed on a stereotaxic frame (Model 940, Kopf, California) on top of a warm plate to keep their body-temperature at 37 °C. They were administered carprofen (5 mg/kg, s.c.), then the skin was retracted, and a small craniotomy was drilled above the injection site. Suspension of either ret-roAAV-hSyn-ZipT-IvfChr (titer: 2.9×10<sup>12</sup> particles/mL) or ret-roAAV-hSyn-IvfChr (titer: 2.5×10<sup>12</sup> particles/mL) was injected unilaterally at a volume of 400 nL and a rate of 2 nL per second, using a nanoliter injector (Nanoject III, Drummond, Broomall, PA). The coordinates for injection in the FN were -5.6 mm anteroposterior, -1.4 mm mediolateral and -6.0 mm ventral, relative to bregma. After virus injection, the injection pipette was kept in place for an additional 10 minutes before being retracted. After they recovered from the surgery on a warm-plate, the animals were put back in their home-cages. They were housed in groups of 4 individuals per cage in a normal 12 h light/ dark cycle with food and water ad-libitum. Four to six weeks after the surgery, the animals were anesthetized with Urethane (Sigma-Aldrich, 1.5 mg/kg for first injection, followed by injection of 1/5 of the initial dose until the reflex disappeared) and head-fixed on a KOPF stereotaxic frame (Model 940, Kopf, California). The reflexes were monitored during the entire procedure. The wound on the skin was reopened and the craniotomy above the injection site was re-open when necessary.

For optical stimulation of the FN motoneurons, a 200 μm diameter optic-fiber connected to a laser source (Doric Lenses, Québec, Canada) was implanted above the facial nucleus at coordinates -5.6 mm anteroposterior, -1.4 mm mediolateral relative to bregma, and -4.1 to -4.5 mm ventral to the brain surface. Pulses of 100 ms of either 450 nm or 638 nm illumination at 20 mW/mm<sup>2</sup> were given at 0.33 Hz through a PulsePal trigger generator (Open Ephys) connected to the laser. The whiskers' movements were recorded at 750 FPS with an Optronis Camera (CR3000, Optronis GMBH, Germany) and acquired through TimeBench (2.6.30, Optronis). After acquisition, the videos were exported as AVI with a 25 AVI framerate.

At the end of the experiment, the animals were sacrificed and the brain extracted and kept in Formalin 10% until slicing. 100 μm sections were sliced with a Vibratom (Leica, VT1200S) and counterstained with DAPI (ThermoFisher, D1306) and eGFP (ThermoFisher, CAB4211) for histological localization of the virus expression, and imaged with a ZEISS Apotome microscope (Axio Imager M2). The experimenters were not blind to the group's status; however, the automated analysis did not require blindness as this would not affect the outcome of the experiment.

## Whiskers tracking and analysis

The whiskers were tracked offline with a custom MatLab tool-box (available on GitHub [GitHub](https://github.com/NabaviLab-Git/WhiskerTrack) - NabaviLab-Git/WhiskerTrack: WhiskerTrack is a MATLAB toolbox to track the whiskers automatically with high accuracy.). In brief, 5 to 10 whiskers per animal were tracked frame by frame by using an automatic search method to fit the best line to the linear part of each whisker. The slope of this line was considered as the angle of its respective whisker. The absolute changes in the angle of each whisker were compared to its baseline and calculated as a Z-Score based on the mean and standard deviation.

The area under curve (AUC) was calculated as  $AUC = Z \times W$  with Z being the averaged Z-Score for the maximum duration of the response and W being the window of the maximum response (200 ms from the onset of the light pulse).

For comparison between ZipT-IvfChr and IvfChr, we averaged the AUC of the responses to 450 nm or 635nm illumination over 3 trials and calculated the ratio of the AUC to 450 nm over 635nm. Statistical significance was calculated with a parametric t-test with MatLab.

## Data Analysis

The data were analyzed and plotted using GraphPad Prism V9 (GraphPad Software, La Jolla, CA, USA). All values are indicated as mean  $\pm$  standard deviation (SD) or mean  $\pm$  standard error of the mean (SEM), as specified. For statistical comparison between groups, data were tested for normal distribution using D'Agostino and Pearson, Shapiro-Wilk or KS normality test. Comparison of the mean between groups has been done with a one-way analysis of variance (ANOVA) followed by Dunnett's multiple comparisons test.

ZEN software (Zeiss) and FIJI (from ImageJ) were used for images processing.

Figure 1A [1B](#) [1C](#) [3E](#) [4D](#) and 5B1 [5C](#) have been created with BioRender.com.

## Data availability

The datasets used and/or analyzed during the current study are available at the following link: <https://github.com/NabaviLab-Git/Dual-color-optical-activation-and-suppression-of-neurons-with-high-temporal-precision> [1](#).


## Acknowledgements

We thank the members of the Nabavi lab, in particular Islam Faress for suggestions, and Mariam Gamaleldin for the preparation of hippocampi organotypic slices. We also thank Dr. Sergio Almeida for his help on HEK 293 cells culture at the early stage of the project, Peter Kerwin from the Philipsborn lab for his help on Drosophila work, and Anne-Katrine Vestergaard, Kathrine Meinecke Christensen and Sanaz Ansarifard for technical assistance. We are thankful to Dr. Thomas Mager and Dr. Ernst Bamberg for their generous gift of vfChrimson DNA study was supported by an ERC starting grant to SN (22736), by the Danish Research Institute of Translational Neuroscience to S.N (19958), by PROMEMO (Center of Excellence for Proteins in Memory funded by the Danish National Research Foundation) to SN (DNRF133), by the Australian Research Council Future Fellowship (FT160100056) to JYL, NIH BRAIN Initiative grant to JYL (R21EY027620), by a Lundbeck NIH Brain Initiative grant to SN and JYL (R360-2021-650 and R273-2017-179), and by a Lundbeck Foundation grant to Andreas Toft Sørensen (R346-2020-1793 (ATS)).

## Additional information

### Authors contribution


The project was designed by SN, JYL and NM-J. The *in vitro* experiments on HEK 293 cells and organotypic slices were done by NM-J, SN and JYL. JP did the recordings on hippocampal acute slices. The *in vivo* experiments were done by NM-J and AM. Analysis of the whisking was done by NM-J and MN. AZ and JYL did HEK 293 cell imaging. NM-J did the surgeries. NK contributed to the



**Figure 1 - figure supplement 1.**

**Red-shifted ChR variants are activated by 470 nm and 590 nm pulses of light.**

(A) Typical photocurrents (left and middle traces) and on-kinetics (right traces) of ChrimsonR and IvfChr in HEK 293 cells with 590 nm and 470 nm illumination at 10 mW/mm<sup>2</sup>. The traces have been recorded 15-s after a 405 nm preconditioning pulse. (B) Comparison of 590/470 nm ratio for ChrimsonR and IvfChr, measured at peak. Each dot represents a cell (mean ± SD). (C) Blue-light illumination of the Chrimson variants induces firing as efficiently as red-light stimulation. This property is represented here with two examples of traces obtained from ChrimsonR expressing CA1 neurons and vfChrimson expressing DG neurons, in response to 10 Hz of 5 ms pulses at 635 nm and 470 nm.



**Figure 2 - figure supplement 1.**

### Improved membrane trafficking of vfChrimson and ChR co-expression designs.

(A) Representative images of various vfChrimson based expression cassette in HEK 293 cells and membrane tethered mCherry. (Left column) the expression pattern of palmitoylated mCherry on the cellular membrane (using a lysine rich and CaaX sequence). (Middle column) images of the citrine attached to vfChrimson in the various designs. (Right column) overlaid images of the two channels for co-expression visualization. vfChrimson is directly tethered to citrine (vfChrimson), or tethered to citrine with addition of trafficking sequence between ChR and citrine (C-vfChrimson), with the addition of N-terminal signalling peptide (from oChIEF, vfChrimson-ts) and with the inclusion of both trafficking sequence and N-terminal signalling peptide (denoted IvfChr). (Bottom panel) the comparison of the Pearson correlation coefficients of mCherry and citrine fluorescence intensities of the different constructs. Statistical tests were performed with Kruskal-Wallis test followed by Dunn's multiple comparison tests between all pairs. IvfChr has the most optimized membrane trafficking followed by the 2A-based ZipV-IvfChr. (B) Left, examples of BiPOLES-vfChrimson and ZipV-IvfChr response to 590 nm light illumination at -40 mV holding potential. Right, peak photocurrent amplitude comparison between BiPOLES-vfChrimson and ZipV-IvfChr adjusted to membrane capacitance. Unpaired t-test was used. (C) Example of membrane enriched expression of 2A-based ZipT-IvfChr and ZipV-IvfChr in organotypic slices. (D) Comparison of the amplitudes of photocurrent of IvfChr and ZipV-IvfChr in HEK 293 cells to 590 nm light illumination.

Figure 2 - figure supplement 2.

### Membrane trafficking of the improved ultrafast vf-Chrimson, IvfChrimson (IvfChr).

(A) Off-time constant of C-ChrimsonR-ts and IvfChr after 1-s of 635 nm illumination at 10mW/mm<sup>2</sup>. (B) Photocurrent amplitudes at the peak of C-ChrimsonR-ts and IvfChr, adjusted to membrane fluorescence (left), and comparison of current density and membrane fluorescence (right). The recordings have been obtained from HEK 293 cells at -60 mV. Each dot represents the values from a single cell. The horizontal black bars represent the mean  $\pm$  SEM for (A) and mean  $\pm$  SD for (B). Statistics for panels (A) \*\*\*\*p<0.0001; unpaired t-test and (B) \*p<0.05; unpaired t-test (Welch correction).






Figure 3 - figure supplement 1.

### Channel properties of the ZipACR variants.

(A) 590/470 nm ratio measured from the ACR variants in HEK 293 cells. Each circle corresponds to value obtained from one cell (mean  $\pm$  SD, n = 3-7). (B) Reversal potential of the selected variants Zip(151T) and Zip(151V) compared to the original ZipACR with cesium (Cs) or potassium gluconate (K) intracellular solution. Each dot corresponds to the value obtained from one cell (mean  $\pm$  SD, n = 5-6). (C) Typical I/V-relationship for the two variants and ZipACR. The recordings have been done with- or without 470 nm LED illumination.






**Figure 3 - figure supplement 2.**

### Light-induced responses and action spectra of ZipACR variants.


(A) Representative responses of the original ZipACR, Zip(151T) and Zip(151V) in HEK 293 cells in response to different light intensities. (B) Normalized photocurrent amplitude in response to 400-650 nm and to 650-400 nm light pulses (as presented in Fig. 3D2). The values have been measured on the photocurrents at the peak. The responses were normalized to the maximum response of each cell across the spectrum. The data points are the mean value  $\pm$  SEM ( $n = 6-7$ ).



**Figure 3 - figure supplement 3.**

**(A) The traces of the two outliers in Fig.3 E2.**


(A1), 470 nm illumination reduced the amplitude of the spikes, but did not abolish the firing. (A2), 470 nm illumination as well as 635 nm light blocked the firing. This cell expressed Zip(151T) channels which was significantly more red-shifted compared to the mean (635/470 nm ratio: 23% for this cell vs 6% for the averaged population). (B) For a range of imposed potentials 470 nm illumination of Zip(151T) does not trigger action potentials. (B1) Typical traces of the DG neurons at various imposed potentials, in response to 10 Hz of 470 nm light illumination. The dotted lines show the potential of the maximal response. (B2) Plot showing the responses in current-clamp of the 10 neurons recorded at -70, -60, -50 and -40 mV imposed potential. Each dot represents a single cell.



**Figure 4 - figure supplement 1.**

### ZipV-IvfChr variant, as for ZipT-IvfChr, is effective in preventing blue-light induced action potentials in-vitro.


(A) Example of the response (A1) of ZipT-IvfChr expressing neurons to 470 nm and 635 nm light pulses, and to a co-illumination protocol (470+635 nm). ZipT-IvfChr is efficient in preventing the firing of APs to 470 nm and to a co-illumination protocol (A2, n = 6). (B) Blue and red light-driven spike fidelity at various light intensities in DG cells (n = 4-8 cells for each opsin; 10 ms pulse width, single pulse). Under 470 nm illumination, neurons expressing ZipV-Ivf-Chr, occasionally fired APs, which may be explained by the variant relatively faster closure rate (see Fig. 3B). (C-D) Comparison of the high-frequency red and blue light-induced spiking (B), and spike probability (C, n = 6-13 for each opsin), of DG cells expressing IvfChr or the Zip-IvfChr variants, with 10 and 20 Hz of light pulses (4-7 mW/mm<sup>2</sup>). Holding potential = -60 mV. (E) I-V curve measured for both variants ZipT-IvfChr and ZipV-IvfChr under 635 nm or 470 nm illumination (n=8 per variant).



**Figure 4 - figure supplement 2.**

#### Desensitization of IvfChr in HEK 293 cells.

(A) Representative responses of IvfChr to 1 sec of 470 nm and 590 nm with or without a preconditioning pulse at 405 nm. Desensitization is much less with 470nm light stimulation. (B) Normalized responses in voltage-clamp to 590 nm pulses of light, at t = 0, t = 15 and t = 60 sec without 405 nm preconditioning. The desensitization of IvfChr to 590 nm light does not recover fully at least within one minute. A pre-reconditioning pulse at 405 nm enhanced the recovery of the 590 nm responses.



**Figure 5 - figure supplement 1.**

### Evaluation of light-gated potassium channels HcKCR1, HcKCR2 and WiChR.

(A) Example of electrophysiological recordings of HEK293 cells expressing HcKCR1, HcKCR2 and WiChR to 1 s of 470 nm light (HcKCR2 and WiChR) or 530 nm light (HcKCR1) at various membrane holding potentials. (B) Reversal potential generated with a ramp protocol. Dark gray light is the ramp profile in the absence of light and light gray light is the ramp profile during light illumination. (C) Measurements of the channels reversal potential. Each dot is a measurement from one cell and the lines represent the mean  $\pm$  S.D. (D) The channel off-rate time constant measured at -10 mV in response to 530 nm light (HcKCR1) and 470 nm light (HcKCR2 and WiChR). Each dot is a measurement from one cell and the lines represent the mean  $\pm$  S.D. (E1) Examples of recordings of HEK293 cells expressing HcKCR1 (left), HcKCR2 (middle) and WiChR (right) to 1 s of 10 mW/mm<sup>2</sup> light at the indicated wavelengths. The cells were held at -10 mV with voltage-clamp. (E2) The amplitudes of light-evoked response of HcKCR1, HcKCR2 and WiChR to 10 mW/mm<sup>2</sup> of 405nm, 470 nm, 530 nm, and 590 nm light is normalized to the peak response of the cells for each of the four tested wavelengths. HcKCR1 has the peak response at 530 nm and HcKCR2 and WiChR have the peak responses at 470 nm.

|                  |                                                               |                      |
|------------------|---------------------------------------------------------------|----------------------|
| ZipACR           | 2A                                                            | Trafficking sequence |
| 151T             | ChIEF/ReaChR N-terminus                                       | citrine              |
| KV2.1 signalling | vfChrimson                                                    | KV2.1 signalling     |
| 1                | M A A L H Q A L E E L G Q G A S R H L L                       | 20                   |
| 1                | ATGGCCGCGCTCCATCAGGCTCTGGAGGAGCTGGGACAGGGAGCCAGGCACCTCCTG     | 60                   |
| 21               | Y R A P E E M V Y G K W V E A N P R C I                       | 40                   |
| 61               | TACAGAGCCCCGAGGAGATGGTGTATGGCAAGTGGGTGGAGGCCAACCTAGGTGCATC    | 120                  |
| 41               | E L H D H S D Y V Y V F Q L C F A V V C                       | 60                   |
| 121              | GAAGTGCACGACCACTCCGACTACGTGTACGTGTTCAGCTGTGCTTCGCCGTGGTGTGC   | 180                  |
| 61               | A C Q V I F M F T R A P N V G W E A I Y                       | 80                   |
| 181              | GCCTGCCAGGTGATCTTCATGTTACACAGGGCCCCAACGTGGGCTGGGAAGCCATCTAT   | 240                  |
| 81               | L P L A E V V T Y S I A A N G E G V L R                       | 100                  |
| 241              | CTGCCCTTGGCCGAGGTGCTGACCTACAGCATGCGCCCAACGGAGAGGGCGTGTGAGG    | 300                  |
| 101              | M A D G R Y F N F A K L A G W A V C C P                       | 120                  |
| 301              | ATGGCCGACGGCAGGTACTTCAATTTGCGCAAACTGGCCGATGGGCCGTGTGCTGCCCC   | 360                  |
| 121              | I M L I Q I G G M A Q I K Y R T I P L N                       | 140                  |
| 361              | ATTATGCTCATCCAGATCGGAGGCATGGCCAGATCAAGTACAGGACCATCCCCCTGAAC   | 420                  |
| 141              | N V V L A A S L N R I F G M A S A I T                         | 160                  |
| 421              | AACGTGGTGTGGCCGCTAGCCTGAATAGGACCATCTTCGGCATGGCCAGCGCCATCACA   | 480                  |
| 161              | A S D P A R W G F Y F C A W I C Y L T E                       | 180                  |
| 481              | GCCTCCGACCTGTAGATGGGCTTTTACTTCTGCGCTGGATCTGTTACCTGACCGAG      | 540                  |
| 181              | V G I T L T I M A V A I S D F S K I K T                       | 200                  |
| 541              | GTCCGCATCACCTGACCATCATGGCCGTGGCCATCAGCGACTTCAGCAAGATCAAGACA   | 600                  |
| 201              | E L G Q W V V G R I O T M R I I F L V A                       | 220                  |
| 601              | GAGCTCGGCCAGTGGTGGTGGGCCGATTCAAACCATGAGGATCATCTTTCTGGTCGCC    | 660                  |
| 221              | W T S F P V V W V L G Y T G F C V I H E                       | 240                  |
| 661              | TGGACCAAGCTTCCCCGCTGCTGGTGGTGGGCTACACCGGATTCTGCGTGATCCACGAG   | 720                  |
| 241              | D Y I A L L Y L F A D L L S K N T W G V                       | 260                  |
| 721              | GACTACATCGCCCTGCTGTATCTGTTCCGCGACCTGCTCTCCAAGAACCTGGGGAGTG    | 780                  |
| 261              | M M W H T T W V K L N G K W D R E F A A                       | 280                  |
| 781              | ATGATGTGGCATACCACTGGGTGAACTCAACGGCAATGGGACAGGGAATTGCGGCC      | 840                  |
| 281              | A G G H E A L K K A L E Q D V E I G A G                       | 300                  |
| 841              | GCCGGAGGACACGAGGCTCTGAAGAAGGCCCTGGAGCAGGACGTGGAATCGGCGCCGGC   | 900                  |
| 301              | E K N Q N Q L T A R Q A A A D S Q P I I                       | 320                  |
| 901              | GAGAAGAACCAGAACCAGCTGACCGCCAGGCAGGCGGCCGCGCAGAGCCAGCCATCTTG   | 960                  |
| 321              | N T K E M A P O S K P P E E L E M S S Y                       | 340                  |
| 961              | AATACCAAGGAGATGGCCCCCAGAGCAAGCCCCCGAGGAGCTGGAGATGAGCAGCATG    | 1020                 |
| 341              | P S P V A P L P A R T E G V I D M R S Y                       | 360                  |
| 1021             | CCGACCCCGTGGCCCCCTGCCCGCCAGAACCAGGGCGTGATAGATATGAGTCTATG      | 1080                 |
| 361              | F S I D S F I S C A T D F P E A T R A                         | 380                  |
| 1081             | AGCAGCATCGACAGCTTCATCTCATGTGCAACTGACTTCCCCGAGGCCACCAGATTCgt   | 1140                 |
| 381              | S G G S G G E G R G S L L T C G D V                           | 400                  |
| 1141             | agcggaggtggtatctgtgtggtgagggcagaggaagtcttctaactgcggtgacgtg    | 1200                 |
| 401              | E E N P G P R T M V S R R P W L L A L A                       | 420                  |
| 1201             | gaggagaaatccggccctcgtagcATGGTGAGCAGAAGACCTTGCTGCTGGCCCTGGCC   | 1260                 |
| 421              | L A V A L A A G S A G A S T G S D A T V                       | 440                  |
| 1261             | CTGGCCGTGGCCCTGGCCGCGCAGCGCCGCGCCAGCACCGGCAGCGACGCCACCGT      | 1320                 |
| 441              | P V A T Q D G P D Y V F H R A H E R M I                       | 460                  |
| 1321             | CCCGTGGCCACCCAGGACGGCCCCGACTACGTGTTCACAGAGCCCACGAGAGAATGTG    | 1380                 |
| 461              | F Q T S Y T L E N N G S V I C I P N N G                       | 480                  |
| 1381             | TTCCAGACCAAGCTACACCTGGAGAACACGGCAGCGTGATCTGCATCCCCAACACGGC    | 1440                 |
| 481              | Q C F C L A W I L H S R G T P G E K I G                       | 500                  |
| 1441             | CAGTGCTTCTGCCTGGCCTGGCTGctgcacagcagaggcacaccaggagaaaaagatcggc | 1500                 |
| 501              | A O V C O W I A P S I A I A L L T F Y G                       | 520                  |
| 1501             | gcccaggtctgccagtggtgattgtttcagcatcgccatcgccctgctgacattctacggc | 1560                 |
| 521              | I S A W K A T C G W E E V Y V C C V E V                       | 540                  |
| 1561             | ttcagcgctggaaggccacttgcggttgaggagggtctacgtctgttgcgtcgaggtg    | 1620                 |
| 541              | L F V T L E I F K E F S S P A T V Y L S                       | 560                  |
| 1621             | ctgttcgtgacctggagatcttcaaggagttcagcagccccgccacagtgtagctgtct   | 1680                 |
| 561              | T G N H A Y C L R Y F E W L L S C P V I                       | 580                  |
| 1681             | accggcaaccagcctattgcctgcgctacttcagtggtgctgctgtcttgcctcgatc    | 1740                 |
| 581              | L I R L S N L S G L K N D Y S K R T M G                       | 600                  |
| 1741             | ctgataCGCctgagcaacctgagcggcctgaagaacgactacagcaagcgaccatgggc   | 1800                 |
| 601              | L I V S C V G M I V F G M A A G L A T D                       | 620                  |
| 1801             | ctgatcgtgtcttgcgtgggaatgatcgtgttcgcatggtggtggtggtggtggtggtg   | 1860                 |
| 621              | W L K W L L Y I V S C I Y G G Y M Y F G                       | 640                  |
| 1861             | tggctcaagtggtgctgtatctcgtgtcttgcacatcagcggtgctacatgtacttcag   | 1920                 |
| 641              | A A K C Y V E A N H S V P K G H C R M V                       | 660                  |
| 1921             | gcccgaagtgtacgtggaagccaaccacagcgtgcctaagggcattgcccagtggtc     | 1980                 |
| 661              | V K L M A Y A F F A S W G M Y P I L W A                       | 680                  |
| 1981             | gtgaagctgatggcctacgtttcttcgcctcttggggcatgtaccaatcctctggtgca   | 2040                 |
| 681              | V G P E G L L K L S P Y A N S I G H S I                       | 700                  |
| 2041             | gtgggaccagaaggactgctgaagctgagcccttacgccaacagcatcgccacagcatc   | 2100                 |

## Supplementary Material 1

### The annotated sequence of the 2A based ZipT-IvfChr design used in this study.

For ZipACR(151V), position 151 of Zi-pACR is valine and encoded by the codon GTC and the sequence of IvfChrimson-citrine starts at polypeptide sequence position 409. Note that the flexible linkers and amino acids introduced by the digestion sites are not annotated.

```

701  C D I I A K E F W T F L A H H L R I K 720
2101 tgcgacatcatcgccaaggagttttggaccttctctggcccaccacctgaggatcaagatc 2160
721  H E H I L I H G D I R K T T K M E I G G 740
2161 cacgagcacatcctgatccacggcgacatccggaagaccaccaagatggagatcggaggc 2220
741  E E V E V E E F V E E E D E D T V L E 760
2221 gaggaggtggaagtggaagagttcgtggaggaggaggacgaggacacagtgcTCGAGAag 2280
761  S R I T S E G E Y I P L D Q I D I N V M 780
2281 agcaggatcaccagcgagggcgagtacatccccctggaccagatcgacatcaacgtgatg 2340
781  V S K G E E L F T G V V P I L V E L D G 800
2341 gtgagcaagggcgaggagctgttcacccgggtggtgcccacctcctggcgagctggacggc 2400
801  D V N G H K F S V S G E G E G D A T Y G 820
2401 gacgtaaacggccacaagttcagcgtgtccggcgagggcgagggcgatgccacctacggc 2460
821  K L T L K F I C T T G K L P V P W P T L 840
2461 aagctgaccctgaagtctcatctgcaccaccggcaagctgcccgctgccctggcccaccctc 2520
841  V T T F G Y G L M C F A R Y P D H M K Q 860
2521 gtgaccaccttcggctacggcctgatgtgtctcgcccgctaccccgaaccacatgaagcag 2580
861  H D F F K S A M P E G Y V Q E R T I F F 880
2581 cagacttcttcaagtcgcccattgcccgaaggtacgtccaggagcgcaccatcttcttc 2640
881  K D D G N Y K T R A E V K F E G D T L V 900
2641 aaggacgacggcaactacaagaccggcgccgaggtgaagttcgagggcgacaccctgggtg 2700
901  N R I E L K G I D F K E D G N I L G H K 920
2701 aaccgcacgagctgaagggcatcgacttcaaggaggacggcaacatcctggggcacaag 2760
921  L E Y N Y N S H N V Y I M A D K Q K N G 940
2761 ctggagtacaactacaacagccacaacgtctatatcatggccgacaagcagaagaacggc 2820
941  I K V N F K I R H N I E D G S V Q L A D 960
2821 atcaaggtgaacttcaagatccgccacaacatcgaggacggcagcgtgcagctcgccgac 2880
961  H Y Q Q N T P I G D G P V L L P D N H Y 980
2881 cactaccagcagaacacccccatcgggcgacggccccgtgctgctgcccgaaccactac 2940
981  L S Y Q S A L S K D P N E K R D H M V I 1000
2941 ctgagctaccagtcgcgcctgagcaagaccccaacgagaagcgcgatcacatggtcctg 3000
1001  L E F V T A A G I T L G M D E L Y K Q S 1020
3001 ctggagttcgtgaccgcgcgggatcactctcgccatggacgagctgtacaagCAGTCC 3060
1021  G P I L N T K E M A P Q S K P P E E L E 1040
3061 CAACCCATACTGAACACGAAGGAAATGGCTCCTCAAAGTAAACCGCCCGAAGAAGTCTCGAa 3120
1041  M S S M P S P V A P L P A R T E G V I 1060
3121 ATGAGTTCAATGCCATCACCTGTGGCTCCGCTTCCCGCCGCACGGAAGGTGTCATCGAC 3180
1061  M R S M S S I D S F I S C A T D F P E A 1080
3181 ATGCGAAGCATGTCTAGTATTGACTCATTTATTAGCTGCGCCACCGATTTCCTGAAGCT 3240
1081  T R E * 1083
3241 ACTCGGTTCTAA 3252

```

## Supplementary Material 1 (continued)

| ZipT-IvfChr | Animal | Trial number | AUC (Red) | Average AUC (Red) | AUC (Blue) | Average AUC (Blue) | Ratio (Blue/Red) |
|-------------|--------|--------------|-----------|-------------------|------------|--------------------|------------------|
|             | #1     | 1            | 0,549     | 0,506             | 0,119      | 0,078              | 0,155            |
|             |        | 2            | 0,509     |                   | 0,080      |                    |                  |
|             |        | 3            | 0,460     |                   | 0,036      |                    |                  |
|             | #2     | 1            | 0,386     | 0,480             | 0,126      | 0,083              | 0,174            |
|             |        | 2            | 0,484     |                   | 0,041      |                    |                  |
|             |        | 3            | 0,570     |                   | 0,083      |                    |                  |
|             | #3     | 1            | 0,372     | 0,250             | 0,037      | 0,005              | 0,018            |
|             |        | 2            | 0,217     |                   | -0,027     |                    |                  |
|             |        | 3            | 0,161     |                   | 0,004      |                    |                  |
|             | #4     | 1            | 0,086     | 0,098             | 0,023      | 0,023              | 0,234            |
|             |        | 2            | 0,120     |                   | 0,007      |                    |                  |
| 3           |        | 0,087        | 0,038     |                   |            |                    |                  |
|             |        |              |           |                   |            | Mean               | 0,145            |
|             |        |              |           |                   |            | SEM                | 0,046            |

| IvfChrimson | Animal | Trial number | AUC (Red) | Average AUC (Red) | AUC (Blue) | Average AUC (Blue) | Ratio (Blue/Red) |
|-------------|--------|--------------|-----------|-------------------|------------|--------------------|------------------|
|             | #1     | 1            | 0,613     | 0,533             | 0,430      | 0,451              | 0,845            |
|             |        | 2            | 0,540     |                   | 0,460      |                    |                  |
|             |        | 3            | 0,447     |                   | 0,463      |                    |                  |
|             | #2     | 1            | 0,463     | 0,482             | 0,395      | 0,312              | 0,649            |
|             |        | 2            | 0,503     |                   | 0,284      |                    |                  |
|             |        | 3            | 0,479     |                   | 0,259      |                    |                  |
|             | #3     | 1            | 0,603     | 0,572             | 0,259      | 0,267              | 0,467            |
|             |        | 2            | 0,541     |                   | 0,275      |                    |                  |
|             |        | 3            | 0,571     |                   | 0,267      |                    |                  |
|             | #4     | 1            | 0,574     | 0,580             | 0,375      | 0,427              | 0,737            |
|             |        | 2            | 0,535     |                   | 0,469      |                    |                  |
|             |        | 3            | 0,631     |                   | 0,438      |                    |                  |
| Mean        |        |              |           |                   |            | 0,674              |                  |
| SEM         |        |              |           |                   |            | 0,080              |                  |

## Supplementary material 2

Area under the curve (AUC) for the whiskers protraction triggered by red or blue-light illumination of the facial nucleus neurons expressing ZipT-IvfChr or IvChrimson.

early stage of the project with surgeries. BE and A-vP performed the experiments on drosophila. RCB and ATS were responsible for virus production. NM-J, AM and JYL designed the figures. NM-J, SN and JYL wrote the manuscript. All authors discussed the results and contributed to the revision of the manuscript.

## References

- Bonaventura Jordi, Eldridge Mark A. G., Hu Feng, Gomez Juan L., Sanchez-Soto Marta, Abramyan Ara M., Lam Sherry, et al. 2019) **High-Potency Ligands for DREADD Imaging and Activation in Rodents and Monkeys** *Nature Communications* **10**:4627 <https://doi.org/10.1038/s41467-019-12236-z>
- Deisseroth Karl, Hegemann Peter 2017) **The Form and Function of Channelrhodopsin** *Science* **357** <https://doi.org/10.1126/science.aan5544>
- Douin Victorine, Bornes Stephanie, Creancier Laurent, Rochaix Philippe, Favre Gilles, Prats Anne-Catherine, Couderc Bettina 2004) **Use and Comparison of Different Internal Ribosomal Entry Sites (IRES) in Tricistronic Retroviral Vectors** *BMC Biotechnology* **4**:16 <https://doi.org/10.1186/1472-6750-4-16>
- Ellendersen Bárður Eyjólfsson, von Philipsborn Anne C 2017) **Neuronal Modulation of D. Melanogaster Sexual Behaviour** *Current Opinion in Insect Science, Neuroscience \* Pheromones* **24**:21–28 <https://doi.org/10.1016/j.cois.2017.08.005>
- Faress Islam, Khalil Valentina, Hou Wen-Hsien, Moreno Andrea, Andersen Niels, Fonseca Rosalina, Piriz Joaquin, Capogna Marco, Nabavi Sadegh 2024) **Non-Hebbian plasticity transforms transient experience into lasting memories** *eLife* **12**:RP91421 <https://doi.org/10.7554/eLife.91421>
- Gharaei Saba, Honnuraiah Suraj, Arabzadeh Ehsan, Stuart Greg J. 2020) **Superior Colliculus Modulates Cortical Coding of Somatosensory Information** *Nature Communications* **11**:1693 <https://doi.org/10.1038/s41467-020-15443-1>
- Govorunova Elena G., Gou Yueyang, Sineshchekov Oleg A., Li Hai, Lu Xiaoyu, Wang Yumei, Brown Leonid S., St-Pierre François, Xue Mingshan, Spudich John L. 2022) **Kalium Channelrhodopsins Are Natural Light-Gated Potassium Channels That Mediate Optogenetic Inhibition** *Nature Neuroscience* **25**:967–74 <https://doi.org/10.1038/s41593-022-01094-6>
- Govorunova Elena G., Sineshchekov Oleg A., Janz Roger, Liu Xiaoqin, Spudich John L. 2015) **Natural Light-Gated Anion Channels: A Family of Microbial Rhodopsins for Advanced Optogenetics** *Science* **349**:647–50 <https://doi.org/10.1126/science.aaa7484>
- Govorunova Elena G., Sineshchekov Oleg A., Rodarte Elsa M., Janz Roger, Morelle Olivier, Melkonian Michael, Wong Gane K.-S., Spudich John L. 2017) **The Expanding Family of Natural Anion Channelrhodopsins Reveals Large Variations in Kinetics, Conductance, and Spectral Sensitivity** *Scientific Reports* **7**:43358 <https://doi.org/10.1038/srep43358>
- Hooks Bryan M., Lin John Y., Guo Caiying, Svoboda Karel 2015) **Dual-Channel Circuit Mapping Reveals Sensorimotor Convergence in the Primary Motor Cortex** *Journal of Neuroscience* **35**:4418–26 <https://doi.org/10.1523/JNEUROSCI.3741-14.2015>
- Inagaki Hidehiko K., Jung Yonil, Hoopfer Eric D., Wong Allan M., Mishra Neeli, Lin John Y., Tsien Roger Y., Anderson David J. 2014) **Optogenetic Control of Drosophila Using a Red-Shifted**

**Channelrhodopsin Reveals Experience-Dependent Influences on Courtship** *Nature Methods* **11**:325–32 <https://doi.org/10.1038/nmeth.2765>

Kato Hideaki E., Kim Yoon Seok, Paggi Joseph M., Evans Kathryn E., Allen William E., Richardson Claire, Inoue Keiichi, et al. 2018) **Structural Mechanisms of Selectivity and Gating in Anion Channelrhodopsins** *Nature* **561**:349–54 <https://doi.org/10.1038/s41586-018-0504-5>

Klapoetke Nathan C., Murata Yasunobu, Kim Sung Soo, Pulver Stefan R., Birdsey-Benson Amanda, Cho Yong Ku, Morimoto Tania K, et al. 2014) **Independent Optical Excitation of Distinct Neural Populations** *Nature Methods* **11**:338–46 <https://doi.org/10.1038/nmeth.2836>

Kleinlogel Sonja, Feldbauer Katrin, Dempski Robert E., Fotis Heike, Wood Phillip G., Bamann Christian, Bamberg Ernst 2011) **Ultra Light-Sensitive and Fast Neuronal Activation with the Ca<sup>2+</sup>-Permeable Channelrhodopsin CatCh** *Nature Neuroscience* **14**:513–18 <https://doi.org/10.1038/nn.2776>

Lin John Y., Knutsen Per Magne, Muller Arnaud, Kleinfeld David, Tsien Roger Y. 2013) **ReaChR: A Red-Shifted Variant of Channelrhodopsin Enables Deep Transcranial Optogenetic Excitation** *Nature Neuroscience* **16**:1499–1508 <https://doi.org/10.1038/nn.3502>

Lin John Y., Lin Michael Z., Steinbach Paul, Tsien Roger Y. 2009) **Characterization of Engineered Channelrhodopsin Variants with Improved Properties and Kinetics** *Biophysical Journal* **96**:1803–14 <https://doi.org/10.1016/j.bpj.2008.11.034>

Mager Thomas, David Lopez, de la Morena, Senn Verena, Schlötte Johannes, D’Errico Anna, Feldbauer Katrin, Wrobel Christian, et al. 2018) **High Frequency Neural Spiking and Auditory Signaling by Ultrafast Red-Shifted Optogenetics** *Nature Communications* **9**:1750 <https://doi.org/10.1038/s41467-018-04146-3>

Mahn Mathias, Gibor Lihi, Patil Pritish, Malina Katayun Cohen-Kashi, Oring Shir, Printz Yoav, Levy Rivka, Lampl Ilan, Yizhar Ofer 2018) **High-Efficiency Optogenetic Silencing with Soma-Targeted Anion-Conducting Channelrhodopsins** *Nature Communications* **9**:4125 <https://doi.org/10.1038/s41467-018-06511-8>

Mahn Mathias, Prigge Matthias, Ron Shiri, Levy Rivka, Yizhar Ofer 2016) **Biophysical Constraints of Optogenetic Inhibition at Presynaptic Terminals** *Nature Neuroscience* **19**:554–56 <https://doi.org/10.1038/nn.4266>

Marshall James H., Kim Yoon Seok, Machado Timothy A., Quirin Sean, Benson Brandon, Kadmon Jonathan, Raja Cephra, et al. 2019) **Cortical Layer-Specific Critical Dynamics Triggering Perception** *Science (New York, N.Y)* **365** <https://doi.org/10.1126/science.aaw5202>

Messier Jessica E., Chen Hongmei, Cai Zhao-Lin, Xue Mingshan 2018) **Targeting Light-Gated Chloride Channels to Neuronal Somatodendritic Domain Reduces Their Excitatory Effect in the Axon** *Edited by Indira M Raman and Gary L Westbrook. eLife* **7**:e38506 <https://doi.org/10.7554/eLife.38506>

Mizuguchi H., Xu Z., Ishii-Watabe A., Uchida E., Hayakawa T. 2000) **IRES-Dependent Second Gene Expression Is Significantly Lower than Cap-Dependent First Gene Expression in a Bicistronic Vector** *Molecular Therapy: The Journal of the American Society of Gene Therapy* **1**:376–82 <https://doi.org/10.1006/mthe.2000.0050>

Mohammad Farhan, Stewart James C., Ott Stanislav, Chlebkova Katarina, Chua Jia Yi, Koh Tong-Wey, Ho Jose, Claridge-Chang Adam 2017) **Optogenetic Inhibition of Behavior with Anion**

**Channelrhodopsins** *Nature Methods* **14**:271–74 <https://doi.org/10.1038/nmeth.41>

Pelletier J., Sonenberg N. 1988) **Internal Initiation of Translation of Eukaryotic MRNA Directed by a Sequence Derived from Poliovirus RNA** *Nature* **334**:320–25 <https://doi.org/10.1038/334320a0>

Philipsborn Anne C. von, Jörchel Sabrina, Tirian Laszlo, Demir Ebru, Morita Tomoko, Stern David L., Dickson Barry J. 2014) **Cellular and Behavioral Functions of Fruitless Isoforms in Drosophila Courtship** *Current Biology: CB* **24**:242–51 <https://doi.org/10.1016/j.cub.2013.12.015>

Philipsborn Anne C. von, Liu Tianxiao, Yu Jai Y., Masser Christopher, Bidaye Salil S., Dickson Barry J. 2011) **Neuronal Control of Drosophila Courtship Song** *Neuron* **69**:509–22 <https://doi.org/10.1016/j.neuron.2011.01.011>

Shemesh Or A., Linghu Changyang, Piatkevich Kiryl D., Goodwin Daniel, Celiker Orhan Tunc, Gritton Howard J., omano Michael F., et al. 2020) **Precision Calcium Imaging of Dense Neural Populations via a Cell-Body-Targeted Calcium Indicator** *Neuron* **107**:470–486 <https://doi.org/10.1016/j.neuron.2020.05.029>

Sreenivasan Varun, Karmakar Kajari, Rijli Filippo M., Petersen Carl C. H. 2015) **Parallel Pathways from Motor and Somatosensory Cortex for Controlling Whisker Movements in Mice** *European Journal of Neuroscience* **41**:354–67 <https://doi.org/10.1111/ejn.12800>

Turecek R., Trussell L. O. 2001) **Presynaptic Glycine Receptors Enhance Transmitter Release at a Mammalian Central Synapse** *Nature* **411**:587–90 <https://doi.org/10.1038/35079084>

Vierock Johannes, Peter Enrico, Grimm Christiane, Rozenberg Andrey, Castro Scalise Alejandro G., Augustin Sandra, Tanese Dimitrii, et al. 2022) **WiChR, a Highly Potassium Selective Channelrhodopsin for Low-Light Two-Photon Neuronal Inhibition** *bioRxiv* <https://doi.org/10.1101/2022.07.02.498568>

Vierock Johannes, Rodriguez-Rozada Silvia, Dieter Alexander, Pieper Florian, Sims Ruth, Tenedini Federico, Bergs Amelie C. F., et al. 2021) **BiPOLES Is an Optogenetic Tool Developed for Bidirectional Dual-Color Control of Neurons** *Nature Communications* **12**:4527 <https://doi.org/10.1038/s41467-021-24759-5>

Waddell Walter H., Schaffer Arnold M., Becker Ralph S. 1973) **Visual Pigments. III. Determination and Interpretation of the Fluorescence Quantum Yields of Retinals, Schiff Bases, and Protonated Schiff Bases** *Journal of the American Chemical Society* **95**:8223–27 <https://doi.org/10.1021/ja00806a002>

Wang Wenjing, Nossoni Zahra, Berbasova Tetyana, Watson Camille T., Yapici Ipek, Lee Kin Sing Stephen, Vasileiou Chrysoula, Geiger James H., Borhan Babak 2012) **Tuning the Electronic Absorption of Protein-Embedded All-Trans-Retinal** *Science* **338**:1340–43 <https://doi.org/10.1126/science.1226135>

Yizhar Ofer, Fenno Lief E., Prigge Matthias, Schneider Franziska, Davidson Thomas J., O’Shea Daniel J., Sohal Vikaas S., et al. 2011) **Neocortical Excitation/Inhibition Balance in Information Processing and Social Dysfunction** *Nature* **477**:171–78 <https://doi.org/10.1038/nature10360>

## Author information

### Noémie Mermet-Joret

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark, Center for Proteins in Memory - PROMEMO, Danish National Research Foundation, University of Tasmania, Hobart, Australia

### Andrea Moreno

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark, Center for Proteins in Memory - PROMEMO, Danish National Research Foundation, University of Tasmania, Hobart, Australia  
ORCID iD: [0000-0001-7348-872X](https://orcid.org/0000-0001-7348-872X)

### Agnieszka Zbela

Tasmanian School of Medicine, College of Health and Medicine, University of Tasmania, Hobart, Australia

### Milad Nazari

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark, Center for Proteins in Memory - PROMEMO, Danish National Research Foundation, University of Tasmania, Hobart, Australia

### Bárður Eyjólfsson Ellendersen

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark

### Raquel Comaposada-Baro

Department of Neuroscience, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

### Nathalie Krauth<sup>†</sup>

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark, Center for Proteins in Memory - PROMEMO, Danish National Research Foundation, University of Tasmania, Hobart, Australia

<sup>†</sup>Present address: Novo Nordisk Foundation Center for Basic Metabolic Research, Copenhagen University, Copenhagen, Denmark and Institute for Neuroscience, Copenhagen University, Copenhagen, Denmark.

### Anne von Philipsborn<sup>#</sup>

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark

#Present address: Université de Fribourg: Fribourg, Switzerland.

**Andreas Toft Sørensen**

Department of Neuroscience, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

**Joaquin Piriz**<sup>&</sup>

Instituto de Fisiología Biología Molecular y Neurociencias (IFIBYNE), Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina

<sup>&</sup>Achucarro Basque Center for Neuroscience, Leioa, Spain, Ikerbasque - Basque Foundation for Science, Bilbao, Spain.

**John Y Lin**<sup>\*</sup>

Tasmanian School of Medicine, College of Health and Medicine, University of Tasmania, Hobart, Australia  
ORCID iD: [0000-0002-1723-4597](https://orcid.org/0000-0002-1723-4597)

**For correspondence:** [john.lin@utas.edu.au](mailto:john.lin@utas.edu.au)

<sup>\*</sup>These authors contributed equally

**Sadegh Nabavi**<sup>\*</sup>

DANDRITE, The Danish Research Institute of Translational Neuroscience, Aarhus University, Aarhus, Denmark, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark, Center for Proteins in Memory - PROMEMO, Danish National Research Foundation, University of Tasmania, Hobart, Australia  
ORCID iD: [0000-0002-3940-1210](https://orcid.org/0000-0002-3940-1210)

**For correspondence:** [snabavi@dandrite.au.dk](mailto:snabavi@dandrite.au.dk)

<sup>\*</sup>These authors contributed equally

**Editors**

Reviewing Editor

**Marcel Goldschen-Ohm**

University of Texas at Austin, Austin, United States of America

Senior Editor

**Kenton Swartz**

National Institute of Neurological Disorders and Stroke, Bethesda, United States of America

**Reviewer #1 (Public review):**

**Summary**

In this manuscript, the authors generate an AAV-deliverable tool that generates action potentials in response to red light, but not blue light, when expressed in neurons. To do this, they screen some red light-excitatory/blue light-inhibitory opsin pairs to find ones that are spectrally and temporally matched. They first show that this works with Chrimson and GtACR2, however, they expand their search after finding that the tau-off (inactivation after

light cessation) kinetics of these two opsins are not well-matched. They directly examine a small set of options based on a literature search and settle on a variant of red light-excitatory Chrimson and blue light-inhibitory ZipACR. To even more closely match the kinetics of this pair, the authors create a structure homology model of the ZipACR retinal binding pocket and use this to guide generation of a small mutant panel, leading to a more optimized ZipACR mutant. They then show that a bicistronically expressed fusion arrangement of these opsins, plus some functional peptides, can drive action potentials up to 20hz with red light and does not do so with blue light, in hippocampal cells transduced by AAV. They also show function *in vivo*, in a mouse, using a physiological readout. They conclude that their new tool may be useful for complex experimental designs requiring multiple optical channels for write-in/read-out.

The major advantage claimed by the authors over existing tools is the temporal time-locking of their inhibitory opsin - this is driven by the contrast between tau-off kinetics of their ZipACR variant compared to gtACR2, which is used by the leading competitor tool (BiPOLES).

### Big thoughts

While the authors were carefully thoughtful about the potential influence of temporal kinetics on the efficiency of a tool such as this one, there were no experiments conducted that make use of the unique properties of this molecular strategy (although the authors state that these experiments are now underway in their lab). They share some examples of how the tool could be useful in the discussion. Where do I think this could be useful?

First, experimental designs that require multiple optical channels of control. This appears to be aligned with the author's thoughts, as they state, correctly, that opsins utilizing retinal as a light-sensing chromophore are universally activated by blue light (the so-called 'blue shoulder'). Therefore, their tool may be useful for stimulating multiple populations using a blue excitatory opsin in neuron A and their tool for red excitation of neuron B - or, in the author's own words, "A potential solution to the problem of cross-talk...". In this manuscript, the authors provide state that there this is possible in theory and that there are no obvious reasons that it would not work, but do not present data that showcases their new tool for this purpose (e.g. Vierock, Johannes, et al. "BiPOLES is an optogenetic tool developed for bidirectional dual-color control of neurons." *Nature communications* 12.1 (2021): 4527. Figure 4f-I; 6). The same set-up could be imagined for green GECI (or equivalent) imaging of cells in the same volume that their tool is being used in - for instance, interleaving red stimulation light and blue imaging light, (perhaps) without the typical concern of imaging light bleed-through activating the opsin itself. I agree that it will likely work for multi-channel control, but only time will tell, at this point.

Second, for high-frequency temporal control over both excitation and inhibition in the same neuron. Red light turns the cell on, and blue light turns the cell off (see, for instance, Zhang, Feng, et al. "Multimodal fast optical interrogation of neural circuitry." *Nature* 446.7136 (2007): 633-639. Figure 2; Vierock as above, Figure 4a,b). Again, here the authors are long on theory ("The new system...can drive time-locked high-frequency action potentials in response to red pulses") and short on explicit data. While they do show that red light = excitation and blue light = inhibition, they neither show 1) all-optical on/off modulation of the same cell; nor 2) high-frequency inhibition or excitation (max stim rate of 20hz, which is the same as the BiPOLES paper used for their LC stimulation paradigm; Vierock, as above, Figure 7a-d). They did provide a response to this critique that data showing excitation and inhibition spread across multiple panels were largely collected from the same cells.

Despite these major shortcomings, the further development and characterization of tandem opsins, such as this one, is of interest to the community. There is on-going work by the BiPOLES team to create new iterations (e.g. Wahid, J., et al. "P-15 BiPOLES2 is a bidirectional optogenetic tool with a narrow activation spectrum and low red-light excitability." *Clinical Neurophysiology* 148 (2023): e16.). The authors have collected a substantial amount of

additional data along the course of review and, even aside from the final tool, the overall data and approaches shown are useful.

<https://doi.org/10.7554/eLife.90327.2.sa1>

### Author response:

The following is the authors' response to the original reviews.

#### **Reviewer #1 (Public Review):**

*Therefore, their tool may be useful for stimulating multiple populations using a blue excitatory opsin in neuron A and their tool for red excitation of neuron B... Yet, there are no data presented that showcases their new tool for this purpose*

We agree with the reviewer that in this manuscript we have not experimentally shown the applicability of our system for dual optical stimulation. However, the suppression of blue-light excitation of ZipV/T-IvfChr-expressing neurons strongly suggests this can be used in experiments exciting populations of neurons similarly shown for BiPOLES. We don't see a theoretical basis where this experiment cannot be done if sufficient cell targeting mechanisms (such as the use of cre-lox or retroAAV) is utilized. We have started several projects pursuing these utilities in the meantime.

*While they do show that red light = excitation and blue light = inhibition, they neither show 1) all-optical on/off modulation of the same cell; nor 2) high-frequency inhibition or excitation (max stim rate of 20hz, which is the same as the BiPOLES paper used for their LC stimulation paradigm; Vierock, as above, Figure 7a-d).*

Regarding point 1, we understand that the reviewer asks if we have optically excited (with red light) and inhibited (with blue light) the same neurons. If so, figure 4B1 (optical excitation of ZipT-IvfCh with red light) and figure 5A (optical inhibition of ZipT-IvfCh with blue light) represent largely the same set of neurons.

Regarding point 2, we respectfully disagree with the reviewer's interpretation of Figure 7a-d) in Vierock et al. As we understand, in this part the authors apply a 20 Hz optical stimulation protocol to the LC neurons in vivo. However, there is no data showing that individual neurons do follow this stimulation protocol. To be clear, we are not saying that BiPOLES cannot drive 20 Hz APs. Very likely it can. It is based on ChrimsonR which is capable of doing so (Klapoetke et al., Figure 2). Although, in this manuscript we have not shown data for optical stimulation above 20Hz, our system is based on vfChrimson, which is known to drive AP of 100Hz and above (Mager et al., figure 2 and 3).

*... they must revise the manuscript to show that their approach is both 1) different in some way when compared to BiPOLES (it is my understanding that they did not do this, as per the supplementary alignment of the BiPOLES sequence and the sequence of the BiPOLES-like construct that they did test) and 2) that the properties that the investigators specifically tailored their construct to have confer some sort of experimental advantage when compared to the existing standard.*

In the latest version of the manuscript, we have compared our ZipV-IvfChr and the BiPOLES construct adapted with vfChrimson (Fig. 2 Suppl 1). The mean photocurrent amplitude of IvfChr in the ZipV-IvfChr construct is ~2.7 x higher than BiPOLES adapted with vfChrimson (14 randomly selected HEK293 cells in each group) (Fig. 2 Suppl 1B). We conducted this experiment in HEK293 cells to ensure accurate voltage-clamping and less biased cell

selection. Even adjusting for the smaller photocurrent of vfChrimson vs ChrimsonR, this would still translate to  $\sim 1.6 \times$  greater photocurrent with ZipV-IvfChr compared to the original BiPOLES utilizing ChrimsonR. We believe the increased efficiency of excitation is an important aspect of adapting vfChrimson for red-light excitation of neurons.

**Reviewer #2 (Public Review):**

*(1) In the Introduction or Discussion, the authors could better motivate the need for a red-shifted actuator that lacks blue crosstalk, by giving some specific examples of how the tool could be productively used, e.g. pairing with another blue-shifted excitatory opsin in a different population, or pairing with a GFP-based fluorescent indicator, e.g. GCaMP. The motivation for the current tool is not obvious to non-experts.*

In the discussion, we now provided examples for potential use of the tool. For example, one of the key aspects that can be manipulated by the existing tool is the induction of spike-timing dependent plasticity with 2 wavelengths of light with blue light channelrhodopsin such as oChIEF is used to evoke presynaptic release and ZipT-IvfChr expressed in postsynaptic neuron. In this situation, the rapid termination of inhibitory response is critical so it does not interfere with the induction of LTP or LTD. Another experiment is the alternate control of projection neurons and interneurons in cortical areas, independent controls of neurons of direct and indirect pathways in the striatum to manipulate behavior.

*(2) Simultaneous excitation and inhibition are not the same as non-excitation. The authors mentioned shunting briefly. Another possible issue is changes in osmotic balance. Activation of a Na<sup>+</sup> channel and a Cl<sup>-</sup> channel will lead to net import of NaCl into the cell, possibly changing osmotic pressure. Please discuss.*

We agree with the notion that osmotic, ionic and pH changes in small neuronal structure can be disruptive to the physiology and this is the reason we developed our approach where the fastest channelrhodopsins are used so we can minimize the channel opening time and the flux of ions through the channels when brief light illuminations are applied. Not only the flux of protons, sodium ions and calcium ions are minimized, the flux of chloride should be minimal as well (as the membrane potential should be close to the reversal potential of chloride reversal potential hence low ion flow). Hence our approach should be minimally disruptive compared to most other existing channelrhodopsin-based approaches when short or minimal light pulses were used in conjunction with our tools. This recommendation is included in the updated manuscript .

*(3) The authors showed that in ZipT-IvfChr, orange light drives excitation and blue light does not. But what about simultaneous blue and orange light? Can the blue light overwhelm the effect of the orange light? Since the stated goal is to open the blue part of the spectrum for other applications, one is now worried about "negative" crosstalk. Please discuss and, ideally, characterize this phenomenon.*

We now have performed this experiment. Simultaneous blue (470nm) and red light (635nm) stimulation does not produce AP (Fig .4 Suppl 1A)). This suggests the inhibitory effect of ACR is more efficient than the excitatory effects of IvfChr due to their higher conductance, this re-emphasizes the rapid termination of the ACR effects is critical for minimal disruption of physiological effects in such pairing strategy.

*(3.1) Does the use of the new tool require careful balancing of the expression levels of the ZipT and the IvfChr? Does it require careful balancing of blue and orange light intensities?*

As with any optogenetic tool, the users should validate the efficacy of the tool in their own system. Our tool solely relies on the balanced expression of the 2A system, the efficiency of the two opsins and their degradation of the time-span of expression. These aspects of the tool would be better addressed in future versions of the tools or improvement of the BiPOLES-type of tandem expression in subsequent versions. From the instrumentation side, the light intensity and differential penetration depth requires careful consideration. However, this holds true in most optogenetic and fluorescence imaging-based approaches as well. In the current update of the manuscript, we have included further discussion on these aspects as well.

*(3.2) Also, many opsins show complex and nonlinear responses to dual-wavelength illumination, so each component should be characterized individually under simultaneous blue + orange light.*

We now have performed this experiment (please see our comment to point 3)

*(3.3) I was expecting to see photocurrents at different holding potentials as a function of illumination wavelength for the coexpressed construct (i.e. to see at what wavelength it switches from being excitatory to inhibitory); and also to see I-V curves of the photocurrent at blue and orange wavelengths for the co-expressed constructs (i.e. to see the reversal potential under blue excitation). Overall, the patch clamp and spectroscopic characterization of the individual constructs was stronger than that of the combined constructs.*

We have added the IV curves for the co-expressed construct at different holding potentials for 470nm and 635nm wavelengths. This shows reverse potential for the two wavelengths that are intended for in vitro and in vivo applications. Performing a similar experiment for a variety of wavelengths would not be as valuable, in part, due to the enormous amount of data generated. As we have shown in the study, the response of any channelrhodopsins vary with different light duration and light intensities in addition to the wavelengths and holding potentials. The results for each recorded cell could include stimulation by different wavelengths, stimulation by different illumination intensities, stimulation with different light duration in addition to different holding potentials. Not only would the results be highly variable from cell-to-cell, there will be potentially hundreds or thousands of combinations to be tested per cell (e.g., 5 light intensities @1, 2.5, 5, 10 and 20 mW/mm<sup>2</sup>, 8 different wavelengths @ 450nm, 475nm, 500nm, 525nm, 550nm, 575nm, 600nm and 625nm, 7 light durations @ 1ms, 5ms, 10ms, 50ms, 100ms, 500ms and 1s, and , and 6 holding potentials @ -80mV, -70mV, -60mV, -40mV, -20mV and 0mV would result in 1680 stimulation conditions per recorded cell). Technically, the significant lowering of membrane resistance when both IvfChr and ZipACR variants are activated simultaneously would compromise the quality of voltage-clamping even in HEK293 cells with series resistance compensation. We have yet to see any other studies that had included such ambitious electrophysiology experiment for the channelrhodopsin characterization, likely due to the feasibility of such experiment.

**Reviewer #3 (Public Review):**

*(1) The enhanced vf-Chrimson could potentially be a highlight of the manuscript, serving broader applications. Yet, gauging the overall improvements of ivf-Chrimson in comparison to other Chrimson variants remains intricate due to several reasons. First, photocurrents from ivf-Chrimson seem smaller than those from C-Chrimson (Supplemental Figure 3), and a direct comparison with standard vf-Chrimson is absent.*

We appreciate the reviewer's positive view of our modified variant. We did not emphasize this particular modification as it was identical to our previous published modification and

similar to that previously published by others (CsChrimson and C1Chrimson). In all these cases, improved membrane expression was consistently detected. We believe that expression data and our comparison of C-Chrimson and IvfChr is sufficient to justify the improved membrane expression and function.

*Second, while membrane expression of ivf-Chrimson appears enhanced in provided brightfield recordings, the quantitative analysis would necessitate confocal microscopy and a membrane marker (Supplemental Figure)*

We have now quantified the results with a membrane palmitoylated mCherry using confocal microscopy shown in Fig 2 Suppl1 A. We measured the Pearson Correlation Coefficient of the mCherry with EGFP or Citrine signal for the 6 constructs (vfChrimson, vfChrimson with trafficking sequence, vfChrimson with N-terminal signaling peptide from oChIEF (C-vfChrimson), vfChrimson with trafficking sequence and N-terminal signaling peptide from oChIEF (IvfChr), BiPOLES with EGFP or citrine and vfChrimson) and the results were identical and consistent with the prior results using epifluorescence microscopy.

*(2) Finally, other N-terminal modified Chrimson variants, like CsChrimson by Klapoetke et al. in 2014 and C1Chrimson by Oda et al. in 2018, have been generated. Comparing ivf-Chrimson to vf-CsChrimson or vf-C1Chrimson would be important to evaluate the benefits of the applied N-terminal modification.*

Our development of IvfChrimson is similar to the approach of vf-CsChrimson and identical to that of vf-C1Chrimson and we do not claim these modifications to be unique or superior. However, we have developed our design independently of these other studies and we have more extensive functional comparison and characterization data of our IvfChrimson variant than the other studies.

*(2.1) The action spectra of ZipACR suggest peak absorption of ZipACR WT and its mutant at 525 - 550 nm (Fig. 3). This is even further red-shifted than previously reported by Govorunova et al. Further action spectra recordings differ for all constructs between recordings initiated with blue or red light (Supplementary Fig. 5). This discrepancy is unexpected and should be discussed.*

We thank the reviewer for the comment, this was a mistake in the traces used for the figure. The example traces were the spectral response measured from the 400 nm to 650 nm instead of the 650 nm to 400 nm order shown in the spectral data. This has now been corrected.

*Additionally, the representative photocurrents of Zip(151V) in Fig. 3D1 do not align with the corresponding action spectrum in Fig. 3D2 as they show maximal photocurrents for 400 nm excitation.*

Please, see point above.

*(3) The authors introduce two different bicistronic expression cassettes-ZipT-IvfChR and ZipV-IvfChR-without providing clear guidelines on their conditions of use. Although the authors assert that ZipT is slower and further red-shifted than ZipV, the differences in the data for both ACR mutants are small and the benefits of the different final constructs should be explained.*

In our testing in neurons, ZipT has less ‘escaped’ spikes after the termination of the light pulses in the cells we have tested. However, this is dependent on the membrane properties such as capacitance and resistance of the cells. ZipV has a faster termination time and in

some situations may be necessary due to its faster termination time and reduced disruption of physiological processes.

We have now included this discussion in our updated manuscript.

*(4) The ZipT/V-IvfChRs are designed as bicistronic constructs; yet, disparities in membrane trafficking and protein degradation between the two channels could lead to divergences in blue and red light photoresponses. For future applicants, understanding the extent of expression ratio variations across cells using the presented expression cassettes could be of significance and should be discussed.*

We now have included this discussion in our responses above.

**Reviewer #1 (Recommendations For The Authors):**

*(1) The Figure 1a mV cartoon traces for chloride are confusing. The chloride currents are depolarizing, not hyperpolarizing. As noted by the authors, these channels largely generate AP blockade through shunting inhibition (division), not hyperpolarization (subtraction).*

The figure has been corrected.

*(2) Figure 2A does not show where the light is applied. Why are some of the bars blue and some of them not filled?*

This has been corrected

*(3) Figure 2C1 does not show where the light is applied. There should be an inset to detail the blue-light-cessation-evoked AP. Also doesn't give the holding potential.*

The requested details are added.

*(4) Figure 2C2 inset is described as showing that "Light-induced currents with 470 nm illumination were initially outward but turned inward immediately following light offset." Is that correct? It looks to me like the current turns inward about half-way through the light pulse and then becomes even stronger after the light turns off. That is also consistent with the CC traces, which appear to show a transition toward depolarization during the light pulse before the AP initiation at light offset.*

Yes, the reviewer's observation is correct. There are blue light-induced outward and inward current peaks at the onset and offset of the light. Accordingly, we have modified the phrasing for Fig. 2C2.

*(5) Figure 3D1 shows that Zip(151V) has a peak current at 400nm, with a steady increase in current from red to blue, however, this is not the case in the summary data in 3D2. It's also not shown in Supplementary Figure 5B. What's going on?*

We apologize for the prior version of the figure associated with the first submission. The example traces from 400nm -> 650 nm were incorrectly included in the figure whereas the 650nm -> 400 nm example traces should be included. This has been corrected.

*(6) Figure 3D1 has no time scale.*

It is now been included

(7) Figure 3E1 should read "Transduced" and not "Transfected"

This has been corrected.

(8) IvfChr fidelity drops off dramatically at 20hz...down to 50% efficiency of generating APs. This is described in the legend as "high frequency". Maybe the cart came before the horse in this figure...as it looks like in panel C that using less light power density improves fidelity in the dual opsin configuration with red light stimulation...why not use that power for the characterization? Did you try any higher frequencies? Or longer pulse widths? This is an important characterization to inform further use of the tool. This shortcoming isn't a cell-intrinsic limitation, as the 470nm stim with IvfChr was 100% successful at both 10hz and 20hz.

It is known that red but not blue light pulses induce desensitization (optical fatigue) in red-shifted ChR variants. Indeed, one can reinstate the response to red light, by giving violet-blue light pulses (Fig 4. Suppl 2). We think this is the reason that the 470nm stimulation was more effective in inducing AP in cells expressing IvfChR. Higher light intensities induce greater desensitization, but are preferred for faster opening of channels and depolarization of neurons. This can explain why, in some situations, lower light intensities were more effective in producing APs when pulse trains were used. We have recordings from cells firing APs at 40Hz (not included). All these cells had high expression levels of the opsin.

(9) Figure 4D: why use 100ms pulse width? How do you know that this isn't causing depol block? Or some of the nefarious concerns that are raised in the discussion, such as "...disrupt[ion of] normal neuronal physiology and signal processing that occurs in millisecond time scale"?

We used 100ms pulse duration to follow the published protocol that this experiment is based on (Lin et al., 2013, Nature Neuroscience).

(10) Figure 4E-bottom: What is the blue peak at light onset? Is the tool driving early activation before silencing?

There seems to be an early, sharp and brief activation by blue light. We don't know the definite cause of this, but we speculate this is driven by blue-light activation of ZipACR and not the IvfChr portion of the construct. The reason is that such a sharp rise is absent when only IvfChr is expressed (Fig. 4E, upper panel). Soma-targeted motif tethered to channelrhodopsins is known to result in preferential expression of channels close to soma but does not exclude the expression of channelrhodopsin in axonal and dendritic compartments, especially when animals are allowed to recover for long period of time after viral injection. We believe that ZipACR at axonal terminals where the chloride concentration is high can still cause blue-light evoked depolarization and transmitter release. We observed this phenomenon in two mice in their first trial. The data for individual trials for each mouse are included in a supplementary table.

(11) Figure 4G: Earlier in this same figure (B2, C), 470nm light was more effective at stimulating IvfChr than 635nm light. Is it unexpected that 638nm light would in this in vivo context be more effective at driving IvfChr responses than 450 nm light (at least as reflected by the AUC measurements)? Does this reflect fiber placement and light penetration/scattering?

The spectral peaks of Chrimson-based variants including vfChrimson are all centered around 600 nm, and at 635 / 638 nm light, the amplitudes of photo-response decline, the channel

onset slows, and the channels suffer greater desensitization. In isolated preparations where the light penetration is similar between 635 / 638 nm and 470 nm, 470 nm responses can outperform 635 / 638 nm responses due to its lack of desensitization and higher consistency in its response. This is also a strong reason that we have developed our current approach. In *in vivo* preparation shown in Fig. 4D-G, the much higher tissue penetration of 638nm light due to reduced absorption and reduced scattering can offset the performance of IvfChr to 450 nm light.

*(12) In the methods, it is noted that different viral batches appear to generate different levels of neuronal toxicity. If that is the case, how did you differentiate between true differences between constructs vs. differential cell health effects?*

For figure 4D-F (whisker movement), we determined virus toxicity using NeuN staining. In slice recordings, we used the electrophysiological property of the neurons to assess their health. For this manuscript, we had one batch of virus that produced toxicity. We did not include any data from this batch.

**Reviewer #2 (Recommendations For The Authors):**

- Define AUC on first use.

It is now defined.

- Figure 3C2: Please explain how the photocurrents were normalized. As presented, it looks like under strong orange light, the ZipACR has higher photocurrent than the ivfChr.

This is due to the fact vfChrimson and other Chrimson-based variants do not fully recover in the dark after 590 nm stimulation. We tested IvfChrimson with both reconditioning light pulse of 405 nm and without 405 nm and we can consistently reach a greater ‘maximal’ response from the same cell after 405 nm reconditioning (see Fig. 4 Suppl 2). We therefore normalize the response to the maximal recorded response of the cell often achieved with 10 or 20 mW/mm<sup>2</sup> 590 nm stimulation after 405 nm reconditioning. We understand this can be confusing and have now replaced the light-intensity response in Fig. 3C2 with the one with 405 nm reconditioning which is easier to interpret for the readers.

- P. 3: "As expected, blue light pulses induce transient membrane suppression..." Unclear what "suppression" means. Shunting? Hyperpolarization?

We rephrased this to “As expected, blue light pulses transiently suppress APs...”

- P. 3: "illumination at 470 nm and 590 nm wavelengths led to similar amounts of courtship song (110.1 {plus minus} 12.8 and 78.5 {plus minus} 11.6, n = 16-17, respectively)". What are the units of "courtship song"?

The unit for courtship song is the number of pulses per 10 seconds. This has been clarified in the figure.

- P. 5: The quantification of photocurrent in terms of pA/pF/A.U. is non-standard. I understand the impetus to normalize by expression to give something proportional to per-molecule conductance, but a user cares about overall photocurrent. Please also give the real photocurrents, either pA or pA/pF.

We have provided the real photocurrent in pA or pA/pF where scientifically appropriate. To avoid selection and experimenter’s bias in our data, we did not set criteria for data

elimination for cells with specific fluorescence intensity or photocurrent amplitude. Some resulting response can range from vary up to 20 folds from the same construct in many experiments. We do not believe that averaging absolute photocurrent amplitude would be justified due to the imbalance of weighing in the results. We do acknowledge that not selecting or eliminating data points would introduce higher noise in recordings with smaller responses but this is preferable over the selection or experimenter bias that is likely to be introduced otherwise.

- Please quote illumination intensities wherever possible.
- P. 7: why was the red light crosstalk into Zip(151T) tested at 635 nm instead of 590 nm? Isn't the relevant parameter 590 nm, since that will be used for the excitatory opsin?

In all our characterizations of the constructs using slice electrophysiology recordings, we used 635nm instead of 590nm. The reason is that compared to 590nm wavelength, at 635nm the photocurrent for Zip(151T) and Zip(151V) is significantly reduced (Fig. 3D1,D2).

- P. 10: "we examined the power at which responses to 470 nm and 635 nm lights induce APs in neurons expressing ZipT-IvfChr, ZipV-IvfChr, or IvfChr", but the preceding sentence says you didn't test the ZipT-IvfChr. This is confusing, please clarify.

The previous paragraph refers to the photocurrent recordings in HEK293 cells where our fast LED based illumination system is limited to 590 nm light, whereas the subsequent paragraph refers to the brain slice neuronal recordings. We have now emphasized the difference of the experiments in the rewrite.

- Fig. 4B1, top: Why don't the blue traces return to the same baseline after the stimulus epochs?

We observed this shift in baseline (~4mV more depolarized) in cells expressing IvfChR (or vfChR) only with blue light stimulation. This was observed in the neurons recorded in the CA1 as well (data not shown). There was no such a change following red light stimulation (Fig. 4B1). Therefore, this should not affect the applicability of our construct. The original paper introducing vfChR did not test the responses of their constructs to blue light. There could be another photocycle state that is activated stronger by 470nm than 590nm and it has a slow off-rate, but this is only a speculation from our side. It must be noted we did not observe such a phenomenon in cells expressing ChrimsonR (Fig. 1 Suppl 1C).

- Fig. S3B, right: The two colors are barely distinguishable on the graph. Consider more distinct colors and/or different symbols.

It has been changed accordingly.

- P. 15: "However, we do not recommend the use of orange light pulses, as we observed a significant photocurrent in this wavelength." Not clear what this is referring to. Which construct? Under which circumstances shouldn't one use orange light pulses? Where's the data showing this?

This is referring to Fig. 3D1,D2 and Figure 4 suppl Fig. 2 which show a normalized ~40-50% photocurrent at 590nm. Now in the text, the reference figures for the data are added.

<https://doi.org/10.7554/eLife.90327.2.sa0>