

Structural basis for kinase inhibition in the tripartite *E. coli* HipBST toxin-antitoxin system

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

August 31, 2023 (this version)

Sent for peer review

July 6, 2023

Posted to bioRxiv

June 19, 2023

René L. Bærentsen, Stine Vang Nielsen, Ragnhild Bager Skjærning, Jeppe Lyngsø, Francesco Bisiak, Jan Skov Pedersen, Kenn Gerdes, Michael A. Sørensen , Ditlev. E. Brodersen 

Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 81, DK-8000 Aarhus C, Denmark • Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 København N • Department of Chemistry and Interdisciplinary Nanoscience Centre (iNANO), Gustav Wieds Vej 14, DK-8000 Aarhus C • Voldmestergade 8, DK-2100 København Ø, Denmark

 https://en.wikipedia.org/wiki/Open_access

 <https://creativecommons.org/licenses/by/4.0/>

Abstract

Many bacteria encode multiple toxin-antitoxin (TA) systems targeting separate, but closely related, cellular functions. The toxin of the *E. coli* *hipBA* system, HipA, is a kinase that inhibits translation via phosphorylation of glutamyl-tRNA synthetase. Enteropathogenic *E. coli* (EPEC) O127:H6 encodes the *hipBA*-like, tripartite TA system; *hipBST*, in which the HipT toxin specifically targets the tryptophanyl-tRNA synthetase, TrpS. Notably, in the tripartite system, the function as antitoxin has been taken over by the third protein, HipS, but the molecular details of how activity of HipT is inhibited remain poorly understood. Here, we show that HipBST is structurally different from *E. coli* HipBA and that the unique HipS protein, which is homologous to the N-terminal subdomain of HipA, inhibits the kinase through insertion of a conserved Trp residue into the active site. We also show how auto-phosphorylation at two conserved sites in the kinase toxin serve different roles and affect the ability of HipS to neutralize HipT. Finally, solution structural studies show how phosphorylation affects overall TA complex flexibility.

eLife assessment

This **valuable** study presents an exhaustive structural analysis of a complete tripartite HipBST toxin-antitoxin system of the Enteropathogenic *E. coli* O127:H6, which represents a fascinating variation on the well-studied HipAB toxin-antitoxin system. The **convincing** data show that major features of the canonical HipAB system have been rerouted to form the tripartite HipBST, revealing a new mode of inhibition of a toxin kinase.

Bacteria employ a wide range of mechanisms to adapt to changes in their environments and imminent threats, such as exposure to antibiotics and phages, including activation of toxin-antitoxin (TA) systems that encode intracellular toxins able to rapidly reduce growth (LeRoux & Laub, 2022). Canonical (type II) TA systems are small, bicistronic loci that encode a protein toxin and its cognate antidote (antitoxin) that tightly interact to form an inactive higher-order complex, usually also capable of controlling transcription via a DNA-binding domain on the antitoxin (Harms *et al.*, 2018). However, there are also examples of tricistronic TA loci, in which the third gene either encodes a chaperone required for folding of the antitoxin and thus, toxin inhibition (Bordes *et al.*, 2016) or an additional, transcriptional regulator (Hallez *et al.*, 2010, Zielenkiewicz & Ceglowski, 2005, Jurenas *et al.*, 2021).

In the widespread and diverse *hipBA* (high persister) system, the HipA toxin is a serine-threonine kinase (STK) (Hanks *et al.*, 1988, Stancik *et al.*, 2018), while the cognate antitoxin, HipB, contains a helix-turn-helix (HTH) motif (Gerdes *et al.*, 2021, Schumacher *et al.*, 2009). For the *hipBA* system from *Escherichia coli* K-12, the toxin (HipA_{Ec}) specifically targets glutamyl-tRNA synthetase (GltX) by phosphorylation of an ATP-binding motif conserved in type I aminoacyl-tRNA synthetases (Eriani *et al.*, 1990, Sekine *et al.*, 2003), thereby inhibiting its activity and blocking translation (Germain *et al.*, 2013). Accumulation of uncharged tRNA^{Glt} subsequently induces the stringent response and cellular dormancy via RelA-mediated (p)ppGpp synthesis on starved ribosomes (Haseltine & Block, 1973, Pacios *et al.*, 2020, Winther *et al.*, 2018). In contrast to most classical type II TA systems, the antitoxin, HipB, does not directly block the HipA active site so inhibition has been proposed to occur by several alternative mechanisms, including inhibition of conformational changes in the HipA kinase required for catalysis, sequestration of HipA on DNA via the HipB DNA binding domain (Schumacher *et al.*, 2009), and allosterically via placement of a C-terminal Trp residue of HipB into a pocket on HipA (Evdokimov *et al.*, 2009). HipA_{Ec} is also regulated by *trans* autophosphorylation at a conserved serine (Ser150) situated in a loop near the active site termed the "Gly-rich loop", which is required for ATP binding and functionally (but not structurally) similar to the P loop of eukaryotic kinases (Huse & Kuriyan, 2002, Schumacher *et al.*, 2012). Two discrete conformations of the Gly-rich loop are observed depending on the phosphorylation state of Ser150 and whether substrate (ATP) is bound or not (Schumacher *et al.*, 2015, Schumacher *et al.*, 2012, Schumacher *et al.*, 2009). In the unphosphorylated state and in the presence of ATP, the flexible Gly-rich loop is found in an inward conformation in which main chain amino groups coordinate several of the ATP phosphate groups and the kinase is active (Schumacher *et al.*, 2009). Phosphorylation of Ser150 results in strong interactions of the phosphate group with several active sites residues, which causes ejection of the Gly-rich loop and prevents binding of ATP, thus inactivating the kinase (Schumacher *et al.*, 2012, Wen *et al.*, 2014).

In addition to the canonical *hipBA* locus, enteropathogenic *E. coli* (EPEC) O127:H6 contains a partially homologous, tricistronic Hip system; *hipBST*, encoding three separate proteins. HipB, by analogy to HipBA, contains a putative DNA-binding HTH domain while HipT is a small HipA-like kinase the specifically targets tryptophanyl-tRNA synthetase, TrpS (Gerdes *et al.*, 2021, Vang Nielsen *et al.*, 2019). Surprisingly, the third protein, HipS, which at the sequence level corresponds to the N-terminal subdomain 1 of HipA (N-subdomain 1), was shown to function as the antitoxin of HipT but the molecular mechanism by which HipS neutralizes HipT is currently not known (Vang Nielsen *et al.*, 2019). Phylogenetic analysis has demonstrated that HipT and HipA toxins are closely related within a very broad superfamily of HipA-homologous kinases (Gerdes *et al.*, 2021). HipT has two phosphoserine positions, Ser57 and Ser59, in its Gly-rich loop, which are both modified by *trans* auto-phosphorylation *in vivo* suggesting that they might play a role in regulating the activity of HipT (Vang Nielsen *et al.*, 2019).

Here, we determine a 2.9 Å crystal structure of a kinase-inactive HipBST^{D233Q} complex, revealing that the overall structure is markedly different from *E. coli* HipBA but similar to a HipBA complex from *S. oneidensis*. In this structure, HipT interacts tightly with HipS and that despite the lack of phosphorylation we find the Gly-rich loop in an outward, inactive conformation similar to phosphorylated HipA. We show that this is likely a result of HipS neutralization of HipT through insertion of a bulky Trp residue into the kinase active site. Next, we show that both of the serine residues in the Gly-loop are essential for HipT toxicity *in vivo* and that auto-phosphorylation of Ser59 prevents neutralization by HipS. We next determine crystal structures of two kinase active HipBST variants revealing that autophosphorylation takes place at both Ser57 and Ser59 despite the lack of *in vivo* toxicity. Surprisingly, phosphorylation of Ser57 or Ser59 does not affect the conformation of HipT inside the HipBST complex, which remains in an inactive state with the Gly-rich loop ejected, which we attribute to the presence of HipS. Finally, we show that the flexibility of the complex in solution varies in a phosphorylation-dependent manner and use a phylogenetic analysis to demonstrate that variations of the HipT Gly-rich loop serine residues correlate with separate clades of proteins.

Results

E. coli HipBST forms an heterohexameric complex with HipT in a canonical inactive conformation

To understand the molecular and structural basis for the functional differences between the HipBA and HipBST toxin-antitoxin systems, we determined the crystal structure of HipBST^{D233Q}, containing the inactive HipT D233Q kinase variant, to 2.9 Å by molecular replacement using HipA_{Ec} as search model (**Figure 1a** and **1b**). The refined structure ($R=19.5\%$, $R_{\text{free}}=22.8\%$) has two copies of each of the three proteins in the asymmetric unit and is generally well-defined for HipB and HipT, while HipS appears to be more flexible. The higher-order HipBST complex structure consists of two HipST toxin:antitoxin complexes separated by a dimer of HipB proteins. This differs markedly from the HipBA_{Ec} complex, in which two HipA toxins are closely packed head-to-tail around two HipB antitoxins (**Figure 1c**, **top**) but is reminiscent of the structure of a HipBA complex from *Shewanella oneidensis* (HipBA_{So}), which was crystallised in complex with DNA (**Figure 1c**, **bottom**) (Wen *et al.*, 2014).

HipB consists of an α -helical bundle of four helices, including the central, DNA-binding HTH motif (residues 62-83), which is exposed at the bridge of the hetero-hexamer (**Figure 1b**, green). The helical bundle is followed by a small β strand that forms an antiparallel sheet by pairing with the corresponding region in the neighbouring HipB protein, thus forming a homodimer. The N-terminus of HipB (residues 1-35) is not visible in the electron density, but AlphaFold2 prediction indicates that it forms a single α -helix (**Figure 1 supplement 1a**) (Jumper *et al.*, 2021). For HipBA_{Ec}, a hydrophobic residue at the C-terminus of HipB was proposed to reach a cleft between the N-subdomain 1 and the core kinase domain of HipA and function as antitoxin by restricting domain movements required for catalysis (**Figure 1c**) (Schumacher *et al.*, 2009). HipB from HipBST does not have such a terminal residue (**Figure 1 supplement 1a**), suggesting a different mechanism of toxin neutralisation. With respect to DNA binding, superposition of HipBST with HipBA_{So} in complex with DNA (**Figure 1 supplement 2a**) shows that HipT has several conserved, positively charged residues near the expected location of the DNA phosphodiester backbone, suggesting that HipBST might bind DNA in a similar way.

HipS interacts directly with HipT and consists of a solvent-facing, five-stranded anti-parallel β sheet of which four strands are located in the N terminus (residues 1-52). A small domain of three helices (residues 53-94) forms the interface to HipT and is followed by the fifth β strand (residues 95-102, **Figure 1b**). Overall, the fold of the HipST heterodimer is very structurally similar to the longer HipA kinase (**Figure 1 supplement 2b**), as expected from sequence analysis (**Figure 1**




Figure 1.

Crystal structure of the *E. coli* O127:H6 HipBST complex.

a. Schematic representation of HipBST and HipBA_{Ec} showing corresponding proteins and domains. Dashed lines indicate similar domains and grey areas represent regions missing domains in the crystal structures: *E. coli* O127:H6 HipBST, this work; *E. coli* HipBA, PDB: 2WIU (Evdokimov *et al.*, 2009). HipS (beige) is structurally similar to the N-subdomain 1 of HipA (dark purple), HipT has an additional N-terminal mini-domain not found in HipA (light blue), while HipA has a short linker between the two N-subdomains (light purple). **b.** *E. coli* O127:H6 HipBST forms a hetero-hexameric complex with HipS (beige) on top of HipT (blue) through dimerisation of HipB (green) also generating a helix-turn-helix DNA-binding motif (HTH, dashed box). **c.** Top, crystal structure of the *E. coli* K-12 HipBA complex (PDB: 2WIU) shown as cartoon with HipA in purple and blue purple (N-terminal subdomain 1) and the HipB homodimer in two shades of green (Evdokimov *et al.*, 2009); bottom, the structure of *S. oneidensis* HipBA (PDB: 4PU3, HipA purple, HipB green) bound to DNA (orange backbone, blue bases) (Wen *et al.*, 2014). **d.** The linker (dark purple) between N-subdomains 1 (N-sub 1) and 2 (N-sub 2) physically linking the two domains in HipA_{Ec}. **e.** The N-terminal mini-domain of HipT (blue), which is absent from HipA.

supplement 1b). Thus, HipS is structurally homologous to the N-subdomain 1 of HipA (Schumacher *et al.*, 2012 [↗](#)), both with respect to its overall fold and to its orientation with respect to HipT (**Figure 1 supplement 2b**). As a result of this, the HipS C-terminus overlaps with an extended linker that bridges N-subdomains 1 and 2 in HipA (**Figure 1d [↗](#) and figure 1 supplement 2b**). HipT corresponds structurally to the core kinase domain of HipA, except for an additional N-terminal mini-domain of unknown significance (residues 1-41, **Figure 1a [↗](#) and 1e [↗](#)**). In this structure, the Gly-rich loop of HipT is found in an outward-facing conformation despite the absence of Gly-rich loop phosphorylation, suggesting that HipT is inactive (**Figure 1 supplement 2c**).

In conclusion, we find that the overall structure of the tripartite HipBST complex is markedly different from the canonical *E. coli* HipBA complex, but structurally similar to a HipBA complex from *S. oneidensis*. Key features of HipB proposed to be involved in the kinase toxin inhibition mechanism in both HipBA_{Ec} and HipBA_{So} are missing. Finally, HipT appears in a canonical inactive conformation despite being unphosphorylated, suggesting a different mechanism of toxin inactivation and inhibition.

HipS inhibits HipT through insertion of a hydrophobic residue near the active site

The observation that HipT is found in a canonical, inactive conformation despite being unphosphorylated led us to hypothesize that HipS might neutralize HipT by direct interaction. This would be consistent with the observation that HipS functionally acts as the antitoxin of the HipBST system *in vivo* (Vang Nielsen *et al.*, 2019 [↗](#)). There are three major areas of contact between the two proteins in the HipBST complex, involving both hydrogen bonds, hydrophobic and charged interactions (**Figure 2a [↗](#)**). Importantly, HipT and HipS were found to interact directly very close to the Gly-rich loop, where Glu63 from HipS points directly into the HipT active site while HipS Trp65 is wedged into a deep cavity formed by the Gly-rich loop in its outward conformation (**Figure 2a [↗](#), right and 2b**). Of these, Glu63 is conserved in the N-subdomain 1 of the longer HipA toxins (**Figure 1 supplement 1b**), and therefore less likely contribute to functional differences between the two systems. Trp65, on the other hand, is found in many HipS orthologues (**Figure 2 supplement 1a**) and moreover, the region surrounding it is structurally different from the corresponding region in HipA_{Ec} (**Figure 2b [↗](#)**). Based on this, we hypothesized that insertion of the bulky Trp residue near the HipT active site might sterically block transition of the Gly-rich loop from its outward to the inward conformation, likely required for ATP binding and kinase activity. To test this, we generated a HipS^{W65A} variant and probed its ability to neutralize HipT toxicity upon ectopic expression of the two proteins using a two-plasmid system in *E. coli* MG1655. Dilution spot assays confirmed that *wt* HipS, but not HipS^{W65A}, can neutralize toxicity of HipT *in vivo* (**Figure 2c [↗](#)**). This result was reproduced by growth assays in liquid culture (**Figure 2 supplement 1b**). In conclusion, we show that HipS interacts directly with the HipT active site and neutralizes the kinase by insertion of a conserved tryptophan residue, likely preventing structural transition from the inactive to the active conformation. This mode of inhibition where the toxin is sterically and physically blocked by the antitoxin thus closely resembles that observed in other toxin-antitoxin systems.

The two phosphoserine positions in HipT are important for toxicity

To understand the functional implications of the two phosphoserine positions in HipT, Ser57 and Ser59 (Vang Nielsen *et al.*, 2019 [↗](#)), we initially decided to investigate their role in toxicity. For this, we designed a set of HipT variants with the serine residues individually substituted for either alanine (A, phosphoablative) to prevent autophosphorylation, or aspartic acid (D, phosphomimetic) to mimic the phosphorylated form. Growth assays in liquid culture showed that both wild-type HipT (S⁵⁷I⁵⁹; SIS), as well as HipT D⁵⁷I⁵⁹ (DIS) and S⁵⁷ID⁵⁹ (SID), caused growth inhibition upon overexpression in *E. coli* MG1655 (**Figure 3a [↗](#)**), demonstrating that the

phosphomimetic mutations do not prevent toxicity, and therefore, kinase activity of HipT towards its target. Moreover, since it has previously been shown that only the HipT Gly-rich loop never is observed in doubly phosphorylated form with both Ser57 and Ser59 modified simultaneously, it is unlikely that the effects are due to autophosphorylation of the remaining serine residue in either case (Vang Nielsen *et al.*, 2019). In addition, while co-expression of HipS neutralized the toxicity of HipT SIS and SID variants, we could not make the antitoxin neutralise the toxicity of the HipT DIS variant, suggesting that Ser57, but not Ser59, negatively impacts the function of HipS as antitoxin. Finally, alanine substitutions at either Ser position almost (S⁵⁷IA⁵⁹; SIA) or completely (A⁵⁷IS⁵⁹; AIS) abolished HipT toxicity (**Figure 3a**). Thus, both serine residues (in either phosphorylation state) are important for HipT toxicity. In summary, we find that introduction of a phosphomimetic mutation at Ser57 negatively affects the function of HipS as antitoxin under conditions where the HipT kinase is toxic. This represents a clear difference from the regulation of HipA, where phosphorylation of Ser150 inactivates the toxin (Schumacher *et al.*, 2012).

The active site conformation of HipT is independent of phosphorylation

To understand how phosphorylation of the two serine residues affects the structure of the HipBST complex, we determined two additional crystal structures in context of the HipT AIS (2.4 Å) and SIA (3.4 Å) variants (**Table 1**), which are non-toxic *in vivo*, but in principle maintain an intact kinase active site. Inspection of the difference maps in the vicinity of the active site revealed that in both cases, additional electron density was present close to the non-mutated serine positions in each structure (**Figure 3 supplement 1**), none of which were not observed for HipBST^{D233Q}. For the HipBST AIS variant, we found a strong signal corresponding to phosphorylation of Ser59 in both copies of HipT in the asymmetric unit (**Figure 3 supplement 1a**), while for the HipBST SIA variant, phosphorylation of Ser57 was present but incomplete and could only be confidently modelled in one of the two HipT molecules in the asymmetric unit (**Figure 3 supplement 1b**). In its phosphorylated state, Ser59 (P-Ser59) forms strong interactions to Lys161, His212, and the catalytic Asp210 inside the active site (**Figure 3b**, **top**), while P-Ser57 is located further away and forms hydrogen bonds to Tyr162 and Asp210 in its phosphorylated state (**Figure 3b**, **bottom**). In both structures, however, the Gly-rich loop maintains a similar outward conformation to the one observed before, likely incompatible with ATP binding. Taken together, the structures suggest that HipT is inactive in the context of the HipBST complex regardless of the phosphorylation state of the serine residues in the Gly-rich loop. This is distinctly different from HipA, where phosphorylation of Ser150 was shown to convert the kinase from an active to an inactive form by causing ejection of the Gly-rich loop (Schumacher *et al.*, 2012). To support these observations, we analysed the purified HipBST SIA and AIS variants on Coomassie-stained Phos-tag SDS-PAGE gels, which can separate different phosphoprotein species, using the inactive HipT^{D210A} variant as control for unphosphorylated HipT. This experiment was carried out both before and after separation on a Heparin column which we found is able to separate complexes based on the phosphorylation state of HipT. This confirmed that HipT AIS primarily is found on the phosphorylated form (**Figure 3c**, **upper band**), while HipT SIA is mostly on the unphosphorylated form (**Figure 3c**, **lower band**).

In summary, we conclude that phosphorylation of neither of the two phosphoserine positions in HipT affects the conformation of the Gly-rich loop in the context of the HipBST complex, and in all cases the loop remains in an outward conformation suggesting the kinase is not in its canonical, activated state. This supports that phosphorylation itself is not responsible for inactivation of the kinase and is consistent with a specific role for HipS in this process. Surprisingly, autophosphorylation still occurs in both the SIA and AIS variants demonstrating that the kinase is still active and consequently, that the lack of toxicity observed for these variants might relate to target binding rather than kinase inactivation. Finally, we note that the position of the Ser59




Figure 3.

The phosphoserine positions in HipT have distinct functional roles.

a. Growth curves of *E. coli* MG1655 harbouring arabinose-inducible, single autophosphorylation variants of HipT; pBAD33::*hipT* (SIS, wt), pBAD33::*hipT*^{S57D} (DIS), pBAD33::*hipT*^{S59D} (SID), pBAD33::*hipT*^{S57A} (AIS), pBAD33::*hipT*^{S59A} (SIA), or the empty pBAD33 vector, in combination with an IPTG-inducible construct of HipS; pNDM220::*hipS*, with expression induced at the indicated time points (ara/IPTG). The curves show mean OD₆₀₀ values from at least two independent experiments with error bars indicating standard deviations (hidden when small). **b.** Overview of the HipT kinase active site in the D233Q mutant as well as HipT^{S57A} (AIS, top) and HipT^{S59A} (SIA, bottom) structures. The phosphate groups on Ser57 (in HipT^{S59A}) and Ser59 (in HipT^{S57A}) are shown in orange and interacting nearby residues are highlighted. Numbers indicate distances in Å. **c.** HipT variants from purified HipBST complex before (-) and after (+) a Heparin-column purification step to separate complexes based on the phosphorylation state visualised on a Phos-tag gel, which separates proteins based on phosphorylation state, and stained by Coomassie Blue. The locations of phosphorylated (P-HipT) and unphosphorylated (HipT) protein species are indicated. The gels are representative of two independent experiments. **d.** Close-up of the HipT S57A active site overlaid with ATP (salmon, semi-transparent) and two Mg²⁺ ions (green, semi-transparent) from the structure of HipA:ATP (PDB: 3DNT) (Schumacher *et al.*, 2009). Relevant active site residues are shown as sticks and the Gly-rich loop is shown in dark red with pSer59 indicated.

phosphate group in the AIS variant (HipT^{S57A}) corresponds closely to that of the gamma-phosphate of ATP in the active site of HipA, suggesting that phosphorylation in fact may stabilise the active site in a similar way to ATP (**Figure 3d** [↗](#)).

HipBST dynamics in solution is affected by the phosphorylation state

To finally understand if the two phosphoserines affect HipBST complex stability, we constructed a kinase-inactive version of the complex containing the HipT^{D210A} active site mutant and analysed three different, phosphomimetic states of the Gly-rich loop (HipT SIS, DIS, and SID) in this context by purifying the HipBST complex using a C-terminal His-tag on HipT. Intriguingly, for the complex purified with the wildtype SIS Gly-rich loop, the band corresponding to HipS was missing (**Figure 4 supplement 1a**). As Asp210 interacts directly with Ser59, this supports that the serine residues are important for stabilisation of the active site and the Gly-rich loop in a conformation compatible with HipS binding as well as activity towards the target. Analysis of the purified samples of the three variants by small-angle X-ray scattering (SAXS) gave very similar intensity curves suggesting that the complexes have similar sizes and shapes in solution, despite the missing HipS protein in the SIS variant (**Figure 4a** [↗](#)). All Guinier plots show a linear behaviour at low angles demonstrating the absence of large aggregates (**Figure 4 supplement 1b**) while the indirect Fourier transformation (**Figure 4 supplement 1c**) and pair distance distribution functions (**Figure 4 supplement 1d**) support similar, overall structures. Interestingly, the maximum diameter of 200 Å is significantly larger than the crystal structure (max. diameter 135 Å), consistent with analysis of the forward scattering that revealed partial oligomerisation of the samples with an average mass corresponding to roughly a dimer of the HipBST heterohexamer. Finally, the normalised Kratky plot agrees with a relatively compact structure without significant random coil content (**Figure 4 supplement 1e**).

We next attempted to model the SAXS data using the crystal structure of HipBST, by including a structural prediction of the missing HipB N-terminus and a hydration layer (**Figure 4** [↗](#)). As expected from the predicted oligomerisation, this model was in relatively poor agreement with the measured data, so we allowed individual domains of the complex to undergo rigid-body motion while imposing C2 symmetry while at the same time refining the degree of oligomerization (Steiner *et al.*, 2018 [↗](#)). For each data set, the model with the best fit (lowest reduced χ^2) to the data out of 10 runs was selected as the representative model. These three models (**Figure 4b** [↗](#)) all gave good fits to the data (**Figure 4a** [↗](#)) with varying degrees of oligomerisation from 1.7-1.8 and an average distance between the heterohexamers ~80 Å. Consistent with the biochemical data, the SAXS data from the HipT SIS variant were matched very well by a model in which HipS was omitted but the disordered N-terminus of HipB was included as a helical element (**Figure 4** [↗](#), **blue**). In this structure, the two HipT modules have rotated slightly to open up the cleft between them as compared to the crystal structure (**Figure 4b** [↗](#), **blue**, **arrows**). On the contrary, data measured for both the SID (**Figure 4a and b** [↗](#), **red**) and DIS (**Figure 4a and b** [↗](#), **orange**) variants both matched the complete HipBST crystal structure (with the N-terminal HipB extension) with only minor adjustments of the domains. Taken together, SAXS analysis suggests that the HipBST complex is dynamic in solution in a phosphoserine-dependent way and that the phosphorylation state of HipT influences the inhibitory function of HipS. This dynamic behaviour likely explains how it is possible for the inactivated HipBST TA complex to undergo auto-phosphorylation *in vivo*.

The dual autophosphorylation sites are conserved among HipT kinases

To understand if Gly-rich loops with two phosphoserine positions are common among HipT-like kinases, we generated a phylogenetic tree of the 48 known HipT orthologues in bacterial genomes (Gerdes *et al.*, 2021 [↗](#)) (**Figure 5a** [↗](#)). Interestingly, investigation of this tree revealed that it largely separates the kinases by the configuration of the Gly-rich loop. The largest branch, which we




Figure 4.

The HipBST complex is dynamic in solution.

a. Experimental SAXS curves (measured x-ray intensity as a function of q , the modulus of the scattering vector) measured for HipBST in the context of HipT D210Q and the variants SIS (blue), SID (red), and DIS (orange). **b.** Structure models of HipBST (for SID, red, and DIS, orange) or HipBT (SIS, blue) as fit to the SAXS scattering data. For each model (SIS, SID, and DIS), the rmsd to the crystal structure is indicated while arrows indicate gross domain movements. Top left, the crystal structure of the HipBST D233Q for reference.

named the SΨS group, contains HipT kinases with two serine residues separated by a small aliphatic amino acid, corresponding to the known autophosphorylation sites in HipT of *E. coli* O127:H6, Ser57 and Ser59 (**Figure 5a** [↗](#)). In many of the remaining HipT orthologues, this region contains various Ser/Thr motifs, including (S/T)ΨT, (S/T)ΨP, SΨQ, and ΨΨT, but invariably with a central, small aliphatic residue (Ψ: Ile, Leu, or Val). Overall, the second Ser/Thr position appears less conserved than the first, and some kinases even contain a proline (P) or glutamine (Q) here, suggesting different regulation patterns for different HipT kinases with Ser57 playing a more important role. This supports the functional analysis showing that Ser57 affects neutralization by HipS *in vivo* and is less prone to autophosphorylation in the SIA variant. The remaining sequence of the Gly-rich loop across a range of HipT kinases is highly similar to HipA_{Ec} (**Figure 5b** [↗](#) and **5c** [↗](#)). Together, this allowed us to generate a consensus motif for the HipT Gly-rich loop, which includes several conserved hydrophobic positions in addition to the two Ser/Thr residues (**Figure 5d** [↗](#)). In summary, phylogenetic analysis reveals that a large fraction of HipT kinases contain a conserved SΨS motif with two potential phosphorylation positions in their Gly-rich loops, supporting that both Ser59, and in particular Ser57, play an important role in the regulation of the HipBST TA system.

Discussion

In this study, we present a detailed structural and functional analysis of the tripartite HipBST toxin-antitoxin system from enteropathogenic *E. coli* O127:H6, which elucidates the neutralising mechanism of the unusual HipS antitoxin and dissects the functional roles of the two phosphoserine residues in HipT, Ser57 and Ser59. The crystal structure of the kinase-inactive HipBST^{D233Q} complex shows that HipS interacts in a similar way to the corresponding N-subdomain 1 in HipA but neutralises HipT by direction interaction involving insertion a large, bulky residue, Trp65, into the kinase active site. Trp65 is conserved in some, but not all HipS proteins, and is located in a motif that differs both structurally and at the sequence level from the corresponding region in HipA, which is otherwise very similar to HipS ([Gerdes et al., 2021](#) [↗](#)). Using multiple experimental approaches, we go on to show that both phosphomimetic and phosphoablative states of HipT impact toxicity and the interaction with the antitoxin HipS. Together, this suggests that the structural basis for the role of HipS as antitoxin in the HipBST system involves preventing the Gly-rich loop from transitioning from the outward, inactive state, to a conformation compatible with ATP binding. In some organisms, the residue corresponding to Trp65 is a proline, which may point to another way of preventing activation via steric hindrance (**Figure 2** [↗](#) **figure supplement 1a**).

Recently, structures of a HipT orthologue from *Legionella pneumophila* (HipT_{Lp}) in complex with the non-hydrolyzable ATP analogue (AMPPNP), HipS_{Lp} in complex with HipT_{Lp}, and a HipS from *Haemophilus influenzae* have been determined ([Lin et al., 2022](#) [↗](#), [Zhen et al., 2022](#) [↗](#), [Koo et al., 2022](#) [↗](#)). HipT_{Lp} belongs to the SΨQ group, and phosphorylation of the Gly-rich loop can thus take place only at the position corresponding to Ser57 in HipT_{Ec}. This allows us to further dissect the functions of the two phosphorylation sites in HipT_{Ec}. Interestingly, and despite being phosphorylated, HipT_{Lp} is found with its active site loop in an inward and active conformation with nucleotide bound. In this conformation, the phosphoserine moiety is stabilized by two conserved arginine residues in a RxDR motif, which is found both in HipT_{Ec} and HipA_{Ec} (**Figure 1 supplement 1**). In HipA, however, the Gly-rich loop does not contact the RxDR motif in either phosphorylation state, suggesting that this interaction is a unique feature of HipT kinases ([Schumacher et al., 2012](#) [↗](#)). Gln56 (SΨQ), which corresponds to Ser59 in HipT_{Ec} and arguably in some respects mimics a phosphorylated serine, interacts with two residues, Asp145 and Lys201. These have chemically similar counterparts in HipT_{Ec} (Gln151 and Arg214) and thus might interact with the phosphorylated Ser59 when the Gly-rich loop is in its inward state, i.e., in the absence of HipS. Interactions at both sites (Ser57 to the RxDR motif, Ser59 to Gln151/Arg214) would stabilize the loop in its inward state and possibly affect the function of HipS as antitoxin.




Figure 5.

Phylogenetic analysis of HipT.

a. Phylogenetic guide tree of 48 HipT orthologues with sequences motifs (potential phosphorylation sites) indicated on the side. The SWS group (red) is by far the largest group followed by the T[IV][TP] group (green), and SIQ group (blue). Sequences used in the alignment in **b** are shown with red letters and the numbers on the tree branches indicate bootstrap confidence levels. **b.** Sequences of the Gly-rich loop (orange background), including the potential phosphorylation sites for selected HipT orthologues compared to HipA from *E. coli* K-12. Known phosphorylation sites in *E. coli* O127:H6 HipT (top) and *E. coli* K-12 HipA (bottom) are indicated with arrows and conserved sequence motifs with bold white text. **c.** Sequence logo for the Gly-rich loop derived from all 48 HipT sequences. **d.** Consensus motif for the HipT Gly-rich loop with known phosphorylation sites in red. Φ indicates a hydrophobic residue, while Ψ are aliphatic residues.

Moreover, since interaction of the phosphoserine with two arginine side chains at Ser57 would be much stronger than to a Gln and an Arg, due to the ionic character of the side chains, this could also explain why P-Ser57 interaction is stronger and thus potentially also why HipS is unable to inactivate the S57D phosphomimetic HipT mutant (DIS).

Finally, we show that the serine residues in the HipT Gly-rich loop (Ser57 and Ser59) are important for toxicity but apparently not for auto-phosphorylation as we observe auto-phosphorylation in both the AIS and SIA variants of HipT. This is compatible with a model in which auto-phosphorylation serves to both control target (TrpS) phosphorylation and HipS binding, which could explain the need for two sites. The molecular mechanism controlling at which site auto-phosphorylation takes place is still not known, but the structures of HipBST AIS and HipBST SIA suggest that subtle changes in the loop can affect this (**Figure 3b** [↗](#)).

Together, our data support a complex network model for the interplay between the active site and the Gly-rich loop depicted in **Figure 6** [↗](#). In this model, residues involved in both ATP binding and catalysis, the phosphoserine residues in the Gly-rich loop, and parts of HipS form a tightly interconnected network of interactions. The result of this complexity is that posttranslational modification and/or mutation of any single residue can have multiple downstream effects. For example, inactivation of the HipT active site through the D210A mutation appears to also affect the position of the Gly-rich loop as well as HipS binding, via direct interactions to the serine residues. Likewise, the position of the Gly-rich loop and binding of HipS are mutually interconnected so that HipS controls whether the loop can transition from an outward to an inward state and conversely, the serine residues and their phosphorylation state likely also control with what affinity HipS binds. This shifts the role of HipS from that of a classical antitoxin that simply blocks and inhibits the toxin active site to that of an allosteric enzyme modulator. We do not yet understand whether auto-phosphorylation can take place with HipS bound or requires its release but note that our model is consistent with both these scenarios. The complexity of the network therefore makes it difficult to associate single roles to specific residues and/or modifications and requires that the full system is considered for each functional state.

Acknowledgements

The authors are indebted to the beamline staff at P14 in EMBL Hamburg, and BioMAX in MaxIV Lund for help during data collection. This project was funded by grants from the Novo Nordisk Foundation (NNF18OC0030646 to D.E.B.) and Danish Natural Research Foundation's Centre of Excellence for Bacterial Stress Response and Persistence (grant number DNRF120).

Competing interests

The authors confirm that there are no competing interests.

Author contributions

Conceptualization, K.G., M.A.S. and D.E.B.; Investigation, R.L.B., S.V.N., F. B., J. S.P. and J.L.; Formal Analysis, R.L.B., S.V.N., R. B. S., F.B., and J.S.P; Writing – Original Draft, R.L.B., S.V.N., and D.E.B.; Writing – Review & Editing, R.L.B., S.V.N., R. B. S., J.L., J.S.P., K.G., M.A.S., and D.E.B; Visualization, R.L.B., S.V.N., M.A.S., and D.E.B.; Funding Acquisition, D.E.B. and K.G.; Resources, J.S.P., K.G., M.A.S. and D.E.B; Supervision, K.G., J.S.P., M.A.S., and D.E.B.




Figure 6.

Model for the active site network of HipT of the HipBST system.

Schematic overview of HipT showing the interactions found in this study. HipT (blue) with the observed outward conformation of the Gly-rich loop (red), and the predicted inward conformation (green). Important residues highlighted are Ser57, Ser59, Asp210 and Asp233 of HipT, and Trp65 of HipS (beige). Shown is also the position of an ATP molecule (orange) together with two Mg-ions (green spheres) based loosely on PDB: 7WCF. The two sites homologous to the sites found in HipT from *L. pneumophila* to interact with the phosphoserine positions (R137/R140 and Q151/R214 in HipT_{Ec}) are also shown since they potentially offer a stabilizing role.

Strains and plasmids

Strains and plasmids used or generated in this study are listed in **Table S2**, and DNA oligonucleotides in **Table S3**. The construct expressing the HipT^{S57A} variant was constructed by PCR mutagenesis (**Table S3**) as previously described (Vang Nielsen *et al.*, 2019 [DOI](#)).

Protein purification and structure determination

Expression of the *E. coli* O127:H6 HipBST complex was done by introduction of the constructs pSVN78 (HipBST^{S57A}), pSVN96 (HipBST^{D233Q}), and pMME3 (HipBST^{S59A}), into *E. coli* BL21 DE3. All constructs encode HipB, HipS, and C-terminal hexa-histidine tagged version of HipT, with genes separated and including individual, optimised Shine-Dalgarno sequences. All constructs contained an isopropyl-D-1 thio-galactopyranoside (IPTG)-inducible promoter and ampicillin resistance gene for selection. For each construct, 2 L cultures of *E. coli* BL21 DE3 grown in LB medium to a cell density of OD₆₀₀ = 0.6 were induced with a final concentration of 1 mM IPTG and left to express overnight at 20°C with vigorous shaking. Cell pellets were resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 20 mM imidazole, 5% glycerol, 5 mM BME, 1 mM PMSF) and lysed by sonication. The lysate was cleared by centrifugation and applied to a 5 mL HisTrap HP column (Cytiva) equilibrated in lysis buffer and subsequently washed in wash buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 40 mM imidazole, 5 mM BME), before eluting with 50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 400 mM imidazole, 5 mM BME. The eluate was applied directly to a 5 mL Heparin HF column equilibrated in 70% Buffer A (50 mM Tris-HCl, pH 7.5, 5 mM BME) and 30% Buffer B (50 mM Tris-HCl, pH 7.5, 1M NaCl, 5 mM BME) attached to an ÄKTA Pure system (Cytiva). This step separated the population into fully formed HipBST complexes that bound to the column, and various partial complexes that were washed off. Final separation was achieved following concentrating the Heparin elution to approximately 8 mg/mL using a 30 kDa MWCO Vivaspin filter (Sartorius), by applying the sample to a Superdex 200 10/300 GL (Cytiva) column equilibrated in gel filtration buffer (20 mM Tris-HCl, pH 7.5, 300 mM NaCl, 5 mM BME). Finally, the sample was concentrated to a protein concentration of 5 mg/mL by spin filtration before crystallisation. Crystals of HipBST mutants grew as clusters of thin plates in 2 µL drops containing a 1:1 ratio of protein to the crystallisation buffer, which consisted of 0.1 M Bicine, pH 9, 8% MPD, set up against a reservoir of 200 µL of the same condition. Crystals were cryoprotected in crystallisation buffer supplemented with 25% MPD before freezing in liquid N₂.

Data collection for HipBST^{S57A} was carried out at the P14 beamline at EMBL/DESY, Hamburg and for HipBST^{S59A} and HipBST^{D233Q} at the BioMAX beamline at MAX IV in Lund, Sweden. For HipBST^{S59A} and HipBST^{D233Q}, 7,200 images were collected with an oscillation of 0.1° and a transmission of 100% while for HipBST^{S57A}, 3,600 images were collected with an oscillation of 0.1° and a transmission of 70%. All data were processed in XDS (Kabsch, 2010 [DOI](#)), using the CC_{1/2} value after scaling to set the diffraction limit (Karplus & Diederichs, 2012 [DOI](#)). The space group was confirmed using Pointless (Evans, 2006 [DOI](#)), and all structures were determined using molecular replacement with Phaser (McCoy *et al.*, 2007 [DOI](#)) from the CCP4 suite (Winn *et al.*, 2011 [DOI](#)). For HipBST^{S57A}, a heavily truncated structure of *E. coli* HipA was used as search model, while for HipBST^{S59A} and HipBST^{D233Q}, the preliminary structure of HipBST^{S57A} was used. Iterative refinement was carried out in Buster (Smart *et al.*, 2012 [DOI](#)) using one big cycle of 20 small cycles of refinement during model building, and 5 big cycles of 100 small cycles of refinement for the subsequent polishing. Non-crystallographic symmetry (NCS) restraints were used throughout since the asymmetric unit (ASU) contained a dimer, but some refinement rounds without NCS were also included to allow the model to adapt to differences between the molecules in the ASU. Automatic water placement was performed in Buster and water molecules were manually pruned by inspecting the electron density. Model building was performed in Coot (Emsley *et al.*, 2010 [DOI](#)). All

structures were validated by the MolProbity server, and Rama-Z scores as calculated by Phenix (Sobolev *et al.*, 2020 [DOI](#)). Final R-work/R-free for HipBST_{D233Q}, HipBST_{S57A}, and HipBST_{S59A}, were 0.19/0.23, 0.21/0.24, and 0.20/0.24, respectively.

Spot assays and growth curves

Cultures of relevant strains of *E. coli* were grown in liquid YT medium or MOPS minimal medium including 0.2% glucose at 37°C with shaking at 160 rpm. YT agar plates were used as solid medium and were incubated at 37°C for approximately 16 h. For selection, media was supplemented with 25 µg/mL chloramphenicol, 30 µg/mL ampicillin, and/or 25 µg/mL kanamycin. For spot assays, *E. coli* cells were grown as overnight cultures, diluted to obtain identical OD₆₀₀, centrifuged at 5,000 rpm for 5 min, washed in phosphate-buffered saline (PBS), and serially diluted before being spotted onto YT agar plates containing the indicated amount of inducer or repressor. Gene expression from plasmids carrying the pBAD promoter was induced by a final concentration of 0.2% arabinose and repressed by 0.2% glucose. Gene expression from plasmids carrying the P_{A1/04/03} promoter was induced by a final concentration of 200 or 500 µM IPTG, as indicated. Growth experiments were done in YT medium with addition of relevant antibiotics, diluted from overnight cultures and grown exponentially for at least 4 h until a constant doubling time was observed. At OD₆₀₀ ~0.2, 0.2% arabinose was added to induce expression of wildtype HipT or autophosphorylation mutants. After another 1.5 h, 200 µM IPTG was added to induce expression of HipS. For each repetition, an independent colony from the strain was used to start separate cultures. No outliers were rejected.

Phos-tag gel

15% Phos-tag acrylamide gels (Wako) were cast according to the manufacturer's guidelines, except that 100 µM Phos-tag acrylamide was added to ensure proper separation between phosphorylated and unphosphorylated HipT. The unphosphorylated inactive HipT^{S57A+D210A} control was expressed from pSNN2. The gel was run at 4 °C until the loading dye reached the bottom of the gel and visualized using standard Coomassie Blue staining as for normal SDS-PAGE gels.

Small-angle X-ray scattering measurements and analysis

HipBST^{D210A}, HipBST^{S57D,D210A}, and HipBST^{S59D,D210A} were expressed from pRBS1, pRBS2 and pRBS3, and purified as described above. SAXS measurements were performed using an optimized NanoSTAR instrument (Bruker AXS), which uses a high brilliance Ga metal-jet X-ray source (Excillum), special long optics, a two-pinhole collimation with a custom 'scatterless' slit pinhole, and a VÅNTEC-2000 (Bruker AXS) microgap 2D gas proportional detector (Lyngsø & Pedersen, 2021 [DOI](#)). Samples and buffer standards were measured in the same flow-through quartz capillary, and the scattering from the buffer was subtracted before the data were converted to absolute scale using the scattering from water. SAXS intensity data, $I(q)$, were analysed as a function of the modulus of the scattering vector $q = 4\pi \sin(\theta) / \lambda$, where 2θ is the scattering angle and λ is the wavelength of the X-rays. Guinier analysis, giving radius of gyration R_g and the forward scattering $I(0)$ the forward scattering, and an indirect Fourier transformation were then calculated (Glatter, 1977 [DOI](#), Pedersen *et al.*, 1994 [DOI](#)), giving the same parameters and also the pair distance distribution function $p(r)$ and the maximum diameter D_{max} of the particles. The predicted molar mass was then calculated by $M = I(0)N_A / (c\Delta\rho_m^2)$, where N_A is Avogadro's number, c is the mass concentration, and $\Delta\rho_m = 2.00 \times 10^{10} \text{ cm g}^{-1}$ is the typical excess scattering length per unit mass for proteins. The data was also plotted in a normalized Kratky plot of $(qR_g)^2 I(q)/I(0)$ versus qR_g to assess compactness and check for contributions from random coil parts. For structural modelling, a AlpheFold2-predicted structure for N-terminus of HipB was added to initially have a molecular model with the correct molecular mass. The predicted SAXS curve of the model was then

calculated as described in (Steiner *et al.*, 2018 [DOI](#)) where the hydration layer is described by dummy atoms. For the calculation, the Debye equation is used (Debye, 1915 [DOI](#)) using an average form factor for all non-hydrogen atoms:

$$I(q) = cM\Delta\rho_m^2 P(q) = cM\Delta\rho_m^2 \exp(-q^2\sigma^2) \sum_{i,j} b_i b_j \frac{\sin(qd_{ij})}{qd_{ij}}$$

which defines the form factor $P(q)$, and where $\sigma = 1.0 \text{ \AA}$, b_i is, respectively, equal to the average excess scattering length of a non-hydrogen atoms for the protein and equal to an average excess scattering length for a hydration dummy atom. The parameter d_{ij} is the distance between the i 'th and the j 'th atom. Poor agreement was observed for all the data sets, so to improve the modelling, rigid-body refinement (RBR) of the structure was performed using three bodies for HipB (residues 1-31, 32-43, and 44-107, respectively), one body for the HipS, and three bodies for HipT (residues 2-59, 60-169, and 170-331, respectively). Distance restraints were then added at the domain boundaries with additional restraints between the domains (Glu A107-Val D101, Pro B55-Gly C154, Trp B 65-Gly C60, Gly D100-Aap F188, Leu D6-Arg F291, Val C147-Gly B94) based on analysis of the crystal structure, and finally a restraint of excluded volume was added. The Guinier and IFT analysis indicated some oligomerization in the samples, and therefore, a structure factor $S(q) = 1 + (N - 1) \sin(qD)/qD$ (Larsen *et al.*, 2020 [DOI](#)) for a dimer was included. In the expression, where N is the average of the number of proteins in the oligomer, and D is the distance between the two copies in the dimer. The structure factor was included in the decoupling approximation (Kotlarchyk & Chen, 1983 [DOI](#)):

$$I(q)/(cM\Delta\rho_m^2) = P(q) + \frac{\langle A(q) \rangle^2}{P(q)} [S(q) - 1]$$

with corresponding amplitude,

$$\langle A(q) \rangle = \exp\left(-\frac{q^2\sigma^2}{2}\right) \sum_i b_i \frac{\sin(qd_{i,CM})}{qd_{i,CM}}$$

where $d_{i,CM}$ is the distance of the i 'th atom from the scattering center of mass. During the optimization of the structure, the two parameters of the structure factor, N and D , were also varied.

Phylogenetic analysis

The previously identified set of 48 HipT orthologues (Gerdes *et al.*, 2021 [DOI](#)) was used for sequence alignment by Clustal Omega (Sievers & Higgins, 2018 [DOI](#)) at www.ebi.ac.uk (<http://www.ebi.ac.uk/> [DOI](#)) and imported into Jalview (Waterhouse *et al.*, 2009 [DOI](#)). The phylogenetic tree was visualized using iTOL (Letunic & Bork, 2019 [DOI](#)). Reconstruction of the phylogenetic tree was accomplished using IQ-TREE that uses the Maximum Likelihood approach and Ultrafast bootstrapping via the CIPRES module in Genious Prime (Minh *et al.*, 2020 [DOI](#)).

Data and software availability

The accession numbers for the structures reported in this paper are PDB: 7AB3 (HipBST SIA), 7AB4 (HipBST AIS), and 7AB5 (HipBST^{D233Q}).

References

- Bordes P., Sala A.J., Ayala S., Texier P., Slama N., Cirinesi A.M., Guillet V., Mourey L., Genevoux P. (2016) **Chaperone addiction of toxin-antitoxin systems** *Nat Commun* **7**
- Debye P (1915) **Zerstreuung von Röntgenstrahlen** *Annalen der Physik* **351**:809–823
- Emsley P., Lohkamp B., Scott W.G., Cowtan K. (2010) **Features and development of Coot** *Acta Crystallogr D Biol Crystallogr* **66**:486–501
- Eriani G., Delarue M., Poch O., Gangloff J., Moras D. (1990) **Partition of tRNA synthetases into two classes based on mutually exclusive sets of sequence motifs** *Nature* **347**:203–206
- Evans P (2006) **Scaling and assessment of data quality** *Acta Crystallogr D Biol Crystallogr* **62**:72–82
- Evdokimov A., Voznesensky I., Fennell K., Anderson M., Smith J.F., Fisher D.A. (2009) **New kinase regulation mechanism found in HipBA: a bacterial persistence switch** *Acta Crystallogr D Biol Crystallogr* **65**:875–879
- Gerdes K., Baerentsen R., Brodersen D.E. (2021) **Phylogeny Reveals Novel HipA-Homologous Kinase Families and Toxin-Antitoxin Gene Organizations** *mBio* **12**
- Germain E., Castro-Roa D., Zenkin N., Gerdes K. (2013) **Molecular mechanism of bacterial persistence by HipA** *Mol. Cell* **52**:248–254
- Glatter O (1977) **A new method for the evaluation of small-angle scattering data** *Journal of applied crystallography* **10**:415–421
- Hallez R., Geeraerts D., Sterckx Y., Mine N., Loris R., Van Melderen L. (2010) **New toxins homologous to ParE belonging to three-component toxin-antitoxin systems in Escherichia coli O157:H7** *Mol. Microbiol* **76**:719–732
- Hanks S.K., Quinn A.M., Hunter T. (1988) **The protein kinase family: conserved features and deduced phylogeny of the catalytic domains** *Science* **241**:42–52
- Harms A., Brodersen D.E., Mitarai N., Gerdes K. (2018) **Toxins, Targets, and Triggers: An Overview of Toxin-Antitoxin Biology** *Mol. Cell* **70**:768–784
- Haseltine W.A., Block R. (1973) **Synthesis of guanosine tetra- and pentaphosphate requires the presence of a codon-specific, uncharged transfer ribonucleic acid in the acceptor site of ribosomes** *Proc. Natl. Acad. Sci. U S A* **70**:1564–1568
- Huse M., Kuriyan J. (2002) **The conformational plasticity of protein kinases** *Cell* **109**:275–282
- Jumper J., Evans R., Pritzel A., Green T., Figurnov M., Ronneberger O., Tunyasuvunakool K., Bates R., Zidek A., Potapenko A., Bridgland A., Meyer C., Kohl S.A.A., Ballard A.J., Cowie A., Romera-Paredes B., Nikolov S., Jain R., Adler J., Back T., Petersen S., Reiman D., Clancy E., Zielinski M., Steinegger M., Pacholska M., Berghammer T., Bodenstein S., Silver D., Vinyals O., Senior A.W., Kavukcuoglu K., Kohli P., Hassabis D. (2021) **Highly accurate protein structure prediction with AlphaFold** *Nature* **596**:583–589

Jurenas D., Fraikin N., Goormaghtigh F., De Bruyn P., Vandervelde A., Zedek S., Jove T., Charlier D., Loris R., Van Melderen L. (2021) **Bistable Expression of a Toxin-Antitoxin System Located in a Cryptic Prophage of Escherichia coli O157:H7** *mBio* **12**

Kabsch W (2010) **Xds** *Acta Crystallogr D Biol Crystallogr* **66**:125–132

Karplus P.A., Diederichs K. (2012) **Linking crystallographic model and data quality** *Science* **336**:1030–1033

Koo J.S., Kang S.-M., Jung W.-M., Kim D.-H., Lee B.-J. (2022) **The Haemophilus influenzae HipBA toxin-antitoxin system adopts an unusual three-component regulatory mechanism** *IUCr* **9**:625–631

Kotlarchyk M., Chen S.H. (1983) **Analysis of small angle neutron scattering spectra from polydisperse interacting colloids** *The Journal of Chemical Physics* **79**:2461–2469

Larsen A.H., Pedersen J.S., Arleth L. (2020) **Assessment of structure factors for analysis of small-angle scattering data from desired or undesired aggregates** *Journal of applied crystallography* **53**:991–1005

LeRoux M., Laub M.T. (2022) **Toxin-Antitoxin Systems as Phage Defense Elements** *Annu. Rev. Microbiol*

Letunic I., Bork P. (2019) **Interactive Tree Of Life (iTOL) v4: recent updates and new developments** *Nucleic Acids Res* **47**:W256–W259

Lin J.D., Stogios P.J., Abe K.T., Wang A., MacPherson J., Skarina T., Gingras A.-C., Savchenko A., Ensminger A.W. (2022) **Lin, J.D., P.J. Stogios, K.T. Abe, A. Wang, J. MacPherson, T. Skarina, A.-C. Gingras, A. Savchenko & A.W. Ensminger, (2022).**

Lyngsø J., Pedersen J.S. (2021) **A high-flux automated laboratory small-angle X-ray scattering instrument optimized for solution scattering** *Journal of applied crystallography* **54**:295–305

McCoy A.J., Grosse-Kunstleve R.W., Adams P.D., Winn M.D., Storoni L.C., Read R.J. (2007) **Phaser crystallographic software** *J Appl Crystallogr* **40**:658–674

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A., Lanfear R. (2020) **IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era** *Mol. Biol. Evol* **37**:1530–1534

Pacios O., Blasco L., Bleriot I., Fernandez-Garcia L., Ambroa A., Lopez M., Bou G., Canton R., Garcia-Contreras R., Wood T.K., Tomas M. (2020) **(p)ppGpp and Its Role in Bacterial Persistence: New Challenges** *Antimicrob. Agents Chemother* **64**

Pedersen J.S., Hansen S., Bauer R. (1994) **The aggregation behavior of zinc-free insulin studied by small-angle neutron scattering** *Eur. Biophys. J* **22**:379–389

Schumacher M.A., Balani P., Min J., Chinnam N.B., Hansen S., Vulic M., Lewis K., Brennan R.G. (2015) **HipBA-promoter structures reveal the basis of heritable multidrug tolerance** *Nature* **524**:59–64

- Schumacher M.A., Min J., Link T.M., Guan Z., Xu W., Ahn Y.H., Soderblom E.J., Kurie J.M., Evdokimov A., Moseley M.A., Lewis K., Brennan R.G. (2012) **Role of unusual P loop ejection and autophosphorylation in HipA-mediated persistence and multidrug tolerance** *Cell Rep* **2**:518–525
- Schumacher M.A., Piro K.M., Xu W., Hansen S., Lewis K., Brennan R.G. (2009) **Molecular mechanisms of HipA-mediated multidrug tolerance and its neutralization by HipB** *Science* **323**:396–401
- Sekine S., Nureki O., Dubois D.Y., Bernier S., Chenevert R., Lapointe J., Vassilyev D.G., Yokoyama S. (2003) **ATP binding by glutamyl-tRNA synthetase is switched to the productive mode by tRNA binding** *EMBO J* **22**:676–688
- Sievers F., Higgins D.G. (2018) **Clustal Omega for making accurate alignments of many protein sequences** *Protein Sci* **27**:135–145
- Smart O.S., Womack T.O., Flensburg C., Keller P., Paciorek W., Sharff A., Vonnrhein C., Bricogne G. (2012) **Exploiting structure similarity in refinement: automated NCS and target-structure restraints in BUSTER** *Acta Crystallogr D Biol Crystallogr* **68**:368–380
- Sobolev O.V., Afonine P.V., Moriarty N.W., Hekkelman M.L., Joosten R.P., Perrakis A., Adams P.D. (2020) **A Global Ramachandran Score Identifies Protein Structures with Unlikely Stereochemistry** *Structure* **28**:1249–1258
- Stancik I.A., Sestak M.S., Ji B., Axelson-Fisk M., Franjevic D., Jers C., Domazet-Loso T., Mijakovic I. (2018) **Serine/Threonine Protein Kinases from Bacteria, Archaea and Eukarya Share a Common Evolutionary Origin Deeply Rooted in the Tree of Life** *J. Mol. Biol* **430**:27–32
- Steiner E.M., Lyngso J., Guy J.E., Bourenkov G., Lindqvist Y., Schneider T.R., Pedersen J.S., Schneider G., Schnell R. (2018) **The structure of the N-terminal module of the cell wall hydrolase RipA and its role in regulating catalytic activity** *Proteins* **86**:912–923
- Vang Nielsen S., Turnbull K.J., Roghanian M., Baerentsen R., Semanjski M., Brodersen D.E., Macek B., Gerde K. (2019) **Serine-Threonine Kinases Encoded by Split hipA Homologs Inhibit Tryptophanyl-tRNA Synthetase** *mBio* **10**
- Waterhouse A.M., Procter J.B., Martin D.M., Clamp M., Barton G.J. (2009) **Jalview Version 2-- a multiple sequence alignment editor and analysis workbench** *Bioinformatics* **25**:1189–1191
- Wen Y., Behiels E., Felix J., Elegheert J., Vergauwen B., Devreese B., Savvides S.N. (2014) **The bacterial antitoxin HipB establishes a ternary complex with operator DNA and phosphorylated toxin HipA to regulate bacterial persistence** *Nucleic Acids Res* **42**:10134–10147
- Winn M.D., Ballard C.C., Cowtan K.D., Dodson E.J., Emsley P., Evans P.R., Keegan R.M., Krissinel E.B., Leslie A.G., McCoy A., McNicholas S.J., Murshudov G.N., Pannu N.S., Potterton E.A., Powell H.R., Read R.J., Vagin A., Wilson K.S. (2011) **Overview of the CCP4 suite and current developments** *Acta Crystallogr D Biol Crystallogr* **67**:235–242
- Winther K.S., Roghanian M., Gerdes K. (2018) **Activation of the Stringent Response by Loading of RelA-tRNA Complexes at the Ribosomal A-Site** *Mol. Cell* **70**:95–105

Zhen X., Wu Y., Ge J., Fu J., Ye L., Lin N., Huang Z., Liu Z., Luo Z.Q., Qiu J., Ouyang S. (2022) **Molecular mechanism of toxin neutralization in the HipBST toxin-antitoxin system of *Legionella pneumophila*** *Nat Commun* **13**

Zielenkiewicz U., Ceglowski P. (2005) **The toxin-antitoxin system of the streptococcal plasmid pSM19035** *J. Bacteriol* **187**:6094–6105

Author information

René L. Bærentsen

Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 81, DK-8000 Aarhus C, Denmark

ORCID iD: [0000-0002-6460-4845](https://orcid.org/0000-0002-6460-4845)

Stine Vang Nielsen

Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 København N

ORCID iD: [0000-0002-4667-7979](https://orcid.org/0000-0002-4667-7979)

Ragnhild Bager Skjerning

Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 81, DK-8000 Aarhus C, Denmark

ORCID iD: [0000-0003-3886-8086](https://orcid.org/0000-0003-3886-8086)

Jeppe Lyngsø

Department of Chemistry and Interdisciplinary Nanoscience Centre (iNANO), Gustav Wieds Vej 14, DK-8000 Aarhus C

ORCID iD: [0000-0002-6301-1300](https://orcid.org/0000-0002-6301-1300)

Francesco Bisiak

Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 81, DK-8000 Aarhus C, Denmark

ORCID iD: [0000-0001-5375-6264](https://orcid.org/0000-0001-5375-6264)

Jan Skov Pedersen

Department of Chemistry and Interdisciplinary Nanoscience Centre (iNANO), Gustav Wieds Vej 14, DK-8000 Aarhus C

ORCID iD: [0000-0002-7768-0206](https://orcid.org/0000-0002-7768-0206)

Kenn Gerdes

Voldmestergade 8, DK-2100 København Ø, Denmark

ORCID iD: [0000-0002-7462-4612](https://orcid.org/0000-0002-7462-4612)

Michael A. Sørensen

Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 København N

For correspondence: mas@bio.ku.dk

ORCID iD: [0000-0001-8931-2999](https://orcid.org/0000-0001-8931-2999)

Ditlev. E. Brodersen

Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 81, DK-8000 Aarhus C, Denmark

For correspondence: deb@mbg.au.dk

ORCID iD: [0000-0002-5413-4667](https://orcid.org/0000-0002-5413-4667)

Editors

Reviewing Editor

Michael Laub

Howard Hughes Medical Institute, Massachusetts Institute of Technology, United States of America

Senior Editor

Jonathan Cooper

Fred Hutchinson Cancer Research Center, United States of America

Reviewer #1 (Public Review):

This work describes a structural analysis of the tripartite HipBST toxin-antitoxin (TA) system, which is related to the canonical two-component HipBA system composed of the HipA serine-threonine kinase toxin and the HipB antitoxin. The crystal structure of the kinase-inactive HipBST complex of the Enteropathogenic *E. coli* O127:H6 was solved and revealed that HipBST forms a hetero-hexameric complex composed of a dimer of HipBST heterotrimers that interact via the HipB subunit. The HipS antitoxin shows a structural resemblance to HipA N-terminal region and the HipT toxin represents to the core kinase domain of HipA, indicating that in HipBST the hipA toxin gene was likely split in two genes, namely hipS and hipT.

The structure also reveals a conserved and essential Trp residue within the HipS antitoxin, which likely prevents the conserved "Gly-rich loop" of HipT from adopting an inward conformation needed for ATP binding. This work also shows that the regulating Gly-rich loop of the HipT toxin contains conserved phosphoserine residues essential for HipT toxicity that are key players within the HipT active site interacting network and which likely control antitoxin binding and/or activity.

Strengths:

The manuscript is well written and the experimental work well executed. It shows that major features of the classical two-component HipAB TA system have somehow been rerouted in the case of the tripartite HipBST. This includes the N-terminal domain of the HipA toxin, which now functions as bona fide antitoxin, and the partly relegated HipB antitoxin, which could only function as a transcription regulator. In addition, this work shows a new mode of inhibition of a kinase toxin and highlights the impact of the phosphorylation state of key toxin residues in controlling the activity of the antitoxin.

Weaknesses:

A major weakness of this work is the lack of data concerning the role of HipB, which likely does not act as an antitoxin. Does it act as a transcriptional regulator of the hipBST operon and to what extent both HipS and HipT contribute to such regulation? These are still open questions.

In addition, there is no in-depth structural comparison between the structure of the HipBST solved in the work and the two recent structures of HipBST from *Legionella*. This is also a major weakness of this work.

Reviewer #2 (Public Review):

The work by Bærentsen et al., entitled "Structural basis for regulation of a tripartite toxin-antitoxin system by dual phosphorylation" deals with the structural aspects of the control of the hipBST TA operon, the role of auto-phosphorylation in the activation and neutralisation of the enzyme and the direct effects of HipS and HipB in neutralisation. This is a follow-up to the Vang Nielsen et al., and Gerdes et al., papers from the same authors on this very unique TA module, that brings forth a thorough and well written dissection of an unusually complex regulatory system.

This is a much improved manuscript, the paper is more focused and the message is now clear.

Author Response:

Reviewer #1 (Public Review):

[...] Strengths:

The manuscript is well written and the experimental work well executed. It shows that major features of the classical two-component HipAB TA system have somehow been rerouted in the case of the tripartite HipBST. This includes the N-terminal domain of the HipA toxin, which now functions as bona fide antitoxin, and the partly relegated HipB antitoxin, which could only function as a transcription regulator. In addition, this work shows a new mode of inhibition of a kinase toxin and highlights the impact of the phosphorylation state of key toxin residues in controlling the activity of the antitoxin.

Weaknesses:

A major weakness of this work is the lack of data concerning the role of HipB, which likely does not act as an antitoxin. Does it act as a transcriptional regulator of the hipBST operon and to what extent both HipS and HipT contribute to such regulation? These are still open questions.

We thank the reviewer for their feedback and will include a supplementary figure (Figure 1 supplement 2) and accompanying text that shows the transcriptional role of HipB, and how HipS and HipT influence this regulatory effect.

*In addition, there is no in-depth structural comparison between the structure of the HipBST solved in the work and the two recent structures of HipBST from *Legionella*. This is also a major weakness of this work.*

A structural comparison to the recent structures from *Legionella* will be included in the discussion, including Figure 6 supplement 1.