

Allosteric regulation of kinase activity in living cells

Shivani Godbole, Nikolay V. Dokholyan 

Department of Pharmacology, Penn State College of Medicine, Hershey, PA 17033-0850, USA • Department of Biomedical Engineering, Penn State University, University Park, PA 16802, USA • Department of Engineering Science and Mechanics, Penn State University, University Park, PA 16802, USA • Department of Biochemistry & Molecular Biology, Penn State College of Medicine, Hershey, PA 17033-0850, USA • Department of Chemistry, Penn State University, University Park, PA 16802, USA

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

October 12, 2023 (this version)

Reviewed preprint version 1

September 21, 2023

Posted to bioRxiv

July 19, 2023

Sent for peer review

March 8, 2023

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The dysregulation of protein kinases is associated with multiple diseases due to the kinases' involvement in a variety of cell signaling pathways. Manipulating protein kinase function, by controlling the active site, is a promising therapeutic and investigative strategy to mitigate and study diseases. Kinase active sites share structural similarities making it difficult to specifically target one kinase, allosteric control allows specific regulation and study of kinase function without directly targeting the active site. Allosteric sites are distal to the active site but coupled via a dynamic network of inter-atomic interactions between residues in the protein. Establishing an allosteric control over a kinase requires understanding the allosteric wiring of the protein. Computational techniques offer effective and inexpensive mapping of the allosteric sites on a protein. Here, we discuss methods to map and regulate allosteric communications in proteins, and strategies to establish control over kinase functions in live cells and organisms. Protein molecules, or “sensors” are engineered to function as tools to control allosteric activity of the protein as these sensors have high spatiotemporal resolution and help in understanding cell phenotypes after immediate activation or inactivation of a kinase. Traditional methods used to study protein functions, such as knockout, knockdown, or mutation, cannot offer a sufficiently high spatiotemporal resolution. We discuss the modern repertoire of tools to regulate protein kinases as we enter a new era in deciphering cellular signaling and developing novel approaches to treat diseases associated with signal dysregulation.

eLife assessment

One of the most promising strategies in development of drugs targeting kinases is provided by using allosteric control that allows specific regulation and study of kinase function without directly targeting the active site. This **important** manuscript provides a **convincing** review of the current repertoire of tools for regulating the activity of protein kinases with the ultimate goal of developing novel approaches in treating diseases associated with signal dysregulation.

Introduction

Protein kinases expedite the cell signaling processes by facilitating the transfer of a phosphate from the nucleotide triphosphate to the target protein. Kinase-mediated phosphorylation is critical for a wide spectrum of cell signaling processes such as transcription, translation, cell proliferation, receptor downregulation, metabolism, and apoptosis¹. The dysregulation of kinases alters cell function and may lead to diseases such as schizophrenia², Alzheimer's disease³, cancer⁴, and cardiovascular diseases⁵. Kinases undergo dynamic conformational changes to elicit their function during cell signaling. All kinases have a conserved residue sequence that forms the nucleotide-binding domain to bind to adenosine triphosphate (ATP) or guanosine triphosphate (GTP)^{1,6}. The binding of the nucleotides is facilitated by phosphorylation of the activation loop⁷. Therapies targeting kinases focus on the nucleotide-binding sites (active/ orthosteric sites) that regulate the kinase function. Therapies targeting orthosteric sites to regulate kinase activity compete with the native ligand (ATP or GTP) to bind to the active site and need to be highly selective for the active site. Due to the conservation of the nucleotide-binding domain across all protein kinases, specifically targeting the active/ orthosteric sites of one type of kinase is not possible.

Kinase activity is also regulated by autoinhibitory domains^{8,9}. Autoinhibitory domains suppress the interaction between the kinases and the effector molecule (nucleotides). The autoinhibitory domains are the sites distal to the active sites that affect the activity of the kinase at the active site through the induction of conformational and dynamic changes in the protein structure^{10–12}. Allosteric regulation does not involve direct competition with the native ligand. Instead, it perturbs active site structure or dynamics. The kinases, especially in diseased conditions, may develop resistance to orthosterically regulated therapies by incorporating mutations at the active site, thus enabling kinases to elicit catalytic activity while disallowing the binding of the orthosteric regulators. Allosteric control of kinase function allows the bypass of this resistance.¹¹

Finding allosteric sites that effectively regulate active sites is critical for establishing allosteric control over a particular kinase. X-ray crystallography¹³ and nuclear magnetic resonance (NMR) spectroscopy¹⁴ techniques are widely used to observe the structures of a protein. The changes occurring in the structure of protein molecules during the allosteric binding of effector molecules can be mapped by comparing structures at different points in time. However, these techniques are expensive and time-consuming. Modern computational approaches offer an inexpensive and fast assessment of protein allosteric “wiring”. However, the computational approaches only offer hypothetical predictions and still need to be tested experimentally. Experimental validation does not require the structural characterization of the protein, instead, a direct test of the predictions in activity assays demonstrates the functioning of the established allosteric control¹⁵. Here, we discuss computational and experimental approaches to map allosteric communications, and tools used to control protein function through allosteric regulation.

Allosteric regulation of protein kinases

Kinases are considered to be biological switches due to their ability to change structural conformation upon activation.¹⁶ Since kinases make up 2% of the protein-coding genome¹⁷ in humans and are implicated in multiple diseases this family of proteins is an important therapeutic target. Protein kinases phosphorylate serines, threonines, or tyrosines by transferring the γ -phosphate of ATP to the hydroxyl group causing structural changes in the protein, eliciting its function as an enzyme.¹⁸ Due to the widespread role of kinases in nature, they are considered to be desirable targets for studying cellular signaling. In eukaryotic cells, across all different types of protein kinases there are close to 250 residues worth of conserved sequences.¹⁶ One conserved

region is attributed to be the nucleotide-binding domain (NBD) which is a cleft that forms the binding pocket for ATP and is located between the N-terminal lobe and the C-terminal lobe of the protein. The flexibility of the protein is maintained by two ensembles of hydrophobic residues or “spines” that connect the α -helix of the C lobe to the β -sheet of the N lobe.¹⁹ These spines are called the “regulatory spine (R spine)”, due to their ability to regulate the assembly and disassembly of the active kinase, and the “catalytic spine (C spine)”, since these residues balance the catalytic activity of the kinases upon ATP binding. The changes in structure of the kinases due to their complex structure-activity relationship are likely to be resolved due to the presence of the spines.²⁰ The dynamic structure of kinases has proved to be of great importance for the functionality of the protein and the NMR structures confirming these dynamic structural changes have underscored the presence and importance of allostery of protein kinases.²¹ (**Figure 1**).

Several approaches have been developed using the pre-existing tools to control protein function for the allosteric regulation of protein kinases. Kinase activity can be controlled by ligands, Zorba et al.²² made use of monobodies²³ (a type of synthetic binding protein) to control allostery in Aurora A (AurA).²⁴ AurA is a protein kinase responsible for the maturation of the centrosome and duplication through its effect on the microtubule spindle formation in mitosis.²⁵ AurA serves to be a major oncogene since its overexpression has been associated with a variety of cancers. The authors used monobodies²⁶ to bind to the allosteric sites of AurA and produce complete inhibition or complete activation without having to actively compete with ATP to bind to the NBD. The effect of the binding of monobodies to AurA, and the activity of the protein, were mapped using X-ray crystallography. The binding of the monobody showed a 20-fold inhibition of the kinases proving its effectiveness as an allosteric inhibitor and demonstrating promising therapeutic potential. Wojcik et al.²⁷ designed high-affinity monobodies to bind to the kinase Bcr-Abl formed by the fusion of breakpoint cluster region (Bcr) and Abelson tyrosine kinase 1 (Abl1). Bcr-Abl activity leads to the development of chronic myelogenous leukemia.²⁸ Studies have shown an interaction between Src homology domain (SH2) and tyrosine kinase, critical to kinase activity and mutations arising at these interfaces (I164E), leads to inhibition of the kinase activity and oncogenesis induced by Bcr-Abl.²⁹ The authors developed monobodies for these critical residues to control the kinase activity of Bcr-Abl allosterically. The monobodies were developed using combinatorial synthesis and the binding affinity was tested using chemical shift perturbation assays that identify binding sites and predict the binding efficiency of ligands. Inhibition of interaction of the SH2 domain and the kinase domain due to monobodies decreased the activity of Bcr-Abl (validated by western blot and immunohistochemistry experiments). Monobodies can be used as a tool for controlling kinase activity, providing an alternative approach to chemogenetic and optogenetic control of protein function as monobodies do not require engineering of the target kinase. (**Figure 2**).

Dagliyan et al.³⁰ developed a rapamycin-binding domain that destabilized the NBD, thereby disrupting the kinase activity in the P21-activated kinase (PAK1). PAK1 is known to play a role in angiogenesis, neurogenesis, and cell motility.³¹ The dysregulation of this protein leads to the development of speech impairment, seizures, and macrocephaly. The authors developed a protein domain that is activated upon dopamine binding to the PAK1 domain. PAK1 is overexpressed in breast cancer and hence is identified as an oncogene.³² The authors tested this rapamycin-PAK construct (RapR-PAK1) in breast cancer cells to observe changes in the kinase activity due to the effect of rapamycin. The construct was successfully able to control the kinase activity of PAK, demonstrating the applicability of ligands to control protein activity allosterically by either inhibiting activity or inducing it.

Karginov et al.³³ used a similar strategy for regulating kinase activity, but instead of using a ligand-based approach, the authors used light as the input signal for controlling activity. The authors built a photoactivatable analog of rapamycin to regulate the dimerization of proteins. The photo-inducible small molecules are generated by the installation of “caging groups”^{34,35} that keep the molecule inactive until removed. Caging groups sequester the site on the ligand




Figure 1

Domains of kinase protein responsible for the regulation of function




Figure 2

A Allosteric activation of kinase (ERK2 kinase, PDB ID: 4GT3) by uniRapR (PDB ID: 7F2J) domain that binds to small molecule (rapamycin) causing activation of the protein. **B** Allosteric inhibition of kinase (ERK2 kinase, PDB ID: 4GT3) by insertion of an optogenetic control protein, LOV2 (PDB ID: 2V0W), causing conformational change upon irradiation by blue light. **C** Activation of kinase using monobodies (PDB ID: 3RZW).

necessary for activation. Dissociation of the caging groups from the activation site through irradiation by light leads to activation of the small molecules. The authors used “caged-rapamycin (pRap)” to regulate both the kinase activity and protein dimerization.

Zhou et al.³⁶ developed an optically-inducible, single-chain protein that could undergo conformational changes upon exposure to light. The authors induced modifications to DronpaN145, a photo-inducible green fluorescent protein to form pdDronpa1³⁷ that could dimerize at the kinase active site in the presence of 500 nm cyan and stymie kinase activity. Upon removal of light, the kinase activity was restored. The authors used dual specificity mitogen-activated protein kinase 1 (MEK1) to test the pdDronpa1 construct. The authors were able to successfully control the kinase activity of MEK1 providing a unique and versatile approach toward optogenetics.

Vishweshwaraiah et al.³⁸ developed a protein-based nanocomputing agent³⁹ containing two inputs that had opposing effects on the kinase activity, forming a logic OR gate to regulate allosteric communications. The authors inserted a chemogenetic (uniRapR⁴⁰) and an optogenetic domain (LOV2⁴¹) within FAK. The uniRapR construct was inserted in the allosteric site of the kinase domain, predicted using Ohm⁴², such that it could activate the protein upon interaction with rapamycin. This activation pathway of the protein upon interaction with rapamycin was predicted by DMD simulations. The LOV2 construct was inserted in loop 2 of the FERM domain (allosteric site on FERM domain was predicted using Ohm) leading to an inhibition of kinase activity by inactivation of the protein⁴³. Inactivation of the protein was predicted using DMD simulations in blue light on/off conditions. The authors observed a decrease in kinase activity upon exposure to blue light and an increase in kinase activity upon exposure to rapamycin upon live cell imaging (changes in phenotype were used to determine kinase activity). This successful study provides a basis for the development of rationally designed nano-computing agents and can be used in diagnostics, drug delivery, regulation of cell signaling, and metabolic flux. In order to demonstrate that the nanocomputing agents are effective in other proteins as well, Chen et al. developed a Src-uniRapR-LOV2 construct that provided a dual control on the function of Src kinase. The uniRapR construct was able to activate the Src kinase while the LOV2 construct rendered the protein inactive upon irradiation by blue light. The on/off protein switch was tested by visualization of the focal adhesion. With rapamycin as the first signal, followed by blue light as the second.⁴⁴ Along with computational methods such as MSA (multiple sequence alignments), MD (molecular dynamics), and DMD simulations (discrete molecular dynamics); molecular docking and *in silico* modeling⁴⁵ paired with mass spectroscopy have also been used to define protein structures and map allosteric sites in proteins. Through microscopic techniques, such as cryo-EM, two-dimensional structures of proteins can be acquired and morphed into three-dimensional models using artificial intelligence⁴⁶. In addition to small molecules, biologics such as antibodies or various input stimuli (such as pH, temperature, or pressure) can be made use of to regulate protein function by exploitation of allosteric communication. Protein allostery proves to be an essential tool in controlling protein function in translational research.

Methods to map allosteric communications

Computer simulations developed to predict allosteric communications offer faster and more cost-effective methods of studying allosteric communication in proteins compared to traditional experimental structural characterization techniques⁴⁷. Experimentally, NMR spectroscopy and X-ray crystallography are used in conjunction to map allostery, Boulton and Melacini⁴⁸ hypothesized that the crystal structures of proteins solved by these techniques would provide information about the changes in the conformation of proteins upon binding of an allosteric regulator. Sites that are susceptible to a conformational change upon binding of the modulator/regulator were termed “hot spots”. The high-resolution images from NMR spectroscopy provided the opportunity to study the conformational dynamics as a result of allostery. By combining the high-resolution images of the protein structure, and the molecular dynamics simulations, predictions can be made about the allosteric communications within proteins. Lisi

and Loria⁴⁹ described allostery using NMR studies paired with the computational predictions in enzyme catalysts; PTP1B (protein tyrosine phosphatase 1B)⁵⁰, which is known to be associated with obesity and diabetes, and PKA-C (protein kinase C)⁵¹, which is responsible for cell signaling and proliferation. Jain et al.⁵² used NMR spectroscopy to configure the active and inactive structures of Cdc42-LOV2 construct. Light oxygen voltage (LOV2) protein is used as an optogenetic tool since it is active under blue light and is used to control protein function allosterically. The authors elucidated the structural changes of the Cdc42-LOV2 occurring upon irradiation by blue light converting the construct from an inactive to an active state. This provides an insight on the allosteric communication within proteins through structural elucidation. NMR spectroscopy is able to map both structural and dynamic changes that occur within an allosteric system, but may not be sufficient to determine the structural and dynamic changes occurring in complex allosteric systems.⁴⁸

Computational methods provide a faster means of deciphering protein interactions. Gadiyaram et al.⁵³ developed a graph spectral analysis of protein structures to predict allostery using non-covalent interactions between amino acid residues to create a protein structure network. The authors put their theories to the test by predicting the side-chain interactions between residues that form the G-protein coupled receptors (GPCRs) (specifically the β 2-adrenergic receptor) and the HIV proteases. Another computational strategy, multiple sequence alignments or MSA⁵⁴ is the process of aligning two or more protein sequences to discover the conserved regions of the proteins. Lockless and Ranganathan⁵⁵ made use of MSA to map allosteric communications in a postsynaptic density protein, Drosophila disc large tumor suppressor, and zonula occludens-1 (PDZ) protein domain. They hypothesized that perturbations in one residue of the protein sequence would cause thermodynamic changes in the residue adjacent to it. If the protein sequences are conserved, similar changes in the vibrational frequencies should be seen in all proteins possessing the conserved residue sequence. The theory had two developmental parameters – conservation of the residues at a given site of the protein as observed in the MSA, and statistical coupling of two residues (nodes) of the given amino acid sequence. This method was validated by analyzing the conserved sequences of the PDZ and the Poxvirus and zinc finger (POZ) protein domains using statistical coupling analysis⁵⁶ and thermodynamic mutation cycles⁵⁷, which measures the strength of the interaction of the residues. Ambroggio and Kulhman⁵⁸ developed an algorithm to engineer proteins that can undergo structural protein-folding changes using MSA. The algorithm Ambroggio and Kulhman developed could predict stable structural conformations of both the changed and native protein instead of one protein conformation favoring the other. To test the algorithm, the authors made use of the Sw2 sequence, which could switch the conformations between the coiled coil and the zinc finger states. The information provided by MSA is susceptible to errors⁵⁹ as the “true alignment” of proteins is not possible limiting the accuracy of allosteric communication predictions using MSA.⁶⁰

The calculations necessary to describe protein structure flexibility and allostery of large molecular systems are computationally intensive. Molecular dynamic (MD) simulations help to overcome this limitation in predicting the interactions between amino acid residues. MD simulations use atomic interaction approximations to simulate atomic motions on the basis of Newtonian physics and reduce the computational complexity of this systems⁶¹. MD simulations are utilized to develop algorithms that predict allosteric communications. The algorithms not only help in predicting allosteric communication pathways within proteins but also help in determining the potential allosteric sites in the protein. Proctor et al.⁶² analyzed the allosteric communications in mutations of cystic fibrosis transmembrane conductance regulator protein (CFTR) and identified the allosteric hot spots using molecular dynamic simulations. These hot spots could be targeted to restore the transcription and translation of CFTR and provide an allosteric therapeutic approach. MD simulations can predict residue interactions within proteins at a high spatiotemporal resolution and provide a faster method of analysis of allosteric communication within a protein compared to experimental methods used for structure elucidation. The timescales play a role in predicting residue interactions since allosteric communications are dynamic in

nature. Short time scales (nanosecond) are not always accurately predicted by MD simulations,⁶³ this can be overcome using coarse-grained simulations (a type of multiscale modeling method that simulates complex behavior of system in a more realistic timescale) or discrete molecular dynamic (DMD) simulations that reach time scales in the range of microseconds.^{64,65} DMD simulations have been successfully used to predict allosteric communications in proteins.^{66,67}

Berezovsky et al.⁶⁸ developed a structure-based statistical mechanical model of allostery (SBSMMA). This model predicts allostery by quantifying the thermodynamic perturbations of proteins (upon ligand binding or mutations in the protein) by estimating the free energy of every residue present in the ligand-bound and ligand-free proteins using an allosteric signaling map (ASM). SBSMMA identifies potential exosites, protein regions that are likely to govern the allosteric regulation through reverse perturbations. AlloSigMA⁶⁹ is a server that provides a quantitative assessment of protein-energy dynamics by estimating the changes in the relative energy of the residues at the allosteric site. AlloMAPS⁷⁰ is a database that provides data on the predicted energy changes and the allosteric pathways of the molecule predicted using SBSMMA.⁶⁸ SBSMMA provides a way to discover allosteric sites on the protein and elucidate allosteric mechanisms within proteins. Tan et al.⁷¹ developed a protocol for designing allosteric modulators using a fragment-based approach based on the knowledge of allosteric interaction sites on a protein using SBSMMA. Fragment-based approach⁷² of drug design is the development of an entire small molecule by joining smaller, low molecular-mass fragments that show similar interactions with the target. This method was developed by Jencks⁷³ and was initially used for building small molecules for kinases. Fragment-based approach can be used to map protein-protein interactions, transcription factors and protein chaperons.⁷⁴

Treating the allosteric phenomenon as signal propagation through heterogeneous media, Wang et al.⁴² designed a physics-based computational approach for mapping allosteric communications called Ohm⁴². The multifaceted platform can be used to predict the allosteric sites along with the possible pathways, identify the critical residues, and predict the possible allosteric correlation between the residues. The platform is built on a perturbation propagation algorithm that measures the perturbation of residues by calculating the probability of interactions between the two residues (obtained by a probability matrix). The predictions were validated by predicting the allosteric communication during phosphorylation of CheY⁷⁵ (a protein found in the flagellar motor of bacteria) and the Caspase-1 protein⁷⁶ (involved in inflammatory and host-defense response). Allosteric communication within proteins can be predicted by a variety of computational methods that provide closely realistic predictions in a short span of time. These predictions can be validated using experimental techniques.

Tools to control allostery

Conceptually, a “sensor” domain is engineered into an allosteric site to establish allosteric control over a protein. This sensor domain responds to stimuli, external or internal, and undergoes conformational changes that propagate through allosteric paths to the active site. The active site then is altered upon the application of the stimuli. Some of the examples are where “sensor” proteins interact with small molecules or light and produce a conformational and dynamic change in the target protein leading to either activation or deactivation of the protein.

Optogenetic control. Photoreceptors are molecules that can change their conformation upon irradiation by light. The light-sensitive domains of the photoreceptors can be used as sensor domains to control protein function⁷⁷. The phytochrome B (PhyB)⁷⁸ is used extensively as a red light photoreceptor, since many optical systems do not have far-infrared light range lasers, red light photoreceptors have limitations to their usage.⁷⁸ Light Oxygen Voltage (LOV2) and Cryptochromes are the two main types of blue light photoreceptors used to control protein function. The (LOV2) domain of phototropin 1⁷⁹ obtained from *Avena sativa* has been shown to produce dramatic conformational changes upon exposure to blue light and is used to control

protein function. Zimmerman, et al.⁸⁰ developed LOV2 photoswitches and validated them by inserting them in various protein domains and determining the activity of the protein upon exposure to blue light. Dagliyan et al.⁸¹ developed a photo-inhibitable Src kinase domain by inserting LOV2 in the L1 loop connecting β -strands of Src kinase causing inhibition of the kinase upon activation of LOV2. However, insertion of the LOV2 domain into the protein without causing structural perturbations in the native protein structure is difficult due to limitations arising from the lack of availability of caging surfaces leading to non-specific activation of the photoreceptors. Caging surfaces or photocages are photosensitive groups that prevent the protein from reacting to small molecules or other proteins that might produce a biological function of the protein. The caging groups ensure activation of proteins only upon irradiation of the proteins by light of suitable wavelength⁸². To overcome this limitation, He et al.⁸³ developed a circular permutant of LOV2 (cpLOV2). The cpLOV2 is constructed by introduction of a glycine and serine-rich linker in the surface-exposed loop. The construct remains compatible with the existing LOV2-based engineered proteins with the advantage of having more caging surfaces thus improving its applicability (**Figure 2**).

Owing to the difficulty of optogenetic tools to control allosteric activity, arising due to the limitations in achieving subcellular control of activity⁸⁴, Shaaya et al.⁸⁵ developed an optogenetic tool, LightR, that enables subcellular control of protein and provides a tight temporal regulation of protein activity. LightR comprises of two tandemly connected Vivid photoreceptor⁸⁶ domains that form an antiparallel homodimer upon irradiation by blue light causing its activation. The use of LightR in controlling allosteric activity was validated in Src kinase, where irradiation with blue light caused activation of Src kinase. LightR provides better spatiotemporal resolution than LOV2 due to the subcellular control of proteins (**Figure 3**).

Chemogenetic control. Although optogenetics provides a high resolution of control of protein activity, optogenetic tools are often challenging to translate from cellular to animal models due to the limitation of the extent of absorption of light by animals that are non-transparent. To overcome this limitation, Dagliyan et al.⁴⁰ developed a protein switch (uniRapR) whose activity can be controlled by a small molecule (rapamycin). uniRapR is a complex of FK506-binding protein (FKBP12) and FKBP12-rapamycin binding protein⁸⁷ built into a single-chain protein. uniRapR binds selectively to rapamycin and upon binding induces a conformational change in the structure. uniRapR can be used to control protein function allosterically. To test the functionality of uniRapR, the authors inserted this construct into Src kinase and focal adhesion kinase (FAK) and compared the changes in phosphorylation of the two enzymes before and after the addition of rapamycin. The activity of uniRapR was further validated using a mutant strain of FAK with a point mutation that caused inhibition of FAK activity, the FAK activity was recovered by uniRapR which validated its use as an allosteric protein function regulator.

Guntas et al.⁸⁸ developed a maltose-binding allosteric regulator using *in vitro* recombination techniques. The wild-type beta-lactamase enzyme (BLA) activity is rendered unaffected by *E. coli* maltose binding protein. Upon recombination of the two genes, TEM-1-beta-lactamase and *E. coli* maltose binding gene⁸⁸, the enzyme activity was seen to be regulated upon the addition of maltose. The change in activity was analyzed by measuring the rate of β -lactam hydrolysis upon an increase in the concentration of maltose. Chemogenetic control of proteins has better *in vivo* resolution since light can only be absorbed up to a certain depth and has a shorter timescale of activation (**Figure 2**).

Conclusion

The human cell kinome consists of 538 kinases that are involved in a variety of cellular processes. Kinases have a conserved active site residue sequence making it challenging to target a specific kinase with ligands acting on the active site, hence, allosteric regulation kinases provides the




Figure 3

Sensor proteins used to control function of kinase protein allosterically. Chemogenetic tools (LightR, LOV2), optogenetic tool (UniRapR, Maltose binding protein), synthetic protein tool (Monobody).

ability to specifically target different types of kinases. Finding the allosteric sites on the proteins and visualizing the changes occurring in the protein conformation is experimentally time-consuming and an expensive endeavor. To overcome this limitation, computational methods are developed that can predict allosteric communication within proteins. Advancements in technology have paved the way to engineer proteins and regulate their activity. Various methods such as chemogenetics, optogenetics, and synthetic proteins are used as allosteric tools to control protein function. Kinases are an important family of proteins as they are implicated in a variety of diseases, hence allosteric regulation of kinases could be used as a potential therapeutic approach. Combining the various allosteric control tools help develop nanocomputing agents with more than one input function leading to a different output function of the protein, providing means of controlling different aspects of protein function. New tools can be developed that react to different inputs such as temperature, pH, and pressure along with the preexisting tools interacting with light or small molecules. These allosteric control sensor proteins provide a better spatiotemporal resolution in controlling protein function and prove to be a potential therapeutic approach targeting diseases that arise from the dysregulation of proteins.

Acknowledgements

We thank Dr. Richard Mailman, Brianna Hnath and Jiaying Chen for valuable discussions. We acknowledge support from the National Institutes for Health (R35 GM134864 to N.V.D.) and the Passan Foundation.

References

1. Cheng H.-C., Qi R. Z., Paudel H., Zhu H.-J (2011) **Regulation and Function of Protein Kinases and Phosphatases** *Enzyme Res* **2011**:1–3
2. Bentea E., et al. (2019) **Kinase network dysregulation in a human induced pluripotent stem cell model of DISC1 schizophrenia.** *Mol Omics* **15**:173–188
3. Ohno M (2014) **Roles of eIF2Î± kinases in the pathogenesis of Alzheimerâ€™s disease** *Front. Mol. Neurosci* **7**
4. Sarkar N., Singh A., Kumar P., Kaushik M (2023) **Protein kinases: Role of their dysregulation in carcinogenesis, identification and inhibition** *Drug Res* **73**:189–199
5. Chen J., et al. (2022) **Protein kinases in cardiovascular diseases** *Chin. Med. J. (Engl)* **135**:557–570
6. Taylor S. S., Kornev A. P (2011) **Protein kinases: evolution of dynamic regulatory proteins** *Trends Biochem. Sci* **36**:65–77
7. Adams J. A (2003) **Activation Loop Phosphorylation and Catalysis in Protein Kinases: Is There Functional Evidence for the Autoinhibitor Model?** *Biochemistry* **42**:601–607
8. Soderling T. R (1990) **Protein kinases. Regulation by autoinhibitory domains** *J. Biol. Chem* **265**:1823–1826
9. Li M., Rehman A. U., Liu Y., Chen K., Lu S (2021) **Dual roles of ATP-binding site in protein kinases: Orthosteric inhibition and allosteric regulation** *Advances in Protein Chemistry and Structural Biology* **124**:87–119
10. Nussinov R., Tsai C.-J (2013) **Allostery in Disease and in Drug Discovery** *Cell* **153**:293–305
11. Cui Q., Karplus M (2008) **Allostery and cooperativity revisited** *Protein Sci* **17**:1295–1307
12. Guo J., Zhou H.-X (2016) **Protein Allostery and Conformational Dynamics** *Chem. Rev* **116**:6503–6515
13. Keedy D. A (2019) **Journey to the center of the protein: allostery from multitemperature multiconformer X-ray crystallography** *Acta Crystallogr. Sect. Struct. Biol* **75**:123–137
14. Grutsch S., Br schweiler S., Tollinger M (2016) **NMR Methods to Study Dynamic Allostery** *PLOS Comput. Biol* **12**
15. Vishweshwaraiah Y. L., Chen J., Dokholyan N. V (2021) **Engineering an Allosteric Control of Protein Function** *J. Phys. Chem. B* **125**:1806–1814
16. Taylor S. S., Keshwani M. M., Steichen J. M., Kornev A. P (2012) **Evolution of the eukaryotic protein kinases as dynamic molecular switches** *Philos. Trans. R. Soc. B Biol. Sci* **367**:2517–2528
17. Manning G., Whyte D. B., Martinez R., Hunter T., Sudarsanam S (2002) **The Protein Kinase Complement of the Human Genome** *Science* **298**:1912–1934

18. Cohen P (2002) **The origins of protein phosphorylation** *Nat. Cell Biol* **4**:E127–E130
19. Johnson L. N., Lewis R. J (2001) **Structural Basis for Control by Phosphorylation** *Chem. Rev* **101**:2209–2242
20. Taylor S. S., Ilouz R., Zhang P., Kornev A. P (2012) **Assembly of allosteric macromolecular switches: lessons from PKA** *Nat. Rev. Mol. Cell Biol* **13**:646–658
21. Masterson L. R., Cembran A., Shi L., Veglia G (2012) **Allostery and Binding Cooperativity of the Catalytic Subunit of Protein Kinase A by NMR Spectroscopy and Molecular Dynamics Simulations** *Advances in Protein Chemistry and Structural Biology* **87**:363–389
22. Zorba A., et al. (2019) **Allosteric modulation of a human protein kinase with monobodies** *Proc. Natl. Acad. Sci* **116**:13937–13942
23. Sha F., Salzman G., Gupta A., Koide S (2017) **Monobodies and other synthetic binding proteins for expanding protein science: Monobodies and Other Synthetic Binding Proteins** *Protein Sci* **26**:910–924
24. Zhou H., et al. (1998) **Tumour amplified kinase STK15/BTAK induces centrosome amplification, aneuploidy and transformation** *Nat. Genet* **20**:189–193
25. Glover D. M., Leibowitz M. H., McLean D. A., Parry H (1995) **Mutations in aurora prevent centrosome separation leading to the formation of monopolar spindles** *Cell* **81**:95–105
26. Koide A., Wojcik J., Gilbreth R. N., Hoey R. J., Koide S (2012) **Teaching an Old Scaffold New Tricks: Monobodies Constructed Using Alternative Surfaces of the FN3 Scaffold** *J. Mol. Biol* **415**:393–405
27. Wojcik J., et al. (2016) **Allosteric Inhibition of Bcr-Abl Kinase by High Affinity Monobody Inhibitors Directed to the Src Homology 2 (SH2)-Kinase Interface** *J. Biol. Chem* **291**:8836–8847
28. Deininger M. W. N., Goldman J. M., Melo J. V (2000) **The molecular biology of chronic myeloid leukemia** *Blood* **96**:3343–3356
29. Grebien F., et al. (2011) **Targeting the SH2-Kinase Interface in Bcr-Abl Inhibits Leukemogenesis** *Cell* **147**:306–319
30. Dagliyan O., et al. (2017) **Engineering Pak1 Allosteric Switches** *ACS Synth. Biol* **6**:1257–1262
31. Hofmann C., Shepelev M., Chernoff J (2004) **The genetics of Pak** *J. Cell Sci* **117**:4343–4354
32. Radu M., Semenova G., Kosoff R., Chernoff J (2014) **PAK signalling during the development and progression of cancer** *Nat. Rev. Cancer* **14**:13–25
33. Karginov A. V., et al. (2011) **Light Regulation of Protein Dimerization and Kinase Activity in Living Cells Using Photocaged Rapamycin and Engineered FKBP** *J. Am. Chem. Soc* **133**:420–423
34. Riggsbee C. W., Deiters A (2010) **Recent advances in the photochemical control of protein function** *Trends Biotechnol* **28**:468–475

35. Mayer G., Heckel A (2006) **Biologically Active Molecules with a “Light Switch”** *Angew. Chem. Int. Ed* **45**:4900–4921
36. Zhou X. X., Fan L. Z., Li P., Shen K., Lin M. Z (2017) **Optical control of cell signaling by single-chain photoswitchable kinases** *Science* **355**:836–842
37. Zhou X. X., Chung H. K., Lam A. J., Lin M. Z (2012) **Optical Control of Protein Activity by Fluorescent Protein Domains** *Science* **338**:810–814
38. Vishweshwaraiah Y. L., Chen J., Chirasani V. R., Tabdanov E. D., Dokholyan N. V (2021) **Two-input protein logic gate for computation in living cells** *Nat. Commun* **12**
39. Dokholyan N. V (2021) **Nanoscale programming of cellular and physiological phenotypes: inorganic meets organic programming** *Npj Syst. Biol. Appl* **7**
40. Dagliyan O., et al. (2013) **Rational design of a ligand-controlled protein conformational switch** *Proc. Natl. Acad. Sci* **110**:6800–6804
41. Lee J., et al. (2008) **Surface Sites for Engineering Allosteric Control in Proteins** *Science* **322**:438–442
42. Wang J., et al. (2020) **Mapping allosteric communications within individual proteins** *Nat. Commun* **11**
43. Lietha D., et al. (2007) **Structural Basis for the Autoinhibition of Focal Adhesion Kinase** *Cell* **129**:1177–1187
44. Chen J., Vishweshwaraiah Y. L., Mailman R. B., Tabdanov E. D., Dokholyan N. V (2023) **A noncommutative combinatorial protein logic circuit controls cell orientation in nanoenvironments** *Sci. Adv* **9**
45. Yuan J., et al. (2022) **In Silico Prediction and Validation of CB2 Allosteric Binding Sites to Aid the Design of Allosteric Modulators** *Molecules* **27**
46. Chirigati F (2021) **Predicting protein structure from cryo-EM data** *Nat. Comput. Sci* **1**:96–96
47. Dokholyan N. V (2016) **Controlling Allosteric Networks in Proteins** *Chem. Rev* **116**:6463–6487
48. Boulton S., Melacini G (2016) **Advances in NMR Methods To Map Allosteric Sites: From Models to Translation** *Chem. Rev* **116**:6267–6304
49. Lisi G. P., Loria J. P (2017) **Allostery in enzyme catalysis** *Curr. Opin. Struct. Biol* **47**:123–130
50. Zhang Z.-Y (2017) **Drugging the Undruggable: Therapeutic Potential of Targeting Protein Tyrosine Phosphatases** *Acc. Chem. Res* **50**:122–129
51. Xiao Y., Liddle J. C., Pardi A., Ahn N. G (2015) **Dynamics of Protein Kinases: Insights from Nuclear Magnetic Resonance** *Acc. Chem. Res* **48**:1106–1114
52. Jain A., Dokholyan N. V., Lee A. L (2023) **Allosteric inactivation of an engineered optogenetic GTPase** *Proc. Natl. Acad. Sci* **120**

53. Gadiyaram V., Vishveshwara S., Vishveshwara S (2019) **From Quantum Chemistry to Networks in Biology: A Graph Spectral Approach to Protein Structure Analyses** *J. Chem. Inf. Model* **59**:1715–1727
54. Sofi M. Y., Shafi A., Masoodi K. Z (2022) **Multiple sequence alignment** *Bioinformatics for Everyone* :47–53 <https://doi.org/10.1016/B978-0-323-91128-3.00011-2>
55. Lockless S. W., Ranganathan R (1999) **Evolutionarily Conserved Pathways of Energetic Connectivity in Protein Families** *Science* **286**:295–299
56. Teşileanu T., Colwell L. J., Leibler S (2015) **Protein Sectors: Statistical Coupling Analysis versus Conservation** *PLOS Comput. Biol* **11**
57. Pagano L., et al. (2021) **Double Mutant Cycles as a Tool to Address Folding, Binding, and Allostery** *Int. J. Mol. Sci* **22**
58. Ambroggio X. I., Kuhlman B (2006) **Computational Design of a Single Amino Acid Sequence that Can Switch between Two Distinct Protein Folds** *J. Am. Chem. Soc* **128**:1154–1161
59. Kemena C., Notredame C (2009) **Upcoming challenges for multiple sequence alignment methods in the high-throughput era** *Bioinforma. Oxf. Engl* **25**:2455–2465
60. Nuin P. A., Wang Z., Tillier E. R (2006) **The accuracy of several multiple sequence alignment programs for proteins** *BMC Bioinformatics* **7**
61. Durrant J. D., McCammon J. A (2011) **Molecular dynamics simulations and drug discovery** *BMC Biol* **9**
62. Proctor E. A., et al. (2015) **Rational coupled dynamics network manipulation rescues disease-relevant mutant cystic fibrosis transmembrane conductance regulator** *Chem. Sci* **6**:1237–1246
63. Hertig S., Latorraca N. R., Dror R. O (2016) **Revealing Atomic-Level Mechanisms of Protein Allostery with Molecular Dynamics Simulations** *PLOS Comput. Biol* **12**
64. Shirvanyants D., Ding F., Tsao D., Ramachandran S., Dokholyan N. V (2012) **Discrete Molecular Dynamics: An Efficient And Versatile Simulation Method For Fine Protein Characterization** *J. Phys. Chem. B* **116**:8375–8382
65. Proctor E. A., Ding F., Dokholyan N. V. (2011) **Discrete molecular dynamics** *WIREs Comput. Mol. Sci* **1**:80–92
66. Bowerman S., Wereszczynski J (2016) **Detecting Allosteric Networks Using Molecular Dynamics Simulation** *Methods Enzymol* **578**:429–447
67. Dokholyan N. V., Buldyrev S. V., Stanley H. E., Shakhnovich E. I (1998) **Discrete molecular dynamics studies of the folding of a protein-like model** *Fold. Des* **3**:577–587
68. Tee W.-V., Tan Z. W., Lee K., Guarnera E., Berezovsky I. N (2021) **Exploring the Allosteric Territory of Protein Function** *J. Phys. Chem. B* **125**:3763–3780
69. Guarnera E., Tan Z. W., Zheng Z., Berezovsky I. N (2017) **AlloSigMA: allosteric signaling and mutation analysis server** *Bioinformatics* **33**:3996–3998

70. Tan Z. W., Tee W.-V., Guarnera E., Booth L., Berezovsky I. N (2019) **AlloMAPS: allosteric mutation analysis and polymorphism of signaling database** *Nucleic Acids Res* **47**:D265–D270
71. Wah Tan Z., Tee W.-V., Berezovsky I. N (2022) **Learning About Allosteric Drugs and Ways to Design Them** *J. Mol. Biol* **434**
72. Scott D. E., Coyne A. G., Hudson S. A., Abell C (2012) **Fragment-Based Approaches in Drug Discovery and Chemical Biology** *Biochemistry* **51**:4990–5003
73. Jencks W. P (1981) **On the attribution and additivity of binding energies** *Proc. Natl. Acad. Sci* **78**:4046–4050
74. Chen L., Cressina E., Leeper F. J., Smith A. G., Abell C (2010) **A Fragment-Based Approach to Identifying Ligands for Riboswitches** *ACS Chem. Biol* **5**:355–358
75. Lee S.-Y., et al. (2001) **Crystal Structure of Activated CheY** *J. Biol. Chem* **276**:16425–16431
76. Clark A. C (2016) **Caspase Allostery and Conformational Selection** *Chem. Rev* **116**:6666–6706
77. Padmanabhan S., Pérez-Castaño R., Osete-Alcaraz L., Polanco M. C., Elías-Arnanz M (2022) **Vitamin B12 photoreceptors** *Vitamins and Hormones* **119**:149–184
78. Ni M., Tepperman J. M., Quail P. H (1999) **Binding of phytochrome B to its nuclear signalling partner PIF3 is reversibly induced by light** *Nature* **400**:781–784
79. Halavaty A. S., Moffat K (2007) **N- and C-Terminal Flanking Regions Modulate Light-Induced Signal Transduction in the LOV2 Domain of the Blue Light Sensor Phototropin 1 from *Avena sativa*** *Biochemistry* **46**:14001–14009
80. Zimmerman S. P., Kuhlman B., Yumerefendi H (2016) **Engineering and Application of LOV2-Based Photoswitches** *Methods in Enzymology* **580**:169–190
81. Dagliyan O., et al. (2016) **Engineering extrinsic disorder to control protein activity in living cells** *Science* **354**:1441–1444
82. Bojtár M., et al. (2020) **Conditionally Activatable Visible-Light Photocages** *J. Am. Chem. Soc* **142**:15164–15171
83. He L., et al. (2021) **Circularly permuted LOV2 as a modular photoswitch for optogenetic engineering** *Nat. Chem. Biol* **17**:915–923
84. Wang J., et al. (2019) **Time-resolved protein activation by proximal decaging in living systems** *Nature* **569**:509–513
85. Shaaya M., et al. (2020) **Light-regulated allosteric switch enables temporal and subcellular control of enzyme activity** *eLife* **9**
86. Nihongaki Y., Suzuki H., Kawano F., Sato M (2014) **Genetically Engineered Photoinducible Homodimerization System with Improved Dimer-Forming Efficiency** *ACS Chem. Biol* **9**:617–621
87. Choi J., Chen J., Schreiber S. L., Clardy J (1996) **Structure of the FKBP12-Rapamycin Complex Interacting with Binding Domain of Human FRAP** *Science* **273**:239–242

88. Guntas G., Ostermeier M (2004) **Creation of an Allosteric Enzyme by Domain Insertion** *J. Mol. Biol* **336**:263–273

Article and author information

Shivani Godbole

Department of Pharmacology, Penn State College of Medicine, Hershey, PA 17033-0850, USA

Nikolay V. Dokholyan

Department of Pharmacology, Penn State College of Medicine, Hershey, PA 17033-0850, USA, Department of Biomedical Engineering, Penn State University, University Park, PA 16802, USA, Department of Engineering Science and Mechanics, Penn State University, University Park, PA 16802, USA, Department of Biochemistry & Molecular Biology, Penn State College of Medicine, Hershey, PA 17033-0850, USA, Department of Chemistry, Penn State University, University Park, PA 16802, USA

For correspondence: dokh@psu.edu

ORCID iD: [0000-0002-8225-4025](https://orcid.org/0000-0002-8225-4025)

Copyright

© 2023, Shivani Godbole & Nikolay V. Dokholyan

This article is distributed under the terms of the [Creative Commons Attribution License \(https://creativecommons.org/licenses/by/4.0/\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Volker Dötsch

Goethe University, Germany

Senior Editor

Volker Dötsch

Goethe University, Germany

Joint Public Review:

This concise review provides a clear and instructive picture of the state-of-the-art understanding of protein kinases' activity and sets of approaches and tools to analyse and regulate it.

Three major parts of the work include: methods to map allosteric communications, tools to control allostery, and allosteric regulation of protein kinases. The work provides an important and timely view of the current status of our understanding of the function of protein kinases and state-of-the-art methods to study its allosteric regulation and to develop allosteric approaches to control it.

Author Response

The following is the authors' response to the original reviews.

We thank the reviewers for their generous comments on the manuscript and have made edits to address their concerns. The manuscript has been restructured and the reference (PMID: 35738428) has been added to the review. We addressed the reviewer's comment below.

Reviewer #1 (Recommendations For The Authors):

Regarding SBSMMA, the authors may complement their discussion by mentioning recent work (PMID: 35738428) where SBSMMA was used to exemplify a potential fragment-based design approach for developing allosteric effectors for kinases.

Thank you for the suggestion, we have added a short summary of the work where SBSMMA is used as a basis for developing small molecules to target kinases using fragment-based design approach.