

Affinity-tagged SMAD1 and SMAD5 mouse lines reveal transcriptional reprogramming mechanisms during early pregnancy

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

February 14, 2024 (this version)

Reviewed preprint version 1

November 22, 2023

Posted to preprint server

September 25, 2023

Sent for peer review

September 11, 2023

Zian Liao, Suni Tang, Kaori Nozawa, Keisuke Shimada, Masahito Ikawa, Diana Monsivais ,
Martin M. Matzuk 

Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA • Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, 77030, USA • Graduate Program of Genetics and Genomics, Baylor College of Medicine, Houston, TX, 77030, USA • Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA • Research Institute for Microbial Diseases, Osaka University, Osaka, 565-0871, Japan

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Endometrial decidualization, a prerequisite for successful pregnancies, relies on transcriptional reprogramming driven by progesterone receptor (PR) and bone morphogenetic protein (BMP)-SMAD1/SMAD5 signaling pathways. Despite their critical roles in early pregnancy, how these pathways intersect in reprogramming the endometrium into a receptive state remains unclear. To define how SMAD1 and/or SMAD5 integrate BMP signaling in the uterus during early pregnancy, we generated two novel transgenic mouse lines with affinity tags inserted into the endogenous SMAD1 and SMAD5 loci (*Smad1^{HA/HA}* and *Smad5^{PA/PA}*). By profiling the genome-wide distribution of SMAD1, SMAD5, and PR in the mouse uterus, we demonstrated the unique and shared roles of SMAD1 and SMAD5 during the window of implantation. We also showed the presence of a conserved SMAD1, SMAD5, and PR genomic binding signature in the uterus during early pregnancy. To functionally characterize the translational aspects of our findings, we demonstrated that SMAD1/5 knockdown in human endometrial stromal cells suppressed expressions of canonical decidual markers (*IGFBP1*, *PRL*, *FOXO1*) and PR-responsive genes (*RORB*, *KLF15*). Here, our studies provide novel tools to study BMP signaling pathways and highlight the fundamental roles of SMAD1/5 in mediating both BMP signaling pathways and the transcriptional response to progesterone (P4) during early pregnancy.

eLife assessment

This study presents two **useful** new mouse models that individually tag proteins from the SMAD family to identify distinct roles during early pregnancy. **Solid** evidence is provided that SMAD1 and SMAD5 target many of the same genomic regions as each other and the progesterone receptor. Given the broad effect of these signaling pathways in multiple systems, these new tools will most likely interest readers across biological disciplines.

Introduction

Infertility is an emerging health issue that affects approximately 15% of couples¹. One in five women aged 15 to 49 years old with no prior births suffers from infertility in the United States². One important factor affecting fertility is failed embryo implantation or subsequent post-implantation loss due to endometrial defects. This is evident from the high number of failed pregnancies, with as many as 15% of pregnancies resulting in early pregnancy losses³. Understanding the molecular mechanism of how the maternal endometrium becomes suitable for embryo implantation and eventual decidualization, will be the key to eradicating global concerns related to infertility and early pregnancy losses.

The transforming growth factor β (TGF β) family plays diverse roles in development, physiology, and pathophysiology^{4,5}, and in particular, signaling pathways downstream of the bone morphogenetic protein (BMP) subfamily are essential for decidual formation^{6,7}. There are more than 30 TGF β family ligands, and these ligands signal through complexes of transmembrane type 1 Activin-Like Kinases (ALK) receptors (ALK1, ALK2, ALK3, ALK6) and transmembrane type 2 receptors (BMPR2, ACVR2A, ACVR2B) and then phosphorylate downstream SMAD1 and SMAD5 proteins. Phosphorylated SMAD1/5 form heteromeric complexes with SMAD4 and translocate to the nucleus to induce specific transcriptional programs. Our laboratory and others have used genetically engineered mouse models with deletions of ligands, receptors, and downstream effectors of BMP signaling pathways to establish that BMP signaling pathways are major regulators of early pregnancy^{6,12}.

A successful pregnancy begins with reciprocal crosstalk between the maternal endometrium and the new blastocyst during the peri-implantation window. Effective implantation requires precise synchronization between the development of the blastocyst and the transformation of the maternal endometrium into a functional decidua. Endometrial stromal fibroblasts undergo the decidualization process in which they differentiate into unique secretory decidual cells that offer a supportive and immune-privileged microenvironment required for embryo implantation and placental development. Decidualizing stromal cells can react to individual embryos in a way that either supports the implantation and subsequent embryonic development or exerts early rejection^{13,14}. Aberrant decidualization processes are observed in patients with recurrent pregnancy loss (RPL), displaying a disordered pro-inflammatory response, decreased induction of decidual marker genes, and abnormal responses to embryonic human chorionic gonadotropin^{13,15,16}. In addition to affecting early pregnancy outcomes, defective decidualization is also involved in the maternal etiology of preeclampsia, causing abnormal placental phenotype^{17,18}. The process of decidualization is tightly regulated by hormone signaling pathways (estrogen, E2, and progesterone, P4), as well as by BMP signaling pathways. Our recent studies found that endometrial *Smad1* deletion had no significant effect on fertility, *Smad5* conditional deletion resulted in subfertility, while double *Smad1/5* conditional deletion led to infertility due to implantation and decidualization defects⁹. The uteri of mice with double

conditional *Smad1/5* deletion also displayed decreased response to P4 during the window of implantation, suggesting synergy between the two pathways. However, the mechanistic genomic actions of SMAD1 and/or SMAD5 in the uterus have not been explored, partly because there are no specific antibodies that distinguish phospho-SMAD1 versus phospho-SMAD5.

In this study, we define how SMAD1/5 instructs the decidualization process using genomic approaches in newly generated transgenic mouse lines. We inspect the potential crosstalk between P4 and BMP signaling pathways mediated by SMAD1/5. Together, our study demonstrates that SMAD1 and SMAD5 exhibit shared and unique genomic binding features and further reveals that SMAD1/5 contributes to the P4 response through transcriptional reprogramming during decidualization.

Materials and methods

Generation of Knock-in Mouse Lines

Smad5^{PA/PA} knock-in (KI) mice were generated using a similar approach as previously described¹⁹. Briefly, single-guide RNA (sg-RNA) was designed to target the regions close to the start codon (Figure 1A,B) and the sgRNA sequence was inserted into the pX459 V2.0 plasmid (#62988, Addgene). The reference plasmids containing PA tag sequence were constructed in pBluescript II SK (+) vector (Agilent, Palo Alto, CA, USA). One µg of guide RNA inserted vector and 1.0 µg of reference plasmid were co-transfected into EGRG01 embryonic stem (ES) cells. Twelve ES clones out of 48 had the expected knock-in allele. ES cell clones that possessed the proper KI allele were injected into ICR embryos and chimeric blastocysts were transferred into pseudopregnant females. Chimeric male mice were mated with B6D2F1 female mice to obtain the PA-tagged SMAD5 KI heterozygous mice. Homozygous *Smad5*^{PA/PA} mice were maintained in the C57BL/6 J × 129S5/SvEvBrd mixed genetic background. To generate *Smad1*^{HA/HA} mice, Cas9 protein (Thermo Fisher, A36497), sg-RNA, and a repair oligo of homology-directed repair (HDR) containing HA-tag and linker sequences were electroporated into zygotes harvested from *in vitro* fertilization using B6D2F1 male and female mice. An ECM830 electroporation system (BTX, Holliston, MA) was used for electroporation. Subsequently, embryos were cultured overnight to the 2-cell stage and then transferred to the oviducts of pseudopregnant CD-1 mice (Center for Comparative Medicine, Baylor College of Medicine). Pups were further screened for successful heterozygous or homozygous knock-in alleles by PCR using primers spanning across the HA tag. Sequences of sgRNA, the single-stranded repair oligo for HDR and primer used for genotype are listed in Supplementary Table 1.

Animal ethics compliance and tissue collection

All mice were housed under standard conditions of a 12-hour light/dark cycle in a vivarium with controlled ambient temperature (70°F ± 2°F and 20-70% relative humidity). All mouse handling and experimental procedures were performed under protocols approved by the Institutional Animal Care and Use Committee of Baylor College of Medicine. All experiments were performed with female mice aged between 7 to 12 weeks with a C57BL/6 J × 129S5/SvEvBrd mixed genetic background. All mice were euthanized using isoflurane induction followed by cervical dislocation, and tissues were snap-frozen in liquid nitrogen.

Cleavage Under Targets and Release Using Nuclease (CUT&RUN) Approach

Nuclei from uterine tissues were purified following a previously published protocol²⁰. The experiments were performed using pooled biological replicates from two mice that were processed as technical replicates throughout the CUT&RUN procedure and analysis. In short, uteri




Figure 1

Mouse models with global HA tagged SMAD1 and PA tagged SMAD5 proteins.

A-B) Schematic approaches for generating *Smad1^{HA/HA}* and *Smad5^{PA/PA}* knock-in mouse lines. Sanger sequencing of the genotyping results are included as validation of knock-in sequence. Black and blue boxes indicate untranslated and coding regions, respectively. **C-D)** Immunoblot (IB) analysis of the immunoprecipitation (IP) of HA tagged SMAD1 and PA tagged SMAD5 proteins from different tissues of the tagged mouse lines. Wild type (WT) mice were used as negative controls. Antibodies used for IB and IP are as labeled. Targeted bands of SMAD1 and SMAD5 are indicated by red arrows.

were harvested from pregnant mice at 4.5 days post coitus and washed with cold swelling buffer (10 mM Tris-HCl pH 7.5, 2 mM MgCl₂, 3 mM CaCl₂, 1X Protease Inhibitor Cocktail (PIC, Roche, 11836170001)) immediately after collection. Then tissue was cut into small pieces (~2-3mm) using scissors, while submerged in cold swelling buffer. Nuclear extract was prepared by dounce homogenization in cold swelling buffer (using a size 7 dounce) and filtered using the cell strainer (100 µm, BD Biosciences). Lysate was centrifuged at 400 g for 10 min, then resuspended in lysis buffer (swelling buffer with 10% glycerol and 1% CA-630, 1X PIC) using end-cut or wide-bore tips and incubated on ice for 5 min. Nuclei were washed twice with lysis buffer and resuspended in lysis buffer. Next, CUT&RUN procedure largely follows a previous protocol²¹. Briefly, around 500,000 nuclei were used per reaction. 10 µl of concanavalin-coated beads (Bangs Labs, BP531) were washed twice in Bead Activation Buffer (20 mM HEPES, pH 7.9, 10 mM KCl, 1 mM CaCl₂, 1 mM MnCl₂) for each reaction. Then, beads were added to nuclei resuspension and incubated for 10 mins at room temperature. After incubation, bead-nuclei complexes were resuspended in 100 µl Antibody Buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 0.5 mM Spermidine, 1X PIC, 0.01% digitonin, and 2mM EDTA) per reaction. One µg of IgG antibody (Sigma, I5006), HA antibody (EpiCypher, 13-2010), PA antibody (Fuji Film, NZ-1) and PR antibody (Cell Signaling, D8Q2J) were added to each group respectively. After overnight incubation at 4 °C, bead-nuclei complexes were washed twice with 200 µl cold Dig-Wash buffer (20 mM HEPES pH=7.5, 150 mM NaCl, 0.5 mM Spermidine, 1X PIC, 0.01% digitonin) and resuspended in 50 µl cold Dig-Wash buffer with 1 µl pAG-MNase (EpiCypher, 15-1016) per reaction. After incubation at room temperature for 10 min, bead-nuclei complexes were washed twice with 200 µl cold Dig-Wash buffer and resuspended in 50 µl cold Dig-Wash buffer, then 1 µl 100 mM CaCl₂ was added to each reaction. The mixture was incubated at 4 °C for 2 hours and the reaction was stopped by adding 50 µl Stop Buffer (340mM NaCl, 20 mM EDTA, 4 mM EGTA, 0.05% Digitonin, 100 ug/mL RNase A, 50 mg/mL glycogen, 0.5 ng E. coli DNA Spike-in (EpiCypher, 18-1401)) and incubate at 37 °C for 10 min. The supernatant was collected and subjected to DNA purification with phenol-chloroform and ethanol precipitation. Sequencing libraries were prepared using NEBNext Ultra II DNA Library Prep Kit (New England BioLabs, E7645) following manufacture's protocol. Paired-end 150 bp sequencing was performed on a NEXTSeq550 (Illumina) platform.

Bioinformatic Analysis for CUT&RUN data and re-analysis of published single-cell RNA seq data

For CUT&Run data, raw data were de-multiplexed by bcl2fastq v2.20 with fastqc for quality control. Clean reads were mapped to reference genome mm10 by Bowtie2, with parameters of --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700. For Spike-in mapping, reads were mapped to E. coli genome U00096.3. Duplicated reads were removed, and only uniquely mapped reads were kept. Spike-in normalization was achieved through multiply primary genome coverage by scale factor (100000 / fragments mapped to E. coli genome). CUT&RUN peaks were called by SECAR²² with the parameters of -norm -stringent -output. Track visualization was done by bedGraphToBigWig²³, bigwig files were imported to Integrative Genomics Viewer for visualization. For peak annotation, common peaks were identified with 'mergePeaks' function in HOMER v4.11²⁴ and then genomic annotation was added by ChIPseeker²⁵. Motif analysis was conducted through HOMER v4.11 with parameter set as findMotifsGenome.pl mm10 -size 200 -mask²⁴. Peak heatmaps were plotted using deepTools 2.4.2 with clustering options set to k-means²⁶. For single-cell RNA seq, raw data was obtained from EMBL-EBI under accession No. E-MTAB-10287. Cells with low coverage (less than 500 genes detected) were filtered, then gene counts were normalized for each cell by converting counts to quantiles and obtaining the corresponding values from a normal distribution. Then normalized cell vectors are concatenated along the gene panel. Plot visualization was conducted through CELLXGENE platform.²⁷

Western Blot Analysis of Immunoprecipitation (IP-WB)

Tissues were pulverized in liquid nitrogen and then lysed using NETN buffer (20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.5 mM EDTA, 10% Glycerol, and 0.5% NP-40). Protein concentration was determined by BCA Protein Assay Kit (Thermo Fisher, 23225). 1.5 mg of total protein lysate was used for IP. IP was performed by adding HA antibody (Cell Signaling, C29F4) or PA antibody (Fuji Film, NZ-1) to the lysate and incubate for 1 hour at 4 °C. Subsequently, protein G magnetic beads (Thermo Fisher, 88847) were added for an additional 1 hour at 4 °C. Then, the beads were washed five times with NETN buffer, and denatured in sample buffer (Thermo Fisher, NP0007) for further analysis by Western blot. For western blot procedures, briefly, denatured protein lysates were run on the 4 to 12%, Bis-Tris protein gels (Thermo Fisher, NP0321BOX) followed by electrophoretic transfer to nitrocellulose membrane. The membrane then went through blocking by 5% milk in Tris-buffered saline with Tween20 (TBST), followed by incubation overnight at 4 °C in the primary antibodies anti-HA (Cell Signaling, C29F4), anti-PA (Fuji Film, NZ-1), anti-SMAD1 (Life technologies, 385400) and anti-SMAD5 (ProteinTech, 12167-1-AP) at 1:1,000 dilution. The next day, membranes were washed three times with TBST, then incubated with horseradish peroxidase-conjugated secondary antibody for 1 h at room temperature, then washed three times with TBST, developed and imaged on iBright Imaging System (FL1500).

Primary endometrial stromal cells isolation/RNAi/decidualization

Studies using human specimens were conducted as indicated in a protocol approved by the Institutional Review Board at Baylor College of Medicine, H-51900. Human endometrial stromal cells were collected from healthy volunteers' menstrual effluent as previously reported^{28,30} (N=3). In brief, samples were collected by participants in a DIVA cup during the 4-8 hours on the first night of menses and stored in DMEM/F12 with 10% FBS, antibiotic/antimycotic and 100 µg/ml Primocin in a cold insulated pack until processing in the laboratory on the day of collection. The effluent was digested with 5 mg/ml collagenase and 0.2 mg/ml DNase I for 20 min at 37 °C, then the cell pellet was collected by centrifuging at 2,500 rpm for 5 min at room temperature. Next, red blood cell lysis was performed by resuspending the cell pellet in 20 ml of 0.2% NaCl for 20 seconds and neutralized with 20 ml of 1.6% NaCl. Then the solution was then centrifuged at 2,500 rpm for 5 min. Five ml complete medium (DMEM/F12 supplement with 10% FBS, 1X Antibiotic-Antimycotic + 100 µg/ml Primocin) was used to resuspend the pellet and the solution was passed through 100 µm and 20 µm cell strainer sequentially. The flowthrough containing the stromal cells was centrifuged for 5 min at 2,500 rpm and the pellet was resuspended in 10 ml complete medium and plated in a 10 cm dish. siRNA knockdown was performed using Lipofectamine RNAiMAX following the manufacturer's protocol. In brief, 0.2 million stromal cells were plated in a 12-well plate one day before transfection. On the day of transfection, 2 µl siRNA (20 µM, Dharmacon, D-001810-10, L-012723-00-0005, L-015791-00-0005) and 3 µl Lipofectamine RNAiMAX were diluted in 50 µl Opti-MEM respectively and then mixed to incubate at room temperature for 15 min. Then, the complex was added dropwise onto the cells. 24 h after transfection, medium was changed to DMEM/F12 supplement with 2% charcoal stripped FBS. Decidualization was induced by the addition of 35 nM estradiol (Sigma, E1024), 1 µM medroxyprogesterone (Sigma, 1378001), and 0.05 mM cyclic adenosine monophosphate (Axxora, JBS-NU-1502L) for 4 days with media changes every 48 hours.

RNA Extraction and RT-qPCR

For mRNA extraction from stromal cells, cells were lysed with TRIzol and processed using the DirectZol kit (Zymo, R2051) following manufacturer's procedures. Approximately 100 ng of mRNA was reverse transcribed into cDNA using iScript cDNA Supermix (Bio-Rad, 1708890) and amplified using specific primers listed in Supplementary Table 1. Primers were amplified using 2X SYBR Green Reagent (Life Technologies, 4364346) using a BioRad CFX384 Touch Real Time PCR Detection System. Data analysis was performed by calculating $\Delta\Delta CT$ value towards the geometric mean of *GAPDH*, *ACTB* and *RPL13A* and then normalized to siCTL. P-value was determined by One-Way ANOVA test with post hoc Tukey test using GraphPad Prism. * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$.

Results

Generation of Mouse Models with Global HA-Tagged SMAD1 and PA-Tagged SMAD5 Proteins

Activation of BMP signaling pathways has been established as one of the hallmarks of the decidualization process^{31,32}. Canonically, SMAD1/5 are regarded as downstream effectors of BMP2 signaling pathways to regulate decidual-specific gene expressions^{6,33}. However, our recent findings demonstrated that SMAD1/5 can also affect the sensitivity of the endometrium towards E2 and P4 stimulation⁹. Since we observed phenotypical differences between uterine-specific single SMAD1 and single SMAD5 deletion mice, it is beneficial to delineate the role of SMAD1 and SMAD5 in mediating P4 responses during early pregnancy. We used CRISPR technology to generate genetically engineered knock-in mice with an HA-tagged *Smad1* allele (herein called *Smad1*^{HA/HA}) and PA-tagged *Smad5* allele (herein called *Smad5*^{PA/PA}) as shown in **Figure 1A**, B. The HA tag and the PA tag were inserted into the N-terminus of the SMAD1 and SMAD5 proteins, respectively. Sanger sequencing was used to confirm genomic insertion (**Figure 1A**, B). To validate the global detection of tagged proteins, we performed immunoprecipitation followed by western blot analysis on different tissues from *Smad1*^{HA/HA} and *Smad5*^{PA/PA} mice. We confirmed the HA and PA antibodies can readily detect HA-tagged SMAD1 and PA-tagged SMAD5 proteins at the predicted size (**Figure 1C**, D). We also demonstrated the molecular size and expression pattern of HA antibody detected SMAD1 protein was comparable to the SMAD1 antibody detected SMAD1 protein across different tissue types. Similarly, PA antibody showed comparable signal intensity to the SMAD5 antibody in detecting SMAD5 protein across different tissue types (**Figure 1C**, D). Thus, we successfully generated viable mouse models with global HA-tagged SMAD1 and PA-tagged SMAD5 proteins.

SMAD1 and SMAD5 Exhibit Shared and Unique Genomic Binding Sites During Decidualization

The BMP signaling pathway regulates multiple key events during early pregnancy⁵, mediated through receptor-regulated SMAD proteins, including SMAD1 and SMAD5. As transducers of the BMP signaling pathway, phosphorylated SMAD1 and SMAD5 form homomeric complexes and then couple with SMAD4 to assemble hetero-oligomeric complexes in the nucleus to execute transcription programs. Our previous studies revealed that conditional ablation of SMAD1 and SMAD5 in the uterus decreased P4 response during the peri-implantation period, suggesting that the transcriptional activities of PR depend on BMP/SMAD1/5 signaling⁹. Furthermore, previous genome-wide PR binding studies show that SMAD1 and SMAD4 binding motifs are enriched in PR binding sites in the uterus³⁴.

To determine the shared and unique transcriptional regulomes of SMAD1 and SMAD5 contributing to the diverse effects of BMP and P4 signaling pathways during decidualization, we first utilized Cleavage Under Targets & Release Using Nuclease (CUT&RUN)²¹ coupled with next generation sequencing to profile genomic loci bound by SMAD1, SMAD5 and PR from mouse uterine tissues. We performed CUT&RUN on the uterine tissues collected at 4.5 days post coitus (dpc), the time when the fertilized embryo reaches the uterus physically and initiates the decidualization program³⁵ (**Figure 2A**). After aligning CUT&RUN reads to the mm10 mouse genome, we called peaks using Sparse Enrichment Analysis for CUT&RUN (SEACR)²². To identify high confidence peaks, background noise was normalized to IgG and the stringent criteria for peak calling in SEACR was used. After merging common peaks from two biological replicates, we identified 118,778 peaks for SMAD1 and 166,025 total peaks for SMAD5. We visualized the enrichment of SMAD1 and SMAD5 peaks to the overall aligned chromatin regions with clustering for preferential enrichment, as shown in **Figure 2B**. We found that 7.55% of SMAD1 peaks and 9.53% of SMAD5 peaks were located within the ± 3 kb of the promoter regions (**Figure 2C**, D). This corresponded to 10,368 genes that were directly bound by SMAD1 at the promoter regions (± 3 kb), whereas

18,270 genes were directly bound by SMAD5 at the promoter regions (± 3 kb). Among these, 4,933 genes were found in common between SMAD1 (47.5%) and SMAD5 (27.0%), while 2,744 and 7,427 genes were found to be uniquely bound by SMAD1 and SMAD5, respectively, providing evidence for the shared and unique functions of SMAD1 and SMAD5 at the transcriptional level (Supplement Figure 1B [↗](#)). Hence, interpreting how the binding events correlate to biological activity requires comparisons with gene expression profiling in a tissue-specific manner.

Identification of Direct Target Genes of SMAD1 and SMAD5 During Early Pregnancy

To pinpoint the direct target genes of SMAD1 and SMAD5, we integrated transcriptomic data from previously published⁹ SMAD1/5 double conditional knockout mice using progesterone receptor cre (SMAD1/5 cKO) (GSE152675) with SMAD1 and SMAD5 genomic data from this paper. We cross-compared the differentially expressed genes in the transcriptomic data to the SMAD1 and SMAD5 bound genes, respectively. Among the 805 significantly up-regulated genes, we identified 449 genes that were both significantly up-regulated upon SMAD1/5 depletion and were directly bound by SMAD1 and SMAD5, whereas 187 of the up-regulated genes were bound by SMAD5 only and 30 were bound only by SMAD1 (Figure 3A [↗](#)). Among the 683 significantly down-regulated genes, we identified 523 genes that were both significantly down-regulated upon SMAD1/5 depletion and were directly bound by SMAD1 and SMAD5, whereas 83 of the down-regulated genes were bound by SMAD5 only and 13 were bound by SMAD1 only (Figure 3B [↗](#), Supplementary Table 2 [↗](#)).

Next, we utilized Binding and Expression Target Analysis (BETA) algorithm³⁶ to perform motif enrichment analysis of the direct target genes to identify putative co-factors working together with SMAD1 and SMAD5 in controlling gene expression (Figure 3C,D [↗](#), Supplement Files 1,2). “Up-targets” represent genes that were up-regulated in the SMAD1/5 cKO mouse uteri and showed either a SMAD1 or a SMAD5 binding site in the genomic profiling data. Similarly, “down-targets” represent genes that were down-regulated in the SMAD1/5 cKO mouse uteri and displayed either a SMAD1 or a SMAD5 binding site. Thus, motifs enriched in the “up-targets” indicate potential repressive SMAD1/5 co-factors while motifs enriched in the “down-targets” indicate potential SMAD1/5 co-activators. Among the “up-targets” of SMAD1, MYB Proto-Oncogene (*Myb*)/MYB Proto-Oncogene Like 1 (*Mybl1*) motif was the most highly enriched with a P-value of 1.85E-02. *Myb* and *Mybl1* transcription factors belong to MYB gene family, which has been well-defined in controlling cell survival, proliferation and differentiation in cancer³⁷. In addition, they have also been reported to be E2 induced in human uterine leiomyoma samples³⁸.

Homeobox containing 1 (*Hmbox1*) and Krüppel-like factor (*Klf*) family members (*Klf4*/*Klf1*/*Klf12*) were also identified as potential repressive co-factors of SMAD1 with P-values of 2.85E-02 and 3.75E-02 respectively (Figure 2C [↗](#)). Of note, *KLF4* has been reported to inhibit the binding activity of estrogen receptor α (Ere) to estrogen response elements in promoter regions³⁹. Among the “up-targets” of SMAD5, EBF Transcription Factor 1 (*Ebf1*) motif was the most enriched with a P-Value of 1.57E-02. Interestingly, *Ebf1* can directly repress the transcription of Forkhead box protein O1 (*Foxo1*)⁴⁰. It is also recognized as downstream effector of steroid hormone receptors in the mouse uterus⁴¹. Additionally, motifs from transcription factors *Zfp128* and *Otx1* were also significantly enriched in the up-regulated genes bound by SMAD5 (Figure 2D [↗](#)). Taken together, our enrichment analysis provided robust evidence for identifying novel co-factors of SMAD1/5, and such co-regulating mechanisms are in line with the unopposed E2 response observed in the SMAD1/5 cKO mice⁹. Furthermore, odd-skipped-related genes (*Osr1* and *Osr2*) were identified as potential co-activators for SMAD1. *Osr2* has been reported to be highly expressed in the human endometrium⁴², and it was also abundantly detected at the protein level in the human decidual tissues⁴³. Decreased OSR2 level was observed in the patients with recurrent spontaneous abortion and knockdown of *OSR2* impairs the decidualization process in the human endometrial stromal cells⁴³. Moreover, *OSR1* has been reported to suppress BMP4 expression, which in turn




Figure 2

Genomic profiling of SMAD1 and SMAD5 binding sites during decidualization *in vivo*.

A) Diagram outlining experimental approaches for tissue collection, processing, and CUT&RUN. **B)** Heatmaps and summary plots showing the enrichment of SMAD1 and SMAD5 binding peaks from one exemplary replicate. Clustering was conducted using k-means algorithm. The colors in the summary plots correspond to clusters labeled in the heatmap below. **C-D)** Feature distribution of the annotated peaks for the SMAD1 (C) binding sites and SMAD5 (D) binding sites.




Figure 3

SMAD1 and SMAD5 show unique direct target genes during early pregnancy.

A-B) Venn diagrams showing the shared and unique direct up-target genes (A) and down-target genes (B) of SMAD1, SMAD5. Numbers indicate genes numbers. **C-D)** Motif enrichment analysis from the up-targets and down-targets for SMAD1 (C) and SMAD5 (D). **E-F)** Dot plot showing Gene Ontology enrichment analysis of shared direct target genes of SMAD1/5 from the up-targets (E) and the down-targets (F), respectively. Dot size represents the gene ratio in the enriched categories compared to background genes, dot colors reflect P-value.

reduced the Wnt/ β -catenin signaling pathways during lung development in xenopus⁴⁴. Apart from Osr family, motifs in the Homeobox genes (HOX) were found to be enriched in the “down-targets” from both SMAD1 and SMAD5 datasets. Specifically, *Hoxa11/Hoxd12/Hoxc10* were predicted to be co-activators for SMAD1 while *Hoxd10* was indicated to be closely interacting with SMAD5. Indeed, HOX genes are critical for endometrial development in normal and disease conditions and are essential during the establishment of pregnancy^{45,48}.

With direct target genes of SMAD1 and SMAD5 identified, we then analyzed the Gene Ontology enrichment for the SMAD1/5 shared up-targets and down-targets, respectively. We found that “up-targets” genes exhibit enrichment for regulation of cell-cell adhesion, cell junction organization and desmosomes organization (**Figure 3E**, Supplementary Table 3). Moreover, among the “down-targets” genes, we found the enrichment for blood vessel / vasculature development and extracellular matrix organization categories (**Figure 3F**, Supplementary Table 4). Indeed, during early pregnancy, the stimulation from corpus-luteum derived P4 enabled the endometrium to be transformed to a receptive state, which allows subsequent embryo attachment and develop through the epithelium into the stromal sections³¹. During this process, apporportioned direct cell-cell contacts are ensured by tight and adherent junctions and such interactions are key in facilitating implantation and embryo invasion. In accordance with our findings, desmosomes and adherens junctions were extensively described to decline in the early pregnancy period, which facilitates the invasion of trophoblast through the epithelial layer^{49,52}. In addition, the stromal compartment of the endometrium also undergoes profound vascular remodeling. Precise regulations of angiogenesis are required to establish an extensive vascular network, which is essential to ensure blood supply and successful embryonic development^{53,54}. Collectively, our findings present evidence that emphasizes the shared roles of SMAD1 and SMAD5 in facilitating endometrial transitions during early pregnancy.

Direct Target Genes of SMAD1 and SMAD5 Maintain the Homeostasis of Uterine Function

To discover novel direct target genes of SMAD1/5, we visualized key genes of interest from the up-targets and down-targets. As shown in **Figure 4A**, data from RNA-seq represents the decrease of several “down-targets” in the SMAD1/5 cKO mouse uteri, including Retinoic Acid-Related Orphan Receptor B (*Rorb*), Follistatin (*Fst*), Lymphoid Enhancer Binding Factor 1 (*Lef1*), and Insulin Like Growth Factor 1 (*Igf1*). Integrative Genomics Viewer (IGV) track view shows the exemplary SMAD1/5 binding activities near the promoter regions of *Rorb* and *Fst* **Figure 4B**, demonstrating that these genes are bona fide direct target genes of SMAD1/5. *Rorb* belongs to the nuclear receptor families in the retinoic acid (RA) signaling pathways⁵⁵ and is considered as a marker for mesenchymal progenitor cells in the stroma compartment of the endometrium⁵⁶. In murine models, deficient RA signaling through the perturbation of RA receptor in the uterus leads to implantation and decidualization failure⁵⁷. *Fst* binds several TGF β family ligands and thereby inhibits TGF β family signaling extracellularly⁵⁸. Under physiological conditions, *Fst* is up-regulated in the decidua during early pregnancy. Conditional deletion of *Fst* in the mouse uterus results in severe subfertility with a phenotype of non-receptive epithelium and poor-differentiated stroma⁵⁹. Notably, RA signaling deficiency also decreases *Fst* levels in the uterus and systematic administration of FST can fully rescue the deficient-decidualization phenotype but not the non-receptive phenotype observed in the RA receptor mutant mice⁵⁷. Our results suggest a direct relationship between BMP and RA signaling pathway, accomplished by SMAD1/5 at the transcriptional level, likely establishing a positive signaling feedback loop. Apart from being a crucial transcriptional activator, SMAD1/5 also plays a role in repressing key gene expression pathways. Shown in **Figure 4C**, upon the deletion of SMAD1/5 in the mouse uteri, several E2-responsive genes were significantly up-regulated, including Fibroblast Growth Factor Receptor 2 (*Fgfr2*), Matrix Metalloproteinase 7 (*Mmp7*) and Wnt Family Member 7B (*Wnt7b*). In addition, *Inhbb*, a downstream target of *Fst*⁵⁹, is also a target gene of SMAD1/5 that resulted in transcriptional repression. SMAD1/5 binding on the *Fgfr2* and *Mmp7* genes are exemplified in an IGV track view in

Figure 4D [↗](#). *Fgfr2* and its ligands regulate epithelial cell proliferation and differentiation. Components of the Fibroblast Growth Factor (*Fgf*) signaling pathway are cyclically expressed in the uterus and act as paracrine and/or autocrine mediators of epithelial-stromal interactions^{60,61}. During early pregnancy in mice, P4 inhibits expression of *Fgfr2* in the stromal cells, which is critical to counteract the E2-driven epithelial proliferation⁶⁰. Similar observations are reported in gilts, where the expression of *Fgfr2* decreased alongside with increased parity of the sows⁶². It is also noteworthy that loss of function of *Fgfr2* in the mouse uterus leads to luminal epithelial stratification and peri-implantation pregnancy loss⁶¹. Moreover, *Mmp7* and *Wnt7b* are up-regulated upon E2 stimulation and participate in the re-epithelialization of the endometrium and implantation process, respectively^{63,65}. In accordance with the phenotype of hyperproliferative endometrial epithelium during early pregnancy observed SMAD1/5 cKO mice, we demonstrated that the suppression of key E2-responsive genes, such as *Fgfr2* and *Mmp7*, by SMAD1/5 maintains the precise balance between E2 and P4.

To explore the major cell types regulated by SMAD1/5 direct targets in human, we profiled the expression levels of the key “up-targets” and “down-targets” in the different cell types of the human endometrium. Using previously published single-cell RNA seq data of human endometrium⁶⁶, we visualized the expression patterns of suppressive targets and activating targets of SMAD1/5 (**Figure 4E** [↗](#)). Apart from the major epithelial and stromal compartments, SMAD1/5 target genes are also widely expressed in the immune cell populations. Such observation reinforced the importance of the BMP signaling pathways in establishing an immune-privileged environment at the maternal-fetal interface⁶⁷.

SMAD1 and SMAD5 Co-regulate PR Target Genes

SMAD1/5 cKO mice were infertile due to endometrial defects and displayed decreased P4 response during the peri-implantation period⁹. Hence, we hypothesized that SMAD1 and SMAD5 act as co-regulators of P4-responsive genes during the window of implantation and are required for endometrial receptivity and decidualization. By determining the genomic co-occupancy of SMAD1, SMAD5 and PR, we aimed to clarify the transcriptional interplay between the BMP and P4 signaling pathways. To this end, we performed additional PR CUT&RUN experiments on the uteri of mice collected at 4.5 dpc and identified 134,737 peaks showing PR binding activities (**Figure 5A** [↗](#)). We identified 7,393 genes that were directly bound by PR at the promoter regions (± 3 kb), among which, 2596 genes were also concurrently bound by both SMAD1 and SMAD5 at the promoter regions (± 3 kb) (**Supplement Figure 1B** [↗](#)).

Next, we performed KEGG pathway enrichment for the genes co-bound by SMAD1, SMAD5 and PR. As expected, pathways critical for decidualization such as relaxin signaling pathways and WNT signaling cascade were identified in the enrichment results (**Figure 5B** [↗](#), Supplementary Table 5). We visualized exemplary genes co-regulated by SMAD1, SMAD5 and PR and presented in the normalized IGV track view (**Figure 5C** [↗](#)). We demonstrated SMAD1, SMAD5 and PR showed co-occupancy at the loci of the SRY-Box Transcription Factor 17 (*Sox17*), Inhibitor of DNA binding 2 (*Id2*), Forkhead box protein O1 (*Foxo1*), Insulin-like growth factor 1 (*Igf1*), Transforming growth factor beta receptor 2 (*Tgfb2*) and RUNX family transcription factor 1 (*Runx1*) (**Figure 5C** [↗](#)). *Sox17* has been reported as one of the direct target genes of PR³⁴ and is essential for uterine functions during implantation and early pregnancy^{68,69}. More recent studies also showed the importance of *Sox17* in regulating uterine epithelial-stromal crosstalk and its indispensable role in female fertility⁷⁰. We provided evidence that *Sox17* is also directly regulated by SMAD1/5 complexes. Our results indicated that *Id2*, considered as canonical direct transcriptional targets of BMP-SMAD signaling^{71,72} is also regulated by PR. We also confirmed that known P4-responsive genes such as *Tgfb2*⁷³ and *Runx1*⁷⁴, as well as decidual markers such as *Foxo1*⁷⁵ and *Igf1*⁷⁶, were co-regulated by SMAD1, SMAD5 and PR (**Figure 5C** [↗](#)).




Figure 4

Direct target genes of SMAD1/5 mediate uterine homeostasis.

A) Bar graph of normalized Fragments Per Kilobase of transcript per Million mapped reads (FPKM) of downregulated transcripts in the Control and SMAD1/5 cKO groups as indicated by the label. Histograms represent average $\pm$ SEM of experiments uteri from Control mice (N=3) and SMAD1/5 cKO mice (N=4). Analyzed by a unpaired t-test. **B**) Integrative Genomics Viewer (IGV) track view of SMAD1, SMAD5 binding activities. Gene loci are as indicated in the figure, genomic coordinates are annotated in mm10. **C**) Bar graph of FPKM of up regulated transcripts in the Control and SMAD1/5 cKO groups as indicated by the label. **D**) IGV track view of SMAD1, SMAD5 binding activities. Gene loci are as indicated in the figure, genomic coordinates are annotated in mm10. **E**) Dot plot showing the gene expression pattern of the key SMAD1/5 direct target genes in different cell types from published human endometrium single-cell RNA-seq dataset.




Figure 5

SMAD1 and SMAD5 co-regulate PR target genes.

A) Heatmaps and summary plots showing the enrichment comparison between SMAD1, SMAD5 and PR binding peaks from one exemplary replicate. Clustering was conducted using k-means algorithm. The colors in the summary plots correspond to clusters labeled in the heatmap below. **B)** Dot plot showing KEGG pathway enrichment analysis for shared genes bound by SMAD1, SMAD5, and PR. **C)** IGV track view of SMAD1, SMAD5 and PR binding activities. Gene loci are as indicated in the figure, genomic coordinates are annotated in mm10. **D)** Table of motif analysis results for shared peaks between SMAD1, SMAD5 and PR, with P-value and motif annotation specified for each motif.

To identify additional transcription factors that are associated with the regulatory interplay between SMAD1/5 and PR during decidualization, we performed unbiased motif analysis on the shared CUT&RUN peaks between SMAD1/5 and PR. We reported the top 10 transcription factors harboring the enriched motifs, including NANOG, Homeobox A protein family (HOXA11 and HOXA9), NK6 homeobox 1 (NKX6.1), TGF β induced factor homeobox 2 (TGIF2), FOS, RUNX family transcription factor 2 (RUNX2), Androgen receptor (AR), SOX17 and Lymphoid enhancer-binding factor 1 (LEF1) (**Figure 5D** [↗](#), Supplementary Table 6). Many of these putative interactors have been reported to interact with the SMAD proteins in other biological processes. For example, NANOG interacts with SMAD1 during mesoderm differentiation⁷⁷ [↗](#), HOXA9 forms heterodimers with SMAD4, leading to BMP-driven initiation of transcription from the mouse *Opn* promoter *in vitro*⁷⁸ [↗](#),⁷⁹ [↗](#). Transcription factor AP-1 family (FOS) and RUNX2, as well as β -catenin/Lef1 complex, increase the effectiveness and specificity of DNA binding activities of SMAD1/5 in response to BMP ligand stimuli⁸⁰ [↗](#),⁸² [↗](#). To further evaluate the key roles of SMAD1/5 as major uterine transcription regulators, we cross-compared the genomic binding sites of SMAD1/5 with known key transcription factors, namely aforementioned SOX17 (**Supplement Figure 1E** [↗](#), Supplementary Table 7), as well as NR2F2 (**Supplement Figure 1F** [↗](#), Supplementary Table 8), an essential regulator of hormonal response⁸³ [↗](#), using our CUT&RUN data sets and published mouse uterine SOX17 and NR2F2 ChIP-seq data sets (GSE118328, GSE232583). Among the annotated genes, 5402 genes are shared between SMAD1/5 and SOX17, and 1922 genes are shared between SMAD1/5 and NR2F2. Such observations indicate a potential co-regulatory mechanism between SMAD1/5 and other key uterine transcription factors in maintaining appropriate uterine functions. Overall, our analyses demonstrate that the transcriptional activity of SMAD1, SMAD5, and PR coordinate the expression of key genes required for endometrial receptivity and decidualization.

Decidualization of Human Endometrial Stromal Cells Requires SMAD1/SMAD5

We next sought to functionally characterize the role of SMAD1/5 during decidualization in human endometrial stromal cells. To do so, we examined the effect of SMAD1/5 perturbations on the decidualization of primary human endometrial stromal cells (EnSCs). EnSCs were transfected with short interfering RNAs (siRNAs) targeting each gene (*SMAD1* and *SMAD5*) and subjected to *in vitro* decidualization by treatment with E2-cAMP and MPA (EPC) for 4.5 days (**Figure 6A** [↗](#)). We hypothesized that the combined SMAD1/5 knockdown would impair the decidualization process significantly compared to cells treated with non-targeting siRNAs. Our results demonstrated that SMAD1/5 knockdown affected decidualization and led to significantly decreased expression of the canonical decidual markers, *PRL* and *IGFBP1* in EnSCs (**Figure 6B** [↗](#)). The PR co-regulator, *FOXO1*⁸⁴ [↗](#), also exhibited a significant decreasing trend in the siSMAD1/5 group. We also examined the expression level of the RA pathway regulator gene, *RORB*, and of the SMAD4-PR target gene, *KLF15*¹² [↗](#), following SMAD1/5 perturbation. We observed a significant decrease in both *RORB* and *KLF15* expression upon SMAD1/5 knockdown during *in vitro* decidualization treatment (**Figure 6C** [↗](#)). Taken together, our findings indicate SMAD1/5 can modulate PR activity during decidualization and that this transcriptional cooperation is required for the *in vitro* decidualization of primary human endometrial stromal cells.

Discussion

SMAD proteins are canonical transcription factors that are activated in response to TGF β family signaling and mediate the biological effects of these pathophysiologically critical ligands⁸¹ [↗](#). While SMAD2 and SMAD3 are downstream of TGF β s, activins, and multiple other family ligands, SMAD1 and SMAD5 preferentially transduce BMP signaling pathways and are regarded as pivotal activators for many physiological processes, including bone development, cardiac conduction




Figure 6

SMAD1 and SMAD5 are required for PR responses during decidualization of human endometrial stromal cells.

A) Schematic approach and timeline outlining in vitro decidualization for endometrial stromal cells (EnSCs). **B-C**) RT-qPCR results showing mRNA levels of *PRL*, *IGFBP1*, *FOXO1*, *RORB* and *KLF15* after SMAD1/5 perturbation using siRNAs. Data are normalized to siCTL-Veh for visualization. Bar graphs represent average $\pm$ SEM of experiments on cells from three different individuals with technical triplicates. Analyzed by a One-Way ANOVA test with post hoc Tukey test.

system development, and embryonic pattern specification^{85,87}. Importantly, SMAD1 and SMAD5 are implicated in diverse female reproductive physiology and pathophysiology processes^{5,9,88-90}.

Due to high structural similarity, SMAD1/5 have been suggested to be redundant from the studies in ovarian biology and chondrogenesis^{89,91}. However, other studies clearly demonstrated that SMAD1/5 have different roles in governing hematopoiesis and uterine functions^{9,92}. The DNA binding activities of SMAD1 and SMAD5 have not been readily distinguished from each other due to anti-phospho antibody limitations. To robustly define the roles of SMAD1/5 in regulating transcriptional programs *in vivo*, we produced two genetically engineered mouse models with global knock-in of an HA tag and a PA tag in the *Smad1* and *Smad5* loci, respectively. We showed that SMAD1 and SMAD5 not only have shared transcriptional activities but also have unique roles in uterine physiology. In agreement with previous studies showing that SMAD1/5 function is partially redundant^{89,91}, we confirmed that SMAD1/5 share a total of 972 direct target genes in the uterus. Furthermore, we demonstrated that 43 genes were uniquely regulated by SMAD1 whereas 270 genes are specifically regulated by SMAD5 only. Our motif analysis also revealed distinct potential co-factors between SMAD1 and SMAD5, providing evidence at the molecular level to mechanistically delineating the distinct roles of SMAD1 and SMAD5 in directing cellular processes in the uterus.

Apart from directly regulating target gene expression, our data demonstrate that SMAD1/5 present as dense genomic occupancies. To date, only a limited amount of transcription factors have been investigated using the CUT&RUN-seq technique from the tissue samples due to antibody compatibility issue. We recognize that the binding sites and gene number identified here are quite high; however, the high density of binding events was also observed in the ENCODE⁹³ chromatin immunoprecipitation followed by sequencing (ChIP-seq) data for SMAD1 and SMAD5 in the human K562 cells, detecting an average of 63,563 peaks for SMAD1 and 109,682 peaks for SMAD5. (Data accessed through GSE95876 and GSE127365 from Gene Expression Omnibus). Multiple aspects can contribute to the observation of dense SMAD1/5 genome occupancies. First, transcription factors (TFs) tend to dwell or “search and bind” throughout the genome⁹⁴. Such events may not yield actual biological effects but rather are due to differences in motif binding affinities⁹⁵. Second, apart from robust binding activities, TFs may not initiate transcription programs owing to the lack of co-factors or favorable conditions to exert their functions⁹⁶.

Additionally, TF binding sites and target genes are unlikely to have a one-to-one relationship. TFs could be positioned from the proximal promoter regions to hundreds of kilobases afar to modulate gene expression. In the meantime, the same binding site could regulate multiple genes by interacting with different promoters in different subpopulations of cells. Lastly, TFs usually direct target gene expression in a cell-type specific manner⁹⁷. Our genomic profiling samples were collected from whole uterus at the time of 4.5 dpc, containing a great range of cell populations, including but not limited to the epithelium (luminal and glandular), stroma (progenitors and differentiated cells), myometrium, endothelium, and immune cell populations. The data is therefore expected to depict the dynamic and complex activities of SMAD1/5 in the entire uterus. Together, the stringent filtering and normalization criteria, comparable peak number to the published dataset and IGV track view visualization collectively validate our CUT&RUN experiments and uncover the enriched regions as robust SMAD1/5 binding events. Our studies also examined the role of SMAD1/5 in mediating progesterone responses at the genomic and transcription levels. Similarly, our analysis was based on data sets generated from the whole mouse uterus, which contains multiple compartments of the uterine structures, including but not limited to epithelium and stroma. Published studies have shown that nuclear SMAD1/5 localize to the stroma and epithelium during the decidualization process at 4.5 dpc, during the window of implantation⁹. Conditional deletion of SMAD1/5 exclusively in the uterine epithelium using lactoferrin-icre (*Ltf-icre*) results in severe subfertility due to impaired implantation and decidual development⁹⁸. Conditional deletion of SMAD1/5/4 exclusively in the cells from mesenchymal

lineage (including uterine stroma) using anti-Mullerian hormone type 2 receptor cre (*Amhr2-cre*) results in infertility with defective decidualization^{89,90}. Given the essential roles of SMAD1/5 in both stroma and epithelium identified by previous studies, we believe that the transcriptional co-regulatory roles of SMAD1/5 and PR reported here using the whole uterus validates a relationship between SMAD1/5 and PR in both the stromal and epithelial compartments. However, it does not rule out potential co-regulatory roles of SMAD1/5 and PR in the myometrium, immune cells, and/or endothelium, given that whole uterus was used. The specific transcriptional evaluations of SMAD1/5 in the stroma versus the epithelium would require future single-cell sequencing (i.e., digital cytometry⁹⁹) and/or spatial transcriptomic analysis.

Although our studies herein confirm that SMAD1 and SMAD5 proteins have distinct transcriptional regulatory activities, our previous studies demonstrated that while SMAD5 can functionally replace SMAD1, SMAD1 cannot replace SMAD5 in the uterus⁹. How this epistatic relationship is established in a tissue-specific manner still needs to be determined by further biochemical investigations. In addition, further studies are needed to uncover whether SMAD1 and SMAD5 response differently upon ligand stimulation in the uterus, and if so, how the preference is achieved. Our study provides versatile *in vivo* genetic tools for these questions and can advance the toolbox for the field studying BMP signaling pathways. Because our mouse models are global knock-in mice, they will not only serve as a powerful tool for studying BMP signaling pathways in the reproductive system but will also promote the study of BMP signaling in other organs and tissues.

BMP signaling pathways are involved in a plethora of cellular processes and appropriate functioning of the BMP pathway depends on the precise crosstalk with other signaling pathways. Coordinated communication with other pathways can yield synergistic effects and lead to a complex regulatory network of biological processes. To be specific, SMAD1/5 mediates the crosstalk with the WNT/ β -catenin pathway. WNT signaling inhibits glycogen synthase kinase 3 β (GSK3 β) activity and prevents SMAD1 from degradation which governs the embryonic pattern formation¹⁰⁰. Also, SMAD1/5 can physically interact with T-cell factor (TCF) or lymphoid enhancer factor (LEF) transcription factors to form transcriptional complexes to activate the transcription of many WNT- and BMP-responsive genes¹⁰¹. In addition, SMAD1 and SMAD5 can directly associate with Notch intracellular domain and enhance known Notch target gene expression by binding to their regulatory DNA sequences¹⁰². Intriguingly, in prostate cells, SMAD1 physically interacts with the androgen receptor (AR) and halts the androgen-stimulated prostate cell growth¹⁰³. Moreover, we provide first-hand evidence showing that BMP signaling pathways converge with RA signaling pathways through the regulation of RORB by SMAD1/5. Further studies will grant a more detailed mechanism of the positive feedback loop between BMP and RA signaling.

Our previous studies suggest that the mouse endometrium presents decreased P4 responsiveness following the conditional deletion of SMAD1/5 in the uterus⁹. In accordance with the phenotypical observation, we offer compelling support in our current study that SMAD1/5 work collectively with PR to regulate their target genes and that SMAD1/5 mediate the crosstalk between BMP and P4 signaling pathways during decidualization, a key process to ensure a successful pregnancy, and ultimately direct the biological transformations of the uterus during early pregnancy. We provide genomic evidence that SMAD1/5 are co-bound at around 35% of PR target genes in the mouse uterus during decidualization. Correspondingly, in a previously published study where they performed PR ChIP-seq in the mouse uterus after P4 stimulation, the SMAD1 motif was the 5th most significantly enriched sequence motifs identified³⁴. In parallel, we also identified nuclear receptor motifs (i.e., PR sequence motifs) enriched in the SMAD1/5 binding sites (**Supplement Figure 1C,D**). From pathway enrichment analysis, we demonstrate that genes with SMAD1/5 and PR bound at the promoter regions are enriched for key pathways in directing the decidualization process, such as WNT and relaxin signaling pathways. Future studies can benefit from analyzing binding events beyond the promoter regions. Profiling the PR genome

occupancy in the SMAD1/5 deficient mice would provide an interesting perspective to reevaluate the major regulatory roles of SMAD1/5 in mediating uterine transcriptomes. In this study, we determined the overlapped transcriptional control between SMAD1/5 and PR at the gene level, and functionally validated the regulatory effect at the transcript level in a human stromal cell decidualization model. While we observe a subset of peak representations that do not overlap at the base pair level in the promoter regions, future functional screenings at the promoter level, such as luciferase reporter assays to assess transcriptional co-activation by SMAD1/5 and PR, will advance this study.

SMADs are known to recruit co-repressors (*i.e.*, Ski¹⁰⁴[\[104\]](#)) or co-activators (*i.e.*, p300¹⁰⁵[\[105\]](#)) to inhibit or activate target gene transcription, less is known about their cell-specific co-factors that confer the precise spatial-temporal control over binding activities to target genes. Our study highlights the potential co-factors by integrating both genomic and transcriptomic data to delineate signaling crosstalks that are responsible for maintaining tissue homeostasis, especially in the female reproductive tract.

Since mice only undergo decidualization upon embryo implantation whilst human stromal cells undergo cyclic decidualization in each menstrual cycle in response to rising levels of progesterone³⁵[\[35\]](#), asynchronous gene responses may occur in comparison between mouse models and human cells. However, cellular transformation during decidualization is conserved between mice and humans³¹[\[31\]](#), which makes findings in the mouse models a valuable and transferable resource to be evaluated in human tissues. Accordingly, our functional validation studies were performed using human endometrial stromal cells induced to decidualize *in vitro* for four days, which models the early phases of decidualization. Additional transcriptomic studies of the SMAD1/5 perturbations in human endometrial stromal cells will be of great resource in understanding the entire SMAD1/5 regulomes in humans.

In summary, our findings and those of others indicate that SMAD1 and SMAD5 not only are signal transducers for BMP signaling pathways, but also engage extensively in the crosstalk with PR signaling pathways. While P4 responses are critical for early pregnancy establishment, abnormal P4 responses are implicated in diseases such as endometriosis and endometrial cancers¹⁰⁶[\[106\]](#),¹⁰⁹[\[109\]](#). Hence, our results which show that BMP and P4 signaling pathways synergize within the endometrium; these key pathways can shed light on the endometrial contribution to conditions that impact reproductive health in women, including early pregnancy loss, endometriosis, and endometrial cancer. Furthermore, we anticipate that the SMAD1/5 knock-in tagged transgenic mouse models developed herein will be useful for studying BMP/SMAD1/5 signaling pathways in other reproductive and non-reproductive tract tissues in the body.

Data availability

Sequencing data and analyses are deposited in the Gene Expression Omnibus under accession number GSE237975. (Reviewer token: gludeoohrczxcj)

Supplementary data

Supplementary Table 1: DNA sequences used in the study.

Supplementary Table 2: Direct target genes of SMAD1/5.

Supplementary Table 3: Results of Gene Ontology enrichment analysis of shared direct target genes of SMAD1/5 from the up-targets.

Supplementary Table 4: Results of Gene Ontology enrichment analysis of shared direct target genes of SMAD1/5 from the down-targets.

Supplementary Table 5: Results of KEGG pathway enrichment analysis for shared genes bound by SMAD1, SMAD5, and PR.

Supplementary Table 6: Results of motif analysis results for shared peaks between SMAD1, SMAD5 and PR.

Supplementary Table 7: Gene overlapping results of SMAD1/5 and SOX17.

Supplementary Table 8: Gene overlapping results of SMAD1/5 and NR2F2.

Supplementary File 1: Complete list of motif enrichment analysis from the up-targets and down-targets for SMAD1.

Supplementary File 2: Complete list of motif enrichment analysis from the up-targets and down-targets for SMAD5.

Supplementary Figures

Author contributions

Zian Liao: Conceptualization, Formal analysis, Methodology, Validation, Writing—original draft. Suni Tang: Data curation, Investigation, Methodology, Visualization, Writing—review & editing. Kaori Nozawa: Data curation, Investigation, Methodology, Writing—review & editing. Keisuke Shimada: Data curation, Methodology, Writing—review & editing. Masahito Ikawa: Project administration, Resources. Diana Monsivais: Conceptualization, Investigation, Supervision, Funding acquisition, Writing – review & editing. Martin M. Matzuk: Conceptualization, Supervision, Funding acquisition, Writing – review & editing.

Acknowledgements


FUNDING

This research is supported by Eunice Kennedy Shriver National Institute of Child Health and Human Development grants HD105800 (D.M.), HD096057 (D.M.), HD032067 (M.M.M.), and HD110038

(M.M.M.). DM is supported by a Next Gen Pregnancy Award from the Burroughs Wellcome Fund (NGP10125).

Conflict of interest


No conflict of interest is declared by the authors.



Supplement Figure 1

Gene numbers with SMAD1/5 promoter binding activities and motif analysis of SMAD1/5 peaks.


A) Venn diagrams showing the shared and unique genes bound by SMAD1 or SMAD5 in the +/- 3kb region of the promoter regions. **B)** Venn diagrams showing the shared and unique genes bound by SMAD1, SMAD5 or PR in the +/- 3kb of the promoter regions. **C-D)** Table of motif analysis results for unique peaks for SMAD1(C) and SMAD5 (D), with P-value and motif annotation specified for each motif. **E)** Venn diagrams showing the shared and unique genes bound by SMAD1, SMAD5 or SOX17 in the +/- 25kb regions near transcription start site. **F)** Venn diagrams showing the shared and unique genes bound by SMAD1, SMAD5 or NR2F2 in the +/- 25kb regions near transcription start site.



Supplement Figure 2

Knockdown effect validation of SMAD1/5 perturbation.

A-B RT-qPCR results showing mRNA levels of SMAD1(A) and SMAD5 (B) after siRNA treatments in the both Veh and EPC conditions. Data are normalized to siCTL for visualization. Bar graphs represent average $\pm$ SEM of experiments on cells from three different individuals with technical triplicates. Analyzed by a One-Way ANOVA test with post hoc Tukey test.




Supplement Figure 3

Genotype of the knock-in mouse lines.

A) Schematic design of the genotype primers for Smad1HA/HA and Smad5PA/PA mouse lines. **B**) Exemplary gel electrophoresis of PCR products derived from homozygous knock-in mice, heterozygous mice, and WT mice using genotyping primers.

Uncropped Western Blots for Figure 1



References

- 1 Boivin J., Bunting L., Collins J. A., Nygren K. G (2007) **International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care** *Hum Reprod* **22**:1506–1512 <https://doi.org/10.1093/humrep/dem046>
- 2 Martinez G. M., Daniels K., Febo-Vazquez I (2018) **Fertility of Men and Women Aged 15–44 in the United States: National Survey of Family Growth, 2011–2015** *Natl Health Stat Report* **113**:1–17
- 3 Wang X., et al. (2003) **Conception, early pregnancy loss, and time to clinical pregnancy: a population-based prospective study** *Fertil Steril* **79**:577–584 [https://doi.org/10.1016/s0015-0282\(02\)04694-0](https://doi.org/10.1016/s0015-0282(02)04694-0)
- 4 Chang H., Brown C. W., Matzuk M. M (2002) **Genetic analysis of the mammalian transforming growth factor-beta superfamily** *Endocr Rev* **23**:787–823 <https://doi.org/10.1210/er.2002-0003>
- 5 Monsivais D., Matzuk M. M., Pangas S. A (2017) **The TGF-beta Family in the Reproductive Tract** *Cold Spring Harb Perspect Biol* **9** <https://doi.org/10.1101/cshperspect.a022251>
- 6 Lee K. Y., et al. (2007) **Bmp2 is critical for the murine uterine decidual response** *Mol Cell Biol* **27**:5468–5478 <https://doi.org/10.1128/MCB.00342-07>
- 7 Monsivais D., et al. (2017) **BMP7 Induces Uterine Receptivity and Blastocyst Attachment** *Endocrinology* **158**:979–992 <https://doi.org/10.1210/en.2016-1629>
- 8 Nagashima T., et al. (2013) **BMPR2 is required for postimplantation uterine function and pregnancy maintenance** *J Clin Invest* **123**:2539–2550 <https://doi.org/10.1172/JCI65710>
- 9 Monsivais D., et al. (2021) **Endometrial receptivity and implantation require uterine BMP signaling through an ACVR2A-SMAD1/SMAD5 axis** *Nat Commun* **12** <https://doi.org/10.1038/s41467-021-23571-5>
- 10 Matzuk M. M., Kumar T. R., Bradley A (1995) **Different phenotypes for mice deficient in either activins or activin receptor type II** *Nature* **374**:356–360 <https://doi.org/10.1038/374356a0>
- 11 Clementi C., et al. (2013) **Activin-like kinase 2 functions in peri-implantation uterine signaling in mice and humans** *PLoS Genet* **9** <https://doi.org/10.1371/journal.pgen.1003863>
- 12 Monsivais D., et al. (2016) **Uterine ALK3 is essential during the window of implantation** *Proc Natl Acad Sci U S A* **113**:E387–395 <https://doi.org/10.1073/pnas.1523758113>
- 13 Teklenburg G., et al. (2010) **Natural selection of human embryos: decidualizing endometrial stromal cells serve as sensors of embryo quality upon implantation** *PLoS One* **5** <https://doi.org/10.1371/journal.pone.0010258>
- 14 Brosens J. J., et al. (2014) **Uterine selection of human embryos at implantation** *Sci Rep* **4** <https://doi.org/10.1038/srep03894>

- 15 Weimar C. H., et al. (2012) **Endometrial stromal cells of women with recurrent miscarriage fail to discriminate between high- and low-quality human embryos** *PLoS One* **7** <https://doi.org/10.1371/journal.pone.0041424>
- 16 Salker M. S., et al. (2011) **Deregulation of the serum- and glucocorticoid-inducible kinase SGK1 in the endometrium causes reproductive failure** *Nat Med* **17**:1509–1513 <https://doi.org/10.1038/nm.2498>
- 17 Garrido-Gomez T., et al. (2017) **Defective decidualization during and after severe preeclampsia reveals a possible maternal contribution to the etiology** *Proc Natl Acad Sci U S A* **114**:E8468–E8477 <https://doi.org/10.1073/pnas.1706546114>
- 18 Garrido-Gomez T., et al. (2020) **Preeclampsia: a defect in decidualization is associated with deficiency of Annexin A2** *Am J Obstet Gynecol* **222**:e371–376 <https://doi.org/10.1016/j.ajog.2019.11.1250>
- 19 Shimada K., et al. (2021) **ARMC12 regulates spatiotemporal mitochondrial dynamics during spermiogenesis and is required for male fertility** *Proc Natl Acad Sci U S A* **118** <https://doi.org/10.1073/pnas.2018355118>
- 20 Fang B., et al. (2014) **Circadian enhancers coordinate multiple phases of rhythmic gene transcription in vivo** *Cell* **159**:1140–1152 <https://doi.org/10.1016/j.cell.2014.10.022>
- 21 Skene P. J., Henikoff S (2017) **An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites** *Elife* **6** <https://doi.org/10.7554/eLife.21856>
- 22 Meers M. P., Tenenbaum D., Henikoff S (2019) **Peak calling by Sparse Enrichment Analysis for CUT&RUN chromatin profiling** *Epigenetics Chromatin* **12** <https://doi.org/10.1186/s13072-019-0287-4>
- 23 Kent W. J., Zweig A. S., Barber G., Hinrichs A. S., Karolchik D (2010) **BigWig and BigBed: enabling browsing of large distributed datasets** *Bioinformatics* **26** <https://doi.org/10.1093/bioinformatics/btq351>
- 24 Heinz S., et al. (2010) **Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities** *Mol Cell* **38**:576–589 <https://doi.org/10.1016/j.molcel.2010.05.004>
- 25 Yu G., Wang L. G., He Q. Y (2015) **ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization** *Bioinformatics* **31**:2382–2383 <https://doi.org/10.1093/bioinformatics/btv145>
- 26 Ramirez F., et al. (2016) **deepTools2: a next generation web server for deep-sequencing data analysis** *Nucleic Acids Res* **44**:W160–165 <https://doi.org/10.1093/nar/gkw257>
- 27 (2023) **Initiative, C. Z. (2023).**
- 28 Warren L. A., et al. (2018) **Analysis of menstrual effluent: diagnostic potential for endometriosis** *Mol Med* **24** <https://doi.org/10.1186/s10020-018-0009-6>
- 29 Nayyar A., et al. (2020) **Menstrual Effluent Provides a Novel Diagnostic Window on the Pathogenesis of Endometriosis** *Front Reprod Health* **2** <https://doi.org/10.3389/frph.2020.00003>

- 30 Martinez-Aguilar R., et al. (2020) **Menstrual blood-derived stromal cells modulate functional properties of mouse and human macrophages** *Sci Rep* **10** <https://doi.org/10.1038/s41598-020-78423-x>
- 31 Gellersen B., Brosens J. J (2014) **Cyclic decidualization of the human endometrium in reproductive health and failure** *Endocr Rev* **35**:851–905 <https://doi.org/10.1210/er.2014-1045>
- 32 Magro-Lopez E., Munoz-Fernandez M. A (2021) **The Role of BMP Signaling in Female Reproductive System Development and Function** *Int J Mol Sci* **22** <https://doi.org/10.3390/ijms222111927>
- 33 Li Q., et al. (2007) **Bone morphogenetic protein 2 functions via a conserved signaling pathway involving Wnt4 to regulate uterine decidualization in the mouse and the human** *J Biol Chem* **282**:31725–31732 <https://doi.org/10.1074/jbc.M704723200>
- 34 Rubel C. A., et al. (2012) **Research resource: Genome-wide profiling of progesterone receptor binding in the mouse uterus** *Mol Endocrinol* **26**:1428–1442 <https://doi.org/10.1210/me.2011-1355>
- 35 Ramathal C. Y., Bagchi I. C., Taylor R. N., Bagchi M. K (2010) **Endometrial decidualization: of mice and men** *Semin Reprod Med* **28**:17–26 <https://doi.org/10.1055/s-0029-1242989>
- 36 Wang S., et al. (2013) **Target analysis by integration of transcriptome and ChIP-seq data with BETA** *Nat Protoc* **8**:2502–2515 <https://doi.org/10.1038/nprot.2013.150>
- 37 Ciciro Y., Sala A (2021) **MYB oncoproteins: emerging players and potential therapeutic targets in human cancer** *Oncogenesis* **10** <https://doi.org/10.1038/s41389-021-00309-y>
- 38 Swartz C. D., Afshari C. A., Yu L., Hall K. E., Dixon D (2005) **Estrogen-induced changes in IGF-I, Myb family and MAP kinase pathway genes in human uterine leiomyoma and normal uterine smooth muscle cell lines** *Mol Hum Reprod* **11**:441–450 <https://doi.org/10.1093/molehr/gah174>
- 39 Akaogi K., et al. (2009) **KLF4 suppresses estrogen-dependent breast cancer growth by inhibiting the transcriptional activity of ERalpha** *Oncogene* **28**:2894–2902 <https://doi.org/10.1038/onc.2009.151>
- 40 Timblin G. A., Schlissel M. S (2013) **Ebf1 and c-Myb repress rag transcription downstream of Stat5 during early B cell development** *J Immunol* **191**:4676–4687 <https://doi.org/10.4049/jimmunol.1301675>
- 41 Pan H., Zhu L., Deng Y., Pollard J. W (2006) **Microarray analysis of uterine epithelial gene expression during the implantation window in the mouse** *Endocrinology* **147**:4904–4916 <https://doi.org/10.1210/en.2006-0140>
- 42 Fagerberg L., et al. (2014) **Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics** *Mol Cell Proteomics* **13**:397–406 <https://doi.org/10.1074/mcp.M113.035600>
- 43 Ma W., et al. (2022) **MAX deficiency impairs human endometrial decidualization through down-regulating OSR2 in women with recurrent spontaneous abortion** *Cell Tissue Res* **388**:453–469 <https://doi.org/10.1007/s00441-022-03579-z>

- 44 Rankin S. A., Gallas A. L., Neto A., Gomez-Skarmeta J. L., Zorn A. M (2012) **Suppression of Bmp4 signaling by the zinc-finger repressors Osr1 and Osr2 is required for Wnt/beta-catenin-mediated lung specification in *Xenopus*** *Development* **139**:3010–3020 <https://doi.org/10.1242/dev.078220>
- 45 Du H., Taylor H. S (2015) **The Role of Hox Genes in Female Reproductive Tract Development, Adult Function, and Fertility** *Cold Spring Harb Perspect Med* **6** <https://doi.org/10.1101/cshperspect.a023002>
- 46 Cakmak H., Taylor H. S (2010) **Molecular mechanisms of treatment resistance in endometriosis: the role of progesterone-hox gene interactions** *Semin Reprod Med* **28**:69–74 <https://doi.org/10.1055/s-0029-1242996>
- 47 Taylor H. S., Bagot C., Kardana A., Olive D., Arici A (1999) **HOX gene expression is altered in the endometrium of women with endometriosis** *Hum Reprod* **14**:1328–1331 <https://doi.org/10.1093/humrep/14.5.1328>
- 48 He B., Ni Z. L., Kong S. B., Lu J. H., Wang H. B (2018) **Homeobox genes for embryo implantation: From mouse to human** *Animal Model Exp Med* **1**:14–22 <https://doi.org/10.1002/ame2.12002>
- 49 Illingworth I. M., et al. (2000) **Desmosomes are reduced in the mouse uterine luminal epithelium during the preimplantation period of pregnancy: a mechanism for facilitation of implantation** *Biol Reprod* **63**:1764–1773 <https://doi.org/10.1095/biolreprod63.6.1764>
- 50 Paria B. C., Zhao X., Das S. K., Dey S. K., Yoshinaga K (1999) **Zonula occludens-1 and E-cadherin are coordinately expressed in the mouse uterus with the initiation of implantation and decidualization** *Dev Biol* **208**:488–501 <https://doi.org/10.1006/dbio.1999.9206>
- 51 Potter S. W., Gaza G., Morris J. E (1996) **Estradiol induces E-cadherin degradation in mouse uterine epithelium during the estrous cycle and early pregnancy** *J Cell Physiol* **169**:1–14 [https://doi.org/10.1002/\(SICI\)1097-4652\(199610\)169:1<1::AID-JCP1>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-4652(199610)169:1<1::AID-JCP1>3.0.CO;2-S)
- 52 Grund S., Grummer R (2018) **Direct Cell(-)Cell Interactions in the Endometrium and in Endometrial Pathophysiology** *Int J Mol Sci* **19** <https://doi.org/10.3390/ijms19082227>
- 53 Schatz F., Guzeloglu-Kayisli O., Arlier S., Kayisli U. A., Lockwood C. J (2016) **The role of decidual cells in uterine hemostasis, menstruation, inflammation, adverse pregnancy outcomes and abnormal uterine bleeding** *Hum Reprod Update* **22**:497–515 <https://doi.org/10.1093/humupd/dmw004>
- 54 Evans J., et al. (2016) **Fertile ground: human endometrial programming and lessons in health and disease** *Nat Rev Endocrinol* **12**:654–667 <https://doi.org/10.1038/nrendo.2016.116>
- 55 Stehlin-Gaon C., et al. (2003) **All-trans retinoic acid is a ligand for the orphan nuclear receptor ROR beta** *Nat Struct Biol* **10**:820–825 <https://doi.org/10.1038/nsb979>
- 56 Spitzer T. L., et al. (2012) **Perivascular human endometrial mesenchymal stem cells express pathways relevant to self-renewal, lineage specification, and functional phenotype** *Biol Reprod* **86** <https://doi.org/10.1095/biolreprod.111.095885>

- 57 Yin Y., Haller M. E., Chadchan S. B., Kommagani R., Ma L (2021) **Signaling through retinoic acid receptors is essential for mammalian uterine receptivity and decidualization** *JCI Insight* **6** <https://doi.org/10.1172/jci.insight.150254>
- 58 Chang C (2016) **Agonists and Antagonists of TGF-beta Family Ligands** *Cold Spring Harb Perspect Biol* **8** <https://doi.org/10.1101/cshperspect.a021923>
- 59 Fullerton P. T., Monsivais D., Kommagani R., Matzuk M. M (2017) **Follistatin is critical for mouse uterine receptivity and decidualization** *Proc Natl Acad Sci U S A* **114**:E4772–E4781 <https://doi.org/10.1073/pnas.1620903114>
- 60 Li Q., et al. (2011) **The antiproliferative action of progesterone in uterine epithelium is mediated by Hand2** *Science* **331**:912–916 <https://doi.org/10.1126/science.1197454>
- 61 Filant J., DeMayo F. J., Pru J. K., Lydon J. P., Spencer T. E (2014) **Fibroblast growth factor receptor two (FGFR2) regulates uterine epithelial integrity and fertility in mice** *Biol Reprod* **90** <https://doi.org/10.1095/biolreprod.113.114496>
- 62 Lim W., Bae H., Bazer F. W., Song G (2017) **Stimulatory effects of fibroblast growth factor 2 on proliferation and migration of uterine luminal epithelial cells during early pregnancy** *Biol Reprod* **96**:185–198 <https://doi.org/10.1095/biolreprod.116.142331>
- 63 Russo L. A., et al. (2009) **Regulated expression of matrix metalloproteinases, inflammatory mediators, and endometrial matrix remodeling by 17beta-estradiol in the immature rat uterus** *Reprod Biol Endocrinol* **7** <https://doi.org/10.1186/1477-7827-7-124>
- 64 Tenvergert E., Gillespie M. W., Kingma J., Klasen H (1992) **Abortion attitudes, 1984-1987-1988: effects of item order and dimensionality** *Percept Mot Skills* **74**:627–642 <https://doi.org/10.2466/pms.1992.74.2.627>
- 65 Hayashi K., et al. (2009) **Wnt genes in the mouse uterus: potential regulation of implantation** *Biol Reprod* **80**:989–1000 <https://doi.org/10.1095/biolreprod.108.075416>
- 66 Garcia-Alonso L., et al. (2021) **Mapping the temporal and spatial dynamics of the human endometrium in vivo and in vitro** *Nat Genet* **53**:1698–1711 <https://doi.org/10.1038/s41588-021-00972-2>
- 67 PrabhuDas M., et al. (2015) **Immune mechanisms at the maternal-fetal interface: perspectives and challenges** *Nat Immunol* **16**:328–334 <https://doi.org/10.1038/ni.3131>
- 68 Guimaraes-Young A., Neff T., Dupuy A. J., Goodheart M. J (2016) **Conditional deletion of Sox17 reveals complex effects on uterine adenogenesis and function** *Dev Biol* **414**:219–227 <https://doi.org/10.1016/j.ydbio.2016.04.010>
- 69 Hirate Y., et al. (2016) **Mouse Sox17 haploinsufficiency leads to female subfertility due to impaired implantation** *Sci Rep* **6** <https://doi.org/10.1038/srep24171>
- 70 Wang X., et al. (2018) **SOX17 regulates uterine epithelial-stromal cross-talk acting via a distal enhancer upstream of Ihh** *Nat Commun* **9** <https://doi.org/10.1038/s41467-018-06652-w>
- 71 Hollnagel A., Oehlmann V., Heymer J., Ruther U., Nordheim A (1999) **Id genes are direct targets of bone morphogenetic protein induction in embryonic stem cells** *J Biol Chem* **274**:19838–19845 <https://doi.org/10.1074/jbc.274.28.19838>

- 72 Miyazono K., Miyazawa K (2002) **Id: a target of BMP signaling** *Sci STKE* 2002, pe40 <https://doi.org/10.1126/stke.2002.151.pe40>
- 73 Holloran S. M., et al. (2020) **Reciprocal fine-tuning of progesterone and prolactin-regulated gene expression in breast cancer cells** *Mol Cell Endocrinol* **511** <https://doi.org/10.1016/j.mce.2020.110859>
- 74 Dinh D. T., et al. (2023) **Progesterone receptor mediates ovulatory transcription through RUNX transcription factor interactions and chromatin remodelling** *Nucleic Acids Res* <https://doi.org/10.1093/nar/gkad271>
- 75 Vasquez Y. M., et al. (2018) **FOXO1 regulates uterine epithelial integrity and progesterone receptor expression critical for embryo implantation** *PLoS Genet* **14** <https://doi.org/10.1371/journal.pgen.1007787>
- 76 Shi J. W., et al. (2022) **An IGF1-expressing endometrial stromal cell population is associated with human decidualization** *BMC Biol* **20** <https://doi.org/10.1186/s12915-022-01483-0>
- 77 Suzuki A., et al. (2006) **Nanog binds to Smad1 and blocks bone morphogenetic protein-induced differentiation of embryonic stem cells** *Proc Natl Acad Sci U S A* **103**:10294–10299 <https://doi.org/10.1073/pnas.0506945103>
- 78 Shi X., Bai S., Li L., Cao X (2001) **Hoxa-9 represses transforming growth factor-beta-induced osteopontin gene transcription** *J Biol Chem* **276**:850–855 <https://doi.org/10.1074/jbc.M005955200>
- 79 Shi X., Yang X., Chen D., Chang Z., Cao X (1999) **Smad1 interacts with homeobox DNA-binding proteins in bone morphogenetic protein signaling** *J Biol Chem* **274**:13711–13717 <https://doi.org/10.1074/jbc.274.19.13711>
- 80 Feng X. H., Derynck R (2005) **Specificity and versatility in tgf-beta signaling through Smads** *Annu Rev Cell Dev Biol* **21**:659–693 <https://doi.org/10.1146/annurev.cellbio.21.022404.142018>
- 81 Massague J., Seoane J., Wotton D (2005) **Smad transcription factors** *Genes Dev* **19**:2783–2810 <https://doi.org/10.1101/gad.1350705>
- 82 Derynck R., Budi E. H (2019) **Specificity, versatility, and control of TGF-beta family signaling** *Sci Signal* **12** <https://doi.org/10.1126/scisignal.aav5183>
- 83 Lee D. K., et al. (2010) **Suppression of ERalpha activity by COUP-TFII is essential for successful implantation and decidualization** *Mol Endocrinol* **24**:930–940 <https://doi.org/10.1210/me.2009-0531>
- 84 Vasquez Y. M., et al. (2015) **FOXO1 is required for binding of PR on IRF4, novel transcriptional regulator of endometrial stromal decidualization** *Mol Endocrinol* **29**:421–433 <https://doi.org/10.1210/me.2014-1292>
- 85 Wu M., Chen G., Li Y. P (2016) **TGF-beta and BMP signaling in osteoblast, skeletal development, and bone formation, homeostasis and disease** *Bone Res* **4** <https://doi.org/10.1038/boneres.2016.9>
- 86 van Weerd J. H., Christoffels V. M (2016) **The formation and function of the cardiac conduction system** *Development* **143**:197–210 <https://doi.org/10.1242/dev.124883>

- 87 Whitman M (1998) **Smads and early developmental signaling by the TGFbeta superfamily** *Genes Dev* **12**:2445–2462 <https://doi.org/10.1101/gad.12.16.2445>
- 88 Middlebrook B. S., Eldin K., Li X., Shivasankaran S., Pangas S. A (2009) **Smad1-Smad5 ovarian conditional knockout mice develop a disease profile similar to the juvenile form of human granulosa cell tumors** *Endocrinology* **150**:5208–5217 <https://doi.org/10.1210/en.2009-0644>
- 89 Pangas S. A., et al. (2008) **Conditional deletion of Smad1 and Smad5 in somatic cells of male and female gonads leads to metastatic tumor development in mice** *Mol Cell Biol* **28**:248–257 <https://doi.org/10.1128/MCB.01404-07>
- 90 Rodriguez A., et al. (2016) **SMAD Signaling Is Required for Structural Integrity of the Female Reproductive Tract and Uterine Function During Early Pregnancy in Mice** *Biol Reprod* **95** <https://doi.org/10.1095/biolreprod.116.139477>
- 91 Retting K. N., Song B., Yoon B. S., Lyons K. M (2009) **BMP canonical Smad signaling through Smad1 and Smad5 is required for endochondral bone formation** *Development* **136**:1093–1104 <https://doi.org/10.1242/dev.029926>
- 92 McReynolds L. J., Gupta S., Figueroa M. E., Mullins M. C., Evans T (2007) **Smad1 and Smad5 differentially regulate embryonic hematopoiesis** *Blood* **110**:3881–3890 <https://doi.org/10.1182/blood-2007-04-085753>
- 93 Consortium E. P (2012) **An integrated encyclopedia of DNA elements in the human genome** *Nature* **489**:57–74 <https://doi.org/10.1038/nature11247>
- 94 Chen J., et al. (2014) **Single-molecule dynamics of enhanceosome assembly in embryonic stem cells** *Cell* **156**:1274–1285 <https://doi.org/10.1016/j.cell.2014.01.062>
- 95 Swinstead E. E., et al. (2016) **Steroid Receptors Reprogram FoxA1 Occupancy through Dynamic Chromatin Transitions** *Cell* **165**:593–605 <https://doi.org/10.1016/j.cell.2016.02.067>
- 96 Chen C. H., et al. (2020) **Determinants of transcription factor regulatory range** *Nat Commun* **11** <https://doi.org/10.1038/s41467-020-16106-x>
- 97 Arvey A., Agius P., Noble W. S., Leslie C (2012) **Sequence and chromatin determinants of cell-type-specific transcription factor binding** *Genome Res* **22**:1723–1734 <https://doi.org/10.1101/gr.127712.111>
- 98 Tang S., Cope D. I., Vasquez Y. M., Monsivais D (2022) **BMP/SMAD1/5 Signaling in the Endometrial Epithelium Is Essential for Receptivity and Early Pregnancy** *Endocrinology* **163** <https://doi.org/10.1210/endocr/bqac043>
- 99 Newman A. M., et al. (2019) **Determining cell type abundance and expression from bulk tissues with digital cytometry** *Nat Biotechnol* **37**:773–782 <https://doi.org/10.1038/s41587-019-0114-2>
- 100 Fuentealba L. C., et al. (2007) **Integrating patterning signals: Wnt/GSK3 regulates the duration of the BMP/Smad1 signal** *Cell* **131**:980–993 <https://doi.org/10.1016/j.cell.2007.09.027>

- 101 Labbe E., Letamendia A., Attisano L (2000) **Association of Smads with lymphoid enhancer binding factor 1/T cell-specific factor mediates cooperative signaling by the transforming growth factor-beta and wnt pathways** *Proc Natl Acad Sci U S A* **97**:8358–8363 <https://doi.org/10.1073/pnas.150152697>
- 102 Zavadil J., Cermak L., Soto-Nieves N., Bottinger E. P (2004) **Integration of TGF-beta/Smad and Jagged1/Notch signalling in epithelial-to-mesenchymal transition** *EMBO J* **23**:1155–1165 <https://doi.org/10.1038/sj.emboj.7600069>
- 103 Qiu T., Grizzle W. E., Oelschlager D. K., Shen X., Cao X (2007) **Control of prostate cell growth: BMP antagonizes androgen mitogenic activity with incorporation of MAPK signals in Smad1** *EMBO J* **26**:346–357 <https://doi.org/10.1038/sj.emboj.7601499>
- 104 Luo K., et al. (1999) **The Ski oncoprotein interacts with the Smad proteins to repress TGFbeta signaling** *Genes Dev* **13**:2196–2206 <https://doi.org/10.1101/gad.13.17.2196>
- 105 Pouponnot C., Jayaraman L., Massague J (1998) **Physical and functional interaction of SMADs and p300/CBP** *J Biol Chem* **273**:22865–22868 <https://doi.org/10.1074/jbc.273.36.22865>
- 106 Brosens J. J., Gellersen B (2006) **Death or survival--progesterone-dependent cell fate decisions in the human endometrial stroma** *J Mol Endocrinol* **36**:389–398 <https://doi.org/10.1677/jme.1.02060>
- 107 Yilmaz B. D., Bulun S. E. (2019) **Endometriosis and nuclear receptors** *Hum Reprod Update* **25**:473–485 <https://doi.org/10.1093/humupd/dmz005>
- 108 Janzen D. M., et al. (2013) **Progesterone receptor signaling in the microenvironment of endometrial cancer influences its response to hormonal therapy** *Cancer Res* **73**:4697–4710 <https://doi.org/10.1158/0008-5472.CAN-13-0930>
- 109 MacLean J. A (2022) **2nd & Hayashi K. Progesterone Actions and Resistance in Gynecological Disorders.** *Cells* **11** <https://doi.org/10.3390/cells11040647>

Article and author information

Zian Liao

Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA, Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, 77030, USA, Graduate Program of Genetics and Genomics, Baylor College of Medicine, Houston, TX, 77030, USA, Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA

Suni Tang

Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA, Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA

Kaori Nozawa

Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA, Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA

Keisuke Shimada

Research Institute for Microbial Diseases, Osaka University, Osaka, 565-0871, Japan
ORCID iD: [0000-0003-3739-7163](https://orcid.org/0000-0003-3739-7163)

Masahito Ikawa

Research Institute for Microbial Diseases, Osaka University, Osaka, 565-0871, Japan
ORCID iD: [0000-0001-9859-6217](https://orcid.org/0000-0001-9859-6217)

Diana Monsivais

Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA,
Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA
For correspondence: diana.monsivais@bcm.edu
ORCID iD: [0000-0001-5660-6392](https://orcid.org/0000-0001-5660-6392)

Martin M. Matzuk

Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA,
Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, 77030,
USA, Graduate Program of Genetics and Genomics, Baylor College of Medicine, Houston, TX,
77030, USA, Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA
For correspondence: mmatzuk@bcm.edu
ORCID iD: [0000-0002-1445-8632](https://orcid.org/0000-0002-1445-8632)

Copyright

© 2023, Liao et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Carmen Williams

National Institute of Environmental Health Sciences, Research Triangle Park, United States of America

Senior Editor

Wei Yan

University of California, Los Angeles, Torrance, United States of America

Reviewer #1 (Public Review):**Summary**

Liao et al leveraged two powerful genomics techniques-CUT&RUN and RNA sequencing-to identify genomic regions bound by and activated or inactivated by SMAD1, SMAD5, and the progesterone receptor during endometrial stromal cell decidualization. Additionally, the authors generated novel knock-in HA-SMAD1 and PA-SMAD5 tagged mice to combat antibody issues facing the field, generating a novel model to advance the study of BMP signaling in the female reproductive tract. During decidualization in a murine model, SMAD1/5 are bound to many genomic sites of genes important in decidualization and pregnancy and coregulated responses with progesterone receptor signaling.

Strengths

The authors utilized powerful next generation sequencing and identified important transcriptional mechanisms of SMAD1/5 and PGR during decidualization in vivo.

Weaknesses

None.

Overall, the manuscript and study are well structured and provide critical mechanistic updates on the roles of SMAD1/5 in decidualization and preparation of the maternal endometrium for pregnancy.

<https://doi.org/10.7554/eLife.91434.2.sa2>

Reviewer #2 (Public Review):

Summary:

Liao and colleagues generated tagged SMAD1 and SMAD5 mouse models and identified genome occupancy of these two factors in the uterus of these mice using the CUT&RUN assay. The authors used integrative bioinformatic approaches to identify putative SMAD1/5 direct downstream target genes and to catalog the SMAD1/5 and PGR genome co-localization pattern. The role of SMAD1/5 on stromal decidualization was assayed in vitro on primary human endometrial stromal cells. The new mouse models offer opportunities to further dissect SMAD1 and SMAD5 functions without the limitation from SMAD antibodies, which is significant. The CUT&RUN data further support the usefulness of these mouse models for this purpose.

Strengths:

The strength of this study is the novelty of new mouse models and the valuable cistromic data derived from these mice. Overall the present manuscript is an excellent resource paper for the field of reproductive biology.

Weaknesses:

The weakness of the present version of the manuscript includes the self-limited data analysis approaches such as the proximal promoter based bioinformatic filter and an outdated method on inferring the cell type composition. Evidence was provided for potential associations between SMAD1/5 and other major transcription factors. However, causal effects of SMAD1/5 on the genome occupancy of other major uterine transcription factors were discussed but not experimentally examined in the present manuscript, which is understandable.

For data in Figure 2B, the current manuscript fails to elaborate the common and distinct features between clusters 1 and 3 as well as the biological significance of having two separate clusters for SMAD1. In addition, Figure S1A shows overlapping genome occupancy between SMAD1 and SMAD5, which is not clearly demonstrated in Figure 2B.

For data in Figure 5A, the result description does not provide adequate information to guide readers to full understanding of the data. The biological meaning behind the three PR clusters is not stated nor speculated. Moreover, Figure 5A and Figure S1B are inherently connected but fail to be adequately described in the main text.

<https://doi.org/10.7554/eLife.91434.2.sa1>

Reviewer #3 (Public Review):

Summary:

As SMAD1/5 activities have previously been indistinguishable, these studies provide a new mouse model to finally understand unique downstream activation of SMAD1/5 target genes, a model useful for many scientific fields. Using CUT&RUN analyses with gene overlap

comparisons and signaling pathway analyses, specific targets for SMAD1 versus SMAD5 were compared, identified, and interpreted. These data validate previous findings showing strong evidence that SMADs directly govern critical genes required for endometrial receptivity and decidualization, including cell adhesion and vascular development. Further, SMAD targets were overlapped with progesterone receptor binding sites to identify regions of potential synergistic regulation of implantation. The authors report strong correlations between progesterone receptor and SMAD1/5 direct targets to cooperatively promote embryo implantation. Finally, the authors validated SMAD1/5 gene regulation in primary human endometrial stromal cells. These studies provide a data-rich survey of SMAD family transcription, defining its role as a governor of early pregnancy.

Strengths:

This manuscript provides a valuable survey of SMAD1/5 direct transcriptional events at the time of receptivity. As embryo implantation is controlled by extensive epithelial to stromal molecular crosstalk and hormonal regulation in space and time, the authors state a strong, descriptive narrative defining how SMAD1/5 plays a central role at the site of this molecular orchestration. The implementation of cutting-edge techniques and models and simple comparative analyses provide a straightforward, yet elegant manuscript.

Although the progesterone receptor exists as a major regulator of early pregnancy, the authors have demonstrated clear evidence that progesterone receptor with SMAD1/5 work in concert to molecularly regulate targets such as Sox17, Id2, Tgfb β 2, Runx1, Foxo1 and more at embryo implantation. Additionally, the authors pinpoint other critical transcription factor motifs that work with SMADs and the progesterone receptor to promote early pregnancy transcriptional paradigms.

Weaknesses:

Although a wonderful new tool to ascertain SMAD1 versus SMAD5 downstream signaling, the importance of these factors in governing early pregnancy is not novel. Furthermore, functional validation studies are needed to confirm interactions at promoter regions. Additionally, the authors presume that all overlapped genes are shared between progesterone receptor and SMAD1/5, yet some peak representations do not overlap. Although, transcriptional activation can occur at the same time, they may not occur in the same complex. Thus, further confirmation of these transcriptional events is warranted.

Since whole murine uterus was used for these studies, the specific functions of SMAD1/5 in the stroma versus the epithelium (versus the myometrium) remain unknown. Further work is needed to delineate binding and transcriptional activation of SMAD1/5 and the progesterone receptor in the uterine compartments.

There are asynchronous gene responses in the SMAD1/5 ablated mouse model compared to the siRNA-treated human endometrial stromal cells. These differences can be confounding. Further investigation is needed to understand the meaning of these differences and as they relate to the entire SMAD transcriptome.

<https://doi.org/10.7554/eLife.91434.2.sa0>

Author Response

We thank you for your careful review of our manuscript and helpful comments and suggestions. We have carefully considered each point and have addressed them by adding changes to the manuscript and figures. The text below detailed our responses and edits.

Reviewer #1 (Public Review):

Summary:

Liao et al leveraged two powerful genomics techniques-CUT&RUN and RNA sequencing-to identify genomic regions bound by and activated or inactivated by SMAD1, SMAD5, and the progesterone receptor during endometrial stromal cell decidualization.

Strengths:

The authors utilized powerful next generation sequencing and identified important transcriptional mechanisms of SMAD1/5 and PGR during decidualization in vivo.

Weaknesses:

Overall, the manuscript and study are well structured and provide critical mechanistic updates on the roles of SMAD1/5 in decidualization and preparation of the maternal endometrium for pregnancy. Please consider the following to improve the manuscript:

- *Figure 4: A and C show bar graphs, not histograms. Please alter this phrasing.*

Figure legends were adjusted as suggested.

- *What post hoc test was performed on qPCR analyses? (Figure 6). It is evident that any assumptions of equal variance need to be negated due to the wide dispersion in experimental response invalidating the assumptions of a one-way ANOVA.*

Yes, a Tukey's post hoc test was performed on the qPCR analyses. To address the reviewer's question regarding equal variance, normality of the dataset was examined by D'agostino & Pearson test in GraphPad Prism. The data demonstrated a normal distribution pattern, thus justifying the one-way ANOVA test.

- *Figure 6: what data points are plotted? Are these technical replicates from individual wells or qPCR technical replicates?*

The dataset represents three technical and three biological data points.

- *Figure 6: Consider changing graph colors to increase visibility of error bars and data points.*

Thank you for this suggestion. The colors of the error bars in Figure 6 have been changed to increase visibility. Additionally, different shapes have been utilized to distinguish between different groups.

- *Figure 6 legend: no histograms are shown in this figure. Refer to all gene names utilizing proper nomenclature and conventions (gene names should be italicized).*

The legend was adjusted as suggested with the correct nomenclature implemented.

- *qPCR analyses: qPCR normalization should be done to at least two internal control genes, preferably three according to the MIQE guidelines (PMID: 19246619).*

As suggested, we have performed additional qPCR analysis with normalization done to three internal controls.

- *Supplement figure 2: graphs are bar graphs, not histograms.*

The legends have been changed as suggested.

Reviewer #2 (Public Review):

Summary:

Liao and colleagues generated tagged SMAD1 and SMAD5 mouse models and identified genome occupancy of these two factors in the uterus of these mice using the CUT&RUN assay. The authors used integrative bioinformatic approaches to identify putative SMAD1/5 direct downstream target genes and to catalog the SMAD1/5 and PGR genome co-localization pattern. The role of SMAD1/5 on stromal decidualization was assayed in vitro on primary human endometrial stromal cells. The new mouse models offer opportunities to further dissect SMAD1 and SMAD5 functions without the limitation from SMAD antibodies, which is significant. The CUT&RUN data further support the usefulness of these mouse models for this purpose.

Strengths:

The strength of this study is the novelty of new mouse models and the valuable cistromic data derived from these mice.

Weaknesses:

The weakness of the present version of the manuscript includes the self-limited data analysis approaches such as the proximal promoter based bioinformatic filter and a missed opportunity to investigate the role of SMAD1/5 on determining the genome occupancy of major uterine transcription regulators.

Thank you for the comments. We addressed the limitation of the promoter-based analysis in the discussion and pointed out the possibility of analyzing additional genomics features (Lines 548-551). Based on the suggestions, we also included an analysis in which we compared SMAD1/5 binding activities in this study to known major uterine transcription regulators' binding activities (namely, SOX17 and NR2F2) using published ChIP-seq data in the mouse uterus. Results from this analysis are discussed in Lines 426-436. Content from the adjusted manuscript is copied below.

Lines 548-551:

“From pathway enrichment analysis, we demonstrate that genes with SMAD1/5 and PR bound at the promoter regions are enriched for key pathways in directing the decidualization process, such as WNT and relaxin signaling pathways. Future studies can benefit from analyzing binding events beyond the promoter regions.”

Lines 426-436:

“To further evaluate the key roles of SMAD1/5 as major uterine transcription regulators, we cross-compared the genomic binding sites of SMAD1/5 with known key transcription factors, namely aforementioned SOX17 (Supplement Figure 1E), as well as NR2F2 (Supplement Figure 1F), an essential regulator of hormonal response, using our CUT&RUN data sets and published mouse uterine SOX17 and NR2F2 ChIP-seq data sets (GSE118328, GSE232583). Among the annotated genes, 5402 genes are shared between SMAD1/5 and SOX17, and 1922 genes are shared between SMAD1/5 and NR2F2. Such observations indicate a potential co-regulatory mechanism between SMAD1/5 and other key uterine transcription factors in maintaining appropriate uterine functions. Overall, our analyses demonstrate that the transcriptional activity of SMAD1, SMAD5, and PR coordinate the expression of key genes required for endometrial receptivity and decidualization.”

Reviewer #3 (Public Review):

Summary:

As SMAD1/5 activities have previously been indistinguishable, these studies provide a new mouse model to finally understand unique downstream activation of SMAD1/5 target genes, a model useful for many scientific fields. Using CUT&RUN analyses with gene overlap comparisons and signaling pathway analyses, specific targets for SMAD1 versus SMAD5 were compared, identified, and interpreted. These data validate previous findings showing strong evidence that SMADs directly govern critical genes required for endometrial receptivity and decidualization, including cell adhesion and vascular development. Further, SMAD targets were overlapped with progesterone receptor binding sites to identify regions of potential synergistic regulation of implantation. The authors report strong correlations between progesterone receptor and SMAD1/5 direct targets to cooperatively promote embryo implantation. Finally, the authors validated SMAD1/5 gene regulation in primary human endometrial stromal cells. These studies provide a data-rich survey of SMAD family transcription, defining its role as a governor of early pregnancy.

Strengths:

This manuscript provides a valuable survey of SMAD1/5 direct transcriptional events at the time of receptivity. As embryo implantation is controlled by extensive epithelial to stromal molecular crosstalk and hormonal regulation in space and time, the authors state a strong, descriptive narrative defining how SMAD1/5 plays a central role at the site of this molecular orchestration. The implementation of cutting-edge techniques and models and simple comparative analyses provide a straightforward, yet elegant manuscript.

Although the progesterone receptor exists as a major regulator of early pregnancy, the authors have demonstrated clear evidence that progesterone receptor with SMAD1/5 work in concert to molecularly regulate targets such as Sox17, Id2, Tgfb2, Runx1, Foxo1 and more at embryo implantation. Additionally, the authors pinpoint other critical transcription factor motifs that work with SMADs and the progesterone receptor to promote early pregnancy transcriptional paradigms.

Weaknesses:

Although a wonderful new tool to ascertain SMAD1 versus SMAD5 downstream signaling, the importance of these factors in governing early pregnancy is not novel. Furthermore, functional validation studies are needed to confirm interactions at promoter regions. Additionally, the authors presume that all overlapped genes are shared between progesterone receptor and SMAD1/5, yet some peak representations do not overlap. Although, transcriptional activation can occur at the same time, they may not occur in the same complex. Thus, further confirmation of these transcriptional events is warranted.

Thank you for the review; we appreciate these valuable comments. Although we used an overlap approach to investigate the gene regulatory networks between SMAD1/5 and PR at the gene level, we functionally validated the regulatory effect in an in vitro decidualization model using a qPCR approach. We acknowledge that gene activations may not occur at the exact same complex, but functional validation screenings at the promoter level are beyond the scope of the study. However, we added the discussion about the possibility of proposed investigations in Lines 553-558. Our current dataset and validation studies support our conclusions with robust evidence. Content from Lines 553-558 is copied below.

Lines 553-558: “In this study, we determined the overlapped transcriptional control between SMAD1/5 and PR at the gene level, and functionally validated the regulatory effect at the transcript level in a human stromal cell decidualization model. While we observe a subset of peak representations that do not overlap at the base pair level in the promoter regions, future functional screenings at the promoter level, such as luciferase reporter assays to assess transcriptional co-activation by SMAD1/5 and PR, will advance this study.”

• Since whole murine uterus was used for these studies, the specific functions of SMAD1/5 in the stroma versus the epithelium (versus the myometrium) remain unknown. Specific roles for SMAD1/5 in the uterine stroma and epithelial compartments still need to be examined. Also, further work is needed to delineate binding and transcriptional activation of SMAD1/5 and the progesterone receptor in stromal versus epithelial uterine compartments.

Thank you for the comments. Indeed, our study was performed in the whole mouse uterus, which includes stroma, epithelium and myometrium. Our previous data shows that nuclear SMAD1/5 are localized to both the stroma and epithelium in the decidua zone during the decidualization process at 4.5 dpc (PMID:34099644). Published in vivo studies also demonstrate the essential role of SMAD1/5 in the uterine epithelium and stroma compartments, respectively (PMIDs:35383354/27335065/17967875). Although we believe the binding/transcriptional activation of SMAD1/5 and PR occurs in both compartments based on the mouse phenotypic data, opportunities for further compartment-specific analysis were granted and discussion regarding such investigations was added (Lines 501-513). Content from Lines 501-513 is copied below.

Lines 501-513:

“Published studies have shown that nuclear SMAD1/5 localize to the stroma and epithelium during the decidualization process at 4.5dpc during the window of implantation. Conditional deletion of SMAD1/5 exclusively in the uterine epithelium using lactoferrin-icre (Ltf-icre) results in severe subfertility due to impaired implantation and decidual development. Conditional deletion of SMAD1/5/4 exclusively in the cells from mesenchymal lineage (including uterine stroma) using anti-Mullerian hormone type 2 receptor cre (Amhr2-cre) results in infertility with defective decidualization. Given the essential roles of SMAD1/5 in both stroma and epithelium identified by previous studies, we believe that transcriptional co-regulation by SMAD1/5 and PR reported here using the whole uterus validates a relationship between SMAD1/5 and PR in both the stromal and epithelial compartments. However, it does not rule out the potential coregulation of SMAD1/5 and PR in the myometrium, immune cells, and/or endothelium, given that whole uterus was used. The specific transcriptional evaluations of SMAD1/5 in the stroma versus the epithelium would require future single-cell sequencing (i.e., digital cytometry) and/or spatial transcriptomic analysis.”

• There are asynchronous gene responses in the SMAD1/5 ablated mouse model compared to the siRNA-treated human endometrial stromal cells. These differences can be confounding, and more clarity is required in understanding the meaning of these differences and as they relate to the entire SMAD transcriptome.

Thank you for the comments. From the mouse models with SMAD1/5 conditional deletions, we observed phenotypic defects at 4.5 dpc, which is the beginning of decidualization in the mouse. Our study used human endometrial stromal cells as a model to validate our findings functionally, aiming to mimic the specific time point during decidualization. Differences between the two models may arise from the strategy used to perturb SMAD1/5; in the mouse, a complete knockout of SMAD1/5 was used, resulting in failed decidualization, while the human endometrial stromal cells used an siRNA knockdown approach, which decreased the

potential for decidualization. As such, this information needs to be considered when evaluating genome-wide effects on the transcriptome. We added a discussion of this point to Lines 564-572. Content from Lines 564-572 is copied below.

Lines 564-572:

“Since mice only undergo decidualization upon embryo implantation whilst human stromal cells undergo cyclic decidualization in each menstrual cycle in response to rising levels of progesterone, asynchronous gene responses may occur in comparison between mouse models and human cells. However, cellular transformation during decidualization is conserved between mice and humans, which makes findings in the mouse models a valuable and transferable resource to be evaluated in human tissues. Accordingly, our functional validation studies were performed using human endometrial stromal cells induced to decidualize in vitro for four days, which models the early phases of decidualization. Additional transcriptomic studies of the SMAD1/5 perturbations in human endometrial stromal cells will be of great resource in understanding the entire SMAD1/5 regulomes in humans.”

Reviewer #1 (Recommendations For The Authors):

- *Minor grammatical errors requiring attention such as inserting punctuation at the end of sentences and including figure legends prior to the end of sentence punctuation.*

Thanks for the comments. Additional proofreading was conducted for the revision.

Reviewer #2 (Recommendations For The Authors):

1. *Between SMAD1 and SMAD5, does losing one SMAD affect the other SMAD's genome occupancy?*

Thanks for the comments. Based on the mouse phenotypic data that conditional deletion of SMAD1 in the uterus does not affect female fertility, while conditional deletion of SMAD5 leads to subfertility, and conditional deletion of both SMAD1 and SMAD5 leads to complete infertility. We believe losing one SMAD will affect the other SMAD's genome occupancy. This point is discussed in Lines 514-517, with contents copied below.

Lines 514-517: “Although our studies herein confirm that SMAD1 and SMAD5 proteins have distinct transcriptional regulatory activities, our previous studies demonstrated that while SMAD5 can functionally replace SMAD1, SMAD1 cannot replace SMAD5 in the uterus. How this epistatic relationship is established in a tissue-specific manner still needs to be determined by further biochemical investigations.”

1. *In light of SMAD1/5 and PGR co-occupied cis-acting elements and coregulating uterine transcriptome, does loss of SMAD1/5 alter the PGR and ESR1 genome occupancy?*

Thanks for the comments. In the SMAD1/5 double conditional knockout mice, we observe the hyposensitivity towards progesterone and unopposed estrogen responses. We hypothesize that loss of SMAD1/5 alters PR genome occupancy and subsequently ER genome occupancy is altered as a secondary effect. To functionally address this question, genomic profiling studies need to be performed in the SMAD1/5 knockout mice, and, ideally, also performed in the PR knockout mice. However, such large-scale studies are beyond the scope of the current study and will not affect our conclusions under physiological conditions. We did include additional discussion regarding this comment in Lines 551-553, with the contents copied below.

Lines 551-553: “Profiling the PR genome occupancy in the SMAD1/5 deficient mice would provide an interesting perspective to reevaluate the major regulatory roles of SMAD1/5 in mediating uterine transcriptomes.”

| 1. *In terms of investigating the impact of SMAD1/5 on cell type composition, perhaps the digital cytometry approach (e.g., PMID: 31061481) could provide unbiased inferences.*

Thank you for the comments. We included expression analysis of a subset of SMAD1/5 direct target genes over different uterine compartments (Figure 4E). We also added the discussion of the opportunities for further compartment-specific analysis, including but not limited to the digital cytometry approach in Lines 506-513, with the contents copied below.

Line 506-513:

“Given the essential roles of SMAD1/5 in both stroma and epithelium identified by previous studies, we believe that the transcriptional co-regulatory roles of SMAD1/5 and PR reported here using the whole uterus validates a relationship between SMAD1/5 and PR in both the stromal and epithelial compartments. However, it does not rule out potential co-regulatory roles of SMAD1/5 and PR in the myometrium, immune cells, and/or endothelium, given that whole uterus was used. The specific transcriptional evaluations of SMAD1/5 in the stroma versus the epithelium would require future single-cell sequencing (i.e., digital cytometry) and/or spatial transcriptomic analysis.”

| 1. *The limitation of focusing on the promoter occupied SMADs should be discussed.*

Additional discussion of the limitation of focusing on the promoter regions was added in Lines 548-551, with contents copied below.

Lines 548-551:

“From pathway enrichment analysis, we demonstrate that genes with SMAD1/5 and PR bound at the promoter regions are enriched for key pathways in directing the decidualization process, such as WNT and relaxin signaling pathways. Future studies can benefit from analyzing binding events beyond the promoter regions.”

| 1. *Methods: The reagent and the condition for PGR CUT&RUN is missing.*

Information added in Line 153.

| 1. *Line 260: Please clarify the statement of "suggesting the transcriptional of PR depends on BMP/SMAD1/5 signaling".*

Thanks for the suggestion. The sentence was rephrased to (Lines 258-261) “Our previous studies revealed that conditional ablation of SMAD1 and SMAD5 in the uterus decreased P4 response during the peri-implantation period, suggesting that the transcriptional activities of PR depend on BMP/SMAD1/5 signaling.”

| 1. *Line 280-289: This statement belongs to the discussion section.*

The statement was moved as suggested.

| 1. *Figure 4E is not cited in the result section.*

Figure 4E was cited in the results section in the revised version. (Line 386)

1. *Figures 3C, 3D, 3E, 3F, 5B and 5D: please include the full lists in the supplemental data so that labs with limited bioinformatic capabilities could use these findings to facilitate scientific discovery.*

Data regarding the aforementioned figures were included in Supplement Tables 3-8 and Supplement Files 1-2.

1. *Figure 2B and Figure 5A: the heatmaps without further grouping on common and distinct genome occupancy among assayed factors provided minimum useful information. Please reconsider the presentation format in order to deliver more meaningful results.*

Figure 2B and Figure 5A were replotted with clustering using the k-means algorithm. Methods and legends were updated accordingly.

Reviewer #3 (Recommendations For The Authors):

To delineate specific roles for SMAD1/5 in the uterine stroma and epithelial compartments, methods such as single cell sequencing or spatial transcriptomic analysis may be warranted.

The manuscript now includes the discussion of future opportunities in investigating the roles of SMAD1/5 in different uterine compartments using single-cell sequencing and/or spatial transcriptomic analysis (Lines 498-513), with contents copied below.

Lines 498-513:

“Our studies also examined the role of SMAD1/5 in mediating progesterone responses at the genomic and transcription levels. Similarly, our analysis was based on data sets generated from the whole mouse uterus, which contains multiple compartments of the uterine structures, including but not limited to epithelium and stroma. Published studies have shown that nuclear SMAD1/5 localize to the stroma and epithelium during the decidualization process at 4.5 dpc, during the window of implantation. Conditional deletion of SMAD1/5 exclusively in the uterine epithelium using lactoferrin-icre (Ltf-icre) results in severe subfertility due to impaired implantation and decidual development. Conditional deletion of SMAD1/5/4 exclusively in the cells from mesenchymal lineage (including uterine stroma) using anti-Mullerian hormone type 2 receptor cre (Amhr2-cre) results in infertility with defective decidualization. Given the essential roles of SMAD1/5 in both stroma and epithelium identified by previous studies, we believe that the transcriptional co-regulatory roles of SMAD1/5 and PR reported here using the whole uterus validates a relationship between SMAD1/5 and PR in both the stromal and epithelial compartments. However, it does not rule out potential co-regulatory roles of SMAD1/5 and PR in the myometrium, immune cells, and/or endothelium, given that whole uterus was used. The specific transcriptional evaluations of SMAD1/5 in the stroma versus the epithelium would require future single-cell sequencing (i.e., digital cytometry) and/or spatial transcriptomic analysis.”