

Sympathetic Motor Neuron Dysfunction is a Missing Link in Age-Associated Sympathetic Overactivity

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

November 2, 2023 (this version)

Posted to bioRxiv

September 29, 2023

Sent for peer review

August 29, 2023

Lizbeth de la Cruz, Derek Bui, Claudia M. Moreno, Oscar Vivas 

Department of Physiology and Biophysics, University of Washington, Seattle, WA • Howard Hughes Medical Institute • Department of Pharmacology, University of Washington, Seattle, WA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Overactivity of the sympathetic nervous system is a hallmark of aging. The cellular mechanisms behind this overactivity remain poorly understood, with most attention paid to likely central nervous system components. In this work, we hypothesized that aging also affects the function of motor neurons in the peripheral sympathetic ganglia. To test this hypothesis, we compared the electrophysiological responses and ion-channel activity of neurons isolated from the superior cervical ganglia of young (12 weeks), middle-aged (64 weeks), and old (115 weeks) mice. Additionally, we assessed whether rapamycin, an anti-aging treatment, reverses the age-related changes in sympathetic motor neurons. These approaches showed that aging does impact the intrinsic properties of sympathetic motor neurons, increasing spontaneous and evoked firing responses. A reduction of KCNQ channel currents emerged as a major contributor to age-related hyperexcitability. The administration of rapamycin in food for 12 weeks in middle-aged mice partially reverted the KCNQ current reduction and hyperexcitability associated with age. Thus, it is essential to consider the effect of aging on motor components of the sympathetic reflex as a crucial part of the mechanism involved in sympathetic overactivity. Further, our data suggest that rapamycin's beneficial anti-aging effects may be partly attributed to its potential to impact sympathetic nervous system components, providing novel insights into therapeutic strategies for age-related conditions.

eLife assessment

This study presents **valuable** observations indicating that the electrophysiological excitability of cultured sympathetic motor neurons progressively increase during aging, and are inversely correlated with the magnitude of KCNQ currents. The alterations in membrane excitability are broadly relevant for those interested in understanding how the nervous system changes during aging. While the data as a whole are **solid** in showing that the excitability of sympathetic neurons increases in neurons cultured from older mice, the mechanism of the underlying changes and in vivo relevance is **incomplete**.

Introduction

This study focuses on the mechanisms underlying the deterioration of the sympathetic nervous system as we age. With advancing age, the sympathetic nervous system tends to become overactive, evidenced by increased electrical activity in sympathetic nerves [Hart et al. \(2009\)](#); [Ito et al. \(1986\)](#); [Narkiewicz et al. \(2005\)](#); [Wallin \(1978\)](#). This overactivity, observed in humans and animal models, leads to heightened release of norepinephrine over organs [Esler et al. \(1995\)](#); [Goldstein et al. \(1983a,b\)](#); [Veith et al. \(1986\)](#); [Ziegler et al. \(1976\)](#), triggering compensatory processes and cellular deterioration [Hogikyan and Supiano \(1994\)](#). Understanding the physiology and pathophysiology of this age-associated overactivity is crucial, as many common age-driven diseases are linked to sympathetic overactivity. For instance, dysregulation of norepinephrine release is associated with age-related hypertension and arrhythmias. However, the neuronal mechanisms underlying the overactivity of the sympathetic nervous system remain to be elucidated.

During execution of sympathetic reflexes, three essential cellular components of the nervous system come into play: sensory neurons for detecting and transmitting signals of internal conditions, neurons of the sympathetic nuclei in the brain for integrating information and responding to internal changes, and the sympathetic motor neurons in the ganglia for receiving and delivering information to target organs (**Figure 1**). Age-associated overactivity of the sympathetic nervous system has been predominantly ascribed to changes in the central system, specifically focusing on the hypothalamic and brainstem nuclei. In the paraventricular nucleus of the hypothalamus, the activation of glutamatergic neurons through leptin pathways has emerged as a significant contributor to sympathetic overactivity [Shi et al. \(2015, 2020\)](#). At the same time, glial senescence and subsequent inflammatory mechanisms in brainstem nuclei also have been linked to sympathetic nervous system overactivity during aging [Balasubramanian et al. \(2021\)](#). However, one key question remains: Is age-related sympathetic overactivity limited solely to changes in the central system?

Aging impacts the morphology and calcium responses of sympathetic motor neurons, the peripheral component of the sympathetic reflex. Notably, sympathetic innervation to the spleen and lymph nodes reduces, while thymus innervation increases with age [Madden et al. \(1997\)](#). Similarly, innervation to vascular smooth muscle changes with age, and the extent and implications of these changes depend on the specific target and neuropeptide content [Andrews \(1996\)](#); [Andrews and Cowen \(1994\)](#); [Burnstock \(1990\)](#). Additionally, aging affects the function of sympathetic motor neurons, as evidenced by altered calcium responses. Old sympathetic motor neurons display reduced expression of SERCA pumps, an essential component for calcium transport into the endoplasmic reticulum [Buchholz et al. \(2007\)](#); [Pottorf et al. \(2000\)](#). The results of these experiments suggest that aging has a distinct influence on the intrinsic function of sympathetic motor neurons, irrespective of any changes in the central system. As a result, the main objective of our study was to examine directly the impact of aging on the intrinsic membrane electrical properties of sympathetic motor neurons (**Figure 1**) and to test if rapamycin, an anti-aging treatment, reverses the age-related changes.

Our work was structured around four main objectives: 1) To standardize the isolation of sympathetic motor neurons from adult to old age in mice, 2) To compare the spontaneous and evoked electrical activity of sympathetic motor neurons at three life stages, 3) To identify potential molecular candidates underlying the observed altered electric properties and neuronal activity, 4) To evaluate if age-related alterations in the electric activity and ion channels are reverted by rapamycin.




Figure 1.

Components of the sympathetic reflex.

Schematic of the components of the sympathetic autonomic reflex: Sensory neurons send information to the central component (pre-ganglionic neurons), where information is processed. Then, sympathetic motor neurons receive information from pre-ganglionic neurons to transmit it to the target organ. This research was focused on evaluating the age-related changes in the function of sympathetic motor neurons.

Sympathetic motor neurons from old mice are healthy in culture

Our first goal was to assess the viability of neurons isolated from young adult and old mice. We isolated sympathetic motor neurons enzymatically from the left and right superior cervical ganglia (SCG) of young adults (12 weeks old) and old animals (115 weeks old, **Figure 2A**). Single neurons exhibited no visible neurite growth before 18 hours (**Figure 2B**, left); however, after 72 hours in culture, both young and old sympathetic motor neurons showed evident neurite growth (**Figure 2B**, right). The mean diameter of the cell soma was found to be similar in young ($19.7 \pm 0.6 \mu\text{m}$) and old ($19.6 \pm 0.6 \mu\text{m}$) neurons after 24 and 72 hours in culture (**Figure 2C**). The dendritic arborization after 24 hours in culture was comparable between young ($37 \pm 2 \mu\text{m}^2$) and old ($42 \pm 3 \mu\text{m}^2$) neurons. However, after 72 hours in culture, the dendritic arborization in old neurons ($105 \pm 7 \mu\text{m}^2$) was more extensive compared to that of young neurons ($74 \pm 5 \mu\text{m}^2$) (**Figure 2D**). We continued the culture for seven days and noted that the neurites grew until they contacted other neurons in the same dish. Furthermore, glial cells became apparent at this stage in cultures of young and old cells. The ability of neurons to regenerate neurites and contact each other was considered an indicator of viability and health. We ruled out the possibility that any functional differences between neurons from animals of different ages were due to potential damage to old neurons during the isolation and culture process. From now on, we report electrical measurements on neurons that had been incubated for approximately just 12–18 h after isolation.

Old sympathetic motor neurons fire action potentials spontaneously

To investigate whether aging alters the function of sympathetic motor neurons, we first compared the spontaneous activity and passive electrical properties of neurons isolated from mice at different ages: 12 weeks (young), 64 weeks (middle age), and 115 weeks (old). For reference, we also provide comparable human ages (**Figure 3A**). Spontaneous activity was recorded using the current clamp modality without applying any holding or current stimulus. Electrical access to the cell was obtained using the perforated patch technique (See Methods for details). Young neurons had a resting membrane potential (RMP) of $-64 \pm 1 \text{ mV}$ and very little spontaneous activity, which aligns with the expected behavior of motor neurons that respond to presynaptic commands (**Figure 3B–C**). Only 3% exhibited spontaneous firing (**Figure 3D**). In contrast, middle-aged neurons showed an RMP of $-58 \pm 1 \text{ mV}$, and 37% of these neurons displayed spontaneous firing (**Figure 3B–D**). The depolarized RMP and the increased number of neurons spontaneously firing persisted in old neurons, where the RMP was $-54 \pm 1 \text{ mV}$ and 58% of neurons exhibited spontaneous firing (**Figure 3B–D**). Despite the higher percentage of old neurons firing spontaneously compared to middle-aged neurons, the mean firing frequency in old neurons (60 ± 20 action potentials (APs)/minute) was significantly lower than that observed in middle-aged neurons (177 ± 41 APs/minute; **Figure 3E**).

We measured the input resistance, another passive membrane property, in cells polarized to a potential of -65 mV by injecting a negative holding current that did not exceed -50 pA . Cells requiring larger holding currents were excluded. From the response to additional negative current pulses (-40 to 0 pA , **Figure 3F**) we calculated the input resistance ($\Delta V/\text{injected current}$). At these membrane potentials, without activation of voltage-activated ion channels, the change in voltage is proportional to the injected current. Input resistance was not affected by age. Young adult neurons ($0.8 \pm 0.09 \text{ G}\Omega$), middle-aged neurons ($1.1 \pm 0.06 \text{ G}\Omega$), and old neurons ($1.1 \pm 0.11 \text{ G}\Omega$) exhibited similar input resistance (**Figure 3G**). In conclusion, our findings suggest that aging leads to a gradual depolarization of the RMP without significant alteration of the input resistance.




Figure 2.

Sympathetic motor neurons from old mice are healthy in culture.

(A) Diagram of the experimental approach: Sympathetic motor neurons were isolated from the superior cervical ganglia (SCG) of 12- and 115-week-old mice. The ganglia did not show morphological differences between ages. (B) Differential Interference Contrast images of sympathetic motor neurons in culture for 18 h and 72 h. Right images at 72 h outline the dendritic area measured. w, weeks. (C) Comparison of soma diameter between neurons isolated from 12 or 115 weeks of age at two-time points in culture. Orange circles showed single cells from 12-week-old mice while purple triangles show single cells from 115-week-old mice. (D) Comparison of the total dendritic area as a proxy for neurite regeneration in culture. Data points are from N = 3 animals, n = 38 cells, from 12 weeks old, and N = 3 animals, n = 39 cells, from 115 weeks old. p-values are shown at the top of the graphs. Red values indicate p-values < 0.05 while black values indicate p-values > 0.05.




Figure 3.

Sympathetic motor neurons from old mice fire action potentials spontaneously.

(A) Schematic of ages in mice, and equivalent in humans, that were used to compare the functional responses of sympathetic motor neurons. (B) Representative membrane potential recordings of spontaneous activity from neurons isolated from 12-, 64-, and 115-week-old mice. (C) Comparison of the Resting Membrane Potential (RMP) between different ages. (D) Comparison of percentage of silent and firing neurons between different ages. (E) Comparison of the number of action potentials (AP) fired spontaneously in one minute between different ages. (F) Top: Representative passive responses to hyperpolarizing stimuli from neurons isolated from 12-, 64-, and 115-week-old mice. Blue traces correspond to 0 pA injection, red traces to -10 pA, and black traces to -40 pA. Bottom: voltage-current relationship of top recordings. (G) Comparison of the input resistance between different ages. Data points are from N = 4 animals, n = 35 cells, from 12-week old, N = 6 animals, n = 65 cells, from 64-week old, and N = 4 animals, n = 32 cells, from 115-week old. p-values are shown at the top of the graphs. Red p-values indicate p-values < 0.05, while black p-values indicate p-values > 0.05.

Sympathetic motor neurons from old mice are more responsive to electrical stimulation

During sympathetic reflexes, the peripheral motor neurons increase their firing frequency in response to commands from the central sympathetic nuclei. These commands can be mimicked as direct electrical stimulation, as shown in [Figure 4A](#). To test whether aging increases the sensitivity of sympathetic motor neurons to electrical stimulation, neurons were injected with 1-second current steps of amplitude ranging from 10 pA to 100 pA starting from a potential of -65 mV ("0 pA current") ([Figure 4B](#)). Young neurons exhibited a characteristic firing pattern demonstrated in the left panel of [Figure 4C](#). With injections of 10 pA, no AP was elicited (red trace), and with a 40 pA stimulus lasting 1 second, only one AP was evoked (black trace), illustrating the phenomenon known as spike adaptation that is typical of young sympathetic motor neurons. In contrast, both middle-aged and old neurons displayed a response. A 10 pA stimulus was sufficient to elicit APs in middle-aged and old neurons ([Figure 4D-E](#)), and in middle-aged neurons, the 40 pA stimulus induced the firing of eleven APs in a typical example ([Figure 4D](#)), and similarly, in old neurons, it led to the firing of ten APs ([Figure 4E](#)).

To assess the impact of age on the excitability of sympathetic motor neurons in a population sample, we compared stimulus-response curves, rheobase values (minimum current needed to elicit at least one AP), and the number of APs elicited with the maximum stimulus. Our results revealed that middle-aged and old neurons fired more APs with each stimulus ([Figure 4F](#)) than young neurons. Older neurons exhibited a reduced threshold for eliciting APs (64 weeks: 20 ± 2 pA; 115 weeks: 23 ± 5 pA) compared to young neurons (34 ± 4 pA, [Figure 4G](#)). Furthermore, older neurons fired more APs with a 100 pA current injection (64 weeks: 15 ± 1 APs; 115 weeks: 15 ± 2 APs) compared to young neurons (12 weeks: 7 ± 1 APs, [Figure 4H](#)). These findings support the concept that aging leads to increased excitability of sympathetic motor neurons.

Analysis of neuronal subpopulations

Sympathetic motor neurons display stereotyped distinct repetitive firing patterns, which classify them into three categories: tonic (Class I), phasic (Class II), and adapting (Class III) [Kim et al. \(2019\)](#); [Malin and Nerbonne 2000](#), 2001); [Springer et al. \(2015\)](#). In our experiments, we classified cells based on their responses to a supra-threshold current injection (20 pA above their rheobase), ensuring that the firing response was not saturated. Representative traces of firing patterns from middle-aged neurons are illustrated in [Figure 5A](#). The increased number of APs at maximal stimulus compared with 20 pA above the rheobase, shows that the firing response was not saturated during classification ([Figure 5B](#)). In agreement with previous reports, subpopulations showed differences in frequency-stimulus curves ([Figure 5C](#)). Also, tonic cells have a more depolarized RMP ([Figure 5D](#), RMP_{tonic} = -53 ± 2 mV, RMP_{phasic} = -61 ± 1 mV, RMP_{adapting} = -60 ± 2 mV), while adapting neurons have a lower input resistance ([Figure 4D](#), tonic = 1.13 ± 0.08 G Ω , Phasic = 1.16 ± 0.09 G Ω , Adapting = 0.55 ± 0.18 G Ω). In general, the responses from the three neuronal subpopulations are consistent with previous reports.

Next, we examined the effect of aging on neuronal subpopulations. We observed an altered subtype distribution in older neurons ([Figure 5E](#), p-value < 0.0001 using a contingency table and Chi-square/Fisher test). The percentage of adapting neurons decreased in middle age (10%) and old (12%) compared to young (29%). The percentage of phasic neurons decreased only in old (42%) compared to middle-aged (56%) and young (55%) neurons. Accordingly, the percentage of tonic neurons increased in middle-aged (34%) and old (46%) compared to young (19%) neurons. These findings highlight how aging impacts the distribution of firing subtypes in sympathetic motor neurons.

While age does impact the distribution of firing subtypes, the question remains whether intrinsic properties are affected within each subpopulation across different ages. Therefore, we analyzed the intrinsic properties across subpopulations between different age groups. The RMP of adapting




Figure 4.

Sympathetic motor neurons from old mice are more responsive to electrical stimulation.

(A) In the context of the sympathetic reflex, motor neurons increase their firing frequency in response to inputs from the sympathetic nuclei in the brain. In this study, motor neuron responses were measured using electrical stimulation at different ages. **(B)** Stimulation protocol to mimic preganglionic input. **(C-E)** Representative voltage responses of sympathetic motor neurons from 12-(C), 64-(D), and 115-week-old mice (E). Blue, red, and black traces are in response to 0, 10, and 40 pA current injection respectively. The dotted line shows 0 mV as reference. **(F)** Comparison of stimulation-response curves of the number of APs vs. injected current between different ages. **(G)** Comparison of the minimum current injected that elicited at least one AP (Rheobase) between different ages. **(H)** Comparison of the number of APs fired at the maximum stimulus (100 pA) between different ages. Data were collected from N = 4 animals, n = 34 cells, from 12 weeks old, N = 6 animals, n = 68 cells, from 64 weeks old, and N = 4 animals, n = 30 cells, from 115 weeks old. p-values are shown at the top of the graphs.




Figure 5.

Analysis of neuronal subpopulations. Aging shifts neuronal population toward tonic firing.

(A) Representative recordings from three different neurons illustrate the variability in the response to show the method used to classify cells by their firing pattern as adapting (left), phasic (center), or tonic (right). Responses to stimulation of 0 (top), 20 pA above the rheobase (middle), and 100 pA (bottom). The classification was based on the response 20 pA above the rheobase. (B) Comparison of the number of APs elicited by current injections of 20 pA more than rheobase or 100 pA between classes. Data are from 64-week-old mice. (C) Comparison of the stimulus-frequency curves between classes. Data are from 64-week-old mice. (D) Comparison of RMP, (left), number of spontaneous APs (middle), and input resistance (right) between classes. All data are from 64-week-old mice. (E) Comparison of the percentage of neuronal firing subtypes between different ages. (F) Comparison of the RMP between ages and divided into neuronal subpopulations. (G) Comparison of the input resistance between ages and divided into neuronal subpopulations. (H) Comparison of the number of spontaneous APs between ages and neuronal subpopulations. (I) Comparison of the number of APs in response to maximum stimulation (100 pA) between ages and neuronal subpopulations. Data points are from N = 4 animals, n = 9, 7, and 3 from single, phasic, and old cells from 12-week-old mice. Data points are from N = 6 animals, n = 17, 38, and 11 from single, phasic, and old cells from 64-week-old mice. Data points are from N = 4 animals, n = 5, 23, and 11 from single, phasic, and old cells from 64-week-old mice. Red values indicate p-values < 0.05 while black values indicate p-values > 0.05.

and phasic firing neurons was more depolarized in middle age and old compared to young neurons (**Figure 5F**). The RMP of tonic neurons tended to become more depolarized with age, but the effect was not significant (**Figure 5F**). Similar to the observation when the entire population was analyzed, the input resistance was not significantly different with age within phasic and adapting neurons. The input resistance of tonic neurons was significantly larger only in old ($1.44 \pm 0.16 \text{ G}\Omega$) neurons compared to young ($0.93 \pm 0.16 \text{ G}\Omega$) neurons (**Figure 5G**). This analysis suggested that, regardless of the subpopulation type, older neurons tend to have more depolarized RMP with no changes in input resistance.

Interestingly, aging leads to an increase in the number of spontaneous APs in phasic and tonic neurons (**Figure 5H**). In the phasic class, middle-aged neurons fired 3 ± 2 APs per minute and old neurons fired 5 ± 3 APs, compared to zero APs in phasic young neurons. In the tonic class, middle-aged neurons fired 179 ± 46 APs and old neurons fired 86 ± 27 APs, in striking contrast to zero APs in tonic young neurons. Next, we compared the maximum number of APs elicited by the strongest stimulus (100 pA, **Figure 5I**). In phasic neurons, the number of APs at 100 pA increased by 53% in middle age (9.5 ± 1.0 APs) and 133% in old (14.5 ± 2.1 APs) compared to young neurons (6.2 ± 1.0 APs). In tonic firing neurons, the number of APs at 100 pA increased by 36% in middle age (25 ± 1.4 APs) and 27% in old (24.1 ± 1.0 AP) compared to young neurons (19.0 ± 1.9 APs).

We conclude that the changes observed when categorizing neuronal classes occur across the entire population. In addition, aging is associated with a shift in the proportion of neurons falling in each class. As a result, we hypothesized that the underlying age-related molecular mechanism is broadly shared among sympathetic motor neurons and plays a role in controlling the firing frequency.

Analysis of neuronal subpopulations

We next directed our attention to identifying molecular candidates underlying changes in membrane excitability. Sympathetic motor neurons, express at least one voltage-gated sodium channel isoform ($\text{Na}_v1.7$) *Schofield et al. (2008)*; *Toledo-Aral et al. (1997)* and several voltage-gated potassium channels *Dixon and McKinnon (1996)*; *Shi et al. (1997)*. Specifically, K_v4 , K_v2 , and K_v7 (KCNQ) channels are crucial in controlling the RMP, rheobase, firing frequency, and spike adaptation *Liu and Bean (2014)*; *Malin and Nerbonne (2000)*, 2001). These channels are of particular interest as potential contributors to the age-related alterations in neuronal excitability that we observed.

Previous work by our group and others demonstrated that cholinergic stimulation (with oxotremorine M, Oxo-M) leads to a decrease in KCNQ currents and increases the excitability of sympathetic motor neurons at young ages. Thus, **Figure 6A** shows a voltage response (measured in current-clamp mode) and a consecutive KCNQ current recording (measured in voltage-clamp mode) in the same neuron upon stimulation of cholinergic type 1 muscarinic receptors. It illustrates the temporal correlation between the decrease of KCNQ current with the increase in excitability and firing of APs *Kruse and Whitten (2021)*. This strong dependence led us to hypothesize that, like Oxo-M, aging decreases KCNQ current, leading to a depolarized RMP and hyperexcitability (**Figure 6B**). For these experiments, we measured the RMP and evoked activity using the perforated patch, followed by the amplitude of KCNQ current using a whole-cell voltage clamp in the same cell. We also measured the membrane capacitance as a proxy for cell size. Interestingly, KCNQ current density was smaller by 29% in middle age ($7.5 \pm 0.7 \text{ pA/pF}$) and by 55% in old ($4.8 \pm 0.7 \text{ pA/pF}$) compared to young ($10.6 \pm 1.5 \text{ pA/pF}$) neurons (**Figure 6C-D**). The average capacitance was similar in young ($30.8 \pm 2.2 \text{ pF}$), middle-aged ($27.4 \pm 1.2 \text{ pF}$), and old ($28.8 \pm 2.3 \text{ pF}$) neurons (**Figure 6E**), suggesting that aging is not associated with changes in cell size of sympathetic motor neurons, and supporting the hypothesis that aging alters the levels of KCNQ current. Interestingly, we observed that KCNQ2 protein levels were 1.5 ± 0.1 -fold higher in old

compared to young neurons (**Figure 6F-G**). The decrease in KCNQ current and the increase in the abundance of KCNQ2 protein suggest a potential compensatory mechanism that occurs during aging, which we are actively investigating in an independent study.

To explore the hypothesis that a reduction in KCNQ current is responsible for the age-associated depolarization of the RMP, we compared these two parameters measured in the same cells. We observed a correlation between KCNQ current and RMP in young (coefficient of determination (r^2) = 0.22, p-value for the correlation fit = 0.007, **Figure 6H** and **6K**) and middle-aged neurons (r^2 = 0.20, p-value for the correlation fit = 0.002, **Figure 6I** and **6K**). In old neurons, the KCNQ current and RMP were no longer correlated (r^2 = 0.05, p-value for the correlation fit = 0.1, **Figure 6J** and **6K**). **Figure 6K** shows the decrease in the coefficient of determination (r^2) with aging. Similarly, the KCNQ current amplitude also correlated well with the number of APs elicited at 100 pA in young and middle-aged neurons but not in old ones. The variance in KCNQ current amplitude explained 32% of the variation in the number of APs in young (r^2 = 0.32, p-value for the correlation fit = 0.001, **Figure 6L**) and 24% in middle-aged neurons (r^2 = 0.24, p-value for the correlation fit = 0.0001, **Figure 6M**). In old neurons, the variance in KCNQ current amplitude explained only 0.05% of the variation in the number of APs (r^2 = 0.05, p-value for the correlation fit = 0.15, **Figure 6N**). **Figure 6O** shows the decrease in the coefficient of determination with aging for the number of APs. These analyses support the hypothesis that a reduction in KCNQ current alters the electrical properties, and in the case of old neurons, the marked decrease in KCNQ current compromises its role in maintaining the RMP and spontaneous firing.

Other voltage-gated sodium and potassium currents are not altered with aging

Loss of voltage-gated potassium channel function, including the current of K_{V2} and K_{V4} , has also been invoked in aging and age-associated memory decline [Fenyves et al. \(2021\)](#); [Frazzini et al. \(2016\)](#); [Navarro-Garcia et al. \(2022\)](#); [Sesti \(2016\)](#); [Simkin et al. \(2015\)](#); [Yu et al. \(2019\)](#). Therefore, we also looked for potential age-associated changes in other voltage-gated potassium currents. We used a recording solution designed to abolish sodium, calcium, and potassium currents mediated by both calcium-activated potassium channels (BK channels) and KCNQ channels. This recording solution contained 100 nM tetrodotoxin (TTX) to suppress voltage-gated sodium channels, 100 μ M Cd^{2+} to inhibit calcium channels and 10 μ M XE-991 to block KCNQ channels. Next, we compared the outward current density before and after application of a cocktail of 100 nM phiroxotoxin and 100 nM guangxitoxin to block K_{V4} and K_{V2} channels. **Figure 7A** and **C** show representative traces and the comparison of the current density between young and old neurons. We did not observe significant differences in the potassium current insensitive to XE-991 between groups (young = 363 ± 20 pA/pF, old = 342 ± 30 pA/pF, **Figure 7A** and **B**) nor in the K_{V2} and K_{V4} sensitive currents (young = 157 ± 31 pA/pF, old = 110 ± 17 pA/pF, **Figure 7C** and **D**).

In nociceptive neurons, similar to sympathetic neurons, modulation of $Na_{V1.7}$ channels has been linked to increased excitability [Akin et al. \(2021\)](#); [Dib-Hajj et al. \(2013\)](#); [Li et al. \(2018\)](#). Additionally, exogenous expression of $Na_{V1.7}$ channels in non-excitatory cells induces cell senescence [Warnier et al. \(2018\)](#), further highlighting a significant connection between aging and voltage-gated sodium channels. To investigate the hypothesis that aging might contribute to hyperexcitability through alterations in sodium currents in sympathetic motor neurons, we conducted experiments using 5 ms voltage steps in a low-sodium recording solution. We found that the sodium current density was similar in young (58.9 ± 5.4 pA/pF), middle-aged (59.0 ± 5.3 pA/pF), and old (50.1 ± 5.7 pA/pF) neurons (**Figure 7E-G**). In conclusion, our data indicate that aging does not alter the Na_V and K_V currents insensitive to XE-991 in sympathetic motor neurons.




Figure 6.

Sympathetic motor neurons from old mice show a KCNQ current reduction.

(A) Recordings illustrating the relevance of KCNQ current for controlling membrane potential and firing in sympathetic motor neurons. Top: voltage response to inhibition of KCNQ channels with 10 μ M Oxo-M, bottom: normalized KCNQ current recording in response to oxo-M in the same cell after going whole cell. **(B)** Schematic representation of the hypothesis that aged cells show reduced activity of KCNQ channels. **(C)** KCNQ current recordings from neurons isolated from mice of different ages (orange 12 weeks old, pink 64 weeks old, and purple 115 weeks old) in response to a voltage step (top). The inset shows an expanded view of the current tail. **(D)** Comparison of KCNQ current density between different ages. Data points are from $N = 5$ animals, $n = 27$ cells, from 12 weeks old, $N = 8$ animals, $n = 62$ cells, from 64 weeks old, and $N = 5$ animals, $n = 24$ cells, from 115 weeks old. **(E)** Comparison of capacitance between ages. **(F)** blot stained for total and KCNQ2 protein collected from superior cervical ganglia from 12- and 115-week-old mice. **(G)** Comparison of fold change of KCNQ2 abundance relative to total protein between ages ($n = 5$ blots from different mice). **(H-J)** Linear correlation between RMP and KCNQ current density in 12-weeks old (H), 64-weeks old (I), and 115-weeks old mice (J). **(K)** Comparison of the determination coefficient for the RMP and KCNQ current between ages. **(L-N)** Linear correlation between the number of APs at 100 pA and KCNQ current density in 12-week-old (L), 64-week-old (M), and 115-week-old mice (N). **(O)** Comparison of the determination coefficient for the maximum #AP and KCNQ current between ages. Data points are from $N = 4$ animals, $n = 26$ cells, from 12 weeks old, $N = 6$ animals, $n = 58$ cells, from 64 weeks old, and $N = 4$ animals, $n = 24$ cells, from 115 weeks old.




Figure 7.

Sympathetic motor neurons from old mice did not show changes in sodium currents or XE-991-insensitive potassium currents.

(A) Left: Schematic of the hypothesis that Kv channels insensitive to XE-991 are reduced in aged cells. Right: Family of outward currents in the presence of KCNQ channel blocker, 100 nM XE-991, in 12-week-old, and 115-week-old mice. Magnitude of voltage step to elicit each trace is shown in black. **(B)** Current-voltage relationship of the steady-state Kv currents insensitive to XE-991 from cells isolated from 12-week-old and 115-week-old mice. **(C)** Left: Schematic of the hypothesis that Kv2 (Guangxitoxin-1E sensitive, Gtx) and Kv4 (Phrixotoxin-1 sensitive, Ptx) channels are reduced in aged cells. Right: Family of Ptx- and Gtx-sensitive outward currents, and in the presence of XE-991, in 12-week-old and 115-week-old mice. Magnitude of stimulus to elicit each trace is shown in black. **(D)** Current-voltage relationship of the steady-state Kv currents sensitive to Ptx- and Gtx from cells isolated from 12-week-old and 115-week-old mice. Data points of Kv currents are from N = 3 animals, n = 12 cells, from 12 weeks old, N = 3 animals, n = 8 cells, from 115 weeks old. **(E)** Left: Schematic of the hypothesis that Na_v channels sensitive to TTX are increased in aged cells. Right: Representative sodium current recordings from neurons isolated from mice of different ages (orange 12-week-old, pink 64-week-old, and purple 115-week-old) in response to a voltage step (top). **(F)** Current-voltage relationship of the peak sodium current for different ages. **(G)** Comparison of sodium current density between different ages. Data points of Kv currents are from N = 3 animals, n = 12 cells, from 12-weeks old, N = 3 animals, n = 14 cells, from 64-weeks old, N = 3 animals, n = 14 cells, from 64-weeks old, and N = 3 animals, n = 13 cells, from 115-weeks old. p-values are shown at the top of the graphs.

Rapamycin treatment partially reverses sympathetic motor neuron dysfunction to levels comparable to those observed in younger neurons

At this point, our analysis of sympathetic motor neurons shows that aging leads to alterations in the intrinsic properties of these neurons, resulting in hyperexcitability. Now, we investigated whether the electric cellular dysfunction associated with aging can be reversed by rapamycin treatment. In vivo, rapamycin treatment extends the lifespan and improves the health of various species, while in vitro studies regulate ion channel function [Harrison et al. \(2009\)](#); [Kaeberlein et al. \(2015\)](#); [Quarles and Rabinovitch \(2020\)](#); [Selvarani et al. \(2021\)](#). For this experiment, 78-week-old mice were fed with food containing either 14 ppm of rapamycin or no drug (**Figure 8A**). The amount of food and water intake was ad libitum. Electrophysiological recordings were conducted after 10–12 weeks of rapamycin or no-drug treatment. A third group of young mice, without any treatment, served as the pre-aging control for comparison. The neurons of rapamycin-treated mice rested at a 5 mV less depolarized potential (-58.4 ± 1.3 mV) than their age-paired controls (-53.8 ± 1.8 mV) but were still 4 mV more depolarized than young (-62.0 ± 1.8 mV) neurons (**Figure 8B**). Treatment with rapamycin also reduced the percentage of neurons firing spontaneously, from 69% to 35% (**Figure 8C**).

The treatment with rapamycin partially reversed the age-induced hyperexcitation to electrical stimulation (**Figure 8D**). Notably, significant changes were observed with low-current stimuli. The rheobase in neurons from rapamycin-treated animals was 11.9 ± 0.9 pA and similar to control neurons (10.8 ± 0.5 pA). Young neurons recorded in the same experiment exhibited a rheobase of 25.9 ± 4.8 pA (**Figure 8E**). The number of APs fired with the maximum stimulus (100 pA) was also similar between rapamycin-treated neurons (13 ± 1.3 AP) and control neurons (14 ± 1.1 AP), whereas young neurons fired only 8 ± 1.6 AP (**Figure 8F**). Rapamycin did not change the size of the cells, measured using the capacitance (control: 23.3 ± 1.3 pF, rapamycin: 25.5 ± 1.8 pF, **Figure 8G**), or the input resistance (control: 0.73 ± 0.90 G Ω , rapamycin: 0.94 ± 0.11 G Ω , **Figure 8H**).

When we assessed the effect of rapamycin on the age-associated reduction of KCNQ current (**Figure 8I**), we found that neurons treated with rapamycin showed more KCNQ current (8.6 ± 0.8 pA/pF) than age-paired control neurons (5.7 ± 0.7 pA/pF), which represented a 38% reversal of the age-associated alteration of KCNQ current (**Figure 8J–K**). Paired young neurons showed a KCNQ current of 13.4 ± 2.3 pA/pF, **Figure 8K**). Together, the findings suggest that rapamycin treatment partially decreases the age-related sympathetic hyperexcitability and increases the KCNQ currents.

Rapamycin regulates sympathetic motor neuron function directly

A crucial aspect to consider with rapamycin treatment in vivo is its potential direct impact on sympathetic motor neurons or indirect effects on autonomic nuclei in the brain or sensory organs, ultimately affecting sympathetic motor neurons. We conducted additional experiments to investigate whether rapamycin directly influences cellular processes in sympathetic motor neurons and can revert age-associated changes. We isolated and cultured neurons from 115-week-old mice and added 100 nM rapamycin to the culture medium overnight (**Figure 9A**) to half of the dishes. The rapamycin-treated neurons rested at 7 mV less depolarized potential (-60.5 ± 1.3 mV) than age-paired control neurons (-53.5 ± 2.0 mV). In vitro treatment with rapamycin also reduced the number of neurons firing spontaneously to 41% ($n = 29$) from 55% of neurons firing spontaneously in the control group; however, this change was not significantly different (one-tailed Fisher's test $p = 0.2$, **Figure 9C**). As observed with the feeding strategy, the in vitro treatment with rapamycin did not change the capacitance (control: 25 ± 2 pF, rapamycin: 27 ± 2 pF). Moreover, the in vitro treatment with rapamycin did not shift the stimulus-response curve of the hyperexcited response to electrical stimulation (**Figure 9D**). The number of APs fired with the maximum stimulus (100 pA) was also similar between rapamycin-treated neurons (15 AP) and




Figure 8.

Rapamycin treatment reverts age-related hyperexcitability and KCNQ current decrease.

(A) Schematic of the hypothesis that rapamycin treatment reverses the development of hyperexcitability. Mice were fed ad libitum with food containing rapamycin for 10-12 weeks. **(B)** Comparison of the RMP in cells isolated from 90-week-old mice fed with or without rapamycin. A paired 12-week-old mouse group was used for reference as young behavior. Data points are from N = 4 animals, n = 25 cells, from 90-week-old mice fed without rapamycin, N = 4 animals, n = 21 cells, from 90-week-old mice fed with rapamycin, and N = 3 animals, n = 17 cells, from 12 weeks old. **(C)** Comparison of the percentage of silent and firing neurons in 90-week-old mice fed with or without rapamycin, and a paired 12-week-old mice group. **(D)** Comparison of the stimulus-response curve between neurons isolated from 90-week-old mice fed with or without rapamycin. The stimulus-response curve from a 12-week-old mice group is shown for comparison. **(E)** Comparison of the rheobase in 90-week-old mice fed with or without rapamycin, and a paired 12-week-old mice group. **(F)** Comparison of the number of APs elicited with a maximum stimulation (100 pA) in mice fed with or without rapamycin, and a paired 12-week-old mice group. **(G)** Comparison of the capacitance in 90-week-old mice fed with or without rapamycin, and a paired 12-week-old mice group. **(H)** Comparison of the input resistance in 90-week-old mice fed with or without rapamycin, and a paired 12-week-old mice group. **(I)** Schematic of the hypothesis that rapamycin treatment reverses the KCNQ current reduction. **(J)** KCNQ2/3 current recordings from neurons isolated from 90-week-old mice fed with or without rapamycin. **(K)** Comparison of KCNQ current density in 90-week-old mice fed with or without rapamycin, and a paired 12-week-old mice group. Data points are from N = 4 animals, n = 23 cells, from 90-week-old mice fed without rapamycin, N = 4 animals, n = 22 cells, from 90-week-old mice fed with rapamycin, and N = 3 animals, n = 13 cells, from 12 weeks old. Red values indicate p-values < 0.05 while black values indicate p-values > 0.05.

control neurons (15 AP, **Figure 9E** [↗](#)). The rheobase in neurons isolated from rapamycin-treated neurons was $19 \text{ pA} \pm 5$ and similar to control neurons ($26 \text{ pA} \pm 5$, **Figure 9F** [↗](#)). In vitro treatment with rapamycin increased KCNQ current to $8.3 \pm 0.9 \text{ pA/pF}$ from $5.4 \pm 0.7 \text{ pA/pF}$ in control neurons (9G). Importantly, when we assessed the effect of acute application of rapamycin while recording KCNQ currents in control neurons, we did not observe any changes in the current amplitude (**Figure 9H** [↗](#)), ruling out potential direct effects of rapamycin on the activity of the channels, and pointing toward another signaling mechanism that requires hours. We concluded that the anti-aging intervention of rapamycin reverts KCNQ current by approximately 35%, acting directly on sympathetic motor neurons, partly reversing the intrinsic hyperexcitability but not fully to young levels.

Discussion

This research investigates the cellular and molecular mechanisms underlying age-associated sympathetic overactivity. Our results support the idea that, alongside age-related central changes, the peripheral component of the sympathetic autonomic reflex is also affected by aging. Key findings are that aging influences the intrinsic membrane properties of sympathetic motor neurons in the following ways: 1) older motor neurons exhibit a more positive RMP and increased rheobase, 2) the percentage of motor neurons displaying spontaneous activity increases with age, 3) older motor neurons respond with higher firing rates to electrical stimulation, 4) older motor neurons show a predominant tonic firing subpopulation, 5) older neurons exhibit reduced KCNQ potassium current, 6) rapamycin treatment recovered excitability levels and increased the KCNQ current in old motor neurons.

Age-related changes in neuronal excitability

The decline in nervous system function during healthy aging has been attributed to alterations in the intrinsic membrane properties of neurons and glial cells (Table 1). These changes in electrical behavior have been observed in various experimental models, ranging from simple organisms like *C. elegans* and *Drosophila* to more complex ones like rodents and monkeys. Notably, age-related hyperexcitability has emerged as a predominant characteristic in neurons across different brain regions (Table 1).

Research on this topic has emphasized the central nervous system. However, recent evidence suggests that aging also impacts the intrinsic properties of peripheral sensory neurons leading to hyperexcitability *Hanani et al. (2023)* [↗](#). In line with this observation, our research reveals increased excitability in old sympathetic motor neurons. Collectively, these findings underscore that aging affects the intrinsic properties of peripheral neurons, challenging the notion that age-related changes are limited to the central nervous system. Hence, it is crucial to reconsider our understanding of aging-related sympathetic overactivity with a holistic perspective. Investigating how aging affects the intrinsic properties of sensory, central, and sympathetic motor neurons throughout a lifetime will be essential research to comprehend the underlying cause-and-effect mechanisms.

The molecular mechanism underlying the age-related excitability changes

The expression, distribution, regulation, and function of ion channels play a pivotal role in determining the intrinsic membrane properties of neurons. The mechanism underlying changes in age-associated excitability encompass alterations in cholinergic and glutamatergic responses and the function and expression of voltage-gated ion channels, such as Ca_v , K_v , and HCN. In recent years, research on aging has revealed a significant impact of ion channels in the aging process. For example, repressing glutamate-gated chloride channels and L-type channels extends longevity in *C.*




Figure 9.

Rapamycin-induced improvement is mediated by a direct effect sympathetic motor neurons.

(A) Schematic of the hypothesis that rapamycin treatment reverses hyperexcitability of aged neurons through a direct effect on sympathetic motor neurons. Isolated cells from 115-week-old mice were divided into two batches, one of them was treated with rapamycin overnight. **(B)** Comparison of the RMP from neurons treated with or without rapamycin from 115-week-old mice. **(C)** Comparison of the percentage of silent and firing neurons treated with or without rapamycin. **(D)** Comparison of the stimulus-response curve between neurons treated with or without rapamycin. **(E)** Comparison of the number of APs elicited with a maximum stimulation (100 pA). **(F)** Comparison of the rheobase. **(G)** Comparison of KCNQ current density. **(H)** Time course of KCNQ current amplitude during acute application of 100 nM rapamycin. Red values indicate p-values < 0.05 while black values indicate p-values > 0.05.

Excitability	Intrinsic properties	Ion channel/mechanisms	Cell type	Research model	Reference
Decrease	RMP-unaffectd Input resistance-unaffectd Rheobase-increase	Decrease of cholinergic response	CA1 pyramidal neurons Slices	Sprague-Dawley rats 3-4 vs 25-32 months old	Potier B et al, Neuroscience, 1992
Decrease	RMP-unaffectd Input resistance-unaffectd Rheobase-increase	-	Neostriatal neurons Slices	Fischer 344 rats 3-5 vs 24-26 months old	Cepeda C et al, Neuroscience, 1992
-	RMP-unaffectd Input resistance-unaffectd Rheobase-unaffectd	Decrease of cholinergic response	CA1, CA3 and fascia dentata	F-344 rats 3 weeks vs 9 months vs 24-27 months old	Shen J and Barnes CA, Neurobiology of Aging, 1996
-	-	Increase of L-type Ca_v channel activity	CA1 hippocampal neurons Zipper slices	F-344 rats 3-6 vs 12-14 vs 23-26 months old	Thibault O and Landfield PW, Science, 1996
-	-	Decrease of NMDA protein levels	CA1 and CA3 hippocampus	Sprague-Dawley rats 3-9 vs 12-17 vs 18-24 vs 25-28 vs 29-31 months old	Wenk GL and Barnes CA, Brain research, 2000
Unaffected	RMP-unaffectd Input resistance-increase	-	Dentate granule cells Slices	Rhesus monkeys 11 vs 24 years old	Luebke JI and Rosene DL, JCN, 2003
Increase	RMP-unaffectd Input resistance-increase Rheobase-decrease	-	Layer 2/3 pyramidal neurons/ prefrontal cortex	Rhesus monkeys 8 vs 22 years old	Chang YM et al, Cerebral Cortex, 2005
Decrease	-	Increase of cAMP signaling and KCNQ/HCN activity	Dorsolateral prefrontal cortical/DELAY neuron slices	Rhesus monkeys 7-9 vs 12-13 vs 17-21 years old	Wang M et al, nature, 2011
-	-	Decrease of KCNQ expression	RT-PCR-whole brain	Drosophila 5, 25, 40, 50 and 60 days of age	Cavaliere S et al, Plos One, 2013
Increase	RMP-unaffectd Input resistance-increase	-	L3 pyramidal neurons/ lateral prefrontal cortex Slices	Rhesus monkeys 8 vs 22 years old	Coskron PJ et al, JCN, 2015
-	RMP-more negative Input resistance-decreases	Increase of inward and outward rectifier K^+ currents	Microglia in striatum, neocortex and entorhinal cortex Slices	C57BL6 mice 2-3 vs 19-24 months old	Schilling T and Eder C, Glia, 2015
Increase	RMP-unaffectd Input resistance-unaffectd	Increase of $K_v4.2/K_v4.3$ expression	CA3 hippocampal neurons Slices	F1 hybrid Fischer 344 x Brown Norway rats 2-5 vs 29-32 months old	Simkin D et al, J Neurosci, 2015
Increase	-	Decrease of BK current	Dorsal root ganglion neuron	Wistar male rats 3 and 18 months	Yu W et al, J Neurol Sci, 2015
Increase	-	Increase of glutamate-gated chloride and L-type calcium channels function		Caenorhabditis elegans	Zullo JM et al, Nature, 2019
Increase	RMP-unaffectd Rheobase-decreases	Increase of I_h	L5 pyramidal neurons Somatosensory cortex Slices	Mice C57BL/6 2-6 vs 18-29 months old	Popescu IR et al, Neurobiology of aging, 2021
Increase	RMP-unaffectd Input resistance-increase Rheobase-decrease	-	CA3 hippocampal interneurons	Sprague Dawley 5 vs 24 months old	Griego E and Galván EJ, Neurobiology of Aging, 2022
Increase	RMP-more positive Input resistance-unaffectd Rheobase-unaffectd	Decrease of KCNQ2/3 current	hypocretin/orexin neurons Hypothalamus	C57BL/6J 2-3 vs 18 months old	Li SB et al, Science, 2022
Increase	RMP-more positive Input resistance-increases Rheobase-decreases	Increase of glial-neuronal coupling	Dorsal root ganglion	Balb/c mice 3 vs 12 vs 17 months old	Hanani et al, Int J Mol Sci, 2023

Figure 10.

Age-related changes in neuronal excitability

elegans [Zullo et al. \(2019\)](#)). Furthermore, inhibiting A-type K^+ channels has shown the potential to revert the intrinsic excitability of aged CA3 pyramidal neurons to a young-like state in rats [Simkin et al. \(2015\)](#).

Our data show a decrease in KCNQ current, suggesting that it is a key mechanism behind the development of a more depolarized RMP and hyperexcitability in old sympathetic motor neurons. Research in *Drosophila* shows a reduction in KCNQ expression in the brain during the organism's lifetime and that KCNQ overexpression in mushroom body neurons reverses age-related memory impairment [Cavaliere et al. \(2013\)](#). Similarly, in macaques, blocking HCN or KCNQ channels partially restores memory-related firing of aged DELAY neurons to more youthful levels [Wang et al. \(2011\)](#). Recent findings by Li et al. demonstrate that aged hypocretin/orexin neurons exhibit hyperexcitability with lower KCNQ expression. Their study shows that selectively disrupting *Kcnq2/3* genes in young hypocretin neurons is sufficient to depolarize these neurons and cause sleep fragmentation, mimicking the sleep instability observed in aged mice [Li et al. \(2022\)](#). Collectively, these data emphasize the significance of KCNQ2/3 current in the development of age-related changes.

While a decrease in the expression of KCNQ during aging has been suggested as a main mechanism, we observed an increase in the abundance of KCNQ2 protein. Nevertheless, the function was diminished in old neurons, suggesting that a compensatory mechanism is being activated in sympathetic motor neurons. It is plausible that aging affects the trafficking of KCNQ channels or the insertion at the plasma membrane. Another alternative is that aging alters cofactors of the channel, such as $\text{PtdIns}(4,5)\text{P}_2$ or even the stability at the plasma membrane required for its proper opening. This mechanism remains elusive and open for exploration.

Rapamycin, an FDA-approved drug for human use, has been shown to extend the lifespan of various species, including mice, flies, worms, and yeast. It effectively ameliorates age-related deterioration and has demonstrated promising results in improving brain function in models of Alzheimer's disease. Rapamycin reduces amyloid beta depositions and misfolded tau proteins, restores cerebral blood flow, prevents neuronal loss, and enhances cognitive function [Kaeberlein and Galvan \(2019\)](#). Our study suggests that rapamycin also benefits the sympathetic nervous system. In our in vitro experiments, rapamycin directly acts on sympathetic motor neurons, reversing hyperexcitability. This leads us to propose that sympathetic hyper-excitability may be caused by mTOR hyper-activation, which has been suggested as a common mechanism for aging's detrimental effects on various organs, including the brain and vasculature.

Our data align with previous research indicating that anti-aging treatments focus on targeting ion channels and excitability in the nervous system to enhance longevity and overall health. For instance, caloric restriction has been shown to prevent the decrease of NMDA and AMPA receptors in the hippocampus of rats [Shi et al. \(2007\)](#). Similarly, curcumin treatment has been found to reduce hyperexcitability in pyramidal neurons in aged rhesus monkeys, leading to improvements in a working memory test [Chang et al. \(2022\)](#). These findings highlight the potential of modulating ion channels and neural excitability as promising strategies for promoting healthy aging and preserving cognitive function.

Limitations and Conclusion

The hypothesis of age-associated hyperexcitability of sympathetic motor neurons should be tested in the neurons of other sympathetic ganglia, including the stellate and celiac. While our study allowed us to determine the effect of aging on sympathetic neurons in isolation from other components of the sympathetic reflex, it did not provide insight into whether the hyperexcitability of motor neurons is a compensatory mechanism responding to earlier changes in the brain or target tissues. This question has posed a long-standing challenge, especially in various pathologies

where sympathetic overactivity, such as arrhythmias and hypertension, play a role. In such cases, sympathetic overactivity could result from deterioration in the target organs or sensory components, such as the heart and carotid body. Further research and exploration are needed to unravel these complex interactions and establish a comprehensive understanding of the mechanisms underlying age-related changes in the sympathetic nervous system.

In conclusion, this study demonstrates that aging directly impacts the intrinsic electrical properties of sympathetic motor neurons. Furthermore, our research postulates that the decrease in KCNQ function underlies the hyperexcitability of sympathetic motor neurons. The administration of rapamycin reverts the age-related neuronal hyperexcitability and KCNQ current decrease. These findings shed light on the mechanisms involved in age-related changes within the sympathetic nervous system and offer a promising avenue for further investigation and potential intervention.

Methods and Materials

Animal Models

Male C57BL/6 WT mice were purchased from the Jackson Laboratory (12 weeks, RRID:IMSR_JAX:000664) or obtained from the NIA-NIH colony (ages 64 weeks and 115 weeks). All animals were kept in an animal facility with controlled conditions and were given standard chow and water ad libitum. The animal handling protocol was approved by the University of Washington Institutional Animal Care and Use Committee.

Sympathetic motor neuron cell culture

Neurons from superior cervical ganglia (SCG) were prepared by enzymatic digestion following a standardized protocol for rats *Vivas et al. (2014)* but reducing the enzyme concentration in half. Isolated neurons were plated on poly-L-lysine (#Cat: P1524-25MG, Sigma) coated glass coverslips and incubated in 5% CO₂ at 37°C in DMEM supplemented with 10% FBS and 0.2% penicillin/streptomycin.

Imaging of neurite growth

Cells were imaged in culture medium at 24 and 72 h and one week after isolation for neurites analysis. Images were taken using the bright field of an LSM 880 confocal microscope (Zeiss).

Electrophysiological Recordings

Voltage responses were recorded using the perforated-patch configuration in the current-clamp mode, whereas KCNQ currents were recorded using the whole-cell configuration in voltage-clamp mode. We used an Axopatch 200B amplifier coupled with an Axon Digidata 1550B data acquisition board (Molecular Devices Electrophysiology) and HEKA EPC 9 amplifier (HEKA Elektronik) to acquire the electrical signals. Patch pipettes had a resistance of 2 – 4 M Ω . A liquid junction potential of 4 mV was calculated using the pCLAMP 10 software and not corrected while recording. Instead, V_{rest} reported in the Results section was corrected during analysis. Voltage responses were sampled at 5 KHz, whereas currents were sampled at 2 KHz. For current recordings in voltage clamp mode, cell capacitance was canceled, and series resistances of <10 M Ω were compensated by 70%. The voltage error due to any remaining series resistance is expected to be <4 mV. The bath solution (Ringer's solution) contained 150 mM NaCl, 2.5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, 10 mM HEPES, and 8 mM glucose, adjusted to pH 7.4 with NaOH. The internal solution used to fill the whole-cell pipettes contained 175 mM KCl, 1 mM MgCl₂, 5 mM HEPES, 0.1 mM K₄BAPTA, 3 mM Na₂ATP, and 0.1 mM Na₃GTP, adjusted to pH 7.2 with KOH. For current-clamp recordings by perforated patch, 60 amphotericin B (Cat #: A4888) was added to the

pipette solution to facilitate electrical access to the cell. The bath solution was perfused at 2 ml/minute, permitting solution exchange surrounding the recording cell with a time constant of 4 s. Sodium currents were recorded 2–8 h after isolation using a low-sodium ringer.

Rapamycin treatment in vivo

Eight mice 76 weeks old were fed for twelve weeks with pellets containing encapsulated rapamycin (eRap, 14 mg/kg; Rapamycin holdings) or Eudagrit S-100 (the polymer used to encapsulate rapamycin). Pellets with drugs were made by first pulverizing regular chow in a food processor until obtaining powder. Then, eRap or Eudagrit was added to the powder and mixed for 5 min using a standard mixer. 1% of agar solution (A7002 from Sigma) was added to the mix. After obtaining a homogeneous blend, food pellets were made by compacting the food. Food was stored in a -20°C cold room and used within two weeks after preparation. All animals were kept in an animal facility with controlled conditions and given food and water ad libitum.

Protein extraction and abundance determination

Protein from sympathetic ganglia was harvested in RIPA buffer (#89900, Thermo Scientific) with Complete, Mini, EDTA-free protease inhibitor cocktail (#11836170001, Roche) for 15 min at 4°C. Post-nuclear supernatant was isolated by centrifuging for 20 minutes at 13,600 g at 4°C. Protein concentration was quantified on a plate reader using the Pierce BCA protein assay kit (#23225, Thermo Scientific). Gel lanes were loaded with 30 µg of total protein. Protein samples were resolved in 4–12% Bis-Tris gels under reducing conditions. Proteins were transferred onto nitrocellulose membranes (0.2 µm; #LC2000, Life Technologies) using the Mini-Bolt system (#A25977, Thermo Scientific). Membranes were blotted using rabbit anti-KCNQ2 (ab22897, Abcam, RRID: AB_775890, 1:500). Blotted bands were detected using HRP conjugated secondary antibodies goat anti-rabbit conjugated with HRP (#1706515, Bio-Rad, 1:15, 000). ImageJ was used to calculate the fluorescence density of each band. The abundance of KCNQ2 was reported as normalized to total protein and relative to the abundance in tissue from young animals.

Reagents

XE-991 (Cat #: X-101), Phrixotoxin-1 (Cat #: STP-700), Guangtoxin-1E (Cat #: STG-200) and Tetrodotoxin (T-550) were obtained from Alomone Labs, and microcapsules of Rapamycin from Rapamycin Holdings.

Data analysis and statistics

We used IGOR Pro (IGOR Software, WaveMetrics, RRID: SCR_000325), Excel (Microsoft), and Prism (GraphPad, RRID: SCR_002798) to analyze data. ImageJ (RRID: SCR_003070) was used to process images. Data were collected from independent experiments from at least three mice and are presented as Mean ± SEM. The statistical analyses were performed using the parametric Student's t-test when comparing two variables and an ordinary one-way ANOVA (Dunnett's multiple comparisons test) when comparing three or more variables. Graph Pad was used to calculate Pearson correlation coefficients using one-tail analysis. A non-parametric statistical test (Mann-Whitney Wilcoxon) was used to test for statistical significance between the percentage of firing and non-firing neurons. p values <0.05 as statistical significance. The number of cells used for each experiment is detailed in each figure legend.

Acknowledgements

This work was supported by grants from the US National Institutes of Health (OV: R35GM142690; CMM: R00 AG056595), the AFAR-Glenn Foundation Junior Faculty Award (CMM), and the AFAR-Sagol Geromics Award (OV). We thank Bertil Hille for his contribution to the editing and writing of

this manuscript. We thank readers Jill Jensen, Duk-Su Koh, Roya Pournejati, and Charles Asbury for their feedback on the manuscript. Lizbeth de la Cruz was awarded the Weill Neurohub Fellowship for the period 2021-2023 to support this research. Claudia Moreno is a Freeman Hrabowski HHMI Scholar.

References

- Akin EJ, Alsaloum M, Higerd GP, Liu S, Zhao P, Dib-Hajj FB, Waxman SG, Dib-Hajj SD (2021) **Paclitaxel increases axonal localization and vesicular trafficking of Nav1.7** *Brain* **144**:1727–1737 <https://doi.org/10.1093/brain/awab113>
- Andrews C. (1996) **Target specific differences in the dendritic morphology and neuropeptide content of neurons in rat SCG during aging.pdf** *J Compar Neurosci*
- Andrews TJ, Cowen T. (1994) **In vivo infusion of NGF induces the organotypic regrowth of perivascular nerves following their atrophy in aged rats** *J Neurosci* **14**:3048–58 <https://doi.org/10.1523/JNEUROSCI.14-05-03048.1994>
- Balasubramanian P, Branen L, Sivasubramanian MK, Monteiro R, Subramanian M. (2021) **Aging is associated with glial senescence in the brainstem - implications for age-related sympathetic overactivity** *Aging (Albany NY)* **13**:13460–13473 <https://doi.org/10.18632/aging.203111>
- Buchholz JN, Behringer EJ, Pottorf WJ, Pearce WJ, Vanterpool CK (2007) **Age-dependent changes in Ca²⁺ homeostasis in peripheral neurones: implications for changes in function** *Aging Cell* **6**:285–96 <https://doi.org/10.1111/j.1474-9726.2007.00298.x>
- Burnstock G. (1990) **Changes in expression of autonomic nerves in aging and disease** *J Auton Nerv Syst* **30**:S25–34 [https://doi.org/10.1016/0165-1838\(90\)90096-2](https://doi.org/10.1016/0165-1838(90)90096-2)
- Cavaliere S, Malik BR, Hodge JJ (2013) **KCNQ channels regulate age-related memory impairment** *PLoS One* **8** <https://doi.org/10.1371/journal.pone.0062445>
- Cepeda C, Lee N, Buchwald NA, Radisavljevic Z, Levine MS (1992) **Age-induced changes in electrophysiological responses of neostriatal neurons recorded in vitro** *Neuroscience* **51**:411–23 [https://doi.org/10.1016/0306-4522\(92\)90325-v](https://doi.org/10.1016/0306-4522(92)90325-v)
- Chang W, Weaver CM, Medalla M, Moore TL, Luebke JI (2022) **Age-related alterations to working memory and to pyramidal neurons in the prefrontal cortex of rhesus monkeys begin in early middle-age and are partially ameliorated by dietary curcumin** *Neurobiol Aging* **109**:113–124 <https://doi.org/10.1016/j.neurobiolaging.2021.09.012>
- Chang YM, Rosene DL, Killiany RJ, Mangiamele LA, Luebke JI (2005) **Increased action potential firing rates of layer 2/3 pyramidal cells in the prefrontal cortex are significantly related to cognitive performance in aged monkeys** *Cereb Cortex* **15**:409–18 <https://doi.org/10.1093/cercor/bhh144>
- Coskren PJ, Luebke JI, Kabaso D, Wearne SL, Yadav A, Rumbell T, Hof PR, Weaver CM (2015) **Functional consequences of age-related morphologic changes to pyramidal neurons of the rhesus monkey prefrontal cortex** *J Comput Neurosci* **38**:263–83 <https://doi.org/10.1007/s10827-014-0541-5>
- Dib-Hajj SD, Yang Y, Black JA, Waxman SG (2013) **The Na(V)1.7 sodium channel: from molecule to man** *Nat Rev Neurosci* **14**:49–62 <https://doi.org/10.1038/nrn3404>

- Dixon JE, McKinnon D. (1996) **Potassium channel mRNA expression in prevertebral and paravertebral sympathetic neurons** *Eur J Neurosci* **8**:183–91 <https://doi.org/10.1111/j.1460-9568.1996.tb01179.x>
- Esler MD, Turner AG, Kaye DM, Thompson JM, Kingwell BA, Morris M, Lambert GW, Jennings GL, Cox HS, Seals DR (1995) **Aging effects on human sympathetic neuronal function** *Am J Physiol* **268**:R278–85 <https://doi.org/10.1152/ajpregu.1995.268.1.R278>
- Fenyves BG, Arnold A, Gharat VG, Haab C, Tishinov K, Peter F, de Quervain D, Papassotiropoulos A, Stetak A. (2021) **Dual Role of an mps-2/KCNE-Dependent Pathway in Long-Term Memory and Age-Dependent Memory Decline** *Curr Biol* **31**:527–539 <https://doi.org/10.1016/j.cub.2020.10.069>
- Frazzini V, Guarnieri S, Bomba M, Navarra R, Morabito C, Mariggio MA, Sensi SL (2016) **Altered Kv2.1 functioning promotes increased excitability in hippocampal neurons of an Alzheimer’s disease mouse model** *Cell Death Dis* **7** <https://doi.org/10.1038/cddis.2016.18>
- Goldstein DS, Lake CR, Chernow B, Ziegler MG, Coleman MD, Taylor AA, Mitchell JR, Kopin IJ, Keiser HR (1983) **Age-dependence of hypertensive-normotensive differences in plasma norepinephrine** *Hypertension* **5**:100–4 <https://doi.org/10.1161/01.hyp.5.1.100>
- Goldstein DS, McCarty R, Polinsky RJ, Kopin IJ (1983) **Relationship between plasma norepinephrine and sympathetic neural activity** *Hypertension* **5**:552–9 <https://doi.org/10.1161/01.hyp.5.4.552>
- Griego E, Galván EJ. (2022) **Biophysical and synaptic properties of regular spiking interneurons in hippocampal area CA3 of aged rats** *Neurobiol Aging* **112**:27–38 <https://doi.org/10.1016/j.neurobiolaging.2021.12.007>
- Hanani M, Spray DC, Huang TY (2023) **Age-Related Changes in Neurons and Satellite Glial Cells in Mouse Dorsal Root Ganglia** *Int J Mol Sci* **24** <https://doi.org/10.3390/ijms24032677>
- Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, Nadon NL, Wilkinson JE, Frenkel K, Carter CS, Pahor M, Javors MA, Fernandez E, Miller RA (2009) **Rapamycin fed late in life extends lifespan in genetically heterogeneous mice** *Nature* **460**:392–5 <https://doi.org/10.1038/nature08221>
- Hart EC, Joyner MJ, Wallin BG, Johnson CP, Curry TB, Eisenach JH, Charkoudian N. (2009) **Age-related differences in the sympathetic-hemodynamic balance in men** *Hypertension* **54**:127–33 <https://doi.org/10.1161/HYPERTENSIONAHA.109.131417>
- Hogikyan RV, Supiano MA (1994) **Arterial alpha-adrenergic responsiveness is decreased and SNS activity is increased in older humans** *Am J Physiol* **266**:E717–24 <https://doi.org/10.1152/ajpendo.1994.266.5.E717>
- Ito K, Sato A, Sato Y, Suzuki H. (1986) **Increases in adrenal catecholamine secretion and adrenal sympathetic nerve unitary activities with aging in rats** *Neurosci Lett* **69**:263–8 [https://doi.org/10.1016/0304-3940\(86\)90491-x](https://doi.org/10.1016/0304-3940(86)90491-x)
- Kaeberlein M, Galvan V. (2019) **Rapamycin and Alzheimer’s disease: Time for a clinical trial?** *Sci Transl Med* **11** <https://doi.org/10.1126/scitranslmed.aar4289>
- Kaeberlein M, Rabinovitch PS, Martin GM (2015) **Healthy aging: The ultimate preventative medicine** *Science* **350**:1191–3 <https://doi.org/10.1126/science.aad3267>

- Kim KW, Kim K, Lee H, Suh BC (2019) **Ethanol Elevates Excitability of Superior Cervical Ganglion Neurons by Inhibiting Kv7 Channels in a Cell Type-Specific and PI(4,5)P2-Dependent Manner** *Int J Mol Sci* **20** <https://doi.org/10.3390/ijms20184419>
- Kruse M, Whitten RJ (2021) **Control of Neuronal Excitability by Cell Surface Receptor Density and Phosphoinositide Metabolism** *Front Pharmacol* **12** <https://doi.org/10.3389/fphar.2021.663840>
- Li SB, Damonte VM, Chen C, Wang GX, Kebschull JM, Yamaguchi H, Bian WJ, Purmann C, Pattani R, Urban AE, Mourrain P, Kauer JA, Scherrer G, de Lecea L. (2022) **Hyperexcitable arousal circuits drive sleep instability during aging** *Science* **375** <https://doi.org/10.1126/science.abh3021>
- Li Y, North RY, Rhines LD, Tatsui CE, Rao G, Edwards DD, Cassidy RM, Harrison DS, Johansson CA, Zhang H, Dougherty PM (2018) **DRG Voltage-Gated Sodium Channel 1.7 Is Upregulated in Paclitaxel-Induced Neuropathy in Rats and in Humans with Neuropathic Pain** *J Neurosci* **38**:1124–1136 <https://doi.org/10.1523/JNEUROSCI.0899-17.2017>
- Liu PW, Bean BP (2014) **Kv2 channel regulation of action potential repolarization and firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal neurons** *J Neurosci* **34**:4991–5002 <https://doi.org/10.1523/JNEUROSCI.1925-13.2014>
- Luebke JI, Rosene DL (2003) **Aging alters dendritic morphology, input resistance, and inhibitory signaling in dentate granule cells of the rhesus monkey** *J Comp Neurol* **460**:573–84 <https://doi.org/10.1002/cne.10668>
- Madden KS, Bellinger DL, Felten SY, Snyder E, Maida ME, Felten DL (1997) **Alterations in sympathetic innervation of thymus and spleen in aged mice** *Mech Ageing Dev* **94**:165–75 [https://doi.org/10.1016/s0047-6374\(96\)01858-1](https://doi.org/10.1016/s0047-6374(96)01858-1)
- Malin SA, Nerbonne JM (2000) **Elimination of the fast transient in superior cervical ganglion neurons with expression of KV4.2W362F: molecular dissection of IA** *J Neurosci* **20**:5191–9
- Malin SA, Nerbonne JM (2001) **Molecular heterogeneity of the voltage-gated fast transient outward K⁺ current, I(Af), in mammalian neurons** *J Neurosci* **21**:8004–14
- Narkiewicz K, Phillips BG, Kato M, Hering D, Bieniaszewski L, Somers VK (2005) **Gender-selective interaction between aging, blood pressure, and sympathetic nerve activity** *Hypertension* **45**:522–5 <https://doi.org/10.1161/01.HYP.0000160318.46725.46>
- Navarro-Garcia JA, Salguero-Bodes R, Gonzalez-Lafuente L, Martin-Nunes L, Rodriguez-Sanchez E, Bada-Bosch T, Hernandez E, Merida-Herrero E, Praga M, Solis J, Arribas F, Bueno H, Kuro OM, Fernandez-Velasco M, Ruilope LM, Delgado C, Ruiz-Hurtado G. (2022) **The anti-aging factor Klotho protects against acquired long QT syndrome induced by uremia and promoted by fibroblast growth factor 23** *BMC Med* **20** <https://doi.org/10.1186/s12916-021-02209-9>
- Popescu IR L, KQ, Ducote AL, Li JE, Leland AE, Mostany R. (2021) **Increased intrinsic excitability and decreased synaptic inhibition in aged somatosensory cortex pyramidal neurons** *Neurobiol Aging* **98**:88–98 <https://doi.org/10.1016/j.neurobiolaging.2020.10.007>
- Potier B, Rascol O, Jazat F, Lamour Y, Dutar P. (1992) **Alterations in the properties of hippocampal pyramidal neurons in the aged rat** *Neuroscience* **48**:793–806 [https://doi.org/10.1016/0306-4522\(92\)90267-6](https://doi.org/10.1016/0306-4522(92)90267-6)

- Pottorf WJ, Duckles SP, Buchholz JN (2000) **SERCA function declines with age in adrenergic nerves from the superior cervical ganglion** *J Auton Pharmacol* **20**:281–90 <https://doi.org/10.1046/j.1365-2680.2000.00194.x>
- Quarles EK, Rabinovitch PS (2020) **Transient and late-life rapamycin for healthspan extension** *Aging (Albany NY)* **12**:4050–4051 <https://doi.org/10.18632/aging.102947>
- Schilling T, Eder C. (2015) **Microglial K(+) channel expression in young adult and aged mice** *Glia* **63**:664–72 <https://doi.org/10.1002/glia.22776>
- Scho?eld GG, Puhl r H L, Ikeda SR (2008) **Properties of wild-type and ?uorescent protein-tagged mouse tetrodotoxin-resistant sodium channel (Na V 1.8) heterologously expressed in rat sympathetic neurons** *J Neurophysiol* **99**:1917–27 <https://doi.org/10.1152/jn.01170.2007>
- Selvarani R, Mohammed S, Richardson A. (2021) **Effect of rapamycin on aging and age-related diseases-past and future** *Geroscience* **43**:1135–1158 <https://doi.org/10.1007/s11357-020-00274-1>
- Sesti F. (2016) **Oxidation of K(+) Channels in Aging and Neurodegeneration** *Aging Dis* **7**:130–5 <https://doi.org/10.14336/AD.2015.0901>
- Shen J, Barnes CA (1996) **Age-related decrease in cholinergic synaptic transmission in three hippocampal sub?elds** *Neurobiol Aging* **17**:439–51 [https://doi.org/10.1016/0197-4580\(96\)00020-6](https://doi.org/10.1016/0197-4580(96)00020-6)
- Shi L, Adams MM, Linville MC, Newton IG, Forbes ME, Long AB, Riddle DR, Brunso-Bechtold JK (2007) **Caloric restriction eliminates the aging-related decline in NMDA and AMPA receptor subunits in the rat hippocampus and induces homeostasis** *Exp Neurol* **206**:70–9 <https://doi.org/10.1016/j.expneurol.2007.03.026>
- Shi W, Wymore RS, Wang HS, Pan Z, Cohen IS, McKinnon D, Dixon JE (1997) **Identi?cation of two nervous system-speci?c members of the erg potassium channel gene family** *J Neurosci* **17**:9423–32
- Shi Z, Li B, Brooks VL (2015) **Role of the Paraventricular Nucleus of the Hypothalamus in the Sympathoexcitatory Effects of Leptin** *Hypertension* **66**:1034–41 <https://doi.org/10.1161/hypertensionaha.115.06017>
- Shi Z, Pelletier NE, Wong J, Li B, Sdrulla AD, Madden CJ, Marks DL, Brooks VL (2020) **Leptin increases sympathetic nerve activity via induction of its own receptor in the paraventricular nucleus** *Elife* **9** <https://doi.org/10.7554/eLife.55357>
- Simkin D, Hattori S, Ybarra N, Musial TF, Buss EW, Richter H, Oh MM, Nicholson DA, Disterhoft JF (2015) **Aging-Related Hyperexcitability in CA3 Pyramidal Neurons Is Mediated by Enhanced A-Type K+ Channel Function and Expression** *J Neurosci* **35**:13206–18 <https://doi.org/10.1523/JNEUROSCI.0193-15.2015>
- Springer MG, Kullmann PH, Horn JP (2015) **Virtual leak channels modulate ?ring dynamics and synaptic integration in rat sympathetic neurons: implications for ganglionic transmission in vivo** *J Physiol* **593**:803–23 <https://doi.org/10.1113/jphysiol.2014.284125>
- Thibault O, Land?eld PW (1996) **Increase in single L-type calcium channels in hippocampal neurons during aging** *Science* **272**:1017–20 <https://doi.org/10.1126/science.272.5264.1017>

Toledo-Aral JJ, Moss BL, He ZJ, Koszowski AG, Whisenand T, Levinson SR, Wolf JJ, Silos-Santiago I, Halegoua S, Mandel G. (1997) **Identification of PN1, a predominant voltage-dependent sodium channel expressed principally in peripheral neurons** *Proc Natl Acad Sci U S A* **94**:1527–32 <https://doi.org/10.1073/pnas.94.4.1527>

Veith RC, Featherstone JA, Linares OA, Halter JB (1986) **Age differences in plasma norepinephrine kinetics in humans** *J Gerontol* **41**:319–24 <https://doi.org/10.1093/geronj/41.3.319>

Vivas O, Kruse M, Hille B. (2014) **Nerve growth factor sensitizes adult sympathetic neurons to the proinflammatory peptide bradykinin** *J Neurosci* **34**:11959–71 <https://doi.org/10.1523/JNEUROSCI.1536-14.2014>

Sa Wallin (1978) **Human muscle nerve sympathetic activity at rest relationship to age.pdf** *J Physiol*

Wang M, Gamo NJ, Yang Y, Jin LE, Wang XJ, Laubach M, Mazer JA, Lee D, Arnsten AF (2011) **Neuronal basis of age-related working memory decline** *Nature* **476**:210–3 <https://doi.org/10.1038/nature10243>

Warnier M, Flaman JM, Chouabe C, Wiel C, Gras B, Griveau A, Blanc E, Foy JP, Mathot P, Saintigny P, Van Coppenolle F, Vindrieux D, Martin N, Bernard D. (2018) **The SCN9A channel and plasma membrane depolarization promote cellular senescence through Rb pathway** *Aging Cell* **17** <https://doi.org/10.1111/accel.12736>

Wenk GL, Barnes CA (2000) **Regional changes in the hippocampal density of AMPA and NMDA receptors across the lifespan of the rat** *Brain Res* **885**:1–5 [https://doi.org/10.1016/S0006-8993\(00\)02792-X](https://doi.org/10.1016/S0006-8993(00)02792-X)

Yu W, Lin X, Gao S, Li C. (2015) **Age-related changes of inactivating BK channels in rat dorsal root ganglion neurons** *J Neurol Sci* **358**:138–45 <https://doi.org/10.1016/j.jns.2015.08.1526>

Yu W, Zhang H, Shin MR, Sesti F. (2019) **Oxidation of KCNB1 potassium channels in the murine brain during aging is associated with cognitive impairment** *Biochem Biophys Res Commun* **512**:665–669 <https://doi.org/10.1016/j.bbrc.2019.03.130>

Ziegler MG, Lake CR, Kopin IJ (1976) **Plasma noradrenaline increases with age** *Nature* **261**:333–5 <https://doi.org/10.1038/261333a0>

Zullo JM, Drake D, Aron L, O’Hern P, Dhamne SC, Davidsohn N, Mao CA, Klein WH, Rotenberg A, Bennett DA, Church GM, Colaiácovo MP, Yankner BA (2019) **Regulation of lifespan by neural excitation and REST** *Nature* **574**:359–364 <https://doi.org/10.1038/s41586-019-1647-8>

Article and author information

Lizbeth de la Cruz

Department of Physiology and Biophysics, University of Washington, Seattle, WA

Derek Bui

Department of Physiology and Biophysics, University of Washington, Seattle, WA

Claudia M. Moreno

Department of Physiology and Biophysics, University of Washington, Seattle, WA, Howard Hughes Medical Institute
ORCID iD: [0000-0001-8397-3649](https://orcid.org/0000-0001-8397-3649)

Oscar Vivas

Department of Physiology and Biophysics, University of Washington, Seattle, WA, Department of Pharmacology, University of Washington, Seattle, WA
For correspondence: vivas@uw.edu
ORCID iD: [0000-0002-0964-385X](https://orcid.org/0000-0002-0964-385X)

Copyright

© 2023, de la Cruz et al.

This article is distributed under the terms of the [Creative Commons Attribution License \(https://creativecommons.org/licenses/by/4.0/\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Murali Prakriya

Northwestern University, United States of America

Senior Editor

John Huguenard

Stanford University School of Medicine, United States of America

Reviewer #1 (Public Review):**Summary:**

The authors study age-related changes in the excitability and firing properties of sympathetic neurons, which they ascribe to age-related changes in the expression of KCNQ (Kv7, "M-type") K⁺ currents in rodent sympathetic neurons, whose regulation by GPCRs has been most thoroughly studied for over 40 years. The authors suggest the ingestion of rapamycin may partially reverse the age-related decrease in M-channel expression. With the rapamycin part included, it is unclear how this work will impact the field of age-related neuronal dysfunction, as the mechanistic information is not strong.

Strengths:

The strengths include the rigor of the current-clamp and voltage-clamp experiments, the lovely, crisp presentation of the data, and the expert statistics. The separation of neurons into tonic, phasic, and adapting classes is also interesting, and informative. The writing is also elegant, and crisp. The above is especially true of the manuscript up until the part dealing with the effects of rapamycin, which becomes less compelling.

Weaknesses:

Where the manuscript becomes less compelling is in the rapamycin section, which does not provide much in the way of mechanistic insights. As such, the effect is more of an epiphenomenon of unclear insight, and the authors cannot ascribe a signaling mechanism to it that is supported by data. Thus, this latter part rather undermines the overall impact and

central advance of the manuscript. The problem is exacerbated by the controversial and anecdotal nature of the entire mTor/aging field, some of whose findings have very unfortunately had to be recently retracted.

I would strongly recommend to the authors that they end the manuscript with their analysis of the role of M current/KCNQ channels in the numerous age-related changes in sympathetic neuron function that they elegantly report, and save the rapamycin, and possible mTor action, for a separate line of inquiry that the authors could develop in a more thorough and scholarly way.

Reviewer #2 (Public Review):

Summary:

This research shows compelling and detailed evidence showing that aging influences intrinsic membrane properties of peripheral sympathetic motor neurons such that they become more excitable. Furthermore, the authors present convincing evidence that the oral administration of the anti-aging drug Rapamycin partially reversed hyperexcitability in aged neurons. This study also investigates the molecular mechanisms underlying age-associated hyperexcitability in mouse sympathetic motor neurons. In that regard, the authors found an age-associated reduction of an outward current having properties similar to KCNQ2/Q3 potassium current. They suggested a reduction of KCNQ2/Q3 current density in aged neurons as a potential mechanism behind their overactivity.

Strengths:

Detailed and rigorous analysis of electrical responses of peripheral sympathetic motor neurons using electrophysiology (perforated patch and whole-cell recordings). Most of the conclusions of this paper are well supported by the data.

Weaknesses:

1. The identity of the age-associated reduced current as KCNQ2/Q3 is not corroborated by pharmacology (blocking the current with the specific blocker XE-991).
2. The manuscript does not include a direct test of the reduction of KCNQ current as the mechanism behind age-induced hyperexcitability.

Reviewer #3 (Public Review):

This is a descriptive study of membrane excitability and Na⁺ and K⁺ current amplitudes of sympathetic motor neurons in culture. The main findings of the study are that neurons isolated from aged animals show increased membrane excitability manifested as increased firing rates in response to electrical stimulation and changes in related membrane properties including depolarized resting membrane potential, increased rheobase, and spontaneous firing. By contrast, neuron cultures from young mice show little to no spontaneous firing and relatively low firing rates in response to current injection. These changes in excitability correlate with significant reductions in the magnitude of KCNQ currents in aged neurons compared to young neurons. Treating cultures with the immunosuppressive drug, rapamycin, which has known antiaging effects in model animals appears to reverse the firing rates in aged neurons and enhance KCNQ current. The authors conclude that aging promotes hyperexcitability of sympathetic motor neurons.

The electrophysiological cataloging of the neuronal properties is generally well done, and the experiments are performed using perforated patch recordings which preserve the internal constituents of neurons, providing confidence that the effects seen are not due to washout of regulators from the cells. The main weakness is that this study is a descriptive tabulation of changes in the electrophysiology of neurons in culture, and the effects shown are correlative

rather than establishing causality. It is difficult to know from the data presented whether the changes in KCNQ channels are in fact directly responsible for the observed changes in membrane excitability. Furthermore, a notable omission seems to be the analysis of Ca^{2+} currents which have been widely linked to alterations in membrane properties in aging. As well, additional experiments in slice cultures would provide greater significance on the potential relevance of the findings for intact preparations. Finally, experiments using KCNQ blockers and activators could provide greater relevance that the observed changes in KCNQ are indeed connected to changes in membrane excitability.