

The larva and adult of *Helicoverpa armigera* use differential gustatory receptors to sense sugars

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

April 26, 2024 (this version)

Reviewed preprint version 1

November 7, 2023

Posted to preprint server

September 21, 2023

Sent for peer review

August 30, 2023

Shuai-Shuai Zhang, Pei-Chao Wang, Chao Ning, Ke Yang, Guo-Cheng Li, Lin-Lin Cao, Ling-Qiao Huang, Chen-Zhu Wang 

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China • Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Almost all herbivorous insects feed on plants and use sucrose as a feeding stimulant, but the molecular basis of their sucrose reception remains unclear. *Helicoverpa armigera* as a notorious crop pest worldwide mainly feeds on reproductive organs of many plant species in the larval stage, and its adult draws nectar. In this study, we determined that the sucrose sensory neurons located in the contact chemosensilla of larval maxillary galeae were 100–1000 times more sensitive to sucrose than those of adult antennae, tarsi and proboscis. Using the *Xenopus* expression system, we discovered that Gr10 highly expressed in the larval sensilla was specifically tuned to sucrose, while Gr6 highly expressed in the adult sensilla responded to fucose, sucrose and fructose. Moreover, using CRISPR/Cas9, we revealed that Gr10 was mainly used by larvae to detect lower sucrose, while Gr6 was primarily used by adults to detect higher sucrose and other saccharides, which results in differences in selectivity and sensitivity between larval and adult sugar sensory neurons. Our results demonstrate the sugar receptors in this moth are evolved to adapt towards the larval and adult foods with different types and amounts of sugar, and fill in a gap in sweet taste of animals.

eLife assessment

This **important** study identifies the gustatory receptors for sugar sensing in the larval and adult forms of the cotton bollworm, which is responsible for the destruction of many food crops world-wide. The authors find that the larval and adult forms utilise different receptors to sense sugars. The data are **convincing** and will be of interest neuroscientists working in sensory coding of sugars and to the pest management field.

Introduction

Over 400,000 species of phytophagous insects live on approximately 300,000 species of vascular plants worldwide (Schoonhoven et al. 2005 [\[1\]](#)). Carbohydrates produced by plant photosynthesis are important nutrients for phytophagous insects. The disaccharide sucrose, the major sugar in plants, is the principal transport form of photoassimilates in higher plants. Sucrose varies considerably in concentration in green leaves, fruits, and flower nectar where it occurs together with its constituent monosaccharides fructose and glucose (Bernays and Chapman 1994 [\[2\]](#)). Sucrose, fructose and glucose are the powerful phagostimulants for most phytophagous insects studied (Bernays and Chapman 1994 [\[2\]](#), Blaney and Simmonds 1988 [\[3\]](#), Chapman 2003 [\[4\]](#), Schoonhoven et al. 2005 [\[1\]](#)).

The gustatory sensory neurons (GSNs) in one or more mouthpart sensilla of phytophagous insects respond to sugars, usually including sucrose (Chapman 2003 [\[4\]](#), Dethier 1973 [\[5\]](#)). In Lepidoptera, caterpillars often have sugar sensitive GSNs in lateral or medial sensilla styloconica in the maxillary galea (Glendinning et al. 2009 [\[6\]](#), Martin and Shields 2012 [\[7\]](#), Schoonhoven and van Loon 2002 [\[8\]](#)), while adults also have sugar sensitive GSNs in contact chemosensilla of the antennae (Jørgensen et al. 2007 [\[9\]](#), Popescu et al. 2013 [\[10\]](#)), tarsi (Schnuch and Seebauer 1998 [\[11\]](#), Zhang et al. 2010 [\[12\]](#)), and proboscis (Blaney and Simmonds 1988 [\[3\]](#), Blaney and Simmonds 1990 [\[13\]](#), Faucheux 2013 [\[14\]](#)). The molecular basis of sugar reception in Lepidopteran insects remains unclear although the sweet taste receptors in mammals and *Drosophila melanogaster* have been extensively studied (Liman et al. 2014 [\[15\]](#), Yarmolinsky et al. 2009 [\[16\]](#)). In mammals, the combination of two taste receptors T1R2 and T1R3 recognizes sugars including sucrose (Damak et al. 2003 [\[17\]](#), Li et al. 2002 [\[18\]](#), Nelson et al. 2001 [\[19\]](#), Zhao et al. 2003 [\[20\]](#)). In *Drosophila*, multiple gustatory receptors (GRs), including DmGr5a, DmGr61a, and DmGr64a–f clusters, are required for the response of adults to multiple sugars (Dahanukar et al. 2001 [\[21\]](#), Dahanukar et al. 2007 [\[22\]](#), Jiao et al. 2007 [\[23\]](#), Jiao et al. 2008 [\[24\]](#), Slone et al. 2007 [\[25\]](#)); DmGr43a specifically responding to fructose is the major larval sugar receptor, but it is only expressed in internal organs and some parts of the brain (Mishra et al. 2013 [\[26\]](#), Miyamoto et al. 2012 [\[27\]](#)). No external sugar-sensing mechanism in *Drosophila* larvae has yet been characterized.

The cotton bollworm *Helicoverpa armigera* (Noctuidae, Lepidoptera) is a worldwide agricultural pest, it causes over \$3 billion in annual global economic loss (Pearce et al. 2017 [\[28\]](#)). The larvae of *H. armigera* mainly feeds on young leaves, flower buds, and fruits of host plants; the adults mainly drink nectar of plants. Larval and adult diets vary dramatically in the variety and concentration of sugars. Sucrose is the major sugar in plant leaves and reproductive organs, whereas flower nectar mainly contains sucrose, fructose and glucose (Nicolson 1998 [\[29\]](#)). Previous studies have shown that one sugar sensitive GSN in *H. armigera* is located in each of two lateral sensilla styloconica in the larva (Tang et al. 2000 [\[30\]](#), Zhang et al. 2013 [\[31\]](#)), and many sugar sensitive GSNs are distributed in the numerous contact chemosensilla of antennae (Jiang et al. 2015 [\[32\]](#)), tarsi (Blaney and Simmonds 1990 [\[13\]](#), Zhang et al. 2010 [\[12\]](#)), and proboscis (Blaney and Simmonds 1988 [\[3\]](#)) in the adult.

To understand the molecular basis of sugar reception in these neurons, we first compared electrophysiological and behavioral responses of *H. armigera* larvae and adults. Second, we analyzed the expression patterns of nine candidate sugar GR genes in larval and adult taste organs. Third, we carried out functional analysis of these GR genes using the ectopic expression system, and revealed the response profiles of *Xenopus* oocytes expressing Gr10, Gr6, and both of them. Finally, we individually knocked out the two GR genes and obtained two homozygous mutants of *H. armigera* by CRISPR/Cas9, and then detected changes in the electrophysiological and behavioral responses of the mutant larvae and adults to sugars. Taking all the results together, we demonstrate that larval and adult *H. armigera* used different GRs to detect sucrose in food.

Results

Electrophysiological responses of contact chemosensilla to sugars

We first investigated the electrophysiological responses of the lateral and medial sensilla styloconica in the larval maxillary galea to eight sugars. These sugars were chosen because they are mostly found in host-plants of *H. armigera* or are representative in the structure and source of sugars. The lateral sensilla styloconica showed significant unicellular responses to 10 mM sucrose and fucose, with the firing rates of 169 ± 8.57 and 43 ± 9.78 spikes/s, respectively (**Figure 1A** [↗](#), **Figure 1—figure supplement 1A**). The threshold concentrations for their responses to sucrose and fucose were 0.1 mM and 10 mM, respectively (**Figure 1B** [↗](#), **Figure 1—figure supplement 1B**). The medial sensilla styloconica had a weak response to xylose (**Figure 1—figure supplement 1C**). We also compared the responses of the lateral sensilla styloconica to 0.05 mM sucrose, 5 mM fucose, and their mixture with the same concentrations, and found that the mixture induced higher responses with only one amplitude spikes and a firing rate of 71 ± 12.36 spikes/s, which was higher than the firing rates induced by 0.05 mM sucrose (39 ± 6.37 spikes/s) and 5 mM fucose (36 ± 8.87 spikes/s) (**Figure 1—figure supplement 1D**), confirming that the same cell responded to sucrose and fucose. We conclude that there is only one GSN sensitive to sucrose and fucose in each maxillary galea of larvae.

We also studied the responses of the contact chemosensilla in the tip of the antennae, foreleg tarsi, and proboscis of female adults to eight sugar compounds at 10 mM and the related dose responses (**Figure 1C–H** [↗](#)). The antennal sensilla responded to sucrose, fucose, and fructose positively (**Figure 1C** [↗](#)), the tarsal sensilla only responded to sucrose and fucose (**Figure 1E** [↗](#)), and the proboscis sensilla responded to sucrose, fucose, and fructose (**Figure 1G** [↗](#)), but their firing rates were between 9–22 spikes/s. All the evoked spike traces had one amplitude, indicating that they were single cell responses (**Figure 1—figure supplement 1A**). The dose–response curves showed that the threshold concentrations of sucrose, fucose and fructose were all 10 mM for the antennal sensilla (**Figure 1D** [↗](#), **Figure 1—figure supplement 1B**), 100 mM, 10 mM and 100 mM for the tarsal sensilla (**Figure 1F** [↗](#), **Figure 1—figure supplement 1B**), and 10 mM, 100 mM and 10 mM for the proboscis sensilla, respectively (**Figure 1H** [↗](#), **Figure 1—figure supplement 1B**).

In short, both larvae and adults of *H. armigera* had sugar GSNs, but their response profiles, intensity and sensitivity differed distinctively. The sugar GSNs in lateral sensilla styloconica of larvae responded strongly to sucrose with high sensitivity, and weakly to fucose; while the sugar GSNs of adults responded to sucrose, fucose, and fructose (except for those on tarsi) with lower intensity and sensitivity. The GSNs of larvae were 100–1000 times more sensitive to sucrose than those of adults.

Behavioral responses of larvae and adults to sugars

To determine how the taste inputs of various sugars cause behavioral responses in insects, the two-choice test was used to examine the feeding behavior of the 5th instar larvae of *H. armigera* to eight sugars at 10 mM. Sucrose and fructose stimulated larval feeding, and their threshold concentrations were 10 mM (**Figure 2A–B** [↗](#), **Figure 2—figure supplement 1**); other tested sugars had no effect on larval feeding (**Figure 2A** [↗](#)).

We also studied the proboscis extension reflex (PER) induced by sugar stimulation in the antennal and tarsal tip sensilla of virgin female adults. In consideration of the sugar content in nectar and the response intensity of the contact chemosensilla in adults, the test concentration of sugars was set at 100 mM. The results showed that the PER percentage reached $64 \pm 4.35\%$, $44 \pm 5.96\%$ and $20 \pm 1.92\%$ when the antennal sensilla were stimulated with sucrose, fucose and fructose (**Figure 2C** [↗](#)), while the PER percentage reached $53 \pm 14.08\%$, $45 \pm 4.44\%$ and $36 \pm 10.19\%$ when the tarsal




Figure 1.

Electrophysiological responses of larval and adult contact chemosensilla in *Helicoverpa armigera* to sugars.

(A) The maxilla morphology (left), the representative spike traces of the responses of lateral sensilla styloconica in larval maxillary galea to eight sugars at 10 mM (middle), and quantifications of the firing rates (right) ($n = 13$). The red arrow marks the recorded lateral sensillum styloconicum. Scale bar represents 20 μ m. (B) Dose-responses of lateral sensilla styloconica in larval maxillary galea to sucrose and fucose (sucrose: $n = 8$; fucose: $n = 6$). (C) The antenna tip morphology (left), the representative spike traces of the responses of sensilla chaetica in adult antennae to eight sugars at 10 mM (middle), and quantifications of the firing rates (right) ($n = 21$). The red arrows mark the recorded sensilla chaetica. Scale bar represents 100 μ m. (D) Dose-responses of sensilla chaetica in adult antennae to sucrose, fucose, and fructose ($n = 21$). (E) The fifth tarsomere morphology (left) (Zhang et al. 2010), the representative spike traces of the responses of sensilla trichodea in adult tarsi to eight sugars at 10 mM (middle), and quantifications of the firing rates (right) ($n = 15$). The red arrows mark the recorded sensilla trichodea. Scale bar represents 100 μ m. (F) Dose-responses of sensilla trichodea in adult tarsi to sucrose, fucose, and fructose ($n = 21$). (G) The proboscis tip morphology (left), the representative spike traces of the responses of sensilla styloconica in adult proboscis to eight sugars at 10 mM (middle), and quantifications of the firing rates (right) ($n = 21$). The red arrows mark the recorded sensilla styloconica. Scale bar represents 200 μ m. (H) Dose-responses to sucrose ($n = 21$), fucose ($n = 21$), and fructose (fructose 0 mM, $n = 18$; fructose 0.01 mM–100 mM, $n = 21$) of sensilla styloconica in adult proboscis. (A to H) Data are mean $\pm$ SEM; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. (A, C, E, G) Data were analyzed by independent-samples t test (compared with control). (B, D, F, H) Data were analyzed by one-way ANOVA with Tukey's HSD test.




Figure 2.

Behavioral responses of *Helicoverpa armigera* larvae and adults to sugars.

(A) Feeding responses and the preference index (PI) value of 5th instar larvae to eight sugars painted on the cabbage leaf discs at 10 mM in two-choice tests. ** $P < 0.01$; ns indicates no significance, $P > 0.05$ (paired t test, $n = 20$). (B) Feeding responses and the PI value of 5th instar larvae to different concentrations of sucrose painted on the cabbage leaf discs in two-choice tests. ** $P < 0.01$, *** $P < 0.001$; ns indicates no significance, $P > 0.05$ (paired t test, $n = 20$). (C) Proboscis extension reflex (PER) in adult females upon antennal stimulation by eight sugars at 100 mM. ** $P < 0.01$; *** $P < 0.001$ (independent-samples t test compared with control, $n = 3$). (D) PER in adult females upon tarsal stimulation by eight sugars at 100 mM. * $P < 0.05$; ** $P < 0.01$ (independent-samples t test, $n = 3$). (E) PER in adult females upon antennal stimulation by different concentrations of sucrose, fucose, and fructose. * $P < 0.05$ (one-way ANOVA with Tukey's HSD test, $n = 3$). (F) PER in adult females upon tarsal stimulation by different concentrations of sucrose, fucose, and fructose ($n = 3$). * $P < 0.05$ (one-way ANOVA with Tukey's HSD test, $n = 3$). (A to F) Data are mean ± SEM. The red arrow indicates the stimulating site.

sensilla were stimulated, respectively (**Figure 2D** [↗](#)). Stimulation of the antennal or tarsal sensilla by other sugar compounds did not induce changes in the PER percentage compared with stimulation of water (**Figure 2C–D** [↗](#)). The dose–response curves showed that the threshold concentrations of sucrose, fucose, and fructose for the antennal sensilla were 100 mM, 100 mM, and 1000 mM (**Figure 2E** [↗](#)), those for the tarsal sensilla were 10 mM, 10 mM, and 100 mM, respectively (**Figure 2F** [↗](#)).

Phylogenetic tree and expression patterns of putative sugar gustatory receptors

On the basis of understanding the electrophysiological and behavioral responses of larvae and adults of *H. armigera* to various sugars, we aimed to reveal the molecular mechanisms of sugar reception in the contact chemosensilla of *H. armigera*. We first analyzed the putative sugar gustatory receptor genes based on the genome data of *H. armigera* (Pearce et al. 2017 [↗](#)), the reported gene sequences of sugar GRs in *H. armigera* and their phylogenetic relationship of *D. melanogaster* sugar gustatory receptors (Jiang et al. 2015 [↗](#), Xu et al. 2012 [↗](#)). All nine putative sugar GR genes in *H. armigera*, *Gr4–12* were validated, and their full-length cDNA sequences were cloned (The GenBank accession number is provided in **Appendix—Table 1**). The lengths of the amino acid sequences of the sugar GRs were consistent with the lengths of the corresponding reported sequences (the similarity was 97%–99%), except for Gr8, Gr9, and Gr10. Gr8 had two more amino acids at the C-terminal than the reported by Xu et al., 2017 [↗](#) with 99% sequence similarity (Xu et al. 2017 [↗](#)). Gr9 had 9 more amino acids at the N-terminal than the reported in the *H. armigera* genome study with 99% sequence similarity (Pearce et al. 2017 [↗](#)), while the sequence length of Gr9 was exactly the same as Gr4 in Jiang et al., 2015 [↗](#) with 99% sequence similarity (Jiang et al. 2015 [↗](#)). Gr10 had 20 more amino acids than the sequence reported in the *H. armigera* genome study (Pearce et al. 2017 [↗](#)) and 186 more amino acids than the sequence reported by Xu et al., 2017 [↗](#) (Xu et al. 2017 [↗](#)), with 97% and 99% sequence similarities, respectively.

We established a phylogenetic tree using sugar GR genes of nine insect species, including *D. melanogaster*, *Aedes aegypti*, and *Anopheles gambiae* in Diptera; *Apis mellifera* in Hymenoptera; and *Danaus plexippus*, *Heliconius melpomene*, *Bombyx mori*, *Plutella xylostella*, and *H. armigera* in Lepidoptera (**Figure 3A** [↗](#)). The phylogenetic analysis showed that candidate sugar GRs of these nine insect species grouped within two clades, the sugar GR clade and the fructose GR clade. All the putative sugar GRs of *H. armigera* were in the sugar GR clade except for HarmGr9, which was in the fructose GR clade (**Figure 3A** [↗](#)).

To determine the expression patterns of nine putative sugar GR genes in larval and adult taste organs, we measured their expression levels in the larval maxillary galea, and the antennae, foreleg tarsi, and proboscis in virgin female and male adults by real-time quantitative PCR (qRT-PCR) (**Figure 3B–3E** [↗](#), **Figure 3—figure supplement 1**). The results showed that the expression patterns of sugar GRs in the same organ of male and female adults were nearly the same (**Figure 3C–3E** [↗](#), **Figure 3—figure supplement 1**). The average expression level of all sugar GRs in the larval maxillary galea was lower than that in the adult organs. In each taste organ, the expression levels of 1–2 sugar GRs were much higher than those of other sugar GRs, and these highly expressed GRs were significantly different between larvae and adults (**Figure 3B–3E** [↗](#), **Figure 3—figure supplement 1**). Among all sugar GRs, *Gr10* was highly expressed in the larval maxillary galea, whereas *Gr6* was highly expressed in the antennae, tarsi, and proboscis of both male and female adults. *Gr5* expression levels greatly varied; they were highest in the proboscis followed by the tarsi, but very low in the antennae and the maxillary galea of the larvae (**Figure 3B–E** [↗](#), **Figure 3—figure supplement 1**).




Figure 3.

The phylogenetic relationship and the expression level of sugar GRs in larval maxilla and female adult antennae, tarsi and proboscis of *Helicoverpa armigera*.

(A) The phylogenetic tree of insect sugar GRs. Diptera (orange): Aaeg, *Aedes aegypti*; Agam, *Anopheles gambiae*; Dmel, *Drosophila melanogaster*. Hymenoptera (blue): Am, *Apis mellifera*. Lepidoptera (black): Bmor, *Bombyx mori*; Dple, *Danaus plexippus*; Hmel, *Heliconius melpomene*; Pxyl, *Plutella xylostella*. Harm (red), *Helicoverpa armigera*. Numbers above branches indicate ultrafast bootstrap approximation (UFBoot). (B) Relative expression levels of sugar GRs in the maxillary galea of 5th instar larvae of *H. armigera* determined by qRT-PCR. (C) Relative expression levels of sugar GRs in the female adult antennae. (D) Relative expression levels of sugar GRs in female adult tarsi. (E) Relative expression levels of sugar GRs in female adult proboscis. (B to E) Data are mean $\pm$ SEM. One-way ANOVA was used, and different letters labeled indicate significant difference (Tukey's HSD test, $P < 0.05$, $n = 3$).

Functional analysis of putative sugar gustatory receptors by ectopic expression

Of all the nine putative sugar GRs in *H. armigera* mentioned above, only Gr9, originally named as Gr4, has been previously identified as a fructose receptor (Jiang et al. 2015 [\[1\]](#)). To explore the functions of the other eight GRs, we first investigated the inward current changes in *Xenopus* oocytes expressing sugar GRs individually in response to the stimulation of 11 sugar compounds at 100 mM using two-electrode voltage-clamp. The results showed that the oocytes expressing Gr10 only responded to sucrose with a current value of $4.10 \pm 0.53 \mu\text{A}$ (Figure 4A–B [\[2\]](#)); the oocytes expressing Gr6 showed responses to fucose and sucrose with a current value of $1.59 \pm 0.08 \mu\text{A}$ and $0.24 \pm 0.03 \mu\text{A}$, respectively (Figure 4A–B [\[2\]](#)). The oocytes expressing remaining six sugar GR genes (Gr4, Gr5, Gr7, Gr8, Gr11, and Gr12) singly had no response to the tested sugar compounds (Figure 4—figure supplement 1A–B). The dose–response curves showed that the threshold concentration of sucrose for the oocytes expressing Gr10 was 100 mM (Figure 4C–D [\[2\]](#)); whereas the threshold concentrations of sucrose and fucose for the oocytes expressing Gr6 were 250 mM and 100 mM, respectively, and 500 mM fructose could also induce an inward current (Figure 4C–D [\[2\]](#)).

Given that there was only one sugar GSN responding to sucrose and fucose in each larval maxillary galea, we speculated that Gr10 should be expressed together with Gr6 in this cell although their expressing levels were different. Therefore, we examined the response profile of the oocytes expressing Gr10 together with Gr6. The results showed that the oocytes co-expressing Gr10 and Gr6 significantly responded to sucrose with a current value of $1.89 \pm 0.58 \mu\text{A}$ at 100 mM, a threshold concentration, and had a weak response to fucose with a threshold concentration of 250 mM (Figure 4 [\[2\]](#)).

Since Gr5 and Gr6 were highly expressed in the proboscis and tarsi (Figure 3D–E [\[3\]](#), Figure 3—figure supplement 1), we suspected that Gr5 and Gr6 might be expressed in the same cells, and then tested the response profile of their co-expression in oocytes. Similar to the oocytes expressing Gr6, the oocytes expressing Gr5 and Gr6 had responses to 100 mM fucose with a current value of $0.11 \pm 0.015 \mu\text{A}$ and to 100 mM sucrose with a current value of $0.03 \pm 0.002 \mu\text{A}$ (Figure 4—figure supplement 1B–C).

In summary, we find that oocytes expressing single Gr10 or Gr6 confer sensitivity to one or more sugars, indicating their direct roles in ligand recognition. Gr10 is tuned to sucrose specifically, Gr6 is responsive to fucose, sucrose and fructose in the order of sensitivity, and the combination of Gr10 and Gr6 is sensitive to sucrose and then fucose, which is in line with the characteristics of the sugar GSNs in larval and adult taste organs where the two GRs are highly expressed respectively.

Establishment of *Gr10* and *Gr6* homozygous mutants

In the above study, we discovered ligands of Gr10 and Gr6 using the ectopic expression system. To validate the function of *Gr10* and *Gr6* *in vivo*, we further constructed *Gr10* and *Gr6* homozygous mutants (*Gr10*^{−/−} and *Gr6*^{−/−}, respectively) using CRISPR/Cas9 techniques (Figure 5 [\[4\]](#)). *Gr10*^{−/−} and *Gr6*^{−/−} generated 4-bp and 17-bp deletions at the sgRNA targeting site (Figure 5 [\[4\]](#)), respectively, which also confirmed the deletion at the mRNA level in *Gr10*^{−/−} and *Gr6*^{−/−} (Figure 5—figure supplement 1). The deletion resulted in truncated Gr10 and Gr6 proteins. The Gr10 sequence was reduced from 460 to 62 amino acids and contained no transmembrane domains, whereas the Gr6 sequence was reduced from 452 to 115 amino acids and contained only two transmembrane domains (Figure 5 [\[4\]](#)).

In order to predict the potential off-target sites of sgRNA of *Gr6* and *Gr10*, we used online software Cas-OFFinder (<http://www.rgenome.net/cas-offinder/> [\[5\]](#)) to blast the genome of *H. armigera*, and the mismatch number was set to less than or equal to 3. According to the predicted results, the *Gr10* sgRNA had no potential off-target region but *Gr6* sgRNA had one. Therefore, we amplified and




Figure 4.

The inward current responses of the *Xenopus* oocytes expressing sugar GRs of *Helicoverpa armigera* to sugar compounds.

(A) The representative traces of the oocytes expressing Gr10, Gr6, and Gr10 + Gr6 to 11 sugar compounds at 100 mM. (B) The responses of the oocytes expressing Gr10, Gr6, and Gr10 + Gr6 to sugars at 100 mM (Gr10, $n = 14$. Gr6, $n = 20$. Gr10 + Gr6, $n = 7$). (C) The representative trace of the oocytes expressing Gr10 to sucrose (the upper), the representative trace of the oocytes expressing Gr6 to sucrose, fucose, and fructose (the middle), the representative trace of the oocytes expressing Gr10 + Gr6 to sucrose and fucose (the lower). (D) The dose-responses of the oocytes expressing Gr10 to sucrose (the upper, $n = 13$); the dose-responses of the oocytes expressing Gr6 to sucrose, fucose, and fructose (the middle, sucrose: $n = 6$; fucose: $n = 6$; fructose: $n = 3$); the dose-responses of the oocytes expressing Gr10 + Gr6 to sucrose and fucose (the lower, sucrose: $n = 4$; fucose: $n = 3$). (B and D) Data are mean $\pm$ SEM, and were analyzed by one-way ANOVA with Tukey's HSD test ($P < 0.05$). Different letters labeled indicate significant differences, $* P < 0.05$. (A to D) Ara, arabinose; Fru, fructose; Fuc, fucose; Gal, galactose; Glu, glucose; Lac, lactose; Mal, maltose; Man, mannose; Suc, sucrose; Tre, trehalose; Xyl, xylose; Rin, ringer solution.




Figure 5.

Establishment of *Gr10* and *Gr6* homozygous mutants (*Gr10*^{-/-} and *Gr6*^{-/-}) of *Helicoverpa armigera* via CRISPR/Cas9.

(A) The cross process of obtaining *Gr10*^{-/-}. (B) The genomic structure of *Gr10*, the single-guide RNA (sgRNA) targeting sequence (in green), and representative chromatograms of direct sequencing of the PCR products obtained from wild types (WT) and *Gr10*^{-/-}, in which 4-bp of the *Gr10* sequence were deleted. (C) The predicted secondary structures of the *Gr10* protein in WT and the truncated *Gr10* protein in *Gr10*^{-/-}. (D) The cross process of obtaining *Gr6*^{-/-}. (E) The genomic structure of *Gr6*, the single-guide RNA (sgRNA) targeting sequence (in green), and representative chromatograms of direct sequencing of the PCR products obtained from WT and *Gr6*^{-/-}, in which 17-bp of the *Gr6* sequence were deleted. (F) The predicted secondary structures of the *Gr6* protein in WT and the truncated *Gr6* protein in *Gr6*^{-/-}. (B and D) Boxes represent exons, black lines represent introns, the green arrowhead indicates the direction of sgRNA; the protospacer adjacent motif (PAM) is in blue. (C and F) The secondary structure of *Gr10* and *Gr6* in WT, *Gr10*^{-/-} and *Gr6*^{-/-} was predicted by <https://dtu.biolib.com/DeepTMHMM>, and the image was constructed by TOPO2 software (<http://www.sacs.ucsf.edu/TOPO2>).

sequenced the potential off-target region of *Gr6*^{-/-} and found there was no frameshift or premature stop codon in the region compared to WT (**Figure 5—figure supplement 2**). It is worth mentioning that there was no potential off-target region of *Gr6* sgRNA or *Gr10* sgRNA in all the other eight sugar receptor genes of *H. armigera*. We further found there was no difference in the response to xylose of the medial sensilla styloconica among WT, *Gr10*^{-/-} and *Gr6*^{-/-} (**Figure 5—figure supplement 2**). Furthermore, WT, *Gr10*^{-/-} and *Gr6*^{-/-} did not show differences in the larval body weight, adult lifespan, and number of eggs laid per female (**Figure 5—figure supplement 2**). All these results suggest that no off-target effects occurred in the study.

Effects of *Gr10* or *Gr6* knockout on sugar reception

After obtaining the two homozygous mutants, we compared electrophysiological responses of the sugar GSNs in larval lateral sensilla styloconica of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} to sucrose and fucose using 10 mM sinigrin as a positive control, which induced the response of a bitter cell in the same sensilla. To 1 mM, 10 mM and 100 mM sucrose, the firing rates of the larval sugar GSNs in *Gr10*^{-/-} were only 40%, 20% and 28% of those in the WT, respectively, whereas those in *Gr6*^{-/-} larvae did not differ from those in the WT larvae (**Figure 6A–B**). To 10 mM and 100 mM fucose, the firing rates of the larval sugar GSNs in *Gr6*^{-/-} were only 20% and 5.8% of those in the WT, respectively, whereas those of *Gr10*^{-/-} did not differ from the WT (**Figure 6A–B**). The electrophysiological responses of the bitter GSNs to sinigrin in the larval lateral sensilla styloconica of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} were all strong and did not differ (**Figure 6A–B**). It is clear that *Gr10* plays a key role in sucrose reception whereas *Gr6* plays a key role in fucose reception of the sugar GSNs in the lateral sensilla styloconica of larvae.

Similarly, we compared the electrophysiological responses of the sugar GSNs in contact chemosensilla of the adult antennae, foreleg tarsi, and proboscis of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} to 10 mM and 100 mM of sucrose and fucose, and 100 mM fructose. Additionally, 10 mM nicotine was used as a positive control, that induced electrophysiological responses of bitter GSNs in the tip sensilla of antennae and tarsi. The firing rates of the sugar GSNs in the antennal sensilla of *Gr6*^{-/-} to 10 mM and 100 mM sucrose, 10 mM and 100 mM fucose, and 100 mM fructose were only 32% and 9%, 22% and 11%, and 29% of those in the WT, respectively (**Figure 6C–D**). The firing rates of the sugar GSNs in antennal sensilla of *Gr10*^{-/-} to sucrose, fucose, or fructose did not differ from those of the WT adults. The antennal sensilla of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} all showed strong responses to nicotine with no differences (**Figure 6C–D**). The response patterns of the sugar GSNs in the foreleg tarsi of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} to 10 mM and 100 mM sucrose, 10 mM and 100 mM fucose, and 100 mM fructose were similar to those of the sugar GSNs in antennae (**Figure 6E–F**). The firing rates of the sugar GSNs in proboscis sensilla of *Gr6*^{-/-} to sucrose, fucose, and fructose were much lower and even close to zero (**Figure 6G–H**). The firing rates of the sugar GSNs in tarsal and proboscis sensilla of *Gr10*^{-/-} to sucrose, fucose, and fructose did not differ from those of the WT adults (**Figure 6E–H**). The tarsal sensilla of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} responded to nicotine with no difference (**Figure 6E–F**).

Effects of *Gr10* and *Gr6* knockout on behavioral responses

To determine the effect of *Gr10* and *Gr6* knockout on feeding responses of *H. armigera* larvae, we compared the larval feeding preferences of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} to sucrose by two-choice tests. The *Gr10*^{-/-} larvae showed no preference for the leaf discs painted with 10 mM sucrose ($P = 0.4258$). However, they still had preference for the discs with 100 mM sucrose ($P = 0.0141$) (**Figure 7A–B**) though their PI value was decreased than those of the WT (**Figure 7B**). The *Gr6*^{-/-} larvae were the same as the WT larvae and retained a significant preference for 10 mM or 100 mM sucrose diets (**Figure 7A–B**). The WT, *Gr10*^{-/-}, and *Gr6*^{-/-} larvae all showed similar preference for 10 mM and 100 mM fructose (**Figure 7—figure supplement 1A–B**).




Figure 6.

Electrophysiological responses of larval and adult contact chemosensilla in WT, *Gr10*^{-/-} and *Gr6*^{-/-} of *Helicoverpa armigera* to sucrose and other compounds.

(A) The representative spike traces of lateral sensilla styloconica in larval maxillary galea. (B) Quantifications of the firing rates of lateral sensilla styloconica in larval maxillary galea (mean ± SEM; WT, *n* = 12; *Gr6*^{-/-}, *n* = 13; *Gr10*^{-/-}, *n* = 13). (C) The representative spike traces of sensilla chaetica in female antennae. (D) Quantifications of the firing rates of sensilla chaetica in female antennae (mean ± SEM; WT, *n* = 20. *Gr6*^{-/-}: nicotine, *n* = 17; other compounds, *n* = 18. *Gr10*^{-/-}, *n* = 20). (E) The representative spike traces of sensilla trichodea in female tarsi. (F) Quantifications of the firing rates of sensilla trichodea in female tarsi (mean ± SEM; WT, *n* = 18; *Gr6*^{-/-}: nicotine, *n* = 14. other compounds, *n* = 18. *Gr10*^{-/-}, *n* = 18). (G) The representative spike traces of sensilla styloconica in female proboscis. (H) Quantifications of the firing rates of sensilla styloconica in female proboscis (mean ± SEM; WT: sucrose 100 mM, *n* = 17; other compounds, *n* = 18. *Gr6*^{-/-}: sucrose 100 mM, *n* = 17; other compounds, *n* = 18. *Gr10*^{-/-}, *n* = 18). (B, D, F, and H) Two-way ANOVA with post hoc Tukey's multiple comparison was used separately for sucrose and fucose, and one-way ANOVA with Tukey's HSD test was used for KCl, sinigrin, fructose, and nicotine. Different letters labeled indicate significant differences (*P* < 0.05). Suc: sucrose; Fuc: fucose; Sin: sinigrin; Fru: fructose; Nic: nicotine.




Figure 7.

Behavioral responses of WT, *Gr10*^{-/-} and *Gr6*^{-/-} larvae and adults of *Helicoverpa armigera* to sugars and plant tissues.

(A) Feeding area of 5th instar larvae in two-choice tests and the PI value to 10 mM sucrose ($n = 20$). ** $P < 0.01$; ns indicates no significance, $P > 0.05$ (paired t test). Suc, sucrose. (B) Feeding area of 5th instar larvae in two-choice tests and the PI value to 100 mM sucrose ($n = 20$). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ (paired t test). Suc, sucrose. (C) Feeding amount of 5th instar larvae on cabbage leaves, corn kernels, pea seeds, pepper fruits, and tomato fruits in no-choice tests ($n = 20$). Data were analyzed by one-way ANOVA with Tukey's HSD test, and different letters labeled on the data of WT, *Gr10*^{-/-} and *Gr6*^{-/-} for each plant tissue indicate significant differences ($P < 0.05$). Proboscis extension reflex (PER) in adult females upon (D) antennal stimulation by sucrose concentrations ($n = 3$), (E) antennal stimulation by fucose concentrations ($n = 3$), (F) antennal stimulation by fructose concentrations ($n = 3$), (G) tarsal stimulation by sucrose concentrations ($n = 3$), (H) tarsal stimulation by fucose concentrations ($n = 3$), and (I) tarsal stimulation by fructose concentrations ($n = 3$). (A to I) Data are mean $\pm$ SEM. (D to I) Data were analyzed by two-way ANOVA with post hoc Tukey's multiple comparison. * $P < 0.05$. The red arrow indicates the stimulating site.

We also compared the feeding amount in 1 hour of the larvae of WT, *Gr10*^{-/-}, and *Gr6*^{-/-} on cabbage leaves, corn kernels, pea seeds, pepper fruits, and tomato fruits by no-choice tests. On each of these plant tissues, the *Gr10*^{-/-} larvae fed significantly less than the WT larvae, but the *Gr6*^{-/-} larvae fed the same amount as the WT larvae (**Figure 7C** [↗](#)).

Moreover, we examined the PER of the WT, *Gr10*^{-/-}, and *Gr6*^{-/-} female adult antennae and tarsi in response to stimulation of sucrose, fucose, and fructose at different concentrations. When antennae or tarsi were stimulated with 10 mM, 100 mM, or 1000 mM sucrose, fucose, or fructose, *Gr6*^{-/-} females showed a considerably lower PER percentage than the WT females and even had little or no PER in response to sucrose and fucose, whereas the PER percentage of *Gr10*^{-/-} females was not different from that of the WT females (**Figure 7D–I** [↗](#)).

In conclusion, *Gr10* plays a key role in sucrose reception by the sugar GSNs in the larval lateral sensilla styloconica, and mediates the feeding preference of larvae to sucrose, whereas *Gr6* plays a key role in sensing sucrose, fucose, and fructose by the sugar GSNs in contact chemosensilla of the adult antennae, foreleg tarsi and proboscis, and mediates the related PER behavior of adults.

Discussion

In this study, we reveal the different molecular bases of sucrose reception between *H. armigera* larvae and adults through ectopic expression and genetic editing of GRs. Gr10, which was highly expressed in the larval maxillary galea, is specifically tuned to sucrose with high sensitivity, whereas Gr6, which was highly expressed in the adult antennae, foreleg tarsi and proboscis, has a broader response spectrum but lower sensitivity to sucrose, indicating that larvae mainly use Gr10 to detect low sucrose in plant tissues, while adults primarily use Gr6 to detect a variety of sugars with high content, including sucrose in nectar (**Figure 8** [↗](#)).

Our findings indicate that the *H. armigera* larvae and adults adopt different sugar-sensing systems to satisfy their requirements for food selection. The larvae mainly feed on young leaves, flower buds, and fruits of host plants. As a universal phagostimulant, sucrose is the major sugar in plants and considerably varies in content in different tissues and developmental stages. Tomato is a principal host plant of *H. armigera*. The sucrose content in its leaves generally ranges from 13.72 to 42.06 mM, but that in its fruits generally ranges from 35.2 to 988.22 mM (Schauer et al. 2005 [↗](#)). The adults mainly feed on plant nectar. Sucrose and some monosaccharides, especially fructose and glucose, are the main components in nectar, where the sugar content is much higher than that in leaves and fruits. The sugar content in the nectar of cotton plants is 10%–34% (which roughly corresponds to 0.5 M–1.7 M) (Vansell 1944 [↗](#), Waller et al. 1981 [↗](#)), and that of soybean plants is even higher, around 23%–60% (which roughly corresponds to 1.2–3 M) (Free 1993 [↗](#)). It is clear that, in their life cycle, a single sugar-sensing mechanism cannot meet the requirements for food selection by this phytophagous insect species. From the perspective of insect and plant co-evolution, using value-based and target-specific sugar-sensing systems is a favorable choice for phytophagous insects to ensure their growth and reproduction on plants.

Although Gr10 plays a major role in sucrose reception in the larval stage, the response of the sugar GSNs in larval lateral sensilla styloconica to sucrose did not completely disappear in *Gr10*^{-/-} (**Figure 6A–B** [↗](#)), and the larvae still showed a behavioral preference for the high concentration of sucrose (100 mM) (**Figure 7B** [↗](#)). It is most likely that low-level expressed Gr6 is functional in the GSNs. The PER percentage of *Gr6*^{-/-} adults was not disappeared when the antennae and tarsi were stimulated by high fructose concentrations (**Figure 7F** [↗](#) and **7I** [↗](#)); this indicates that fructose-tuned GRs other than Gr6 were involved. Previous studies have shown that Gr9 (Gr4 in Jiang et al., 2015 [↗](#)), as an ortholog of DmelGr43a and BmorGr9, is a fructose-tuned GR (Ai et al. 2022 [↗](#), Jiang et al. 2015 [↗](#), Sato et al. 2011 [↗](#), Xu et al. 2012 [↗](#)). From our results, knocking out *Gr10* or *Gr6* is unlikely to be compensated by overexpression of other sugar GRs. One of our recent studies




Figure 8.

Two gustatory receptors in the cotton bollworm, *Helicoverpa armigera* mainly mediate taste sensation of sugars in the larval and adult foods

showed that *Orco* knockout had no significant effect on the expression of most OR, IR and GR genes in adult antennae of *H. armigera* although some genes were up- or down-regulated to a certain extent (Fan et al. 2022 [DOI](#)). To elucidate the precise molecular mechanisms of sugar reception in *H. armigera* is necessary to compare a series of single, double and even multiple Gr knock-out lines and investigate the downstream effects of the GRs.

The larval maxillary galea with only two sensilla styloconica is an ideal system for studying the mechanisms of sugar sensing. In each maxillary galea of *H. armigera*, the only sugar GSN sensitive to sucrose and fucose is located in the lateral sensillum styloconicum. Based on our results, we speculate that Gr10 and Gr6 are expressed in this GSN with different levels, and mainly determine the response profile of the cell. The firing rate of the cell in *Gr10*^{-/-} to 1mM, 10 mM and 100 mM sucrose was significantly lower than that in WT, but there was no difference between WT and *Gr6*^{-/-}, indicating that Gr10 mainly determine the sensitivity of this cell to sucrose (**Figure 6A-B** [DOI](#)). In contrast, the firing rate of the cell in *Gr6*^{-/-} to 10 mM and 100 mM fucose was significantly lower than that in the WT, but there was no difference between WT and *Gr10*^{-/-}, indicating that Gr6 was also functional in larvae (**Figure 6A-B** [DOI](#)). This can be well explained by the response profiles of the oocytes expressing Gr10, Gr6, and their combination (**Figure 4** [DOI](#)).

Sugars are common phagostimulants in animals, but the molecular basis of sugar perception differs among animals. In mammals, the taste receptors T1R2 and T1R3 combine to complete the perception of all carbohydrate compounds, including sucrose, and both receptors are indispensable (Damak et al. 2003 [DOI](#), Li et al. 2002 [DOI](#), Nelson et al. 2001 [DOI](#), Zhao et al. 2003 [DOI](#)). In *D. melanogaster* adults, multiple sugar GRs are involved in sugar reception, different GRs expressed in sweet taste neurons may function as multimers (Dahanukar et al. 2001 [DOI](#), Dahanukar et al. 2007 [DOI](#), Fujii et al. 2015 [DOI](#), Jiao et al. 2007 [DOI](#), Jiao et al. 2008 [DOI](#), Slone et al. 2007 [DOI](#)). Gr5a and Gr64a are widely used for responses to complementary subsets of sugars in labellar sweet taste neurons (Dahanukar et al. 2007 [DOI](#)), and Gr64f is also required broadly as a coreceptor for the detection of sugars (Jiao et al. 2008 [DOI](#)). The recent successful ectopic expression of single sugar GRs in olfactory neurons confirmed that each sugar GR is directly involved in ligand recognition (Freeman et al. 2014 [DOI](#)). Sweet sensing neurons or receptors have not been identified in the larval peripheral taste system (Kwon et al. 2011 [DOI](#)). Sugar perception in *Drosophila* larvae has been assigned to the fructose receptor *Gr43a*, which, however, is expressed in pharyngeal sensory organs, foregut, and in the brain, but absent from external taste neurons (Mishra et al. 2013 [DOI](#)). This is most likely due to the fact that fruit fly larvae prefer to feed on rotting fruits at the optimum stage of fermentation, which contain fructose as the main sugar. Anyway, the sucrose sensing neurons were recently identified in the larval primary taste center (Maier et al. 2021 [DOI](#)). In honeybees, AmGr1 acts as a GR for certain sugars other than fructose, and AmGr2 acts as a coreceptor with AmGr1 (Jung et al. 2015 [DOI](#)). AmGr3 is tuned to fructose and is phylogenetically closely related to the DmGr43a (Değirmenci et al. 2020 [DOI](#), Takada et al. 2018 [DOI](#)). The whitefly *Bemisia tabaci* consistently choose diets containing higher sucrose concentrations. It has been shown that BtabGr1 displayed significant sucrose specificity when expressed in *Xenopus* oocytes, and silencing of this GR significantly interfered with the ability of *B. tabaci* adults to detect sucrose in the phloem sap (Harari et al. 2023 [DOI](#)).

The most studied phytophagous insects for sugar perception are Lepidopterans. BmGr9 in *B. mori* (Morinaga et al. 2022 [DOI](#), Sato et al. 2011 [DOI](#)), HarmGr9 in *H. armigera* (named as HarmGr4 by Jiang et al., 2015 [DOI](#)) (Jiang et al. 2015 [DOI](#), Xu et al. 2012 [DOI](#)), PxGr43a-1 in *P. xylostella* (Liu et al. 2020 [DOI](#)), and SlGr8 in *Spodoptera litura* (Liu et al. 2019 [DOI](#)) were all the orthologs of DmGr43a in *D. melanogaster* (Miyamoto et al. 2012 [DOI](#)), and tuned to fructose without complexing with other GRs. BmGr4 was recently identified as a receptor for sucrose and glucose and may function as a chemosensor for nutritional compounds in both midgut and brain (Mang et al. 2022 [DOI](#)). In this study, we find *Gr10* and *Gr6* were highly expressed in the larval and adult taste organs, respectively. The experiments on ectopic expression in *Xenopus* oocytes shows that Gr10 specifically responded to sucrose, Gr6 responded to fucose and sucrose, and weakly to fructose,

and the combination of Gr10 and Gr6 responded to sucrose and weakly to fucose (**Figure 4A–B**). These results indicate that Gr10 and Gr6 are capable of forming functional receptors singly, and the function of the two co-expressed receptors is not equal to the sum of the functions of the two receptors (**Figure 4**). When further tracing the genomic locations of the two GR genes, we confirm that *Gr10* and *Gr6* together with other 6 sugar GR genes (except for *Gr9*) are tandemly duplicated on the same scaffold of *H. armigera* (Pearce et al. 2017). The sugar GR gene cluster and interactions between GRs could provide a degree of evolutionary and regulatory flexibility that may meet the needs of the insect to sense subtle sugar components and concentrations in food sources in a changing environment (Delventhal and Carlson 2016). In this study, the expression patterns of 9 sugar GRs in three taste organs of adult *H. armigera* show that there is a disparity in GRs, specifically GR5 and GR6, between the female antenna, tarsi and proboscis, which may be an evolutionary adaptation reflecting subtle differentiation in the function of these taste organs in adult foraging. Antennae and tarsi play a role in the exploration of potential sugar sources, while the proboscis plays a more precise role in the final decision to feed.

In this work, we report for the first time the molecular basis of sucrose reception in external taste neurons of a Lepidopteran insect, and discover that different taste receptors are used for food selection in *H. armigera* larval and adult stages. These findings greatly improve our understanding of “sweet” taste of phytophagous insects. According to the phylogenetic tree of sugar GRs in insects, *Gr10* is clustered in the same branch with BmorGr5 of *Bombyx mori*, whereas *Gr6* is clustered in another branch with BmorGr6 (**Figure 3A**). GRs closely associated with *Gr10* and *Gr6* are also found in other Lepidopteran species such as *D. plexippus*, *H. melpomene*, and *P. xylostella* (**Figure 3A**). Therefore, we speculate that similar sugar-sensing mechanisms may also exist in other Lepidopteran species, which is worth verifying in these species in the future. Knockout of *Gr10* or *Gr6* led to a significant decrease in sugar sensitivity and food preference of the larvae and adults of *H. armigera*, respectively, which is bound to bring adverse consequences to the survival and reproduction of the insects. Therefore, studying the molecular mechanisms underlying sugar perception in phytophagous insects would provide new insights into the behavioral ecology of this highly diverse and important group of insects, and may also provide new means for agricultural pest control by blocking or disrupting sugar receptors.

Materials and Methods

Insects

Helicoverpa armigera were collected from Xuchang (Henan Province, China) and raised in an indoor environment ($27 \pm 1^\circ\text{C}$, 70%–80% humidity) with a 16 : 8 h light : dark cycle. The larvae fed on artificial diet (mainly including wheat bran, wheat germ, and tomato paste) in a glass tube (I.D., 2.5 cm; length, 8 cm). When the larvae reached the 3rd instar, they were transferred to a new glass tube and raised individually until pupation. Female and male moths were raised separately until they became sexually mature and then moved to a cage for mating. Since the taste system of the female is similar to that of the male and the nutrient requirements are more critical for females (Jiang et al. 2015, Zhang et al. 2010), the virgin female adults were used in the related experiments.

Xenopus laevis

Xenopus laevis were fostered at $20 \pm 1^\circ\text{C}$ in the Animal Laboratory Center, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences. They fed on artificial frog food (mainly including fish meal, yeast extract powder, high-gluten flour). All experiments were approved by the Animal Care and Use Committee of the Institute of Zoology, Chinese Academy of Sciences and followed the Guide for the Care and Use of Laboratory Animals (IOZ17090-A).

Scanning electron microscopy

Just as our previous study on tarsal contact chemosensilla in fore legs of female moths (Zhang et al. 2010), the samples of antennae or proboscis in adult female *H. armigera* were mounted directly on stainless steel sample buds and sputter-coated with a 10 nm thick layer of gold. Photomicrographs were obtained with a scanning electron microscope (HITACHI SU8010) in the Institute of Microbiology, Chinese Academy of Sciences.

Tip-recording

The tip-recording technique based on a previously described methods (Hodgson et al. 1955, van Loon 1990, Zhang et al. 2010) was used to record the electrophysiological responses of lateral and medial sensilla styloconica in larval maxillary galea, sensilla chaetica in adult antennae, sensilla trichodea in adult tarsi, and sensilla styloconica in adult proboscis of *H. armigera*. A glass capillary filled with the test compound solution, into which a silver wire was inserted, was placed in contact with the contact chemosensillum to be recorded as the recording electrode. The tip diameter of the electrode was about 50 μm , suitable for recording from a single sensillum. Action potentials (spikes) generated during the first second after stimulus onset were amplified by a preamplifier and was sampled with a computer equipped with a Metrabyte DAS16 A/D conversion board. The amplifier used an AD 515-K (Analog Devices) integrated circuit in the first stage, yielding < 1 pA input bias current, 1015 Ohm and 0.8 pF input impedance. An interface (GO-box) was used for signal conditioning. This involved a second-order band pass filter (~ 3 dB frequencies: 180 and 1700 Hz). Stimuli were dissolved in 2 mM KCl, which was used as the control. Two pentoses (arabinose and xylose), four hexoses (fructose, fucose, galactose and glucose), and two disaccharides (sucrose and trehalose) (**Key resources table**) were used as sugar stimuli. Each sugar first tested at 10 mM to screen active compounds, which were then tested in a range of concentrations in dose-response assays (0.01 mM, 0.1 mM, 1 mM, 10 mM and 100 mM). To avoid possible adaptation of the contact chemosensilla tested, the interval between two successive stimulations was at least 3 min. Digitized traces were analyzed by means of SAPID Tools software version 3.5 (Smith et al. 1990) and Autospike 3.7 (Syntech, Buchenbach, Germany). The spikes were sorted on the basis of typical biphasic waveforms and spike amplitudes, and the frequency of the same type of spikes was counted from the first second after stimulation.

In recording of electrophysiological responses of larvae, the first day of larvae in the 5th instar were used. To guarantee all test caterpillars were in the same stage, we first selected 4th instar larvae which were preparing to undergo ecdysis, and then transferred them to a glass tube with green pepper. The test larva was starved for approximately 2 h before recording. The head was cut and fixed in a silver holder which was connected to a preamplifier with a copper miniconnector. The lateral and medial sensilla styloconica in larval maxillary galea were exposed for testing electrophysiological responses. About 12–13 replicates were set for each treatment.

The virgin female moths on the second day after eclosion were used for recording electrophysiological responses (Jiang et al. 2015, Zhang et al. 2010). They were only fed with ddH₂O. When testing sensilla in a tarsus, the foreleg was cut between coxa and femur, and the silver holder connected to the amplifier was inserted into the femur. When testing sensilla in the antennae and proboscis, the head and thorax without all legs were reserved. The silver holder was inserted into the thorax. The orientation of the taste organ was gently adjusted to allow the sensilla to be touched by the electrode. To guarantee the sensilla was steadily exposed, the distal part of antenna or proboscis was fixed on a small platform with double sided adhesive tape. For each individual, three sensilla on the top area of antennae, foreleg tarsi, or proboscis were chosen for testing electrophysiological responses, and about 15–21 replicates were used for each treatment.

Two-choice tests

We determined the behavioral responses to sugars of the *H. armigera* larvae using the two-choice test as previously described with small modifications (Chen et al. 2022 [↗](#)). We used the leaf of cabbage (*Brassica oleracea*, Shenglv-7) as the substrate since it is not a favorite food for larvae. Circular cabbage leaf discs of 1 cm in diameter were prepared. The upper surface of the treated leaf disc was painted with the 35.2 μ L sugar solution with 0.01% Tween 80 (CAS: 9005-55-6, Sigma-Aldrich, St Louis, USA) using a paintbrush, and the control leaf disc was painted with 35.2 μ L ddH₂O with 0.01% Tween 80. The 0.01% Tween 80 was added to reduce the surface tension of the cabbage leaves and increase the malleability of the test sugar solution (Blaney and Simmonds 1988 [↗](#)). The feeding preference of larvae to eight sugars including arabinose, fructose, fucose, galactose, glucose, sucrose, trehalose and xylose (**Key resources table**) was tested at 10 mM. 0.01 mM, 0.1 mM, 1 mM, 10 mM, and 100 mM sucrose and fructose were used in a dose-response assay. The intact and the remaining leaf discs were scanned by DR-F120 scanner (Canon, Tokyo, Japan), and the area of intact and remaining leaf discs were calculated by ImageJ software (NIH) (Abràmoff et al. 2004 [↗](#), Chen et al. 2022 [↗](#)). The feeding area = the area of intact leaf disc – the area of remaining leaf disc. The preference index (PI) was calculated according to the following formula: $PI = \frac{\text{the feeding area of test leaf discs} - \text{the feeding area of control leaf discs}}{\text{the feeding area of test leaf discs} + \text{the feeding area of control leaf discs}}$. 20 replicates were run.

Adult proboscis extension reflex (PER) tests

We determined the behavioral responses to sugars of the *H. armigera* female adults using the proboscis extension reflex (PER) test as previously described with some modifications (Jiang et al. 2015 [↗](#)). Arabinose, fructose, fucose, galactose, glucose, sucrose, trehalose, and xylose (**Key resources table**) were first tested at 100 mM to screen active compounds, which were then tested in a range of concentrations in dose-response assays (0.01 mM, 0.1 mM, 1 mM, 10 mM, 100 mM, and 1000 mM). In brief, the virgin females on the first day after eclosion were used. Each moth was harnessed in a 1 mL pipette with a tip removed, from which its head protruded, and its antennae, proboscis and forelegs were permitted to move freely. They were allowed to adapt to the environment for about 12 h, and satiated with ddH₂O until the trial. The testing moth was stimulated to the distal part of one antenna with a given test solution in an absorbent cotton ball, and the PER in response to the stimulation was recorded. The interval between two successive stimulations was 5 min, during which the distal part of the antenna was washed, and the insect was fed with ddH₂O if needed. Four hours later, two foreleg tarsi of the moth were simultaneously stimulated with the test solutions in the same way as the antenna was stimulated, and the PER was recorded. 20 to 30 individual moths were used for each set of tests. The experiments were repeated three times with groups of insects from three different rearing batches.

RNA extraction and cDNA synthesis

The galea of larvae, and the antennae, foreleg tarsi, proboscis, and abdomen of both male and female adults were removed and directly placed into 1.5 mL Eppendorf tubes (Axygen, Tewksbury, MA, USA) containing 1 mL QIAzol Lysis Reagent (QIAGEN, Hilden, Germany); then, the extraction of total RNA was carried out following the manufacturer's protocols. After extraction, the RNA quality and concentration were determined with a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). First, for cDNA synthesis 2 μ g total RNA was mixed with 1 μ L oligo-dT (Promega, Madison, WI, USA); then, the mixture was heated at 70°C 5 min and followed by an ice bath for 5 min. M-MLV reverse transcriptase (Promega, Madison, WI, USA) was added; subsequently, the mixture was incubated at 42°C for 90 min. The products were dissolved in ddH₂O and stored at –20°C for later use.

Gene cloning of sugar GRs

We designed specific primers (**Appendix—Table 2**) according to the reported sugar GR sequence of *H. armigera* (the GenBank accession number is provided in **Appendix—Table 2**) for PCR cloning (Pearce et al. 2017 [Xu et al. 2017](#)). The GenBank accession number of *Gr9* (named as *Gr4* in Jiang et al., 2015 [Jiang et al., 2015](#)) nucleic acid sequence and protein sequence were JX982536.2 and AKG90011 in Jiang et al., 2015 [Jiang et al., 2015](#), respectively (Jiang et al. 2015 [Jiang et al., 2015](#)). The Q5 High-fidelity DNA Polymerase (New England Biolabs, Beverly, MA, USA) was used to clone sugar sequences except *Gr10*. The PCR instrument was Applied Biosystems (Thermo Fisher Scientific, Woodlands, Singapore) and the sequence amplification procedure was as follows: 98°C for 10 s, 40 cycles (60°C for 30 s, 72°C for 1 min), and final extension at 72°C for 2 min. TransStart FastPfu DNA Polymerase (TransGen Biotech, Beijing, China) was used to clone the *Gr10* sequence. The amplification procedure was as follows: 95°C for 1 min, 40 cycles (95°C for 20 s, 50°C for 20 s, 72°C for 45 s), and 72°C for 5 min. PCR products were purified, subcloned into pGEM-T vector according to the manufacturer's instructions (Promega), and sequenced. *Gr4* (402 amino acids) and *Gr7* (428 amino acids) were obtained from the tissues of male and female *H. armigera* tarsi, respectively. *Gr5* (482 amino acids), *Gr6* (452 amino acids), and *Gr10* (460 amino acids) were all obtained from the tissues of female *H. armigera* proboscis. *Gr8* (427 amino acids) and *Gr9* (474 amino acids) was obtained from the tissues of female *H. armigera* antennae. Finally, *Gr11* (435 amino acids) and *Gr12* (434 amino acids) were obtained from the tissues of female and male *H. armigera* abdomens, respectively. No alternatively spliced transcript variants were found in the gene cloning of all the sugar GRs studied.

Xu et al., 2017 [Xu et al. 2017](#) reported that *Gr10* was a pseudogene because the nucleotide sequence changed at positions 521–524 compared with the sequence reported in Pearce et al., 2017 [Pearce et al. 2017](#), and it terminated early during translation. The *Gr10* sequence that we obtained had no termination on the same position. Aligning the *Gr10* sequences of Pearce et al., 2017 [Pearce et al. 2017](#) and Xu et al., 2017 revealed that *Gr10* in Xu et al., 2017 [Xu et al., 2017](#) had 22 more amino acids at the C-terminus than that in Pearce et al., 2017 [Pearce et al., 2017](#). Therefore, we designed upstream and downstream primers of *Gr10* from the N-terminus in Pearce et al., 2017 [Pearce et al., 2017](#) and the C-terminus in Xu et al., 2017 [Xu et al., 2017](#). Finally, we obtained the full-length sequence of *Gr10* with 460 amino acids.

Phylogenetic analysis of sugar GRs

The phylogenetic tree was built with *H. armigera* and sugar GRs of eight other insect species: Diptera, *Drosophila melanogaster* (Robertson et al. 2003 [Robertson et al. 2003](#)), *Aedes aegypti* (Kent et al. 2008 [Kent et al. 2008](#)), and *Anopheles gambiae* (Hill et al. 2002 [Hill et al. 2002](#)); Hymenoptera, *Apis mellifera* (Robertson and Wanner 2006 [Robertson and Wanner 2006](#)); and Lepidoptera, *Danaus plexippus* (Zhan et al. 2011 [Zhan et al. 2011](#)), *Heliconius melpomene* (Briscoe et al. 2013 [Briscoe et al. 2013](#)), *Bombyx mori* (Guo et al. 2017 [Guo et al. 2017](#)), and *Plutella xylostella* (Engsontia et al. 2014 [Engsontia et al. 2014](#)). Amino acid sequences were aligned with MAFFT version 7.455 (Rozewicki et al. 2019 [Rozewicki et al. 2019](#)), and gap sites were removed with trimAI version 1.4 (Capella-Gutierrez et al. 2009 [Capella-Gutierrez et al. 2009](#)). The phylogenetic tree was constructed using the maximum likelihood method in IQ-tree version 6.8 (Nguyen et al. 2015 [Nguyen et al. 2015](#)). The bootstrap algorithm was based on the Jones–Taylor–Thornton (JTT) + F + G4 model and run 5000 times. Visualization and later modification of phylogenetic trees were performed in FigTree version 1.4.4 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Quantitative real-time PCR

The relative expression levels of sugar GRs in *H. armigera* were analyzed by quantitative real-time PCR (qRT-PCR) (Livak and Schmittgen 2001 [Livak and Schmittgen 2001](#)). The specific primers (**Appendix—Table 2**) were designed by Primer3plus (<http://www.primer3plus.com/cgi-bin/dev/primer3plus.cgi>). qRT-PCR was conducted with a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA, USA). All reactions included 10 µL SYBR Premix Ex Taq (TaKaRa, Shiga, Japan); 0.4 µL (10 mM) primer (forward and reverse), 2 µL template DNA, 0.4 µL ROX Reference Dye II Dye, 6.8 µL ddH₂O

(20 μ L total). The procedure was as follows: 95°C for 30 s; 40 cycles of 95°C for 5 s and 60°C for 34 s; 95°C for 15 s; 60°C for 1 min; and 95°C for 15 s. 18S ribosomal RNA (GenBank number: KT343378.1) was used as the standard gene because the Ct value of this gene was not different between different tissues (Sharath Chandra et al. 2014 [\[1\]](#)). The relative expression levels of the specific genes were quantified by using the $2^{-\Delta Ct}$ method, where ΔCt is the Ct value of GR genes subtracted from that of 18S. Three biological replicates were run for all experiments, and there were 30–100 insects in each replicate.

Ectopic expression and functional analysis of sugar GRs

The sugar GRs sequences which were used for ectopic expression and functional identification need to contain restriction enzyme sites and Kozak sequences (the primer sequences of *H. armigera* sugar GRs for functional identification are provided in **Appendix—Table 2**). The full-length sequence of the sugar receptor was cloned from cDNA and transferred into pGEM-T vector (Promega, Madison, WI, USA). After cleavage by appropriate endonucleases, it was linked to pCS2⁺ vector. pCS2⁺ vector was linearized and cRNA was synthesized according to the protocol of the mMESSAGE mMACHINE SP6 Kit (Ambion, Austin, TX, USA). The products were dissolved in ddH₂O, diluted to 3000 ng/ μ L, and then stored at –80°C until use.

Each of Gr4, Gr5, Gr6, Gr7, Gr8, Gr10, Gr11, Gr12, Gr5 + Gr6, and Gr10 + Gr6 was expressed in oocytes. The acquisition of *X. laevis* oocytes was performed following a previously described protocol (Nakagawa and Touhara 2013 [\[2\]](#)). *X. laevis* were anesthetized by a mixture of ice and water for 30 min; then, the abdomen was cut to obtain oocytes. The *Xenopus* oocytes were treated with 2 mg/mL collagenase type I (Sigma-Aldrich, St Louis, MO, USA) in a calcium-free saline buffer (82.5 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 5 mM HEPES, pH = 7.5; the reagents are provided in **Key resource table**) at room temperature for 25 min. The oocytes were microinjected with 27.6 nL (80 ng) cRNA of each GR by SMART TouchTM (World Precision Instruments, Sarasota, FL, USA). For Gr10 + Gr6 or Gr5 + Gr6, a cRNA mixture with a ratio of 1:1 was used. The oocytes microinjected with RNase-free water were used as the control. Then, the oocytes were incubated for 3–4 days at 16°C in the bath solution (96 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 1.8 mM CaCl₂, 5 mM HEPES, pH 7.5; reagents are provided in **Key resource table**).

The whole-cell current was recorded with a two-electrode voltage clamp, the procedures as previously described (Jiang et al. 2015 [\[3\]](#)). The glass electrode, filled with 3M KCl, was inserted into the oocytes and maintained at 0.2–2.0 M Ω . The signal was amplified with an OC-725C amplifier (Warner Instruments, Hamden, CT, USA) at a holding potential at –80 mV all the time, low-pass filtered at 50 Hz, and digitized at 1 kHz. The sugar compounds (**Key resource table**) were dissolved in 1 $\times$ Ringer solution (96 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 1.8 mM CaCl₂, 5 mM HEPES, pH 7.5) as the test solution. The oocytes were stimulated by test solution via the perfusion system for 20 s each time, and then washed by 1 $\times$ Ringer solution (Nakagawa and Touhara 2013 [\[2\]](#)). The concentration of 100 mM for each chemical was randomly used at first, and then concentration gradients (ranging from 0.01 mM, 0.1 mM, 1 mM, 10 mM, 50 mM, 100 mM, 250 mM, and 500 mM) were recorded later when a clear current response was detected. Data were obtained and analyzed using Digidata 1322A and pCLAMP software version 10.4.2.0 (Axon Instruments Inc., Foster City, CA, USA).

In vitro synthesis of single-guide RNA

Gr10 and Gr6 single-guide RNA (sgRNA) position screening was conducted at <http://www.rgenome.net/cas-designer/> [\[4\]](#). The sgRNA target site of Gr10 was designed on exon 4 (**Figure 5B** [\[5\]](#)), and the sgRNA target site of Gr6 was designed on exon 2 (**Figure 5E** [\[6\]](#)). The Gr10 and Gr6 sgRNA primers (**Appendix—Table 5**) were designed based on the principles provided by the GeneArtTM Precision gRNA Synthesis Kit manual (Invitrogen, Vilnius, Lithuania). The sgRNAs were synthesized by the GeneArtTM gRNA Prep Kit (Invitrogen, Vilnius, Lithuania), and purified by the GeneArtTM gRNA

Clean Up Kit (Invitrogen, Vilnius, Lithuania). The products were dissolved in RNase-free water and then stored at -80°C until use. TrueCut™ Cas9 protein 2 (Invitrogen, Vilnius, Lithuania) was diluted to 500 ng/mL and stored at -20°C .

Embryo microinjection

The fresh laid eggs of *H. armigera* were placed on microscope slides and fixed on double-sided adhesive tape. These operations were finished within 2 h after oviposition. Microinjection into these pretreated eggs was performed with a mixture of *Gr10* (or *Gr6*) sgRNA and Cas9 protein, with final concentrations at 250 ng/ μL and 100 ng/ μL , respectively. Microinjection was finished using a PLI-100A PICO-INJECTOR (Warner Instruments, Hamden, Connecticut, USA). The injected eggs were placed in a 10-cm Petri dish, covered with wet filter paper, and incubated at 25°C until hatching.

DNA extraction and mutagenesis detection

To screen *Gr10* (or *Gr6*) homozygous mutants, the tarsi on hindlegs were used for DNA extraction with the Animal Tissue PCR Kit (TransGen Biotech, Beijing, China). Then, we used PCR to amplify the sequences that covered *Gr10* or *Gr6* sgRNA targeting sites. The reactions were 50 μL total and included: 2 μL template, 1 μL forward primer, 1 μL reverse primer (mutation detection primer sequences are provided in **Appendix—Table 2**), 25 μL Premix *Taq*™ (Takara, Dalian, China), and 19 μL ddH₂O. The procedure was as follows: 94°C for 3 min, 40 cycles [94°C for 30 s, 55°C (*Gr6*), or 60°C (*Gr10*) for 30 s], 72°C for 1 min, and finally 72°C for 5 min. The PCR products were sent to Beijing Genomics Institute (BGI) or the SinoGenoMax for Sanger sequencing, and the sequence results were analyzed by SnapGene® software version 4.3.8 (from Insightful Science; available at snapgene.com) and SeqMan software version 7.1 (DNASTAR, Madison, WI, USA).

Screening homozygous mutants

Gr10 and *Gr6* homozygous mutant lines (*Gr10*^{-/-} and *Gr6*^{-/-}, respectively) were established. A mixture of *Gr10* sgRNA and Cas9 protein was co-injected into each of 732 eggs that were laid within two hours. After hatching, the larvae were raised on artificial diet, and finally 125 G0 adults were obtained. Among 125 G0 adults, 15 G0 harbored multiple targeted mutations. Each mutated G0 was backcrossed with the wild type to obtain G1 offspring. From 57 G1 adults, one DNA strands containing 4 bp-deleted modifications, were in-crossed to establish stable lines; for the expanding mutant population, 4 bp-deleted homozygotes of G2 moths were in-crossed to generate homozygotes G3 moths (**Figure 5A**). Similarly, a mixture of *Gr6* sgRNA and Cas9 protein was co-injected into each of 695 eggs. After hatching, the larvae were raised on artificial diet, and finally 132 G0 adults were obtained. Among 132 G0 adults, 12 G0 harbored multiple targeted mutations. From 66 G1 adults, one DNA strands containing 17 bp-deleted modifications, were in-crossed to establish stable lines; for the expanding mutant population, 17 bp-deleted homozygotes of G2 moths were in-crossed to generate homozygotes G3 moths (**Figure 5D**). The related mutation rates are provided in **Appendix—Table 2**.

To confirm the deletion in *Gr10*^{-/-} and *Gr6*^{-/-} at mRNA level, the total RNA of the female adult proboscis of WT, *Gr10*^{-/-} and *Gr6*^{-/-} were extracted, and the cDNA was synthesized following the previous gene cloning protocol. The specific primers (**Appendix—Table 2**) were designed based on the DNA exon sequences of *Gr10* and *Gr6*, which covered the DNA-confirmed sgRNA targeting sites. During mRNA sequence cloning, the PCR reaction conditions are largely the same as those used for DNA sequence confirmation, with little modification (the annealing temperature reduced to 50°C). The nucleic acid sequences of the identified *Gr10* and *Gr6* were aligned manually by tools at the T-Coffee web server (Notredame et al. 2000).

Off-target effects detection

The potential off-target sites were predicted as previously described (Chen et al. 2020 [\[1\]](#)). The online software Cas-OFFinder (<http://www.rgenome.net/cas-offinder/> [\[2\]](#)) was used to blast the *Gr10* and *Gr6* sgRNAs against the *H. armigera* genome (Pearce et al. 2017 [\[3\]](#)), with mismatch number ≤ 3 . According to the blast results, *Gr10* has no potential off-target site, and one sequence (GenBank accession number: LOC126054697) in the presence of *Gr6* sgRNA showed a potential off-target region. The specific primers (Appendix—Table 2) were designed based on the DNA sequence of LOC126054697 covering the potential off-target sites. The DNA extraction and mutagenesis detection methods were described previously. Five individual replicates were performed.

To further illustrate that the mutation strains did not exhibit off-target effects, the electrophysiological responses of larvae and the biological parameters were compared among WT, *Gr10*^{-/-}, and *Gr6*^{-/-}. The larval medial sensilla styloconica to xylose, the body weight of the fifth instar larvae on the first day, the adult lifespan, and the number of eggs laid were recorded. Five to sixteen individuals were recorded.

Effects of *Gr10* or *Gr6* knockout on sugar reception in larvae and adults

We recorded the electrophysiological responses of contact chemosensilla in the wild type (WT), *Gr10*^{-/-}, and *Gr6*^{-/-} by the tip-recording technique as described earlier. Concentration of 1 mM, 10 mM, and 100 mM for sucrose and fucose were used as stimuli for lateral sensilla styloconica in larvae. As the response thresholds of sugar GSNs in adults were higher than those in larvae, concentrations of 10 mM and 100 mM for sucrose and fucose and 100 mM for fructose were used for adult contact chemosensilla. 2 mM KCl was used as the control for both larval and adult contact chemosensilla. 10 mM sinigrin and 10 mM nicotine were used as a positive control (induced the response of a bitter cell in the same sensilla) for larval and adult contact chemosensilla, respectively. All stimuli were tested randomly, and 12–20 replicates were set for each experiment.

Effects of *Gr10* or *Gr6* knockout on behavioral responses of larvae to sugars and plant tissues

The feeding preference of WT, *Gr10*^{-/-}, and *Gr6*^{-/-} larvae to 10 mM and 100 mM sucrose was investigated in two-choice tests as described earlier. 20 replicates were set for each experiment. We also performed the no-choice tests to measure the feeding amount of WT, *Gr10*^{-/-}, and *Gr6*^{-/-} larvae in 1 h on leaves of cabbage (*Brassica oleracea* ssp. capitata, Shenglv-7), corn kernels of maize (*Zea mays*, Jingke-968), seeds of pea (*Pisum sativum*, Changshouren), fruits of pepper (*Capsicum frutescens*, Changjian), and fruits of tomato (*Solanum lycopersicum*, Ailsa Craig-LA2838A). Fifth instar larvae on the first day after molting were starved for 2.5 h before testing. Each plant tissue was weighed and put into a Petri dish (10 cm diameter × 2 cm depth; Corning, Wujiang, China), and then one larva was introduced. The larva was removed after 1 h, and the rest of the plant tissue was weighed. Petri dishes with plant tissues but no caterpillar was used to measure water evaporation of the plant tissues. The feeding amount of larvae is calculated based on the weight changes of the plant tissue before and after being fed. 20 replicates were set for each experiment.

Effects of *Gr10* or *Gr6* knockout on behavioral responses of adults to sugars

The behavioral responses of the female adults of WT, *Gr10*^{-/-}, and *Gr6*^{-/-} to sugars were determined by using the PER test as described earlier. sucrose, fucose, and fructose at the concentrations of 10 mM, 100 mM, and 1000 mM were tested randomly to antennae and tarsi. H₂O

was used as the control. All the experiments were repeated three times, and 20 individual moths were used for each set of tests.

Statistical analysis

Statistical analyses were performed by SPSS 20 (IBM, Chicago, IL, USA) and GraphPad Prism 8.2.1 (Dotmatics, San Diego, CA, USA). For the tip-recording data and the PER data, the comparison between test compounds and the control was performed using two-tailed independent-samples *t* test. For the two-choice feeding test, paired-samples *t* test was used. The data of dose-response curves in tip-recording, PER experiments and two-electrode voltage-clamp recording, and gene relative expression levels and feeding amount in the no-choice test were analyzed by one-way ANOVA with Tukey's HSD tests. The comparison among mutation strains and WT in electrophysiological and PER experiments were analyzed by two-way ANOVA with post hoc Tukey's multiple comparison. All analytical methods were tested at $P < 0.05$ for significant differences.

Acknowledgements

We thank our colleagues Zi-Lin Li, Qing Zhang, Yan Chen, Jun Yang, Xin-Lin Gong, Yan-Yan Jia for assistances in PER experiments, mutant line screening and scanning electron microscopy. This work was supported by National Key R&D Program of China (Grant No. 2022YFD1400800), the National Natural Science Foundation of China (Grant No. 32130090), and the Beijing Natural Science Foundation (Grant No. 6224062).

Author Contributions

Conceptualization, C.Z.W. and S.S.Z.; Methodology, S.S.Z., P.C.W., L.Q.H., C.N., G.C.L., and C.Z.W.; Software, S.S.Z., P.C.W., C.N., G.C.L., and C.Z.W.; Validation, S.S.Z. and C.Z.W.; Formal Analysis, S.S.Z., C.N., G.C.L., and C.Z.W.; Investigation, S.S.Z., P.C.W., K.Y., L.L.C., L.Q.H., and C.Z.W.; Resources, S.S.Z., P.C.W., L.Q.H., and C.Z.W.; Data curation, S.S.Z., C.N., P.C.W., G.C.L., and C.Z.W.; Writing – original draft, C.Z.W. and S.S.Z.; Writing – review & editing, C.Z.W., S.S.Z.; Visualization, S.S.Z. and C.Z.W.; Supervision, C.Z.W.; Project administration, C.Z.W. and L.Q.H.; Funding acquisition, C.Z.W. and K.Y.

Competing interests

The authors declare no competing interests.

Data availability

All data and materials generated in this study are available in the manuscript, supplementary materials, Appendix, and source data. Sequence data that support the finding of this study have been deposited in GenBank with the accession numbers (see **Appendix—Table 2**).

References

1. Abràmoff MD, Magalhães PJ, Ram SJ (2004) **Image processing with ImageJ** *Biophotonics international* **11**:36–42
2. Ai D, Dong C, Yang B, Yu C, Wang G (2022) **A fructose receptor gene influences development and feed intake in *Helicoverpa armigera*** *Insect Science* **29**:993–1005 <https://doi.org/10.1111/1744-7917.12984>
3. Bernays EA, Chapman RF (1994) **Host-Plant Selection by Phytophagous Insects**
4. Blaney WM, Simmonds MSJ (1988) **Food selection in adults and larvae of three species of *Lepidoptera*: a behavioral and electrophysiological study** *Entomologia Experimentalis Et Applicata* **49**:111–121 <https://doi.org/10.1111/j.1570-7458.1988.tb02482.x>
5. Blaney WM, Simmonds MSJ (1990) **A behavioural and electrophysiological study of the role of tarsal chemoreceptors in feeding by adults of *Spodoptera*, *Heliothis virescens* and *Helicoverpa armigera*** *Journal of Insect Physiology* **36**:743–756 [https://doi.org/10.1016/0022-1910\(90\)90048-K](https://doi.org/10.1016/0022-1910(90)90048-K)
6. Briscoe AD, et al. (2013) **Female behaviour drives expression and evolution of gustatory receptors in butterflies** *PLoS Genetics* **9** <https://doi.org/10.1371/journal.pgen.1003620>
7. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T (2009) **trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses** *Bioinformatics* **25**:1972–1973 <https://doi.org/10.1093/bioinformatics/btp348>
8. Chapman RF (2003) **Contact chemoreception in feeding by phytophagous insects** *Annual Review of Entomology* **48**:455–484 <https://doi.org/10.1146/annurev.ento.48.091801.112629>
9. Chen W, Dong YH, Saqib HSA, Vasseur L, Zhou WW, Zheng L, Lai YF, Ma XL, Lin LY, Xu XJ (2020) **Functions of duplicated glucosinolate sulfatases in the development and host adaptation of *Plutella xylostella*** *Insect Biochemistry and Molecular Biology* **119** <https://doi.org/10.1016/j.ibmb.2020.103316>
10. Chen Y, Wang PC, Zhang SS, Yang J, Li GC, Huang LQ, Wang CZ (2022) **Functional analysis of a bitter gustatory receptor highly expressed in the larval maxillary galea of *Helicoverpa armigera*** *PLoS Genetics* **18** <https://doi.org/10.1371/journal.pgen>
11. Dahanukar A, Foster K, van der Goes van Naters WM, Carlson JR. (2001) **A Gr receptor is required for response to the sugar trehalose in taste neurons of *Drosophila*** *Nat Neurosci* **4**:1182–1186 <https://doi.org/10.1038/nn765>
12. Dahanukar A, Lei YT, Kwon JY, Carlson JR (2007) **Two Gr genes underlie sugar reception in *Drosophila*** *Neuron* **56**:503–516 <https://doi.org/10.1016/j.neuron.2007.10.024>
13. Damak S, Rong M, Yasumatsu K, Kokrashvili Z, Varadarajan V, Zou S, Jiang P, Ninomiya Y, Margolskee RF (2003) **Detection of sweet and umami taste in the absence of taste receptor T1r3** *Science* **301**:850–853 <https://doi.org/10.1126/science.1087155>

14. Değirmenci L, Geiger D, Rogé Ferreira FL, Keller A, Krischke B, Beye M, Steffan-Dewenter I, Scheiner R (2020) **CRISPR/Cas 9-mediated mutations as a new tool for studying taste in Honeybees** *Chemical Senses* **45**:655–666 <https://doi.org/10.1093/chemse/bjaa063>
15. Delventhal R, Carlson JR (2016) **Bitter taste receptors confer diverse functions to neurons** *Elife* **5** <https://doi.org/10.7554/eLife.11181>
16. Dethier VG (1973) **Electrophysiological studies of gustation in Lepidopterous larvae** *Journal of comparative physiology* **82**:103–134 <https://doi.org/10.1007/BF00696148>
17. Engsontia P, Sangket U, Chotigeat W, Satasook C (2014) **Molecular evolution of the odorant and gustatory receptor genes in Lepidopteran insects: implications for their adaptation and speciation** *Journal of Molecular Evolution* **79**:21–39 <https://doi.org/10.1007/s00239-014-9633-0>
18. Fan XB, Mo BT, Li GC, Huang LQ, Guo H, Gong XL, Wang CZ (2022) **Mutagenesis of the odorant receptor co-receptor (Orco) reveals severe olfactory defects in the crop pest moth *Helicoverpa armigera*** *BMC Biology* **20** <https://doi.org/10.1186/s12915-022-01411-2>
19. Faucheux MJ (2013) **Sensillum types on the proboscis of the Lepidoptera: a review** *Annales De La Societe Entomologique De France* **49**:73–90 <https://doi.org/10.1080/00379271.2013.767514>
20. Free JB (1993) **Insect Pollination of Crops**
21. Freeman EG, Wisotsky Z, Dahanukar A (2014) **Detection of sweet tastants by a conserved group of insect gustatory receptors** *PNAS* **111**:1598–1603 <https://doi.org/10.1073/pnas.1311724111>
22. Fujii S, Yavuz A, Slone J, Jagge C, Song X, Amrein H (2015) **Drosophila sugar receptors in sweet taste perception, olfaction, and internal nutrient sensing** *Current Biology* **25**:621–627 <https://doi.org/10.1016/j.cub.2014.12.058>
23. Glendinning JI, Foley C, Loncar I, Rai M (2009) **Induced preference for host plant chemicals in the tobacco hornworm: contribution of olfaction and taste** *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **195**:591–601 <https://doi.org/10.1007/s00359-009-0434-7>
24. Guo H, et al. (2017) **Expression map of a complete set of gustatory receptor genes in chemosensory organs of *Bombyx mori*** *Insect Biochemistry and Molecular Biology* **82**:74–82 <https://doi.org/10.1016/j.ibmb.2017.02.001>
25. Harari OA, Dekel A, Wintraube D, Vainer Y, Mozes-Koch R, Yakir E, Malka O, Morin S, Bohbot JD (2023) **A sucrose-specific receptor in *Bemisia tabaci* and its putative role in phloem feeding** *iScience* **26** <https://doi.org/10.1016/j.isci.2023.106752>
26. Hill CA, Fox AN, Pitts RJ, Kent LB, Tan PL, Chrystal MA, Cravchik A, Collins FH, Robertson HM, Zwiebel LJ (2002) **G protein-coupled receptors in *Anopheles gambiae*** *Science* **298**:176–178 <https://doi.org/10.1126/science.1076196>
27. Hodgson ES, Lettvin JY, Roeder KD (1955) **Physiology of a primary chemoreceptor unit** *Science* **122**:417–418 <https://doi.org/10.1126/science.122.3166.417-a>

28. Jiang XJ, Ning C, Guo H, Jia YY, Huang LQ, Qu MJ, Wang CZ (2015) **A gustatory receptor tuned to D-fructose in antennal sensilla chaetica of *Helicoverpa armigera*** *Insect Biochemistry and Molecular Biology* **60**:39–46 <https://doi.org/10.1016/j.ibmb.2015.03.002>
29. Jiao Y, Moon SJ, Montell C (2007) **A *Drosophila* gustatory receptor required for the responses to sucrose, glucose, and maltose identified by mRNA tagging** *PNAS* **104**:14110–14115 <https://doi.org/10.1073/pnas.0702421104>
30. Jiao YC, Moon SJ, Wang XY, Ren QT, Montell C (2008) **Gr64f is required in combination with other gustatory receptors for sugar detection in *Drosophila*** *Current Biology* **18**:1797–1801 <https://doi.org/10.1016/j.cub.2008.10.009>
31. Jørgensen K, Almaas TJ, Marion-Poll F, Mustaparta H (2007) **Electrophysiological characterization of responses from gustatory receptor neurons of sensilla chaetica in the moth *Heliothis virescens*** *Chemical Senses* **32**:863–879 <https://doi.org/10.1093/chemse/bjm057>
32. Jung JW, Park KW, Ahn YJ, Kwon HW (2015) **Functional characterization of sugar receptors in the western honeybee, *Apis mellifera*** *Journal of Asia-Pacific Entomology* **18**:19–26 <https://doi.org/10.1016/j.aspen.2014.10.011>
33. Kent LB, Walden KK, Robertson HM (2008) **The Gr family of candidate gustatory and olfactory receptors in the yellow-fever mosquito *Aedes aegypti*** *Chemical Senses* **33**:79–93 <https://doi.org/10.1093/chemse/bjm067>
34. Kwon JY, Dahanukar A, Weiss LA, Carlson JR (2011) **Molecular and cellular organization of the taste system in the *Drosophila* larva** *Journal of Neuroscience* **31**:15300–15309 <https://doi.org/10.1523/jneurosci.3363-11.2011>
35. Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E (2002) **Human receptors for sweet and umami taste** *PNAS* **99**:4692–4696 <https://doi.org/10.1073/pnas.072090199>
36. Liman ER, Zhang YV, Montell C (2014) **Peripheral coding of taste** *Neuron* **81**:984–1000 <https://doi.org/10.1016/j.neuron.2014.02.022>
37. Liu XL, Sun SJ, Hou W, Zhang J, Yan Q, Dong SL (2020) **Functional characterization of two spliced variants of fructose gustatory receptor in the diamondback moth, *Plutella xylostella*** *Pesticide biochemistry and physiology* **164**:7–13 <https://doi.org/10.1016/j.pestbp.2019.12.002>
38. Liu XL, Yan Q, Yang YL, Hou W, Miao CL, Peng YC, Dong SL (2019) **A gustatory receptor GR8 tunes specifically to D-fructose in the common cutworm *Spodoptera litura*** *Insects* **10** <https://doi.org/10.3390/insects10090272>
39. Livak KJ, Schmittgen TD (2001) **Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT Method** *Methods* **25**:402–408 <https://doi.org/10.1006/meth.2001.1262>
40. Maier GL, Komarov N, Meyenhofer F, Kwon JY, Sprecher SG (2021) **Taste sensing and sugar detection mechanisms in *Drosophila* larval primary taste center** *Elife* <https://doi.org/10.7554/eLife.67844>

41. Mang D, Mayu K, Toyama T, Yamagishi T, Sato R (2022) **BmGr4 responds to sucrose and glucose and expresses in tachykinin-related peptide-secreting enteroendocrine cells** *Insect Biochemistry and Molecular Biology* **150** <https://doi.org/10.1016/j.ibmb.2022.103858>
42. Martin TL, Shields VDC (2012) **Detection of alkaloids and carbohydrates by taste receptor cells of the galea of gypsy moth larvae, *Lymantria dispar* (L.)** *Arthropod-Plant Interactions* **6**:519–529 <https://doi.org/10.1007/s11829-012-9209-0>
43. Mishra D, Miyamoto T, Rezenom YH, Broussard A, Yavuz A, Slone J, Russell DH, Amrein H (2013) **The molecular basis of sugar sensing in *Drosophila* larvae** *Current Biology* **23**:1466–1471 <https://doi.org/10.1016/j.cub.2013.06.028>
44. Miyamoto T, Slone J, Song XY, Amrein H. (2012) **A fructose receptor functions as a nutrient sensor in the *Drosophila* brain** *Cell* **151**:1113–1125 <https://doi.org/10.1016/j.cell.2012.10.024>
45. Morinaga S, Nagata K, Ihara S, Yumita T, Niimura Y, Sato K, Touhara K (2022) **Structural model for ligand binding and channel opening of an insect gustatory receptor** *Journal of Biological Chemistry* **298** <https://doi.org/10.1016/j.jbc.2022.102573>
46. Nakagawa T, Touhara K, Touhara K (2013) **Functional assays for insect olfactory receptors in *Xenopus* oocytes. Pages 107–119** *Pheromone Signaling Totowa*
47. Nelson G, Hoon MA, Chandrashekar J, Zhang Y, Ryba NJ, Zuker CS (2001) **Mammalian sweet taste receptors** *Cell* **106**:381–390 [https://doi.org/10.1016/s0092-8674\(01\)00451-2](https://doi.org/10.1016/s0092-8674(01)00451-2)
48. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. (2015) **IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies** *Molecular Biology* **32**:268–274 <https://doi.org/10.1093/molbev/msu300>
49. Nicolson SW (1998) **The importance of osmosis in nectar secretion and its consumption by insects** *American Zoologist* **38**:418–425 <https://doi.org/10.1093/icb/38.3.418>
50. Notredame C, Higgins DG, Heringa J (2000) **T-Coffee: A novel method for fast and accurate multiple sequence alignment** *Journal Of Molecular Biology* **302**:205–217 <https://doi.org/10.1006/jmbi.2000.4042>
51. Pearce SL, et al. (2017) **Genomic innovations, transcriptional plasticity and gene loss underlying the evolution and divergence of two highly polyphagous and invasive *Helicoverpa* pest species** *BMC biology* **15** <https://doi.org/10.1186/s12915-017-0402-6>
52. Popescu A, Couton L, Almaas TJ, Rospars JP, Wright GA, Marion-Poll F, Anton S (2013) **Function and central projections of gustatory receptor neurons on the antenna of the noctuid moth *Spodoptera littoralis*** *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **199**:403–416 <https://doi.org/10.1007/s00359-013-0803-0>
53. Robertson HM, Wanner KW (2006) **The chemoreceptor superfamily in the honey bee, *Apis mellifera*: expansion of the odorant, but not gustatory, receptor family** *Genome Research* **16**:1395–1403 <https://doi.org/10.1101/gr.5057506>
54. Robertson HM, Warr CG, Carlson JR (2003) **Molecular evolution of the insect chemoreceptor gene superfamily in *Drosophila melanogaster*** *PNAS* **100**:14537–14542 <https://doi.org/10.1073/pnas.2335847100>

55. Rozewicki J, Li S, Amada KM, Standley DM, Katoh K (2019) **MAFFT-DASH: integrated protein sequence and structural alignment** *Nucleic Acids Research* **47**:W5–W10 <https://doi.org/10.1093/nar/gkz342>
56. Sato K, Tanaka K, Touhara K (2011) **Sugar-regulated cation channel formed by an insect gustatory receptor** *PNAS* **108**:11680–11685 <https://doi.org/10.1073/pnas.1019622108>
57. Schauer N, Zamir D, Fernie AR (2005) **Metabolic profiling of leaves and fruit of wild species tomato: a survey of the *Solanum lycopersicum* complex** *Journal Of Experimental Botany* **56**:297–307 <https://doi.org/10.1093/jxb/eri057>
58. Schnuch M, Seebauer H (1998) **Sugar cell responses to lactose and sucrose in labellar and tarsal taste hairs of *Musca domestica*** *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **182**:767–775 <https://doi.org/10.1007/s003590050221>
59. Schoonhoven LM, van Loon JJA. (2002) **An inventory of taste in caterpillars: each species its own key** *Acta Zoologica Academiae Scientiarum Hungaricae* **48**:215–263
60. Schoonhoven LM, van Loon JJA, Dicke M (2005) **Insect-Plant Biology**
61. Sharath Chandra G, Asokan R, Manamohan M, Krishna Kumar NK, Sita T (2014) **Evaluation of reference genes for quantitative real-time PCR normalization in cotton bollworm *Helicoverpa armigera*** *Molecular Biology (Mosk)* **48**:813–822
62. Slone J, Daniels J, Amrein H (2007) **Sugar receptors in *Drosophila*** *Current Biology* **17**:1809–1816 <https://doi.org/10.1016/j.cub.2007.09.027>
63. Smith JJB, Mitchell BK, Rolseth BM, Whitehead AT, Albert PJ (1990) **SAPID tools - microcomputer programs for analysis of multiunit nerve recordings** *Chemical Senses* **15**:253–270 <https://doi.org/10.1093/chemse/15.3.253>
64. Takada T, Sasaki T, Sato R, Kikuta S, Inoue MN (2018) **Differential expression of a fructose receptor gene in honey bee workers according to age and behavioral role** *Archives of Insect Biochemistry and Physiology* **97**:1–9 <https://doi.org/10.1002/arch.21437>
65. Tang DL, Wang CZ, Luo L, Qin JD (2000) **Comparative study on the responses of maxillary sensilla styloconica of cotton bollworm *Helicoverpa armigera* and Oriental tobacco budworm *H. assulta* larvae to phytochemicals** *Science China Life Sciences* **43**:606–612 <https://doi.org/10.1007/Bf02882281>
66. van Loon JJA. (1990) **Chemoreception of phenolic acids and flavonoids in larvae of two species of *Pieris*** *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **166**:889–899 <https://doi.org/10.1007/bf00187336>
67. Vansell GH (1944) **Cotton nectar in relation to bee activity and honey production** *Journal of Economic Entomology* **37**:528–530 <https://doi.org/10.1093/jee/37.4.528>
68. Waller GD, Wilson FD, Martin JH (1981) **Influence of phenotype, season, and time-of-day on nectar production in cotton** *Crop Science* **21**:507–511 <https://doi.org/10.2135/cropsci1981.0011183X002100040007x>
69. Xu W, Liu N, Liao Y, Anderson A (2017) **Molecular characterization of sugar taste receptors in the cotton bollworm *Helicoverpa armigera*** *Genome* **60**:1037–1044 <https://doi.org/10.1139/gen-2017-0086>

70. Xu W, Zhang HJ, Anderson A (2012) **A sugar gustatory receptor identified from the foregut of cotton bollworm *Helicoverpa armigera*** *Journal of Chemical Ecology* **38**:1513–1520 <https://doi.org/10.1007/s10886-012-0221-8>
71. Yarmolinsky DA, Zuker CS, Ryba NJP (2009) **Common sense about taste: from mammals to insects** *Cell* **139**:234–244 <https://doi.org/10.1016/j.cell.2009.10.001>
72. Zhan S, Merlin C, Boore JL, Reppert SM (2011) **The monarch butterfly genome yields insights into long-distance migration** *Cell* **147**:1171–1185 <https://doi.org/10.1016/j.cell.2011.09.052>
73. Zhang HJ, Faucher CP, Anderson A, Berna AZ, Trowell S, Chen QM, Xia QY, Chyb S (2013) **Comparisons of contact chemoreception and food acceptance by larvae of polyphagous *Helicoverpa armigera* and oligophagous *Bombyx mori*** *Journal of Chemical Ecology* **39**:1070–1080 <https://doi.org/10.1007/s10886-013-0303-2>
74. Zhang YF, van Loon JJA, Wang CZ (2010) **Tarsal taste neuron activity and proboscis extension reflex in response to sugars and amino acids in *Helicoverpa armigera* (Hübner)** *Journal of Experimental Biology* **213**:2889–2895 <https://doi.org/10.1242/jeb.042705>
75. Zhao GQ, Zhang Y, Hoon MA, Chandrashekar J, Erlenbach I, Ryba NJ, Zuker CS (2003) **The receptors for mammalian sweet and umami taste** *Cell* **115**:255–266 [https://doi.org/10.1016/s0092-8674\(03\)00844-4](https://doi.org/10.1016/s0092-8674(03)00844-4)

Article and author information

Shuai-Shuai Zhang

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Pei-Chao Wang

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Chao Ning

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Ke Yang

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Guo-Cheng Li

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Lin-Lin Cao

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Ling-Qiao Huang

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China

Chen-Zhu Wang

State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China, Chinese Academy of Sciences Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

For correspondence: czwang@ioz.ac.cn

Copyright

© 2023, Zhang et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Sonia Sen

Tata Institute for Genetics and Society, Bangalore, India

Senior Editor

K VijayRaghavan

National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, India

Reviewer #1 (Public Review):

Summary:

The process of taste perception is significantly more intricate and complex in Lepidopteran insects. This investigation provides valuable insights into the role of Gustatory receptors and their dynamics in the sensation of sucrose, which serves as a crucial feeding cue for insects. The article highlights the differential sensitivity of Grs to sucrose and their involvement in feeding and insect behavior.

Strengths:

To support the notion of the differential specificity of Gr to sucrose, this study employed electrophysiology, ectopic expression of Grs in *Xenopus*, genome editing, and behavioral studies on insects. This investigation offers a fundamental understanding of the gustation process in lepidopteran insects and its regulation of feeding and other gustation-related physiological responses. This study holds significant importance in advancing our comprehension of lepidopteran insect biology, gustation, and feeding behavior.

Weaknesses:

While this manuscript demonstrates technical proficiency, there exists an opportunity for additional refinement to optimize comprehensibility for the intended audience. Several crucial sugars have been overlooked in the context of electrophysiology studies and should be incorporated. Furthermore, it is imperative to consider the potential off-target effects of Gr knock-out on other Gr expressions. This investigation focuses exclusively on Gr6 and Gr10, while neglecting a comprehensive narrative regarding other Grs involved in sucrose sensation.

<https://doi.org/10.7554/eLife.91711.2.sa1>

Reviewer #2 (Public Review):**Summary:**

To identify sugar receptors and assess the capacity of these genes the authors first set out to identify behavioral responses in larva and adult as well as physiological response. They used phylogenetics and gene expression (RNAseq) to identify candidates for sugar reception. Using first an in vitro oocyte system they assess the responses to distinct sugars. A subsequent genetic analysis shows that the Gr10 and Gr6 genes provide stage specific functions in sugar perception.

Strengths:

A clear strength of the manuscript is the breadth of techniques employed allowing a comprehensive study in a non-canonical model species.

Weaknesses:

There are no major weaknesses in the study for the current state of knowledge in this species. Since it is much basic work to establish a broader knowledge, context with other modalities remain unknown. It might have been possible to probe certain context known from the fruit fly, which would have strengthened the manuscript.

<https://doi.org/10.7554/eLife.91711.2.sa0>

Author response:

The following is the authors' response to the original reviews.

We have revised the manuscript mainly in the following aspects: (1) the data of electrophysiological and behavioral responses of larvae and adults to trehalose have been added, and the related figures and texts have been modified accordingly; (2) the photos of taste organs of larvae and adults indicating the position of recorded sensilla have been added; (3) the potential off-target effects of GR knock-out on other GR expressions has been

carefully explained and revised in the relevant text; (4) the abstract has been revised to present the findings more technically in a limited number of words; (5) some details of experiments in Materials and Methods and some new literatures have been added; (6) a new figure (Figure 8) summarizing the main findings of the study has been added.

In the following, we respond to the reviewers' comments and suggestions one by one. We hope that our answers will satisfy you and the three reviewers. We are also very happy to get further valuable advices from you.

Public Reviews:

Reviewer #1 (Public Review):

Summary:

The process of taste perception is significantly more intricate and complex in Lepidopteran insects. This investigation provides valuable insights into the role of Gustatory receptors and their dynamics in the sensation of sucrose, which serves as a crucial feeding cue for insects. The article highlights the differential sensitivity of Grs to sucrose and their involvement in feeding and insect behavior.

Strengths:

*To support the notion of the differential specificity of Gr to sucrose, this study employed electrophysiology, ectopic expression of Grs in *Xenopus*, genome editing, and behavioral studies on insects. This investigation offers a fundamental understanding of the gustation process in lepidopteran insects and its regulation of feeding and other gustation-related physiological responses. This study holds significant importance in advancing our comprehension of lepidopteran insect biology, gustation, and feeding behavior.*

Thank you for your recognition of our research.

Weaknesses:

While this manuscript demonstrates technical proficiency, there exists an opportunity for additional refinement to optimize comprehensibility for the intended audience. Several crucial sugars have been overlooked in the context of electrophysiology studies and should be incorporated. Furthermore, it is imperative to consider the potential off-target effects of Gr knock-out on other Gr expressions. This investigation focuses exclusively on Gr6 and Gr10, while neglecting a comprehensive narrative regarding other Grs involved in sucrose sensation.

We accept the reviewer's suggestion. Because trehalose is a main sugar in insect blood, and it is converted by insects after feeding on plant sugars, we have added the new data on electrophysiological and behavioral responses of larvae and adults of *Helicoverpa armigera* to trehalose (see Figure 1-2, Figure 1-figure supplement 1, Figure 2-figure supplement 1). Now, the total eight sugars include 2 pentoses (arabinose and xylose), 4 hexoses (fructose, fucose, galactose and glucose), and 2 disaccharides (sucrose and trehalose), which were chosen because they are mainly present in host-plants of *H. armigera* and/or representative in the structure and source of sugars.

We fully agree to the reviewer's opinion and have already taken the potential off-target effects of CRISPR/Cas9 knockout of Gr on other GR expressions into consideration. To predict the potential off-target sites of sgRNA of Gr6 and Gr10 establishing homozygous mutants using CRISPR/Cas9 technology, we first use online software CasOFFinder (<http://www.rgenome.net/cas-offinder/>) to blast the genome of the wild type cotton bollworm and set the

mismatch number less than or equal to 3. We found that Gr10 sgRNA had no potential potential off-target site, and the sgRNA of Gr6 had only one potential off-target site. Therefore, we designed primers according to the sequence of potential off-target sites of Gr6 sgRNA, and conducted PCR using genomic DNA of homozygous mutant as a template, performed Sanger sequencing on the PCR products obtained, and found that the potential off-target sites of Gr6 sgRNA were no different from those of the wild type. Particularly, concerning the sgRNA of Gr6 and Gr10 may produce off-target effects on other sugar receptor genes of *H. armigera*, we conducted the same off-target site analysis with the designed sgRNA on each of the other eight sugar receptor genes, and found that there were no off-target sites on these receptor genes (see Line254-256).

Reviewer #2 (Public Review):

Summary:

To identify sugar receptors and assess the capacity of these genes the authors first set out to identify behavioral responses in larvae and adults as well as physiological response. They used phylogenetics and gene expression (RNAseq) to identify candidates for sugar reception. Using first an in vitro oocyte system they assess the responses to distinct sugars. A subsequent genetic analysis shows that the Gr10 and Gr6 genes provide stage specific functions in sugar perception.

Strengths:

A clear strength of the manuscript is the breadth of techniques employed allowing a comprehensive study in a non-canonical model species.

Thank you for your recognition of our research.

Weaknesses:

There are no major weaknesses in the study for the current state of knowledge in this species. Since it is much basic work to establish a broader knowledge, context with other modalities remains unknown. It might have been possible to probe certain contexts known from the fruit fly, which would have strengthened the manuscript.

Thank you so much for your suggestion. According to this suggestion, we further added some sentences probing sugar sensing and behaviors of fruit fly larvae in the Introduction and discussion sections (Line 68-71 in Introduction section, Line 395-399 in Discussion section).

Reviewer #3 (Public Review):

In this study, the authors combine electrophysiology, behavioural analyses, and genetic editing techniques on the cotton bollworm to identify the molecular basis of sugar sensing in this species.

*The larval and adult forms of this species feed on different plant parts. Larvae primarily consume leaves, which have relatively lower sugar concentrations, while adults feed on nectar, rich in sugar. Through a series of experiments-spanning electrophysiological recordings from both larval and adult sensillae, qPCR expression analysis of identified GRs from these sensillae, response profiles of these GRs to various sugars via heterologous expression in *Xenopus* oocytes, and evaluations of CRISPR mutants based on these parameters-the authors discovered that larvae and adults employ distinct GRs for sugar sensing. While the larva uses the highly sensitive GR10, the adult uses the less sensitive and broadly tuned GR6. This differential use of GRs are in keeping with their behavioral ecology.*

The data are cohesive and consistently align across the methodologies employed. They are also well presented and the manuscript is clearly written.

Recommendations for the authors:

While appreciating the quality of the work and its presentation, we have a few comments for the authors, should they wish to consider them, that would significantly improve the presentation of the work.

Title: Could the authors please revisit their title to better reflect the main finding of their work?

The title has been changed into “The larva and adult of *Helicoverpa armigera* use differential gustatory receptors to sense sugars”.

Text: There are a few comments related to the text, and these are listed below:

*(1) Could the authors place their work in the context of what's known about sugar sensing in *Drosophila* larva and adult?*

In the Introduction section, we added the status of research on sugar perception in *Drosophila* larvae, pointing out “No external sugar-sensing mechanism in *Drosophila* larvae has yet been characterized.” (Line 70-71); in the Discussion section, the research progress of sugar sensing in *Drosophila* adults and larvae was also summarized (Line 397-399).

(2) For each results section, could the authors please include a sentence or two that interprets the data in the context of previously presented data?

We accept the reviewer's suggestion. In order to make it easy for readers to follow up, we included a sentence interprets the above data at the beginning of each part of the Results on the premise of avoiding duplication.

(3) Could the authors please provide details of the generation and screening of the CRISPR mutants?

We have added more details on mutant establishment and screening in the Materials and Methods section (Line 722-726, 729-732).

Figures: Could the authors please include images and schematics wherever possible? For example, a schematic depicting the position of the sense organs and one summarising the main findings of the studies.

In Figure 1 we added the photo of each taste organ, on which the recorded sensilla were indicated. We also added a new figure, Figure 8, summarizing the main findings of the study.

Choice of Sugars: Could the authors please justify their choice of sugars they have used in the analyses?

In the first paragraph of the Results section of the article, we further explain the reasons for using the sugars in the study. “We first investigated the electrophysiological responses of the lateral and medial sensilla styloconica in the larval maxillary galea to eight sugars. These sugars were chosen because they are mostly found in host-plants of *H. armigera* or are representative in the structure and source of sugars.”

In addition to this, there are several specific comments in the detailed reviewers comments below, which the authors could consider responding to.

Reviewer #1 (Recommendations For The Authors):

The article titled "Sucrose taste receptors exhibit dissimilarities between larval and adult stages of a moth" by Shuai-Shuai Zhang and colleagues provides an intriguing analysis. The authors have conducted a meticulously planned and executed study. However, I do have some inquiries.

*(1) What precisely does the term "differ" signify in the title? It can be expounded upon in terms of differing in expression or sensitivity. The title could benefit from being more informative. The authors should appropriately specify the insect species in the title of the paper. This would make it more comprehensible to readers. Merely mentioning the term "moth" does not provide any information about the model organism. Hence, it would be preferable to mention *Helicoverpa armigera* instead of using the generic term "moth" in the title.*

Thank you for your suggestions. We considered it better to emphasize that the receptors for sucrose are different, and we have accepted the suggestion of adding the name of the animal. The title has been changed into "The larva and adult of *Helicoverpa armigera* use differential gustatory receptors to sense sugars".

(2) The abstract is written in a simple and easily understandable manner, but it overlooks important findings from a technical standpoint.

We add some key experimental techniques to illustrate some important findings in the Abstract.

(3). Almost all herbivorous insects are said to consume plants and utilize sucrose as a stimulus for feeding, as stated by the authors. Sucrose, glucose, and fructose sugar are among the commonly observed stimulants for feeding in numerous insects. It would be appropriate to incorporate not only sucrose but also glucose and fructose as feeding stimulants for almost all herbivorous insects.

Thank you for your suggestion. Sucrose is the major sugar in plants, and its concentration varies greatly from tissue to tissue, while the concentration of the hexose sugars is much lower and the concentration does not change much. In Line 48, we state that sucrose, glucose, and fructose are feeding stimuli for herbivorous insects. From the previous studies, it seems that sucrose is the strongest, followed by fructose, and finally glucose. The cotton bollworm larvae showed no electrophysiological and behavioral response to glucose.

(4) The reason why trehalose is not considered in the electrophysiology analysis is unclear. Given that trehalose is a major sugar in insects and plants, it would be intriguing to include it in the analysis.

We have accepted the reviewer's suggestion, and supplemented the electrophysiological responses of taste organs in larvae and adults of *Helicoverpa armigera* to trehalose (Figure 1, Figure 1-Figure Supplement 1), and also tested the behavioral responses of the larvae and adults to trehalose (Figure 2, Figure 2-Figure Supplement 1). Therefore, all the related figures have been changed.

(5) The author's intention regarding the co-receptor relationship between Gr5 and Gr6 (line 211) is unclear. If this is indeed the case, then the reason for considering Gr5 in

| *further studies remains uncertain.*

We have changed the sentence as follows: “Since Gr5 was highly expressed with Gr6 in the proboscis and tarsi (Figure 3D-3E, Figure 3—figure supplement 1), we suspected that Gr5 and Gr6 might be expressed in the same cells, and then tested the response profile of their co-expression in oocytes.”

| *(6) The homologous nature of Grs is emphasized by the authors. It is not specified how the author ensured that the guide RNA targeting Gr6 or Gr10 did not result in off-target effects on other Grs.*

Thank you so much for your suggestion. We have rewritten the relevant paragraph (Line 238-251), detailing our tests and the results on the potential off-target effects of knocking out GRs by CRISPR/Cas9: “In order to predict the potential off-target sites of sgRNA of Gr6 and Gr10, we used online software Cas-OFFinder (<http://www.rgenome.net/cas-offinder/>) to blast the genome of *H. armigera*, and the mismatch number was set to less than or equal to 3. According to the predicted results, the Gr10 sgRNA had no potential off-target region but Gr6 sgRNA had one. Therefore, we amplified and sequenced the potential off-target region of Gr6^{-/-} and found there was no frameshift or premature stop codon in the region compared to WT (Figure 5—figure supplement 2). It is worth mentioning that there was no potential off-target region of Gr6 and Gr10 sgRNA in other sugar receptor genes of *H. armigera*, Gr4, Gr5, Gr7, Gr8, Gr9, Gr11 and Gr12. We further found there was no difference in the response to xylose of the medial sensilla styloconica among WT, Gr10^{-/-} and Gr6^{-/-} (Figure 5—figure supplement 2). Furthermore, WT, Gr10^{-/-} and Gr6^{-/-} did not show differences in the larval body weight, adult lifespan, and number of eggs laid per female (Figure 5—figure supplement 2). All these results suggest that no off-target effects occurred in the study.”

| *(7) Is it possible that knocking out Gr10 is not compensated for by the overexpression of Gr6 or other sucrose sensing Grs? Similarly, would the vice versa scenario hold true?*

In the Discussion section, we have added some sentences to discuss this issue: “From our results, knocking out Gr10 or Gr6 is unlikely to be compensated by overexpression of other sugar GRs. One of our recent studies showed that Orco knockout had no significant effect on the expression of most OR, IR and GR genes in adult antennae of *H. armigera*, but some genes were up- or down-regulated (Fan et al., 2022).”

| *(8) What was the rationale for selecting nine candidate GR genes for expression analysis?*

Based on the reviewer's suggestion, we expanded the relevant paragraphs to illustrate the rationale for selecting nine candidate GR genes for expression analysis: “To reveal the molecular basis of sugar reception in the taste sensilla of *H. armigera*, we first analyzed the putative sugar gustatory receptor genes based on the reported gene sequences of GRs in *H. armigera* and their phylogenetic relationship of *D. melanogaster* sugar gustatory receptors (Jiang et al., 2015; Pearce et al., 2017; Xu et al., 2017). Nine putative sugar GR genes, Gr4–12 were identified, and their full-length cDNA sequences were cloned (The GenBank accession number is provided in Appendix—Table S1).” (Line 155-161)

| *(9) What is the potential reason for the difference between the major larval sugar receptors of Drosophila and Lepidopterans?*

The difference between the major larval sugar receptors of *Drosophila* and *Lepidopterans* is probably due to differences in the food their larvae feed on. Fruit fly larvae feed on rotten fruit, the main sugar of which is fructose. The larvae of *Lepidoptera* mainly feed on plants, and the main sugar is sucrose. In the Discussion section, we have added a sentence “This is

most likely due to fruit fly larvae feeding on rotten fruits, which contain fructose as the main sugar.” (Line 399-401)

(10) There is a disparity in GRs, specifically GR5 and GR6, between the female antenna, proboscis, and tarsi. What could be the possible justification and significance of this?

Thank you so much for this question. We have added a sentence in the Discussion section, “In this study, the expression patterns of 9 sugar GRs in three taste organs of adult *H. armigera* show that there is a disparity in GRs, specifically GR5 and GR6, between the female antenna, tarsi and proboscis, which may be an evolutionary adaptation reflecting subtle differentiation in the function of these taste organs in adult foraging. Antennae and tarsi play a role in the exploration of potential sugar sources, while the proboscis plays a more precise role in the final decision to feed.” (Line 433-438)

(11) I suggest that a visual representation illustrating the positioning of GSNs, particularly the lateral and medial sensilla, in both larva and adult stages would enhance the correlation with the results.

In Figure 1 we added the photo of each taste organ and the position of the recorded sensilla, and also added a new figure, Figure 8 summarizing the main findings of the studies.

(12) Further experiments can be conducted to elucidate the precise molecular mechanisms, particularly the downstream effects of GRs, in order to establish the specificity of GRs more convincingly.

Thank you so much for your suggestion. We have discussed the further experiments in the Discussion section, “To elucidate the precise molecular mechanisms of sugar reception in *H. armigera* is necessary to compare a series of single, double and even multiple Gr knock-out lines and investigate the downstream effects of the GRs.” (Line 363-369)

(13) Figure 6 caption: In Figure 6 (D to I), the percentage of PER is depicted. There is redundancy in the Y-axis title (Percentage of PER) and the legend. This appears to be repetitive. I suggest that it would be better to include the Y-axis title only in Figure D or in Figures D and G.

We accept the suggestion. Figure 7 (not Figure 6) has been revised accordingly.

(14) In Figures 6A and 6C, there is inconsistency in the colors used for WT, Gr6, and Gr10. This could potentially confuse the reader. I recommend using the same colors in both figures instead of using a blue color. Please specify how the authors calculated the feeding area in Figure 6.

We accept the reviewer's suggestion and have changed the color of Figure 7A, B. We have also added the detail method for calculating feeding area (Line 541-545).

(15) In Two-choice tests, why did the authors use 0.01% Tween 80? Please provide comments on this.

Use of 0.01% Tween 80 is to reduce the surface tension and increase the malleability of the solution. We have given detailed explanation in the Method section and cite the reference. (Line 538-540)

(16) It would be valuable if the authors could comment on the prospects of this study, considering that GRs play a vital role in controlling behavior and developmental

pathways. What are the potential consequences of blocking or disrupting these receptors in terms of behavioral and developmental phenotypic deformities? Could this potentially lead to increased insect mortality?

Thank you so much for your suggestions. In the last paragraph of the Discussion section, we have added the following perspectives, “Knockout of Gr10 or Gr6 led to a significant decrease in sugar sensitivity and food preference of the larvae and adults of *H. armigera*, respectively, which is bound to bring adverse consequences to survival and reproduction of the insects. Therefore, studying the molecular mechanisms underlying sugar perception in phytophagous insects may provide new insights into the behavioral ecology of this important and highly diverse group of insects, and measures blocking or disrupting sugar receptors could also have applications to control agricultural pests and improve crop yields worldwide” (Line 449-456).

Reviewer #2 (Recommendations for The Authors):

There are a few comments, that I feel would be beneficial to be addressed.

- *The authors used 7 different sugars for their experimental approach. While I agree that this is a sufficiently large collection for a study, I was wondering why they specifically chose these sugars; an explanatory section might be helpful for a reader to follow the reasoning.*

According to reviewer 1's suggestion, we increased trehalose to 8 sugars in experiments. Trehalose is a main sugar in insect blood. It is converted by insects after feeding on plant sugars. The 8 sugars were chosen because they are present in host-plants of *H. armigera* or are representative in the structure and source of sugars. They contain 2 pentoses (arabinose and xylose), 4 hexoses (fructose, fucose, galactose and glucose), and 2 disaccharides (sucrose and trehalose).

- *It might be beneficial to provide some broader overview on the gustatory system in the cotton bollworm, particularly at the larval stage since this may not be common knowledge. Along these lines eg. the complexity of sensilla types, organs and overall number (or estimation) of neurons might be good to know, a graphical representation of the sense organs might be informative.*

In the Introduction section, we give a more specific description on sugar sensitive GSNs in the taste system of the larva and adult of *H. armigera*, and cite the corresponding references.

- *Concerning phylogeny of GRs, it might be relevant to know how complete the genome information is and some more general background on GR diversity in the cotton bollworm.*

We agree to your opinion. According to this idea, we got the putative sugar GRs from the previously published genome (Pearce et al. 2017) and the related annotation of GRs (Jiang et al. 2015, Xu et al. 2012). We have made a more detailed explanation about this in the new version of the manuscript, “We first analyzed the putative sugar gustatory receptor genes based on the genome data of *H. armigera* (Pearce et al. 2017), the reported gene sequences of sugar GRs in *H. armigera* and their phylogenetic relationship of *D. melanogaster* sugar gustatory receptors (Jiang et al. 2015, Xu et al. 2012). All nine putative sugar GR genes in *H. armigera*, Gr4–12 were validated, and their full-length cDNA sequences were cloned (The GenBank accession number is provided in Appendix—Table S1).” (Line 155-161).

- *Generation of mutants based on CRISPR is intriguing and a powerful step. While the techniques are well described in the method section, there is no information concerning efficiency or broader feasibility of the approach. I feel it would be quite interesting to learn about how feasible or laborious the approach is to generate mutants (e.g. number of initial injected eggs, the resulting F0 offspring, number of back-crosses, number of screened F1s).*

In the Materials and Methods section, we have added specific success rates for each step in the process of building the two mutants (Line 722-726, 729-732).

Reviewer #3 (Recommendations For The Authors):

I want to congratulate the authors on this very nice study and have only minor comments for them.

*(1) It would be very nice to include pictures of the larva and adult of *H. armigera*. It would also help to have schematics of where the sensilla they are recording from are.*

We have added photos of four taste organs on which the recorded sensilla were indicated (Figure 1), and picture of the larva and adult on which the stimulating site was indicated (Figure 2).

(2) A schematic summarising their findings, including the relevance to the animal's behavioural ecology, will greatly improve interpretations for the broader audience.

A schematic summarizing the findings has been added.

(3) The manner in which PIs are represented in figure 2A, B (among others) is confusing. Can the authors please plot the PI and not the feeding area? From the PI values listed beside the plot, it actually suggests that the larvae don't really show a preference. Could the authors please comment on this?

Yes, sucrose has a significant stimulating effect on larva feeding, but the effect is not as large as the predicted based on the sensitivity of the sensillum, the main reasons are as follows: (1) there are many factors affecting larva feeding, sucrose is only one of them; (2) due to the substrate leaf discs also contain sugar, the effect of newly added sucrose may be reduced. After careful consideration, we think it is better to display the feeding area and PI together so that readers have a complete understanding of the data.

(4) The heterologous expression experiments suggest that co-expression of GR6 with either GR10 or GR5 somehow suppress the response of the GR6 alone to fucose. Am I reading the data correctly? Why would this be? Perhaps the authors could discuss this. In this context, it would help to reproduce all the GR6 data together.

Your interpretation is reasonable to a certain extent. The result of co-injection might be that Gr10 or Gr5 inhibited the response of Gr6. However, there is another possibility that the amount of Gr6 sRNA was diluted by co-injection of two GRs, resulting in a reduced response of Gr6 to fucose.

(5) In general, for each results section, it would help to have a sentence or two that interprets the data in the context of previously presented data. This would help the reader digest the data and interpret it as they read along. Currently, the authors summarise the observations and leave all the interpretation to the discussion section.

We accept the suggestion. In each part of the results, we have added a sentence to explain the above data, which will help readers to clarify the context of the research more easily.

| (6) *Is the GR6 data in 4C not lined up correctly?*

Yes, it is right.

| (7) *Line 228 suggests that the mutants were validating with qPCRs - I don't see that data.*

The mutants were not validating with qPCR. We used the ordinary PCR technology at the mRNA level to verify whether the related sequences were really deleted in the mutants.