

PI3K α inhibition blocks osteochondroprogenitor specification and the hyper-inflammatory response to prevent heterotopic ossification

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

October 9, 2023 (this version)

Posted to bioRxiv

September 14, 2023

Sent for peer review

August 29, 2023

José Antonio Valer, Alexandre Deber, Marius Wits, Carolina Pimenta-Lopes, Marie-José Goumans, José Luis Rosa, Gonzalo Sánchez-Duffhues, Francesc Ventura 

Departament de Ciències Fisiològiques, Universitat de Barcelona, IDIBELL, C/ Feixa Llarga s/n 08907 Hospitalet de Llobregat, SPAIN • Department of Cell and Chemical Biology, Leiden University Medical Center, Einthovenweg 20, 2333 ZC, Leiden, the Netherlands • Nanomaterials and Nanotechnology Research Center (CINN-CSIC), Health Research Institute of Asturias (ISPA), 33011 Oviedo, Asturias, Spain

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Heterotopic ossification (HO) occurs following mechanical trauma and burns, or congenitally in patients suffering from fibrodysplasia ossificans progressiva (FOP). Recently, we demonstrated that inhibitors of phosphatidylinositol 3-kinase α (PI3K α) may be a useful therapy for patients undergoing HO. In this study, using the already marketed BYL719/Alpelisib/Piqray drug, we have confirmed further these results, detailed the underlying mechanisms of action, and optimized the timing of the administration of BYL719. We found that BYL719 effectively prevents HO even when administered up to three to seven days after injury. We demonstrate in cell cultures and in a mouse model of HO that the major actions of BYL719 are on-target effects through the inhibition of PI3K α , without directly affecting ACVR1 or FOP-inducing ACVR1^{R206H} kinase activities. *In vivo*, we found that a lack of PI3K α in progenitors at injury sites is sufficient to prevent HO. Moreover, time course assays in HO lesions demonstrate that BYL719 not only blocks osteochondroprogenitor specification, but also reduces the inflammatory response. BYL719 inhibits the migration, proliferation and expression of pro-inflammatory cytokines in monocytes and mast cells, suggesting that BYL719 hampers the hyper-inflammatory status of HO lesions. Altogether, these results highlight the potential of PI3K α inhibition as a safe and effective therapeutic strategy for HO.

eLife assessment

This paper presents **valuable** insights into the implication of PI3K α in heterotopic ossification (HO), and illustrates a potential therapeutic efficacy of BYL719 in suppressing HO within a murine model of Fibrodysplasia Ossificans Progressiva. While certain data are novel and **compelling**, others exhibit redundancies with prior publications and are **inadequate** in terms of methodology and presentation. Additionally, elucidating the precise molecular mechanisms of BYL719's action is imperative for a comprehensive understanding.

Introduction

Heterotopic ossification (HO) is a disorder characterized by ectopic bone formation at extraskeletal sites, including skeletal muscle and connective tissues. Trauma-induced HO develops as a common post-operative complication after orthopedic surgeries (e.g., hip arthroplasty), blast injuries, deep burns and nervous system injuries (Hwang et al., 2022 [\[1\]](#)). In addition to trauma-induced HO, fibrodysplasia ossificans progressiva (FOP) is a rare congenital autosomal dominant disorder also involving HO (Bravenboer et al., 2015 [\[2\]](#)). Ectopic bones form progressively through endochondral ossification, mostly in episodic flare-ups associated with inflammation (Towler and Shore, 2022 [\[3\]](#)). These cumulative osteogenic events lead to reduced mobility resulting from ankylosing joints. Affected individuals have a shorter life-span, most commonly due to thoracic insufficiency syndrome (Kaplan et al., 2010 [\[4\]](#); Pignolo et al., 2016 [\[5\]](#)).

HO reflects a shift from a normal tissue repair process to an aberrant reactivation of bone-forming developmental programs that require acute inflammation and the excessive expansion of local progenitors, followed by inappropriate differentiation of these progenitors into chondroblasts and osteoblasts which finally results in bone formation (Hwang et al., 2022 [\[1\]](#)). In both trauma-induced HO and FOP, pathology appears following the excessive activation of receptors sensitive to Transforming growth factor- β (TGF β) superfamily family members (Lees-Shepard et al., 2018 [\[6\]](#); Wang et al., 2018 [\[7\]](#)). FOP arises from gain-of-function mutations in the bone morphogenetic protein (BMP) type I receptor *ACVR1*, with the most common mutation being c.617G>A, R206H (Shore et al., 2006 [\[8\]](#)). Higher SMAD 1/5 mediated signaling of mutated *ACVR1*/ALK2 has been partially attributed to a loss of auto-inhibition of the receptor and mild hypersensitivity to BMP ligands (Chaikuad et al., 2012 [\[9\]](#); Groppe et al., 2011 [\[10\]](#); Van Dinther et al., 2010 [\[11\]](#)). Signaling by mutated *ACVR1* still relies on ligand-induced heterotetrameric clustering but, unlike wild type *ACVR1*, does not require *ACVR2A/B* kinase activity (Agnew et al., 2021 [\[12\]](#); Ramachandran et al., 2021 [\[13\]](#)). More importantly, mutations alter their signaling specificity, abnormally transducing SMAD 1/5 and altering the non-canonical BMP signals (e.g., p38 or PI3K) in response to activin A (Hatsell et al., 2015 [\[14\]](#); Hino et al., 2015 [\[15\]](#); Valer et al., 2019a [\[16\]](#)). Accordingly, evidence supports that activin A is necessary and sufficient for HO in FOP (Hatsell et al., 2015 [\[14\]](#); Lees-Shepard et al., 2018 [\[6\]](#); Upadhyay et al., 2017 [\[17\]](#)). In contrast, activin A does not drive post-traumatic HO (Hwang et al., 2020 [\[18\]](#)), and non-genetically-driven HO mostly arises from excessive BMP and TGF β signaling, with functional redundancy between different type I receptors of the TGF β -superfamily (Agarwal et al., 2017 [\[19\]](#); Patel et al., 2022 [\[20\]](#); Sorkin et al., 2020 [\[21\]](#); Wang et al., 2018 [\[7\]](#)).

Evidence points to mesenchymal fibroadipogenic precursors (FAPs), which are widely distributed in muscle and other connective tissues, as the key cell-of-origin that aberrantly undergo chondrogenesis and further ectopic bone formation (Dey et al., 2016 [\[22\]](#); Eisner et al., 2020 [\[23\]](#); Lees-Shepard et al., 2018). However, either FOP or non-genetic HO also require local tissue destruction and inflammation, which indicate that FAPs should be invariably primed for ectopic bone formation by this inflammatory microenvironment (Barruet et al., 2018 [\[24\]](#); Convente et al., 2018 [\[25\]](#); Hwang et al., 2022 [\[1\]](#); Matsuo et al., 2019 [\[26\]](#); Pignolo et al., 2013 [\[27\]](#); Sorkin et al., 2020 [\[21\]](#)).

Inflammation in HO is characterized by an initial acute response following injury with the activation of innate immunity and the influx of neutrophils and monocytes (Convente et al., 2018 [\[25\]](#); Hwang et al., 2022 [\[1\]](#)). Additionally, B and T cells of the adaptive system are also recruited (Chakkalakal et al., 2012 [\[28\]](#); Convente et al., 2018 [\[25\]](#)). However, mice lacking B or T lymphocytes exhibited no delay in the development of heterotopic ossification (HO) after injury, indicating that these cells may play a subsequent role in the dissemination of bone lesions (Kan et al., 2009 [\[29\]](#)). During intermediate and late inflammatory stages, the recruitment of monocytes, macrophages and mast cells occurs, which in turn exerts autocrine and paracrine effects on nearby FAPs (Chakkalakal et al., 2012 [\[28\]](#); Convente et al., 2018 [\[25\]](#); Sorkin et al., 2020 [\[21\]](#); Tu et al., 2022 [\[30\]](#)).

Highlighting its relevance in HO, the depletion of mast cells and macrophages profoundly impairs genetic and non-genetic HO (Convente et al., 2018 [↗](#); Torossian et al., 2017 [↗](#); Tu et al., 2022 [↗](#)). BMP receptors are robustly expressed in monocytes and macrophages and the expression of *ACVR1^{R206H}* has been shown to extend inflammatory responses in patient-derived macrophages (Barruet et al., 2018 [↗](#); Matsuo et al., 2021 [↗](#), 2019 [↗](#)). From the plethora of cytokines secreted by monocytes, macrophages and mast cells, both activin A and TGF- β stand out as extremely relevant for HO (Alessi Wolken et al., 2017 [↗](#); Hattell et al., 2015 [↗](#); Lees-Shepard et al., 2018 [↗](#); Patel et al., 2022 [↗](#); Sorkin et al., 2020 [↗](#); Upadhyay et al., 2017 [↗](#)).

Genetic and pharmacological studies have indicated that osteochondroprogenitor specification and maturation depend on phosphatidylinositol 3-kinase- α (PI3K α) (Ford-Hutchinson et al., 2007 [↗](#); Fujita et al., 2004 [↗](#); Gámez et al., 2016 [↗](#); Ikegami et al., 2011 [↗](#)). We found that PI3K signaling was also linked to HO, since inhibitors of PI3K α (BYL719/Alpelisib/Piqray) prevented HO in mouse models without major side-effects (Valer et al., 2019a [↗](#)). Mechanistically, PI3K α inhibitors hamper canonical and non-canonical BMP signaling, decreasing total and phosphorylated SMAD1/5 levels and reducing transcriptional responsiveness to BMPs/Activin A in mesenchymal progenitors (Gámez et al., 2016 [↗](#); Valer et al., 2019b [↗](#)). In the current study, we demonstrate that the pharmacological and genetic inhibition of PI3K α in HO progenitors at injury sites reduces HO *in vivo*. Moreover, envisioning a future translation into the clinic, we have optimized the administration of BYL719 and found that the delayed administration of BYL719, up to seven days after inflammatory injury, still prevents HO in mice. In addition, we found that BYL719 blocks osteochondroprogenitor specification and effectively reduces the essential inflammatory response. Altogether, the data presented here show the potent therapeutic effect of PI3K α inhibition on HO in mouse models.

Results

Delayed initiation of treatment with PI3K α inhibitor effectively prevents heterotopic ossification

Heterotopic ossification takes place following a precise temporal pattern of progenitor activation and the recruitment of distinct cell types at the ossification centers which causes early divergence from normal skeletal muscle repair program. We previously found that the pharmacological administration of BYL719 prevents HO in a mouse model of HO (Valer et al., 2019a [↗](#)). Given that BYL719 is marketed for the treatment of cancer and overgrowth syndrome, it is a promising therapeutic molecule in pathological HO. To further determine a potential therapeutic window for BYL719 and better understand the cellular targets of BYL719, we evaluated the effect of BYL719 when administered intermittently and several days after HO induction. For this, we used a conditional mouse model (*ACVR1^{Q207Dfl/fl}*) in which we combine the overexpression of Cre recombinase through adenoviral particles with the injection of cardiotoxin intramuscularly in the hindlimb (Fukuda et al., 2006 [↗](#)). In this model, we administered BYL719 (i.p. 25 mg/kg) or vehicle control intermittently starting one, three or seven days after HO induction. We also implemented a different regimen involving the administration of BYL719 only for the initial three days following the injury (**Figure 1A** [↗](#)). None of the treatments led to significant changes in the weight of the mice (Figure 1-Figure Supplement 1). HO volume was analyzed by micro computed tomography (μ CT) twenty-three days post-injury. The early administration of the treatment with BYL719 at day one or three after the onset of HO resulted in significantly lower HO; late administration seven days after the induction of HO was also partially effective (**Figure 1B and C** [↗](#)). Discontinuation of the treatment after three days did not prevent HO at day 23. These results indicate that the inhibition of PI3K α by BYL719 after injury prevents HO and this protection is still effective if the treatment is started several days after injury, expanding its therapeutic window.




Figure 1.

(A) Heterotopic ossification was induced at P7 injecting adenovirus-Cre (Ad.Cre) and cardiotoxin in the mice hindlimb. The pattern of 5 administrations of DMSO or BYL719 (25 mg/kg) is indicated by grey dots started at P8 (start day 1), P10 (delayed start day 3), and P14 (delayed start day 7). One group only received the administration of DMSO or BYL719 (25 mg/kg) at P8, P9, and P10 (first 3 days post-HO-induction), as indicated by grey dots. The final time-point, P30, was conserved between experimental groups. (B) Quantification of heterotopic ossifications bone volume (mm³) of each experimental group. Colored symbols indicate the presence of heterotopic ossifications. Black symbols indicate the absence of heterotopic ossifications. Data shown are of each individual mouse with group median. *P<0.05, **P<0.01, Kruskal-Wallis test with Dunn's multiple comparison test. (C) Representative 3D microtomography images of the injected hindlimbs of 3 different mice for each experimental group. White arrows show heterotopic ossification.

Deficiency of PI3Kα at injury sites is sufficient to partially prevent heterotopic ossification

Since other small molecules designed to target PI3Kα have reported partial off-target effects on kinases other than PI3Kα (Furet et al., 2013 [↗](#); Jamieson et al., 2011 [↗](#)), we aimed to confirm that the results obtained with BYL719 were due to the specific inhibition of PI3Kα. Therefore, we developed a conditional mouse model ($ACVR1^{Q207D/f/f};p110\alpha^{f/f}$) in which the Cre recombinase drives the expression of $ACVR1^{Q207D}$ and deletion of the catalytic subunit of PI3Kα ($p110\alpha$) after intramuscular hindlimb combined injection of adenovirus-Cre and cardiotoxin to trigger HO. With this approach, Cre mediates the expression of $ACVR1^{Q207D}$ and the deletion of PI3Kα in the same cells, whereas cells recruited afterwards would be wild type for both genes. As expected, μ CT analysis performed 23 days after injury showed extensive HO in mice expressing $ACVR1^{Q207D}$, which was prevented by intermittent treatment with BYL719 initiated one day after injury (**Figure 2A** [↗](#), **2B** [↗](#) and **2D** [↗](#)). Genetic deletion of $p110\alpha$ in mice expressing $ACVR1^{Q207D}$ led to a reduction in HO, which was further enhanced by BYL719 (**Figure 2B** [↗](#)). Importantly, no significant changes in the weight of the mice were observed, confirming the absence of severe toxicity (Figure 2-Figure Supplement 1A). Histomorphometrically, $ACVR1^{Q207D}$ mice wild type for PI3Kα developed islands and/or bone spurs with a well-organized bone structure. However, in PI3Kα deficient mice $ACVR1^{Q207D}$ expression only led to minor ectopic calcifications that were already surrounded by fully regenerated muscle tissue on the 23rd day after injury (**Figure 2D** [↗](#), Figure 2-Figure Supplement 1B). Accordingly, analysis of the ratio bone volume/tissue volume (BV/TV) and the correlation between bone volume and BV/TV of individual mice was clearly different between $p110\alpha$ genotypes, irrespective of BYL719 treatment (**Figure 2C** [↗](#), Figure 2-Figure Supplement 1C). These results demonstrate that the abrogation of PI3Kα activity (either genetic or pharmacologically) in cells at injury sites is relevant for their inhibitory effects in HO. This might be attributed to the direct effects of BYL719 on progenitor cell expansion and chondroblast specification, as well as its possible ability to reduce the inflammatory response required for HO.

Genetic or pharmacological inhibition of PI3Kα prevents osteochondroprogenitor specification

Next, we aimed to assess the effects of PI3Kα inhibition in osteochondroprogenitor cells. For this, we isolated bone marrow-derived mesenchymal stem cells (BM-MSCs) from $p110\alpha^{f/f}$ mice and transduced them *ex-vivo* with retrovirus expressing *Acvr1* (wild type and $R206H$) with or without Cre recombinase. The exogenous mRNA gene expression of both *Acvr1* and $Acvr1^{R206H}$ was similar (**Figure 3A** [↗](#)). As anticipated, transduction with Cre viruses significantly reduced the expression levels of $p110\alpha$ to approximately 40%. This is consistent with an estimated transduction efficiency of nearly 60% (**Figure 3B** [↗](#)). The overexpression of $Acvr1^{R206H}$ increased basal and activin dependent expression of canonical (*Id1* and *Sp7*) and non-canonical (*Ptgs2*) BMP target genes (**Figure 3C** [↗](#)), known markers and drivers of osteochondrogenic differentiation (Nakashima et al., 2002 [↗](#); Wang et al., 2013 [↗](#)). Both pharmacological and genetic inhibition of PI3Kα reduced canonical and non-canonical $ACVR1^{R206H}$ transcriptional responses induced by recombinant activin A. Of note, stimulation with activin A increased its own transcription (*Inhba*) in $Acvr1^{R206H}$ -transduced BM-MSCs, which was also inhibited by BYL719 (**Figure 3C** [↗](#)). Therefore, BYL719 may also prevent osteochondroprogenitor differentiation indirectly, via transcriptional inhibition of *INHBA*/Activin A expression. These results demonstrate a direct effect of BYL719 on osteochondroprogenitor cells and suggest that the effects of BYL719 on $ACVR1$ -downstream signaling are mainly on target, due to PI3Kα kinase inhibition.

It is known that 2-aminothiazole-derived PI3Kα inhibitors could also inhibit $ACVR1$ kinase activity (Jamieson et al., 2011 [↗](#)). Therefore, we investigated whether BYL719 effects could be explained by the reduction in $ACVR1$ kinase activity. For this, we made use of a target kinase engagement approach. Upon transient transfection of Nanoluciferase-tagged TGFβRs, we incubated the cells




Figure 2.

(A) Heterotopic ossification was induced at P7 injecting adenovirus-Cre (Ad.Cre) and cardiotoxin in the mice hindlimb. Starting at P8, mice were treated with DMSO or BYL719 (25 mg/kg) as indicated by grey dots for 5 administrations. (B) Quantification of heterotopic ossifications bone volume (mm³) of each experimental group. Colored symbols indicate the presence of heterotopic ossifications. Black symbols indicate the absence of heterotopic ossifications. Data shown are of each individual mouse with group median. *P<0.05, **P<0.01, Kruskal-Wallis test with Dunn's multiple comparison test. (C) Quantification of the ratio of bone volume per tissue volume (BV/TV) within heterotopic ossifications of each experimental group, including only mice with detected HO. Data shown are of each individual mouse with the group median. **P<0.01, ***P<0.001, Kruskal-Wallis test with Dunn's multiple comparison test. (D) Representative 3D frontal microtomography images of the injected hindlimbs of mice for each experimental group and a detailed close-up image for each selected mouse. White arrows show heterotopic ossification in the close-up images.




Figure 3.

mRNA expression of *Acvr1* (A) or *p110α* (B) genes in BM-MSCs from *p110α^{fl/fl}* mice, infected with virus expressing wild-type *Acvr1* (WT) or *Acvr1^{R206H}* (RH) and/or Cre co-infection. (A) Comparison to the endogenous expression level of *Acvr1* gene in mock-transfected BM-MSCs is shown as a dotted horizontal line. Data are shown as mean $\pm$ SD ($n = 12$ per group). Unpaired t -test between transfected groups. (B) Data are shown as mean $\pm$ SD ($n = 3$ per group). *** $P < 0.001$, two-way ANOVA with Tukey's multiple comparisons test. (C) mRNA expression of canonical (*Id1* and *Sp7*), non-canonical (*Ptgs2*) target genes, and activin A (*Inhba*) in BM-MSCs *p110α^{fl/fl}* transfected with *Acvr1* (wild type or R206H) with or without Cre recombinase. Cells were NT (not treated), or treated with BYL719 (2μM) and/or activin A (2nM). Expression data was normalized to those of control cells which were transfected only with *Acvr1* WT without any treatment, shown as a dotted horizontal line. Asterisks (*) refer to the differences between *Acvr1* RH cells compared to control cells or *Acvr1* RH cells without Cre recombinase treated with BYL719 or activin A. Hash signs (#) refer to the differences between *Acvr1* RH cells without Cre recombinase treated with activin A compared to the same condition with Cre recombinase, or compared with *Acvr1* RH cells treated with BYL719 and/or activin A. Data are shown as mean $\pm$ SD ($n = 6$ per group). * or # $P < 0.05$, ** or ## $P < 0.01$, *** or ### $P < 0.001$, two-way ANOVA with Tukey's multiple comparisons test.

with BYL719 and/or an ATP-like tracer analogue. In the absence of an ATP competitor molecule, the tracer analogue and Nanoluciferase enzyme stay in close proximity, thereby allowing for bioluminescence resonance energy transfer (BRET) between the Nanoluciferase (energy donor, emitting at 460 nm) and the tracer molecule (energy acceptor). Once excited, the acceptor molecule emits fluorescence (610 nm) (**Figure 4A and B**). BYL719 incubated at 1 and 10 μM failed to significantly inhibit the emission of the BMP type I receptors ALK1, ALK2 (ACVR1), ALK3, ALK4, ALK5 and the mutant receptor ACVR1/ALK2^{R206H} (**Figure 4C**). In addition, we demonstrated that BYL719 does not target the activity of the type II kinase receptors ACVR2A, ACVR2B, BMPR2 and TGFBR2. We confirmed these results using *in vitro* kinase activity assays with recombinant ACVR1^{R206H}. BYL719 exhibited no major effects on ACVR1^{R206H} activity up to 10 μM (**Figure 4D**).

Given that BYL719 does not directly target the activity of ACVR1 or related receptors, we established an *in vitro* model using human MSCs (hMSCs) stably overexpressing ACVR1 or ACVR1^{R206H} to profile further the main targeted pathways. In this model, recombinant Activin A was able to induce SMAD1/5 activation and the expression of ACVR1 downstream targets *ID1* and *ID3* in cells over-expressing ACVR1^{R206H} (Figure 5-Figure Supplement 1A and B). In these cells, coincubation with BYL719 (10 μM) was able to repress Activin A-induced Alkaline phosphatase expression and activity upon 7 and 11 days of culture, respectively (**Figure 5A, B** and Figure 5-Figure Supplement 1C). Furthermore, we performed 3D chondrogenic differentiation assays. Coincubation with BYL719 (10 μM) almost completely inhibited the formation of a chondrogenic glycosaminoglycans-rich matrix in response to Activin A after 3 weeks of culture (**Figure 5C and D**) and partially abrogated the expression of the osteochondrogenic genes *COL1A1* and *ALPL* in early chondrogenic differentiation (**Figure 5E**). These results confirmed that BYL719 administration blocks osteochondrogenic specification irrespective of the stage of differentiation.

To identify the immediate targets of BYL719, we performed bulk RNA sequencing (GSE237512) where we preincubated hMSCs-ACVR1^{R206H} with BYL719 for 30 min, which was followed by stimulation with recombinant Activin A (50 ng/ml) for 1 hour (**Figure 6A**). Consistent with our results on osteogenic progenitor cells, preincubation with BYL719 (1 μM) resulted in the robust regulation of genes involved in ossification and osteoblast differentiation (including *PTGS2*, *MSX2*, *SOX9*, and *BMP2*), as identified through enrichment analysis of the GO terms GO:0001503 ($\text{padj}=3.83\text{e}^{-5}$) and GO:0001649 ($\text{padj}=3.10\text{e}^{-5}$), respectively (**Figure 6B and C**), indicating that BYL719 mainly targets osteochondral specification processes in these cells. Moreover, a number of inflammatory cytokine genes (e.g. *IL6* and *IL15*) and inflammatory signaling pathways (TNF and NF κ B), were downregulated by BYL719 (Figure 6-Figure Supplement 1A and B). Therefore, we hypothesized that, apart from preventing the activation of osteogenic and chondrogenic differentiation pathways in progenitor cells, part of the mechanism by which BYL719 prevents HO *in vivo* may be due to the modulation of inflammatory responses.

PI3K α inhibition reduces proliferation, migration and inflammatory cytokine expression in monocytes, macrophages and mast cells

Next, we used several models of disease-relevant immune cells to study the effect of pharmacological PI3K α inhibition. Activation of ACVR1 signaling with BMP6 did not significantly modify the proliferation rate of any of the cell lines tested, that is, monocytes (THP1), macrophages (RAW264.7) and mast cells (HMC1) (**Figure 7A, B** and **C**). However, incubation with BYL719 *in vitro* strongly reduced their growth rate at 2 μM and almost completely arrested their proliferation at 10 μM . It is well known that monocytes are actively recruited from circulation to the ossification sites early after injury (Sorkin et al., 2020). Therefore, we also analyzed chemotactic migration of monocytes in transwell assays. Whereas monocytes actively migrated towards FBS as a




Figure 4.

Transforming growth factor- β receptor (TGF β R) kinase assays. (A) A schematic depiction of nano-bioluminescence resonance energy transfer (nanoBRET) target engagement assays. TGF β R-Nanoluciferase fusion proteins are expressed in live COS-1 cells and an ATP-like tracer analogue causes proximity-based BRET from the Nanoluciferase donor to the fluorescent tracer acceptor. Competition of the tracer with a possible ATP-pocket inhibitor would result in decreased nanoBRET. (B) The nanoBRET emission spectra consist of the Nanoluciferase donor (460nm) and the fluorescent acceptor (610nm). The nanoBRET ratio is shown in millibRET units (mBU) by dividing the acceptor emission by the donor emission times 1000. (C) NanoBRET target engagement analyses of ALK1, ALK2 (ACVR1), ALK2^{R206H}, ALK3, ALK4, ALK5, TGFBR2, ACVR2A, ACVR2B and BMPR2 testing 1 or 10 μ M BYL719 with $n=4$. As controls, LDN193189 (0.5 μ M), SB431542 (10 μ M), and ML347 (10 μ M) were used. Data are shown as mean $\pm$ SD. One-way ANOVA with Dunnett's multiple comparisons test. (D) Casein phosphorylation by ACVR1^{R206H} kinase. Phosphorylation was performed in the presence of ACVR1^{R206H} kinase and increasing concentrations of the PI3K α inhibitor BYL719. Quantification of kinase activity. Data shown as mean $\pm$ SD ($n=4$ independent experiments). One-way ANOVA with Tukey's multiple comparisons test. * $P<0.05$, ** $P<0.01$, *** $P<0.001$.




Figure 5.

Analyses of chondrogenic progenitor specification (using differentiation medium; DM) of hMSC-ACVR1/ALK2^{WT} or hMSC-ACVR1/ALK2^{R206H} treated with BYL719 (10 μM) with and without Activin A (50 ng/ml), compared to growth medium (GM). (A) Representative images of the ALP staining after one week of chondrogenic progenitor specification. The images were obtained at 5x magnification and the scale bar represents 1000 μm. (B) Quantification of the ALP staining by densitometry (n=3 per group). (C) Representative images of Alcian Blue staining of micromass cultures after 3 weeks of chondrogenic progenitor specification. The images were taken at 2.5x magnification and the scale bar represents 500 μm. (D) Quantification of the solubilized Alcian Blue staining via the measurement of absorbance (595nm) (n=3 per group). (E) RT-qPCR results of COL1A1 and ALPL after 3 days of chondrogenic progenitor specification (n=3 per condition). Data are shown as mean ± SD. In all figure panels, significance is detailed between DM and DM with BYL719, and between DM with Act A and DM with Act A and BYL719. * P<0.05, ** P<0.01, *** P<0.001, two-way ANOVA with Tukey's multiple comparisons test.




Figure 6.

Bulk RNA sequencing of hMSC-ACVR1/ALK2^{R206H} upon overnight starvation, 30 minutes pre-treatment with 1 μM BYL719 and 1 hour Activin A (50 ng/ml) stimulation including continued BYL719 treatment. (A) Schematic depiction of the experimental setup. (B) Gene Set Enrichment Analysis (GSEA) with the groups described in (B), showing the significant enrichment plots of the gene ontology sets ossification (GO: 0001503, padj=3.83e⁻⁵) and osteoblast differentiation (GO:0001649, padj=3.10e⁻⁵). (C) A heatmap of the top 40 most relevant genes within the ossification GO geneset derived from the leading-edge subset of the enrichment plot (n=4 per group). The statistics are described in the methods section.

chemotactic agent, the incubation of monocytes with BYL719 deeply diminished their migratory ability (**Figure 7D**). These results suggest that BYL719 is able to diminish monocyte recruitment, while reducing also the expansion of inflammatory cells at injury sites.

Upon tissue injury, a multitude of pro-inflammatory and anti-inflammatory cytokines are released at the site of injury. This triggers the recruitment and infiltration of various cell types, including immune cells and osteochondrogenic progenitors. We examined the expression of distinct cytokines upon ACVR1 stimulation with BMP6 in the presence of BYL719 in monocytic, macrophagic and mast cell lines. In monocytes, BYL719 inhibited the expression of the pro-inflammatory cytokine *TNF- α* and the expression of activin A (*INHBA*), whereas the expression of transforming growth factor *TGF- β 1* was increased by BMP6 and not modified by the addition BYL719 (**Figure 7E**). *CCL2* (MCP-1), a chemokine inducing the recruitment of monocytes and macrophages into inflamed tissues as well as the M2 polarization of macrophages (Sierra-Filardi et al., 2014) was significantly induced by BMP6 and partially reduced by BYL719 (**Figure 7E**). In macrophages, BYL719 reduced the expression of pan-macrophage markers *Adgre1* (F4/80), *Cd68* and the M1 macrophage marker *Cd80* (**Figure 7F**). Conversely, BYL719 increased the expression of the M2 macrophage marker *Il10*, regardless of the presence or absence of BMP6. In the human mast cell line HMC1, BYL719 inhibited the expression of the pro-inflammatory cytokine *IL6*, while the expression of *TGF β 1*, *TNF α* , and activin A (*INHBA*) remained unchanged upon the addition of either BMP6 or BYL719 (**Figure 7G**). Altogether, these data suggest that PI3K α inhibition reduces the expression of certain pro-inflammatory cytokines and might skew macrophages towards M2. Moreover, the reduction on activin A expression by BYL719 in inflammatory cells and mesenchymal progenitors could be relevant for HO in FOP.

PI3K α inhibition reduces the hyperinflammatory response in FOP

To confirm whether BYL719 can modulate excessive inflammation at the injured sites, we examined its impact on monocytes, macrophages and mast cells *in vivo*. These cell populations are known to increase in number two to four days after injury (Convente et al., 2018; Sorkin et al., 2020), which coincides with the effects observed following early administration of BYL719 (Convente et al., 2018; Sorkin et al., 2020). Hindlimb muscle samples were collected at two, four, nine, sixteen and twenty-three days post-induction of HO with Cre viruses and cardiotoxin in *ACVR1^{Q207Dfl/fl}* mice (**Figure 8A**). The group of treated animals received BYL719 intermittent treatment starting one day after injury. After sixteen days, structures of mineralized bone became detectable in vehicle-treated mice (Figure 8-Figure Supplement 1A and B). The number of F4/80-positive monocytes/macrophages was high two days after injury and remained elevated in the HO lesions throughout the ossification process (**Figure 8B and C**). BYL719-treated animals showed an almost 50% reduction in the number of F4/80 positive cells at day two post-injury and maintained the relative difference with vehicle-treated mice afterwards. Consistent with previous studies, we observed an increased number of mast cells at injury sites. Their numbers peaked at four days and only dropped twenty-three days after injury in vehicle-treated mice (**Figure 8D and E**). In BYL719-treated mice, mast cells were increased up to day 9, but their number was completely normalized to preinjury levels already at day 16 (**Figure 8D and E**). This observation correlates with the complete muscle regeneration observed in BYL719-treated mice compared to untreated mice, which are actively developing ossifications at that time (Figure 8-Figure Supplement 2). These results confirm a functionally relevant dual role of PI3K α inhibition, by preventing the differentiation of osteochondrogenic progenitor cells and also by reducing the local inflammatory response.




Figure 7.

(A, B, C) Proliferation assays of THP1 (A), RAW264.7 (B), and HMC-1 (C) cells. Cells were cultured for 6 days and in control conditions, with BMP6 (2nM) and/or BYL719 (2 or 10μM). Areas under the proliferation curves were compared by one-way ANOVA with Dunnett's multiple comparisons test against the control. An asterisk (*) refers to significant differences between control and single treatments (BMP6 or BYL719). A hash sign (#) refers to significant differences between control and combined treatments. Data shown as mean ± SD (n = 4 per group). ** or ## P<0.01, *** or ### P<0.001. (D) Migration assay of THP1 cells with control conditions, or with FBS or BMP6 as chemotactic agents with or without BYL719 treatment. Data are shown as mean ± SD (n = 4 per group). ***P<0.001, two-way ANOVA with Tukey's multiple comparisons test. (E, F, G) Gene expression assays of THP1 (E), RAW264.7 (F) and HMC-1 (G) cells. Cells were treated for 48 hours with BYL719 (2μM) and/or BMP6 (2nM). Data are shown as mean ± SD (n = 4 per group). *P<0.05, **P<0.01, ***P<0.001, one-way ANOVA with Dunnett's multiple comparisons test, significance shown between control group and other groups.




Figure 8.

(A) Heterotopic ossification was induced at P7 injecting adenovirus-Cre (Ad.Cre) and cardiotoxin in the mice hindlimb. Starting at P8, mice were treated with DMSO or BYL719 (25 mg/kg) as indicated by grey dots. The final time-point changed between experimental groups, stopping at P9 (2 days post-HO-induction), P11 (4 days post-HO-induction), P16 (9 days post-HO-induction), P23 (16 days post-HO-induction), and P30 (23 days post-HO-induction). (B) The quantification of F4/80-positive monocytes/macrophages per field view from immunohistochemistry staining, with 20x amplification as shown in the representative images depicted in [Figure 8C](#). Five images were randomly acquired per mice, from 2 mice per group, for a total of 10 quantified images per group. F4/80 positive cells were detected as cells with dark brown staining. Data are shown as mean $\pm$ SD. * P <0.05, ** P <0.01, *** P <0.001, two-way ANOVA with Sidak's multiple comparisons test, comparing both groups on each day (n =10). (C) Representative images from the quantified immunohistochemistry staining for F4/80 to detect monocytes/macrophages. Representative images are shown for each final time-point at day 2, 4, 9, 16, and 23 post-HO-induction. All images were obtained at 20x, with a representative scale bar at 100 μ m. (D) Quantification of CEM-positive mast cells per field view from CEM staining, with 20x amplification as shown in the representative images depicted in [Figure 8E](#). Three images were obtained per mice, from 4 mice per group, for a total of 12 quantified images per group. Mast cells were detected as cells with bright blue staining. Data are shown as mean $\pm$ SD. *** P <0.001, two-way ANOVA with Sidak's multiple comparisons test, comparing both groups in each day (n =12). (E) Representative images from the quantified C.E.M. staining to detect mast cells, highlighted with black arrows. Representative images are shown for each final time-point at day 2, 4, 9, 16, and 23 post-HO-induction. All the images were obtained at 20x magnification, with a representative scale bar at 100 μ m.

We previously showed that pharmacological PI3K α inhibition using BYL719 has the potential to suppress HO by increasing SMAD1/5 degradation, reducing transcriptional responsiveness to BMPs, and blocking non-canonical responses such as the activation of AKT/mTOR (Valer et al., 2019a [DOI](#)). In this manuscript, we confirmed these results and further expanded the insights into the molecular and cellular mechanisms responsible for the therapeutic effect of PI3K inhibition. Here, we optimized the therapeutic window for BYL719 administration and found that the delayed administration of BYL719 effectively prevents HO. Moreover, *in vivo* simultaneous genetic inhibition of *p110 α* and the activation of mutated *ACVR1* at injury sites partially prevents HO, and we demonstrate that genetic or pharmacological inhibition of PI3K α inhibits osteochondroprogenitor specification. We determined that BYL719 does not inhibit recombinant *ACVR1*^{R206H} kinase activity *in vitro* and does not behave as an ATP competitor inhibitor for none of the TGF- β kinase receptors. Finally, the administration of BYL719 prevented an exacerbated inflammatory response *in vivo*, possibly due to specific effects observed on immune cell populations.

While FAPs play a key role in bone formation in different types of HO, inflammation is also closely associated with all types of HO (Barruet et al., 2018 [DOI](#); de Ruiter et al., 2021; Matsuo et al., 2019 [DOI](#)). For instance, the depletion of mast cells and macrophages has been shown to reduce bone volume in FOP mouse models (Convente et al., 2018 [DOI](#)). Moreover, Activin A, TGF- β and other cytokines secreted by FAPs, monocytes, macrophages and mast cells are essential for FOP and non-genetic HO (Alessi Wolken et al., 2017 [DOI](#); Hatsell et al., 2015 [DOI](#); Lees-Shepard et al., 2018 [DOI](#); Patel et al., 2022 [DOI](#); Sorkin et al., 2020 [DOI](#); Upadhyay et al., 2017 [DOI](#)). In addition, *ACVR1* antibodies can activate *ACVR1*^{R206H} even in the absence of Activin A, but muscular trauma is still required to induce HO (Aykul et al., 2022 [DOI](#); Lees-Shepard et al., 2022 [DOI](#)), indicating the involvement of additional immune factors. Our study finds that BYL719, a PI3K α inhibitor, reduces the expression of certain pro-inflammatory cytokines, effectively blocks the proliferation of monocytes, macrophages and mast cells and reduces the migratory potential of monocytes. This likely contributes to a decreased number of monocytes and macrophages at injury sites and throughout the *in vivo* ossification process. It is known that excessive pro-inflammatory cytokine expression, including activin A, by monocytes and macrophages is observed in all types of HO and is induced by *ACVR1*^{R206H} (Matsuo et al., 2021 [DOI](#)). Non-canonical signaling pathways (mTOR, p38, TAK1, NF- κ B) are also affected by *ACVR1*^{R206H} in immune cell types (Barruet et al., 2018 [DOI](#); Hwang et al., 2022 [DOI](#)). M1-type macrophages initiate acute inflammatory responses but later transition to M2-type macrophages associated with anti-inflammatory and reparative functions. Therefore, modulating macrophage phenotype towards regeneration may dampen HO (Sorkin et al., 2020 [DOI](#)). BYL719 reduces monocyte/macrophage numbers, migration, and pro-inflammatory cytokine expression, potentially altering their polarization. It also reduces activin A expression and promotes a shift towards the M2 phenotype. BYL719 can help to reduce the hyper-inflammatory state in HO, inhibit FAP expansion, and favor myogenic regeneration of muscle tissue (Stanley et al., 2022 [DOI](#)).

Our observation that the delayed administration of BYL719 is still effective has pathophysiological and therapeutic derivatives. Muscle injury induces sequential changes in HO progression: early immune cell infiltration (days 1-3), a fibroproliferative and late inflammatory phase, with an expansion of FAPs and a spectrum of monocytes, macrophages and mast cells as highly secretory cells (days 3-7), followed by chondrogenesis (day 7-14) and osteogenesis (days 14-23) (Hwang et al., 2022 [DOI](#)). Starting BYL719 administration 3 days post-injury is still fully therapeutically effective and even partially at 7 days post-injury. In addition, whereas untreated mice developed cartilage and bony lesions by days 9 and 16 post-injury, respectively, BYL719-treated animals fully regenerated muscle by day 16 post-injury. Altogether, this evidence suggests that the major effects of BYL719 occur between 3 and 16 days post-injury. We found that BYL719 completely blocks chondrogenesis in cultured hMSCs irrespective of the differentiation duration. In addition, in the histological analyses of these BYL719-treated mice that do not develop HO, we cannot find any sign

of cartilage formation on either the 9th, 16th or 23rd days post-injury, which confirms that BYL719 is also able to block chondrogenesis in mice *in vivo*. This suggests that intervention with BYL719 over a temporal window after a HO flare might be sufficient to inhibit endochondral ossification. In addition, although surgical resection is not recommended in FOP patients and has a risk of recurrence in resected non-genetic HO, it could be envisaged using as preventive therapy in subjects undergoing surgeries to remove ectopic bone.

Treatment options for HO and FOP are currently limited, primarily consisting of anti-inflammatories such as corticosteroids and NSAIDs. Promising pharmacological treatments have progressed to II/III clinical trials. Palovarotene, a retinoic acid receptor- γ agonist, has been approved in Canada for patients over 10 years old (NCT03312634). However, a previous clinical trial with palovarotene was paused for children due to concerns about premature growth plate closure, which is debated in FOP mice (Chakkalakal et al., 2016 [\[1\]](#); Rosen et al., 2018 [\[2\]](#)). Antibodies targeting activin A (Garetosmab) showed effectiveness in a phase II trial, but serious adverse effects led to a temporary hold on the trial (NCT03188666). The outcome of a clinical trial with Rapamycin has not been publicly disclosed and a case report indicated limited benefits in classical FOP patients (UMIN000028429) (Kaplan et al., 2018 [\[3\]](#)). BYL719 (Alpelisib/Piqray®) was recently licensed in combination with hormone therapy for a PIK3CA-mutated breast cancer (Cardoso et al., 2020 [\[4\]](#)). BYL719 has been shown to be also clinically effective in adult patients with PIK3CA-related overgrowth syndrome (PROS) and children under 1 year of age (Madsen and Semple, 2022 [\[5\]](#); Morin et al., 2022 [\[6\]](#)). In both scenarios, BYL719 had no major adverse effect when patients were treated for more than one year (in adults at a dose of 300mg daily for oncological or PROS purposes and at a dose of 25 mg daily in infants with PROS) (Morin et al., 2022 [\[6\]](#)). Thus, implementing BYL719 for treatment of HO, at least during a restricted temporal window (i.e. surgery to remove ectopic bone in HO, or during flare-ups in FOP individuals), may be a valid therapeutic option for FOP patients.

Materials and Methods

Murine bone marrow mesenchymal stem cells isolation

Murine bone marrow-derived mesenchymal stem cells (BM-MSCs) were isolated from 6-8 weeks-old C57BL6/J mice for the floxed p110 α allele (Beatriz Gámez et al., 2016 [\[7\]](#)) as previously described (Soleimani and Nadri, 2009 [\[8\]](#)). Mice were euthanized and femurs were dissected and stored in DMEM with 100 U/ml penicillin/streptomycin (P/S). Soft tissues were cleaned and femur ends were cut under sterile conditions. Bone marrow was flushed with media using a 27-gauge needle. The resulting cell suspension was filtered through a 70- μ m cell strainer and seeded in a 100-mm cell culture dish. Non-adherent cells were discarded after 3 hours. Media was slowly replaced every 12 hours for up to 72 hours. Then, media was replaced every 2 days until the culture reached 70% confluence. Then, cells were lifted by incubation with 0.25% trypsin/0.02% EDTA for 5 minutes at room temperature. Lifted cells were cultured and expanded.

Production of human fetal mesenchymal stem cells (hMSCs) expressing ACVR1

Bone marrow-derived hMSCs were transduced by lentiviral delivery encoding ACVR1^{WT}-nanoluc or ACVR1^{R206H}-nanoluc. The nanoluc fusion inserts (Promega, NV2341/NV2381) were subcloned into a PLV-CMV-IRES vector. First, an eGFP was amplified with PCR containing PstI, SalI and XbaI restriction sites and subcloned in the PLV using PstI and XbaI. Next, the ACVR1-nanoluc inserts were subcloned in the PLV by restriction of PstI and SalI, exchanging the eGFP. Subsequently, lentiviral particles were produced using HEK293t cells and cells were transduced.

Cell culture

Murine BM-MSCs were cultured in DMEM supplemented with 10% FBS, 2 mM glutamine, 1 mM sodium pyruvate, and 100 U/ml P/S and incubated at 37°C with 5% CO₂. The human monocyte cell line THP1 was cultured in RPMI-1640 supplemented with 10% FBS, 2 mM glutamine, 1 mM sodium pyruvate, 100 U/ml P/S, 0.1 mM NEAA, 10 mM HEPES and 50 μM 2-mercaptoethanol. Mouse macrophage cell line RAW264.7 was cultured in DMEM supplemented with 10% FBS, 2 mM glutamine, 1 mM sodium pyruvate and 100 U/ml P/S. The human mast cell line HMC1 was cultured in IMDM supplemented with 10% FBS, 2 mM glutamine, 1 mM sodium pyruvate and 100 U/ml P/S. The hMSCs were cultured in αMEM supplemented with 10% FBS, 1 ng/ml human bFGF and 100 U/ml P/S (Growth Medium; GM).

Proliferation and chemotactic migration assays

For proliferation assays, 3000 cells/well were seeded in 12-well culture plates in correspondent culture medium with 2 nM BMP6 and/or 2 μM or 10 μM BYL719 (Chemietek). Photos of several fields of each well were taken at 24, 48, 72 and 144 hours using a Leica DM-IRB inverted microscope. Cell counting was performed through ImageJ software.

The chemotactic migration assay of THP1 was performed through the InnoCyte Monocyte Migration Assay (Merck Millipore). Cells were serum-starved for 3h prior to the experiment and, when needed, treated with 2 μM BYL719 for 30 minutes before the experiment started. Then, 10⁵ cells/100 μL were added to each upper chamber. Lower chambers were charged with 10% FBS-supplemented RPMI-1640 (positive control), FBS-depleted RPMI-1640 (negative control) or 2 nM BMP6 (R&D), respectively. Cells migrated for 2h and were then stained with calcein-AM and pipetted into a 96-well conical bottom plate (Sigma-Aldrich). A plate reader was used to measure fluorescence from the top of the plate at an excitation wavelength of 485 nm and an emission wavelength of 520 nm. Unless otherwise stated, BYL719 at 2 or 10 μM (Chemietek) and 2 nM BMP6 (R&D) was used for 6 days (proliferation) or 4 hours (migration) in HMC1, THP1 and RAW264.7 cells.

Retroviral transduction and gene expression analysis

BM-MSCs were infected with mock virus (pMSCV-GFP) or with viruses expressing wild type *Acvr1* (WT), *Acvr1*^{R206H} (RH) and/or pMSCV-Cre. Plasmids with WT and RH *Acvr1* forms were kindly provided by Dr. Petra Seemann and were subcloned into pMSCV vector. *Acvr1* and p110α expression levels were analyzed by qRT-PCR. Cells were treated with BYL719 (2 μM) and/or activin A (2 nM) for 48 hours in complete media without FBS. THP1, RAW264.7, and HMC-1 cells lines, cells were treated in the media indicated for each cell line with BYL719 (2 μM) and/or BMP6 (2 nM) for 48 hours.

Total RNA for RT-PCR was extracted from all cellular models using TRIsure reagent. At least 2 μg of purified RNA was reverse-transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). Quantitative PCRs were carried out on ABI Prism 7900 HT Fast Real-Time PCR System with Taqman 5'-nuclease probe method and SensiFAST Probe Hi-ROX Mix. All transcripts were normalized using *Tbp* as an endogenous control.

NanoBRET target engagement assays

COS-1 cells were cultured in DMEM (Gibco, 11965092) supplemented with 10% FBS and 100 U/ml P/S. These cells were transfected in 70% confluent 6-wells TC-treated plates with 2 μg TGFβ receptor-Nanoluciferase constructs (constructs were provided by Promega and Promega R&D) using a 1 μg DNA: 2 μl PEI (1 mg/ml) ratio. The transfected cells were reseeded in white 384-wells TC-treated assay plates one day before the nanoBRET readout in DMEM supplemented with 1% FBS in a quantity of 15*10³ cells in 40 μl per well. Two hours before measurement, the dedicated tracer (Promega) and the test compounds were incubated to the cells at 37°C. Immediately after

intracellular TE Nano-Glo Substrate/Inhibitor (Promega, N2160) addition, the wells were measured by 450-80BP and 620-10BP filters using the ClarioSTAR (BMG Labtech). Subsequent nanoBRET ratios were measured by the formula: acceptor emission (620-10 nm)/donor emission (450-80 nm) * 1000 (milliBRET units, mBU).

Chondrogenic progenitor differentiation

The hMSCs were cultured in α MEM supplemented with 10% FBS, 1 ng/ml human bFGF and 100 U/ml P/S (Growth Medium; GM). Chondrogenic differentiation medium (DM) consisted of DMEM supplemented with 10 μ l/ml Insulin-Transferrin-Selenium 200 μ M Ascorbic Acid, 0.1 μ M Dexamethasone, 350 μ M L-proline, 1 mM Sodium Pyruvate, 1.25 μ g/ml BSA and 100 U/ml P/S. During differentiation, the medium was refreshed twice a week for the duration of the experiment. For the ALP staining and ALP activity experiments, the hMSC-ACVR1 lines were seeded at 2×10^4 cells per well in 48-wells plates and differentiated for 7 and 11 days, respectively. For the Alcian Blue experiments, 3D micromass cultures were made by carefully seeding 3×10^5 cells per 10 μ l droplets in a 24-wells plate and incubated for 2 hours at 37°C prior to GM addition. Chondrogenic progenitor differentiation was started 1 day after micromass seeding and continued for 3 weeks. For the analysis of early chondroprogenitor specification, the hMSC-ACVR1 lines were differentiated for only 3 days where after RT-qPCR analysis was performed.

Construct	Cat. #	Tracer (conc.)	Cat. #	Compounds (conc.)
<i>ALK1-nluc</i>	NV2391	K-11 (0.31 μ M)	N2652	BYL719 (1 and 10 μ M) LDN193189 (0,5 μ M)
<i>ALK2-nluc</i>	NV2341	K-11 (0.16 μ M)	N2652	BYL719 (1 and 10 μ M) LDN193189 (0,5 μ M)
<i>ALK2 R206H-nluc</i>	NV2381	K-11 (0.16 μ M)	N2652	BYL719 (1 and 10 μ M) LDN193189 (0,5 μ M)
<i>ALK3-nluc</i>	NV2471	K-11 (0.63 μ M)	N2652	BYL719 (1 and 10 μ M) LDN193189 (0,5 μ M)
<i>ALK4-nluc</i>	NV1021	K-5 (2 μ M)	N2482	BYL719 (1 and 10 μ M) SB431542 (10 μ M)
<i>ALK5-nluc</i>	Promega R&D	K-14 (0.5 μ M)	Promega R&D	BYL719 (1 and 10 μ M) SB431542 (10 μ M)
<i>ACVR2A-nluc</i>	Promega R&D	PBI6948 (1 μ M)	Promega R&D	BYL719 (1 and 10 μ M) ML347 (10 μ M)
<i>ACVR2B-nluc</i>	Promega R&D	PBI6948 (1 μ M)	Promega R&D	BYL719 (1 and 10 μ M) ML347 (10 μ M)
<i>TGFBR2-nluc</i>	Promega R&D	K-11 (0.32 μ M)	N2652	BYL719 (1 and 10 μ M) ML347 (10 μ M)
<i>BMPR2-nluc</i>	Promega R&D	PBI7394 (0.5 μ M)	Promega R&D	BYL719 (1 and 10 μ M) ML347 (10 μ M)

Alkaline phosphatase (ALP) staining, ALP activity assay and Alcian Blue staining

For the ALP staining, the progenitors were washed with PBS, fixated using 3.7% formalin for 5 minutes at RT, washed twice with PBS and stained in ALP solution containing 2 mg Naphthol AS-MX, 6 mg Fast Blue, 5 ml 0.2M Tris (pH 8.9), 100 μ l MgSO_4 and dH_2O up to 10 ml. Images were acquired using the Leica DMi8 with a 10x magnification. Densitometry analysis of the whole well (from total plate scan) was performed using ImageJ (v1.53t).

ALP lysates were obtained by washing twice with PBS, freezing at -80°C for 1 hour, and lysing on ice for 1 hour using 100 μl ALP buffer (100 μM MgCl_2 , 10 μM ZnCl_2 , 10 mM Glycine (pH 10.5)) plus 0.1% Triton X-100 per well. ALP activity was quantified by adding 20 μl lysate and 80 μl 6 mM PNPP in ALP buffer to a clear 96-well plate incubated at RT until the samples turned yellow. Next, the samples were measured at an absorbance of 405 nm.

The micromass pellets were stained with Alcian Blue as described before (Sánchez-Duffhues et al., 2019 [DOI](#)). Images were acquired using the Leica DMI8 with 5x magnification. To quantify the chondroprogenitor specification, the Alcian Blue was solubilized by 250 μl of 6M guanidine hydrochloride and absorbance was measured at 595 nm.

Bulk RNA sequencing and analyses

The hMSCs-ACVR1^{R206H} were seeded at 100% confluency and serum starved overnight with DMEM without supplements. Afterwards, the cells were pre-treated with 1 μM of BYL719 in starvation medium for 30 minutes and stimulated with 50 ng/ml Activin A for 1 hour. RNA was isolated using the ReliaPrep RNA Miniprep Systems (Promega) and sequenced using the Illumina NovaSeq 6000 (Illumina) platform. The reads were mapped using the Hisat2 (v2.0.5) package and the total and normalized counts were measured using featureCounts (v1.5.0-p3). Differential expression analysis was performed using the DESeq2 R package (v1.20.0), in which the p-values were adjusted following the Benjamini and Hochberg approach. Genes with an adjusted p-value < 0.05 were considered differentially expressed. Subsequent Gene Ontology analysis was performed using the clusterProfiler R package and the GO terms were considered significant if the p-values < 0.05. Gene Set Enrichment Analysis was performed using the GSEA analysis tool (*gsea-msigdb.org* (<https://gsea-msigdb.org/>) [DOI](#)). Heatmaps were generated using heatmap2 function within the R gplots package in Galaxy (*usegalaxy.eu* (<https://usegalaxy.eu/>) [DOI](#)). The datasets are publicly available (see GSE237512).

Western blot

Cells were lysed using RIPA lysis buffer and quantified using the Pierce BCA Protein assay (Thermo) following standard protocols. The protein samples were loaded using β -ME containing sample buffer and run using SDS-PAGE and transferred to PVDF membranes. Primary antibodies against SMAD1 (Cell Signaling, 6944), p-SMAD1/5 (Cell Signaling, 9516), and vinculin (Sigma, V9131) were used at 1:1000 dilution in 5% BSA in TBST. Binding was detected with HRP-conjugated secondary antibodies and visualized by Brightfield ECL (Thomas Scientific) on the ChemiDoc (BioRad).

In vitro recombinant kinase activity assay

Fifty ng of recombinant ACVR1^{R206H} (ab167922, Abcam) was diluted in 30 μl of 25mM Tris/HCl (pH7.5), 10mM MgCl_2 , 10mM MnCl_2 and 2mM DTT. The reaction was started by the addition of 5 μM ATP-Mg, 0.05 $\mu\text{Ci}/\mu\text{l}$ ATP [γ -³²P] and 0.5mg/ml of dephosphorylated casein (Sigma) and performed for 30 min at 30°C. The reaction was stopped and samples were processed by SDS-PAGE. After drying the gel, phosphorylation was determined by autoradiography.

Heterotopic ossification mouse model

To study heterotopic ossification *in vivo*, we used the Cre-inducible constitutively active ACVR1^{Q207D} (CAG-Z-EGFP-caALK2) mouse model as previously described (Fukuda et al., 2006 [DOI](#); Shimono et al., 2011 [DOI](#); Yu et al., 2008 [DOI](#)). Mice ACVR1Q207D^{fl/fl} p110 α ^{fl/fl} were obtained by crossing the detailed homozygous ACVR1Q207D^{fl/fl} with the p110 α ^{fl/fl} homozygous mutant mice carrying loxP sites flanking exons 18 and 19 of the p110 α alleles (Beatriz Gámez et al., 2016 [DOI](#)). Heterozygous mice were crossed until ACVR1Q207D^{fl/fl} p110 α ^{fl/fl} was obtained.

To induce heterotopic ossification in P7 ACVR1^{Q207D} mice, 1×10^8 pfu of adenovirus-Cre (Ad-CMV-Cre, Viral Vector Production Unit, UAB) and 0.3 μ g of cardiotoxin in 10 μ l 0.9% NaCl volume were injected into the left hindlimb. Control groups had the same procedure without Ad-Cre in the injection. On P8, mice started receiving either placebo (intraperitoneal administration (i.p.) of DMSO) or BYL719 (i.p. of 25 mg/kg), both diluted in 0.5% carboxymethylcellulose sodium. Mice were housed under controlled conditions (12-hour light/12-hour dark cycle, 21°C, 55% humidity) and fed *ad libitum* with water and a 14% protein diet (Teklad2014, Envigo). Mice were regularly weighted over the whole period. Heterotopic ossification induction was performed blinded to mouse and group identity. All procedures were approved by the Ethics Committee for Animal Experimentation of the Generalitat de Catalunya.

Micro-computed tomography analysis

The caudal half of mice was collected and fixed in 4% paraformaldehyde (PFA) for 48 hours at 4°C. Samples were conserved in PBS and high-resolution images were acquired using a computerized microtomography imaging system (Skyscan 1076, Bruker microCT), in accordance with the recommendations of the American Society of Bone and Mineral Research (ASBMR). Samples were scanned in air at 50 kV and 200 μ A with an exposure time of 800 ms, using a 1-mm aluminum filter and an isotropic voxel size of 9 μ m. Two-dimensional images were acquired every 1° for 180° rotation and subsequently reconstructed, analyzed for bone parameters, and visualized by NRecon v1.6, CT-Analyser v1.13 and CTVox v3.3 programs (Bruker), respectively. For heterotopic ossification, manual VOIs comprising heterotopic ossifications were employed and a binary threshold was established at 25-255.

Histology

Whole legs were fixed in 4% PFA for 48 hours at 4°C, decalcified in 16% EDTA pH 7.4 for 6 weeks, and embedded in paraffin. Five μ m sections were cut and stained with Fast Green/Safranin O, or Masson's Trichrome. Images were obtained with brightfield Eclipse E800 (Nikon). To detect monocyte/macrophage cells, paraffined slides were immunostained with Anti-F4/80 antibody [SP115] (ab111101). Briefly, after deparaffination and rehydration, an antigen retrieval step with citrate buffer at pH6 in a decloaking chamber was applied to the slides. Slides were cooled down and rinsed with 1x TBS and endogenous peroxidases were blocked with 70% methanol, 28% distilled water, and 2% hydrogen peroxide for 5 minutes at room temperature. After a blocking step, F4/80 primary antibody incubation at concentration 1:100 overnight at 4 °C was applied. Envision Dual Link conjugated with HRP was applied for 40 minutes at room temperature, and detection was performed with diaminobenzidine (DAB) for 3 minutes. Nuclear staining with Lilly's hematoxylin was applied to the slides for 30 seconds. Images of stained slides were obtained with brightfield Eclipse E800 (Nikon). Images were randomly acquired with 20x magnification from the region of interest (injected muscle, with or without visible heterotopic ossification). Five images were obtained per mice, from 2 mice per group, for a total of 10 quantified images per group. F4/80 positive cells were detected as cells with dark brown staining. To detect mast cells, paraffined slides were stained with C.E.M. staining using the Eosinophil-Mast cell staining kit (ab150665, Abcam). The recommended staining procedure was applied to the deparaffinized sections. Images of stained slides were obtained with brightfield Eclipse E800 (Nikon). Images were randomly acquired with 20x magnification from the region of interest (injected muscle, with or without visible heterotopic ossification). Three images were obtained per mice, from 4 mice per group, for a total of 12 quantified images per group. Mast cells were detected as cells with bright blue staining.

Statistical analysis

Unless stated otherwise in each figure legend, the results were expressed as mean $\pm$ SD. The median was shown in heterotopic ossification datasets where non-parametric tests were applied. Each figure legend has detailed information explaining the statistical test used to compare

between groups, the compared groups, the explanation of the symbols used to show significance, usually $*P<0.05$, $**P<0.01$, $***P<0.001$, and the sample size of the experiment. Heterotopic ossification, evaluation, and quantification was performed blinded to mouse and group identity. Microscopy and histology representative images were selected from the total quantified images. Microscopy and histology quantified fields were randomly acquired from the plate or the region of interest. Statistical tests were performed on GraphPad Prism 9.5.

Acknowledgements

We thank E. Adanero, E. Castaño for technical assistance. We acknowledge the support from Promega R&D. Alexandre Deber is a recipient of a F.P.I. fellowship from the Ministry of Science and Innovation (MCIN/AEI/ 10.13039/501100011033). Carolina Pimenta-Lopes is the recipient of a F.P.U. fellowship from the Spanish Ministry of Education. This research was supported by grants PID2022-141212OA-I00, PDC2021-121776-I00 and PID2020-117278GB-I00 from MCIN/AEI/10.13039/501100011033, co-funded by FEDER ‘Una manera de hacer Europa’ or ‘NextGenerationEU’/PRTR; a grant 202038-30 from La Marató de TV3 and grants from IFOPA (ACT for FOP) and FOPItalia. MW and MJG are sponsored by the Netherlands Cardiovascular Research Initiative (the Dutch Heart Foundation, Dutch Federation of University Medical Centers, the Netherlands Organization for Health Research and Development, and the Royal Netherlands Academy of Sciences), PHAEDRA-IMPACT (DCVA) and DOLPHIN-GENESIS (CVON). GSD is also sponsored by the Spanish Ministry of Science through the Ramón y Cajal grant RYC2021-030866-I and the BHF-DZHK-DHF, 2022/23 award PROMETHEUS.

Author contributions

J.A.V., A.D., M.W., M.J.G., J.L.R., G.S-D. and F.V. designed the study, and analyzed and interpreted the results. J.A.V., A.D., M.W. and C.P-L. performed the experiments. J.A.V., A.D., G.S-D. and F.V. wrote the manuscript. All authors contributed to the manuscript. G.S-D. and F.V. supervised the project.

Competing interests

The authors declare no conflicts of interest.

References

- Agarwal S, Loder SJ, Breuler C, Li J, Cholok D, Brownley C, Peterson J, Hsieh HH, Drake J, Ranganathan K, Niknafs YS, Xiao W, Li S, Kumar R, Tompkins R, Longaker MT, Davis TA, Yu PB, Mishina Y, Levi B (2017) **Strategic Targeting of Multiple BMP Receptors Prevents Trauma-Induced Heterotopic Ossification** *Molecular Therapy* **25**:1974–1987 <https://doi.org/10.1016/j.ymthe.2017.01.008>
- Agnew C, Ayaz P, Kashima R, Loving HS, Ghatpande P, Kung JE, Underbakke ES, Shan Y, Shaw DE, Hata A, Jura N (2021) **Structural basis for ALK2/BMP2 receptor complex signaling through kinase domain oligomerization** *Nat Commun* **12** <https://doi.org/10.1038/s41467-021-25248-5>
- Alessi Wolken DM, Idone V, Hatsell SJ, Yu PB, Economides AN (2017) **The obligatory role of Activin A in the formation of heterotopic bone in Fibrodysplasia Ossificans Progressiva** *Bone* **109**:210–217 <https://doi.org/10.1016/j.bone.2017.06.011>
- Aykul S, Huang L, Wang L, Das NM, Reisman S, Ray Y, Zhang Q, Rothman N, Nannuru KC, Kamat V, Brydges S, Troncione L, Johnsen L, Yu PB, Fazio S, Lees-Shepard J, Schutz K, Murphy AJ, Economides AN, Idone V, Hatsell SJ (2022) **Anti-ACVR1 antibodies exacerbate heterotopic ossification in fibrodysplasia ossificans progressiva (FOP) by activating FOP-mutant ACVR1** *Journal of Clinical Investigation* **132** <https://doi.org/10.1172/JCI153792>
- Barruet E, Morales BM, Cain CJ, Ton AN, Wentworth KL, Chan T V., Moody TA, Haks MC, Ottenhoff TH, Hellman J, Nakamura MC, Hsiao EC (2018) **NF- κ B/MAPK activation underlies ACVR1-mediated inflammation in human heterotopic ossification** *JCI Insight* **3** <https://doi.org/10.1172/jci.insight.122958>
- Bravenboer N, Micha D, Triffitt JT, Bullock AN, Ravazollo R, Boccardi R, Di Rocco M, Netelenbos JC, Ten Dijke P, Sánchez-Duffhues G, Kaplan FS, Shore EM, Pignolo RJ, Seemann P, Ventura F, Beaujat G, Eekhoff EMW, Pals G. (2015) **Clinical Utility Gene Card for: Fibrodysplasia ossificans progressiva** *European Journal of Human Genetics* **23** <https://doi.org/10.1038/ejhg.2014.274>
- Cardoso F, Paluch-Shimon S, Senkus E, Curigliano G, Aapro MS, André F, Barrios CH, Bergh J, Bhattacharyya GS, Biganzoli L, Boyle F, Cardoso M-J, Carey LA, Cortés J, el Saghir NS, Elzayat M, Eniu A, Fallowfield L, Francis PA, Gelmon K, Gligorov J, Haidinger R, Harbeck N, Hu X, Kaufman B, Kaur R, Kiely BE, Kim S-B, Lin NU, Mertz SA, Neciosup S, Offersen B v, Ohno S, Pagani O, Prat A, Penault-Llorca F, Rugo HS, Sledge GW, Thomssen C, Vorobiof DA, Wiseman T, Xu B, Norton L, Costa A, Winer EP. (2020) **5th ESO-ESMO international consensus guidelines for advanced breast cancer (ABC 5)** *Ann Oncol* **31**:1623–1649 <https://doi.org/10.1016/j.annonc.2020.09.010>
- Chaikuad A, Alfano I, Kerr G, Sanvitale CE, Boergermann JH, Triffitt JT, Von Delft F, Knapp S, Knaus P, Bullock AN. (2012) **Structure of the bone morphogenetic protein receptor ALK2 and implications for fibrodysplasia ossificans progressiva** *Journal of Biological Chemistry* **287**:36990–36998 <https://doi.org/10.1074/jbc.M112.365932>

Chakkalakal SA, Uchibe K, Convente MR, Zhang D, Economides AN, Kaplan FS, Pacifici M, Iwamoto M, Shore EM (2016) **Palovarotene Inhibits Heterotopic Ossification and Maintains Limb Mobility and Growth in Mice With the Human ACVR1(R206H) Fibrodysplasia Ossificans Progressiva (FOP) Mutation** *J Bone Miner Res* **31**:1666–75 <https://doi.org/10.1002/jbmr.2820>

Chakkalakal SA, Zhang D, Culbert AL, Convente MR, Caron RJ, Wright AC, Maidment A DA, Kaplan FS, Shore EM (2012) **An Acvr1 R206H knock-in mouse has fibrodysplasia ossificans progressiva** *Journal of Bone and Mineral Research* <https://doi.org/10.1002/jbmr.1637>

Convente MR, Chakkalakal SA, Yang E, Caron RJ, Zhang D, Kambayashi T, Kaplan FS, Shore EM. (2018) **Depletion of Mast Cells and Macrophages Impairs Heterotopic Ossification in an Acvr1R206H Mouse Model of Fibrodysplasia Ossificans Progressiva** *Journal of Bone and Mineral Research* **33** <https://doi.org/10.1002/jbmr.3304>

Dey D, Bagarova J, Hatsell SJ, Armstrong KA, Huang L, Ermann J, Vonner AJ, Shen Y, Mohedas AH, Lee A, Eekhoff EMW, Van Schie A, Demay MB, Keller C, Wagers AJ, Economides AN, Yu PB. (2016) **Two tissue-resident progenitor lineages drive distinct phenotypes of heterotopic ossification** *Sci Transl Med* **8** <https://doi.org/10.1126/scitranslmed.aaf1090>

Eisner C, Cummings M, Johnston G, Tung LW, Groppa E, Chang C, Rossi FMV (2020) **Murine Tissue-Resident PDGFR α + Fibro-Adipogenic Progenitors Spontaneously Acquire Osteogenic Phenotype in an Altered Inflammatory Environment** *Journal of Bone and Mineral Research* **35**:1525–1534 <https://doi.org/10.1002/jbmr.4020>

Ford-Hutchinson AF, Ali Z, Lines SE, Hallgrímsson B, Boyd SK, Jirik FR (2007) **Inactivation of Pten in osteo-chondroprogenitor cells leads to epiphyseal growth plate abnormalities and skeletal overgrowth** *Journal of Bone and Mineral Research* <https://doi.org/10.1359/jbmr.070420>

Fujita T, Azuma Y, Fukuyama R, Hattori Y, Yoshida C, Koida M, Ogita K, Komori T (2004) **Runx2 induces osteoblast and chondrocyte differentiation and enhances their migration by coupling with PI3K-Akt signaling** *Journal of Cell Biology* <https://doi.org/10.1083/jcb.200401138>

Fukuda T, Scott G, Komatsu Y, Araya R, Kawano M, Ray MK, Yamada M, Mishina Y (2006) **Generation of a mouse with conditionally activated signaling through the BMP receptor, ALK2** *Genesis* <https://doi.org/10.1002/dvg.20201>

Furet P, Guagnano V, Fairhurst RA, Imbach-Weese P, Bruce I, Knapp M, Fritsch C, Blasco F, Blanz J, Aichholz R, Hamon J, Fabbro D, Caravatti G (2013) **Discovery of NVP-BYL719 a potent and selective phosphatidylinositol-3 kinase alpha inhibitor selected for clinical evaluation** *Bioorg Med Chem Lett* **23**:3741–3748 <https://doi.org/10.1016/j.bmcl.2013.05.007>

Beatriz Gámez, Rodríguez-Carballo E, Graupera M, Rosa JL, Ventura F. (2016) **Class I PI-3-Kinase Signaling Is Critical for Bone Formation Through Regulation of SMAD1 Activity in Osteoblasts** *Journal of Bone and Mineral Research* **31**:1617–1630

Gámez B., Rodríguez-Carballo E, Graupera M, Rosa JL, Ventura F (2016) **Class I PI-3-Kinase Signaling Is Critical for Bone Formation Through Regulation of SMAD1 Activity in Osteoblasts** *Journal of Bone and Mineral Research* **31** <https://doi.org/10.1002/jbmr.2819>

Groppe JC, Wu J, Shore EM, Kaplan FS (2011) **In vitro analyses of the dysregulated R206H ALK2 kinase-FKBP12 interaction associated with heterotopic ossification in FOPCells** *Tissues Organs* :291–295 <https://doi.org/10.1159/000324230>

Hatsell SJ, Idone V, Wolken DMA, Huang L, Kim HJ, Wang L, Wen X, Nannuru KC, Jimenez J, Xie L, Das N, Makhoul G, Chernomorsky R, D'Ambrosio D, Corpina RA, Schoenherr CJ, Feeley K, Yu PB, Yancopoulos GD, Murphy AJ, Economides AN (2015) **ACVR1 R206H receptor mutation causes fibrodysplasia ossificans progressiva by imparting responsiveness to activin A** *Sci Transl Med* **7**:303–303 <https://doi.org/10.1126/scitranslmed.aac4358>

Hino K, Ikeya M, Horigome K, Matsumoto Y, Ebise H, Nishio M, Sekiguchi K, Shibata M, Nagata S, Matsuda S, Toguchida J (2015) **Neofunction of ACVR1 in fibrodysplasia ossificans progressiva** *Proceedings of the National Academy of Sciences* **112**:15438–15443 <https://doi.org/10.1073/pnas.1510540112>

Hwang CD, Pagani CA, Nunez JH, Cherief M, Qin Q, Gomez-Salazar M, Kadaikal B, Kang H, Chowdary AR, Patel N, James AW, Levi B (2022) **Contemporary perspectives on heterotopic ossification** *JCI Insight* **7** <https://doi.org/10.1172/jci.insight.158996>

Hwang C, Pagani CA, Das N, Marini S, Huber AK, Xie LQ, Jimenez J, Brydges S, Lim WK, Nannuru KC, Murphy AJ, Economides AN, Hatsell SJ, Levi B (2020) **Activin A does not drive post-traumatic heterotopic ossification** *Bone* **138** <https://doi.org/10.1016/j.bone.2020.115473>

Ikegami D, Akiyama H, Suzuki A, Nakamura T, Nakano T, Yoshikawa H, Tsumaki N (2011) **Sox9 sustains chondrocyte survival and hypertrophy in part through Pik3ca-Akt pathways** *Development* <https://doi.org/10.1242/dev.057802>

Jamieson S, Flanagan JU, Kolekar S, Buchanan C, Kendall JD, Lee W-J, Rewcastle GW, Denny WA, Singh R, Dickson J, Baguley BC, Shepherd PR (2011) **A drug targeting only p110 α can block phosphoinositide 3-kinase signalling and tumour growth in certain cell types** *Biochemical Journal* <https://doi.org/10.1042/BJ20110502>

Kan L, Liu Y, McGuire TL, Berger DMP, Awatramani RB, Dymecki SM, Kessler JA (2009) **Dysregulation of Local Stem/Progenitor Cells as a Common Cellular Mechanism for Heterotopic Ossification** *Stem Cells* **27**:150–156 <https://doi.org/10.1634/stemcells.2008-0576>

Kaplan FS, Zasloff MA, Kitterman JA, Shore EM, Hong CC, Rocke DM (2010) **Early mortality and cardiorespiratory failure in patients with fibrodysplasia ossificans progressiva** *Journal of Bone and Joint Surgery - Series A* **92** <https://doi.org/10.2106/JBJS.I.00705>

Kaplan FS, Zeitlin L, Dunn SP, Benor S, Hagin D, al Mukaddam M, Pignolo RJ. (2018) **Acute and chronic rapamycin use in patients with Fibrodysplasia Ossificans Progressiva: A report of two cases** *Bone* **109**:281–284 <https://doi.org/10.1016/j.bone.2017.12.011>

Lees-Shepard JB, Stoessel SJ, Chandler JT, Bouchard K, Bento P, Apuzzo LN, Devarakonda PM, Hunter JW, Goldhamer DJ (2022) **An anti-ACVR1 antibody exacerbates heterotopic ossification by fibro-adipogenic progenitors in fibrodysplasia ossificans progressiva mice** *Journal of Clinical Investigation* **132** <https://doi.org/10.1172/JCI153795>

Lees-Shepard JB, Yamamoto M, Biswas AA, Stoessel SJ, Nicholas SAE, Cogswell CA, Devarakonda PM, Schneider MJ, Cummins SM, Legendre NP, Yamamoto S, Kaartinen V, Hunter JW, Goldhamer DJ (2018) **Activin-dependent signaling in fibro/adipogenic progenitors causes fibrodysplasia ossificans progressiva** *Nat Commun* **9** <https://doi.org/10.1038/s41467-018-02872-2>

- Madsen RR, Semple RK (2022) **PIK3CA-related overgrowth: silver bullets from the cancer arsenal?** *Trends Mol Med* **28**:255–257 <https://doi.org/10.1016/j.molmed.2022.02.009>
- Matsuo K, Chavez RD, Barruet E, Hsiao EC (2019) **Inflammation in Fibrodysplasia Ossificans Progressiva and Other Forms of Heterotopic Ossification** *Curr Osteoporos Rep* **17**:387–394 <https://doi.org/10.1007/s11914-019-00541-x>
- Matsuo K, Lepinski A, Chavez RD, Barruet E, Pereira A, Moody TA, Ton AN, Sharma A, Hellman J, Tomoda K, Nakamura MC, Hsiao EC (2021) **ACVR1R206H extends inflammatory responses in human induced pluripotent stem cell-derived macrophages** *Bone* **153** <https://doi.org/10.1016/j.bone.2021.116129>
- Morin G, Degrugillier-Chopinet C, Vincent M, Fraissenon A, Aubert H, Chapelle C, Huguin C, Dubos F, Catteau B, Petit F, Mezel A, Domanski O, Herbreteau G, Alesandrini M, Boddaert N, Boutry N, Broissand C, Han TK, Branle F, Sarnacki S, Blanc T, Guibaud L, Canaud G (2022) **Treatment of two infants with PIK3CA-related overgrowth spectrum by alpelisib** *J Exp Med* **219** <https://doi.org/10.1084/jem.20212148>
- Nakashima K, Zhou X, Kunkel G, Zhang Z, Deng JM, Behringer RR, De Crombrughe B. (2002) **The novel zinc finger-containing transcription factor Osterix is required for osteoblast differentiation and bone formation** *Cell* [https://doi.org/10.1016/S0092-8674\(01\)00622-5](https://doi.org/10.1016/S0092-8674(01)00622-5)
- Patel NK, Huber AK, Levi B (2022) **Macrophage TGFβ signaling is critical for wound healing with heterotopic ossification after trauma**
- Pignolo RJ, Bedford-Gay C, Liljesthröm M, Durbin-Johnson BP, Shore EM, Rocke DM, Kaplan FS (2016) **The Natural History of Flare-Ups in Fibrodysplasia Ossificans Progressiva (FOP): A Comprehensive Global Assessment** *Journal of Bone and Mineral Research* **31**:650–656 <https://doi.org/10.1002/jbmr.2728>
- Pignolo RJ, Shore EM, Kaplan FS (2013) **Fibrodysplasia ossificans progressiva: diagnosis, management, and therapeutic horizons** *Pediatr Endocrinol Rev* **10**:437–448
- Ramachandran A, Mehić M, Wasim L, Malinova D, Gori I, Blaszczyk BK, Carvalho DM, Shore EM, Jones C, Hyvönen M, Tolar P, Hill CS (2021) **Pathogenic ACVR1 R206H activation by Activin A-induced receptor clustering and autophosphorylation** *EMBO J* **40** <https://doi.org/10.15252/emboj.2020106317>
- Rosen CJ, Lees-Shepard JB, Nicholas S-AE, Stoessel SJ, Devarakonda PM, Schneider MJ, Yamamoto M, Goldhamer DJ (2018) **Palovarotene reduces heterotopic ossification in juvenile FOP mice but exhibits pronounced skeletal toxicity** <https://doi.org/10.7554/eLife.40814.001>
- Sánchez-Duffhues G, Williams E, Benderitter P, Orlova V, van Wijhe M, de Vinuesa A Garcia, Kerr G, Caradec J, Lodder K, de Boer HC, Goumans MJ, Eekhoff EMW, Morales-Piga A, Bachiller-Corral J, Koolwijk P, Bullock AN, Hoflack J, ten Dijke P (2019) **Development of Macrocyclic Kinase Inhibitors for ALK2 Using Fibrodysplasia Ossificans Progressiva-Derived Endothelial Cells** *JBM Plus* **3** <https://doi.org/10.1002/jbm4.10230>
- Shimono K, Tung W, Macolino C, Chi AH-T, Didizian JH, Mundy C, Chandraratna RA, Mishina Y, Enomoto-Iwamoto M, Pacifici M, Iwamoto M (2011) **Potent inhibition of heterotopic ossification by nuclear retinoic acid receptor-γ agonists** *Nat Med* <https://doi.org/10.1038/nm.2334>

- Shore EM, Xu M, Feldman GJ, Fenstermacher DA, Brown MA, Kaplan FS (2006) **A recurrent mutation in the BMP type I receptor ACVR1 causes inherited and sporadic fibrodysplasia ossificans progressiva** *Nat Genet* **38**:525–527 <https://doi.org/10.1038/ng1783>
- Sierra-Filardi E, Nieto C, Domínguez-Soto Á, Barroso R, Sánchez-Mateos P, Puig-Kroger A, López-Bravo M, Joven J, Ardavin C, Rodríguez-Fernández JL, Sánchez-Torres C, Mellado M, Corbí ÁL (2014) **CCL2 Shapes Macrophage Polarization by GM-CSF and M-CSF: Identification of CCL2/CCR2-Dependent Gene Expression Profile** *The Journal of Immunology* **192**:3858–3867 <https://doi.org/10.4049/jimmunol.1302821>
- Soleimani M, Nadri S (2009) **A protocol for isolation and culture of mesenchymal stem cells from mouse bone marrow** *Nat Protoc* **4**:102–106 <https://doi.org/10.1038/nprot.2008.221>
- Sorkin M, Huber AK, Hwang C, Iv WFC, Menon R, Li J, Vasquez K, Pagani C, Patel N, Li S, Visser ND, Niknafs Y, Loder S, Scola M, Nycz D, Gallagher K, McCauley LK, Xu J, James AW, Agarwal S, Kunkel S, Mishina Y, Levi B (2020) **. by monocytes in a mouse model of aberrant** *Nat Commun* <https://doi.org/10.1038/s41467-019-14172-4>
- Stanley A, Tichy ED, Kocan J, Roberts DW, Shore EM, Mourkioti F (2022) **Dynamics of skeletal muscle-resident stem cells during myogenesis in fibrodysplasia ossificans progressiva** *NPJ Regen Med* **7**:1–15 <https://doi.org/10.1038/s41536-021-00201-8>
- Torossian F, Guerton B, Anginot A, Alexander KA, Desterke C, Soave S, Tseng HW, Arouche N, Boutin L, Kulina I, Salga M, Jose B, Pettit AR, Clay D, Rochet N, Vlachos E, Genet G, Debaud C, Denormandie P, Genet F, Sims NA, Banzet S, Levesque JP, Lataillade JJ, Le Bousse-Kerdilès MC (2017) **Macrophage-derived oncostatin M contributes to human and mouse neurogenic heterotopic ossifications** *JCI Insight* **2** <https://doi.org/10.1172/jci.insight.96034>
- Towler OW, Shore EM (2022) **BMP signaling and skeletal development in fibrodysplasia ossificans progressiva (FOP)** *Developmental Dynamics* **251**:164–177 <https://doi.org/10.1002/dvdy.387>
- Tu B, Li J, Sun Z, Zhang T, Liu H, Yuan F, Fan C (2022) **Macrophage-Derived TGF- β and VEGF Promote the Progression of Trauma-Induced Heterotopic Ossification** *Inflammation* <https://doi.org/10.1007/s10753-022-01723-z>
- Upadhyay J, Xie LQ, Huang L, Das N, Stewart RC, Lyon MC, Palmer K, Rajamani S, Graul C, Lobo M, Wellman TJ, Soares EJ, Silva MD, Hesterman J, Wang L, Wen X, Qian X, Nannuru K, Idone V, Murphy AJ, Economides AN, Hatsell SJ (2017) **The Expansion of Heterotopic Bone in Fibrodysplasia Ossificans Progressiva Is Activin A-Dependent** *Journal of Bone and Mineral Research* **32**:2489–2499 <https://doi.org/10.1002/jbmr.3235>
- Valer JA, Sánchez-de-Diego C, Gámez B, Mishina Y, Rosa JL, Ventura F. (2019) **Inhibition of phosphatidylinositol 3-kinase α (PI3K α) prevents heterotopic ossification** *EMBO Mol Med* <https://doi.org/10.15252/emmm.201910567>
- Valer JA, Sánchez-de-Diego C, Gámez B, Mishina Y, Rosa JL, Ventura F. (2019) **Inhibition of phosphatidylinositol 3-kinase α (PI3K α) prevents heterotopic ossification** *EMBO Mol Med* **11** <https://doi.org/10.15252/emmm.201910567>

Van Dinther M, Visser N, De Gorter DJJ, Doorn J, Goumans MJ, De Boer J, Ten Dijke P. (2010) **ALK2 R206H mutation linked to fibrodysplasia ossificans progressiva confers constitutive activity to the BMP type I receptor and sensitizes mesenchymal cells to BMP-induced osteoblast differentiation and bone formation** *Journal of Bone and Mineral Research* **25**:1208–1215 <https://doi.org/10.1002/jbmr.091110>

Wang JH, Liu YZ, Yin LJ, Chen L, Huang J, Liu Y, Zhang RX, Zhou LY, Yang QJ, Luo JY, Zuo G wei, Deng ZL, He BC. (2013) **BMP9 and COX-2 form an important regulatory loop in BMP9-induced osteogenic differentiation of mesenchymal stem cells** *Bone* **57**:311–321 <https://doi.org/10.1016/j.bone.2013.08.015>

Wang X, Li F, Xie L, Crane J, Zhen G, Mishina Y, Deng R, Gao B, Chen H, Liu S, Yang P, Gao M, Tu M, Wang Y, Wan M, Fan C, Cao X (2018) **Inhibition of overactive TGF- β attenuates progression of heterotopic ossification in mice** *Nat Commun* **9** <https://doi.org/10.1038/s41467-018-02988-5>

Wang X, Li F, Xie L, Crane J, Zhen G, Mishina Y, Deng R, Gao B, Chen H, Liu S, Yang P, Gao M, Tu M, Wang Y, Wan M, Fan C, Cao X. (1970) **progression of heterotopic ossi fi cation in mice** *Nat Commun* <https://doi.org/10.1038/s41467-018-02988-5>

Yu PB, Deng DY, Lai CS, Hong CC, Cuny GD, Bouxsein ML, Hong DW, McManus PM, Katagiri T, Sachidanandan C, Kamiya N, Fukuda T, Mishina Y, Peterson RT, Bloch KD (2008) **BMP type I receptor inhibition reduces heterotopic ossification** *Nat Med* <https://doi.org/10.1038/nm.1888>

Article and author information

José Antonio Valer

Departament de Ciències Fisiològiques, Universitat de Barcelona, IDIBELL, C/ Feixa Llarga s/n 08907 Hospitalet de Llobregat, SPAIN

Alexandre Deber

Departament de Ciències Fisiològiques, Universitat de Barcelona, IDIBELL, C/ Feixa Llarga s/n 08907 Hospitalet de Llobregat, SPAIN

Marius Wits

Department of Cell and Chemical Biology, Leiden University Medical Center, Einthovenweg 20, 2333 ZC, Leiden, the Netherlands

Carolina Pimenta-Lopes

Departament de Ciències Fisiològiques, Universitat de Barcelona, IDIBELL, C/ Feixa Llarga s/n 08907 Hospitalet de Llobregat, SPAIN

Marie-José Goumans

Department of Cell and Chemical Biology, Leiden University Medical Center, Einthovenweg 20, 2333 ZC, Leiden, the Netherlands
ORCID iD: [0000-0001-9344-6746](https://orcid.org/0000-0001-9344-6746)

José Luis Rosa

Departament de Ciències Fisiològiques, Universitat de Barcelona, IDIBELL, C/ Feixa Llarga s/n 08907 Hospitalet de Llobregat, SPAIN

Gonzalo Sánchez-Duffhues

Department of Cell and Chemical Biology, Leiden University Medical Center, Einthovenweg 20, 2333 ZC, Leiden, the Netherlands, Nanomaterials and Nanotechnology Research Center (CINN-CSIC), Health Research Institute of Asturias (ISPA), 33011 Oviedo, Asturias, Spain

Francesc Ventura

Departament de Ciències Fisiològiques, Universitat de Barcelona, IDIBELL, C/ Feixa Llarga s/n 08907 Hospitalet de Llobregat, SPAIN

For correspondence: fventura@ub.edu

Copyright

© 2023, Valer et al.

This article is distributed under the terms of the [Creative Commons Attribution License \(https://creativecommons.org/licenses/by/4.0/\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Sakae Tanaka

University of Tokyo, Japan

Senior Editor

Mone Zaidi

Icahn School of Medicine at Mount Sinai, United States of America

Reviewer #1 (Public Review):

Summary:

In the present study, the authors examined the possibility of using phosphatidyl-inositol kinase 3-kinase alpha (PI3Ka) inhibitors for heterotopic ossification (HO) in fibrodysplasia ossificans progressiva (FOP). Administration of BYL719, a chemical inhibitor of PI3Ka, prevented HO in a mouse model of FOP that expressed a mutated ACVR1 receptor. Genetic ablation of PI3Ka (p110a) also suppressed HO in mice. BYL719 blocked osteochondroprogenitor specification and reduced inflammatory responses, such as pro-inflammatory cytokine expression and migration/proliferation of immune cells. The authors claimed that inhibition of PI3Ka is a safe and effective therapeutic strategy for HO.

Strengths:

This manuscript reports an interesting finding that BYL719 inhibits HO in a mouse model of FOP.

Weaknesses:

The molecular mechanisms of BYL719 were still unclear because BYL719 affected multiple events and many types of cells. Additional experimental data would be needed to show more clearly how PI3Ka regulates HO.

Reviewer #2 (Public Review):

Summary:

The authors in this study previously reported that BYL719, an inhibitor of PI3K α , suppressed heterotopic ossification in mice model of a human genetic disease, fibrodysplasia ossificans progressive, which is caused by the activation of mutant ACVR1/R206H by Activin A. The aim of this study is to identify the mechanism of BYL719 for the inhibition of heterotopic ossification. They found that BYL719 suppressed heterotopic ossification in two ways: one is to inhibit the specification of precursor cells for chondrogenic and osteogenic differentiation and the other is to suppress the activation of inflammatory cells.

Strengths:

This study is based on the authors' previous reports and the experimental procedures including the animal model are established. In addition, to confirm the role of PI3K α , the authors used the conditional knock-out mice of the subunit of PI3K α . They clearly demonstrated the evidence indicating that the targets of PI3K α are not members of TGFBR by a newly established experimental method.

Weaknesses:

Overall, the presented data were closely related to those previously published by the authors' group or others, and there were very few new findings.

Heterotopic ossification in the mice model was not stable and was inappropriate for scientific evaluation.

The method for chondrogenic differentiation was not appropriate, and the scientific evidence of successful differentiation was lacking.

The design of the gene expression profile comparison was not appropriate and failed to obtain the data for the main aim of this study.

The experiments of inflammatory cells were performed in cell lines without ACVR1/R206H mutation, and therefore the obtained data were not precisely related to the inflammation in FOP.