

The Intricate Relationship of G-Quadruplexes and Pathogenicity Islands: A Window into Bacterial Pathogenicity

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

December 5, 2023 (this version)

Posted to bioRxiv

September 13, 2023

Sent for peer review

September 13, 2023

Bo Lyu 

Division of Plant Science and Technology, University of Missouri, Columbia, MO 65211, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The dynamic interplay between guanine-quadruplex (G4) structures and pathogenicity islands (PAIs) represents a captivating area of research with implications for understanding the molecular mechanisms underlying pathogenicity. This study conducted a comprehensive analysis of a large-scale dataset from reported 89 pathogenic strains to investigate the potential interactions between G4 structures and PAIs. G4 structures exhibited an uneven and non-random distribution within the PAIs and were consistently conserved within the same pathogenic strains. Additionally, this investigation identified positive correlations between the number and frequency of G4s and the GC content across different genomic features, including the genome, promoters, genes, tRNA, and rRNA regions, indicating a potential relationship between G4 structures and the GC-associated regions of the genome. The observed differences in GC content between PAIs and the core genome further highlight the unique nature of PAIs and underlying factors, such as DNA topology. High-confidence G4 structures within regulatory regions of *Escherichia coli* were identified, modulating the efficiency or specificity of DNA integration events within PAIs. Collectively, these findings pave the way for future research to unravel the intricate molecular mechanisms and functional implications of G4-PAI interactions, thereby advancing our understanding of bacterial pathogenicity and the role of G4 structures in infectious diseases.

eLife assessment

This **fundamental** study explores the relationship between guanine-quadruplex structures and pathogenicity islands in 89 pathogenic strains. Guanine-quadruplex structures were found to be non-randomly distributed within pathogenicity islands and conserved within the same strains. Positive correlations were observed between Guanine-quadruplex structures and GC content across various genomic features, suggesting a link between these structures and GC-rich regions. These **compelling** findings shed light on the molecular mechanisms of Guanine-quadruplex structure-pathogenicity island interactions and will be of interest to all microbiologists.

Introduction

The discovery of the DNA double helix by Watson and Crick in 1953 revolutionized our understanding of genetics and laid the foundation for the modern field of molecular biology (1). Nonetheless, the intricate nature of DNA continues to surprise us even today. One such captivating feature is the DNA guanine (G)-quadruplex (G4) structure, a unique arrangement that defies the conventional double helix (2, 3). A G4 consists of four guanine bases and stabilized by Hoogsteen hydrogen bonding. These stacked tetrads are interconnected by loop regions, which can vary in length and sequence, adding further complexity to the structure (Fig. 1). It is important to consider the inherent directionality of nucleic acids, with all four strands having the possibility to run in the same 5' to 3' direction, referred to as “parallel”, or alternatively, they can run in different directions, known as “antiparallel”. G4 regions can be very stable *in vitro*, particularly in the presence of K^+ (4).

Pathogenicity islands (PAIs) are genomic regions that contribute to the virulence and pathogenic potential of various microorganisms (5, 6). PAIs often exhibit close proximity to tRNA genes, suggesting a putative mechanism where tRNA genes act as anchor points for the integration of foreign DNA acquired through horizontal gene transfer (Fig. 1E). One notable feature is their variable GC content, which tends to deviate from the average GC content of the genome in various organisms, such as *Streptomyces* (7), *Salmonella* (8), and *Yersinia* (9). One question raised in PAI is that PAIs often exhibit distinct base composition (G+C contents) compared to the core genome. The underlying reasons for this variation remain unknown, but the preservation of a genus-or species-specific base composition represents a noteworthy characteristic of bacteria (5). Herbert Schmidt et al. (2004) proposed a hypothetical mechanism to explain the observed variation, suggesting that factors such as DNA topology and codon message in the virulence regions present could contribute to the preservation of the distinct base composition (5). Hopefully, the availability of genome sequences from pathogenic bacteria and their non-pathogenic counterparts currently presents an exceptional opportunity to explore the intricate structure variance and underlying mechanisms within PAIs.

The possibility of interactions between G4 structures and pathogens has been suggested, although this field of study is still in its nascent phase. Some studies observed that bacterial genomes possess G4-forming sequences within their genome regions (10, 11). G4 structures are formed by G-rich DNA sequences, and their stability is influenced by the G+C content and arrangement of G tetrads. Interestingly, PAIs often exhibit an altered GC content, putatively contributing to the propensity of G4 structure formation within these regions. The G4 structures in PAIs might modulate the accessibility of transcription factors, DNA-binding proteins, or RNA polymerase in pathogens, as documented in eukaryotes (2, 12), thereby influencing the expression of virulence-associated genes. The formation of G4 structures within PAIs may serve as an additional layer of regulation that fine-tunes the expression of genes critical for pathogenesis. Hence, the investigation of G4 structures within PAIs may open new avenues for the development of therapeutic strategies aimed at disrupting the regulatory mechanisms of pathogenicity-associated genes.




Figure 1.

Structural and Functional Aspects of G-quadruplex (G4) Structures and Pathogenicity Islands (PAIs).

(A) Schematic representation of a guanine tetrad stabilized by Hoogsteen base pairing and a positively charged central ion, illustrating the key elements of G4 structures. (B) Structural heterogeneity of G4s. G-quadruplex structures exhibit polymorphism and can be categorized into different families, such as parallel or antiparallel, based on the orientation of the DNA strands. They can fold either intramolecularly or intermolecularly, leading to diverse structural configurations. (C) General sequence formula for G4s, highlighting the repeated occurrence of guanine-rich sequences that form G-quartets. (D) Regulatory roles of G-quadruplexes in transcription. G4 structures can regulate transcription by blocking RNA polymerase from binding to promoter sequences or aiding in single-stranded DNA (ssDNA) formation, thereby enhancing transcription. (E) General structure of PAI. PAIs are characteristic regions of DNA found within the genomes of pathogenic bacteria, distinguishing them from nonpathogenic strains of the same or related species. Repeat sequences are DNA segments duplicated within the PAI and can serve as recognition sites for various enzymes involved in the integration and excision of the PAI from the bacterial chromosome. tRNA genes act as anchor points for the insertion of foreign DNA acquired through horizontal gene transfer. Virulence genes encode proteins or factors that play crucial roles in the virulence and pathogenicity of the bacterium, contributing to adhesion, invasion, immune evasion, toxin production, or other pathogenic mechanisms. Insertion elements include transposons, bacteriophages, or plasmids, enabling the PAI to be transferred between bacterial cells and potentially disseminated to different strains or species.

Results and Discussion

Genomic information, PAI patterns, and the presence of G4s in 89 reported pathogenic strains

A dataset of PAIs was compiled from 89 reported pathogenic strains, encompassing 222 distinct types of PAIs. Those pathogens that share closely related PAIs on phylogenetic branches, such as LEE in *E. coli* strains (Fig. 2A), may have acquired these elements through recent common ancestors or horizontal gene transfer events (5, 13, 14). Additional information, including the genome length (bp), G+C content (%), rRNA density, tRNA density, and PAI length (bp), was present and showed conserved patterns in the same species (Fig. 2A & Table S1). PAIs commonly exhibit mosaic-like patterns, exemplified by the presence of distinct PAIs like FPI in *Francisella tularensis*, SaPI_{bov} in *Staphylococcus aureus*, and Hrp PAI in *Xanthomonas campestris* (Fig. 2B). Many PAIs were present associated with tRNAs, such as the insertions of tRNA^{Thr}, tRNA^{Phe}, and tRNA^{Gly} in *E. coli* strains (Fig. 2B & Table S2). The presence of PAIs distributes in similar genomic regions across different pathogens or strains, showing non-random patterns and functionally clustered. Employing the G4Hunter search algorithm, the study identified a total of 225,376 putative G4 sequences in these 89 pathogenic genomes (Table S1). The heatmap showed that the number of G4 structures was diverse in the pathogen genomes (Fig. 2C), similar to previous reports that showed that the G4 structures display uneven distribution patterns in eukaryotic and prokaryotic genomes and are conserved in evolutionary groups (15–17).

Interaction between PAIs and G4s in different genomic features

The analysis of G4 structures across all pathogen species demonstrated a positive correlation between the number of G4s and the GC content in various genomic features, including the whole genome, gene, promoter, rRNA, and tRNA regions (Fig. 2D). Furthermore, the density of G4s, measured as the frequency of predicted G4-forming sequences per 1000 base pairs (bp), also showed a positive correlation with the GC content across the analyzed genomic elements (Fig. 2E). The presence of G4 structures within the promoter, rRNA, and tRNA regions may have functional implications for the regulation of DNA replication, ribosome biogenesis, protein synthesis, and other RNA-related processes (18–20). Throughout evolution, there seems to be a greater enrichment of G4 structures in regulatory genes compared to other genes, enabling intricate control of gene expression in cascade signal transduction pathways (21).

This study observed that the GC contents in the genome region were significantly higher compared to the corresponding PAIs region that was classified into five parts according to the genome datasets (Fig. 3A–E & Table S1). This suggests that the genomic regions surrounding the PAIs (i.e., core genome) tend to have a higher GC content than PAI regions, which was consistent with the fact that PAIs often exhibit distinct base composition compared with the core genome (5). The variation was explained by the presence of G4 sequences within the PAIs, whereas the results were surprising. This study observed a distinct pattern in the frequency of G4 structures within different regions of the PAIs. This differential distribution of G4 structures suggests that (i) specific segments (e.g., <30% and >60% GC contents) within the PAIs may be more prone to induce G4 formation discrepancy, as these segments showed unstable gene structure; (ii) the variation of base composition between core genome and PAIs is partially correlated with the presence of G4s; (iii) the frequency of G4s in PAIs present stable as the core genome in the most situation; (iv) an alternative hypothesis, other factors, such as i-motif (i.e., the anti-G4 structure) and CpG island, may work synergistically with G4 and potentially contribute the base composition variation (22, 23).




Figure 2.

Analysis of Pathogenicity Islands (PAIs) and G-quadruplexes (G4s) in Pathogen Genomes.

(A) Phylogenetic analysis of pathogen genomes based on 89 bacterial strains, showing the evolutionary relationships among species. Additional genomic information, including genome size, GC content, rRNA density, tRNA density, and PAI length, is provided. The same color indicates the same species. (B) Genomic location of specific PAIs in bacterial genomes, divided into ten regions. PAIs are represented by green triangles, and their names are indicated. The tRNA insertion sites are also marked. (C) Heatmap illustrating the relative abundance of G4s in bacterial genomes, divided into ten regions. Red indicates a higher relative abundance, while blue indicates a lower relative abundance. (D & E) Correlation analysis between the number of G4s, the frequency of G4s and GC content in various genomic features, including the whole genome, genes, promoters, rRNA, and tRNA.




Figure 3.

Putative Origin and Functional Annotation of G-quadruplexes (G4s) within Pathogenicity Islands (PAIs) in *Escherichia coli*.

(A-E) Comparison of GC content (left panel) and GC frequency (right panel) between the genome and pathogenicity islands (PAIs), categorized into five regions (20%-30%, 30%-40%, 40%-50%, 50%-60%, and 60%-70%). */**/**/* indicates significant difference ($p < 0.05/0.01/0.001/0.0001$). (F) Hypotheses on the origin of G4 structures within PAIs, involving gene horizontal transfer mechanisms (conjugation, transduction, and transformation). (G) Evolutionary relatedness of 10 types of PAIs (categorized into six main categories) in *E. coli* strains. (H & I) Examples of G4 structures within PAIs in *E. coli* strains. The grey bar represents the virulence region, the red box indicates a virulence gene, the blue box represents an insertion site region or repeat, the green box denotes an integrase, the purple triangle indicates a tRNA insertion site, and the yellow triangle indicates an effector. (J & K) Functional annotation analysis of G4-covered genes within PAIs in two *E. coli* strains, including biological process (BP), cellular component (CC), and molecular function (MF) categories.

Putative origin, transfer mechanisms, and functions of G4s in PAIs

To understand the origin of G4 structures within PAIs, the study hypothesized that these G4 sequences could be acquired through three types of horizontal gene transfer mechanisms: conjugation, transformation, and transduction (**Fig. 3F**). These mechanisms serve as means for genetic material exchange between different organisms. Considering the presence of G4 sequences within the PAIs, it is plausible that these sequences are transferred along with the PAIs through these horizontal gene transfer mechanisms. Later, the study used *E. coli* as an example to investigate the potential regulatory role and function of genes covered by G4 structures in PAIs. *E. coli* contains at least ten types of PAI in different strains, and one of the well-known PAI is LEE (**Fig. 3G**), harboring genes responsible for causing attaching and effacing lesions ([14](#), [24](#)). One stable G4 structure with a G4hunter score of 1.6 was identified at position 37,085 in the LEE PAI of *E. coli* str. O103:H2 12009 (**Fig. 3H**), located between an IS element and a tRNA insertion site. The tRNA region generally contains a higher G4 frequency compared with transfer-messenger RNA (tmRNA) and rRNA regions in the bacterial genome ([16](#)). Interestingly, this G4 structure found in *E. coli* str. O103:H2 12009 was present in close proximity to a tRNA region, suggesting a potential regulatory role of G4s in the tRNA gene, or upstream- and downstream-genes that are responsible for LEE virulence. Additionally, another stable G4 sequence with a score of 1.381 was discovered at position 12,457 in the *E. coli* str. CFT073 PAI II to provide one more evidence of G4 in PAI regions (**Fig. 3I**).

Functional enrichment analysis was conducted to explore the putative functions of G4-covered genes in two *E. coli* strains (Table S3&S4). The results revealed that the genes covered by G4 structures were predominantly involved in genetic information processes, including DNA binding, DNA integration, and nucleic acid metabolism processes (**Fig. 3J & K**). This suggests that G4 structures may play a regulatory role in these essential cellular processes, especially gene expression and DNA-related functions. For instance, G4 structures in the promoter regions of certain transcription factors may influence their binding affinity to DNA and subsequently affect downstream gene expression patterns ([25](#), [26](#)). These elements frequently utilize DNA integration mechanisms mediated by integrases, recombinases, or transposases to transfer or incorporate genetic material into the bacterial genome ([27](#), [28](#)).

Overall, the conserved evolutionary relatedness of PAIs, the detection of stable G4 structures in specific genomic positions, and the enrichment of G4-covered genes in genetic information processes collectively support the hypothesis that G4 structures may have regulatory functions in key biological processes in pathogens.

Methods

Selection and extraction of DNA Sequences

A total of 89 genomes corresponding to the identified pathogens from the Pathogenicity Island Database (PAIDB) were included in the study. The complete bacterial genomic DNA sequences and their corresponding annotation files in .gff and .fna formats were obtained from the Genome database of the National Center for Biotechnology Information (NCBI). To ensure the reliability and completeness of the dataset, only completely assembled genomes were included in the analysis. To avoid redundancy and incomplete sequences, one representative genome was selected for each species or strain. The selection of representative genomes was based on a careful examination of the supplementary material (Table S1) accompanying the study. The study employed TBtools (<https://cj-chen.github.io/tbtools>), a versatile bioinformatics tool, to investigate the presence of G-quadruplex (G4) sequences in specific genomic features. The gene regions, promoters (2 kb upstream of the genes), tRNA regions, and rRNA regions were extracted from the

selected genomes using TBtools. PAI regions were downloaded following previously documented information in PAIDB (Table S1&S2). Default thresholds and parameters were applied during extraction to maintain consistency across all genomes.

Data process and detection of G4s in genomic features

The G4Hunter algorithm, a widely used tool for G4 prediction, was employed for the identification of G4 motifs in the genomic sequences. The G4Hunter parameters were set to a window size of “25” and a G4 score threshold of 1.2, which ensured the identification of potential G4 sequences. The analysis was performed for each species group, where the genomic DNA size and the number of putative G4 sequences found were determined. The study quantified the predicted number of putative G4-forming sequences within different genomic features, including the whole genome, gene, promoter, tRNA, rRNA, and PAI regions. The density of G4 motifs was determined by dividing the number of G4 sequences by the total length of the genome, while the length ratio of G4 motifs was calculated by dividing the total length of the G4 sequences by the total length of the genome.

Relationship between G4s and PAIs

The heatmap was used to show the distribution of G4 motifs in the genome divided by ten parts as PAI regions using R package “pheatmap”. The correlation between the number of G4 structures and the GC content was analyzed across various genomic elements, including the whole genome, gene, promoter, rRNA, and tRNA regions. The analysis utilized the *R*-squared value (R^2) to determine the strength of the correlation. The significance of the correlation was assessed using *p*-values. The GC content in the genome regions and corresponding PAI regions was compared and classified into different ranges to explore the variation in base composition. GraphPad Prism (V.5.02, GraphPad Software, Inc) was employed to conduct Normality and Lognormality Tests. The K-S test and F-test were used to assess normal distribution and variances, and Student’s t-test was used to identify significant difference.

Phylogenetic Tree Construction

The exact Taxonomy ID (taxid) for each analyzed group was obtained from the NCBI Taxonomy Database using the Taxonomy Browser. The Neighbor-Joining (NJ) method was employed to construct the phylogenetic trees for the analyzed groups. The phylogenetic trees were generated using MEGA11 software (www.megasoftware.net [↗](#)), which offers robust algorithms and comprehensive tools for phylogenetic analysis. To assess the reliability and statistical support of the phylogenetic tree branches, bootstrap analysis was performed. One thousand bootstrap replicates were used to estimate the confidence levels of the branching patterns in the phylogenetic trees. The phylogenetic trees, along with the bootstrap support values, were displayed and visualized using the Interactive Tree Of Life (ITOL) platform (<https://itol.embl.de/> [↗](#)).

Gene functional annotation

The gene sequences covered by G4 structures within PAIs were subjected to gene ontology annotation. The gene sequences were translated into protein sequences using the Expasy online toolkit (<https://web.expasy.org/translate/> [↗](#)). This tool performs the translation based on the standard genetic code, converting the DNA nucleotide sequence into its corresponding amino acid sequence. The GO annotation database assigned GO terms to the protein sequences based on their predicted functions and known biological process (BP), molecular function (MF), and cellular component (CC). Fisher’s exact test was employed to determine the statistical significance of the enrichment results. The obtained *p*-values indicated the overrepresentation of specific GO terms, with lower *p*-values suggesting higher significance.

Statistics and reproducibility

All genomic data utilized in this study, including the species-specific datasets, were obtained from publicly available sources. Statistical analyses, such as the Kruskal-Wallis test, F test, Student's t-test, Wilcoxon test, and chi-square test, were performed using GraphPad Prism software. The samples used in the statistical analyses corresponded to the genomic data, PAIs, or specific genes under investigation.

Acknowledgements

The sincere appreciation extends to Dr. Sung Ho Yoon and his colleagues for their dedicated efforts in identifying PAIs and establishing the Pathogenicity Island Database for public analysis. Their commitment to advancing the field of pathogen genomics has greatly facilitated this research. This study would like to thank Prof. Qisheng Song and Dr. Jingjing Li for their insightful suggestions and constructive comments regarding the exploration of G4 structures in genomes. Their expertise and guidance have significantly enriched the understanding of the potential roles and implications of G4 structures in the context of PAIs. The study is grateful to Dr. Minyu Zhou for her valuable discussions and exchange of ideas regarding the interplay between G4 structures and pathogenicity islands.

Competing Interests

The author declares no competing interests.

Funding information

This research received no external funding.

Credit author statement

Bo Lyu: writing original draft, data analysis, and visualization.

Ethical approval


Ethical review and approval were not required for this study.

Availability of data and materials

The original reported PAIs datasets analyzed in this study are available from the publication with the DOI: 10.1093/nar/gku985. Additionally, supplementary materials accompanying this article provide further PAIs data analyzed in the study. These supplementary materials contain detailed information and datasets that support the findings and conclusions presented in this research. Researchers and interested individuals can access supplementary materials to explore and validate the results obtained in this study.

Consent for publication

Not applicable.



References

1. Watson J. D., Crick F. (1953) **A structure for deoxyribose nucleic acid**
2. Rhodes D., Lipps H. J. (2015) **G-quadruplexes and their regulatory roles in biology** *Nucleic acids research* **43**:8627–8637
3. Spiegel J., Adhikari S., Balasubramanian S. (2020) **The structure and function of DNA G-quadruplexes** *Trends in Chemistry* **2**:123–136
4. Stegle O., Payet L., Mergny J.-L., MacKay D. J., Huppert J. L. (2009) **Predicting and understanding the stability of G-quadruplexes** *Bioinformatics* **25**:i374–i1382
5. Schmidt H., Hensel M. (2004) **Pathogenicity islands in bacterial pathogenesis** *Clinical microbiology reviews* **17**:14–56
6. Groisman E. A., Ochman H. (1996) **Pathogenicity islands: bacterial evolution in quantum leaps** *Cell* **87**:791–794
7. Kers J. A., et al. (2005) **A large, mobile pathogenicity island confers plant pathogenicity on Streptomyces species** *Molecular Microbiology* **55**:1025–1033
8. Kombade S., Kaur N. (2021) **in Salmonella spp.-A Global Challenge.** (IntechOpen
9. Carniel E. (1999) **The Yersinia high-pathogenicity island** *Int Microbiol* **2**:161–167
10. Yadav P., et al. (2021) **G-quadruplex structures in bacteria: Biological relevance and potential as an antimicrobial target** *Journal of Bacteriology* **203**
11. Harris L. M., Merrick C. J. (2015) **G-quadruplexes in pathogens: a common route to virulence control?** *PLoS pathogens* **11**
12. Varshney D., Spiegel J., Zyner K., Tannahill D., Balasubramanian S. (2020) **The regulation and functions of DNA and RNA G-quadruplexes** *Nature reviews Molecular cell biology* **21**:459–474
13. Hartland E. L., Leong J. M. (2013) **E. L. Hartland, J. M. Leong. (Frontiers Media SA, 2013), vol. 3, pp. 15. Frontiers Media SA 3**
14. Jores J., Rumer L., Wieler L. H. (2004) **Impact of the locus of enterocyte effacement pathogenicity island on the evolution of pathogenic Escherichia coli** *International Journal of Medical Microbiology* **294**:103–113
15. Du Z., Zhao Y., Li N. (2009) **Genome-wide colonization of gene regulatory elements by G4 DNA motifs** *Nucleic acids research* **37**:6784–6798
16. Bartas M., et al. (2019) **The presence and localization of G-quadruplex forming sequences in the domain of bacteria** *Molecules* **24**
17. Lombardi E. P., Londoño-Vallejo A., Nicolas A. (2019) **Relationship between G-quadruplex sequence composition in viruses and their hosts** *Molecules* **24**

18. Ivanov P., et al. (2014) **G-quadruplex structures contribute to the neuroprotective effects of angiogenin-induced tRNA fragments** *Proceedings of the National Academy of Sciences* **111**:18201–18206
19. Mestre-Fos S., et al. (2019) **G-quadruplexes in human ribosomal RNA** *Journal of molecular biology* **431**:1940–1955
20. Zybailov B., Sherpa M., Glazko G., Raney K., Glazko V. (2013) **G4-quadruplexes and genome instability** *Molecular Biology* **47**:197–204
21. Wu F., et al. (2021) **Genome-wide analysis of DNA G-quadruplex motifs across 37 species provides insights into G4 evolution** *Communications biology* **4**
22. Deaton A. M., Bird A. (2011) **CpG islands and the regulation of transcription** *Genes Dev* **25**:1010–1022
23. Sushmita N., et al. (2020) **I-motif DNA: significance and future prospective** *Exploratory Animal and Medical Research* **10**:18–23
24. Franzin F. M., Sircili M. P. (2015) **Locus of enterocyte effacement: a pathogenicity island involved in the virulence of enteropathogenic and enterohemorrhagic Escherichia coli subjected to a complex network of gene regulation** *Biomed Res Int* **2015**
25. Niu K., et al. (2018) **BmILF and i-motif structure are involved in transcriptional regulation of BmPOUM2 in Bombyx mori** *Nucleic Acids Research* **46**:1710–1723
26. Xiang L., et al. (2022) **DNA G-quadruplex structure participates in regulation of lipid metabolism through acyl-CoA binding protein** *Nucleic Acids Research* **50**:6953–6967
27. Arkhipova I. R., Rice P. A. (2016) **Mobile genetic elements: in silico, in vitro, in vivo** *Mol Ecol* **25**:1027–1031
28. Wozniak R. A., Waldor M. K. (2010) **Integrative and conjugative elements: mosaic mobile genetic elements enabling dynamic lateral gene flow** *Nature Reviews Microbiology* **8**:552–563

Article and author information

Bo Lyu

Division of Plant Science and Technology, University of Missouri, Columbia, MO 65211, USA

For correspondence: bl3pt@missouri.edu

ORCID iD: [0000-0002-2744-730X](https://orcid.org/0000-0002-2744-730X)

Copyright

© 2023, Bo Lyu

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Bavesh Kana

University of the Witwatersrand, South Africa

Senior Editor

Bavesh Kana

University of the Witwatersrand, South Africa

Reviewer #1 (Public Review):

Summary:

This study explores the relationship between guanine-quadruplex (G4) structures and pathogenicity islands (PAIs) in 89 pathogenic strains. G4 structures were found to be non-randomly distributed within PAIs and conserved within the same strains. Positive correlations were observed between G4s and GC content across various genomic features, suggesting a link between G4 structures and GC-rich regions. Differences in GC content between PAIs and the core genome underscored the unique nature of PAIs. High-confidence G4 structures in *Escherichia coli*'s regulatory regions were identified, influencing DNA integration within PAIs. These findings shed light on the molecular mechanisms of G4-PAI interactions, enhancing our understanding of bacterial pathogenicity and G4 structures in infectious diseases.

Strengths:

The findings of this study hold significant implications for our understanding of bacterial pathogenicity and the role of guanine-quadruplex (G4) structures.

Molecular Mechanisms of Pathogenicity: The study highlights that G4 structures are not randomly distributed within pathogenicity islands (PAIs), suggesting a potential role in regulating pathogenicity. This insight into the uneven distribution of G4s within PAIs provides a basis for further research into the molecular mechanisms underlying bacterial pathogenicity.

Conservation of G4 Structures: The consistent conservation of G4 structures within the same pathogenic strains suggests that these structures might play a vital and possibly conserved role in the pathogenicity of these bacteria. This finding opens doors for exploring how G4s influence virulence across different pathogens.

Unique Nature of PAIs: The differences in GC content between PAIs and the core genome underscore the unique nature of PAIs. This distinction suggests that factors such as DNA topology and G4 structures might contribute to the specialized functions and characteristics of PAIs, which are often associated with virulence genes.

Regulatory Role of G4s: The identification of high-confidence G4 structures within regulatory regions of *Escherichia coli* implies that these structures could influence the efficiency or specificity of DNA integration events within PAIs. This finding provides a potential mechanism by which G4s can impact the pathogenicity of bacteria.

Weaknesses:

No weaknesses were identified by this reviewer.

Overall, the study provides fundamental insights into the pathogenicity island and conservation of G4 motifs.

Reviewer #2 (Public Review):

Summary:

In the manuscript entitled "The Intricate Relationship of G-Quadruplexes and Pathogenicity Islands: A Window into Bacterial Pathogenicity" Bo Lyu explored the interactions between guanine-quadruplex (G4) structures and pathogenicity islands (PAIs) in 89 bacterial genomes through a rigorous computational approach. This paper handles an intriguing and complex topic in the field of pathogenomics. It has the potential to contribute significantly to the understanding of G4-PAI interactions and bacterial pathogenicity.

Strengths:

- The chosen research area.
- The summarizing of the results through neat illustrations.

Weaknesses:

This reviewer did not find any significant weaknesses.

Reviewer #3 (Public Review):

The main problem with the work is that the results are only descriptive and do not allow any inferences or conclusions about the importance of the function of G4 structures. The discussion and conclusions are poor. The results are preliminary and in order to try to make the analysis more interesting, it should be further extended and the data must be explored in a much greater depth.