

Expression of modified FcγRI enables myeloid cells to elicit robust tumor-specific cytotoxicity

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

April 29, 2024 (this version)

Reviewed preprint version 1

December 27, 2023

Posted to preprint server

October 2, 2023

Sent for peer review

September 10, 2023

Leen Farhat-Younis, Manho Na, Amichai Zarfin, Aseel Khateeb, Nadine Santana-Magal, Alon Richter, Amit Gutwillig, Diana Rasoulouniriana, Annette Gleiberman, Lir Beck, Tamar Giger, Avraham Ashkenazi, Adi Barzel, Peleg Rider, Yaron Carmi 

Department of Pathology, School of Medicine, Tel Aviv University, Israel • Department of Human Molecular Genetics and Biochemistry, Tel Aviv University, Israel • Department of Molecular Cell Biology, Weizmann Institute, Israel • Department of Cell and Developmental Biology, School of Medicine, Tel Aviv University, Israel • Department of Biochemistry Molecular Biology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Israel

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Despite the central role of T cells in tumor immunity, attempts to harness their cytotoxic capacity as a therapy have met limited efficacy, partially as a result of the suppressive microenvironment which limits their migration and activation. In contrast, myeloid cells massively infiltrate tumors and are well adapted to survive these harsh conditions. While they are equipped with cell-killing abilities, they often adopt an immunosuppressive phenotype upon migration to tumors. Therefore, the questions of how to modify their activation programming against cancer, and what signaling cascades should be activated in myeloid cells to elicit their cytotoxicity have remained unclear.

Here, we found that activation of IgM-induced signaling in myeloid cells results in secretion of lytic granules and massive tumor cell death. These findings open venues for designing novel immunotherapy by equipping monocytes with chimeric receptors that target tumor antigens and consequently, signal through IgM receptor. Nonetheless, we found that myeloid cells do not express the antibody-derived portion used to recognize the tumor antigen due to the induction of an ER stress response. To overcome this limitation, we designed chimeric receptors that are based on the high-affinity FcγRI for IgG. Incubation of macrophages expressing these receptors along with tumor-binding IgG induced massive tumor cell killing and secretion of reactive oxygen species and Granzyme B.

Overall, this work highlights the challenges involved in genetically reprogramming the signaling in myeloid cells and provides a framework for endowing myeloid cells with antigen-specific cytotoxicity.

eLife assessment

The findings are **fundamental** for understanding IgM signaling in myeloid cells. The work is **compelling** in its ability to manipulate and harness myeloid cells to further anti-tumor immunity.

Introduction

Clinical and experimental data have highlighted the important role of the immune cell function and composition in determining tumor progression or eradication (1, 2). While a high prevalence of tumor-infiltrating T cells is associated with improved prognosis and survival (3, 4), infiltration of myeloid cells is often associated with poor prognosis and tumor refractory (5, 6). Based on these findings, vast scientific effort is executed to increase the host's T cell response to cancer cells (7, 8). In most cases however, tumor-reactive T cells generated spontaneously by the host bear low to moderate affinity (10^{-4}M - 10^{-6}M) (9, 10), which is below the activation threshold needed for therapeutic effect, and attempts to increase TCR affinity using genetical engineering are often limited by induction of cross-reactivity (11, 12).

To overcome these limitations, T cells can be genetically modified to express Chimeric Antigen receptor (CAR) consisting of Single Chain Fragment Variable (scFv) for tumor antigen recognition fused to T cell signaling domains (e.g., CD3 ζ , CD28) (13, 14). Engineered cells are reinfused into the patient, providing the host's immune system with T cells capable of specifically recognizing tumor antigens in an MHC-independent fashion. Indeed, this strategy benefits from overcoming the complex, multistep, and highly regulated activation process of T cells (15, 16) and has been proven remarkably successful in treating liquid tumors.

However, the harsh microenvironment characterizing solid tumors manifests a major limitation to T cell-based treatments. Hypoxic and acidic conditions at tumor sites, along with immunosuppressive capacity of tumor cells limit T cell infiltration, survival, and cytotoxicity (17-19). In contrast to T cells, myeloid cells migrate efficiently into the tumor mass and are well adapted to survive and function under this harsh environment (4, 5). However, while myeloid cells are potentially able to produce cytotoxic compounds, they acquire an immunosuppressive phenotype once in the vicinity of the tumor cells and promote tumor growth (20, 21). Hence, equipping myeloid cells with means to release cytotoxic compounds following recognition of tumor antigens may provide a new therapeutic strategy. While recognition of tumors can be facilitated through scFv, which signaling chains should be fused are not clear. Early attempts to generate CAR myeloid cells have fused to CD3- ζ chain, which may function in these cells as Fc γ chain (22, 23). Yet, what signaling cascades should be induced in myeloid cells to elicit their cytotoxicity has remained unclear.

Here, we found that activation of the IgM receptor signaling in myeloid cells induces massive killing of tumor cells. However, we found that in contrast to lymphoid cells, which can ectopically express scFv, myeloid cells identify it as a misfolded protein and rapidly degrade it. Instead, we modified the high-affinity Fc γ RI, which normally binds IgG, to transmit IgM receptor-driven signals, while recognition of tumor antigens is mediated by IgG tumor-binding antibodies. Indeed, incubation of mRNA-engineered myeloid cells with tumor cells resulted in massive tumor cell killing only in the presence of IgG tumor-binding antibodies. Overall, this work suggests a novel mean to endow myeloid cells with antigen-specific killing abilities and harness their capacity to migrate and survive in the harsh conditions at the tumor microenvironment.

Material and Method

Mice

Wild-type (WT) C57BL/6 mice were purchased from Envigo (Jerusalem, Israel). 129S1 mice were purchased from Jackson laboratories (Bar Harbor, Maine, USA). C57BL/6-Tg(TcraTcrb)1100Mjb/J OT-I mice were kindly gifted from Prof. Carmit Levy, Tel Aviv University, CD45.1 congenic mice were kindly gifted from Prof. Adi Barezel, Tel Aviv University, and B6.Cg-Tg(TcraTcrb)425Cbn/J OT-II mice were generously gifted from Prof. Jakub Abramson, Weizmann Institute, and were housed and maintained in a specific pathogen-free (SPF) conditions animal facility in Tel-Aviv University according to the American Association for the Accreditation of Laboratory Animal. Male and female 8-12 weeks-old mice were used in all experiments. Animal experiments were approved by the Tel-Aviv University ethics committee.

Cell lines

Human Embryonic Kidney (HEK)-293FT cells were purchased from ThermoFisher Scientific (Waltham, MA, USA). DC 2.4 were a kind gift from Dr. Kenneth Rock Gross from UMass Chan Medical School. RAW 264.7 and Jurkat cell lines were purchased directly from ATCC. THP-1 were kind gifts from Prof. Mordechai (Motti) Gerlic from the Department of Clinical Microbiology and Immunology at Tel Aviv University. B16F10 (CRL-6475) cells and 4T1 (CRL-2539) cells were directly purchased from ATCC. Cells were cultured in DMEM supplemented with 10% heat-inactivated FBS, 2 mM L-glutamine, and 100 µg/mL penicillin/streptomycin (all from Biological Industries, Beit Haemek, Israel). All cells were routinely tested for mycoplasma (EZ-PCR Mycoplasma Test Kit, Biological Industries).

Primary cells

For bone-marrow DC (BMDC), C57BL/6 mice were sacrificed using a CO₂ chamber. Femurs, hip bones, and tibias were obtained and kept in DMEM supplemented with 10% FBS. Bones were washed twice with Phosphate Buffered Saline (Gibco, NY, USA) and sterilized for 10 seconds in a 70% ethanol solution. The cleaned bones were crushed with a sterile mortar and pestle in full medium and the cell mixture was filtered through a 40µm cell strainer. Cells were then centrifuged and resuspended in complete DMEM supplemented with 50 ng/mL GM-CSF and 10 ng/mL IL-4 (PeproTech, Rehovot, Israel) and plated at a concentration of 1×10^6 /mL for 6-7 days.

Transient transfection

Liposomes-based transfection

For transfecting HEK293 cells, jetOPTIMUS reagent (Polyplus Transfection, Strasbourg, France) was used according to manufacturer protocol. For DNA transfection of macrophages, LipofectamineTM 2000 (Invitrogen, Waltham, MA, USA) and jetPEI-Macrophage (Polyplus Transfection) were used.

Electroporation

3×10^6 cells were suspended in 0.25 mL Opti-MEM medium, mixed with the relevant plasmids, and placed on ice for 20 min in 4 mm cuvettes. Cell electroporation was performed using Gene Pulser XcellTM electroporation system (Bio-Rad, Hercules, CA, USA). electroporation protocol: voltage - 250V, capacitance 900 µF, resistance $\infty \Omega$. Immediately following pulsation, cells were washed with pre-warmed DMEM media, centrifuged, and plated in culture plates.

Production of Lentiviral particles

8×10^6 HEK-293FT cells were plated on a 10 cm plate pre-coated with 200 $\mu\text{g/mL}$ poly-L-lysine and left to adhere overnight and reach a confluence of $\sim 80\%$. pLVX plasmids containing either receptor of interest tagged with a fluorescent protein or wasabi/tdTomato control under an EF1 promoter were mixed with psPAX2 (a gift from Didier Trono, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland; Addgene plasmid 12260) and pCMV-VSV-G (a gift from Bob Weinberg, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA; Addgene plasmid 8454) at a molar ratio of 3:2:1, and cells were transfected using jetOPTIMUS reagent (Polyplus Transfection). After 24 hours, the medium was replaced with complete DMEM supplemented with 0.075% sodium bicarbonate (Biological Industries). Virus-containing medium was harvested after 24 hours and 48 hours. Following collection, the supernatant was passed through a $0.45\mu\text{m}$ membrane filter to rid it of cellular debris. In order to produce high viral titer stock, viral-containing media was concentrated via centrifugation in Amicon Ultra-15 centrifugal filter unit (100KDa) (Sigma Aldrich, St. Louis, MO, USA).

Tumor cell lines viral infection

For transduction of tumor cells, cells were incubated with viruses and 100 $\mu\text{g/mL}$ polybrene (Sigma Aldrich) for 30 minutes followed by 30 minutes of centrifugation before medium was replaced. Following three days, cells that expressed HER2^+ /tdTomato were sorted by FACS Aria II.

Mouse IgG and IgM purification

Mouse antibodies were obtained from pooled 5 mL mouse blood obtained from Inferior Vena Cava. Whole blood was left on ice for 20 minutes to allow blood to clot. Blood was then centrifuged at 600 RCF for 10 minutes and serum was collected and re-centrifuged at 20,000 RCF for an additional 10 minutes. Serum was filtered through $0.1\mu\text{M}$ and total the IgG and IgM were purified using protein-G and 2-mercaptopyridine columns, respectively (GE Healthcare, Chicago, IL, USA). The levels of purified IgG and IgM were measured with specific ELISA kits (Bethyl, Montgomery, TX, USA) according to manufacturer's instructions.

Antibody-tumor lysate immune complexes (Ig-IC) and antibody-bound tumor cells

When obtained from surgical resections, tumor cells were initially isolated after enzymatic digestion and sorted as $\text{FSC}^{\text{hi}}/\text{CD45}^{\text{neg}}$ cells prior to their fixation and staining. For tumor-antibody complexes, tumor cells were fixed in 2% paraformaldehyde, washed extensively, and incubated with 2-5 μg allogeneic IgG per 1×10^5 tumor cells, and then washed to remove excess antibodies. DC activation with the above Ig-IC was repeated in at least 5 independent experiments in biological duplicates.

In vivo tumor vaccination models

For tumor recurrence studies, 2×10^5 tumor cells were injected s.c. above the right flank, and the size of growing tumors was measured using calipers. When tumors reached 16-25 mm^2 for B16F10 tumors, mice were anesthetized, and visible macroscopic tumor was surgically removed. Resected tumors were enzymatically digested with 0.1mg/mL of DNase I (Sigma Aldrich) and 5mg/mL collagenase IV (Sigma Aldrich) in HBSS. Cells were then fixed in 2% paraformaldehyde for 10 min, washed extensively in PBS, and coated for 30 min with syngeneic or allogeneic antibodies. Antibody-coated tumor cells were then washed and injected into tumor-resected mice (2×10^6 per mouse) or were added to DC cultures. After overnight incubation, DC were washed, and 2.5×10^6 were injected s.c. into tumor-resected mice one day after the tumors were removed, adjacent to the site of tumor resection. For prophylactic vaccination assays, blood, and tumor DC were incubated overnight with B16F10-IC, DC were then washed and 6×10^6 cells per mouse were

injected s.c. After 5 days 2.5×10^4 B16F10 cells were injected s.c. above the right flank and tumor growth was monitored. Experiments were repeated independently at least 3 times with 4-5 mice per group.

For treatment of established melanoma, C57Bl/6 female mice bearing approximately 20 mm² B16F10 tumors were injected s.c. with 2×10^6 BMDC transfected with modified FcγRI and with i.p. injection 250 μg per mouse of TA99 antibody (BioXcell, Lebanon, NH, USA). Mice were treated twice, one week apart and tumor size was measured using a caliper twice a week.

Kinase phosphorylation measurements

For measurement of kinase phosphorylation by FACS, 1×10^5 mouse MoDC were generated from C57Bl/6 bone marrow. Thereafter, 1×10^5 B16F10 tumor-IC were added to DC cultures for 1 min and 15 min Fluorochrome-conjugated antibodies against phospho-p38 (Thr180/Tyr182), phospho-JNK (Ser63), phospho-ERK1/2 (p44) (pT202/pY204) and phospho-Akt (pY473) were purchased from Cell Signaling Technologies (Danvers, MA, USA). DC protein phosphorylation experiments were repeated 2-4 times.

NO and Granzyme B measurements

BMDC were activated with IC composed of B16F10 cells coated with either IgG or IgM isolated from the 129S1 mice. GrB was measured using mouse granzyme B ELISA kit (Abcam, Cambridge, UK) and NO was measured using Griess reagent system (Promega, Madison, WI, USA) according to manufacturer's protocol.

Mass Spectrometry

Sample preparation

DC 2.4 were transfected with either αCD19 scFv-GFP or GFP plasmids using Lipofectamine™ 2000 transfection reagent. 4-6 hours post-transfection, when scFv expression was at its maximal level during the expression window (determined by prior calibration experiments), cells were harvested and sorted based on GFP expression using BD FACSAria™ III Cell Sorter. GFP-positive cells collected were resuspended in cell lysis buffer based on PBS + protease inhibitor cocktail (NEB#5871) and lysis was performed using cell sonication (Sonics Vibra Cell VCX 130 Digital Ultrasonic Processor). Sonication protocol: time: 100sec, pulse on: 10sec, pulse off: 10sec, amplitude 25%.

Protein complex isolation

Immunoprecipitation was done using GFP-trap magnetic agarose beads (chromotek, Planegg, Germany) according to manufacturer protocol.

Mass spectrometry – (this part of the experiment was performed through a collaboration with Prof. Tami Gieger's lab, Tel Aviv University)

Sample preparation

Samples were washed 6 times with washing buffer containing 150mM NaCl and 50mM Tris-HCL (pH 7.5). In every wash, samples were gently rotated in an automatic rotator for 1 minute. Then samples were collected using two rounds of 50ul elution buffer. First samples were incubated in an elution buffer containing fresh 2M urea, 50mM Tris-HCL (pH7.5), and 1mM DTT for 2 hours at room temperature and then collected in a new tube. Second, on the left beads, a second elution buffer containing 2M urea, 50mM Tris-HCL (pH7.5), and 5mM Choloroacetmide was added for 10 minutes and collected the liquid to the same tube as collected with the first elution and wait 30

minutes. Then we added 5ug/ml trypsin (Promega) for overnight incubation. Samples were collected in PCR tubes, vacuum dried, and re-suspended in 5ul buffer containing 2% ACN and 0.1% trifluoroacetic acid (TFA).

Lc ms

LC-MS/MS runs were performed on the EASY-nLC1000 UHPLC (ThermoFisher Scientific) coupled to Q-Exactive HF mass spectrometers (ThermoFisher Scientific). Peptides were separated with a 75uM X 50 cm EASY-spray column (ThermoFisher Scientific) using a water-acetonitrile gradient of 120 minutes, with a flow rate of 300 ml/min at 40°C. The resolutions of the MS and MS/MS spectra were 60,000 and 15,000 respectively. MS raw files of all samples were jointly analyzed by MaxQuant 2 version 1.6.2.6. MS/MS spectra were referenced to the Uniprot Mus musculus proteome ().

Statistical analysis

Analyses were performed using the Perseus software 3 version 1.6.2.3.. For the student's t-test results we use a cut-off of FDR < 0.1, S0 = 0.1.

Rt pcr

RAW 264.7 cells were transfected with either αCD19 scFv – GFP, GFP control, or additional control containing α chain of FcγRI receptor (CD64) attached to GFP. Transfections were done using JetOPTIMUS according to manufacturer's protocol (Polyplus Transfection). 24 hours post-transfection, cells were collected from growth plate using trypsin (Promega). Cells lysis was performed and RNA was purified using RNA NucleSpin RNA manufacturer isolation protocol (Macherey-Nagel, Dueren, Germany). To evaluate RNA sample quality, gel electrophoresis was performed to detect ribosomal RNA integrity. RNA concentration and purity were tested using NanoDrop One/One^C Microvolume UV-Vis Spectrophotometer (ThermoFisher Scientific). – all RNA samples used were at purity levels 260/280 > 2.0, 260/230 between 2.0-2.2. cDNA was synthesized using qScript cDNA Synthesis Kit (Quantabio, Beverly, MA, USA)

GFP	Forward primer	AAGTTCAGCGTGTCGGCGGA
	Reverse primer	AAGCACTGCACGCCGTAGGT
Neomycin	Forward primer	TGAAGCGGGAAGGGACTGGC
	Reverse primer	CGAATGGGCAGGTAGCCGGA
Mus musculus β-actin	Forward primer	GTCCACCTTCCAGCAGATGT
	Reverse primer	GCTCAGTAACAGTCCGCT

Quantitative Real-time PCR was conducted using PerfeCTa SYBR[®] Green FastMix (Quantabio) using stepOne[™] Real-Time PCR system. All experiments were done in triplicates.

mRNA synthesis

For in vitro capped RNA synthesis pGEM4Z plasmid (kindly gifted from Prof. Eli Gilboa, University of Miami health system) containing T7 promoter along with insert αCD19 scFv – GFP or GFP for control was used as template.

For in vivo assay, we used βGlobin5'UTR (kindly gifted from Dr. Gal Cafri, Sheba medical center, Israel)

Synthesis was performed using AmpliCap-Max[™] T7 High Yield Message Maker Kit (Cellscript, Maddison, WI, USA). The synthesis reaction product was tested for integrity via gel electrophoresis. mRNA was purified using NucleSpin RNA.

mRNA Transfection

5.0×10^4 RAW264.7 macrophages were plated on 24-well plate a day before the transfection. Synthesized mRNAs were transfected into RAW264.7 using jetMESSENGER RNA transfection kit (Polyplus Transfection) according to the manufacturer's protocol. Transfected cells were analyzed 16-24 hours' post-transfection.

For in vivo experiments, bone marrow cells were isolated from naïve 12 weeks CD45.1 female mice and incubated for seven days with GM-CSF and IL-4. 2.5×10^6 BMDC were collected in 200 μ L OptiMEM and electroporated in 0.2 cuvettes using BioRad GenePulsar Xcell with 5 μ g mRNA and 400V square wave for one millisecond, as previously described (24 [24](#)).

T-cell proliferation assay

RAW 264.7 cells were transfected with OVA constructs using LipofectamineTM 2000. Splenic T cells were isolated from OTI and OTII mice using Ficoll-paque plus density gradient (BD Biosciences, San Jose, CA, USA) followed by incubation with biotin-conjugated anti-CD8⁺ and anti-CD4⁺ magnetic beads and magnetic isolation using streptavidin-magnetic beads (BioLegend, Carlsbad, CA, USA). T cells were then labeled with CFSE, washed extensively three times in complete media, and co-cultured at ratios of 1:5 and 1:10 with transfected RAW264.7 cells. After three days, CFSE dilution in T cells was determined by CytoFLEX5.

Confocal microscopy

For confocal microscopy imaging, cells were cultured on glass-bottom confocal plates (Cellvis, Mountain View, CA, USA). Cells were live-imaged or fixed and permeabilized with 2% PFA for 20 minutes. Fixed cultures were washed twice with PBS. For immunofluorescence, fixed cultures were blocked overnight with 5% BSA (ThermoFisher Scientific) and stained with 1:100 or 1:200 diluted primary antibodies. Cells were then washed with PBS containing 1% BSA and stained with a secondary antibody, diluted 1:100 or 1:200. Stains used included: Hoechst 33342 (ThermoFisher Scientific). ER staining kit - Cytopainter (Abcam). Anti-GPR78 BiP antibody (Abcam), G3BP1 antibody (mab#17798, Cell signaling technologies), and anti-mouse CD64 (Fc γ RI) (Biolegend). DAPI (Abcam). All specimens were imaged by ZEISS LSM800 (ZEISS) and analysed by ZEN 2.3 (ZEISS) and ImageJ.

Structured illumination microscopy

μ -Slide 8 well High Grid 500 (ibidi - Munich, Germany) were pre-coated with trastuzumab (Roche) and left to incubate overnight at 4 °C then washed thrice with PBS. transfected RAW 264.7 cells were cocultured with 4T1 cell line expressing human HER2. Immunological synapse was visualized using 3i Marianas Spinning Disk Confocal (Denver, CO, USA). For Granzyme B detection, cells were fixated using PFA 4% and washed to allow permeabilization and stained using Alexa Fluor® 647 anti-human/mouse Granzyme B Antibody (Biolegend 515405).

Flow cytometry

Flow cytometry was performed on CytoFLEX5 (Beckman Coulter, Brea, CA, USA) using anti-mouse CD64 (Fc γ RI) (clone: X54-5/7.1), anti-mouse CD3 (clone: 17-A2) anti-mouse CD45 (clone: 30-F11), Annexin V, and propidium iodide (PI) (all purchased from Biolegend). Data sets were analyzed using FlowJo software (Tree Star, Inc.).

Killing assay

For measuring tumor cell killing, 10^4 4T1 H2B-tdTomato, which are genetically modified to express human EGFR or HER2 on cancer cell membranes, were cultured in 96 well plates for 2 hours to adhere. Transfected myeloid cells were added at E:T ratios of 4:1 and 8:1, with or without 10 μ g/mL

of Trastuzumab (Roche, Basel, Switzerland), Cetuximab (Merck, Darmstadt, Germany), and Rituximab (Genentech, San Francisco, CA, USA). Cells were imaged by IncuCyte S3 imager (Sartorius, Göttingen, Germany), and the numbers of target cells were calculated by incuCyte software. In some experiments, cells were stained with Annexin V and propidium iodide (PI) (Biolegend) according to manufacturer's instructions, and specific tumor cell lysis was measured by CytoFLEX5.

Constructs subunits and design process

Kits and materials used for cloning

NucleoSpin Plasmid easy pure (Macherey-Nagel), NucleoSpin Gel and PCR clean-up (Macherey-Nagel), NucleoBond Xtra Maxi (Macherey-Nagel), Zymoclean Gel DNA recovery kit (Zymo Research, Irvine, CA, USA). **Cloning:** DH5α Competent cells (ThermoFisher scientific) were grown in LB broth with agar (Miller) (Sigma Aldrich) or LB broth (Miller) (Merck) supplemented with Ampicillin, sodium salt or Kanamycin sulfate Bio Basic, Markham, Ontario, Canada). **Enzymes and reagents:** Restriction enzymes, T4 DNA ligase, KLD enzyme mix, and Phusion High Fidelity DNA polymerase were purchased from New England Biolabs (Ipswich, MA, USA), Taq DNA polymerase mix (PCRBIO, Wayne, PA, USA), RepliQa HiFi Assembly Mix (Quantabio). **Gel electrophoresis:** SeaKem LE agarose (Lonza, Basel, Switzerland), Ethidium Bromide (Merck), TAE buffer (Sartorius, Göttingen, Germany)

Statistical analyses

All statistical analyses and graphs were performed by GraphPad Prism 9. Each experimental group consisted of at least three mice and repeated at least three independent times. Significance of results was determined using the nonparametric one-way ANOVA with Tukey's and Sidak correction, when multiple groups are analyzed, two-way ANOVA with Tukey's and Sidak correction for multiple comparisons.

Illustrations

Experimental illustration shown in figures were created using <https://BioRender.com> 

Results

IgM-induced signaling elicits cytotoxic response in macrophages and can be integrated to a CAR design

We have previously demonstrated that MHC-matched allogeneic tumors, which spontaneously regressed, are coated with IgG and IgM antibodies soon after tumor initiation and are found in proximity to tumor-infiltrating myeloid cells (25 ). While we characterized the therapeutic role of tumor-binding IgG and their interactions with dendritic cells (DC) (26 ), the role of IgM in facilitating DC-mediated immunity has remained unclear. To address that, we initially compared their capacity to induce tumor immunity in a prophylactic tumor immunization assay. To this end, we incubated monocyte-derived dendritic cells (MoDC) with immune complexes (IC) composed of B16F10 tumor cells coated with allogenic IgG or IgM and injected them subcutaneously (s.c) to syngeneic mice (illustrated in Fig. 1A ). After two rounds of immunization, five days apart, mice were challenged with B16F10 cells and tumor size was monitored over time. We found that incubation of MoDC with IgG-IC, but not with IgM-IC, induced T cell immunity and prevented tumor growth (Fig. 1B ). We noticed however, that incubation *in vitro* of MoDC with tumor cells coated with allogenic IgM but not with IgG, resulted in a massive killing of the tumor cells (Fig. 1C ). To assess the underlying killing mechanisms, we measured the levels of nitric oxide (NO)

Construct subunits design	Construct preparation
(ssCD8 α) – TA99 scFv – CH1 – hinge – CH2 – CH3 – transmembrane + intracellular portion of CD209/ C5aR – mCherry	All subunits were designed and designed and ordered in gBlock format from Integrated DNA Technologies (IDT) (Coralville, IA, USA). Subunits were added to pcDNA 3.1 (+) using Gibson assembly, and cloned to pmCherry- N1.
(ssCD8 α) – TA99 scFv – CH3 – transmembrane + intracellular portion of CD209/ C5aR- mCherry	Construct was created by elimination of CH1- hinge-CH2 subunits through invert PCR followed by kinase, ligase, DpnI (KLD) enzyme mix.
pLVX: TA99 scFv – CH3 – (TM+IC) C5aR - mCherry	pLVX-IRES-Hyg vector was restricted using SpeI/NotI. Receptor insert including TA99 scFv - heavy chain - C5aR (transmembrane + intracellular) - mCherry was restricted from mCherry vector via restriction enzymes NheI/NotI followed by ligation using 4T DNA ligase.
(ssCD8 α) – CH1 – hinge – CH2 – CH3 – transmembrane + intracellular portion of CD209/ C5aR- mCherry	Constant heavy sequence was isolated and amplified using invert PCR for scFv removal followed by KLD enzymes.
(ssCD8 α) – TA99 scFv - transmembrane and intracellular portion of either CD209 or C5aR - mCherry	Removal of constant heavy subunits was done by invert PCR amplification. Followed by KLD enzyme reaction.
(ssCD8 α) – TA99 scFv – CD8 α hinge – CD8 α transmembrane – mCherry	Extracellular portion was isolated via restriction with BamHI/EcoRI and inserted into pmCherryN1 vector followed by ligation with 4T DNA ligase.
(ssCD8 α) – CD8 α hinge - a transmembrane and intracellular portion of either CD209 or C5aR - mCherry	scFv portion was removed via invert PCR using Fw primer that included a tail of CD8 α hinge sequence as an extracellular remnant. The PCR product was used in KLD enzymes mix reaction.
TA99 scFv - GFP	TA99 scFv sequence was restricted from pemCherry-N1 plasmid using EcoRI/XhoI enzymes and inserted to peGFP-N1 plasmid linearized with same enzymes.
GFP - T2A - TA99 scFv - mCherry	T2A sequence was designed and ordered in linear gBlock sequence, flanked by restriction enzymes: BglII/NheI. The sequence was inserted to peGFP-C1 vector. GFP-T2A sequence was isolated via restriction with NheI/XhoI and inserted to plasmid

Table 1.

scFv construct design and preparation process

	which included an insert of TA99 scFv – mCherry.
GFP – TA99 scFv – mCherry	Removal of T2A sequence was performed by invert PCR followed by KLD enzyme mix reaction.
(ssCD8 α) – α CD19 scFv – CD8 α hinge – CD8 α transmembrane – mCherry	(ssCD8 α) – TA99 scFv – CD8 α hinge – CD8 α transmembrane – mCherry was used as backbone. TA99 was removed by invert PCR amplification exclusion of α CD19 scFv was amplified using plasmid pHR-PGK-antiCD19-synNotch-GalVP64 (plasmid#79125) which was purchased from Addgene (Watertown, MA, USA). Both backbone and insert were flanked with XhoI/EcoRI.
α CD19 scFv - GFP	α CD19 scFv from plasmid#79125 (Addgene) was amplified using primers with tails encoding restriction sites for XhoI/EcoRI enzymes. peGFP-N1 backbone was restricted using same enzymes.
pGEM4Z: T7 promoter - α CD19 scFv – GFP – polyA tail	α CD19 scFv was cloned into pGEM4Z using restriction enzymes NheI + NotI. pGEM4Z GFP was restricted using: XbaI + NheI.
pLVX: α CD19 scFv - GFP	α CD19 – GFP was isolated from peGFP-N1 backbone using enzymes NheI/NotI and inserted into pLVX backbone linearized using SpeI/NotI.
GFP - α CD19 scFv	α CD19 scFv sequence was isolated and amplified using primers which included overhang sequence of restriction sites for enzymes XhoI/EcoRI. peGFPC1 was restricted with same enzymes followed by ligation.
TA99 variable heavy chain-GFP	Variable heavy chain was isolated and amplified using invert PCR for variable light chain removal followed by KLD enzyme mix.
TA99 variable light chain-GFP	Variable light chain was isolated and amplified using invert PCR for variable heavy chain removal followed by KLD enzyme mix.
α CD19 variable heavy chain-GFP	Variable heavy chain was isolated and amplified using invert PCR for variable light chain removal followed by KLD enzyme mix.
α CD19 variable light chain-GFP	Variable light chain was isolated and amplified using invert PCR for variable heavy chain removal followed by KLD enzyme mix.
α CD19 mutated VL (linear)	G-block of mutated α CD19 variable light chain.

Table 1. (continued)

	Two-point mutations included replacement of Cysteine amino acid with glycine in positions 24 and 89, flanked by restriction enzymes XhoI/EcoRI was designed and orders from IDT. PeGFP-N1 plasmid was restricted with same enzymes.
α CD19 mutated VH (linear)	gBlock of mutated α CD19 variable heavy chain. Two-point mutations included replacement of Cysteine amino acid with glycine in positions 22 and 95, flanked by restriction enzymes XhoI/EcoRI was designed and orders from IDT. PeGFP-N1 plasmid was restricted with same enzymes.
Immunoglobulin loop fragment GFP	DNA sequence was ordered in linear gBlock from IDT. Insert of interest was flanked by XhoI/EcoRI and inserted into peGFP-N1 backbone linearized with both enzymes. Final amino acid sequence included: MVTISCRASQDISKYL-GGGGSGGGGSGGGGS-EQEDIATYFCQGN
(ssCD8 α) – CD64 – transmembrane + intracellular portion of C5 α /CD209/ μ – GFP	CD64 sequence was purchased in gBlock format from IDT, subunits were added in sequence keeping reading frame in check using homology sequences into peGFP-N1.
(ssCD8 α) – CD64 – transmembrane + intracellular portion of C5 α R/CD209/ μ – T2A - GFP	T2A sequence was ordered in linearized DNA gBlock from IDT. Sequence was flanked by homology sequence which overlapped linearized plasmid ends created by restriction with enzymes BamHI/XmaI.
pLVX: (ssCD8 α) – CD64 – transmembrane + intracellular portion of C5 α R/CD209 – T2A GFP	Receptor sequence including all subunits - GFP was isolated from peGFPN1 plasmid using restriction enzymes NheI/NotI. pLVX backbone was linearized using SpeI/NotI.
CD64 (extracellular + transmembrane) – GFP (used for RT PCR expression control)	Transmembrane + intracellular sequences were removed from (ssCD8 α) – CD64 – transmembrane + intracellular sequence – GFP using invert PCR. Followed by KLD enzyme's reaction.
Alpha mu in β Globin5'UTR	Plasmid was ordered with insert in place through A2S technologies Ltd (Yavne, Israel)
anti TRP1 murine TCR	Minigene was cloned to pcDNA3-EGFP backbone plasmid using restriction enzymes.
TA99 scFv – OVA ₂₅₇₋₂₆₄ – G4S linker -	DNA sequence including all subunits was
OVA ₃₂₃₋₃₃₉	designed and ordered in gBlock via IDT. Sequence was inserted into pcDNA 3.1 (+) using restriction enzymes.
OVA ₂₅₇₋₂₆₄ – G4S linker - OVA ₃₂₃₋₃₃₉	Control construct to scFv-OVA was prepared by removal of scFv sequence using invert PCR amplification followed by a KLD reaction according to protocol.

Table 1. (continued)

and granzyme B (GrB) in the supernatants of overnight cultures. Consistent with their killing rates, significantly higher levels of both GrB and NO were detected upon incubation with allogenic IgM-IC (**Fig. 1D**), indicating that both secretion of reactive oxygen species and lysosome deposition are involved in the killing. To further assess what signaling cascade is induced by IgM-IC, we tested the levels of major inflammatory cytokines and phosphorylated enzymes. Activation with IgG-IC induced classical signaling through Fcγ receptors, characterized by high levels of TNFα and IL12, and activation of enzymes in the MAP kinase pathway (**Fig. 1E–1F**). The phenotype of IgM-IC activated MoDC was more complicated and characterized by low levels of TNFα and high IL-12, suggesting that Akt was only partially phosphorylated. Flow cytometric analysis indicated only neglectable levels of Akt phosphorylation, while ERK and p38 phosphorylation was comparable to that induced by IgG-IC along with higher levels of phospho-JNK (**Fig. 1E–1F**). This somewhat unique signaling cascade most likely indicates a combination of several signaling chains recruited by IgM, including Fcγ and C1q receptors.

To harness this signaling cascades to induce tumor cytotoxicity, we designed Chimeric Antigen Receptors (CAR) in which the antigen recognition region, a single chain fragment variable (scFv) derived from TA99 antibody, targets the melanoma antigen gp75, as previously reported (27). The scFv, consisting of Variable light (V_L) and variable heavy (V_H) chains was attached to a spacer region of Constant Heavy subunits 1-3 (CH_{1-3}). These subunits were fused to several potential IgM signaling chains in the transmembrane and intracellular regions and fused to a fluorescence tag to aid detection of expression (**Fig. 1G**). Next, we transfected HEK 293FT cells with the CAR constructs to test their expressions and cellular localization using confocal microscopy and flow cytometry. Indeed, all constructs were expressed on cell membrane within 24 hours (**Fig. 1H–1I**).

scFv is not expressed by myeloid cells

Next, we sought to test the construct's expressions and functionality in myeloid cells. To this end, we transfected two established murine cell lines of dendritic cells (DC2.4) and macrophages (RAW 264.7) with our three constructs and with mCherry as a control plasmid. In both cell lines, about 30 percent of the cells expressed mCherry. In sharp contrast, however, we were unable to detect any construct expression in myeloid cells by either confocal microscopy or flow cytometric analysis (**Fig. 2A–2C** and Supplementary Fig. S1A). To test if this phenomenon would also occur in human cells, we transfected THP-1 monocyte cell line in comparison to Jurkat T cell line. Consistent with our results using mouse cells, we could not detect any CAR expression in THP-1 cells, while in Jurkat cells their expression was comparable to mCherry transfection rates (**Fig. 2D–2E**).

Since we could not detect expression of our constructs in myeloid cells, we sought to characterize the subunit leading to inhibition of expression by generating various constructs, in which different segments of CAR are omitted. Since myeloid cells were reported to express TRIM21 E3 ligases that bind C_H1 and C_H2 fragments of internalized IgG (28), we next generated a CAR construct with a minimal spacer region by elimination of Constant Heavy domain 1 and 2 (C_H1 , C_H2) and tested the expression of these constructs in DC2.4 cell line. However, consistent with our previous results, we were unable to detect any protein expression (Supplementary Fig. S1B–S1C). We then removed the rest of the spacers, generating a construct consisting only of the scFv region and an intracellular signaling domain. As with our previous finding, no expression in myeloid cell lines was detected (**Fig. 2F–2G**). Similarly, removal of the intracellular portion did not alter the expression patterns of TA99-derived scFv, suggesting that the scFv domains may be the element that prevents its expression (**Fig. 2G–2H**). To verify that these expression patterns do not stem from the use of mCherry, we have also fused these constructs to GFP. Nonetheless, in both cases, the presence of scFv prevented protein expression in macrophages and DC (Supplementary Fig. S1C).




Figure 1

IgM-induced signaling elicits cytotoxic response in macrophages and can be integrated to a CAR design.

(A) Illustration of experimental setting¹. (B) BL6F10 tumor size (mm²) in mice following prophylactic immunization with MoDC pulsed with tumor cells coated with allogeneic IgG or IgM (n=4). (C) Mean percentages of BL6F10 melanoma cells stained for Annexin V/PI incubated with allogeneic IgG and IgM following incubation with MoDC (n=5). (D-E) Mean levels of Granzyme B (GrB) and proinflammatory cytokines (TNF- α , IL-12) secretion (ng/ml) in the supernatants of MoDC following overnight activation with IgG and IgM immune complexes. (n=5). (F) Mean fluorescent intensity (MFI) of MAPK enzymes in MoDC following activation for 20 min with IgG and IgM tumor immune complexes (n=5). (G) Illustration representing CAR-macrophage design¹. (H) Confocal microscopy images of HEK293FT cells 24 hours post-transfection with CAR plasmids and membranous wasabi. (I) Representative FACS analysis of HEK293FT cells 24 hours post-transfection with CAR plasmids. Results are from one representative experiment out of at least three performed. Statistical significance was calculated using non-parametric t-test (*** denote p<0.001, **** denote p<0.0001). ¹Illustrations were created using




Figure 2

scFvis not expressed by myeloid cells.

(A) Confocal microscopy images of DC 2.4 cells 24 hours post-transfection with CAR plasmids and membranous wasabi. (B) Representative FACS analysis of DC 2.4 cells 24 hours post-transfection with CAR plasmids. (C) Percentages of transfected cells 24 hours following transfection (n=4). (D) Confocal microscopy images of THP-1 cells 72 hours post-lentiviral infection with CAR-C5u-mCherry and tdTomato plasmids. (E) Percentages of transfected human cell lines 72 hours following transduction (n=4). (F-G) Representative confocal microscopy (F) and mean percentages (G) of cells expressing chimeric molecules 24 hours after transfection. (H-I) Representative confocal microscopy (H) and mean percentages of cells (I) expressing chimeric molecules 24 hours following transfection (n=4). Results are from one representative experiment out of at least three performed. Statistical significance was calculated using non-parametric t-test (*** denote p<0.001, **** denote p<0.0001). Illustrations were created using BioRender.com

To test if this phenomenon is restricted to TA99 CAR or reflects a broader phenomenon, we sought to compare TA99 CAR to a well-established α CD19-derived scFv. Towards this end, we generated a plasmid that contains the CD8 signal peptide followed by α CD19-derived scFv and fused to CD8 spacer and transmembrane portion, and an identical one that lacks the scFv part (**Fig. 2H** [↗](#)). Interestingly, while we detected low levels of α CD19-derived scFv, they were localized in intracellular compartments and not on the cell membrane (supplementary Fig. S1D). Removal of scFv domain enabled expression of the inserted protein in the Golgi and cell membrane of both macrophages and DC cells (**Fig. 2H** [↗](#)-**2I** [↗](#) and Supplementary Fig. S1E).

Both VH and VL domains prevent the expression of scFv in myeloid cells

We next aimed to test which portion of scFv prevents its membranal expression. Initially, we designed two plasmids in which all other signaling subunits were removed, leaving only the TA99- or α CD19-derived scFv fragment. As with our previous findings, here too we were not able to detect intracellular TA99- nor α CD19-derived scFv expression (**Fig. 3A** [↗](#)-**3B** [↗](#)). To test the possibility that the location of the fluorescent tag inhibits scFv expression, we transfected DC2.4 with a GFP fused to the C-terminus of α CD19-derived scFv. Similarly, this protein did not lead to scFv expression (Supplementary Fig. S2A). Next, we generated plasmids consisting of minimal subunits, consisting of only the variable light or variable heavy domains of TA99 and α CD19. While expression levels were somewhat higher in these constructs, compared to the full length of scFv, they were still benign and significantly lower compared to GFP only (**Fig. 3C** [↗](#)-**3E** [↗](#)).

As a result, we sought to further investigate whether the primary protein structure of amino acid sequence or the tertiary immunoglobulin structure of the scFv subunits prevent the expression. To this end we inserted point mutations that switched cysteine amino acids to glycine, creating a linear protein by preventing disulfide bonds. Interestingly, linearization of α CD19 variable light chain (or heavy chain) had only a marginal effect on its expression patterns and resulted in reduced and compartmentalized expressions (**Fig. 3F** [↗](#)-**3H** [↗](#)). In addition, we generated a plasmid that consisted of the first and last 15 amino acids of variable light chain which includes the cysteine amino acids for disulfide bond formation. This fragment was almost inert, and its expression patterns were comparable to that of GFP only (Supplementary Fig. S2B). We also tested whether specific fragments within the variable light chain sequence prevented its expression. Dividing the light chains of α CD19 scFv into three sequences resulted in a high expression in myeloid cells, albeit tending to induce different expression patterns of cellular compartmentalization (**Figure 3G** [↗](#)-**3H** [↗](#)). Taken jointly, our results suggest that the linear form of either the variable light or the heavy chain of antibodies results in a poor and compartmentalized protein expression compared to the full length of ScFv to a lesser extent.

scFv induce ER stress in myeloid cells

To elucidate the level at which scFv constructs are silenced, we sought to transfect cells of myeloid lineage with mRNA composed of α CD19 scFv fused to a green fluorescent protein tag. Nonetheless, this form of transfection did not lead to expression of scFv in myeloid cells (**Fig 4A** [↗](#)). Additionally, we measured the mRNA levels of GFP mRNA in mouse myeloid cells following transfection with α CD19 scFv-GFP vectors, in comparison to GFP only or to Fc receptor fused to GFP (which are robustly expressed in these cells). Similar levels of GFP mRNA were detected in all transfected cells (**Fig 4B** [↗](#)). Based on these findings we speculate that the negative regulation on scFv expression is not evident at the mRNA level.

To further assess this possibility, we utilized 2A ribosomal skipping peptide in order to test if scFv is regulated at the post-translational level. Hence, we generated plasmids containing GFP fluorescence tag followed by T2A skipping peptide, succeeded with TA99 scFv fused to mCherry, all in one reading frame. This design allowed us to test if the GFP and scFv expression are coupled, thus indicating that the regulation is made on the mRNA transcript and prior to protein




Figure 3

Both VH and VL domains prevent expression of ScFv in myeloid cells.

(A) Confocal microscopy images of DC 2.4 cells 24 hours post transfection with uCD19-scFv GFP plasmid.¹ (B) Geometric mean of GFP positive cells 24hr post transfection with different uCD19- and TA99-ScFv GFP constructs in DC2.4. (n=3) (C) Confocal microscopy images of HEK 293FT and DC 2.4 cells 24 hours post transfection with uCD19-variable light chain GFP plasmid.¹ (D) Confocal microscopy images of HEK 293FT and DC 2.4 cells 24 hours post transfection with uCD19-variable heavy chain GFP plasmid.¹ (E) Mean percentages of cells expressing scFv fragments 24 h post transfection (n=). (F) left: illustration of mutated variable light chain. right: confocal microscopy images of HEK 293FT and DC 2.4 cells 24 hours post transfection with uCD19-mutated (linear) variable light chain. (G) Representative confocal images of HEK293FT and DC2.4 cells 24 hours post transfection with 1/3 fragments of uCD19-variable light chain GFP plasmid (H) Mean percent ages of GFP positive cells 24hr post transfection with different fragments of scFv-GFP in DC2.4 and RAW264.7 (n=4). Results are from one representative experiment out of at least three performed. Statistical significance was calculated using non-parametric t test (***) denote p<0.001, **** denote p<0.0001). ¹Illustrations were created using BioRender.com




Figure 4

scFv fragments induce ER stress in myeloid cells.

(A) Confocal microscopy imaging of RAW 264.7 24hr post-transfection with linear mRNA vectors translating to GFP and uCD19-scFv GFP. **(B)** qPCR data showing relative mRNA levels in RAW 264.7 transfected with GFP, Fe receptor-GFP, and uCD19-ScFv GFP. (n=3) **(C)** Upper: illustration of plasmid subunits. Lower: confocal microscopy imaging of HEK293FT and RAW 264.7 24hr post-transfection with T2A ribosomal skipping plasmid including ScFv. **(D)** Upper: Illustration of plasmid subunits. Lower: confocal microscopy imaging of HEK293FT and RAW 264.7 24hr post-transfection with plasmid containing no T2A. **(E)** Confocal microscopy images of RAW264.7 cells at 6 hours and 24 hours post-transfection with uCD19-ScFv GFP plasmid. **(F)** Volcano plot showing differentially expressed proteins uCD19 ScFv GFP/GFP in DC 2.4 cells. **(G)** Confocal microscopy images of DC2.4 stained with an ER stain, 24 hours post-transfection with GFP, membranous TA99-ScFv GFP. **(H)** Confocal microscopy images of DC2.4 stained for BiP 24 hours post-transfection. **(I)** Mean levels of phospho-JNK 6 hours following transfection. **(J)** Percentage of cells expressing GFP or TRP-TCR 24 hours following transfection. Results are from one representative experiment out of at least three performed. Statistical significance was calculated using non-parametric t-test (***)

translation. Alternatively, expression of one fluorescent tag would indicate that the regulation is made at the protein level following translation. While we detected both tags (GFP, mCherry) in HEK293 cells, only GFP was expressed in myeloid cells, but not the mCherry which was fused to scFv (**Fig. 4C**). Eliminating the T2A skipping peptide from this construct, generating a plasmid containing GFP tag fused to TA99 scFv succeeded by mCherry, resulted in abrogated and reduced expression levels of both fluorescent tags in myeloid cells (**Fig. 4D**). These results therefore suggest that scFv expression is regulated on the protein level.

Consistent with this notion, we found that scFv is expressed in myeloid cell at earlier time points, starting at four hours post-transfection and peaking at about 6 hours (**Fig. 4E**). In order to determine the mechanism through which scFv is degraded, we next performed proteomics of immunoprecipitated scFv. Hence, macrophages were transfected with GFP only, or GFP fused to α CD19 scFv. After 6 hours, GFP-expressing cells were sorted and lysed, and associated proteins were pulled down using anti-GFP conjugated beads. Eluted proteins were then sent to analyses by mass spectrometry (illustrated in Supplementary Fig.S3A). Analysis of protein interactors with scFv in myeloid cells indicated that scFv fraction was statistically enriched with proteins associated with ER stress that could promote scFv degradation (**Fig. 4F**). To further corroborate that possibility, we assessed the co-localization of scFv with key proteins in this pathway. Confocal microscopy corroborated that the majority of α CD19 scFv were localized at the endoplasmic reticulum (**Fig. 4G**) and co-localized with BIP (**Fig. 4H**) and G3BP1(Supplementary Fig.S3B), which are master regulators of ER stress. Lastly, flow cytometric analysis indicated higher levels of phospho-JNK (**Fig. 4I**). Next, we tested if TCR fragments, which have been shown to induce ER stress in cells lacking the corresponding chaperons for their folding, will be expressed in myeloid cells. Consistent with this notion, we could not detect expression of alpha chain from TRP1-reactive TCR (29) following transfection of RAW264.7 cells (**Fig. 4J** and Supplementary Fig. S3C). Lastly, we assessed whether scFv fragments are presented on MHC-I and MHC-II. To this end, we transfected Raw 265.7 cells with MHC-I peptide OVA₂₅₇₋₂₆₄ followed by MHC-II OVA₃₂₃₋₃₃₉ peptide alone or fused to the C-terminal of α CD19 scFv. After 16 hours, we incubated T cells from OT-I and OT-II mice and tested their rates of proliferation (Supplementary Fig. S3D). Yet, only negligible differences were observed in T cells incubated with RAW264 cells expressing the different constructs (Supplementary Fig. S3E).

FcγRI provides a scaffold for incorporating IgM-induced signaling in myeloid cells and endows them with tumor cell-specific killing ability

Taking our findings into consideration, we sought to adopt an alternative approach; instead of equipping monocytes with scFv segment, we fused the α chain of the high-affinity Fc IgG receptor (FcγRI) as extracellular region fused to signaling chain (**Fig. 5A**). These construct's recognition of the target cells was designed to be mediated by tumor-binding antibodies. First, we tested the expression levels of the naive and the chimeric FcγRI constructs in RAW264.7 cells. Flow cytometric analysis 24h post-transfection indicated that both constructs were successfully expressed at comparable levels (**Fig. 5B-5C**). Confocal microscopy further indicated that both constructs expressed on the cell membrane and Golgi. (**Fig. 5D**). Similar expression patterns were also observed in infected murine bone marrow cells (**Fig. 5E**).

Consequently, we sought to characterize the effect of tumor-binding antibodies on macrophage cytotoxic activity. To this end, we co-incubated tdTomato-labeled 4T1 tumor cell lines expressing human HER2⁺ with RAW264.7 cells that constitutively express the chimeric receptors with and without the anti-HER2⁺ antibody *trastuzumab*. Confocal microscopy indicated that RAW264.7 alone had no cytotoxic activity following a 24h incubation. In contrast, the addition of *trastuzumab* induced a polarized accumulation of GrB in the synapse between macrophages and tumor cells. Significantly higher levels of GrB in the immune synapse were counted in cells expressing C5aR chimeric receptor (**Fig. 5F-5G** and Supplementary Fig. S4A). To quantify the killing capacity of




Figure 5

FeyRI can provide a scaffold for incorporating IgM-induced signaling in myeloid cells and endows them with tumor cell-specific killing ability.

(A) Illustration of chimeric Fey receptor design.¹ (B-C) Representative FACS plots (B) and mean percentages (C) of RAW264.7 cells expressing chimeric Fcγ receptors 24 h after transfection (n=3). (D) Confocal microscopy images of RAW264.7 cells 24 hours post transfection with Fey receptors tagged with GFP and membrane tagged tdTomato. (E) Mean percentages of BMDc 72 h post lentivirus transduction with Fey receptors (n=4). (F) Confocal microscopy staining of GrB in RAW 264.7 cells co-cultured overnight with 4T1 cells expressing human HER2+. (G) Mean counts of GrB in the synapse between transduced RAW264.7 cells and the tumor cells (n=18). (H) IncuCyte analysis of human HER2+ 4T1 cells growth following incubation with transduced RAW 264.7 cells (n=6). (I) Super-resolution microscopy of GFP-tagged chimeric FcγR and mCherry-tagged gamma chain. (J) Confocal microscopy staining of GrB in RAW 264.7 cells co-cultured overnight with 4T1 cells expressing human HER2+. (K) IncuCyte analysis of human HER2+ 4T1 cells growth following incubation with transduced RAW 264.7 cells (n=6). (L) mean percentages of CD3+ out of CD45+ cells in B16F10 tumors from day 26. (n=4) Results are from one representative experiment out of at least three performed. Statistical significance was calculated using non-parametric t test (** denote p<0.01, **** denote p<0.0001).¹ Illustrations

this construct, we next transfected RAW264.7 macrophages with mRNA which encode these modified receptors, and incubated them with fluorescently labeled HER2⁺ expressing 4T1 tumor cells under IncuCyte. Consistent with GrB polarization, transfected macrophages alone were almost completely inert whereas the addition of *trastuzumab* promoted tumor cell killing (**Fig. 5H**). Importantly, while killing rates of macrophages expressing C5aR chimeric receptor were higher, they were also induced by free and irrelevant antibody (e.g., rituximab) (**Fig. 5H**).

To assess the underlying mechanism, we analyzed the transfected RAW264.7 cells under high-resolution microscopy. In steady state, cells transfected with the native FcγRI had a clear separation between the alpha chain (antibody binding chain) and their signaling gamma chain. The addition of free *trastuzumab* did not alter their membranal architecture. Incubation with plastic-immobilized *trastuzumab* resulted in polarization of the alpha chain toward the plastic and co-localization with the gamma chain. The architecture of C5aR chimeric receptor completely abrogated the membrane organization of the signaling chains, as the two co-localized even under steady state (**Fig. 5I** and Supplementary Fig. S4B). We then speculated that the fusion of such signaling chain to the alpha chain could mitigate the effects of tonic signaling by attaching IgM receptor signaling chain, instead of the gamma chain, to the transmembrane domain (illustrated in Supplementary Fig. 4C). Indeed, this architecture can be highly expressed in macrophage membrane and did not recruit or abrogate the gamma chain upon addition of antibodies, while maintaining alpha chain clustering to immobilized *trastuzumab* (Supplementary Fig. S4D-S4F). This construct promotes polarized expression of GrB at the immunological synapse and induces an antibody-specific tumor cell lysis (**Fig. 5J** and supplementary Fig S4G). Lastly, we tested whether myeloid cells expressing modified FcγRI can affect tumor growth *in vivo*. To this end, mice were challenged with B16F10 tumor cells and to grow for 8 days until they reached an average size of 20 mm². Next, mRNA transfect BMDC were injected s.c. once a week alone or in combination with i.p. injections of TA99 antibody against the melanoma antigen TRP1. Indeed, significant inhibition of tumor growth was observed in mice treated with modified FcγRI and TA99, but not with sham transfected BMDC or without TA99 antibodies (**Fig. 5K** and Supplementary Fig. 4H-4I). analysis of immune cell infiltration indicated higher rates of T cells in tumors treated with modified DC and TA99. (**Fig. 5L**) suggesting this treatment also potentiate the hosts immune response.

Discussion

The microenvironment of most solid tumors is often enriched with myeloid cells at different stages of maturity and polarization and the general notion suggests that they often adopt a suppressive phenotype that supports tumor progression and escape (4, 21, 30). The inherent plasticity of these cells, however, enables them to change their phenotype upon injection of adjuvant or immune-stimulatory agents (31–33). Countless experimental and clinical data have been tested for their capacity to convert tumor-infiltrating myeloid cells phenotype to an anti-tumor, including TLR agonists (34–37), immune complexes (25, 38, 39), immune checkpoints (40, 41) and co-stimulatory antibodies (42). While they showed improved T cell infiltration and overall lower tumor burden, they did not meet the expectations. It is noteworthy that most of these attempts are based on the notion that CD8⁺ T cells are the main effector arm against cancer and that the purpose of myeloid cell activation is to support T cell-mediated killing (32, 33). Nonetheless, myeloid cells are highly equipped with cell-killing mechanisms, including antibody-mediated cellular toxicity (ADCC) and secretion of reactive oxygen species (43). Here, we demonstrated that activating the IgM-receptor signaling in myeloid cells using tumor-binding IgG induces oxidative burst, Granzyme release, and massive tumor cell lysis.

While genetic engineering of immune cells to treat cancer has proven to be successful in lymphoid cells (16 [↗](#), 44 [↗](#)), cells of the myeloid lineage resist gene modification. Noteworthy, these cells are traditionally thought to be programmed to respond to foreign genetic materials. Early attempts to overcome this inherent barrier were achieved by Biglari et al. using adenoviral vectors. They demonstrated that the fusion of scFv against CEA to FcγR can be expressed in monocytes (45 [↗](#)). Although this construct leads to granzyme release in T cells, its function in monocytes remains obscure. Further attempts to equip macrophages with specific tumor antigen recognition were done by Morrissey et al. They fused scFv derived from αCD19 antibody to different signaling chains that facilitate phagocytosis (46 [↗](#)). Interestingly, most of these receptors do not appear to be expressed on the cell membrane. Moreover, CAR-merTK, which appears on the cell membrane, was functionally inert and did not induce phagocytosis.

Recently, Klichinsky et al. have also attempted to genetically modify macrophages using an adenoviral vector expressing αCD19 and αHER2 Chimeric Antigen Receptor fused to CD3ζ – CD28 signaling domains. They suggested that this strategy can be applied to kill HER2-expressing tumor cells through phagocytosis. While the authors demonstrated expression by flow cytometry, they did not attempt to identify the membranous localization in macrophages. It may also be that the use of Adeno-associated viruses (AAV) as transduction vectors enable overcoming the inflammatory cascade induced in myeloid cells as a result of liposome-based transfection or lentiviral transduction, thus enabling expression. With that being said, the authors concluded that the improved survival induced in mice was due to M1-like phenotype induced in transduced macrophages by the virus, rather than the CAR construct (22 [↗](#)).

Why scFv induces ER stress specifically in myeloid cells remains unclear. Activation of ER stress by misfolded proteins in macrophages plays a dominant role in many human pathologies, yet it also occurs in other cell types. One possible explanation is that these cells lack the machinery needed for successful folding of antibody-like structures. Along these lines, many non-immune cells such as HeLa or HEK293 cells initiate ER-stress response upon transfection of fragments derived from immune-associated receptors such as the alpha chain of the T cell receptor (47 [↗](#)).

Overall, while we identify FcγRI as a possible construct that could endow myeloid cells with tumor-specific recognition abilities, the main goal of the present work is to highlight the unique challenges and considerations that are involved in genetically reprogramming the signaling in myeloid cells. We demonstrate that myeloid cells cannot express scFv, most likely since they lack the corresponding chaperons required for its folding. Yet, more work is needed before this could be promoted as an efficient therapy for solid tumors.

References

1. Coussens LM, Zitvogel L, Palucka AK (2013) **Neutralizing tumor-promoting chronic inflammation: a magic bullet?** *Science* **339**
2. Fridman WH, Pages F, Sautes-Fridman C, Galon J (2012) **The immune contexture in human tumours: impact on clinical outcome** *Nat Rev Cancer* **12**:298–306
3. Thorsson V, Gibbs DL, Brown SD, Wolf D, Bortone DS, Ou Yang TH, et al. (2018) **The Immune Landscape of Cancer** *Immunity* **48**:812–30
4. Bruni D, Angell HK, Galon J (2020) **The immune contexture and Immunoscore in cancer prognosis and therapeutic efficacy** *Nature Reviews Cancer* **20**:662–80
5. Greten FR, Grivennikov SI. (2019) **Inflammation and Cancer: Triggers, Mechanisms, and Consequences** *Immunity* **51**:27–41
6. Hanahan D, Coussens LM (2012) **Accessories to the Crime: Functions of Cells Recruited to the Tumor Microenvironment** *Cancer Cell* **21**:309–22
7. Fritz JM, Lenardo MJ (2019) **Development of immune checkpoint therapy for cancer** *The Journal of experimental medicine* **216**:1244–54
8. Waldman AD, Fritz JM, Lenardo MJ (2020) **A guide to cancer immunotherapy: from T cell basic science to clinical practice** *Nat Rev Immunol* **20**:651–68
9. Hoffmann MM, Slansky JE (2020) **T-cell receptor affinity in the age of cancer immunotherapy** *Mol Carcinog* **59**:862–70
10. Reuben A, Zhang J, Chiou SH, Gittelman RM, Li J, Lee WC, et al. (2020) **Comprehensive T cell repertoire characterization of non-small cell lung cancer** *Nat Commun* **11**
11. Johnson LA, Morgan RA, Dudley ME, Cassard L, Yang JC, Hughes MS, et al. (2009) **Gene therapy with human and mouse T-cell receptors mediates cancer regression and targets normal tissues expressing cognate antigen** *Blood* **114**:535–46
12. Casucci M, Hawkins RE, Dotti G, Bondanza A (2015) **Overcoming the toxicity hurdles of genetically targeted T cells** *Cancer Immunol Immunother* **64**:123–30
13. Newick K, O'Brien S, Moon E, Albelda SM (2017) **CAR T Cell Therapy for Solid Tumors** *Annu Rev Med* **68**:139–52
14. Rafiq S, Hackett CS, Brentjens RJ (2020) **Engineering strategies to overcome the current roadblocks in CAR T cell therapy** *Nat Rev Clin Oncol* **17**:147–67
15. June CH, Maus MV, Plesa G, Johnson LA, Zhao Y, Levine BL, et al. (2014) **Engineered T cells for cancer therapy** *Cancer Immunol Immunother* **63**:969–75
16. Gross G, Eshhar Z (2016) **Therapeutic Potential of T Cell Chimeric Antigen Receptors (CARs) in Cancer Treatment: Counteracting Off-Tumor Toxicities for Safe CAR T Cell Therapy** *Annual review of pharmacology and toxicology* **56**:59–83

17. Joyce JA, Fearon DT (2015) **T cell exclusion, immune privilege, and the tumor microenvironment** *Science* **348**
18. Hanley CJ, Thomas GJ (2020) **T-cell tumour exclusion and immunotherapy resistance: a role for CAF targeting** *Br J Cancer* **123**:1353–5
19. Lim AR, Rathmell WK, Rathmell JC (2020) **The tumor microenvironment as a metabolic barrier to effector T cells and immunotherapy** *Elife* **9**
20. Grivennikov SI, Greten FR, Karin M (2010) **Immunity, inflammation, and cancer** *Cell* **140**:883–99
21. Mantovani A, Sica A (2010) **Macrophages, innate immunity and cancer: balance, tolerance, and diversity** *Current opinion in immunology* **22**:231–7
22. Klichinsky M, Ruella M, Shestova O, Lu XM, Best A, Zeeman M, et al. (2020) **Human chimeric antigen receptor macrophages for cancer immunotherapy** *Nat Biotechnol* **38**:947–53
23. Sloas C, Gill S, Klichinsky M (2021) **Engineered CAR-Macrophages as Adoptive Immunotherapies for Solid Tumors** *Front Immunol* **12**
24. Cafri G, Sharbi-Yunger A, Tzehoval E, Alteber Z, Gross T, Vadai E, et al. (2015) **mRNA-transfected Dendritic Cells Expressing Polypeptides That Link MHC-I Presentation to Constitutive TLR4 Activation Confer Tumor Immunity** *Mol Ther* **23**:1391–400
25. Carmi Y, Spitzer MH, Linde IL, Burt BM, Prestwood TR, Perlman N, et al. (2015) **Allogeneic IgG combined with dendritic cell stimuli induce antitumour T-cell immunity** *Nature* **521**
26. Carmi Y, Prestwood TR, Spitzer MH, Linde IL, Chabon J, Reticker-Flynn NE, et al. (2016) **Akt and SHP-1 are DC-intrinsic checkpoints for tumor immunity** *JCI Insight* **1**
27. Ma L, Dichwalkar T, Chang JYH, Cossette B, Garafola D, Zhang AQ, et al. (2019) **Enhanced CAR-T cell activity against solid tumors by vaccine boosting through the chimeric receptor** *Science* **365**
28. Mallery DL, McEwan WA, Bidgood SR, Towers GJ, Johnson CM, James LC (2010) **Antibodies mediate intracellular immunity through tripartite motif-containing 21 (TRIM21)** *Proc Natl Acad Sci U S A* **107**:19985–90
29. Dougan SK, Dougan M, Kim J, Turner JA, Ogata S, Cho HI, et al. (2013) **Transnuclear TRP1-specific CD8 T cells with high or low affinity TCRs show equivalent antitumor activity** *Cancer Immunol Res* **1**:99–111
30. Gajewski TF, Schreiber H, Fu YX (2013) **Innate and adaptive immune cells in the tumor microenvironment** *Nat Immunol* **14**:1014–22
31. Quatromoni JG, Eruslanov E (2012) **Tumor-associated macrophages: function, phenotype, and link to prognosis in human lung cancer** *Am J Transl Res* **4**:376–89
32. Ricketts TD, Prieto-Dominguez N, Gowda PS, Ubil E (2021) **Mechanisms of Macrophage Plasticity in the Tumor Environment: Manipulating Activation State to Improve Outcomes** *Front Immunol* **12**

33. Bercovici N, Guerin MV, Trautmann A, Donnadieu E (2019) **The Remarkable Plasticity of Macrophages: A Chance to Fight Cancer** *Front Immunol* **10**
34. Cheng H, Sun L, Shen D, Ren A, Ma F, Tai G, et al. (2020) **Beta-1,6 glucan converts tumor-associated macrophages into an M1-like phenotype** *Carbohydr Polym* **247**
35. Dudek AZ, Yunis C, Harrison LI, Kumar S, Hawkinson R, Cooley S, et al. (2007) **First in human phase I trial of 852A, a novel systemic toll-like receptor 7 agonist, to activate innate immune responses in patients with advanced cancer** *Clin Cancer Res* **13**:7119–25
36. Miura K, Ishioka M, Minami S, Horie Y, Ohshima S, Goto T, Ohnishi H (2016) **Toll-like Receptor 4 on Macrophage Promotes the Development of Steatohepatitis-related Hepatocellular Carcinoma in Mice** *J Biol Chem* **291**:11504–17
37. Rodell CB, Arlauckas SP, Cuccarese MF, Garriss CS, Li R, Ahmed MS, et al. (2018) **TLR7/8-agonist-loaded nanoparticles promote the polarization of tumour-associated macrophages to enhance cancer immunotherapy** *Nat Biomed Eng* **2**:578–88
38. Gao CH, Dong HL, Tai L, Gao XM (2018) **Lactoferrin-Containing Immunocomplexes Drive the Conversion of Human Macrophages from M2-into M1-like Phenotype** *Front Immunol* **9**
39. Ackerman SE, Pearson CI, Gregorio JD, Gonzalez JC, Kenkel JA, Hartmann FJ, et al. (2021) **Immune-stimulating antibody conjugates elicit robust myeloid activation and durable antitumor immunity** *Nat Cancer* **2**:18–33
40. Arlauckas SP, Garriss CS, Kohler RH, Kitaoka M, Cuccarese MF, Yang KS, et al. (2017) **In vivo imaging reveals a tumor-associated macrophage-mediated resistance pathway in anti-PD-1 therapy** *Sci Transl Med* **9**
41. Gordon SR, Maute RL, Dulken BW, Hutter G, George BM, McCracken MN, et al. (2017) **PD-1 expression by tumour-associated macrophages inhibits phagocytosis and tumour immunity** *Nature* **545**
42. Beatty GL, Chiorean EG, Fishman MP, Saboury B, Teitelbaum UR, Sun W, et al. (2011) **CD40 agonists alter tumor stroma and show efficacy against pancreatic carcinoma in mice and humans** *Science* **331**
43. Mantovani A, Marchesi F, Malesci A, Laghi L, Allavena P (2017) **Tumour-associated macrophages as treatment targets in oncology** *Nat Rev Clin Oncol* **14**:399–416
44. Kalos M, Levine BL, Porter DL, Katz S, Grupp SA, Bagg A, June CH (2011) **T cells with chimeric antigen receptors have potent antitumor effects and can establish memory in patients with advanced leukemia** *Science translational medicine* **3**:95–95
45. Biglari A, Southgate TD, Fairbairn LJ, Gilham DE (2006) **Human monocytes expressing a CEA-specific chimeric CD64 receptor specifically target CEA-expressing tumour cells in vitro and in vivo** *Gene Ther* **13**:602–10
46. Morrissey MA, Williamson AP, Steinbach AM, Roberts EW, Kern N, Headley MB, Vale RD (2018) **Chimeric antigen receptors that trigger phagocytosis** *Elife* **7**
47. Osowski CM, Urano F (2011) **Measuring ER stress and the unfolded protein response using mammalian tissue culture system** *Methods Enzymol* **490**:71–92

Article and author information

Leen Farhat-Younis

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Manho Na

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Amichai Zarfin

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Aseel Khateeb

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Nadine Santana-Magal

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Alon Richter

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Amit Gutwillig

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Diana Rasoulouniriana

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Annette Gleiberman

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Lir Beck

Department of Human Molecular Genetics and Biochemistry, Tel Aviv University, Israel

Tamar Giger

Department of Molecular Cell Biology, Weizmann Institute, Israel

Avraham Ashkenazi

Department of Cell and Developmental Biology, School of Medicine, Tel Aviv University, Israel

Adi Barzel

Department of Biochemistry Molecular Biology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Israel

Peleg Rider

Department of Pathology, School of Medicine, Tel Aviv University, Israel

Yaron Carmi

Department of Pathology, School of Medicine, Tel Aviv University, Israel

For correspondence: aron.carmi@gmail.com

ORCID iD: [0000-0002-0972-0616](https://orcid.org/0000-0002-0972-0616)

Copyright

© 2023, Farhat-Younis et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Neeha Zaidi

Johns Hopkins University, Baltimore, United States of America

Senior Editor

Tadatsugu Taniguchi

University of Tokyo, Tokyo, Japan

Reviewer #2 (Public Review):

Summary:

While a significant portion of immunotherapy research has focused on the pivotal role of T cells in tumor immunity, their effectiveness may be limited by the suppressive nature of the tumor environment. On the other hand, myeloid cells are commonly found within tumors and can withstand these adverse conditions. However, these cells often adopt an immunosuppressive phenotype when infiltrating tumors. Therefore, manipulating myeloid cells could potentially enhance the anti-tumor potential of immunotherapy. In this manuscript, Farhat-Younes and colleagues have demonstrated that activating the IgM receptor signaling in myeloid cells induces an oxygen burst, the secretion of Granzyme B, and the lysis of adjacent tumor cells. Furthermore, they have outlined a strategy to utilize these features to generate CAR macrophages. However, they have identified a limitation: the expression of scFv in myeloid cells induces ER stress and the degradation of misfolded proteins. To address this issue, chimeric receptors were designed based on the high-affinity FcγRI for IgG. When macrophages transfected with these receptors were exposed to tumor-binding IgG, extensive tumor cell killing, and the release of reactive oxygen species and Granzyme B were observed.

Strengths:

In general, I consider this work to be significant, and the results are compelling. It emphasizes the specific considerations and requirements for successful manipulation in myeloid cells, which could further advance the field of cellular engineering for the benefit of immunotherapy

Following the revision of the original manuscript, I can clearly state that my concerns have been addressed and the article has been improved.

<https://doi.org/10.7554/eLife.91999.2.sa0>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

Farhat-Younis and colleagues demonstrate tumor-specific IgM's capacity to induce tumor cell death in monocyte-derived dendritic cell cultures. They subsequently designed a chimeric receptor based on high-affinity FcγRI. However, the authors found that the transfection process was more efficient when either the variable light or heavy chain was transfected individually rather than the entire scFv. This scFv construct led to an endoplasmic reticulum (ER) stress response and scFv degradation. A considerable portion of the manuscript is dedicated to the negative scFv expression results. The authors pivoted to a modified FcγRI capable of transmitting IgM signals. This represents a tremendous amount of work in the development of this chimeric receptor, the critical experiment showing efficacy in vivo was not presented, and instead various in vitro assays are shown. Thus, this manuscript will markedly benefit from showing improved responses to tumors in vivo when macrophages express FcγRI-IgM.

We deeply thank the reviewer for his thoughtful comments and overall favorable review of our manuscript.

1. In a mouse tumor model, the authors demonstrated that monocyte-derived dendritic cells (MoDCs) treated with IgG immune complexes (ICs) were more effective at preventing tumor growth compared to those treated with IgM ICs (as shown in Figure 1B). In Figure 1C, their in vitro experiments revealed that IgM resulted in tumor cell death, as well as increased production of nitric oxide (NO) and granzyme B. How do the authors reconcile IgG IC-treated MoDCs performing better in preventing tumors in vivo than IgM IC-treated MoDCs, despite the in vitro results with IgM-ICs. The authors speculate that IgG IC-treated MoDCs might trigger T cell immunity but do not show T cell involvement.

We apologize for not making this point clearer. We have extensively studied this phenomenon and published two papers that detailed the underlying mechanism in two consecutive papers (PMID: 27812544, PMID: 25924063). Briefly, we showed that DC activated with IgM-IC DC undergo cell death concomitantly to their release of lytic granules and lysis of tumor cells. As a result, they do not migrate to the lymph nodes where they should induce reactive T cell clones. In contrast, DC activated with IgG-IC do not elicit in vitro cytotoxicity but rather process the IC to present its derived antigens of MHC-II. We addressed that issue in the revised version and cited the relevant paper to further clarify it.

(2) The authors report distinct functional consequences of MoDCs incubated with tumor-IgG complexes and tumor IgM complexes. Tumor growth was inhibited and T cell immunity induced with the former. The latter, however, elicited robust anti-tumor killing. What happens if MoDCs are incubated with both IgG and IgM complexes? If this combined treatment induces effective killing and T cell memory, would this impact the design of the chimeric receptor to include IgG responsiveness as well?

This is a very interesting point. As mentioned above, our previous publications strongly suggest that tumor binding IgG and IgM induce different processes in myeloid cells. Yet, since MoDC naturally express the high affinity receptors for IgG FcγRI, we speculate that treating tumor-bearing mice modified monocyte, alone or in combination with tumor-binding IgG, would shed some light into that. Indeed, such treatment elicit a strong T cell immunity in these mice and the data was added to Supplementary Data Figure S4J. With that being said, a complete analysis of this question is very complicated and extent beyond the scope of this work. We would like to emphasize that the purpose of this work is to highlight some of the challenges unique to genetic manipulation in myeloid cells and to suggest one alternative scaffold for integrating signaling in these cells. We do not argue that the specific solution

presented here is the most potent one and more work is required before promoting such treatment into the clinic. We have added a sentence to the Discussion section that stress that issue.

(3) In Figure 5H, the authors demonstrate the ability of the chimeric receptor construct to deplete tumor cells in vitro. The ms would improve if the authors could show the chimeric receptor construct results in tumor cell death and/or prevention in an in vivo model. Similarly, if combined stimulation with IgG and IgM complexes enhances tumor response, this should be incorporated into the therapeutic strategy.

This is a wonderful suggestion. To address that, we challenged C57Bl/6 mice with B16F10 melanoma and allowed them to grow until it reached a palpable size of approximately 25 mm². Concomitantly, we cultured bone marrow dendritic cells from syngeneic mice and transfected them with a linear mRNA of the alpha/mu construct. Tumor bearing mice were then treated with alpha/mu and sham transduced BMDC alone, or in combination with antibody against the melanoma antigen Trp1 (TA99). The results were added as Figure 5K and to Supplementary Figure S4h-S4I.

Reviewer #2 (Public Review):

Summary:

While a significant portion of immunotherapy research has focused on the pivotal role of T cells in tumor immunity, their effectiveness may be limited by the suppressive nature of the tumor environment. On the other hand, myeloid cells are commonly found within tumors and can withstand these adverse conditions. However, these cells often adopt an immunosuppressive phenotype when infiltrating tumors. Therefore, manipulating myeloid cells could potentially enhance the anti-tumor potential of immunotherapy.

In this manuscript, Farhat-Younes and colleagues have demonstrated that activating the IgM receptor signaling in myeloid cells induces an oxygen burst, the secretion of Granzyme B, and the lysis of adjacent tumor cells. Furthermore, they have outlined a strategy to utilize these features to generate CAR macrophages. However, they have identified a limitation: the expression of scFv in myeloid cells induces ER stress and the degradation of misfolded proteins. To address this issue, chimeric receptors were designed based on the high-affinity FcγRI for IgG. When macrophages transfected with these receptors were exposed to tumor-binding IgG, extensive tumor cell killing, and the release of reactive oxygen species and Granzyme B were observed.

Strengths:

In general, I consider this work to be significant, and the results are compelling. It emphasizes the specific considerations and requirements for successful manipulation in myeloid cells, which could further advance the field of cellular engineering for the benefit of immunotherapy

We thank the reviewer for his thoughtful comments and overall appreciation of our findings.

Weaknesses:

Nevertheless, there are several minor issues that should be addressed:

(1) TCR fragments are commonly used to induce ER stress in non-immune cells. Therefore, it would be interesting to investigate whether TCR fragments can be expressed in myeloid cells and if they induce ER stress. Addressing this issue would support the

notion that these cells lack the ER chaperones required for folding immunoglobulin variable chains.

This is a wonderful suggestion. To assess that possibility, we cloned the alpha chain of anti-Trp1 TCR and transfected RAW 264.7 macrophages. Importantly, we could not detect expression on this construct in macrophages, further supporting our findings with scFv in these cells. We added this result to Figure 4J and Supplementary Figure S3C.

(2) It would be valuable to determine whether, after the degradation of scFv fragments by myeloid cells, they are presented on MHC-I and MHC-II.

This is a very interesting point. To address that, we generated a genetic construct where we fused the anti-CD19 scFv to a polypeptide composed from the MHCI and the MHCII fragments of Ova Albumin. Next, DC 2.4 were transfected with this construct and measured their capacity to stimulate the proliferation of CD8+ T cells from OT-I and CD4+ from OT-II mice. DC transfected with this construct efficiently stimulated the proliferation of both T cells, suggesting that both Ova fragments are indeed presented on MHCI and MHCII. Nonetheless, DC transfected with polypeptide of MHCI and MHCII fragments of Ova Albumin only (with no scFv), were almost equally effective in stimulating OT-I and OT-II T cell proliferation. We added that result to Supplementary Figure S3D-S3E.

(3) Some methodological details, such as the vaccination protocol and high-resolution microscopy procedures, are missing from the text.

We thank the reviewer for pointing out these issues. We added the missing details to the revised version of the manuscript.